

***Interactive comment on “Shallow water carbonate platforms (Late Aptian, Southern Apennines) in the context of supraregional to global changes” by A. Raspini***

**Dr. Raspini**

raspini@igg.cnr.it

Received and published: 2 February 2012

I thank Prof. Föllmi for his comment to my manuscript. You may find the final version of my manuscript (attached as Supplement) revised following most of the suggestions of all the referees.

General remarks The overall stratigraphic frame provided by Di Lucia et al. (this volume; 2009) for the “Orbitolina level” of the Apennine carbonate platform is strikingly in contrast with the points that I report in my manuscript. In fact: 1) I basically ground my stratigraphic, geochemical and paleoecologic discussion on the solid assumption that the “Orbitolina level” of the Apennine carbonate platform is Late Aptian in age

(Gargasian; see Cherchi et al., 1978; Bravi & De Castro, 1995; but also Schroeder et al., 2010, among many others), confirmed by my own experience with Mesorbitolinid stratigraphy in Aptian sediments of the Central-Southern Apennines since the Nineties (Raspini, 1996, 1998, 2001; D'Argenio et al., 1999); 2) my paleoenvironmental and paleoecologic interpretations are rooted in a stratigraphic model which includes sea level oscillations and paleoclimatic controls on the evolving trophic levels as the main governors for the sedimentation of the "Orbitolina level". As consequence, the main features of the Gargasian hydrosphere-biosphere system, as deduced by local carbon isotopic curves coupled to secular fluctuations of Ca/Mg ratio and pCO<sub>2</sub> (among others), have been taken into consideration and discussed consistently.

Finally, I think that, in all aspects of life, the great (and legitimate) desire to achieve success and "make history" may lead to unexpected positive results. But, if one is in a hurry, and the desire is too great, results may become counterproductive. In my view, this is what happened to the paper of Di Lucia et al. (this volume).

Specific answers Following the referee's request, I have added the section 3.1 (Remarks on the age of the "Orbitolina level") to the final version of the manuscript (attached as Supplement), as reported below: "Recently, the "Orbitolina level" of the Southern Apennines has been attributed to the Bedoulian, based on the  $\delta^{13}\text{C}$  record of shallow carbonate sections, supported by a numerical age of 122.9 Ma (122.1-123.5) obtained from the <sup>87</sup>Sr/<sup>86</sup>Sr ratio of a sample collected 1 m above the first marls with Mesorbitolina (Di Lucia et al., this volume). According to these authors, the orbitolinid-rich marly level represents the base of the "Selli level" black shales of epicontinental and oceanic basins. The "Orbitolina level" of the Southern Apennines is rich in Mesorbitolina texana and subordinately Mesorbitolina parva (Cherchi et al., 1978). Although M. parva possibly appeared in the latest Bedoulian (Velić, 2007; Schroeder et al., 2010), the F.O. of M. texana occurred in the Late Aptian (Gargasian), as recently reported by Schroeder et al. (2010): it, therefore, remains one of the tie-points for the calibration of the Late Aptian biostratigraphy of central and southern Tethyan carbonate

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive  
Comment

platforms (Simmons, et al., 2000; Bachmann and Hirsch, 2006; Velić, 2007; Chihaoui et al., 2010; Embry et al., 2010; Heldt et al., 2010; Vincent et al., 2010). In this paper the “Orbitolina level” is considered a Gargasian litho and biostratigraphic marker in the carbonate platform successions of the Southern Apennines, in which it marks the first occurrence of *M. texana* and *M. parva*. This age is further supported by the following observations: 1 - In the lower part of the Tobenna-Faito composite section a progressive decrease of the accommodation space on the platform is recorded; it culminates with the Sequence Boundary Zone (SBZ1) above the “Orbitolina level”, as testified by the maximum abundance of characean-rich sediments (see section 5.2 and Fig. 4 below). According to D’Argenio et al. (1999), this ~8 m-thick deposits represents about 800 ka. An overall comparable trend of facies evolution was also observed in coeval sediments cropping out at Serra Sbragavitelli now located more than 100 km from the Faito section and encasing the orbitolinid-rich biostratigraphic marker (D’Argenio et al., 1999). If the “Orbitolina level” marked the onset of the OAE1a in the Apenninic platform, it would imply that: i) about two-third of the Selli-equivalent deposits at the Tobenna-Faito settled during the sea level lowstand, not during the rising of the sea level as reported in the literature (Schlanger & Jenkyns, 1976; Jenkyns, 1980; Haq et al., 1988; Bralower et al., 1994; Erbacher et al., 1996; Erba et al., 1999; Wissler et al., 2004; Emeis & Weissert, 2009; Jenkyns, 2010; Skelton & Gili, 2011, and many others); ii) considering the cyclostratigraphic studies carried out on the Aptian carbonate platform successions encasing the “Orbitolina level” (Raspini, 1998, 2001; D’Argenio et al., 1999) and the related C-isotopic record recently published (D’Argenio et al., 2004), at Tobenna-Faito the negative isotopic spike corresponding to the segment C3 of Menegatti et al. (1998) would be recorded in sediments deposited more than 400 ka from the beginning of the Selli event (bed-scale cycle 45 in Fig. 4), instead of marking its onset (cf. also Mehay et al., 2009; Erba et al., 2010). This would also occur in the Serra Sbragavitelli section (D’Argenio et al., 2004). 2 - According to Ferreri et al. (1997) and D’Argenio et al. (2004), carbon isotope curves extracted from several shallow water carbonate sections cropping out in the southern Apennines evidence

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

that the positive  $\delta^{13}\text{C}$  spike reflecting the Selli event clearly underlies the “Orbitolina level”. This level is placed close to the boundary of *G. algerianus*/*G. ferreolensis* zone, confirming a Gargasian age as proposed by Cherchi et al. (1978). 3 – The recent astronomical tuning of the Aptian Stage (Huang et al., 2010) constrains the Selli event between 124.55 and 123.16 Ma, while for Di Lucia et al. (this volume) the “Orbitolina level” (that should represent the base of the Selli event) is dated at 122.9 Ma.”

All of the above considerations claim a Gargasian age for the “Orbitolina level” of the southern Apennines (see also my reply to the comments by Di Lucia to my manuscript at <http://www.solid-earth-discuss.net/3/901/2011/sed-3-901-2011-discussion.html>). This is also supported by the following observations: 1) *Archaeoalveolina reicheli* cannot be considered an independent tie-point of Gargasian age as indicated by Di Lucia et al. (this volume). In fact, *A. reicheli* has been reported from the Gargasian of Algeria and north Tunisia (Bismuth, 1973; Fourcade & Raoult, 1973), from the Late Gargasian and Clansayesian of the eastern Algeria (Masse & Chiki-Aouimeur, 1982) or dated as Late Aptian/Early Albian in the Southern Apennines, Croatia and even in Central Tunisia (De Castro, 1991; Chiocchini et al., 1994; Bravi & De Castro, 1995; Husinec et al., 2000; 2009; Chihaoui et al., 2010). This conflicting age is likely the reason for Tethyan range charts of *A. reicheli* have not yet been established (see Heldt et al., 2010). 2) As stated in a recently discussed but not yet published Doctoral thesis “The “Orbitolina level” of the southern Apennines cannot be correlated with the “Couches supérieures à orbitolines” of the French Vercors and the “Upper Orbitolina Beds” of the Helvetic Alps (Linder et al., 2006; Föllmi and Gainon, 2008). Instead, a correlation with the “Niveau Fallot” of the Vocontian Basin (Friedrich et al., 2003), recently proposed by Raspini (2011), is compatible with our chemostratigraphic data” (Trecalli, 2011, p. 94-95). These statements overturn the interpretation and conclusions by Di Lucia et al. (this volume) on the orbitolinid-rich level of the Southern Apennines, lending explicit and definitive support to my manuscript. Consequently, the stratigraphic conclusions on the “Orbitolina level”, interpreted as the base of the basinal Selli OAE by Di Lucia et al. (this vol., but see also Di Lucia, 2009), have to be

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



considered unrealistic and therefore rejected.

I really do not know what the referee is referring when he writes “. . .the 4‰ positive shift was correlated with a minor, 1‰ shift near the base of the Late Aptian. . . in corresponding pelagic sections.”. The only 4‰ positive shift in the  $\delta^{13}\text{C}$  record reported by D’Argenio et al. (2004) and in my manuscript is above the “Orbitolina level” (Fig. 6). The correlation between this 4‰ positive shift (trend) and a minor (1‰ shift is with the Resolution Guyot (Cretaceous shallow-water carbonates; Jenkyns & Wilson, 1999). In the manuscript the correlation between the  $\sim 4\%$  positive shift of the Apenninic carbonate sections fits well with the corresponding positive shifts of about 4‰ in the (hemi)pelagic record represented by Piobbico-Cismon and Peregrina Canyon deposits (Fig. 6). In general, however, as the referee well knows, an offset of  $\delta^{13}\text{C}$  values such as the one between Apenninic and Pacific shallow water sections (or between deposits of shallow and deeper-water settings) could be related to different diagenetic conditions between compared sections (see also Vincent et al., 2010, their Fig. 9) and/or reflect the productivity- and depth-dependent uptake of  $^{12}\text{C}$  and  $^{13}\text{C}$  in carbonates (e.g., Bodin et al. 2009; Meyer et al. 2011). Finally, a climatic cooling can cause major oceanographic changes such as a destabilization in the oceanic density structure or ocean overturns that bring deep  $^{13}\text{C}$ -depleted water to the surface (Wilde & Berry, 1984; Glumac & Walker, 1998).

Regardless of diagenetic effects the oxygen isotope absolute values locally suffered, I think that the very good reproducibility of  $\delta^{18}\text{O}$  main trends of the studied sections at a regional scale (Fig. 5B) and the comparison with reference curves (Fig. 7) suggest that the main trends can be plausibly regarded as indicative of excursions of the isotopic composition of the sea water and, consequently, that the “Orbitolina level” deposited during a long-term global climate cooling, similarly to the Niveau Fallot in the hemipelagic setting (Takashima et al., 2007). This is in agreement with the sedimentological and cyclostratigraphical results summarized in Fig. 4 in which a progressive longer-term sea-level lowering is clearly recorded in the corresponding deposits (see

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

also Raspini, 1998).

In my manuscript I also utilised previously published isotopic data (D'Argenio et al., 2004) which have a low resolution; this is one of the reasons why the research presented here has to be considered a fundament for future, more conceptual work that I hope to realize (I am working to do it soon). Therefore, more detailed sampling of both the already studied and new coeval sections are required to reinforce the conclusions of the manuscript. Nevertheless, this does not mean that everything must be directly supported by geochemical evidences. Similarly to the more widely accepted interpretation (Vilas et al., 1995; Pittet et al., 2002; Bachmann & Hirsch, 2006; Burla et al., 2008; Embry et al., 2010; Schroeder et al., 2010 and many others), the exceptional concentration of conical foraminifera featuring the marly "Orbitolina level" of the southern Apennines can be well explained as mainly influenced by a variation of nutrient levels in the surficial waters (which were advantageous to the development of the orbitolinids) respect to the general paleoenvironmental conditions that favour the settlement of the under- and overlying carbonates.

Finally, I must say to be saddened and astonished by the "subliminal message" coming out from the referee's comment to my manuscript. In my view, a referee should evaluate a paper considering, for example, its organization, if the collected data are coherent with their interpretation, and so on, without suggesting, d'emblée, which manuscript he tends to favour. If different interpretations exist on the same topic, I think it is better make all of them known, giving the readers the chance to fully evaluate them and finally choose the version they prefer.

#### References cited

Bachmann M. & Hirsch. F. (2006) Lower Cretaceous carbonate platform of the eastern Levant (Galilee and the Golan Heights): stratigraphy and second-order sea-level change. *Cret. Res.*, 27, 487-512.

Bismuth H. (1973) Réflexions stratigraphiques sur l'Albo-Aptien dans la région des

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



djebels Douleb et Semmama et son environnement (Tunisie du Centre-Nord). Ann. Des Mines et de la Géologie, 6, Livre Jubilaire M. Solignac, 179-212.

Bodin S., Fiet N., Godet A., Matera V., Westermann S., Clément A., Janssen N.M.M., Stille P., Föllmi K.B. (2009) Early Cretaceous (late Berriasian to early Aptian) palaeoceanographic change along the northwestern Tethyan margin (Vocontian Trough, southeastern France):  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$  and Sr-isotope belemnite and whole-rock records. Cret. Res., 30, 1247-1262.

Bralower, T.J., Arthur, M.A., Leckie, R.M., Sliter, W.V., Allard, D., Schlanger, S.O., 1994. Timing and paleoceanography of oceanic dysoxia/anoxia in the Late Barremian to Early Aptian (Early Cretaceous). Palaios, 9, 335-369.

Bravi S. & De Castro P. (1995) The Cretaceous fossil fishes level of Capo d'Orlando, near Castellammare di Stabia (NA): biostratigraphy and depositional environment. Mem. Sci. Geol. 47, 45-72.

Burla S., Heimhofer U., Hochuli P.A., Weissert H. & Skelton P. (2008) Changes in sedimentary patterns of coastal and deep-sea successions from the North Atlantic (Portugal) linked to Early Cretaceous environmental change. Palaeogeogr. Palaeoclimatol. Palaeoecol., 257, 38-57.

Cherchi A., De Castro P. & Schroeder R. (1978) Sull'età dei livelli a Orbitolinidi della Campania e delle Murge baresi (Italia meridionale). Boll. Soc. Nat., Napoli, 87, 363-385.

Chihaoui A., Jaillard E., Latil J.-L., Zghal I., Susperregui A.-S., Tuir J. & Ouali, J. (2010) Stratigraphy of the Hameima and lower Fahdene Formations in the Tadjerouine area (Northern Tunisia). J. Af. Earth Sci., 58, 387-399.

Chiocchini M., Farinacci A., Mancinelli A., Molinari V. & Potetti M. (1994) Biostratigrafia a foraminiferi, dasycladali e calpionelle delle successioni carbonatiche mesozoiche dell'Appennino centrale (Italia). In: Mancinelli A. (ed.) Biostratigrafia dell'Italia

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Centrale. Studi Geol. Camerti, Spec. Vol. A, 9-129.

D'Argenio B., Ferreri V., Raspini A., Amodio S. & Buonocunto F.P. (1999) Cyclostratigraphy of a carbonate platform as a tool for high precision correlation. *Tectonophysics*, 315, 357-384.

De Castro P. (1991) Mesozoic. In: Barattolo F., De Castro P. & Parente M., 5th International Symposium on Fossil Algae. Field-trip guide-book, 21-38.

Di Lucia M. (2009) Il record dei cambiamenti globali nelle piattaforme carbonatiche del Cretacico medio dell'Appennino meridionale. Doctorate Thesis, Università di Napoli Federico II, 125 p.

Di Lucia M., Mutti M. & Parente M. (submitted) Bio-chemostratigraphy of the Barremian–Aptian shallow-water carbonates of the southern Apennines (Italy): pinpointing the OAE1a in a Tethyan carbonate platform. *Solid Earth Discuss.*, 3, 789–838, 2011 [www.solid-earth-discuss.net/3/789/2011/doi:10.5194/sed-3-789-2011](http://www.solid-earth-discuss.net/3/789/2011/doi:10.5194/sed-3-789-2011).

Embry J.C., Vennin E., Van Buchem F.S.P., Schroeder R., Pierre C. & Aurell M. (2010) Sequence stratigraphy and carbon isotope stratigraphy of an Aptian mixed carbonate-siliciclastic platform to basin transition (Galve sub-basin, NE Spain). In: Van Buchem, F.S.P., Gerdes, K.D., and Esteban, M. (eds) *Mesozoic and Cenozoic Carbonate Systems of the Mediterranean and the Middle East: Stratigraphic and Diagenetic Reference Models*. Geol. Soc., London, Engineering Geology Spec. Publ., 329, 113-143.

Emeis K.-C. & Weissert H. (2009) Tethyan–Mediterranean organic carbon-rich sediments from Mesozoic black shales to sapropels. *Sedimentology*, 56, 247-266.

Erba E, Bottini C, Weissert HJ, Keller CE (2010) Calcareous nannoplankton response to surface-water acidification around Oceanic Anoxic Event 1a. *Science*, 329, 428-432.

Erbacher J., Thurow J. & Littke R. (1996) Evolution patterns of radiolaria and organic matter variations: A new approach to identify sea-level changes in mid-Cretaceous pelagic environments. *Geology*, 24, 499-502.

SED

3, C650–C662, 2012

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper





Ferreri V., Weissert H., D'Argenio B. & Buonocunto F.P. (1997) Carbon isotope stratigraphy: a tool for basin to carbonate platform correlation. *Terra Nova*, 9, 57-61.

Föllmi K.B. & Gainon F. (2008) Demise of the northern Tethyan Urganian carbonate platform and subsequent transition towards pelagic conditions: the sedimentary record of the Col de la Plaine Morte area, central Switzerland. *Sed. Geol.*, 205, 142-159.

Fourcade E. & Raoult J.-F. (1973) Crétacé du Kef Hahouner et position stratigraphique de "Ovalveolina" reicheli De Castro (série septentrionale du mole néritique du Costantinois, Algérie). *Rev. Micropal.*, 15, 227-246.

Friedrich O., Reichelt K., Herrle J.O., Lehmann J., Pross J. & Hemleben C. (2009) Formation of the Late Aptian Niveau Fallot black shales in the Vocontian Basin (SE France): evidence from foraminifera, palynomorphs, and stable isotopes. *Mar. Micropaleontol.*, 49, 65-85.

Glumac B. & Walker K.R. (1998). A Late Cambrian positive carbon-isotope excursion in the southern Appalachians: Relation to biostratigraphy, sequence stratigraphy, environments of deposition, and diagenesis. *Jour. Sed. Res.*, 68, 1212-1222.

Haq B.U., Hardenbol J., Vail P.R. (1988) Mesozoic and Cenozoic chronostratigraphy and cycles of sea-level change. In: Wilgus C.K., Hastings B.S., Kendall C.G.St.C., Posamentier H.W., Ross C.A & Van Wagoner J.C. (eds) *Sea-level changes – An integrated approach*. *SEPM Spec. Publ.*, 42, 71-108.

Heldt M., Lehmann J., Bachmann M., Negra H. & Kuss J. (2010) Increased terrigenous influx but no drowning: palaeoenvironmental evolution of the Tunisian carbonate platform margin during the Late Aptian. *Sedimentology*, 57, 695-719.

Huang C., Hinnov L., Fischer A.G., Grippo A. & Herbert T. (2010) Astronomical tuning of the Aptian Stage from Italian reference sections. *Geology*, 38, 899-902.

Husinec A., Velić I., Fucek L., Vlahović I., Maticec, D., Ostrić N. & Korbar T. (2000) Mid Cretaceous orbitolinid (Foraminiferida) record from the islands of Cres and Losinj

(Croatia) and its regional stratigraphic correlation. *Cret. Res.*, 21, 155-171.

Husinec A., Velić, I. & Sokač B. (2009) Diversity patterns in Mid-Cretaceous benthic foraminifers and dasycladalean algae of the southern part of the Mesozoic Adriatic Platform, Croatia. In: Demchuk T.D. & Gary A.C. (eds) *Geologic Problem Solving with Microfossils: A Volume in Honor of Garry D. Jones*, SEPM, Spec. Publ. 93, 153-170.

Jenkyne H.C. (1980) Cretaceous anoxic events: from continents to oceans. *J. Geol. Soc.*, London, 137, 171-188.

Jenkyne H.C. (2010) Geochemistry of oceanic anoxic events. *Geochem. Geophys. Geosys.*, 11, doi:10.1029/2009GC002788.

Jenkyne H.C. & Wilson P.A. (1999) Stratigraphy, paleoceanography and evolution of Cretaceous Pacific guyots: relics from a greenhouse earth. *Am. J. Sci.*, 299, 341-392.

Linder P., Gigandet J., Hüsler J.L., Gainon, F. & Föllmi K.B. (2006) The early Aptian Grönten 20 Member: description of a new lithostratigraphic unit of the Helvetic Garschella Formation, *Ecl. Geol. Helv.*, 99, 327-341.

Masse J.P. & Chiki-Aouimeur F. (1982) La plate-forme carbonatée de l'Ouenza (Sud-Constantinois, Algérie). Organisation et dynamique durant l'Aptien supérieur. *Géologie Méditerranéenne*, 9, 259-267.

Méhay S., Keller C.E., Bernasconi S.M., Weissert H., Erba E., Bottini C. & Hochuli, P.A. (2009) A volcanic CO<sub>2</sub> pulse triggered the Cretaceous Oceanic Anoxic Event 1a and a biocalcification crisis. *Geology* 37, 819-822.

Menegatti A., Weissert H., Brown R.S., Tyson R.V., Farrimond P., Strasser A. & Caron M (1998) High-resolution  $\delta^{13}\text{C}$  stratigraphy through the early Aptian "Livello Selli" of the Alpine Tethys. *Paleoceanography* 13, 530-545.

Meyer K.M., Yu M., Jost A.B., Kelley B.M., Paine J.L. (2011)  $\delta^{13}\text{C}$  evidence that high primary productivity delayed recovery from end-Permian mass extinction. *Earth Plan*

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Sci Lett. doi: 10.1016/j.epsl.2010.12.033.

Pittet B., Van Buchem F.S.P., Hillgärtner H., Razin P., Grötsch J. & Droste H. (2002) Ecological succession, palaeoenvironmental change, and depositional sequences of Barremian-Aptian shallow-water carbonates in northern Oman. *Sedimentology*, 49, 555-581.

Raspini A. (1996) *Sedimentologia e ciclostratigrafia del Cretacico inferiore in facies di piattaforma carbonatica dell'Appennino centro-meridionale*. Doctorate Dissertation, Univ. Bologna.

Raspini A. (1998) Microfacies analysis of shallow water carbonates and evidence of hierarchically organized cycles. *Aptian of Monte Tobenna, Southern Apennines, Italy. Cretac. Res.*, 19, 197-223.

Raspini A. (2001) Stacking pattern of cyclic carbonate platform strata. Lower Cretaceous of southern Apennines, Italy. *J. Geol. Soc., Lond.*, 158, 353-366.

Schroeder R., Van Buchem F.S.P., Cherchi A., Baghbani D., Vincent B., Immenhauser A. & Granier B. (2010) Revised orbitolinid biostratigraphic zonation for the Barremian-Aptian of the eastern Arabian Plate and implications for regional stratigraphic correlations. *GeoArabia Spec. Publ.*, 4, 49-96.

Schlanger S.O., Jenkyns H.C. (1976) Oceanic anoxic events: causes and consequences. *Geol Mijnb* 55: 179-211.

Simmons M.D., Whittaker J.E., & Jones R.W. (2000) Orbitolinids from Cretaceous sediments of the Middle East – a revision of the F.R.S. Henson and Associates Collection. In: Hart M.B., Kaminski M.A. & Smart C.W. (eds), *Proceedings of the Fifth International Workshop on Agglutinated Foraminifera*, Grzybowski Foundation Spec. Publ., 7, 411-437.

Skelton P.W. & Gili E. (2011) Rudists and carbonate platforms in the Aptian: a case study on biotic interactions with ocean chemistry and climate. *Sedimentology*, doi:

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



10.1111/j.1365-3091.2011.01292.x.

Takashima R., Sano S.-I., Iba Y. & Nishi H. (2007) The first Pacific record of the Late Aptian warming event. *Jour. Geol. Soc.*, London, 164, 333-339.

Trecalli A. (2011) The record of the Early Toarcian and Early Aptian oceanic anoxic events in the Apenninic Carbonate Platform (Southern Italy). Unpublished Doctoral tesi, Università “Federico II” di Napoli.

Velić I.(2007): Stratigraphy and palaeobiogeography of Mesozoic benthic foraminifera of the Karst Dinarides (SE Europa), *Geol. Croatica*, 60, 1-113.

Vilas L., Masse J.P. & Arias C. (1995) Orbitolina episodes in carbonate platform evolution: the early Aptian model from SE Spain. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 119, 35-45.

Vincent B., van Buchem F.S.P., Bulot L.G., Immenhauser A., Caron M., Baghbani D. & Huc, A.Y. (2010) Carbon-isotope stratigraphy, biostratigraphy and organic matter distribution in the Aptian - Lower Albian successions of southwest Iran (Dariyan and Kazhdumi formations). *GeoArabia, Spec. Pub.*, 4, 139-197.

Wilde P. & Berry W.B.N. (1984) Destabilization of the oceanic density structure and its significance to marine “extinction” events. *Palaeogeog. Palaeoclimat. Palaeoecol.*, 48, 143-162.

Wissler L., Weissert H., Buonocunto F.P., Ferreri V. & D’Argenio B (2004) Calibration of the Early Cretaceous Time Scale: a combined chemostratigraphic and cyclostratigraphic approach to the Barremian–Aptian interval, Campania Apennines and Southern Alps (Italy). In: D’Argenio B. & Fischer A.G., Premoli Silva I., Weissert H. & Ferreri V. (eds) *Cyclostratigraphy; Approaches and cases histories*. *SEPM Spec. Publ.*, 81, 123-133.

Please also note the supplement to this comment:

C661

SED

3, C650–C662, 2012

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



<http://www.solid-earth-discuss.net/3/C650/2012/sed-3-C650-2012-supplement.pdf>

---

Interactive comment on Solid Earth Discuss., 3, 901, 2011.

**SED**

3, C650–C662, 2012

---

Interactive  
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C662

