

This discussion paper is/has been under review for the journal Solid Earth (SE).
Please refer to the corresponding final paper in SE if available.

Grassland fire effect on soil organic carbon reservoirs in semiarid environment

A. Novara¹, L. Gristina¹, J. Rühl¹, S. Pasta¹, G. D'Angelo¹, T. La Mantia¹, and P. Pereira²

¹Dipartimento di Scienze Agrarie e Forestali, University of Palermo, 90128 Palermo, Italy

²Department of Environmental Policy, Mykolas Romeris University, Ateities g. 20, 08303 Vilnius, Lithuania

Received: 18 April 2013 – Accepted: 25 April 2013 – Published: 3 July 2013

Correspondence to: A. Novara (agata.novara@unipa.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The aim of this work was to investigate the effect of a experimental fire, used for grassland management, on soil organic carbon (SOC) reservoirs. The study was carried out on *Hyparrhenia hirta* (L.) Stapf (*Hh*) grassland and *Ampelodesmos mauritanicus* (Desf.) T. Durand and Schinz (*Am*) grasslands, located in the north of Sicily. Soil samples were collected at 0–5 cm before and after experimental fire and SOC was measured. During grassland fire soil surface temperature was monitored. Biomass of both grasses was analyzed in order to determine dry weight and its chemical composition. The results showed that SOC varied significantly with vegetation cover, while it is not affected in the short period by grassland fire. *Am* grassland stored more SOC compared with *Hh* grassland thanks to lower content in biomass of labile carbon pool. No significant difference was observed in SOC before and after fire which could be caused by several factors: first, in both grassland types the measured soil temperature during fire was low due to thin litter layers; second, in semiarid environment higher mineralization rate results in lower soil carbon labile pool; and third, the C stored in the finest soil fractions, physical protected, is not affected by fire.

1 Introduction

Human societies are using the fire for millenia, fire is part of the Earth System and was tool for many societies (Pyne, 2001). Fire is regarded as an active ecological agent able to mobilize nutrients and restore soil fertility (Snyman, 2003) but, also, as a primary cause of soil degradation due to nutrients loss for volatilization, leaching and erosion, especially in severe wildfires. It is, in fact, considered a major disturbance in many ecosystem, which lead to important shifts in the soil properties and vegetation (Certini, 2005; Granged et al., 2011). One of the most common effect of fire is the alteration in composition and amount of soil organic matter (Knicker, 2007, Terefe et al., 2008). Several studies recorded a decrease (Fernàndez et al., 1997; Novara et al.,

SED

5, 883–895, 2013

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2011) in soil organic carbon (SOC) after fire, while results of other studies showed no significantly changing or even increase of previous SOC content (Kavdir et al., 2005). These discrepancies occur due to the large amount of controlling factors and therefore the effect of fire is highly variable in space. Among these factors, fire intensity, fire severity, fire regimen, type of burned vegetation, connectivity, distribution of fuel on soil surface, type of ash produced and dispersion, topography, soil properties, aspect, regional climate and meteorological conditions in the immediate period after the fire play a key role to determine SOC alteration and accumulation in soils (Certini, 2005; Pereira et al., 2010; Pereira et al., 2013).

10 In semiarid areas fire is one of the common management tools used by shepherd to enhance pasture regrowth. In fact, the recovery of vegetation canopy after fire in the Mediterranean area can be quite rapid due to adaption of plant communities to the disturbances caused by fire as observed in several studies (Trabaud, 1981; Barberis et al., 2003; Pausas and Verdù 2005). It is known, moreover, that fire is considered an important factor for arid and semiarid grasslands because it avoids invasion of trees and shrubs with implications on soil carbon storage (Briggs et al., 2005). Despite the importance of fire on grassland ecosystems (Bond et al., 2005), its impact on SOC is not well understood in the immediate period after the fire in the Mediterranean grasslands (Snyman, 2003). The aim of this work is to quantify SOC stock change as a result of a experimental fire of two of the most widespread types of Mediterranean grasslands (Brullo et al., 2010; Díez-Garretas and Asensi, 1999) and, therefore, to establish if this practice could be used sustainably as a management tool for grazing recovery.

2 Materials and methods

25 The field studies were carried out in the province of Palermo, Sicily (Italy) (350 m a.s.l.) (Fig. 1). Local soil type is an *Eutric Cambisol* according to WRB (WRB, 2006) with sand and clay contents of 18 % and 46 %, respectively. The climate is Mediterranean, with mean annual rainfall of 580 mm and yearly average temperature of 16 °C.

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



An experimental fire was conducted on July and September 2009 on five (replicas) delimited square areas (50 × 50 cm) in two different grassland types, dominated by *Hyparrhenia hirta* (L.) Stapf (*Hh*) grassland and *Ampelodesmos mauritanicus* (Desf.) T. Durand and Schinz (*Am*) grassland, respectively. Each sampling square was about 2 m distant from the neighbor square. In order to simulate a natural wildfire, burning was allowed to take its natural course until it extinguished itself. The fire was generated with a match, starting from leeward in each plot. Soil surface temperature during the burning was measured using a thermocouple system. In each selected area three soil samples were collected at 0–5 cm depth before and immediately after fire. SOC content was measured using a CHN-Elemental Analyzer.

For the $\delta^{13}\text{C}$ analysis, an EA-IRMS (elemental analyzer isotope ratio mass spectrometry) was used. The International Atomic Energy Agency (IAEA), Vienna, distribute IAEA-CH-6 as a reference standard material. The results of the isotope analysis are expressed as a δ value (‰) relative to the international Pee Dee Belemnite standard as follows:

$$\delta(\text{‰}) = \frac{R_s - R_{st}}{R_{st}} \times 1000 \quad (1)$$

where $\delta = \delta^{13}\text{C}$, $R = 13\text{C}/12\text{C}$, s = sample, and st = standard.

Dry biomass weight and its chemical composition (ADF acid detergent fiber, NDF neutral detergent fiber, Cellulose, Hemicellulose, Lignin, Ash) were determined on three 0.5 m² square area subsamples for each grassland types.

Data analysis was conducted using the SAS statistical package (SAS Inst., 2002). Normal distribution of data was verified and analysis of variance (ANOVA) was conducted. Significant differences were considered at a $p < 0.05$.

3 Results and discussion

SOC ranged from 20.3 to 37.0 g kg⁻¹ and from 15.4 to 32.5 g kg⁻¹ before and after experimental fire, respectively, in soil covered by *Hh*, and from 32.5 to 38.2 g kg⁻¹ and from 38.3 to 49.1 g kg⁻¹ before and after experimental fire, respectively, in soil covered by *Am*. The experimental fire did not had significant differences in SOC in both grassland types (Fig. 2). Similarly to SOC results, $\delta^{13}\text{C}$ was not affected significantly by fire. The average by time of $\delta^{13}\text{C}$ values measured in *Hh* grassland were $-25.418 \pm 0.25 \text{‰}$ and $-25.161 \pm 0.40 \text{‰}$ in soil sampled before and after fire, respectively; while in *Am* grassland were $-26.873 \pm 0.16 \text{‰}$ and $-26.98 \pm 0.31 \text{‰}$ before and after fire, respectively. Our results are in agreement with similar observations reported by other authors (Granged et al., 2011b) who found no change in SOC content before and after prescribed fire. In our experiment we reproduced the same environmental conditions of a wildfire. We can consider our experimental fire as a moderate fire severity, similar to prescribed fire described by before mentioned authors. In our experiment the time of combustion was 13 ± 2 min and 7 ± 1 min for *Hh* and *Am*, respectively (Fig. 3). The maximum temperature measured at soil surface was around 480 °C in both grassland. Temperatures over 200 °C persisted for 5 min and 3 min for *Hh* and *Am*, respectively. The burning time and intensity was lowest due to low amount of fuel in both grasslands. Mediterranean environmental conditions involve high organic matter mineralization rates and, thus, negligible amounts of litter biomass stock. The lower temperature arose during a low severity fires do not entail dramatic effects on SOC stock (Übeda et al., 2005). The loss of organic carbon by burning can occur even at relatively low temperatures such as 200 °C, but total combustion is only observed at high temperatures 450–500 °C, (De Bano et al., 1998). When comparing the two grasslands, SOC amount and the effect of fire on SOC stock was different. The lower SOC content was measured under *Hh* grassland, which also recorded the lower biomass yield. The above ground biomass estimated is 4.76 Mg ha⁻¹ and 11.60 Mg ha⁻¹ of dry matter for *Hh* and *Am* grassland, respectively.

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Even if the SOC change before and after fire was not statistically significant, after fire SOC content decreased of 11.5 % in *Hh* and increased of 27.9 % in *Am* grassland. The increase of SOC after fire could occur due to external inputs of charred material and ash, as commonly is observed in low severity fires. In particular, the burned material returns to soil as particles smaller than 2 mm in the form of ash, which are mixed in the top horizon, and which cause a net increase of SOC content (Gonzalez-Perez et al., 2004). The reason for the slight SOC increase after fire only in *Am* grassland may depend on different characteristics of the two considered grasses. Firstly, *Am* biomass contains more lignin and cellulose than *Hh* biomass (Table 1), and, thus, more recalcitrant compounds that under low temperature do not completely volatilize. Secondly, *Am* has a densely caespitose habit: this feature impedes a complete burning and favors the retention of not completely burnt plant residues. The ash of *Hh* is, instead, more light and quickly blown away by wind. Thirdly, biomass of *Am* contains siliceous compounds that obstruct burning.

4 Conclusions

Data here reported confirm that the use of experimental fire to favour plant recovery in *Hh* and *Am* grassland does not affect SOC stock. Our findings showed that in the studied grasslands the fire did not cause significant differences in SOC content, even if, these grasslands did not burn from many years. Our study shows that it is possible to adopt the system of controlled burning to maintain grassland formations, however, this management tool must be adopted only after thorough phytosociological analyses of local vegetation patterns and dynamics and after detailed planning.

Acknowledgements. Work was financially supported by the Italian government through the PRIN project “The impacts of secondary succession processes on carbon storage in soil and biomass and on biodiversity and the role of dispersal centers and vectors for recolonization processes”.

References

- Barberis, A., Dettori, S., and Filigheddu, M. R.: Management problems in Mediterranean cork-oak forests: Post-fire recovery, *J. Arid Environ.*, 54, 565–569, 2003.
- Bond, W. J., Woodward, F. I., and Midgley, G. F.: The global distribution of ecosystems in a world without fire, *New phytol.*, 165, 525–538, 2005.
- Braun-Blanquet J.: *Plant sociology*, Mc Graw-Hill, New York-London, 1932.
- Briggs, J. M., Knapp, A. K., Blair, J. M., Heisler, J. L., Hoch, G. A., Lett, M., S., and McCarron, J., K.: An ecosystem in transition: Causes and consequences of the conversion of mesic grassland to shrubland, *BioScience*, 55, 243–254, 2005.
- Brullo, C., Brullo, S., Giusso del Galdo, G., Guarino, R., Minissale, P., Scuderi, L., Siracusa, G., Sciandrello, S., and Spampinato, G.: The *Lygeo-Stipetea* class in Sicily, *Annali di Botanica*, 1-28, ISSN:0365-0812, 2010.
- Certini, G.: Effects of fire on properties of forest soils: a review, *Oecologia* 143, 1–10, 2005.
- DeBano, L. F., Neary, D. G., and Ffolliott, P. F.: *Fire's Effects on Ecosystems*, Wiley, New York, 1998.
- Díez-Garretas B. and Asensi A.: Syntaxonomic analysis of the *Andropogon*-rich grasslands (*Hyparrhenetalia hirtae*) in the Western Mediterranean Region, *Folia Geobotanica et Phytotaxonomica*, 34, 307–320, 1999.
- Fernández, I., Cabaneiro, A., and Carballas, T.: Organic matter changes immediately after a wildfire in an Atlantic forest soil and comparison with laboratory soil heating. *Soil Biol. Biochem.* 29, 1–11, 1997.
- González-Pérez, J. A., González-Vila, F. J., Almendros, G., and Knicker, H.: The effect of fire on soil organic matter, a review, *Environment International* 30, 855–870, 2004.
- Granged, A. J. P., Jordán, A., Zavala, L. M., Muñoz-Rojas, M., and Mataix-Solera, J.: Short-term effects of experimental fire for a soil under eucalyptus forest (SE Australia), *Geoderma*, 167–168, 125–134, 2011a.
- Granged, A. J. P., Zavala, L. M., Jordán, A., and Barcenas-Moreno, G.: Post-fire evolution of soil properties and vegetation cover in a Mediterranean heathland after experimental burning: A 3-year study, *Geoderma*, 164, 85–94, 2011b.
- Gristina, L., La Mantia, T., Pasta, S., and Poma I.: An analysis of carbon storage in protected areas with relation to agroforestry management and biodiversity, in: *The Soils of Tomorrow:*

SED

5, 883–895, 2013

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soils Changing in a Changing World, edited by: Dazzi, C. and Costantini, E., *Advances in GeoEcology*, Vol. 39, Catena Verlag, ISBN 978-3-923381-56-2: 540-554, 2008.

Kavdir, Y., Ekinci, H., Yüksel, O., and Mermut, A. R.: Soil aggregate stability and ¹³C CP/MAS-NMR assessment of organic matter in soils influenced by forest wildfires in Canakkale, Turkey. *Geoderma* 129, 219–229, 2005.

Knicker, H.: How does fire affect the nature and stability of soil organic nitrogen and carbon? A review, *Biogeochemistry*, 85, 91–118, 2007.

La Mantia T., Carimi F., Di Lorenzo R., and Pasta S.: The agricultural heritage of Lampedusa (Pelagie Archipelago, South Italy) and its key role for cultivar and wildlife conservation, *Italian Journal of Agronomy*, Vol. 6, 17, 106–110, 2011.

Novara, A., Gristina, L., Bodi, B. M., and Cerdà, A.: The impact of fire on redistribution of soil organic matter on a Mediterranean hillslope under maquia vegetation type, *Land degrade, development*, doi:10.1002/ldr.1027, 2011.

Pausas, J. G. and Verdu, M.: Plant persistence traits in fire-prone ecosystems of Mediterranean basin: a phylogenetic approach, *Oikos*, 109, 196–202, 2005.

Pereira, P., Cerdà, A., Ubeda, X., Mataix-Solera, J., Arcenegui, V., and Zavala, L.: Modelling the impacts of wildfire on ash thickness in a short-term period, *Land degrade, development*, doi:10.1002/ldr.2195, 2013.

Pereira, P., Bodi, M., Ubeda, X., Cerdà, A., Mataix-Solera, J., Balfour, V., and Woods, S.: Las cenizas y el ecosistema suelo, in: *Actualización en métodos y técnicas para el estudio de los suelos afectados por incendios forestales*, edited by: Cerdà, A., and Jordan, A., 345–398, Càtedra de Divulgació de la Ciència, Universitat de Valencia, 2010.

Pyne, S. J.: *Fire: A Brief History*, Seattle, WA: University of Washington Press, 2001.

SAS Institute: *The SAS System for Microsoft Windows*, Release 8.2. SAS Inst.: Cary, NC2002.

Snyman, HA.: Short-term response of rangeland following an unplanned fire in terms of soil characteristics in a semiarid climate of South Africa. *J. Arid Environ.* 55, 160-180, 2003.

Terefe, T., Mariscal-Sancho, I., Peregrina, F., and Espejo, R.: Influence of heating on various properties of six Mediterranean soils, A laboratory study, *Geoderma*, 143, 273–280, 2008.

Trabaud, L.: *Man and fire: impacts on Mediterranean vegetation*, edited by: di Castri, F., Goodallm, D. W., and Specht, R. L., *Mediterranean-type shrublands*, Elsevier, Amsterdam, 523–537, 1981.

Ùbeda, X., Lorca, M., Outeiro, L. R., Bernia, S., and Castellnou, M.: Effect of prescribed fire on soil quality in Mediteanean grassland (prades Mountains, north-east Spain), *International Journal of Wildland Fire*, 14, 379–384, 2005.

5 WRB: World Reference base for soil resources 2006, 2 Edn., World Soil Resources Reports, No 103, Food and Agriculture Organization of the United Nation, Rome, 2006.

SED

5, 883–895, 2013

Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Grassland fire effect

A. Novara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 1.** Biomass composition (% of dry biomass) of *Am* and *Hh*. Values in parenthesis are standard deviations. Abbreviations: ADL = acid detergent lignin, NDF = neutral detergent fiber.

Grassland	ADL	Cellulose	NDF	Hemicellulose	Ash	Aboveground biomass (Mg ha ⁻¹)	C Biomass (g kg ⁻¹)
<i>Am</i>	6.91 (0.58)	37.72 (1.58)	73.03 (2.65)	23.99 (1.32)	4.02 (1.10)	4.76	43.8
<i>Hh</i>	5.98 (0.68)	34.00 (1.20)	72.01 (1.53)	28.26 (1.76)	4.34 (1.49)	11.60	45.8




Fig. 1. Localization of the sampling area.

Grassland fire effect

A. Novara et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Fig. 2. Soil organic carbon before and after fire in *Hh* and *Am* grassland.

SED

5, 883–895, 2013

Grassland fire effect

A. Novara et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Grassland fire effect

A. Novara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Soil temperature during fire under *Hh* (blu line) and *Am* (red line) grassland.