

1 **SHORT COMMUNICATION**

2 **Grassland fire effect on soil organic carbon reservoirs in a semiarid environment**

3 Agata Novara⁽¹⁾, Luciano Gristina⁽¹⁾, Juliane Rühl⁽¹⁾, Salvatore Pasta⁽¹⁾, Giuseppe D'Angelo⁽¹⁾
4 Tommaso La Mantia⁽¹⁾, Paulo Pereira⁽²⁾

5 1 Dipartimento di Scienze agrarie e forestali – University of Palermo - 90128 Palermo – Italy

6 2 Environmental Research Centre, Mykolas Romeris University, Ateities g. 20, LT-08303 Vilnius,
7 Lithuania (paulo@mruni.eu).

8 Corresponding author: Agata Novara agata.novara@unipa.it

9

10 **Abstract**

11 The aim of this work was to investigate the effect of an experimental fire, used for grassland
12 management, on soil organic carbon (SOC) reservoirs. The study was carried out on *Hyparrhenia*
13 *hirta* (L.) Stapf (*Hh*) grassland and *Ampelodesmos mauritanicus* (Desf.) T. Durand & Schinz (*Am*)
14 grasslands, located in the north of Sicily. Soil samples were collected at 0-5 cm before and after
15 experimental fire and SOC was measured. During grassland fire soil surface temperature was
16 monitored. Biomass of both grasses was analyzed in order to determine dry weight and its
17 chemical composition. The results showed that SOC varied significantly with vegetation type, while
18 it is not affected in the short period by grassland fire. *Am* grassland stored more SOC compared
19 with *Hh* grassland thanks to lower content in biomass of labile carbon pool. No significant
20 difference was observed in SOC before and after fire which could be caused by several factors:
21 first, in both grassland types the measured soil temperature during fire was low due to thin litter
22 layers; second, in semiarid environment higher mineralization rate results in lower soil carbon
23 labile pool; and third, the SOC stored in the finest soil fractions, physically protected, is not
24 affected by fire.

25 Keywords: SOC, experimental fire, grassland, Mediterranean environment

26

27 **Introduction**

28 Fire is part of the Earth System and has been for millennia a tool for many societies (Pyne, 2001).
29 Fire is regarded as an active ecological agent able to mobilize nutrients and restore soil fertility
30 (Snyman, 2003) but, also as a primary cause of soil degradation due to nutrients loss for
31 volatilization, leaching and erosion, especially in severe wildfire. It is, in fact, considered a major
32 disturbance in many ecosystems, which lead to important shifts in the soil properties and
33 vegetation (Certini, 2005; Granged et al., 2011a). One of the most common effects of fire is the
34 alteration in composition and amount of soil organic matter (Knicker, 2007, Terefe et al., 2008).
35 Several studies recorded a decrease (Fernández et al., 1997; Novara et al., 2011) in soil organic
36 carbon (SOC) after fire, while results of other studies showed no significantly changing or even
37 increase of previous SOC content (Kavdir et al., 2005). These discrepancies occur due to the large
38 amount of controlling factors and therefore the effect of fire is highly variable in space and time.
39 Among these factors, fire intensity, fire severity, fire regimen, type of burned vegetation,
40 connectivity, distribution of fuel on soil surface, type of ash produced and dispersion, topography,
41 soil properties, aspect, regional climate and meteorological conditions in the immediate period
42 after the fire play a key role to determine SOC alteration and accumulation in soils (Certini, 2005;
43 Pereira et al., 2010; Pereira et al., 2013).

44 In semiarid areas fire is one of the common management tools used by shepherds to enhance
45 pasture regrowth. In fact, the recovery of vegetation canopy after fire in the Mediterranean area
46 can be quite rapid due to adaption of plant communities to the disturbances caused by fire as
47 observed in several studies (Trabaud, 1981; Barberis et al., 2003; Pausas and Verdù 2005). It is
48 known, moreover, that fire is considered an important factor for arid and semiarid grasslands
49 because it avoids invasion of trees and shrubs with implications on soil carbon storage (Briggs et
50 al., 2005). Despite the importance of fire on grassland ecosystems (Bond et al., 2005), its impact
51 on SOC is not well understood in the immediate period after the fire in the Mediterranean
52 grasslands (Snyman, 2003). The aim of this work is to quantify SOC stock change as a result of an

53 experimental fire of two of the most widespread types of Mediterranean grasslands (Brullo et al.,
54 2010; Díez-Garretas and Asensi, 1999) and, therefore, to establish if this practice could be used
55 sustainably as a management tool for grazing recovery.

56 **Materials and Methods**

57 The field studies were carried out in the province of Palermo, Sicily (Italy) (350 m a.s.l.) (Fig. 1).
58 Local soil type is an *Eutric Cambisol* according to WRB (WRB, 2006) with sand and clay contents of
59 18% and 46%, respectively. The climate is Mediterranean, with mean annual rainfall of 580 mm
60 and yearly average temperature of 16 °C.

61 An experimental fire was conducted on July and September 2009 on five (replicas) delimited
62 square areas (50×50 cm) in two different grassland types, dominated by *Hyparrhenia hirta* (L.)
63 Stapf (Hh) and *Ampelodesmos mauritanicus* (Desf.) T. Durand & Schinz (Am). Each sampling
64 square was about 2m distant from the neighbor square. In order to simulate a natural wildfire,
65 burning was allowed to take its natural course until it extinguished itself. The fire was generated
66 with a match, starting from leeward in each plot. Soil surface temperature during the burning was
67 measured using a thermocouple system (type K Inconel 600 insulated). In each selected area
68 three soil samples were collected at 0-5 cm depth before and immediately after fire. On three one
69 meter square in both grasslands (dominated by *Hyparrhenia hirta* or *Ampelodesmos mauritanicus*)
70 all plants were cut, oven dried for 3-4 days at 60-65 °C, and weighted. SOC content was measured
71 using a CHN-Elemental Analyzer. For the $\delta^{13}\text{C}$ analysis, an EA-IRMS (elemental analyzer isotope
72 ratio mass spectrometry) was used. The International Atomic Energy Agency (IAEA), Vienna,
73 distribute IAEA-CH-6 as a reference standard material. The results of the isotope analysis are
74 expressed as a δ value (‰) relative to the international Pee Dee Belemnite standard as follows:

$$\delta(\text{‰}) = \frac{R_s - R_{st}}{R_{st}} * 1000$$

77 where $\delta = \delta^{13}\text{C}$, R = $^{13}\text{C}/^{12}\text{C}$, s = sample, and st = standard.

78 Dry biomass weight and its chemical composition (ADF acid detergent fiber, NDF neutral detergent
79 fiber, Cellulose, Hemicellulose, Lignin, Ash) were determined on three 0.5 m² square area
80 subsamples for each grassland types.

81 Data analysis was conducted using the SAS statistical package (SAS Inst., 2002). Normal
82 distribution of data was verified previously to statistical data comparisons and analysis of variance
83 (ANOVA) was conducted. Significant differences were considered at a $p < 0.05$.

84

85 **Results and discussion**

86


87 SOC ranged from 20.3 to 37.0g kg⁻¹ and from 15.4 to 32.5 g kg⁻¹ before and after experimental
88 fire, respectively, in soil covered by *Hh*, and from 32.5 to 38.2 g kg⁻¹ and from 38.3 to 49.1 g kg⁻¹
89 before and after experimental fire, respectively, in soil covered by *Am*. The experimental fire did
90 not have significant differences in SOC in both grassland types (Fig. 2). Similarly to SOC results, δ
91 ¹³C was not affected significantly by fire. The average by time of δ ¹³C values measured in *Hh*
92 grassland were $-25.418 \pm 0.25\text{‰}$ and $-25.161 \pm 0.40\text{‰}$ in soil sampled before and after fire,
93 respectively; while in *Am* grassland were $-26.873 \pm 0.16\text{‰}$ and $-26.98 \pm 0.31\text{‰}$ before and after
94 fire, respectively. Our results are in agreement with similar observations reported by other authors
95 (Granged et al., 2011b) who found no change in SOC content before and after prescribed fire. ~~We~~
96 ~~reproduced the same environmental conditions of a wildfire. We can consider our experimental fire~~
97 ~~as a moderate fire severity, similar to prescribed fire described by previous mentioned authors.~~
98 The time of combustion was 12 ± 2 minutes and 7 ± 1 minutes for *Hh* and *Am*, respectively (Fig. 3).
99 The maximum temperature measured at soil surface was around 480 °C in both grasslands.
100 Temperatures over 200 °C persisted for 5 minutes and 3 minutes for *Hh* and *Am*, respectively. The
101 burning time and intensity was low due to low amount of fuel in both grasslands. Mediterranean
102 environmental conditions involve high organic matter mineralization rates and, thus, negligible
103 amounts of litter biomass stock. The low temperatures registered during low severity fires does
104 not have important



105 effects on SOC stock (Úbeda et al., 2005). The loss of organic carbon by burning can occur even
106 at relatively low temperatures such as 200 °C, but total combustion is only observed at high
107 temperatures 450-500°C (De Bano et al., 1998). When comparing the two grasslands, SOC
108 amount and the effect of fire on SOC stock was different. The lower SOC content was measured
109 under *Hh* grassland, which also recorded the lower biomass yield. The above ground biomass
110 estimated is 4.76 Mg ha⁻¹ and 11.60 Mg ha⁻¹ of dry matter for *Hh* and *Am* grassland, respectively.
111 Even if the SOC change before and after fire was not statistically significant, after fire SOC content
112 decreased of 11.5% in *Hh* and increased of 27.9 % in *Am* grassland. The increase of SOC after fire
113 could occur due to external inputs of charred material and ash, as commonly is observed in low
114 severity fires due fuel and organic matter incomplete combustion. In particular, the burned
115 material returns to soil as particles smaller than 2 mm in the form of ash, which are mixed in the
116 top horizon, and which cause a net increase of SOC content (Gonzalez-Perez et al., 2004). The
117 reason for the slight SOC increase after fire only in *Am* grassland may depend on different
118 characteristics of the two considered grasses. Firstly, *Am* biomass contains more lignin and
119 cellulose than *Hh* biomass (Table 1), and, thus, more recalcitrant compounds that under low
120 temperature do not completely volatilize. Secondly, *Am* has a densely caespitose habit: this feature
121 impedes a complete burning and favors the retention of not completely burnt plant residues. The
122 ash of *Hh* is, instead, lighter and quickly eroded by wind. This is clear evidence that *Hh* grassland
123 burned at higher severity, despite the similar temperatures observed. Previous studies observed
124 that fire severity is different according the burned specie (Pereira et al., 2011). Thirdly, biomass of
125 *Am* contains siliceous compounds that obstruct burning.


126 **Conclusions**

127 Data here reported confirm that the use of experimental fire to favour plant recovery in *Hh* and
128 *Am* grassland does not affect SOC stock, even if these grasslands did not burn for many years. Our
129 study shows that it is possible to adopt the system of controlled burning to maintain grassland
130 formations, however, this management tool must be adopted only after thorough phytosociological

131 analyses of local vegetation patterns and dynamics and after detailed planning of grazing after fire
132 (~~Naveh, 1974; Montserrat et al., 2001~~) 

133

134 **Acknowledgement**

~~135 Work was financially supported by the Italian government through the PRIN project "The impacts
136 of secondary succession processes on carbon storage in soil and biomass and on biodiversity and
137 the role of dispersal centers and vectors for recolonization processes".~~ 

138 **References**

- 139 Barberis, A., Dettori, S., and Filigheddu, M.R.: Management problems in Mediterranean cork-oak
140 forests: Post-fire recovery, *J. Arid Environmen.*, 54, 565–569, 2003.
- 141 Bond, W.J., Woodward, F.I., and Midgley, G.F.: The global distribution of ecosystems in a world
142 without fire. *New phytol.*, 165, 525–538, 2005.
- 143 Braun-Blanquet J.: *Plant sociology*. Mc Graw-Hill, New York-London, 1932.
- 144 Briggs, J.M., Knapp, A.K., Blair, J.M., Heisler, J.L., Hoch, G.A., Lett, M.S., and McCarron, J.K.: An
145 ecosystem in transition: Causes and consequences of the conversion of mesic grassland to
146 shrubland. *BioScience*, 55, 243–254, 2005.
- 147 Brullo, C., Brullo, S., Giusso del Galdo, G., Guarino, R., Minissale, P., Scuderi, L., Siracusa, G.,
148 Sciandrello, S., and Spampinato, G.: The Lygeo-Stipetea class in Sicily.- *Annali di Botanica*,
149 pp. 1-28, 2010.
- 150 Certini, G.: Effects of fire on properties of forest soils: a review. *Oecologia*, 143, 1-10, 2005.
- 151 DeBano, L.F., Neary, D.G., and Ffolliott, P.F.: *Fire's Effects on Ecosystems*. Wiley, New York, 1998.
- 152 Díez-Garretas B., and Asensi A.: Syntaxonomic analysis of the Andropogon-rich grasslands
153 (*Hyparrhenetalia hirtae*) in the Western Mediterranean Region.- *Folia Geobotanica et*
154 *Phytotaxonomica*, 34, 307–320, 1999.

155 Fernandez, I., Cabaneiro, A., and Carballas, T.: Organic matter changes immediately after a
156 wildfire in an Atlantic forest soil and comparison with laboratory soil heating. *Soil Biol.*
157 *Biochem.*, 29, 1–11, 1997.

158 Gonzalez-Perez, J.A., Gonzalez-Vila, F.J., Almendros, G., and Knicker, H.: The effect of fire on soil
159 organic matter-a review. *Environmen. Int.*, 30, 855–870, 2004

160 Granged, A.J.P., Jordán A., Zavala, L.M., Muñoz-Rojas, M., and Mataix-Solera, J.: Short-term
161 effects of experimental fire for a soil under eucalyptus forest (SE Australia), *Geoderma*,
162 167–168, 125–134, 2011a.

163 Granged, A.J.P. Zavala, L.M, Jordán A., and Barcenas-Moreno, G.: Post-fire evolution of soil
164 properties and vegetation cover in a Mediterranean hethland after experimental burning: A
165 3-year study, *Geoderma*, 164, 85-94. 2011b.

166 Kavdir, Y., Ekinci, H., Yüksel, O., and Mermut, A.R.: Soil aggregate stability and ¹³C CP/MAS-NMR
167 assessment of organic matter in soils influenced by forest wildfires in Canakkale, Turkey,
168 *Geoderma*, 129, 219–229, 2005.

169 Knicker, H., 2007.: How does fire affect the nature and stability of soil organic nitrogen and
170 carbon? A review, *Biogeochemistry* 85, 91–118, 2007.

171 La Mantia T., Carimi F., Di Lorenzo R. and Pasta S.: The agricultural heritage of Lampedusa
172 (Pelagie Archipelago, South Italy) and its key role for cultivar and wildlife conservation.
173 *Italian Jurnal of Agronomy*, 17, 106–110, 2011.

174 Montserrat, V., Lloret, F., Ogheri, E., Terradas, J.: Positive fire–grass feedback in Mediterranean
175 Basin woodlands 2001. *Forest Ecol. Manag.* 147, 3–14, 2001.

176 Naveh, Z.: Effects of fire in the mediterranean region. In Kozlowski T.T., Ahlgern CE. „Fire and
177 Ecosystem, Academic Press, New York, 401-434,1974

178 Novara, A., Gristina, L., Bodì, B.M, Cerdà, A.: 2011. The impact of fire on redistribution of soil
179 organic matter on a Mediterranean hillslope under maquia vegetation type. *Land Degrad.*
180 *Dev.*, 22, 530–536, 2011.

- 181 Pausas, J.G., Verdu, M.: 2005. Plant persistence traits in fire-prone ecosystems of Mediterranean
182 basin: a phylogenic approach, *Oikos*, 109, 196–202., 2005.
- 183 Pereira, P., Bodi, M., Úbeda, X., Cerdà, A., Mataix-Solera, J., Balfour, V., and Woods, S.: Las
184 cenizas y el ecosistema suelo, In: Cerdà, A. Jordan, A. (eds) Actualización en métodos y
185 técnicas para el estudio de los suelos afectados por incendios forestales, 345-398. Càtedra
186 de Divulgació de la Ciència. Universitat de València, 2010.
- 187 Pereira, P., Cerdà, A., Úbeda, X., Mataix-Solera, J. Arcenegui, V., and Zavala, L.: Modelling the
188 impacts of wildfire on ash thickness in a short-term period. *Land Degrad. Dev.*
189 DOI:10.1002/ldr.2195, 2013.
- 190 Pereira, P., Úbeda, X. Martin, D.A., Mataix-Solera, J., and Guerrero, C.: Effects of a low prescribed
191 fire in ash water soluble elements in a Cork Oak (*Quercus suber*) forest located in
192 Northeast of Iberian Peninsula, *Env. Res.*, 111, 237–247, 2011.
- 193 Pyne, S. J.: *Fire: A Brief History*. Seattle, WA: University of Washington Press, 2001.
- 194 SAS Institute.: *The SAS System for Microsoft Windows*. Release 8.2. SAS Inst.: Cary, NC2002.
- 195 Snyman, HA.: Short-term response of rangeland following an unplanned fire in terms of soil
196 characteristics in a semiarid climate of South Africa. *J. Arid Environ.*, 55, 160–180, 2003.
- 197 Terefe, T., Mariscal-Sancho, I., Peregrina, F., Espejo, R.: Influence of heating on various
198 properties of six Mediterranean soils. A laboratory study, *Geoderma*, 143 (3-4), 273–280,
199 2008.
- 200 Trabaud, L.: Man and fire: impacts on Mediterranean vegetation. In: di Castri, F., Goodallm, D.W.,
201 Specht, R.L. (Eds.), *Mediterranean-type shrublands*. Elsevier, Amsterdam, pp. 523–537,
202 1981.
- 203 Úbeda, X., Lorca, M., Outeiro, L.R., Bernia, S., Castellnou, M.: Effect of prescribed fire on soil
204 quality in Mediteanean grassland (prades Mountains, north-east Spain), *Int. J. Wildland*
205 *Fire*, 14, 379–384, 2005.
- 206 WRB,: *World Reference base for soil resources 2006*, 2nd edition. World Soil Resources Reports, No
207 103. Food and Agriculture Organization of the United Nation, Rome, 2006.

208

209 **Table caption**

210 Table 1. Biomass composition (% of dry biomass) of *Hyparrhenia Hirta* (Hh) and *Ampelodesmos*
211 *mauritanicus* (Am). Values in parenthesis are standard deviations. Abbreviations: ADF = acid
212 detergent fiber, NDF = neutral detergent fiber.

213

Grassland	ADF	Cellulose	NDF	Hemicellulose	Ash	Aboveground biomass (Mg ha ⁻¹)	C Biomass (g kg ⁻¹)
<i>Am</i>	6.91 (0.58)	37.72 (1.58)	73.03 (2.65)	23.99 (1.32)	4.02 (1.10)	4.76	43.8
<i>Hh</i>	5.98 (0.68)	34.00 (1.20)	72.01 (1.53)	28.26 (1.76)	4.34 (1.49)	11.60	45.8

214

215


216 **Figure Caption**

217 Fig.. Localization of the study area (black point) in Sicily (rectangle in inset) and with respect to
218 the next urban settlement Cefalù. Grey scale represents altitudinal gradient (1 shade of grey = 200
219 m).

220

221


222



223

224

225 Fig. 2. Soil organic carbon before and after fire in *Hyparrhenia Hirta* (Hh) and *Ampelodesmos*
226 *mauritanicus* (Am) grassland.



227

228

229 Figure 3- Soil temperature during fire under *Hyparrhenia Hirta* (Hh) (blu line) and *Ampelodesmos*
230 *mauritanicus* (Am) (red line) grassland

231

232

