

1 **SHORT COMMUNICATION**

2 **Grassland fire effect on soil organic carbon reservoirs in a semiarid environment**

3 Agata Novara<sup>(1)</sup>, Luciano Gristina<sup>(1)</sup>, Juliane Rühl<sup>(1)</sup>, Salvatore Pasta<sup>(1)</sup>, Giuseppe D'Angelo<sup>(1)</sup> Tommaso La  
4 Mantia<sup>(1)</sup>, Paulo Pereira<sup>(2)</sup>

5 1 Dipartimento di Scienze agrarie e forestali – University of Palermo - 90128 Palermo – Italy

6 2 Environmental Research Centre, Mykolas Romeris University, Ateities g. 20, LT-08303 Vilnius, Lithuania  
7 (paulo@mruni.eu).

8 Corresponding author: Agata Novara agata.novara@unipa.it

9

10 **Abstract**

11 The aim of this work was to investigate the effect of an experimental fire, used for grassland management,  
12 on soil organic carbon (SOC) stocks. The study was carried out on *Hyparrhenia hirta* (L.) Stapf (*Hh*)  
13 grassland and *Ampelodesmos mauritanicus* (Desf.) T. Durand & Schinz (*Am*) grasslands, located in the north  
14 of Sicily. Soil samples were collected at 0-5 cm before and after experimental fire and SOC was measured.  
15 During grassland fire soil surface temperature was monitored. Biomass of both grasses was analyzed in order  
16 to determine dry weight and its chemical composition. The results showed that SOC varied significantly with  
17 vegetation type, while it is not affected in the short period by grassland fire. *Am* grassland stored more SOC  
18 compared with *Hh* grassland thanks to lower content in biomass of labile carbon pool. No significant  
19 difference was observed in SOC before and after fire which could be caused by several factors: first, in both  
20 grassland types the measured soil temperature during fire was low due to thin litter layers; second, in  
21 semiarid environment higher mineralization rate results in lower soil carbon labile pool; and third, the SOC  
22 stored in the finest soil fractions, physically protected, is not affected by fire.

23 **Keywords:** SOC, experimental fire, grassland, Mediterranean environment

24

## 25 **Introduction**

26 Fire is part of the Earth System and has been for millennia a tool for many societies (Pyne, 2001). Fire is  
27 regarded as an active ecological agent able to mobilize nutrients and restore soil fertility (Snyman, 2003) but,  
28 also as a primary cause of soil degradation due to nutrients loss for volatilization, leaching and erosion,  
29 especially in severe wildfires. It is, in fact, considered a major disturbance in many ecosystems, which lead  
30 to important shifts in the soil properties and vegetation (Certini, 2005; Granged et al., 2011a). One of the  
31 most common effects of fire is the alteration in composition and amount of soil organic matter (Knicker,  
32 2007, Terefe et al., 2008). Several studies recorded a decrease (Fernández et al., 1997; Novara et al., 2011) in  
33 soil organic carbon (SOC) after fire, while results of other studies showed no significantly changing or even  
34 increase of previous SOC content (Kavdir et al., 2005). These discrepancies occur due to the large amount of  
35 controlling factors and therefore the effect of fire is highly variable in space and time. Among these factors,  
36 fire intensity, fire severity, fire regimen, type of burned vegetation, connectivity, distribution of fuel on soil  
37 surface, type of ash produced and dispersion, topography, soil properties, aspect, regional climate and  
38 meteorological conditions in the immediate period after the fire play a key role to determine SOC alteration  
39 and accumulation in soils (Certini, 2005; Pereira et al. 2010; Pereira et al., 2013).

40 In semiarid areas fire is one of the common management tools used by shepherds to enhance pasture  
41 regrowth. In fact, the recovery of vegetation canopy after fire in the Mediterranean area can be quite rapid  
42 due to adaption of plant communities to the disturbances caused by fire as observed in several studies  
43 (Trabaud, 1981; Oba, 1990; Woube, 1998; Barberis et al., 2003; Pausas and Verdú 2005). It is known,  
44 moreover, that fire is considered an important factor for arid and semiarid grasslands because it avoids  
45 invasion of trees and shrubs with implications on soil carbon storage (Briggs et al., 2005). Despite the  
46 importance of fire on grassland ecosystems (Bond et al., 2005), its impact on SOC is not well understood in  
47 the immediate period after the fire in the Mediterranean grasslands (Snyman, 2003). The aim of this work is  
48 to quantify SOC stock change as a result of an experimental fire of two of the most widespread types of  
49 Mediterranean grasslands (Brullo et al., 2010; Díez-Garretas and Asensi, 1999) and, therefore, to establish if  
50 this practice could be used as a sustainably management tool for grazing recovery (Álvarez-Martínez et al.,  
51 2013).

52 **Materials and Methods**

53 The field studies were carried out in the province of Palermo, Sicily (Italy) (350 m a.s.l.) (Fig. 1). Local soil  
54 type is an *Eutric Cambisol* according to WRB (WRB, 2006) with sand and clay contents of 18% and 46%,  
55 respectively. The climate is Mediterranean, with mean annual rainfall of 580 mm and yearly average  
56 temperature of 16 °C.

57 An experimental fire was conducted on July and September 2009 on five (replicas) delimited square areas  
58 (50×50 cm) in two different grassland types, dominated by *Hyparrhenia hirta* (L.) Stapf (Hh) and  
59 *Ampelodesmos mauritanicus* (Desf.) T. Durand & Schinz (Am). Each sampling square was about 2m distant  
60 from the neighbor square. In order to simulate a natural wildfire, burning was allowed to take its natural  
61 course until it extinguished itself. The fire was generated with a match, starting from leeward in each plot.  
62 Soil surface temperature during the burning was measured using a thermocouple system (type K Inconel 600  
63 insulated). In each selected area three soil samples were collected at 0-5 cm depth before and immediately  
64 after fire. On three one meter square in both grasslands (dominated by *Hyparrhenia hirta* or *Ampelodesmos*  
65 *mauritanicus*) all plants were cut, oven dried for 3-4 days at 60-65 °C, and weighted. SOC content was  
66 measured using a CHN-Elemental Analyzer. For the  $\delta^{13}C$  analysis, an EA-IRMS (elemental analyzer  
67 isotope ratio mass spectrometry) was used. The International Atomic Energy Agency (IAEA), Vienna,  
68 distribute IAEA-CH-6 as a reference standard material. The results of the isotope analysis are expressed as a  
69  $\delta$  value (‰) relative to the international Pee Dee Belemnite standard as follows:

70

$$\delta(\text{‰}) = \frac{R_s - R_{st}}{R_{st}} * 1000$$

71

72 where  $\delta = \delta^{13}C$ , R =  $^{13}C/^{12}C$ , s = sample, and st = standard.

73 Dry biomass weight and its chemical composition (ADF acid detergent fiber, NDF neutral detergent fiber,  
74 Cellulose, Hemicellulose, Lignin, Ash) were determined on three 0.5 m<sup>2</sup> square area subsamples for each  
75 grassland types.

76 Data analysis was conducted using the SAS statistical package (SAS Inst., 2002). Normal distribution of data  
77 was verified previously to statistical data comparisons and analysis of variance (ANOVA) was conducted.

78 Significant differences were considered at a  $p < 0.05$ .

79

## 80 **Results and discussion**

81

82 SOC ranged from 20.3 to 37.0g kg<sup>-1</sup> and from 15.4 to 32.5 g kg<sup>-1</sup> before and after experimental fire,  
83 respectively, in soil covered by *Hh*, and from 32.5 to 38.2 g kg<sup>-1</sup> and from 38.3 to 49.1 g kg<sup>-1</sup> before and after  
84 experimental fire, respectively, in soil covered by *Am*. The experimental fire did not have significant  
85 differences in SOC in both grassland types (Fig. 2). Similarly to SOC results,  $\delta^{13}\text{C}$  was not affected  
86 significantly by fire. The average by time of  $\delta^{13}\text{C}$  values measured in *Hh* grassland were  $-25.418 \pm 0.25\text{‰}$   
87 and  $-25.161 \pm 0.40\text{‰}$  in soil sampled before and after fire, respectively; while in *Am* grassland were  $-26.873$   
88  $\pm 0.16\text{‰}$  and  $-26.98 \pm 0.31\text{‰}$  before and after fire, respectively. Our results are in agreement with similar  
89 observations reported by other authors (Granged et al., 2011b) who found no change in SOC content before  
90 and after prescribed fire. The experimental fire has a moderate fire severity, similar to prescribed fire  
91 described by Granged et al. (2011b). The time of combustion was  $12 \pm 2$  minutes and  $7 \pm 1$  minutes for *Hh* and  
92 *Am*, respectively (Fig. 3). The maximum temperature measured at soil surface was around 480 °C in both  
93 grasslands. Temperatures over 200 °C persisted for 5 minutes and 3 minutes for *Hh* and *Am*, respectively.  
94 The burning time and intensity was low due to low amount of fuel in both grasslands. Mediterranean  
95 environmental conditions involve high organic matter mineralization rates and, thus, negligible amounts of  
96 litter biomass stock. The low temperatures registered during low severity fires does not have important  
97 effects on SOC stock (Úbeda et al., 2005). The loss of organic carbon by burning can occur even at relatively  
98 low temperatures such as 200 °C, but total combustion is only observed at high temperatures 450-500°C (De  
99 Bano et al., 1998). When comparing the two grasslands, the SOC amount and the effect of fire on SOC stock  
100 was contrasted. The lower SOC content was measured under *Hh* grassland, which also recorded the lower  
101 biomass yield. The above ground biomass estimated is 4.76 Mg ha<sup>-1</sup> and 11.60 Mg ha<sup>-1</sup> of dry matter for *Hh*  
102 and *Am* grassland, respectively.

103 Even if the SOC change before and after fire was not statistically significant, after fire SOC content  
104 decreased of 11.5% in *Hh* and increased of 27.9 % in *Am* grassland. The increase of SOC after fire could  
105 occur due to external inputs of charred material and ash, as commonly is observed in low severity fires due  
106 fuel and organic matter incomplete combustion. In particular, the burned material returns to soil as particles

107 smaller than 2 mm in the form of ash, which are mixed in the top horizon, and which cause a net increase of  
108 SOC content (González-Pérez et al., 2004). The reason for the slight SOC increase after fire only in *Am*  
109 grassland may depend on different characteristics of the two considered grasses. Firstly, *Am* biomass  
110 contains more lignin and cellulose than *Hh* biomass (Table 1), and, thus, more recalcitrant compounds that  
111 under low temperature do not completely volatilize. Secondly, *Am* has a densely caespitose habit: this feature  
112 impedes a complete burning and favors the retention of not completely burnt plant residues. The ash of *Hh*  
113 is, instead, lighter and quickly eroded by wind (Cerdà and Doerr, 2007). This is clear evidence that *Hh*  
114 grassland burned at higher severity, despite the similar temperatures observed. Previous studies observed that  
115 fire severity is different according the burned specie (Pereira et al., 2011). Thirdly, biomass of *Am* contains  
116 siliceous compounds that obstruct burning.

## 117 **Conclusions**

118 Data here reported confirm that the use of experimental fire to favour plant recovery in *Hh* and *Am* grassland  
119 does not affect SOC stock, even if these grasslands did not burn for many years. Our study shows that it is  
120 possible to adopt the system of controlled burning to maintain grassland formations, however, this  
121 management tool must be adopted only after thorough phytosociological analyses of local vegetation patterns  
122 and dynamics and after detailed planning of grazing after fire.

123

## 124 **Acknowledgement**

125 This research was financially supported by the MIUR through the PRIN project “CARBOTREES”.

## 126 **References**

- 127 Álvarez-Martínez, J., Gómez-Villar, A., and Lasanta, T.: The use of goats grazing to restore pastures invaded  
128 by shrubs and avoid desertification: a preliminary case study in the spanish cantabrian mountains,  
129 Land degradation & development 2013, DOI: 10.1002/ldr.2230.
- 130 Barberis, A., Dettori, S., and Filigheddu, M.R.: Management problems in Mediterranean cork-oak forests:  
131 Post-fire recovery, J. Arid Environmen., 54, 565–569, 2003.

- 132 Bond, W.J., Woodward, F.I., and Midgley, G.F.: The global distribution of ecosystems in a world without  
133 fire. *New phytol.*, 165, 525–538, 2005.
- 134 Braun-Blanquet J.: *Plant sociology*. Mc Graw-Hill, New York-London, 1932.
- 135 Briggs, J.M., Knapp, A.K., Blair, J.M., Heisler, J.L., Hoch, G.A., Lett, M.S., and McCarron, J.K.: An  
136 ecosystem in transition: Causes and consequences of the conversion of mesic grassland to shrubland.  
137 *BioScience*, 55, 243–254, 2005.
- 138 Brullo, C., Brullo, S., Giusso del Galdo, G., Guarino, R., Minissale, P., Scuderi, L., Siracusa, G., Sciandrello,  
139 S., and Spampinato, G.: The Lygeo-Stipetea class in Sicily.- *Annali di Botanica*, pp. 1-28, 2010.
- 140 Cerdà, A. and Doerr, S.H.: Soil wettability, runoff and erodibility of major dry-Mediterranean land use types  
141 on calcareous soils. *Hydrological Processes*, 21, 2325-2336, 2007, doi:  
142 10.1016/j.catena.2008.03.010.
- 143 Certini, G.: Effects of fire on properties of forest soils: a review. *Oecologia*, 143, 1-10, 2005.
- 144 DeBano, L.F., Neary, D.G., and Ffolliott, P.F.: *Fire's Effects on Ecosystems*. Wiley, New York, 1998.
- 145 Díez-Garretas, B., and Asensi A.: Syntaxonomic analysis of the *Andropogon*-rich grasslands  
146 (*Hyparrhenetalia hirtae*) in the Western Mediterranean Region.- *Folia Geobotanica et*  
147 *Phytotaxonomica*, 34, 307–320, 1999.
- 148 Fernández, I., Cabaneiro, A., and Carballas, T.: Organic matter changes immediately after a wildfire in an  
149 Atlantic forest soil and comparison with laboratory soil heating. *Soil Biol. Biochem.*, 29, 1–11, 1997.
- 150 González-Perez, J.A., González-Vila, F.J., Almendros, G., and Knicker, H.: The effect of fire on soil organic  
151 matter-a review. *Environmen. Int.*, 30, 855–870, 2004.
- 152 Granged, A.J.P., Jordán A., Zavala, L.M., Muñoz-Rojas, M., and Mataix-Solera, J.: Short-term effects of  
153 experimental fire for a soil under eucalyptus forest (SE Australia), *Geoderma*, 167–168, 125–134,  
154 2011a.
- 155 Granged, A.J.P., Zavala, L.M, Jordán A., and Barcenás-Moreno, G.: Post-fire evolution of soil properties and  
156 vegetation cover in a Mediterranean hethland after experimental burning: A 3-year study, *Geoderma*,  
157 164, 85-94. 2011b.
- 158

- 159 Kavdir, Y., Ekinçi, H., Yüksel, O., and Mermut, A.R.: Soil aggregate stability and <sup>13</sup>C CP/MAS-NMR  
160 assessment of organic matter in soils influenced by forest wildfires in Canakkale, Turkey, *Geoderma*,  
161 129, 219–229, 2005.
- 162 Knicker, H.: How does fire affect the nature and stability of soil organic nitrogen and carbon? A review,  
163 *Biogeochemistry* 85, 91–118, 2007.
- 164 La Mantia T., Carimi F., Di Lorenzo R. and Pasta S.: The agricultural heritage of Lampedusa (Pelagic  
165 Archipelago, South Italy) and its key role for cultivar and wildlife conservation. *Italian Journal of*  
166 *Agronomy*, 17, 106–110, 2011.
- 167 Novara, A., Gristina, L., Bodì, B.M, Cerdà, A.: 2011. The impact of fire on redistribution of soil organic  
168 matter on a Mediterranean hillslope under maquia vegetation type. *Land Degrad. Dev.*, 22, 530–536,  
169 2011.
- 170 Oba, G.; Effects of wildfire on a semidesert riparian woodland along the Turkwel river, Kenya, and  
171 management implications for turkana pastoralists *Land Degradation & Development*, 2 (4) 247–259,  
172 1990.
- 173 Pausas, J.G., Verdú, M.: 2005. Plant persistence traits in fire-prone ecosystems of Mediterranean basin: a  
174 phylogenetic approach, *Oikos*, 109, 196–202., 2005.
- 175 Pereira, P., Bodi, M., Úbeda, X., Cerdà, A., Mataix-Solera, J., Balfour, V., and Woods, S.: Las cenizas y el  
176 ecosistema suelo, In: Cerdà, A. Jordan, A. (eds) *Actualización en métodos y técnicas para el estudio*  
177 *de los suelos afectados por incendios forestales*, 345-398. Càtedra de Divulgació de la Ciència.  
178 Universitat de Valencia, 2010.
- 179 Pereira, P., Cerdà, A., Úbeda, X., Mataix-Solera, J. Arcenegui, V., and Zavala, L.: Modelling the impacts of  
180 wildfire on ash thickness in a short-term period. *Land Degrad. Dev.* DOI:10.1002/ldr.2195, 2013.
- 181 Pereira, P., Úbeda, X. Martin, D.A., Mataix-Solera, J., and Guerrero, C.: Effects of a low prescribed fire in  
182 ash water soluble elements in a Cork Oak (*Quercus suber*) forest located in Northeast of Iberian  
183 Peninsula, *Env. Res.*, 111, 237–247, 2011.
- 184 Pyne, S. J.: *Fire: A Brief History*. Seattle, WA: University of Washington Press, 2001.
- 185 SAS Institute.: *The SAS System for Microsoft Windows*. Release 8.2. SAS Inst.: Cary, NC2002.

186 Snyman, HA.: Short-term response of rangeland following an unplanned fire in terms of soil characteristics  
 187 in a semiarid climate of South Africa. *J. Arid Environ.*, 55, 160–180, 2003.

188 Terefe, T., Mariscal-Sancho, I., Peregrina, F., Espejo, R.: Influence of heating on various properties of six  
 189 Mediterranean soils. A laboratory study, *Geoderma*, 143 (3-4), 273–280, 2008.

190 Trabaud, L.: Man and fire: impacts on Mediterranean vegetation. In: di Castri, F., Goodallm, D.W., Specht,  
 191 R.L. (Eds.), *Mediterranean-type shrublands*. Elsevier, Amsterdam, pp. 523–537, 1981.

192 Úbeda, X., Lorca, M., Outeiro, L.R., Bernia, S., Castellnou, M.: Effect of prescribed fire on soil quality in  
 193 Mediteanean grassland (prades Mountains, north-east Spain), *Int. J. Wildland Fire*, 14, 379–384,  
 194 2005.

195 Woube, M.: Effect of fire on plant communities and soils in the humid tropical savannah of Gambela,  
 196 Ethiopia, *Land Degradation & Development* 9 (3), 275–292, 1998.

197 WRB.: World Reference base for soil resources 2006, 2nd edition. *World Soil Resources Reports*, No  
 198 103. Food and Agriculture Organization of the United Nation, Rome, 2006.

199 **Table caption**

200 Table 1. Biomass composition (% of dry biomass) of *Hyparrhenia Hirta* (Hh) and *Ampelodesmos*  
 201 *mauritanicus* (Am). Values in parenthesis are standard deviations. Abbreviations: ADF = acid detergent  
 202 fiber, NDF = neutral detergent fiber.

203

Grassland	ADF	Cellulose	NDF	Hemicellulose	Ash	Aboveground biomass (Mg ha <sup>-1</sup> )	C Biomass (g kg <sup>-1</sup> )
<i>Am</i>	6.91 (0.58)	37.72 (1.58)	73.03 (2.65)	23.99 (1.32)	4.02 (1.10)	4.76	43.8
<i>Hh</i>	5.98 (0.68)	34.00 (1.20)	72.01 (1.53)	28.26 (1.76)	4.34 (1.49)	11.60	45.8

204

205




206 **Figure Caption**

207 Fig.. Localization of the study area (black point) in Sicily (rectangle in inset) and with respect to the next  
208 urban settlement Cefalù. Grey scale represents altitudinal gradient (1 shade of grey = 200 m).

209

210


211



212

213

214 Fig. 2. Soil organic carbon before and after fire in *Hyparrhenia Hirta* (Hh) and *Ampelodesmos mauritanicus*  
215 (Am) grassland.



216

217

218 Figure 3- Soil temperature during fire under *Hyparrhenia Hirta* (Hh) (blu line) and *Ampelodesmos*  
219 *mauritanicus* (Am) (red line) grassland

220

221

