

# Jurassic–Paleogene Intra–Oceanic Magmatic Evolution of the Ankara Mélange, North–Central Anatolia, Turkey

Ender Sarifakioglu<sup>1</sup>, Yildirim Dilek<sup>2</sup> and Mustafa Sevin<sup>1</sup>

<sup>1</sup>General Directorate of Mineral Research & Exploration, Department of Geology, TR–06520 Ankara, Turkey, esarifakioglu@mta.gov.tr

<sup>2</sup>Department of Geology & Env. Earth Science, Miami University, Oxford, OH 45056, USA, dileky@miamioh.edu

## Abstract

Oceanic rocks in the Ankara Mélange along the Izmir–Ankara–Erzincan suture zone (IAESZ) in North–Central Anatolia include locally coherent ophiolite complexes (~179 Ma and ~80 Ma), seamount or oceanic plateau volcanic units with pelagic and reefal limestones ( $96.6 \pm 1.8$  Ma), metamorphic rocks with ages of  $187.4 \pm 3.7$  Ma,  $158.4 \pm 4.2$  Ma, and  $83.5 \pm 1.2$  Ma, and subalkaline to alkaline volcanic and plutonic rocks of an island arc origin (~67–63 Ma). All but the arc rocks occur in a shaly–graywacke and/or serpentinite matrix, and are deformed by south–vergent thrust faults and folds that developed in the Middle to Late Eocene due to continental collisions in the region. Ophiolitic volcanic rocks have mid–ocean ridge (MORB) and island arc tholeiite (IAT) affinities showing moderate to significant large ion lithophile elements (LILE) enrichment and depletion in Nb, Hf, Ti, Y and Yb, which indicate the influence of subduction–derived fluids in their melt evolution. Seamount/oceanic plateau basalts show ocean island basalt (OIB) affinities. The arc–related volcanic rocks, lamprophyric dikes and syeno–dioritic plutons exhibit high–K shoshonitic to medium–to high–K calc–alkaline compositions with strong enrichment in LILE, rare earth elements (REE) and Pb, and initial  $\varepsilon_{\text{Nd}}$  values between +1.3 and +1.7. Subalkaline arc volcanic units occur in the northern part of the mélange, whereas the younger alkaline volcanic rocks and intrusions (lamprophyre dikes and syeno–dioritic plutons) in the southern part. The Early to Late Jurassic and Late Cretaceous epidote–actinolite, epidote–chlorite and epidote–glaucophane schists represent the metamorphic units formed in a subduction channel in the Northern Neotethys. The Middle to Upper Triassic neritic limestones spatially associated with the seamount volcanic rocks indicate that the Northern Neotethys was an open ocean with its MORB–type oceanic lithosphere by the Early Triassic. The Latest Cretaceous–Early Paleocene island arc volcanic, dike and plutonic rocks with subalkaline to alkaline geochemical affinities represent intraoceanic magmatism that developed on and across the subduction–accretion complex above a N–dipping, southward–rolling subducted lithospheric slab within the Northern Neotethys. The Ankara Mélange thus exhibits the record of ~120–130 million years of oceanic magmatism in geological history of the Northern Neotethys.

**Keywords:** Ankara Mélange (Turkey), Northern Neotethys, seamount volcanics, suprasubduction zone ophiolites, subduction–accretion complex, island arc magmatism

## 1. Introduction

In the circum-Mediterranean mountain chains belonging to the Alpine-Himalayan system, subduction-related tectonic mélanges during pre-collisional stages are described, but they generally overprinted by arc-continent and continent-continent collisions (Festa et al., 2010 and references therein). In northern Turkey, the 2600-km-long IAESZ extends from west to east, connecting Vardar Suture in west and Sevan-Akera Suture Zone in east. The ophiolitic mélanges and ophiolite slabs are observed along this zone. At the central part of IAESZ, in the vicinity of Ankara, Kırıkkale, Çankırı and Çorum, the Ankara Mélange, first described by Bailey and McCallien (1950), is a well-known subduction-accretion type mélange of the world. They defined metamorphic, limestone and ophiolitic rock blocks in age from Paleozoic to Mesozoic in the mélange. In previous works, three major tectonic units are identified in the Ankara Mélange from the northwest to the southeast. These are metamorphic block mélange, limestone block mélange and ophiolitic mélange (Norman, 1984; Akyürek et al., 1984; Koçyiğit, 1991; Tüysüz et al., 1995; Tankut et al., 1998). The metamorphic block mélange contains a chaotic mixture of variably metamorphosed sedimentary, basic-ultrabasic, and pyroclastic rocks in age from Permian to Triassic while the limestone block mélange consists of neritic to pelagic limestone blocks in age ranging from Permian through Albian in a shale-graywacke matrix. The ophiolitic mélange includes several kilometer-size thrust sheets of mantle peridotite, oceanic basic crustal rock, and blocks of serpentinite, massive to pillow basaltic lava flows, radiolarite, chert and neritic-pelagic limestone in pelitic and/or serpentinite matrix (Norman, 1984; Akyürek et al., 1984; Koçyiğit, 1991; Tüysüz et al., 1995). Tankut et al. (1998) emphasized that the ophiolitic mélange unit of the Ankara Mélange is represented by two major mappable coherent units as ophiolitic fragments and volcanic seamounts. They determined that N-MORB character of the Neo-Tethyan oceanic crust along with its seamounts was overprinted by a chemical signature related to subduction zone processes and associated magmatism prior to their incorporation into the subduction-accretion mélange.

In this paper we document the internal structure of the Ankara Mélange along the IAESZ in North-Central Anatolia, and present new geochemical and geochronological data from various magmatic rock assemblages that make up distinct tectonic units in this mélange. Our geochemical data and interpretations indicate that all units within the Ankara Mélange are intraoceanic in origin and appear to have formed during the seafloor spreading, seamount volcanism and island arc magmatism stages of the Northern Neotethys. We also present new Pb-Sr-Nd isotopic compositional data, and radiometric data belonging to both magmatic arc rocks and basic rocks from the Ankara Mélange. Thus, the Ankara Mélange

displays a complete record of ~120–130 m.y. of intraoceanic magmatism that took place prior to the continental collisional events in Anatolia in the Eocene.

## 2. Regional Geology

The IAESZ forms the tectonic boundary between the Pontide tectonic belt (including the Sakarya Continent) in the north and the Anatolide–Tauride block including the Central Anatolian Crystalline Complex (CACC) in the south. The suture zone is marked by ophiolite units, ophiolitic mélanges, and seamount fragments. The Sakarya Continent in the Pontides represents the southern margin of Eurasia (Figs. 1 and 2). Carboniferous (330–310 Ma), high-grade metamorphic rocks (gneiss, migmatite, amphibolite and marble), currently exposed in the Kazdağ, Söğüt, Devrekani, and Pulur massifs (from west to east), make up the continental basement of the Sakarya Continent (Topuz et al., 2004, 2006; Okay et al., 2006; Nzegge et al., 2006). These metamorphic basement rocks are intruded by the Carboniferous (295 Ma) granitoids (Çoğulu et al., 1965; Delaloye and Bingöl, 2000). The Triassic Karakaya Complex, representing a subduction–accretion complex, tectonically overlies the crystalline basement units (Tekeli, 1981). It includes the Lower Karakaya, which comprises metabasite, marble and phyllite rocks, and the Upper Karakaya consisting mainly of unmetamorphosed clastic and basic volcanic rocks with blocks of Carboniferous and Permian neritic limestones (Bingöl et al., 1975; Okay et al., 2002; Okay and Göncüoğlu, 2004).

The CACC consists mainly of Paleozoic–Mesozoic metamorphic massifs (Kırşehir, Akdağ, and Niğde massifs) and Cretaceous–Paleocene granitoids (Fig. 2). The metamorphic massifs comprise metacarbonate, metapelite and amphibolite–gneiss rocks that are the products of varied P/T conditions of metamorphism (Whitney and Dilek 1998). The Late Cretaceous granitoids and the Eocene–Upper Miocene volcanic rocks crosscut and overlie (respectively) the crystalline basement units of the CACC (Güleç, 1994; Boztuğ, 2000; Kadioğlu et al., 2003, 2006; İlbeyli et al., 2004). The Late Cretaceous plutons are composed of *Granite*, *Monzonite* and *Syenite Supersuites* with ages of  $77.7\pm0.3$  Ma,  $70\pm1.0$  Ma and  $69.8\pm0.3$  Ma, respectively (Kadioğlu et al., 2006). They display a chemical progression from high-K calc-alkaline and high-K shoshonitic to alkaline compositions, representing the development of within-plate magmatism across the CACC with time (Kadioğlu et al., 2006).

## 3. Internal Structure and Tectonic Units of the Ankara Mélange

The most important component of the IAESZ in its central segment in northern Anatolia is the Ankara Mélange, extending from Ankara in the west to Çorum in the east (Fig. 2). The

Ankara Mélange is a well known subduction–accretion complex (Bailey and McCallien, 1950, 1953), consisting of blocks of Paleozoic limestone and metamorphic rocks, Jurassic–Cretaceous ophiolitic units, and Jurassic–Cretaceous seamount volcanic assemblages in a shaly–graywacke and/or serpentinite matrix (Figs. 3a, 3b and 4; Norman, 1984; Akyürek et al., 1984; Koçyiğit, 1991; Tüysüz et al., 1995; Tankut et al., 1998; Dilek and Thy, 2006; Dangerfield et al., 2011).

Megablocks and imbricated thrust sheets of oceanic rocks occur as mappable units enveloped in a pelitic (clayey, sandy–silty), serpentinite or volcanic matrix within the Ankara Mélange (Fig. 5). In some of these blocks or thrust sheets the mafic–ultramafic rock units and the associated sedimentary rocks make up coherent ophiolite complexes (e.g. the Eldivan ophiolite) (Figs. 6 and 7), representing the Neotethyan oceanic lithosphere. Plagiogranite dikes intruding the serpentinitized peridotites near Eldivan (Çankırı) revealed U–Pb zircon ages of 179 Ma (Dilek and Thy, 2006), indicating that part of the Neotethyan oceanic crust preserved in the mélange is as old as the Early Jurassic. The radiolarian fauna in the chert blocks have yielded Late Carnian–Middle Norian, and Middle Jurassic to Middle Cretaceous ages (Sarifakioglu et al., unpublished data). However, the whole–rock  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of basaltic pillow lava from an ophiolitic thrust sheet farther south in the Ankara Mélange has revealed an age of  $80.3\pm7.6$  Ma, indicating that Late Cretaceous oceanic crustal rocks also exist within the mélange (Table 1).

The Senomanian–Santonian flyschoidal sedimentary rocks with pebblestone, sandstone, mudstone and clayey limestone with interbedded chert layers unconformably rest on the ophiolitic rocks (Figs. 3a and 4). However, Kimmeridgian–Hauterivian flyschoidal sedimentary rocks cover the ophiolitic pillow lavas farther south in the Ankara Mélange (Sarifakioglu et al., unpublished data). The Upper Santonian–Maastrichtian, thin– to medium–layered clayey to sandy limestone and volcanic detrital rocks rest unconformably on these flyschoidal sedimentary and ophiolitic rocks around Yapraklı (Çankırı) and Laloğlu (Çorum), and represent the forearc basin strata (Figs. 6 and 7). The ophiolitic, flyschoidal and forearc basin rocks are imbricated along south–directed thrust faults (Sarifakioglu et al., 2011).

Blocks (km–size) of alkaline volcanic and pyroclastic rocks, debris flow deposits, and coarse–grained reefal limestones representing seamount and/or oceanic plateau fragments also occur in the Ankara Mélange (Fig. 8). We have obtained Middle–Upper Triassic and Cretaceous biostratigraphic ages from the reefal limestones overlying the seamount volcanic units, and  $^{40}\text{Ar}/^{39}\text{Ar}$  whole–rock ages of  $96.6\pm1.8$  Ma from the alkaline pillow lavas that are



stratigraphically associated with the pink colored pelagic limestones (Sarifakioglu et al., unpublished data). Rojay et al. (2004) obtained the Late Barremian–Early Aptian biostratigraphic ages from the reefal limestones resting on the pillow lavas with ocean island basalt (OIB) geochemical affinities. Blocks of these neritic carbonates and the underlying alkaline pillow lavas are also embedded in a turbiditic sequence consisting of chert and volcanic rock clasts in a fine-grained sandstone matrix. Volcanic debris flow deposits also occur within the turbiditic sequence.

In addition to the blocks of ophiolitic, seamount and oceanic plateau rocks, the Ankara Mélange also contains blocks of metamorphic rocks, mainly epidote–glaucophane, epidote–chlorite, and epidote–actinolite schists (Fig. 6). The geochemical fingerprinting of these rocks suggests that their protoliths were made of seamount volcanics and ophiolitic basic rocks, and related sediments. Detailed descriptions and documentation of these metamorphic rocks will be presented elsewhere. We interpret these metamorphic rocks to have formed in an intra–oceanic subduction zone. The  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  dating of the epidote–glaucophane, epidote–chlorite and epidote–actinolite schists revealed the cooling ages of  $83.5\pm1.2$  Ma,  $158.4\pm4.2$  Ma, and  $187.4\pm3.7$  Ma, respectively whereas phyllite, actinolite schist and amphibole–epidote schist yielded  $119.8\pm3.3$  Ma,  $177.4\pm5.8$  Ma,  $256.9\pm8.0$  Ma, respectively (Tables 2 and 3).

Overlying the Ankara Mélange tectonically or unconformably are volcanic and volcanoclastic rocks of an island arc origin (Figs. 9 and 10). Nearly 20 km north of Kalecik subalkaline to alkaline volcanic rocks (Dönmez et al., 2009), intercalated with clayey and sandy limestone, calcareous sandstone, pebblestone, sandstone and shale, overlie the Ankara Mélange units and the flyschoidal sedimentary rocks (Hakyemez et al., 1986; Rojay and Süzen, 1997). The volcanic rocks are locally overlain by the Upper Cretaceous reefal limestones and sandstones containing rudist fossils (Fig. 10a and b). Both pillowed and massive lava flows with cooling joints occur (Figs. 9c and 10d); the massive lava flows contain cm–size augite and leucite phenocrysts. Mafic dikes locally crosscut the volcanoclastic rocks of the arc sequence (Fig. 9d). The  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  whole–rock dating of an arc–related pillow lava has yielded an age of  $67.8\pm4.9$  Ma (Table 4a).

Lamprophyre dikes and a syeno–diorite pluton of an island arc origin are intruded into the ophiolitic and seamount rocks and the mélange matrix along the Kizilirmak River near and east of Kalecik (Fig. 11). The brownish grey colored lamprophyric dikes continue along–strike for 200 to 1000 m, and are displaced by local thrust faults. The  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  whole–rock dating

and the  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  biotite age from the lamprophyric dikes revealed ages of  $67.2\pm1.2$  Ma and  $63.6\pm1.2$  Ma, respectively (Table 4b and c).

We have also obtained an  $^{40}\text{Ar}$ – $^{39}\text{Ar}$  biotite age of  $75.9\pm1.3$  Ma from a syeno–dioritic pluton, approximately 1 km in diameter, indicating that the arc magmatism started as early as the Campanian and that it progressed with alkaline volcanism and dike emplacement throughout the Maastrichtian and Early Paleocene (Table 4d). Andesitic lavas and volcaniclastic and pyroclastic rocks are intercalated with the Upper Cretaceous–lower Paleocene turbiditic rocks in the region. These turbiditic and flyschoidal rocks contain volcanic pebbles in the lower stratigraphic levels and grade upwards into sandstone and shale. The Paleocene rocks (Dizilitaslar Formation) are conformably overlain by the lower to Middle Eocene sandstone, shale, clayey limestone and marl units that collectively make up the Mahmutlar Formation (Akyürek et al., 1984). All these Paleogene sedimentary rocks were deformed by south–vergent thrust faults and folds, indicating that they underwent N–S–directed contractional deformation in the Middle to Late Eocene.

#### **4. Petrography**

In this section we describe the primary and secondary mineral assemblages and the textures of the main lithological types associated with the Neotethyan oceanic crust, seamount volcanic units, and island arc assemblages (e.g. volcanic rocks, lamprophyre dikes and syeno–diorite plutons) that we investigated in the study area.

##### **4.1. Basalt**

The seamount-related alkaline basaltic rocks consist mainly of plagioclase (55–60%) and clinopyroxene (approximately 40%), displaying an intergranular texture (Fig. 12a). Some of the basalt samples contain olivine phenocrysts (about 15%) ranging in size from 0.2 mm to 2 mm. Clinopyroxene grains (titanaugites) are partially altered into chlorite, olivine to serpentine and iddingsite, and plagioclase to sericite and chlorite. Apatite and opaque minerals (Fe–Ti oxide) occur as accessory minerals. Amygdals are filled with secondary carbonate and chlorite minerals.

Tholeiitic basaltic rocks of the Neotethyan oceanic crust comprise microlitic plagioclase and clinopyroxene crystals in a fine–grained texture (Fig. 12b). They are partially or completely spilitized, with plagioclase replaced by albite, sericite, chlorite and epidote (saussuritization), whereas clinopyroxene replaced by actinolite (uralitization) and chlorite. The glassy material

in the matrix is transformed into chlorite. Leucoxene and opaque minerals are present as accessories. Vesicles in the basaltic lavas are filled by secondary carbonate and chlorite.

The island-arc basaltic rocks consist mainly of plagioclase (about 55%) and clinopyroxene (45%) crystals in the porphyritic textures with chloritized glassy and microcrystalline groundmass. Clinopyroxene (diopside) grains range in size from 0.2 mm to 2 mm in length (Fig. 12c and d), and locally display twinning. The plagioclases are partly altered to chlorite and carbonate minerals. Accessory minerals are made of fine crystalline Fe–Ti oxides. Basaltic andesites contain plagioclase, clinopyroxene, minor olivine and biotite within porphyritic and glomeroporphyritic textures (Fig. 12e). Ferromagnesian minerals are locally 1.5 cm-long. Fe–Ti oxide minerals are accessories. The groundmass consists of plagioclase microlites, and chloritized and/or devitrified glass. Basaltic lavas include vesicles filled by secondary carbonate, chlorite, and zeolite.

#### **4.2. Basanite**

The ultrabasic volcanic rocks consist of clinopyroxene, plagioclase and minor olivine occurring as euhedral and subhedral grains in a hyalomicrolitic, porphyritic texture. Plagioclase forms microlites or micro–phenocrysts, and is commonly altered to clay minerals. Clinopyroxene is mainly augite, and displays zoning and twinning. Olivine is surrounded by a groundmass that is made entirely of serpentine minerals. Small analcime crystals occur as a replacement of leucite between plagioclase and clinopyroxene crystals within the groundmass.

#### **4.3. Tephrite**

This fine–grained basaltic rock comprises clinopyroxene (augite), leucite, rare olivine and black mica (phlogopite) crystals within a hyalomicrolitic or porphyritic texture. Plagioclase microlites, ultra fine–grained clinopyroxene, phlogopite, leucite and glassy material form the groundmass, whereas clinopyroxene and leucite occur as euhedral to subhedral microphenocrysts. The leucite contents in the leucite-tephrite rock are up to ~25% (Fig. 12d). Small, anhedral or subhedral opaque minerals are found as accessory minerals.

Some tephrites display characteristic features of phonolitic tephrite with feldspar crystals (plagioclase > K–feldspar) and mafic minerals (phlogopite, hornblende) in a microcrystalline porphyritic texture. Plagioclase is partially altered to sericite and chlorite, whereas sanidine is partially altered to sericite and clay minerals. Leucite occurs as subhedral grains, is mostly altered to sanidine microlites, zeolite and clay minerals, and is surrounded by small

phlogopite flakes. Euhedral apatite crystals and anhedral opaque minerals (Fe–Ti oxides) are present as accessories.

#### **4.4. Lamprophyre**

These alkaline dike rocks consist mainly of small prismatic clinopyroxene (diopside), minor phlogopite and leucite pseudomorphs embedded in a groundmass composed of feldspars (orthoclase > plagioclase), analcime crystals and glassy material (Fig. 12f and g). Both plagioclase and orthoclase are partly or completely altered to carbonate, clay and zeolite minerals; phlogopite is replaced by chlorite along its rims. Small, interstitial apatite laths are enclosed in the orthoclase crystals. In addition, euhedral prismatic apatite crystals up to 0.7 mm in length are also present in the groundmass. Opaque minerals occur as accessory crystals.

#### **4.5. Syeno–diorite**

The main minerals in this intrusive rock include feldspar (plagioclase  $\geq$  orthoclase), clinopyroxene, hornblende and biotite (Fig. 12h). Subhedral to anhedral plagioclase crystals form a granular texture; some large orthoclase crystals (~2.5 cm) locally give the rock a porphyry texture. Plagioclase grains (An<sub>28</sub>–An<sub>48</sub>) are locally surrounded by orthoclase. K–feldspar grains display a perthitic texture. Subhedral to anhedral clinopyroxene (diopside), hornblende and biotite crystals show partial chloritization. Subhedral hornblende crystals have opacite rims around them as a result of metasomatism during their reaction with melt (Plechov et al., 2008). The subhedral prismatic apatite and anhedral granular opaque minerals are present as accessories.

### **5. Analytical Methods**

We analyzed fifty–one (51) rock samples for major, trace, and rare–earth element chemistry at ACME Analytic Laboratory (Canada). Inductively coupled plasma–emission spectroscopy has been used for major–element analysis, and inductively coupled plasma–mass spectroscopy has been used for the analysis of both trace elements and rare–earth elements (REE). The results of these analyses are presented in Tables 5, 6, 7, 8 and 9.

<sup>40</sup>Ar/<sup>39</sup>Ar age dating was done at the Geochronology and Isotopic Geochemistry Laboratory of Activation Laboratories Ltd. (Actlabs), Ancaster, Ontario, Canada. We obtained <sup>40</sup>Ar/<sup>39</sup>Ar ages of biotite separates from two samples of the arc rocks. In addition, whole rock fractions of five rock samples were analyzed. The samples wrapped in Al foil was loaded in evacuated

and sealed quartz vial with K and Ca salts and packets of LP-6 biotite interspersed with the samples to be used as a flux monitor. The sample was irradiated in the nuclear reactor for 48 hours. The flux monitors were placed between every two samples, thereby allowing precise determination of the flux gradients within the tube. After the flux monitors were run, J values ( $n \times 10^{-10}$  cc STP) were then calculated for each sample, using the measured flux gradient. LP-6 biotite has an assumed age of 128.1 Ma. The neutron gradient did not exceed 0.5% on sample size. The Ar isotope composition was measured in a Micromass 5400 static mass spectrometer. 1200°C blank of  $^{40}\text{Ar}$  did not exceed  $n \times 10^{-10}$  cc STP.

Argon is extracted from the sample as degassing at  $\sim 100^\circ\text{C}$  during two days in double vacuum furnace at  $1700^\circ\text{C}$ . Argon concentration is determined using isotope dilution with  $^{38}\text{Ar}$  spike, which is introduced to the sample system prior to each extraction. The obtained pure Ar is introduced into custom build magnetic sector mass spectrometer (Reinolds type) with Varian CH5 magnet. Measurement Ar isotope ratios is corrected for mass-discrimination and atmospheric argon assuming that  $^{36}\text{Ar}$  is only from the air. After each analysis the extraction temperature is elevated to  $1800^\circ\text{C}$  for few minutes. Then, Aliquot of the sample is weighted into graphite crucible with lithium metaborate/tetraborate flux and fused using LECO induction furnace for K-analysis. The fusion bead is dissolved with acid. Standards, blanks and sample are analyzed on Thermo Jarrell Ash Enviro II ICP Spectrometer.

The Sr, Nd, and Pb isotopic compositions of six samples from the alkaline lamprophyric dikes have been determined at the ACT Analytical Laboratories Ltd., Canada (Table 9). The Sr isotope analysis was performed with a Triton multi-collector mass-spectrometer in static mode. The weighted average of 15 SRM-987 Sr-standard runs yielded  $0.710258 \pm 9$  (2s) for  $^{87}\text{Sr}/^{86}\text{Sr}$ . Sm and Nd were separated by extraction chromatography on hexyl di-ethyl hydrogen phosphate-covered Teflon powder. The analysis was performed on a Triton multi-collector mass spectrometer in static mode.  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios are relative to the value of 0.511860 for the La Jolla standard. Pb was separated using the ion-exchange technique with Bio-Rad 1x8. Pb isotope compositions were analyzed on Finnigan MAT-261 multicollector mass spectrometer. The measured Pb isotope ratios were corrected for mass fractionation calculated from replicate measurements of Pb isotope composition in the National Bureau of Standards SRM – 982 standards. External reproducibility of lead isotope ratios –  $^{206}\text{Pb}/^{204}\text{Pb} = 0.1\%$ ,  $^{207}\text{Pb}/^{204}\text{Pb} = 0.1\%$ ,  $^{208}\text{Pb}/^{204}\text{Pb} = 0.2\%$  – on the  $2\sigma$  level has been demonstrated through multiple analyses of standard BCR-1.

## 6. Geochemistry

We report below on the geochemistry of the representative samples of oceanic basaltic rocks in the Ankara Mélange, as well as the lamprophyric dikes, a syeno–dioritic pluton, and alkaline lavas that crosscut and/or cover the blocks of volcanic and volcanoclastic rocks, serpentinite, radiolarian chert, and shale in the Ankara Mélange.

## 6.1. Oceanic Basaltic Rocks

The  $\text{Na}_2\text{O}+\text{K}_2\text{O}$  values of basaltic blocks of the Neotethyan oceanic crust range from 1 wt% to 4.28 wt%, with the  $\text{K}_2\text{O}$  values much lower than those of  $\text{Na}_2\text{O}$  (Table 5). The Na enhancement of two samples (CE.07, CE.08) may be a result of spilitization caused by low–grade hydrothermal ocean floor metamorphism. Similarly, the total alkali values from the seamount volcanic blocks vary between 4.72 and 8.14 wt%, with the  $\text{Na}_2\text{O}$  values (3.78–6.79 wt%) much higher than that of oceanic crust (Table 6).

On the total alkali vs. silica (TAS) diagram the tholeiitic–calcalkaline volcanic and isolated dike rocks from the Tethyan oceanic crust fall in the field of basalt and basaltic andesite, whereas the samples of seamount alkaline rocks plot in the basanite, tephrite ( $\text{SiO}_2 = 39.77\text{--}46.36$  wt%), trachyte ( $\text{SiO}_2 = 68.47$  wt%), trachybasalt ( $\text{SiO}_2 = 50.15$  wt%) and foidite ( $\text{SiO}_2 = 39.77$  wt%) fields (Fig. 13a and b). The oceanic basalt samples have lower  $\text{TiO}_2$  values (0.26–1.74 wt%) in comparison to the alkaline, seamount volcanic rocks (1.64–2.46 wt%), except for a volcanic sample with tholeiitic OIB (Ocean–Island basalt) characteristics. On a Ti–Zr–Y discrimination diagram (Pearce and Cann, 1973), the oceanic basalt samples plot in the MORB (mid–ocean ridge basalt) and island arc tholeiite (IAT) fields, whereas the seamount volcanic rocks generally fall in the within–plate alkali basalt field (except a trachyte sample; Fig. 13c). On a Ti–V diagram (Shervais, 1982), the samples of oceanic basaltic rocks mostly plot in the MORB field ( $\text{Ti/V} = 22.6\text{--}28.9$ ), whereas four samples have island arc tholeiite to boninitic affinities ( $\text{Ti/V} = 5.4\text{--}25.55$ ) (Fig. 13d). The samples of silica–undersaturated, seamount volcanic rocks display an OIB–character with high Ti/V ratios (62.6–261.2).

The N–MORB normalized multi–element diagrams of the representative samples of basalts of oceanic crust and seamount volcanic rocks are shown in Fig. 13e. Basaltic samples of both MORB and SSZ (suprasubduction zone) affinities show enrichment in their LILE (the large ion lithophile elements: Rb, Ba, K, Sr, Cs, Th) contents. The HFSE (high field strength elements: Nb, Ta, Zr, Hf, Ti, Y) and REE (rare earth elements) contents of the MORB–type basaltic rocks display a slight increase, whereas the SSZ–related basaltic rocks (four samples) exhibit depletion in HFSE and REE. The LILE, HFSE, LREE (light–REE) contents

of the seamount volcanic rocks are extremely enriched relative to the HREE (heavy-REE) values. Also, the Th/Yb (2.8–5.6) and Nb/Yb (27.6–54.8) values of the seamount volcanic rocks are high in comparison to those of the Neotethyan oceanic basalt samples (Th/Yb=0.2–1.1; Nb/Yb=0.7–2.7). However, the alkaline lava samples have the ratios of Nb/Y>1.5 and Zr/Nb<6 that are typical for within-plate basalts (Edwards et al., 1991). The seamount volcanic rocks have Nb/Y ratios of 2.3–3.1 and Zr/Nb ratios of 3.1–4.1, indicating OIB-like geochemical characteristics, whereas the oceanic crust basalt samples have Nb/Y (0.1–0.4) and Zr/Nb (8.1–32) values.

## 6.2. Island Arc Rocks

A small syeno-diorite pluton, a suite of volcanic rocks, and lamprophyric dikes in the Kalecik (Ankara) area collectively represent the products of island arc magmatism. These arc rocks mostly plot in the alkaline field on a TAS diagram (Fig. 14a and b). The alkaline rock samples with medium to high Al<sub>2</sub>O<sub>3</sub> contents (10–19 wt%) represent both silica-saturated and silica-undersaturated rock units (Tables, 7, 8 and 9). The lamprophyric dikes have picrobasalt, trachybasalt, trachyandesite, tephrite and phonotephrite compositions, whereas the volcanic rocks display basalt, basanite, tephrite, leucite tephrite and foidite compositions. The samples from small alkaline intrusions fall into the syeno-diorite field in the TAS diagram (Fig. 14b; Cox et al., 1979). The Late Cretaceous–Early Paleocene volcanic rocks (andesite, dacite, rhyolite), found nearly 60 km SW of Kalecik, show subalkaline (tholeiitic and calc-alkaline) compositions, except for a few trachytbasalt and trachyandesite samples (Fig. 14a, c and d; Dönmez et al., 2009).

The alkaline volcanic rocks mostly display high-K shoshonitic compositions in the K<sub>2</sub>O vs. SiO<sub>2</sub> diagram (Fig. 14d; Peccerillo and Taylor, 1976). Some volcanic and dike rocks also plot in the fields of medium– high-K, calc-alkaline series. Although some alkaline volcanic rocks show medium-K calc-alkaline characteristics as a result of hydrothermal alteration (LOI/loss on ignition>2wt%), they have high-K shoshonitic affinity since the leucite bearing, silica-undersaturated alkaline rocks experienced analcimization resulting in low K<sub>2</sub>O values in favor of Na<sub>2</sub>O values. On the Hastie et al. (2007), and Pearce (1982) diagrams, which utilize the immobile elements and the ratios of immobile elements (Th vs. Co, and Ce/Yb vs. Ta/Yb), the arc-related plutonic, volcanic and dike rocks generally display high-K (K<sub>2</sub>O/Na<sub>2</sub>O = 1.5–3.4) and shoshonitic characteristics (Fig. 14e and f). However, seven samples from the volcanic rocks and lamprophyre dikes contain high K<sub>2</sub>O/Na<sub>2</sub>O ratios (18.16–24.52) showing ultrapotassic (K<sub>2</sub>O/Na<sub>2</sub>O>3) characteristics.

When plotted on MgO vs. major element diagrams, the analyzed samples mainly exhibit negative correlations, except on the  $\text{Fe}_2\text{O}_3$  and  $\text{TiO}_2$  plots, which show positive correlations (Fig. 15). Based on the MgO vs. trace element variation diagrams (Fig. 15), Co shows a positive trend while Ba, Rb, Sr, Th and Zr all exhibit negative trends. These major and trace element trends can be explained by fractionation of clinopyroxene, feldspar, black mica (biotite, phlogopite), Fe–Ti oxides and apatite. However, the scatter in Fig. 15 may also be caused by the alteration of the arc rocks and/or the involvement of subducted sediments in their melt regime. The rock samples from the small syeno–diorite pluton with metaluminous characteristics plot in the VAG (volcanic arc granites) field (Fig. 16a, b and c). The Ti–Zr–Y and Ti–V diagrams (Pearce and Cann, 1973; Shervais, 1982) show that the alkaline basic samples and the subalkaline volcanic rocks (Dönmez et al., 2009) from the southwestern part of the study area all plot in the arc field (Fig. 16d and e), whereas the  $\text{TiO}_2$ – $\text{Al}_2\text{O}_3$  and Y–Zr diagrams (Muller et al., 1992) show that these samples fall into the arc field (Fig. 16f and g). The analyzed alkaline rocks display shoshonitic characteristics in the Th/Yb vs. Nb/Yb diagram (Pearce, 2008), and their Hf/Th ratios are rather low ranging from 0.11 to 0.57, consistent with their shoshonitic affinity. The island–arc tholeiitic (IAT) basaltic rocks have  $\text{Hf/Th} > 3$ , whereas the calc–alkaline volcanic rocks have  $\text{Hf/Th} < 3$  (Wood, 1980). Their Th enrichment and increased Th/Yb ratios along the mantle metasomatism trend indicate the influence of subduction–derived fluids in their magma source (Fig. 16h; Pearce, 2008). The samples derived from the blocks of N–MORB–, SSZ– and OIB–like oceanic basalts in the Ankara Mélange typically plot within the MORB–OIB mantle array (Fig. 16h).

The primitive mantle–normalized, multi–element diagrams of the representative samples from the high–K shoshonitic arc rocks around Kalecik (Ankara), Yapraklı (Çankırı) and Laloğlu (Çorum) are plotted in Fig. 17a. The trace element patterns of all the analyzed alkaline rocks display strong enrichment of the LILE, LREE and also Pb, U in comparison to HFSE (Nb, Ta, Zr, Hf, Ti, Y), which show negative anomalies indicating subduction zone influence (Kempton et al., 1991). The high Ba/Ta ( $> 450$ ) and Ba/Nb ( $> 28$ ) ratios are characteristic features of subduction–related magmas (Fitton et al., 1988). The very high ratios of Ba/Ta (383–5255), Ba/Nb (64–538), and relatively high Zr/Nb (5–22), Th/Yb (2–14), Zr/Y (3–7) and La/Yb (9–36) have been attributed to a mantle source, which was enriched by a subduction component (Frey et al., 1978; Fitton et al., 1988; Maury et al., 1992; Schiano et al., 1995). However, some of the lamprophyre dike samples (DM.2, DM.6, DM.8, DM.9, DM.10) contain La/Yb ratios of 30, indicating highly undersaturated magmas for their origin. Also, the alkaline rocks with  $\text{Mg\#} < 61$ , except for one sample ( $\text{Mg\#} = 71$ ),  $[\text{MgO}/(\text{MgO} + 0.8 \cdot \text{FeO total})]$ , imply that none of these shoshonitic rocks represents primary mantle-derived



subduction-related magmas. However, their chondrite-normalized REE patterns (Fig. 17b) show LREE enrichment, flat HREE ( $\text{La}/\text{Sm}_n=2.18\text{--}5.71$ ;  $\text{Gd}/\text{Lu}_n=1.69\text{--}4.14$ ;  $\text{La}/\text{Lu}_n=6.57\text{--}24.72$ ), and minor negative Eu anomalies ( $\text{Eu}/\text{Eu}^*=0.77\text{--}0.95$ ). These geochemical characteristics are compatible with those defining subduction-related, arc volcanic assemblages (Tatsumi et al., 1986; Kelemen et al., 1993; Hawkesworth et al., 1993; Pearce and Peate, 1995).

The high-K shoshonitic lamprophyric dikes are characterized by intermediate  $^{143}\text{Nd}/^{144}\text{Nd}$  ( $0.512674\text{--}0.512690$ ) and  $^{87}\text{Sr}/^{86}\text{Sr}$  ( $0.704697\text{--}0.704892$ ) isotopic compositions. The initial  $\epsilon_{\text{Nd}}$  values range from +1.3 to +1.7, whereas the modern  $\epsilon_{\text{Nd}}$  values vary between +0.7 and +1.0 indicating a relatively enriched mantle source. Their Pb isotope ratios range from 19.332 to 19.939 for  $^{206}\text{Pb}/^{204}\text{Pb}$ , 15.655 to 15.691 for  $^{207}\text{Pb}/^{204}\text{Pb}$ , and 39.192 to 39.612 for  $^{208}\text{Pb}/^{204}\text{Pb}$ . The high  $^{206}\text{Pb}/^{204}\text{Pb}$ , and relatively high  $^{143}\text{Nd}/^{144}\text{Nd}$  and  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios seem to be compatible with a mantle source that is enriched by slab-derived fluids and/or subducted pelagic sediments. Rock (1977; 1984) described shoshonitic lamprophyres (minette, kersantite, vogesite, spessartite) as mildly potassic alkaline rocks ( $\text{Na}<\text{K}$ ;  $\text{SiO}_2 \approx 53$  wt%), indicating their magma source to be hybrids between basic magma and granitic residua or crustal sediments. Also, alkaline lamprophyres (camptonite, monchiquite, sannaite) with mantle-type  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios derived from a lamprophyre magma by hydrous crystallization of basaltic magma (Rock, 1977).

## 7. Discussion

### 7.1. Source Characteristics

The subduction-accretion complex represented by the Ankara Mélange contains blocks of oceanic lithosphere showing geochemical affinities ranging from MORB to IAT and calc-alkaline. The SSZ-type ophiolite assemblages in the melange display both IAT-like and boninitic geochemical signatures. The ophiolitic units with an IAT-like chemistry are the manifestation of partial melting of the upper mantle peridotites, which were modified by incompatible element-enriched hydrous fluids (or melt) released from the subducting Tethyan oceanic slab. The ophiolitic units with MORB-like signatures represent the products of a depleted mantle source. Some of the samples with MORB-like chemistry plot within or near the IAT field (Figs. 13c and 16h) indicating that their magmas were influenced by subduction-derived fluids. These ophiolitic rocks are the oldest units as constrained by the volcanic stratigraphy and crosscutting relationships. Some doleritic dikes and basaltic rocks in the ophiolites show boninitic affinities, consistent with their formation in a forearc setting

(Dilek and Furnes, 2011; Sarifakioglu et al., 2011). Collectively, the ophiolitic units in the Ankara mélangé display a geochemical progression that is typical of the development of forearc oceanic crust in the early stages of subduction-induced magmatism, as also documented from other Tethyan ophiolites (Dilek and Furnes, 2009, 2011; Dilek and Thy, 2009; Pearce and Robinson, 2010; Sacconi et al., 2011; Moghadam et al., 2013).

Seamount volcanic rocks occurring in the Ankara mélangé have OIB-like geochemical features, showing tholeiitic to alkaline affinities (Fig. 13e) with enrichment in incompatible elements and LREEs. The tholeiitic OIB affinity of some of the seamount volcanic rocks may have resulted from the interaction of plume-derived melts with MORB-type melts near a seafloor spreading system. The depletion of the OIB-type volcanic rocks in immobile elements (especially Ti) suggests mixing of the plume and MORB-type melts during seamount evolution.

The high-K alkaline rocks exhibit LILE and HFSE enrichments and negative Nb, Ta, Hf, Zr, Ti anomalies, indicating strong subduction influence in their melt evolution (Fig. 17a). The high ratios of LILE/HFSE ( $Ba/Nb = 64\text{--}538$ ;  $Ba/Ta = 383\text{--}5255$ ;  $Rb/Nb = \sim 2\text{--}20$ ), LREE/HFSE ( $La/Nb = 1.8\text{--}7.2$ ;  $La/Ta = 48\text{--}188$ ;  $La/Sm_n \sim 4$ ), LILE/LREE ( $Th/La = 0.16\text{--}0.49$ ) and  $Zr/Nb$  ( $5\text{--}22$ ), and the large negative Nb-Ta anomaly in the multi-element diagrams all point to a melt source affected by subduction-generated fluids and/or crustally contaminated magmas. The observed high  $Ba/Nb$  ( $64\text{--}538$ ),  $La/Yb$  ( $9\text{--}36$ ),  $Sr/Nd$  ( $14\text{--}45$ ) and  $Ce/Yb$  ( $20\text{--}73$ ) ratios, and low  $Nb/U$  ( $2\text{--}7$ ),  $Ba/La$  ( $20.02\text{--}59.83$ ),  $U/Th$  ( $0.13\text{--}0.50$ ) and  $Ce/Pb$  ( $\sim 2\text{--}20$ ) values indicate that the mantle melt source may have been modified by some melts derived from relatively incompatible element-rich, subducted pelagic and/or terrigenous sediments. In contrast, the high  $Ce/Pb$  ( $25\text{--}5$ ) and  $Nb/U$  ( $47\text{--}10$ ) ratios observed in the OIB-type seamount volcanic rocks indicate that the magmas of these rocks were not modified by subducted sediments (Hoffman et al., 1986).

Enrichments in Cs, Rb, Ba, Th, U, K, La, Ce and Pb of the alkaline rocks suggest that their melt source was modified by subducted slab material (mainly fluids, and pelagic and/or terrigenous sediments). Slab-derived fluids helped to form hydrous and K-rich minerals, such as amphibole, apatite and phlogopite with high  $Rb/Sr$  ( $0.04\text{--}0.71$ ) and  $K/Ti$  ( $3.77\text{--}16.62$ ) ratios relative to MORB- and OIB-like magmas, and resulted in a positive correlation between  $Ba/Nb$  and  $La/Nb$  ratios (Fig. 18a). Also, the high La ( $18.4\text{--}69.2$  ppm) contents and  $La/Yb$  ratios ( $9.5\text{--}34.6$ ) reflect that the high-K magmas may have been produced by small degrees of partial melting of a subduction-metasomatised mantle source (Fig. 18b).

470 As illustrated in the  $^{143}\text{Nd}/^{144}\text{Nd}$  vs.  $^{87}\text{Sr}/^{86}\text{Sr}$  diagram (Fig. 19a), six lamprophyre samples plot  
 471 on the mantle array defining a subduction component during the evolution of their magmas.  
 472 We also show in this diagram, for comparison, the Late Cretaceous–Early Tertiary volcanic  
 473 rocks from the southern part of Central Anatolia and the Eastern Pontides, and the Cenozoic  
 474 volcanic units in Western Anatolia (Alpaslan et al., 2004; 2006; Eyüboğlu, 2010; Altunkaynak  
 475 and Dilek, 2006 and references therein). The relatively high Pb (up to 34 ppm in some  
 476 samples) and  $^{87}\text{Sr}/^{86}\text{Sr}$  contents, and the Rb/Sr ratios (0.02–0.71) of the lamprophyre rocks  
 477 also indicate the effects of subducted oceanic sediments added to the mantle melt source  
 478 (Pearce and Peate, 1995). In the  $^{206}\text{Pb}/^{204}\text{Pb}$  vs.  $^{208}\text{Pb}/^{204}\text{Pb}$ ,  $^{206}\text{Pb}/^{204}\text{Pb}$  vs.  $^{207}\text{Pb}/^{204}\text{Pb}$ ,  
 479  $^{87}\text{Sr}/^{86}\text{Sr}$  vs.  $^{206}\text{Pb}/^{204}\text{Pb}$  and  $^{143}\text{Nd}/^{144}\text{Nd}$  vs.  $^{206}\text{Pb}/^{204}\text{Pb}$  variation diagrams, the data points lie  
 480 above the Northern Hemisphere Reference Line (NHRL), and the radiogenic isotope data fall  
 481 close to the fields of MORB, enriched lithospheric mantle source (EMII) and oceanic  
 482 sediments. These features collectively suggest that the magmas of the lamprophyre rocks  
 483 were derived from a MORB-like mantle source that was enriched by subducted terrigenous  
 484 and carbonate sediments (Fig. 19b–e). However, the post-collisional Late Cretaceous–Early  
 485 Tertiary volcanic rocks in the Ulukisla basin in the southern part of Central Anatolia have  
 486 higher  $^{87}\text{Sr}/^{86}\text{Sr}$  and lower  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios than those of the lamprophyres in the Ankara  
 487 Mélange, indicating an EMI with recycled, continent-derived material. The late Cretaceous  
 488 high-K volcanic rocks representing active continental margin arc units in the Eastern  
 489 Pontides with low  $^{87}\text{Sr}/^{86}\text{Sr}$  reflect a mantle source enriched by continental crustal rocks. The  
 490  $^{143}\text{Nd}/^{144}\text{Nd}$ ,  $^{87}\text{Sr}/^{86}\text{Sr}$ ,  $^{206}\text{Pb}/^{204}\text{Pb}$ ,  $^{208}\text{Pb}/^{204}\text{Pb}$  and  $^{207}\text{Pb}/^{204}\text{Pb}$  values reflecting subduction  
 491 enrichment and crustal contamination of the source of the post-collisional, Middle Eocene  
 492 volcanic units in Central Anatolia and the Tertiary volcanic suites in western Anatolia have  
 493 been explained by slab breakoff-induced asthenospheric upwelling and associated partial  
 494 melting of the orogenic lithospheric mantle (Alpaslan et al., 2004, 2006; Altunkaynak and  
 495 Dilek, 2006, 2013; Dilek and Altunkaynak, 2007; Keskin et al., 2008; Gündoğdu–Atakay,  
 496 2009; Sarifakioglu et al., 2013).

497 The depletion of HFSE with respect to LREE enrichment, and high LILE/HFSE and  
 498 radiogenic isotope ratios suggest that the high-K shoshonitic rocks are likely to have formed  
 499 by small degrees of partial melting of a lithospheric mantle modified by slab-derived hydrous  
 500 fluids.

## 7.2. Tectonic Model

The Ankara Mélange displays a heterogeneous structural architecture containing oceanic and crustal rocks with different internal structure, stratigraphy and geochemical compositions. The oldest ophiolitic rocks in the Ankara Mélange appear to have formed in a SSZ setting within the Northern Tethys around 180 Ma (Dilek and Thy, 2006; Sarifakioglu et al., 2011). The ~80 Ma ( $80.3 \pm 7.6$  Ma) ophiolitic rocks in the same mélange also indicate that oceanic crust formation in the Northern Tethys was still in operation in the Late Cretaceous (Table 1).

We obtained Middle–Upper Triassic biostratigraphic age data from the neritic limestones that are spatially associated with the seamount volcanic rocks, indicating that an oceanic lithosphere of the Late Triassic and older ages must have existed in this ocean to make up the substratum of the seamounts. Thus, we know that the northern branch of Neotethys was already a wide–open ocean with its MORB–type oceanic lithosphere between the Pontide block to the north and the Anatolide–Tauride micro–continent to the south in the Early Triassic (or even before). The ophiolitic mélange units in the Kırıkkale–Ankara–Çankırı–Çorum area are unconformably overlain by basal volcanic conglomerates of an arc origin. The overlying volcanosedimentary units contain clayey– and sandy–limestone, limey sandstone, and sandstone–claystone alternating with volcanoclastic rocks. These rock types and their internal stratigraphy suggest their deposition in a frontal arc–forearc basin. The clayey limestones are intruded by dikes and sills and have Late Santonian, and Campanian–Maastrichtian ages based on their fossil contents (Sarifakioglu, unpublished data). The radiometric age data from an alkaline basaltic rock (YK.4) and a syeno–diorite intrusion (YK.438) give ages of  $67.8 \pm 4.9$  Ma and  $75.9 \pm 1.3$  Ma, respectively (Table 4a and d), constraining the timing of intra–oceanic arc magmatism as the Latest Cretaceous.

In general, subalkaline (tholeiitic and calcalkaline) volcanic arc rocks occur in the northern part of the study area, whereas the younger alkaline volcanic and plutonic rocks in the south. We interpret this spatial and temporal relationship to have resulted from a southward progression of the arc magmatism from subalkaline to alkaline affinities through time due to arc rifting above the southward retreating Tethyan subduction system (Fig. 20). We, therefore, think that the arc–related late alkaline dikes and plutons were emplaced on and across the evolving subduction–accretion complex above the north–dipping, southward rolling Tethyan slab.

The high-K and shoshonitic Eocene dikes and lavas in the Ankara mélange formed from melts derived from partial melting of the metasomatized arc mantle that was triggered by the influx of slab breakoff-induced asthenospheric flow. This slab breakoff was a result of an arc-continent (Central Anatolian Crystalline Complex – CACC) collision, followed by the continent-continent collision (Sakarya and CACC) in the Early to Middle Eocene.

## 8. Conclusions

1. Blocks of Middle-Late Triassic seamount and Upper Permian metamorphic rocks occurring in the Ankara Mélange represent an intra-oceanic subduction-accretion complex that developed in the Northern Tethys during the late Paleozoic through Cretaceous.
2. Thrust sheets and/or megablocks containing SSZ ophiolite units with Liassic and Cretaceous ages were incorporated into this subduction-accretion complex during the early Late Cretaceous.
3. The Late Cretaceous tholeiitic to calc-alkaline volcanic rocks are the products of an intra-oceanic island arc system. The tholeiitic and calc-alkaline arc rocks show enrichment in incompatible elements due to the influence of slab-derived fluids. The shoshonitic arc rocks representing the latest stage of island arc magmatism were produced by partial melting of a subduction-enriched mantle source.

## Acknowledgements

This study was supported by a grant from the General Directorate of Mineral Research and Exploration of Turkey (MTA, Ankara; Project no: 2007–2008.30.1601.d–f). We thank Dr. Yakov Kapusta (ACTLabs, Canada) for the radiometric and isotopic analyses of our rock samples in a timely fashion, and for his help with the interpretation of the obtained data. We extend our appreciation to the referee, Assoc. Prof. Ibrahim Uysal (Blacksea Technical University, Turkey) and Prof. Dr. Harald Furnes (Bergen University, Norway). Special thanks are due to Dr. Francesca Funicello for her editorial suggestions, which improved the manuscript. We would like to acknowledge the Head of the Geological Research Department, Dr. Erol Timur, in MTA-Ankara for his continued support of our *Ophiolite Inventory Project* in Turkey over the last two years.

## References

- Akyürek, B., Bilginer, E., Akbaş, B., Hepşen, N., Pehlivan, Ş., Sunu, O., Soysal, T., Dağ, Z., Çatal, E., Sözeri, B., Yıldırım, H. and Hakyemez, Y.: Fundamental geological characteristics of Ankara–Elmadağ–Kalecik area. *Journal of Geological Engineering (Jeoloji Mühendisliği Dergisi)*, Chamber of Geological Engineers of Turkey (TMMOB Jeoloji Mühendisleri Odası), 20, 31–46, 1984 (in Turkish with English abstract).
- Alpaslan, M., Boztug, D., Frei, R., Temel, A. and Kurt, M.A.: Geochemical and Pb–Sr–Nd isotopic composition of the ultrapotassic volcanic rocks from the extension–related Çamardı–Ulukisla basin, Nigde Province, Central Anatolia, Turkey. *Journal of Asian Earth Sciences* 27, 613–627, 2006.
- Alpaslan, M., Frei, R., Boztug, D., Kurt, M.A. and Temel, A.: Geochemical and Pb–Sr–Nd isotopic constraints indicating an enriched–mantle source for Late Cretaceous to Early Tertiary volcanism, Central Anatolia, Turkey. *International Geology Review* 46, 1023–1041, 2004.
- Altunkaynak, Ş. and Dilek, Y.: Timing and nature of postcollisional volcanism in Western Anatolia and geodynamic implications, in: *Postcollisional Tectonics and Magmatism in the Mediterranean Region and Asia*, edited by: Dilek, Y. and Pavlides, S., Geological Society of America Special Paper 409, 321–351, 2006.
- Altunkaynak, Ş. and Dilek, Y.: Eocene mafic volcanism in northern Anatolia: its causes and mantle sources in the absence of active subduction. *International Geology Review* 55 (13), 1641–1659, 2013.
- Bailey, E.B. and McCallien, W.S.: The Ankara Mélange in Central Anatolia. *Bulletine of Mineral Research and Exploration Institute, Maden Tetkik ve Arama Enstitüsü, Ankara, Turkey*, 40, 12–22, 1950.
- Bailey, E.B. and McCallien, W.S.: The Ankara Mélange and Anatolia Thrust. *Nature* 166, 938–941, 1953.
- Bingöl, E., Akyürek, B. and Korkmazer, B.: Geology of Biga Peninsula and characteristics of Karakaya formation. *Geoscience Congress of 50<sup>th</sup> Anniversary of the Turkish Republic, Ankara, 1975*, the Publications of MTA General Directorate, 70–77 (in Turkish with English abstract), 1975.

- 593 Boztug, D.: S–I–A type intrusive associations: geodynamic significance of synchronism  
594 between metamorphism and magmatism in Central Anatolia, Turkey, in: Tectonics and  
595 Magmatism in Turkey and the Surrounding Area, edited by: Bozkurt, E., Winchester, J.  
596 A., and 10 Piper, J. D. A., Geological Society of London Special Publication, 173, 441–  
597 458, 2000.
- 598 Çoğulu, E., Delaloye, M. and Chessex, R.: Sur l'âge de quelques roches plutoniques acides  
599 dans region d'Eskisehir, Turquie, Archives des Sciences, Genève, 18, 692–699, 1965.
- 600 Cox, K.G., Bell, J.D. and Pankhurst, R.J.: The interpretation of igneous rocks. Allen and  
601 Unwin, London, 450p, 1979.
- 602 Dangerfield, A., Harris, R. Sarifakioglu, E. and Dilek, Y.: Tectonic evolution of the Ankara  
603 Mélange and associated Eldivan ophiolite near Hançili, Central Turkey. Geological  
604 Society of America Special Paper 480, 143–169, 2011.
- 605 Delaloye, M. and Bingöl, E.: Granitoids from western and Northwestern Anatolia:  
606 Geochemistry and modelling of geodynamic evolution. International Geology Review  
607 42, p. 241–268, doi:10.1080/00206810009465081, 2000.
- 608 Dilek, Y. and Altunkaynak, Ş.: Cenozoic Crustal Evolution and Mantle Dynamics of Post–  
609 Collisional Magmatism in Western Anatolia. International Geology Review 49, 431–  
610 453, 2007.
- 611 Dilek, Y. and Thy, P.: Age and petrogenesis of plagiogranite intrusions in the Ankara  
612 Mélange, central Turkey. Island Arc 15, 44–57, 2006.
- 613 Dilek, Y. and Furnes, H.: Ophiolite genesis and global tectonics: Geochemical and tectonic  
614 fingerprinting of ancient oceanic lithosphere. Geological Society of America Bulletin  
615 123, 387–411, doi: 10.1130/B30446.1, 2011.
- 616 Dilek, Y. and Furnes, H.: Structure and geochemistry of Tethyan ophiolites and their  
617 petrogenesis in subduction rollback systems. Lithos 113, 1–20, 2009.
- 618 Dilek, Y. and Thy, P.: Island arc tholeiite to boninitic melt evolution of the Cretaceous Kizildag  
619 (Turkey) ophiolite: Model for multi-stage early arc-forearc magmatism in Tethyan  
620 subduction factories. Lithos 113, 68–87, 2009.
- 621 Dönmez, M., Akçay, A.E., Türkecan, A., Evcimen, Ö., Atakay, E. and Görmüş, T.: Tertiary  
622 Volcanites in Ankara surroundings. Mineral Research and Exploration Institute of

- 623 Turkey (MTA) Report, No. 11164, 116 pp., Ankara, Turkey, 2009 (in Turkish,  
624 unpublished).
- 625 Eyüboğlu, Y.: Late cretaceous high-K volcanism in the eastern pontide orogenic belt:  
626 implications for the geodynamic evolution of NE Turkey. *International Geology Review*  
627 52(2-3), 142-186, 2010.
- 628 Festa, A., Pini, A., Dilek, Y. and Codegone, G.: Mélanges and mélanges-forming processes:  
629 a historical overview and new concepts. *International Geology Review*, 52, 1040-1105,  
630 2010.
- 631 Fitton, J.G., James, D., Kempton, P.D., Ormerod, D.S. and Leeman, W.P.: The role of  
632 lithospheric mantle in the generation of Late Cenozoic basic magmas in the Western  
633 United States. *Journal of Petrology, Special Lithosphere Issue*, 331–349, 1988.
- 634 Frey, F.A., Green, D.H. and Roy, S.D.: Integrated models of basalt petrogenesis: A study of  
635 Quartz tholeiite to olivine melilitites from southeastern Australia utilizing geochemical  
636 and experimental data. *Journal of Petrology* 19, 463–513, 1978.
- 637 Güleç, N.: Rb–Sr isotope data from the Agacoren granitoid (East of Tuz Gölü):  
638 Geochronological and genetical implications. *Turkish Journal of Earth Sciences* 3, 39–  
639 43, 1994.
- 640 Gündoğdu–Atakay, E.: Geological and petrological characteristics of volcanic rocks at  
641 southwest of Çorum and geoarc heological studies at Alaca höyük excavation site. PhD  
642 Thesis, Ankara University, Ankara, Turkey, 194pp., 2009.
- 643 Hakyemez, Y., Barkurt, M.Y., Bilginer, E., Pehlivan, Ş., Can, B., Dağ, Z. and Sözeri, B.:  
644 The geology of Yapraklı–Ilgaz–Çankırı–Çandır surroundings. *Mineral Research and*  
645 *Exploration Institute of Turkey (MTA) Report*, No. 7966, Ankara, Turkey, 1986 (in  
646 Turkish, unpublished).
- 647 Hastie, A.R., Kerr, A.C., Pearce, J.A. and Mitchell, S.F.: Classification of Altered Volcanic  
648 Island Arc Rocks using Immobile Trace Elements: Development of the Th–Co  
649 Discrimination Diagram. *Journal of Petrology* 48 (12), 2341–2357, 2007.
- 650 Hawkesworth, C.J., Gallagher, K., Hergt, J.M. and McDermott, F.: Mantle and slab  
651 contributions in arc magmas. *Annual Review of Earth and Planetary Sciences* 21, 175–  
652 204, 1993.



653 Hoffman, A.W., Jochum, K.P., Seufert, M. and White, W.M.: Nb and Pb in oceanic basalts:  
654 New constraints on mantle evolution. *Earth and Planetary Science Letters* 90, 421–  
655 436, 1986.

656 Ilbeyli, N., Pearce, J.A., Thirwall, M.F. and Mitchell, J.G.: Petrogenesis of collision-related  
657 plutonics in Central Anatolia, Turkey. *Lithos* 72, 163–182, 2004.

658 Irvine, T.N. and Baragar, W.R.A.: A guide to the chemical classification of the common  
659 volcanic rocks. *Canadian Journal of Earth Sciences*, 8, 523–548, doi:10.1139/e71-055,  
660 1971.

661 Kadioglu, Y.K., Dilek, Y. and Foland, K.A.: Slab breakoff and syncollisional origin of the Late  
662 Cretaceous magmatism in the Central Anatolian Crystalline Complex, Turkey, in Dilek,  
663 Y., and Pavlides, S., eds., *Postcollisional tectonics and magmatism in the*  
664 *Mediterranean Region and Asia*. Geological Society of America, Special Paper 409,  
665 381–415, doi:10.1130/2006/2409(19), 2006.

666 Kadioglu, Y.K., Dilek, Y., Güleç, N. and Foland, K.A.: Tectonomagmatic evolution of bimodal  
667 plutons in the Central Anatolian Crystalline Complex, Turkey. *Journal of Geology* 111,  
668 671–690, 2003.

669 Kelemen, P.B., Shimizu, N. and Dunn, T.: Relative depletion of niobium in some arc magmas  
670 and the continental crust: partitioning of K, Nb, La and Ce during melt/rock reaction in  
671 the upper mantle. *Earth and Planetary Science Letters* 120, 111–134, 1993.

672 Kempton, P.D., Fitton, J.G., Hawkesworth, C.J. and Ormerod, D.S.: Isotopic and trace  
673 element constraints on the composition and evolution of the lithosphere beneath the  
674 southern United States. *Journal of Geophysical Research* 96, 13713–13735, 1991.

675 Keskin, M., Genç, C.Ş. and Tüysüz, O.: Petrology and geochemistry of post-collisional  
676 Middle Eocene volcanic units in North-Central Turkey: Evidence for magma generation  
677 by slab breakoff following the closure of the Northern Tethys Ocean. *Lithos* 104,  
678 267–305, 2008.

679 Koçyiğit, A.: An example of an accretionary forearc basin from northern Central Anatolia and  
680 its implications for the history of the Neo-Tethys in Turkey. Geological Society of  
681 America, Special Paper 103, 22–36, 1991.

682 Le Bas, M.J., Le Maitre, R.W., Streckeisen, A. and Zanettin, B.: Achemical classification of  
683 volcanic rocks based on total Alkali–Silica content. *Journal of Petrology* 27, 745–750,  
684 1986.

685 Maury, R.C., Defant, M.J. and Joron, J.L.: Metasomatism of arc mantle inferred from trace  
686 elements in Philippine xenoliths. *Nature* 360, 661–663, doi: 10.1038/360661a0, 1992.

687 Moghadam, H. S., Stern R.J., Chiaradia, M. and Rahgoshay, M.: Geochemistry and tectonic  
688 evolution of the Late Cretaceous Gogher-Baft ophiolite, Central Iran. *Lithos*, 168, 33–  
689 47, doi:10.1016/j.lithos.2013.01.013, 2013.

690 MTA: The geological map of Turkey, 1/500 000 scaled, Geological Survey of Turkey, The  
691 General Directorate of Mineral Research and Exploration, Ankara, Turkey, 2001.

692 Muller, D., Rock, N.M.S. and Groves, D.I.: Geochemical discrimination between shoshonitic  
693 and potassic volcanic rocks in different tectonic settings: a pilot study. *Mineralogy and*  
694 *Petrology* 46, 259–289, 1992.

695 Norman, T.N.: The role of the Ankara Melange in the development of Anatolia (Turkey), in:  
696 The Geological Evolution of the Eastern Mediterranean, edited by: Dixon, J. E., and  
697 Robertson, A. H. F., Geological Society of London Special Publication, 17, 441–447,  
698 1984.

699 Nzegge, O. M., Satır, M., Siebel, W. and Taubald, H.: Geochemical and isotopic constraints  
700 on the genesis of the Late Palaeozoic Deliktas, and Sivrikaya granites from the  
701 Kastamonu granitoid belt (Central Pontides, Turkey). *Neues Jahrbuch Mineralogische*  
702 *Abhandlungen* 183, p. 27–40, 2006.

703 Okay, A.I. and Göncüoğlu, M.C.: Karakaya Complex: A review of data and concepts. *Turkish*  
704 *Journal of Earth Sciences* 13, p. 77–95, 2004.

705 Okay, A.I., Monod, O. and Monié, P.: Triassic blueschists and eclogites from northwest  
706 Turkey: vestiges of the Paleo–Tethyan subduction. *Lithos* 64, 155–178, 2002.

707 Okay, A.I., Tüysüz, O., Satır, M., Özkan–Altınır, S., Altınır, D., Sherlock, S. and Eren, R.H.:  
708 Cretaceous and Triassic subduction–accretion HP–LT metamorphism and continental  
709 growth in the Central Pontides, Turkey. *Geological Society of America Bulletin* 118, p.  
710 1247–1269, doi:10.1130/B25938.1, 2006.

711 Pearce, J.A.: Trace element characteristics of lavas from destructive plate boundaries. In:  
 712 Thorpe, R.S. (ed), Andesites: orogenic andesites and related rocks. John Wiley &  
 713 Sons, Chichester, 525–548, 1982.

714 Pearce, J.A.: Geochemical fingerprinting of oceanic basalts with applications to ophiolite  
 715 classification and the search for Archean oceanic crust. *Lithos* 100, 14–48, 2008.

716 Pearce, J.A. and Cann, J.R.: Tectonic setting of basic volcanic rocks determined using trace  
 717 element analyses. *Earth and Planetary Scientific Letters* 19, 290–300, 1973.

718 Pearce, J.A. and Peate, D.W.: Tectonic implications of the composition of volcanic arc  
 719 magmas. *Annual Review of Earth and Planetary Sciences* 23, 251–285, doi:  
 720 10.1146/annurev.earth.23.050195.001343, 1995.

721 Pearce, J. A. and Robinson, P.T.: The Troodos ophiolitic complex probably formed in a  
 722 subduction initiation, slab edge setting. *Gondwana Research* 18 60–81.  
 723 10.1016/j.gr.2009.12.003, 2010.

724 Pearce, J.A., Harris, B.W., and Tindle, A.G.: Trace element discrimination diagrams for the  
 725 tectonic interpretation of granitic rocks. *Journal of Petrology* 25, 956–983, 1984.

726 Peccerillo, A. and Taylor, S.R.: Geochemistry of Eocene calc–alkaline volcanic rocks from  
 727 the Kastamonu area, northern Turkey. *Contributions to Mineralogy and Petrology* 58,  
 728 63–81, 1976.

729 Plechov, P.Y., Tsai, A.E., Shcherbakov, V.D. and Dirksen, O.V.: Opacitization conditions of  
 730 hornblende in Bezmyannyi volcano andesites (March 30, 1956 eruption). *Petrology* 1,  
 731 19–35, 2008.

732 Rock, N.M.S.: The nature and origin of lamprophyres: some definitions, distinctions, and  
 733 derivations. *Earth-Science Reviews* 13, 123–169, 1977.

734 Rock, N.M.S.: The nature and origin of calc-alkaline lamprophyres: minettes, vogesites,  
 735 kersantites and spessartites. *Transactions of the Royal Society of Edinburgh: Earth*  
 736 *Sciences* 74 (04), 193–227, 1984.

737

- 738 Rojay, B. and Süzen, L.: Tectonostratigraphic evolution of the Cretaceous dynamic basins on  
739 accretionary ophiolitic melange prism, SW of Ankara Region. Turkish Association of  
740 Petroleum Geologists Bulletin 9, 1–12, 1997.
- 741 Rojay, B., Altıner, D., Altıner Özkan, S., Önen, A.P., James, S. and Thirlwall, M.F.:  
742 Geodynamic significance of the Cretaceous pillow basalts from North Anatolian  
743 Ophiolitic Mélange (Central Anatolia, Turkey): geochemical and paleontological  
744 constraints. *Geodinamica Acta* 17/5, 349–361, 2004.
- 745 Saccani, E., Beccaluva, L., Photiades, A. and Zeda, O.: Petrogenesis and tectono–magmatic  
746 significance of basalts and mantle peridotites from the Albanian–Greek ophiolites and  
747 sub–ophiolitic mélanges. New constraints for the Triassic–Jurassic evolution of the  
748 Neo–tethys in the Dinaride sector. *Lithos* 124, 227–242, 2011.
- 749 Sarıfakioğlu, E., Dilek, Y. and Winchester, J.A.: Late Cretaceous Subduction Initiation and  
750 Paleocene–Eocene Slab Breakoff Magmatism in South–Central Anatolia, Turkey,  
751 *International Geology Review* 55 (1), 66–87, 2013.
- 752 Sarıfakioğlu, E., Sevin, M., Esirtgen, E., Bilgiç, T., Duran, S., Parlak, O., Karabalık, N.,  
753 Alemdar, S., Dilek, Y. and Uysal, I.: The geology of ophiolitic rocks around Çankırı–  
754 Çorum Basen: petrogenesis, tectonics and ore deposits, Mineral Research and  
755 Exploration Institute of Turkey (MTA) Report, No. 11449, 196 pp., 2011 (in Turkish,  
756 unpublished).
- 757 Schiano, P., Clocchiatti, R., Shimizu, N., Mury, R.C., Johum, K.P. and Hofmann, A.W.:  
758 Hydrous, silica rich melts in the subarc mantle and their relationship with erupted arc  
759 lavas. *Nature* 377, 595–600, doi:10.1038/377595a0, 1995.
- 760 Shand, S.J.: *Eruptive Rocks*: London: Thomas Murby & Co, p. 1–360, 1927.
- 761 Shervais, J.W.: Ti–V plots and the petrogenesis of modern and ophiolitic lavas. *Earth and*  
762 *Planetary Scientific Letters* 59, 101–118, 1982.
- 763 Sun, S.S. and McDonough, W.F.: Chemical and isotopic systematics of oceanic basalts:  
764 Implications for mantle composition and processes. In: Saunders, A.D. and Norry, M.J.  
765 (eds.), *Magmatism in the ocean basins*. Geological Society of London Special  
766 Publication 42, 313–345, 1989.

- 767 Tankut, A., Dilek, Y. and Önen, P.: Petrology and geochemistry of the Neo-Tethyan  
768 volcanism as revealed in the Ankara melange, Turkey. *Journal of Volcanology and*  
769 *Geothermal Research* 85, 265–284, 1998.
- 770 Tatsumi, Y., Hamilton, D. and Nesbitt, R.W.: Chemical characteristics of fluid phase released  
771 from a subducted lithosphere and origin of arc magmas. Evidence from high pressure  
772 experiments and natural rocks. *Journal of Volcanology and Geothermal Research* 29,  
773 293–309, 1986.
- 774 Tekeli, O.: Subduction complex of pre-Jurassic age, northern Anatolia, Turkey. *Geology* 9,  
775 68–72, 1981.
- 776 Topuz, G., Altherr, R., Satır, M. and Schwarz, M.: Low grade metamorphic rocks from the  
777 Pulur complex, NE Turkey: Implications for pre-Liassic evolution of the Eastern  
778 Pontides. *International Journal of Earth Sciences* 93, p. 72–91, doi:10.1007/s00531–  
779 003–0372–5, 2004.
- 780 Topuz, G., Okay, A.I., Altherr, R., Meyer, H.P. and Nasdala, L.: Partial high-pressure  
781 aragonitization of micritic limestones in an accretionary complex, Tavşanlı Zone, NW  
782 Turkey. *Journal of Metamorphic Geology* 24, p. 603–613, 2006.
- 783 Tüysüz, O., Dellaloğlu, A.A. and Terzioğlu, N.: A magmatic belt within the Neo-tethyan  
784 suture zone and its role in the tectonic evolution of northern Turkey. *Tectonophysics*  
785 243, 173–191, 1995.
- 786 Uğuz, M. F., Sevin, M. and Duru, M.: 1/500 000 scale geological maps of Turkey, Sinop  
787 Quadrangle (MTA General Directorate, Ankara), 2002.
- 788 White, W.M.: Sources of oceanic basalts: radiogenic isotopic evidence. *Geology* 13, 115–  
789 118, 1985.
- 790 Whitney, D.L. and Dilek, Y.: Metamorphism during crustal thickening and extension in central  
791 Anatolia: The Nigde metamorphic core complex. *Journal of Petrology* 39, 1385–1403,  
792 1998.
- 793 Wood, D.A.: The application of a Th–Hf–Ta diagram to problems of tectonomagmatic  
794 classification and to establishing the nature of crustal contamination of basaltic lavas of  
795 the British Tertiary Volcanic Province. *Earth and Planet Scientific Letters* 56, 11–30,  
796 1980.

797 Zindler, A. and Hart, S.R.: Chemical geodynamics. Annual Review of Earth and Planet  
798 Science Letters 14, 493–571, 1986.

## FIGURE CAPTIONS

**Figure 1.** Simplified ophiolite map of Turkey showing the distribution of the suture zones and some of the major tectonic entities in Turkey (from MTA, 2001). Pontide tectonic belt including the Sakarya Continent. The inset box refers to the map area in Fig. 2.

**Figure 2.** Geological map of the Çankırı–Çorum area along the IAESZ in north-central Turkey (modified after Uğuz et al., 2002). NAF: North Anatolian Fault.

**Figure 3a.** Geological map of the Kalecik area, east of Ankara, showing the distribution of the ophiolitic, turbiditic and island-arc rock units in the Ankara Mélange in north-central Turkey.

**Figure 3b.** The generalized tectonostratigraphic columnar section showing the igneous pseudostratigraphy and internal structure of the Eldivan ophiolite, the Ankara Mélange and the island-arc magmatic rocks, their tectonic basement, and sedimentary cover.

**Figure 4.** Geological map of the northern part of the Kalecik area (modified after Hakyemez et al., 1986).

**Figure 5.** View of the Ankara Mélange and the Karakaya Complex (Sakarya Continent). Key to lettering: AOM = Ankara Mélange,  $\beta$  = basalt, KC = Karakaya Complex, ms = mudstone, pg = plagiogranite, sp = serpentinized peridotites.

**Figure 6.** Simplified geological map of the Yapraklı–Çankırı area, showing the distribution of the ~180 Ma Neotethyan ophiolitic units, ophiolitic mélange and island-arc rocks.

**Figure 7.** Geological map of the Lalöglü (Çorum) area, showing the Neotethyan Eldivan ophiolite and the island-arc rock units.

**Figure 8.** (a) Neritic limestone covering the seamount volcanic-volcaniclastic rocks in the Ankara Mélange. (b) Seamount pillow lavas in the Ankara Mélange. NL= neritic limestone.

**Figure 9.** (a) Limestone-volcanic sandstone intercalation in the island-arc sequence. (b) A mafic dike (island-arc origin) crosscutting the pelagic limestone rocks. (c) Alkaline basaltic rocks with columnar joint structures. (d) Arc volcaniclastic rocks intruded by basaltic to andesitic dikes.

**Figure 10.** (a) Upper Cretaceous reefal limestone with rudist fossils unconformably overlying the arc volcanic rocks. (b) Reefal limestone underlain by volcanic sandstone. (c) Alkaline

pillow lavas overlain by volcanic sandstone-pebblestone. (d) Alkaline pillow lavas with radial joint structures. All rocks in a through d represent the island-arc units.

**Figure 11.** Lamprophyric dikes crosscutting various lithological units in the Ankara Mélange.

**Figure 12.** Photomicrographs of (a) A seamount alkaline basalt sample. (b) Doleritic dike rock of the 180 Ma Neotethyan oceanic crust. (c) Island-arc alkaline basalt sample in cross-polarized light. (d) Island-arc alkaline basalt sample in plane-polarized light. (e) Island-arc basaltic andesite dike, showing a glomeroporphyritic texture. (f) Lamprophyric dike rock with small prismatic cpx (diopside) in a feldspar + phlogopite groundmass (plane-polarized light). (g). Lamprophyric dike rock with small prismatic cpx (diopside and phlogopite). (h) Syenodioritic pluton rock with plagioclase (altered to clay minerals) and biotite + hornblende and minor cpx (cross-polarized light).

**Figure 13.** Geochemical classification of ophiolitic and seamount volcanic rocks. (a) Total alkali vs.  $\text{SiO}_2$  diagram (Le Bas et al., 1986). (b) AFM diagram (Irvine and Baragar, 1971). (c) Ti–Zr–Y discrimination diagram (Pearce and Cann, 1973). (d) Ti–V diagram (Shervais, 1982). (e) N-MORB-normalized multi-element diagrams of the most representative samples (normalization values from Sun and McDonough, 1989). Key to lettering: A = andesite, B = basalt, BA = basaltic andesite, BS = basanite, BTA = basaltic trachyandesite, D = dacite, F = foidite, PC = picrobasalt, PH = phonolite, PHTP = phonotephrite, R = rhyolite, T = trachyte, TA = trachyandesite, TB = trachybasalt, TD = trachydacite, TP = tephrite. IB = alkali–subalkali subdivision from Irvine and Baragar (1971).

**Figure 14.** Geochemical classification of island-arc rocks. (a) Total alkali vs.  $\text{SiO}_2$  diagram (Le Bas et al., 1986). (b) TAS diagram (Cox et al., 1979) for syeno-dioritic pluton rocks. (c) Alk–MgO–FeO<sub>t</sub> diagram (Irvine and Baragar, 1971) of the subalkaline arc volcanic units (Dönmez et al., 2009, and this study). (d)  $\text{K}_2\text{O}$  vs.  $\text{SiO}_2$  diagram (Peccerillo and Taylor, 1976). (e) Th vs. Co diagram (Hastie et al., 2007). Ce/Yb vs. Ta/Yb diagram (Pearce, 1982).

**Figure 15.** Major oxides and trace elements vs. MgO variation diagrams for various alkaline island-arc units.

**Figure 16.** (a) A/CNK, molar  $\text{Al}_2\text{O}_3/(\text{CaO}+\text{Na}_2\text{O}+\text{K}_2\text{O})$  vs. A/NK, molar  $\text{Al}_2\text{O}_3/(\text{Na}_2\text{O}+\text{K}_2\text{O})$  diagram (Shand, 1927). (b, c) trace element discrimination diagrams (Nb–Y and Rb vs. Y+Nb) for syenodioritic pluton rocks (fields from Pearce et al., 1984; VAG = volcanic arc granites, WPG = within-plate granites, ORG = ocean ridge granites. SYN-COLG = syn-collisional granites. (d) Ti–Zr–Y diagram. (e) Ti–V diagram. (f)  $\text{TiO}_2$  vs.  $\text{Al}_2\text{O}_3$  diagram. (g) Y




860 vs. Zr diagram. (h) Th/Yb vs. Nb/Yb diagram (fields after Pearce and Cann, 1973; Shervais,  
861 1982; Muller et al., 1992; Pearce, 2008).

862 **Figure 17.** (a) Primitive mantle-normalized multi-element diagrams for the high-K shoshonitic  
863 arc rocks (normalization values from Sun & McDonough, 1989). (b) Chondrite-normalized  
864 REE patterns of the same rocks (normalization values from Sun & McDonough, 1989).


865 **Figure 18.** (a) Ba/Nb vs. La/Nb diagram for the high-K island arc rocks. The data for N-  
866 MORB, OIB and PM are from Sun & McDonough (1989). (b) La/Yb vs. La diagram for the  
867 island-arc rock units, illustrating the effects of partial melting and fractionation in their melt  
868 evolution.

869 **Figure 19.** Isotope variation diagrams for the Upper Cretaceous–Lower Paleocene high-K  
870 island-arc rocks. (a)  $^{143}\text{Nd}/^{144}\text{Nd}$  vs.  $^{87}\text{Sr}/^{86}\text{Sr}$  diagram. (b)  $^{206}\text{Pb}/^{204}\text{Pb}$  vs.  $^{208}\text{Pb}/^{204}\text{Pb}$  diagram.  
871 (c)  $^{206}\text{Pb}/^{204}\text{Pb}$  vs.  $^{207}\text{Pb}/^{204}\text{Pb}$  diagram. (d)  $^{87}\text{Sr}/^{86}\text{Sr}$  vs.  $^{206}\text{Pb}/^{204}\text{Pb}$  diagram. (e)  $^{143}\text{Nd}/^{144}\text{Nd}$   
872 vs.  $^{206}\text{Pb}/^{204}\text{Pb}$  diagram. Compositional fields for the upper and lower crust, MORB (mid-  
873 ocean ridge basalt), HIMU (enriched mantle in U and Th relative to Pb), OIB (ocean island  
874 basalt), EMI (enriched mantle I) and EMII (enriched mantle II) are from Zindler and Hart  
875 (1986). The field for *Oceanic Islands* is from White (1985). NHRL = Northern Hemisphere  
876 Reference Line.



877 **Figure 20.** Sequential tectonic diagrams depicting the intra-oceanic magmatic evolution of  
878 the Ankara Mélange in the Northern Neotethys during the Jurassic – Paleocene. A.  
879 Suprasubduction zone generation of the oldest Neotethyan oceanic crust (~180 Ma) in the  
880 upper plate of a North-dipping intra-oceanic subduction zone, and seamount construction  
881 (SM1 and SM2) in the downgoing oceanic plate. High-grade metamorphic rock blocks and  
882 turbiditic sandstone-mudstone sequences in the Ankara Mélange formed in the subduction  
883 channel (blue in color) and the accretionary prism, respectively. B. Accretion of Seamount-1  
884 into the accretionary complex and related deformation in the subduction-accretion system. C.  
885 Slab rollback and associated extension and SSZ oceanic crust formation (~85-80 Ma) in the  
886 upper plate. Accretion of Seamount-2 into the accretionary complex, and the lateral growth  
887 and deformation in the subduction-accretion system. D. Island arc construction and  
888 magmatism on and across the pre-existing SSZ oceanic lithosphere and the subduction-  
889 accretion complex (i.e. Ankara Mélange units). With continued slab retreat, arc magmatism  
890 shifts southward following the migrating trench, and becomes more alkaline in time,  
891 producing lamprophyric and syeno-dioritic intrusions. See text for further explanation.



**Figure 1.**




**Figure 2.**





**Figure 3a.**






**Figure 3b.**




**Figure 4.**



**Figure 5.**





**Figure 6.**








Figure 7.



**Figure 8.**






910



911

**Figure 9.**






**Figure 10.**




**Figure 11.**






**Figure 12.**




**Figure 13.**






**Figure 14.**








**Figure 15.**




**Figure 16.**




**Figure 17.**



**Figure 18.**



**Figure 19.**



**Figure 20.**


934 **TABLES**

935 **Table 1.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for a basaltic rock sample (YK-11) from the youngest SSZ ophiolite in the Ankara Mélange, Turkey.

**Sample: YK-11 (whole rock): Basalt, J=0.004426±0.000051**

| T°C  | $^{40}\text{Ar}_{\text{cc}}(\text{STP})$ | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K | $\sum^{39}\text{Ar}$<br>(%) | Age (Ma) | $\pm 1\sigma$ | $\pm 1\sigma$ |
|------|--|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|------|-----------------------------|----------|---------------|---------------|
| 500  | $20.81 \times 10^{-9}$                   | 29.6                            | 0.1           | 0.0418                          | 0.0029        | 0.479                           | 0.011         | 0.0620                          | 0.0049        | 1.72 | 40.5                        | 88.0     | 11.0          |               |
| 600  | $9.45 \times 10^{-9}$                    | 22.5                            | 0.1           | 0.0364                          | 0.0037        | 0.584                           | 0.011         | 0.0438                          | 0.0033        | 2.10 | 64.8                        | 74.4     | 7.6           |               |
| 700  | $12.18 \times 10^{-9}$                   | 79.8                            | 0.6           | 0.0733                          | 0.0129        | 0.627                           | 0.036         | 0.2306                          | 0.0075        | 2.26 | 73.6                        | 90.6     | 16.5          |               |
| 800  | $9.18 \times 10^{-9}$                    | 184.7                           | 9.4           | 0.1457                          | 0.0530        | 0.751                           | 0.145         | 0.5966                          | 0.0347        | 2.70 | 76.5                        | 65.7     | 38.7          |               |
| 1000 | $9.00 \times 10^{-9}$                    | 58.5                            | 0.7           | 0.0714                          | 0.0153        | 1.630                           | 0.046         | 0.1440                          | 0.0125        | 5.87 | 85.3                        | 123.2    | 27.4          |               |
| 1130 | $8.12 \times 10^{-9}$                    | 32.0                            | 0.2           | 0.0332                          | 0.0070        | 1.131                           | 0.019         | 0.0811                          | 0.0055        | 4.07 | 100.0                       | 62.9     | 12.6          |               |

**Age Spectrum:** The sample yielded age spectrum with well behaved plateau, characterized by 73.6% of  $^{39}\text{Ar}$ , Age value of  $80.3 \pm 7.6$  Ma. On the Inverse Isochrone Plot points form linear regression characterized by age value of  $75.8 \pm 7.4$  and  $(^{40}\text{Ar}/^{36}\text{Ar})_0 = 300 \pm 8$






936 **Table 2a.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for an epidote-glaucophane schist rock from a metamorphic block in the Ankara Mélange, Turkey.

**Sample: YK-6: epidote-glaucophane schist,  $J=0.004420\pm0.000051$**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K  | $\sum^{39}\text{Ar}$ (%) | Age (Ma)    | $\pm 1\sigma$ | $\pm 1\sigma$ |
|------|------------------------------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|-------|--------------------------|-------------|---------------|---------------|
| 500  | $35.70 \times 10^{-9}$             | 48.05                           | 0.1           | 0.0463                          | 0.003         | 1.5385                          | 0.0082        | 0.1299                          | 0.002         | 5.54  | 6.4                      | 75.4        | 4.6           |               |
| 600  | $32.83 \times 10^{-9}$             | 18.75                           | 0.02          | 0.0257                          | 0.0005        | 1.221                           | 0.0031        | 0.0274                          | 0.0012        | 4.4   | 21.5                     | 83.1        | 2.9           |               |
| 700  | $38.21 \times 10^{-9}$             | 17.65                           | 0.01          | 0.0255                          | 0.0007        | 0.5962                          | 0.0014        | 0.0229                          | 0.0006        | 2.15  | 40.1                     | <b>84.7</b> | <b>1.7</b>    |               |
| 800  | $40.74 \times 10^{-9}$             | 19.89                           | 0.02          | 0.0236                          | 0.001         | 1.3996                          | 0.0017        | 0.027                           | 0.0008        | 5.04  | 57.8                     | 92.7        | 2.1           |               |
| 900  | $21.78 \times 10^{-9}$             | 19.32                           | 0.03          | 0.0277                          | 0.0008        | 10.821                          | 0.0144        | 0.0325                          | 0.0012        | 38.96 | 67.5                     | 75.8        | 2.9           |               |
| 1000 | $16.15 \times 10^{-9}$             | 14.81                           | 0.03          | 0.0253                          | 0.0016        | 12.5237                         | 0.0228        | 0.0168                          | 0.0017        | 45.09 | 76.9                     | 76.8        | 4             |               |
| 1130 | $35.36 \times 10^{-9}$             | 13.2                            | 0.01          | 0.0244                          | 0.0004        | 9.9743                          | 0.0097        | 0.0145                          | 0.0009        | 35.91 | 100                      | 69.8        | 2.2           |               |

**Age Spectrum:** The sample yielded age spectrum with two 3 steps plateaus, characterized accordingly by 40.1% of  $^{39}\text{Ar}$ , Age value of  $83.5 \pm 1.7$  Ma and 42.2 % of  $^{39}\text{Ar}$ , Age value of  $72.2 \pm 2.2$  Ma. On the Inverse Isochrone Plot points form two linear regression characterized by age value of  $87.8 \pm 2.5$  and  $(^{40}\text{Ar}/^{36}\text{Ar})_0 = 285 \pm 5$ . The presence of two age plateaus evidence to isotope heterogeneity of YK 6.






937 **Table 2b.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for an epidote-chlorite schist rock from a metamorphic block in the Ankara Mélange, Turkey.

**Sample: YK-7: epidote-chlorite schist, J=0.004121±0.000044**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | ±1σ | $^{38}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | $^{37}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | $^{36}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | Ca/K   | $\sum^{39}\text{Ar}$ (%) | Age (Ma)     | ±1σ        | ±1σ |
|------|------------------------------------|---------------------------------|-----|---------------------------------|--------|---------------------------------|--------|---------------------------------|--------|--------|--------------------------|--------------|------------|-----|
| 500  | $33.53 \times 10^{-9}$             | 57.6                            | 0.2 | 0.051                           | 0.0027 | 4.4795                          | 0.0134 | 0.1488                          | 0.0027 | 16.13  | 16.1                     | 98.8         | 5.7        |     |
| 600  | $31.15 \times 10^{-9}$             | 38.4                            | 0.1 | 0.0304                          | 0.0018 | 3.5391                          | 0.0081 | 0.0525                          | 0.0019 | 12.74  | 38.7                     | <b>162.5</b> | <b>4.2</b> |     |
| 700  | $42.75 \times 10^{-9}$             | 44.5                            | 0.1 | 0.0355                          | 0.0018 | 5.4612                          | 0.0142 | 0.0761                          | 0.0023 | 19.66  | 65.3                     | 156.8        | 4.9        |     |
| 800  | $19.66 \times 10^{-9}$             | 31                              | 0.1 | 0.0271                          | 0.0028 | 1.8875                          | 0.0089 | 0.0323                          | 0.0027 | 6.8    | 82.9                     | 153.1        | 5.8        |     |
| 900  | $12.92 \times 10^{-9}$             | 50.5                            | 0.4 | 0.0507                          | 0.0062 | 3.436                           | 0.0365 | 0.1139                          | 0.0087 | 12.37  | 90                       | 121          | 17.9       |     |
| 1000 | $6.24 \times 10^{-9}$              | 60.7                            | 1   | 0.0518                          | 0.0203 | 11.168                          | 0.1806 | 0.1419                          | 0.0162 | 40.2   | 92.8                     | 134.7        | 32.7       |     |
| 1130 | $22.81 \times 10^{-9}$             | 88.4                            | 0.5 | 0.0743                          | 0.0064 | 56.2813                         | 0.3389 | 0.2421                          | 0.0062 | 202.61 | 100                      | 121.4        | 12.4       |     |


**Age Spectrum:** The sample yielded age spectrum with 3 steps plateau, characterized by 66.7% of  $^{39}\text{Ar}$ , Age value of  $158.4 \pm 4.2$  Ma. On the Inverse Isochrone Plot points form linear regression characterized by age value of  $166.9 \pm 5.9$  and  $(^{40}\text{Ar}/^{36}\text{Ar})_0 = 272 \pm 8$ .



939 **Table 2c.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for an epidote-actinolite schist rock from a metamorphic block in the Ankara Mélange, Turkey.

**Sample: YK-1: epidote-actinolite schist,  $J=0.004428\pm0.000051$**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K   | $\sum^{39}\text{Ar}$ (%) | Age (Ma)     | $\pm 1\sigma$ | $\pm 1\sigma$ |
|------|------------------------------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|--------|--------------------------|--------------|---------------|---------------|
| 500  | $28.20 \times 10^{-9}$             | 195.111                         | 1.614         | 0.1185                          | 0.0072        | 8.7082                          | 0.0754        | 0.5725                          | 0.0095        | 31.35  | 4.1                      | 196.1        | 17.7          |               |
| 600  | $35.22 \times 10^{-9}$             | 109.08                          | 0.736         | 0.0809                          | 0.0035        | 12.8319                         | 0.0878        | 0.2835                          | 0.007         | 46.19  | 13.2                     | 191.6        | 14.5          |               |
| 700  | $46.14 \times 10^{-9}$             | 52.526                          | 0.083         | 0.0345                          | 0.0021        | 5.2249                          | 0.0101        | 0.0848                          | 0.0016        | 18.81  | 38.1                     | 207          | 4             |               |
| 800  | $22.10 \times 10^{-9}$             | 43.305                          | 0.132         | 0.032                           | 0.0025        | 3.3143                          | 0.0128        | 0.0652                          | 0.003         | 11.93  | 52.5                     | 182.5        | 6.8           |               |
| 900  | $28.97 \times 10^{-9}$             | 56.607                          | 0.158         | 0.0384                          | 0.0023        | 16.1294                         | 0.0458        | 0.1083                          | 0.0028        | 58.07  | 67                       | <b>186.6</b> | <b>6.3</b>    |               |
| 1000 | $30.80 \times 10^{-9}$             | 41.871                          | 0.112         | 0.0328                          | 0.0018        | 21.4028                         | 0.0573        | 0.0558                          | 0.0027        | 77.05  | 87.8                     | 192.1        | 6             |               |
| 1130 | $30.24 \times 10^{-9}$             | 69.813                          | 0.261         | 0.0523                          | 0.0025        | 34.5698                         | 0.1292        | 0.1345                          | 0.0038        | 124.45 | 100                      | 225.6        | 8.2           |               |



940

941 **Table 3.** Whole-rock K/Ar age data from metamorphic rock blocks in the Ankara Mélange, Turkey.

| Sample no. | Rock                     | %K   | $^{40}\text{Ar}/^{36}\text{Ar}$ | $^{40}\text{Ar}_{\text{rad}}$ , nl/g | % $^{40}\text{Ar}_{\text{air}}$ | error | Age, Ma   |
|------------|--------------------------|------|---------------------------------|--------------------------------------|---------------------------------|-------|-----------|
| CE.981     | Phyllite                 | 1.68 | 883.2                           | 7.932                                | 33.5                            | 4.5   | 119.8±3.3 |
| CE.228     | Actinolite schist        | 0.36 | 347.8                           | 2.558                                | 85.1                            | 0.6   | 177.4±5.8 |
| CE.976     | Amphibole-epidote schist | 0.22 | 405.9                           | 2.316                                | 72.9                            | 1.3   | 256.9±8.0 |


942

943 **Table 4a.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for an island-arc basaltic rock (Sample No. YK-4) in the Ankara Mélange, Turkey.

**Sample: YK-4 (whole rock): Basalt, J=0.004353±0.000050**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | ±1σ  | $^{38}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | $^{37}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | $^{36}\text{Ar}/^{39}\text{Ar}$ | ±1σ    | Ca/K  | $\Sigma^{39}\text{Ar}$ (%) | Age (Ma) | ±1σ  |
|------|------------------------------------|---------------------------------|------|---------------------------------|--------|---------------------------------|--------|---------------------------------|--------|-------|----------------------------|----------|------|
| 500  | $44.60 \times 10^{-9}$             | 168.69                          | 0.97 | 0.1269                          | 0.007  | 7.5932                          | 0.0458 | 0.539                           | 0.0065 | 27.34 | 6.9                        | 72.4     | 12.8 |
| 600  | $14.83 \times 10^{-9}$             | 32.9                            | 0.17 | 0.0413                          | 0.0033 | 10.1845                         | 0.0528 | 0.0795                          | 0.0051 | 36.66 | 18.7                       | 72.4     | 11.4 |
| 700  | $13.82 \times 10^{-9}$             | 26.82                           | 0.1  | 0.0333                          | 0.0048 | 4.2753                          | 0.017  | 0.0621                          | 0.0036 | 15.39 | 32.2                       | 65.2     | 8    |
| 800  | $24.70 \times 10^{-9}$             | 26.65                           | 0.08 | 0.0331                          | 0.0022 | 1.935                           | 0.0067 | 0.0635                          | 0.0029 | 6.97  | 56.5                       | 60.8     | 6.6  |
| 900  | $18.59 \times 10^{-9}$             | 18.29                           | 0.04 | 0.021                           | 0.0013 | 1.3887                          | 0.0051 | 0.0299                          | 0.0022 | 5     | 83.1                       | 72.8     | 4.9  |
| 1000 | $7.83 \times 10^{-9}$              | 31.49                           | 0.17 | 0.0381                          | 0.0053 | 1.4804                          | 0.0211 | 0.0611                          | 0.0055 | 5.33  | 89.7                       | 102.6    | 12.1 |
| 1130 | $11.68 \times 10^{-9}$             | 29.58                           | 0.12 | 0.0279                          | 0.0033 | 3.5167                          | 0.016  | 0.054                           | 0.0041 | 12.66 | 100                        | 104      | 9.2  |

**Age Spectrum:** The sample yielded age spectrum with 3 steps plateau, characterized by 64.4% of  $^{39}\text{Ar}$ , Age value of  $67.8 \pm 4.9$  Ma. On the Inverse Isochrone Plot points form linear regression characterized by age value of  $68.1 \pm 4.4$  and  $(^{40}\text{Ar}/^{36}\text{Ar})_0 = 296.1 \pm 3.5$ .




944

945 **Table 4b.**  $^{40}\text{Ar}/^{39}\text{Ar}$  biotite age data for a lamprophyre dike (Sample No. YK-19) from the island-arc unit in the Ankara Mélange, Turkey.

**Sample: YK-19 (biotite): Lamprophyre,  $J=0.007143 \pm 0.000133$**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K  | $\sum^{39}\text{Ar}$ (%) | Age (Ma) | $\pm 1\sigma$ |
|------|------------------------------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|-------|--------------------------|----------|---------------|
| 500  | $3.9 \times 10^{-9}$               | 92.802                          | 0.0393        | 0.015480                        | 0.004274      | 0.61970                         | 0.05770       | 0.028643                        | 0.001314      | 22.30 | 0.6                      | 10.50    | 5.0           |
| 625  | $7.7 \times 10^{-9}$               | 83.584                          | 0.0116        | 0.015744                        | 0.001217      | 0.31455                         | 0.01480       | 0.015871                        | 0.001284      | 1.13  | 2.0                      | 46.7     | 4.80          |
| 750  | $37.9 \times 10^{-9}$              | 69.914                          | 0.0044        | 0.013301                        | 0.000117      | 0.05874                         | 0.00360       | 0.008418                        | 0.000210      | 0.21  | 10.40                    | 57.1     | 1.30          |
| 850  | $45.7 \times 10^{-9}$              | 64.526                          | 0.0017        | 0.013498                        | 0.000076      | 0.04725                         | 0.00102       | 0.004948                        | 0.000237      | 0.17  | 21.20                    | 63.2     | 1.40          |
| 950  | $51.6 \times 10^{-9}$              | 65.506                          | 0.0033        | 0.013393                        | 0.000248      | 0.08738                         | 0.00394       | 0.004805                        | 0.000142      | 0.31  | 33.3                     | 64.9     | 1.30          |
| 1050 | $112.6 \times 10^{-9}$             | 62.462                          | 0.0035        | 0.013738                        | 0.000060      | 0.07037                         | 0.00091       | 0.004305                        | 0.000083      | 0.25  | 61.0                     | 63.0     | 1.20          |
| 1130 | $156.9 \times 10^{-9}$             | 61.778                          | 0.0020        | 0.013593                        | 0.000030      | 0.07395                         | 0.00067       | 0.003986                        | 0.000051      | 0.27  | 100.0                    | 63.3     | 1.20          |

**Age Spectrum:** The sample yielded age spectrum with four steps Plateau characterized by 89.6% of  $^{39}\text{Ar}$ , Age value of  $63.6 \pm 1.2$  Ma. On the Inverse Isochrone Plot plateau points form linear trend, characterized by age value of  $57.5 \pm 4.1$  Ma, MSWD = 1.5




946

947 **Table 4c.** Whole-rock  $^{40}\text{Ar}/^{39}\text{Ar}$  age data for a lamprophyre dike (Sample No. YK-20) from the island-arc unit in the Ankara Mélange, Turkey.

**Sample: YK-20 (whole rock): Lamprophyre,  $J=0.007258 \pm 0.000137$**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K | $\Sigma^{39}\text{Ar}$ (%) | Age (Ma) | $\pm 1\sigma$ |
|------|------------------------------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|------|----------------------------|----------|---------------|
| 550  | $114.7 \times 10^{-9}$             | 48.724                          | 0.0048        | 0.014063                        | 0.000040      | 0.28860                         | 0.00023       | 0.004714                        | 0.000071      | 1.04 | 16.20                      | 45.0     | 0.9           |
| 625  | $78.3 \times 10^{-9}$              | 71.168                          | 0.0064        | 0.014657                        | 0.000137      | 128.516                         | 0.00083       | 0.006715                        | 0.000045      | 4.63 | 23.70                      | 66.0     | 1.20          |
| 700  | $73.1 \times 10^{-9}$              | 63.482                          | 0.0041        | 0.014148                        | 0.000062      | 115.422                         | 0.00187       | 0.003713                        | 0.000142      | 4.16 | 31.70                      | 67.5     | 1.40          |
| 775  | $85.2 \times 10^{-9}$              | 62.554                          | 0.0032        | 0.013660                        | 0.000068      | 0.51460                         | 0.00080       | 0.003174                        | 0.000096      | 1.85 | 41.0                       | 68.3     | 1.30          |
| 850  | $68.1 \times 10^{-9}$              | 59.659                          | 0.0028        | 0.013744                        | 0.000112      | 0.19993                         | 0.00023       | 0.004958                        | 0.000090      | 0.72 | 48.9                       | 58.0     | 1.10          |
| 950  | $54.6 \times 10^{-9}$              | 58.431                          | 0.0024        | 0.014092                        | 0.000162      | 0.29838                         | 0.00075       | 0.004897                        | 0.000152      | 1.07 | 55.3                       | 56.7     | 1.20          |
| 1050 | $210.7 \times 10^{-9}$             | 50.825                          | 0.0020        | 0.013886                        | 0.000024      | 0.61827                         | 0.00022       | 0.005145                        | 0.000055      | 2.23 | 83.8                       | 46.0     | 0.9           |
| 1130 | $121.2 \times 10^{-9}$             | 51.432                          | 0.0013        | 0.014028                        | 0.000073      | 144.061                         | 0.00118       | 0.005433                        | 0.000056      | 5.19 | 100.0                      | 45.7     | 0.9           |

**Age Spectrum:** The sample yielded complex age spectrum with noticeable hump after low temperature step containing three steps intermediate plateau followed by high temperature two steps intermediate Plateau. Intermediate plateaus are characterized accordingly by 24.8% of  $^{39}\text{Ar}$ , Age value of  $67.2 \pm 1.2$  Ma and 44.7% of  $^{39}\text{Ar}$ , Age value of  $45.9 \pm 0.9$  Ma.




948 **Table 4d.**  $^{40}\text{Ar}/^{39}\text{Ar}$  biotite age data for a syeno-diorite plutonic rock (Sample No. YK-438) from the island-arc unit in the Ankara Mélange, Turkey.

**Sample: YK-438 (biotite): Syeno-diorite,  $J=0.004553 \pm 0.000054$**

| T°C  | $^{40}\text{Ar}_{\text{cc}}$ (STP) | $^{40}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{38}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{37}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | $^{36}\text{Ar}/^{39}\text{Ar}$ | $\pm 1\sigma$ | Ca/K | $\sum^{39}\text{Ar}$ (%) | Age (Ma) | $\pm 1\sigma$ |
|------|------------------------------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|---------------------------------|---------------|------|--------------------------|----------|---------------|
| 500  | $10.2 \times 10^{-9}$              | 61.084                          | 0.0090        | 0.01989                         | 0.00107       | 0.7800                          | 0.0026        | 0.01226                         | 0.00144       | 2.81 | 9.0                      | 20.30    | 3.46          |
| 600  | $14.4 \times 10^{-9}$              | 82.535                          | 0.0136        | 0.01627                         | 0.00075       | 15.022                          | 0.0075        | 0.00992                         | 0.00159       | 5.41 | 18.4                     | 43.20    | 3.79          |
| 700  | $17.3 \times 10^{-9}$              | 121.050                         | 0.0098        | 0.00960                         | 0.00130       | 21.617                          | 0.0064        | 0.01488                         | 0.00069       | 7.78 | 26.1                     | 62.22    | 1.78          |
| 800  | $25.9 \times 10^{-9}$              | 128.671                         | 0.0263        | 0.01743                         | 0.00152       | 0.9964                          | 0.0146        | 0.00617                         | 0.00198       | 3.59 | 37.0                     | 88.51    | 4.68          |
| 900  | $39.8 \times 10^{-9}$              | 126.457                         | 0.0087        | 0.01645                         | 0.00019       | 0.8052                          | 0.0025        | 0.00343                         | 0.00027       | 2.9  | 53.9                     | 93.11    | 1.25          |
| 1000 | $48.8 \times 10^{-9}$              | 126.405                         | 0.0147        | 0.01668                         | 0.00067       | 0.4739                          | 0.0055        | 0.00371                         | 0.00111       | 1.71 | 74.8                     | 92.41    | 2.77          |
| 1130 | $52.0 \times 10^{-9}$              | 111.241                         | 0.0102        | 0.01900                         | 0.00068       | 21.389                          | 0.0021        | 0.00361                         | 0.00065       | 7.7  | 100.0                    | 80.77    | 1.78          |

**Age Spectrum:** The sample yielded age spectrum with three steps plateau characterized by 48.6 % of  $^{39}\text{Ar}$ , Age value of  $92.7 \pm 1.2$  Ma. On the Inverse Isochrone Plot points don't form linear regression.



**Table 5.** Major, trace element and REE data for a selected group of volcanic and dike rocks from the Neotethyan ophiolitic units in the Ankara Mélange (first nine samples from Tankut et al., 1998).

| Sample no                      | BM1    | BM3               | BM5    | 95GK4    | 95GK6    | 95GKE4   | 96GKE51  | 96GKE57  | 96GKE58B | CE.07  | CE.08  |
|--------------------------------|--------|-------------------|--------|----------|----------|----------|----------|----------|----------|--------|--------|
| Rock-type                      | Basalt | Basaltic andesite | Basalt | Dolerite | Dolerite | Dolerite | Dolerite | Dolerite | Dolerite | Basalt | Basalt |
| <b>Oxide, wt %</b>             |        |                   |        |          |          |          |          |          |          |        |        |
| SiO <sub>2</sub>               | 50.53  | 51.34             | 50.47  | 51.08    | 49.45    | 51.53    | 48.93    | 50.45    | 47.08    | 55.50  | 54.94  |
| TiO <sub>2</sub>               | 1.12   | 0.26              | 1.13   | 1.66     | 1.68     | 1.82     | 0.67     | 1.74     | 1.55     | 0.82   | 0.81   |
| Al <sub>2</sub> O <sub>3</sub> | 15.96  | 17.16             | 15.93  | 13.45    | 13.52    | 12.53    | 11.00    | 12.97    | 11.18    | 18.43  | 18.66  |
| Fe <sub>2</sub> O <sub>3</sub> | 10.01  | 6.80              | 9.99   | 11.91    | 13.23    | 12.69    | 9.07     | 12.31    | 12.21    | 8.12   | 8.47   |
| MnO                            | 0.14   | 0.12              | 0.14   | 0.23     | 0.22     | 0.23     | 0.18     | 0.23     | 0.22     | 0.60   | 0.63   |
| MgO                            | 6.32   | 8.17              | 5.89   | 7.04     | 6.41     | 6.67     | 10.54    | 6.30     | 8.67     | 3.54   | 3.81   |
| CaO                            | 4.88   | 11.07             | 5.33   | 9.18     | 10.91    | 7.93     | 15.30    | 8.66     | 13.11    | 1.27   | 1.17   |
| Na <sub>2</sub> O              | 3.31   | 2.30              | 3.34   | 3.79     | 2.89     | 3.89     | 0.43     | 3.41     | 1.22     | 7.62   | 7.50   |
| K <sub>2</sub> O               | 0.28   | 0.28              | 0.29   | 0.34     | 0.36     | 0.39     | 0.07     | 0.37     | 0.12     | 0.17   | 0.19   |
| P <sub>2</sub> O <sub>5</sub>  | 0.13   | 0.07              | 0.12   | 0.17     | 0.21     | 0.22     | 0.06     | 0.21     | 0.23     | 0.09   | 0.10   |
| LOI                            | 6.81   | 2.05              | 6.94   | 1.2      | 1.04     | 0.85     | 3.12     | 1.69     | 3.06     | 3.70   | 3.50   |
| Total                          | 99.49  | 99.62             | 99.57  | 100.05   | 99.92    | 99.75    | 99.35    | 98.34    | 98.65    | 99.85  | 99.84  |
| <b>Trace, ppm</b>              |        |                   |        |          |          |          |          |          |          |        |        |
| Cr                             | 53.00  | 166.00            | 36.00  | 70.00    | 84.00    | 130.00   | 624.00   | 134.00   | 103.00   | 27.40  | 20.55  |
| Ni                             | 10.00  | 101.00            | 8.00   | 21.00    | 25.00    | 48.00    | 91.00    | 59.00    | 37.00    | 13.20  | 11.70  |
| Co                             |        |                   |        |          |          |          |          |          |          | 27.90  | 26.10  |
| Sc                             |        |                   |        | 35       | 38       | 32       | 41       | 35       | 45       |        |        |
| Rb                             | 11.00  | 9.00              | 12.00  | 4.00     | 4.00     | 5.00     | 2.00     | 7.00     | 0.00     | 2.10   | 2.40   |
| Ba                             | 242.00 |                   | 270.00 | 60.00    | 503.00   | 188.00   | 22.00    | 185.00   | 52.00    | 130.00 | 110.00 |
| Sr                             | 441.00 | 100.00            | 420.00 | 174.00   | 510.00   | 402.00   | 24.00    | 208.00   | 71.00    | 472.80 | 494.80 |
| Cs                             |        |                   |        |          |          |          | 0.64     | 24.94    | 3.91     | 0.40   | 0.50   |
| Th                             | 0.47   | 0.60              | 0.44   | 0.00     | 1.00     | 3.00     | 2.00     | 4.00     | 3.00     | 0.60   | 0.60   |
| U                              |        |                   |        | 0.21     | 0.26     | 0.29     | 0.14     | 0.28     | 0.28     | 0.30   | 0.30   |
| Nb                             | 3.00   | 2.60              | 4.00   | 3.70     | 5.70     | 4.00     | 1.30     | 3.40     | 4.80     | 1.30   | 1.40   |
| Ta                             | 0.00   | 0.00              | 0.00   | 0.24     | 0.29     | 0.25     | 0.15     | 0.18     | 0.29     | 0.10   | 0.10   |
| Zr                             | 96.00  | 21.00             | 98.00  | 101.00   | 126.00   | 119.00   | 39.00    | 91.00    | 113.00   | 37.90  | 43.60  |
| Hf                             | 1.70   | 0.30              | 2.20   | 2.77     | 3.10     | 3.14     | 1.16     | 2.64     | 3.13     | 1.40   | 1.40   |
| Y                              | 25.00  | 7.00              | 25.00  | 37.00    | 40.00    | 41.00    | 18.00    | 39.00    | 39.00    | 14.50  | 15.50  |
| V                              |        | 288               |        | 363      | 383      | 378      | 273      | 462      | 337      | 202.00 | 190.00 |
| Pb                             |        |                   |        |          | 1        |          | 1        |          | 3        | 5.9    | 5.3    |
| <b>REE, ppm</b>                |        |                   |        |          |          |          |          |          |          |        |        |
| La                             | 3.90   | 2.90              | 4.30   | 7.42     | 9.25     | 9.17     | 3.03     | 8.19     | 9.66     | 4.00   | 2.90   |
| Ce                             | 10.70  | 3.70              | 8.20   | 16.67    | 20.29    | 19.96    | 6.18     | 17.66    | 21.46    | 8.30   | 6.90   |
| Pr                             |        |                   |        | 2.31     | 2.81     | 2.73     | 0.83     | 2.43     | 2.90     | 1.31   | 1.05   |
| Nd                             | 5.70   | 2.20              | 6.90   | 11.64    | 13.66    | 13.37    | 4.44     | 12.10    | 14.01    | 5.90   | 5.90   |
| Sm                             | 2.90   | 0.50              | 3.00   | 4.07     | 4.64     | 4.71     | 1.67     | 4.41     | 4.75     | 1.71   | 1.53   |
| Eu                             | 0.90   | 0.18              | 1.00   | 1.43     | 1.59     | 1.29     | 0.70     | 1.42     | 1.52     | 0.49   | 0.46   |
| Gd                             |        |                   |        | 5.14     | 5.71     | 5.73     | 2.31     | 5.85     | 6.04     | 2.08   | 2.00   |
| Tb                             | 0.60   | 0.15              | 0.70   | 1.02     | 1.13     | 1.13     | 0.46     | 1.10     | 1.13     | 0.41   | 0.40   |
| Dy                             |        |                   |        | 6.78     | 7.42     | 7.42     | 3.12     | 7.39     | 7.34     | 2.54   | 2.59   |
| Ho                             |        |                   |        | 1.47     | 1.60     | 1.63     | 0.70     | 1.62     | 1.58     | 0.57   | 0.61   |
| Er                             |        |                   |        | 4.30     | 4.59     | 4.78     | 2.03     | 4.58     | 4.39     | 1.77   | 1.84   |
| Tm                             |        |                   |        | 0.61     | 0.65     | 0.68     | 0.30     | 0.67     | 0.65     | 0.29   | 0.31   |
| Yb                             | 3.10   | 0.96              | 2.90   | 3.71     | 3.94     | 4.18     | 1.89     | 4.09     | 3.90     | 1.80   | 1.90   |
| Lu                             | 0.49   | 0.16              | 0.50   | 0.58     | 0.63     | 0.65     | 0.30     | 0.63     | 0.61     | 0.28   | 0.32   |



953 **Table 6.** Major, trace element and REE data for a selected group of seamount volcanic rocks  
954 from the Ankara Mélange.

| Sample no                      | 2007KM327 | DM19         | KM24     | KM27     | KM28     | KM121     | KM126      | CM38     |
|--------------------------------|-----------|--------------|----------|----------|----------|-----------|------------|----------|
| Rock-type                      | Foidite   | Trachybasalt | Basanite | Basanite | Trachyte | Tephrrite | Tephryrite | Basanite |
| <b>Oxide, wt %</b>             |           |              |          |          |          |           |            |          |
| SiO <sub>2</sub>               | 39.77     | 50.15        | 41.08    | 44.55    | 68.47    | 46.36     | 44.83      | 44.26    |
| TiO <sub>2</sub>               | 2.16      | 2.46         | 1.64     | 2.17     | 0.61     | 2.10      | 2.46       | 2.15     |
| Al <sub>2</sub> O <sub>3</sub> | 15.84     | 15.90        | 13.21    | 15.83    | 15.16    | 16.70     | 15.78      | 15.75    |
| Fe <sub>2</sub> O <sub>3</sub> | 11.67     | 9.10         | 7.75     | 10.41    | 3.17     | 12.04     | 12.27      | 10.54    |
| MnO                            | 0.33      | 0.19         | 0.12     | 0.16     | 0.04     | 0.18      | 0.18       | 0.30     |
| MgO                            | 8.10      | 3.43         | 3.91     | 4.91     | 0.54     | 3.12      | 3.93       | 9.48     |
| CaO                            | 13.95     | 5.64         | 13.78    | 7.8      | 1.48     | 8.59      | 9.60       | 4.63     |
| Na <sub>2</sub> O              | 0.57      | 4.74         | 5.25     | 4.15     | 6.79     | 4.18      | 3.78       | 3.98     |
| K <sub>2</sub> O               | 1.56      | 1.91         | 0.45     | 2.1      | 1.35     | 1.81      | 1.78       | 0.74     |
| P <sub>2</sub> O <sub>5</sub>  | 0.51      | 0.77         | 0.464    | 0.653    | 0.484    | 0.89      | 0.84       | 0.68     |
| LOI                            | 5.3       | 5.5          | 11.8     | 7.1      | 1.7      | 3.8       | 4.3        | 7.1      |
| Total                          | 99.81     | 99.77        | 99.46    | 99.84    | 99.77    | 99.73     | 99.72      | 99.61    |
| <b>Trace, ppm</b>              |           |              |          |          |          |           |            |          |
| Cr                             | 342.45    | 13.70        | 232.87   | 198.62   | 13.70    | 13.70     | 20.55      | 219.17   |
| Ni                             | 128.00    | 21           | 113      | 106      | 20       | 20        | 20         | 130.70   |
| Co                             | 40.10     | 24.3         | 30.6     | 36.4     | 2.2      | 26.10     | 32.50      | 39.30    |
| Sc                             | 25.00     | 15           | 18       | 20       | 3        | 5         | 7          | 21       |
| Rb                             | 26.80     | 39.8         | 7.7      | 43.5     | 33.4     | 18.7      | 18.1       | 13.50    |
| Ba                             | 280       | 135          | 151      | 251      | 379      | 426       | 401        | 371.00   |
| Sr                             | 223.30    | 368.6        | 529.6    | 588.4    | 294.3    | 577.9     | 547.0      | 740.00   |
| Cs                             | 0.20      | 1.00         | 0.1      | 1.00     | 0.7      | 0.10      | 0.10       | 0.30     |
| Th                             | 5.90      | 7.6          | 4.9      | 5.8      | 10.5     | 8.8       | 8.0        | 6.30     |
| U                              | 1.40      | 1.6          | 1.5      | 1.2      | 1.4      | 1.9       | 1.6        | 1.60     |
| Nb                             | 55.40     | 74.5         | 47.2     | 60.5     | 96       | 88.3      | 82.0       | 62.40    |
| Ta                             | 3.00      | 4.5          | 2.8      | 3.6      | 6        | 5.1       | 5.2        | 3.70     |
| Zr                             | 200.40    | 291.3        | 187.2    | 240      | 389.5    | 273.4     | 257.0      | 252.10   |
| Hf                             | 4.90      | 6.9          | 4.6      | 6.2      | 10.2     | 5.7       | 5.9        | 6.50     |
| Y                              | 22.70     | 24.3         | 20.6     | 25       | 34       | 29.2      | 29.0       | 25.60    |
| V                              | 207       | 136          | 157      | 199      | 14       | 90        | 117        | 168      |
| Pb                             | 4         | 1.5          | 5.6      | 4.9      | 7.2      | 5         | 7.9        | 1.8      |
| <b>REE, ppm</b>                |           |              |          |          |          |           |            |          |
| La                             | 37.00     | 54.4         | 33       | 43.4     | 60.7     | 63.4      | 61.1       | 42.30    |
| Ce                             | 73.00     | 122.0        | 64.8     | 83.5     | 114.7    | 128.8     | 122.9      | 84.90    |
| Pr                             | 8.80      | 13.61        | 7.8      | 9.71     | 14.25    | 13.73     | 13.57      | 9.87     |
| Nd                             | 34.30     | 54.0         | 29.1     | 37.3     | 49       | 50.7      | 51.0       | 39.90    |
| Sm                             | 6.11      | 9.76         | 5.21     | 6.56     | 8.05     | 9.06      | 9.04       | 6.64     |
| Eu                             | 2.03      | 3.17         | 1.74     | 2.11     | 2.19     | 2.84      | 2.99       | 2.16     |
| Gd                             | 5.78      | 8.46         | 4.65     | 5.81     | 6.67     | 8.33      | 8.33       | 6.21     |
| Tb                             | 0.88      | 1.17         | 0.76     | 0.93     | 1.15     | 1.18      | 1.21       | 0.93     |
| Dy                             | 4.49      | 5.56         | 4.01     | 4.89     | 6.34     | 5.88      | 5.90       | 4.72     |
| Ho                             | 0.82      | 0.87         | 0.76     | 0.93     | 1.21     | 1.07      | 1.06       | 0.91     |
| Er                             | 2.11      | 1.94         | 2        | 2.48     | 3.42     | 2.72      | 2.63       | 2.68     |
| Tm                             | 0.33      | 0.25         | 0.29     | 0.36     | 0.51     | 0.36      | 0.36       | 0.38     |
| Yb                             | 1.85      | 1.36         | 1.71     | 2.07     | 3.27     | 2.18      | 2.17       | 2.10     |
| Lu                             | 0.28      | 0.17         | 0.25     | 0.31     | 0.47     | 0.30      | 0.30       | 0.33     |

955

956 **Table 7.** Major, trace element and REE data for a selected group of island-arc volcanic and  
957 syeno-diorite rocks from the Ankara Mélange.

| Sample no                      | 04.NAM        | 05.NAM        | 06.NAM        | DM35     | DM 36    | DM37    | CE960    | 08CM01           | 08CM07   | KM54     |
|--------------------------------|---------------|---------------|---------------|----------|----------|---------|----------|------------------|----------|----------|
| Rock-type                      | Syeno-diorite | Syeno-diorite | Syeno-diorite | Tephrite | Tephrite | Foidite | Basanite | Leucite Tephrite | Basanite | Basanite |
| <b>Oxide, wt %</b>             |               |               |               |          |          |         |          |                  |          |          |
| SiO <sub>2</sub>               | 50.31         | 49.35         | 49.11         | 41.65    | 41.86    | 39.81   | 42.41    | 48.29            | 45.00    | 46.34    |
| TiO <sub>2</sub>               | 0.59          | 0.61          | 0.62          | 0.76     | 0.74     | 0.70    | 0.89     | 0.67             | 0.79     | 0.93     |
| Al <sub>2</sub> O <sub>3</sub> | 18.69         | 18.75         | 18.39         | 15.90    | 15.55    | 12.84   | 16.89    | 18.21            | 14.57    | 14.36    |
| Fe <sub>2</sub> O <sub>3</sub> | 8.45          | 8.77          | 8.42          | 10.82    | 9.31     | 8.90    | 12.38    | 9.15             | 11.21    | 11.03    |
| MnO                            | 0.17          | 0.19          | 0.19          | 0.15     | 0.15     | 0.14    | 0.25     | 0.25             | 0.21     | 0.18     |
| MgO                            | 2.25          | 2.45          | 2.39          | 7.23     | 7.35     | 10.81   | 7.5      | 4.23             | 6.28     | 6.33     |
| CaO                            | 6.38          | 7.14          | 6.82          | 3.51     | 4.08     | 5.36    | 11.17    | 7.91             | 12.22    | 9.64     |
| Na <sub>2</sub> O              | 2.68          | 2.57          | 2.35          | 0.57     | 0.46     | 0.21    | 2.48     | 3.78             | 2.56     | 2.79     |
| K <sub>2</sub> O               | 5.32          | 5.07          | 5.68          | 5.99     | 6.15     | 5.15    | 0.62     | 2.91             | 2.15     | 3.09     |
| P <sub>2</sub> O <sub>5</sub>  | 0.48          | 0.49          | 0.49          | 0.44     | 0.41     | 0.24    | 0.42     | 0.46             | 0.33     | 0.44     |
| LOI                            | 4.30          | 4.20          | 5.10          | 12.7     | 13.6     | 15.4    | 4.7      | 3.7              | 4.2      | 4.5      |
| Total                          | 99.61         | 99.58         | 99.63         | 99.69    | 99.68    | 99.61   | 99.71    | 99.56            | 99.52    | 99.67    |
| <b>Trace, ppm</b>              |               |               |               |          |          |         |          |                  |          |          |
| Cr                             | 47.94         | 82.19         | 47.94         | 13.70    | 20.55    | 342.45  | 21       | 14.00            | 130.00   | 61.64    |
| Ni                             | 20            | 20            | 20            | 23       | 41       | 156     | 43       | 5.00             | 24.50    | 39.00    |
| Co                             | 20.60         | 22.80         | 20.70         | 25.7     | 28.0     | 38.4    | 43.1     | 24.80            | 37.10    | 36.20    |
| Sc                             | 11            | 11            | 11            | 20       | 21       | 37      | 30       | 20.00            | 48.00    | 39.00    |
| Rb                             | 140.2         | 138.5         | 148.2         | 184.1    | 172.6    | 162.6   | 18.3     | 52.50            | 29.40    | 195.80   |
| Ba                             | 1685          | 1557          | 1520          | 783      | 739      | 887     | 577      | 1883             | 1051     | 840      |
| Sr                             | 845.2         | 915.3         | 759.4         | 257.5    | 331.3    | 362.2   | 563.5    | 742.60           | 779.80   | 934.80   |
| Cs                             | 1.1           | 2.1           | 3.5           | 6        | 4.5      | 3.4     | 0.9      | 3.10             | 1.20     | 1.50     |
| Th                             | 11.9          | 13.9          | 11.1          | 10.1     | 10.3     | 7.5     | 4.7      | 18.00            | 5.90     | 5.30     |
| U                              | 3.2           | 3.1           | 3.0           | 3.3      | 3.7      | 1.8     | 1.5      | 2.40             | 1.70     | 2.20     |
| Nb                             | 8.9           | 9.4           | 7.6           | 9.1      | 8.7      | 4.2     | 4        | 12.30            | 5.10     | 3.40     |
| Ta                             | 0.4           | 0.4           | 0.3           | 0.3      | 0.4      | 0.1     | 0.3      | 0.50             | 0.20     | 0.20     |
| Zr                             | 76.1          | 86.0          | 72.9          | 72.8     | 74.4     | 49.2    | 68.9     | 87.80            | 57.70    | 75.20    |
| Hf                             | 1.6           | 2.2           | 2.0           | 2.0      | 1.7      | 1.4     | 2        | 2.50             | 1.60     | 2.00     |
| Y                              | 23.6          | 23.0          | 20.3          | 14.8     | 16.5     | 12.5    | 21.4     | 20.90            | 18.70    | 22.00    |
| V                              | 169           | 182           | 166           | 279      | 275      | 260     | 339      | 273              | 301      | 302      |
| Pb                             | 4             | 21.8          | 24.9          | 3.9      | 7.4      | 6.1     | 5.2      | 7.3              | 5.5      | 9        |
| <b>REE, ppm</b>                |               |               |               |          |          |         |          |                  |          |          |
| La                             | 33.7          | 36.6          | 31.9          | 26.0     | 25.7     | 18.8    | 20       | 36.70            | 20.70    | 18.40    |
| Ce                             | 60.0          | 63.7          | 55.4          | 53.2     | 53.5     | 37.7    | 42.6     | 66.70            | 40.00    | 41.90    |
| Pr                             | 6.72          | 7.12          | 6.20          | 5.85     | 5.63     | 3.93    | 5.52     | 7.87             | 4.89     | 5.51     |
| Nd                             | 25.3          | 25.7          | 24.9          | 23.0     | 22.5     | 15.7    | 25       | 32.80            | 20.70    | 22.90    |
| Sm                             | 4.91          | 4.88          | 4.41          | 4.42     | 4.29     | 3.21    | 5.27     | 5.81             | 4.18     | 5.12     |
| Eu                             | 1.32          | 1.31          | 1.24          | 1.18     | 1.21     | 0.91    | 1.57     | 1.53             | 1.33     | 1.47     |
| Gd                             | 4.70          | 4.51          | 4.18          | 3.89     | 4.02     | 3.21    | 5.12     | 5.23             | 4.31     | 5.17     |
| Tb                             | 0.70          | 0.71          | 0.64          | 0.57     | 0.60     | 0.47    | 0.79     | 0.76             | 0.66     | 0.80     |
| Dy                             | 3.89          | 4.11          | 3.73          | 2.83     | 3.28     | 2.52    | 4.03     | 3.88             | 3.54     | 4.34     |
| Ho                             | 0.72          | 0.75          | 0.67          | 0.56     | 0.60     | 0.46    | 0.76     | 0.73             | 0.69     | 0.81     |
| Er                             | 2.02          | 2.14          | 1.92          | 1.60     | 1.61     | 1.24    | 2.07     | 2.05             | 2.00     | 2.14     |
| Tm                             | 0.30          | 0.30          | 0.28          | 0.21     | 0.23     | 0.17    | 0.32     | 0.28             | 0.29     | 0.34     |
| Yb                             | 2.16          | 2.05          | 1.89          | 1.49     | 1.47     | 1.10    | 1.82     | 1.84             | 1.65     | 1.94     |
| Lu                             | 0.28          | 0.33          | 0.25          | 0.22     | 0.23     | 0.16    | 0.28     | 0.27             | 0.26     | 0.30     |
| Mg#                            | 35            | 36            | 36            | 57       | 61       | 71      | 55       | 48               | 53       | 53       |
| KO/NaO                         | 1.99          | 1.97          | 2.42          | 10.51    | 13.37    | 24.52   | 0.25     | 0.77             | 0.84     | 1.11     |

958

959 **Table 8.** Major, trace element and REE data for a selected group of island-arc volcanic rocks from the Ankara Mélange.

| Sample no                      | CE.962   | CE.964   | CS.07    | CS.11    | CE.96  | CE.98   | CS.99   | MS.34    | MS.35    | MS.36                   | COR.6    | COR.7         | COR.9         | COR.10   |
|--------------------------------|----------|----------|----------|----------|--------|---------|---------|----------|----------|-------------------------|----------|---------------|---------------|----------|
| Rock-type                      | Basanite | Basanite | Basanite | Basanite | Basalt | Basalt  | Basalt  | Basanite | Basanite | Basaltic-trachyandesite | Basanite | Trachy basalt | Trachy basalt | Basanite |
| <b>Oxide, wt %</b>             |          |          |          |          |        |         |         |          |          |                         |          |               |               |          |
| SiO <sub>2</sub>               | 42.28    | 42.78    | 44.35    | 42.69    | 51.27  | 48.78   | 48.28   | 44.49    | 44.56    | 54.26                   | 44.13    | 48.08         | 46.26         | 42.86    |
| TiO <sub>2</sub>               | 0.88     | 0.9      | 0.78     | 0.82     | 0.78   | 1.74    | 1.71    | 1.00     | 0.92     | 0.50                    | 0.84     | 0.84          | 0.85          | 0.93     |
| Al <sub>2</sub> O <sub>3</sub> | 16.57    | 16.84    | 16.10    | 16.55    | 15.91  | 14.85   | 14.89   | 17.60    | 15.63    | 17.93                   | 17.26    | 17.21         | 18.47         | 17.03    |
| Fe <sub>2</sub> O <sub>3</sub> | 12.07    | 12.26    | 11.44    | 11.98    | 9.57   | 12.69   | 12.69   | 9.35     | 10.31    | 6.34                    | 9.40     | 10.17         | 9.49          | 10.21    |
| MnO                            | 0.27     | 0.24     | 0.21     | 0.20     | 0.14   | 0.20    | 0.20    | 0.16     | 0.19     | 0.14                    | 0.18     | 0.19          | 0.16          | 0.20     |
| MgO                            | 8        | 7.37     | 7.67     | 6.68     | 6.44   | 5.78    | 5.72    | 6.82     | 8.84     | 2.44                    | 7.21     | 4.22          | 4.13          | 4.74     |
| CaO                            | 10.83    | 10.42    | 10.47    | 9.48     | 7.64   | 8.26    | 8.54    | 9.17     | 9.84     | 5.97                    | 8.84     | 6.9           | 8.19          | 13.37    |
| Na <sub>2</sub> O              | 2.7      | 2.47     | 3.29     | 3.88     | 4.73   | 3.84    | 3.48    | 2.09     | 2.15     | 3.75                    | 2.89     | 4.32          | 2.35          | 1.28     |
| K <sub>2</sub> O               | 0.73     | 1.57     | 0.51     | 1.06     | 0.12   | 0.59    | 0.75    | 2.39     | 2.33     | 4.10                    | 2.82     | 2.51          | 3.01          | 2.41     |
| P <sub>2</sub> O <sub>5</sub>  | 0.38     | 0.43     | 0.33     | 0.39     | 0.06   | 0.18    | 0.16    | 0.31     | 0.31     | 0.27                    | 0.39     | 0.46          | 0.34          | 0.44     |
| LOI                            | 5        | 4.3      | 4.5      | 6.0      | 3.1    | 2.7     | 3.1     | 6.2      | 4.5      | 4                       | 5.6      | 4.6           | 6.4           | 6.1      |
| Total                          | 99.71    | 99.58    | 99.65    | 99.69    | 99.81  | 99.57   | 99.47   | 99.64    | 99.61    | 99.67                   | 99.56    | 99.48         | 99.67         | 99.62    |
| Mg#                            | 57       | 54       | 57       | 52       | 57     | 47      | 47      | 59       | 63       | 43                      | 60       | 45            | 46            | 48       |
| <b>Trace, ppm</b>              |          |          |          |          |        |         |         |          |          |                         |          |               |               |          |
| Cr                             | 21       | 14       | 68       | 14       | 14     | 41      | 41      | 21       | 82       | 14                      | 55       | 14            | 27            | 41       |
| Ni                             | 29       | 37       | 31.00    | 19.70    | 20.40  | 18.60   | 18.00   | 12.20    | 23.00    | 5.10                    | 16.7     | 1.9           | 15.1          | 17.5     |
| Co                             | 44.5     | 43.7     | 44.10    | 43.40    | 31.80  | 40.10   | 40.70   | 33.20    | 40.60    | 14.70                   | 32.4     | 26.5          | 24.1          | 40.2     |
| Sc                             | 32       | 31       | 41       | 34       | 35     | 37      | 37      | 34       | 49       | 11                      | 31       | 15            | 26            | 33       |
| Rb                             | 46.5     | 65.1     | 11.00    | 42.50    | 3.20   | 8.30    | 10.00   | 58.50    | 55.70    | 129.20                  | 67.7     | 38.3          | 92.4          | 59.3     |
| Ba                             | 636      | 2044     | 596.00   | 613.00   | 36.00  | 882.00  | 1437.00 | 796.00   | 700.00   | 1225.00                 | 885      | 2166          | 902           | 877      |
| Sr                             | 497.4    | 620.5    | 471.90   | 321.40   | 43.40  | 1129.10 | 1385.20 | 497.00   | 460.50   | 546.70                  | 966.6    | 875.8         | 579.5         | 660.2    |
| Cs                             | 1.3      | 1        | 2.90     | 1.50     | 32.70  | 0.70    | 0.40    | 2.90     | 2.70     | 3.70                    | 4.4      | 2.1           | 2.8           | 2.5      |
| Th                             | 3.8      | 4.3      | 3.70     | 3.90     | 0.30   | 0.90    | 0.90    | 12.70    | 11.00    | 22.90                   | 14.3     | 12.1          | 14.3          | 15.8     |
| U                              | 1        | 1.2      | 1.50     | 1.10     | 0.10   | 0.20    | 0.30    | 3.20     | 2.90     | 5.70                    | 3.6      | 3             | 3.1           | 4.1      |
| Nb                             | 3.1      | 3.8      | 4.90     | 5.40     | 4.70   | 7.40    | 7.10    | 7.40     | 6.30     | 10.30                   | 5.9      | 6.3           | 6.4           | 5.9      |
| Ta                             | 0.2      | 0.3      | 2.40     | 1.60     | 3.30   | 3.50    | 3.90    | 0.90     | 1.00     | 1.50                    | 0.3      | 0.3           | 0.4           | 0.3      |
| Zr                             | 63.3     | 65       | 50.60    | 55.20    | 46.10  | 107.30  | 104.80  | 110.90   | 106.30   | 175.80                  | 126.4    | 94.4          | 108.3         | 113.3    |
| Hf                             | 2.2      | 1.9      | 1.50     | 1.60     | 1.50   | 2.90    | 2.90    | 2.90     | 2.90     | 4.10                    | 3.1      | 2.6           | 2.6           | 2.3      |
| Y                              | 19.6     | 20.7     | 18.20    | 19.80    | 20.30  | 34.00   | 34.10   | 25.80    | 24.80    | 27.10                   | 25.3     | 24.3          | 20.4          | 23.8     |
| V                              | 357      | 357      | 365      | 339      | 335    | 379     | 385     | 364.00   | 401.00   | 157.00                  | 292      | 282           | 280           | 388      |
| Pb                             | 3.6      | 4.8      | 5.40     | 8.30     | 0.7    | 0.4     | 0.6     | 8.9      | 8.9      | 7                       | 13.4     | 17.5          | 13.1          | 18.8     |

960

961 **Table 8. Continued.**

| Sample no                          | CE.962   | CE.964   | CS.07    | CS.11    | CE.96  | CE.98  | CS.99  | MS.34    | MS.35    | MS.36                   | COR.6    | COR.7         | COR.9         | COR.10   |
|------------------------------------|----------|----------|----------|----------|--------|--------|--------|----------|----------|-------------------------|----------|---------------|---------------|----------|
| Rock-type                          | Basanite | Basanite | Basanite | Basanite | Basalt | Basalt | Basalt | Basanite | Basanite | Basaltic-trachyandesite | Basanite | Trachy basalt | Trachy basalt | Basanite |
| <b>REE, ppm</b>                    |          |          |          |          |        |        |        |          |          |                         |          |               |               |          |
| La                                 | 17.3     | 17.6     | 14.90    | 16.80    | 3.00   | 8.40   | 8.30   | 37.60    | 33.30    | 60.20                   | 43.1     | 36.2          | 41.1          | 43.8     |
| Ce                                 | 37.6     | 39.5     | 31.60    | 35.20    | 7.70   | 20.90  | 20.40  | 73.30    | 67.00    | 109.70                  | 90.9     | 74.4          | 82.0          | 92.2     |
| Pr                                 | 5.16     | 5.27     | 4.16     | 4.63     | 1.10   | 2.90   | 2.84   | 8.63     | 8.13     | 11.97                   | 9.92     | 8.46          | 8.73          | 9.80     |
| Nd                                 | 22.7     | 23       | 17.60    | 20.10    | 5.20   | 14.50  | 13.70  | 34.50    | 32.80    | 43.60                   | 35.3     | 34.6          | 31.1          | 34.8     |
| Sm                                 | 4.99     | 5.22     | 4.14     | 4.70     | 1.86   | 3.92   | 3.95   | 6.88     | 6.66     | 7.35                    | 7.23     | 6.81          | 5.85          | 6.93     |
| Eu                                 | 1.51     | 1.53     | 1.27     | 1.38     | 0.74   | 1.26   | 1.43   | 1.94     | 1.80     | 1.87                    | 1.96     | 1.86          | 1.55          | 1.79     |
| Gd                                 | 4.87     | 4.94     | 3.97     | 4.36     | 2.49   | 5.00   | 5.13   | 6.21     | 5.88     | 5.83                    | 6.42     | 6             | 5.30          | 6.23     |
| Tb                                 | 0.72     | 0.73     | 0.63     | 0.70     | 0.52   | 0.96   | 0.97   | 0.95     | 0.92     | 0.90                    | 0.94     | 0.92          | 0.76          | 0.89     |
| Dy                                 | 3.92     | 4.06     | 3.23     | 3.47     | 3.17   | 5.54   | 5.76   | 4.85     | 4.66     | 4.68                    | 4.80     | 4.83          | 3.66          | 4.48     |
| Ho                                 | 0.71     | 0.75     | 0.63     | 0.68     | 0.70   | 1.23   | 1.23   | 0.96     | 0.86     | 0.91                    | 0.89     | 0.88          | 0.76          | 0.84     |
| Er                                 | 1.88     | 1.96     | 1.75     | 1.79     | 2.17   | 3.59   | 3.50   | 2.46     | 2.35     | 2.58                    | 2.44     | 2.62          | 2.06          | 2.28     |
| Tm                                 | 0.3      | 0.31     | 0.25     | 0.29     | 0.33   | 0.55   | 0.56   | 0.39     | 0.36     | 0.42                    | 0.35     | 0.36          | 0.30          | 0.34     |
| Yb                                 | 1.78     | 1.78     | 1.56     | 1.67     | 2.11   | 3.35   | 3.34   | 2.38     | 2.23     | 2.67                    | 2.27     | 2.33          | 1.86          | 2.19     |
| Lu                                 | 0.27     | 0.27     | 0.24     | 0.25     | 0.33   | 0.52   | 0.51   | 0.36     | 0.33     | 0.42                    | 0.35     | 0.36          | 0.29          | 0.33     |
| K <sub>2</sub> O/Na <sub>2</sub> O | 0.27     | 0.64     | 0.16     | 0.27     | 0.03   | 0.15   | 0.22   | 1.14     | 1.08     | 1.09                    | 0.98     | 0.58          | 1.28          | 1.88     |

962 **Table 9.** Major, trace element and REE data for a selected group of lamprophyric dike rocks from the Ankara Mélange.

| Sample no                      | DM2      | DM3      | DM4      | DM5      | DM5A     | DM6           | DM7            | DM7A     | DM8           | DM9            | DM10           | DM17     | CE1206       | CE1207       | CE2          | CE1210        | 25BM11          |
|--------------------------------|----------|----------|----------|----------|----------|---------------|----------------|----------|---------------|----------------|----------------|----------|--------------|--------------|--------------|---------------|-----------------|
| Rock-type                      | Tephrite | Tephrite | Tephrite | Tephrite | Tephrite | Trachy-basalt | Phono-tephrite | Tephrite | Trachy-basalt | Phono-tephrite | Phono-tephrite | Tephrite | Picro-basalt | Picro-basalt | Picro-basalt | Trachy-basalt | Trachy-andesite |
| <b>Oxide, wt %</b>             |          |          |          |          |          |               |                |          |               |                |                |          |              |              |              |               |                 |
| SiO <sub>2</sub>               | 45.25    | 43.02    | 47.41    | 45.84    | 46.23    | 48.59         | 50.29          | 50.49    | 49.34         | 47.48          | 47.02          | 46.45    | 41.94        | 47.34        | 44.89        | 51.33         | 58.15           |
| TiO <sub>2</sub>               | 0.69     | 0.76     | 0.58     | 0.64     | 0.66     | 0.73          | 0.62           | 0.62     | 0.70          | 0.73           | 0.74           | 0.78     | 1.12         | 0.83         | 1.26         | 1.24          | 0.43            |
| Al <sub>2</sub> O <sub>3</sub> | 11.20    | 10.93    | 15.79    | 16.98    | 17.00    | 10.34         | 14.94          | 14.73    | 12.10         | 14.37          | 14.66          | 11.40    | 12.91        | 15.66        | 13.36        | 15.62         | 17.43           |
| Fe <sub>2</sub> O <sub>3</sub> | 11.16    | 12.10    | 9.54     | 10.86    | 11.03    | 11.46         | 10.85          | 10.72    | 10.92         | 11.07          | 11.13          | 12.59    | 10.8         | 9.79         | 11.12        | 7.33          | 5.81            |
| MnO                            | 0.23     | 0.22     | 0.23     | 0.21     | 0.21     | 0.22          | 0.20           | 0.18     | 0.21          | 0.24           | 0.23           | 0.22     | 0.2          | 0.21         | 0.22         | 0.08          | 0.14            |
| MgO                            | 5.41     | 4.98     | 2.90     | 4.66     | 4.87     | 6.02          | 4.02           | 4.13     | 4.49          | 3.43           | 3.50           | 5.09     | 7.71         | 4.02         | 8.23         | 5.99          | <b>1.88</b>     |
| CaO                            | 14.50    | 15.89    | 12.25    | 10.25    | 9.57     | 13.68         | 7.30           | 7.53     | 13.50         | 8.44           | 8.53           | 12.26    | 16.47        | 9.11         | 14.37        | 8.22          | 4.04            |
| Na <sub>2</sub> O              | 0.67     | 0.28     | 0.36     | 1.63     | 1.68     | 0.56          | 2.84           | 2.78     | 1.96          | 2.64           | 2.95           | 1.58     | 1.62         | 2.97         | 1.70         | 3.95          | 5.56            |
| K <sub>2</sub> O               | 5.47     | 5.54     | 6.96     | 4.67     | 4.99     | 5.73          | 5.53           | 5.50     | 4.11          | 6.34           | 5.82           | 5.40     | <b>0.72</b>  | 4.5          | 1.01         | 1.85          | 4.32            |
| P <sub>2</sub> O <sub>5</sub>  | 0.89     | 0.80     | 0.45     | 0.44     | 0.40     | 1.08          | 0.96           | 0.93     | 0.89          | 0.78           | 0.78           | 0.90     | 0.49         | 0.64         | 0.55         | 0.75          | 0.29            |
| LOI                            | 4        | 4.9      | 2.9      | 3.4      | 3.0      | 1.1           | 2.0            | 1.9      | 1.3           | 3.7            | 3.9            | 2.8      | 5.8          | 4.6          | 2.9          | 3.4           | 1.6             |
| Total                          | 99.5     | 99.46    | 99.42    | 99.61    | 99.60    | 99.49         | 99.53          | 99.53    | 99.52         | 99.27          | 99.28          | 99.48    | 99.78        | 99.67        | 99.64        | 99.76         | 99.64           |
| <b>Trace, ppm</b>              |          |          |          |          |          |               |                |          |               |                |                |          |              |              |              |               |                 |
| Cr                             | 13.70    | 13.70    | 13.70    | 13.70    | 13.70    | 13.70         | 13.70          | 13.70    | 13.70         | 13.70          | 13.70          | 13.70    | 89           | 21           | 82           | <b>158</b>    | 68              |
| Ni                             | 20       | 20       | 20       | 20       | 20       | 20            | 20             | 20       | 20            | 20             | 20             | 20       | 48           | 28           | 67           | 86            | 45              |
| Co                             | 33.0     | 39.4     | 26.9     | 34.8     | 34.5     | 37.9          | 34.1           | 33.8     | 32.2          | 28.5           | 28.9           | 41.2     | 35.5         | 24.2         | 39.7         | 24.9          | 12.50           |
| Sc                             | 30       | 34       | 10       | 25       | 27       | 45            | 24             | 25       | 30            | 21             | 22             | 36       | 41           | 22           | 39           | 19            | 7               |
| Rb                             | 51.4     | 52.6     | 94.6     | 76.9     | 81.5     | 46.9          | 65.0           | 67.7     | 29.7          | 47.3           | 43.1           | 50.0     | 20.8         | 75.5         | 13.8         | 29.9          | 76.5            |
| Ba                             | 1861     | 1881     | 2899     | 1224     | 1228     | 1760          | 1183           | 1233     | 1846          | 3229           | 3172           | 2019     | 180          | 1109         | 257          | 475           | 1456            |
| Sr                             | 697.0    | 864.1    | 762.5    | 701.6    | 811.3    | 823.9         | 1483.3         | 1401.6   | 758.9         | 1287.1         | 1335.2         | 790.0    | 1073         | 1116         | 805.4        | 1006          | 679.1           |
| Cs                             | 0.1      | 0.1      | 0.2      | 0.3      | 0.2      | 0.1           | 0.1            | 0.1      | 0.1           | 0.4            | 0.4            | 0.1      | 0.3          | 0.6          | 0.1          | 0.2           | 0.1             |
| Th                             | 24.4     | 14.0     | 19.7     | 8.9      | 9.1      | 21.9          | 14.2           | 14.4     | 23.1          | 34.6           | 33.6           | 13.2     | 7.3          | 14.9         | 6.9          | 5.4           | 25.0            |
| U                              | 6.0      | 5.6      | 7.4      | 2.9      | 3.1      | 5.4           | 7.1            | 6.9      | 6.0           | 10.1           | 10.5           | 5.1      | 2.4          | 4.9          | 2.5          | 1.0           | 6.9             |
| Nb                             | 18.3     | 22.7     | 18.4     | 7.8      | 7.9      | 17.0          | 13.7           | 13.9     | 18.6          | 34.1           | 33.9           | 21.7     | 9.7          | 13.1         | 15.1         | 7.4           | 10.7            |
| Ta                             | 0.6      | 0.7      | 0.7      | 0.3      | 0.3      | 0.7           | 0.5            | 0.5      | 0.6           | 1.1            | 1.2            | 0.7      | 0.6          | 0.9          | 1.0          | 1             | 0.6             |
| Zr                             | 111.4    | 111.2    | 117.3    | 65.0     | 62.3     | 103.2         | 95.1           | 93.8     | 112.6         | 188.1          | 186.3          | 101.6    | 92.9         | 163.5        | 111.9        | 121.9         | 199.7           |
| Hf                             | 2.8      | 2.9      | 2.8      | 1.6      | 1.4      | 2.7           | 2.5            | 2.3      | 2.8           | 4.7            | 4.4            | 2.6      | 2.5          | 4.1          | 2.8          | 3.1           | 4.3             |
| Y                              | 23.7     | 24.5     | 22.1     | 16.2     | 15.1     | 24.4          | 21.5           | 20.4     | 24.8          | 34.2           | 34.4           | 23.7     | 21.9         | 34.8         | 24.7         | 16.9          | 28.9            |
| V                              | 305      | 403      | 405      | 307      | 299      | 329           | 262            | 269      | 347           | 329            | 329            | 370      | 300          | 245          | 330          | 162           | 116             |
| Pb                             | 32.3     | 17.0     | 11.1     | 16.8     | 15.5     | 14.8          | 9.6            | 8.8      | 9.4           | 12.6           | 11.3           | 20.2     | 2.3          | 6.6          | 0.9          | 1.30          | 33.9            |
| Mg#                            | 49       | 45       | 38       | 46       | 47       | 51            | 42             | 43       | 45            | 38             | 38             | 44       | 59           | 45           | 59           | 62            | 39              |
| KO/NaO                         | 8.16     | 19.79    | 19.33    | 2.87     | 2.97     | 10.23         | 1.95           | 1.98     | 2.10          | 2.40           | 1.97           | 3.42     | 0.44         | 1.52         | 0.59         | 0.47          | 0.78            |

| Sample no                            | DM2      | DM3      | DM4      | DM5      | DM5A     | DM6           | DM7            | DM7A     | DM8           | DM9            | DM10           | DM17     | CE1206       | CE1207       | CE2          | CE1210        | 25BM11          |
|--------------------------------------|----------|----------|----------|----------|----------|---------------|----------------|----------|---------------|----------------|----------------|----------|--------------|--------------|--------------|---------------|-----------------|
| Rock-type                            | Tephrite | Tephrite | Tephrite | Tephrite | Tephrite | Trachy-basalt | Phono-tephrite | Tephrite | Trachy-basalt | Phono-tephrite | Phono-tephrite | Tephrite | Picro-basalt | Picro-basalt | Picro-basalt | Trachy-basalt | Trachy-andesite |
| <b>REE, ppm</b>                      |          |          |          |          |          |               |                |          |               |                |                |          |              |              |              |               |                 |
| La                                   | 60.0     | 41.8     | 49.5     | 26.4     | 25.0     | 56.6          | 32.2           | 30.7     | 58.4          | 83.6           | 81.2           | 41.4     | 26.3         | 43.2         | 29.3         | 33.3          | 69.2            |
| Ce                                   | 120.4    | 87.4     | 97.8     | 52.4     | 49.7     | 115.9         | 69.8           | 67.3     | 119.6         | 159.0          | 156.7          | 86.5     | 53.1         | 88.9         | 58.1         | 70.7          | 124.9           |
| Pr                                   | 13.07    | 9.79     | 10.41    | 5.75     | 5.52     | 13.12         | 7.99           | 7.71     | 13.19         | 16.87          | 16.92          | 9.76     | 6.83         | 10.9         | 7.52         | 8.76          | 13.24           |
| Nd                                   | 49.9     | 39.3     | 38.7     | 22.2     | 22.3     | 53.9          | 32.7           | 32.0     | 51.5          | 63.2           | 63.9           | 39.6     | 28.2         | 42.5         | 29.8         | 33.1          | 46.9            |
| Sm                                   | 9.36     | 8.07     | 7.17     | 4.49     | 4.29     | 9.83          | 6.15           | 5.93     | 9.48          | 11.52          | 11.64          | 7.84     | 5.98         | 8.52         | 6.96         | 5.67          | 7.82            |
| Eu                                   | 2.26     | 2.02     | 1.86     | 1.20     | 1.14     | 2.35          | 1.53           | 1.50     | 2.27          | 2.84           | 2.88           | 2.00     | 1.72         | 2.49         | 1.98         | 1.59          | 1.93            |
| Gd                                   | 7.94     | 7.09     | 6.23     | 4.01     | 3.88     | 8.37          | 5.32           | 5.22     | 7.97          | 10.04          | 10.03          | 7.19     | 5.68         | 8.09         | 6.36         | 4.56          | 6.22            |
| Tb                                   | 1.04     | 0.97     | 0.88     | 0.59     | 0.56     | 1.07          | 0.77           | 0.76     | 1.07          | 1.36           | 1.36           | 0.95     | 0.86         | 1.21         | 0.93         | 0.67          | 0.90            |
| Dy                                   | 4.95     | 4.71     | 4.31     | 2.95     | 2.97     | 4.94          | 3.94           | 3.91     | 4.88          | 6.88           | 6.66           | 4.72     | 4.06         | 6.33         | 4.68         | 3.28          | 4.98            |
| Ho                                   | 0.82     | 0.82     | 0.73     | 0.55     | 0.55     | 0.79          | 0.74           | 0.73     | 0.84          | 1.16           | 1.19           | 0.80     | 0.79         | 1.2          | 0.81         | 0.61          | 0.91            |
| Er                                   | 2.05     | 2.12     | 2.00     | 1.50     | 1.45     | 2.10          | 1.96           | 1.94     | 2.08          | 3.08           | 3.02           | 2.10     | 2.09         | 3.27         | 2.30         | 1.64          | 2.63            |
| Tm                                   | 0.29     | 0.29     | 0.27     | 0.22     | 0.21     | 0.28          | 0.29           | 0.27     | 0.29          | 0.43           | 0.44           | 0.30     | 0.32         | 0.52         | 0.33         | 0.25          | 0.42            |
| Yb                                   | 1.65     | 1.79     | 1.83     | 1.41     | 1.33     | 1.69          | 1.74           | 1.76     | 1.69          | 2.67           | 2.80           | 1.73     | 1.76         | 2.94         | 1.89         | 1.47          | 2.66            |
| Lu                                   | 0.25     | 0.27     | 0.26     | 0.21     | 0.20     | 0.25          | 0.27           | 0.26     | 0.25          | 0.41           | 0.41           | 0.26     | 0.27         | 0.46         | 0.27         | 0.22          | 0.43            |
| <sup>87</sup> Sr/ <sup>86</sup> Sr   |          | 0.704786 |          |          | 0.704892 |               |                | 0.704697 | 0.704720      |                |                | 0.704797 |              |              | 0.704820     |               |                 |
| <sup>143</sup> Nd/ <sup>144</sup> Nd |          | 0.512681 |          |          | 0.512686 |               |                | 0.512690 | 0.512682      |                |                | 0.512680 |              |              | 0.512674     |               |                 |
| <sup>206</sup> Pb/ <sup>204</sup> Pb |          | 19.540   |          |          | 19.332   |               |                | 19.939   | 19.604        |                |                | 19.594   |              |              | 19.418       |               |                 |
| <sup>207</sup> Pb/ <sup>204</sup> Pb |          | 15.662   |          |          | 15.655   |               |                | 15.691   | 15.675        |                |                | 15.659   |              |              | 15.664       |               |                 |
| <sup>208</sup> Pb/ <sup>204</sup> Pb |          | 39.376   |          |          | 39.192   |               |                | 39.612   | 39.536        |                |                | 39.407   |              |              | 39.297       |               |                 |