

This discussion paper is/has been under review for the journal Solid Earth (SE).
Please refer to the corresponding final paper in SE if available.

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi¹, L. Gholami^{1,*}, E. Sharifi Moghadam¹, and A. Khaledi Darvishan¹

¹Department of Watershed Management Engineering, Faculty of Natural Resources, Tarbiat Modares University, P. O. Box 46417-76489, Noor, Iran

*present at: Department of Rangeland and Watershed Management, Faculty of Natural Resources, Sari Agricultural Sciences and Natural Resources University, Sari, Iran

Received: 10 September 2014 – Accepted: 10 September 2014 – Published: 9 October 2014

Correspondence to: S. H. R. Sadeghi (sadeghi@modares.ac.ir)

Published by Copernicus Publications on behalf of the European Geosciences Union.

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Amendments can control the runoff and soil loss by protecting soil surface. However, scale effects on runoff and soil loss control has not been considered yet. The present study has been formulated to determine the efficiency of two plot sizes of 6 and 0.25 m² covered by straw mulch with rate of 0.5 kg m⁻² in changing the time to runoff, runoff coefficient, sediment concentration and soil loss under laboratory conditions. The study has been conducted for a sandy-loam soil taken from summer rangeland, Alborz Mountains, Northern Iran under simulated rainfall intensities of 50 and 90 mm h⁻¹ and in 3 replicates. The results of the study showed that the straw mulch had more significant effect in in reducing runoff coefficient, sediment concentration and soil loss at 0.25 m² plot scale. The maximum effectiveness in time to runoff for both the scales, observed in rainfall intensity of 90 mm h⁻¹. The maximum increasing and decreasing rates in time to runoff and runoff coefficient observed in the rainfall intensity of 90 mm h⁻¹ with the amounts of 367.92 and 96.71 % for 0.25 m² plot and the amounts of 110.10 and 15.08 % for 6 m² plot respectively. The maximum change of soil loss in both the intensities of 50 and 90 mm h⁻¹ occurred at 0.25 m² plot with the amount of 100 % whereas at 6 m² plot, decreasing rates of soil loss for in both the intensities of 50 and 90 mm h⁻¹ were 46.74 and 63.24 %, respectively.

1 Introduction

There are various methods for soil conservation but biological methods in bare and degraded slopes need long time for establishment (Adekalu et al., 2007; Smets et al., 2008a). In this context, various natural and organic mulches viz. crop residues, leaf litter, woodchips, bark chips, biological geotextiles, gravel and crushed stones (Ruy, 2006; Smets et al., 2008a; Ruiz-Sinoga et al., 2010) have been applied for soil conservation. Mulches have extraordinary potential in soil erosion control (Morgan, 1986) and runoff reduction (Poesen and Lavee, 1991). However, establishment of degraded areas

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and bare slopes by vegetation cover takes long time (Adekalu et al., 2007; Smets et al., 2008a). The effect of mulches depends on many factors including raindrops erosivity, soil condition, steepness and length of slope, and the mulch rate and type (Amimoto, 1981; Cogo et al., 1984; Poesen and Lavee, 1991; Morgan, 1995; Auerswald et al., 2003; Adekalu et al., 2007; Kukal and Sarkar, 2010; Jordán et al., 2010; Choi et al., 2012; Gholami et al., 2013).

Although, there are a lot of studies about soil amendments as soil conservation e.g. Fernández et al. (2012), Jiménez et al. (2012), García-Moreno et al. (2013), Robichaud et al. (2013) and Martins et al. (2014) but the effects of study scale on effectiveness of various mulch covers has been rarely considered. The present study has been conducted to determine the effects of spatial scale on the effectiveness of rice straw mulch on runoff and soil loss for a sandy-loam soil taken from summer rangeland in the Alborz Mountains, Northern Iran. The study was taken place under laboratory conditions with simulated rainfall intensities of 50 and 90 mm h⁻¹ in two scales of 0.25 and 6 m² with constant gradient of 30 % at rainfall simulation and erosion laboratory of Tarbiat Modares University, Noor, Iran. There are a few studies about spatial scale variations mulches on runoff and soil loss. Poesen et al. (1994) reviewed the effects of rock fragments on soil erosion and stated the spatial scale has an important impact on the soil erosion. They showed that at the microplot scale, 4 mm² to 1 m², sediment yield reached a maximum value with 0 % rock fragment cover and reached minimum value with 100 % rock fragment cover. At the mesoplot scale (i.e. interrill areas), negative, positive as well as convex upward relationships with cover percentages have been observed, depending on the fine earth structure, on the vertical position in the topsoil, on the size of rock fragments and on the surface slope. Finally, at the macroplot scale (i.e. interrill and rill areas), 10–10 000 m², sediment yield decreased exponentially with rock fragment cover. Cerdan et al. (2002) investigated scale effect (plot to catchment) on runoff in agricultural areas of Normandy, France. Three database of 450 m² plots, a 90 ha catchment and an 1100 ha catchment were selected to collect runoff data. Between the three scales, a significant decrease in the runoff coefficient was observed

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as the area increased. Mingguo et al. (2007) also studied the effect of vegetation on runoff-sediment yield relationship at different spatial scales (plot to watershed) in hilly areas of the Loess Plateau, North China and found that the vegetation could reduce runoff and soil loss in both scales but the reduction rate of sediment was more than runoff at plot scale. Smets et al. (2008a) reviewed the impact of plot length on the effectiveness of different soil-surface covers in reducing runoff and soil loss. The results indicated that for plot lengths < 11 m, there was a large variation in the runoff and erosion-reducing effectiveness of each soil cover, depending on various factors. Smets et al. (2008b) also examined the spatial scale effects on the effectiveness of organic mulches in reducing soil erosion at field and laboratory experiments (plot length ranges between 0.1 and 30.5 m). Results verified the effectiveness of mulches in reducing soil erosion by water in various scales. In addition, they reported a positive linear relation between the erosion-reducing effectiveness of an organic mulch cover and plot length. On short plots, the response of a soil surface cover on runoff and soil loss was immediately observed. Nevertheless, on longer plots, the runoff and soil loss response could be compensated due to the longer plot length. Liu et al. (2012) evaluated the effects of rice straw mulch and plastic film mulching at plot scale and 2 years in the Xiaofuling watershed in the Danjiangkou Reservoir area, China. The straw mulch treatment significantly decreased the sediment yield from 18 to 22 %. The results showed that the straw mulch was beneficial for controlling runoff and sediment.

Scrutinizing the available literatures showed that although there are lots of references on using straw as mulch for runoff and soil erosion control, but there was no literature in regards to report the effectiveness of straw mulch in various plot scales. The present study was therefore planned to determine the efficiency of two plot sizes covered by straw mulch changing the important runoff and soil loss components under laboratory conditions.

2 Materials and methods

2.1 Erosion plots and rainfall simulator

The laboratory experiments were conducted by using two sets of 6m × 1m and 0.5m × 0.5m plots installed in the rainfall simulator laboratory, Faculty of Natural Resources of Tarbiat Modares University (TMU), located in Noor Mazandaran Province, Northern Iran. The experiments were carried out to study the effect of rice straw mulch on runoff and soil loss processes by using simulated rainfall in intensities of 50 and 90 mm h⁻¹ and in 3 replicates. The rainfall simulator consists of a 4000 L water tank and 27 precalibrated nozzles in three parallel lines designed to simulate raindrops of 1.3 mm average size. The drops fall from a height between 4 and 6 m at the upper and lower parts of the plot, respectively, reaching a 7 m s⁻¹ speed (Duiker et al., 2001) the study plot.

2.2 Soil preparation, placement and rice straw mulch application

A sandy-loam topsoil was collected 0–20 cm (Kukul and Sarkar, 2010) the Alborz Mountains, Northern Iran. The soil was transport to the lab and air-dried up to optimum moisture content to maintain the relative stability of soil aggregates and decrease breaking down the aggregates in sieving process (Khaledi Darvishan et al., 2013). The coarse rock fragments and plant residues were removed from the soil through passing from 8 mm sieve to obtain maximum homogeneity in soil profile (Hawke et al., 2006).

Three layers of mineral pumice grains with different sizes and total thickness of 15 cm were used as a filter layer and placed at the bottom of the plots in order to simulate natural drainage condition and decreasing plot weight (Defersha et al., 2011). A 15 cm-thick soil layer was then placed on the top and separated from the mineral pumice by a sheet of porous jute (Defersha et al., 2011). The soil was ultimately compacted by a small PVC roller (a hand-made roller and filled with cement and sand) to achieve the bulk density of 1.38 g cm⁻³ almost equal to that measure for the soil under natural

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

The present study was done to determine the efficiency of scale change in changing time to runoff, runoff coefficient and soil loss for a rangeland sandy-loam soil in Northern Iran. The experiments were conducted in 6 and 0.25 m² plots and under lab conditions with simulated rainfall intensities of 50 and 90 mm h⁻¹.

3.1 Runoff

The amounts of time to runoff and runoff coefficient before and after conservation treatment in each plot output and each scale are shown in Table 1. The percentage of changes in study variables in treated plots and in comparison with control plots have been summarized in Table 2. Figures 2 and 3 also show the average rates of time to runoff and coefficient in the both scales.

3.2 Soil loss

The sediment concentration and soil loss amounts before and after conservation treatment in each scale have been shown in Table 3. The relative effectiveness of straw mulch on sediment concentration and soil loss for two scales has also been summarized in Table 4. Figures 4 and 5 accordingly show the average rates of sediment concentration and soil loss in two study plots. Also the results of statistical analysis based on the GLM has been summarized in Table 5.

4 Discussion

4.1 Runoff

Tables 1 and 2 showed that the straw mulch increased time to runoff compared to untreated plots except in rainfall intensity of 50 mm h⁻¹ for 0.25 m² plot and also de-

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



creased runoff coefficient in both the scales. It might be due to water storing effects of straw and also increasing ponding time on the plot surface. This finding is in the same line with that reported by Poesen and Lavee (1991), Mingguo et al. (2007) and Smets et al. (2008a, b). Though the maximum change effectiveness in time to runoff, for two scales, could be found in rainfall intensity of 90 mm h^{-1} . These effects were more serious in 0.25 m^2 plot with rate of $+367.92 \%$. While, 6 m^2 plot compared to 0.25 m^2 plot could reduce the time to runoff in rainfall intensity of 50 mm h^{-1} with rate of $+106.15 \%$. Scrutinizing Table 2 and Figs. 2 and 3 also verified the varying effect of straw mulch on runoff coefficient from -10.43 to -96.71% in two scales. The minimum and the maximum effects were also in rainfall intensities of 50 in 6 m^2 plot with rate of -10.43% and 90 in 0.25 m^2 plot with rate of 96.71% mm h^{-1} , respectively. The 0.25 m^2 plot had the maximum reduction in runoff coefficient for rainfall intensities of 50 and 90 mm h^{-1} . These results showed that the 0.25 m^2 plot had the maximum impact on decreasing runoff coefficient and increasing time to runoff except in case of rainfall intensity of 50 mm h^{-1} . It verified that the straw mulch pieces could store more runoff leading to more infiltration as already reported by Poesen and Lavee (1991), Choi et al. (2012) and Liu et al. (2012). The results showed that there were large variation in the runoff coefficient (Smets et al., 2008a) and time to runoff on 0.25 m^2 plots compared to those recoded for 6 m^2 plots in various rainfall intensities. In this study the effectiveness of mulch in reducing runoff was influenced by the plot size. So that, with increasing plot size the runoff amount increased while the Poesen et al. (1994), Cerdan et al. (2002) and Smets et al. (2008a, b) showed that runoff amount decreased with increasing plot size. The differences between mulch type, application manner and density as well as soil type and rainfall intensity could be supposed as potential reasons behind the disagreement. But, according to McGregor et al. (1988), plot border effects on runoff rates were much more important in small plots compared to large ones.

4.2 Soil loss

Tables 3 and 4 showed that the conservation treatment essentially reduced soil loss which is consistent with Poesen and Lavee (1991) and Fernández and Vega (2014). Sediment concentration also decreased in treated plots as similarly reported by Poesen and Lavee (1991) and Smets et al. (2008a, b). This indicated that the flow could not get enough power to detach particles. A similar finding has been reported by Poesen and Lavee (1991).

Table 4 showed that the amounts of sediment concentration at two study scales changed from -43.47 to -100% . The maximum change occurred at 0.25 m^2 plot in both the intensities of 50 and 90 mm h^{-1} (Figs. 4 and 5). So that, the soil loss was found negligible after mulching in small plot of 0.25 m^2 (Poesen et al., 1994). The results also showed that the soil loss reduced at 0.25 and 6 m^2 plots and also the variation ranged from -58.69 to -100% (Table 4). Poesen et al. (1994) found that the soil loss reduced by 100% in small plots 1 m^2 of with cover 100% . It was also observed that both the study variables got the maximum effect in small plot of 0.25 m^2 in view point of decreasing sediment concentration and soil loss. It has also been verified by Mingguo et al. (2007) that the soil loss by water erosion in laboratory condition reduced as plot size decreased. Poesen and Lavee (1994) and Smets et al. (2008a, b) also stated that the soil loss by water erosion was influenced by the plot length. They showed that the small plots with mulch cover were significantly less effective in reducing relative soil loss compared to longer plots. Whereas, this study stated that the small plot with straw mulch was more effective in reducing runoff and soil loss amounts (Mingguo et al., 2007). Therefore, effectiveness of mulch cover in reducing runoff and soil loss by water erosion decreased with increasing plot size. These results were not consistent with Poesen et al. (1994) and Smets et al. (2008a, b), whereas it agreed Mingguo et al. (2007).

Poesen et al. (1994), Cerdan et al. (2002), Boix-Fayos et al. (2006) and Smets et al. (2008a, b) showed that plot length (or spatial scale) can be important in vari-

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ations of runoff or soil loss rates and in the effectiveness of surface covers. These results were found to be important in designing runoff production and erosion plots and modeling runoff and soil loss rates (Smets et al., 2008a).

According to Table 5, changing plot size could have significant effect ($P > 0.01$) on time to runoff and coefficient, sediment concentration and soil loss. The runoff coefficient ($p = 0.00$), sediment concentration ($p = 0.00$) and soil loss ($p = 0.02$) were significantly influenced by plot size as well as conservation treatment of rice straw mulch. The interaction effect of plot size and conservation treatment on runoff coefficient, sediment concentration and soil loss were also significant with respective p values of 0.001, 0.002 and 0.02. However time to runoff was only influenced by plot size.

5 Conclusions

The present study was conducted to study the effects of plot size treated by rice straw mulch on runoff and soil loss control under two rainfall intensities of 50 and 90 mm h⁻¹. It can be concluded from the results that the straw mulching at rate of 0.5 kg m⁻² and two plot scales of 0.25 and 6 m² could increase the time to runoff and decrease runoff coefficient, sediment concentration and soil loss. The straw mulch increased time to runoff compared to untreated plots except in rainfall intensity of 50 mm h⁻¹ for 0.25 m² plot and also decreased runoff coefficient in both the scales. The maximum change effectiveness in time to runoff, for two scales, could be found in rainfall intensity of 90 mm h⁻¹. The maximum change of soil loss occurred at 0.25 m² plot in both the intensities of 50 and 90 mm h⁻¹. The results showed that the 0.25 m² plot had the better effectiveness in reducing runoff coefficient, sediment concentration and soil loss. The results of the study clearly proved the different responses of the plots in regards to runoff soil loss components which can be practically applied at time of setting up experimental studies. The results further showed that the plots are mainly advised to be used for comparative studies rather those leading to accurate data on larger scale outcomes. Though, further studies with deeper insights to hydraulic aspects of

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the phenomena and more different plot sizes and under various rainfall intensities and conservation measures are essentially needed to draw more reliable conclusion.

References

- Adekalu, K. O., Olorunfemi, I. A., and Osunbitan, J. A.: Grass mulching effect on infiltration, surface runoff and soil loss of three agricultural soils in Nigeria, *Bioresource Technol.*, 98, 912–917, 2007.
- Amimoto, P. Y.: Erosion and Sediment Control Handbook, California Department of Conservation Report No. EPA 4 40/3-78-003, 1981.
- Auerswald, K., Kainz, M., and Fiener, P.: Soil erosion potential of organic versus conventional farming evaluated by USLE modeling of cropping statistics for agricultural districts in Bavaria, *Soil Use Manage.*, 19, 305–311, 2003.
- Biro, K., Pradhan, B., Buchroithner, M., Makeschin, F.: Land use/land cover change analysis and its impact on soil properties in the Northern part of Gadarif region, Sudan, *Land Degrad. Dev.*, 24, 90–102, 2013.
- Boix-Fayos, C., Martínez-Mena, M., Arnau-Rosalén, E., Calvo-Cases, A., Castillo, V., and Albaladejo, J.: Measuring soil erosion by field plots: understanding the sources of variation, *Earth-Sci. Rev.*, 78, 267–85, 2006.
- Cerdà, A., Giménez-Morera, A., and Bodí, M. B.: Soil and water losses from new citrus orchards growing on sloped soils in the western Mediterranean basin, *Earth Surf. Proc. Land.*, 34, 1822–1830, 2009.
- Cerdà, A., Hooke, J., Romero-Díaz, A., Montanarella, L., and Lavee, H.: Soil erosion on Mediterranean type-ecosystems, *Land Degrad. Dev.*, 21, 71–74, doi:10.1002/ldr.968, 2010.
- Cerdan, O. Y., Le Bissonnais, V., Souchere, P. M., and Lecomte, V.: Sediment concentration in interrill flow: interactions between soil surface conditions, vegetation and rainfall, *Earth Surf. Proc. Land.*, 27, 193–205, 2002.
- Choi, J., Shin, M. H., Yoon, J. S., and Jang, J. R.: Effect of Rice Straw Mulch on Runoff and NPS Pollution Discharges from a Vegetable Field, *International Conference of Agriculture Engineering*, 8–12 July, Spain, 4 pp., 2012.
- Cogo, N. P., Moldenhauer, W. C., and Foster, G. R.: Soil loss reductions from conservation tillage practices, *Soil Sci. Soc. Am. J.*, 48, 368–373, 1984.

SED

6, 2915–2938, 2014

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Das, D. K. and Agrawal, R. P.: Physical properties of soils, in: Fundamentals of Soil Science, J. Indian Soci. Soil Sci., New Delhi, 283–295, 2002.
- Defersha, M. B., Quraishi, S., and Melesse, A.: The effect of slope steepness and antecedent moisture content on interrill erosion, runoff and sediment size distribution in the highlands of Ethiopia, *Hydrol. Earth Syst. Sci.*, 15, 2367–2375, doi:10.5194/hess-15-2367-2011, 2011.
- Duiker, S. W., Flanagan, D. C., and Lal, R.: Erodibility and infiltration characteristics of five major soils of southwest Spain, *Catena*, 45, 103–121, 2001.
- Fernández, C. and Vega, J. A.: Efficacy of bark strands and straw mulching after wildfire in NW Spain: effects on erosion control and vegetation recovery, *Ecol. Eng.*, 63, 50–57, 2014.
- Fernández, C., Vega, J. A., Jiménez, E., Vieira, D. C. S., Merino, A., Ferreiro, A., and Fonturbel, T.: Seeding and mulching+seeding effects on post-fire runoff, soil erosion and species diversity in Galicia (NW Spain), *Land Degrad. Dev.*, 23, 150–156, 2012.
- García-Moreno, J., Gordillo-Rivero, A., Zavala, L. M., Jordán, A., and Pereira, P.: Mulch application in fruit orchards increases the persistence of soil water repellency during a 15-years period, *Soil Till. Res.*, 130, 62–68, 2013.
- García-Orenes, F., Cerdà, A., Mataix-Solera, J., Guerrero, C., Bodí, M. B., Arcenegui, V., Zornoza, R., and Sempere, J. G.: Effects of agricultural management on surface soil properties and soil-water losses in eastern Spain, *Soil Till. Res.*, 106, 117–123, 2009.
- García-Orenes, F., Guerrero, C., Roldán, A., Mataix-Solera, J., Cerdà, A., Campoy, M., Zornoza, R., Bárcenas, G., and Caravaca, F.: Soil microbial biomass and activity under different agricultural management systems in a semiarid Mediterranean agroecosystem, *Soil Till. Res.*, 109, 110–115, 2010.
- García-Orenes, F., Roldán, A., Mataix-Solera, J., Cerdà, A., Campoy, M., Arcenegui, V., and Caravaca, F.: Soil structural stability and erosion rates influenced by agricultural management practices in a semi-arid Mediterranean agro-ecosystem, *Soil Use Manage.*, 28, 571–579, 2012.
- Gholami, L., Sadeghi, S. H. R., and Homaei, M.: Straw mulching effect on splash erosion, runoff and sediment yield from eroded plots, *Soil Sci. Soc. Am. J.*, 77, 268–278, 2013.
- Giménez Morera, A., Ruiz Sinoga, J. D., Cerdà, A.: The impact of cotton geotextiles on soil and water losses in Mediterranean rainfed agricultural land, *Land Degrad. Dev.*, 210–217, 2010.
- Hawke, R. M., Price, A. G., and Bryan, R. B.: The effect of initial soil water content and rainfall intensity on near-surface soil hydrologic conductivity: a laboratory investigation, *Catena*, 65, 237–246, 2006.

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jordán, A. and Zavala, L. M.: Gil J. Effects of mulching on soil physical properties and runoff under semi-arid conditions in southern Spain, *Catena*, 81, 77–85, 2010.
- Jordán, A., Zavala, L. M., and Muñoz-Rojas, M.: Mulching, effects on soil physical properties in: *Encyclopedia of Agrophysics*, edited by: Glinski, J., Horabik, J., and Lipiec, J., Springer, Berlin, 492–496, 2011.
- Khaledi Darvishan, A. V., Sadeghi, S. H. R., Homaei, M., and Arabkhedri, M.: Measuring sheet erosion using synthetic color-contrast aggregates, *Hydrol. Process.*, 9, doi:10.1002/hyp.9956, 2013.
- Kukal, S. S. and Sarkar, M.: Splash erosion and infiltration in relation to mulching and polyvinyl alcohol application in semi-arid tropics, *Arch. Agronom. Soil Sci.*, 56, 697–705, 2010.
- Kukal, S. S. and Sarkar, M.: Laboratory simulation studies on splash erosion and crusting in relation to surface roughness and raindrop size, *J. Indian Soci. Soil Sci.*, 59, 87–93, 2011.
- Lieskovský, J. and Kenderessy, P.: Modelling the effect of vegetation cover and different tillage practices on soil erosion in vineyards: a case study in Vráble (Slovakia) using watem/sedem, *Land Degrad. Dev.*, 25, 288–296, 2014.
- Liu, Y., Tsoa, Y., Wana, K. Y., Zhanga, G. S., Liub, D. B., Xiongb, G. Y., and Chena, F.: Runoff and nutrient losses in citrus orchards on sloping land subjected to different surface mulching practices in the danjiangkou reservoir area of China, *Agr. Water Manage.*, 110, 34–40, 2012.
- Mandal, D. and Sharda, V. N.: Appraisal of soil erosion risk in the Eastern Himalayan region of India for soil conservation planning, *Land Degrad. Dev.*, 24, 430–437, 2013.
- McGregor, K. C., Bengtson, R. L., and Mutchler, C. K.: Effects of surface straw on interrill runoff and erosion of grenada silt loam, *T. ASAE*, 31, 111–116, 1988.
- Mingguo, Zh., Qiangguo, C., and Hao, Ch.: Effect of vegetation on runoff-sediment yield relationship at different spatial scales in hilly areas of the Loess Plateau, North China, *Acta Ecologica Sin.*, 27, 3572–3581, 2007.
- Morgan, R. P. C.: *Soil Erosion and Conservation*, Longman Scientific and Technical, Burnt Mile, Harlow, UK, 1986.
- Morgan, R. P. C.: *Soil Erosion and Conservation*, Longman, Essex, England, 1995.
- Poesen, J. W. A. and Lavee, H.: Effects of size and incorporation of synthetic mulch on runoff and sediment yield from interrills in a laboratory study with simulated rainfall, *Soil Till. Res.*, 21, 209–223, 1991.
- Poesen, J. W., Torri, D., and Bunte, K.: Effects of rock fragments on soil erosion by water at different spatial scales: a review, *Catena*, 23, 141–66, 1994.

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



- Prats, S. A., Macdonald, L. H., Monteiro, M., Coelho, C. O. A., and Keizer, J. J.: Effectiveness of forest residue mulching in reducing post-fire runoff and erosion in a pine and a eucalypt plantation in north-central Portugal, *Geoderma*, 191, 115–124, 2012.
- Prats, S. A., Martins, M. A. D. S., Malvar, M. C., Ben-Hur, M., and Keizer, J. J.: Polyacrylamide application versus forest residue mulching for reducing post-fire runoff and soil erosion, *Sci. Total Environ.*, 468–469, 464–474, 2014.
- Prokop, P. and Poreba, G. J.: Soil erosion associated with an upland farming system under population pressure in Northeast India, *Land Degrad. Dev.*, 23, 310–321, 2012.
- Robichaud, P. R., Lewis, S. A., Wagenbrenner, J. W., Ashmun, L. E., and Brown, R. E.: Post fire mulching for runoff and erosion mitigation, Part I: Effectiveness at reducing hillslope erosion rates, *Catena*, 105, 75–92, 2013.
- Romkens, M. J. M., Helming, K., and Prasad, S. N.: Soil erosion under different rainfall intensities, surface roughness and soil water regimes, *Catena*, 46, 103–123, 2001.
- Ruiz-Sinoga, J. D., Romero-Diaz, A., Ferre-Bueno, E., and Martínez-Murillo, J. F.: The role of soil surface conditions in regulating runoff and erosion processes on a metamorphic hillslope (southern Spain) soil surface conditions, runoff and erosion in southern Spain, *Catena*, 80, 131–139, 2010.
- Ruy, S., Findeling, A., and Chadoeuf, J.: Effect of mulching techniques on plot scale runoff: FDTF modeling and sensitivity analysis, *J. Hydrol.*, 326, 277–294, 2006.
- Smets, T., Poesen, J., and Bochet, E.: Impact of plot length on the effectiveness of different soil-surface covers in reducing runoff and soil loss by water, *Prog. Phys. Geog.*, 32, 654–677, 2008a.
- Smets, T., Poesen, J., and Knapen, A.: Spatial scale effects on the effectiveness of organic mulches in reducing soil erosion by water, *Earth-Sci. Rev.*, 89, 1–12, 2008b.
- SPSS Inc.: Released, PASW Statistics for Windows, Version 18.0., SPSS Inc, Chicago, 2009.
- Zhao, G., Mu, X., Wen, Z., Wang, F., and Gao, P.: Soil erosion, conservation and Eco environment changes in the Loess Plateau of China, *Land Degrad. Dev.*, 24, 499–510, 2013.

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Time to Runoff and coefficient before and after conservation treatment in study scales.

Plot Area (m ²)	Rainfall Intensity (mm h ⁻¹)	Time to Runoff (s)		Runoff Coefficient (%)	
		Control	Treated	Control	Treated
0.25	50	420.00	480.00	24.56	2.03
		609.6	368.4	19.60	2.94
		432.00	372.00	23.86	2.07
	90	69.00	480.00	34.18	1.30
		120.00	564.00	49.56	1.18
		126.00	300.00	37.91	1.39
6	50	38.51	72.52	69.35	60.20
		30.27	68.11	68.45	62.95
		34.34	70.44	69.48	62.48
	90	23.15	56.11	79.42	66.85
		30.32	52.27	78.32	72.18
		26.70	57.28	77.65	60.90

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Changes (%) in time to runoff and coefficient in treated plots with rice straw mulch.

Plot Area (m ²)	Variable	Rainfall intensity (mm h ⁻¹)	
		50	90
0.25	Time to Time	–13.06	+367.92
	Coefficient	–89.34	–96.71
6	Time to Time	+106.15	+110.10
	Coefficient	–10.43	–15.08

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Sediment concentration and soil loss measured at the outlet of the study plots before and after applying conservation treatment.

Plot Area (m ²)	Rainfall Intensity (mm h ⁻¹)	Sediment Concentration (g L ⁻¹)		Soil Loss (g)	
		Control	Treated	Control	Treated
0.25	50	2.04	0.00	1.61	0.00
		1.13	0.00	0.98	0.00
		1.88	0.00	1.54	0.00
	90	2.69	0.00	3.78	0.00
		1.56	0.00	3.42	0.00
		2.00	0.00	3.27	0.00
6	50	6.13	3.87	226.27	131.38
		7.43	3.69	266.64	128.94
		8.27	4.70	302.82	161.62
	90	10.28	4.39	756.69	286.37
		10.71	4.47	787.94	315.10
		10.15	4.01	738.20	239.42

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 4. Reduction rates (%) in average sediment concentration and soil loss in treated plots with rice straw mulch.

Plot Area (m ²)	Variable	Rainfall intensity (mm h ⁻¹)	
		50	90
0.25	Sediment Concentration	–100	–100
	Soil Loss	–100	–100
6	Sediment Concentration	–43.47	–58.69
	Soil Loss	–46.74	–63.24

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Results of GLM test for plot size and conservation treatment effects on the quantitative characteristics of runoff and soil loss.

Source	Dependent Variable	Type III Sum of Squares	df	Mean Square	F	Significant level
Plot	Time to Runoff (s)	595 564.22	1	595 564.22	40.92	0.00
	Runoff Coefficient (%)	16 413.83	1	16 413.83	381.42	0.00
	Sediment Concentration (g L ⁻¹)	185.59	1	185.59	194.67	0.00
	Soil Loss (g)	780 024.69	1	780 024.69	38.46	0.00
Treatment	Time to Runoff (s)	40 142.53	1	40 142.53	2.76	0.11
	Runoff Coefficient (%)	2317.91	1	2317.91	53.86	0.00
	Sediment Concentration (g L ⁻¹)	63.64	1	63.64	66.75	0.00
	Soil Loss (g)	139 578.68	1	139 578.68	6.88	0.02
Plot Treatment	Time to Runoff (s)	14 704.47	1	14 704.47	1.01	0.33
	Runoff Coefficient (%)	616.72	1	616.72	14.33	0.001
	Sediment Concentration (g L ⁻¹)	11.48	1	11.48	12.04	0.002
	Soil Loss (g)	135 178.56	1	135 178.56	6.67	0.02

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.




Figure 1. A general view of treated plots of 6 m^2 (a), the runoff collection system at 6 m^2 plot outlet (b) and 0.25 m^2 (c) with rice straw mulch under the lab condition.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.




Figure 2. Average time to runoff for two study scales and two rainfall intensities.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.




Figure 3. Average runoff coefficient for two study scales and two rainfall intensities.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.




Figure 4. Average sediment concentration for two study scales and two rainfall intensities.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Scale effect on runoff and soil loss control using rice straw mulch under laboratory conditions

S. H. R. Sadeghi et al.




Figure 5. Average soil loss for two study scales and two rainfall intensities.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)