

Interactive
Comment

Interactive comment on “Low titanium magmatism in northwest region of Paraná continental flood basalts (Brazil): volcanological aspects” by F. B. Machado et al.

F. B. Machado et al.

fabio brazmachado@gmail.com

Received and published: 1 November 2014

Thank you for your comments.

There is no study on volcanic aspects on the North part of PCFB. For the first time, strong evidences of the presence of a humid environment at the beginning of the volcanism (and not only desartic) are reported, as well as the interaction features between lavas and sediments. In addition, for the first time in literature, the occurrence of sand filled cracks is described for the whole PCFB. This is very important to know the onset conditions of this huge volcanism in Northern Paraná Basin, which, until now are poorly understood. It is also remarkable that, only in this area, the volcanic activity started with

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



basaltic flows of Ribeira type, according to the chemical classification criteria proposed by Peate et al. (1992), which is scarce in the province.

Previous studies about the lava-sediment interaction were focused on very well preserved expositions from the South area of the province, where there are found several outcrops (e.g. Luchetti et al., 2014). In contrast, the investigated region has a completely different geomorphology, encompassing several geological faults, rare outcrops and significant erosive (tropical climate) processes. It is important to emphasize that, those features make much more difficult to recognize important volcanological characteristics, in comparison to regions where the eruptions occurred recently and/or the rocks are fresh. Therefore, the geochemical signatures of the rocks are essential in order to stablish the accurate stratigraphy of the flows. Although it is not possible to remove completely the geochemistry of the discussion, it will be minimized.

In fact, the distinction among HTi and LTi basaltic rocks using titanium contents as the unique criterion is very understandable for those who are not working in this specific issue. This classification originated in the eighties, mainly based on the contents of major and minor elements (e.g. Bellieni et al., 1983; 1984a, b, 1986a, b; Mantovani et al., 1985; Piccirillo & Melfi, 1988; Piccirillo et al., 1987, 1989; Marques et al., 1989). During this period, a large number of analyzes of incompatible trace elements and isotopic ratios were obtained, allowing refining the characterization of the basalts, which were referred by different names. The data also indicated a geochemical provinciality of the magmatism and Peate et al. (1992) proposed classification criteria to distinguish the different magma types, and gave names to the identified geochemical groups, as it can be seen in the table presented below (Fig. 1). It is possible to verify that there is some overlapping on the titanium contents, on other minor and trace elements, as well as on their ratios. Thus, for a complete characterization, it is necessary to take account the groups the geographical localization of the basalts (Northern: Ribeira, Paranapanema and Pitanga; Southern: Esmeralda, Gramado and Urubici). For the SE amended version of the manuscript, only the proposition of Peate et al. (1992) will be

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



used.

Following your comments and those of referee#2, the discussion about mineral chemistry and temperature of the magmas will be removed from the paper.

For Specific comments: Your corrections, comments and suggestions for the improvement of the paper were very well appreciated. All of them will be rigorously followed.

References

BELLIENI, G., COMIN-CHIARAMONTI, P., MARQUES, L.S., MARTINEZ, L.A., MELFI, A.J., NARDY, A.J.R., PICCIRILO, E.M., STOLVA, D., 1986a. Continental flood Basalts from Central-Western Regions of the Paraná Plateau (Paraguay and Argentina): Petrology and Petrogenetic Aspects. *Neues Jahr. Miner. Abh.*, v. 154, n 2, p. 11–139. BELLIENI, G.; BROTZU, P.; COMIN-CHIARAMONTI, P., ERNESTO, M.; MELFI, A.J.; PACCA, I.G.; PICCIRILO, E.M.; STOLVA, D., 1983. Petrological and Paleomagnetic Data on the Plateau Basalts to Rhyolite sequences of the Southern Paraná Basin (Brazil). *Anais da Academia Brasileira de Ciências*, v. 55, p. 355–383. BELLIENI, G.; COMIN-CHIARAMONTI, P.; MARQUES, L.S., MELFI, A.J., STOLVA, D., 1984b. Low-pressure evolution of basalt sills from bore-holes in the Paraná Basin, Brazil. *TMPM*, v. 33, p. 25-47. BELLIENI, G.; COMIN-CHIARAMONTI, P.; MARQUES, L.S.; MELFI, A.J.; PICCIRILO, E. M.; A.J., NARDY, A.J.R.; ROISENBERG, A. 1984a. High- and low-Ti flood basalts from the Paraná plateau (Brazil): petrology and geochemical aspects bearing on their mantle origin. *Neues Jahr. Miner. Abh.*, v. 150, p. 272–306. BELLIENI, G.; COMIN-CHIARAMONTI, P.; MARQUES, L.S.; MELFI, A.J.; NARDY, A.J.R.; PAPATRECHAS, C.; PICCIRILLO, E. M.; ROISENBERG, A. 1986b. Petrogenetic aspects of acid and basaltic lavas from the Paraná plateau (Brazil): geological, mineralogical and petrochemical relationships. *Journal of Petrology*, v. 27, p.915-944. LUCHETTI, A. C. F. ; NARDY, A. J. R. ; MACHADO, F. B. ; MADEIRA, J. E. O. ; ARNOSIO, J. M., 2014. New insights on the occurrence of peperites and sedimentary deposits within the silicic volcanic sequences of the Paraná Magmatic Province, Brazil.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



Solid Earth, v. 5, p. 121-130. MANTOVANI, M.S.M.; MARQUES, L.S.; SOUZA, M.A.; ATALLA, L.; CIVETA, L.; INONOCENTI, F., 1985. Trace Element and Strontium Isotope Constrains of the Origin and Evolution of Paraná Coantinal Flood Basalts of Santa Catarina State (Southern Brazil). *Journal of Petrology*, v. 26, p. 187-209. MARQUES, L.S.; PICCIRILO, E.M.; MELFI, A.J.; COMIN-CHIARAMONTI, P.; BELLINI, G., 1989. Distribuição de terras raras e outros elementos traços em basaltos da Bacia do Paraná. *Geochimica Brasiliensis*, v. 3, p. 33-50. PEATE, D. W.; HAWKESWORT, C. J.; MANTOVANI, M. S. M., 1992. Chemical stratigraphy of the Paraná lavas (South America): classification of magma types and their spatial distribution. *Bull. Volcanol*, v. 55, p. 119-139. PICCIRILLO, E.M. & MELFI, A.J., 1988. The Mesozoic Flood Volcanism of the Paraná Basin: Petrogenetic and Geophysical Aspects. São Paulo, Brasil: IAG-USP, 600 p. PICCIRILLO, E.M.; CIVETTA, L.; PETRINI, R.; LONGINELLI, A.; BELLINI, G.; COMIN-CHIARAMONTI, P.; MARQUES, L.S.; MELFI, A.J. 1989. Regional variations within the Paraná flood basalts (southern Brazil): evidence for subcontinental mantle heterogeneity and crustal contamination. *Chemical Geology*, v. 75, p. 103-122. PICCIRILLO, E.M.; RAPOSO, M.I.B.; MELFI, A.J.; COMIN-CHIARAMONTI, P.; BELLINI, G.; CORDANI, U.G.; KAWASHITA, K. 1987. Bimodal fissural volcanic suítes from the Paraná Basin (Brazil): K-Ar age, Sr-isotopes and geochemistry. *Geochimica Brasiliensis*, v. 1, p. 53-69.

Interactive comment on *Solid Earth Discuss.*, 6, 2215, 2014.

SED

6, C1233–C1237, 2014

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C1236



Table 2. Classification criteria for basalt magma types

	'High-Ti'			'Low-Ti'		
	Urubici	Pitanga	Paranapanema	Ribeira	Esmeralda	Gramado
SiO ₂	> 49	> 47	48 – 53	49 – 52	48 – 55	49 – 60
TiO ₂	> 3.3	> 2.8	1.7– 3.2	1.5 – 2.3	1.1– 2.3	0.7 – 2.0
P ₂ O ₅	> 0.45	> 0.35	0.2– 0.8	0.15– 0.50	0.1– 0.35	0.05– 0.40
Fe ₂ O ₃ (t)	< 14.5	12.5–18	12.5– 17	12 – 16	12 – 17	9 – 16
Sr	>550	>350	200 –450	200 –375	<250	140 –400
Ba	>500	>200	200 –650	200 –600	90 –400	100 –700
Zr	>250	>200	120 –250	100 –200	65 –210	65 –275
Ti/Zr	> 57	> 60	> 65	> 65	> 60	< 70
Ti/Y	>500	>350	>350	>300	<330	<330
Zr/Y	> 6.5	> 5.5	4.0– 7.0	3.5 – 7.0	2.0– 5.0	3.5 – 6.5
Sr/Y	> 14	> 8	4.5– 15	5 – 17	< 9	< 13
Ba/Y	> 14	> 9	5 – 19	6 – 19	< 12	< 19

Fig 1 - Extracted from Peate et al. (1992).

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper