Manuscript prepared for Solid Earth with version 5.0 of the LATEX class copernicus.cls. Date: 21 May 2014

Pacific Plate slab pull and intraplate deformation in the early Cenozoic

Butterworth, N.P.¹, Müller, R.D.¹, Quevedo, L.¹, O'Connor, J.M.², Hoernle, K.³, and Morra, G.⁴

¹EarthByte Group, School of Geosciences, The University of Sydney, New South Wales, 2006, Australia
 ²GeoZentrum Nordbayern, Erlangen and Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany
 ³GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany
 ⁴Department of Physics and School of Geosciences, University of Louisiana at Lafayette, 70504, LA, USA

Correspondence to: N. P. Butterworth (nathaniel.butterworth@sydney.edu.au)

Abstract. Large tectonic plates are known to be susceptible to internal deformation, leading to a range of phenomena including intraplate volcanism. However, the space and time dependence of intraplate deformation and its relationship with changing plate boundary configurations, subducting slab geometries, and absolute plate motion is poorly understood. We utilise a buoyancy driven Stokes

- 5 flow solver, BEM-Earth, to investigate the contribution of subducting slabs through time on Pacific Plate motion and plate-scale deformation, and how this is linked to intraplate volcanism. We produce a series of geodynamic models from 62 to 42 Ma in which the plates are driven by the attached subducting slabs and mantle drag/suction forces. We compare our modelled intraplate deformation history with those types of intraplate volcanism that lack a clear age progression. Our models
- 10 suggest that changes in Cenozoic subduction zone topology caused intraplate deformation to trigger volcanism along several linear seafloor structures, mostly by reactivation of existing seamount chains, but occasionally creating new volcanic chains on crust weakened by fracture zones and extinct ridges. Around 55 Ma subduction of the Pacific-Izanagi ridge reconfigured the major tectonic forces acting on the plate by replacing ridge push with slab pull along its north-western perimeter,
- 15 causing lithospheric extension along pre-existing weaknesses. Large scale deformation observed in the models coincides with the seamount chains of Hawaii, Louisville, Tokelau, and Gilbert during our modelled time period of 62 to 42 Ma. We suggest that extensional stresses between 72 and 52 Ma are the likely cause of large parts of the formation of the Gilbert chain and that localised extension between 62 and 42 Ma could cause late-stage volcanism along the Musicians Volcanic Ridges. Our
- 20 models demonstrate that early Cenozoic changes in Pacific plate driving forces only cause relatively minor changes in Pacific absolute plate motion directions, and cannot be responsible for the Hawaii-

Emperor Bend (HEB), confirming previous interpretations that the 47 Ma HEB does not primarily reflect an absolute plate motion event.

1 Introduction

- 25 The origin of intraplate volcanism without age progression and far away from plate boundaries is poorly understood (Lee and Grand, 2012; Koppers, 2011). Intraplate volcanism can be viewed as being due to hotspots within tectonic plates, which may be caused by a range of processes including mantle plumes, small-scale convection, or lithospheric extension of plates (Ito and van Keken, 2007). In an effort to categorize these phenomena Courtillot et al. (2003) distinguished three categories of
- 30 hotspots. The first is the classic Wilson-Morgan type mantle plume (Morgan, 1971; Wilson, 1963), a thermal anomaly rising through the mantle due to the density difference between the core-mantle boundary and the surface. These are often long-lived and have a relatively stable source location. The second type is similar, but originates from the bottom of the transition zone, associated with superswells (Koppers et al., 2003; Romanowicz and Gung, 2002), and is comparatively short-lived.
- 35 The third type (Liu and Stegman, 2012; Ito and van Keken, 2007; Hirano et al., 2006; Koppers et al., 2003) is the most broadly classified hotspot, potentially caused by many factors, and the use of the term hotspot to describe this type of volcanism can be a misnomer. The melting anomaly may not be necessarily hot (Bonatti, 1990) and may not be a singular spot (Sandwell and Fialko, 2004). It has been suggested (Ballmer et al., 2013; Conrad et al., 2011) that shear mantle flow within the
- 40 asthenosphere mostly explains this type of intraplate volcanism. However, lithospheric extension driven by plate boundary forces, plate motion, and small-scale covection may be causing intraplate volcanism as well (Ballmer et al., 2009; Koppers et al., 2003; Sandwell et al., 1995). Lithospheric cracking due to plate flexure (Hirano et al., 2006) and thermal contraction (Sandwell and Fialko, 2004) is also a possible contributor to surface volcanics. The cracking hypothesis presumes pre-
- 45 existing partial melt below the surface that may be erupted when stress is applied (Ballmer et al., 2009; Hieronymus and Bercovici, 2000). Intraplate magmatism may occur in conjunction with classic hotspot volcanism, and may be associated with highly strained areas overlapping pre-existing zones of weakness (Davis et al., 2002; Staudigel et al., 1991) or may create new weak zones that give rise to volcanism. Most intraplate volcanism occurs along pre-existing tectonic fabric or around bickly stranged lithearthere (Clauserd and Carthoult, 2008a)
- 50 highly stressed lithosphere (Clouard and Gerbault, 2008a).

Here we investigate how intraplate deformation in the oceanic lithosphere may be caused by subduction driven plate dynamics, how this deformation might contribute to the occurrence and timing of volcanic melting anomalies, and to what extent intraplate volcanism may leave the lithosphere more susceptible to the passage of future melts (Hillier, 2007), focussing on the Pacific Plate evo-

55 lution in the early Cenozoic. This time period captures a major tectonic plate reorganisation seen in several oceanic regions between 53–50 Ma (Cande and Stegman, 2011; Whittaker et al., 2007) during a period of heterogeneous plate tessellation (Morra et al., 2013) whereby the ratio of large to small plates is low. Large plate accelerations have been shown to lead to increased volcanic flux (Hieronymus and Bercovici, 2000; Anderson, 1994); therefore the early Cenozoic is a good candidate

- for a relatively active volcanic period. A good example for such volcanism is the Cenozoic diffuse 60 alkaline magmatic province (DAMP), which formed in the southwest Pacific after 50 Ma (Finn et al., 2005). We analyse changes in plate motion around the Pacific ocean basin by considering slab-pull and mantle drag/suction forces and compare the results with absolute plate reconstructions (Seton et al., 2012; Doubrovine et al., 2012; Chandler et al., 2012; Wessel and Kroenke, 2008) and the occurrence of intraplate volcanics (e.g. Clouard and Bonneville, 2005).
- 65

2 Model setup

We apply a novel workflow utilising a Stokes flow solver, BEM-Earth (Quevedo et al., 2012a; Butterworth et al., 2012; Morra et al., 2012, 2007), to analyse the coupled plate-mantle dynamics in the Late Cretaceous and early Cenozoic. Our model is driven by upper mantle slab-pull buoyancy

70 forces (Faccenna et al., 2012) and by induced slab-suction from the down-going plates as described in Conrad and Lithgow-Bertelloni (2004, 2002). A BEM-Earth simulation requires a set of rheological isosurfaces (here represented by the lithospheric plates, the core, and the external Earth surface). Each isosurface bounds a homogenous region characterised by an effective density and viscosity. In our models these are defined by the surface location of the Pacific, Izanagi, Farallon, and Kula plates and their attached lithosphere and subducting slabs (Fig. 1). 75

The location of the plates and subducting slabs, used as an initial model starting condition, are determined using tectonic reconstructions from Seton et al. (2012) as implemented in the GPlates software (Boyden et al., 2011). We use reconstructed topologically closed plate boundaries through time, along with modelled plate lithospheric thickness to produce a three-dimensional representation

- 80 of the Pacific Plate through time. Oceanic lithosphere thickness is derived from the reconstruction model along with oceanic paleo-age grids with a $1^{\circ} \times 1^{\circ}$ resolution (Müller et al., 2013). Ages are input into a half-space cooling model to determine plate thickness, as used previously in BEM-Earth (Morra et al., 2012, Appendix B).
- We seek to simulate the effect of slab pull and slab suction on the Pacific Plate. For this we assume that the convergence history between a subducting and overriding plate is related to the amount of 85 slab material contributing to the pull force. We utilise 10 million years of subduction history, from plate kinematic reconstructions, to provide an estimate of slab material that drives the slab-pull force in BEM-Earth. A 10 million year interval reflects the approximate time for a slab to subduct to the lower mantle and thus estimates the portion of the slab in the upper-mantle contributing to the slab-
- pull (Conrad and Lithgow-Bertelloni, 2002, 2004; Billen, 2008). To determine initially subducted 90 slab morphology we advect subducting plates into the mantle using surface plate kinematics based on

published plate rotations (Seton et al., 2012) starting 10 Myr before the geodynamic model starting time. The absolute and convergent velocities are determined for each point along the reconstructed subduction zone for each time period (Quevedo et al., 2012b). Absolute motion of the trench defines

- 95 the surface rotation of the slab's position (horizontal component of slab dip) and the convergence rate between the subducting and overriding plate define the upper mantle slab sinking rate (vertical component of slab dip). The initial dip of the slab does not affect the BEM-Earth simulation (Morra et al., 2012), but rather having the correct amount of upper mantle slab material (Quevedo et al., 2012b; Billen, 2008; Schellart, 2004) is important for simulating driving forces. Nevertheless, re-
- 100 sulting dips of slabs from a present day model run are comparable with Slab1.0 (Hayes et al., 2012, Appendix A). At each reconstructed point, we calculate the volume of the slab driving the model, as the convergence rate times the lithospheric thickness added up in 1 Myr time steps for the 10 Myr prior to the model start time. In summary, the slab structure is constructed from the relative motion of the plates in the plate reconstructions for the 10 Myr prior to the model being run. For example,
- 105 the 52 Ma model, uses subduction history between 62–52Ma to generate the slab structure driving this geodynamic model.

The resulting modelled plate consists of a mantle viscosity and density structure that is postprocessed to ensure smooth non-overlapping 3-D surfaces. This is required to maintain a consistent isostatic equilibrium between the model isosurfaces (Butterworth et al., 2012; Morra et al., 2012).

- 110 Rheology of the plate is defined by an isosurface bounding a region of homogenous density and viscosity (as described in Table 1). We approximate the fluid dynamics of subduction by considering the mantle and the lithosphere as regions of homogeneous density and viscosity, disregarding other chemical and rheological inhomogeneities. We assume a simple temperature-independent rheology for such multiphase flow, and model only the fundamental forces controlling the process (Quevedo)
- et al., 2012a), which we take to be: the buoyancy resulting from the different densities between the flow phases; the viscous drag that might hamper or assist plate motion; and the viscous resistance to bending and stretching. The simplified rheology structure is free to deform, and is a simple, yet fair representation for modelling plate-scale lithospheric processes (Capitanio et al., 2010; Li and Ribe, 2012). Each subducting plate is embedded in a homogenous mantle fluid surrounded by an adaptive
- 120 external surface. We use a mantle without viscosity layering to simplify our model, however the role of mantle layering would influence the trench and slab morphology as well as the plateness of a large plate like the Pacific (Morra et al., 2012). Viscosity layering of the mantle would decrease the importance of slab suction relative to slab pull but is probably not sufficient to significantly affect plate motions (Conrad and Lithgow-Bertelloni, 2004). The viscosity contrast between each
- 125 isosurface is fixed for the simulation, thus thermal effects are not considered in the model.

Resolution of the model is determined by the size of triangular elements (panels) constituting each rheological isosurface, which is ~ 50 km. Evolution of the model is driven by the negative buoyancy of the already-subducted lithosphere. As no lithosphere is being replenished at the mid ocean ridges,

Table 1. Reference model parameters.

Parameter	Symbol	Non-dimensional value	Natural value	Units
Earth Radius	$r_{\rm E}$	1	6371	km
Mantle Viscosity	$\eta_{ m m}$	1	10^{21}	Pas
Mantle Density	$ ho_{ m m}$	50	3300	${\rm kgm^{-3}}$
Slab Viscosity	$\eta_{ m s}$	$100 \times \eta_{\rm m}$	10^{23}	Pas
Slab Density	$ ho_{ m s}$	80	3330	${\rm kgm^{-3}}$

we only run the model for a few million years at a time to obtain the intraplate deformation and

- 130 velocities of the plates. The tapered lithospheric thickness at the ridges of the model isosurfaces prompts a ridge-push force to contribute to plate motion, however, the force is diminished by a "surface contact layer" (Butterworth et al., 2012; Morra et al., 2012). The contact layer keeps the plates in isostatic equilibrium by preventing the slab from detaching from the external Earth surface boundary and sinking vertically; rather the subducting plate advances in a more realistic fashion.
- 135 There are several methods for providing this balancing buoyancy force in numerical models (Ribe, 2010; Stegman et al., 2010; Morra et al., 2007). Here we use a "lubrication layer" method, where the Earth surface boundary is described as an adaptive surface, whose dynamic behaviour is controlled partially by the distance between the model isosurfaces. The ridge-push force contributes less than 10 % (Lithgow-Bertelloni and Richards, 1998) to forces driving plate motions, in BEM-Earth models
- 140 we find this force contributes less than 5 % due to the contact layer overwhelming the interaction (Butterworth et al., 2012).

2.1 Plate deformation

We extend the work of Clouard and Gerbault (2008a) into a 3-D spherical domain, where we examine intraplate deformation driven by plate-scale tectonics and its relationship to volcanism. However,

- 145 we use a dynamic simulation with no external velocity forcing. The natural strains, ϵ , are calculated for each model panel through time using $\epsilon = \frac{L}{\ell} - 1$, where L and ℓ are the original and final lengths of model panels respectively. Principal axis stresses are then computed from the natural strains and strain rosette gauge transformation tensors. The von Mises criterion, which is an effective or equivalent stress that can be used as an indication for likely regions of deformation (Gueydan et al.,
- 150 2008), is defined in three dimensions as $\sigma_e = \frac{1}{\sqrt{2}}\sqrt{(\sigma_1 \sigma_2)^2 + (\sigma_2 \sigma_3)^2 + (\sigma_3 \sigma_1)^2}$, where σ is the principal stress in each of the 3 axes (Boresi and Schmidt, 2003). The von Mises yielding criterion is a measure of 'likelihood' for a plate to fracture (possibly expressed through volcanism in the geological record) especially if an area of high von Mises values has pre-existing weakness or is under stress from some other mechanism.
- 155 The plate with a simplified rheology is free to deform due to forces driving the natural evolution



Fig. 1. The initial starting model used for input into BEM-Earth. Each isosurface bounds a region of discrete viscosity and density as described in Table 1. The external Earth surface has been peeled back to show the other core and plate isosurfaces, with the view centred near the North Pole. The modelled plates, here at 42 Ma, are the Kula, Farallon, and Pacific, . The blue mesh is indicative of model resolution and shows the panels that are free to deform.

and transmission of stresses in the plate. Areas of likely deformation are determined over a phase of steady model evolution after a period of initialisation. This delay in measurement allows the model to equilibrate. However, results are found to be similar when the von Mises criteria for deformation is measured early or later in the simulation. Intraplate stresses are much higher in our model than
in reality, since we do not use a highly viscous plate core or model the plastic deformations that play a role in real plates. The constitutive relationship of the material that we use for the plate can be regarded as unrealistic, but models of the semi-rigid rheology of the plates as part of a global model has so far only been applied to present-day plate motions (e.g. Stadler et al., 2010), never to sequences of models for the past. Pre-existing zones of weakness (e.g. fracture zones) likely act as

165 conduits for melting anomalies (Davis et al., 2002), but plate-scale deformation due to subduction processes may provide the stresses required to promote volcanics. Volcanism is not expressed in the models, but we use ${}^{40}\text{Ar}/{}^{39}\text{Ar}$ dates for the relatively few samples available from seafloor volcanism edifices, to see if a link can be established in some places between areas likely to deform due to subduction forcing and spatio-temporal localisation of hotspot melting anomalies.

170 3 Model results

We run four subduction driven models which start with surface reconstructions at 62, 52, 47, and 42 Ma and include the previous 10 million years of subduction material as an initial condition. The resulting model von Mises deformation is correlated with age-dated volcanic structures, and the model kinematics are compared with alternative plate model reconstructions. Slab-pull due to slab

material mechanically attached to subducting plates and the resulting slab-suction induced by these down-going slabs are the only significant model driving forces (Morra et al., 2012; Butterworth et al., 2012). For each model and kinematic reconstruction, we report the mean direction of motion of the Pacific. Although the rigidity (the plateness) of the geodynamic modelled plate is low, we compare plate motions with the rigid kinematic models, as a simple test for how well we are reproducing
absolute plate motions. Our models cannot capture changes over time periods in between model

runs, but rather represent the average motion change between modelled intervals.

3.1 62 Ma reconstruction

The reconstructed Pacific Plate at 62 Ma (Fig. 2) has only one subducting slab mechanically attached to it, along the East Junction subduction zone, located to the north of Australia between the Tethys and Panthalassa (Seton and Müller, 2008). At this time, the Junction slab attached to the Pacific

- and Panthalassa (Seton and Müller, 2008). At this time, the Junction slab attached to the Pacific consists of $\sim 4\%$ of all global upper-mantle slab material (Appendix B). The subducting plates, Izanagi, Kula, and Farallon, that surround the Pacific have subduction zones with over 70% of global slab material driving them. Reconstructed Pacific Plate velocities from Seton et al. (2012) show the plate heading toward the north-west (303°). The Doubrovine et al. (2012) reconstructions have the
- 190 Pacific moving generally toward the north (15°). The Izanagi, Kula, and Farallon plates maintain the dominant subducting slabs around the Pacific. The direction of movement of the modelled Pacific (287°) is more in line with those predicted by the Seton et al. (2012) reconstructions. We observe the areas likely to deform on the plate at 62 Ma to be contrasted across the plate with a large zone of focussed deformation running from the centre of the Pacific to the north-west intersection with the
- 195 Izanagi plate.

3.2 52 Ma reconstruction

Between 62 and 52 Ma the Pacific Plate model undergoes a relatively major change in its kinematics and topology (Fig. 3). The Izanagi plate is now fully subducted and its subducting slab is mechanically attached to the north-west portion of the Pacific Plate. The exact timing and location of Izanagi

ridge subduction is not well resolved (Whittaker et al., 2007; Seton et al., 2012). In our model the

Izanagi ridge subducts roughly parallel to the trench. The western flank of the Izanagi ridge flank is still contributing to the pulling force acting on the Pacific plate at 52 Ma after ridge subduction. Previous modelling (Burkett and Billen, 2009) suggests ridge subduction is not a pre-requisite for a loss of slab-pull.

- Subducting slabs attached to the Pacific Plate now account for $\sim 24\%$ of global material being subducted. The Farallon and Kula plates have down-going material accounting for $\sim 24\%$ and $\sim 12\%$ of global material respectively. This induces high strain just behind the subduction zone in the down-going Pacific Plate. This high-strain region feeds into the same north-west trending feature seen at 62 Ma that tapers off toward the centre of the plate. There are also smaller zones of high-
- 210 strain scattered around the plate. The modelled velocities show the Pacific moving westerly (280°), and rotating clockwise. As subduction is now the major driver of the Pacific Plate, the model velocity vectors are more consistent with the direction of the velocities of Seton et al. (2012) (293°). In the model the velocities are exaggerated close to the subduction zone. The Doubrovine et al. (2012) velocities have increased in magnitude and show the Pacific moving more westerly (337°), similar
- 215 to the Seton et al. (2012) reconstructions during this epoch. Doubrovine et al. (2012) reconstructions favour a more northward trend to our model velocities and the Seton et al. (2012) reconstructions.

3.3 47 Ma reconstruction

220

The Pacific Plate approaches periods of rapid change in the Seton et al. (2012) reconstruction between 52 and 42 Ma. We run a model in the intervening period at 47 Ma to capture this change. At this time (Fig. 4) the major subduction zone attached to the Pacific is only along the west and north-

- west region, now accounting for $\sim 15\%$ of global slab material, topologically similar to the 52 Ma model. The other major dynamic influences come from the Kula and Farallon plates and their attached subducting slabs, which account for $\sim 16\%$ and $\sim 33\%$ of global slab material respectively. The Pacific Plate in our model is now moving in a predominantly westerly direction (283°) and has
- slowed corresponding with a decrease in assumed depth of initial slab material. The large subducting slab attached to the Pacific Plate is the main driver of plate motion in this epoch. The model and reconstructed velocity vectors of Seton et al. (2012) (288°) agree well overall. Doubrovine et al. (2012) plate velocities trend more to the north (341°) and the magnitude has decreased significantly since 52 Ma. Deformation in the plate has similar style to the 52 Ma model, with patches of high-strain appearing over the plate or propagating from the edges; however, the large region of
 - deformation propagating from the north-west subduction zone has been greatly dissipated.

3.4 42 Ma reconstruction

The modelled Pacific at 42 Ma (Fig. 5) is kinematically and topologically similar to the 47 Ma model. The same subduction zones continue to drive the Pacific as from the 47 Ma model epoch. The mean model direction (288°) has the same north-west trend as the Seton et al. (2012) reconstructed



Fig. 2. Pacific Plate reconstruction and model output at 62 Ma. Paleo-plate reconstructed positions (Seton et al., 2012) are outlined in black, with attached subduction zones in red. Velocities of kinematic plate reconstruction models are shown by the black (Seton et al., 2012) and grey arrows (Doubrovine et al., 2012). Our subduction model derived velocity vectors are shown by red arrows. The yellow features are the reconstructed positions of age-dated igneous provinces that have appeared in the 10 Myr preceding the model (see discussion for references). Igneous structures are outlined in green. Significant locations are labelled. Orange lines are the reconstructed fracture zone locations (Matthews et al., 2011). The brown outlines represent the reconstructed positions of the present day coastlines. The aqua to magenta logarithmic colour scale represents the von Mises criterion of our model, with aqua representing minimal potential for plate deformation and magenta representing the maximal potential for deformation. The smooth, homogenous style of deformation at the borders of divergent and passive margins is a post-processing artefact due to the interpolation of the grid of the von Mises criterion values onto the reconstructed edge of the Plate boundary.

plate motion vectors (294°), whereas Doubrovine et al. (2012) reconstructions show the Pacific undergoing an absolute motion change, trending from north at 47 Ma to now north-west (310°). The areas of high-strain are maintained in similar locations to 47 Ma.



Fig. 3. Same as Fig. 2 but for the Pacific Plate reconstruction and model output at 52 Ma.

4 Discussion

240 4.1 Kinematic vs. geodynamic model plate motions

The plates in our geodynamic model are primarily driven by slab material pulling on a given attached plate. Contributions of induced mantle flow, expressed as a suction force, are secondary to this, but can still be appreciable, depending on the location of the slabs relative to the plates (Morra et al., 2012). The slab-suction force is driven from the slabs attached to the down-going plates, but we do

not model the effect of tractions that may be induced by other density heterogeneities in the mantle (Ricard et al., 1993). The role of slab-suction is most evident in the motion of the 62 Ma Pacific,



Fig. 4. Same as Fig. 2 but for the Pacific Plate reconstruction and model output at 47 Ma.

where the plate has no major subduction zones, but continues to move towards the north-west as predicted by kinematic reconstructions (Seton et al., 2012). At times when massive subduction zones bound the Pacific Plate, motion in our models is expected to better predict plate reconstructions,
because the models are primarily driven by the pull of slabs mechanically attached to the down-going plate. Thus we find the direction of motion of the Pacific plate is closer to resembling kinematically derived plate motions of Seton et al. (2012) during these times. The geodynamic model uses non-dimensional rheological parameters; therefore renormalisation of plate velocities to Earth-like values is carried out in a post-processing step. We use the same model time renormalisation as Morra et al.

255 (2010), $\eta/(\Delta \rho \cdot g \cdot r)$, where η is the mantle viscosity, $\Delta \rho$ is the differential density between the



Fig. 5. Same as Fig. 2 but for the Pacific Plate reconstruction and model output at 42 Ma.

external surface and the mantle, g is gravity, and r is the radius of the Earth. We find a mantle-Earth differential density of $\Delta \rho = 60 \text{ kg m}^{-3}$, and a mantle viscosity of $\eta = 1 \times 10^{21} \text{ Pa s}$, as listed in Table 1, best match kinematic reconstruction velocity magnitudes.

260

The correspondence between the model velocities favouring Seton et al. (2012) over Doubrovine et al. (2012) derived velocities, are in-part because the Doubrovine et al. (2012) plate model has been constructed to include Pacific hotspots that mimic the Hawaiian-Emperor bend. However, because this hotspot track is likely uniquely related to mantle flow, perhaps involving ridge-plume capture (Tarduno et al., 2009), we find our subduction driven model better fits Seton et al. (2012) (based on the hotspot moving reference frame of O'Neill et al., 2005) reconstructions. Figure 7 shows the

265 motion paths of the fixed Hawaiian and Louisville hotspots through time relative to the Pacific Plate as reconstructed by different models.

Furthermore, between 83.5 and 45 Ma the Seton et al. (2012) and Doubrovine et al. (2012) plate motions are constrained using different plate circuits. Between 50–70 Ma there is a large transition in the absolute plate motion of the Doubrovine et al. (2012) model attributed to the fast motion of

270 the Indian plate. This could be a point of model velocity mismatch, as our models do not include plates far from the Pacific.

To constrain the reliability of our modelled plate motions we compute best-fit Euler poles from our model velocity fields using the Matlab Euler Pole Calculator (EPC) from Goudarzi et al. (2014) (averaged from the motion of the entire non-rigid Pacific plate). We use the resulting stage poles

- 275 (Table 2) to compute finite reconstruction poles (Table 3) by adding our model stage rotations to the 42 Ma finite rotation for the absolute motion of the Pacific Plate from Seton et al. (2012) as our model does not provide rotations for times after 42 Ma (Fig. 6). We compare the resulting model pole path with those of four published plate reconstructions (Fig. 6). For the time period modelled here (62–42 Ma) Wessel and Kroenke (2008) determine plate motions by assuming fixed hotspots
- in the Pacific. Alternatively Doubrovine et al. (2012) applies a moving Pacific, Atlantic, and Indian hotspot model, with rotations of the Pacific linked through a plate circuit to the absolute reference frame. Euler poles determined from Chandler et al. (2012) and Wessel and Kroenke (2008) are strikingly similar in absolute motion through time as both models rely on a Pacific hotspot reference frame. They deviate from each other after 47 Ma as Chandler et al. (2012) interprets the rapid change
 in Pacific plate motion expressed in the Hawaiian-Emperor bend as being due to a slowdown in drift
- of the Hawaiian plume.

Each of the four published plate reconstruction models implies some change in Pacific plate motion between 62 and 52 Ma. The disparity of the Wessel and Kroenke (2008) Euler poles from the other models (Fig. 6) highlights the impact of including fixed Pacific hotspots in plate reconstruc-

290 tions without considering differential hotspot motion or seamounts offset from hotspot locations. Synthetic hotspot tracks for all models, except for the Wessel and Kroenke (2008) model, which fits the geometry of the Hawaii-Emperor and Louisville chains very closely, are shown on Fig.7. The pole path associated with our dynamically modelled Pacific displays a kink at 52 Ma (Fig. 6)

- however, this kink is expressed only as a very subtle variation in direction of Pacific plate motion

- 295 around the Hawaii-Emperor bend and therefore mismatches the Emperor Chain severely (Fig. 7). In addition the Louisville chain is also not well matched by our model and in both cases the mismatches are similar to those of the Seton et al. (2012) model. Our dynamically subduction-driven motions of the Pacific Plate are agnostic of mantle plumes and plume drift, as they are purely reliant on a given set of plate boundary topologies and reconstructions, as the plate model used determines the amount
- 300 and location of slab material in the upper mantle as initial condition for our geodynamic models. The overall good correspondence between the absolute plate velocities from (Seton et al., 2012) and

our slab driven model provides the insight that a combined relative/absolute plate motion model built without relying on Pacific hotspot tracks, as is the case for Seton et al.'s (2012) model, and particularly not the Hawaiian-Emperor chain, predicts geodynamically plausible absolute Pacific plate

- 305 velocities. This is consistent with the view that the Hawaiian-Emperor bend reflects the slowdown of the drift of the Hawaiian plume (Tarduno, 2007; Tarduno et al., 2009) with a change in absolute (or relative) plate motion playing a minor, ancillary role. This explains why subduction-driven plate models cannot be used to predict tracks for hotspots which have not been stationary during the model time interval. This is clearly the case for the Hawaii hotspot during Emperor chain formation (Tar-
- 310 duno, 2007; Tarduno et al., 2009): For the time interval of interest here (62-42 Ma) in Doubrovine et al.'s (2012) model the Hawaii hotspot moves roughly 380 km to the southeast whereas Louisville is predicted to move about 90 km in a similar direction. Even though these estimates are model-dependent, at least for the Hawaii hotspot there is overwhelming, and converging, observational and model-based evidence that points to a relatively fast southward motion of the Hawaii plume
- 315 during Emperor chain formation. Our model provides additional evidence for this inference, as we demonstrate that even the relatively major changes in subduction zone geometry and slab pull and suction forces experienced by the Pacific Plate do not predict anything close to the change in Pacific plate motion direction that would be required to produce the Hawaii-Emperor bend (Fig. 7). Our conclusions differ from those of Faccenna et al. (2012), who modelled Pacific plate driving forces
- 320 by computing slab pull from a combined plate kinematic and half-space lithospheric cooling model, and obtained larger changes in Pacific plate motion direction than we do.Faccenna et al. (2012) use the age of seafloor along the trench to determine lithospheric thickness, and multiply the thickness by a fixed slab length of 700 km to estimate subducted slab material in the upper mantle. Rather than assuming any particular slab length we use 10 Myr of the convergence history fromSeton et al.
- 325 (2012) combined with associated oceanic paleo-age grids, from which we obtain age and thickness of subducting lithosphere in 1 million year time intervals (see Section 2 and Appendix B). Our methodology and the resulting non-uniform distribution of mass along subductions zones results in a pulling-force significantly different from Faccenna et al.'s (2012) and partially explains the differences in plate motions between their model and ours. Furthermore, their analytical model does not
- 330 account for slab suction, whereas our model does. The mean velocity magnitudes of our modelled Pacific plate (5–10 km/Myr) are in a similar range as those computed byFaccenna et al. (2012); however the directional differences are as large as 30° likely owing to the differences in model setup as discussed above.

4.2 Subduction zone topologies driving plate deformation

335 Location and amount of slab material along subduction zones determines the direction and magnitude of plate motion. Tectonic plates do not move completely rigidly but are free to deform according to the interaction and relative contribution of the slab-pull force, induced suction forces, and basal



Fig. 6. Finite reconstruction pole locations for the absolute motion of the Pacific Plate at each of the model times. The five different absolute reference frames are coloured as in the legend. Stars represent the 42 Ma Euler poles and progressive points are the 47, 52 and 62 Ma poles. Present day continents are overlain in peach for reference. The BEM-Earth reconstruction poles are derived from the addition of the stage rotations for each geodynamic model run to the 42 Ma Pacific absolute motion pole from Seton et al. (2012). Projected error ellipses are determined from each model's published covariance matrices.

Table 2.	Stage rotations	for absolute	Pacific p	plate motion	from the	geodynamic	model.
	0					0 2	

Time Period	Latitude	Longitude	Angle
47–42 Ma	69.4049	-24.2841	-1.8404
52–47 Ma	65.3369	-30.538	-2.0658
62–52 Ma	60.2093	-47.641	-4.9922
72–62 Ma	63.3748	-52.2041	-2.6762

drag forces over the entire plate. In our models magnitudes of the non-dimensional deformation, represented by the von Mises values, are derived from the change in length of the plate panels and

340

345

the interval of time steps that the model is run for. The higher a given von Mises value, the more likely the plate will yield at that location; however, in reality this also dependent on the rheological properties of the lithosphere.

At 62 Ma there is an absence of any major subduction zones driving Pacific Plate motion and deformation. For this modelled time significant deformation is due to nearby slabs (Kula, Izanagi, Farallon) being strongly coupled to the plate through slab suction. The flow cell set-up by the Izanagi

Table 3. Finite poles of rotation for the geodynamic model.

350

Time	Latitude	Longitude	Angle
42 Ma	65.9747	314.4962	31.3938
47 Ma	66.3074	312.0884	33.6642
52 Ma	69.5168	298.9913	37.9722
62 Ma	63.8622	299.4621	46.4814

slab is dominant in controlling Pacific Plate kinematics at this time, because the trench-perpendicular length of the Izangai plate is relatively small (Morra et al., 2012). Induced flow in the model results in minimal surface uplift, so radial stress is not apparent. Instead, deformation is caused by induced flow, dragging sections of the plate with spatially varying tractions. As a result, deformation due to induced upwellings is minimally constrained.

A major tectonic plate reorganisation between 53–50 Ma has been mapped both in the Southeast Indian Ocean and parts of the Pacific Ocean (Whittaker et al., 2007; Cande and Stegman, 2011), who proposed a link between this event and the subduction of the Pacific-Izanagi ridge (Whittaker et al., 2007). This tectonic reconfiguration is captured between the 62 and 52 Ma models, particularly by

- 355 a speedup of the Pacific Plate. Between these times there is a significant increase in deformation across the entire Pacific Plate (Figs. 2 and 3). By 52 Ma the Pacific-Izanagi ridge is fully subducted and the volume of slab material controlling the pull force on the Pacific is at the maximum of all the epochs modelled (Appendix B). The model velocities capture a significant change in absolute plate motion during this time interval. The 52 Ma model reflects a peak amplitude in total lithospheric
- 360 deformation over the Plate compared to the other modelled times. However, we do not see a marked increase in volcanic flux at this time (Hillier, 2007; Clouard and Bonneville, 2005). This is in contrast to an expected increase in volcanism during such a period of rapid plate motion change (Anderson, 1994; Hieronymus and Bercovici, 2000).

Between 52 and 47 Ma the Junction Plate in the western Pacific has fully subducted leaving
a smoother plate boundary between the Pacific and Philippine Plates. In this time period the amount of total global slab material directly pulling the Pacific Plate has reduced from ~ 24 % to ~ 15 %. Changing motions in the Pacific during the Cenozoic have previously been shown to be driven by the variations in slab-pull (Faccenna et al., 2012) and slab-suction (Conrad and Lithgow-Bertelloni, 2004). And, asymmetric distribution of slab material along the subduction zones partially controls

370 the location of intraplate deformation (Clouard and Gerbault, 2008a).

The plate topology, subduction zone, and slab material configurations driving the 47 and 42 Ma models are relatively similar. In turn, the patterns of deformation across the Pacific Plate are similar.

4.3 Plate deformation correlated with magmatic events and evidence for non-plume related intraplate volcanism

- 375 A variety of age-dated volcanic structures formed across the Pacific Plate between 72 and 42 Ma (Fig. 7), which can be compared with the deformation predicted by our models (Fig. 18). The complexity and abundance of seafloor features are highlighted by the use of the vertical gravity gradient in Fig. 7, however seafloor structures that have been sampled and dated to this time period are scarce (Hillier, 2007). Plate deformation does not necessarily generate volcanism, just as volcanism can
- 380 take place without plate deformation, e.g. small-scale convection that is not directly related to deformation. Although plate deformation may induce upwelling and decompression melting, the plate deformation (cracking) can also serve to simply facilitate the rise of melts that already exist or are produced by other mechanisms.

The northernmost area of Pacific intraplate volcanism in Fig. 7 are the Emperor Seamounts (Duncan and Keller, 2004), a product of plume-plate interaction (Sharp and Clague, 2006). However,

- throughout its formation history, subduction driven plate deformation is seen to overlap with the chain. A region of high-strain, between $15-30^{\circ}$ N, encompasses the southern part of the Emperor chain in the 52 Ma model run (Fig. 3). Rather than produce new volcanism, deformation induced during the 52 Ma time period may impose small stress-bends in the pre-existing linear chain (Kop-
- 390 pers and Staudigel, 2005). The 47 Ma model (Fig. 4) continues to show plate-strain, with a diminished magnitude, overlapping with the Hawaiian-Emperor chain. Volcanism is active during the 42 Ma model time period in Hawaii (Sharp and Clague, 2006) and the chain continues to show age-progressive volcanism after the bend at 47 Ma. But the 42 Ma model shows minimal plate deformation correlated with the location of the chain.
- 395 In the north-east of the Pacific Plate lies the Musicians Volcanic Ridges, that have active volcanism prior to our model run epochs. They formed from a hotspot interacting with a spreading ridge between 96 to 75 Ma (Pringle, 1993; Kopp et al., 2003). In the 62 and 52 Ma models in the Musicians seamounts there is increased deformation indicated by the model (Fig. 3). The 47 and 42 Ma models continue to show deformation around the Musicians ridge. The coincidence of von Mises deformation with the extinct ridge may incite late-stage volcanism in the province. 400

The Louisville seamount chain in the South Pacific documents a history of volcanism from 82 to 42 Ma (Koppers et al., 2010, 2011), associated with classical hotspot activity (Koppers et al., 2004). There is plate deformation toward the north of the chain in the 42 Ma model (Fig. 5). However, this deformation is likely not the main driver of magmatism, but would aid in decompressional melting

405 of the hotspot material.

385

The Austral seamounts show volcanism between 62 and 52 Ma (Clouard and Bonneville, 2005). This region of seamounts is influenced by many hotspots (Clouard and Bonneville, 2005) and also shows correlation with highly deformed lithosphere at 52 Ma (Fig. 3). This suggests correlation between hotspots located under lithosphere weakened by previous volcanism (Hillier, 2007).

- 410 In the western Pacific (Fig. 7) there are several clusters of seamounts that together encompass the Western Pacific Seamount Province. This province shows weak age-progression in some areas (Ito and van Keken, 2007), suggesting some formation mechanism other than a plume. Koppers et al. (2003) show that there are in fact some age progressions in this region, but overall is a rather complex area with a spike in volcanism lasting until \sim 70 Ma. This province includes the Japanese
- 415 Seamounts (Ozima et al., 1983) in the north, the Mid-Pacific Mountains (Pringle, 1993) that show weak age progression in the central region, and the Magellan seamounts to the south. Between 82 and 62 Ma there are only four Japanese seamounts displaying volcanism, with no mapped features correlating with our modelled plate-strain.

The Line Islands in the central Pacific are considered to have formed through volcanism due to 420 lithospheric extension (Davis et al., 2002). Their temporal appearance between 80 and 68 Ma is not correlated with subduction driven deformation observed in the 62 Ma model. The Line Islands show reduced volcanism after this time until 55 Ma without any deforming regions coinciding with their formation. The lack of coincidence between subduction driven deformation and Line Islands volcanism, suggest the magmatism is influenced by another process. Lithospheric extension in this 425 region is possibly related to the upwarping of the Superswell in the eastern South Pacific (Davis et al., 2002).

The initial formation of the Gilbert Ridge can be extended back in time along the Marshall Islands to around 100 Ma (Konter et al., 2008; Koppers et al., 2003). Basement near the ridge was likely preconditioned to volcanism (Koppers et al., 2007) because of the emplacement of volcanic

- 430 sills during the formation of the Ontong-Java and Hikurangi Plateaus around 125 Ma. The location of the Gilbert chain follows a likely zone of weakness extending north from the Manahiki-Chizca ridge, and running parallel to existing fracture zones (Fig. 2). The Gilbert ridge has been shown to have poor age progression and also shows signs of stress bends at times after formation (Koppers et al., 2007). Volcanism is expressed from 72 to 62 Ma at the Gilbert ridge and later at the Tuvalu-
- 435 Ellice and Samoan seamount chain (Koppers and Staudigel, 2005). This chain trends and extends in the same north-west direction as the zone of high-strain banding in the central Pacific seen in Fig. 2. It has been shown that seamount chains are generally aligned with the direction of the most tensile principal tectonic stress (Hieronymus and Bercovici, 2000). The modelled Pacific Plate is experiencing tensional deformation along a zone aligned with the orientation of this feature (Fig. 2).
- 440 The timing of the modelled lithospheric deformation correlates with the formation age of 67 Ma (Koppers and Staudigel, 2005), with continued stress likely influencing a long slow stress-bend after the Gilbert chain's formation. Deformation of the Pacific lithosphere overprinting pre-existing weaknesses seems to have been significant enough to activate the Gilbert chain around this time. Melt material may have already existed in the upper-mantle from the events around 100 Ma, only
- 445 requiring the changing tectonic stresses, incited by the subducting Izanagi slab, to initiate surface volcanism.

The Tokelau seamounts and Phoenix Islands are formed between 72 and 62 Ma in the eastern Pacific (Koppers and Staudigel, 2005), away from any significant tectonic deformation. The Tokelau seamounts are volcanically active between the 62 and 47 Ma models, correlating with a well-defined

450

region of high-strain (Figs. 3–5). Formation of these structures are likely influenced by lithospheric extension (Koppers and Staudigel, 2005) on crust weakened by nearby fracture zones.

The Tarava seamounts become active for the 42 Ma model and are thought to have formed from a hotspot influenced by lithospheric stress (Clouard et al., 2003). This is consistent with the regional deformed area overlapping the location of the reconstructed sample location (Fig. 5). An

455 unknown source mechanism has produced volcanic activity along the poorly sampled and dated nearby Tuamotus seamounts, around 40–50 Ma, visible in the the 47 Ma and 42 Ma models (Figs. 4 and 5). There is minimal correlation between modelled plate deformation and the formation of these seamounts.

It has previously been shown that present day subduction zone forces on the Pacific Plate lead to internal deformation (Clouard and Gerbault, 2008a,b). Extensional mechanisms and lithospheric thickness variations can contribute to enhanced volcanism on ridges and hotspots, but neither mechanism is likely the sole source of seamount chains (Pilger, 2008). Lithospheric weaknesses (e.g. fracture zones, pre-existing magmatism) that override a source of melt material, possibly derived from mantle plumes too weak to penetrate the surface, may be perturbed by tectonic stresses due to plate

- 465 motion changes, inturn exciting surface eruption. Age-dated Late Cretaceous and early Cenozoic seafloor structures across the Pacific show signs of Pacific-wide plate deformation. Intraplate volcanism sampled across the Pacific is partially indicative of a proposed global reorganisation (Whittaker et al., 2007; Cande and Stegman, 2011) between about 62 and 47 Ma and may be considered as a proxy for stress on the plate (Clouard and Gerbault, 2008a). Because the stress state of the
- 470 lithosphere, plate deformation, and subsequent volcanism are inherently mixed, but are not necessarily mutually exclusive, it is difficult to extract a definitive location of volcanism based on stress or deformation alone, or vice-versa. This is even more so where there is pre-existing hotspot volcanism in reactivated seamount chains. However, regions likely to deform predicted by our models gives an indication of potential sites of intraplate volcanism that are related to anomalously stressed
- 475 lithosphere.

4.4 Modelled lithospheric structure influencing plate deformation

There is competition between thermal contraction that strengthens the lithosphere during cooling as the lithosphere gets older and thicker (Koppers and Watts, 2010) and load-induced stress relaxation that weakens it. However, it is found that plate thickness seems to have no noticeable effect on plate

480 deformation, driven by large-scale convection in our models, regardless of the tectonic configuration. This is likely because the plate rheology is homogenous and there is no heat flow in the model that would otherwise create a weakened crust (Shaw and Lin, 1996). Also the thickness variations are



Fig. 7. Present day vertical gravity gradient of the Pacific Plate (Sandwell and Smith, 2009). Present day locations of volcanics dated between 72 and 42 Ma are binned in 10 Myr increments and coloured as in the legend. The computed motion paths for fixed Hawaiian and Lousiville hotspots are shown back to 62 Ma, for each of the different models, as listed in the legend. Dots along the motion path show the trail's position at 42, 47, 52, and 62 Ma respectively.

comparatively small to the major convective cells in the model domain.

485

The rheology of the modelled subducting plate and the mantle will influence the deformation of the plate. In the models, decreasing the lithosphere viscosity to $50 \times \eta_m$ or increasing it to $200 \times \eta_m$ will make the model evolve slower or faster, respectively. However, regions of deformation remain broadly consistent between models with a different viscosity lithosphere. Intraplate deformation resulting from the stresses imposed by the subduction and mantle drag/suction forces are a robust prediction of the model. Whether the stresses are sufficient to cause intraplate deformation depends

490 on the actual rheological parameters of the Pacific Plate. A subsequent increase in deformation may result in volcanism only if melt material is available, and the lithosphere has pre-existing lines of weakness and/or is weak enough to fracture (Ballmer et al., 2009; Hieronymus and Bercovici, 2000).

5 Conclusions

- During the Late Cretaceous and early Cenozoic the Pacific Plate underwent a major tectonic shift in its primary driving forces; the plate changed from being entirely surrounded by mid-ocean ridges to a topology characerised by a progressive increase in subduction zones particularly along its western and northern perimeter. In our models prior to 52 Ma, the Pacific was controlled by the Izanagi, Kula, and Farallon subducting plates surrounding it, and to a lesser extent by a small subducting slab attached at the East Junction subduction zone. From 52 Ma onwards, following the subduction of the
- 500 Pacific-Izanagi ridge, the Pacific Plate was primarily controlled by slab-pull in the north-west. The absolute motions of the Pacific derived from subduction driven forces correspond well with other modelled plate reconstructions (Seton et al., 2012; Chandler et al., 2012), when model assumptions and simplifications are taken into account. Slab pull and suction combined control the deformation in the attached subducting plate. We find the regions of highest deformation occur directly adjacent to
- 505 the most voluminous subducting slabs. Several areas of deformation across the Pacific can be linked to age-dated intraplate volcanism. The seamount chains of Hawaii-Emperor, Louisville, and Tokelau are subject to lithospheric deformation occurring during the early Cenozoic. Plate-scale extensional stresses between our modelled time intervals correlate with a large section of the location and timing of formation of the Gilbert chain, suggesting an origin largely due to lithospheric extension at this
- 510 time. The Musicians Volcanic Ridges, which likely formed by traditional plume mechanisms, spatially correlate with modelled lithospheric stress and is a likely candidate for late-stage volcanism between 52 and 42 Ma. Our simplified 3-D subduction simulations suggest stress-induced deformation in the Pacific during the Late Cretaceous and early Cenozoic is partially controlled by plate-scale kinematics. Our dynamic models, combined with kinematically reconstructed absolute plates mo-
- 515 tions, confirm the view that the Hawaiian-Emperor bend is largely not due to a change in absolute plate motion, but that it mainly reflects the slowdown of the Hawaiian plume drift.

Acknowledgements. Thanks to C. Heine and S. Williams for fruitful discussions, and to A. A. P. Koppers and two anonymous referees for reviewing the manuscript.

Appendix A

520 Comparison of initial model condition with Slab1.0

We use 10 Myr of subduction history to build the slabs attached to the geodynamic model subducting plates. The kinematics of the plate model (Seton et al., 2012) determines the total volume of material, dip, and depth of the slab. We find these dips to be comparable to the available Slab1.0 (Hayes et al., 2012) present day slabs (Figs. 8–13).

525 Appendix B

Volume of subducted material

thickness of the lithosphere, z, is determined as

Plate convergent velocities are determined for each point along the reconstructed subduction zone for each time period. Oceanic lithosphere thickness is derived from the paleo-reconstruction model (Seton et al., 2012) along with sampling age grids with a $1^{\circ} \times 1^{\circ}$ resolution (Müller et al., 2013). Using a Half-Space Cooling model truncated at 95 km (after Chapter 4.2 Schubert et al., 2001) the

$$z = \operatorname{erf}^{-1}\left(\frac{T_{\rm l} - T_{\rm o}}{T_{\rm m} - T_{\rm o}}\right) 2\sqrt{\kappa} \sqrt{\operatorname{age}},\tag{B1}$$

where, erf⁻¹ is the inverse of the error function, T_l = 1300 °C and is the isotherm of the lithosphere, T_o = 0 °C +and is the surface temperature, T_m = 1600 °C and is the temperature of the mantle,
535 κ = 8 × 10⁻⁸ m s⁻¹ and is the thermal diffusivity constant, and age is the age of the lithosphere sampled from the age grids. We calculate the volume of the slab as the convergence rate times the lithospheric thickness times each subduction segment length (from the resolution of the plate model).

Figures 14–17 show the amount of globally subducted material for each of the model time periods.

Appendix C

540 Von Mises deformation through time

For each point of volcanism from Figure 7 we determine the upper 95 percentile of von Mises deformation occurring within a 1 degree radius around the point at the time of volcanism (Fig. 18). The reconstructed position of the volcanic structures and the diffuse location of the von Mises contribute to the spatial error between the datasets.



Fig. A 8. Top-down view of the Nazca Plate at present day for geodynamic model input. The green to purple colored topology represents the depth of the Nazca slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model. The Slab1.0 interpretation is made slightly transparent to see the extent of modelled plate.



Fig. A 9. North side view of the Nazca Plate at present day for geodynamic model input. The green to purple colored topology represents the depth of the Nazca slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model. The Slab1.0 interpretation is made slightly transparent to see the extent of modelled plate.



Fig. A 10. South side view of the Nazca Plate at present day for geodynamic model input. The green to purple colored topology represents the depth of the Nazca slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model. The Slab1.0 interpretation is made slightly transparent to see the extent of modelled plate.



Fig. A 11. Top-down view of the Pacific Plate at present day for geodynamic model input, localized over the Tonga-Kermadec subduction zone. The green to purple colored topology represents the depth of the Kermadec slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model. The modelled plate is made slightly transparent to see the extent of the Slab1.0 interpretation.



Fig. A 12. Side view of the Pacific Plate at present day for geodynamic model input, looking from the south at the Tonga-Kermadec subduction zone. The green to purple colored topology represents the depth of the Kermadec slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model. The modelled plate is made slightly transparent to see the extent of the Slab1.0 interpretation.



Fig. A 13. View of the Pacific Plate at present day for geodynamic model input, looking from beneath the plate at the Tonga-Kermadec subduction zone. The green to purple colored topology represents the depth of the Kermadec slab from the Slab1.0 (Hayes et al., 2012) interpretation. The black mesh over the plate indicates the resolution of the model.



Fig. B 14. Integrated volume of subducted material between 52 and 42 Ma. The color scale represents the volume of material. The total amount of material for this time period is 3.7×10^7 km³.



Fig. B 15. Integrated volume of subducted material between 57 and 47 Ma. The color scale represents the volume of material. The total amount of material for this time period is 3.9×10^7 km³.



Fig. B 16. Integrated volume of subducted material between 62 and 52 Ma. The color scale represents the volume of material. The total amount of material for this time period is 3.9×10^7 km³.



Fig. B 17. Integrated volume of subducted material between 72 and 62 Ma. The color scale represents the volume of material. The total amount of material for this time period is 4.6×10^7 km³.



Fig. C 18. Von Mises criteria for yielding at the time of volcanism for the structures discussed in the text. The von Mises shown here are the upper 95 percentile of a 1 degree radius around each point of volcanism shown in figures 2 to 5. This is overlain on the vertical gravity gradient highlighting Pacific seafloor structures for reference to Figure 7.

545 References

Anderson, D.: Superplumes or supercontinents?, Geology, 22, 39–42, doi:10.1130/0091-7613(1994)022<0039:SOS>2.3.CO;2, 1994.

Ballmer, M., van Hunen, J., Ito, G., Bianco, T., and Tackley, P.: Intraplate volcanism with complex agedistance patterns: a case for small-scale sublithospheric convection, Geochem. Geophy. Geosy., 10, Q06015,

550 doi:10.1029/2009GC002386, 2009.

Ballmer, M. D., Conrad, C. P., Smith, E. I., and Harmon, N.: Non-hotspot volcano chains produced by migration of shear-driven upwelling toward the East Pacific Rise, Geology, 41, 479–482, doi:10.1130/G33804.1, 2013.

Billen, M.: Modeling the dynamics of subducting slabs, Annu. Rev. Earth Pl. Sc., 36, 325–56, doi:10.1146/annurev.earth.36.031207.124129, 2008.

- 555 Bonatti, E.: Not so hot "hot spots" in the oceanic mantle, Science, 250, 107–111, doi:10.1126/science.250.4977.107, 1990.
 - Boresi, A. P. and Schmidt, R. J.: Advanced Mechanics of Materials, John Wiley and Sons, 2003.
 - Boyden, J., Müller, R., Gurnis, M., Torsvik, T., Clark, J., Turner, M., Ivey-Law, H., Watson, R., and Cannon, J.: Next-generation plate-tectonic reconstructions using GPlates, in: Geoinformatics: Cyberinfrastructure for
- 560 the Solid Earth Sciences, edited by: Keller, G. R. and Baru, C., Chap. 7, Cambridge University Press, 95– 116, doi:10.1017/CBO9780511976308.008, 2011.
 - Burkett, Erin R. and Billen, Magali I.: Dynamics and implications of slab detachment due to ridge-trench collision, Journal of Geophysical Research: Solid Earth, 114, B12402, doi:10.1029/2009JB006402, 2009.
 Butterworth, N., Quevedo, L., Morra, G., and Müller, R.: Influence of overriding plate geometry and rheology

on subduction, Geochem. Geophy. Geosy., 13, Q06W15, doi:10.1029/2011GC003968, 2012.

- Cande, S. C. and Stegman, D. R.: Indian and African plate motions driven by the push force of the Réunion plume head, Nature, 475, 47–52, doi:10.1038/nature10174, 2011.
 - Capitanio, F., Stegman, D., Moresi, L., and Sharples, W.: Upper plate controls on deep subduction, trench migrations and deformations at convergent margins, Tectonophysics, 483, 80–92,
- 570 doi:10.1016/j.tecto.2009.08.020, 2010.
 - Chandler, M. T., Wessel, P., Taylor, B., Seton, M., Kim, S.-S., and Hyeong, K.: Reconstructing Ontong Java Nui: implications for Pacific absolute plate motion, hotspot drift and true polar wander, Earth Planet. Sc. Lett., 331–332, 140–151, doi:10.1016/j.epsl.2012.03.017, 2012.

Clouard, V. and Bonneville, A.: Ages of seamounts, islands, and plateaus on the Pacific plate, Geol. Soc. Am.

575 S., 388, 71–90, doi:10.1130/0-8137-2388-4.71, 2005.

Clouard, V. and Gerbault, M.: Break-up spots: Could the Pacific open as a consequence of plate kinematics?, Earth Planet. Sc. Lett., 265, 195–208, doi:10.1016/j.epsl.2007.10.013, 2008a.

Clouard, V. and Gerbault, M.: Reply to "Break-up spots: Could the Pacific open as a consequence of plate kinematics?" Comment by R. Pilger, Earth Planet. Sc. Lett., 275, 196–199, doi:10.1016/j.epsl.2008.08.008, 2008b.

- 580
 - Clouard, V., Bonneville, A., and Gillot, P.-Y.: The Tarava Seamounts: a newly characterized hotspot chain on the South Pacific Superswell, Earth Planet. Sc. Lett., 207, 117–130, doi:10.1016/S0012-821X(02)01143-3, 2003.
 - Conrad, C. and Lithgow-Bertelloni, C.: How mantle slabs drive plate tectonics, Science, 298, 207,

585 doi:10.1126/science.1074161, 2002.

- Conrad, C. and Lithgow-Bertelloni, C.: The temporal evolution of plate driving forces: importance of "slab suction" versus "slab pull" during the Cenozoic, J. Geophys. Res., 109, B10407, doi:10.1029/2004JB002991, 2004.
- Conrad, C. P., Bianco, T. A., Smith, E. I., and Wessel, P.: Patterns of intraplate volcanism controlled by asthenospheric shear, Nat. Geosci., 4, 317–321, doi:10.1038/ngeo1111, 2011.
- Courtillot, V., Davaille, A., Besse, J., and Stock, J.: Three distinct types of hotspots in the Earth's mantle, Earth Planet. Sc. Lett., 205, 295–308, doi:10.1016/S0012-821X(02)01048-8, 2003.
- Davis, A. S., Gray, L. B., Clague, D. A., and Hein, J. R.: The Line Islands revisited: new ⁴⁰Ar/³⁹Ar geochronologic evidence for episodes of volcanism due to lithospheric extension, Geochem. Geophy. Geosy., 3, 1018,

590

600

Doubrovine, P., Steinberger, B., and Torsvik, T.: Absolute plate motions in a reference frame defined by moving hot spots in the Pacific, Atlantic, and Indian oceans, J. Geophys. Res., 117, B09101, doi:10.1029/2011JB009072, 2012.

Duncan, R. A. and Keller, R. A.: Radiometric ages for basement rocks from the Emperor Seamounts, ODP Leg 197, Geochem. Geophy. Geosy., 5, Q08L03, doi:10.1029/2004GC000704, 2004.

- Faccenna, C., Becker, T. W., Lallemand, S., and Steinberger, B.: On the role of slab pull in the Cenozoic motion of the Pacific plate, Geophys. Res. Lett., 39, L03305, doi:10.1029/2011GL050155, 2012.
 - Finn, C. A., Müller, R. D. and Panter, K. S.: A Cenozoic diffuse alkaline magmatic province (DAMP) in the southwest Pacific without rift or plume origin, Geochem. Geophy. Geosy., 6, Q02005,
- doi:10.1029/2004GC000723, 2005.
 - Goudarzi, M. A., Cocard, M. and Santerre, R.: EPC: Matlab software to estimate Euler pole parameters, GPS Solutions, 18, 153—162, doi:10.1007/s10291-013-0354-4, 2014.

Gueydan, F., Morency, C., and Brun, J.-P.: Continental rifting as a function of lithosphere mantle strength, Tectonophysics, 460, 83–93, doi:10.1016/j.tecto.2008.08.012, 2008.

- 610 Hayes, G. P., Wald, D. J., and Johnson, R. L.: Slab1.0: a three-dimensional model of global subduction zone geometries, J. Geophys. Res.-Sol. Ea., 117, B01302, doi:10.1029/2011JB008524, 2012.
 - Hieronymus, C. and Bercovici, D.: Non-hotspot formation of volcanic chains: control of tectonic and flexural stresses on magma transport, Earth Planet. Sc. Lett., 181, 539–554, doi:10.1016/S0012-821X(00)00227-2, 2000.
- 615 Hillier, J.: Pacific seamount volcanism in space and time, Geophys. J. Int., 168, 877–889, doi:10.1111/j.1365-246X.2006.03250.x, 2007.

Hirano, N., Takahashi, E., Yamamoto, J., Abe, N., Ingle, S. P., Kaneoka, I., Hirata, T., Kimura, J.-I., Ishii, T., Ogawa, Y., Machida, S., and Suyehiro, K.: Volcanism in response to plate flexure, Science, 213, 1426–1428, doi:10.1126/science.1128235, 2006.

- 620 Ito, G. and van Keken, P. E.: Hotspots and melting anomalies, in: Treatise on Geophysics, edited by: Bercovici, D., vol. 7, Mantle Dynamics, Elsevier, 371–435, 2007.
 - Konter, J. G., Hanan, B. B., Blichert-Toft, J., Koppers, A. A. P., Plank, T., and Staudigel, H.: One hundred million years of mantle geochemical history suggest the retiring of mantle plumes is premature, Earth Planet. Sc. Lett., 275, 285–295, doi:10.1016/j.epsl.2008.08.023, 2008.

⁵⁹⁵ doi:10.1029/2001GC000190, 2002.

625 Kopp, H., Kopp, C., Morgan, J. P., Flueh, E. R., Weinrebe, W., and Morgan, W.: Fossil hot spot-ridge interaction in the Musicians Seamount Province: geophysical investigations of hot spot volcanism at volcanic elongated ridges, J. Geophys. Res., 108, 2160, doi:10.1029/2002JB002015, 2003.

Koppers, A. A. P.: Mantle plumes persevere, Nat. Geosci., 4, 816–817, doi:10.1038/ngeo1334, 2011.

- Koppers, A. A. P. and Staudigel, H.: Asynchronous bends in Pacific seamount trails: a case for extensional
 volcanism?, Science, 307, 904–907, doi:10.1126/science.1107260, 2005.
 - Koppers, A. A. P. and Watts, A. B.: Intraplate seamounts as a window into deep Earth processes, Oceanography, 23, 42–57, doi:10.5670/oceanog.2010.61, 2010.

Koppers, A. A. P., Staudigel, H., Pringle, M., and Wijbrans, J.: Short-lived and discontinuous intraplate volcanism in the South Pacific: hot spots or extensional volcanism?, Geochem. Geophy. Geosy., 4, 1089, doi:10.1029/2003GC000533, 2003.

Koppers, A. A. P., Duncan, R., and Steinberger, B.: Implications of a nonlinear ⁴⁰Ar/³⁹Ar age progression along the Louisville seamount trail for models of fixed and moving hot spots, Geochem. Geophy. Geosy., 5, Q06L02, doi:10.1029/2003GC000671, 2004.

635

Koppers, A. A. P., Staudigel, H., Morgan, J. P., and Duncan, R.: Nonlinear ⁴⁰Ar/³⁹Ar age systematics along the

- 640 Gilbert Ridge and Tokelau Seamount Trail and the timing of the Hawaii-Emperor Bend, Geochem. Geophy. Geosy., 8, Q06L13, doi:10.1029/2006GC001489, 2007.
 - Koppers, A. A. P., Yamazaki, T., and Geldmacher, J.: Louisville Seamount Trail: implications for geodynamic mantle flow models and the geochemical evolution of primary hotspots, IODP Scientific Prospectus, 330, doi:10.2204/iodp.sp.330.2010, 2010.
- 645 Koppers, A. A. P., Gowen, M. D., Colwell, L. E., Gee, J. S., Lonsdale, P. F., Mahoney, J. J., and Duncan, R.: New ⁴⁰Ar/³⁹Ar age progression for the Louisville hot spot trail and implications for inter–hot spot motion, Geochem. Geophy. Geosy., 12, Q0AM02, doi:10.1029/2011GC003804, 2011.

Lee, C.-T. A. and Grand, S. P.: Intraplate volcanism, Nature, 482, 314–315, doi:10.1038/482314a, 2012.

- Li, Z.-H. and Ribe, N. M.: Dynamics of free subduction from 3-D boundary element modeling, J. Geophys.
 Res., 117, B06408, doi:10.1029/2012JB009165, 2012.
 - Lithgow-Bertelloni, C. and Richards, M.: The dynamics of Cenozoic and Mesozoic plate motions, Rev. Geophys., 36, 27–78, doi:10.1029/97RG02282, 1998.
 - Liu, L. and Stegman, D. R.: Origin of Columbia River flood basalt controlled by propagating rupture of the Farallon slab, Nature, 482, 386–390, doi:10.1038/nature10749, 2012.
- 655 Matthews, K., Müller, R., Wessel, P., and Whittaker, J.: The tectonic fabric of the ocean basins, J. Geophys. Res., 116, B12109, doi:10.1029/2011JB008413, 2011.
 - Morgan, W.: Convection plumes in the lower mantle, Nature, 230, doi:10.1038/230042a0, 1971.
 - Morra, G., Chatelain, P., Tackley, P., and Koumoutsakos, P.: Large scale three-dimensional boundary element simulation of subduction, in: Computational Science ICCS 2007, edited by: Shi, Y., Albada, G., Don-
- 660 garra, J., and Sloot, P., Lecture Notes in Computer Science, Vol. 4489, 1122–1129, Springer, Berlin, Heidelberg, 1122–1129, doi:10.1007/978-3-540-72588-6_178, 2007.
 - Morra, G., Yuen, D.A., Boschi, L., Chatelain, P., Koumoutsakos, P., Tackley, P.J.:The fate of the slabs interacting with a density/viscosity hill in the mid-mantle, Physics of the Earth and Planetary Interiors, 180, 271–282, doi:10.1016/j.pepi.2010.04.001, 2010.

- 665 Morra, G., Quevedo, L., and Muller, R. D.: Spherical dynamic models of top-down tectonics, Geochem. Geophy. Geosy., 13, 1–66, doi:10.1029/2011GC003843, 2012.
 - Morra, G., Seton, M., Quevedo, L., and Müller, R. D.: Organization of the tectonic plates in the last 200 Myr, Earth Planet. Sc. Lett., 373, 93–101, doi:10.1016/j.epsl.2013.04.020, 2013.
- Müller, R., Dutkiewicz, A., Seton, M., and Gaina, C.: Seawater chemistry driven by supercontinent assembly,
 breakup, and dispersal, Geology, 41, 907–910, doi:10.1130/G34405.1, 2013.
 - O'Neill, C., Müller, D., and Steinberger, B.: On the uncertainties in hot spot reconstructions and the significance of moving hot spot reference frames, Geochem. Geophy. Geosy., 6, Q04003, doi:10.1029/2004GC000784, 2005.
 - Ozima, M., Kaneoka, I., Saito, K., Honda, M., Yanagisawa, M., and Takigami, Y.: Summary of geochrono-
- 675 logical studies of submarine rocks from the western Pacific Ocean, in: Geodynamics of the Western Pacific-Indonesian Region, edited by: Hilde, T. W. C., and Uyeda, S., AGU, vol. 11, 137–142 doi:10.1029/GD011p0137, 1983.
 - Pilger, R.: Discussion of "Break-up spots: could the Pacific open as a consequence of plate kinematics?" by Clouard and Gerbault, Earth Planet. Sc. Lett., 275, 193–195, doi:10.1016/j.epsl.2008.08.005, 2008.
- 680 Pringle, M. S.: Age progressive volcanism in the Musicians seamounts: a test of the hot spot hypothesis for the late Cretaceous Pacific, in: The Mesozoic Pacific: Geology, Tectonics, and Volcanism: A Volume in Memory of Sy Schlanger, Geophys. Monogr. Ser, AGU, Washington, D.C., vol. 77, 187–216, doi:10.1029/GM077p0187, 1993.
 - Quevedo, L., Hansra, B., Morra, G., Butterworth, N., and Müller, R. D.: Oblique mid ocean ridge subduc-
- 685 tion modelling with the parallel fast multipole boundary element method, Comput. Mech., 51, 455–463, doi:10.1007/s00466-012-0751-5, 2012a.
 - Quevedo, L., Morra, G., and Müller, R. D.: Global paleo-lithospheric models for geodynamical analysis of plate reconstructions, Phys. Earth Planet. In., 212–213, 106–113, doi:10.1016/j.pepi.2012.09.007, 2012b.
- Ribe, N. M.: Bending mechanics and mode selection in free subduction: a thin-sheet analysis, Geophys. J. Int.,
 180, 559–576, doi:10.1111/j.1365-246X.2009.04460.x, 2010.
 - Ricard, Y., Richards, M., Lithgow-Bertelloni, C., and Le Stunff, Y.: A geodynamic model of mantle density heterogeneity, J. Geophys. Res.-Sol. Ea., 98, 21895–21909, doi:10.1029/93JB02216, 1993.
 - Romanowicz, B. and Gung, Y.: Superplumes from the core-mantle boundary to the lithosphere: implications for heat flux, Science, 296, 513–516, doi:10.1126/science.1069404, 2002.
- 695 Sandwell, D. and Fialko, Y.: Warping and cracking of the Pacific plate Warping and cracking of the Pacific plate by thermal contraction, J. Geophys. Res., 109, B10411, doi:10.1029/2004JB003091, 2004.
 - Sandwell, D. and Smith, W. H. F.: Global marine gravity from retracked Geosat and ERS-1 altimetry: ridge segmentation versus spreading rate, J. Geophys. Res., 114, B01411, doi:10.1029/2008JB006008, 2009.
 - Sandwell, D., Winterer, E., Mammerickx, J., Duncan, R., Lynch, M., Levitt, D., and Johnson, C.: Evidence for
- diffuse exension of the Pacific plate from Pukapuka ridges and cross-grain gravity lineations, J. Geophys.
 Res., 100, 15087–15099, doi:10.1029/95JB00156, 1995.
 - Schellart, W.: Quantifying the net slab pull force as a driving mechanism for plate tectonics, Geophys. Res. Lett., 31, L07611, doi:10.1029/2004GL019528, 2004.
 - Schubert, G., Turcotte, D. L., and Olson, P.: Mantle Convection in the Earth and Planets, Cambridge University

705 Press, 2001.

- Seton, M. and Müller, R.: Reconstructing the junction between Panthalassa and Tethys since the Early Cretaceous, Eastern Australian Basins III. Petroleum Exploration Society of Australia. Special Publication, 263– 266, 2008.
- Seton, M., Müller, R., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., Talsma, A., Gurnis, M., Turner, M.,
- Maus, S., and Chandler, M.: Global continental and ocean basin reconstructions since 200 Ma, Earth-Sci. Rev., 113, 212–270, doi:10.1016/j.earscirev.2012.03.002, 2012.
 - Sharp, W. and Clague, D.: 50-Ma initiation of Hawaiian-Emperor bend records major change in Pacific plate motion, Science, 313, 1281, doi:10.1126/science.1128489, 2006.
- Shaw, W. J. and Lin, J.: Models of ocean ridge lithospheric deformation: dependence on crustal thickness, spreading rate, and segmentation, J. Geophys. Res., 101, 17977–17993, doi:10.1029/96JB00949, 1996.
 - Stadler, G., Gurnis, M., Burstedde, C., Wilcox, L.C., Alisic, L. and Ghattas, O.: The dynamics of plate tectonics and mantle flow: From local to global scales, Science, 329, 1033, doi:10.1126/science.1191223, 2010.
 - Staudigel, H., Park, K.-H., Pringle, M., Rubenstone, J., Smith, W., and Zindler, A.: The longevity of the South Pacific isotopic and thermal anomaly, Earth Planet. Sc. Lett., 102, 24–44, doi:10.1016/0012-821X(91)90015-
- 720 A, 1991.
 - Stegman, D., Farrington, R., Capitanio, F., and Schellart, W.: A regime diagram for subduction styles from 3-D numerical models of free subduction, Tectonophysics, 483, 29–45, doi:10.1016/j.tecto.2009.08.041, 2010.
 - Tarduno, J., Bunge, H.-P., Sleep, N., and Hansen, U.: The Bent Hawaiian-Emperor hotspot track: inheriting the mantle wind, Science, 324, 50–53, doi:10.1126/science.1161256, 2009.
- 725 Tarduno, J. A.: On the motion of Hawaii and other mantle plumes, Chem. Geol., 241, 234–247, doi:10.1016/j.chemgeo.2007.01.021, the Great Plume Debate: Testing the Plume Theory, 2007.
 - Wessel, P. and Kroenke, L. W.: Pacific absolute plate motion since 145 Ma: an assessment of the fixed hot spot hypothesis, J. Geophys. Res., 113, B06101, doi:10.1029/2007JB005499, 2008.
- Whittaker, J. M., Muller, R. D., Leitchenkov, G., Stagg, H., Sdrolias, M., Gaina, C., and Goncharov, A.:
 Major Australian-Antarctic plate reorganization at Hawaiian-Emperor Bend Time, Science, 318, 83–86, doi:10.1126/science.1143769, 2007.

Wilson, J.: A possible origin of the Hawaiian Islands, Can. J. Phys., 41, 863-870, doi:10.1139/p63-094, 1963.