

Abstract

In the changing world, the prevalence of land degradation is becoming a serious problem worldwide especially in countries with arid and semiarid rangelands. There are many techniques to assess rangeland degradation but most of them rely on classic science. So a study was conducted to find out how indigenous people assess rangeland degradation and how their ecological knowledge can be used for rangeland degradation assessment. We interviewed pastoralists of two sites (Mirza-Baylu and Dasht) where part of both areas is located in Golestan National Park (NE Iran). A structured questionnaire was designed based on some indicators taken from literature and also primary discussions with pastoralists in order to evaluate land degradation. A qualitative Likert scale was used for scoring rangeland degradation indicators. The results revealed that pastoralist pay first attention to edaphic indicators than vegetative and other indicators. There were significant differences between inside and outside of the park in rangeland degradation indicators for both sites. The results show that the rangelands outside the park in both sites were degraded compare to inside the park especially in the areas near to villages. It can be concluded that pastoralists own a vast amount of knowledge on the vegetation and grazing animal habits that can be used in rangeland degradation assessment and it is necessary to document their ecological indigenous knowledge and involve them in rangeland degradation assessment process.

1 Introduction

Rangelands are the vastest terrestrial ecosystems on the earth, covering close to 40 % of the world landscape, of which more than 80 % located in arid and semiarid zones. Soil is the most important component of rangeland ecosystems that has an interdisciplinary nature and is associated with biodiversity, biogeochemical cycling, hydrology, human health and social sciences (Brevik et al., 2015). Rangeland soils moreover offer services to the human societies and makes the Earth System stable (Keesstra et al.,

SED

7, 2999–3019, 2015

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2012; Berendse et al., 2012). Unfortunately, rangelands have undergone (and continue to undergo) rapid transformations as a result of factors such as overgrazing, deforestation, woody-plant encroachment, and invasion by non-native plant species (Wilcox and Thurow, 2006). Each of these factors has led to the reduction in the quantity or nutritional quality of the vegetation available for grazing that called rangeland degradation. And this resulted also in higher soil and water losses (Cerdà, 1998; Kröpfl et al., 2013; Li et al., 2013).

It is believed that livestock grazing is associated with rangeland degradation. Grazing is the most important factor affecting vegetation and soil in all rangelands of the world, having critical impacts on the rangeland biodiversity and species composition (Sharafatmandrad et al., 2014; Angassa, 2014), biological groups (Sharafatmandrad et al., 2014; Tarhouni et al., 2015), structure (Eckert and Spencer, 1987; Noy-Meir, 1979, 1993; Walker and Noy-Meir, 1982), goods and services (Papanastasis et al., 2015), function (White, 1979; Sousa, 1984; Hobbs and Huenneke, 1992), soil erosion (Tadesse and Penden, 2002; Palacio et al., 2014; Mekuria and Aynekulu, 2013), nutrient cycling (Frank et al., 1998; Ritchie and Tilman, 1995; Fernandez et al., 2008) and hydrological processes (Cerdà and Lavee, 1999; Hiernaux et al., 1999; Sharafatmandrad et al., 2010). However, there are evidences that grazing management activities, not grazing, is the main cause of rangeland degradation in arid and semi-arid environments (Gulelat, 2002). Pastoralism is a traditional range management activity, which focuses mostly on the natural forage rather than cultivated fodder (Sandford, 1983). Pastoralists usually own a vast amount of knowledge on their grazing lands, attained through long experiences and observations in herding practices (Oba and Kotile, 2001; Mapinduzi et al., 2003). To combat rangeland degradation, it is recommended that rangeland management systems integrate community perceptions and practices (Khwarae, 2006). Thus, the indigenous knowledge of the local communities can be used in conjunction with technical knowledge to manage natural resources (Khwarae, 2006). In many developing countries where rangelands are a dominant land type and critically important in livelihoods of a significant portion of the population, severe range-

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



land degradation can create significant social, economic, and environmental problems (Bedunah and Angerer, 2012). So scientific and indigenous knowledge should be integrated so that local communities be able to realize their capacity for monitoring and responding to the land degradation and environmental changes (Stringer and Reed, 2007). The resulting system for environmental management would improve the communities' livelihood and decrease rangeland degradation at the same time (Khwarae, 2006). That is interesting that pastoralists and ecologists are unanimous on most of the rangeland degradation indicators.

If we want rangeland degradation indicators to be applicable in land management, they must then be easy to use by local communities, accurate to assess environmental sustainability and result in conservation (Reed et al., 2008). In the other hand, the involvement of the pastoralists in planning and implementing land conservation programs require conservationists and technicians to be aware of environmental indicators used by pastoralists for assessing rangeland degradation. There are too many studies that conveying combination of local and scientific ecological knowledge may contribute to easy and accurate monitoring and management of natural resource changes by local communities (e.g., Folke et al., 2002; Thomas and Twyman, 2004; Fraser et al., 2006; Reed et al., 2007, 2008).

Bottom-up or local participation approaches implicate that pastoralists have accumulated a wealth of knowledge over time, based on long-term experiences that can complement scientific knowledge in environmental assessment and conservation (Richards, 1980). Recently it has become known that indigenous knowledge and local management play an important role in natural resource conservation (Warren, 1992; Berkes et al., 2000) and combat land degradation. Additionally, there is growing interest on how indigenous ecological knowledge and management practices can be used in collaboration with standard scientific methods for improved understanding of the environment and its changes (Dahlberg, 2000; Reed et al., 2007).

The history of pastoralism in Iran goes back to several thousand years ago, but indigenous ecological knowledge of pastoralism is neglected in most studies related

to the rangelands. So our main objective was to evaluate pastoralism' knowledge of rangeland degradation assessment, based on their perceptions and experiences. Our research questions were as follow:

- (a) What are the pastoralist' land evaluation criteria?
- (b) What indicators do the pastoralists use for degradation assessment? And
- (c) is there a possibility to combine scientific land degradation indicators with the pastoralists one to assess rangeland degradation?

2 Materials and methods

2.1 Study area

The sites used for this study are parts of Golestan National Park located in Golestan Province in north-eastern Iran (37.31–37.17° N, 53.04–55.43° E). The park was established in 1957 as the first National Park and Biosphere Reserve of the Middle East. Golestan National Park spans an area of 87 242 ha, comprising Caspian forests, steppe rangelands and the Juniper woodlands. The two rangeland sites selected for this study were Mirza-Baylu (37°19'29"–37°21'35" N and 56°13'56"–56°19'20" E; 1248–1310 m.a.s.l.) and Dasht (37°18'12"–37°19'37" N and 56°13'–56°1'33" E; 993–1058 m.a.s.l.). In each site, under grazing parts outside the park are separated from the enclosure parts inside the park by a narrow road (Fig. 1).

The Mirza-Baylu site is located at the eastern the park and is mostly flat, with slopes less than 5%, and some hilly lands occur just in a few parts. In this site, outside the Park, there is a village known Robot-e Qarebil, 5 km away from the Mirza-Baylu site. The mean annual temperature is 12.9 °C. The study site receives about 236 mm of annual precipitation. The site is dominated by relatively pure stands of the dwarf shrub *Artemisia sieberi* accompanied by some grasses. There are some saline parts in the site that are mostly occupied by halophytes such as *Salsola dendroides*, *Phragmites*

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



australis, *Suaeda physophora* and *Anabasis aphylla*. Also some rare species can be seen in the plains (e.g. *Diaphanoptera stenocalycina*).

The Dasht site is located in the southern part of the Golestan National Park. Most of the site is hilly but there are also a few flat areas. The mean annual precipitation and the mean annual temperatures are 191 mm and 11 °C respectively. The vegetation of this semi-steppe rangeland is consisting of grasses and shrubs, dominated by grasses *Bromus danthonia*, *Festuca ovina*, *Eremopyrum bonaepartis* and *Phleum paniculatum* and dwarf shrubs *Acantholimon pterostegium* and *Artemisia kopedaghensis*.

Regarding to grazing, inside the park is only grazed by wildlife but outside the park is grazed by the pastoralists herds consist of sheep and a few goats from early morning till afternoon. The herds are in their own fields all the seasons specified by the Department of Natural Resources. Dry forages and agricultural residuals (straw and hay) are used as winter forage for livestock in the both study sites.

2.2 Land degradation assessment

2.2.1 Selection of indicators

The pastoralists' first encounter is generally plagued by suspicion and fear because of government regulatory restrictions on rangeland use. As a first step, we tried to build a foundation of trust by connecting with educated ones, volunteering our personal information, showing interest in the pastoralism and lifestyle that were completely effective. The pastoralists have been then informed how important their indigenous knowledge is and no research in the region will be fulfilled without their viewpoints and help. So we go through the explaining the research and its objectives and make sure that the pastoralists are convinced how effective would be the results in their profession, economic status, rangelands health assessment and management.

Descriptive research was used to obtain information. So data were collected using both the documentary and field survey. By being present between the pastoralists, we have tried to gather data through participation and using Focus Group Discus-

SED

7, 2999–3019, 2015

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sion (FGD) technique, directive interviews and the narrative threads of the key figures and experienced persons. Through meetings and individual and group interviews, pastoralists were asked about ecological knowledge on rangeland degradation indicators and assessment. The most important part of the study was to discuss with interviewees about the importance of the pastoralist ecological knowledge on recognition of rangeland degradation and its assessment. To understand how pastoralists assess degradation at landscape level, the key questions were: what are the pastoralists' land evaluation criteria, what indicators did the pastoralists use for degradation assessment, and what are the roles of the degradation assessment in rangeland health assessment and restoration?

To understand pastoralists' perceptions of land degradation and its influence on rangeland conservation, the questions posed were: what do pastoralists think of a "good" or "bad" rangeland, and what indicators do the pastoralists use as signs of rangeland change from "good" to "bad", for the purposes of rangeland health and management (Roba, 2008)? The results of meetings and interviews were used to identify indicators related to rangeland degradation.

The indicators taken from the literature were discussed in the pastoralists' interviews with their own language and terminology so that they could understand the exact concept of the indicators. As it was expected, most of them had the same indicators as taken from literatures but with their own language. So the duplicates were removed and the new ones were added to the list. A structured questionnaire was designed based on the identified indicators to obtain data on rangeland degradation according to the Land Degradation Assessment in Drylands (LADA, 2011). So the indicators were assessed on a 5-point Likert-type scale ranging from very poor (0–20) to very good (80–100). Qualitative scale was used because of being more intuitive and easier to understand for pastoralists but it was necessary to convert it to a quantitative scale to compare inside and outside the park.

2.2.2 Field assessment

For the Mirza-Baylu site, 28 × 8 m² quadrats were randomly located throughout the region, 12 quadrats inside and 16 quadrats outside the Park. For the Dasht site, 22 × 8 m² quadrats were randomly located throughout the region, 15 quadrats inside and 18 quadrats outside the Park. Each quadrats was assessed by the 3 selected pastoralists (i.e. there were 3 replications). In total, 84 and 99 questionnaires were respectively filled for the Mirza-Baylu (36 inside and 48 outside the Park) and Dasht (45 inside and 54 outside the Park) sites. So the pastoralists were ranked the rangeland degradation indicators in each quadrats.

2.3 Data analyses

Each pastoralist was considered as a replication. The mean scores for each indicator was calculated and used to compare inside and outside the park. Comparisons were based on quantitative scale. Two-sample *t* tests were used for each site separately to determine if degradation indicators differed between two sites pastoralists. Indicators with significant differences were then compared for both sites as total to see if there is any differences between inside and outside the park. Statistical Package for the Social Sciences (SPSS, Version 18) was used for data analysis.

3 Results

Based on literature and indigenous ecological knowledge of pastoralists, 18 degradation indicators were identified and used for questionnaires preparation (Table 1). From 18 indicators, 56 % were classified as vegetative indicators, 33 % as edaphic indicators and 11 % as other indicators (i.e. indicators related to different aspects of rangelands apart from vegetation and soil).

Regarding the Likert scale, plots in Mirza-Baylu site had higher mean scores inside the park (3.249) than outside the park (3.026). According to pastoralists' assessment,

SED

7, 2999–3019, 2015

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



good to bad; simple indicators) are good because they can be easily adapted and used even by non-experts (Kapalanga, 2008). This helps in comparing areas, involves stakeholders as much as possible, and aids in land use and restoration planning and projects prioritizing (Kapalanga, 2008).

5 Conclusions

The traditional knowledge of local pastoralists in the both study sites was useful and important in the management of rangeland resources. Pastoralists have a wealth of interests for emphasizing on their own indicators to be more practical for the rangeland assessments. The pastoralists have a broad knowledge base covering materials from rangelands vegetation and animal habits to land characteristics. Controlling degradation in grazing lands without considering the people who have a substantial role in that will be imperfect. So matching the scientific land degradation indicators with the ones pastoralists are believed in and understand, can lead to the successfully control of land degradation. Involvement of pastoralists and documenting their knowledge on rangelands can provide useful bases for the sustainable utilization and conservation of natural rangelands. It is believed that such plans that are based on indigenous knowledge can be easily accepted by local people.

References

- Abate, T., Ebro, A., and Nigatu, L.: Traditional rangeland resource utilization practices and pastoralists' perceptions on land degradation in south-east Ethiopia, *Trop. Grasslands*, 44, 202–212, 2010.
- Angassa, A.: Effects of grazing intensity and bush encroachment on herbaceous species and rangeland condition in southern Ethiopia, *Land Degrad. Dev.*, 25, 438–451, 2014.

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bedunah, D. J. and Angerer, J. P.: Rangeland degradation, poverty, and conflict: how can rangeland scientists contribute to effective responses and solutions?, *Rangeland Ecol. Manage.*, 65, 606–612, doi:10.2111/REM-D-11-00155.1, 2012.

Berendse, F., van Ruijven, J., Jongejans, E., and Keesstra, S. D.: Loss of plant species diversity reduces soil erosion resistance of embankments that are crucial for the safety of human societies in low-lying areas, *Ecosystems*, 18, 881–888, doi:10.1007/s10021-015-9869-6, 2015.

Berkes, F., Colding, J., and Folke, C.: Rediscovery of traditional ecological knowledge as adaptive management, *Ecol. Appl.*, 10, 1251–1262, doi:10.1890/1051-0761(2000)010[1251:ROTEKA]2.0.CO;2, 2000.

Brevik, E. C., Cerdà, A., Mataix-Solera, J., Pereg, L., Quinton, J. N., Six, J., and Van Oost, K.: The interdisciplinary nature of soil, *Soil*, 1, 117–129, doi:10.5194/soil-1-117-2015, 2015.

Cerdà, A.: Changes in overland flow and infiltration after a rangeland fire in a Mediterranean scrubland, *Hydrol. Process.*, 12, 1031–1042, doi:10.1002/(SICI)1099-1085(19980615)12:7<1031::AID-HYP636>3.0.CO;2-V, 1998.

Cerdà, A. and Lavee, H.: The effect of grazing on soil and water losses under arid and Mediterranean climates, *Implications for desertification*, Pirineos, 159, 153–154, 1999.

Dahlberg, A. C.: Interpretations of environmental change and diversity: a critical approach to indications of degradation – the case of Kalakamate, Northeast Botswana, *Land Degrad. Dev.*, 11, 549–562, doi:10.1002/1099-145X(200011/12)11:6<549::AID-LDR413>3.0.CO;2-5, 2000.

Eckert, R. E. and Spencer, J. S.: Growth and reproduction of grasses heavily grazed under rest-rotation management, *J. Range Manage.*, 40, 156–159, 1987.

Fernandez, D. P., Neff, J. C., and Reynolds, R. L.: Biogeochemical and ecological impacts of livestock grazing in semi-arid southeastern Utah, USA, *J. Arid Environ.*, 72, 777–791, 2008.

Fernandez-Gimenez, M. E.: The role of Mongolian nomadic pastoralists' ecological knowledge in rangeland management, *Ecol. Appl.*, 10, 1318–1326, doi:10.1890/1051-0761(2000)010[1318:TROMNP]2.0.CO;2, 2000.

Folke, C., Carpenter, S., Elmqvist, T., Gunderson, L., Holling, C. S., and Walker, B. H.: Resilience and sustainable development: building adaptive capacity in a world of transformations, *Ambio*, 31, 437–440, 2002.

Frank, D. A., McNaughton, S. J., and Tracy, B. F.: The ecology of the Earth's grazing ecosystems, *Bioscience*, 48, 513–521, 1998.

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fraser, E. D. G., Dougill, A. J., Mabee, W. E., Reed, M. S., and McAlpine, P.: Bottom up and top down: analysis of participatory processes for sustainability indicator identification as a pathway to community empowerment and sustainable environmental management, *Environ. Manage.*, 78, 114–127, 2006.

Gulelat, W.: Household Herd Size Among Pastoralists in Relation to Overstocking and Rangeland Degradation, International Institute for Geoinformation Science and Earth Observation, Enschede, the Netherlands, 2002.

Hiernaux, P., Biolders, C. L., Valentin, C., Bationo, A., and Fernandez-Rivera, S.: Effects of livestock grazing on physical and chemical properties of sandy soils in Sahelian rangelands, *J. Arid Environ.*, 41, 231–245, 1999.

Hobbs, R. J. and Huenneke, L. F.: Disturbance, diversity and invasion: implications for conservation, *Conserv. Biol.*, 6, 324–337, 1992.

Kapalanga, T. S.: A Review of Land Degradation Assessment Methods, Land Restoration Training Program, Keldnaholt, 112 Reykjavic, Iceland, 2008.

Keesstra, S. D., Geissen, V., van Schaik, L., Mosse, K., and Piirani, S.: Soil as a filter for groundwater quality, *Curr. Opinions Environ. Sustain.*, 4, 507–516, doi:10.1016/j.cosust.2012.10.007, 2012.

Khwarae, G.: Community Perceptions of Rangeland Degradation and Management Systems in Loologane and Shadishadi, Kweneng North, Botswana, Department of International Environment and Development Studies, Norway, 2006.

Kröpfl, A. I., Cecchi, G. A., Villasuso, N. M., and Distel, R. A.: Degradation and recovery processes in semi-arid patchy rangelands of northern Patagonia, Argentina, *Land Degrad. Dev.*, 24, 393–399, doi:10.1002/ldr.1145, 2013.

LADA: LADA Project Document, Field Manual for Local Level Land Degradation Assessment in Dry Lands, FAO, available at: http://www.fao.org/fileadmin/templates/nr/kagera/Documents/LADA_manuals/MANUAL2_final_draft.pdf, 2011.

Li, X.-L., Gao, J., Brierley, G., Qiao, Y.-M., Zhang, J., and Yang, Y.-W.: Rangeland degradation on the Qinghai-Tibet plateau: implications for rehabilitation, *Land Degrad. Dev.*, 24, 72–80, doi:10.1002/ldr.1108, 2013.

Mapinduzi, A. L., Oba, G., Weladji, R. B., and Colman, J. E.: Use of indigenous ecological knowledge of the Maasai pastoralists for assessing rangeland biodiversity in Tanzania, *African J. Ecol.*, 41, 329–336, doi:10.1111/j.1365-2028.2003.00479.x, 2003.

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mekuria, W. and Aynekulu, E.: Exclosure land management for restoration of the soils in degraded communal grazing lands in Northern Ethiopia, *Land Degrad. Dev.*, 24, 528–538, doi:10.1002/ldr.1146, 2013.

Noy-Meir, I.: Structure and function of desert ecosystems, *Israel J. Bot.*, 28, 1–19, 1979.

5 Noy-Meir, I.: Compensating growth of grazed plants and its relevance to the use of rangelands, *Ecol. Appl.*, 3, 32–34, 1993.

Oba, G.: Harnessing pastoralists' indigenous knowledge for rangeland management: three African case studies, in: *Pastoralism: Research, Policy and Practice*, 2012, 2, 1–25, doi:10.1186/2041-7136-2-1, 2012.

10 Oba, G. and Kotile, D. G.: Assessments of landscape level degradation in southern Ethiopia: pastoralists vs. ecologists, *Land Degrad. Dev.*, 12, 461–475, doi:10.1002/ldr.463, 2001.

Palacio, R. G., Bisigato, A. J., and Bouza, B. J.: Soil erosion in three grazed plant communities in northeastern Patagonia, *Land Degrad. Develop.*, 25, 594–603, doi:10.1002/ldr.2289, 2014.

15 Papanastasis, V. P., Bautista, S., Chouvardas, D., Mantzanas, K., Papadimitriou, M., Mayor, A. G., Koukioumi, P., Papaioannou, A., and Vallejo, R. V.: Comparative assessment of goods and services provided by grazing regulation and reforestation in degraded Mediterranean rangelands, *Land Degrad. Dev.*, doi:10.1002/ldr.2368, in press, 2015.

Reed, M. S., Dougill, A. J., and Taylor, M. J.: Integrating local and scientific knowledge for adaptation to land degradation: Kalahari rangeland management options, *Land Degrad. Dev.*, 18, 249–268, doi:10.1002/ldr.777, 2007.

20 Reed, M. S., Dougill, A. J., and Baker, T. R.: Participatory indicator development: what can ecologists and local communities learn from each other?, *Ecol. Appl.*, 18, 1253–1269, doi:10.1890/07-0519.1, 2008.

25 Richards, P.: Community environmental knowledge in African rural development, in: *Indigenous Knowledge Systems and Development*, edited by: Brokensha, D., Warren, D. M., and Werner, O., University Press of America, New York, 181–194, 1980.

Ritchie, M. E. and Tilman, D.: Responses of legumes to herbivores and nutrients during succession on a nitrogen-poor soil, *Ecology*, 76, 2648–2655, 1995.

30 Roba, H. G.: *Global Goals, Local Actions: A Framework for Integrating Indigenous Knowledge and Ecological Methods for Rangeland Assessment and Monitoring in Northern Kenya*, PhD thesis, Department of International Environment and Development Studies, Norwegian University of Life Sciences (UMB), Norway, 2008.

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Sandford, S.: Management of Pastoral Development in the Third World, Overseas Development Institute, London, 1983.
- Sharafatmandrad, M., Mesdaghi, M., Bahremand, A., and Barani, H.: The role of litter in rainfall interception and maintenance of superficial soil water content in an arid rangeland in Khabr National Park in south-eastern Iran, *Arid Land Res. Manag.*, 24, 213–222, 2010.
- Sharafatmandrad, M., Sepehry, A., and Barani, H.: Plant species and functional types' diversity in relation to grazing in arid and semi-arid rangelands, Khabr National Park, Iran, *J. Rangeland Sci.*, 4, 203–214, 2014.
- Sousa, W. P.: The role of disturbance in natural communities, *Ann. Rev. Ecol. Sys.*, 15, 353–391, 1984.
- Stringer, L. C. and Reed, M. S.: Land degradation assessment in southern Africa: integrating local and scientific knowledge bases, *Land Degrad. Dev.*, 18, 99–116, 2007.
- Tadesse, G. and Penden, D.: Livestock grazing impact on vegetation, soil and hydrology in a tropical highland watershed, in: *Integrated water and land management research and capacity building priorities*, edited by: McCornick, P. G., Karma, A. B., and Tadesse, G., *Proceedings of MoWR/EARO/IWMI/ILRI International Workshop, Addis Ababa, Ethiopia, 2–4 December 2002*, ILRI, 87–97, 2002.
- Tarhouni, M., Hmida, W. B., and Neffati, M.: Long-term changes in plant life forms as a consequence of grazing exclusion under arid climatic conditions, *Land Degrad. Dev.*, doi:10.1002/ldr.2407, in press, 2015.
- Thomas, D. S. G. and Twyman, C.: Good or bad rangeland? Hybrid knowledge, science and local understandings of vegetation dynamics in the Kalahari, *Land Degrad. Dev.*, 15, 215–231, doi:10.1002/ldr.610, 2004.
- Turner, N. J., Ignace, B., and Ignace, R.: Traditional ecological knowledge and wisdom of aboriginal peoples in British Columbia, *Ecol. Appl.*, 10, 1275–1287, doi:10.1890/1051-0761(2000)010[1275:TEKAWO]2.0.CO;2, 2000.
- Walker, B. H. and Noy-Meir, I.: Aspects of the stability and resilience of savanna ecosystems, in: *Ecology of Tropical Savannas*, edited by: Huntley, B. J. and Walker, B. H., Springer, Berlin, 556–590, 1982.
- Warren, D.: Indigenous knowledge, biodiversity conservation and development, Keynote address at the International Conference on Conservation of Biodiversity in Africa: Local Initiatives and Institutional Roles, 30 August–3 September 1992, Nairobi, Kenya, 102, 1992.

White, P. S.: Pattern, process, and natural disturbance in vegetation, *Bot. Rev.*, 45, 229–299, 1979.

Wilcox, B. P. and Thurow, T. L.: Preface “Emerging issues in rangeland ecohydrology”, *Hydrol. Process.*, 20, 3155–3157, 2006.

SED

7, 2999–3019, 2015

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Identified rangeland degradation indicators based on literature and indigenous ecological knowledge. Indicators related to each category is shown with a check mark.

Attributes	Indicators	Literature	Mirza-Baylu Pastoralists	Dasht Pastoralists
Vegetation	Decrease of vegetation productivity	✓	✓	✓
	Loss of phytodiversity	✓		✓
	Removal of palatable plants	✓		✓
	Increase in poisonous plants	✓	✓	✓
	Decrease of shrubs	✓		
	Increase in annual plants	✓		
	Increase in the distance between plants	✓		
	Decrease of plants height	✓		
Soil	Loss of litter mass	✓		✓
	Soil salination	✓	✓	
	Decrease of ground cover	✓	✓	✓
	Increase in bare soil	✓	✓	✓
	Soil muddiness		✓	
	Decrease of soil sandiness		✓	
	Decrease of soil infiltration	✓	✓	✓
	Increase in soil looseness		✓	✓
Other	Increased risk of wildfires			✓
	Increased risk of pest damage		✓	

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Comparison of all identified rangeland degradation indicators between inside and outside the park for each site separately.

Indicators	Sites					
	Mirza-Baylu			Dasht		
	Inside the park	Outside the park	<i>P</i>	Inside the park	Outside the park	<i>P</i>
Decrease of vegetation productivity	3.35 ± 0.64	3.24 ± 0.75	0.85	3.44 ± 1.036	2.64 ± 0.69	0.04
Loss of biodiversity	3.22 ± 0.56	3.15 ± 0.91	0.58	3.51 ± 0.92	2.55 ± 0.73	0.009
Removal of palatable plants	3.26 ± 0.7	3.28 ± 0.92	0.64	3.55 ± 1.06	2.57 ± 0.66	0.02
Increase in poisonous plants	3.35 ± 1.32	4.42 ± 0.81	0.02	4.35 ± 0.51	4.55 ± 0.24	0.38
Decrease of shrubs	2.99 ± 0.79	3.39 ± 0.86	0.1	2.66 ± 0.56	2.52 ± 0.58	0.58
Increase in annual plants	2.99 ± 0.77	2.24 ± 0.77	0.02	3.11 ± 1.47	2.59 ± 0.69	0.63
Decrease of ground cover	3.55 ± 0.53	2.71 ± 0.79	0.003	2.79 ± 1.25	2.15 ± 0.74	0.19
Increase in the distance between plants	3.59 ± 0.52	2.86 ± 0.72	0.005	3.44 ± 0.95	2.68 ± 0.47	0.03
Soil salination	4.35 ± 0.23	4.62 ± 0.57	0.02	4.19 ± 0.88	4.73 ± 0.40	0.06
Loss of litter mass	2.97 ± 0.73	2.17 ± 0.85	0.008	3.27 ± 1.21	2.33 ± 0.76	0.03
Increase in bare soil	3.95 ± 0.71	3.24 ± 0.77	0.01	3.86 ± 1.05	2.796 ± 0.69	0.007
Soil muddiness	3.57 ± 1.01	3.84 ± 0.89	0.5	3.14 ± 1.04	2.68 ± 1.20	0.31
Decrease of plants height	2.89 ± 0.37	2.84 ± 0.63	0.76	3.15 ± 0.74	2.73 ± 0.44	0.08
Decrease of soil sandiness	2.53 ± 1.38	2.51 ± 1.21	0.85	3.13 ± 0.95	3.04 ± 0.94	0.82
Decrease of soil infiltration	3.71 ± 0.56	3.42 ± 0.73	0.32	3.13 ± 1.05	3.02 ± 1.01	0.56
Increase in soil looseness	2.19 ± 0.84	2.29 ± 0.78	0.77	2.28 ± 0.94	2.33 ± 0.87	0.8
Increased risk of wildfires	2.17 ± 0.59	1.08 ± 0.15	2.58 × 10 ⁻⁶	2.79 ± 1.53	1.99 ± 0.78	0.3
Increased risk of pest damage	4.08 ± 0.26	4.15 ± 0.99	0.17	3.66 ± 0.59	3.93 ± 0.54	0.22

Rangeland degradation assessment

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Comparison of significant degradation indicators between inside and outside the park for both sites as total.

Indicators	Rank	CV inside the Park	CV outside the Park	p
Decrease of ground cover	1	0.003	0.29	0.15
Increase in the distance between plants	2	0.005	0.25	0.14
Loss of litter mass	3	0.008	0.39	0.24
Increase in bare soil	4	0.01	0.24	0.18
Soil salination	5	0.02	0.12	0.05
Increase in annual plants	6	0.02	0.34	0.26
Increase in poisonous plants	7	0.02	0.18	0.39
Increase in bare soil	1	0.007	0.25	0.27
Loss of biodiversity	2	0.009	0.28	0.26
Removal of palatable plants	3	0.02	0.25	0.29
Loss of litter mass	4	0.03	0.33	0.27
Increase in the distance between plants	5	0.03	0.18	0.27
Decrease of plant production	6	0.04	0.26	0.29

**Rangeland
degradation
assessment**

B. Behmanesh et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 1. Map of study area in Golestan National Park, Golestan Province, Iran. Dasht site was located in the southern park and Mirza-baylu site was located in the eastern park. The points are sampling plots.