

This discussion paper is/has been under review for the journal Solid Earth (SE).
Please refer to the corresponding final paper in SE if available.

CO₂ emission and structural characteristics of two calcareous soils amended with municipal solid waste and plant residue

N. Yazdanpanah

Department of Water Engineering, Kerman Branch, Islamic Azad University, Kerman, Iran

Received: 24 October 2015 – Accepted: 29 October 2015 – Published: 9 November 2015

Correspondence to: N. Yazdanpanah (najmeyazdanpanah@yahoo.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

This investigation examines the effect of different amendments on selected soil physical and biological properties over a twenty four month period in two cropland fields. Urban municipal solid waste (MSW) compost and alfalfa residue (AR) were used as different organic amendments at the rates of 0 (control), 10 and 30 Mgha⁻¹ to a clay loam soil and a loamy sand soil in a semiarid region. Result showed that the soil improvement was controlled by the application rate and decomposability of amendments and soil type. The addition of organic amendments to the soils improved aggregate stability and consequently enhanced total porosity, especially macro pores fraction. The increased soil organic carbon (SOC) and total porosity values as compared to the control treatment were greater in the loamy sand soil than in the clay loam soil. Moreover, compared to the microbial respiration of control plots, the application of MSW resulted in higher values of microbial respiration in the clay loam soil than in the loamy sand soil, whereas the reverse order was found for AR. Linear and power functions were provided for the relationships between microbial respiration and SOC in the loamy sand and clay loam soils, respectively. Also, CO₂ emission was stimulated significantly as power functions of the total porosity and the ratio of macro to micro pores. However, the soil microbial respiration and carbon storage improved aggregate stability and pore size distribution, as a response, soil porosity especially macro pores fraction controlled CO₂ flux.

1 Introduction

The maintenance of soil organic carbon (SOC) is important for the sustainable productivity of agroecosystems (González et al., 2010). In this issue, the carbon sequestration is of importance for the earth system (Jaijarree et al., 2014; Parras-Alcántara and Lozano-García, 2014; Bruun et al., 2015; de Graaff et al., 2015). Because most soils of arid and semiarid regions are poor in organic carbon (Tejada and González, 2003), the

SED

7, 3151–3177, 2015

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SED

7, 3151–3177, 2015

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



application of organic amendments is a management strategy to improve soil properties with relevant economic benefits for farmers (Bronick and Lal, 2005; González et al., 2010). Therefore, organic amendment has been widely used to increase the content of SOC (Fernández et al., 2009; Benbouali et al., 2013; Mahmoodabadi et al., 2013).

5 In this regard, the application of crop residues and manure has been suggested to improve soil quality and to support the sustainable production in agricultural land (Yu and Jia, 2014; Kaleeem Abbasi et al., 2015; Musinguzi et al., 2015; Turgut, 201). Because, the amount of livestock manure as traditional organic product is limited, crop residue as an exogenous source of organic matter has been widely used for the remediation of soil (Benbouali et al., 2013; Mahmoodabadi and Heydarpour, 2014; Novara et al., 2015). The application of organic municipal solid waste (MSW) compost is an alternative approach in agricultural land (Aggelides and Londra, 2000; Ferreras et al., 2006). Exponential growth of population and urbanization coupled with the improvement of living standards have resulted in an increase in the amount of urban MSW generation throughout the world (Karak et al., 2012), so that the global generation of MSW exceeds 2 billion Mg yr⁻¹. Therefore, in recent years the composted urban MSW has been added to agricultural soils for waste disposal and to improve soil quality (García-Gil et al., 2000). Soil application of organic amendments, such as animal manure, crop residue and MSW compost provides management strategies to compensate the removal of organic carbon from the soils. Besides, the use of organic amendments in the soils reduces the serious environmental problems caused by residue accumulation (Tejada and González, 2003; Ferreras et al., 2006).

20 The addition of organic amendments can improve soil physical, chemical and biological properties (Yazdanpanah et al., 2013). Soil organic matter plays an important role in improving soil structure through maintaining aggregate stability (Cerdà, 1998a; Benbouali et al., 2013). In arid and semiarid regions, an increase in SOC results in the formation of more stable aggregates (Bronick and Lal, 2005) and in the reduction of soil erosion (Mahmoodabadi et al., 2014a, b; Mahmoodabadi and Arjmand Sajjadi, 2016). Therefore, the organic sources have been used in these areas to reduce soil

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



degradation (Hueso-González et al., 2014; Srivastava et al., 2014). On the other hand, soil aggregate stability influences several aspects related to the soil behavior, such as pore size distribution, water infiltration and runoff generation (Cerdà, 2000; Maza-heri and Mahmoodabadi, 2012; Sirjani and Mahmoodabadi, 2014; Arjmand Sajjadi and Mahmoodabadi, 2015a, b). In fact, there is an interaction between soil physical and biological properties following the application of an organic amendment. For instance, a positive relationship was found between soil porosity and microbial respiration (Marinari et al., 2000). In general, the application of organic amendments can stimulate soil microbial respiration (Ferrerias et al., 2006), in order that a higher respiration occurs in those treatments applied at a higher rate (Marinari et al., 2000; González et al., 2010). An increase in the microbial respiration of an amended soil may cause to the improvement of soil aggregate stability and porosity (Balashov et al., 2010). Also, the influence of organic amendments on improving soil aggregate stability not only depends on the quantity but also on the quality of adding organic materials specially their rate of decomposability and their capacity to induce soil microbial activity (Benbouali et al., 2013). Apart from the type and application rate of organic inputs, soil texture plays an important role in carbon stock. Mahmoodabadi and Heydarpour (2014) found that CO₂ emission to the atmosphere is much more in a coarse-textured soil compared to a fine-textured soil. Furthermore, cover and also vegetation are some other parameters controlling the content of soil organic carbon (Cerdà, 1998b; Cerdà and Doerr, 2005; Jiménez et al., 2013; Mahmoodabadi and Cerdà, 2013; Cerdà et al., 2015).

In most previous studies carried out about the effect of organic amendments on soil properties under field conditions, only one agricultural field (soil type) has been examined. In the present study, two contrasting agricultural fields (two soils with different textures) were examined. Furthermore, little is known about the interaction between microbial respiration and structural porosity in soils with different degrees of aggregate stability, especially in semiarid region soils. Therefore, the aims of the present work were (1) to attribute soil microbial respiration to aggregate stability and porosity fractions (i.e. macro and micro pores) in response to the type and application rate

of organic amendments under field conditions, and (2) to compare the effect of two types of organic sources including alfalfa residue (AR) and urban MSW compost on soil CO₂ emission from two different cropland soils. The findings of the present experiment should therefore enhance our understanding of the interrelationship between the microbial respiration and soil structural characteristics in contrasting soils.

2 Materials and methods

2.1 Experimental sites description

This research was conducted in two different agricultural fields both located in a same semiarid climate conditions in Kerman province, central Iran (30°14' N and 57°06' E). The first experimental field is placed on a clay loam soil (Haplocalcids) and the second is located on a loamy sand soil established on Aeolian deposits (Torripsamments), hereafter called “clay loam soil” and “loamy sand soil”, respectively. A long-term mean precipitation of the area is 140 mm yr⁻¹, which mainly occurs in winter and the average annual temperature is 16.5 °C. Both the experimental fields had been under agricultural cropping for more than 10 years, with a conventional management. Irrigation has been performed with water having an electrical conductivity of 1.1 dS m⁻¹ and sodium adsorption ratio of 0.73. Prior to the start of the experiment, the fields were rested under fallow for 2 years and were not fertilized to make them more homogeneous. Some selected properties of the soils before the amendments incorporation are presented in Table 1. Particle size distribution of the soils is different, in order that the loamy sand soil contains much more sand particles, nevertheless it has less value of mean weight diameter (MWD) of aggregates. As a key parameter, the content of organic carbon is higher in the clay loam soil (2.9 g kg⁻¹) than in the loamy sand soil (1.33 g kg⁻¹). However, the soils are poor in organic matter, they have considerable amounts of CaCO₃ (16.2 and 21.5 %) as is common in the region.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Organic amendments

In this study, two organic inputs from different sources as well as C : N ratios including urban MSW compost and alfalfa residue (AR) were applied. The urban MSW compost was obtained from the organic solid waste of Kerman Municipality. Alfalfa residue was used as a green manure, which is commonly associated with organic farming and can play an important role in sustainable cropping systems. Chemical composition of the organic amendments was measured. Electrical conductivity (EC) and pH were measured 24 h after 1 h shaking of 1 g samples in vials with 5 mL distilled water. The amounts of organic carbon and total nitrogen were measured by the Walkley and Black (1934) and Kjeldahl methods, respectively (Pansu and Gautheyrou, 2006). The measured chemical composition of amendments is presented in Table 2. As is clear, AR has higher organic carbon and lower amount of total nitrogen compared to MSW. The obtained C : N ratio of AR and MSW is 22.3 and 13.6, respectively (Table 2).

2.3 Experimental design

The experiment was established in a randomized complete block design with five treatments each at three replicates on two separate agricultural fields. For each field (soil texture), 15 experimental plots of 3 m × 5 m were established, so that totally 30 plots were prepared. The applied treatments were: (1) control, without any amendment application (C), (2) municipal solid waste at a rate of 10 Mg ha⁻¹ (MSW10), (3) municipal solid waste at a rate of 30 Mg ha⁻¹ (MSW30), (4) alfalfa residue at a rate of 10 Mg ha⁻¹ (AR10), and (5) alfalfa residue at a rate of 30 Mg ha⁻¹ (AR30). The rates of organic amendments were based on dry matter. The cured amendments were passed through a 5 mm mesh screen before soil incorporation (Aggelides and Londra, 2000). Then, the organic amendments were manually spread uniformly on the surface of the specified plots and they were incorporated into the top 15 cm of the soil profile. Similar procedures were followed for the two fields. During the experiment, no crop was planted and no fertilizer was applied.

2.4 Measurement of soil properties

Twenty four months after the application of amendments, soil samples were taken after mixing the four subsamples from each plot at depths of 0–15 cm. All samples were immediately stored in sealed plastic bags in a cooler and transported to laboratory.

The air dried soil samples crushed to pass through a 2 mm sieve and, some physical and chemical properties were measured. Soil microbial respiration was measured on fresh soil samples (Benbouali et al., 2013).

To study the effects of applied organic amendments on biological properties, soil microbial respiration was measured by an incubation-alkaline absorption method (Yazdanpanah et al., 2013). The production of CO₂ was measured as indicator of soil microbial activity. Soil samples (100 g) at 75 % of water holding capacity were incubated at 25 °C 7d in hermetic flasks; the CO₂ evolved was trapped in excess of 0.5 N NaOH. The alkali was titrated to the phenolphthalein with HCl in the presence of BaCl₂ to precipitate the carbonate. The CO₂ evolved was calculated by difference between samples and blanks without soil (Ferrerias et al., 2006; Yazdanpanah et al., 2013).

Soil organic carbon was measured as described by Walkley and Black (1934). Briefly, organic matter from the soil (1 g) was oxidized with K₂Cr₂O₇ 1 N (10 mL) in concentrated sulphuric acid for 30 min, followed by titration of the excess of K₂Cr₂O₇ with ferrous- ammonium sulfate 0.5 N and N-phenyl anthranilic acid to indicate the end point (González et al., 2010).

A part of soil samples was provided for measurement of aggregate stability as the proportion of aggregates that were stable to water. For this purpose, each soil sample was initially pre-wetted with distilled water. After the soil sample was saturated, it was placed on a 0.25 mm mesh sieve, and was sieved in distilled water at 30 oscillations per min for 2 min. The remaining soil on each sieve was dried in oven (105 °C), and then sand and aggregates were separated. After sand content correction, the fraction of aggregates larger than 0.25 mm was expressed as WSA > 0.25 mm (Six et al., 2001; Mahmoodabadi and Ahmadbeygi, 2013). Sample pre-treatments with water allow the

SED

7, 3151–3177, 2015

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



evaluation of the main factors involved in structural stability. The pre-treatment with water, since it do not allow air expulsion from the aggregate, indicated that aggregate disruption due to the wetting process occurred (Ferrerias et al., 2006).

Total porosity and included components (i.e. macroporosity and microporosity) were calculated according to Celik et al. (2004). At first, soil bulk density (BD) was determined on undisturbed samples using cylinder method, being made of 100 cm³ cylinders, after drying 24 h in 105 °C ovens. Then, total porosity (F) was calculated based on $F = 1 - (BD/2.65)$. Macroporosity fraction was determined from the volumetric water content using a pressure membrane apparatus at field capacity. Microporosity was calculated as the difference between total porosity and macroporosity values (Celik et al., 2004).

2.5 Statistical analysis

To study the effects of applied treatments on the selected soil properties, the obtained data were subjected to analysis of variances (ANOVA) procedure for a randomized complete block design with three replications. Comparison of means was performed by the Duncan multiple range test at 95 % level of probability. The soil microbial respiration was related to SOC, total porosity as well as macro/micro porosity ratio by regression functions. All the statistical analyses were performed in the SAS system (SAS Institute, 1990).

3 Results

3.1 Soil organic carbon

The effect of treatments on SOC was significant ($p < 0.05$) in loamy sand and clay loam soils (Table 3). The addition of AR and MSW caused to significant increments ($p < 0.05$) in SOC compared to the control plots, with larger increases at higher rate of application (i.e. 30 Mg ha⁻¹). Also, the soils amended with AR exhibited significantly

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



($p < 0.05$) higher levels of SOC than those amended with MSW. The SOC concentration under the application rate of 10 Mg ha^{-1} MSW and AR was about 1.5 and 1.7 times higher in the loamy sand soil and about 1.1 and 1.3 times higher in the clay loam soil than those obtained for unamended soils, respectively. The comparison with the control plots demonstrated that the addition of 30 Mg ha^{-1} MSW and AR resulted in 2.6 and 3.0 times higher SOC in the loamy sand soil and 1.5 and 1.7 times higher SOC in the clay loam soil, respectively. This means that as compared with the control plots, the increased SOC concentrations due to AR incorporation into both soils were more than the values resulted from MSW application. Also, with respect to control, SOC was enhanced following the application of organic amendments in the loamy sand soil more than did in the clay loam soil.

3.2 Soil microbial respiration

The effect of treatments on the microbial respiration of soils with different textures is provided in Table 3. As is clear, the application of alfalfa residue (AR) and municipal solid waste (MSW) led to significant ($p < 0.05$) increases in the soil microbial respiration compared to the control plots. Also, there were significant ($p < 0.05$) differences in the microbial respiration between MSW and AR for each soil. The maximum CO_2 flux was found in plots amended with 30 Mg ha^{-1} MSW, so that the highest values for loamy sand and clay loam soils were respectively 182.1 and $261.5 \mu\text{g CO}_2 \text{ g}^{-1}$ soil. The values of microbial respiration for 10 and 30 Mg ha^{-1} application rates of MSW were respectively 2 and 3 times higher in the loamy sand soil and 2.1 and 3.3 times higher in the clay loam soil than those values obtained for unamended soils. The addition of 10 and 30 Mg ha^{-1} AR stimulated the CO_2 emission by 1.5 and 2.5 times higher in the loamy sand soil and by 1.2 and 1.9 times higher in the clay loam soil with respect to the control plots, respectively. In fact, comparison between the amendment treatments suggests that for both the soils, those plots amended with MSW showed significantly ($p < 0.05$) higher values of microbial respiration than those amended with AR. Also, compared to

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the soil microbial respiration of control plots, the application of MSW caused to greater increments in the microbial respiration in the clay loam soil than in the loamy sand soil, whereas the reverse order was found for AR. On the other hand, higher rate of organic amendments added to the soils (i.e. 30 Mg ha⁻¹), greater increase in CO₂ production in relation to the control was found.

3.3 Soil aggregate stability

The applied treatments showed significant influences ($p < 0.05$) on the percentage of water stable aggregates of both the soils (Table 3). In the loamy sand soil, the addition of organic amendments, especially MSW increased the aggregate stability values compared to the control plots. Following 10 and 30 Mg ha⁻¹ application rates of MSW, the aggregate stability levels were respectively 1.6 and 1.9 times higher than those found for the control plots, while for the corresponding application rates of AR, a moderate increase was observed. In the clay loam soil, plots amended with both rates of MSW and those amended with 30 Mg ha⁻¹ AR showed a significant increment ($p < 0.05$) in aggregate stability. In this soil, the aggregate stability values for 10 and 30 Mg ha⁻¹ application rates of MSW was respectively 1.1 and 1.2 times higher than those for unamended soil. Furthermore, the values of aggregate stability for the clay loam soil were in general, much more than those for the loamy sand soil, although as compared to the control plots, the loamy sand soil experienced higher increases in aggregate stability.

3.4 Soil porosity components

At the end of field experiment, the applied treatments showed significant effects ($p < 0.05$) on the total porosity of soils (Table 3). In the loamy sand soil, MSW and AR had similar effects on total porosity, whereas a different pattern was observed in the clay loam soil. Depends on the type and application rates of amendments added to the soils, the total porosity of loamy sand soil varied from 0.32 to 0.47 cm³ cm⁻³ and in the clay loam soil it ranged from 0.41 to 0.59 cm³ cm⁻³. The total porosity values produced

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by 10 and 30 Mg ha⁻¹ application rates of MSW were respectively 1.3 and 1.5 times higher in the loamy sand soil and 1.2 and 1.4 times higher in the clay loam soil than those observed in the control plots. The addition of 10 and 30 Mg ha⁻¹ AR resulted in increases in the total porosity by 1.2 and 1.5 times in the loamy sand soil and by 1.1 and 1.3 times in the clay loam soil as compared to the control plots, respectively. This result indicates that the increased total porosity with respect to the control plots was more pronounced in the loamy sand soil than in the clay loam soil.

In addition to the total porosity, the macroporosity of soils was influenced significantly ($p < 0.05$) by the organic amendment treatments (Table 3). The incorporation of amendments into the soils, especially at the higher rate (30 Mg ha⁻¹) enhanced significantly ($p < 0.05$) the fraction of soil volume allocated to macro pores. In general, due to different treatments studied, the macro pores fraction varied from 11 to 25 cm³ cm⁻³ in the loamy sand soil and from 12 to 28 cm³ cm⁻³ in the clay loam soil. The soils amended with MSW gave significantly ($p < 0.05$) higher levels of macroporosity than those treated with AR. On the other hand, the maximum value of macroporosity in both the soils was found under the 30 Mg ha⁻¹ application rate of MSW. Considering the different texture of soils, due to MSW application the increased macroporosity in relation to the control plots was similar in both the soils, whereas the application of AR led to higher increments in the macroporosity in the loamy sand soil than that in the clay loam soil.

The result also indicated that all the treatments except AR at the application rate of 30 Mg ha⁻¹, did not show any significant effect on the soil microporosity (Table 3). In other words, only the application of 30 Mg ha⁻¹ AR caused to a significant increase ($p < 0.05$) in the soil microporosity compared to the control plots. In all cases, the microporosity of clay loam soil was in general greater than that of the loamy sand soil.

4 Discussion

Results of this study indicated that in general, application of MSW and AR had positive effects on structural and biological properties of the soils with different textures. The use of organic amendments found to increase the SOC concentration with larger increments at the higher application rate. This result is consistent with the findings of other researchers (e.g. Ferreras et al., 2006; González et al., 2010), who found increases in the final SOC content after the application of organic sources. It is apparent from the result that at the end of experiment, soils amended with AR showed higher SOC concentrations than those amended with MSW. Regarding the differences between chemical composition of amendments, the organic carbon content of AR (468 g kg^{-1}) was more than the organic carbon of MSW (394 g kg^{-1}), meanwhile the C : N ratio of AR-treated soils was more than the C : N ratio of MSW-treated soils (Table 2). In other words, depends on the chemical composition and C : N ratio of amendments, different amounts of organic carbon had been finally added to the soils.

On the other hand, an increase in the applications rate of amendments stimulated the soils microbial respiration, which is in agreement with what reported by others (e.g. Marinari et al., 2000; Ferreras et al., 2006; Yazdanpanah et al., 2013). The use of organic amendments has been reported previously to increase SOC and improve microbial activity (Marinari et al., 2000). As the soil microbial respiration was measured through CO_2 production, it is a direct indicator of microbial activity and indirectly reflects the availability of organic substrates (Ferreras et al., 2006). This investigation aimed to provide the relationship between microbial respiration and SOC of soils. Figure 1 shows the relationship between SOC concentration and CO_2 emission from loamy sand and clay loam soils treated with the two types of organic amendments. As is obvious, the soil microbial respiration increases significantly with increasing SOC as linear (Fig. 1a) and power (Fig. 1b) relationships for loamy sand and clay loam soils, respectively. Higher values of microbial respiration were found in plots amended with MSW compared to those treated with AR. Moreover, MSW compost caused to higher increases

SED

7, 3151–3177, 2015

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CO₂ emission and structural characteristics

N. Yazdanpanah

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in CO₂ emission as a function of SOC than did AR. This can be partly attributed to this fact that most of the carbon supplied by MSW comprises easily degradable material to be used as energy and nutrient source for soil microorganisms, resulting in the increased soil microbial respiration (Ferrerias et al., 2006; Mahmoodabadi and Heydarpour, 2014). It is well known that the microbial decomposition processes are controlled by the substrate quality (e.g. lignin and polyphenol content) and the availability of labile carbon (Koranda et al., 2013). In some studies, the C : N ratio has been attributed to the rate of decomposability of organic inputs, i.e. the lower ratio can be associated with the higher rate of carbon mineralization and CO₂ emission (Majumder et al., 2008). It can be assumed that compared to MSW, the application of AR with less easily degradable components may cause to the formation of more stable soil organic complexes, resulting in more resistant against the microbial decomposition (Majumder et al., 2008; Mahmoodabadi and Heydarpour, 2014). Therefore, AR shows slower microbial decomposition as well as lower mineralization rate (Liu et al., 2010; Cely et al., 2014). In other words, the different levels of organic carbon added to the soils are likely to be influenced by the biochemical composition and the decomposability of amendments (Yazdanpanah et al., 2013).

At the beginning of experiment, the clay loam soil used in this study had higher content of organic carbon than the loamy sand soil (Table 1). In spite of greater percentage of clay particles and much less sand particles in the clay loam soil, the mean weight diameter (MWD) of clay loam soil (0.27 mm) was more than that of the loamy sand soil (0.18 mm). When the soil microbial respiration was assessed in relation to control, MSW caused to higher increases in the CO₂ emission from clay loam soil than from loamy sand soil, whereas the reverse order was observed for AR. Alternatively, as compared to control, the increased SOC concentration in the loamy sand soil was higher than in the clay loam soil. This means that apart from soil texture, the aggregate size distribution plays an important role in the carbon stock and microbial activity. It can be assumed that the decomposition rate of organic matter can vary in soils among different aggregate size classes (Mangalassery et al., 2013). Mangalassery et al. (2013) in two

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



contrasting soil textures found that both texture and aggregate size significantly influenced CO₂ emission. Mahmoodabadi and Heydarpour (2014) found that CO₂ emission from a coarse-textured soil is relatively higher than a fine-textured soil. Sey et al. (2008) reported that higher CO₂ emitted from micro aggregates (< 0.25 mm) compared to macro aggregate (> 0.25 mm). In a clay loam soil, Drury et al. (2004) found a decrease in CO₂ production with increasing aggregate size. In contrast, Strong et al. (2004) found faster decomposition rate of carbon in a soil with relatively larger pore sizes. Overall, our result are conditioned by the chemical composition of amendment, the rate of application and the soil texture and aggregate size distribution, which have been supported by other studies (Tejada and González, 2003; Ferreras et al., 2006; Mahmoodabadi and Heydarpour, 2014).

The addition of organic amendments to the soils also improved the aggregate stability and consequently increased the soils total porosity especially macro pores fraction. The result indicated that the capability of different organic amendments in the improving soil structural stability depends on the dose of application, the rate of decomposability, the capacity of microbial respiration and the texture of soils, which corresponds to previous observations (Benbouali et al., 2013; Yazdanpanah and Mahmoodabadi, 2013). The added organic carbon in general is necessary for the flocculation of soil particles to form more stable aggregates. The incorporation of organic amendments into the soils increased the cohesion of aggregates (Ferreras et al., 2006), with more significant effect in plots amended with the higher rate (i.e. 30 Mgha⁻¹). Similarly, Bronick and Lal (2005) found parallel increases in SOC concentration and aggregate stability following the poultry manure application. In contrary, some researchers (e.g. Celik et al., 2004) reported that organic amendments increased the SOC concentration, but did not show any significant effect on the aggregate stability.

The higher aggregate stability observed in amended soils was concurrent with the improvement of total porosity and the remarkable increasing in macro pores fraction. Several authors have previously reported that the organic matter from amendment incorporation improved pore size distribution (Marinari et al., 2000; Tejada and González,

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2003). Figure 2 shows the relationship between the total porosity and microbial respi-
 ration of loamy sand and clay loam soils after the application of different levels of MSW
 and AR. For both the organic amendments used, good relationships were obtained, so
 that the CO₂ emission was stimulated significantly ($p < 0.01$) as power functions of the
 total porosity of soils. Similar result was reported by Marinari et al. (2000) who found
 positive linear correlations between soil porosity, microbial activity and CO₂ production
 in organic and mineral treatments. Aggelides and Londra (2000) demonstrated that the
 organic amendment application considerably improved soil physical properties through
 increasing the total porosity and changing the distribution of pore sizes in loamy and
 clay textured soils. In some studies, the effect was significant in micro pores fraction,
 as Celik et al. (2004) found that the organic treatments had positive effects on microp-
 orosity compared to control.

The concurrent improvement in the aggregate stability and soil porosity due to the
 amendments addition was more pronounced in the macro pores than in the micro
 pores fraction. In other words, the macro pores fraction was much more sensitive to
 the amendments application than the micro pores. This finding has been approved
 by Jarvis (2007) who characterized the macro pores by high temporal variability. The
 macro pores fraction in general contributes to ease the aeration of soil and conse-
 quently affects on the soil microbial respiration. Therefore, in the present study the ratio
 of macroporosity to microporosity as a soil structural indicator was related to the mi-
 crobial respiration. As can be seen in Fig. 3, there are significant relationships (power)
 between the ratio of macroporosity to microporosity and CO₂ production from the soils
 treated with different levels of MSW and AR.

Overall, an interrelationship was found between the soil microbial respiration and the
 structural characteristics such as aggregate stability and porosity fractions. In fact, the
 microbial respiration and SOC content of soils can be linked to the soils aggregate sta-
 bility and pore size distribution. The increased microbial respiration and SOC content
 in the amended soils can be contribute to the improvement of soil aggregate stability
 (Balashov et al., 2010). Mangalassery et al. (2013) found that CO₂ flux was affected

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arjmand Sajjadi, S. and Mahmoodabadi, M.: Aggregate breakdown and surface seal development influenced by rain intensity, slope gradient and soil particle size, *Solid Earth*, 6, 311–321, doi:10.5194/se-6-311-2015, 2015a.

Arjmand Sajjadi, S. and Mahmoodabadi, M.: Sediment concentration and hydraulic characteristics of rain-induced overland flows in arid land soils, *J. Soil. Sediment.*, 15, 710–721, doi:10.1007/s11368-015-1072-z, 2015b.

Balashov, E., Kren, J., and Prochazkova, B.: Influence of plant residue management on microbial properties and water-stable aggregates of two agricultural soils, *Int. Agrophys.*, 24, 9–13, 2010.

Benbouali, E. H., Hamoudi, S. A. E. A., and Larich, A.: Short- term effect of organic residue incorporation on soil aggregate stability along gradient in salinity in the lower cheliff plain (Algeria), *Afr. J. Agric. Res.*, 8, 2144–2152, 2013.

Bronick, C. J. and Lal, R.: Manuring and rotation effects on soil organic carbon concentration for different aggregate size fraction on two soils in northeastern Ohio, USA, *Soil Till. Res.*, 81, 239–252, 2005.

Bruun, T. B., Elberling, B., de Neergaard, A., and Magid, J.: Organic carbon dynamics in different soil types after conversion of forest to agriculture, *Land Degrad. Dev.*, 26, 272–283, doi:10.1002/ldr.2205, 2015.

Celik, I., Ortas, I., and Kilic, S.: Effects of compost, mycorrhiza, manure and fertilizer on some physical properties of a Chromoxerert soil, *Soil Till. Res.*, 78, 59–67, 2004.

Cely, P., Tarquis, A. M., Paz-Ferreiro, J., Méndez, A., and Gascó, G.: Factors driving the carbon mineralization priming effect in a sandy loam soil amended with different types of biochar, *Solid Earth*, 5, 585–594, doi:10.5194/se-5-585-2014, 2014.

Cerdà, A.: Soil aggregate stability under different Mediterranean vegetation types, *Catena*, 32, 73–86, 1998a.

Cerdà, A.: The influence of aspect and vegetation on seasonal changes in erosion under rainfall simulation on a clay soil in Spain, *Can. J. Soil Sci.*, 78, 321–330, doi:10.4141/S97-060, 1998b.

Cerdà, A.: Aggregate stability against water forces under different climates on agriculture land and scrubland in southern Bolivia, *Soil Till. Res.*, 36, 1–8, 2000.

Cerdà, A. and Doerr, S. H.: Influence of vegetation recovery on soil hydrology and erodibility following fire: an 11 year investigation, *Int. J. Wildland Fire*, 14, 423–437, doi:10.1071/WF05044, 2005.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cerdà, A., González-Pelayo, O., Giménez-Morera, A., Jordán, A., Pereira, P., Novara, A., Bre-
vik, E. C., Prosdocimi, M., Mahmoodabadi, M., Keesstra, S., García Orenes, F., and Rit-
sema, C.: The use of barley straw residues to avoid high erosion and runoff rates on persim-
mon plantations in Eastern Spain under low frequency – high magnitude simulated rainfall
5 events, *Soil Res.*, in press, 2015.

de Graaff, M. A., Adkins, J., Kardol, P., and Throop, H. L.: A meta-analysis of soil biodiversity
impacts on the carbon cycle, *Soil*, 1, 257–271, doi:10.5194/soil-1-257-2015, 2015.

Drury, C. F., Yang, X. M., Reynolds, W. D., and Tan, C. S.: Influence of crop rotation and ag-
gregate size on carbon dioxide production and denitrification, *Soil Till. Res.*, 79, 87–100,
10 2004.

Fernández, J. M., Plaza, C., García-Gil, J. C., and Polo, A.: Biochemical properties and barley
yield in a semiarid Mediterranean soil amended with two kinds of sewage sludge, *Appl. Soil
Ecol.*, 42, 18–24, 2009.

Ferreras, L., Gomez, E., Toresani, S., Firpo, I., and Rotondo, R.: Effect of organic amendments
15 on some physical, chemical and biological properties in a horticultural soil, *Bioresource
Technol.*, 97, 635–640, 2006.

García-Gil, J. C., Plaza, C., Soler-Rovira, P., and Polo, A.: Long-term effects of municipal
solid waste compost application on soil enzyme activities and microbial biomass, *Soil Biol.
Biochem.*, 32, 1907–1913, 2000.

González, M., Gomez, E., Comese, R., Quesada, M., and Conti, M.: Influence of organic
20 amendments on soil quality potential indicators in an urban horticultural system, *Bioresource
Technol.*, 101, 8897–8901, 2010.

Hueso-González, P., Martínez-Murillo, J. F., and Ruiz-Sinoga, J. D.: The impact of organic
amendments on forest soil properties under Mediterranean climatic conditions, *Land Degrad.
25 Dev.*, 25, 604–612, doi:10.1002/ldr.2296, 2014.

Jaiarree, S., Chidthaisong, A., Tangtham, N., Polprasert, C., Sarobol, E., and Tyler, S. C.: Car-
bon budget and sequestration potential in a sandy soil treated with compost, *Land Degrad.
25 Dev.*, 25, 120–129, doi:10.1002/ldr.1152, 2014.

Jarvis, N. J.: A review of non-equilibrium water flow and solute transport in soil macropores:
30 principles, controlling factors and consequences for water quality, *Eur. J. Soil Sci.*, 58, 523–
546, 2007.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jiménez, M. N., Fernández-Ondoño, E., Ripoll, M. Á., Castro-Rodríguez, J., Huntsinger, L., and Navarro, F. B.: Stones and organic mulches improve the *Quercus ilex* L. afforestation success under Mediterranean climatic conditions, *Land Degrad. Dev.*, doi:10.1002/ldr.2250, 2013.

5 Kaleem Abbasi, M., Mahmood Tahir, M., Sabir, N., and Khurshid, M.: Impact of the addition of different plant residues on nitrogen mineralization–immobilization turnover and carbon content of a soil incubated under laboratory conditions, *Solid Earth*, 6, 197–205, doi:10.5194/se-6-197-2015, 2015.

Karak, T., Bhagat, R. M., and Bhattacharyya, P.: Municipal solid waste generation, composition, and management: the world scenario, *Crit. Rev. Env. Sci. Tec.*, 42, 1509–1630, 2012.

10 Koranda, M., Kaiser, C., Fuchslueger, L., Kitzler, B., Sessitsch, A., Zechmeister-Boltenstern, S., and Richter, A.: Seasonal variation in functional properties of microbial communities in beech forest soil, *Soil Biol. Biochem.*, 60, 95–104, 2013.

Liu, E., Yan, C., Mei, X., He, W., Bing, S. H., Ding, L., Liu, S., and Fan, T.: Long-term effect of chemical fertilizer effects on soil organic matter fraction and microbes under a wheat-maize cropping system in northern China, *Geoderma*, 149, 318–324, 2010.

15 Mahmoodabadi, M. and Ahmadbeygi, B.: Dry and water-stable aggregates in different cultivation systems of arid region soils, *Arabian Journal of Geosciences*, 6, 2997–3002, doi:10.1007/s12517-012-0566-x, 2013.

Mahmoodabadi, M. and Arjmand Sajjadi, S.: Effects of rain intensity, slope gradient and particle size distribution on the relative contributions of splash and wash loads to rain-induced erosion, *Geomorphology*, 253, 159–167, doi:10.1016/j.geomorph.2015.10.010, 2016.

20 Mahmoodabadi, M. and Cerdà, A.: WEPP calibration for improved predictions of interrill erosion in semi-arid to arid environments, *Geoderma*, 204–205, 75–83, doi:10.1016/j.geoderma.2013.04.013, 2013.

25 Mahmoodabadi, M. and Heydarpour, E.: Sequestration of organic carbon influenced by the application of straw residue and farmyard manure in two different soils, *Int. Agrophys.*, 28, 169–176, doi:10.2478/intag-2014-0005, 2014.

Mahmoodabadi, M., Yazdanpanah, N., Rodríguez Sinobas, L., Pazira, E., and Neshat, A.: Reclamation of calcareous saline sodic soil with different amendments (I): redistribution of soluble cations within the soil profile, *Agr. Water Manage.*, 120, 30–38, doi:10.1016/j.agwat.2012.08.018, 2013.

30

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mahmoodabadi, M., Ghadiri, H., Rose, C., Yu, B., Rafahi, H., and Rouhipour, H.: Evaluation of GUEST and WEPP with a new approach for the determination of sediment transport capacity, *J. Hydrol.*, 513, 413–421, doi:10.1016/j.jhydrol.2014.03.060, 2014a.

Mahmoodabadi, M., Ghadiri, H., Yu, B., and Rose, C.: Morpho-dynamic quantification of flow-driven rill erosion parameters based on physical principles, *J. Hydrol.*, 514, 328–336, doi:10.1016/j.jhydrol.2014.04.041, 2014b.

Majumder, B., Mandal, B., Bandyopadhyay, P. K., Gangopadhyay, A., Man, P. K., Kuudu, A. L., and Mazumdar, D.: Organic amendments influence soil organic carbon pools and rice-wheat productivity, *Soil Sci. Soc. Am. J.*, 72, 775–685, 2008.

Mangalassery, S., Sjögersten, S., Sparkes, D. L., Sturrock, C. J., and Mooney, S. J.: The effect of soil aggregate size on pore structure and its consequence on emission of greenhouse gases, *Soil Till. Res.*, 132, 39–46, 2013.

Marinari, S., Masciandaro, G., Ceccanti, B., and Grego, S.: Influence of organic and mineral fertilisers on soil biological and physical properties, *Bioresource Technol.*, 72, 9–17, 2000.

Mazaheri, M. R. and Mahmoodabadi, M.: Study on infiltration rate based on primary particle size distribution data in arid and semiarid region soils, *Arabian Journal of Geosciences*, 5, 1039–1046, doi:10.1007/s12517-011-0497-y, 2012.

Musinguzi, P., Ebanyat, P., Tenywa, J. S., Basamba, T. A., Tenywa, M. M., and Mubiru, D.: Precision of farmer-based fertility ratings and soil organic carbon for crop production on a Ferralsol, *Solid Earth*, 6, 1063–1073, doi:10.5194/se-6-1063-2015, 2015.

Novara, A., Rühl, J., La Mantia, T., Gristina, L., La Bella, S., and Tuttolomondo, T.: Litter contribution to soil organic carbon in the processes of agriculture abandon, *Solid Earth*, 6, 425–432, doi:10.5194/se-6-425-2015, 2015.

Pansu, M. and Gautheyrou, J.: *Handbook of Soil Analysis, Mineralogical, Organic and Inorganic Methods*, Springer, Heidelberg, Germany, 993 pp., 2006.

Parras-Alcántara, L. and Lozano-García, B.: Conventional tillage versus organic farming in relation to soil organic carbon stock in olive groves in Mediterranean rangelands (southern Spain), *Solid Earth*, 5, 299–311, doi:10.5194/se-5-299-2014, 2014.

SAS Institute: *SAS/STAT Users Guide, Statistics Version, 6th edn.*, SAS Institute, Cary, NC, USA, 1990.

Sey, B. K., Manceur, A. M., Whalen, J. K., Gregorich, E. G., and Rochette, P.: Small-scale heterogeneity in carbon dioxide, nitrous oxide and methane production from aggregates of a cultivated sandy-loam soil, *Soil Biol. Biochem.*, 40, 2468–2473, 2008.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Sirjani, E. and Mahmoodabadi, M.: Effects of sheet flow rate and slope gradient on sediment load, *Arabian Journal of Geosciences*, 7, 203–210, doi:10.1007/s12517-012-0728-x, 2014.
- Six, J., Guggenberger, G., Paustian, K., Haumaier, L., Elliott, E. T., and Zech, W.: Sources and composition of soil organic matter fractions between and within soil aggregates, *Eur. J. Soil Sci.*, 52, 607–618, 2001.
- 5 Srivastava, P. K., Gupta, M., Shikha, N., Singh, N., and Tewari, S. K.: Amelioration of sodic soil for wheat cultivation using bioaugmented organic soil amendment, *Land Degrad. Dev.*, doi:10.1002/ldr.2292, 2014.
- Strong, D. T., De Wever, H., Merckx, R., and Recous, S.: Spatial location of carbon decomposition in the soil pore system, *Eur. J. Soil Sci.*, 55, 739–750, 2004.
- 10 Tejada, M. and González, J. L.: Effects of the application of a compost originating from crushed cotton gin residues on wheat yield under dryland conditions, *Eur. J. Agron.*, 19, 357–368, 2003.
- Turgut, B.: Comparison of wheat and safflower cultivation areas in terms of total carbon and some soil properties under semi-arid climate conditions, *Solid Earth*, 6, 719–725, doi:10.5194/se-6-719-2015, 2015.
- 15 Walkley, A. and Black, I. A.: An examination of the degtjareff method for determining soil organic matter, and proposed modification of the chromic acid titration method, *Soil Sci.*, 37, 29–38, 1934.
- 20 Yazdanpanah, N. and Mahmoodabadi, M.: Reclamation of calcareous saline- sodic soil using different amendments: Time changes of soluble cations in leachate, *Arabian Journal of Geosciences*, 6, 2519–2528, doi:10.1007/s12517-011-0505-2, 2013.
- Yazdanpanah, N., Pazira, E., Neshat, A., Mahmoodabadi, M., and Rodríguez Sinobas, L.: Reclamation of calcareous saline sodic soil with different amendments (II): Impact on nitrogen, phosphorous and potassium redistribution and on microbial respiration, *Agr. Water Manage.*, 120, 39–45, doi:10.1016/j.agwat.2012.08.017, 2013.
- 25 Yu, Y. and Jia, Z. Q.: Changes in soil organic carbon and nitrogen capacities of *Salix cheilophila* Schneid along a revegetation chronosequence in semi-arid degraded sandy land of the Gonghe Basin, Tibet Plateau, *Solid Earth*, 5, 1045–1054, doi:10.5194/se-5-1045-2014, 2014.
- 30

CO₂ emission and structural characteristics

N. Yazdanpanah

Table 1. Some physical and chemical properties of the soils with different textures.

Soil property	Loamy sand	Clay loam
Clay (< 0.002 mm) (%)	5.8	31.0
Silt (0.05–0.002 mm) (%)	10.0	40.8
Sand (2–0.05 mm) (%)	84.2	28.2
MWD ^a (mm)	0.18	0.27
Bulk density (Mg m ⁻³)	1.76	1.53
EC ^b (dS m ⁻¹)	0.28	2.45
pH	6.8	7.2
OC ^c (g kg ⁻¹)	1.33	2.9
CaCO ₃ (%)	16.2	21.5

^a MWD: mean weight diameter.^b EC: electrical conductivity.^c OC: organic carbon.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. The chemical composition of two types of amendments used in the experiment.

Amendment	OC ^a (g kg ⁻¹)	Total N ^b (g kg ⁻¹)	C : N	Ash (g kg ⁻¹)	EC (1 : 5) ^c (dS m ⁻¹)	pH (1 : 5)
Alfalfa residue	468	21	22.3	320	4.8	7.6
Municipal solid waste	394	29	13.6	540	4.0	6.9

^a OC: organic carbon.

^b Total N: total nitrogen (Kjeldahl).

^c EC: electrical conductivity.

CO₂ emission and structural characteristics

N. Yazdanpanah

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Mean values of the selected soil properties for each soil texture treated with different rates of organic amendments.

Soil texture	Treatment	OC ^a (g kg ⁻¹)	WSA ^b (%)	Total porosity (cm ³ cm ⁻³)	Micro porosity (cm ³ cm ⁻³)	Macro porosity (cm ³ cm ⁻³)	Respiration (μg CO ₂ g ⁻¹ Soil)
Loamy sand	C	1.32e	9.4d	0.32c	0.21b	0.11e	61.5e
	MSW 10	1.95d	15.2b	0.40b	0.22b	0.19c	122.9c
	MSW 30	3.46b	18.1a	0.47a	0.22b	0.25a	182.1a
	AR 10	2.27c	10.3cd	0.37b	0.21b	0.15d	93.7d
	AR 30	3.98a	11.6c	0.47a	0.25a	0.22b	154.1b
Clay loam	C	2.92e	44.4c	0.41d	0.28b	0.12e	78.6e
	MSW 10	3.31d	50.6b	0.50bc	0.29b	0.21b	163.3b
	MSW 30	4.29b	52.9a	0.59a	0.31b	0.28a	261.5a
	AR 10	3.88c	45.8c	0.44c	0.30b	0.14d	96.0d
	AR 30	4.96a	51.5ab	0.53b	0.35a	0.18c	151.4c

Means for treatments in the same soil texture followed by the same letter are not significantly different (Duncan $p < 0.05$), $n = 3$. C: control; MSW: municipal solid waste, AR: alfalfa residue. Suffixes 10 and 30 represent the application rates of 10 and

30 Mg ha⁻¹, respectively.

^a OC: organic carbon.

^b WSA: water stable aggregates.

CO₂ emission and structural characteristics

N. Yazdanpanah




Figure 1. Relationship between the soil organic carbon and microbial respiration of loamy sand (a) and clay loam (b) soils treated with different levels of municipal solid waste (MSW) and alfalfa residue (AR).




Figure 2. Relationship between the total porosity and soil microbial respiration of loamy sand (a) and clay loam (b) soils treated with different levels of municipal solid waste (MSW) and alfalfa residue (AR).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Relationship between the ratio of macroporosity to microporosity and soil microbial respiration of loamy sand **(a)** and clay loam **(b)** soils treated with different levels of municipal solid waste (MSW) and alfalfa residue (AR).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

