

**Delamination  
beneath Apennines**

I. Koulakov et al.

# Subduction or delamination beneath Apennines? Evidences from regional tomography

I. Koulakov<sup>1,2</sup>, A. Jakovlev<sup>1,2</sup>, I. Zabelina<sup>1,2</sup>, F. Roure<sup>3,4</sup>, S. Cloetingh<sup>4</sup>,  
S. El Khrepy<sup>5,6</sup>, and N. Al-Arifi<sup>6</sup>

<sup>1</sup>Trofimuk Institute of Petroleum Geology and Geophysics SB RAS, Prospekt Koptyuga, 3, 630090, Novosibirsk, Russian Federation

<sup>2</sup>Novosibirsk State University, Novosibirsk, Russia, Pirogova 2, 630090, Novosibirsk, Russia

<sup>3</sup>IFP-Energies Nouvelles, Rueil-Malmaison

<sup>4</sup>Tectonics Group, Utrecht University, the Netherlands

<sup>5</sup>King Saud University, Riyadh, Saudi Arabia, P.O. Box 2455, Riyadh 11451, Saudi Arabia

<sup>6</sup>National Research Institute of Astronomy and Geophysics, NRIAG, 11421, Helwan, Egypt

Received: 24 December 2014 – Accepted: 14 January 2015 – Published: 24 February 2015

Correspondence to: I. Koulakov (ivan.science@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

In this study we present a new regional tomography model of the upper mantle beneath Italy and surrounding areas derived from inversion of travel times of  $P$  and  $S$  waves from the updated ISC catalogue. Beneath Italy we identify a high-velocity anomaly which behaves as a long narrow “sausage” with a steeply dipping part down to a depth of 400 km and then expanding horizontally over approximately 400 km. Rather than to interpret it as a remnant of the former Tethyan oceanic slab, we consider that it is made up of the infra continental lithospheric mantle of Adria, which is progressively delaminated, whereas its overlying crust becomes progressively accreted into the Apenninic tectonic wedge.

## 1 Introduction

The Mediterranean region is located in the convergence zone between the African and European plates which is characterized by a very complex interaction of different tectonic regimes including subduction, collision, spreading and shear zones (e.g., Fac-cenna et al., 2004, 2014). In some parts of the region, such as in the Italian Peninsula and its surroundings, all of these regimes are confined to a limited area that results in very complex geological and tectonic structures (Fig. 1).

Today for instance, even the nature of the lithosphere and crust beneath the deep Ionian Basin remains debated, being either made up of a remnant of the former Mesozoic Tethys ocean or by the distal portions of the stretched African continental crust (Dercourt et al., 1985, 1993; Stampfli and Borel, 2004; Roure et al., 2012).

When going back in the past, the paleogeographic reconstructions of the various platform and basinal domains of Adria, with the Albanides, and Hellenides in the east, and the Apennines and Sicily in the west, are also increasingly ambiguous. Crucial in this respect is whether the Mesozoic Ionian allocthonous units of the Albanides and Hellenides, and coeval Mesozoic basinal series from the LagoNegro and Imerese units

SED

7, 859–884, 2015

## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the Southern Apennines and Sicily, are considered as lateral equivalents of the modern, still deep water Ionian basin (Roure et al., 2004, 2012).

Seismic tomography is one of the key tools which are used to resolve the mechanisms of deep tectonic processes. However, despite the large number of different tomography models, knowledge on the major tectonic processes is often ambiguous and contradictory. In this paper we present a new 3-D seismic model of the upper mantle beneath the Italian region constructed based on generally same calculation schemes as in Koulakov et al. (2009), but using a considerably larger dataset. Based on the distributions of  $P$  and  $S$  velocity anomalies, we propose a new interpretation for the recent tectonic history of the Italian region.

For a long time, the European part of Mediterranean has been an attractive region for seismic tomography studies thanks to relatively dense distribution of seismic stations, intensive seismicity and fairly heterogeneous deep structure controlled by complex geodynamic processes. In the early nineties, the regional mantle structure beneath Europe was studied by the use of travel times of body waves (Spakman, 1990, Spakman et al., 1993, Spakman and Wortel, 2004) and surface waves (Zielhuis and Nolet, 1994). The global models by Bijwaard et al. (1998) and Bijwaard and Spakman (2000) have provided the compatible resolution for the European region with the existing regional models. Later, Piromallo and Morelli (2003) presented another  $P$  velocity regional model for the entire European region based on body waves. More recently Koulakov et al. (2009) have constructed  $P$  and  $S$  tomography models for the upper mantle beneath Europe which took into account a newly obtained crustal model by Tesauro et al. e.g. (2008). In parallel there were several models of Europe created based on surface wave data (Boschi et al., 2004, 2009; Marone et al., 2004; Legendre et al., 2012). Adjoint tomography has been used for studying the European region by Zhu et al., (2012). Most of these models are generally consistent with each other, especially for the uppermost mantle. They identify highly contrasting features which detect general lithospheric structures known from geology. However, there are some differences between results obtained based on body and surface waves. Surface wave

SED

7, 859–884, 2015

## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tomography usually provides stronger amplitudes of anomalies and smoother lateral and sharper vertical variations.

The Italian Peninsula and surrounding regions have been studied using the regional data of the Italian permanent and temporary networks in many different studies. *P* wave velocity models of the crust and uppermost mantle based on travel times of body waves from regional events were the focus of many investigations, mostly performed by Italian scientists (e.g. Amato et al., 1993; Alessandrini et al., 1995; Selvaggi and Chiarabba, 1995; Di Stefano et al., 1999; Cimini and De Gori, 2001; Orecchio et al., 2011; Gualtieri et al., 2014).

Both regional and local scale tomography models provide valuable information on the deep processes which explains many observations on surface tectonics. Together with various geological and geophysical data, the tomography results are used for geodynamic interpretation and to reconstruct the scenarios of plate interactions in the European regions. One of the key regions is the Calabrian–Tyrrhenian system where a complex shape of the plate boundary passing through Calabria, Apennines, Alps and Dinarides is identified (Fig. 1). The existence of deep seismicity down to ~ 400 km beneath Calabria and the Tyrrhenian Sea and of the active volcanism of the Eolian Arc suggests that the subduction processes are active here (e.g., Isacks and Molnar, 1971; Selvaggi and Chiarabba, 1995). It was first proposed by Malinverno and Ryan (1986) that the loop shaped boundary, leading to back-arc extension in the Tyrrhenian sea (Spadini et al. 1995), was formed due to strongly curved subduction occurring in the narrow zone in front of Apennines. However, such a strong bending of the subducting plate appears to be mechanically not plausible, as was shown by analogue experiments by Faccenna et al. (1996). Alternatively, Faccenna et al. (2004) have proposed that the complex boundary shape in the Apennines region was formed due to the initial evolution of the western Mediterranean subduction zone (WMSZ) followed by trench retreat and back-arc extension. According to their model, the counterclockwise rotation of the Apennines block started at ~ 35 Ma due to the opening of the Ligurian-Provençal basin, as also proposed by Mattei et al., (2002), and then continued due to the spread-

## SED

7, 859–884, 2015

### Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing of the Tyrrhenian basin. However, the final stage, which resulted in the origin of the narrow subduction zone remains not completely clear. Spakman and Wortel (2004) have drawn attention to the segmented nature of the down going slabs in the western Mediterranean region whereas Govers and Wortel (2005) have pointed out that STEP faults are playing an important role in the dynamics of the underlying subduction dynamics. In this paper we present an alternative scenario based on new tomography models of  $P$  and  $S$  velocities. We also provide some additional testing results to show that derived seismic structures related to the Apennines region are robust.

**2 Data and tomography algorithm**

To construct the upper mantle velocity structure beneath central Mediterranean, we use  $P$  and  $S$  travel times from the ISC catalogue in a time period from 1964 to 2012. The calculations were performed using generally same computing schemes as in Koulakov et al. (2009) but with the use of a large data amount accumulated in the ISC catalogue since 2007 (the last year of data in the previous study). Here we use more than 11 million travel times ( $\sim 8.9 \times 10^6$  of  $P$  and  $2.3 \times 10^6$  of  $S$  picks) from more than 150 000 events (72 picks per event). In the previous study by Koulakov et al. (2009), the number of data was several times smaller.

Prior to use the ISC data for tomography, they were reprocessed using the location tools and outlier analysis developed by Koulakov and Sobolev (2006) which resulted in rejection of almost 30 % of the data. All data corresponding to seismic rays travelling inside the selected window, at least partly, are selected for the inversion. They include data from  $\sim 66$  000 earthquakes located in the study region (red dots in Fig. 2a) recorded by the worldwide stations (blue triangles in Fig. 2b), as well as data from  $\sim 90$  000 teleseismic events (grey dots in Fig. 2b) recorded by 1915 seismic stations located inside the study area (black triangles in Fig. 2a). The travel times of all seismic rays in the European region were corrected for the crustal model developed by Tesauro et al. (2008).

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The inversion is performed in three circular windows covering the study area having a diameter of 1500 km (Fig. 2a), which appears to be the most appropriate size to study the upper mantle, as estimated by Koulakov and Sobolev (2006). In this study we use the windows, which cover much larger area than the region of interest of this study, in order to avoid boundary effects that may appear at the margin of a circle due to smearing of outside anomalies. Such a window-by-window approach allows more optimal definition of inversion parameters depending on the data amount in each window, and it provides higher resolution than a global inversion for the entire region. After performing the inversions, the results in all windows are combined in a single model.

The algorithm of tomographic inversion was developed by Koulakov et al. (2002) and then significantly modified by Koulakov and Sobolev (2006). The parameterization of the velocity models is performed using the nodes installed according to the ray density on horizontal levels at depths of 10, 25, 50, 100, 150, 220, 290, 360, 430, 500, 570, 640, 710, 780, 850, 930, and 1000 km. The minimum grid spacing in horizontal levels is set at 30 km. No nodes are installed in areas where there is no data. To avoid an effect of the grid orientation upon the results, the inversions were performed in four grids with different basic orientations (0, 22, 45 and 67°) and then combined in a single model. The inversion was performed simultaneously for the  $P$  and  $S$  velocity anomalies, source parameters ( $dx$ ,  $dy$ ,  $dz$  and  $dt$ ) and station corrections. The matrix inversion was performed using the LSQR algorithm designed by Paige and Saunders (1982) and Nolet (1987). The same tomography code was used for studying various regions of the world including several subduction zones, such as the Kurile-Kamchatka and Aleutian (Koulakov et al., 2011), Mariana and Izu-Bonin arcs (Jaxybulatov et al., 2013), Taiwan (Koulakov et al., 2014). In all subduction zones, both  $P$  and  $S$  velocity models revealed consistent images of the slabs coinciding with the locations predicted by other studies and marked with deep seismicity.

### 3 Tomography results

The computed  $P$  and  $S$  velocity anomalies, which are considered as the main result of our calculations, are presented in horizontal and vertical sections in Figs. 3 and 4. The anomalies are given in percent in respect to the 1-D reference model AK 135 (Kennett et al., 1995). The model was computed down to 1000 km, while here we show only upper mantle structures down to 700 km.

One of the key parameters in inversion which strongly affects the solution is the value of damping which controls the smoothing of the resulting anomalies. Figure 5 presents an example of solution with weaker damping, as compared to the model in Figs. 3 and 4. It can be seen that this solution adds some smaller patterns and increase the amplitudes of anomalies, especially in the lower part of the model. Although this model appears to be stable, and the size of anomalies remains larger than those resolved in synthetic tests (see next paragraphs), we prefer staying at a conservative side and present a smoothed model as the main result of this study.

This model appears to be generally consistent with the results by Koulakov et al. (2009) and other tomography models. For example, vertical sections in Fig. 2.8a in (Spakman and Wortel, 2004) looks very similar to our result for the  $P$  velocity in vertical section A1-B1 having approximately same location.

Before discussing the observed seismic anomalies, we present a series of synthetic tests which were specially performed to assess the robustness of the obtained model in the studied region, which appears to be very important, regarding some inconsistencies with previously published models. First of all, we present the results of the traditional checkerboard test which consists of a reconstruction of periodic rectangular anomalies with the amplitude of  $\pm 4\%$  based on the existing data configurations. In the example presented in Fig. 6, we present results of two checkerboard models with different sizes of  $P$  and  $S$  patterns. In the upper two rows, the horizontal sizes of  $P$  and  $S$  anomalies are  $1^\circ$  by  $1^\circ$  and  $2^\circ$  by  $2^\circ$ , respectively. In the second case (lower two rows in Fig. 6), the sizes of anomalies are  $1^\circ$  by  $1^\circ$  and  $2^\circ$  by  $2^\circ$ , for  $P$  and  $S$  mod-

SED

7, 859–884, 2015

## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



els, respectively. In all cases the sign of anomalies changed with depth every 200 km (200, 400, 600 km etc). The checkerboard anomalies were set in the entire earth; thus the ray paths corresponding to remote stations or sources were affected by the outside anomalies. The data were perturbed with random noise of 0.3 and 0.6 s of mean magnitude. The results of the checkerboard reconstructions for the  $P$  and  $S$  models are presented in Fig. 6 in horizontal sections at 100, 300 and 500 km (middle depths of the checkerboard levels). For the  $P$  anomalies with  $1^\circ$  size patterns, the correct reconstruction occurs in selected areas with the maximum amount of data. In deeper sections, the resolution is much poorer compared to the result at 100 km depth. For the  $2^\circ$  size anomalies, the  $P$  model is reconstructed almost perfectly in the onshore areas and in the Adriatic Sea. In this test, the  $P$  anomalies almost do not lose their amplitude with depth. In the offshore areas, the anomalies are strongly smeared, which is related to lack of available data. The resolution of the  $S$  model appears to be poorer mostly because of a smaller amount of data. In addition, most of the  $S$  rays correspond to short rays traveling in the uppermost layers; for the deeper parts of the model, the  $S$  ray coverage is even poorer. We observe that in the case of  $2^\circ$  size of patterns, the anomalies below 200 km depth are smeared and lose their amplitudes. Nevertheless, their locations in space are correct, which is important for qualitative interpretation of the results.

The second test shown in Fig. 7 is aimed at studying the capacity of the algorithm to resolve a sausage-shaped anomaly similar to that we obtain in the observed data inversion. We define the synthetic anomalies along the Profile A1-B1, same as used for presenting the main model in Fig. 4, as free shaped polygonal horizontal prisms with a thickness of 200 km in the direction across the profile. We considered two anomalies corresponding to the Calabrian slab and to the Apennine collision belt. It can be seen that all these anomalies are robustly resolved in  $P$  and  $S$  models in correct depth intervals, although in the case of the  $S$  model, the amplitude of the resolved slab-related anomaly is much weaker compared to the true model. This test is especially important



this body in space, whereas the direction of the flat descending plate mostly follows the dipping attitude of the plate surface.

The deeper anomaly is seen in Section A1-B1 as a large high-velocity body located between 500 and 700 km depth. It is only observed in the  $P$  model; the  $S$  anomalies tend to be negative in this depth interval. We propose that this pattern represents the Alpine-Tethys remnant which subducted in previous stages of closing the paleocean (as proposed, for example, by Spakman and Wortel, 2004). If the time of stagnation was sufficiently large, no thermal anomaly, which affects the  $S$  velocity, is left, but there might be still compositional factors, which increase the  $P$  velocity.

The present configuration of anomalies beneath central Mediterranean originated from a complex series of different geodynamic episodes which occurred in Western Mediterranean in Cenozoic. According to the reconstruction proposed by Faccenna et al. (2004) shown in a simplified form in Fig. 8, the present configuration of the contact zone between African and European plates evolved from a classical subduction which occurred at about 35 Ma along the present coast of Spain and France (Fig. 8a). The back-arc processes caused the opening of the Ligurian-Provençal basin (Fig. 8b) and subsequently the Tyrrhenian basin (Fig. 8c) which resulted in counterclockwise rotation of the suture zone, which presently forms the Apennines, to approximately 90–100°. During this rotation, along this suture zone, an ocean-ocean type of subduction took place and resulted in arc volcanism which is still recorded by calc-alkaline material. At some moment, this subduction reached the margin of the Adriatic Shelf represented by a transition from oceanic to continental type of the lithosphere; the latter is more buoyant than the oceanic lithosphere. As a result, the oceanic lithosphere in the west was overriding the continental lithosphere of the Adriatic Shelf, and the subduction was transformed into a collision of continental type which caused shortening of the crust in both sides of the suture zone. This has led to active mounting building and formation of the Apennine crust. We assume that the subducted oceanic lithosphere from the Adriatic side was detached and sank. The observed in our tomography model high  $P$  velocity anomaly below 500 km depth might be the trace of the remnant lithosphere.

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the recent past, the continental crust of the western portion of the Adriatic plate became progressively accreted into the tectonic wedge, leading to the delamination and detachment of its lithospheric mantle which behaved as a subducting slab and remained below the Apennines (Roure et al., 2012), as shown in the scheme in Fig. 9.

We propose that the observed high-velocity sausage shaped anomaly beneath the Apennines represents the sunken part of the Adriatic lithosphere. Actually, its overall lateral and vertical lengths are quite compatible with the restored palinspastic surfaces of the Apenninic upper crustal and sedimentary units. Balanced cross-sections in the Southern Apennines account for instance for more than 200 km of shortening among the Apulian and Apenninic platforms and intervening Lago Negro basal units are currently accreted into the tectonic wedge (Casero et al., 1991; Roure et al., 1991, 2012).

In Fig. 9 we present our interpretation of the main structures below the Apennines in present time and give a hypothetical reconstruction to the recent past. Sections correspond to the location of the Profiles A1-B1 and A2-B2. In Section A1-B1 parallel to the Apennines, reference points are used for the interpretation: a is located in the continental part of Europe; B marks the suture zone in the Alps; C is a point in Po plain in Northern Italy; D marks the southernmost part of the Apennines, E marks the Calabrian arc, F is a point in the African Plate.

According to the GPS data (e.g., Hollenstein et al., 2003, Nocquet and Calais, 2003), in present days, there is no considerable convergence between Calabria and the African plate (points D and F). The present displacements of Tunisia, Sicily and Calabria are not significantly different. At the same time, most scientists accept the retreat of the Calabrian arc. Thus, the point E tends to be replaced from D to E (Fig. 9). Between the southern and northern Apennines (points D and C), the geodetic observations record considerable shortening. The same conclusion follows from the analysis of stress patterns based on focal mechanisms (Serpelloni et al., 2007) which indicate the extension in SW–NE direction and compression in NW–SE direction. Between the northern Apennines and continental Europe there is an obvious shortening which re-

sulted in the Plio-Quaternary overthrusting of the Northern Apennines towards the Po Plain, following pre-Messinian episodes of mountain building in the facing southwestern segment of the Alps (Roure et al., 1989, 1996; Turrini et al., 2014).

To explain the detachment of the Adriatic lithospheric mantle, the key process is the NW–SE shortening of Apennines between points C and D. The crust, which is a weaker part of the lithosphere, has been compressed and reduced its length. For the lithospheric mantle part, shortening was less plausible; instead it has been detached and formed a curved structure sank into the asthenosphere. In the tomography image we observe that the NW part of this body has been completely detached, whereas the SE end of it is still attached to the Calabrian arc. Vertical sinking of the “Calabrian sausage” causes the breaking of the oceanic lithosphere in the African side, and thus leads to the retreat of the contact zone between the newly formed Tyrrhenian Sea and the Ionian-African foreland lithosphere.

## 5 Conclusions

Based on an analysis of the  $P$  and  $S$  tomographic models of the upper mantle beneath the Central Mediterranean region derived from the tomography inversion of the ISC data, we present a new interpretation for the existence of the high-velocity anomaly beneath the Apennines, which was previously interpreted as a subduction zone of the former Tethyan oceanic lithosphere. We found that this anomaly behaves as a long narrow “sausage” with a steeply dipping part down to a depth of 400 km and then expanding horizontally over approximately 400 km. In cross section, this anomaly appears to be 150–200 km thick. We propose that this pattern represents the detached part of the Adriatic mantle lithosphere which was delaminated due to the final episode of the collision along the Apennines. The sinking of this segment was due to the negative buoyancy of the lithosphere material and was additionally triggered by NW–SE shortening of Apennines. The NW part of this “sausage” was detached, whereas the SE end of it is still connected to the Ionian-African foreland lithosphere.

## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



*Author contributions.* I. Koulakov, A. Jakovlev and I. Zabelina performed all tomographic calculations and figures preparation. I. Koulakov, S. El Khrepy and F. Roure provided geodynamical interpretation of presented results. I. Koulakov prepared manuscript with contributions of all co-authors.

5 *Acknowledgements.* Ivan Koulakov is supported by Integration Project of SB RAS #76, and Irina Zabelina is supported by RFBR Grant # 14-17-00430. The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding the work through the international research group project No IRG14-21. This study is based on freely available data catalogue provided by the International Seismological Center.

## 10 References

Alessandrini, B., Beranzoli, L., and Mele, F. M.: 3-D crustal  $P$  wave velocity tomography of the Italian region using local and regional seismicity data, *Ann. Geophys.-Italy*, 38, 189–211, 1995.

15 Amato, A., Alessandrini, B., Cimini, G., Frepoli, A., and Selvaggi, G.: Active and remnant subducted slabs beneath Italy: evidence from seismic tomography, *Ann. Geofis.*, 36, 201–214, 1993.

Bijwaard, H., Spakman, W., and Engdahl, E. R.: Closing the gap between regional and global travel time tomography, *J. Geophys. Res.-Sol. Ea.*, 103, 30055–30078, 1998.

Bijwaard, H., and Spakman, W.: Nonlinear global  $P$  wave tomography by iterated linearised inversion, *Geophys. J. Int.*, 141, 71–82, 2000.

20 Boschi, L., Ekström, G., and Kustowski, B.: Multiple resolution surface wave tomography: the Mediterranean basin, *Geophys. J. Int.*, 157, 293–304, 2004.

Boschi, L., Fry, B., Ekström, G., and Giardini, D.: The European upper mantle as seen by surface waves, *Surv. Geophys.*, 30, 463–501, 2009.

25 Casero, P., Roure, F., and Vially, R.: Tectonic framework and petroleum potential of the southern Apennines. Generation, accumulation, and production of Europe's hydrocarbons, *Sp. Pub. EAPG*, 1, 381–387, 1991.

Cimini, G. B. and De Gori, P.: Nonlinear  $P$  wave tomography of subducted lithosphere beneath central-southern Apennines (Italy), *Geophys. Res. Lett.*, 28, 4387–4390, 2001.

## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Cimini, G. B. and Marchetti, A.: Deep structure of peninsular Italy from seismic tomography and subcrustal seismicity, *Ann. Geophys.-Italy*, 49, 331–345, 2006.
- Dercourt, J., Ricou, L., Vrielynck, B., Atlas Tethys Palaeoenvironmental Maps, Gauthier-Villars, Paris, 14, 307, 1993
- 5 Dercourt, J., Zonenshain, L., Ricou, L., Kazmin, V., Le Pichon, X., Knipper, A., Grandjacquet, C., Sborshchikov, I., Boulin, J., and Sorokhtin, O.: Presentation de 9 cartes paleogeographiques au 1/20000000 000eme s' etendant de l'Atlantique au Pamir pour la periode du Lias a l'Actuel. *Bulletin De La Société Géologique De France*, 1, 637–652, 1985.
- 10 Di Stefano, R., Chiarabba, C., Lucente, F., and Amato, A.: Crustal and uppermost mantle structure in Italy from the inversion of *P* wave arrival times: geodynamic implications, *Geophys. J. Int.*, 139, 483–498, 1999.
- Faccenna, C., Davy, P., Brun, J., Funicello, R., Giardini, D., Mattei, M., and Nalpas, T.: The dynamics of back-arc extension: an experimental approach to the opening of the Tyrrhenian Sea, *Geophys. J. Int.*, 126, 781–795, 1996.
- 15 Faccenna, C., Piromallo, C., Crespo-Blanc, A., Jolivet, L., and Rossetti, F.: Lateral slab deformation and the origin of the western Mediterranean arcs, *Tectonics*, 23, TC1012, doi:10.1029/2002TC001488, 2004.
- Faccenna, C., Becker, T. W., Auer, L., Billi, A., Boschi, L., Brun, J. P., Capitanio, F. A., Funicello, F., Horvath, F., Jolivet, L., Piromallo, C., Leigh Royden, L., Rossetti, F., and Serpelloni, E.: Mantle dynamics in the Mediterranean, *Rev. Geophys.*, 52, 283–332, doi:10.1002/2013RG000444, 2014.
- 20 Govers, R. and Wortel, M.: Lithosphere tearing at STEP faults: response to edges of subduction zones, *Earth Planet. Sc. Lett.*, 236, 505–523, 2005.
- Gualtieri, L., Serretti, P., and Morelli, A.: Finite-difference *P* wave travel time seismic tomography of the crust and uppermost mantle in the Italian region, *Geochem. Geophys. Geosy.*, 15, 69–88, 2014.
- 25 Hollenstein, C., Khale, H. G., Geiger, A., Jenny, S., Goes, S., and Giardini, D.: New GPS constraints on the Africa-Eurasia plate boundary zone in southern Italy, *Geophys. Res. Lett.*, 30, 1935, doi:10.1029/2003GL017554, 2003.
- 30 Isacks, B. and Molnar, P.: Distribution of stresses in the descending lithosphere from a global survey of focal mechanism solutions of mantle earthquakes, *Rev. Geophys.*, 9, 103–174, 1971.

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jaxybulatov, K., Koulakov, I., and Dobretsov, N. L.: Segmentation of the Izu-Bonin and Mariana slabs based on the analysis of the Benioff seismicity distribution and regional tomography results, *Solid Earth*, 4, 59–73, doi:10.5194/se-4-59-2013, 2013.

Kennett, B., Engdahl, E., and Buland, R.: Constraints on seismic velocities in the Earth from traveltimes, *Geophys. J. Int.*, 122, 108–124, 1995.

Koulakov, I. and Sobolev, S. V.: Moho depth and three-dimensional P and S structure of the crust and uppermost mantle in the Eastern Mediterranean and Middle East derived from tomographic inversion of local ISC data, *Geophys. J. Int.*, 164, 218–235, 2006.

Koulakov, I., Tychkov, S., Bushenkova, N., and Vasilevsky, A.: Structure and dynamics of the upper mantle beneath the Alpine–Himalayan orogenic belt, from teleseismic tomography, *Tectonophysics*, 358, 77–96, 2002.

Koulakov, I., Kaban, M., Tesauro, M., and Cloetingh, S.: *P* and *S* velocity anomalies in the upper mantle beneath Europe from tomographic inversion of ISC data, *Geophys. J. Int.*, 179, 345–366, 2009.

Koulakov, I. Yu., Dobretsov, N., Bushenkova, N., and Yakovlev, A.: Slab shape in subduction zones beneath the Kurile–Kamchatka and Aleutian arcs based on regional tomography results, *Russ. Geol. Geophys.*, 52, 650–667, 2011.

Koulakov, I., Wu, Y., Huang, H., Dobretsov, N., Jakovlev, A., Zabelina, I., Jaxybulatov, K., and Chervov, V.: Slab interactions in the Taiwan region based on the *P* and *S* velocity distributions in the upper mantle, *J. Asian Earth Sci.*, 79, 53–64, 2014.

Legendre, C., Meier, T., Lebedev, S., Friederich, W., and Viereck-Götte, L.: A shear wave velocity model of the European upper mantle from automated inversion of seismic shear and surface waveforms, *Geophys. J. Int.*, 191, 282–304, 2012.

Loiselet, C., Braun, J., Husson, L., Le Carlier de Veslud, C., Thieulot, C., Yamato, P., and Grujic, D.: Subducting slabs: Jellyfishes in the Earth's mantle, *Geochem. Geophys. Geosy.*, 11, Q08016, doi:10.1029/2010GC003172, 2010.

Malinverno, A., Ryan, W. B.: Extension in the Tyrrhenian Sea and shortening in the Apennines as result of arc migration driven by sinking of the lithosphere, *Tectonics*, 5, 227–245, 1986.

Marone, F., Van Der Lee, S., and Giardini, D.: Three-dimensional upper-mantle *S* velocity model for the Eurasia–Africa plate boundary region, *Geophys. J. Int.*, 158, 109–130, 2004.

Mattei, M., Cipollari, P., Cosentino, D., Argentieri, A., Rossetti, F., Speranza, F., Di and Bella, L.: The Miocene tectono-sedimentary evolution of the southern Tyrrhenian Sea: stratigraphy,

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



structural and palaeomagnetic data from the on-shore Amantea basin (Calabrian Arc, Italy), Basin Res., 14, 147–168, 2002.

Nocquet, J. and Calais, E.: Crustal velocity field of western Europe from permanent GPS array solutions, 1996–2001, Geophys. J. Int., 154, 72–88, 2003.

5 Nolet, G.: Seismic wave propagation and seismic tomography, in: Anonymous Seismic tomography, Springer, 1–23, 1987.

Orecchio, B., Presti, D., Totaro, C., Guerra, I., and Neri, G.: Imaging the velocity structure of the Calabrian Arc region (southern Italy) through the integration of different seismological data, Bollettino Di Geofisica Teorica ed Applicata, 52, 625–638, 2011.

10 Paige, C. C. and Saunders, M. A.: LSQR: an algorithm for sparse linear equations and sparse least squares, ACM T. Math. Software, 8, 43–71, 1982.

Piromallo, C. and Morelli, A.: P wave tomography of the mantle under the Alpine-Mediterranean area, J. Geophys. Res.-Sol. Ea., 108, 2065, doi:10.1029/2002JB001757, 2003.

15 Roure, F., Polino, R., and Nicolich, R.: Poinçonnement, rétrocharrriages et chevauchements post-basculément dans les Alpes occidentales: Évolution intracontinentale d'une chaîne de collision, CR ACAD Sci. II B, 309, 283–290, 1989.

Roure, F., Casero, P., and Vially, R.: Growth processes and melange formation in the southern Apennines accretionary wedge, Earth Planet. Sc. Lett., 102, 395–412, 1991.

20 Roure, F., Choukroune, P., and Polino, R.: Deep seismic reflection data and new insights on the bulk geometry of mountain ranges, CR ACAD Sci. II A, 322, 345–359, 1996.

Roure, F., Nazaj, S., Mushka, K., Fili, I., Cadet, J., and Bonneau, M.: Kinematic evolution and petroleum systems, an appraisal of the outer Albanides, in: Thrust Tectonics and Hydrocarbon Systems, edited by: McClay, K. R., AAPG Mem. 82, Ch. 24, 474–493, 2004.

25 Roure, F., Casero, P., and Addoum, B.: Alpine inversion of the North African margin and delamination of its continental lithosphere, Tectonics, 31, TC3006, doi:10.1029/2011TC002989, 2012.

Selvaggi, G. and Chiarabba, C.: Seismicity and  $P$  wave velocity image of the southern Tyrrhenian subduction zone, Geophys. J. Int., 121, 818–826, 1995.

30 Serpelloni, E., Vannucci, G., Pondrelli, S., Argnani, A., Casula, G., Anzidei, M., Baldi, P., and Gasperini, P.: Kinematics of the western Africa-Eurasia plate boundary from focal mechanisms and GPS data, Geophys. J. Int., 169, 1180–1200, 2007.

Spadini, G., Cloetingh, S., and Bertotti, G.: Thermo-mechanical modeling of the Tyrrhenian Sea: lithospheric necking and kinematics of rifting, Tectonics, 14, 629–644, 1995.

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Spakman, W.: Tomographic images of the upper mantle below central Europe and the Mediterranean, *Terra Nova*, 2, 542–553, 1990.
- Spakman, W., van der Lee, S., van der Hilst, R.: Travel-time tomography of the European-Mediterranean mantle down to 1400 km, *Phys. Earth Planet. In.*, 79, 3–74, 1993.
- 5 Spakman, W. and Wortel, R.: A tomographic view on western Mediterranean geodynamics, in: anonymous *The TRANSMED atlas, The Mediterranean region from crust to mantle*, Springer, 31–52, 2004.
- Stampfli, G. M. and Borel, G. D.: The TRANSMED transects in space and time: constraints on the paleotectonic evolution of the Mediterranean domain, in: Anonymous *The TRANSMED Atlas. The Mediterranean Region from Crust to Mantle*, Springer, 53–80, 2004.
- 10 Tesauro, M., Kaban, M. K., and Cloetingh, S. A.: EuCRUST-07: A new reference model for the European crust, *Geophys. Res. Lett.*, 35, L05313, doi:10.1029/2007GL032244, 2008.
- Turrini, C., Lacombe, O., and Roure, F.: Present-day 3-D structural model of the Po Valley basin, Northern Italy, *Mar. Petrol. Geol.*, 56, 266–289, 2014.
- 15 Zhu, H., Bozdağ, E., Peter, D., and Tromp, J.: Structure of the European upper mantle revealed by adjoint tomography, *Nat. Geosci.*, 5, 493–498, 2012.
- Zielhuis, A. and Nolet, G.: Shea  $r$  wave velocity variations in the upper mantle beneath central Europe, *Geophys. J. Int.*, 117, 695–715, 1994.



**Figure 1.** Topography/bathymetry map of the Tyrrhenian Sea and Calabrian regions. Major structural units and sense of transport along main thrust fronts and extensional detachments are shown according to Faccenna et al. (2004). Red lines with arrows indicate thrust and subduction zones.

**Delamination  
beneath Apennines**


I. Koulakov et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Delamination  
beneath Apennines**

I. Koulakov et al.



**Figure 2.** Distribution of data used for tomography in the study region **(a)** and globally **(b)**. Circles indicate the target areas where the inversions were performed. Black and blue triangles denote stations inside and outside the target circles, respectively. Red and grey dots are the events inside and outside target circles, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Delamination  
beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Figure 3.** *P* and *S* velocity anomalies in horizontal sections.

## Delamination beneath Apennines

I. Koulakov et al.



**Figure 4.** *P* and *S* velocity anomalies in vertical sections. Locations of the profiles are shown in Fig. 3. Dots indicate projections of the earthquakes located at distances of less than 50 km from the profile. Exaggerated relief along the profiles is presented above each plot. Dotted line indicates the intersection with another profile.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Delamination beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 5.** Inversion results with the use of smaller damping. Two horizontal and one vertical sections are shown for the  $P$  and  $S$  models. Indications are the same as in Figs. 3 and 4.

Delamination  
beneath Apennines

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 6.** Checkerboard tests for two different  $P$  and  $S$  velocity models. In all cases, the signs of anomalies change at 200, 400, 600 km etc. Dotted lines mark boundaries of the synthetic anomalies.

**Delamination  
beneath Apennines**

I. Koulakov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 7.** Synthetic test with a synthetic model of realistic configuration. The shape of the synthetic model is highlighted with a contour. The thickness of the anomaly in the direction across the section is 200 km.



**Figure 8.** Simplified plate reconstruction in western Mediterranean based on Faccenna et al. (2004) with our modifications. Dark blue indicate areas of back arc spreading. Yellow is Adriatic plate with transitional ocean-to-continent structure. Dark brown color in Apennines and Alps highlights shortening areas. Indications: Lig – Ligurian Sea, CR – Corsica, SA – Sardinia, Tyr – Tyrrhenian Sea, Io – Ionian Sea, AD – Adriatic Sea.

**Delamination  
beneath Apennines**


I. Koulakov et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Delamination  
beneath Apennines

I. Koulakov et al.



**Figure 9.** Schematic representation of the origin of the Calabrian Sausage Background in plots with sections of present configuration corresponds to  $P$  velocity anomalies in vertical Sects. 1 and 2. Black arrow schematically indicate the displacements of blocks; red arrow denotes the direction of the Calabrian trench retreat. Abbreviations: Ap – Apennines; Io – Ionian Sea; Tyr – Tyrrhenian Sea; AD – Adriatic Sea; Sa – Sardinia; Lig – Ligurian Sea.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

