

1 **Evaluating management-induced soil salinization in golf courses in semi-arid**
2 **landscapes**

3 Joseph Young¹, Theophilus K. Udeigwe^{1*}, David C. Weindorf¹, Tarek Kandakji¹, Prativa
4 Gautam¹, and Mahmoud M.A. Mahmoud²

5 ¹Department of Plant and Soil Science, Texas Tech University, Lubbock, TX 79409, USA

6 ²Soil, Water & Environment Research Institute (SWERI), Agricultural Research Center (ARC),
7 Giza, Egypt

8

9

10

11

12

13

14

15

16

17

18

19

20 *Corresponding author: Theophilus K. Udeigwe

21 Address: Department of Plant and Soil Science, Texas Tech University

22 2907 15th Street, Lubbock, TX 79409, USA.

23 Tel: 1-806-834 6664

24 Fax: 1-806-742-0775

25 E-mail: theo.udeigwe@ttu.edu

26

Abstract

27
28 Site-specific information on land management practices are often desired to make better
29 ~~assertions~~ assessments of their environmental impacts. A study was conducted in Lubbock, TX,
30 in the Southern High Plains of the United States, an area characterized by semi-arid climatic
31 conditions, to (1) examine the potential management-induced alteration in soil salinity indicators
32 in golf course facilities, and (2) develop predictive relationships for a more rapid soil salinity
33 examination within these urban landscape soils using findings from portable x-ray fluorescence
34 (PXRF) spectrometer. Soil samples were collected from managed (well irrigated) and non-
35 managed (non-irrigated) areas of seven golf course facilities at 0–10, 10–20, and 20–30 cm
36 depths, and analyzed for a suite of chemical properties. Among the extractable cations, sodium
37 (Na) was significantly ($p < 0.05$) higher in the managed zones of all the golf facilities. Soil
38 electrical conductivity (EC), exchangeable sodium percentage (ESP), and sodium adsorption
39 ratio (SAR), parameters often used in characterizing soil salinity and sodicity, were in most part
40 significantly ($p < 0.05$) higher in the managed areas. Water quality report collected over a 22-
41 year period (1991–2013, all years not available) indicated a gradual increase in pH, EC, SAR,
42 total alkalinity, and extractable ions, thus, supporting the former findings. Findings from the
43 PXRF suggested possible differences in chemical species and sources that contribute to salinity
44 between the managed and non-managed zones. PXRF quantified Cl and S, and to a lesser extent
45 Ca, individually and collectively explained 23–85% of the variability associated with soil salinity
46 at these facilities.

47
48 **Keywords:** salinity, sodicity, Ogallala, drought, water quality, management practices, chemical
49 properties.

50 **1 Introduction**

51 Soil salinization is a global environmental problem that has gained a lot of research
52 attention over the years (Pitman and Läuchli, 2002; Martinez-Beltran and Manzur, 2005; Herrero
53 and Pérez-Coveta, 2005; Fan, 2012). Site-specific research on soil salinization are often needed
54 because generalization of findings could be misleading. The Southern High Plains (SHP) of the
55 United States (US), an area characterized by semi-arid climatic conditions (Peel et al., 2007), is
56 noted for complex environmental challenges such as drought, dust, wind erosion, soil
57 salinization, and nutrient deficiency. Nevertheless, in this region lie very important economic
58 cities, such as Lubbock that substantially contributes to US cotton production (USDA-NASS,
59 2014). Lubbock, located in the northwestern part of Texas, among other environmental
60 challenges is currently plagued by extreme water scarcity, attributed to low precipitation (a 30-
61 year average annual precipitation of approximately 470 mm) and the declining local aquifer, the
62 Ogallala. Recent observations have also shown an increasing pollutant concentration in well
63 waters (Scanlon et al., 2005), therefore, a concern over the water quality of the aquifer. Thus, the
64 intensification of agricultural and municipal activities could have a substantial impact on water
65 quantity and soil quality in this region.

66 Given the chemical properties of soils in the semi-arid and arid regions, which are
67 typified by high pH (>7.0) and ~~the relatively higher level~~limited leaching of soluble salts (IUSS
68 Working Group, 2006), poor management practices could ~~lead to~~induce soil salinization. In most
69 semi-arid and arid regions of the world, the unavailability of sufficient rainfall is often associated
70 with impaired soil quality as salts tend to accumulate in the soil as a result of limited leaching
71 (Pariante, 2001). This could result to soil salinization; a process where salts build up in the soil to
72 a potentially toxic level (Pitman and Lauchli, 2002; Rengasamy, 2006). Such altered chemical

73 properties could affect the soil hydraulic properties and its susceptibility to erosion (Morgan,
74 2009), environmental fate of soil pollutants (Du et al., 2009), and nutrient availability to
75 agronomic crops (Havlin et al., 2005). Poor quality irrigation water could also worsen such
76 scenarios as more contaminants from the water are continuously added; a typical case being the
77 declining Ogallala aquifer, which has been noted as a potential source of arsenic (As) and nitrate
78 (NO_3^-) to irrigated agricultural soils in the SHP (Hudak, 2000; Scanlon et al., 2005). Although a
79 common topic, but there are still very limited scientific reference materials on soil salinization in
80 agricultural and urban landscapes in the study area.

81 The first approach to addressing environmental degradations resulting from
82 contaminations is usually the identification of the major contributors. Evidently, in this region,
83 management (irrigation)-induced soil salinization has received less attention, particularly within
84 urban landscape facilities such as golf courses, despite its severity. Golf courses are major users
85 of irrigation water per unit area; a typical 18-hole golf facility in Southwest region could use an
86 average of approximately 1200 mmyr^{-1} of water (USGA, 2012) compared to 600 mmyr^{-1} for a
87 fully irrigated cotton in the same region (Snowden et al., 2013). Thus, in assessing the potential
88 impact of impaired water quality on soil and other environmental media in any setting, it will be
89 logical to examine the contributions of major irrigation water users in that given region of
90 interest. With the increasing severity of environmental degradation in the SHP region, it will be
91 of great interest to attempt to extend the applications of modern tools such as the portable x-ray
92 fluorescence (PXRF) for a more rapid investigation of environmental contamination, particularly
93 relating to soil salinization in golf course facilities. This tool is gaining importance in the fields
94 of soil and environmental sciences ([Kilbride et al., 2006](#); [Jang, 2010](#); McWhirt et al., 2012;
95 Gardner et al., 2013; Hu et al., 2014; Weindorf et al., 2014). Swanhart et al. (2015) demonstrated

96 the utility of using PXRF for soil salinity determination. This approach was further refined by
97 Aldabaa et al. (2015) who coupled PXRF data with visible near infrared diffuse reflectance
98 spectra as well as hyperspectral satellite data for improved measurement of salinity in playas of
99 West Texas, USA. This tool has also been extended to gypsum determination in arid soils
100 (Weindorf et al., 2013). The proposed study serves as an attempt to extend the application of the
101 PXRF to soil salinity examination in urban landscapes of the semi-arid climates. The main
102 advantage of this tool is its ability to quantify elements in an environmental medium such as soil
103 with minimal need for sample preparation, thereby, saving time and labor (Kalnicky and Singhvi,
104 2001) compared to the traditional wet chemistry techniques. Likewise, other tools for field-scale
105 salinity measurement such as electromagnetic induction, will not provide information on the
106 chemical species contributing or controlling salinity.

107 We hypothesize that there will be significant differences in key chemical properties
108 between managed and non-managed areas of golf course facilities. This was deduced from the
109 fact that in addition to the unique management practices of golf course facilities such as
110 perennial monoculture, less soil pulverization, and extended irrigation window, the managed
111 zones are frequently irrigated and would reflect the state of the irrigation water quality. Given the
112 semi-arid climatic condition of the study area and the characteristically alkaline nature of the
113 soils, these hypothesized differences could be more obvious in their salinity and/or sodicity
114 properties. Thus, the objective of this study were to (1) examine the possible management-
115 induced changes in soil chemical properties, particularly those significant to salinization, within
116 golf course facilities in a semi-arid climate, and (2) develop predictive relationships for a more
117 rapid soil salinity examination within these urban landscape soils using findings from PXRF.

118 **2 Materials and methods**

119 **2.1 Study site description**

120 This study was conducted in Lubbock, Texas, USA. Lubbock lies within 33° 34' N and
121 101° 53' W and sits on an elevation of 990 m.a.s.l. (USGS, 2014). This area is characterized by
122 semi-arid climatic conditions. Mean weather parameters recorded in year 2013 when soil
123 sampling was conducted were 320 mm (for precipitation), 61°F (ambient air temperature), 53%
124 (relative humidity), and 18.2 mph (wind speed) (NOAA, 2015). Geological materials are
125 composed mainly of Quaternary aeolian sand and loess (Nordstrom and Hotta, 2004). To achieve
126 our objectives, seven golf course facilities spread all over the city were selected for this study.
127 Each facility has been under management for at least 12 years. Figure 1 shows the locations of
128 the selected facilities, which are designated as A, B, C, D, E, F, and G. Using web soil survey,
129 soil types at the sites were broadly identified to belong to the Amarillo series (Fine-loamy,
130 mixed, superactive, thermic Aridic Paleustalfs) and Acuff series (Fine-loamy, mixed,
131 superactive, thermic Aridic Paleustolls). The average golf course contains 10 to 12 ha of irrigated
132 fairways. All managed fairways were planted with hybrid bermudagrass (*Cynodon dactylon* (L.)
133 Pers. x *C. transvaalensis* Burt-Davy) while the non-managed areas were composed of poorly
134 managed grass cover, native vegetation, or bare soil.

135 **2.2 Soil sampling and handling**

136 The fairways which are consistently irrigated were designated as the “managed areas”,
137 whereas adjacent areas of similar soil types that are not irrigated or managed were designated as
138 the “non-managed areas” in each facility. In each area, 3-4 core samples were randomly
139 collected using a 30 cm long x 6 cm wide (diameter) core sampler and then separated into three
140 depths of 0–10, 10–20, and 20–30 cm, then samples from same depth were combined to get a
141 representative sample. Soil sampling was conducted once within months of June and July in

142 2013. Sampling was conducted only once since the aim of the study was to evaluate the resultant
143 cumulative effect of many years (> 12 years) of management practices on soil chemical
144 properties of interest. Collected soil samples were then transported to the lab, air dried, ground
145 and passed through a 2 mm sieve.

146 **2.3 Soil characterization**

147 Soil samples were analyzed for a suite of chemical properties. Soil electrical conductivity
148 (EC) and pH were measured in a 1:2 solid (soil) to water suspension (Rhoades, 1996). Total
149 carbon (C) and nitrogen (N) were analyzed using a TruSpec C/N analyzer (LECO, St Joseph,
150 MI). Organic matter (OM) was determined using a modified Walkley and Black method (Nelson
151 and Sommers, 1996), using sodium (Na) dichromate and read on Gilford unit. Percent calcium
152 carbonate (CaCO_3) was determined by the tensimeter approach (Soil Survey Staff, 1996); a
153 modification of the pressure calcimeter approach (Loeppert and Suarez, 1996). Exchangeable
154 Na, calcium (Ca), magnesium (Mg) and potassium (K) were measured in ammonium acetate
155 extract (Soil Survey Staff, 2009) using atomic absorption spectrometer (AAS) (Spectra AA 220,
156 Varian, Palo Alto, California). Exchangeable sodium percentage (ESP) was calculated using
157 measured exchangeable cation values (Sparks, 2003). Sodium adsorption ratio (SAR) was
158 determined using the established relationship between ESR and SAR of saturated extract
159 developed by US Salinity Laboratory (Richards, 1954; Sparks, 2003). For the purpose and scope
160 of this study, water extractable chloride (Cl^-) and bicarbonate (HCO_3^-) were measured in 1:5
161 soil water extract and Cl^- concentration determined by titration with 0.005M silver nitrate
162 (AgNO_3) standard following Mohr titration approach (Soil Survey Staff, 1996), and HCO_3^- by
163 titration with 0.01M sulfuric acid (H_2SO_4) (Soil Survey Staff, 1996).

164 **2.4 PXRF scanning**

165 Collected samples were scanned using a PXRF (DP-6000 Delta Premium, Olympus,
166 Waltham, MA, USA) equipped with a Rh-X-ray tube which is operated at 10–40 kV with
167 integrated silicon drift detector (165 eV) (USEPA, 2007). The tool was operated in the Soil
168 Mode to measure a suite of elements, among which only Cl, K, S, and Ca were selected for our
169 purpose. Importantly, PXRF is not able to quantify Na, given its small, stable electron cloud. Soil
170 mode consist of three beams operating sequentially, each set to scan for 30 s for a total scan time
171 of 90 s per sample. Calibration of the instrument was conducted before sample analysis using a
172 316 alloy chip fitted to the aperture. Each soil sample was scanned in triplicate and the average
173 value reported. The data on elemental concentration and limit of detection (LOD) (three times
174 the standard error) were obtained and compiled.

175 **2.5 Water quality**

176 Water quality reports were obtained from the various golf course facilities, where
177 available. Since the facilities pump from the same groundwater source, the available reports were
178 enough to achieve the objectives of this study. In summary, 12 years (1991–2013, not all years
179 were included) of data were provided by one of the facilities, 2 years by another (2009–2010), 1
180 year each (2011 and 2013) by the remaining two facilities. The data sets broken down by water
181 sources were: well (12 years of data), retention pond (3 years) and recycled wastewater (1 year).
182 Water quality parameters reported include, EC, pH, SAR, Na, Mg, K, Ca, HCO_3^- , S in SO_4^{2-} ,
183 Cl^- , and total alkalinity.

184 **2.6 Statistical analyses**

185 All statistical analyses were performed using the Statistical Analysis Software (SAS 9.3,
186 SAS Institute, Cary, NC). Differences among means were examined using PROC GLM. Single
187 and multiple linear regression analyses using the stepwise technique were performed using the

188 PROC REG procedure to establish the relationships among parameters.

189 3 Results and discussions

190 3.1 Soil pH, CaCO₃, and OM

191 Soil pH, %CaCO₃ and %OM between the managed and non-managed areas of each golf
192 course facility [are summarized in Table 1](#). Salinity parameters will be discussed separately (see
193 Section 3.2). The results indicated little differences in mean pH (range= 7.7-8.8, mean= 8.25, $n =$
194 42) between managed and non-managed areas of all the courses examined (Table 1). The
195 differences in means between managed and non-managed areas at each facility ranged between
196 0.1-0.3 pH units and there was no consistent trend observed between the areas. However, these
197 differences were significant ($p < 0.05$) in three (B, C and F) of the seven facilities. Percent
198 CaCO₃ (range= 0.09–15.7 %, mean= 3.01, $n = 42$) showed no definite trend with depth and no
199 consistent differences between managed and non-managed areas (Table 1). Although not
200 significantly different, %CaCO₃ was higher in the non-managed zones of 4 (D, E, F, and G) of
201 the seven courses examined. Organic matter (range= 0.2–3.3 %, mean= 0.9, $n = 42$) tended to be
202 higher in the managed areas as was observed in six (A, B, C, D, F and G) of the seven sites,
203 although these were not statistically significant. The higher values observed in the managed
204 zones could be attributed to more biomass (Havlin et al., 2005) resulting from better
205 management. The exact same trend observed for soil OM also reflected in the soil TC as well as
206 TN which could be influenced by N fertilizer additions and N in irrigation water. Apart from
207 OM, TC, and TN, there was no consistent trend between managed and non-managed areas at
208 these set of facilities examined. The lack of significant differences between managed and non-
209 managed zones for most of the examined soil properties reported here somewhat indicates there
210 are no major external sources of these introduced through irrigation or other management

211 activities.

212 3.2 Extractable ions and salinity parameters

213 The differences in selected extractable ions and some salinity indicators between
214 managed and non-managed sites at each golf course [are summarized in Table 2](#). Among the
215 extractable cations (Ca, K, Mg, and Na), Na (range = 37–407mgkg⁻¹, mean = 67mgkg⁻¹, *n* = 42)
216 was significantly higher (*p* < 0.05) in the managed zone of each facility. This finding could
217 somewhat be attributed to the Na contained in the irrigation water originating mainly from
218 groundwater sources (see Section 3.3) because Na is not typically added through fertilization.
219 Exchangeable Ca (range= 1360–5477 mgkg⁻¹, mean= 2968 mgkg⁻¹, *n* = 42) was higher in the
220 non-managed zones of six (A to F) of the seven facilities, and this finding was significant (*p* <
221 0.05) at 3 of the facilities. This observed difference could be attributed to the possible leaching of
222 Ca (possibly in the form of sulfates and chlorides) from the more frequently irrigated areas.
223 Extractable Mg (range= 145–1381mgkg⁻¹, mean = 738mgkg⁻¹, *n* = 42) and K (range = 215–1491
224 mgkg⁻¹, mean = 587mgkg⁻¹, *n* = 42) were found to be higher in the irrigated areas of six of the
225 seven and five of the seven examined facilities, respectively, with significant differences (*p* <
226 0.05) observed in some facilities (Table 2). The higher levels of these elements in the managed
227 areas is likely due to their addition to the soil from irrigation water (see Section 3.3) because they
228 are not typically added through fertilization in this region. In general, the chloride salts of Ca are
229 more soluble than those of Mg and K, while the sulfate salts of Mg and K are more soluble than
230 those of Ca, and carbonate salts are generally insoluble (Clugston and Flemming, 2000). Thus,
231 using their solubility characteristics, it could be inferred that Na, Mg, and K in these soils could
232 be more of carbonate salts because they will be less soluble and thus mildly leached by irrigation
233 water. Conversely, Ca which tended to be more susceptible to leaching from these irrigated

234 zones could be predominantly in the form of chloride salts of Ca. The slight positive relationship
235 ($R^2 = 0.65$, $p < 0.05$, $n = 42$) observed between Na and Cl^- could suggest the presence of
236 chloride salts of Na as well.

237 The water extractable anions examined revealed that HCO_3^- (range = 77.6–326 mg kg^{-1} ,
238 mean = 170, $n = 42$) and Cl^- (range = 0–604 mg kg^{-1} , $n = 42$) were mostly higher in the managed
239 areas compared to the non-managed areas, some of which were significantly different (Table 2).
240 The only exception was HCO_3^- in facility F. The higher levels of these anions in the managed
241 zones of these facilities could be attributed to their addition to the soil from irrigation water
242 sources. The dominant anions in the soil solution of most semi-arid salty soils are Cl^- , SO_4^{2-} ,
243 HCO_3^- (at pH values of 6.0–8.0) and some NO_3^- (Dierickx, 2013). Thus, significant increases in
244 these ions could reflect a shift toward soil salinization. In this study, less emphasis was placed on
245 soil SO_4^{2-} , NO_3^- , and PO_4^{3-} concentrations because these are commonly added through
246 fertilization and thus, possible contribution from irrigation sources would not be easily
247 quantified.

248 The potential contribution of the management practices to salinity and sodicity could be
249 evidenced from the examination of the soil EC (range = 0.15–2.28 dSm^{-1} , mean = 0.643, $n = 42$),
250 ESP (range = 0.80–7.10 %, mean = 3.20 %, $n = 42$) and SAR (range = 1.40–6.04, mean = 3.12, n
251 = 42) values. It is apparent that the practices at the facilities and possibly irrigation water tended
252 to increase the salinity and sodicity properties of these soils. This is supported by the
253 significantly higher EC, ESP and SAR values generally observed in the managed areas of these
254 facilities (Table 2). A comparison was made among depths to examine the distribution of EC,
255 ESP, and SAR between all managed and non-managed sites (Fig. 2). When all the managed
256 zones were grouped and compared against the non-managed zone, at each depth, the salinity

257 parameters were significantly higher in the managed zones, suggesting the effects were similar
258 within all the depths examined. Besides irrigation, this shift toward salinization is further
259 supported by the semi-arid condition of the study site, characterized by low rainfall and less
260 leaching of the soluble salts, leading to their build up in the top soil.

261 3.3 Influence of local aquifer water quality

262 The water quality reports obtained from the facilities are summarized in Table 3. Of
263 interest, the concentration of each parameter examined (with the exception of pH) was on the
264 average approximately 2-folds-times higher in the well water compared to the retention pond,
265 which is mainly a collection of runoff and rain water (Table 3). These differences could be most
266 likely attributed to the inherently low pollutant concentration in rain water, filtration of pollutants
267 as it flows over vegetation on its way to the pond, and further settling of pollutants and uptake by
268 vegetation in the reservoir. The concentrations of the examined parameters in the effluent treated
269 water were 2–11 folds-times higher than those of the well water. Using the water quality
270 information, pollutant addition to soil from the water sources could be estimated. For instance,
271 using the average values of contaminants in the well water, approximately 5.60 g Cl^- , 7.60 g
272 SO_4^{-2} , 9.0 g HCO_3^- and 3.80 g Na^+ will be added to 1.0 kg of the receiving soil over a 10-year
273 period if a field receives approximately 1200 ~~mm~~mm~~year~~year⁻¹ of irrigation water from well sources
274 in this area. The limited rainfall and thus minimal leaching of salts in the semi-arid and arid areas
275 could make the situation described above more realistic.

276 The well water quality, which is a better representation of that of the local aquifer was
277 further examined. The available data were grouped into three sets: 1991–1993, 2004–2008, and
278 2009–2013, and the average values for each parameter in a set calculated. A striking feature
279 observed was the gradual increase in pH, EC, SAR, total alkalinity, Na, K, HCO_3^- and Cl^- over

280 the years (Fig. 3), suggesting that the declining aquifer (Terrel [and Johnson, 1999](#); [Terrel et al.,](#)
281 2002) could be associated with an increase in contaminant concentration, particularly salts.
282 Using the mean values, the $\text{Na}^+:\text{Ca}^{+2}$ ratio of approximately 2:1 in the well water sources
283 ~~justifies~~ likely explains the higher SAR and ESP in the managed areas that are irrigated using
284 water from well sources. This ratio is higher than those of recycled wastewater (1.5:1) and ditch
285 water (1:1) reported by Qian and Mecham (2005) that still ~~impacted~~ led to higher SAR in soils
286 after years of irrigation in Denver and Fort Collins, Colorado. Thus, our findings suggest that
287 continuous irrigation with well water could increase the salt contents of the receiving soils
288 overtime, a situation that is already apparent in the managed zones of the facilities examined in
289 this study as discussed under Section 3.2. The water quality data and the observed differences in
290 salinity parameters between managed (irrigated) and non-managed (non-irrigated) areas establish
291 a possible influence of the aquifer water quality on soil quality at these facilities.

292 **3.4 Application of PXRF to salinity prediction**

293 The PXRF quantified Ca, Cl, K, and S were individually and collectively used to explain
294 the variability associated with salinity, approximated using EC. The findings are presented in
295 Table 4. As evidenced from the R^2 values, when all the sites were considered ($n = 42$),
296 approximately 70% of the variability associated with salinity was explained by the Cl alone, 82%
297 by Cl and S, and 85% by Cl, S, and K. The findings here suggest the likely contributions of salts
298 of Cl^- and SO_4^{-2} to soil salinity within these facilities. When studied individually, the strengths
299 of these relationships were notably higher within the managed area ($n = 21$) compared to the
300 non-managed area ($n = 21$) (Table 4) as evidenced from R^2 average of 0.72 (managed) vs. 0.29
301 (non managed) for all the relationships examined. The stronger relationships observed within the
302 managed group support part of our hypothesis that salinity (EC) is influenced by anthropogenic

303 sources, thus, the more the input of these cation and anion-forming elements through irrigation,
304 fertilization, etc., the higher the salinity. The weaker relationships observed within the non-
305 managed group suggest a limited influence of the anthropogenic sources of the elements
306 (particularly Cl and S), further suggesting that salinity could be controlled by other parameters
307 that were not accounted for by the PXRF. From this information, it can be inferred that the
308 chemistry of salinity, i.e., the elemental species contributing to it, could be different within the
309 managed and the non-managed groups. This is an important piece of information that was
310 rapidly obtained using the PXRF. Overall, the relationships developed when all data points
311 (managed and non managed) were collectively considered suggest that the PXRF could be used
312 for rapid ~~in situ~~ prediction ~~of~~ and examination of chemistry of salinity in the semi-arid urban
313 soils, an application that could be extended to other semi-arid regions. Although, it is important
314 to note that the capability of this tool is still limited since the contributions of some elements
315 such as Na, and anions such as HCO_3^- and CO_3^{-2} cannot be ascertained yet.

316 **4 Conclusions**

317 The impacts of management practices on environmental quality could vary with climate
318 and thus, site specific investigations are often desired because extending findings from one
319 practice and location to others could be misleading. Thus, this study, serves as an initial probe
320 into the potential management-induced changes in soil chemical properties with a focus on
321 salinity in golf courses in Lubbock, TX, located in the SHP of the USA. This is an area
322 characterized by semi-arid climatic conditions, typified by drought, wind erosion, and potential
323 for soil salinization, ~~etc.~~ Evaluation of soil chemical properties of managed (irrigated) and non-
324 managed (non-irrigated) areas at seven different golf course facilities and information on well
325 water quality revealed possible differences in soil properties. The major findings are summarized

326 as thus: (1) among the exchangeable cations, Na was significantly higher in all the managed and
327 well irrigated zones of all the seven golf courses ~~in the semi-arid region of the USA~~, suggesting
328 the addition of Na salts (possibly in the forms of carbonates and chlorides) to irrigated soils from
329 irrigation water sources. (2) Irrigation tended to increase the salinity and sodicity properties of
330 the soils as evidenced from the significantly higher Soil EC, ESP, and SAR observed in majority
331 of the managed areas compared to the non-managed areas. This finding was supported by the
332 water quality data of the local aquifer which showed an increase in pH, EC, SAR, total alkalinity,
333 and extractable ions over the years. (3) PXRF quantified Cl and S, and to a lesser extent Ca,
334 individually and collectively explained most of the variability associated with salinity within the
335 soils of these facilities. The strengths of the relationships were generally higher in the managed
336 area.

337 Although in the SHP and other semi-arid and arid regions, the emphasis is more on water
338 quantity; however, it is important to point out that salt build up can affect water ~~quantity~~ quality,
339 by altering the hydrological properties of soils such as hydraulic conductivity, infiltration,
340 permeability, water holding capacity and thus water availability to crops. This study was an
341 initial investigation into an observed environmental issue and findings will support future
342 research effort in the subject area.

343 **Author contributions**

344 J. Young and T. K. Udeigwe planned and implemented the study. T. Kandakji and P. Gautam
345 assisted in field and laboratory activities. M. A. Mahmoud suggested ideas and assisted in
346 various aspect of the project as needed. D.C. Weindorf provided assistance with the portable x-
347 ray fluorescence spectrometer.

348

349 **References**

- 350 Aldabaa, A. A. A., Weindorf, D. C., Chakraborty, S., Sharma, A., and Li, B.: Combination of
351 proximal and remote sensing methods for rapid soil salinity quantification, *Geoderma*,
352 239–240, 34–46, 2015.
- 353 Clugston, M. and Flemming, R.: *Advanced Chemistry*, Oxford University Press, Oxford, UK,
354 2000.
- 355 Dierickx, W.: The salinity and alkalinity status of arid and semi-arid lands, *Land Use, Land*
356 *Cover and Soil Sciences V*, 163–189, 2013.
- 357 Du Laing, G., Rinklebe, J., Vandecasteele, B., Meers, E., and Tack, F. M. G.: Trace metal
358 behaviour in estuarine and riverine floodplain soils and sediments: a review, *Sci. Total*
359 *Environ.*, 407, 3972–3985, 2009.
- 360 Fan, X., Pedroli, B., Liu, G., Liu, Q., Liu, H., and Shu, L.: Soil salinity development in the
361 yellow river delta in relation to groundwater dynamics, *Land Degrad. Dev.*, 23, 175–189,
362 2012.
- 363 Gardner, D., Weindorf, D. C., and Flynn, M.: Presence of chromium, copper, and arsenic in
364 schoolyard soils, *Soil Horizons*, 54, 1–5, doi:10.2136/sh12-12-0032, 2013.
- 365 Havlin, J. L., Beaton, J. D., Tisdale, S. L., and Nelson, W. L.: *Soil fertility and fertilizers: an*
366 *introduction to nutrient management*, Vol. 515, Pearson Prentice Hall, Upper Saddle
367 River, NJ, 2005.
- 368 Herrero, J. and Pérez-Coveta, O.: Soil salinity changes over 24 years in a Mediterranean irrigated
369 district, *Geoderma*, 125, 287–308, 2005.
- 370 Hu, W., Huang, B., Weindorf, D. C., and Chen, Y.: Metals analysis of agricultural soils via
371 portable X-ray fluorescence spectrometry, *B. Environ. Contam. Tox.*, 92, 420–426, 2014.

372 Hudak, P. F.: Distribution and sources of arsenic in the southern high plains aquifer, Texas,
373 USA, *J. Environ. Sci. Heal. A*, 35, 899–913, 2000.

374 IUSS Working Group WRB: World Reference Base for Soil Resources, 2nd edn., World Soil
375 Resources Reports No. 103, FAO, Rome, ISBN 92-5-105511-4, 2006.

376 [Jang, M.: Application of portable X-ray fluorescence \(pXRF\) for heavy metal analysis of soils in](#)
377 [crop fields near abandoned mine sites, *Environ. Geochem. Hlth*, 32\(3\), 207-216, 2010.](#)

378 [Kilbride, C., Poole, J. and Hutchings, T. R.: A comparison of Cu, Pb, As, Cd, Zn, Fe, Ni and Mn](#)
379 [determined by acid extraction/ICP–OES and ex situ field portable X-ray fluorescence](#)
380 [analyses, *Environ. Pollut*, 143\(1\), 16-23, 2006.](#)

381 Loeppert, R. H. and Suarez, D. L.: [Carbonate](#) and gypsum, in: *Methods of Soil Analysis, Part 3,*
382 *Chemical Methods*, edited by: Bartels, J. M. and Bigham, J. M., Soil Science Society of
383 America, Inc., Madison, USA, SSSA Book Series 5, 437–474, 1996.

384 Martinez-Beltran, J. and Manzur, C. L.: Overview of salinity problems in the world and FAO
385 strategies to address the problem, in: *Proceedings of the international salinity forum,*
386 *Riverside, California*, 311–313, 2005.

387 McWhirt, A., Weindorf, D. C., and Zhu, Y.: Rapid analysis of elemental concentrations in
388 compost via portable x-ray fluorescence spectrometry, *Compost Sci. Util.*, 20, 185–193,
389 2012.

390 Morgan, R. P. C.: *Soil Erosion and Conservation*, Blackwell Publishing, Oxford, UK, 2009.

391 Nelson, D. W. and Sommers, L. E.: Total carbon, organic carbon, and organic matter, in:
392 *Methods of Soil Analysis, Part 3, Chemical Methods*, edited by: Sparks D. L., Soil
393 Science Society of America Book Ser. 5, SSSA and ASA, Madison, Wis., 961–1010,
394 1996.

395 [Sarah, P.: Soluble salts dynamics in the soil under different climatic regions. *Catena*. 43\(4\). 307-](#)
396 [321, 2001.](#)

397 Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger
398 climate classification, *Hydrol. Earth Syst. Sci.*, 11, 1633–1644, doi:10.5194/hess-11-
399 1633-2007, 2007.

400 Pitman, M. G. and Läuchli, A.: Global impact of salinity and agricultural ecosystems, in:
401 Salinity: Environment-Plants-Molecules, Springer Netherlands, Part A, 3–20,
402 doi:10.1007/0-306-48155-3_1, 2002.

403 Qian, Y. L. and Mecham, B.: Long-term effects of recycled wastewater irrigation on soil
404 chemical properties on golf course fairways, *Agron. J.*, 97, 717–721, 2005.

405 Rengasamy, P.: World salinization with emphasis on Australia, *J. Exp. Bot.*, 57, 1017–1023,
406 2006.

407 Rhoades, J. D.: Salinity: electrical conductivity and total dissolved solids, in: *Methods of Soil*
408 *Analysis, III, Chemical Methods*, edited by: Sparks, D. L., SSSA, Madison, WI, 417–
409 435, 1996.

410 Richards, L. A.: *Diagnosis and Improvement of Saline and Sodic Soils*, USDA Agric. Handb.60,
411 USDA, Washington, DC, 1954.

412 Scanlon, B. R., Nicot, J. P., Reedy, R. C., Tachovsky, J. A., Nance, S. H., Smyth, R. C., Keese,
413 K., Ashburn, R. E., and Christian, L.: Evaluation of arsenic contamination in Texas, The
414 Univ. of Texas at Austin, Bureau of Economic Geology, final report prepared for Texas
415 Commission on Environmental Quality, under umbrella contract no. 582-4-56385 and
416 work order no. UT-08-5-70828, 177, p. 167, 2005.

417 Snowden, C., Ritchie, G., Cave, J., Keeling, W., and Rajan, N.: Multiple irrigation levels affect
418 boll distribution, yield, and fiber micronaire in cotton, *Agron. J.*, 105, 1536–1544, 2013.

419 Soil Survey Staff: Chemical analyses, calcium carbonate (4E) HCl treatment (4E1) manometer,
420 electronic (4E1) < 20mm basis (4E), in: *Soil Survey Laboratory Manual, USDA-NRCS*
421 *SSRI #42, Version 4.0, 269–273, 1996.*

422 Soil Survey Staff: *Soil Survey Field and Laboratory Methods Manual, Soil Survey Investigations*
423 *Report No. 51, Version 1.0, edited by: Burt, R., U.S. Department of Agriculture, Natural*
424 *Resources Conservation Service, 2009.*

425 Sparks, D. L.: *Environmental Soil Chemistry, Academic Press, San Diego, California, 2003.*

426 Swanhart, S., Weindorf, D. C., Chakraborty, S., Bakr, N., Zhu, Y., Nelson, C., Shook, K., and
427 Acree, A.: Measuring soil salinity via portable x-ray fluorescence spectrometry, *Soil Sci.*,
428 in press, 2015.

429 Terrell, B. L. and Johnson, P. N.: Economic impact of the depletion of the Ogallala aquifer: a
430 case study of the southern high plains of Texas, in: *American Agricultural Economics*
431 *Association annual meeting, Nashville, TN, 8–11 August 1999.*

432 Terrell, B. L., Johnson, P. N., and Segarra, E.: Ogallala aquifer depletion: economic impact on
433 the Texas high plains, *Water Policy*, 4, 33–46, 2002.

434 U.S. Environmental Protection Agency: *Method 6200: Field Portable X-Ray Fluorescence*
435 *Spectrometry for the Determination of Elemental Concentrations in Soil and Sediment,*
436 *USEPA, 2007.*

437 USDA-NASS: Cotton ginning 2013 summary, available at: [http://usda.mannlib.cornell.edu/
438 usda/current/CottGinnSu/CottGinnSu-05-09-2014.pdf](http://usda.mannlib.cornell.edu/usda/current/CottGinnSu/CottGinnSu-05-09-2014.pdf) (last access: 9 July 2013), 2014

439 USGA: How Much Water Does Golf Use and Where Does It Come From?, available
440 at: [http://www.usga.org/uploadedFiles/USGAHome/Course_Care/Golf_and_the_Environment/ Water/214418%20Lyman,%20Greg%20-%20How%20Much%20Water%20Does%20 Golf%20Use.pdf](http://www.usga.org/uploadedFiles/USGAHome/Course_Care/Golf_and_the_Environment/Water/214418%20Lyman,%20Greg%20-%20How%20Much%20Water%20Does%20Golf%20Use.pdf) (last access: 11 April
441
442 2014), 2012.
443
444 USGS: National Elevation Dataset, available at: <http://ned.usgs.gov/> (last access: 28 March
445 2014), 2014.
446 [NOAA: NOAA Online Weather Data \(Lubbock Area\), available: http://www.weather.gov/](http://www.weather.gov/climate/xmacis.php?wfo=lub)
447 [climate/xmacis.php?wfo=lub](http://www.weather.gov/climate/xmacis.php?wfo=lub), accessed Feb. 22, 2015.
448 Weindorf, D. C., Zhu, Y., Chakraborty, S., Bakr, N., and Huang, B.: Use of portable X-ray
449 fluorescence spectrometry for environmental quality assessment of peri-urban
450 agriculture, *Environ. Monit. Assess.*, 184, 217–227, 2012.
451 Weindorf, D. C., Herrero, J., Castañeda, C., Bakr, N., and Swanhart, S.: Direct soil gypsum
452 quantification via portable x-ray fluorescence spectrometry, *Soil Sci. Soc. Am. J.*, 77,
453 2071–2077, 2013.
454 Weindorf, D. C., Bakr, N., and Zhu, Y.: Advances in portable x-ray fluorescence (PXRF) for
455 environmental, pedological, and agronomic applications, *Adv. Agron.*, 128, 1–45, 2014.

456
457
458
459
460
461

462

463 **Table 1:** Selected soil properties examined at the managed and non-managed areas of the seven
 464 **golf facilities in Lubbock, TX, USA**

<u>Golf Course</u>	<u>Management</u>	<u>Irrigation Source</u>	<u>pH</u>	<u>CaCO₃</u>	<u>TC</u>	<u>TN</u>	<u>OM</u>
				-----%-----			
<u>A</u>	<u>Managed</u>	<u>Well</u>	<u>8.2a</u>	<u>1.1a</u>	<u>1.88a</u>	<u>0.19a</u>	<u>1.5a</u>
	<u>Non-managed</u>		<u>8.1a</u>	<u>0.2b</u>	<u>0.77a</u>	<u>0.07a</u>	<u>0.7a</u>
<u>B</u>	<u>Managed</u>	<u>Well</u>	<u>8.3b</u>	<u>8.8a</u>	<u>2.31a</u>	<u>0.14a</u>	<u>1.0a</u>
	<u>Non-managed</u>		<u>8.4a</u>	<u>4.5a</u>	<u>1.20b</u>	<u>0.07a</u>	<u>0.5a</u>
<u>C</u>	<u>Managed</u>	<u>Well</u>	<u>8.2b</u>	<u>1.9a</u>	<u>1.68a</u>	<u>0.13a</u>	<u>1.2a</u>
	<u>Non-managed</u>		<u>8.5a</u>	<u>1.2a</u>	<u>0.89a</u>	<u>0.07a</u>	<u>0.5a</u>
<u>D</u>	<u>Managed</u>	<u>Well</u>	<u>8.6a</u>	<u>0.5a</u>	<u>0.87a</u>	<u>0.08a</u>	<u>0.8a</u>
	<u>Non-managed</u>		<u>8.6a</u>	<u>0.7a</u>	<u>0.47a</u>	<u>0.03a</u>	<u>0.3a</u>
<u>E</u>	<u>Managed</u>	<u>Well</u>	<u>8.2a</u>	<u>4.6a</u>	<u>2.24a</u>	<u>0.18a</u>	<u>1.1a</u>
	<u>Non-managed</u>		<u>7.9a</u>	<u>6.4a</u>	<u>2.96a</u>	<u>0.22a</u>	<u>1.3a</u>
<u>F</u>	<u>Managed</u>	<u>Well & RW</u>	<u>8.0a</u>	<u>0.7a</u>	<u>1.91a</u>	<u>0.18a</u>	<u>1.3a</u>
	<u>Non-managed</u>		<u>8.1b</u>	<u>1.6a</u>	<u>1.57a</u>	<u>0.13a</u>	<u>0.9a</u>
<u>G</u>	<u>Managed</u>	<u>Well & RP</u>	<u>8.1a</u>	<u>4.2a</u>	<u>2.86a</u>	<u>0.21a</u>	<u>1.5a</u>
	<u>Non-managed</u>		<u>8.3a</u>	<u>4.7a</u>	<u>1.48a</u>	<u>0.06a</u>	<u>0.7a</u>

TC, Total Carbon; TN, Total Nitrogen; OM, Organic Matter; RW, Recycled Wastewater; RP, Retention Pond. Mean values in a column within a golf course with the same letter are not statistically different (Fisher's LSD, $\alpha = 0.05$)

465

466

467

468

469

470 **Table 2:** A summary of extractable ions and soil salinity parameters for the managed and non-managed areas of the seven golf
471 facilities in the study in Lubbock, TX, USA
472

Golf Course	Management	Irrigation Source	NH ₄ -Acetate Extractable				Water Extractable		EC μs/cm	ESP %	SAR
			Na ⁺	Ca ²⁺	Mg ²⁺	K ⁺	HCO ₃ ⁻	Cl ⁻			
			-----mg kg ⁻¹ -----				-----mg kg ⁻¹ -----				
A	Managed	Well	271a	2165a	810a	534a	253a	5.9	445a	5.8a	5.0a
	Non-managed		42.0b	2259a	160b	321b	90.3b	nd	199b	1.3b	1.8b
B	Managed	Well	322a	2757b	1058a	633a	170a	307.7	1561a	5.4a	4.7a
	Non-managed		47b	3684a	569b	386b	125a	nd	417b	0.8b	1.4b
C	Managed	Well	309a	2355a	1109a	600a	186a	236.7	1187a	5.7a	5.1a
	Non-managed		68b	2786a	806b	520a	125a	nd	219b	1.3b	1.8b
D	Managed	Well	132a	1610b	657a	380a	160a	88.8	426a	3.9a	3.6a
	Non-managed		65.2b	2328a	293b	253a	125a	nd	221b	1.9b	2.2b
E	Managed	Well	264a	2732b	826a	441b	192a	88.8a	815a	5.1a	4.5a
	Non-managed		107b	5134a	912a	888a	176a	71.0a	699a	1.3b	1.7b
F	Managed	Well & RW	255a	2428a	776a	381b	189a	166a	991a	5.4a	4.7a
	Non-managed		114b	3038a	667a	786a	214a	76.9b	605a	2.1b	2.3b
G	Managed	Well & RP	270a	4401a	1140a	1272a	230a	59.2	810a	3.3a	3.2a
	Non-managed		78b	3872a	551b	822b	144a	nd	409b	1.3b	1.7b

EC, Electrical Conductivity; ESP, Exchangeable Sodium Percentage; RW, Recycled Wastewater; RP, Retention Pond; SAR, Sodium Adsorption Ratio (estimated using calculated Exchangeable Sodium Ratio); nd, not detected. Mean values in a column within a golf course with the same letter are not statistically different (Fisher's LSD, $\alpha = 0.05$).

479 **Table 4:** Regression equation and coefficients for the relationships between electrical
 480 conductivity (EC) and the selected PXRF quantified elements within managed and non-managed
 481 facilities of the golf courses in Lubbock, TX, USA

All facilities (n = 42)		
Parameter (s)	Equation	R ²
Cl	EC = 0.0015Cl + 0.2476	0.70 ^c
S	EC = 0.0007S - 0.5716	0.63 ^c
Ca	EC = 0.00001Ca + 0.3813	0.23
K	EC = 0.0341K + 160.28	0.06
Cl + S	EC = 0.001Cl + 0.0004S - 0.3063	0.82 ^c
Cl + K	EC = 0.0015Cl + 0.00004K - 0.2876	0.77 ^c
Cl + Ca	EC = 0.0014Cl + 0.000006Ca + 0.1490	0.75 ^c
Cl + S + K	EC = 0.0012Cl + 0.00003S + 0.00003K + -0.5931	0.85 ^c
Cl + S + Ca	EC = 0.0010Cl + 0.0004S + 0.000003Ca - 0.3004	0.83 ^c
Cl + Ca + K	EC = 0.0014Cl + 0.000003Ca + 0.00003K - 0.2085	0.78 ^c
Managed (n = 21)		
Parameter (s)	Equation	R ²
Cl	EC = 0.0017Cl + 0.1987	0.85 ^c
S	EC = 0.0007S - 0.4108	0.52 ^b
Ca	EC = 0.00002Ca + 0.5547	0.43 ^b
K	EC = 0.0444K + 275.83	0.09
Cl + S	EC = 0.0014Cl + 0.0002S - 0.1399	0.89 ^c
Cl + K	EC = 0.0017Cl + 0.1987K - 0.1459	0.88 ^c
Cl + Ca	EC = 0.0015Cl + 0.000005Ca + 0.1796	0.87 ^c
Cl + S + K	EC = 0.0014Cl + 0.0002S + 0.00002K - 0.4151	0.91 ^c
Cl + S + Ca	EC = 0.0013Cl + 0.0002S + 0.000004Ca - 0.1250	0.91 ^c
Cl + Ca + K	EC = 0.0015Cl + 0.0000003Ca + 0.00002K - 0.0718	0.89 ^c
Non managed (n = 21)		
Parameter (s)	Equation	R ²
Cl	EC = 0.0003Cl + 0.3598	0.03
S	EC = 0.0005S - 0.3238	0.39 ^b
Ca	EC = 0.000005Ca + 0.2884	0.10
K	EC = 0.0392K - 171.05	0.26
Cl + S	EC = 0.00035Cl + 0.000516S - 0.3803	0.43 ^b
Cl + K	EC = 0.0004Cl + 0.000004K - 0.2820	0.33 ^a
Cl + Ca	EC = 0.00037Cl + 0.000006Ca + 0.2321	0.15
Cl + S + K	EC = 0.00042Cl + 0.0004S + 0.00002K - 0.5226	0.49 ^b
Cl + S + Ca	EC = 0.0003Cl + 0.0005S - 0.0000007Ca - 0.3946	0.43 ^b
Cl + Ca + K	EC = 0.0005Cl + 0.000000008Ca + 0.00004K - 0.2804	0.33

^a = Significant at 0.05 probability level; ^b = Significant at 0.01 probability level;

^c = Significant at 0.001 probability level; EC in dS m⁻¹; Cl, S, Ca, and K in mg kg⁻¹

482

483 **Figure Captions**

484 **Figure 1:** Map showing the study area located in Lubbock, Texas, USA and the locations of the
485 seven golf facilities. The facilities are designated as A, B, C, D, E, F, and G.

486 **Figure 2:** Differences in selected soil chemical properties examined within depths between
487 managed and non-managed zones of all seven golf courses examined in Lubbock, TX, USA.
488 Each data point represents the average of seven points. Mean values within a soil depth with the
489 same letter are not statistically different (Fisher's LSD, $\alpha = 0.05$).


490 EC, Electrical Conductivity; ESP, Exchangeable Sodium Percentage; SAR, Sodium Adsorption
491 Ratio.

492 **Figure 3:** The observed trend in selected water quality parameters from 1991-2013. _____
493 Water samples were obtained from well sources utilized by a golf course in Lubbock, TX, USA.
494 For each parameter, data were averaged over 1991-1993 (n = 9), 2004-2008 (n = 6) and 2009-
495 2013 (n = 6), error bars are for the standard deviations. Mean values within a parameter with the
496 same letter are not statistically different (Fisher's LSD, $\alpha = 0.05$).

497
498
499
500
501
502
503
504
505
506



Figure 1:



1
2
3

Figure 2:




Figure 3:

17

18

