

1 **Effect of soil coarseness on soil base cations and available**
2 **micronutrients in a semi-arid sandy grassland**

3 Linyou Lü^{1,2}, Ruzhen Wang^{1,*}, Heyong Liu^{1,3}, Jinfei Yin^{1,4}, Jiangtao Xiao^{1,4},
4 Zhengwen Wang¹, Yan Zhao², Guoqing Yu², Xingguo Han¹, Yong Jiang¹

5 ¹ State Key Laboratory of Forest and Soil Ecology, Institute of Applied Ecology,
6 Chinese Academy of Sciences, Shenyang 110016, China

7 ² Institute of Sandyland Improvement and Utilization, Liaoning Academy of
8 Agricultural Sciences, Fuxin 123000, China

9 ³ Key Laboratory of Regional Environment and Eco-remediation, College of
10 Environment, Shenyang University, Shenyang 110044, China

11 ⁴ University of Chinese Academy of Sciences, Beijing 10049, China

12 * Corresponding author: Tel.: +86 24 83970603; fax: +86 24 83970300.

13 E-mail address: ruzhenwang@iae.ac.cn (Ruzhen Wang)

14

15 **Abstract**

16 Soil coarseness is the main process decreasing soil organic matter and threatening the
17 productivity of sandy grasslands. Previous studies demonstrated negative effect of soil
18 coarseness on soil carbon storage, but less is known about how soil base cations
19 (exchangeable Ca, Mg, K, and Na) and available micronutrients (available Fe, Mn, Cu,
20 and Zn) response to soil coarseness. In a semi-arid grassland of northern China, a field
21 experiment was initiated in 2011 to mimic the effect of soil coarseness on soil base
22 cations and available micronutrients by mixing soil with different mass proportions of
23 sand: 0% coarse elements (C0), 10% (C10), 30% (C30), 50% (C50), and 70% (C70).
24 Soil coarseness significantly increased soil pH in three soil depths of 0-10 cm, 10-20
25 cm and 20-40 cm with the highest pH values detected in C50 and C70 treatments. Soil
26 fine particles (smaller than 0.25 mm) significantly decreased with the degree of soil
27 coarseness. Exchangeable Ca and Mg concentrations significantly decreased with soil
28 coarseness degree by up to 29.8% (in C70) and 47.5% (in C70), respectively, across
29 three soil depths. Soil available Fe, Mn and Cu significantly decreased with soil
30 coarseness degree by 62.5%, 45.4% and 44.4%, respectively. As affected by soil
31 coarseness, the increase of soil pH, decrease of soil fine particles (including clay), and
32 decline in soil organic matter were the main driving factors for the decrease of
33 exchangeable base cations (except K) and available micronutrients (except Zn)
34 through soil profile. Developed under soil coarseness, the loss and redistribution of
35 base cations and available micronutrients along soil depths might pose threat to
36 ecosystem productivity of this sandy grassland.

37 *Key words:* soil degradation, dryland ecosystem, sandy soil, soil texture, Calcium,

38 Manganese

39

40 1 Introduction

41 Dryland ecosystems, accounting for 41% of the total land area of the world, are prone
42 to desertification which would result in soil coarseness (Cerdàs et al., 2014; Wang et al.,
43 2015a). Dryland ecosystems represent 25% of land surface area in Latin America with
44 75% of them having desertification problems (Torres et al., 2015). Desertified land
45 area has been reported to reach 45.6 million km² (Torres et al., 2015) and accounted
46 for 74% of total dryland area (61.5 million km²) with more than 100 countries and
47 8.5 × 10⁸ people being affected (Miao et al., 2015; Vieira et al., 2015; Wang et al.,
48 2015a). Desertification exerts large impact on social and economic resources (Beyene,
49 2015; Escadafal et al., 2015). Areas susceptible to desertification tend to be marked by
50 socioeconomic inequality and have low human development index in Brazil (Vieira et
51 al., 2015). Desertification was reported to cause economic losses of up to €6 billion in
52 Northern China in the year of 2005 (Miao et al., 2015).

53 In China, most of the grasslands have undergone degradation and desertification
54 with 50% distributed in the agro-pastoral transition zone of northern China (Wang et
55 al., 2015a; Yan and Cai, 2015). Continuous grazing and intense cultivation can reduce
56 vegetation cover and litter accumulation which exposes the ground surface to wind
57 erosion in the erosion-prone sandy lands (Su et al., 2005). Desertification and wind
58 erosion processes in fragile arid and semi-arid rangelands have contributed to
59 increased soil coarseness (Yan and Cai, 2015). Together with reduction of plant cover,
60 increased soil coarseness contribute to loss of agricultural productivity, environmental
61 deterioration, and associated social and economic disruptions (Vieira et al., 2015; Xie

62 et al., 2015). In this case, it is urgent to combat desertification and study the causes,
63 processes, consequences, and mechanisms of soil coarseness (Xu et al., 2012;
64 Weinzierl et al., 2015).

65 Soil base cations are not only essential nutrient cations for both plants and soil
66 microbes, but also serve as one of the main mechanisms of soil acid buffering
67 capacity (Lu et al., 2014) ~~as well as~~ and are a good indicator of soil fertility (Zhang et
68 al., 2013). Micronutrient availabilities essentially affect terrestrial net primary
69 production, plant quality, and consequently food and forage supply worldwide (Cheng
70 et al., 2010; Marques et al., 2015). Current research about desertification and soil
71 coarseness mainly focus on its effects on degradation of forest and grasslands due to
72 logging and overgrazing (Conte et al., 1999; Cao et al., 2008), C and N depletion in
73 soils and plant components (Zhou et al., 2008; Bisaro et al., 2014), soil compaction
74 and erosion risk (Allington and Valone, 2010), and soil physical properties of particle
75 size distributions (Su et al., 2004; Huang et al., 2007). However, less is known about
76 the changes in soil base cations and availabilities of micronutrient during dryland
77 desertification and soil coarseness.

78 Soil coarseness is suggested to cause a decrease of soil silt and clay contents
79 (Zhou et al., 2008), ~~decline in~~ decrease of soil C and nutrient (such as N and P)
80 concentrations (Xie et al., 2015), and losses in species diversity and productivity
81 (Zhao et al., 2006; Huang et al., 2007). As biogeochemical cyclings of base cations
82 and micronutrients are largely controlled by soil organic matter (SOM) (complexation
83 and chelation) (Sharma et al., 2004) and properties of soil mineral (reversible sorption

84 and desorption processes) (Jobb gy et al., 2004), decrease of SOM and soil fine
85 particles would potentially decrease soil base cations and micronutrient availability.
86 Also, the changes in SOM along soil depth could shape the vertical distribution of
87 base cations and available micronutrients (Sharma et al., 2004).

88 The Horqin Sandy Land, or Horqin Sandy Grassland is an important part of Inner
89 Mongolia grassland and one of the main sandy areas in northern China covering
90 approximately 43,000 km² (Li et al., 2004). With windy and dry winters and springs
91 in Horqin region, the soils are prone to aeolian soil erosion and soil coarseness
92 especially when natural sandy grassland is converted into farmland (Li et al., 2004).
93 To examine the effect of soil coarseness during desertification on the concentrations
94 of soil base cation (exchangeable Ca, Mg, K, and Na) and available micronutrient (Fe,
95 Mn, Cu, and Zn) of this region, we set up a field experiment in Zhanggutai by mixing
96 the soil with different mass proportions of sand: 10% (light soil coarseness), 30%
97 (moderate soil coarseness), 50% (heavy soil coarseness), and 70% (severe soil
98 coarseness). We hypothesized that both soil base cations and available micronutrients
99 would decrease with the increasing degree of soil coarseness due to the decrease of
100 SOM and soil fine particles. We also expected that soil base cations and available
101 micronutrients would decrease with soil depth.

102

103 **2 Materials and methods**

104 **2.1 Study area**

105 The study was conducted at the Desertified Grassland Restoration Research Station

106 maintained by Institute of Sand Fixation and Utilization, Liaoning Academy of
107 Agricultural Sciences. The study site (42°43'N and 122°22'E, elevation 226.5 m a.s.l.)
108 was located in the southeast of Horqin Sandy Land, near Zhanggutai Town, Zhangwu
109 County, Liaoning Province, China (Fig. 1). Productive grasslands from Zhanggutai
110 County have undergone severe desertification due to intense cultivation, overgrazing
111 and increased population (Li et al., 2000; Chen et al., 2005). The soils are susceptible
112 to wind erosion as a result of decrease in plant cover and high annual wind velocity
113 (varies from 3.4 to 4.1 m s⁻¹) with frequent occurrence of gales (wind speed > 20 m s⁻¹)
114 (Li et al., 2000). The mean annual temperature is 6.2 °C and mean annual
115 precipitation is about 450 mm which defines the area as semi-arid~~being a semi-arid-~~
116 ~~region~~ (Chen et al., 2005). The frost-free period lasts approximately 150 days (Chen
117 et al., 2005). Soil texture of the experiment site is sandy soil with 99.32 ± 0.13 % sand,
118 0.45 ± 0.14 % silt, and 0.23 ± 0.02 % clay (means ± standard deviation, data
119 measured from control soil). The soil type is classified as a *Aeolic Eutric Arenosol*
120 according to the FAO classification (IUSS Working Group WRB, 2014). Arenosols
121 are mainly developed in sand dune areas which are featured by a sandy texture and
122 low soil organic carbon (SOC) concentrations and prone to soil coarsening (Barthold
123 et al., 2013). This area constitutes an agro-pastoral ecotone which is severely
124 degraded due to excessive cultivation and grazing (Chen et al., 2005).

125

126 **2.2 Experimental design**

127 In May 2011, a complete randomized design was applied to the site. Within a 24 m ×

128 29 m area (696 m²), thirty 4 m × 4 m plots were established for five treatments with
129 six replicates per treatment. The soil within this experimental area is homogenous.
130 Adjacent plots were separated by 1 m buffer zone and PVC plates to prevent water
131 and nutrient exchanges. A certain mass proportion of 2 mm- sieved river sand
132 (siliceous, pH 7.5 ± 0.2) was mixed with native soil for each of three depths (0-20 cm,
133 20-40 cm, and 40-60 cm). Three soil depths were considered in this study to study
134 effect of soil coarseness on soil properties of plant root layer (0-20 cm), transition
135 layer of plant roots (20-40 cm), and transition layer of soil genesis (40-60 cm). To mix
136 the river sand and native soils evenly in each plot, soils of each depth were dug out
137 and mixed with the sand by agitators in the same mass proportion separately. The soils
138 were refilled back to the field in respective depths after mixture. The proportions are 0,
139 10%, 30%, 50%, and 70% to mimic different soil coarseness degrees or intensities:
140 control grassland without soil coarseness (C0), light soil coarseness (C10), moderate
141 soil coarseness (C30), heavy soil coarseness (C50), and severe soil coarseness (C70),
142 respectively. In August 2012, 0-5 cm soils of all plots were taken out and autoclaved
143 at 105 °C for 3 h to deactivate the seeds and then refilled back. This was adequate to
144 prevent the reproduction of original plants. In July 2013, plant community was
145 transplanted from local natural grassland according to its species composition by
146 point quadrats (Goodall, 1952). At the start of this experiment, the plant community
147 composition was the same for all the treatment plots. Plant community composition
148 was investigated in a permanent quadrat of 1 m × 1 m at August of 2014 and 2015
149 (unpublished data). The plant community at the site is dominated by *Carex duriuscula*.

150 The chemical characteristics of the 0-10 cm soil are given in Table 1.

151

152 **2.3 Soil sampling and chemical analysis**

153 In October 2015 (i.e. after 2 years of plant community settled), a composite soil
154 sample was taken from three randomly selected locations within each plot from three
155 soil layers of 0-10 cm, 10-20 cm, and 20-40 cm, respectively. Fresh soil samples were
156 sieved through 2 mm screen and visible plant roots were taken out. After
157 transportation to laboratory, the soils were air-dried and a subsample of the soil was
158 ground for C and N analysis.

159

160 **2.3.1 Soil pH and particle size distribution**

161 Soil pH was determined in a 1:2.5 (w/v) soil-to-water extract of soil samples from all
162 treatments with a PHS-3G digital pH meter (Precision and Scientific Corp., Shanghai,
163 China). Soil particle size distribution was determined by the pipette method in a
164 sedimentation cylinder, using Na-hexamethaphosphate as the dispersing agent (Zhao
165 et al., 2006). Proportion of soil fine particles (<0.25mm) were calculated by summing
166 up the proportions of fine sand, silt and clay in this study.

167

168 **2.3.2 Soil base cations (Ca, Mg, K, Na) and available micronutrients (Fe, Mn, Cu, 169 Zn)**

170 Soil base cations were determined using the CH₃COONH₄-extraction method
171 according to Ochoa-Hueso et al. (2014). Briefly, 2.5 g of soil sample was extracted

172 with 1 M CH₃COONH₄ (pH 7.0) with a soil:extractant ratio of 1:20 (w/v) and shaken
173 at 150 rpm for 30 min. After filtration with Whatman no. 2V filter paper, the
174 concentrations of soil base cations were determined by atomic absorption
175 spectrometer (AAS, Shimazu, Japan).

176 Available Fe, Mn, Cu and Zn were extracted by diethylenetriaminepentaacetic
177 acid (DTPA) according to method of Lindsay and Norvell (1978). Briefly, 10 g of soil
178 samples was mixed with 20 ml 0.005 M DTPA + 0.01 M CaCl₂ + 0.1 M
179 triethanolamine (TEA) (pH 7.0). The slurry was shaken at 180 rpm for 2 h and filtered
180 through Whatman no. 2V filter paper. The concentrations of available micronutrients
181 were analyzed by AAS.

182

183 **2.4 Statistical analyses**

184 The normality of data was tested using the Kolmogorov-Smirnov test, and
185 homogeneity of variances using Leven's test. Effects of soil coarseness on soil pH,
186 fine particles, base cations and available micronutrients were determined by one-way
187 ANOVA. Multiple comparisons with Duncan design were performed to determined
188 difference in soil parameters among soil coarseness degrees. Pearson correlation
189 analysis was used to examine the relationship among soil parameters. Multivariate
190 linear regression analyses (stepwise removal) were conducted to determine variables
191 that made significant contributions to variance of soil base cations and available
192 micronutrients. All statistical analyses were performed in SPSS 16.0 (SPSS, Inc.,
193 Chicago, IL, U.S.A) and statistical significance was accepted at P < 0.05.

194

195 **3 Results**

196 **3.1 Soil pH**

197 Soil coarseness significantly increased soil pH by up to 8.8% across three soil depths
198 (Fig. 2a; Table 2). For both 0-10 cm and 10-20 cm soils, the highest soil pH was
199 detected in C70 (7.3 and 7.4, respectively) and C50 (7.2 and 7.3, respectively) soils,
200 which were followed by C30 and C10 soils (Fig 2a). Significant and positive overall
201 effect of soil depth was detected on soil pH (Fig. 2a; Table 2). For both C0 and C50
202 treatments, soil pH of 10-20 cm and 20-40 cm was significantly higher as compared
203 to that of 0-10 cm soil (Fig. 2a). Soil pH in 10-20 cm of C10 and C30 was
204 significantly higher than that in 0-10 cm of C10 and C30, respectively (Fig. 2a).
205 Significant interactive effect of soil coarseness and soil depth was found on soil pH
206 (Table 2).

207 Proportions of soil fine particles (< 0.25 mm) were determined in 0-10 cm soil.
208 Soil fine particles significantly decreased with soil coarseness degree by 6.3% (for
209 treatment of C10), 17.7% (C30), 34.1% (C50), and 55.6% (C70) as compared to C0
210 (Fig. 2b). The lowest proportion of soil fine particles was detected in C70 (39.1%),
211 and followed by C50 (58.1%) (Fig. 2b). Proportion of clay particles significantly
212 decreased under soil coarseness (Fig. S1).

213

214 **3.2 Soil base cations**

215 Across three soil depths, soil coarseness significantly decreased both exchangeable Ca

216 and Mg concentrations by up to 29.8% and 47.5%, respectively, as compared to C0
217 (Fig. 3a,b). Both exchangeable Ca and Mg concentrations were the lowest in the C70
218 and followed by C50 as compared to C0 in all soil depths (Fig. 3a,b). Soil depth
219 significantly decreased soil exchangeable Mg, while showed no effect on
220 exchangeable Ca (Table 2). Both soil coarseness and soil depth had no impact on soil
221 exchangeable K (Fig. 3c). At 0-10 cm, C50 and C70 significantly decreased soil
222 exchangeable Na by 22.3% and 24.2%, respectively, as compared to C0 (Fig. 3d). Soil
223 exchangeable Na did not change with soil depth (Fig. 3d, Table 2).

224

225 **3.3 Soil available micronutrients**

226 Soil available Fe significantly decreased with soil coarseness degree by as much as
227 17.1% in C10, 22.0% in C30, 36.6% in C50 and 62.5% in C70 across three soil
228 depths (Fig. 4a). Soil coarseness significantly decreased soil available Mn for 0-10 cm
229 (by up to 17.3% in C70) and 10-20 cm (by up to 45.4% in C70) soils (Fig. 4b). Both
230 soil available Fe and Mn significantly decreased with soil depth (Fig. 4a,b; Table 2).
231 Significant negative desertification effect was detected on soil available Cu by 14.7%
232 - 44.4% as compared to C0 in 0-10 cm soil (Fig. 4c). For both C30 and C50
233 treatments, soil available Cu concentration of 10-20 cm soil was significantly higher
234 than that in 0-10 cm and 20-40 cm soils. Soil available Zn concentration was not
235 affected by soil coarseness but it decreased with soil depth (Fig. 4d; Table 2).

236

237 **3.4 Regression analyses between soil parameters**

238 All regression analysis were conducted for 0-10 cm soil as the data of fine particles (<
239 0.25 mm) were only available for 0-10 cm soil. At 0-10 cm soil, soil pH significantly
240 and negatively correlated with exchangeable Ca, Mg and Na, and with available Fe,
241 Mn and Cu (Table 3). Soil fine particles (< 0.25 mm) significantly and positively
242 correlated with exchangeable Ca, exchangeable Mg, exchangeable Na, available Fe,
243 available Mn, and available Cu (Table 3). The SOC significantly and positively
244 correlated with exchangeable Ca, Mg, Na, available Fe, Mn, and Cu (Table 3).

245 According to multiple regression models, change of soil fine particles explained
246 65.5%, 75.7%, 31.4%, 24.0% of variations in exchangeable Ca, Mg, Na, and available
247 Mn (Table 3). Soil pH explained 75.7% of variation in available Fe (Table 2). The
248 SOC explained 59.3% of variation in available Cu (Table 2).

249

250 **4 Discussion**

251 **4.1 Effect of soil coarseness on soil base cations and available micronutrients**

252 Significant decrease in exchangeable Ca and Mg concentrations in three soil depths
253 and exchangeable Na in 0-10 cm soil as affected by soil coarseness partially
254 supported our first hypothesis. The decrease of exchangeable Ca, Mg and Na might be
255 due to increase of soil pH under soil coarseness as suggested by the significant and
256 negative correlation between soil pH and exchangeable Ca, Mg and Na (Table 3).
257 Indeed, with the increase of soil pH, soil base cations (such as Ca^{2+} and Mg^{2+}) and
258 available micronutrients (Fe^{2+} , Mn^{2+} and Cu^{2+}) would precipitate with OH^- (McLean,
259 1982) resulting in the decrease of soil base cations and available micronutrients under

260 soil coarseness.

261 Soil fine particles (< 0.25 mm), especially clay inside these fine particles were
262 suggested to provide additional binding surfaces for exchangeable base cations and
263 available micronutrients (Beldin et al., 2007). Confirmed by the negative correlation
264 of soil fine particles with both base cations (exchangeable Ca, Mg and Na) available
265 micronutrients (Fe, Mn and Cu), the decrease of soil fine particles and clay content
266 (Fig. S1) might also contribute to lower base cations and available micronutrients
267 under soil coarseness. Consistent with our findings, previous studies also suggested
268 that the decrease of soil fine particles and increase of soil coarseness resulted in loss
269 of SOM as well as reduction in the nutrient storage (Lopez, 1998; Zhao et al., 2006;
270 Zhou et al., 2008).

271 As the essential role of SOM in retaining base cations and micronutrients by its
272 functional groups (Oorts et al., 2003), significantly lower soil base cations and
273 micronutrients would possibly due to lower C (the largest component of SOM)
274 concentration in coarsen soils. This can be further enhanced by the significant positive
275 correlation of soil C with both base cations and micronutrients (Table 3). Consistently,
276 Vittori Antisari et al. (2013) reported that humified organic compounds in soil could
277 retain base cations and decrease their leaching from soils. As compared to higher soil
278 coarseness degree, higher soil microbial activities (unpublished data) under conditions
279 of lower soil coarseness degree could promote humification or microbial-processing
280 of the SOM (Wang et al., 2015b), potentially increasing the availability of functional
281 groups to complex with the base cations and micronutrients. Additionally, higher net

282 primary production and plant nutrient demands would induce the activation of base
283 cation and micronutrients from the soils under lower soil coarseness degree (Burke et
284 al., 1999). Due to the fact of reduction in ecosystem productivity under cation
285 deficiencies (Lawrence et al., 1995; Cheng et al., 2010), the loss of base cations and
286 available micronutrients as developed under soil coarseness could constrain both plant
287 growth and pasture productivity of this nutrient poor sandy ecosystem.

288

289 **4.2 Effect of soil depth on base cations and available micronutrients**

290 The hypothesized decrease of exchangeable base cations and available micronutrients
291 with soil depth was partially supported as only exchangeable Mg (Fig. 3b), available
292 Fe (Fig. 4a), Mn (Fig. 4b) and Zn (Fig. 4d) decreased with soil depth. Vertical
293 distribution of soil nutrients can be influenced by two opposite processes, leaching
294 and biological cycling (such as plant absorption) (Truggill, 1988). Being a ubiquitous
295 process in ecosystems, plant absorption of nutrients can transport soil elements
296 aboveground and return the litterfall to soil surface (Stark, 1994). Especially in this
297 sandy land or desertified grassland, plants tend to accumulate SOM or nutrients to
298 form 'island of fertility' (Cao et al., 2008). In these sandy soils, leaching is also an
299 essential process in shaping the vertical distribution of soil nutrients (Truggill, 1988).
300 As leaching moves nutrients downward while biological cycling moves them upward
301 (Jobb gy and Jackson, 2001), the unchanged Ca, K and Na concentrations might be
302 the combining effects of leaching and biological cycling. This area experiences
303 freeze-thaw cycles for at least 4-5 months per year (Alamusa et al., 2014).

304 Freeze-thaw cycles might promote the leaching of exchangeable Ca, K and Na from
305 surface to subsoil resulting in the unchanged base cations along soil profile. Our
306 results are in contrast with previous studies suggesting that ecosystem were more
307 capable to retain K than other base cations (Nowak et al., 1991; Jobb gy and Jackson,
308 2001). In this case, it is obvious that many environmental factors, like soil types and
309 plant community composition can be drivers for the vertical distribution of base
310 cations and micronutrients (Burke et al., 1999; Van der Ploeg et al., 2012). The
311 leaching of base cations to subsoils might enhance mineral weathering process and
312 pedogenesis by forming kaolinite in topsoils as rapid removing of water-soluble
313 elements (such as exchangeable Ca and Na) (Chadwick and Chorover, 2001).
314 Stronger effect of plant absorption than leaching might contribute to the shallower
315 distribution of exchangeable Mg (Fig. 3b), available Fe (Fig. 4a), Mn (Fig. 4b) and Zn
316 (Fig. 4d). The dominant role of plant cycling in determining the vertical distribution
317 of Mg, Fe, Mn and Zn might illustrate that these elements were scarcer and more
318 limiting nutrients for plant growth in this semi-arid sandy ecosystem (Jobb gy and
319 Jackson, 2001).

320

321 **5 Conclusions**

322 The results showed that grassland soil coarseness decreased soil base cations of
323 exchangeable Ca, Mg and Na as well as available micronutrients of Fe, Mn and Cu.

324 The loss of SOM, decrease of soil fine particles, and increase of soil pH were the
325 main driving factors for the decrease of base cations and micronutrient availability as

326 affected by soil coarseness. Unchanged concentrations of exchangeable Ca, K and Na
327 along the soil depth might result from the balance between plant cycling and leaching
328 effects. The dominant role of plant cycling over leaching shaped the shallower
329 distribution of exchangeable Mg as well as available Fe, Mn and Zn. The reduction
330 and re-distribution of soil base cations and available micronutrients would potentially
331 influence soil fertility and plant productivity in this desertified grassland ecosystem.

332

333 **Author contribution**

334 Z. Wang, G. Yu, and X. Han designed the experiments; and L. Lü and Y. Zhao carried
335 them out. H. Liu and J. Yin help to do the laboratory analysis. L. Lü and R. Wang
336 prepared the manuscript with contributions from all authors. Y. Jiang helped to revise
337 the manuscript. Mr. Xiao created the figure of our experimental location.

338

339 **Acknowledgments**

340 This work was financially supported by the National Natural Science Foundation of
341 China (41371251).

342

343 **References**

344 Alamusu, Niu C., Zong Q.: Temporal and spatial changes of freeze-thaw cycles in
345 Ulan'aodu Region of Horqin Sandy Land, Northern China in a changing climate,
346 Soil Sci. Soc. Am. J., 78, 89-96, 2014.

347 Allington, G., Valone, T.: Reversal of desertification: the role of physical and

348 chemical soil properties, *J. Arid Environ.*, 74, 973-977, 2010.

349 Antisari, L. V., Carbone, S., Gatti, A., Vianello, G., Nannipieri, P.: Toxicity of metal
350 oxide (CeO₂, Fe₃O₄, SnO₂) engineered nanoparticles on soil microbial biomass and
351 their distribution in soil, *Soil Biol. Biochem.*, 60, 87-94, 2013.

352 Barbero-Sierra, C., Marques, M. J., Ruiz-Pérez, M., Escadafal, R., Exbrayat, W.: How
353 is desertification research addressed in Spain? Land versus soil approaches, *Land*
354 *Degrad. Dev.*, 26, 423-432, 2015.

355 [Barthold, F. K., Wiesmeier, M., Breuer, L., Frede, H.-G., Wu, J., Blank, F. B.: Land](#)
356 [use and climate control the spatial distribution of soil types in the grasslands of](#)
357 [Inner Mongolia, *J. Arid Environ.*, 88, 194-205, 2013.](#)

358 Beldin, S. I., Caldwell, B. A., Sollins, P., Sulzman, E. W., Lajtha, K., Crow, S. E.:
359 Cation exchange capacity of density fractions from paired conifer/grassland soils,
360 *Biol. Fertil. Soils*, 43, 837-841, 2007.

361 Beyene, F.: Incentives and Challenges in Community - Based Rangeland Management:
362 Evidence from Eastern Ethiopia, *Land Degrad. Dev.*, 26, 502-509, 2015.

363 Bisaro, A., Kirk, M., Zdruli, P., Zimmermann, W.: Global drivers setting
364 desertification research priorities: insights from a stakeholder consultation forum,
365 *Land Degrad. Dev.*, 25, 5-16, 2014.

366 [Burke, I. C., Lauenroth, W. K., Riggle, R., Brannen, P., Madigan, B., Beard, S.:](#)
367 [Spatial variability of soil properties in the shortgrass steppe: the relative importance](#)
368 [of topography, grazing, microsite, and plant species in controlling spatial patterns.](#)
369 [*Ecosystems*, 2, 422-438, 1999.](#)

370 Cao, C., Jiang, D., Teng, X., Jiang, Y., Liang, W., Cui, Z.: Soil chemical and
371 microbiological properties along a chronosequence of *Caragana microphylla* Lam.
372 plantations in the Horqin sandy land of Northeast China. *Appl. Soil Ecol.*, 40, 78-85,
373 2008.

374 Cerdà A., Gallart, F., Li, J., Papanastasis, V. P., Parmenter, R. R., Turnbull, L.,
375 Parsons, A. J., Wainwright, J.: Long-range ecogeomorphic processes, in:
376 Self-organized ecogeomorphic systems: confronting models with data for
377 land-degradation in drylands, edited by: Müller, E. N., Wainwright, J., Parsons, A.
378 J., Turnbull, L., Springer, Dordrecht 103 – 139, 2014.

379 Cheng, L., Zhu, J., Chen, G., Zheng, X., Oh, N. H., Ruffy, T., Hu, S.: Atmospheric
380 CO₂ enrichment facilitates cation release from soil, *Ecol. Lett.*, 13, 284-291, 2010.

381 Chen, F., Zeng, D., Narain, S. A., Chen, G.: Effects of soil moisture and soil depth on
382 nitrogen mineralization process under Mongolian pine plantations in Zhanggutai
383 sandy land, PR China, *J. Forest. Res.*, 16, 101-104, 2005.

384 Chadwick, O.A., Chorover, Jon: The chemistry of pedogenic thresholds, *Geoderma*,
385 100, 321-353, 2001.

386 Conte, C. A.: The forest becomes desert: Forest use and environmental change in
387 Tanzania's West Usambara mountains, *Land Degrad. Dev.*, 10, 291-309, 1999.

388 Escadafal, R., Barbero-Sierra, C., Exbrayat, W., Marques, M. J., Akhtar-Schuster, M.,
389 El Haddadi, A., Ruiz, M.: First appraisal of the current structure of research on land
390 and soil degradation as evidenced by bibliometric analysis of publications on
391 desertification, *Land Degrad. Dev.*, 26, 413-422, 2015.

392 Fleskens, L., Stringer, L. C.: Land management and policy responses to mitigate
393 desertification and land degradation. *Land Degrad. Dev.*, 25, 1-4, 2014.

394 Goodall, D. W.: Some considerations in the use of point quadrats for the analysis of
395 vegetation, *Aust. J. Biol. Sci.*, 5, 1-41, 1952.

396 Huang, D., Wang, K., Wu, W.: Dynamics of soil physical and chemical properties and
397 vegetation succession characteristics during grassland desertification under sheep
398 grazing in an agro-pastoral transition zone in Northern China, *J. Arid Environ.*, 70,
399 120-136, 2007.

400 IUSS Working Group WRB: World Reference Base for Soil Resources 2014.
401 International soil classification system for naming soils and creating legends for
402 soil maps, World Soil Resources Reports No. 106. FAO, Rome, 2014.

403 Jobb gy, E. G., Jackson, R. B.: The distribution of soil nutrients with depth: global
404 patterns and the imprint of plants, *Biogeochemistry*, 53, 51-77, 2001.

405 Lawrence, G. B., David, M. B., Shortle, W.C.: A new mechanism for calcium loss in
406 forest-floor soils, *Nature*, 378, 162-165, 1995.

407 Li, F., Zhao, L., Zhang, H., Zhang, T., Shirato, Y.: Wind erosion and airborne dust
408 deposition in farmland during spring in the Horqin Sandy Land of eastern Inner
409 Mongolia, China, *Soil and Tillage Research*, 75, 121-130, 2004.

410 Li, F., Cook, S., Geballe, G. T., Burch, Jr. W. R., Rainwater harvesting agriculture: an
411 integrated system for water management on rainfed land in China's semiarid areas,
412 *AMBIO: A Journal of the Human Environment*, 29, 477-483, 2000.

413 Lindsay, W. L., Norvell, W. A., Development of a DTPA soil test for zinc, iron,

414 manganese, and copper, *Soil Sci. Soc. Am. J.*, 42, 421-428, 1978.

415 Lopez, M.: Wind erosion in agricultural soils: an example of limited supply of
416 particles available for erosion, *Catena*, 33, 17-28, 1998.

417 Lu, X., Mao, Q., Gilliam, F. S., Luo, Y., Mo, J: Nitrogen deposition contributes to soil
418 acidification in tropical ecosystems, *Global Change Biol.*, 20, 3790-3801, 2014.

419 Marques, M. J., Bienes, R., Cuadrado, J., Ruiz-Colmenero, M., Barbero-Sierra, C.,
420 Velasco, A.: Analysing perceptions attitudes and responses of winegrowers about
421 sustainable land management in central Spain, *Land Degrad. Dev.*, 26, 458-467,
422 2015.

423 McLean, E. O.: Soil pH and lime requirement, in: *Methods of soil analysis, Part 2,*
424 *Chemical and microbiological properties*, edited by: Page, A. L., Miller, R. H.,
425 Keeney, D. R., American Society of Agronomy, Madison, WI, USA, 199-224,
426 1982.

427 Miao, L., Moore, J. C., Zeng, F., Lei, J., Ding, J., He, B., Cui, X.: Footprint of
428 research in desertification management in China, *Land Degrad. Dev.*, 26, 450-457,
429 2015.

430 Nowak, C., Downard, R., White, E.: Potassium trends in red pine plantations at Pack
431 Forest, New York, *Soil Sci. Soc. Am. J.*, 55, 847-850, 1991.

432 Ochoa-Hueso, R., Bell, M. D., Manrique, E.: Impacts of increased nitrogen deposition
433 and altered precipitation regimes on soil fertility and functioning in semiarid
434 Mediterranean shrublands, *J. Arid Environ.*, 104, 106-115, 2014.

435 Oorts, K., Vanlauwe, B., Merckx, R.: Cation exchange capacities of soil organic

436 matter fractions in a Ferric Lixisol with different organic matter inputs, Agr.
437 Ecosyst. Environ., 100, 161-171, 2003.

438 Schlesinger, W. H., Reynolds, J., Cunningham, G. L., Huenneke, L., Jarrell, W.,
439 Virginia, R., Whitford, W.: Biological feedbacks in global desertification, Science,
440 247, 1043-1048, 1990.

441 Sharma, B., Arora, H., Kumar, R., Nayyar, V.: Relationships between soil
442 characteristics and total and DTPA-extractable micronutrients in Inceptisols of
443 Punjab. Commun. Soil Sci. Plan., 35, 799-818, 2004.

444 Stark, J. M.: Causes of soil nutrient heterogeneity at different scales. in: Exploitation
445 of Environmental Heterogeneity by Plants: Ecophysiological Processes Above and
446 Below Ground, edited by Caldwell, M. M, Pearcy, R., J. Wiley & Sons, NY, pp.
447 255-284, 1994.

448 Torres, L., Abraham, E. M., Rubio, C., Barbero, C., Ruiz, M.: Desertification research
449 in Argentina, Land Degrad. Dev., 26, 433-440, 2015.

450 Van der Ploeg, M. J., Appels, W. M., Cirkel, D. G., Oosterwoud, M.R., Witte, J.-P. M.,
451 Van der Zee, S. E. A. T. M.: Microtopography as a driving mechanism for
452 ecohydrological processes in shallow groundwater systems, Vadose Zone J., 11, No.
453 3, 2012.

454 Vieira, R. M. S. P., Tomasella, J., Alvalá R. C. S., Sestini, M. F., Affonso, A. G.,
455 Rodriguez, D. A., Barbosa, A. A., Cunha, A. P. M. A., Valles, G. F., Crepani, E., de
456 Oliveira, S. B. P., de Souza, M. S. B., Calil, P. M., de Carvalho, M. A., Valeriano, D.
457 M., Campello, F. C. B., Santana, M. O.: Identifying areas susceptible to

458 desertification in the Brazilian northeast, *Solid Earth*, 6, 347-360, 2015.

459 Wang, T., Xue, X., Zhou, L., Guo, J.: Combating aeolian desertification in northern
460 China, *Land Degrad. Dev.*, 26, 118-132, 2015a.

461 Wang R., Dungait, J. A. J., Creamer, C. A., Cai, J., Li, B., Xu, Z., Zhang, Y., Ma, Y.,
462 Jiang, Y.: Carbon and nitrogen dynamics in soil aggregates under long-term
463 nitrogen and water addition in a temperate steppe, *Soil Sci. Soc. Am. J.*, 79,
464 527-535, 2015b.

465 Weinzierl, T., Wehberg, J., Böhner, J., Conrad, O.: Spatial Assessment of Land
466 Degradation Risk for the Okavango River Catchment, Southern Africa, *Land
467 Degrad. Dev.*, in press, 2015. DOI: 10.1002/ldr.2426

468 Xie, L., Zhong, J., Cao, F., Li, J., Wu, L.: Evaluation of soil fertility in the succession
469 of karst rocky desertification using principal component analysis, *Solid Earth*, 6,
470 515-524, 2015.

471 Xu, Q., Wang, T., Cai, C., Li, Z., Shi, Z.: Effects of soil conservation on soil
472 properties of citrus orchards in the Three - Gorges Area, China, *Land Degrad. Dev.*,
473 23, 34-42, 2012.

474 Yan, X., Cai, Y.: Multi-scale anthropogenic driving forces of karst rocky
475 desertification in Southwest China, *Land Degrad. Dev.*, 26, 193-200, 2015.

476 Zhang, Y., Xu, Z., Jiang, D., Jiang, Y.: Soil exchangeable base cations along a
477 chronosequence of *Caragana microphylla* plantation in a semi-arid sandy land,
478 China, *J. Arid Land*, 5, 42-50, 2013.

479 Zhao, H., Zhou, R., Zhang, T., Zhao, X.: Effects of desertification on soil and crop

480 growth properties in Horqin sandy cropland of Inner Mongolia, north China, Soil
481 Till. Res., 87, 175-185, 2006.
482

483 **Tables**

484

485 **Table 1** Mean and range of soil chemical characteristics for 0-10 cm soil in different
 486 soil coarseness degrees from 0% sand addition (C0) to 70% (C70).

	Range of mean from C0 to C70
Soil organic carbon (g kg soil ⁻¹)	4.1-2.7
Total nitrogen (g kg soil ⁻¹)	0.48-0.22
Dissolved organic carbon (mg kg soil ⁻¹)	66.3-53.9
Microbial biomass C (mg kg soil ⁻¹)	104.3-60.4
Electric conductivity (μs cm ⁻¹)	54.7-44.7

487

488 **Table 2** Results (*F* values) of two-way ANOVAs on the effect of soil depth (D),
 489 treatments of soil coarseness degrees (T), and their interactions on soil base cations
 490 (exchangeable Ca, Mg, K, and Na) and available micronutrients (Fe, Mn, Cu, and Zn).

	pH	Ca	Mg	K	Na	Fe	Mn	Cu	Zn
D	17.33**	0.99	4.54*	0.74	0.22	135.67**	99.63**	9.69**	22.13**
T	31.74**	50.26**	57.22**	1.61	2.35	41.51**	12.11**	10.60**	0.62
DXT	4.12**	1.49	3.81**	0.64	5.02**	2.97**	2.49*	3.40**	0.83

491 * Significance level at $P < 0.05$.

492 ** Significance level at $P < 0.01$.

493

494

495 **Table 3** Regression statistics relating soil base cations (exchangeable Ca, Mg, K, and
 496 Na) and available micronutrients (Fe, Mn, Cu, and Zn) to soil pH, soil fine particles
 497 (<0.25 mm) and soil organic carbon (SOC).

	Soil pH	< 0.25mm	SOC	Multiple
Ca	-0.67	0.81**	0.73	0.81
Mg	-0.75	0.87**	0.72	0.87
K	—	—	—	—
Na	-0.54	0.56**	0.56	0.56
Fe	-0.87**	0.80	0.74	0.87
Mn	-0.42	0.49**	0.45	0.49
Cu	-0.68	0.72	0.77**	0.77
Zn	—	—	—	—

498 Values are *R* statistics for significant ($P < 0.05$) linear regressions. Multiple is *R*
 499 values for multiple regressions (stepwise removal) of soil base cation and
 500 micronutrients with soil pH, <0.25 mm fine particles, and SOC. ** indicates variables
 501 that make significant contributions to the multiple linear regressions.
 502

503 **Figure Legends**

504 **Fig. 1** Location of the experimental site.

505

506 **Fig. 2** Soil pH values for three soil depths (a) and proportion of soil fine particles (<
507 0.25 mm) for 0-10 cm soil in different soil coarseness degrees of 0% sand addition
508 (C0), 10% (C10), 30% (C30), 50% (C50) and 70% (C70). Data represent mean \pm SE
509 (n=6). Letters indicate significant differences among treatments (lowercase letters)
510 and differences among soil depths when averaging across all treatments (capital
511 letters).

512

513 **Fig. 3** Soil base cations of exchangeable Ca (a), Mg (b), K (c) and Na (d) for three
514 soil depths in different soil coarseness degrees of 0% sand addition (C0), 10% (C10),
515 30% (C30), 50% (C50) and 70% (C70). Data represent mean \pm SE (n=6). Letters
516 indicate significant differences among treatments (lowercase letters) and differences
517 among soil depths when averaging across all treatments (capital letters). Bars without
518 letters above denote no significance is detected among treatments.

519

520 **Fig. 4** Soil available micronutrients of available Fe(a), Mn (b), Cu (c) and Zn (d) for
521 three soil depths in different soil coarseness degrees of 0% sand addition (C0), 10%
522 (C10), 30% (C30), 50% (C50) and 70% (C70). Data represent mean \pm SE (n=6).
523 Letters indicate significant differences among treatments (lowercase letters) and
524 differences among soil depths when averaging across all treatments (capital letters).

525 | Bars without letters above denote no significance is detected among treatments.

526

527 **Fig. S1** Proportion of soil clay particles for 0-10 cm soil in different soil coarseness


528 degrees of 0% sand addition (C0), 10% (C10), 30% (C30), 50% (C50) and 70% (C70).

529 Data represent mean \pm SE (n=6). Letters indicate significant differences among

530 treatments.


531

532 **Fig. 1**



533

534 **Fig. 2**




535

536

537

538 **Fig. 3**




539

540

541

542 **Fig. 4**




543

544

545 **Fig. S1**

546



547