

Cooperative Effects of Field Traffic and Organic Matter Treatments on Some Compaction-Related Soil Properties

Metin Mujdeci^{1*}, Ahmet Ali Isildar¹, Veli Uygur¹, Pelin Alaboz¹, Husnu Unlu², Huseyin Senol¹

¹Suleyman Demirel University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition, Isparta, Turkey.

²Suleyman Demirel University, Faculty of Agriculture, Department of Horticultural Sciences, Isparta, Turkey.

5 Correspondence to: Metin Mujdeci (metinmujdeci@sdu.edu.tr)

Abstract. Soil compaction is common problem of mineral soils under conventional tillage practices. Organic matter addition is an efficient way of reducing the effects of field traffic in soil compaction. The aim of this study was to investigate the effects of number of tractor passes (1, 3 and 5) on depth dependent (0-10 and 10-20 cm) penetration resistance, bulk density and porosity of a clay textured soil (Typic Xerofluvent) under organic vegetable cultivation practiced in 2010-2013 growing
10 seasons treated with farmyard manure (35 t FYM ha⁻¹), green manure (GM) (common vetch, *Vicia sativa* L.) and conventional tillage (C). The number of tractor passes resulted in increases in bulk density and penetration resistance (C>GM>FYM) whereas the volume of total and macro pores decreased. The maximum penetration resistance (3.60 MPa) was recorded in C treatment with 5 passes in 0-10 cm depth whereas the minimum (1.64 MPa) was observed for FYM treatment with 1 pass in 10-20 cm depth. The highest bulk density was determined as 1.61 g cm⁻³ for C treatment with 5
15 passes in 10-20 cm depth, the smallest value was 1.25 g cm⁻³ in the FYM treatment with only 1 pass in 0-10 cm depth. The highest total and macro pores volume were determined as 0.53 and 0.16 cm³ cm⁻³, respectively, in 0-10 cm depth of FYM treatment with 1 pass. The volume of micro pores (0.38 cm³ cm⁻³) was higher in 0-10 cm depth of FYM treatment with 3 passes. It can be concluded that organic pre-composted organic amendment rather than green manure was likely to be more efficient in **mitigating** compaction problem in soils.

20

Key Words: soil penetration resistance, farmyard manure, green manure, bulk density, porosity

1. Introduction

25 Agriculture development resulted in the increase of the food supply to the humankind, but also the increase in soil and water losses, reduction of the vegetation cover and degradation of the soil (Cerdeira, 2000). One of the consequences of the agriculture use and abuse is the increase in the soil compaction. Soil compaction that can be defined as a soil degradation process in which an applied pressure to a soil causes re-organisation to get closer of soil grains resulting in porosity and pore volume ratio is regarded as the most serious environmental problem resulting in conventional agriculture
30 (McGarry, 2003). Since the farmers have difficulty in locating and rationalizing this type of degradation without making any

measurement this problem can be more deleterious in conventional agriculture. In addition, compaction induced shallow plant rooting and poor plant growth reduce crop yield for deep rooting plants and vegetative cover to protect soil from erosion (Ni et al., 2015; Ola et al., 2015; Shaw et al., 2016). Compaction can increase runoff and erosion from sloping land or waterlogged soils in flatter food slopes depending on reduced water infiltration through soil surface (Al-Dousari et al., 2000; USDA-NRCS, 2012; Pulido et al., 2016). Intensive agricultural related-soil compaction may be regarded as one of the significant reasons for land degradation (Cerdeira, 2000; Barbero-Sierra et al., 2015; Wang et al., 2015; Yan et al., 2015) and the elevated risks about food security, water scarcity, climate change, biodiversity loss and health threats which were pointed out as soil related challenges for sustainable society (Keesstra et al., 2016).

The most significant cause of soil compaction is field traffic. Meanwhile the close relation between field traffic density and frequency and crop type should also be taken into account (De Oliveira et al., 2015; Gelaw et al., 2015). Thus, more than 80% of a corn and soybean field has been under tyre pressure in a growth season (Erbach, 1986). 90%, 35% and 60% of field under wheel pressure during seed bed preparation, harvesting and baling practices, respectively, in cereal cultivation (Munsuz, 1985). Soil aeration, infiltration, and hydraulic conductivity parameters of soils which are closely related to differential porosity show decreases related to increased-field traffic (Seker and Isildar, 2000; Aksakal, 2004). In order to prevent from such adverse effects, a decreased field traffic is necessary along with tillage soil at optimum moisture content and increasing the organic matter content of soil with farmyard manure, compost, green manure, etc. addition. Stabilization and fortification of soil aggregates with organic matter addition can increase compaction resistance of soils (Cochrane and Aylmore, 1994; Thomas et al., 1996; Aksakal et al., 2016), and enhance to a differing degree the compaction related attributes such as bulk density, pore-size distribution, infiltration, etc. in soils (Sparovek et al., 1999; Carter, 2002; Aksakal et al., 2016). The changes in soil organic carbon stock under different management systems (Munoz et al., 2015) can also influence total soil quality in soils. For example, Parras-Alcantra et al. (2015) reported a higher organic farming induced-stratification ratio deeper in the surface horizon comparing to conventional agricultural system. Gelaw et al. (2015) pointed out that the management of soil differently affected the partition of organic matter in different aggregate sizes which in turn influenced bulk density and water stable aggregates. Despite the specific effect of field traffic was not elucidated in these studies (Gelaw et al., 2015; Parras-Alcantra et al., 2015), the management systems are closely related to traffic density and soil physical attributes. The organic matter is also relevant on the soil behavior but also at the atmospheric level and at the whole behavior of the Earth System as can control the Carbon cycle (Novara et al., 2013; Kaleem Abbasi et al., 2015; Peng et al., 2015).

Keesstra et al. (2016) pointed out the significance of raising public/farmers' awareness about key attributes of soil organic matter to function and sustain ecosystem services. Similarly many researcher reported that soil physical and chemical properties in terms of fertility and sustainability of agriculture may be enhanced, to a large extent, by regular organic matter application (Aggelides and Londra, 2000; Alagoz et al., 2006; Mamman et al., 2007; Celik et al., 2010; Gulser and Candemir, 2012). The above mentioned literature points out that the nature and extent of compaction induced soil degradation can be exaggerated by the lack of organic matter. Artificial loosening of soils by deep ripping is common

suggested practice for elimination of the deleterious effects of compaction but its effect is not long-lasting (Hamza and Anderson, 2003; Hamza and Anderson, 2008; Arslan, 2006). Therefore, the aim of this study was to investigate the effects of both annual addition of different organic matter (farmyard manure and green manuring) and field traffic density on penetration resistance, bulk density, porosity in a clay textured soil with low organic matter content.

5 2. Materials and Methods

Study area and experimental design

This study was carried out in experimental field of Agricultural Research and Application Centre of Suleyman Demirel University in 2010-2013 years. Some chemical and physical properties of soil were given in Table 1.

Organic vegetables were cultivated with farmyard manure (FYM), green manure (GM) and conventional tillage without any organic matter addition (C) treatments. The field experiment was set up in completely randomised design with three replications. FYM application was executed in between 21st of May and 7th of June at 35 t ha⁻¹ (FYM consist of 45% dry matter) and then thoroughly mixed into surface layer (10-15 cm) of soils by rototiller. Common vetch (*Vicia sativa* L.) was sowed in the second week of March and allowed to grow till flowering stage, between 21st of May and 7th of June then it was incorporated into the soil by firstly chisel (20-25 cm tillage depth) and followed by rototillers (10-15 cm tillage depth). Both GM and FYM treatments were performed at the same time. The field was tilled by means of disk-harrow at 10-15 cm depth (16.09.2013) then the field was sprinkler-irrigated on 27th of September 2013. In order to increase the susceptibility of soils to compaction, field traffic was realised at soil water content just below the field capacity (0.23 g g⁻¹) on 5th of November 2013. A tractor (80-66s fiat, manufactured in 1998) having 85 horse power (HP) and 3460 kg weight, including the operator weight, was used to resemble field traffic. Tractor passes (1, 3 and 5 times) were performed at 5 km h⁻¹ speed on the same track. One disturbed and three undisturbed soil cores per plot were taken from 0-10 and 10-20 cm depths before tractor passes and only undisturbed core samples after each passing treatment. Penetration resistance which is indication of soil compaction was measured with a penetrometre (Eijkelkamp penetrograph) equipped with 1 cm² cone attachment. Penetration resistance were performed in 15 replications per treatment at 2 m interval and 0-10 and 10-20 cm depth. The averages of resistance values obtained for 0-10 and 10-20 cm depth were evaluated. Some physical and chemical properties of experimental soil, FYM and plots before passing were given in Table 1 and 2. Particle size distribution was determined by means of Bouyoucos hydrometer method (Bouyoucos, 1962) and bulk density by undisturbed soil cores (Blake and Hartge, 1986). For this analysis soil cores were oven-dried and weighed then calculated by dividing the dry weight to core volume. Total porosity was determined in undisturbed water-saturated core samples of 100 cm³. Micro porosity was accounted as the volume of water at field capacity, measured by using a pressure membrane apparatus at 0.033 MPa suction. Macro porosity was calculated from the difference between total porosity and micro porosity (Danielson and Sutherland, 1986). Wet oxidation method of modified-Walkley and Black and dry ashing method at 400°C for 16 h in an oven were used for organic matter content of soil and FYM, respectively (Burt, 2004). Soil pH and electrical conductivity (EC) were measured in 1:1

soil: distilled water suspension whereas pH and EC of FYM were determined in 1:2.5 FYM: distilled water suspension (Burt, 2004). Carbonate equivalent was determined by a volumetric method using Scheibler calcimeter (Kacar, 2009).

Statistical analysis

The data were subjected to ANOVA using MINITAB 16 statistical package programme (Minitab, 2010). The mean separation between the treatments was performed by **least significant difference** (LSD) test at 95% confidence level.

Table 1. Some chemical and physical properties of experimental soil and FYM.

	Clay (g kg ⁻¹)	Silt (g kg ⁻¹)	Sand (g kg ⁻¹)	Texture class	Organic matter (g kg ⁻¹)	CaCO ₃ (%)	pH	EC (dS m ⁻¹)
Soil								
0-10 cm	425.1	394.5	180.4	C	15.5	24.1	7.44	0.34
10-20 cm	412.9	399.9	187.2	C	15.6	24.2	7.38	0.38
FYM	-	-	-	-	451	-	7.70	3.58

10

Table 2. Porosity, bulk density, organic matter content of the plots before passing.

Treatments	Depth (cm)	Total porosity (cm ³ cm ⁻³)	Macro porosity (cm ³ cm ⁻³)	Micro porosity (cm ³ cm ⁻³)	Bulk density (g cm ⁻³)	Organic matter (g kg ⁻¹)
C	0-10	0.54	0.22	0.32	1.28	15.50
	10-20	0.52	0.19	0.33	1.33	15.60
FYM	0-10	0.61	0.24	0.37	1.14	28.00
	10-20	0.58	0.22	0.36	1.22	21.50
GM	0-10	0.59	0.23	0.36	1.20	17.40
	10-20	0.55	0.20	0.35	1.25	18.30

3. Results and Discussion

15 3.1 Soil penetration resistance

Organic matter amendments significantly reduced penetration resistance in both depths; however the effect of FYM treatment was higher than GM treatment (Fig. 1, Table 3). This finding is in accordance with the previous studies (Celik et

al., 2010; Gulser and Candemir, 2012; Xin et al., 2016). Incorporation of organic matter to clay-textured soils can strengthen the aggregates by weakening cohesion forces and interfering with the formation of crust and large aggregate (Aksakal et al., 2012). The larger amounts of organic matter addition may mediate the formation of clay-organic matter complexes which in fact reduces the penetration resistance on one hand and conserves organic matter against microbial decay on the other hand.

5 In this respect, Blanco-Moure et al. (2016) investigated the effect of soil texture on carbon and organic matter distribution among different fractions under different tillage and management practices. They found that soil clay had critical role in chemical stabilization of organic matter through clay-organic complexes in the soils. Czyz and Dexter (2016) pointed out the relation between the magnitude of clay-soil complex and porous and open nature of structure. Thus, stable organic matter sources such as FYM resulted in desirable penetration resistance for plant growth under changing field traffic.

10 Table 3. Main effects of organic matter incorporation, depth and passing number on measured parameters.

Treatments	Penetration resistance (MPa)	Bulk density (g cm^{-3})	Total porosity ($\text{cm}^3 \text{cm}^{-3}$)	Micro porosity ($\text{cm}^3 \text{cm}^{-3}$)	Macro porosity ($\text{cm}^3 \text{cm}^{-3}$)
Organic matter					
C	2.46 a	1.53 a	0.424 c	0.336 c	0.087 b
FYM	2.02 c	1.40 c	0.471 a	0.364 a	0.107 a
GM	2.27 b	1.47 b	0.445 b	0.354 b	0.091 b
Depth					
0-10 cm	2.44 a	1.43 b	0.458 a	0.354 a	0.104 a
10-20 cm	2.06 b	1.50 a	0.435 b	0.349 b	0.086 b
Passes					
1	1.91 c	1.38 c	0.480 a	0.349 b	0.131 a
3	2.10 b	1.48 b	0.441 b	0.360 a	0.082 b
5	2.73 a	1.54 a	0.418 c	0.345 c	0.073 c

Different letters in the same column indicates differences between the treatment means for each main effect.

Increasing number of passing irrespective from the organic matter treatments and soil depth increased **soil compaction measured by penetration resistance** (Table 3). The effect of field traffic on penetration resistance, as expected, was more negative in **0-10 cm depths** (Table 3). Accordingly, Carman (1994) and Seker and Isildar (2000) were determined higher compaction ratio in the 0-10 and 0-15 cm surface layers, respectively. A penetration resistance value as high as 3.60 MPa in the surface layer caused by 5 times passing in control treatment (**Fig 1**) with no organic matter may have significant inverse effects on infiltration, percolation, and runoff induced erosion under intensive precipitation events on slopy lands (**Kozłowski, 1999; Seker and Isildar, 2000**). In this study, we also determined this well-known manner that surface soil become more compact with field traffic and the severity of the problem may be overcome by organic matter addition into soil **or by adoption soil management system with decreased annual traffic**. Penetration resistance value in 10-20 cm

depths obtained for C and GM treatments after 5 passes was over 2 MPa value which considered limit value by USDA (1993) as a critical physical quality parameter in conventional agricultural practices. This critical value can change depending on the soil tillage systems, for example in minimum tillage practices where chisel is used for soil tillage it is 3 MPa and in no-till system it is 3.5 MPa (De Moraes et al., 2014). Critical penetration resistance value inhibiting the root development is accepted as 3 MPa (Busscher and Sojka, 1987; Hakansson and Lipiec, 2000, Aksakal et al., 2011). Soil management systems can differ soil organic carbon contents (Munoz-Rojas et al., 2015) and field traffic density which ultimately degraded soil physical traits for optimal plant growth such as water stable aggregates and bulk density (Gelaw et al. 2015). In fact, these tendency in soil physical traits can lead a more compaction both in surface and subsurface soil layer as in our case.

10

3.2. Bulk density

The main effect of organic matter treatments on bulk density was statistically significant ($P < 0.01$). Both FYM and GM incorporation into soil were distinctly different than the control (Table 3). FYM amendment reduced the bulk density to as low as 1.40 g cm^{-3} . Similar to our findings, organic matter amendment-induced-decreases (Haynes and Naudi, 1998; Chaudhari et al., 2013; Gulser and Candemir, 2012) and field traffic related increases (Seker and Isildar, 2000; Patel and Mani, 2011) in bulk density have been frequently reported in the literature. The magnitude of above mentioned changes were depth dependent. In terms of plant growth, Aksakal et al. (2016) reported the enhancing effect of increasing vermicompost treatment rate on bulk density for three soils with differing clay content. In general, mixing of soil with less dense organic material results in decreased particle density in soils amended with organic manures (Haynes & Naidu, 1998). However, their efficiency to improve bulk density for plant growth is related to the magnitude and quality of organic residues (Aksakal et al., 2016). Green manure which is largely decomposed off by leaving relatively smaller extent of organic matter have limited influence on soil physical attributes (Sauerbeck, 1982). Since FYM is more stable in terms of decomposition resistance than the GM, more organic compounds accumulated in soils (Table 2) treated with FYM. This in fact mediated the formation of aggregates resistant to soil traffic and therefore minimum bulk density was obtained in FYM plots. The main effect of soil depth irrespective to passing number and organic matter addition was significant (Table 3). Surface layer had smaller bulk density. The minimum bulk density (1.25 g cm^{-3}) was obtained in 0-10 cm depth for one pass in FYM treatments whereas the maximum one (1.61 g cm^{-3}) was recorded in 10-20 cm depth in C treatment with 5 passes (Fig. 2).

The depth of soil compaction in soil profile is dependent on the axle load, soil moisture content, tire size, contact pressure, traffic density, and soil organic matter content related attributes such as aggregation, aggregate stability, porosity, etc. (Hamza and Anderson, 2005). The greater axle loads and wetter soils consolidates the deeper in the soil profile. Since in our case these two dependents were taken constant, the effects of organic matter treatments are rather apparent. The stronger structures induced by larger amount and decomposition resistant organic substances scattered the force to a larger area which minimized the compaction-induced bulk density differences in FYM treated plots at any given pass number. At any steady

35

state condition in terms of organic matter addition each treatment may be regarded as fixed condition but with differing passes number such as in our case for any treatment, the increasing number of traffic density resulted in increases in the bulk density deeper in the profile (Fig. 2). Parras-Alcantra et al. (2015) similarly reported that organic farming comparing to conventional tillage significantly improved soil organic carbon stocks of soil which resulted in a decrease of soil bulk density in the soil profile as deep as 76.1 cm. Vice versa, organic matter loss upon conversion of forest soil to ag-lands have made soil progressively bulkier for vertisols and ultisols in 20 years (Bruun et al., 2015) which in turn explain the significance of organic matter to manage bulk density.




Figure 1. The effects number of passes on penetration resistance in different depths (passing×depth×treatment). Different letters written above columns indicate the difference in the treatment means at $P < 0.05$.




Figure 2. Number of passes and soil depth dependent bulk density changes induced by organic matter treatments (passing×treatment, depth×treatment; $P < 0.01$). Different letters written above columns indicate the difference in the treatment means at $P < 0.05$.


3.3. Porosity

The main effects of organic matter incorporation, depth and passes number on porosity were significant (Table 3). The effects of treatments in relation to soil depth and field traffic density on porosity in descending order were FYM>GM>C. The main overall effect of depth on total porosity was detrimental in 10-20 cm depth where a significant decrease was observed. The initial average ($0.565 \text{ cm}^3 \text{ cm}^{-3}$) of total porosity calculated from Table 2 showed a detrimental decrease down to 0.418 and $0.441 \text{ cm}^3 \text{ cm}^{-3}$ after 5 and 3 passes (Table 3). As all plots considered, the maximum and minimum values of total porosity were 0.53 and $0.39 \text{ cm}^3 \text{ cm}^{-3}$ respectively (Fig 3). The effects of organic matter treatments, depth and field traffic density on micro porosity was statistically significant (Table 3). Microporosity and total porosity parameters similarly responded to organic matter treatments. The volume of micropores were significantly higher in surface layer than the one observed in 10-20 cm (Table 3). Maximum micro porosity ($0.38 \text{ cm}^3 \text{ cm}^{-3}$) was recorded in FYM treatment in 0-10 cm depth with 3 passes whereas the minimum ($0.32 \text{ cm}^3 \text{ cm}^{-3}$) was determined in C treatment in the surface layer (0-10 cm) with 5 passes (Fig. 4). There was an increase in the micro porosity of control plot upon one pass comparing to its corresponding initial porosity, in contrast a decrease was recorded for annually FYM and GM incorporated plots. As three passes were done, it increased in all treatments including the control. Initial micro porosity value and the one after 5 passes were nearly the same in C treatment but they dropped below even the initial porosity for FYM and GM treatments. Despite the organic matter incorporation was made only four years the enhancement was recorded about porosity parameters. The enhancement induced by organic matter amendments in long term studies at various locations was even more astonishing and show similar manner to our findings. For example, Rasool et al. (2008) and Arthur et al. (2013) reported increases in total porosity and water retention which is related to pore nature of soil with increase in organic matter content depending on agricultural practices or organic matter amendments. The FYM and GM, to relatively smaller extent, promotes total and micro porosity in the current study. Organic matter application to soils promotes the development of better soil structure by binding the soil particles with polysaccharides and bacterial exudates which results in decreased bulk density and hence the porosity (Bhatia and Shukla, 1982). As the level of soil compaction increased, the amount of water held in high matric potentials decreases whereas the magnitude of water held at low matric potentials is to increase (Gupta et al., 1989) due to conversion of some macropores into micropores by compression stress. Similarly, Seker and Isildar (2000) reported an increase in the pore volume holding plant available water after 4 passes. The descending order of the treatments was FYM>GM>C for both of the depths and each pass-treatment and as pointed out by Celik et al. (2004) micro porosity increased upon organic matter amendments.

Macro porosity which is critical for soil aeration and soil water circulation were changed as a function of soil depth and field traffic density and organic matter amendments (Table 3). The main effect of organic matter was in descending order as FYM>GM>C. In this study, organic matter amendments significantly improved macro porosity ($P<0.01$). However, field traffic at 3 and 5 passes (Table 3) reduced the macropores volume below the critical level of $0.1 \text{ cm}^3 \text{ cm}^{-3}$ (Hakansson and Lipiec, 2000) and covered on the conditioner effects of organic amendments. The maximum macro porosity ($0.16 \text{ cm}^3 \text{ cm}^{-3}$)

was observed in 0-10 cm depth of FYM treatment at one pass whereas the minimum was recorded for C treatment in 10-20 cm depth at 5 passes (Fig 5).

Enhancement in soil structure traits by reduction of aggregate wettability (Zhang and Hartge, 1992) strength of aggregate stability with organic matter incorporation partially eliminated the effects of field traffic on macro porosity after four consecutive years of FYM and GM application. The organic matter bound-ambiguity was attributed to type of organic matter, C/N ratio and the degree of resistance to decomposition. Readily decomposable soil organic matter reported to be more relevant than total organic matter in mechanical characterization of the soil (Ball et al., 2000). For example the less humified organic matters, such as green manure, was reported to highly efficient to increase aggregate porosity (Zhang, 1994) but this effects was found to be not durable and resistant to field traffic as compared to the effects of FYM in the current study. Since, the overall effect of GM to increase soil organic matter are much lesser than the one maintained by FYM which more stable and resistant to microbial decay such behaviour is likely in organic matter poor soils.



15 Figure 3. Number of passes and soil depth dependent total porosity changes induced by organic matter treatments (passing×depth×treatment; $P < 0.01$). Different letters written above columns indicate the difference in the treatment means at $P < 0.05$.




Figure 4. Number of passes and soil depth dependent micro porosity changes induced by organic matter treatments (passing×depth; $P < 0.05$, depth×treatment; $P < 0.01$) Different letters written above columns indicate the difference in the treatment means at $P < 0.05$.

5




Figure 5. Number of passes and soil depth dependent macro porosity changes induced by organic matter treatments (passing× depth, depth×treatment, passing×treatment; $P < 0.01$). Different letters written above columns indicate the difference in the treatment means at $P < 0.05$.

10

4. Conclusions

- 15 The effect of field traffic density on soil compaction was found to be dependent to addition and type of organic matter treatment. The overall effects of organic treatments irrespective from soil depth on penetration resistance and bulk density were in descending order as $C > GM > FYM$ whereas it was $FYM > GM > C$ for total and micro porosity. Macro porosity appeared to be higher at minimum field traffic for FYM treatment in the surface layer. It can be concluded that the use of

organic matter enhance soil conditions by influencing the soil water holding and circulation characteristics, aeration, penetration resistance and bulk density which have implications on plant root growth.

References

5

Aggelides, S. M. and Londra, P.A.: Effects of compost produced from town wastes and sewage sludge on the physical properties of a loamy and a clay soil, *Biores. Technol.*, 71, 253-259, 2000.

Aksakal, E.L., Oztas, T., and Oztas, M.: Time-dependent changes in distribution patterns of soil bulk density and penetration resistance in a rangeland under overgrazing. *Turkish J. Agric. Forest.*, 35, 195–204, 2011.

10 Aksakal, E.L., Angin, I., and Oztas, T.: Effects of diatomite on soil physical properties. *Catena*, 88, 1–5, 2012.

Aksakal, E.L., Sari, S., Angin, I.: Effects of vermicompost application on soil aggregation and certain physical properties. *Land Degrad. Develop.*, 27 (4), 983-995, 2016.

Aksakal, E.L.: Soil compaction and its importance for agriculture. *Ataturk Univ. J. Fac. Agric.*, 35, 3-4, 247-252, 2004.

15 Alagoz, Z., Yilmaz, E., and Okturen, F.: Effects of organic material addition on some physical and chemical properties of soils, *Akdeniz Univ. J. Fac. Agric.*, 19(2), 245-254, 2006.

Al-Dousari, A. M., Misak, R., and Shahid, S.: Soil compaction and sealing in Al Salmi Area, Western Kuwait. *Land Degrad. Develop.*, 11, 401-418, 2000.

[Arslan, S.: An Alternative Method for Preventing Soil Compaction: Controlled Traffic. *KSU. Journal of Science and Engineering* 9\(1\), 2006.](#)

20 Arthur, E., Schjonning, P., Moldrup, P., Tuller, M., and de Jonge, L. W.: Density and permeability of a loess soil: Long-term organic matter effect and the response to compressive stress, *Geoderma*, 193-194, 236- 245, 2013.

Ball, B.C., Campbell, D.J., and Hunter, E.A.: Soil compactibility in relation to physical and organic properties at 156 sites in UK. *Soil Tillage Res.*, 57, 83–91, 2000.

25 [Barbero-Sierra, C., Marques, M.J., Ruiz-Pérez, M., Escadafal, R., Exbrayat, W.: How is Desertification Research Addressed in Spain? Land Versus Soil Approaches. *Land Degrad. Develop.*, 26 \(5\), 423-432, 2015.](#)

Bhatia, K.S. and Shukla, K.K.: Effect of continuous application of fertilizers and manure on some physical properties of eroded alluvial soil, *J. Indian Soc. Soil Sci.*, 30, 33–36, 1982.

Blake, G.R. and Hartge, K.H.: Bulk density, in: *Methods of Soil Analysis, Part 1, Physical and Mineralogical Methods*, (Ed: A. Klute), Agr. Monogr. 9, ASA and SSSA, Madison WI., USA, 363-375, 1986.

30 Blanco-Moure, N., Gracia, R., Bielsa, A. C., and López M. V.: Soil organic matter fractions as affected by tillage and soil texture under semiarid Mediterranean conditions. *Soil Tillage Res.*, 155 (Special Issue), 381-389, 2016.

Bouyoucos, G.J.: Hydrometer method improved for making particle size analyses of soils. *Agron. J.*, 54, 464-465, 1962.

- Bruun, T. B., Elberling, B., de Neergaard, A., and Magid, J.: Organic carbon dynamics in different soil types after conversion of forest to agriculture, *Land Degrad. Develop.*, 26 (3), 272-283, 2015.
- Burt, R. (Ed.): *Soil Survey Laboratory Methods Manual*, Soil Survey Investigations Rep. 42, version 4.0, USDA–NRCS, 2004.
- 5 Busscher, W.J. and Sojka, R.E.: Enhancement of subsoiling effect on soil strength by conservation tillage, *Transac. Am. Soc. Agric. Engin.*, 30, 4, 888-892, 1987.
- Carman, K.: Tractor forward velocity and tine load effects on soil strength, *J. Terramech.*, 31, 11-20, 1994.
- Carter, M.R.: Soil quality for sustainable land management: organic matter and aggregation interactions that maintain soil functions, *Agron. J.*, 94, 38–47, 2002.
- 10 Celik, I., Gunal, H., Budak, M., and Akpinar C.: Effects of long-term organic and mineral fertilizers on bulk density and penetration resistance in semi-arid Mediterranean soil conditions, *Geoderma*, 160, 236-243, 2010.
- Celik, I., Ortas, I., and Kilic, S.: Effects of compost, mycorrhiza, manure and fertilizer on some physical properties of a Chromoxerert soil. *Soil Tillage Res.*, 78, 59–67, 2004.
- Cerdà A.: Aggregate stability against water forces under different climates on agriculture land and scrubland in southern
- 15 Bolivia. *Soil Tillage Res.*, 57, 159–166, 2000.
- Chaudhari, P.R., V. Ahire, D.V., Ahire, V.D., Chkravarty, M., and Maity, S.: Soil bulk density as related to soil texture, organic matter content and available total nutrients of Coimbatore soil. *Int. J. Sci. Res. Pub.*, 3(2), 1-8, 2013.
- Cochrane, H.R. and Aylmore, L.A.G.: The effects of plant roots on soil structure, in: *Proceedings of 3rd Triennial Conference “Soils 94”*, 207–212, 1994.
- 20 Czyz, E.A. and Dexter, A.R.: Estimation of the density of the clay-organic complexes in soil. *Int. Agrophys.*, 30 (1), 19-23, 2016.
- Danielson, R.E. and Sutherland, P.L.: Porosity, in: *Methods of Soil Analysis, Part 1, Physical and Mineralogical Methods*. 2nd ed. (Ed: A. Klute) *Agron. Monogr. 9*. ASA and SSSA, Madison WI., USA, 443-461, 1986.
- De Moraes, M. T., Debiassi, H., Carlesso, R., Franchini, J.C., and Da Silva, V.R.: Critical limits of soil penetration resistance
- 25 in a rhodic eutrudox, *R. Bras. Ci. Solo*, 38, 288-298, 2014.
- De Oliveira, S. P., de Lacerda, N. B., Blum, S. C., Escobar, M. E. O., and de Oliveira, T. S.: Organic carbon and nitrogen stocks in soils of northeastern Brazil converted to irrigated agriculture, *Land Degrad. Develop.*, 26 (1), 9-21, 2015.
- Erbach, D.: Farm equipment and soil compaction. *ASAE Paper No: 860730*, 1986.
- Gelaw, A. M., Singh, B. R., Lal R.: Organic carbon and nitrogen associated with soil aggregates and particle sizes under
- 30 different land uses in Tigray, Northern Ethiopia, *Land Degrad. Develop.*, 26 (7), 690-700, 2015.
- Gulser, C. and Candemir, F.: Changes in penetration resistance of a clay field with organic waste applications. *Eurasian J. Soil Sci.*, 1, 16–21, 2012.
- Gupta, S.C., Sharma, P.P., and DeFranchi, S.A.; Compaction effects on soil structure, in: *Advances in Agronomy*, Ed. N.C. Brady, 42, 311-338, 1989.

- Hakansson, I. and Lipiec, J.: A review of the usefulness of relative bulk density values in studies of soil structure and compaction. *Soil Tillage Res.*, 53, 71-85, 2000.
- Hamza, M.A. and Anderson, W.K.: Responses soil properties and grain yields to deep ripping and gypsum application in a compacted loamy sand soil contrasted with a sandy clay loam soil in Western Australia. *Australian Journal of Agricultural Res.* 54, 273-282, 2003.
- Hamza, M.A. and Anderson, W.K.: Soil compaction in cropping systems: a review of the nature, causes and possible solutions. *Soil Tillage Res.*, 82, 121–145, 2005.
- Hamza, M.A. and Anderson, W.K.: Combinations of ripping depth and tine spacing for compacted sandy and clayey soils. *Soil & Tillage Res.*, 99, 213–220, 2008.
- Haynes, R.J. and Naidu, R.: Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. *Nutr. Cyc. Agroecosys.*, 51, 123–137, 1998.
- Kacar, B.: *Soil Analysis*, Nobel publications, Ankara (in Turkish), 2009.
- Kaleem Abbasi M., Mahmood Tahir M., Sabir N., Khurshid M. Impact of the addition of different plant residues on nitrogen mineralization-immobilization turnover and carbon content of a soil incubated under laboratory conditions. *Solid Earth*, 6 (1), 197-205, 2015.
- Keesstra, S.D., Bouma, J., Wallinga, J., Tittonell, P., Smith, P., Cerdà, A., Montanarella, L., Quinton, J.N., Pachepsky, Y., van der Putten, W.H., Bardgett, R.D., Moolenaar, S., Mol, G., Jansen, B., and Fresco, L.O.: The significance of soils and soil science towards realization of the United Nations Sustainable Development Goals. *Soil*, 2, 111–128, 2016.
- Kozłowski, T.T.: *Soil Compaction and Growth of Woody Plants*. *Scand. J. For. Res.* 14, 596-619, 1999.
- Mamman, E., Ohu, J. O., and Crowther, T.: Effect of soil compaction and organic matter on the early growth of maize (*Zea mays*) in a vertisol, *Int. Agrophys.*, 21, 367-375, 2007.
- McGarry, D.: Tillage and soil compaction, in: *Conservation Agriculture*, L. Garcia-Torres et al. (eds.), 307-316, 2003.
- Minitab.: *Minitab 16 Statistical Software*, Minitab Inc., State College, Pennsylvania, USA, 2010.
- Muñoz-Rojas, M., Jordán, A., Zavala, L. M., De la Rosa D., Abd-Elmabod, S. K., and Anaya-Romero, M.: Impact of land use and land cover changes on organic carbon stocks in mediterranean soils (1956-2007). *Land Degrad. Develop.*, 26 (2), 168-179, 2015.
- Munsuz, N.: *Soil Mechanics and Technology*, Publications of Ankara Univ. Faculty of Agriculture, 922, Text book, 260, Ankara (in Turkish), 1985.
- Ni, J., Luo, D.H., Xia, J., Zhang, Z.H., and Hu, G.: Vegetation in karst terrain of southwestern China allocates more biomass to roots. *Solid Earth*, 6 (3), 799-810, 2015.
- Novara A., Gristina L., Guaitoli F., Santoro A., Cerdà A. Managing soil nitrate with cover crops and buffer strips in Sicilian vineyards. *Solid Earth*, 4 (2), 255-262, 2013.

- Ola, A., Dodd, I.C., and Quinton, J.N.: Can we manipulate root system architecture to control soil erosion? *Soil* 1, 603-612, 2015.
- Parras-Alcántara, L., Díaz-Jaimes, L., and Lozano-García, B.: Organic farming affects C and N in soils under olive groves in mediterranean areas. *Land Degrad. Develop.*, 26 (8), 800-806, 2015.
- 5 Patel, S.K. and Mani, I.: Effect of multiple passes of tractor with varying normal load on subsoil compaction. *J. Terramech.*, 48, 277–284, 2011.
- Peng F., Quangang Y., Xue X., Guo J., Wang T. Effects of rodent-induced land degradation on ecosystem carbon fluxes in an alpine meadow in the Qinghai-Tibet Plateau, China. *Solid Earth*, 6 (1), 303-310, 2015.
- Pulido, M., Schnabel, S., Contador, J. F. L., Lozano-Parra, J., and González, F.: The impact of heavy grazing on soil quality and pasture production in rangelands of SW Spain. *Land Degrad. Develop.*, DOI: 10.1002/ldr.2501, 2016.
- 10 Rasool, R., Kukal, S. S., and Hira, G. S.: Soil organic carbon and physical properties as affected by long-term application of FYM and inorganic fertilizers in maize-wheat system, *Soil Tillage Res.*, 101 (1-2), 31-36, 2008.
- Sauerbeck, D.R.: Influence of crop rotation, manurial treatment and soil tillage on the organic matter content of German soils. In: *Soil Degradation*, Boels D, Davies DB & Johnston AE (eds), Proceedings of the EEC Seminar held in Wageningen, Rotterdam, A A Balkema, Netherlands, pp 163–179, 1982.
- 15 Seker, C. and Isildar, A.A.: Effects of wheel traffic porosity and compaction of soil profile. *Turk. J. Agric. Forest.*, 24, 71–77, 2000.
- Shaw, E.A., Deneff, K., Milano de Tomasel, C., Cotrufo, M.F., and Wall, D.H.: Fire affects root decomposition, soil food web structure, and carbon flow in tallgrass prairie. *Soil*, 2, 199-210, 2016.
- 20 Sparovek, G., Lambais, M.R., Silva, A.P., and Tormena, C.A.: Earthworm (*Pontoscolex corethrurus*) and organic matter effects on the reclamation of an eroded oxisol. *Pedobiologia*, 43, 698–704, 1999.
- Thomas, G.W., Haszler, G.R., and Blevins, R.I.: The effect of organic matter and tillage on maximum compactibility of soils using the proctor test. *Soil Sci.*, 161, 502–508, 1996.
- USDA, Soil Survey Manual, Soil Survey Division Staff, Washington, DC, USA, 1993.
- 25 USDA-NRCS: Soil Compaction. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1187272.pdf. 27.05.2016, 2012.
- Xin, X., Zhang, J., Zhu, A., and Zhang, C.: Effects of long-term (23 years) mineral fertilizer and compost application on physical properties of fluvo-aquic soil in the North China Plain. *Soil Tillage Res.*, 156, 166–172, 2016.
- Wang, T., Xue, X., Zhou, L., Guo, J.: Combating Aeolian Desertification in Northern China. *Land Degrad. Develop.*, 26 (2), 118-132, 2015.
- 30 Yan, X., and Cai, Y.L.: Multi-Scale Anthropogenic Driving Forces of Karst Rocky Desertification in Southwest China. *Land Degrad. Develop.*, 26 (2), 193-200, 2015.
- Zhang, H. and Hartge, K.H.: Effect of differently humified organic matter on the aggregate stability by reducing aggregate wettability. *Z. Pflanzenernahr. Bodenkd.*, 155, 143–149, 1992.

Zhang, H.: Organic matter incorporation affects mechanical properties of soil aggregates. *Soil Tillage Res.*, 31, 263–275, 1994.