

Mid/Late Devonian-Carboniferous collapse basins on the Finnmark Platform and in the southwesternmost Nordkapp basin, SW Barents Sea

5 Jean-Baptiste P. Koehl^{1,2}, Steffen G. Bergh^{1,2}, Tormod Henningsen¹, Jan Inge Faleide^{2,3}

¹Department of Geosciences, University of Tromsø, N-9037 Tromsø, Norway.

²Research Centre for Arctic Petroleum Exploration (ARCEX), University of Tromsø, N-9037 Tromsø, Norway.

³Department of Geosciences, University of Oslo, P.O. Box 1047 Blindern, NO-0316 Oslo, Norway.

Correspondence to: Jean-Baptiste Koehl (jean-baptiste.koehl@uit.no)

10
Abstract. The SW Barents Sea margin experienced a pulse of extensional deformation in the Middle-Late Devonian through the Carboniferous, after the Caledonian Orogeny terminated. These events marked the initial stages of formation of major offshore basins such as the Hammerfest and Nordkapp basins. We mapped and analyzed three major fault complexes, i) the Måsøy Fault
15 Complex, ii) the Rolvsøya fault, iii) the Troms-Finnmark Fault Complex. We discuss the formation of the Måsøy Fault Complex as a possible extensional splay of an overall NE-SW trending, NW-dipping, basement-seated Caledonian shear zone, the Sørøya-Ingøya shear zone, which was partly inverted during the collapse of the Caledonides and accommodated top-to-the-NW normal displacement in Mid/Late Devonian-Carboniferous times. The Troms-Finnmark Fault Complex
20 displays a zigzag-shaped pattern of NNE-SSW and ENE-WSW trending extensional faults before it terminates to the north as a WNW-ESE trending, NE-dipping normal fault that separates the southwesternmost Nordkapp basin in the northeast from the Finnmark Platform west and the Gjesvær Low in the southwest. The WNW-ESE trending, margin-oblique segment of the Troms-Finnmark Fault Complex is considered to represent the offshore prolongation of a major
25 Neoproterozoic fault complex, the Trollfjorden-Komagelva Fault Zone, which is made of WNW-ESE trending, subvertical faults that crop out on the island of Magerøya in NW Finnmark. Our results suggest that the Trollfjorden-Komagelva Fault Zone dies out to the northwest before reaching the Finnmark Platform west. We propose an alternative model for the origin of the WNW-ESE trending fault segment of the Troms-Finnmark Fault Complex as a possible hard-linked,
30 accommodation cross-fault that developed along the Sørøya-Ingøya shear zone. This brittle fault

decoupled the Finnmark Platform west from the southwesternmost Nordkapp basin and merged with the Måsøy Fault Complex in Carboniferous times. Seismic data over the Gjesvær Low and southwesternmost Nordkapp basin show that the low-gravity anomaly observed in these areas may result from the presence of Mid/Late Devonian sedimentary units resembling Middle Devonian, spoon-shaped, late/post-orogenic collapse basins in western and mid Norway. We propose a model for the formation of the southwesternmost Nordkapp basin and its counterpart Devonian basin in the Gjesvær Low by exhumation of narrow, ENE-WSW to NE-SW trending basement ridges along a bowed portion of the Sørøya-Ingøya shear zone in the Mid/Late Devonian-early Carboniferous. Exhumation may have involved part of a large-scale metamorphic core complex that potentially included the Lofoten Ridge, the West Troms Basement Complex and the Norsel High. Finally, we argue that the Sørøya-Ingøya shear zone truncated and decapitated the Trollfjorden-Komagelva Fault Zone during the Caledonian Orogeny and that the western continuation of the Trollfjorden-Komagelva Fault Zone was mostly eroded and potentially partly preserved in basement highs in the SW Barents Sea.

45

1. Introduction

The SW Barents Sea margin is located near the Iapetus suture zone that formed when Laurentia collided with Fennoscandia to produce the Caledonian Orogeny (Ramberg et al., 2008; Gernigon et al., 2014). This suture and possibly related deep-seated shear zones, which accommodated e.g. thrust nappe emplacement during the Caledonian Orogeny, are now covered by late Paleozoic to Cenozoic sedimentary basins that formed during multiple episodes of extension. These repeated extension events led to the breakup of the North Atlantic Ocean and formation of a transform plate margin at the boundary between the Mid-Norwegian and SW Barents Sea margins (Faleide et al., 1993, 2008; Blystad et al., 1995; Doré et al., 1997; Bergh et al., 2007; Hansen et al., 2012; Gernigon et al., 2014). The rift-margin along the SW Barents Sea, offshore Western Troms and NW Finnmark (Figure 1), consists of the Finnmark Platform and an adjacent, glacial sediment-free strandflat, and of deep offshore basins such as the Hammerfest and Nordkapp basins (Gabrielsen et al., 1990). These basins are bounded by major NE-SW trending extensional faults such as the Troms-Finnmark Fault Complex (TFFC; Gabrielsen et al., 1990; Smelror et al., 2009; Indrevær et al., 2013), the Måsøy Fault Complex (MFC; Gabrielsen et al., 1990; Gudlaugsson et al., 1998), and potential basement-seated ductile detachments (Figure 1).

60

The study area also includes a deep Paleozoic basin that is located southwest of the Nordkapp Basin and east of the Hammerfest Basin, and which is bounded to the southwest by the WNW-ESE trending segment of the TFFC and to the southeast by the MFC (Figure 1).

65 The SW Barents Sea margin off Western Troms and NW Finnmark is segmented by margin-oblique, NNW-SSE to WNW-ESE trending transfer fault zones, e.g. Senja Fracture Zone and Fugløya transfer zone (Indrevær et al., 2013), which may represent analogs of the onshore, Neoproterozoic, WNW-ESE trending Trollfjorden-Komagelva Fault Zone (TKFZ) in eastern Finnmark (Siedlecki, 1980; Herrevold et al., 2009) and to the Kokelv Fault on the Porsanger
70 Peninsula (Figure 1; Gayer et al., 1985; Lippard & Roberts, 1987; Rice, 2013). The TKFZ is believed to continue farther west, off the coast, where it is thought to interact with and merge into the WNW-ESE trending fault segment of the TFFC (Gabrielsen, 1984; Vorren et al., 1986; Townsend, 1987; Gabrielsen & Færseth, 1989; Gabrielsen et al., 1990; Roberts et al., 2011; Bergø, 2015; Lea, 2015). Onshore-nearshore, margin-parallel fault complexes include the Langfjord-
75 Vargsund fault (LVF; Figure 1) trending NE-SW and possibly representing an analog to the TFFC and MFC. The geometric interaction, timing and controlling effects of the TFFC, MFC, TKFZ and LVF, and adjacent offshore basins and ridges are not yet resolved. In particular, the presence of potential Caledonian structures in the deeper portion of the Finnmark Platform, e.g. in the footwall of the TFFC (cf. Johansen et al., 1994; Gudlaugsson et al., 1998) is further explored in the present
80 contribution.

 The goal of this paper is to contribute to the understanding of tectonic and sedimentary processes in the Arctic in the Late Devonian-Carboniferous. To achieve this, we demonstrate the presence of an overall NE-SW trending, NW-dipping, basement-seated, low-angle shear zone on the Finnmark Platform, the Sørøya-Ingøya shear zone (SISZ; Figure 1), and to discuss its role
85 played in shaping the SW Barents Sea margin during late/post-orogenic collapse of the Caledonides in late Paleozoic times and its influence on the formation and evolution of Devonian-Carboniferous collapse basins. We mapped and analyzed basin-bounding brittle faults on the Finnmark Platform and in the southwesternmost Nordkapp basin (named the easternmost Hammerfest basin in Omosanya et al., 2015), such as the TFFC and the MFC (Figure 1), to evaluate the impact of the
90 SISZ on post-Caledonian brittle faults. We aim at showing the importance of structural inheritance by examining the relationship between Precambrian-Caledonian structural grains, post-Caledonian fault trends and offshore sedimentary basin geometries. Minor Carboniferous grabens and half-

grabens on the Finnmark Platform (e.g. the Sørvær Basin; Figure 1), which are thought to have formed during early stages of extension shortly after the end of the Caledonian Orogeny (Lippard & Roberts, 1987; Olesen et al., 1990; Johansen et al., 1994; Bugge et al., 1995; Gudlaugsson et al., 1998; Roberts et al., 2011), are of particular importance to the present work. We further investigate the presence of possible Devonian sedimentary deposits on the Finnmark Platform and in the southwesternmost Nordkapp basin and tentatively interpret them as potential analogs to Middle Devonian basins in western Norway (Séranne et al., 1989; Chauvet & Séranne, 1994; Osmundsen & Andresen, 2001) and mid-Norway (Braathen et al., 2000). In this context, NE-SW to ENE-WSW trending basement ridges in the footwall of the TFFC and on the northern flank of the southwesternmost Nordkapp basin are described and analyzed, and we compare them to adjacent basement highs such as the Norsel High (Figure 1; Gabrielsen et al., 1990; Gudlaugsson et al., 1998), the West Troms Basement Complex (Zwaan, 1995; Bergh et al., 2010) and the Lofoten Ridge (Blystad et al., 1995; Bergh et al., 2007; Hansen et al., 2012). Finally, we propose a model of exhumation of these ENE-WSW to NE-SW trending basement ridges as a metamorphic core complex (cf. Lister & Davis, 1989) using shear zones in Lofoten-Vesterålen as onshore analogs for the SISZ (Steltenpohl et al., 2004; Osmundsen et al., 2005; Steltenpohl et al., 2011).

110 2. Geological setting

The bedrock geology of the SW Barents Sea margin (Figure 1) consists of (i) an Archean to Paleoproterozoic basement suite, the West Troms Basement Complex (Zwaan, 1995; Bergh et al., 2010), (ii) locally preserved autochthonous Neoproterozoic cover sequences (Kirkland et al., 2008), (iii) a series of Caledonian thrust nappes (Andersen, 1981; Ramsay et al., 1985; Corfu et al., 2014), and (iv) late Paleozoic to Cenozoic sedimentary sequences offshore (Faleide et al., 1993, 2008; Gudlaugsson et al., 1998; Worsley, 2008; Smelror et al., 2009; Figure 1). Archean to Paleoproterozoic basement rocks are mostly exposed in major horsts and ridges in Western Troms (Bergh et al., 2010; Indrevær et al., 2013; Indrevær & Bergh, 2014), whereas Neoproterozoic and Caledonian rocks dominate in the eastern part of Troms and in NW Finnmark (Kirkland et al., 2008; Corfu et al., 2014; Indrevær & Bergh, 2014; Figure 1). In offshore areas adjacent to Western Troms and NW Finnmark, extensive post-Caledonian normal faulting led to the formation of large sedimentary basins that are filled with thick, late Paleozoic to Cenozoic deposits related to the post-orogenic collapse of the Caledonides and to the opening of the NE Atlantic Ocean (Faleide et al.,

1993, 2008; Gudlaugsson et al., 1998; Worsley, 2008; Smelror et al., 2009). Late Paleozoic-
125 Cenozoic sedimentary units are missing in onshore areas of Troms and Finnmark likely due to
erosion and/or non-deposition (Ramberg et al., 2008; Smelror et al., 2009).

2.1. Onshore Precambrian and Caledonian geology

130 2.1.1. Precambrian basement rocks

The Western Troms margin is characterized by Archean to Paleoproterozoic basement
rocks of the West Troms Basement Complex (Bergh et al., 2010) that are preserved and exposed
in a horst block formed during post-Caledonian extension (Indrevær et al., 2013). The West Troms
Basement Complex consists of tonalitic, trondhjemitic and granitic gneisses, metasupracrustal
135 rocks and mafic and felsic igneous rocks (Corfu et al., 2003; Bergh et al., 2010). These rocks were
deformed during the Svecofennian orogeny, which resulted in the formation of NW-SE trending
steep foliation, ductile shear zones and upright and vertical macrofolds, only weakly reworked
during the Caledonian Orogeny (Corfu et al., 2003; Bergh et al., 2010).

In NW Finnmark, Paleoproterozoic basement rocks occur in several tectonic windows of
140 the Caledonides, e.g. Repparfjord-Komagfjord and Alta-Kvænangen tectonic windows (Zwaan &
Gautier, 1980; Pharaoh et al., 1982; 1983; Bergh & Torske, 1988; Figure 1), and consist of low-
grade supracrustal metavolcanics and metasedimentary rocks of the Raipas Group. These
Greenstone belts formed as NW-SE trending rift basins in the Archean?-Paleoproterozoic during
the opening of the Kola Ocean (Bergh & Torske, 1986; 1988), although more recent studies
145 tentatively reinterpret these rocks as foreland basin deposits derived from the Svecokarelian
Orogeny (Torske & Bergh, 2004). A thin cover of Neoproterozoic to Cambrian (para-)
autochthonous metasedimentary rocks occurs on top of Paleoproterozoic basement rocks in
Finnmark (Siedlecki, 1980; Ramsay et al., 1985; Andresen et al. 2014; Corfu et al., 2014). Other
Neoproterozoic-Ordovician units in eastern Finnmark include metasedimentary rocks of the
150 Barents Sea and Tanafjorden-Varangerfjorden regions (Siedlecki, 1980; Siedlecka & Roberts,
1992) which are exposed on the Varanger Peninsula (Figure 1).

The Timanian Orogeny produced major NW-SE trending folds (Roberts & Siedlecka,
2002) and WNW-ESE trending fault complexes like the TKFZ (Jonhson et al., 1978; Herrevold et
al., 2009). The TKFZ was mapped as a narrow, single-segment fault strand all the way along the

155 Kola Peninsula in Russia in the east, where it merges with the Sredni-Rybachy Fault Zone (Roberts
et al., 1997; Roberts et al., 2011), to the Barents shelf in the west (Gabrielsen, 1984; Gabrielsen &
Færseth, 1989; Gabrielsen et al., 1990; Roberts et al. 2011). We present an alternative model in
which the TKFZ splays into multiple fault segments and dies out between the Varanger Peninsula
and the Barents shelf. On the Varanger Peninsula, the TKFZ is well displayed on satellite and DEM
160 images, but is generally poorly exposed. In map view, the TKFZ is irregular, with different
structural segments and branching subsidiary faults both across- and along-strike, locally showing
duplex structures (Siedlecka & Siedlecki, 1967; Siedlecka, 1975). The TKFZ formed along the
southwestern boundary of the Timanian Orogeny in the late Cryogenian-Ediacaran (Roberts &
Siedlecka, 2002; Siedlecka et al., 2004), and was later reactivated as a strike-slip fault during the
165 Caledonian Orogeny when it accommodated significant lateral displacement constrained to 200-
250 km of dextral strike-slip movement (Bylund, 1994; Rice, 2013).

2.1.2. Caledonian nappes

Coastal areas of NW Finnmark are dominated by Caledonian thrust sheets of the Kalak
170 Nappe Complex and Magerøy Nappe (Ramsay et al., 1985; Ramberg et al., 2008; Corfu et al.,
2014), formed in the Neoproterozoic through Silurian (Figure 1). The Kalak Nappe Complex is
composed of amphibolite facies schists, metapsammities and paragneisses, and comprises several
allochthonous thrust sheets with Proterozoic basement rocks, clastic metasedimentary rocks, and
plutonic rocks of the Seiland Igneous Province (Corfu et al., 2014). A major thrust defines the
175 contact with the underlying pre-Caledonian basement (Ramsey et al., 1985). Dominant structures
include a gently NW-dipping foliation, NNE-SSW trending, east-verging, asymmetrical recumbent
folds and low-angle thrusts that accommodated top-to-the-ESE shortening (Townsend, 1987a;
Kirkland et al., 2005). The Kalak Nappe Complex was previously considered to represent an exotic
terrane accreted on the Laurentian margin of Rodinia prior to the rifting of the Iapetus Ocean, and
180 to have later been thrust over Baltica during the Caledonian Orogeny (Kirkland et al., 2008).
However, paleocurrent and geochronological data suggest these rocks to be of Baltican origin
(Roberts, 2007; Zhang et al., 2016).

The Seiland Igneous Province corresponds to a large, late Neoproterozoic mafic and
ultramafic intrusion linked to the early-mid rifting stages of the Iapetus Ocean (Elvevold et al.,
185 1994; Corfu et al., 2014). Recent geophysical studies by Pastore et al. (2016) show that the base of

the Seiland Igneous Province defines two deep-reaching roots located below the islands of Seiland and Sørøya constraining the thickness of the Kalak Nappe Complex in this area to a maximum of 10 km. On the Porsanger and Varanger Peninsula, ENE-WSW to NNE-SSW trending, Ediacaran metadolerite dyke swarms are particularly common, and they are as well associated to the rifting
190 of the Iapetus Ocean (cf. Roberts, 1972; Siedlecka et al., 2004; Nasuti et al., 2015).

The Kalak Nappe Complex is structurally overlain by the Magerøy Nappe, which consists of Late Ordovician to early Silurian greenschist facies metasedimentary and metaplutonic rocks (Andersen, 1981; 1984; Corfu et al., 2014) that crop out on the island of Magerøya (Figure 1). The Magerøy Nappe is characterized by asymmetrical, NNE-SSW trending, east-verging folds and low-angle, NW- and SE-dipping thrusts similar in trend to those observed within the Kalak Nappe
195 Complex (Andersen, 1981), and is intruded by granitic and gabbroic plutons, e.g. the Silurian Honningsvåg Igneous Complex (Corfu et al., 2006) and the Finnvik Granite (Andersen, 1981). Remnants of the Magerøy Nappe thrust units are also found in northeastern Sørøya and on the Porsanger Peninsula (Kirkland et al., 2005; 2007; Corfu et al., 2014; Figure 1).

200

2.2. Post-Caledonian brittle faults and basins

2.2.1. Post-Caledonian offshore basins

The SW Barents Sea margin was subjected to multiple episodes of extensional faulting after
205 the end of the Caledonian Orogeny, starting with the collapse of the Caledonides in the Mid/Late Devonian-early Carboniferous, lasting until the early/mid Permian, although evidence of this stage is only preserved onshore western and mid-Norway (Séranne et al., 1989; Chauvet & Séranne, 1994; Braathen et al., 2000; Osmundsen & Andresen, 2001). During this period, basement ridges in Lofoten-Vesterålen (Klein & Steltenpohl, 1999; Klein et al., 1999; Steltenpohl et al., 2004; 2011;
210 Figure 1) and in mid-Norway (Osmundsen et al., 2005; Figure 1) were exhumed as metamorphic core complexes, synchronously with the development of large half-graben basins such as the Vøring and Møre basins in mid-Norway (Blystad et al., 1995) and the Hammerfest, Nordkapp and Ottar basins in the SW Barents Sea (Gabrielsen et al., 1990; Breivik et al., 1995; Gudlaugsson et al., 1998; Indrevær et al., 2013; Figure 1). The main rifting events occurred in the Late Jurassic and
215 peaked in the Early Cretaceous, when major offshore basins such as the Tromsø and Harstad basins formed. The rifting ended with full breakup of the North Atlantic Ocean and formation of a

transform plate margin in the SW Barents Sea at the Paleocene-Eocene transition (Faleide et al., 1993; 2008).

Off the coasts of Western Troms and NW Finnmark, the SW Barents Sea margin is
220 characterized by a relatively shallow area, the Finnmark Platform (Gabrielsen et al., 1990; Figure
1), which is thought to have remained relatively stable since late Paleozoic times. For example, the
inner part of the Finnmark Platform, here referred to as the Finnmark Platform east (Figure 1), was
only affected by the formation of minor Carboniferous, ENE-WSW to NE-SW trending half-
graben and graben structures (Bugge et al., 1995; Samuelsen et al., 2003; Rafaelsen et al., 2008;
225 Figure 1). In the hanging-wall of the MFC on the western part of the Finnmark Platform, the
Finnmark Platform west (Figure 1), shows a prominent gravity low, the Gjesvær Low, which was
ascribed to the presence of low-density Caledonian rocks (Johansen et al., 1994; Gernigon et al.,
2014). We explore and argue for an alternative explanation, i.e. the presence of Devonian collapse
basin deposits draped against a low-angle extensional detachment of the SISZ, similar to the
230 Nordfjord-Sogn Detachment Zone, a late-orogenic shear zone that bounds the Middle Devonian
Hornelen, Kvamshesten and Solund sedimentary basins onshore western Norway (Séranne et al.,
1989; Chauvet & Séranne, 1994; Wilks & Cuthbert, 1994; Osmundsen & Andersen, 2001). Ductile
detachment surfaces of comparable size, showing analog kinematics and contemporaneous timing
of activity as the Nordfjord-Sogn Detachment Zone are documented as far north as the Lofoten-
235 Vesterålen Margin (Klein & Steltenpohl, 1999; Klein et al., 1999; Steltenpohl et al., 2004; 2011),
but Devonian collapse basin sedimentary rocks and extensional detachments have not yet been
reported along the margins of Western Troms and NW Finnmark.

2.2.2. Post-Caledonian faults

240 Multiple studies have reported post-Caledonian brittle faults onshore coastal areas in
Lofoten-Vesterålen, Western Troms and NW Finnmark (Roberts, 1971; Worthing, 1984; Lippard
& Roberts, 1987; Townsend, 1987a; Rykkelid, 1992; Lippard & Prestvik, 1997; Roberts &
Lippard, 2005; Bergh et al., 2007; Hansen et al., 2012; Indrevær et al., 2013; Davids et al., 2013).
A common feature is the presence of rhombic, zigzag-shaped fault trends similar in geometry to
245 offshore basin-bounding faults. Dominant fault-fracture trends of the margin strike NNE-SSW,
ENE-WSW and NW-SE, respectively (Bergh et al., 2007; Eig, 2008; Eig & Bergh, 2011; Hansen
et al., 2012; Hansen & Bergh, 2012; Indrevær et al., 2013). Typical examples are basin-bounding,

250 NNE-SSW and ENE-WSW trending brittle normal faults that are part of the Vestfjorden-Vanna Fault Complex, which bounds the offshore Vestfjorden Basin southeast of the Lofoten islands and which can be traced northward to Western Troms (Indrevær et al., 2013; Figure 1), whereas the NNW-SSE to WNW-ESE trend typically reflects margin-oblique, transform fault trends (Faleide et al., 2008). An analog to the onshore Vestfjord-Vanna fault complex in NW Finnmark is the Langfjorden-Vargsundet fault (Figure 1), described by Zwaan & Roberts (1978) and Worthing (1984) as a major NE-SW trending, NW-dipping normal fault where rocks from the Kalak Nappe
255 Complex and the Seiland Igneous Province in the northwest are juxtaposed against Precambrian basement rocks of the Repparfjord-Komagfjord and Alta-Kvænangen tectonic windows in the southeast (Figure 1).

The NW Finnmark margin is located along the northeastward prolongation of the Lofoten-Vesterålen and Western Troms segments of the Norwegian continental shelf (Figure 1). Similar
260 fault sets and trends as in Lofoten-Vesterålen exist in Finnmark and their interaction is thought to partly have controlled the rhombic geometry of many offshore sedimentary basins (Bergh et al., 2007; Indrevær et al., 2013). A typical example along the Western Troms and NW Finnmark margins is the NW-dipping TFFC, which bounds the Harstad Basin to the east and the Hammerfest Basin to the southeast (Gabrielsen et al., 1990; Indrevær et al., 2013). The TFFC defines a system
265 of irregular branching faults trending NNE-SSW and ENE-WSW and terminating as a WNW-ESE trending fault zone northwest of the island of Magerøya where it merges with the NE-SW trending, NW-dipping MFC at the southeastern boundary of the Nordkapp Basin (Gabrielsen et al., 1990) and of the triangular-shaped southwesternmost Nordkapp basin (Omosanya et al., 2015; Figure 1). We address a possible genetic relationship and structural inheritance of the post-Caledonian MFC
270 with the Caledonian SISZ and argue that the MFC may have initiated as an extensional splay during the reactivation of the SISZ as an extensional detachment during the late/post-orogenic collapse of the Caledonides. Furthermore, we tentatively link basement ridges such as the Norsel High in the footwall of the Nysleppen Fault Complex (Gabrielsen et al., 1990) to bowed segments of the SISZ (Figure 1).

275

2.2.3. Post-Caledonian transfer zones

The Norwegian continental shelf is segmented by transfer fault zones of which the largest is the offshore De Geer Zone (Faleide et al., 1984; 2008; Cianfarra & Salvini, 2015), which main

280 fault segment is the Hornsund Fault Zone, an offshore NNW-SSE trending fault that runs parallel
to the west coast of Spitsbergen and separates the SW Barents Sea margin from the Lofoten-
Vesterålen Margin (Figure 1). In the south, the De Geer Zone proceeds through the Senja Fracture
Zone and into the Senja Shear Belt onshore the island of Senja (Figure 1). Olesen et al. (1993;
1997) suggested shifts of polarity of the Vestfjorden-Vanna Fault Complex along the Senja
Fracture Zone, and they argued that the formation of the Senja Fracture Zone offshore was
285 controlled by a major onshore basement weakness zone, the Bothnian-Senja Fault Complex (Figure
1), which provided suitably oriented basement heterogeneities for the development of a transfer
zone (e.g. Doré et al., 1997). Similarly, Indrevær et al. (2013) proposed the existence of a fault
array termed the Fugløya transfer zone to explain offsets and shifts of polarity along the
Vestfjorden-Vanna Fault Complex farther northeast in Western Troms (Figure 1). The Fugløya
290 transfer zone trends N-S/NNW-SSE and continues onshore Western Troms, where it merges with
the NW-SE trending Bothnian-Kvænangen Fault Complex, and offshore where it is thought to
merge into the TFFC and the Ringvassøy-Loppa Fault Complex (Indrevær et al., 2013; Figure 1).

Analogously in NW Finnmark, the WNW-ESE trending TKFZ seems to merge into a basin-
bounding fault, in this case the WNW-ESE trending, NE-dipping fault segment of the TFFC
295 (Gabrielsen, 1984; Gabrielsen & Færseth, 1989; Roberts et al., 2011). In nearshore areas of NW
Finnmark, the TKFZ is thought to proceed offshore and seems to correlate with a large escarpment
north of Magerøya and into the Barents Sea (Vorren et al., 1986; Townsend, 1987b). In the area
where it terminates, it merges and links up with the TFFC to form triangular-shaped mini-basins
(Gabrielsen, 1984; Gabrielsen & Færseth, 1989; Roberts et al., 2011). We explore an alternative
300 origin for the WNW-ESE trending fault segment of the TFFC and further examine its interaction
with the onshore-nearshore TKFZ, which potentially acted as a transfer fault after the Caledonian
Orogeny and contributed to offset the LVF onshore Magerøya and in adjacent coastal areas (Koehl
et al., submitted; Figure 1). Other major WNW-ESE trending faults exist offshore, northeast of the
Varanger Peninsula, and these bound the Tiddlybanken Basin, a large WNW-ESE trending basin
305 that formed in Carboniferous times (Mattingsdal et al., 2015; Figure 1).

2.2.4. Absolute age dating of post-Caledonian faulting

The absolute age of post-Caledonian brittle faults onshore NW Finnmark is poorly
constrained although a few contributions provide valid insights (Lippard & Prestvik, 1997; Davids

310 et al., 2013; Torgersen et al., 2014; Koehl et al., 2016). Torgersen et al. (2014) performed K/Ar
dating of brittle fault gouge in the footwall of the LVF and obtained dominantly Carboniferous to
early Permian ages, as well as a subsidiary Early Cretaceous age for one of the faults. Roberts et
al. (1991) and Lippard & Prestvik (1997) presented indirect evidence of early Carboniferous
dolerite dykes emplaced along and cementing WNW-ESE trending brittle fault segments of the
315 TKFZ onshore Magerøya, thus providing a minimum estimate for the latest stage of faulting along
this fault. These dykes produce high positive aeromagnetic anomalies (Nasuti et al., 2015) and may
be used to further identify brittle faults in NW Finnmark. Late Devonian dolerite dykes emplaced
along brittle faults that trend NE-SW and N-S have been identified and dated in eastern Varanger
Peninsula (Guise & Roberts, 2002) and on the Kola Peninsula (Roberts & Onstott, 1995). By
320 comparison, Davids et al. (2013) obtained Late Devonian-early Carboniferous ages from K/Ar
dating of illite clay minerals for early extensional faulting along the Vestfjorden-Vanna Fault
Complex and related faults in Lofoten-Vesterålen and Western Troms.

2.3. Offshore sedimentary successions and well-ties

325
Deep fault-bounded basins formed along the SW Barents Sea margin during successive
extension events in late Paleozoic-early Cenozoic times, and these basins contain important
sedimentary successions for hydrocarbon exploration. We particularly focus on the late Paleozoic
succession (Figure 3) which sedimentary rocks were deposited on top of eroded Precambrian and
330 Caledonian basement rocks (cf. Townsend, 1987a; Johansen et al., 1994; Bugge et al., 1995;
Zwaan, 1995; Gudlaugsson et al., 1998; Samuelsberg et al., 2003; Bergh et al., 2010). Late
Paleozoic sedimentary deposits in the study area were penetrated by only a few exploration wells
to which we tied our seismic interpretation (Figure 2). Overlying Mesozoic to Cenozoic
sedimentary units were not investigated and are better described in Omosanya et al. (2015).

335
The nature and age of basement rocks along the SW Barents Sea margin remain relatively
complex to resolve since only a handful of wells drilled through the thick post-Caledonian
sedimentary cover. Nevertheless, wells 7128/4-1 and 7128/6-1 penetrated quartzitic
metasedimentary rocks on the Finnmark Platform east (Figure 2) and these are believed to correlate
with upper Proterozoic rocks involved in Caledonian thrusting in northern Finnmark (Røe &
340 Roberts, 1992).

Devonian sedimentary rocks are yet to be reported in North Norway and along the SW Barents Sea margin (Figure 3). However, Devonian sedimentary deposits are present in western Norway (Osmundsen & Andersen, 2001) where they represent a several km-thick succession made up with clastic deposits that notably include rhythmic sandstone and coarse-grained conglomerate units. These were deposited in the hanging-wall of major, low-angle extensional shear zones, e.g. the Nordfjord-Sogn Detachment Zone (Séranne et al., 1989; Wilks & Cuthbert, 1994; Osmundsen & Andersen, 2001).

Lower Carboniferous sedimentary rocks of the Billefjorden Group directly overlie basement rocks on the Finnmark Platform east as evidenced by exploration wells 7128/4-1 and 7128/6-1 (Larssen et al., 2002; Figure 2 & Figure 3). These rocks mostly correspond to fluvial clastic deposits interbedded with coal-bearing sedimentary rocks that correlate with contemporaneous deposits onshore Bjørnøya (Cutbill & Challinor, 1965; Gjelberg, 1981, 1984) and Spitsbergen (Cutbill & Challinor, 1965; Cutbill et al., 1976; Gjelberg, 1984). The total thickness of Billefjorden Group sedimentary deposits evidenced by exploration wells on horst-blocks on the Finnmark Platform east ranges from 350 m to 450 m. However, in the hanging-wall of a minor normal fault interpreted by Bugge et al. (1995) near the coast of northern Finnmark (Figure 2), shallow drill-cores 7127/10-U-2 and 7127/10-U-3 indicate that the thickness of lower Carboniferous sedimentary rocks reaches a thickness > 600 m within a NE-SW trending mini-basin on the Finnmark Platform east near the coast of the Nordkinn Peninsula (cf. star symbol in Figure 1 & Figure 2). In the Serpukhovian, fluvial sediments of the Billefjorden Group were gradually replaced by shallow marine sediments of the Gipsdalen Group from which they are generally separated by a mid-Carboniferous (Serpukhovian) unconformity (Cutbill et al., 1976; Gjelberg, 1984; Bugge et al., 1995) potentially related to a global sea-level fall (Saunders & Ramsbottom, 1986).

Shallow marine sedimentary deposits of the Gipsdalen Group are widespread along the SW Barents Sea margin and have proven prolific for hydrocarbon exploration (Larssen et al., 2002; Figure 3). Thus, this sedimentary succession benefits from a relatively high number of well penetrations and, as a result, its lateral facies and thickness variations are well-constrained (Gjelberg & Steel, 1981, 1983; Samuelsen et al., 2003; Rafaelsen et al., 2008). The Gipsdalen Group was notably penetrated by wells 7128/4-1 and 7128/6-1 on the Finnmark Platform east, by well 7120/12-4 on the Finnmark Platform west and by well 7124/3-1 on the northern flank of

southwesternmost Nordkapp basin (Larssen et al., 2002; Figure 2). This succession consists of alluvial clastic sedimentary rocks that are progressively replaced upwards by shallow marine platform carbonates interbedded with clastic and evaporite deposits (McCann & Dallmann, 1996).
375 In well 7124/3-1 (Figure 2), Asselian evaporite deposits typically include thin layers of anhydrite and gypsum, but thicker, halite-rich end-members are found along the flanks of the Nordkapp Basin and southwesternmost Nordkapp basin where large pillows of upper Carboniferous-lower Permian salt were observed (Gabrielsen et al., 1992; Jensen & Sørensen, 1992; Koyi et al., 1993; Nilsen et al., 1995; Gudlaugsson et al., 1998; Koehl et al., 2017). In the Nordkapp Basin, pre-Permian
380 deposits may in places reach a thickness of up to 7-8 km (Gudlaugsson et al., 1998). These deposits are composed of thick clastic sedimentary rocks and of upper Carboniferous to lower Permian evaporite deposits characterized by mobile salt that was involved in salt tectonism in the southwesternmost Nordkapp basin (Gudlaugsson et al., 1998; Koehl et al. 2017) and in the Nordkapp Basin (Gabrielsen et al., 1992; Jensen & Sørensen, 1992; Koyi et al., 1993; Nilsen et al.,
385 1995).

3. Methods & databases

3.1. Seismic data and well-ties

390 The seismic interpretation shown in this study is based on publicly available 2D and 3D data from the DISKOS database, thus providing reasonably tight 2D data coverage. However, only one seismic 3D survey was available in the study area. The interpretation of seismic data aims at providing good constraints for the extent and geometry of offshore brittle faults and for offshore stratigraphy on the Finnmark Platform and in the southwesternmost Nordkapp basin. The present
395 study uses ties to wells 7120/12-4, 7128/4-1 and 7128/6-1 and 7124/3-1 based on publicly available well data (www.npd.no) and private well-tie seismograms and to shallow drill-cores 7127/10-U-2 and 7127/10-U-3 from Bugge et al. (1995; Figure 2). Seven seismic profiles from the BSS01 2D seismic survey were used to analyze and describe offshore basin and fault geometries and provide
400 the basis for discussion about the late Paleozoic evolution of the SW Barents Sea margin. Note that none of the seismic profiles used was depth-converted. Therefore, all relevant estimates of fault offsets and stratigraphic seismic unit thicknesses will be described in seconds (s) two-way time

(TWT). In addition, we analyzed two time-slice from 3D seismic survey MC3D-MFZ02 to constrain fault interaction in map-view.

405

3.2. Aeromagnetic anomaly data

The offshore aeromagnetic data used in this study correspond to a compilation of the BASAR project of the Geological Survey of Norway (NGU) published by Gernigon & Brønner (2012) and Gernigon et al. (2014; Figure 4). The dataset is composed of tilt derivatives of aeromagnetic data and has been used to delineate possible magmatic intrusions (dykes) emplaced along brittle faults (cf. Nasuti et al., 2015) and abrupt changes of lithology generally recorded across major faults, thus, contributing to the mapping of post-Caledonian offshore brittle faults along the SW Barents Sea margin. However, data uncertainties arise from the fact that significantly different rock types may yield similar aeromagnetic responses. A crucial example in northern Finnmark is the similar high positive narrow aeromagnetic anomalies produced both by sub-vertical folded beds of metasedimentary rocks (Roberts & Siedlecka, 2012; Roberts & Williams, 2013) and dolerite dykes intruded along brittle faults (Nasuti et al., 2015; Figure 4). In order to distinguish such features, we carefully analyzed onshore geology in coastal areas of NW Finnmark and the results of exploration wells on the Finnmark Platform and adjacent offshore basins.

415

420

4. Results

4.1. Seismic interpretation of offshore basins and faults

425

4.1.1. Seismic units and stratigraphy

On seismic data (Figure 5), basement rocks typically show chaotic internal reflection patterns, which complicate the task of identifying intra-basement structures and basins, and individualize layered sedimentary sequences. However, km-thick layers bearing strong basement fabrics such as widespread gently dipping foliation or pronounced mylonitic fabric commonly found along large shear zones may turn out to be resolvable at seismic scale (see chapter 4.1.2.; Fountain et al., 1984; Reeve et al., 2013; Phillips et al., 2016; Fazlikhani et al., 2017). For instance, we observed a several km-thick, curved, shallow-dipping layer that is characterized by moderate-amplitude reflections, which are parallel to the layer's upper and lower boundaries (see "Sørøya-

430

435 Ingøya shear zone” reflections in Figure 5c-g). We interpret these pronounced internal fabrics as
widespread mylonitic foliation separated by internal thrusts within a large-scale shear zone.
Numerous smaller basement shear zones may be present below late Paleozoic-Cenozoic
sedimentary rocks on the Finnmark Platform west, and these correspond to steeply to moderately
dipping fabrics made of sub-parallel, moderate- to high-amplitude reflections (cf. Figure 5b, e, f, g
440 & Figure 6a-c).

Potential Devonian sedimentary deposits along the SW Barents Sea are sparse and as a
result their seismic character is not well constrained (Figure 3). This sedimentary succession has
not been drilled, which makes its interpretation on seismic data rather speculative. However, we
believe that the best two candidates to represent Devonian sedimentary deposits analog to those in
445 western and mid-Norway (Braathen et al., 2000; Osmundsen & Andersen, 2001; Fazlikhani et al.,
2017) are located at the base of the southwesternmost Nordkapp basin and on the Finnmark
Platform west near the Gjesvær Low (Figure 1). In the southwesternmost Nordkapp basin, possible
Devonian sedimentary strata are located at a deep level (below 4 seconds TWT) and their seismic
signature is thus largely masked by overlying sedimentary successions (Figure 5c & d). By contrast,
450 on the Finnmark Platform west (Figure 5e), potential Devonian sedimentary rocks are relatively
shallower, which makes their seismic pattern easier to distinguish from underlying basement rocks,
and from overlying Carboniferous sedimentary deposits and seismic artifacts (Figure 5e).
Devonian sedimentary rocks on the Finnmark Platform west display relatively low seismic
amplitudes, partly similar to analog deposits in the North Sea (cf. seismic facies 1 in Fazlikhani et
455 al., 2017). The internal reflection pattern is rather chaotic apart from a few discrete, shallow-
dipping, moderate-amplitude reflections that converge towards each other upwards and that we
interpret as major sedimentary sequence boundaries (cf. dotted white reflections in Figure 5e &
Figure 6b & c). Furthermore, Devonian sedimentary deposits are likely separated from underlying
basement rocks by an angular unconformity that appears as arcuate, high-amplitude seismic
460 reflections (“Base Devonian” reflection in Figure 5e and Figure 6b & c). We interpret these arcuate,
high-amplitude seismic reflections as an erosional unconformity.

Lower Carboniferous sedimentary deposits of the Billefjorden Group, composed of thick
clastic sedimentary deposits interbedded with occasional coal-bearing sedimentary rocks (Figure
3), may produce high-amplitude seismic reflections related to their organic-rich content (Figure 5a
& b). Such sedimentary strata are present on the Finnmark Platform east, where they appear to
465

thicken to the southeast near the coasts of NW Finnmark (Figure 6d), whereas they are rather sparse on the Finnmark Platform west, i.e. eroded or never deposited (Figure 5e & f). On the Finnmark Platform east, the transition from basement rocks (cf. “Top basement” reflection in Figure 5a & b) to lower Carboniferous sedimentary rocks is difficult to interpret on seismic sections. This is
470 attributable to the strong similarities between high seismic amplitudes displayed locally both by basement rock fabrics such as major shear zones (cf. yellow dotted lines in Figure 5b) and by lower Carboniferous coal-bearing sedimentary deposits. Low amplitude reflections also show identical chaotic patterns in both basement rocks and clastic sedimentary rocks of the Billefjorden Group (Figure 5a & b). In the southwesternmost Nordkapp basin, lower Carboniferous sedimentary strata
475 are believed to be present although their seismic signature certainly appears to be affected by overlying upper Carboniferous evaporite deposits (Figure 5c & d). The boundary between lower Carboniferous sedimentary deposits and potential underlying Devonian sedimentary rocks was not identified in the southwesternmost Nordkapp basin. Nevertheless, the maximum thickness of Billefjorden Group sedimentary strata on the Finnmark Platform east is ca. 600 m (Bugge et al.,
480 1995), and this suggests that the several km-thick succession below the mid-Carboniferous reflection and above a thick shear zone in the southwesternmost Nordkapp basin is composed of lower Carboniferous sedimentary rocks probably complemented by thick Devonian sedimentary deposits (Figure 5c & d). Alternatively, sedimentary deposits of the Billefjorden Group directly overlie basement rocks.

485 On the Finnmark Platform (Figure 1 & Figure 2), the base of the upper Carboniferous sedimentary succession is difficult to identify (cf. “mid-Carboniferous” reflection in Figure 3 & Figure 5). In places, it appears as a linear, moderate to low amplitude seismic reflection that separates subparallel reflections of lower and upper Carboniferous sedimentary rocks, whereas in other places the reflection is irregular and truncates high-amplitude coal-bearing sedimentary
490 deposits of the Billefjorden Group, and/or high-amplitude reflections produced by basement rocks (Figure 6a), and/or low-amplitude reflections in Devonian sedimentary strata (Figure 6b & c). Nevertheless, this reflection generally corresponds to an angular unconformity (e.g. Figure 6a-c & e) and is therefore interpreted to correspond to a regional erosion surface.

495 In the southwesternmost Nordkapp basin, the base of upper Carboniferous sedimentary deposits (cf. “mid-Carboniferous” reflection in Figure 3 and Figure 5c & d) appears as a clear, discrete high-amplitude reflection. The strong acoustic impedance contrast producing the high

seismic amplitude for the mid-Carboniferous reflection most likely arises from the presence of upper Carboniferous evaporite deposits partly composed of mobile salt (halite), which is significantly less dense than regular sedimentary rocks, (cf. “Top upper Carboniferous evaporites” reflection in Figure 3 and Figure 5c & d). This evaporite succession was identified by Gudlaugsson et al. (1998) and is restricted to basinal areas located northwest of the MFC and north of the TFFC (Figure 1 & Figure 2). It is characterized by a highly variable thickness, which is due to the presence of lensoidal bodies bounded to the top and bottom by high-amplitude reflections on the basin edges and to the occurrence of thick bodies made of chaotic reflection patterns near the center of the basin (Figure 5c). We interpret the lensoidal bodies on the basin edges as pillows of mobile salt and the chaotic bodies near the basin center as small salt diapirs based on similarities with large salt diapirs and evaporite deposits observed in the Nordkapp Basin (Gabrielsen et al., 1992; Jensen & Sørensen, 1992; Koyi et al., 1993; Nilsen et al., 1995). We consider that the presence of analog, late Paleozoic evaporite deposits in the southwesternmost Nordkapp basin and in the Nordkapp Basin (Jensen & Sørensen, 1992; Koyi et al., 1993; Gudlaugsson et al., 1998) and the absence of such deposits in the Hammerfest Basin constitute strong arguments to justify a change of name for the “easternmost Hammerfest basin” (Omosanya et al., 2015) into the “southwesternmost Nordkapp basin”. However, this basin shows large amount of normal displacement along its southern boundary fault, the NW-dipping MFC, which is opposite to the Nordkapp Basin where basin subsidence was dominantly accommodated along the SE-dipping Nysleppen Fault Complex (Figure 1). Hence, despite their similarities, the Nordkapp Basin and the southwesternmost Nordkapp basin should be treated as two separate basins.

Non-evaporitic, upper Carboniferous and Permian sedimentary deposits are characterized by subparallel, flat-lying to shallow-dipping, homogeneous, moderate to low-amplitude seismic reflections (see Figure 5). Permian deposits are relatively thin on the Finnmark Platform and are sometimes difficult to distinguish from upper Carboniferous deposits (Figure 5a, b, e, f & g). In the southwesternmost Nordkapp basin, however, late Paleozoic sedimentary deposits are thicker and individual units are therefore easier to identify at seismic scale. Thus, we interpreted a thin unit characterized by high-amplitude reflections (cf. “Base Asselian” and “Top Asselian evaporites” reflections in Figure 3 and Figure 5c & d) as Asselian (earliest Permian) evaporite deposits that were evidenced by exploration well 7124/3-1 on the northern flank of the southwesternmost Nordkapp basin (Figure 2, Figure 5c & d). Where present, this thin Asselian evaporite succession

530 defines the base of the Permian sedimentary succession and therefore serves as an upper boundary for the Carboniferous succession (cf. “Base Asselian” reflection in Figure 5c & d). However, Asselian evaporites are too thin and too discontinuous to be seismically resolvable on the Finnmark Platform (Bugge et al., 1995). Occasionally, Asselian evaporites are truncated by chaotic reflections of small salt diapirs sourced from deeper upper Carboniferous evaporites in the southwesternmost Nordkapp basin (Figure 5c).

535 The Base Triassic reflection (see Figure 5) defines the (near-) top of the late Paleozoic sedimentary succession and is easily interpreted through the whole Barents Sea as it corresponds to a high-amplitude reflection that represents the top of a regionally widespread carbonate unit (Bugge et al., 1995). Other important seismic reflections interpreted in the present study include the Base Cretaceous, Base Paleocene, the Upper Regional Unconformity (URU), which corresponds to a major erosional unconformity and represents the base of Quaternary sediment 540 cover (Solheim & Kristoffersen, 1984), and the seabed reflection (Figure 5). These reflections are penetrated by a large number of exploration wells and shallow drill-cores both on the Finnmark Platform and in the southwesternmost Nordkapp basin, where they all display consistently high seismic amplitudes (Faleide et al., 1984; Bugge et al., 1995; Gudlaugsson et al., 1998; Omosanya et al., 2015).

545

4.1.2. Structural architecture of the Finnmark Platform and of the southwesternmost Nordkapp basin

In this section, we describe the most important structural elements of the Finnmark Platform and of the southwesternmost Nordkapp basin (cf. Figure 1 & Figure 2) based on interpreted key 550 seismic sections (Figure 5). We also highlight the most dominant fault trends and their interaction with major structures such as the TFFC, MFC, TKFZ and SISZ to form offshore sedimentary basins.

Faults and shear zones within basement rocks

555 We identified a several km-thick, curved (in cross-section), shallow-dipping layer of moderate-amplitude reflections that we interpreted to represent a large-scale basement-seated shear zone, which we name the SISZ. The upper boundary surface of the SISZ (Figure 7) appears to be relatively shallow in coastal areas on the Finnmark Platform west. On the Finnmark Platform west,

the SISZ dominantly dips to the NW but switches to a dominant dip to the northeast on the
560 Finnmark Platform east. In the footwall of the MFC and in the southwestern part of the Finnmark
Platform west, the SISZ occurs at relatively shallow depth (< 1.5 s TWT). There it is believed to
have been deeply eroded and is now overlain by a very thin sedimentary cover (cf. Figure 5c-f &
Figure 6d). The SISZ shows significant lateral thickness variations that range from 2.0-2.5 seconds
(TWT) near the coastline and in the footwall of the TFFC to 0.5 second (TWT) below the MFC
565 and the TFFC (Figure 5f). The SISZ deepens to the northwest towards the center of the Finnmark
Platform west before bending upwards in the footwall of the TFFC (Figure 5e & f). The SISZ then
curves down where the listric TFFC merges with the shear zone at depth, thus delineating an
elongated, NE-SW trending ridge in the footwall of the TFFC (cf. “basement ridges” in Figure 1
and Figure 5e & f). A similar pattern is observed in the southwesternmost Nordkapp basin where
570 the SISZ deepens to the northwest before curving up near the center of the basin and merging with
the N-boundary fault of the southwesternmost Nordkapp basin, the Rolvsøya fault, hence giving
this basin a characteristic “U” shape in cross-section (Figure 5c & d). The SISZ also curves down
in the footwall of the Rolvsøya fault and defines a second elongated, ENE-WSW trending ridge (cf.
“basement highs” in Figure 1). Importantly, the two basement ridges located in the footwall of the
575 TFFC and of the Rolvsøya fault (“basement highs” in Figure 1), respectively, are separated by a
narrow trough that is bounded to the southwest by the WNW-ESE trending segment of the TFFC
(Figure 7). Apart from this narrow trough, the attitude of the SISZ is uniform along NE-SW
transects on the Finnmark Platform west and within the southwesternmost Nordkapp basin with a
gentle dip to the northeast (Figure 5g).

580 Notably, the spoon-shaped geometry of the SISZ, with asymmetric, NE-SW trend,
northeastwards-broadening, NE-plunge (Figure 7) appears to coincide with a basement gravity low
on the Finnmark Platform west: the Gjesvær Low (Johansen et al., 1994; Gernigon et al., 2014;
Figure 1) The geometry of the SISZ also matches the trend and shape of the southwesternmost
Nordkapp basin (Figure 1 & Figure 7). Farther south, along the coasts of Western Troms and
585 westwards below the Hammerfest Basin, the low quality of available seismic data did not allow us
to trace the SISZ more precisely (Figure 7). On the Finnmark Platform east, the SISZ bends from
NE-SW into a more WNW-ESE trend and changes dip from gentle to steep to the northeast (Figure
7), and as a result, the SISZ becomes too deep to interpret on seismic data in the northeastern part

of the Finnmark Platform east (Figure 7). The multiple changes of trend, dip direction, dip angle
590 and thickness of the SISZ gives the shear zone a spoon-shaped geometry (Figure 7).

On the Finnmark Platform west, subsidiary steep, SE-dipping high-amplitude reflections
occur in basement rocks and these are truncated by the mid-Carboniferous reflection and the Base
Devonian erosional unconformity in the footwall of the TFFC (see yellow dotted lines in Figure
5e-g). Despite dipping southeast, these reflections resemble the dominant reflection pattern
595 observed within the SISZ (Figure 5e & f). Thus, we interpret them as SE-dipping, mylonitic shear
zones (yellow dotted lines in Figure 5e-g). The upper boundary of one of these SE-dipping shear
zones coincides with an abrupt seismic facies change on the Finnmark Platform west, from
moderately dipping, moderate-amplitude reflections in the west to gently dipping/sub-horizontal,
low-amplitude seismic reflections in the east (Figure 5g). This change also coincides with a ca. one
600 second (TWT) deepening of the upper boundary of the SISZ towards the northeast (Figure 5g), and
with a small normal offset of a lensoidal, eastwards-thickening layer of sub-horizontal reflections
located above the SISZ (cf. dotted black lines in Figure 5g). We interpret these changing attributes
to be related to the presence of a NNE-SSW trending, ESE-dipping brittle fault that flattens and
soles into the SISZ and which may have developed along a pre-existing, steep ductile shear zone
605 (yellow dotted lines in Figure 5g).

Similar NE-SW trending but NW-dipping shear zones may exist in basement rocks on the
Finnmark Platform east, for example in the form of steeply dipping, high-amplitude seismic
reflections truncated by the mid-Carboniferous reflection (cf. yellow dotted lines in Figure 5b and
Figure 6a). These reflections differ from gently dipping, high-amplitude reflections of lower
610 Carboniferous coal-bearing sedimentary deposits (Figure 6a) and rather resemble the SISZ
reflection pattern, though these are located well above the presumed continuation of the SISZ
(Figure 5e & f). We therefore interpret these steep reflections as a NE-SW trending, NW-dipping
shear zone similar to the SISZ (Figure 5b).

615 *Faults within late Paleozoic sedimentary successions*

Faults bounding Paleozoic sedimentary strata and basins include the major TFFC and MFC
and numerous faults on the Finnmark Platform. The TFFC is made of alternating ENE-WSW and
NNE-SSW trending, NW-dipping, listric fault segments that form a zigzag pattern and that separate
the Hammerfest Basin in the northwest from the Finnmark Platform west in the southeast (Figure

620 1 & Figure 5e & f; Gabrielsen et al., 1990; Indrevær et al., 2013). Seismic data below ENE-WSW and NNE-SSW trending fault segments of the TFFC show that these fault segments merge with and sole into shallow-dipping reflections of the SISZ at depth (Figure 5e & f). At the northeast termination of the Hammerfest Basin, the TFFC bends 90 degrees clockwise and continues to the southeast as a WNW-ESE trending, NE-dipping, listric fault (Figure 1, Figure 2 & Figure 5g). At 625 depth, this fault merges with the SISZ (cf. Figure 5g) near a narrow trough in the top surface of the SISZ, separating two elongated NE-SW to ENE-WSW trending basement ridges in the footwall of the TFFC and of the Rolvsøya fault (cf. “basement highs” in red in Figure 1 & Figure 7). In map-view, the WNW-ESE trending, NE-dipping fault segment of the TFFC bends anticlockwise into the main fault segment of the MFC, which corresponds to a linear, NE-SW trending, NW-dipping 630 fault (Figure 1, Figure 2 and Figure 8a & b). The interaction of these two faults in map-view gives the Finnmark Platform west and the southwesternmost Nordkapp basin triangular shapes (Figure 2 & Figure 8a & b). The main fault segment of the MFC defines the southeastern boundary of the southwesternmost Nordkapp basin (Figure 1, Figure 2 & Figure 5c & d) and of a ca. 25-30 km wide graben structure on the Finnmark Platform west that is believed to be partly filled with 635 Devonian sedimentary deposits (Figure 1, Figure 2 & Figure 5e & f). Northeastwards, the main segment of the MFC (Figure 5c-f) is replaced by several minor fault segments of the MFC with limited vertical throw (Figure 5a & b) that defines the southeastern boundary of the Nordkapp Basin (Figure 1 and Figure 5a & b). The southwesternmost Nordkapp basin is bounded to the north by an E-W to ENE-WSW trending, south-dipping, listric normal fault, the Rolvsøya fault, which 640 flattens at depth and merges into gently dipping reflections of the SISZ (Figure 5c & d). The Rolvsøya fault separates the southwesternmost Nordkapp basin from the Ottar Basin to the northwest and from the Nordkapp Basin to the northeast (Figure 1 & Figure 2).

Late Paleozoic grabens on the Finnmark Platform east display fault patterns that are analogous to those that shape the southwesternmost Nordkapp basin and the Finnmark Platform 645 west (Figure 1 & Figure 2). Numerous steeply dipping, listric normal faults made of alternating, zigzag-shaped, ENE-WSW and NNE-SSW trending fault segments bound relatively narrow, few km-wide graben and half-graben structures that are filled with wedge-shaped, late Paleozoic sedimentary successions (Figure 2 & Figure 5a & b). In particular, one of these zigzag-shaped faults trends NE-SW to NNE-SSW, dips to the northwest and can be traced for about 60 km from 650 the northern coast of Magerøya onto the Finnmark Platform east (Figure 1 & Figure 2).

Southwestwards, this fault roughly aligns with a similarly shaped and oriented, NW-dipping onshore-nearshore fault complex syntetic to the TFFC described as the LVF (Figure 2 & Figure 5a & b; Zwaan & Roberts, 1978; Lippard & Roberts, 1987; Roberts & Lippard, 2005; Koehl et al., submitted). We tentatively interpret the ca. 60 km-long, zigzag-shaped brittle fault on the Finnmark Platform east, northeast of Magerøya, as the northeastward continuation of the LVF on the
655 Finnmark Platform east (Figure 1, Figure 5a & b and Figure 6a).

Below the minor northern fault segments of the MFC, we identified a large NE-SW trending, SE-dipping fault that is antithetic to the MFC (Figure 5a & b). Due to the rather low quality of seismic data at large depths, the interaction of the northern fault segments of the MFC with the antithetic SE-dipping fault is difficult to evaluate. Our data indicate that the northern fault segments of the MFC crosscuts the NE-SW trending, SE-dipping fault in the southwest (Figure 5b), whereas farther northeast, along strike, the northern fault segments of the MFC seem to sole into upper Carboniferous evaporite deposits (Figure 5a).
660

665 **4.1.3. Fault-controlled thickness variations**

In the following chapter, fault offsets and thickness variations in the sedimentary successions across brittle faults will be described as a basis to infer timing and sense of shear for brittle faults on the Finnmark Platform and in the southwesternmost Nordkapp basin. Regional stratigraphic thickness maps (Figure 9a-c) show that late Paleozoic sedimentary strata on the Finnmark Platform east thicken from < 0.1 second (TWT) in the southeast to a maximum thickness of ca. 2 seconds (TWT) in the footwall of the MFC (see also Figure 5a & b). This gradual thickness increase contrasts with the abrupt thickness increase of Devonian-Carboniferous sedimentary strata in the hanging-wall of major normal faults, e.g. the WNW-ESE trending segment of the TFFC and the main segment of the MFC (Figure 9a-b), thus separating depositional versus tectonic thickness changes.
670
675

Intra-basement thickness changes

The dominant shear zone system within basement rocks on the Finnmark Platform west is the SISZ (Figure 5c-g, Figure 6b-c & e-f and Figure 7). A pronounced intra-basement unit made of sub-horizontal, high-amplitude reflections occurs above the SISZ (Figure 5g). The top reflection of the SISZ and the overlying intra-basement unit are offset by a NNE-SSW trending, gently east-
680

dipping fault, which is accompanied by thickness increase of the intra-basement unit across the east-dipping fault (cf. black dotted line in Figure 5g & Figure 6e). This fault is interpreted to have a top-to-the-E, normal sense of shear (cf. dotted black lines in Figure 5g & Figure 6e), and is itself truncated by the subhorizontal mid-Carboniferous reflection, which constrains its activity to the Mid/Late Devonian-early Carboniferous (Figure 5g).

Fault-controlled thickness changes in Devonian-Carboniferous strata

In the southwesternmost Nordkapp basin, the Devonian-lower Carboniferous sedimentary succession (Figure 5c & d) appears to be thickest at the intersection of the TFFC and MFC (Figure 9a), where vertical displacement along the MFC and TFFC is estimated to be ca. 1.5 second (TWT), based on offset of the mid-Carboniferous reflection (cf. Figure 5d). The overlying upper Carboniferous succession displays a similar attitude as shown by the broad thickening of similar sedimentary strata at the intersection of the TFFC and MFC (Figure 9b). These observations suggest that the WNW-ESE trending segment of the TFFC and the main segment of the MFC potentially formed simultaneously in Devonian times and acted as syn-sedimentary normal faults that contributed to the thickening of Devonian-lower Carboniferous and upper Carboniferous sedimentary deposits within the southwesternmost Nordkapp basin (Figure 5c & d). In this scenario, the Rolvsøya fault likely limits the extent of thickened Devonian-lower Carboniferous and upper Carboniferous sedimentary strata to the north. If we consider the thickness of the seismic package limited upwards by the mid Carboniferous reflection and downwards by the top reflection of the SISZ in the footwall of the Rolvsøya fault, the maximum thickness of Devonian and lower Carboniferous sedimentary rocks on the northern flank of the basin does not exceed ca. 1 second (TWT). This thickness estimate is significantly thinner than what is observed within the southwesternmost Nordkapp basin, where the Devonian-lower Carboniferous succession reaches a maximum thickness of ca. 2-2.5 seconds (TWT; see Figure 5c & d). By analogy, the thickness of upper Carboniferous sedimentary strata on the northern flank of the southwesternmost Nordkapp basin decreases from ca. 1.5 seconds (TWT) to ca. 0.5-1 second across the Rolvsøya fault (Figure 5c & d and Figure 9b). Hence, the Rolvsøya fault was active and largely contributed to sediment thickening within the southwesternmost Nordkapp basin during the Mid/Late Devonian-Carboniferous.

On the Finnmark Platform west, potential Devonian sedimentary rocks are characterized by low-amplitude, chaotic reflections within which we observed distinct, shallow-dipping, moderate-amplitude reflections that we interpreted as major sedimentary sequence boundaries (cf. white dotted lines in Figure 5e and Figure 6b & c). These shallow-dipping reflections diverge from each other downwards and define gently dipping, wedge-shaped layers of low-amplitude, chaotic reflections that thicken downwards against arcuate, high-amplitude basement reflections that represent an erosional unconformity (cf. “Base Devonian” reflection in Figure 5e), and to the northwest against an ENE-WSW trending, SE-dipping normal fault (Figure 5e and Figure 6b & c). We interpret these sedimentary sequences separated by shallow-dipping, moderate-amplitude reflections to represent growth strata deposited along an active ENE-WSW trending, SE-dipping normal fault, which is parallel to SE-dipping basement shear zones (Figure 5e and Figure 6b & c). In addition, the main fault segment of the MFC shows decreasing amount of vertical displacement to the southwest, accompanied by a simultaneous thickness decrease in the upper Carboniferous succession along strike (Figure 9b), before the MFC eventually dies out on the Finnmark Platform west (Figure 1, Figure 2 & Figure 5e & f). Analogously, upper Carboniferous sedimentary deposits on the Finnmark Platform west display a wedge shape that is thickest in the southeast, near the MFC, and gradually thins towards the TFFC in the northwest (Figure 5e & f, Figure 9b). This upper Carboniferous sedimentary wedge likely formed by syn-tectonic sedimentary growth along the main fault segment of the MFC.

On the Finnmark Platform east, the offshore portion of the LVF (cf. Figure 1 & Figure 5a & b) downthrows the mid-Carboniferous reflection by ca. 0.5 second (TWT) to the northwest (Figure 5b) and bounds a NE-SW trending graben structure filled with thickened lower Carboniferous and upper Carboniferous sedimentary strata (cf. Figure 5a & b). In this graben structure, the lower Carboniferous and upper Carboniferous sedimentary successions thicken against the LVF (Figure 5b), while thickness variations become negligible farther north where the LVF dies out (Figure 1 & Figure 5a). Consequently, similar thickness increases of lower Carboniferous and upper Carboniferous sedimentary strata elsewhere within graben and half-graben structures on the Finnmark Platform east suggest that syn-tectonic sediment deposition along the LVF and analog ENE-WSW to NNE-SSW trending faults mostly occurred in Carboniferous times. Furthermore, in the footwall of the northern fault segments of the MFC, we recorded anomalously thick upper Carboniferous succession (Figure 9b) with a thickness

comparable to what is observed within the southwesternmost Nordkapp basin (Figure 9b). This succession shows a half-ellipsoid shape in map-view with a NE-SW trending major axis parallel to the MFC (Figure 9b). We therefore argue that this thickness change on the Finnmark Platform east is the result of syn-tectonic sediment deposition in the hanging-wall of a NE-SW trending, SE-dipping fault antithetic to the MFC (Figure 5a & b). We suggest that the half-ellipsoid shape of the thickened upper Carboniferous sedimentary deposits on the Finnmark Platform east reflects large offset near the center of the SE-dipping fault, and decreasing vertical throw towards the fault-tips, a feature characterizing syn-sedimentary, rift-related normal faults (Figure 9b).

By contrast, depositional sediment wedges may as well occur on the Finnmark Platform east, and they differ from fault-controlled thickness changes. One example is the ca. 600 m-thick lower Carboniferous succession evidenced by shallow drilling between the Nordkinn Peninsula and Magerøya (cf. “star” symbol in Figure 1; Bugge et al., 1995), which we re-interpreted as a prograding Carboniferous sedimentary system (Figure 6d). The apparent thickening of the lower Carboniferous succession near the coasts of NW Finnmark is more likely to be related to sedimentary processes during the formation of large clinofolds in a prograding sedimentary system (Figure 6d) than to syn-tectonic deposition in the hanging-wall of a NE-SW trending, NW-dipping fault.

760

Fault-controlled thickness changes in Permian strata

In the southwesternmost Nordkapp basin and on the Finnmark Platform east and west, the Permian sedimentary succession is thin and shows a relatively constant thickness compared to the underlying Devonian-lower Carboniferous and upper Carboniferous successions (Figure 5a-d and Figure 9a-c). However, the Base Asselian and Base Triassic reflections marking the lower and upper boundary of the Permian succession show some offsets across the main fault segment of the MFC, WNW-ESE trending segment of the TFFC and Rolvsøya fault, thus accounting for minor thickness variations in the Permian succession across these faults (Figure 5c, d & g and Figure 9c). We interpret these small offsets and thickness variations as the product of minor faulting activity in the Permian and mild Mesozoic reactivation of these faults, thus suggesting that the main tectonic activity along these faults was essentially restricted to the Mid/Late Devonian-late Carboniferous (Figure 5c, d & g). Moreover, on the Finnmark Platform west and east, most brittle

770

faults die out within the upper Carboniferous succession and only a few faults crosscut the Permian succession with limited amount of offset (Figure 5a, b, e, & f).

775

Fault-controlled thickness changes in Mesozoic-Cenozoic strata

Most faults observed within the late Paleozoic succession on the Finnmark Platform east and west and in the southwesternmost Nordkapp basin die out in the upper part of the succession before reaching the Base Triassic reflection (Figure 5). A few exceptions exist where the MFC and the WNW-ESE trending segment of the TFFC show small offsets of Mesozoic sedimentary strata (Figure 5c-g). The weak influence of these faults compared to offsets observed within late Paleozoic successions (Figure 5c-g) suggests that, at least some major faults were mildly reactivated in Mesozoic times but, in general, most brittle faults on the Finnmark Platform east and west and in the southwesternmost Nordkapp basin remained inactive after Carboniferous times.

785

4.2. Offshore aeromagnetic data

To better verify our 2D interpretation of faults and basin architectures on the Finnmark Platform and in the southwesternmost Nordkapp basin, we compare and tie our results using high-resolution, offshore aeromagnetic data from Gernigon et al. (2014; Figure 4). Aeromagnetic anomalies, when combined with seismic interpretation, may provide useful results allowing to identify brittle faults and offset patterns (cf. Indrevær et al., 2013).

790

On the Finnmark Platform east, offshore aeromagnetic data (Figure 4; Gernigon et al., 2014) show multiple narrow, NNE-SSW trending, high-positive aeromagnetic anomalies that bend into NW-SE/NNW-SSE orientations near the center of the Nordkapp Basin, which Gernigon et al. (2014) interpreted as arc-shaped prolongations of Caledonian nappes. A more detailed analysis of these aeromagnetic data reveals a set of triangular to rhomboidal, high-negative aeromagnetic anomalies, the largest of which was observed northeast of the island of Magerøya (dashed white lines in Figure 4). This high-negative anomaly is bounded to the northeast and to the northwest by narrow, linear, NNE-SSW to NE-SW trending, high-positive aeromagnetic anomalies (dashed white lines in Figure 4). On seismic data, the locations of these linear, high-positive aeromagnetic anomalies coincide with SE-dipping normal fault for the northwestern anomaly, and the NW-dipping, zigzag-shaped LVF for the southeastern anomaly (cf. black arrows in Figure 5a & b).

800

805 These two faults bound a triangular-shaped basin filled up with thickened Carboniferous
sedimentary deposits (cf. Figure 5a & b), which shape and extent mimic those of the triangular,
high-negative anomaly observed on aeromagnetic data northeast of Magerøya (Figure 4). Such
triangular-shaped, high-negative aeromagnetic anomaly may thus be indicators of offshore
Carboniferous sedimentary basins.

810 Similarly, on the Finnmark Platform west, a large NE-SW trending, linear, high-positive
aeromagnetic anomaly is observed in the footwall of the TFFC (dotted white lines in Figure 4),
where it extends northeastwards into the footwall of the Rolvsøya fault (Figure 4). This NE-SW
trending, high-positive aeromagnetic anomaly coincides with a NE-SW trending basement ridge in
the footwall of the TFFC on the Finnmark Platform west and with the location of an ENE-WSW
815 trending basement ridge in the footwall of the Rolvsøya fault (Figure 1 and Figure 5c-f). We
interpret this NE-SW trending, high-positive aeromagnetic anomaly to highlight a significant
compositional difference between highly-magnetic basement rocks in NE-SW and ENE-WSW
trending basement ridges and poorly magnetic, adjacent basement rocks on the Finnmark Platform
west and in the southwesternmost Nordkapp basin (Figure 1, Figure 4 and Figure 5c-f).

820 5. Discussion

Our regional and detailed seismic studies of basin-boundary faults such as the TFFC, MFC,
Rolvsøya fault and TKFZ on the Finnmark Platform and adjacent southwesternmost Nordkapp
basin show multiple links and interactions. We focus the discussion on the interaction of these
825 faults and associated minor faults on Late-Devonian-Carboniferous (half-) graben basins. We
specifically discuss how deep-seated ductile Caledonian shear zones, i.e. the Sørøya-Ingøya shear
zone and basement ridges may have been exhumed and thus enabled to control post-Caledonian
brittle faulting and formation of Late-Devonian-Carboniferous basins as collapse basins. In
combinaison, the structural architecture, timing of faulting and fault-controlled thickness variations
830 on the Finnmark Platform and in the southwesternmost Nordkapp basin provide the framework to
discuss the evolution of the SW Barents Sea margin from the Mid/Late Devonian to the Permian.

5.1. Interaction of the main segment of the Måsøy Fault Complex with the Sørøya- Ingøya shear zone

835

The linear, NE-SW trending geometry of the main segment of the MFC in map-view (Figure 1 & Figure 2) strongly differs from the dominant ENE-WSW to NNE-SSW trending, zig-zag pattern typically observed for post-Caledonian faults in Mid-Norway (Blystad et al., 1995), Lofoten-Vesterålen (Bergh et al., 2007; Eig, 2008; Hansen et al., 2012), Western Troms (Indrevær et al., 2013) and NW Finnmark (Koehl et al., submitted). Notably, the anomalously linear fault segment of the MFC trends fully parallel to and soles into high-amplitude, NW-dipping seismic reflections of the SISZ on the Finnmark Platform and in the southwesternmost Nordkapp basin (Figure 5c-f). This obvious merging of main segment of the MFC into the basement-seated SISZ (Figure 5c-f) suggests it formed as a brittle splay fault along an inverted portion of the SISZ, likely during the collapse of the Caledonides in the Mid/Late Devonian (Gudlaugsson et al., 1998). We suggest a similar interpretation for the Rolvsøya fault, which also flattens and merges into a bowed portion of the SISZ (Figure 5c & d), and for the northwest-boundary fault of the Devonian graben on the Finnmark Platform west soling into a minor, SE-dipping shear zone (Figure 5e & Figure 6b & c). These faults are thought to have remained active through the late Carboniferous as suggested by potential syn-tectonic sediment thickening within the upper Carboniferous succession (Figure 9b) but most likely ceased before the Permian as supported by the relatively constant thickness of Permian sedimentary strata throughout the study area (Figure 9c).

By analogy, in the North Sea, Phillips et al. (2016) successfully tied the southernmost onshore occurrence of the Karmøy Shear Zone, a major Caledonian shear zone, to a thick seismic unit made up with sub-parallel, high-amplitude reflections similar to those ascribed to the SISZ in the footwall of the main segment of the MFC (Figure 5d-f). Phillips et al. (2016) argue that the Åsta Fault, a large N-S trending, W-dipping, post-Caledonian fault in the North Sea, formed during a phase of extensional reactivation of the Karmøy Shear Zone. Similarly, in western Norway, Wilks & Cuthbert (1994) proposed that the Hornelen Basin formed along a brittle fault that splayed upwards from the Nordfjord-Sogn Detachment Zone during Middle Devonian late-orogenic extension.

5.2. Formation of the WNW-ESE trending fault segment of the Troms-Finnmark Fault Complex as a hard-linked accommodation cross-fault

865

Our data (Figure 9a & b) show abrupt fault-controlled thickening of the Devonian-lower Carboniferous and upper Carboniferous sedimentary successions just northeast of the WNW-ESE trending segment of the TFFC into the southwesternmost Nordkapp basin. On the Finnmark Platform west, potential Devonian sedimentary rocks are truncated upwards by the mid- Carboniferous reflection (Figure 5e & f). We propose that the absence of high-amplitude, coal-bearing sedimentary deposits of the Billefjorden Group (lower Carboniferous) on the Finnmark Platform west is related to a major episode of eustatic sea-level fall in the Serpukhovian (Saunders & Ramsbottom, 1986), which may have contributed to expose lower Carboniferous sedimentary rocks in this area to coastal erosion. Hence, part of the thickening of the Devonian-lower Carboniferous succession across the WNW-ESE trending segment of the TFFC might be related to extensive erosion of the Finnmark Platform west in mid-Carboniferous times. In addition, the clear deepening (plunge) to the northeast of the spoon-shaped trough formed by the three-dimensionally folded and bowed geometry of the SISZ (Figure 7) suggests that the thickening of Devonian-lower Carboniferous sedimentary strata into the southwesternmost Nordkapp basin (Figure 9a) is also partly controlled by the shape and attitudes of the underlying SISZ. Finally, the thickened sediment depocenter observed in the southwesternmost Nordkapp basin at the intersection of the TFFC and the MFC (Figure 9a & b) is at least partly related to syn-sedimentary normal faulting along the WNW-ESE trending fault segment of the TFFC and along the main segment of the MFC. This most likely indicates that the TFFC and the MFC had already merged and acted as a single fault zone during sediment deposition in the southwesternmost Nordkapp basin from the end of the Serpukhovian and potentially from Devonian times. We propose that the WNW-ESE trending fault segment of the TFFC acted as an accommodation cross-fault, as defined in Sengör (1987), that transferred displacement between the NNE-SSW trending segment of the TFFC and the main segment of the MFC, defining a step synthetic with the deepening direction of the spoon-shaped trough formed by the geometry of the SISZ (Figure 7). This interpretation is based on the dominant dip-slip kinematic of the WNW-ESE trending segment of the TFFC and on its sub-parallel strike to the dominant WNW-ESE trending extension direction inferred along the SW Barents Sea margin during late Paleozoic times (Bergh et al., 2007; Eig & Bergh, 2011; Hansen & Bergh, 2012). Further, we infer that the strike and location of the WNW-ESE trending segment of the TFFC was controlled by the geometry of the underlying SISZ (see below), which dips gently to the northeast on the Finnmark Platform west and in the southwesternmost Nordkapp basin and

may therefore have favored the formation of a NE-dipping fault at this location (Figure 5g & Figure 7).

Alternatively, Lea (2016) proposed that the WNW-ESE trending fault segment of the TFFC
900 corresponds to a breached relay-ramp fault between the NNE-SSW trending fault segment of the
TFFC and the MFC. However, this model implies that this portion of the TFFC would have
accommodated significantly fewer displacement than the two faults it links (i.e. the NNE-SSW
trending segment of the TFFC and the MFC), which is clearly not the case. The offset of the mid-
Carboniferous reflection and the thickness increase both of the Middle/Upper Devonian-lower
905 Carboniferous and upper Carboniferous sedimentary successions across the WNW-ESE trending
segment of the TFFC are comparable to the offset and thickness increase observed across the main
fault segment of the MFC (Figure 5c, d & g), and the TFFC and MFC seem to have evolved
synchronously in the late Paleozoic.

910 **5.3. Devonian collapse basins on the Finnmark Platform west (Gjesvær Low) and in the southwesternmost Nordkapp basin**

Devonian sedimentary rocks in the SW Barents Sea may exist on the Finnmark Platform
west and in the southwesternmost Nordkapp basin (Figure 5c-g). The most probable occurrence is
915 on the Finnmark Platform west, in the hanging-wall of the main segment of the MFC (Figure 1,
Figure 5e). The presumed Devonian seismic unit corresponds to a suite of low-amplitude
reflections crosscut by a few, moderate-amplitude reflection that dip gently to the northeast (Figure
5e). The main argument for a Devonian succession is that these reflections are remarkably different
from the typical seismic patterns observed for lower Carboniferous sedimentary deposits and
920 basement rocks. Lower Carboniferous sedimentary deposits of the Billefjorden Group are
characterized by high-amplitude reflections produced by coal-bearing sedimentary rocks (Figure
5a & b and Figure 6a), while basement rocks are mostly associated with thick packages of chaotic
seismic reflections (Figure 6a-c) and thick layers of moderate- to high-amplitude, sub-parallel
seismic reflections that we interpreted as basement-seated shear zones (e.g. the SISZ; Figure 6f).
925 Another argument in favor of Devonian sedimentary deposits is the presence of a NE-SW trending
gravimetric low on the Finnmark Platform west: the Gjesvær Low (Johansen et al., 1994).
Devonian sedimentary rocks in Svalbard show an average density of ca. 2.4 g.cm^{-3} associated to

depths of 0-8 km (i.e. average depth of 4 km; Manby & Lyberis, 1992), which is less dense than metamorphosed Caledonian rocks ($2.6-3.0 \text{ g.cm}^{-3}$) and Carboniferous sedimentary rocks on the Finnmark Platform west ($< 2.5 \text{ g.cm}^{-3}$; Johansen et al., 1994). However, taking into account the effect of burial up to a depth of 5-6 km on the Finnmark Platform (Johansen et al., 1994) and the resulting density increase for Devonian sedimentary deposits with an approximate rate of ca. $0.15 \text{ g.cm}^{-3}.\text{km}^{-1}$ (cf. “all rocks density-depth gradient” in table 3 in Maxant 1980), Devonian sedimentary rocks on the Finnmark Platform may reach densities $2.55-2.7 \text{ g.cm}^{-3}$. Thus, the occurrence of the Gjesvær low can be explained by the presence of intermediate-density Devonian sedimentary rocks below the mid-Carboniferous reflection (Figure 5e & Figure 6b & c). This is as well in accordance with density variations and related estimates of Johansen et al. (1994) in the Gjesvær low.

In addition, the discrete, moderate-amplitude, NW-dipping reflections observed within the presumed Devonian sedimentary strata on the Finnmark Platform west may represent syn-tectonic sedimentary growth strata (Figure 5e & Figure 6b & c). These strata are located above and thickened against arcuate, high-amplitude reflections that we interpreted as a major erosional unconformity that truncates SE-dipping Caledonian basement shear zones subparallel to the SISZ (dotted yellow lines in Figure 6b & c). We consider the Devonian sedimentary rocks on the Finnmark Platform west to have been deposited in wedge-shaped late/post-Caledonian extensional basins due to reactivation of a set of partly eroded, exhumed, SE-dipping Caledonian shear zones (dotted yellow lines in Figure 5e & Figure 6b & c). In mid-Norway, Braathen et al. (2000) reported a similar setting of Middle Devonian sedimentary basins located above Caledonian shear zones and folded nappe stack, and proposed that Middle Devonian basins in western and mid-Norway formed during extensional reactivation of these Caledonian shear zones. Such a model is further supported by the similar geometry of the Devonian sedimentary growth strata on the Finnmark Platform west to the highly tilted geometry of sedimentary strata in Middle Devonian basins in western Norway (cf. Séranne & Seguret, 1987; Séranne et al., 1989; Wilks & Cuthbert, 1994; Osmundsen & Andersen, 2001). Moreover, the Devonian sedimentary basins on the Finnmark Platform west (Figure 5e and Figure 6b & c) and in the southwestermost Nordkapp basin (Figure 5c & d) define NE-SW trending graben structures with comparable, $< 50 \text{ km}$ -wide sizes to those of the Middle Devonian Hornelen, Kvamsheten and Solund basins in western Norway (Séranne & Seguret, 1987; Osmundsen & Andersen, 2001).

In the southwesternmost Nordkapp basin, the presence of Middle/Upper Devonian sedimentary rocks is more speculative and is mostly based on the maximum thickness of lower Carboniferous sedimentary deposits registered in the SW Barents Sea, which is ca. 600 m on the Finnmark Platform east (Bugge et al., 1995; Figure 6d). Assuming a seismic velocity $< 6 \text{ km.s}^{-1}$ for lower Carboniferous coal-bearing sedimentary deposits, a thickness of 600 m would account for only part (maximum 0.2 s) of the 2-2.5 second-thick (TWT) seismic unit observed below the mid-Carboniferous reflection in the southwesternmost Nordkapp basin (Figure 5c & d). If basement rocks were present at the base of the southwesternmost Nordkapp basin, they would most likely produce a seismic reflection pattern similar to that of the sub-parallel, high-amplitude reflections of the underlying SISZ (Figure 5c & d) or potentially form an unconformity to the overlying late Paleozoic sedimentary rocks. We therefore believe that the southwesternmost Nordkapp basin, which is bounded below by the SISZ, giving the basin a peculiar “U” shape in cross-section (Figure 5c & d), is composed of thick Middle/Upper Devonian sedimentary deposits overlain by lower Carboniferous sedimentary strata below the mid-Carboniferous reflection. Based on the brittle extensional reactivation, bowed geometry and controlling effect of the basement-seated SISZ (Figure 7), we suggest deposition of Devonian sedimentary rocks within a late/post-Caledonian, spoon-shaped, collapse basin formed along bowed, inverted portions of the Caledonian SISZ (Figure 5c-f), thus representing analogs to Middle Devonian collapse basins in western and mid-Norway (Séranne et al., 1989; Wilks & Cuthbert, 1994).

5.4. Formation of NE-SW to ENE-WSW trending basement ridges as exhumed metamorphic core complexes

We have argued for an upward-bowed seismic geometry of the SISZ (Figure 5c-f) into which major fault complexes such as the TFFC, the MFC and the Rolvsøya fault merge and sole into (Figure 5c-f). In map-view (Figure 7) and cross-section (Figure 5c-f), the bowed geometry of the SISZ defines two, ENE-WSW to NE-SW trending ridges of basement rocks on the northwestern flanks of presumed Devonian basins. These basement ridges correlate well by displaying high-positive gravimetric (fig. 5 in Olesen et al., 2010 and fig. 5 in Gernigon et al., 2014) and high-positive aeromagnetic anomalies (Figure 4; Gernigon et al. 2014) that suggest these ridges are made of basement lithologies significantly different from adjacent basement rocks on

990 the Finnmark Platform west and in the southwesternmost Nordkapp basin. These basement ridges
seem to align with high-positive gravimetric anomalies that coincide with the NE-SW trending
Norsel High (Gabrielsen et al., 1990) along the northwestern flank of the Nordkapp Basin in the
northeast. Farther southwest, these basement ridges coincide with the NE-SW trending West Troms
Basement Complex in Western Troms (Bergh et al., 2010; Figure 1) and the NE-SW trending
995 Lofoten Ridge in Lofoten-Vesterålen (Bergh et al., 2007; Figure 1), among which at least the
Lofoten Ridge was exhumed as a metamorphic core complex (Klein & Steltenpohl, 1999; Klein et
al., 1999; Steltenpohl et al., 2004; 2011) along inverted Caledonian shear zones such as the
Eidsfjord and Fiskefjord shear zones (Steltenpohl et al., 2011). By comparison, the SISZ seems to
coincide with high-positive aeromagnetic anomalies on the Finnmark Platform west that follow the
1000 trace of the MFC (Indrevær et al., 2013) and continue past the southwestern fault-tip of the MFC
(Gernigon & Brønner, 2012). The aeromagnetic anomalies visible on the dataset Gernigon &
Brønner (2012) appear to line up with aeromagnetic anomalies onshore the island of Vannøya, in
the northeasternmost part of the West Troms Basement Complex (Figure 1), and these onshore
anomalies correlate with NE-SW trending, SE-dipping basement shear zones that were reactivated
1005 as extensional brittle faults (Paulsen et al. pers. comm.).

These data indicate that SE-dipping portions of the SISZ propagated southwest of the MFC
fault-tip on the Finnmark Platform west and possibly merged onshore Vannøya in Western Troms
with a suite of NE-SW trending, SE-dipping shear zones. As a consequence, the basement ridges
observed on the Finnmark Platform west and along the northern flank of the southwesternmost
1010 Nordkapp basin may have formed as part of a large metamorphic core complex, which included
the Lofoten Ridge, the West Troms Basement Complex and possibly also the Norsel High (Figure
1), exhumed along inverted Caledonian shear zones, e.g. SISZ on the SW Barents shelf and the
analogous Eidsfjord and Fiskefjord shear zones in Lofoten-Vesterålen (Steltenpohl et al., 2011).

The timing, nature of uplift and processes of core complex exhumation can be inferred from
1015 thickness variations of the SISZ in cross-section, for example, thickest in the footwall of the MFC
and TFFC and thinnest below these two fault complexes (Figure 5f). We link these thickness
variations to excisement and incisement processes (cf. Lister & Davis, 1989) along the SISZ during
core complex exhumation, after the embrittlement of the SISZ (Figure 10). A model of Devonian
late/post-orogenic extension is proposed, when inversion of the SISZ as a low-angle, top-to-the-
1020 NW extensional detachment and extensive erosion of the Caledonides contributed to crustal

thinning (Figure 10a). Rapid thinning through extension and erosion above the upper part of the SISZ may have triggered early exhumation of basement rocks on the Finnmark Platform west and along the northern flank of the southwesternmost Nordkapp basin (Figure 10a), causing the upper part of the SISZ to bow upwards (Figure 10b). Continued crustal extension and continental erosion further enhanced exhumation of basement rocks below the upper part of the SISZ, leading the bowed portion of the SISZ to become unsuitably oriented to accommodate top-to-the-NW extensional displacement and thus become inactive (dashed black line in Figure 10c). Further extension likely triggered upwards splaying of the SISZ into its hanging-wall, becoming suitable again to accommodate top-to-the-NW extension displacement (Figure 10c). This upward splay-faulting process is referred to as excisement by Lister & Davis (1989) and we tentatively apply this process to explain the observed thickening of the SISZ in the footwall of the MFC (Figure 5f). Further extension/erosion-related crustal thinning along the SISZ may have initiated exhumation of basement rocks along progressively deeper parts of the SISZ (Figure 10b & c), causing bend-up of the SISZ at deeper crustal levels (Figure 10d). Extreme bowing of lower portions of the SISZ led to opposite top-to-the-SE transport direction on the Finnmark Platform west and in the southwesternmost Nordkapp basin (Figure 10d), which contributed to exhume NE-SW to ENE-WSW trending ridges of basement rocks in the footwall of the NNE-SSW trending segment of the TFFC (Figure 5e & f) and in the footwall of the Rolvsøya fault (Figure 5c & d), thus forming a large spoon-shaped trough where Devonian sedimentary rocks deposited (Figure 7 and Figure 10d & e). Incisement (downward splaying; cf. Lister & Davis, 1989) may have occurred below the basement ridges during continued top-to-the-NW extension along the SISZ (Figure 10e) and possibly contributed to thicken the SISZ in the footwall of the TFFC (Figure 5e & f) and of the Rolvsøya fault (Figure 5c & d), resulting in the current geometry of the SISZ (Figure 10f).

By comparison, in northeast Greenland, Sartini-Rideout et al. (2006) and Hallett et al. (2014) proposed that ultra-high pressure (UHP) basement rocks were exhumed along large, mylonitic, Caledonian shear zones in Late Devonian-early Carboniferous times (ca. 370-340 Ma). The study of Sartini-Rideout et al. (2006) also shows that the last stages of exhumation were accommodated by steep, brittle normal faults that strike parallel to major Caledonian shear zones, i.e. similar to the main segment of the MFC striking parallel to the SISZ along the SW Barents Sea margin (Figure 1). In addition, results from sediment provenance and geochronological studies by McClelland et al. (2016) in Carboniferous basins in northeast Greenland showed that the

exhumation of UHP basement rocks as elongated ridges could have formed a regional Serpukhovian erosional unconformity, contemporaneous with the mid-Carboniferous (Serpukhovian?) unconformity observed on the Finnmark Platform east (Figure 5a & b and Figure 1055 6a; Bugge et al., 1995) and on the Finnmark Platform west (Figure 5e-g & Figure 6b & c), and in agreement with eustatic sea-level fluctuations at that time (Saunders & Ramsbottom, 1986). In late Paleozoic times, the northeast Greenland margin was located close to its conjugate counter-part of the SW Barents Sea margin and these two areas were most likely subjected to similar regional stresses and closely related sea-level fluctuations. Therefore, we suggest that the mid-1060 Carboniferous unconformity reflection observed in the SW Barents Sea (cf. Figure 5 & Figure 6a-c; Bugge et al., 1995), formed as a response to major eustatic sea-level fall in the early Serpukhovian (Saunders & Ramsbottom, 1986) and due to large-scale exhumation of basement rocks in Late Devonian-early Carboniferous times. Exhumation occurred along inverted Caledonian shear zones (e.g. SISZ) and brittle splay faults such as the main segment of the MFC, 1065 the NNE-SSW trending segment of the TFFC, the Rolvsøya fault and the NNE-SSW trending, SE-dipping fault that bounds potential Devonian sedimentary strata on the Finnmark Platform west (Figure 5). The timing of exhumation is constrained to the end of the Serpukhovian based on deposition of thick alluvial and shallow marine upper Carboniferous sedimentary deposits of the Gipsdalen Group (Figure 3) on top of the mid-Carboniferous unconformity in the SW Barents Sea 1070 (Figure 5) during an eustatic sea-level rise near the end of the Serpukhovian (Saunders & Ramsbottom, 1986).

In Lofoten-Vesterålen, Steltenpohl et al. (2004) inferred a Late Devonian age for the exhumation of metamorphic core complexes and this age was refined by recent $^{40}\text{Ar}/^{39}\text{Ar}$ isotopic results (Steltenpohl et al., 2011), which constrained extensional reactivation of the Eidsfjord shear 1075 zone to the Early Devonian. In the SW Barents Sea, much work is needed to better constrain the timing of late/post-Caledonian extension and collapse basin formation. Nonetheless, we believe that the Early Devonian age obtained in Lofoten-Vesterålen (Steltenpohl et al., 2011) represents a reasonable estimate for the onset of crustal thinning in the SW Barents Sea (Figure 10a-c). Additionally, Late Devonian-early Carboniferous timing of exhumation for basement rocks in 1080 northeast Greenland and formation of a regional mid-Carboniferous (Serpukhovian) unconformity (Sartini-Rideout et al., 2006; Hallett et al., 2014; McClelland et al., 2016) corresponds to a realistic approximation for the final stages of late/post Caledonian extension, ending with the formation of

Devonian-Carboniferous collapse basins along exhumed, NE-SW to ENE-WSW trending basement ridges (Figure 10d-f).

1085

5.5. Interaction of the Trollfjorden-Komagelva Fault Zone with the Troms-Finnmark and Måsøy fault complexes

The prolongation of the TKFZ from onshore areas in eastern Finnmark to offshore areas of the SW Barents Sea has been a matter of debate. Most studies tend to connect the onshore TKFZ with the offshore WNW-ESE trending fault segment of the TFFC (Gabrielsen, 1984; Gabrielsen & Færseth, 1989; Roberts et al., 2011; Bergø, 2016; Lea, 2016). Our data, however, suggest that the TKFZ dies out near the coasts of NW Finnmark (present contribution and Koehl et al., submitted), and in this section we review and discuss new evidence obtained from the interpretation of offshore seismic and aeromagnetic data.

First, the TKFZ was described onshore eastern Finnmark as a major sub-vertical fault that accommodated dominantly strike-slip movement (Roberts, 1972; Rice, 2013). Farther west, the TKFZ crops out onshore Magerøya, where it is made of numerous, high-frequency, subparallel, subvertical, WNW-ESE trending faults and fractures that accommodated at least small-scale post-Caledonian, strike-slip to oblique-slip displacement (Koehl et al., submitted). By contrast, seismic interpretation of the WNW-ESE trending fault segment of the TFFC shows that this fault exhibits a typical, high-angle (ca. 70°) normal fault geometry and accommodated significant amount of post-Caledonian normal dip-slip displacement (Figure 5g), thus contrasting significantly with the geometry of the TKFZ. Second, the imaginary prolongation of the TKFZ from the island of Magerøya to the WNW, onto the Finnmark Platform, would crosscut the Finnmark Platform west nearly 23 km southwest of the observed trace of the WNW-ESE trending segment of the TFFC (Figure 5g). This represents a significant mismatch that is far too important to represent minor dextral strike-slip offset of the TKFZ across the main fault segment of the MFC, which dominantly accommodated normal dip-slip motions (Figure 5c-f). Third, the interpretation of 3D seismic data at the intersection of the MFC and TFFC reveals that the footwall of the MFC is largely intact and seismically unaffected by brittle faults (Figure 8). There is no evidence of intense fracturing as typically observed along the TKFZ onshore Magerøya (Koehl et al., submitted). We therefore believe that the TKFZ and the WNW-ESE trending fault segment of the TFFC represent two

distinct faults. This suggests that the TKFZ dies out instead of propagating onto the Finnmark
1115 Platform. This is also supported by the absence of WNW-ESE trending faults offshore (Figure 1
& Figure 2). For example, the Austhavet fault previously interpreted near the coasts of Finnmark
(Townsend, 1987b; Lippard & Roberts, 1987; Roberts et al., 2011) was re-interpreted as seismic
artifacts related to the Djuprenna trough, a large glacial trough that trends parallel to the
northeastern coast of Finnmark (Ottesen et al., 2008; Rise et al., 2015). This re-interpretation is
1120 supported by shallow drillings on the Finnmark Platform east, which show no sign of fault-related
offset in this part of the Finnmark Platform east (cf. fig. 4 in Bugge et al., 1995). Similarly, our
mapping and regional analysis of brittle faults on the Finnmark Platform show very few
occurrences of WNW-ESE trending faults (Figure 1 & Figure 2).

Onshore studies (Koehl et al., submitted) show an increased number of large-scale WNW-
1125 ESE trending fault segments and splays along the TKFZ, varying from a single-segment fault
onshore the Varanger Peninsula in eastern Finnmark to multiple segments onshore and nearshore
Magerøya (Figure 1), suggesting that the island of Magerøya is located near the fault-tip process
zone (Shipton & Cowie, 2003; Braathen et al., 2013) of the TKFZ, and that the TKFZ therefore
dies out to the west before reaching the Finnmark Platform and the southwesternmost Nordkapp
1130 basin (Figure 1; Koehl et al. submitted). Nearby Magerøya and the Nordkinn Peninsula, high-
resolution aeromagnetic data reveal the presence of highly magnetic dolerite dykes along WNW-
ESE trending fault segments of the TKFZ (Roberts et al., 1991; Nasuti et al., 2015; Koehl et al.,
submitted). These narrow, high-positive aeromagnetic anomalies also die out westwards (Gernigon
& Brönnert, 2012; Gernigon et al., 2014), therefore supporting that the dolerite dykes and, thus, the
1135 TKFZ die out before reaching the Finnmark Platform (Koehl et al., submitted).

We explore an alternative model to the fault-tip process zone of Koehl et al. (submitted) in
which we argue that the Precambrian, orogen-parallel, WNW-ESE trending TKFZ was not suitably
oriented to be reactivated as a major thrust or strike-slip fault during the Caledonian Orogeny. Our
data indicate that if the TKFZ extended farther west prior to the onset of Caledonian deformation,
1140 it was certainly truncated and decapitated by large-scale, top-to-the-SE movement along the SISZ
and associated NE-SW trending, Caledonian thrusts and shear zones. This is supported by
dominant top-to-the-SE transport direction inferred along Caledonian thrusts onshore NW
Finnmark (Townsend, 1987a). Thus, we propose that the western continuation of the TKFZ below
the Finnmark Platform may have been thrust southeastwards along the SISZ and is now eroded.

1145 However, if the TKFZ ever extended westwards, portions of its western prolongation might be preserved in offshore basement highs such as the Loppa and Veslemøy highs (Figure 1). More work is needed on this hypothesis, but a possible insight is the recent observation of subvertical, WNW-ESE trending brittle faults analog to the TKFZ in basement rocks of the Veslemøy High (Kairanov et al., 2016).

1150

5.6. Late Paleozoic evolution of the SW Barents Sea margin

Based on the seismic data and discussions from previous chapters we now address the tectonic evolution of the Finnmark Platform and the southwesternmost Nordkapp basin in the late Paleozoic (Figure 11). The main structural element discussed in our model is the SISZ and we link its influence on (i) the development of the southwesternmost Nordkapp basin, (ii) the geometry of the TFFC and MFC, (iii) the deposition of Mid/Late Devonian sedimentary rocks in the southwesternmost Nordkapp basin and on the Finnmark Platform west, (iv) transfer faults such as the TKFZ and (v) the deposition of syn-tectonic sedimentary wedges along steep normal faults that bound triangular-shaped, Carboniferous basins.

1160

The trend and dominant northwestern dip of the SISZ (Figure 5c-g & Figure 6f) suggest that it formed as a large thrust that accommodated top-to-the-SE tectonic transport during the Caledonian Orogeny. The SISZ has a bow-shaped, three-dimensionally folded geometry that coincides with basement ridges in the footwall of the TFFC and of the Rolvsøya fault (Figure 5c-f, Figure 7 and Figure 11). We propose that the SISZ and potential other Caledonian shear zones along the SW Barents Sea margin were inverted as low-angle extensional shear zones during late/post-Caledonian orogenic extension and subsequent collapse. This is based on analog examples in northeast Greenland (Sartini-Rideout et al., 2006; Hallett et al., 2014; McClelland et al., 2016), western Norway (Séranne & Seguret, 1987; Séranne et al., 1989; Wilks & Cuthbert, 1994; Osmundsen & Andersen, 2001), mid-Norway (Braathen et al., 2000) and Lofoten-Vesterålen (Klein & Steltenpohl, 1999; Klein et al., 1999; Steltenpohl et al., 2004; 2011; Osmundsen et al., 2005). Extensional reactivation of such ductile shear zones along the Barents Sea margin may have initiated in the Early Devonian, as in Lofoten-Vesterålen (Steltenpohl et al., 2011), through orogenic collapse dominated by top-to-the-NW movement along the SISZ.

1170

1175 Exhumation of the SISZ and underlying basement ridges as a metamorphic core complex
was probably triggered by extensional reactivation of the SISZ combined to continental erosion,
leading to crustal thinning. Reactivation of these exhumed basement ridges occurred by normal
faulting along new, steep, brittle faults such as the main segment of the MFC and the NNE-SSW
trending fault segment of the TFFC (cf. Figure 5f), likely due to incisement and excisement
1180 processes (Figure 10; Lister & Davis, 1989). These processes also contributed to the progressive
exhumation of ENE-WSW and NE-SW trending basement ridges along bowed portions of the SISZ
(cf. Figure 5c-f and Figure 11a & b). We believe that these ridges were part of a larger-scale NE-
SW trending metamorphic core complex that included the Norsel High and the two basement ridges
located in the footwall of the TFFC and the Rolvsøya fault. Farther south, this core complex may
1185 be linked to the West Troms Basement Complex (Bergh et al., 2010) and the Lofoten Ridge
(Blystad et al., 1995; Figure 11). Such a regional link is favored by the alignment of NE-SW
trending, high-positive gravimetric anomalies that characterize these ridges (Olesen et al., 2010;
Gernigon et al., 2014). The timing of final core complex exhumation can be constrained to
Mid/Late Devonian-early Carboniferous and possibly linked to the regional Serpukhovian
1190 unconformity on the Finnmark Platform (cf. Figure 5a, b, e, f & g and Figure 6a-c; Bugge et al.,
1995), in accordance with Sartini-Rideout et al. (2006) and Hallett et al. (2014) in northeast
Greenland.

The exhumation of basement ridges as metamorphic core complexes along the inverted
SISZ and subsequent normal faulting along the MFC and TFFC created a deep, spoon-shaped
1195 topographic depression on the Finnmark Platform west and in the southwesternmost Nordkapp
basin (Figure 5c, d, e & g, Figure 6b & c and Figure 11a). These depressions were filled with thick
Devonian clastic deposits analog to those observed in Middle Devonian collapse basins in western
Norway (Séranne et al., 1989; Osmundsen & Andersen, 2001), and with lower Carboniferous coal-
bearing and clastic sedimentary rocks of the Billefjorden Group (Figure 3) deposited
1200 unconformably above Devonian strata (cf. Figure 11b). These collapse basins are also likely
responsible for the gravimetric low observed on the Finnmark Platform west, the Gjesvær Low
(Figure 4).

On the Finnmark Platform west, the final stages of core complex exhumation and a major
phase of eustatic sea-level fall in the Serpukhovian (Saunders & Ramsbottom, 1986) led to
1205 extensive erosion of Devonian and lower Carboniferous sedimentary rocks, therefore explaining

the absence of lower Carboniferous sedimentary deposits and the erosional truncation of Devonian sedimentary strata along this part of the margin (Figure 5e-g & Figure 6b & c). On the Finnmark Platform east, lower Carboniferous sedimentary rocks are preserved as minor syn-tectonic sedimentary wedges within small triangular grabens and half-grabens that correlate with aeromagnetic lows (dashed white line in Figure 4). These grabens are bounded by zigzag-shaped, Late Devonian-Carboniferous normal faults such as the LVF (Figure 5a & b and Figure 6a; Koehl et al., submitted), which coincide with narrow, high-positive aeromagnetic anomalies (cf. Figure 4 and black vertical arrows in Figure 5a & b). In addition, triangular basins like the graben bounded by the LVF and the southwesternmost Nordkapp basin were partly offset and segmented by WNW-ESE trending transfer faults that accommodated small amount of strike-slip motion. Examples include the TKFZ onshore NW Finnmark, which may offset the LVF in a right-lateral fashion (Koehl et al., submitted), and accommodation cross-faults (Sengör, 1987) that accommodated large amount of orogen-parallel extension through normal dip-slip movement, e.g. the WNW-ESE trending fault segment of the TFFC (Figure 5g and Figure 9a).

In the late Serpukhovian, a regional episode of eustatic sea-level rise (Saunders & Ramsbottom, 1986) flooded the Finnmark Platform east and west and allowed the deposition of sedimentary rocks of the upper Carboniferous Gipsdalen Group (Figure 3). These rocks occur as syn-tectonic sedimentary wedges that thicken in the hanging-wall of basin-bounding normal faults such as the LVF on the Finnmark Platform east and the main segment of the MFC on the Finnmark Platform west (Figure 5a, b, e & f and Figure 11c). Similarly, in the southwesternmost Nordkapp basin, which may have remained flooded through the entire phase of eustatic sea-level fall and core complex exhumation, thick, partly evaporitic, upper Carboniferous sedimentary rocks were deposited in the basin and these are thickest at the intersection of the TFFC and the MFC (Figure 5c, d & g and Figure 9b). Thus, the thickening of upper Carboniferous strata probably reflects significant syn-sedimentary normal faulting along these two faults, which may have acted as a single fault during in the final stage of extension in the late Carboniferous (Figure 11c).

Most faults on the Finnmark Platform east and west and in the southwesternmost Nordkapp basin appear to die out below the Base Asselian reflection and those that propagate through the Base Asselian reflection show limited amount of offset within Permian and Mesozoic-Cenozoic sedimentary strata (Figure 5). Moreover, the Permian sedimentary succession shows a rather constant thickness through the entire study area (Figure 5 & Figure 9c). Thus, we argue that

late/post-Caledonian extensional faulting linked to the collapse of the Caledonides essentially took place in the Mid/Late Devonian-Carboniferous and came to a halt towards the end of this Period (Figure 11d). This presumed timing is consistent with recent K/Ar radiometric dating of brittle fault gouges in Western Troms (Davids et al., 2013) and in NW Finnmark (Torgersen et al., 2014; Koehl et al., 2016), as well as with radiometric dating of dolerite dykes in NW Finnmark (Lippard & Prestvik, 1997), eastern Finnmark (Guise & Roberts, 2002) and on the Kola Peninsula in Russia (Roberts & Onstott, 1995), which constrain significant extensional faulting activity onshore northern Norway and adjacent areas in Russia to the Late Devonian-early/mid Permian. Minor reactivation of major fault complexes occurred in the Mesozoic-Cenozoic and are most likely associated with the rifting of the NE Atlantic (Faleide et al., 2008).

6. Conclusions

- 1) The atypically linear, NE-SW trending, main fault segment of the Måsøy Fault Complex formed as a brittle splay of the inverted Caledonian Sørøya-Ingøya shear zone through excisement processes during the collapse of the Caledonides in the Mid/Late Devonian-early Carboniferous and was active until the end of the late Carboniferous.
- 2) The WNW-ESE trending fault segment of the Troms-Finnmark Fault Complex developed as a hard-linked, accommodation cross-fault that accommodated orogen-parallel, late/post-orogen extension in the Mid/Late Devonian-Carboniferous. This fault merged with the main fault segment of the Måsøy Fault Complex and the two faults acted as a single fault at least during the late Carboniferous, but potentially from Devonian-early Carboniferous times, and accommodated the deposition of thick Devonian-lower Carboniferous and partly evaporitic upper Carboniferous deposits in the southwesternmost Nordkapp basin before faulting came to a halt towards the end of the late Carboniferous.
- 3) Low-gravity anomalies in the Gjesvær Low and southwesternmost Nordkapp basin may result from the presence of a thick, Mid/Late Devonian, spoon-shaped sedimentary basin that developed along an inverted, bowed portion of the Sørøya-Ingøya shear zone during the collapse of the Caledonides and that display a geometry similar to those of Middle Devonian, late/post-orogenic collapse basins in western and mid-Norway.

- 1270 4) The ENE-WSW and NE-SW trending basement ridges in the footwall of the Troms-Finnmark Fault Complex and on the northern flank of the southwesternmost Nordkapp basin formed through incisement processes and were exhumed along a bowed portion of the Sørøya-Ingøya shear zone during the collapse of the Caledonides in the Mid/Late Devonian-early Carboniferous. These basement ridges are thought to be part of a large-scale, margin-parallel, NE-SW trending, metamorphic core complex that includes a succession of aligned basement highs such as the Lofoten Ridge, the West Troms Basement Complex and the Norsel High. Core complex exhumation is believed to have stopped by the end of the Serpukhovian when a major eustatic sea-level rise flooded the Finnmark Platform, leading to the deposition of widespread upper Carboniferous sediments.
- 1275
- 1280 5) The Sørøya-Ingøya shear zone is thought to have truncated and decapitated Precambrian faults such as the Trollfjorden-Komagelva Fault Zone through top-to-the-SE thrusting during the Caledonian Orogeny and subsequent late/post-orogenic extension. We nevertheless believe that preserved segments of these Precambrian faults might be preserved offshore on basement highs such as the Loppa and Veslemøy highs. However, more work is required in order to map and evaluate the impact of these WNW-ESE trending, subvertical Precambrian faults on the SW Barents Sea margin.

1285

Data availability

The seismic data analyzed in this study are part of the DISKOS database and are publicly accessible from any Norwegian academic institution. Aeromagnetic data discussed in the present contribution are from Gernigon et al. (2014).

1290

Author contribution

Jean-Baptiste Koehl interpreted the seismic and aeromagnetic data and is the main contributor to the writing process (work-load ca. 45 %). Professor Steffen Bergh provided significant input to the “Introduction” and “Geological Setting” sections as well as detailed critical reviews of the whole manuscript (work-load ca. 30 %). Tormod Henningsen helped initiating the project and provided help with seismic well-ties and regional seismic interpretation (work-load ca.

1295

15 %). Professor Jan-Inge Faleide provided help with the writing process and helped improve the margin evolution model (work-load ca. 10 %).

1300 **Acknowledgements**

The present study is part of the ARCEX project (Research Centre for Arctic Petroleum Exploration), which is funded by the Research Council of Norway (grant number 228107) together with ten academic and eight industry partners. We would like to thank all the persons from these
1305 institutions that are involved in this project. We acknowledge the NPD, the NTNU and Schlumberger for sharing seismic data from the DISKOS database, with special thanks to Dicky Harishidayat for taking care of the transfer of the data. We thank Tom Arne Rydningen from the University of Tromsø for reviewing parts of the manuscript. We would also like to thank Hanne-Kristin Paulsen from the University of Tromsø as well as Gwenn Péron-Pinvidic and Per-Terje
1310 Osmundsen from the NGU for fruitful discussions.

References

- Andersen, T. B.: The structure of the Magerøy Nappe, Finnmark, North Norway, *Nor. geol. unders.*, 363, 1-23, 1981.
1315
- Andersen, T. B.: The stratigraphy of the Magerøy Supergroup, Finnmark, north Norway, *Nor. geol. unders.*, 395, 25-37, 1984.
- Andersen T. B., Austrheim, H., Sturt, B.A., Pedersen, S. and Kjærø, K.: Rb-Sr whole rock ages from Magerøy, North Norwegian Caledonides, *Norsk Geologisk Tidsskrift*, 62, 79-85,
1320 1982.
- Andresen, A., Agyei-Dwarko, N. Y., Kristoffersen, M. and Hanken, N-M.: A Timanian foreland basin setting for the late Neoproterozoic-Early Palaeozoic cover sequences (Dividal Group) of northeastern Baltica, in: *New Perspectives on the Caledonides of Scandinavia and Related Areas*, Corfu, F., Gasser, D. and Chew, D.M. (eds), Geological Society, London,
1325 Special Publications, 390, 157-175, 2014.
- Bergh, S. G. and Torske, T.: The Proterozoic Skoadduvarri Sandstone Formation, Alta, Northern Norway: A tectonic fan-delta complex, *Sedimentary Geology*, 47, 1-25, 1986.

- 1330 Bergh, S. G. and Torske, T.: Palaeovolcanology and tectonic setting of a Proterozoic metatholeiitic sequence near the Baltic Shield Margin, northern Norway, *Precambrian Research*, 39, 227-246, 1988.
- Bergh, S. G., Eig, K., Kløvjan, O. S., Henningsen, T., Olesen, O. and Hansen, J-A.: The Lofoten-Vesterålen continental margin: a multiphase Mesozoic-Palaeogene rifted shelf as shown by offshore-onshore brittle fault-fracture analysis, *Norwegian Journal of Geology*, 87, 29-58, 2007.
- 1335 Bergø, E.: Analyses of Paleozoic and Mesozoic brittle fractures in West-Finnmark, Unpublished Master's Thesis, University of Tromsø, 128 pp., 2016.
- Blystad, P., Brekke, H., Færseth, R. B., Larsen, B. T., Skogseid, J. and Tørudbakken, B.: Structural elements of the Norwegian continental shelf, Part II: The Norwegian Sea Region. *Norwegian Petroleum Directorate Bulletin*, 8, 45 pp., 1995.
- 1340 Braathen, A., Osmundsen, P-T., Nordgulen, Ø., Roberts D. and Meyer, G. B.: Orogen-parallel extension of the Caledonides in northern Central Norway: an overview, *Norwegian Journal of Geology*, 82, 225-241, 2000.
- Braathen, A., Osmundsen, P-T., Hauso, H., Semshaug, S., Fredman, N. and Buckley, S. J.: Fault-induced deformation in poorly consolidated, siliciclastic growth basin: A study from the Devonian in Norway, *Tectonophysics*, 586, 112-129, 2013.
- 1345 Breivik, A. J., Gudlaugsson, S. T. and Faleide, J-I.: Ottar Basin, SW Barents Sea: a major Upper Palaeozoic rift basin containing large volumes of deeply buried salt, *Basin Research*, 7, 299-312, 1995.
- Bugge, T., Mangerud, G., Elvebakk, G., Mørk, A., Nilsson, I., Fanavoll, S. and Vigran, J. O.: The Upper Palaeozoic succession on the Finnmark Platform, Barents Sea, *Norsk Geologisk Tidsskrift*, 75, 3-30, 1995.
- 1350 Bylund, G.: Palaeomagnetism of the Late Precambrian Vadsø and Barents Sea Groups, Varanger Peninsula, Norway, *Precambrian Research*, 69, 81-93, 1994.
- Chauvet, A. and Séranne, M.: Extension-parallel folding in the Scandinavian Caledonides: implications for late-orogenic processes, *Tectonophysics*, 238, 31-54, 1994.
- 1355 Cianfarra, P. and Salvini, F.: Lineament Domain of Regional Strike-Slip Corridor: Insight from the Neogene Transtentional De Geer Transform Fault in NW Spitsbergen, *Pure Appl. Geophys.*, 172, 1185-1201, 2015.

- 1360 Corfu, F., Armitage, P. E. B., Kullerud, K. and Bergh, S.G.: Preliminary U-Pb geochronology in the West Troms Basement Complex, North Norway: Archaean and Palaeoproterozoic events and younger overprints, *Nor. geol. unders. bull.*, 441, 61-72, 2003.
- Corfu, F., Torsvik, T. H., Andersen, T. B., Ashwal, L. D., Ramsay, D. M. and Roberts, R. J.: Early Silurian mafic-ultramafic and granitic plutonism in contemporaneous flysch, Magerøy, northern Norway: U-Pb ages and regional significance, *Journal of the Geological Society*, London, 163, 291-301, 2006.
- 1365 Corfu, F., Andersen, T. B. and Gasser, D.: The Scandinavian Caledonides: main features, conceptual advances and critical questions, in: *New Perspectives on the Caledonides of Scandinavia and Related Areas*, Corfu, F., Gasser, D. and Chew, D. M. (eds), Geological Society, London, Special Publications, 390, 9-43, 2014.
- 1370 Crider, J. G.: Oblique slip and the geometry of normal-fault linkage: mechanics and a case study from the Basin and Range in Oregon, *Journal of Structural Geology*, 23, 1997-2009, 2001.
- Cutbill, J. L. and Challinor, A.: Revision of the Stratigraphical Scheme for the Carboniferous and Permian of Spitsbergen and Bjørnøya, *Geol. Mag.*, 102, 418-439, 1965.
- Cutbill, J. L., Henderson, W. G. and Wright, N. J. R.: The Billefjorden Group (Early Carboniferous) of central Spitsbergen, *Norsk Polarinstitutt Skrifter*, 164, 57-89, 1976.
- 1375 Davids, C., Wemmer, K., Zwingmann, H., Kohlmann, F., Jacobs, J. and Bergh, S. G.: K-Ar illite and apatite fission track constraints on brittle faulting and the evolution of the northern Norwegian passive margin, *Tectonophysics*, 608, 196-211, 2013.
- Doré, A. G., Lundin, E. R., Fichler, C. and Olesen, O.: Patterns of basement structure and reactivation along the NE Atlantic margin, *Journal of the Geological Society*, London, 154, 85-92, 1997.
- 1380 Eig, K.: Onshore and offshore tectonic evolution of the Lofoten passive margin, North Norway, PhD thesis, University of Tromsø, 256 pp, 2008.
- Eig, K. and Bergh, S. G.: Late Cretaceous-Cenozoic fracturing in Lofoten, North Norway: Tectonic significance, fracture mechanisms and controlling factors, *Tectonophysics*, 499, 190-215, 2011.
- 1385 Elvevold, S., Reginiussen, H., Krogh, E. J. and Bjørklund, F.: Reworking of deep-seated gabbros and associated contact metamorphosed paragneisses in the south-eastern part of the Seiland Igneous Province, northern Norway, *Journal of Metamorphic Geology*, 12, 539-556, 1994.

- 1390 Faleide, J-I., Gudlaugsson, S. T. and Jacquart, G.: Evolution of the western Barents Sea, *Marine and Petroleum Geology*, 1, 123-150, 1984.
- Faleide, J-I., Vågnes, E. and Gudlaugsson, S. T.: Late Mesozoic-Cenozoic evolution of the southwestern Barents Sea in a regional rift-shear tectonic setting, *Marine and Petroleum Geology*, 10, 186-214, 1993.
- 1395 Faleide, J-I., Tsikalas, F., Breivik, A. J., Mjelde, R., Ritzmann, O., Engen, Ø., Wilson, J. and Eldholm, O.: Structure and evolution of the continental margin off Norway and the Barents Sea, *Episodes*, 31, 82-91, 2008.
- Fazlikhani, H., Fossen, H., Gawthorpe, R. L., Faleide, J-I. and Bell, R. E.: Basement structure and its influence on the structural configuration of the northern North Sea rift, *Tectonics*, 36, 1-27, 2017.
- 1400 Fountain, D. M., Hurich, C. A. and Smithson, S. B.: Seismic reflectivity of mylonite zones in the crust, *Geology*, 12, 195-198, 1984.
- Gabrielsen, R. H.: Long-lived fault zones and their influence on the tectonic development of the southwestern Barents Sea, *Journal of the Geological Society, London*, 141, 651-662, 1984.
- 1405 Gabrielsen R. H. and Færseth, R. B.: The inner shelf of North Cape, Norway and its implications for the Barents Shelf-Finnmark Caledonide boundary. A comment, *Norsk Geologisk Tidsskrift*, 69, 57-62, 1989.
- Gabrielsen R. H., Færseth, R. B., Jensen, L. N., Kalheim, J. E. and Riis, F.: Structural elements of the Norwegian continental shelf, Part I: The Barents Sea Region, *Norwegian Petroleum Directorate Bulletin*, 6, 33 pp., 1990.
- 1410 Gabrielsen R. H., Kløvjan, O. S., Rasmussen, A. and Stølan, T.: Interaction between halokinesis and faulting: structuring of the margins of the Nordkapp Basin, Barents Sea region, in: *Structural and Tectonic Modelling and its Application to Petroleum Geology*, Larsen, R. M., Brekke, H., Larsen, B. T. and Talleraas, E. (eds), NPF Special Publication, 1, 121-131, 1992.
- 1415 Gayer, R. A., Hayes, S. J. and Rice, A. H. N.: The structural development of the Kalak Nappe Complex of Eastern and Central Porsangerhalvøya, Finnmark, Norway, *Nor. geol. unders. bull.*, 400, 67-87, 1985.

- 1420 Gernigon L. & M. Brönnner. 2012. Late Palaeozoic architecture and evolution of the southwestern Barents Sea: insights from a new generation of aeromagnetic data. *Journal of the Geological Society*, London, vol. 127, pp. 1-11.
- Gernigon, L., Brönnner, M., Roberts, D., Olesen, O., Nasuti, A. and Yamasaki, T.: Crustal and basin evolution of the southwestern Barents Sea: From Caledonian orogeny to continental breakup, *Tectonics*, 33, 347-373, 2014.
- 1425 Gjelberg, J. G.: Upper Devonian (Famennian) – Middle Carboniferous succession of Bjørnøya, a study of ancient alluvial and coastal marine sedimentation, *Norsk Polarinstitutt Skrifter*, 174, 67 pp., 1981.
- Gjelberg, J. G.: Early-Middle Carboniferous sedimentation on Svalbard. A study of ancient alluvial and coastal marine sedimentation in rift- and strike-slip basins, PhD thesis, University of
1430 Bergen, 306 pp., 1984.
- Gjelberg, J. G. and Steel, R. J.: An outline of Lower-Middle Carboniferous sedimentation on Svalbard: Effects of tectonic, climatic and sea level changes in rift basin sequences. In: *Geology of the North Atlantic Borderlands*, Kerr, J. W. and Ferguson, A. J. (eds), *Can. Soc. Of Petrolm Geol. Mem.*, 7, 543-561, 1981.
- 1435 Gjelberg, J. G. and Steel, R. J.: Middle Carboniferous marine transgression, Bjørnøya, Svalbard: facies sequences from an interplay of sea level changes and tectonics, *Geological Journal*, 18, 1-19, 1983.
- Gudlaugsson, S. T., Faleide, J-I., Johansen, S. E. and Breivik, A. J.: Late Palaeozoic structural development of the South-western Barents Sea, *Marine and Petroleum Geology*, 15, 73-
1440 102, 1998.
- Guise, P. G. and Roberts, D.: Devonian ages from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of plagioclase in dolerite dykes, eastern Varanger Peninsula, North Norway, *Nor. geol. unders.*, 440, 27-37, 2002.
- Hallett, B. W., McClelland, W. C. and Gilotti, J. A.: The Timing of Strike-Slip Deformation Along the Storstrømmen Shear Zone, Greenland Caledonides: U-Pb Zircon and Titanite
1445 Geochronology, *Geoscience Canada*, 41, 19-45, 2014.
- Hansen, J-A. and Bergh, S. G.: Origin and reactivation of fracture systems adjacent to the Mid-Norwegian continental margin on Hamarøya, North Norway: use of digital geological mapping and morphotectonic lineament analysis, *Norwegian Journal of Geology*, 92, 391-403, 2012.

- 1450 Hansen, J-A., Bergh, S. G. and Henningsen, T.: Mesozoic rifting and basin evolution on the Lofoten and Vesterålen Margin, North-Norway; time constraints and regional implications, *Norwegian Journal of Geology*, 91, 203-228, 2012.
- Herrevold, T., Gabrielsen, R. H. and Roberts, D.: Structural geology of the southeastern part of the Trollfjorden-Komagelva Fault Zone, Varanger Peninsula, Finnmark, North Norway, 1455 *Norwegian Journal of Geology*, 89, 305-325, 2009.
- Indrevær, K. and Bergh, S. G.: Linking onshore-offshore basement rock architecture and brittle faults on the submerged strandflat along the SW Barents Sea margin, using high-resolution (5 x 5 m) bathymetry data, *Norwegian Journal of Geology*, 94, 1-34, 2014.
- Indrevær, K., Bergh, S. G., Koehl, J-B., Hansen, J-A., Schermer, E. R. and Ingebrigtsen, A.: Post-1460 Caledonian brittle fault zones on the hyperextended SW Barents Sea margin: New insights into onshore and offshore margin architecture, *Norwegian Journal of Geology*, 93, 167-188, 2013.
- Jensen, L. N. and Sørensen, K.: Tectonic framework and halokinesis of the Nordkapp Basin, Barents Sea, in: *Structural and Tectonic Modelling and its Application to Petroleum Geology*, Larsen, R. M., Brekke, H., Larsen, B. T. and Talleraas, E. (eds.), Norwegian Petroleum Society (NPF), Special Publications, 1, 109-120, 1992. 1465
- Johansen, S. E., Henningsen, T., Rundhovde, E., Sæther, B. M., Fichler, C. and Rueslåtten, H. G.: Continuation of the Caledonides north of Norway: seismic reflectors within the basement beneath the southern Barents Sea, *Marine and Petroleum Geology*, 11, 190-201, 1994.
- 1470 Kairanov, B., Escalona, A., Norton, I., Nadeau, P. and Lawver, L.: Early Cretaceous-Cenozoic relation between the Tromsø Basin and NE Greenland; Implications for margin evolution, in: *Onshore–Offshore Relationships on the North Atlantic margins*, Norsk Geologi Forening Abstracts and Proceedings, 1, 2016.
- Kirkland, C. L., Daly, J. S. and Whitehouse, M. J.: Early Silurian magmatism and the Scandian 1475 evolution of the Kalak Nappe Complex, Finnmark, Arctic Norway, *Journal of the Geological Society, London*, 162, 985-1003, 2005.
- Kirkland, C. L., Daly, J. S. and Whitehouse, M. J.: Provenance and Terrane Evolution of the Kalak Nappe Complex, Norwegian Caledonides: Implications for Neoproterozoic Paleogeography and Tectonics, *The Journal of Geology*, 115, 21-41, 2007.

- 1480 Kirkland, C. L., Daly, J. S. and Whitehouse, M. J.: Basement-cover relationships of the Kalak Nappe Complex, Arctic Norwegian Caledonides and constraints on Neoproterozoic terrane assembly in the North Atlantic region, *Precambrian Research*, 160, 245-276, 2008.
- Klein, A. C. and Steltenpohl, M. G.: Basement-cover relations and late- to post-Caledonian extension in the Leknes group, west-central Vestvågøy, Lofoten, north Norway, *Norsk Geologisk Tidsskrift*, 79, 19-31, 1999.
- 1485 Klein, A. C., Steltenpohl, M. G., Hames, W. E. and Andresen, A.: Ductile and brittle extension in the southern Lofoten archipelago, north Norway: implications for differences in tectonic style along an ancient collisional margin, *American Journal of Science*, 299, 69-89, 1999.
- Koehl, J-B., Wemmer, K. and Bergh, S. G.: Shallow faulting and westwards migration of rifting from mid-Permian times in North Norway, evidence from K/Ar radiometric dating, in: *Onshore–Offshore Relationships on the North Atlantic margins*, Norsk Geologi Forening Abstracts and Proceedings, 1, 2016.
- 1490 Koehl, J-B., Kvanli, T-E., Rowan, M. G. and Faleide, J-I.: Late Carboniferous salt tectonics in the southwesternmost Nordkapp basin, Poster presentation, ARCEX Annual Conference 2017, 10-11 May, Malangen, Norway, 2017.
- 1495 Koehl, J-B. P., Bergh, S. G., Osmundsen, P-T., Redfield, T. F., Indrevær, K., Lea, H. and Bergø, E.: Late Devonian-Carboniferous faulting in NW Finnmark and controlling fabrics, *Norwegian Journal of Geology*, submitted.
- Koyi, H., Talbot, C. J. and Tørudbakken, B. O.: Salt diapirs of the southwest Nordkapp Basin analogue modelling, *Tectonophysics*, 228, 167-187, 1993.
- 1500 Larssen, G. B., Elvebakk, G., Henriksen, S. E., Nilsson, I., Samuelsen, T. J., Svånå, T. A., Stemmerik, L. and Worsley D.: Upper Palaeozoic lithostratigraphy of the Southern Norwegian Barents Sea, *Norwegian Petroleum Directorate Bulletin*, 9, 76 pp., 2002.
- Lea, H.: Analysis of Late palaeozoic-Mesozoic brittle faults and fractures in West-Finnmark: geometry, kinematics, fault rocks and the relationship to offshore structures on the Finnmark Platform in the SW Barents Sea, Unpublished Master's Thesis, University of Tromsø, 129 pp., 2016.
- 1505 Lippard, S. J. and Prestvik, T.: Carboniferous dolerite dykes on Magerøy: new age determination and tectonic significance, *Norsk Geologisk Tidsskrift*, 77, 159-163, 1997.

- 1510 Lippard, S. J. and Roberts, D.: Fault systems in Caledonian Finnmark and the southern Barents Sea, *Nor. geol. unders. bull.*, 410, 55-64, 1987.
- Lister, G. S. and Davis, G. A.: The origin of metamorphic core complexes and detachment faults formed during Tertiary continental extension in the northern Colorado River region, U.S.A., *Journal of Structural Geology*, 11, 65-94, 1989.
- 1515 Manby, G. and Lyberis, N.: Tectonic evolution of Devonian Basin of northern Svalbard, *Norsk Geologisk Tidsskrift*, 72, 7-19, 1992.
- Mattingsdal, R., Høy, T., Simonstad, E. and Brekke, H.: An updated map of structural elements in the southern Barents Sea, Norwegian Petroleum Directorate, 2015.
- Maxant, J.: Variation of density with rock type, depth, and formation in the Western Canada basin
1520 from density logs, *Geophysics*, 45, 1061-1076, 1980.
- McCann, A. J. and Dallmann, W. K.: Reactivation of the long-lived Billefjorden Fault Zone in north central Spitsbergen, Svalbard, *Geological Magazine*, 133, 63-84, 1996.
- McClelland, W. C., Gilotti, J. A., Ramarao, T., Stemmerik, L. and Dalhoff, F.: Carboniferous basin in Holm Land records local exhumation of the North-East Greenland Caledonides:
1525 Implications for the detrital zircon signature of a collisional orogeny, *Geosphere*, 12, 925-947, 2016.
- Nasuti, A., Roberts, D. and Gernigon, L.: Multiphase mafic dykes in the Caledonides of northern Finnmark revealed by a new high-resolution aeromagnetic dataset, *Norwegian Journal of Geology*, 95, 251-263, 2015.
- 1530 Nilsen, K. T., Vendeville, B. C. and Johansen, J-T.: Influence of Regional Tectonics on Halokinesis in the Nordkapp Basin, Barents Sea, in: *Salt tectonics a global perspective*, Jackson, M. P. A., Roberts, D. G. and Snelson, S. (eds), AAPG Memoir, 65, 413-436, 1995.
- Olesen, O., Roberts, D., Henkel, H., Lile, O. B. and Torsvik, T. H.: Aeromagnetic and gravimetric interpretation of regional structural features in the Caledonides of West Finnmark and North
1535 Troms, northern Norway, *Nor. geol. unders. bull.*, 419, 1-24, 1990.
- Olesen, O., Torsvik, T. H., Tveten, E. and Zwaan, K. B.: The Lofoten-Lopphavet Project – an integrated approach to the study of a passive continental margin, Summary report, 54 pp., 1993.
- Olesen, O., Torsvik, T. H., Tveten, E., Zwaan, K. B., Løseth, H. and Henningsen, T.: Basement
1540 structure of the continental margin in the Lofoten-Lopphavet area, northern Norway:

- constraints from potential field data, on-land structural mapping and palaeomagnetic data, *Norsk Geologisk Tidsskrift*, 77, 15-30, 1997.
- 1545 Olesen, O., Brønner, M., Ebbing, J., Gellein, J., Gernigon, L., Koziel, J., Lauritsen, T., Myklebust, R., Pascal, C., Sand, M., Solheim, D. and Usov, S.: New aeromagnetic and gravity compilations from Norway and adjacent areas: methods and applications, in: *From Mature Basins to New Frontiers*, Vining, B. A. and Pickering, S. C. (eds), *Petroleum Geology, Proceedings of the 7th Petroleum Geology Conference*, 559-586, 2010.
- 1550 Omosanya, K. O., Johansen, S. E. and Harishidayat, D.: Evolution and character of supra-salt faults in the Easternmost Hammerfest Basin, SW Barents Sea, *Marine and Petroleum Geology*, 66, 1013-1028, 2015.
- Osmundsen, P-T. and Andersen, T. B.: The middle Devonian basins of western Norway: sedimentary response to large-scale transtensional tectonics, *Tectonophysics*, 332, 51-68, 2001.
- 1555 Osmundsen, P.T., Braathen, A., Sommaruga, A., Skilbrei, J. R., Nordgulen, Ø., Roberts, D., Andersen, T. B., Olesen, O. and Mosar, J.: Metamorphic core complexes along the Mid-Norwegian margin: an overview and some current ideas. In: *Onshore-offshore relations on the Mid-Norwegian margin*, Wandaas, B., Gradstein, F. et al. (eds), *Norwegian Petroleum Society, Special Publications*, 12, 29-41, 2005.
- 1560 Ottesen, D., Stokes, C. R., Rise, L. and Olsen, L.: Ice-sheet dynamics and ice streaming along the coastal parts of northern Norway, *Quaternary Science Reviews*, 27, 922-940, 2008.
- Pastore, Z., Fichler, C. and McEnroe, S. A.: The deep crustal structure of the mafic-ultramafic Seiland Igneous Province of Norway from 3-D gravity modelling and geological implications, *Geophys. J. Int.*, 207, 1653-1666, 2016.
- 1565 Pharaoh, T. C., Macintyre, R. M. and Ramsay, D. M.: K-Ar age determinations on the Raipas suite in the Komagfjord Window, northern Norway, *Norsk Geologisk Tidsskrift*, 62, 51-57, 1982.
- Pharaoh, T. C., Ramsay, D. M. and Jansen, Ø.: Stratigraphy and Structure of the Northern Part of the Repparfjord-Komagfjord Window, Finnmark, Northern Norway, *Nor. geol. unders.*, 377, 1-45, 1983.


- 1570 Phillips, T., Jackson, C. A-L., Bell, R. E., Duffy, O. B. and Fossen, H.: Reactivation of intrabasement structures during rifting: A case study from offshore southern Norway, *Journal of Structural Geology*, 91, 54-73, 2016.
- Rafaelsen, B., Elvebakk, G., Andreassen, K., Stemmerik, L., Colpaert, A. and Samuelsen, T. J.: From detached to attached carbonate buildup complexes – 3D seismic data from the Upper Palaeozoic, Finnmark Platform, southwestern Barents Sea, *Sedimentary Geology*, 206, 17-32, 2008.
- 1575
- Ramberg, I. B., Bryhni, I., Nøttvedt, A. and Rangnes, K.: The making of a land. *Geology of Norway*, The Norwegian geological Association, Oslo, 2008.
- Reeve, M. T., Bell, R. E. and Jackson, C. A. L.: Origin and significance of intra-basement seismic reflections offshore western Norway, *Journal of the Geological Society, London*, 171, 1-4, 2013.
- 1580
- Rice, A. H. N.: Restoration of the External Caledonides, Finnmark, North Norway, in: *New Perspectives on the Caledonides of Scandinavia and Related Areas*, Corfu, F., Gasser, D. and Chew, D. M. (eds), Geological Society, London, Special Publications, 390, 271-299, 2013.
- 1585
- Rise, L., Bellec, V. K., Chand, S. and Bøe R.: Pockmarks in the southwestern Barents Sea and Finnmark fjords, *Norwegian Journal of Geology*, 94, 263-282, 2015.
- Roberts, D.: Patterns of folding and fracturing in north-east Sørøy, *Nor. geol. unders.*, 269, 89-95, 1971.
- 1590
- Roberts, D.: Tectonic deformation in the Barents Sea region of Varanger Peninsula, Finnmark, *Nor. geol. unders.*, 282, 1-39, 1972.
- Roberts, D.: Palaeocurrent data from the Kalak Nappe Complex northern Norway: a key element in models of terrane affiliation, *Norwegian Journal of Geology*, 87, 319-328, 2007.
- Roberts, D. and Lippard, S. J.: Inferred Mesozoic faulting in Finnmark: current status and offshore links, *Nor. geol. unders. bull.*, 443, 55-60, 2005.
- 1595
- Roberts, D. and Onstott, T. C.: $^{40}\text{Ar}/^{39}\text{Ar}$ laser microprobe analyses and geochemistry of dolerite dykes from the Rybachi and Sredni Peninsulas, NW Kola, Russia, *Nor. geol. unders. Special Publication*, 7, 307-314, 1995.

- 1600 Roberts, D. and Siedlecka, A.: Timanian orogenic deformation along the northeastern margin of Baltica, Northwest Russia and Northeast Norway, and Avalonian-Cadomian connections, *Tectonophysics*, 352, 169-184, 2002.
- Roberts, D. and Siedlecka, A.: Provenance and sediment routing of Neoproterozoic formations on the Varanger, Nordkinn, Rybachi and Sredni peninsulas, North Norway and Northwest Russia: a review, *Nor. geol. unders. bull.*, 452, 1-19, 2012.
- 1605 Roberts, D. and Williams, G. D.: Berggrunnskart KJØLLEFJORD 2236-4 M 1:50 000, foreløpig utgave. *Nor. geol. unders.*, 2013.
- Roberts, D., Mitchell, J. G. and Andersen, T. B.: A post-Caledonian dyke from Magerøy North Norway: age and geochemistry, *Norwegian Journal of Geology*, 71, 289-294, 1991.
- Roberts, D., Olesen, O. and Karpuz, M. R.: Seismo- and neotectonics in Finnmark, Kola Peninsula and the southern Barents Sea. Part 1: Geological and neotectonic framework, *Tectonophysics*, 270, 1-13, 1997.
- 1610 Roberts, D., Chand, S. and Rise, L.: A half-graben of inferred Late Palaeozoic age in outer Varangerfjorden, Finnmark: evidence from seismic reflection profiles and multibeam bathymetry, *Norwegian Journal of Geology*, 91, 191-200, 2011.
- 1615 Røe, S-L. and Roberts, D.: Basement core sandstones in well 7128/6-1, southern Barents Sea, and correlation with onshore upper Proterozoic sequences of northeastern Finnmark, Unpublished report, University of Bergen, Geological Institute, Bergen, Norway, 33 pp., 1992.
- Rykkelid, E.: E69 "FATIMA" – Geologisk undersøkelse av tunnel under Magerøysundet, *Veglaboratoriet, Statens Vegvesen*, 1992.
- 1620 Samuelsberg, T. J., Elvebakk, G. and Stemmerik, L.: Late Paleozoic evolution of the Finnmark Platform, southern Norwegian Barents Sea, *Norwegian Journal of Geology*, 83, 351-362, 2003.
- Sartini-Rideout, C., Gilotti, J. A. and McClelland, W. C.: Geology and timing of dextral strike-slip shear zones in Danmarkshavn, North-East Greenland Caledonides, *Geological Magazine*, 143, 431-446, 2006.
- 1625 Saunders, W. B. and Ramsbottom, W. H. C.: The mid-Carboniferous eustatic event, *Geology*, 14, 208-212, 1986.

- 1630 Sengör, A. M. C.: Cross-faults and differential stretching of hanging walls in regions of low-angle normal faulting: examples from western Turkey, In: Continental Extensional Tectonics, Coward, M. P., Dewey, J. F. and Hancock P. L. (eds), Geological Society Special Publication, 28, 575-589, 1987.
- Séranne, M., Chauvet, A., Seguret, M. and Brunel, M.: Tectonics of the Devonian collapse-basins of western Norway, Bull. Soc. Géol. Fr., 8, 489-499, 1989.
- 1635 Shipton, Z. K. and Cowie, P. A.: A conceptual model for the origin of fault damage zone structures in high-porosity sandstones, Journal of Structural Geology, 25, 333-344, 2003.
- Siedlecka, A.: Late Precambrian Stratigraphy and Structure of the North-Eastern Margin of the Fennoscandian Shield (East Finnmark – Timan Region), Nor. geol. unders., 316, 313-348, 1975.
- 1640 Siedlecka, A. and Roberts, D.: The bedrock geology of Varanger Peninsula, Finnmark, North Norway: an excursion guide, Nor. geol. unders. Special Publication, 5, 1-45, 1992.
- Siedlecka, A. and Siedlecki, S.: Some new aspects of the geology of Varanger peninsula (Northern Norway), Nor. geol. unders., 247, 288-306, 1967.
- 1645 Siedlecka, A., Roberts, D., Nystuen, J. P. and Olovyanishnikov, V. G.: Northeastern and northwestern margins of Baltica in Neoproterozoic time: evidence from the Timanian and Caledonian Orogens, Geological Society, London, Memoirs, 30, 169-190, 2004.
- Siedlecki, S.: Geologisk kart over Norge, berggrunnskart Vadsø – M 1:250 000. Nor. geol. unders., 1980.
- 1650 Smelror, M., Petrov, O. V., Larssen, G. B. and Werner, S. C.: Geological history of the Barents Sea, Nor. geol. unders., Trondheim, 135 pp., 2009.
- Solheim, A. and Kristoffersen, Y.: The physical environment in the Western Barents Sea, 1: 5,000,000. Sediments above the upper regional unconformity: thickness, seismic stratigraphy and outline of the glacial history, Norsk Polarinstittutt Skrifter, 179B, 1-26, 1984.
- 1655 Steltenpohl, M. G., Hames, W. E. and Andresen, A.: The Silurian to Permian history of a metamorphic core complex in Lofoten, northern Scandinavian Caledonides, Tectonics, 23, 1-23, 2004.

- Steltenpohl, M. G., Moecher, D., Andresen, A., Ball, J., Mager, S. and Hames, W. E.: The Eidsfjord shear zone, Lofoten-Vesterålen, north Norway: An Early Devonian, paleoseismogenic low-angle normal fault, *Journal of Structural Geology*, 33, 1023-1043, 2011.
- 1660 Torgersen, E., Viola, G., Zwingmann, H. and Harris, C.: Structural and temporal evolution of a reactivated brittle-ductile fault – Part II: Timing of fault initiation and reactivation by K-Ar dating of synkinematic illite/muscovite, *Earth and Planetary Science Letters*, 407, 221-233, 2014.
- 1665 Torske, T. and Bergh, S. G.: The Caravari Formation of the Kautokaino Greenstone Belt, Finnmark, North Norway; a Palaeoproterozoic foreland basin succession, *Nor. geol. unders. bull.*, 442, 5-22, 2004.
- Townsend, C.: Thrust transport directions and thrust sheet restoration in the Caledonides of Finnmark, North Norway, *Journal of Structural Geology*, 9, 345-352, 1987a.
- 1670 Townsend, C.: The inner shelf of North Cape, Norway and its implications for the Barents Shelf-Finnmark Caledonide boundary, *Norsk Geologisk Tidsskrift*, 67, 151-153, 1987b.
- Vorren, T., Kristoffersen, Y. and Andreassen, K.: Geology of the inner shelf west of North Cape, Norway, *Norsk Geologisk Tidsskrift*, 66, 99-105, 1986.
- Wilks, W. J. and Cuthbert, S. J.: The evolution of the Hornelen Basin detachment system western Norway: implications for the style of late orogenic extension in the southern Scandinavian Caledonides, *Tectonophysics*, 238, 1-30, 1994.
- 1675 Worsley, D.: The post-Caledonian development of Svalbard and the western Barents Sea, *Polar Research*, 27, 298-317, 2008.
- Worthing, M. A.: Fracture patterns on Eastern Seiland, North Norway and their possible Relationship to Regional Faulting, *Nor. geol. unders. bull.*, 398, 35-41, 1984.
- 1680 Zhang, W. Roberts, D. and Pease, V.: Provenance of sandstones from Caledonian nappes in Finnmark, Norway: Implications for Neoproterozoic-Cambrian palaeogeography, *Tectonophysics*, 691, 198-205, 2016.
- Zwaan, K. B.: Geology of the West Troms Basement Complex, northern Norway, with emphasis on the Senja Shear Belt: a preliminary account, *Nor. geol. unders. bull.*, 427, 33-36, 1995.
- 1685 Zwaan, K. B. and Gautier, A. M.: Alta and Gargia. Description of the geological maps (AMS-M711) 1834 I and 1934 IV 1:50 000, *Nor. geol. unders.*, 357, 1-47, 1980.

Zwaan, K. B. & Roberts, D.: Tectonostratigraphic Succession and Development of the Finnmarkian Nappe Sequence, North Norway, *Nor. geol. unders.*, 343, 53-71, 1978.



1695 **Figure 1: Regional structural map of the SW Barents Sea margin (based on Bergh et al. 2007, Faleide et al. 2008, Hansen et al. 2012 and Indrevær et al. 2013 and Koehl et al. submitted). The onshore geology is from the NGU and Ramberg et al. (2008). Dashed black frame locates Figure 2. The black star marks the location of the speculated half-graben structure described in Bugge et al. (1995), which we reinterpret as a prograding sedimentary system unconformably resting on basement rocks. Location of the Barents Sea shown as a black frame in lower left inset map. Abbreviations: AFC = Asterias Fault Complex; AW = Alta-Kvænangen tectonic window; BFC = Bjørnøyrenna Fault Complex; BSFC = Bothnian-Senja Fault Complex; BKFC = Bothnian-Kvænangen Fault Complex; FPe = Finnmark Platform east; FPw = Finnmark Platform west; FTZ = Fugløya transfer zone; GL = Gjesvær Low; Ig = Ingøya; KF = Kokelv Fault; LR = Lofoten Ridge; LVF = Langfjord-Vargsund fault; Ma = Magerøya; MFC = Måsøy Fault Complex; NFC = Nysleppen Fault Complex; NH = Norsel High; NP = Nordkinn Peninsula; PP = Porsanger Peninsula; Rf = Rolvsøya fault; RLFC = Ringvassøya-Loppa Fault Complex; RW = Repparfjord-Komagfjord tectonic window; SB = Sørvær Basin; SFZ = Senja Fracture Zone; SISZ = Sørøya-Ingøya shear zone; sNB = southwesternmost Nordkapp basin; SSB = Senja Shear Belt; SØ = Sørøya; TB = Tiddlybanken Basin; TFFC = Troms-Finnmark Fault Complex; TKFZ = Trollfjorden-Komagelva Fault Zone; Va = Vannøya; VP = Varanger Peninsula; VVFC = Vestfjorden-Vanna fault complex.**

1700

1705





Figure 2: Regional structural map summarizing the architecture of the Finnmark Platform east (FPe) and west (FPw) and of the southwesternmost Nordkapp basin (sNB). The figure includes a time map of the interpreted mid-Carboniferous reflection. Grey lines show the location of seismic profiles displayed in Figure 5a-g, the red line displays the location of the seismic section shown in Figure 6d and the red frame indicates the location of seismic Z-slices described in Figure 8. White dots show the location of exploration wells and shallow drill-cores while purple blobs represent major salt diapirs in the southernmost part of the Nordkapp Basin (NBSW). See Figure 1 for abbreviations.

1710



1715 **Figure 3: Simplified stratigraphic chart of late Paleozoic sedimentary successions on the Finnmark Platform and in the southwesternmost Nordkapp basin. From left to right, columns indicate the unit name, the successions dominant lithologies and types of succession boundaries (undulating lines = erosional unconformity; straight line = conformity; dashed lines = uncertain), interpreted seismic reflections (cf. Figure 4 & Figure 5) and the units age. Lithological legend at the bottom.**

1720







Figure 4: Zoom in offshore tilt-derivative aeromagnetic data published by Gernigon et al. (2014). The white dashed line on the Finnmark Platform east represents a triangular- to rhomboid-shaped aeromagnetic low that coincides with a Carboniferous basin bounded by zig-zag-shaped brittle faults (e.g. LVF). The dotted white lines on the Finnmark Platform west and on the northern flank of the southwesternmost basin represent ENE-WSW to NE-SW trending ridges of magnetic basement rocks. The dashed black line represents a linear, NE-SW trending, high positive aeromagnetic anomaly that has been tied to the occurrence of the main segment of the MFC (cf. Indrevær et al. 2013). Dolerite dykes intruded along WNW-ESE trending fault segments of the TKFZ are shown by dotted black lines. Dashed red lines represent inferred from Gernigon et al. (2014). See Figure 1 for abbreviations.

1725









1735 **Figure 5: Examples of interpreted seismic profiles from the BSS-01 survey (2D) which locations are displayed in Figure 2. Brittle faults are shown in black and depth is in seconds (s) TWT. See Figure 1 for abbreviations; a) Interpreted seismic section that shows a system of Carboniferous horst-graben structures on the Finnmark Platform east; b) Seismic profile showing increased normal displacement across the NW-dipping LVF compared with (a) and thickening of the Carboniferous sedimentary succession within the graben bounded by the LVF. Note the insignificant amount of the displacement accommodated by the northern segment of the MFC in (a) and (b). Black arrows mark brittle faults that bound a triangular-shaped, negative aeromagnetic anomaly (cf. dashed white line in Figure 4); c) Seismic profile showing a highly thickened Carboniferous succession and potential Devonian-lower Carboniferous sedimentary rocks in the southwesternmost Nordkapp basin. Note the large offset accommodated by the main segment of the MFC and the peculiar “U” shape of the southwesternmost Nordkapp basin. Also displayed is a lateral projection of exploration well 7124/3-1; d) Interpreted seismic section that shows the listric geometries of the main segment of the MFC and of the Rolvsøya fault; e) Seismic section showing potential Devonian sedimentary rocks deposited in a NE-SW trending graben above a set of minor, SE-dipping shear zones on the Finnmark Platform west; f) Seismic section showing the listric geometries of the TFFC and MFC, which both seem to sole into the SISZ; g) NE-SW trending seismic cross-section across the Finnmark Platform west and the southwesternmost Nordkapp basin showing the gentle dip of the SISZ to the northeast and a gradual thinning of the upper Carboniferous sedimentary succession towards the southwest. A major NNE-SSW trending, SE-dipping brittle fault seems to offset the SISZ and an intra-basement reflection on the Finnmark Platform west before being truncated by the mid-Carboniferous reflection. The vertical red arrow shows the location of the imaginary prolongation of the TKFZ on the Finnmark Platform west as a comparison with the actual location of the WNW-ESE trending fault segment of the TFFC, which are separated by a distance of ca. 23 km.**

1740

1745

1750






1755

1760

1765


Figure 6: Zoom in seismic sections on the Finnmark Platform east and west. The locations of (a), (b), (c), (e), & (f) are displayed as white frames in Figure 5 and the location of (d) is shown as a red line in Figure 2. See Figure 1 for abbreviations and Figure 5 for seismic reflection legend; a) Interpreted seismic section across the Finnmark Platform east. White arrows represent high-amplitude lower Carboniferous and basement seismic reflections that are truncated upwards (toplaps) by the mid-Carboniferous reflection. Note the contrast between low-amplitude upper Carboniferous-Permian reflections, gently dipping, high-amplitude lower Carboniferous reflections and steeply dipping, high-amplitude basement reflections that possibly belong to a basement-seated shear zone (yellow dotted lines); b) uninterpreted and c) interpreted seismic zoom of a section across presumed Devonian sedimentary rocks and SE-dipping basement shear zones (yellow dotted lines) on the Finnmark Platform west; d) Interpreted seismic section from the IKU-87-BA (2D) survey showing a thick lower Carboniferous succession made up of large clinoforms (thin black lines) on the Finnmark Platform east (location in Figure 2). Note the presence of seismic artifacts in the southeast, including several multiples and NW-dipping diffraction rays; e) Interpreted seismic section across the Finnmark Platform west that displays NE-dipping basement shear zones (yellow dotted lines) including the SISZ (yellow dashed lines); f) Seismic zoom in the SISZ in the footwall of the main segment of the MFC on the Finnmark Platform west. The SISZ is composed of NW-dipping, moderate- to high-amplitude reflections that

1770 dip more gently than the MFC but that are steeper than basement reflections in the southeast. Note the significant thickness variations of the SISZ, thick in the footwall of the MFC and thin below the MFC.




1775


Figure 7: Time surface map of the top reflection of the SISZ and major brittle faults in the SW Barents Sea. Note the spoon-shaped depression formed by the SISZ on the Finnmark Platform west and southwesternmost Nordkapp basin, the abrupt change to a northeastward dip on the Finnmark Platform east, and the two narrow, NE-SW and ENE-WSW trending ridges in the footwall of the TfFc and of the Rolvsøya fault.



1780 **Figure 8:** (a) Intra-Permian seismic time-slice within 3D seismic survey MC3D-MFZ02 in the southwesternmost Nordkapp basin. Dashed black lines correspond to interpreted brittle faults; (b) Seismic time-slice within 3D seismic survey MC3D-MFZ02 near the interpreted mid-Carboniferous reflection in the southwesternmost Nordkapp basin. Black dashed lines represent interpreted brittle faults. See Figure 2 for location.




1785 **Figure 9: Thickness maps in milliseconds (ms) two-way time (TWT) of late Paleozoic sedimentary successions on the Finnmark Platform and in the southwesternmost Nordkapp basin. Color scale in (a); a) Thickness map of the Devonian-lower Carboniferous succession on the Finnmark Platform west and in the southwesternmost Nordkapp basin. The succession is thickest in the southwesternmost Nordkapp basin and represents the thickest sedimentary unit of the basin. Note that in this part of the margin, the SISZ and basin-bounding faults were used as base Devonian reflections. On the Finnmark Platform west, lower Carboniferous sedimentary rocks are missing but Devonian sedimentary deposits are**
1790 **possibly preserved in an ENE-WSW trending graben adjacent to the southwesternmost Nordkapp basin and bounded to the southeast by the MFC; b) Thickness map of the upper Carboniferous sedimentary succession showing gradual thickening of upper Carboniferous sedimentary rocks in the southwesternmost Nordkapp basin, on the Finnmark Platform west in the hanging-wall of the MFC, and on the Finnmark Platform east in the hanging-wall of the LVF and of a SE-dipping fault that parallels the MFC; c) Thickness map of the Permian succession showing very thin Permian sedimentary deposits and very mild thickness variations within the Permian sedimentary succession throughout the study area.**
1795



1800
 1805
 1810
 1815

Figure 10: Evolutionary model explaining thickness variations along the SISZ. Note that the timing of (a) to (e) is tentative. Dashed red lines in (a) to (e) correspond to tectonically active portions of the SISZ whereas dashed black lines show inactive portions. Red lines in (c), (d) & (e) show presumed normal faults. Thick vertical red arrows indicate exhumation of basement rocks along the SISZ. The model is adapted to the geometry of the SISZ observed below the Finnmark Platform west (see f); a) Extensional reactivation (thin red arrow) of the SISZ in Early Devonian times. Rapid crustal thinning and possible erosion along the upper part of the SISZ triggers exhumation of basement rocks near the coasts of NW Finnmark (thick red arrow); b) In the Early-Middle Devonian, continued extension and erosion further thin the crust and exhume basement rocks in the footwall of the SISZ, leading the upper part of the SISZ to bow. Incremental crustal thinning due to continued extensional reactivation of the SISZ and continental erosion triggers exhumation of basement rocks along lower portions of the SISZ (left-hand side, thick red arrow); c) In Mid/Late Devonian times, bowed portions of the SISZ become inactive and excision (i.e. upwards splaying; cf. Lister & Davis 1989) of the SISZ into its hanging-wall leads to thickening of the upper portion of the SISZ. Continued extension and erosion (i.e. crustal thinning) trigger bending of the lower part of the SISZ (thick red arrow) above which brittle normal faults may have formed and localized the deposition of Devonian sedimentary deposits (orange); d) Further exhumation of basement rocks along lower portions of the SISZ in the Late Devonian-early Carboniferous leads to extreme bending of the SISZ, to antithetic top-to-the-SE extensional faulting, and to early Carboniferous syn-tectonic sedimentation (green); e) Towards the end of the early Carboniferous, the lower portion of the SISZ is thickened due to incision (i.e. downward splaying; cf. Lister & Davis 1989) of the SISZ into bow-shaped portions in its footwall. Core complex exhumation ceased in the Serpukhovian and a major sea-level fall exposed the Finnmark Platform to continental erosion (green line representing the mid-Carboniferous reflection); f) Present times seismic

expression of thickness variations along the SISZ (dashed yellow) on the Finnmark Platform west. See Figure 5 for seismic reflections color schemes.



1820 **Figure 11: Map-view figures summarizing the late Paleozoic tectono-sedimentary evolution of the Finnmark Platform and**
 1825 **southwesternmost Nordkapp basin (sNB). The tectonic evolution of onshore and nearshore faults in NW Finnmark is from**
Koehl et al. (submitted). Abbreviations as in Figure 1. a) In the Early to Mid/Late Devonian, major Caledonian thrusts (e.g.
 1830 **SISZ) were inverted as low-angle extensional shear zones and exhumed metamorphic core complexes in the footwall of the**
TFFC and of the Rolvsøya fault. Thick Devonian sedimentary rocks were deposited within a spoon-shaped trough created
by the geometry of the SISZ; b) Core complex exhumation continued through the early Carboniferous, though mostly
accommodated by high-angle normal faults, which formed as brittle splays along Caledonian thrusts and shear zones (e.g.
MFC, TFFC, Rolvsøya fault and LNF). Core complex exhumation ceased by the end of the Serpukhovian and the WNW-
ESE trending fault segment of the TFFC formed as an accommodation cross-fault that decoupled the Finnmark Platform
west from the southwesternmost Nordkapp basin, thus contributing to preserve thick Devonian and lower Carboniferous
sedimentary successions in the southwesternmost Nordkapp basin while these sedimentary rocks were almost completely
eroded on the Finnmark Platform west. Minor graben and half-graben structures formed on the Finnmark Platform east.
 1835 **Precambrian, WNW-ESE to NNW-SSE trending fault zones such as the TKFZ segmented the margin and acted as minor**
transfer faults that accommodated limited amount of lateral displacement. Lateral movements along these faults ceased in
the early Carboniferous; c) In the late Carboniferous, inverted Caledonian thrusts and shear zones became inactive and
were truncated by high-angle splay-faults that accommodated the deposition of syn-tectonic sedimentary wedges on the
Finnmark Platform east and west, and of thick, partly evaporitic deposits in the southwesternmost Nordkapp basin; d) By
the end of the Carboniferous, active brittle faulting came to a halt and the Finnmark Platform and the southwesternmost
Nordkapp basin are believed to have remained tectonically quiet.