



1 **Influence of slope aspect on the microbial properties of rhizospheric**
2 **and non-rhizospheric soil on the Loess Plateau, China**

3
4 Ze Min Ai ^{1,2,3}, Jiao Yang Zhang ^{1,2,3}, Hong Fei Liu ⁴, Sha Xue ^{1,2*}, Guo Bin Liu ^{1,2}

5
6
7
8
9
10
11 ¹ State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, Institute of Soil and
12 Water Conservation, Northwest A&F University, Yangling, 712100, People's Republic of China

13 ² Institute of Soil and Water Conservation, Chinese Academy of Sciences & Ministry of Water
14 Resources, Yangling, Shaanxi 712100, People's Republic of China

15 ³ University of Chinese Academy of Sciences, Beijing 100101, People's Republic of China

16 ⁴ College of Forestry, Northwest A&F University, Yangling, Shaanxi 712100, People's Republic of
17 China

18
19
20
21
22
23
24 * Corresponding author.

25 E-mail address: xuesha100@163.com.

26 *Corresponding address: Xinong Rd. 26, Institute of Soil and Water Conservation, Northwest A&F
27 University, Yangling, Shaanxi 712100, People's Republic of China.

28
29
30
31
32
33
34



35 **Abstract.** Slope aspect is an important topographic factor, but its effect on the microbial properties of
36 grassland rhizospheric soil (RS) and non-rhizospheric soil (NRS) remain unclear. A field experiment
37 was conducted at the Ansai Research Station on the Loess Plateau in China to test the influence of
38 slope aspects (south-facing, north-facing, and northeast-facing slopes, all with *Artemisia sacrorum* as
39 the dominant species) on RS and NRS microbial biomass carbon (MBC) and phospholipid fatty acid
40 (PLFA) contents, and the rhizospheric effect (RE) of various microbial indices. MBC content differed
41 significantly among the slope aspects in RS but not in NRS, and RE for MBC content in the
42 south-facing slope was larger than that in the north-facing slope. RS total, bacterial, and gram-positive
43 bacterial PLFA contents in the south-facing slope were significantly lower than those in the north- and
44 northeast-facing slopes, and RS gram-negative bacterial (G⁻) and actinomycete PLFA contents in the
45 south-facing slope were significantly lower than those in the north-facing slope. Differently, NRS total,
46 bacterial, and G⁻ PLFA contents in the north-facing slope were significantly higher than those in the
47 south- and northeast-facing slopes, and NRS fungal and actinomycete PLFA contents in the north- and
48 south-facing slopes were significantly higher than those in the northeast-facing slope. RE for all PLFA
49 contents except fungal in the northeast-facing slope were higher than those in the south-facing slope.
50 Slope aspect significantly but differentially affected the microbial properties in RS and NRS, and the
51 variable influence was due an evident RE for most microbial properties.

52 **Keywords:** topographic factor, rhizospheric effect, phospholipid fatty acid, fungi, bacteria,
53 actinomycete

54 1 Introduction

55 As an important topographic factor, slope aspect can affect the amount of solar radiation received and
56 the angle between the ground and wind direction, which is defined as the orientation faced by a slope
57 (Selvakumar et al., 2009). Solar radiation influences ecologically critical factors of local microclimates
58 and determines soil temperature, evaporation capacity, and soil-moisture content (Carletti et al.,
59 2009; Bennie et al., 2008). Slope aspect can substantially affect soil-moisture content, water budget, and
60 soil temperatures (Sidari et al., 2008; Wang et al., 2011; Carletti et al., 2009; Sariyildiz et al., 2005).
61 South-facing slopes in the Northern Hemisphere, which receives the more solar radiation than
62 north-facing slopes, are typically hot, dry, and subject to rapid changes in seasonal and diurnal
63 microclimates. North-facing slopes are the opposite, which receive the least insolation, are cool, moist,
64 and subject to slow changes in seasonal and daily microclimates (Sariyildiz et al., 2005). The effect of
65 slope aspect on basic soil properties (pH, bulk density, and texture), nutrient (carbon, nitrogen, and
66 phosphorus) contents, microbial biomass, and enzymatic activities have been studied (Ai et al.,
67 2017a; Ascher et al., 2012; Gilliam et al., 2014; Huang et al., 2015; Sidari et al., 2008; Qin et al., 2016).
68 North-facing slopes have more microbial biomass carbon (MBC), bacteria, and actinomycetes than
69 south-facing slopes (Ascher et al., 2012; Huang et al., 2015); in contrast, other studies have found that
70 MBC and total and fungal phospholipid fatty acid (PLFA) contents were significantly higher in
71 south-facing than north-facing slopes (Huang et al., 2015; Sidari et al., 2008; Gilliam et al., 2014).
72 Gilliam et al. (2014) found that bacterial biomass did not vary with slope aspect. The different results
73 of these studies may have been due to the differences in plant species (trees vs shrubs), soil properties,
74 climatic conditions, and research methods. Previous studies mainly focused on trees and shrubs, but the
75 influence of slope aspect on grassland soil microorganisms is still unclear, even though the grassland
76 ecosystem is an important component of terrestrial ecosystems.



77 The rhizosphere is commonly defined as the narrow zone of soil adjacent to and influenced by
78 plant roots (Chen et al., 2002). The rhizosphere contains root exudates, i.e. leaked and secreted
79 chemicals, sloughed root cells, and plant debris (Warembourg et al., 2003). Microbial activity is
80 therefore high in rhizospheric soil (RS) and clearly distinct from the activity in non-rhizospheric soil
81 (NRS) due to differences in nutrient availability, pH, and redox potential (Hinsinger et al., 2009).
82 Microbial content is higher in RS than NRS (Buyer et al., 2002; Marschner et al., 2002), which is
83 known as the rhizospheric effect (RE). The effect of slope aspect on RS and NRS microbial biomass
84 and composition has not been extensively studied. Knowledge of the influence of slope aspect on the
85 differences between RS and NRS microbial communities could provide new insights into topographical
86 influences of RE on local micro-ecosystemic environments.

87 Soil microbial communities play important roles in soil quality and ecosystemic processes,
88 including nutrient cycling, decomposition of organic matter, bioremediation of structural formation,
89 and even plant interactions (Harris, 2009). These communities are closely associated with their
90 surroundings, rapidly responding to changes and environmental stresses. Soil microbes are thus
91 commonly used as sensitive indicators of change to soil quality under environmental stresses. Soil
92 respiration is widely used for measuring microbial activity (e.g. basal respiration) or determining the
93 potential microbial activity in soil (e.g. substrate-induced respiration) (Nannipieri et al., 1990; Wardle,
94 1995). Various microbial PLFAs represent the different nutritional requirements of the microbial groups.
95 Bacteria and fungi form most of the microbial biomass and represent the main drivers of organic-matter
96 turnover (Bååth and Anderson, 2003). Moreover, different kinds of bacteria produce different PLFAs:
97 Gram-negative (G^-) and Gram-positive (G^+) bacterial PLFA contents are usually considered indicators
98 of chemolithotrophic and heterotrophic bacterial communities, respectively. G^- bacteria are mainly
99 associated with roots and thus decompose low-molecular-weight organic molecules (Griffiths et al.,
100 1999), whereas G^+ bacteria decompose more complex materials, such as organic matter and litter
101 (Kramer and Gleixner, 2006). These microbial indices are all sensitive bio-indicators that can be used
102 to estimate soil quality and the effect of slope aspect on RS and NRS microbial communities. Soil
103 ecologists have long been interested in the response of microbial communities to environmental factors
104 for understanding the underlying mechanisms determining the content and composition of microbial
105 biomass. Microbial communities have a close relationship with pH, carbon (organic and water-soluble
106 organic carbon), nitrogen (total nitrogen, ammonium and nitrate nitrogen, and water-soluble
107 ammonium and nitrate nitrogen), and phosphorus (total and available phosphorus) (Bardelli et al.,
108 2017; Huang et al., 2014; Nilsson et al., 2005; Ma et al., 2015). The effect of slope aspect on the main
109 soil nutrient factors that affect RS and NRS microbial communities, however, remains unclear.

110 The Chinese government introduced the Grain for Green Project in the 1990s to control soil
111 erosion and improve the ecological environment of the Loess Plateau by converting large areas of
112 sloping cropland to forest and grassland. *Artemisia sacrorum*, a perennial herb with multiple branches,
113 well-developed root suckers, and high seed production and fertility, is widely distributed on the plateau
114 (Wang and Liu, 2002), especially in the converted grassland. We selected *A. sacrorum* as a typical
115 grassland plant of this region to study the effect of slope aspect on the MBC, total, fungal, bacterial,
116 and actinomycete PLFA contents in RS and NRS and the differences of their REs. We also identified
117 the main RS and NRS environmental factors affecting microbial content and composition. We tested
118 three slope aspects (south-facing, north-facing, and northeast-facing slopes) with the same
119 rehabilitation age on the Loess Plateau in China. We tested the following hypotheses: (1) slope aspect
120 would significantly but differentially affect the MBC, total, fungal, bacterial, and actinomycete PLFA



121 contents and their REs; and (2) soil carbon (C) and nitrogen (N) would have the larger effect on the RS
122 and NRS microbial communities, and different C and N compounds would have different effects.

123 2 Materials and methods

124 2.1 Study site

125 A field experiment was conducted at the Ansai Research Station (ARS) of the Chinese Academy of
126 Sciences (36°51'30"N, 109°19'23"E; 1068–1309 m a.s.l.), northern Loess Plateau, China. The mean
127 annual temperature in the study area is 8.8 °C, and the mean annual precipitation is approximately 505
128 mm, with >70% concentrated from July to September. Annual evaporation ranges from 1500 to 1800
129 mm. Three grassland areas abandoned in the same year were selected for the experiment. Details on
130 soil properties were described in Ai et al. (2017a).

131 2.2 Experimental design and soil sampling

132 The slopes of the three grassland areas were south-facing (S15°W), northeast-facing (N75°W), and
133 north-facing (N57°E). The three study areas were selected in September 2014 after consultation with
134 ARS researchers and reviewing relevant land documents. The basic characteristics are shown in Table 1.
135 Three replicate 10 × 10 m plots were established at each site (*A. sacrorum* was the dominant plant at
136 each site). Each plot was first surveyed for latitude, longitude, elevation, slope aspect, and slope
137 gradient. Three 1 × 1 m quadrats were then randomly set in each plot to characterise the vegetation, e.g.
138 plant species, coverage, and number. The plants were removed, and the soil strongly adhering to the
139 roots, i.e. RS, was collected (0–20 cm soil layer). Soil was also sampled from the same layer at
140 locations approximately 15 cm from the plant roots (i.e. NRS). Each NRS sample was a composite of
141 subsamples collected at five points (the four corners and the centre of the plot). A total of 18 soil
142 samples (3 sites × 3 plots per site × 2 soil types) were collected, and each was divided into two
143 subsamples: one subsample was placed in a cool container, and the other was placed into a cloth bag.
144 The samples were then taken to the laboratory, and gravel and coarse fragments were removed. The
145 container samples were homogenised and sieved to 2 mm and were also divided into two subsamples:
146 one subsample was stored at -80 °C, and the other was stored at 4 °C until analysis. The samples in the
147 cloth bags were air-dried and sieved to 0.25 and 1 mm prior to analysis.

148 **Table 1.** Characteristics of the sampling sites.

Slope aspect	Latitude (°N)	Longitude (°E)	Altitude (m)	Plant community
S15°W	36.85	109.31	1269	<i>A. sacrorum</i> + <i>Bothriochloa ischaemum</i>
N75°W	36.85	109.31	1275	<i>A. sacrorum</i> + <i>Phragmites australis</i>
N57°E	36.85	109.31	1278	<i>A. sacrorum</i> + <i>Artemisia capillaries</i>

149 2.3 Laboratory analysis

150 The samples stored at 4 °C were used for determining MBC content, basal respiration (BR), and
151 substrate-induced respiration (SIR). Microbial biomass was measured by chloroform fumigation
152 (Vance et al., 1987). The soil samples that were fumigated for 24 h at 25.8 °C with CHCl₃ (ethanol free)
153 after the fumigation and non-fumigation treatments, and then were extracted with 100 ml of 0.5 M
154 K₂SO₄ by horizontal shaking for 1 h at 200 rpm and then filtered. The amount of K₂SO₄-extracted
155 organic C was determined by a liquiTOCII analyser (Elementar, Hanau, Germany) and MBC content



156 was calculated using a k_{EC} factor of 0.38 (Vance et al., 1987). BR and SIR were measured by an
 157 infrared gas analyser (QGS-08B, Beijing, China) (Hueso et al., 2011). The metabolic quotient was
 158 calculated as BR per unit MBC (BR/MBC) (Anderson and Domsch, 1993).

159 The soil stored at $-80\text{ }^{\circ}\text{C}$ was used for the determination of PLFA contents. The structures of the
 160 microbial communities were determined using a method (Bligh and Dyer, 1959) modified by Bardgett
 161 et al. (1996). Briefly, fatty acids were extracted from 3.0 g of freeze-dried soil using a solution
 162 containing citrate buffer, chloroform, and methanol. The PLFAs were separated from neutral and
 163 glycolipid fatty acids by solid-phase-extraction chromatography. After mild alkaline methanolysis, the
 164 PLFAs were analysed using a gas chromatograph (GC7890A, Agilent Technologies Inc., Wilmington,
 165 USA) equipped with MIDI Sherlock software (Version 4.5; MIDI Inc., Newark, USA). An external
 166 standard of 19:0 methyl ester was used for quantification (Frostegård et al., 1993), and the amounts
 167 were expressed as nmol g^{-1} for dry soil.

168 According to Zelles (1999), specific PLFA signatures can serve as indicators of specific microbial
 169 groups. Total PLFAs were obtained by summing the contents of all fatty acids detected in each sample.
 170 The classification PLFA was shown in Table 2.

171 **Table 2.** The characterization of the microbial phospholipid fatty acids.

Microbial group	Specific PLFA markers
Gram-positive bacteria	11:0 anteiso, 12:0 anteiso, 13:0 iso, 13:0 anteiso, 14:0 iso, 14:0 anteiso, 15:0 iso, 15:0 anteiso, 15:1 iso w6c, 15:1 iso w9c, 16:0 iso, 16:0 anteiso, 17:0 iso, 17:0 anteiso, 18:0 iso, 19:0 iso, 19:0 anteiso, 22:0 iso
Gram-negative bacteria	12:1 w4c, 12:1 w8c, 14:1 w5c, 14:1 w8c, 14:1 w9c, 15:1 w5c, 15:1 w7c, 15:1 w8c, 16:1 w7c DMA, 16:1 w7c, 16:1 w9c DMA, 17:0 cyclo w7c, 17:1 w5c, 17:1 w7c, 17:1 w8c, 18:1 w5c, 18:1 w6c, 18:1 w7c, 18:1 w8c, 18:1 w9c, 19:0 cyclo w6c, 19:0 cyclo w7c, 19:1 w6c, 19:1 w8c, 20:1 w6c, 20:1 w9c, 21:1 w3c, 21:1 w5c, 21:1 w6c, 22:1 w3c, 22:1 w5c, 22:1 w6c, 22:1 w8c, 22:1 w9c, 24:1 w9c, 19:0 cyclo 9,10 DMA
Fungi	16:1w5c, 18:2w6c
Actinomycetes	16:0 10-methyl, 17:0 10-methyl, 17:1 w7c 10-methyl, 18:0 10-methyl, 18:1 w7c 10-methyl, 19:1 w7c 10-methyl, 20:0 10-methyl

172 The concentrations of soil organic carbon (SOC), total nitrogen (TN), and total phosphorus at the
 173 sites have been reported by Ai et al. (2017a). Soil pH and available phosphorus (SAP), ammonium N
 174 (NH_4), nitrate N (NO_3), water-soluble organic C (WSOC), water-soluble NH_4 (WNH_4), and
 175 water-soluble NO_3 (WNO_3) contents were measured as described by Ai et al (2017b).

176 2.4 Calculations and statistical analysis

177 RE was calculated as: $\text{RE}=\text{Rs}/\text{NRs}$, where Rs is a microbial property in RS, and NRs is a microbial
 178 property in NRS (Mukhopadhyay et al., 2016). All data were analysed using one-way ANOVAs,
 179 followed by Duncan's tests at a probability level of $P<0.05$ for multiple comparisons. All statistical
 180 analyses were performed using SPSS 20.0 (SPSS Inc., Chicago, USA), and structural equation models
 181 (SEMs) were analysed using the AMOS SPSS expansion pack. Redundancy analysis (RDA) was
 182 performed using CANOCO 5.0 (Biometris, Wageningen, The Netherlands). The graphs were plotted
 183 using SigmaPlot 12.5 (Systat Software, San Jose, USA).

184 3 Results



185 3.1 MBC content, respiration, and BR/MBC

186 RS MBC content did not differ significantly among the slope aspects, but NRS MBC content in the
 187 north-facing slope was higher than those in the south- and northeast-facing slopes (Fig. 1A). RE in the
 188 south-facing slope was highest among the slope aspects (Fig. 3). Slope aspect did not affect BR,
 189 BR/MBC, or SIR in either RS or NRS (Fig. 1B and Table 3). The RE for BR did not differ significantly
 190 among the slope aspects (Fig. 3). The RE for SIR in the south-facing slope was higher than that in the
 191 north-facing slope.

192 3.2 Microbial PLFA contents and composition

193 The microbial PLFA contents in RS differed significantly among the slope aspects. Total PLFA
 194 contents in the north- and northeast-facing slopes were 115 and 88% higher, respectively, than that in
 195 the south-facing slope (Fig. 2A). Fungal PLFA content did not differ significantly among the slope
 196 aspects. Bacterial PLFA content was similar to the trend for total PLFA content, with the lowest content
 197 in the south-facing slope. In contrast to total PLFA content, the ratio of fungal PLFA content to
 198 bacterial PLFA content (F/B ratio) in the south-facing slope was significantly higher than those in the
 199 north- and northeast-facing slopes (Table 3). Both G^+ and G^- PLFA contents had trends similar to that
 200 of the bacterial PLFA content, with the lowest contents in the south-facing slope (Fig. 2A). The ratio of
 201 G^+ PLFA content to G^- PLFA content (G^+/G^- ratio) did not differ significantly among the slope aspects
 202 (Table 3). With a trend similar to that of G^- PLFA content, actinomycete PLFA content in the
 203 north-facing slope was 102% higher than that in the south-facing slope.

204 **Table 3.** Microbial respiratory quotients (BR/MBC), ratios of fungal PLFA content to bacteria PLFA
 205 content (F/B), and ratios of G^+ PLFA content to G^- PLFA content (G^+/G^-) in the rhizospheric and
 206 non-rhizospheric soils.

Slope aspect	Rhizospheric soil			Non-rhizospheric soil		
	BR/MBC (10^3 h^{-1})	F/B ratio	G^+/G^- ratio	BR/MBC (10^3 h^{-1})	F/B ratio	G^+/G^- ratio
South-facing	3.03 ± 0.49a	0.07 ±	2.16 ±	2.47 ± 0.52a	0.07 ±	1.45 ±
		0.00a	0.58a		0.00a	0.23a
North-facing	2.67 ± 0.41a	0.03 ±	1.55 ±	2.00 ± 0.26a	0.04 ±	1.20 ±
		0.00b	0.29a		0.00b	0.10a
Northeast-facing	2.77 ± 0.23a	0.04 ±	1.54 ±	2.47 ± 0.35a	0.05 ±	1.33 ±
		0.00b	0.16a		0.00ab	0.06a

207 The composition of the NRS PLFA contents also differed significantly among the slope aspects.
 208 Total PLFA content in the north-facing slope was 50 and 62% higher than those in the south- and
 209 northeast-facing slopes, respectively (Fig. 2B). Bacterial PLFA content had a trend similar to that of
 210 total PLFA content, with the highest content in the north-facing slope. Fungal PLFA content in the
 211 south- and north-facing slopes was significantly higher than that in the northeast-facing slope. The F/B
 212 ratio in the south-facing was substantially higher than that in the north-facing slope (Table 3). G^- PLFA
 213 content had a trend similar to that of bacterial PLFA content, and G^+ PLFA content did not differ
 214 significantly among the slope aspects (Fig. 2B). The G^+/G^- ratio did not differ significantly among the
 215 slope aspects (Table 3). Actinomycete PLFA content had a trend similar to that of fungal PLFA content,
 216 with the higher contents in the south- and north-facing slopes, which were 49 and 117% higher,
 217 respectively, than that in the northeast-facing slope (Fig. 2B).

218 The REs for total, bacterial, G^+ , G^- , and actinomycete PLFA contents differed significantly among



219 the slope aspects, but not the RE for fungal PLFA content (Fig. 3). The REs for total, G^+ , G^- , and
220 bacterial PLFA contents in the northeast-facing slope were highest among the slope aspects. The RE for
221 actinomycete PLFA content in the northeast-facing slope was highest among the slope aspects.

222 3.3 Redundancy analysis (RDA)

223 The constrained RDAs indicated that environmental factors affected RS microbial characteristics (Fig.
224 4A). The total variation was 6.10, and the explanatory variables accounted for 96.8%. The first two
225 axes (RDA1 and RDA2) explained 89.6% of the total variance, wherein 84.1% was attributed to RDA1
226 and 5.5% to RDA2. WSOC content was the most significant of the seven environmental factors and
227 explained 63.6% ($P=0.006$) of the total variance. The slope aspect was the next most significant
228 environmental variable and explained 62.8% ($P=0.004$), followed by NH_4 (58.6%, $P=0.004$), SAP
229 (45.7%, $P=0.022$), and WNH_4 (45.2%, $P=0.032$) contents.

230 The constrained RDAs indicated that environmental factors affected NRS microbial characteristics
231 (Fig. 4B). The total variation was 2.97, and the explanatory variables accounted for 94.2%. RDA1 and
232 RDA2 explained 81.6% of the total variance, 68.3% for RDA1 and 13.3% for RDA2. WNO_3 content
233 was the most significant of the seven environmental factors and explained 34.7% ($P=0.04$) of the total
234 variance. Slope aspect (33.7%, $P=0.054$) and WNH_4 content (32.8%, $P=0.092$) also played important
235 roles.

236 3.4 Path analysis

237 The final SEM based on all indices adequately fitted the data to describe the effects of the
238 environmental factors on RS microbial characteristics ($\chi^2=0.506$; $P=0.918$; RMSEA, $P<0.001$;
239 standardised path coefficients are shown in Fig. 5A). The final model accounted for 99% of the
240 variation in RS WSOC content, with 71% of the variation in bacterial PLFA content, 78% of the
241 variation in G^+ PLFA content, and 72% of the variation in total PLFA content. Slope aspect was
242 positively correlated with WSOC content ($P<0.001$). WSOC content was negatively correlated with
243 bacterial PLFA ($P<0.001$), G^+ PLFA ($P<0.001$), and total PLFA ($P<0.001$) contents.

244 All indices adequately fitted the data to describe the effects of the environmental factors on NRS
245 microbial characteristics ($\chi^2=3.222$; $P=0.521$; RMSEA, $P<0.001$; standardised path coefficients are
246 shown in Fig. 5B). The model was able to explain 59% of the variation in WNH_4 content, 58% of the
247 variation in MBC content, 55% of the variation in G^- PLFA content, and 45% of variation in total PLFA
248 content. Slope aspect was strongly positively correlated with WNH_4 content ($P<0.001$). WNH_4 content
249 was strongly negatively correlated with MBC ($P<0.001$), G^- PLFA ($P<0.05$), and total PLFA ($P<0.05$)
250 contents.

251 4 Discussion

252 4.1 MBC, respiration, and BR/MBC

253 Soil microbial biomass is closely associated with soil-moisture content (Zhang et al., 2005; Drenovsky
254 et al., 2010; Ma et al., 2015). The north-facing slope contained more moisture than the south-facing
255 slope (Sariyildiz et al., 2005), so microbial activity in the north-facing slope was higher than that in the
256 south-facing slope. NRS MBC content in the north-facing slope was significantly higher than that in
257 the south-facing slope in our study, supporting our hypothesis 1 and in agreement with other studies
258 (Huang et al., 2015; Sidari et al., 2008). Carletti et al. (2009), however, reported an opposite trend: soil



259 MBC was higher in a south-facing slope. This disparity may have been due to the differences in plant
260 species, soil type, and regional climate (Gilliam et al., 2014). We also found that NRS MBC content in
261 the northeast-facing slope was lower than that in the north-facing slope, inconsistent with Huang et al.
262 (2015), whose study area had the same soil and climatic conditions as ours. We speculate that the
263 different result may mainly due to the different plant species: studied, the effect of shrubland plants
264 (Huang et al., 2015) on NRS may be different from grassland plants (ours). As plant shade can affect
265 soil microbial activity (Blok et al., 2010), different shading can cause different MBC contents.

266 Plant roots release a high amount of exudates, such as sugars, amino acids, organic acids,
267 hormones, and enzymes (Zhang et al., 2012; Grayston et al., 1997). In contrast, soil with a low amount
268 of shading is prone to desiccation (Wang et al., 2008), and the quantity of exudates released by plant
269 roots is low, which may lead to lower activities of the microorganisms. RS MBC content therefore
270 should be higher than NRS MBC content, consistent with our results. In our study, the RE for MBC
271 content in the south-facing slope was significantly higher than that in the north-facing slope.
272 South-facing slopes in the Northern Hemisphere receive more sunlight, which would have a greater
273 impact on the soil micro-environmental light than that in north-facing slopes between RS and NRS.
274 The RDA and path analysis found that NRS WSOC, WNO_3 , and WNH_4 contents were well correlated
275 with MBC content (Figs. 4B and 5B), supporting our hypothesis 2 and in agreement with other studies
276 (Haynes, 2000; Huang et al., 2014).

277 Neither RS nor NRS BR, SIR, and BR/MBC differed significantly among the slope aspects,
278 indicating that the actual microbial activities, potential microbial activities, and bioenergetic status of
279 the microbial biomass (Nannipieri et al., 1990; Wardle, 1995; Sinha et al., 2009) were similar among the
280 slope aspects in the study area. The RE of SIR in the south-facing slope was 96% higher than that in
281 the north-facing slope, indicating that the effect of slope aspect on the RS and NRS SIRs was more
282 evident in the south-facing slope than that in the north-facing slope, even though the influence of slope
283 aspect on SIR was not significant either in RS or NRS.

284 4.2 PLFA contents and composition

285 4.2.1 Fungal and bacterial PLFA contents and composition

286 NRS fungal PLFA content in the northeast-facing slope was lower than those in the south- and
287 north-facing slopes, however, RS fungal PLFA content did not differ significantly among the slope
288 aspects. Previous studies have reported different results: Huang et al. (2015) and Gilliam et al. (2014)
289 found that slope aspect significantly affected the fungal community, and fungal abundance was lower
290 in north-facing slope; Bardelli et al. (2017) found that fungal abundance did not differ significantly
291 between north- and south-facing slopes. These different results may due to the differences in plant
292 species (e.g. herbs vs shrubs), soil conditions, climate, and research methods (Gilliam et al., 2014). The
293 different responses of RS and NRS fungal PLFA contents meant that rhizospheres could form an
294 environment that negates the effect of slope aspect on fungal communities more than in
295 non-rhizospheric zones. SOC and TN can supply the microbial biomass with enough C, N, and energy
296 resources to support microbial growth (Jia et al., 2005), so the solubility of SOC (WSOC) and TN
297 (WNO_3 , WNH_4) would be closely associated with the fungal community. The RDA showed that NRS
298 WSOC and WNO_3 were well correlated with fungal PLFA content (Fig. 4B), supporting our hypothesis
299 2 and agreed by previous studies (Haynes, 2000; Nilsson et al., 2005; Huang et al., 2014).

300 The effect of slope aspect on PLFA content differed between bacteria and fungi. Both RS and NRS
301 bacterial PLFA contents in the south-facing slope were lower than those in the north-facing slope,



302 suggesting more soil moisture in the north-facing slope suitable for the growth of bacteria, in
303 agreement with some studies (Huang et al., 2015; Ascher et al., 2012) but not others (Gilliam et al.,
304 2014; Bardelli et al., 2017). The effect of slope aspect on the bacterial community would therefore
305 become significant due to the plant species, soil type, and climatic conditions. The RE for bacterial
306 PLFA content in the northeast-facing slope was significantly higher than that in the south-facing slope,
307 indicating that the environmental conditions of the rhizosphere helped the bacterial community to resist
308 environmental pressure. The RDA indicated that the RS WNH_4 , WSOC, and SAP contents were well
309 correlated with the bacterial PLFA content, and the NRS WNH_4 content was well correlated with the
310 bacterial PLFA content (Fig. 4A, B). The path analysis indicated that RS WSOC content was the main
311 factor influencing the bacterial PLFA content and mainly affected the G^+ PLFA content (Fig. 5A), in
312 agreement with another study (Fierer et al., 2003), but the NRS WNH_4 content mainly affected the G^-
313 PLFA content (Fig. 5B). These results indicated that the RS and NRS bacterial PLFA contents were
314 affected by different soil nutrient factors.

315 Soil moisture is an important environmental factor affecting the composition of microbial
316 communities, the higher amounts of soil moisture in north-facing slopes (Sariyildiz et al., 2005) can
317 lead to lower F/B ratios (Brockett et al., 2012; Drenovsky et al., 2010; Ma et al., 2015). In this paper, the
318 F/B ratio was highest in the north-facing slope and lowest in the south-facing slope for both RS and
319 NRS, consistent with previous studies (Huang et al., 2015; Gilliam et al., 2014). The higher amount of
320 soil moisture in the north-facing slope would reduce soil aeration, lower oxygen levels would create an
321 environment favourable for facultative and obligate anaerobic bacteria (Drenovsky et al., 2004).
322 Drought stress in the south-facing slope would likely facilitate the survival of fungi, because soil fungi
323 rely on more aerobic conditions and are more tolerant of drought due to their filamentous nature
324 (Zhang et al., 2005).

325 The significant difference in the RE for bacterial PLFA content was not obvious for fungal PLFA
326 content, so the RE was much weaker in the fungal than the bacterial community, consistent with Buyer
327 et al. (2002). These results indicated that RE had a large effect on the structures of the fungal and
328 bacterial communities. RE was significantly affected by slope aspect for both the G^+ and G^- PLFA
329 contents, and their REs were consistent with the RE of the total bacterial PLFA content. The G^+/G^-
330 ratio can indicate the dominance of bacteria in soil microbial communities (Tscherko et al., 2004; Zhang
331 et al., 2015). Neither the RS nor the NRS G^+/G^- ratio was affected by slope aspect, indicating that slope
332 aspect did not significantly affect the dominant bacterial community in either RS or NRS.

333 4.2.2 Actinomycete and total PLFA contents

334 RS and NRS actinomycete PLFA contents were significantly affected by slope aspect, supporting our
335 hypothesis 1. Actinomycetes and G^+ bacteria have similar life habits, so wetter soils are more enriched
336 in actinomycetes (Zhang et al., 2005; Drenovsky et al., 2010; Ma et al., 2015). RS actinomycete PLFA
337 content in the north-facing slope was therefore higher than that in the south-facing slope, and the
338 northeast-facing slope had more moderate growth conditions for actinomycetes compared with the
339 north-facing and south-facing slopes. NRS actinomycete PLFA content, however, was lower in the
340 northeast-facing slope than that in the south-facing slope. This difference may have been due to RE,
341 because RE in the northeast-facing slope was significantly higher than those in the other slopes. RE
342 will affect soil nutrients more in RS than NRS (Zhang et al., 2012; Grayston et al., 1997). The RDA
343 indicated that the RS but not NRS actinomycete PLFA content was well correlated with WSOC and
344 WNH_4 contents, supporting our hypothesis 2 (Fig. 4A, B).



345 The G⁺ and actinomycete PLFA contents accounted for more than 50% of total PLFA content in
346 both RS (57–59%) and NRS (54–58%), so the distribution of total PLFA content in our study area
347 depended mainly on the G⁺ and actinomycete PLFA contents. Drier soils tend to be more enriched in G⁺
348 bacteria and fungi, whereas wetter soils tend to be more enriched in G⁺ bacteria and actinomycetes
349 (Zhang et al., 2005; Drenovsky et al., 2010; Ma et al., 2015), so total PLFA contents in both RS and
350 NRS were highest in the north-facing slope. Total PLFA content, however, was higher in the
351 northeast-facing slope than that in the south-facing slope for RS, and did not differ significantly
352 between the northeast-facing and south-facing slopes in NRS. These differences in total PLFA content
353 between RS and NRS may have been mostly due to RE. The shading by herbs in the northeast-facing
354 slope may make RS was suitable for microbial life as in the north-facing slope, whereas NRS in the
355 northeast-facing slope was not suitable for microbial life as in the south-facing slope without plant
356 shading. The path analysis indicated that WSOC content had a significant effect on RS total PLFA
357 content and that WNH₄ content had a significant effect on NRS total PLFA content (Fig. 5A, B), as
358 expected (Haynes, 2000; Huang et al., 2014; Nilsson et al., 2005). These results supported our
359 hypothesis 1 and 2, but hypothesis 1 were inconsistent with Huang et al. (2015) who found a significantly
360 higher total PLFA content in the south-facing slope than in other slopes. RE may be one of the main
361 reasons, because shrub shading (Huang et al., 2015) clearly differs from herb shading (our study),
362 which could be caused a different RE (Blok et al., 2010).

363 5 Conclusions

364 This study provides experimental evidence that slope aspect can markedly but differentially affect
365 MBC and PLFA contents in RS and NRS, and the different influences can produce an evident RE; the
366 RE for most microbial properties was higher in the northeast-facing slope. WSOC content was well
367 correlated with RS microbial properties, and WNH₄ content was well correlated with NRS microbial
368 properties, likely due to RE. Studies of the influence of slope aspect on soil microbial communities
369 should therefore consider REs. This study provides new insights into the influences of topographic
370 factors affecting the mechanisms driving the structure of microbial communities in a
371 micro-ecosystemic environment. Further field investigation on different plant species, however, is
372 needed to determine the role of RE under the effect of slope aspect in micro-ecosystemic environments.

373 Author contributions

374 GbL and SX provided research ideas and designed the experiments. They were also responsible for the
375 revision of the paper. SX, ZmA, JyZ and HfL participated in the soil sample collection, ZmA, JyZ and
376 HfL contributed to the soil analysis. ZmA analyzed the data and wrote the paper.

377

378 *Acknowledgements.* We thank the National Natural Science Foundation of China (41371510, 41671513,
379 41471438) and the West Young Scholars Project of The Chinese Academy of Sciences (XAB2015A05)
380 for funding this work.

381 References

382 Ai, Z. M., He, L. R., Xin, Q., Yang, T., Liu, G. B., and Xue, S.: Slope aspect affects the non-structural



- 383 carbohydrates and C: N: P stoichiometry of *Artemisia sacrorum* on the Loess Plateau in China,
384 Catena, 152, 9-17, 2017a.
- 385 Ai, Z. M., Zhang, J. Y., Liu, H. F., Xin, Q., Xue, S., and Liu, G. B.: Soil nutrients influence the
386 photosynthesis and biomass in invasive *Panicum virgatum* on the Loess Plateau in China,
387 Plant and Soil, 418, 153-164, 2017b.
- 388 Anderson, T.-H., and Domsch, K.: The metabolic quotient for CO₂ (qCO₂) as a specific activity
389 parameter to assess the effects of environmental conditions, such as pH, on the microbial
390 biomass of forest soils, Soil Biology and Biochemistry, 25, 393-395, 1993.
- 391 Ascher, J., Sartori, G., Graefe, U., Thornton, B., Ceccherini, M., Pietramellara, G., and Egli, M.: Are
392 humus forms, mesofauna and microflora in subalpine forest soils sensitive to thermal
393 conditions?, Biology and Fertility of Soils, 48, 709-725, 2012.
- 394 Bååth, E., and Anderson, T.-H.: Comparison of soil fungal/bacterial ratios in a pH gradient using
395 physiological and PLFA-based techniques, Soil Biology and Biochemistry, 35, 955-963, 2003.
- 396 Bardelli, T., Gómez-Brandón, M., Ascher-Jenull, J., Fornasier, F., Arfaioli, P., Francioli, D., Egli, M.,
397 Sartori, G., Insam, H., and Pietramellara, G.: Effects of slope exposure on soil
398 physico-chemical and microbiological properties along an altitudinal climosequence in the
399 Italian Alps, Science of The Total Environment, 575, 1041-1055, 2017.
- 400 Bardgett, R. D., Hobbs, P. J., and Frostegård, Å.: Changes in soil fungal: bacterial biomass ratios
401 following reductions in the intensity of management of an upland grassland, Biology and
402 Fertility of Soils, 22, 261-264, 1996.
- 403 Bennie, J., Huntley, B., Wiltshire, A., Hill, M. O., and Baxter, R.: Slope, aspect and climate: Spatially
404 explicit and implicit models of topographic microclimate in chalk grassland, Ecological
405 Modelling, 216, 47-59, 2008.
- 406 Bligh, E. G., and Dyer, W. J.: A rapid method of total lipid extraction and purification, Canadian
407 journal of biochemistry and physiology, 37, 911-917, 1959.
- 408 Blok, D., Heijmans, M. M., SCHAEPMAN - STRUB, G., Kononov, A., Maximov, T., and Berendse, F.:
409 Shrub expansion may reduce summer permafrost thaw in Siberian tundra, Global Change
410 Biology, 16, 1296-1305, 2010.
- 411 Brockett, B. F. T., Prescott, C. E., and Grayston, S. J.: Soil moisture is the major factor influencing
412 microbial community structure and enzyme activities across seven biogeoclimatic zones in
413 western Canada, Soil Biology and Biochemistry, 44, 9-20, 2012.
- 414 Buyer, J. S., Roberts, D. P., and Russek-Cohen, E.: Soil and plant effects on microbial community
415 structure, Canadian Journal of Microbiology, 48, 955-964, 2002.
- 416 Carletti, P., Vendramin, E., Pizzeghello, D., Concheri, G., Zanella, A., Nardi, S., and Squartini, A.: Soil
417 humic compounds and microbial communities in six spruce forests as function of parent
418 material, slope aspect and stand age, Plant and Soil, 315, 47-65, 10.1007/s11104-008-9732-z,
419 2009.
- 420 Chen, C., Condon, L., Davis, M., and Sherlock, R.: Phosphorus dynamics in the rhizosphere of
421 perennial ryegrass (*Lolium perenne* L.) and radiata pine (*Pinus radiata* D. Don.), Soil Biology
422 and Biochemistry, 34, 487-499, 2002.
- 423 Drenovsky, R., Vo, D., Graham, K., and Scow, K.: Soil water content and organic carbon availability
424 are major determinants of soil microbial community composition, Microbial Ecology, 48,
425 424-430, 2004.
- 426 Drenovsky, R. E., Steenwerth, K. L., Jackson, L. E., and Scow, K. M.: Land use and climatic factors



- 427 structure regional patterns in soil microbial communities, *Global Ecology and Biogeography*,
428 19, 27-39, 2010.
- 429 Fierer, N., Schimel, J. P., and Holden, P. A.: Variations in microbial community composition through
430 two soil depth profiles, *Soil Biology and Biochemistry*, 35, 167-176, 2003.
- 431 Frostegård, Å., Tunlid, A., and Bååth, E.: Phospholipid fatty acid composition, biomass, and activity of
432 microbial communities from two soil types experimentally exposed to different heavy metals,
433 *Applied and Environmental Microbiology*, 59, 3605-3617, 1993.
- 434 Gilliam, F. S., Håll, R., Chudomelová M., McCulley, R. L., and Nelson, J. A.: Variation in vegetation
435 and microbial linkages with slope aspect in a montane temperate hardwood forest, *Ecosphere*,
436 5, 1-17, 2014.
- 437 Grayston, S., Vaughan, D., and Jones, D.: Rhizosphere carbon flow in trees, in comparison with annual
438 plants: the importance of root exudation and its impact on microbial activity and nutrient
439 availability, *Applied Soil Ecology*, 5, 29-56, 1997.
- 440 Griffiths, B. S., Bonkowski, M., Dobson, G., and Caul, S.: Changes in soil microbial community
441 structure in the presence of microbial-feeding nematodes and protozoa, *Pedobiologia*, 43,
442 297-304, 1999.
- 443 Harris, J.: Soil microbial communities and restoration ecology: facilitators or followers?, *Science*, 325,
444 573-574, 2009.
- 445 Haynes, R.: Labile organic matter as an indicator of organic matter quality in arable and pastoral soils
446 in New Zealand, *Soil Biology and Biochemistry*, 32, 211-219, 2000.
- 447 Hinsinger, P., Bengough, A. G., Vetterlein, D., and Young, I. M.: Rhizosphere: biophysics,
448 biogeochemistry and ecological relevance, *Plant and Soil*, 321, 117-152, 2009.
- 449 Huang, X., Liu, S., Wang, H., Hu, Z., Li, Z., and You, Y.: Changes of soil microbial biomass carbon
450 and community composition through mixing nitrogen-fixing species with *Eucalyptus*
451 *urophylla* in subtropical China, *Soil Biology and Biochemistry*, 73, 42-48, 2014.
- 452 Huang, Y.-M., Liu, D., and An, S.-S.: Effects of slope aspect on soil nitrogen and microbial properties
453 in the Chinese Loess region, *Catena*, 125, 135-145, 2015.
- 454 Hueso, S., Hernández, T., and García, C.: Resistance and resilience of the soil microbial biomass to
455 severe drought in semiarid soils: The importance of organic amendments, *Applied Soil*
456 *Ecology*, 50, 27-36, 2011.
- 457 Jia, G. M., Cao, J., Wang, C. Y., and Wang, G.: Microbial biomass and nutrients in soil at the different
458 stages of secondary forest succession in Ziwulin, northwest China, *Forest Ecology and*
459 *Management*, 217, 117-125, 2005.
- 460 Kramer, C., and Gleixner, G.: Variable use of plant- and soil-derived carbon by microorganisms in
461 agricultural soils, *Soil Biology and Biochemistry*, 38, 3267-3278, 2006.
- 462 Ma, L. N., Guo, C. Y., Lü X. T., Yuan, S., and Wang, R. Z.: Soil moisture and land use are major
463 determinants of soil microbial community composition and biomass at a regional scale in
464 northeastern China, *Biogeosciences*, 12, 2585-2596, 2015.
- 465 Marschner, P., Marino, W., and Lieberei, R.: Seasonal effects on microorganisms in the rhizosphere of
466 two tropical plants in a polyculture agroforestry system in Central Amazonia, Brazil, *Biology*
467 *and Fertility of Soils*, 35, 68-71, 2002.
- 468 Mukhopadhyay, S., Masto, R. E., Cerdà A., and Ram, L. C.: Rhizosphere soil indicators for carbon
469 sequestration in a reclaimed coal mine spoil, *Catena*, 141, 100-108, 2016.
- 470 Nannipieri, P., Grego, S., and Ceccanti, B.: Ecological significance of the biological activity in soil,



- 471 Soil Biochemistry, 6, 293-355, 1990.
- 472 Nilsson, L. O., Giesler, R., Bååth, E., and Wallander, H.: Growth and biomass of mycorrhizal mycelia
473 in coniferous forests along short natural nutrient gradients, *New Phytologist*, 165, 613-622,
474 2005.
- 475 Qin, Y. Y., Feng, Q., Holden, N. M., and Cao, J. J.: Variation in soil organic carbon by slope aspect in
476 the middle of the Qilian Mountains in the upper Heihe River Basin, China, *Catena*, 147,
477 308-314, 2016.
- 478 Sariyildiz, T., Anderson, J., and Kucuk, M.: Effects of tree species and topography on soil chemistry,
479 litter quality, and decomposition in Northeast Turkey, *Soil Biology and Biochemistry*, 37,
480 1695-1706, 2005.
- 481 Selvakumar, G., Joshi, P., Mishra, P. K., Bisht, J. K., and Gupta, H. S.: Mountain aspect influences the
482 genetic clustering of psychrotolerant phosphate solubilizing Pseudomonads in the Uttarakhand
483 Himalayas, *Current Microbiology*, 59, 432-438, 2009.
- 484 Sidari, M., Ronzello, G., Vecchio, G., and Muscolo, A.: Influence of slope aspects on soil chemical and
485 biochemical properties in a *Pinus laricio* forest ecosystem of Aspromonte (Southern Italy),
486 *European Journal of Soil Biology*, 44, 364-372, 2008.
- 487 Sinha, S., Mastro, R., Ram, L., Selvi, V., Srivastava, N., Tripathi, R., and George, J.: Rhizosphere soil
488 microbial index of tree species in a coal mining ecosystem, *Soil Biology and Biochemistry*, 41,
489 1824-1832, 2009.
- 490 Tschirko, D., Hammesfahr, U., Marx, M.-C., and Kandeler, E.: Shifts in rhizosphere microbial
491 communities and enzyme activity of *Poa alpina* across an alpine chronosequence, *Soil Biology
492 and Biochemistry*, 36, 1685-1698, 2004.
- 493 Vance, E. D., Brookes, P. C., and Jenkinson, D. S.: An extraction method for measuring soil microbial
494 biomass C, *Soil Biology and Biochemistry*, 19, 703-707, 1987.
- 495 Wang, G. L., and Liu, G. B.: Study on the Interspecific Association of the *Artemisia sacrorum*
496 Community in Loess Hilly Region, Grassland of China, 2002.
- 497 Wang, L., Wang, Q., Wei, S., Shao, M. a., and Li, Y.: Soil desiccation for Loess soils on natural and
498 regrown areas, *Forest Ecology and Management*, 255, 2467-2477, 2008.
- 499 Wang, L., Wei, S., Horton, R., and Shao, M. a.: Effects of vegetation and slope aspect on water budget
500 in the hill and gully region of the Loess Plateau of China, *Catena*, 87, 90-100, 2011.
- 501 Wardle, D. A.: Impacts of disturbance on detritus food webs in agro-ecosystems of contrasting tillage
502 and weed management practices, *Advances in Ecological Research*, 26, 105-185, 1995.
- 503 Warembourg, F., Roumet, C., and Lafont, F.: Differences in rhizosphere carbon-partitioning among
504 plant species of different families, *Plant and Soil*, 256, 347-357, 2003.
- 505 Zelles, L.: Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of
506 microbial communities in soil: a review, *Biology and Fertility of Soils*, 29, 111-129, 1999.
- 507 Zhang, C., Liu, G. B., Xue, S., and Zhang, C. S.: Rhizosphere soil microbial properties on abandoned
508 croplands in the Loess Plateau, China during vegetation succession, *European Journal of Soil
509 Biology*, 50, 127-136, 2012.
- 510 Zhang, C., Liu, G. B., Xue, S., and Wang, G. L.: Changes in rhizospheric microbial community
511 structure and function during the natural recovery of abandoned cropland on the Loess Plateau,
512 China, *Ecological Engineering*, 75, 161-171, 2015.
- 513 Zhang, W., Parker, K., Luo, Y., Wan, S., Wallace, L., and Hu, S.: Soil microbial responses to
514 experimental warming and clipping in a tallgrass prairie, *Global Change Biology*, 11, 266-277,



515 2005.
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558



559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602




Fig. 1 Microbial biomass carbon (MBC) content, basal respiration (BR), and substrate-induced respiration (SIR) in the rhizospheric and non-rhizospheric soils. Error bars are standard errors (n=3). Different letters above the bars indicate significant differences at $P=0.05$.



603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646




Fig. 2 Effects of slope aspect on PLFA contents in rhizospheric and non-rhizospheric soils. Error bars are standard errors (n=3). Different letters above the bars indicate significant differences at $P=0.05$.



647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690




Fig. 3 The rhizospheric effect of microbial biomass carbon, basal respiration, substrate-induced respiration, and PLFA contents. Error bars are standard errors (n=3). Different letters above the bars indicate significant differences at $P=0.05$.




Fig. 4 Bidimensional graph for a redundancy analysis of the relationships between microbial properties and environmental factors in the rhizospheric (A) and non-rhizospheric (B) soils.

Note: RS, rhizospheric soil; NRS, non-rhizospheric soil; RQ, respiratory quotient; F/B, F/B ratio; G⁺/G⁻, G⁺/G⁻ ratio.

691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734



735
 736




Fig. 5 Structural equation models of the effect of slope aspect on microbial properties in the rhizospheric (A) and non-rhizospheric (B) soils. Numbers on the arrows are standardised path coefficients (equivalent to correlation coefficients). Solid lines indicate significant standardised path coefficients ($P < 0.05$). Circles indicate error terms (e1–e4). Percentages near the endogenous variables indicate the variance explained by the model.