

1 Tie-points for Gondwana reconstructions from a structural interpretation of the  
2 Mozambique Basin, East Africa, and the Riiser-Larsen Sea, Antarctica

3

4 **Jennifer Klimke<sup>1,\*</sup>, Dieter Franke<sup>1</sup>, Estevão Stefane Mahanjane<sup>2</sup> and German**  
5 **Leitchenkov<sup>3,4</sup>**

6 <sup>1</sup>Federal Institute for Geosciences and Natural Resources (BGR), Stilleweg 2, 30655  
7 Hannover, Germany

8 <sup>2</sup>Institute National Petroleum (INP), Av. Fernão Magalhaes N. 34, 2nd Floor, PO Box 4724,  
9 Maputo, Mozambique

10 <sup>3</sup>All-Russia Research Institute for Geology and Mineral Resources of the World Ocean, 1  
11 Angliysky Ave, St. Petersburg 190121, Russia

12 <sup>4</sup>St.-Petersburg State University, 13B Universitetskaya Emb., St. Petersburg, 199034, Russia

13

14 \*Corresponding author now at: University of Hamburg, Bundesstraße 55, 20146 Hamburg;  
15 Tel.: +49 40 42838 9466; Jennifer.Klimke@uni-hamburg.de; Jennifer.Klimke@gmx.net

16

## 17 **ABSTRACT**

18 Movements within early East Gondwana dispersal are poorly constrained and there is debate  
19 about conjugate geologic structures and the timing and directions of the rifting and earliest  
20 seafloor spreading phases. We present a combined structural interpretation of multichannel  
21 reflection seismic profiles from offshore northern Mozambique (East Africa), and the  
22 conjugate Riiser Larsen Sea (Antarctica). We find similar structural styles at the margins of  
23 both basins. At certain positions at the foot of the continental slope, close to the continent-  
24 ocean transition, the basement is intensely deformed and fractured, a structural style very  
25 untypical for rifted continental margins. Sediments overlying the fractured basement are

26 deformed and reveal toplap and onlap geometries, indicating a post-breakup deformation  
27 phase. We propose this unique deformation zone as tie-point for Gondwana reconstructions.  
28 Accordingly, we interpret the western flank of Gunnerus Ridge, Antarctica, as a transform  
29 margin, similar to the Davie Ridge offshore Madagascar, implying that they are conjugate  
30 features. As the continental slope deformation is post-rift, we propose a two-phase opening  
31 scenario. A first phase of rifting and early seafloor spreading, likely in NW-SE direction, was  
32 subsequently replaced by a N-S directed transform deformation phase, overprinting the  
33 continent-ocean transition. From previously identified magnetic chrons and the sediment  
34 stratigraphy, this change of the spreading directions from NW-SE to N-S is suggested to have  
35 occurred by the Late Middle Jurassic. We suggest that the second phase of deformation  
36 corresponds to the strike-slip movement of Madagascar and Antarctica and discuss  
37 implications for Gondwana breakup.

## 38 **1. INTRODUCTION**

39 The Mozambique Basin off East Africa and the conjugate Riiser-Larsen Sea off Antarctica  
40 (Fig. 1) resulted from the Middle Jurassic separation of East Gondwana (Madagascar,  
41 Antarctica, India and Australia) from West Gondwana (South America and Africa). However,  
42 a consistent reconstruction of prerift configurations relies on the knowledge of the crustal  
43 types and the location and structural style of the continent-ocean boundaries. Therefore, the  
44 early movements within Gondwana are poorly constrained and there is a debate about the  
45 timing and directions of the earliest rifting and spreading phases (e.g. Cox, 1992; Davis et al.,  
46 2016; Eagles and König, 2008; Jokat et al., 2003; Leinweber and Jokat, 2012; Marks and  
47 Tikku, 2001; Martin and Hartnady, 1986; Nguyen et al., 2016; Phethean et al., 2016; Reeves,  
48 2014, Reeves et al., 2016; Roeser et al., 1996; Smith and Hallam, 1970; Torsvik and Cocks,  
49 2013). The Mozambique Basin is of special importance for Gondwana reconstructions, as two  
50 end-members of rifted margins, a volcanic rifted and a transform margin can be studied in the  
51 immediate vicinity. In the Mozambique Basin, the transition from the SW-NE trending rifted

52 margin to the N-S trending transform margin along the Davie Ridge (Fig. 1) remains poorly  
53 studied. Existing studies focus mostly on the sedimentary infill of the Mozambique Basin  
54 (e.g. Castelino et al., 2015; Mahanjane, 2014; Salman and Abdula, 1995), or on the crustal  
55 structure in the western and central parts of the Mozambique Basin (e.g. Leinweber et al.,  
56 2013; Mahanjane, 2012; Müller and Jokat, 2017; Mueller et al., 2016). While it is generally  
57 accepted that the Riiser-Larsen Sea is the conjugate of the Mozambique Basin (e.g. Jokat et  
58 al., 2003; Nguyen et al., 2016), it remains much less well studied in spite of an available set of  
59 modern geophysical data (e.g. Hinz et al., 2004; Leitchenkov et al., 2008; Roeser et al., 1996).  
60 In this study, we present a combined structural interpretation of new and previously published  
61 multichannel reflection seismic profiles from different datasets. We concentrate on offshore  
62 Mozambique (East Africa), in the vicinity of the Davie Ridge, and the conjugate Riiser Larsen  
63 Sea (Antarctica) at the transition from the rifted margin to the Gunnerus Ridge (Fig. 1) and  
64 compare the structural configuration of the basement and the earliest postrift sediments.  
65 The main outcome of this study is the identification of a zone of deformed and faulted  
66 basement at the foot of the continental slope at both margins. The sediments overlying the  
67 deformation zone are deformed, revealing a post-breakup deformation phase. We provide  
68 evidence that these unique structures can serve as tie-point for Gondwana reconstructions.  
69 This leads to a two-phase opening scenario for the conjugate Mozambique Basin and Riiser  
70 Larsen Sea.

## 71 **2.1 BREAKUP OF EAST AND WEST GONDWANA**

72 Several plate kinematic models describe the breakup of Gondwana along the East African  
73 margin (e.g. Cox, 1992; Davis et al., 2016; Gaina et al., 2013, 2015; Eagles and König, 2008;  
74 Leinweber and Jokat, 2012; Nguyen et al., 2016; Reeves et al., 2016). It is generally accepted  
75 that breakup of Gondwana along the East African margin took place in the Early Jurassic, at  
76 about 170-180 Ma (e.g. Gaina et al., 2013, 2015; Leinweber and Jokat, 2012; Leinweber et  
77 al., 2013; Nguyen et al., 2016; Reeves et al., 2016). While earlier studies proposed that the

78 Mozambique Basin and West Somali Basin opened in a generally N-S direction, more recent  
79 plate tectonic reconstructions argue for an almost simultaneous opening of both basins in  
80 NW-SE direction (e.g. Gaina et al., 2013; Reeves et al., 2016). There is also debate about the  
81 timing and directions of the earliest rifting and spreading phases. A change of the spreading  
82 direction has been suggested to have occurred at ~159 Ma (Leinweber and Jokat, 2012), ~153  
83 Ma (Reeves et al., 2016), or ~150 Ma (Phethean et al., 2016).

84 Oceanic crust generated by seafloor spreading between Africa and Antarctica has been dated  
85 by the identification of marine magnetic anomalies. Recent studies, using new geophysical  
86 data, tentatively identify M41n (~165 Ma; Leinweber and Jokat, 2012) or M38n.2n (~164 Ma;  
87 Müller and Jokat, 2017; magnetic polarity timescale of Ogg, 2012) as the oldest magnetic  
88 anomaly in the Mozambique Basin. This makes the Mozambique Basin/Riiser-Larsen Sea  
89 considerably older than proposed in previous studies (M2 to M22, ~148-127 Ma; Simpson et  
90 al., 1979; Segoufin, 1978).

91 In the conjugate Riiser-Larsen Sea, Leinweber and Jokat (2012) identify M25n (~154 Ma) as  
92 the oldest magnetic anomaly (Fig. 1), extending the model of Bergh (1977) and confirming  
93 previous interpretations of Roeser et al. (1996) and Leitchenkov et al. (2008), who identified  
94 M0 to M24 (~152-125 Ma). However, well-defined magnetic anomalies older than M25n  
95 were not yet identified (Leinweber and Jokat, 2012; Leitchenkov et al., 2008; Roeser et al.,  
96 1996), although it is implied that spreading started before M25n (Leinweber and Jokat, 2012).

97 There is general agreement that, by the Late Jurassic, seafloor spreading was underway in the  
98 Mozambique and Riiser Larsen Sea Basins (e.g. Coffin and Rabinowitz, 1987; Eagles and  
99 König, 2008; Rabinowitz et al., 1983; Segoufin and Patriat, 1980; Simpson et al., 1979).

## 100 **2.2 ENIGMATIC CRUSTAL BLOCKS IN THE MOZAMBIQUE BASIN AND** 101 **RIISER-LARSEN SEA**

102 The Mozambique Basin and the West Somali Basin are separated by a bathymetric elevation  
103 rising 1-2 km above the surrounding seafloor that is referred to as the Davie Ridge (Fig. 1). It

104 has been widely accepted that the Davie Ridge is located at the trace of a fossil transform fault  
105 that accommodated the motion of Madagascar/Antarctica with respect to Africa. This  
106 transform is thought to have been active from the Late Middle Jurassic (~160-165 Ma) to the  
107 Early Cretaceous (~125-135 Ma) (e.g. Coffin and Rabinowitz, 1987; Segoufin and Patriat,  
108 1980). Although the presence of a transform continental margin and its expression, the Davie  
109 Ridge, has been questioned in the West Somali Basin (Klimke and Franke, 2016), offshore  
110 west Madagascar this structure is obvious. The Gunnerus Ridge in the Riiser-Larsen Sea may  
111 be the prolongation of the shear zone offshore Madagascar that accommodated the southward  
112 drift of Madagascar relative to Africa (Nguyen et al., 2016). (Fig. 1). Its western flank has  
113 been interpreted as a strike-slip fault delineating a transform margin (e.g. Leitchenkov et al.,  
114 2008). The Gunnerus Ridge has been the subject of seismic and potential field studies in the  
115 last decades (e.g. Leitchenkov et al., 2008; Roeser et al., 1996; Saki et al., 1987). Based on its  
116 top basement seismic velocities of 5.8-6.1 km/s and dredged granitoid and gneissic rock  
117 samples, the Gunnerus Ridge has been ascribed a continental origin (Leitchenkov et al., 2008;  
118 Saki et al., 1987).

119 Other prominent crustal features in the Mozambique Basin and the Riiser-Larsen Sea are the  
120 Beira High and the Astrid Ridge, respectively (Fig. 1). Both, structural interpretation  
121 (Mahanjane, 2012) and seismic velocities derived from refraction seismic data (Müller et al,  
122 2016) indicate that Beira High is made up of stretched and highly intruded continental crust.  
123 The Astrid Ridge in the western Riiser-Larsen Sea (Fig. 1) is separated into a northern and a  
124 southern part by the Astrid Fracture Zone (e.g. Bergh, 1987; Leitchenkov et al., 2008). While  
125 Bergh (1987) proposed that the Astrid Ridge is an entirely magmatic structure, Roeser et al.  
126 (1996) proposed that N-S striking strong magnetic anomalies over the western flank of the  
127 southern part of Astrid Ridge originate from seaward-dipping reflectors and that this part is  
128 made up of continental crust.

### 129 **3. MATERIALS AND METHODS**

130 In this study, we use several marine reflection seismic datasets acquired by different institutes  
131 in the Mozambique Channel and the Riiser-Larsen Sea (Fig. 1).

132 The **BGR14** dataset was acquired by the Federal Institute for Geosciences and Natural  
133 Resources (BGR) during a cruise of R/V Sonne in 2014. For a detailed description of the  
134 acquisition parameters and seismic processing, the reader is referred to Klimke et al. (2016).

135 In this study, we present a yet unpublished profile striking E-W, crossing the Mozambique  
136 Basin into the Morondava Basin offshore Madagascar (Fig. 1). For the seismostratigraphic  
137 interpretation of the areas in the Morondava Basin and the Davie Ridge, we use the  
138 stratigraphic interpretation established in Franke et al. (2015) and Klimke et al. (2016). For  
139 the Mozambique Basin, we use results from previous offshore studies (e.g. Castelino et al.,  
140 2015; Franke et al., 2015; Mahanjane, 2014).

141 We present two out of eight profiles of the **Mbwg00** dataset acquired by Western Geophysical  
142 in 2000, which run NW-SE and SW-NE in the Mozambique Channel (Fig. 1). This dataset is  
143 part of the National Petroleum Institute of Mozambique archive and has recently been  
144 presented by Mahanjane (2014). Here, we present one previously published profile  
145 (Mahanjane, 2014) with the focus on the continental slope and additionally show one  
146 previously unpublished profile of this dataset. For the interpretation of the sedimentary  
147 successions, we base on the stratigraphic framework established in Castelino et al. (2015),  
148 Franke et al. (2015) and Mahanjane (2014).

149 The **RAE43** reflection seismic dataset in the Riiser Larsen Sea was acquired by Polar Marine  
150 Geosurvey Expedition during a survey with the R/V Akademik Alexander Karpinsky in 1998.  
151 For a detailed description of the used equipment, the acquisition parameters, and the  
152 processing, the reader is referred to Leitchenkov et al. (2008). In this study, we show two  
153 reinterpreted profiles of this dataset (Fig. 1) using as a basis the stratigraphic framework of  
154 Leitchenkov et al. (2008).

155 The seismic profiles shown in this paper are located in the northeastern part of the  
156 Mozambique Basin, off East Africa, and in the eastern part of the Riiser-Larsen Sea, off  
157 Antarctica (Fig. 1) and thus cover parts of two conjugate margins resulting from the  
158 separation of Antarctica from Africa. Two profiles (Fig. 2 and Fig. 3) are oriented in a NW-  
159 SE direction, parallel to the spreading direction and run from the continental slope towards the  
160 abyssal plain in the Mozambique Basin and Riiser-Larsen Sea. Profile C (Fig. 4 and Fig. 5)  
161 trends NW-SE and runs from the Mozambique margin towards the Davie Ridge, while  
162 Profiles D and E (Figs. 6, 7 and 8) are oriented in E-W direction, crossing the Davie Ridge  
163 and Gunnerus Ridge, respectively.

#### 164 **4. COMMON CHARACTERISTICS OF CONJUGATE MARGIN SECTIONS: THE** 165 **TIE-POINT**

166 We identify an untypical yet similar structural style of the continent-ocean transition at both,  
167 the Mozambique and the Riiser-Larsen Sea continental margins. The continental slopes dip  
168 steeply at angles of  $\sim 6^\circ$ - $7^\circ$  at the Mozambique margin (Fig. 2A) and  $\sim 5^\circ$  in the Riiser-Larsen  
169 Sea (Fig. 2B). The top basement reflection is clearly imaged below the slopes and increases in  
170 depth from  $\sim 1$ s (TWT) to  $\sim 7$ s (TWT) over distances of  $\sim 50$ - $70$  km. At the foot of the  
171 continental slope, at depths of  $\sim 7$  s (TWT), there is a distinct zone of highly deformed  
172 continental basement on both continental margins (Fig. 2A, offset range: 50-70 km; Fig. 2B,  
173 offset range: 160-190 km). In the deformed zone, the basement is intensely faulted over  
174 distances of about 30 km (Fig. 2). On Profile A (Fig. 3A), the basement is uplifted by  
175 apparent high-angle reverse faulting and the sedimentary cover is folded at the tip of the main  
176 fault strands. Internal horizons are heavily deformed and dissected by faults (e.g. Fig. 3A,  
177 offset range: 50-70 km). The unconformity MJ seals the deformation, which has, according to  
178 our seismostratigraphic concept, an age of the transition from the Middle to the Late Jurassic.  
179 The sedimentary unit underlying horizon MJ is characterized by subparallel reflectors with

180 low amplitudes. The seismic transparency of this unit allows a clear along-margin distinction  
181 from younger, reflective deposits (e.g. Fig. 2).

182 The same kind of deformation is identified on the conjugate continental slope in the Riiser-  
183 Larsen Sea (Figs. 2B and 3B, offset range: 160-190 km). Again, the basement is dissected by  
184 high-angle faults at the foot of the continental slope. A similar package of post-rift sediments  
185 is affected by folding to form a gentle anticline, altogether resembling the observed  
186 deformation pattern in the Mozambique Basin (Figs. 2A and 3A). The overall geometries  
187 resemble positive flower structures developed along strike-slip faults (e.g. Harding, 1985,  
188 1990; Sylvester, 1988) (Fig. 2 and Fig. 3).

189 Further northeast in the Mozambique Basin (Fig. 4), the basement deformation is  
190 characterized by steep and very closely spaced faults (Fig. 4, offset range: 40-50 km).  
191 Faulting increases towards the SE (Fig. 5, offset range: 50-60 km) where internal reflections  
192 have been heavily deformed and rotated to form gentle synclines. In contrast to the area  
193 further west at the continental slope of the Mozambique margin, which is characterized by  
194 compressional deformation (Fig. 2), the horizontal component of motion across the faults in  
195 the SE (Fig. 5) is extensional, and the overall geometry resembles negative flower structures  
196 (Harding, 1985, 1990; Sylvester, 1988). Profile D in the Mozambique Basin (Fig. 6) shows  
197 that the basement is transparent in the deformed zone (profile distance: 25-45 km), possibly  
198 due to intense faulting.

199 Seaward of the deformation zone along both continental margins, oceanic crust is interpreted  
200 that is characterized by high-amplitude, low-frequency, multi-reflector bands in depths of 7-9  
201 s (TWT) (Figs. 2, 4, 6, 7 and 8). Locally, closely spaced diffractions are distinct (Figs. 2, 4, 6,  
202 7 and 8), both features being typical for oceanic crust (Klimke et al., 2016). The interpretation  
203 of oceanic crust seaward of the deformation zone is well in line with refraction seismic  
204 experiments and gravity modelling by Leinweber et al. (2013), refraction seismic experiments  
205 supported by 2D magnetic modelling of Müller and Jokat (2017) and magnetic anomaly

206 identifications by Leinweber and Jokat (2012) and Müller and Jokat (2017) in the  
207 Mozambique Basin. Normal faults dissecting the oceanic crust with throws of ~250 ms  
208 (TWT) in the Mozambique Basin (Figs. 2A and 4) and up to ~1s (TWT) in the Riiser-Larsen  
209 Sea (Figs. 2B, 7 and 8) are distinct. The faults are spaced at 5-15 km (Fig. 2A, offset range:  
210 90-190 km; Fig. 4, offset range: 70-180 km; Fig. 6, offset range: 70-100 km) and 10-40 km  
211 (Fig. 2B, offset range: 30-110 km; Fig. 7, offset range: 0-300 km), respectively. The  
212 abundance of the faults is increasing significantly in the vicinity of the Davie Ridge (from ~15  
213 km to 5 km) and the Gunnerus Ridge (from ~40 km to ~10 km).

214 At both margins, unconformity MJ, which is sealing the deformation, is terminating seawards  
215 against oceanic crust, which likely formed during the Jurassic Magnetic Quiet Zone (Middle  
216 to the Late Jurassic). An extrapolation of identified magnetic anomalies (Figs. 1 and 9;  
217 Leinweber and Jokat, 2012; Müller and Jokat, 2017) to the NE Mozambique Basin (Fig. 9),  
218 indicates that the sedimentary unit below horizon MJ terminates against oceanic crust at  
219 approximately the position of magnetic anomaly M38n.2n (~164 Ma). The extrapolation of  
220 the magnetic anomalies was done by noting the distance of magnetic anomaly M38n from the  
221 continent-ocean transition in the Mozambique Basin (Fig. 9). This is well in line with our  
222 stratigraphic concept and we propose that the deformation is Middle Jurassic in age and was  
223 finished at the transition from the Middle to Late Jurassic. The deformation of the earliest,  
224 likely Middle Jurassic sediments observed at both continental margins is characterized by  
225 onlap and toplap geometries, where the MJ horizon acts as an unconformity sealing the  
226 deformation. In the Mozambique Basin, the top of the Middle Jurassic sediments has been  
227 eroded resulting in toplap structures of older sediments against the MJ horizon (Fig. 3A,  
228 offset: 60 km). In the Riiser-Larsen Sea, the Middle Jurassic sediments have been folded  
229 upward in conjunction with the basement (Fig. 3B, offset range: 160-190 km) and subsequent,  
230 likely Late Jurassic sediments onlap the MJ horizon (Fig. 3B, profile distance: 170 km).

231 According to Leinweber et al. (2013) and Müller and Jokat (2017), the continent-ocean

232 transition at the Mozambique margin is located very close to the Zambezi coast and is  
233 characterized by high-velocity lower crustal bodies and seaward-dipping reflectors, typical for  
234 volcanic rifted margins.

235 This previously identified position of the continent-ocean transition corresponds in our  
236 reflection seismic profiles to the area of the deformed basement (Figs. 2, 4, 6).  
237 Geographically, the deformed basement zone is distinct in the eastern parts of the oceanic  
238 basins, close to the Davie Ridge and the Gunnerus Ridge (Fig. 9). The zone is clearly depicted  
239 on several profiles over distances of 100-200 km in E-W direction along the margins (Fig. 9).

## 240 **5. DISCUSSION**

### 241 **5.1 LANDWARD EXTENT OF OCEANIC CRUST**

242 Both, the Mozambique Basin and the Riiser-Larsen Sea show a steeply dipping continental  
243 slope with angles of  $5^{\circ}$ - $7^{\circ}$  with a zone of deformed basement situated at the foot of the  
244 continental slope. Seaward of the deformed zone oceanic crust is interpreted, which is highly  
245 dissected by normal faults. The abundance of the faults, with throws of up to 1s (TWT),  
246 increases towards the Davie Ridge and the Gunnerus Ridge.

247 At both margins, magnetic anomaly M25n (~154-155 Ma) is located ~250-280 km seaward of  
248 the coast (Fig. 1), which implies symmetric spreading. If the interpretation of magnetic  
249 anomaly M38n.2n (~164 Ma; Müller and Jokat, 2017) is correct, oceanic crust older than  
250 ~155 Ma (M25n) should be found also in the Riiser-Larsen Sea. A comparably wide strip of  
251 oceanic crust with ages of ~155-166 Ma fits well in the area in between magnetic anomaly  
252 M25n and the here identified zone of deformed basement at the base of the continental slope  
253 (section 4). This implies a considerably more southern position of the continent-ocean  
254 transition than previously anticipated for the Riiser-Larsen Sea (Fig. 9). Gravity modelling  
255 derived crustal thicknesses of 5-6 km (Leitchenkov et al., 2008) are in accordance with this  
256 concept. The crustal thickness remains relatively constant west of the Gunnerus Ridge and  
257 increases from 5-6 km to 10 km only near the Astrid Ridge (Fig. 16 in Leitchenkov et al.,

258 2008). Based on these observations, we suggest to relocate the continent-ocean transition in  
259 the Riiser-Larsen Sea to the zone of deformed basement at the continental slope (Fig. 9).  
260 Along the Davie Ridge and the Gunnerus Ridge, the transition from continental to oceanic  
261 crust is abrupt. At the western flank of the Gunnerus Ridge, the continent-ocean transition is  
262 ~40-50 km wide and at the Davie Ridge, it does not exceed 10-20 km. This is typical for  
263 transform continental margin settings, where the transition from continental to oceanic crust  
264 occurs over distances of not more than 50-80 km (e.g. Bird, 2001). Gravity modelling of  
265 profiles crossing the Gunnerus Ridge by Leitchenkov et al. (2008) and Roeser et al. (1996)  
266 confirm the abrupt continent-ocean transition. Thus, we propose that the western margin of  
267 Gunnerus Ridge is a transform margin, similar to Davie Ridge. As the abundance of normal  
268 faults increases significantly in the vicinity of the Davie Ridge and Gunnerus Ridge (Fig. 4  
269 and Fig. 7), we suggest that the oceanic crust has been affected by intense shear motions  
270 during spreading.

## 271 **5.2 IMPLICATIONS FOR GONDWANA BREAKUP**

272 As origin of the distinct basement deformation at the continent-ocean transition in the eastern  
273 parts of both Mozambique Basin and the Riiser-Larsen Sea we propose intense strike-slip  
274 shearing. From the reflection seismic data, we interpret oceanic crust just seaward of the  
275 deformed basement. Thus, there was a short period of seafloor spreading preceding the N-S  
276 directed strike-slip movement. From the orientation, this is in agreement with plate tectonic  
277 reconstructions which propose an early, NW-SE directed phase of rifting and seafloor  
278 spreading in the Mozambique Basin/Riiser-Larsen Sea (e.g. Eagles and König, 2008; Gaina et  
279 al., 2013; Reeves et al., 2016), followed by a change of spreading directions from NW-SE to  
280 N-S. According to our seismostratigraphic concept, the change in spreading directions from  
281 NW-SE to N-S likely occurred in the Late Middle Jurassic, shortly before the formation of the  
282 sealing unconformity, dated at the transition from Middle to Late Jurassic. At the latter time,  
283 seafloor spreading likely has formed such a wide oceanic domain that strike-slip movements

284 did no longer affect the rifted continental margins in Mozambique and Antarctica. A change  
285 in the elongation of the early mid-oceanic ridge corresponding to the proposed variation in the  
286 spreading direction has so far not been reported. This may be difficult to identify as this early  
287 oceanic basement has been intensively deformed by subsequent shear movements. The major  
288 portion of shearing certainly occurred along the Davie Ridge and the Gunnerus Ridge that in  
289 our view represent transform margins on their western flanks in the Mozambique Basin and  
290 the Riiser-Larsen Sea (Fig. 9). However, the reflection seismic data reveal that the shearing  
291 processes affected oceanic crust located as far as 200 km away from the main transform faults  
292 (Fig. 9), indicating also a longer-lasting process. Klimke et al. (2016) observed similar  
293 structures in extended basement to the east of Davie Ridge in the West Somali Basin (Fig. 9).  
294 The observed faults are steeply dipping wrench faults that were active during the southward  
295 movement of Madagascar along the Davie Ridge. Here, a prominent unconformity of inferred  
296 Early Cretaceous age marks the end of wrench deformation (Klimke et al., 2016).

297 Westward of the study area, the Beira High (Fig. 1) is suggested to have separated from  
298 Africa during the initial opening of the Mozambique Basin (e.g. Nguyen et al., 2016). As  
299 significant differences in the amount of stretching are observed below the margins of Beira  
300 High, some authors propose a rift jump during the early rifting stage from the northwestern to  
301 the southeastern edge of Beira High (e.g. Mahanjane, 2012; Müller et al., 2016). Mahanjane  
302 (2012) observes two rift phases in reflection seismic data covering the Beira High and  
303 postulates a two break-up stages concept. Our observed two-phase break-up scenario (Fig. 10)  
304 concurs well with the proposed rift jump model (e.g. Mahanjane, 2012; Müller et al., 2016).

305 We suggest that the “ridge jump” from the northwestern to the southern edge of Beira High is  
306 associated with the change in spreading direction from NW-SE to N-S direction, initiating the  
307 strike-slip movement of Madagascar and Antarctica (Fig. 10). However, the structure of the  
308 eastern margin of Beira High remains elusive, the nature of this continent-ocean transition is  
309 unclear.

310 Our proposed model for the initial opening of the Mozambique Basin/Riiser-Larsen Sea  
311 implies that the Gunnerus Ridge was located at the southwestern flank of Madagascar in order  
312 to be aligned with the Davie Ridge. This brings the Astrid Ridge, regardless of its crustal  
313 nature and formation age, to the western flank of Beira High (Fig. 10), indicating that they are  
314 conjugate features (Nguyen et al., 2016).

## 315 **6. CONCLUSIONS**

316 1) In reflection seismic profiles, we identify a symmetric zone of deformed and faulted  
317 basement at the foot of the continental slope at the continental margins of the  
318 northeastern Mozambique Basin and the conjugate eastern Riiser-Larsen Sea.

319 2) The architecture and style of the observed deformation zone, which is unique at rifted  
320 margins, represents a mirror image between both conjugate margins and is proposed as  
321 a tie point for Gondwana reconstructions. Strike-slip shearing is proposed as the origin  
322 of the deformed continental slope.

323 3) Sediments overlying the basement deformation zone at the foot of the continental  
324 slope are deformed with onlap and toplap geometries, indicating a post-breakup  
325 deformation phase. For the unconformity sealing the strike-slip deformation we  
326 estimate an age at about the transition from Middle to Late Jurassic. The structural  
327 configuration indicates a first phase of rifting and early seafloor spreading that  
328 subsequently has been overprinted by Late Middle Jurassic strike-slip deformation and  
329 the formation of a transform boundary at the expenses of the original continent-ocean  
330 transition.

331 4) From the structural configuration, the Gunnerus Ridge, Antarctica, is conjugate to the  
332 Davie Ridge, offshore Mozambique/Madagascar. A major transform fault is proposed  
333 at the western margin of the Gunnerus Ridge, similar to the Davie Ridge. Strike-slip  
334 deformation affected not only the rims of Davie and Gunnerus Ridge, but also the  
335 adjacent oceanic crust up to a distance of 200 km from the main transform fault. In the

336 eastern Riiser-Larsen Sea, oceanic crust likely extends further south than previously  
337 proposed.

338 5) In the here proposed breakup scenario, a first, likely NW-SE directed extensional  
339 phase resulted in localized seafloor spreading in the Mozambique Basin/Riiser-Larsen  
340 Sea Basin in the Middle Jurassic. A second, Late Middle Jurassic phase, likely in  
341 association with a ridge-jump, initiated the generally N-S opening of the oceanic  
342 basin. The second phase represents the southward displacement of East Gondwana,  
343 with strike-slip movement of Madagascar and Antarctica against Africa and the  
344 development of transform margins along Gunnerus Ridge and Davie Ridge.

345

#### 346 **DATA AVAILABILITY**

347 All reflection seismic profiles of the BGR14 dataset can be accessed via Geo-Seas  
348 (<http://www.geo-seas.eu>). The reflection seismic dataset (RAE43) located in the Riiser-Larsen  
349 Sea has been made available through Antarctic Seismic Data Library System (SDLS) and can  
350 be accessed via <http://sdls.ogs.trieste.it/>. Two profiles of the Mbwg00 dataset located in the  
351 Mozambique Channel are commercial seismic lines, original data of which cannot be made  
352 available.

353

#### 354 **COMPETING INTERESTS**

355 The authors declare that they have no conflict of interest.

356

#### 357 **ACKNOWLEDGEMENTS**

358 The German Ministry for Research and Education (BMBF) funded the cruise SO231 and  
359 supported the study through grant 03G0231A. BGR staff is thanked for helping in data  
360 acquisition and processing. We thank the National Petroleum Institute of Mozambique for  
361 allowing publication of two seismic profiles of the Mbwg00 dataset. The reflection seismic

362 dataset (RAE43) located in the Riiser-Larsen Sea has been made available through Antarctic  
363 Seismic Data Library System (SDLS). Reviews by Carmen Gaina and an anonymous  
364 reviewer greatly improved the manuscript. We are particularly thankful for supportive  
365 comments and suggestions by Topical Editor Federico Rossetti.

366

## 367 REFERENCES

- 368 Amante, C. and Eakins, B.W., 2009. ETOPO1 1 Arc-Minute Global Relief Model:  
369 Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS  
370 NGDC-24, National Geophysical Data Center, NOAA. doi:10.7289/V5C8276M  
371 [11.02.2016].
- 372 Bergh, H.W., 1987. Underlying fracture zone nature of Adrid Ridge off Antarctica's Queen  
373 Maud Land. *Journal of Geophysical Research* 92 (B1), 475-484.
- 374 Bergh, H.W., 1977. Mesozoic sea floor off Dronning Maud Land, Antarctica. *Nature* 269,  
375 686-687.
- 376 Bird, D., 2001. Shear margins: Continent-ocean transform and fracture zone boundaries: The  
377 Leading Edge, 150-159.
- 378 Castelino, J.A., Reichert, C., Klingelhoefer, F. and Aslanian, D., 2015. Mesozoic and Early  
379 Cenozoic sediment influx and morphology of the Mozambique Basin. *Marine and  
380 Petroleum Geology* 66, 890-905.
- 381 Coffin, M.F., and Rabinowitz, P.D., 1987. Reconstruction of Madagascar and Africa:  
382 Evidence from the Davie Fracture Zone and Western Somali Basin. *Journal of  
383 Geophysical Research: Solid Earth* 92, 9385-9406.
- 384 Cox, K.G., 1992. Karoo igneous activity, and the early stages of the break-up of  
385 Gondwanaland. From: Storey, B.C., Alabaster, T. and Pankhurst, R.j. (eds), 1992.  
386 *Magmatism and the Causes of Continental Break-up*. Geological Society Special  
387 Publication 68, 137-148.
- 388 Davis, J.K., Lawver, L.A., Norton, I.O. and Gahagan, L.M., 2016. New Somali Basin  
389 magnetic anomalies and a plate model for the early Indian Ocean. *Gondwana Research*  
390 34, 16-28.
- 391 Eagles, G., and König, M., 2008. A model of plate kinematics in Gondwana breakup.  
392 *Geophysical Journal International* 173, 703-717.
- 393 Franke, D., Jokat, W., Ladage, S., Stollhofen, H., Klimke, J., Lutz, R., Mahanjane, E.S.,  
394 Ehrhardt, A. and Schreckenberger, B., 2015. The offshore East African Rift System:  
395 Structural framework at the toe of a juvenile rift. *Tectonics* 34, 2086-2104,  
396 doi:10.1002/2015TC003922.
- 397 Gaina, C., Van Hinsbergen, D.J.J. and Spakman, W., 2015. Tectonic interactions between  
398 India and Arabia since the Jurassic reconstructed from marine geophysics, ophiolite  
399 geology, and seismic tomography. *Tectonics* 34, 875-906.
- 400 Gaina, C., Torsvik, T.H., van Hinsbergen, D.J.J., Medvedev, S., Werner, S.C., and Labails,  
401 C., 2013. The African Plate: A history of oceanic crust accretion and subduction since  
402 the Jurassic. *Tectonophysics* 604, 4-25.

- 403 Harding, T.P., 1985. Seismic characteristics and identification of negative flower structures,  
404 positive flower structures, and positive structural inversion. *Bulletin of the American*  
405 *Association of Petroleum Geologists* 69 (4), 582-600.
- 406 Harding, T.P., 1990. Identification of wrench faults using subsurface structural data: criteria  
407 and pitfalls. *Bulletin of the American Association of Petroleum Geologists* 74, 1590-  
408 1609. Hinz, K., Neben, S., Gouseva, Y.B. and Kudryavtsev, G.A., 2004. A  
409 Compilation of Geophysical Data from the Lazarev Sea and the Riiser-Larsen Sea,  
410 Antarctica. *Marine Geophysical Researches* 25, 233-245.
- 411 Jokat, W., Boebel, T., König, M. and Meyer, U., 2003. Timing and geometry of early  
412 Gondwana breakup. *Journal of Geophysical Research* 108 (B9), 2428.
- 413 Klimke, J., Franke, D., Gaedicke, C., Schreckenberger, B., Schnabel, M., Stollhofen, H.,  
414 Rose, J. and Chaheire, M., 2016. How to identify oceanic crust – evidence for a  
415 complex break-up in the Mozambique Channel, off East Africa, *Tectonophysics* 693,  
416 Part B, 436-452, <http://dx.doi.org/10.1016/j.tecto.2015.10.012>.
- 417
- 418 Klimke, J. and Franke, D., 2016. Gondwana breakup: no evidence for a Davie Fracture Zone  
419 offshore northern Mozambique, Tanzania and Kenya. *Terra Nova* 28, 233-244.
- 420 Leinweber, V.T. and Jokat, W., 2012. The Jurassic history of the Africa-Antarctica corridor –  
421 new constraints from magnetic data on the conjugate continental margins.  
422 *Tectonophysics* 530-531, 87-101.
- 423 Leinweber, V.T., Klingelhoefer, F., Neben, S., Reichert, C., Aslanian, D., Matias, L., Heyde,  
424 I., Schreckenberger, B. and Jokat, W., 2013. The crustal structure of the Central  
425 Mozambique continental margin – Wide-angle seismic, gravity and magnetic study in  
426 the Mozambique Channel, Eastern Africa. *Tectonophysics* 599, 170-196.
- 427 Leitchenkov, G., Guseva, J., Gandyukhin, V., Griukurov, G., Kristoffersen, Y., Sand, M.,  
428 Golynsky, A and Aleshkova, N., 2008. Crustal structure and tectonic provinces of the  
429 Riiser-Larsen Sea area (East Antarctica): results of geophysical studies. *Marine*  
430 *Geophysical Research* 29, 135-158.
- 431 Mahanjane, E.S., 2012. A geotectonic history of the northern Mozambique Basin including  
432 the Beira High – A contribution for the understanding of its development. *Marine and*  
433 *Petroleum Geology* 36, 1-12.
- 434 Mahanjane, E.S., 2014. The Davie Ridge and adjacent basins in the offshore Mozambique  
435 Margin – A new insights for the hydrocarbon potential. *Marine and Petroleum*  
436 *Geology* 57, 561-571.
- 437 Marks, K.M. and Tikku, A.A., 2001. Cretaceous reconstructions of East Antarctica, Africa  
438 and Madagascar. *Earth and Planetary Science Letters* 186, 479-495.
- 439 Martin, A.K. and Hartnady, C.J.H., 1986. Plate tectonic development of the South West  
440 Indian Ocean: A revised reconstruction of East Antarctica and Africa. *Journal of*  
441 *Geophysical Research* 91 (B5), 4767-4786.
- 442 Müller, C.O. and Jokat, W., 2017. Geophysical evidence for the crustal variation and  
443 distribution of magmatism along the central coast of Mozambique, *Tectonophysics*  
444 712-713, 684-713.
- 445 Müller, C.O., Jokat, W. and Schreckenberger, B., 2016. The crustal structure of Beira High,  
446 central Mozambique – Combined investigation of wide-angle seismic and potential  
447 field data. *Tectonophysics* 683, 233-254.

448 Nguyen, L.C., Hall, S.A., Bird, D.E. and Ball, P.J., 2016. Reconstruction of the East Africa  
449 and Antarctica continental margins. *Journal of Geophysical Research: Solid Earth* 121,  
450 4156-4179.

451 Ogg, J.G., 2012. Geomagnetic polarity time scale. In: Gradstein, F.M., Ogg, J.G., Schmitz,  
452 M., Ogg, G. (Eds.). *The Geologic Time Scale 2012*. Elsevier, pp. 731-791.

453 Phethean, J.J.J., Kalnins, L.M., van Hunen, J., Biffi, P.G., Davies, R.J. and McCaffrey,  
454 K.J.W., 2016. Madagascar's escape from Africa: A high-resolution plate  
455 reconstruction for the Western Somali Basin and implications for supercontinent  
456 dispersal. *Geochemistry, Geophysics, Geosystems* 17, 5036-5055.

457 Rabinowitz, P.D., Coffin, M.F., and Flavey, D., 1983. The separation of Madagascar and  
458 Africa. *Science* 220, 67-69.

459 Reeves, C.V., Teasdale, J.P. and Mahanjane, E.S., 2016. Insight into the Eastern Margin of  
460 Africa from a new tectonic model of the Indian Ocean. From: Nemcok, M., Rybár, S.,  
461 Sinha, S.T., Hermeston, S.A. and Ledvényiová, L. (eds). *Transform margins:  
462 Development, Controls and Petroleum Systems*. Geological Society, London, Special  
463 Publications 431.

464 Reeves, C., 2014. The position of Madagascar within Gondwana and its movements during  
465 Gondwana dispersal. *Journal of African Earth Sciences* 94, 45-57.

466 Roeser, H.A., Fritsch, J. and Hinz, K., 1996. The development of the crust of Dronning Maud  
467 Land, East Antarctica. From: Storey, B.C., King, E.C. and Livermore, R.A. (eds),  
468 1996. *Weddell Sea Tectonics and Gondwana Break-up*. Geological Society Special  
469 Publication 108, 243-264.

470 Saki, T., Tamura, Y., Tokuhashi, S., Kodato, T., Mizukoshi, I. and Amano, H., 1987.  
471 Preliminary report of geological and geophysical surveys off Dronning Maud Land,  
472 East Antarctica. *Proceedings of the Nat Inst Polar Res (NIPR) Symposium: Ant.  
473 Geoscience* 1, 23-40, Tokyo.

474 Salman, G. and Abdula, I., 1995. Development of the Mozambique and Ruvuma sedimentary  
475 basins, offshore Mozambique. *Sedimentary Geology* 96 (1-2), 7-41.

476 Segoufin, J., 1978. Anomalies magnétiques mésozoïques dans le bassin de Mozambique. *C.  
477 R. Acad. Sci. D* 287, 109-112.

478 Segoufin, J., and Patriat, P., 1980. Existence d'anomalies mesozoïques dans le bassin de  
479 Somalie. Implications pour les relations Afrique-Antarctique-Madagascar. *C.R. Acad.  
480 Sci. Paris* 291, 85-88.

481 Simpson, E.S.W., Sclater, J.G., Parsons, B., Norton, I., Meinke, L., 1979. Mesozoic magnetic  
482 lineations in the Mozambique Basin. *Earth Planet. Sci. Lett.* 43, 260-264.

483 Smith, A.G. and Hallam, A., 1970. The fit of the Southern Continents. *Nature* 225, 139-144.

484 Sylvester, A.G., 1988. Strike-slip faults. *Geological Society of America Bulletin* 100, 1666-  
485 1703.


486 Torsvik, T.H., and Cocks, L.R.M., 2013. Gondwana from top to base in space and time.  
487 *Gondwana Research* 24, 999-1030.

488

489


490

491  
 492  
 493  
 494  
 495  
 496  
 497  
 498  
 499




500  
 501 FIGURE 1: Bathymetric map of the Africa-Antarctic corridor, the Mozambique Basin and the  
 502 Riiser-Larsen Sea (ETOPO1 1 arc-minute global relief model; Amante and Eakins, 2009) A).

503 The yellow flow lines indicate the motion between Africa and Antarctica according to Eagles  
504 and König (2008). Red boxes indicate the study area in the Mozambique Basin and the Riiser-  
505 Larsen Sea. AR= Astrid Ridge, EB= Enderby Basin, GR= Gunnerus Ridge, MB=  
506 Mozambique Basin, RLS= Riiser-Larsen Sea WSB= West Somali Basin. B). Black and  
507 orange lines indicate the locations of the reflection seismic profiles of the BGR14 and  
508 Mbwg00 datasets. Locations of Profiles A, C and D (Figs. 2A, 4 and 6) are highlighted with  
509 red lines. The location of Beira High is from Mahanjane (2012). Magnetic isochrons (yellow  
510 lines) and oceanic fracture zones (dashed white lines) compiled from Leinweber and Jokat  
511 (2012) and Müller and Jokat (2017). The location of Davie Ridge is marked with solid white  
512 line. BH= Beira High. C) Thick black lines indicate the location of the reflection seismic  
513 profiles of the RAE43 dataset. Position of Profiles B and E (Figs. 2B and 7) are highlighted  
514 with red lines. Magnetic isochrons (yellow) and fracture zones (dashed white lines) are  
515 compiled from Leinweber and Jokat (2012) and Leitchenkov et al. (2008). Continent-ocean  
516 transition as interpreted from Leitchenkov et al. (2008) is indicated with a thick orange line.




517  
518 **FIGURE 2:** Migrated profile Mbwg00-511 in the Mozambique Basin (A) and stacked profile  
519 RAE4303 in the Riiser-Larsen Sea (B) (line locations in inset maps and Fig. 1). In the lower  
520 panels, the stratigraphic interpretation according to Castelino et al. (2015), Leitchenkov et al.


521 (2008) and Mahanjane (2014) is presented. At the foot of the continental slope, at the  
 522 continent-ocean transition, a 20-30 km wide zone of deformed and fractured basement is  
 523 distinct. Postrift sediments, overlying the deformation zone have been affected by the  
 524 deformation. This deformation zone is proposed as tie-point for Gondwana reconstructions.



525  
 526 FIGURE 3: Close-up view of the zone of deformed basement in the Mozambique Basin (A)  
 527 and Riiser-Larsen Sea (B) presented in Fig. 2. The lower panels show the interpreted sections  
 528 of the profiles. The basement is distinctively deformed and faulted. Overlying postrift  
 529 sediments are deformed and indicate toplap (A) and onlap (B) geometries. Unconformity MJ,  
 530 with an inferred age at the transition from the Middle to Late Jurassic, seals the deformation.  
 531 Blue sedimentary horizons indicate Late Jurassic and Middle Cretaceous strata, the green  
 532 horizon is near the top Late Cretaceous.




533  
 534 FIGURE 4: Migrated section of profile Mbwg00-510 (line location in inset map and Fig. 1).  
 535 The lower panel shows the section overlain by the stratigraphic interpretation according to  
 536 Castelino et al. (2015), Franke et al. (2015) and Mahanjane (2014). The profile runs from the  
 537 continental slope to the Davie Ridge offshore Madagascar. The zone of deformed basement is  
 538 observed at the foot of the continental slope. The Davie Ridge appears as bathymetric high,  
 539 rising 1 s (TWT) above the surrounding seafloor. At the foot of the western flank of Davie  
 540 Ridge, a zone of deeply buried, compressed sediments is observed that might have been  
 541 thrust onto the oceanic crust during southward motion of Madagascar.  
 542




543

544 FIGURE 5: Close-up view of the zone of deformed basement in the Mozambique Basin  
 545 presented in Fig. 4. The lower panel shows the interpreted section of the profile. The  
 546 basement is deformed by steeply dipping, fan-like normal faults that at depths converge into a  
 547 single, subvertical fault (green, profile distance: 40-60 km). The overall geometry of the  
 548 deformation resembles a negative flower structure. Unconformity MJ has an inferred age at  
 549 the transition from the Middle to Late Jurassic and the blue sedimentary horizons indicate  
 550 Late Jurassic and Middle Cretaceous strata.


551




552  
 553 FIGURE 6: Pre-stack migrated section of profile BGR14-305 (line location in inset map and  
 554 in Fig. 1). The lower panel shows the interpreted section according to the seismostratigraphic  
 555 concepts of Castellino et al. (2015), Franke et al. (2015), Klimke et al. (2016) and Mahanjane  
 556 (2014). The profile runs from the continental slope offshore northern Mozambique across the  
 557 Davie Ridge into the Morondava Basin offshore Madagascar. The zone of deformed basement  
 558 is observed at the foot of the continental slope (offset range: 30-50 km), where the basement  
 559 is not imaged, which is probably due to the intense faulting of the basement. The Davie Ridge  
 560 shows a clear morphological expression in the center of the profile. The shear zone, however,  
 561 is much wider and, including the Davie Ridge, it is characterized by three prominent crustal  
 562 blocks. Overall, the deformation zone extends over ~120 km, perpendicular to the shear  
 563 movement. The westernmost block consists of deeply buried, compressed sediments that  
 564 might have been thrust onto the oceanic crust during southward motion of Madagascar.



565  
 566 FIGURE 7: Stacked section of profile RAE4307 (line location in inset map and in Fig. 1). The  
 567 lower panel shows the stratigraphic interpretation according to Leitchenkov et al. (2008). The  
 568 profile runs from the Riiser-Larsen Sea across the Gunnerus Ridge into the Enderby Basin.  
 569 The Gunnerus Ridge rises  $\sim 4$  s (TWT) above the surrounding seafloor. The transition from  
 570 continental to oceanic crust along the Gunnerus Ridge is very abrupt ( $\sim 30$ - $40$  km). The  
 571 oceanic crust of the Riiser-Larsen Sea is dissected by normal faults. The abundance of the  
 572 faults increases significantly towards the Gunnerus Ridge.




573  
 574 FIGURE 8: Closeup view of the faulted oceanic basement presented in Fig. 7 in the Riiser-  
 575 Larsen, Sea, close to the Gunnerus Ridge. The lower panel shows the interpreted section of the  
 576 profile.



577  
 578 FIGURE 9: Sketch map illustrating the location of the deformed basement observed at the  
 579 foot of the continental slope in the Mozambique Basin (A) and the Riiser-Larsen Sea (B).  
 580 Solid lines indicate the location of Profiles A to E (Figs. 2, 4, 6 and 7). Blue and black lines

581 highlight magnetic anomalies and fracture zones in the Mozambique Basin and Riiser-Larsen  
 582 Sea according to Leinweber and Jokat (2012), Leitchenkov et al. (2008) and Müller and Jokat  
 583 (2017). The continent-ocean transition (COT) as proposed in this study is shown in green. The  
 584 continent-ocean transition according to Leitchenkov et al. (2008) in the Riiser-Larsen Sea is  
 585 shown in purple. Orange lines indicate wrench faulting in the West Somali Basin (Klimke et  
 586 al., 2016). Blue dots mark onlap locations of horizon MJ against oceanic basement. Dashed  
 587 blue line marks the extrapolation of magnetic chron M38n (Müller and Jokat, 2017) to the  
 588 study area in the Mozambique Basin. The extrapolation was done by noting the distance of  
 589 magnetic chron M38n from the continent-ocean transition. Yellow and rose hatched areas  
 590 mark the location of transpressional (yellow) and transtensional (rose) deformation. The  
 591 location of Davie Ridge is marked with thick dashed red line. The location of SDRs in the  
 592 Mozambique Basin is compiled from Leinweber et al. (2013) and Müller and Jokat (2017).



593  
 594 FIGURE 10: Schematic sketch of the initial opening of the Mozambique Basin/Riiser-Larsen  
 595 Sea. A) In the Middle Jurassic, NW-SE directed rifting and seafloor spreading between Africa  
 596 and Antarctica initiates with the possible formation of localized spreading centers close to the

597 present-day shoreline. B) By the Late Jurassic, the spreading center has jumped to the south  
598 and the NW-SE extensional phase has been replaced by N-S oriented seafloor spreading. At  
599 the eastern margin of the evolving Mozambique Basin/Riiser Larsen Sea Basin, a transform  
600 deformation phase, overprinting the previous continent-ocean transition, accommodates the  
601 extension. The transform fault develops along the conjugate western flanks of the Davie  
602 Ridge and the Gunnerus Ridge. The positions of Madagascar, Antarctica and India have been  
603 adopted from Nguyen et al. (2016). Locations of SDRs, lava flows and intrusions in the  
604 Mozambique Basin are taken from Mahanjane (2012) and Müller and Jokat (2017). Magnetic  
605 chrons taken from Leinweber and Jokat (2012) and Müller and Jokat (2017). Thick green  
606 lines mark the basement deformation zone presented in this study. Dashed green line marks  
607 the continent-ocean transition (COT) of this study. Dashed purple line is the continent-ocean  
608 transition (COT) of Leitchenkov et al. (2008). SW-propagating, NW-SE oriented oblique  
609 rifting and seafloor spreading between Madagascar and Africa according to Klimke and  
610 Franke (2016). AR= Astrid Ridge, BH= Beira High, GR= Gunnerus Ridge.

611