Deeply subducted continental fragments: II. Insight from petrochronology in the central Sesia Zone (Western Italian Alps)

Francesco Giuntoli¹, Pierre Lanari¹, Marco Burn¹, Barbara Eva Kunz¹& Martin Engi¹

¹Institute of Geological Sciences, University of Bern, Baltzerstrasse 1+3, 3012 CH-Bern

5 Correspondence to: Francesco Giuntoli (francesco.giuntoli@plymouth.ac.uk)

10

15

Abstract. Subducted continental terranes commonly comprise an assembly of subunits that reflect the different tectonometamorphic histories they experienced in the subduction zone. Our challenge is to unravel how, when and in which part of the subduction zone these subunits were juxtaposed. Petrochronology offers powerful tools to decipher pressure-temperature-time (P-T-t) histories of metamorphic rocks that preserve a record of several stages of transformation. A major issue is that the driving forces for re-equilibration at high pressure are not well understood. For example, continental granulite terrains subducted to mantle depths frequently show only partial and localized eclogitization. The Sesia Zone (NW Italy) is exceptional because it comprises several continental subunits in which eclogitic rocks predominate and high-pressure (HP) assemblages almost completely replaced the Permian granulite protoliths. This field-based study comprises both main complexes of the Sesia terrane, covering some of the recently recognized tectonic subunits involved in its assembly, hence our data constrain the HP-tectonics that formed the Sesia Zone. We used a petrochronological approach consisting of petrographic and microstructural analysis linked with thermodynamic modeling and U-Th-Pb age dating to reconstruct the P-T-t trajectories of these tectonic subunits. Our study documents when and at what conditions re-equilibration took place. Results constrain the main stages of mineral growth and deformation, associated with fluid influx

- 20 that occured in the subduction channel. In the Internal Complex (IC), pulses of fluid percolated at eclogite facies conditions, between 77 and 55 Ma with the HP-conditions reaching ~2 GPa and 600-670 °C. By contrast the External Complex (EC) records a lower pressure peak of ~0.8 GPa for 500 °C, at ~63 Ma. The juxtaposition of the two complexes occurred during exhumation, probably at ~0.8GPa and 350°C; the timing is constrained between 46 and 38 Ma. Mean vertical exhumation velocities are constrained between 0.9 and 5.1 mm/year for the IC, up to its juxtaposition with the EC. Exhumation to the
- 25 surface occurred before 32 Ma, as constrained by the overlying Biella Volcanic Suite, at a mean vertical velocity between 1.6 and 4 mm/year. These findings constrain the processes responsible for the assembly and exhumation of high pressure continental subunits, thus adding to our understanding of how continental terranes behave during subduction.

1 Introduction

The behaviour of continental crust subducted to high-pressure (HP) conditions is far from fully understood (Kylander-Clark et al., 2008;Rubatto and Hermann, 2001;Brun and Faccenna, 2008;Malusà et al., 2011;Angiboust et al., 2016). Seismic tomography beneath collisional orogens shows that large slab parts reached mantle depths (e.g. Lippitsch et al.,

- 5 2003;Replumaz et al., 2010;Zhao et al., 2015), but the source of such remnants is hard to assess because the rocks cannot be directly investigated. However, where orogens contain exhumed continental HP-rocks, these offer opportunities to investigate some of the tectono-metamorphic processes involved, notably those responsible for the return flow of continental fragments back to the surface.
- In recent studies on the Western Alps, the combination of tectonic and numerical modeling studies (e.g. Yamato et al., 2008;Faccenda et al., 2009) with the reconstruction of pressure-temperature-time (P-T-t) paths (e.g. Regis et al., 2014;Rubatto et al., 2011) has led to two possible end-members scenarios for the subduction of continental domains to HP conditions. Either the units essentially experienced tectonic slicing, followed by accretion to overlying continental units, thus assembling complexes composed of different tectonometamorphic slices (e.g. Regis et al., 2014;Manzotti et al., 2014;Vitale Brovarone et al., 2013;Angiboust et al., 2014). Alternatively, units were eroded or ablated in the subduction channel, which
- 15 lost coherence and experienced substantial mixing, leading to diverse and complex P-T-t paths (e.g. Tao and O'Connell, 1992;Stöckhert and Gerya, 2005;Warren et al., 2008;Keppie et al., 2009;Roda et al., 2012). The distinction between these two end-members scenarios is important to understand the evolution of deeply subducted continental domains and possibly their paleogeographic provenance.
- Crucial questions related to continental HP-units in orogens include: How rapidly were they subducted and to what depth?
 When and how fast were they exhumed? What P-T-t trajectories did they experience? When and how did fluids affect these continental fragments? To shed light on these questions, the sequence of metamorphic stages recorded in suitable samples needs to be analysed in detail (Engi et al., 2017). P-T-t paths promise insight into details of these fundamental tectonic processes. Petrochronological studies in the Western Alps have helped to constrain exhumation rates of Alpine HP and UHP continental domains. Calculated rates vary from a few mm (e.g. Zucali et al., 2002;Regis et al., 2014 for the internal part of
- 25 the Sesia Zone) to a few centimetres per year (e.g. Rubatto and Hermann, 2001 for Dora Maira UHP massif). In detail, some studies found that, after a first phase of rapid exhumation up to lower crustal levels, exhumation rates markedly decreased to a few millimetres per year (Rubatto and Hermann, 2001;Yamato et al., 2008).
 - Generally speaking, how much of the subduction history is recorded in a sample is related to what processes triggered mineral recrystallization or equilibration. Several studies (e.g. Erambert and Austrheim, 1993;Austrheim, 1987;Rubie, 1986;Oliver, 1996;Etheridge et al., 1983;Engi et al., in review;Pennacchioni, 1996) proposed that fluid influx triggers deformation and mineral reactions in deeply subducted high-grade (granulite and amphibolite facies) domains. It is of interest to know when this happened in the P-T-t evolution, notably whether it occurred early, during subduction or only late, upon exhumation (e.g. Konrad-Schmolke et al., 2011).
- 30

10

We present a field-based study emphasizing P-T-t data and discuss their implications on such first order questions. Datasets from a well defined multidimensional analysis, i.e. pressure-temperature-time-deformation-space, are essential as a reference frame for numerical modelling studies. Our petrochronological approach highlights the complex and heterogeneous tectonometamorphic evolution in a polydeformed continental terrane. A key requirement in this approach is to establish reliable links between age data (t) and the P-T conditions of mineral formation (e.g. Schenk, 1989;Buick and Holland, 1989;Scott and St-Onge, 1995;Liati and Gebauer, 1999;Rubatto and Hermann, 2001;Foster et al., 2004;Janots et al., 2009;Gasser et al., 2012;Donaldson et al., 2013;Rubatto et al., 2011;Regis et al., 2014;Mottram et al., 2014). In this study, mutual inclusions relationships of the main mineralogical phases in the datable accessory minerals were used, along with microstructural analyses, to bracket mineral age data to P-T conditions. The study area is the central Sesia Zone, located in the Western Alps (Valle d'Aosta, Italy). P-T-t data are reconstructed for the eclogite facies IC and, for the first time, for the epidote-blueschist facies EC. These data allow us to constrain the juxtaposition of the two complexes, which occurred under HP greenschist facies conditions, and to determine exhumation rates for the IC and for the assembly of the Sesia Zone.

2 Geological setting

- The Alps are an orogen that has developed since the Cretaceous due to the subduction of the European Plate below the
 Adriatic Plate, with subsequent continental collision (e.g. Dewey et al., 1989;Rosenbaum et al., 2002;Handy et al., 2010;Beltrando et al., 2010;Schmid et al., 2017). The Sesia Zone (SZ) is located in the Western Alps (Fig. 1a) and regarded as a rifted portion of the Adriatic Margin that experienced subduction to HP conditions (e.g. Dal Piaz, 1999). At present, it is bounded by two tectonic lineaments: the Insubric Line, to the SE, and the Gressoney Shear Zone, to the NW. The Insubric line is a major fault system of Oligocene-Neogene age that separates the SZ from the Southern Alpine domain, which
 represents the Adriatic Margin and shows but a weak Alpine imprint at sub-greenschist facies. An important complex in the
- Southern Alps is the Ivrea Zone, consisting of amphibolite to granulite facies meta-sediments, mantle peridotites and mafic intrusives. It is proposed to represent a cross section through the lower pre-Alpine crust (e.g. Bertolani, 1959, 1954;Rivalenti et al., 1975;Zingg, 1983;Quick et al., 2003). Analogies between the pre-Alpine metamorphism of the Internal Complex (Giuntoli and Engi, 2016) of the SZ and the Ivrea Zone have been repeatedly emphasized (e.g. Dal Piaz et al.,
- 25 1972;Compagnoni et al., 1977). The Gressoney Shear Zone is a greenschist shear zone (e.g. Wheeler and Butler, 1993;Babist et al., 2006) that separates the SZ from units derived from the Piedmont-Ligurian ocean that experienced blueschist to eclogite facies metamorphism during the Alpine orogeny (e.g. Bucher et al., 2005;Negro et al., 2013).

Recently, the Central Sesia Zone has been subdivided into an Internal Complex (IC) and an External Complex (EC; Giuntoli and Engi, 2016; Fig. 1b-c). The IC corresponds, as a whole, to the Eclogitic Micaschist Complex of Williams and
Compagnoni (1983) and Passchier et al. (1981). In detail, however, the IC comprises several eclogitic sheets with a thickness of few kilometres, each of them characterized by an alternance of bands and elongate bodies of micaschist, eclogite, ortho-

and paragneisses, and leucogneiss. The main foliation of this complex is of eclogite facies. Several deformation phases locally overprint this foliation at retrograde blueschist and greenschist facies conditions (details in Giuntoli and Engi, 2016).

The EC corresponds to the Gneiss Minuti and the 2DK complexes of previous authors. It comprises km-thick tectonic sheets of leuco- to mesocratic orthogneiss with minor paragneiss, calc-micaschist, impure quartzite and marble separated by lenses
preserving Permian high-temperature (HT) metamorphic relics (2DK; Giuntoli and Engi, 2016;Carraro et al., 1970;Dal Piaz et al., 1971). The main composite foliation reflects HP greenschist, at most epidote blueschist facies; these are the highest Alpine metamorphic conditions recorded by this complex. The HP-foliation is affected by subsequent greenschist facies deformation. Juxtaposition of the two complexes involved a greenschist facies shear zone (Barmet Shear Zone in Giuntoli and Engi, 2016); the subunits in each complex are thin sheets that range in thickness from 0.5 to 3 km.

- 10 Based on P-T-t paths determined by petrochronological techniques, Rubatto et al. (2011) and Regis et al. (2014) recognized two tectonometamorphic subunits in the IC: A more internal subunit, called Druer slice, experienced eclogite facies condition with pressure of 1.9-2 GPa and temperatures of ~550 °C at around 85 Ma, then followed by exhumation. A more external subunit, called Fondo slice, experienced a first stage at eclogite facies, with pressures of 1.7-1.8 GPa and temperatures between 520 and 550°C at around 75 Ma, then followed by an intermediate lower pressure stage (P<1.6 GPa and T<520 °C) at around 68 Ma, and a second high pressure stage (P=1.4-2 GPa and ~550 °C) at around 65 Ma, followed by</p>
- retrograde decompression. For the EC in the central Sesia Zone, no P-T-t data are available so far. A number of studies have produced additional age data for the Sesia Zone, using various methods, notably for the U-Th-Pb, Rb-Sr, and Ar-Ar systems; results are summarized in Table 1 of Compagnoni et al. (2014). These age data span from 85 to 62 Ma for the eclogite facies metamorphism in the IC, but apart from those detailed above, none of the datasets report
- 20 detailed links to petrogenetic conditions. Few age data are available for the EC, ranging from 46 to 38 Ma, and these are generally linked to the greenschist facies imprint (Compagnoni et al., 2014 and references therein).

3 Sampling strategy and petrochronological approach

To document the polyphase history of the Sesia Zone, we reconstructed detailed P-T-t paths for five samples taken in the IC and two samples in the EC. Of over a hundred samples checked, a very small percentage fulfilled the requirements for such a study. In the EC, in particular, suitable material to quantify P-T-t conditions by present methods turned out to be very rare. This is mostly due to the fact that orthogneiss, the dominant rock type in the EC, is lacking in minerals suitable to obtain the Alpine P-T-t path, as discussed by Giuntoli and Engi (2016). Nevertheless, the samples analysed provide constraints to derive a P-T-t path for the EC as well, allowing us to determine when and at what conditions the two complexes were juxtaposed.

- 30 All samples were taken in key areas of the mapped structures and were collected oriented, in order to keep the link between the meso- and microstructural evidence. Samples were carefully studied by optical and (where needed) scanning electron microscopy (SEM) to reconstruct their microstructural and metamorphic evolution. Once a relative chronology was
 - 4

established, selected growth zones of mineral phases were analysed by electron probe micro-analyser (EPMA) as a basis to perform thermodynamic modelling. P-T data were linked to fabric elements, using textural criteria. Geochronology was performed by LA-ICP-MS, targeting specific growth zones of accessory phases that were separated after noting their microstructural and geochemical context. It turned out to be critical to analyse each growth zone separately to link the age (t) to a specific metamorphic stage (P-T). Observations and P-T-t data derived from each sample are then compared and correlated within the sample series and then related to observations made in the field data and in microscopy. This detailed petrochronological approach (e.g. Engi et al., 2017) is particularly effective when applied to a study area that has been mapped and structurally characterized in detail, as in the present case.

4 Petrography and mineral chemistry

10 4.1 Methods

5

4.1.1 SEM

Back-scattered electron images (BSE) were acquired using the Zeiss EVO50 SEM at the Institute of Geological Sciences (University of Bern) using an accelerating voltage of 15 to 25 KeV, a beam current of 500 pA and a working distance of 10 mm. Cathodoluminescence (CL) pictures where obtained with the same operative conditions, but with 10 KeV accelerating voltage and a working distance of 9.5 mm.

4.1.2 EPMA analyses

EPMA analyses were performed using a JEOL JXA-8200 superprobe at the Institute of Geological Sciences (University of Bern). Point mode analyses and X-ray compositional maps were acquired both using wavelength dispersive spectrometers (WDS). For X-ray mapping the procedure described in Lanari et al. (2013) was followed. It consists in measuring point 20 mode analyses first and then acquiring X-ray compositional maps on the same area. For point analyses, analytical conditions were 15 KeV accelerating voltage, 10 to 20 nA specimen current, 40 s dwell times (including 2×10 s of background measurement) and a beam ø from 1 to 5 µm. Lower current and higher beam size were used for minerals containing Ca, Na and K such as phengite and plagioclase. Nine oxide compositions were measured, using synthetic and natural standards: wollastonite / orthoclase / almandine (SiO₂), anorthite / almandine (Al₂O₃), anorthite (CaO), almandine (FeO), forsterite / 25 spinel (MgO), orthoclase / phlogopite (K₂O), albite (Na₂O), ilmenite (TiO₂), and tephroite (MnO). For X-ray maps, analytical conditions were 15 KeV accelerating voltage, 100 nA specimen current and dwell times of 150-250 ms. Nine elements (Si, Ti, Al, Fe, Mn, Mg, Na, Ca and K) were measured at the specific wavelength in two passes. Intensity X-ray maps were standardized to concentration maps of oxide weight percentage using spot analyses as internal standard. X-ray maps were processed using XMapTools 2.2.1 (Lanari et al., 2014).

15

4.2 Results: Sample description

15

25

Five samples were analysed from the IC and two samples from the EC. These were collected from internal (SE) to external areas (NW) of the Sesia Zone. A brief account is given here, with characteristic images shown in Fig. 2. Supplement S1 contains detailed descriptions and GPS locations.

- 5 The samples of the IC (FG1324, FG1315, FG12157, FG1347, and FG1249) are micaschists with eclogite facies assemblages comprising quartz, phengite, garnet, ± paragonite ± glaucophane ± omphacite ± chloritoid, with accessory allanite ± rutile. The main fabric in all of these samples is an eclogite facies foliation (Fig. 2a). Evidence of several deformation stages occurring before or after the main eclogite facies foliation is preserved in several samples as microlithons, commonly of phengite, omphacite, glaucophane or chloritoid oriented at high angles relative to the main foliation, which wraps around
- 10 them or is overgrown by them (Fig. 2b). Further evidence of several metamorphic stages occurring at eclogite facies conditions is reflected in growth zones of garnet (Giuntoli et al., in review). Pre-Alpine relics include garnet cores (Fig. 2c) and zircon (cores ± first rims; chapters 6.4 and 6.5).

Retrograde stages of blueschist or greenschist facies assemblages related to decompression are locally present in samples. The blueschist facies stage produced pleochroic crossite rims around glaucophane (Fig. 2d). The greenschist facies stage produced symplectites of actinolite \pm albite \pm chlorite around glaucophane and omphacite, chlorite at the expense of garnet,

- epidote/clinozoisite rims around allanite, and titanite rims formed around rutile (Fig. 2e-f). Sample FG1420, collected in the EC, is a garnet orthogneiss that shows a HP greenschist foliation marked by phengite, chlorite, and titanite; the foliation wraps garnet porphyroblasts that preserve a relic internal foliation (Fig. 2g). Permian
- magmatic relics of pleochroic allanite are surrounded by an Alpine corona of epidote grains (Fig. 2h). Some hundred meters
 to the north, another sample (FG12107) of the EC was collected. This is a leucogneiss characterized by the same metamorphic fabric and parageneses as the previous sample, except that garnet and magmatic allanite are missing.

4.3 Results: Growth zones of garnet and phengite

Garnet and phengite display features in the IC samples that differ from those in the EC samples. To highlight and describe these, mostly two samples are compared in the following paragraphs: FG1249 (IC) and FG1420 (EC). A more complete account of garnet textures and mineral inclusions in the IC is shown in Giuntoli et al. (in review).

- In the IC samples, garnet consists of a core followed by several rims with a grain size up to several millimeters (Fig. 3a, b). The compositional map of the grossular end-member fraction (X_{Grs}) in sample FG1249 shows a porphyroclastic core (Alm₇₂Prp₁₈Grs₅Sps₅; Table 1) with internal fractures sealed by garnet of higher X_{Grs} (Fig. 3b). A first rim (rim1-Alm₇₆Prp₁₅Grs₉) overgrows the core and displays higher grossular contents. This rim1 is followed by rim2, which again records an increase of X_{Grs} (Alm₆₂Prp₂₀Grs₁₈). Rim2 resorbs parts of rim1 externally, as well as parts of rim1 internally and
- core. Rim3 is peripheral and shows the highest Ca contents (Alm₅₈Prp₁₉Grs₂₃).
 - 6

Sample FG1315 is characterized by a porphyroclastic core ($Alm_{69}Prp_{28}Grs_4$) with lobate edges and resorption features (details in Giuntoli et al. submitted and Engi et al. in prep.) surrounded by several rim generations: rim1 ($Alm_{61}Prp_{21}Grs_{19}$), rim2 ($Alm_{65}Prp_{24}Grs_{11}$) and rim3 ($Alm_{70}Prp_{24}Grs_6$). Atoll garnets, a few hundred microns in size, are observed in this sample. The shells of the atoll garnet have similar zoning patterns and compositions as the rim generations just described. In

- 5 sample FG12157 the garnet core (Alm₇₀Prp₂₆Grs₄) is rimmed by two growth zones: rim1 (Alm₆₄Prp₂₀Grs₁₆) and rim2 (Alm₅₉Prp₂₄Grs₁₇). In sample FG1347, the garnet core (Alm₆₉Prp₂₈Grs₃) is enclosed by three rims (rim1-Alm₆₆Prp₂₃Grs₁₁, rim2- Alm₆₈Prp₂₆Grs₆, rim3-Alm₇₀Prp₂₆Grs₄). The exception is sample FG1324, in which garnet shows a single growth zone of homogeneous composition (Alm₇₀Prp₁₈Grs₁₁Sps₁).
- In the EC, sample FG1420 shows garnet with completely different features. As shown in Fig. 3d, the X_{Sps} map highlights
 concentric zoning (values of Alm₅₄Prp₂Grs₃₆Sps₈ for the core, Alm₅₇Prp₃Grs₃₇Sps₄ for rim1, Alm₆₀Prp₃Grs₃₅Sps₂ for rim2 and Alm₆₃Prp₄Grs₃₂Sps₁ for rim3), with no visible resorption features (further compositional end-member maps are shown in Supplement S2).

To link the growth zones of garnet to the main assemblage observed in the mineral matrix, microstructural relations, overprinting criteria and mutual inclusions were employed, based on optical microscopy, SEM, and compositional maps. In

- 15 particular, garnet in sample FG1249 contains inclusions of paragonite, phengite, and quartz between rim1 and rim2 (Fig. 3a). Rutile inclusions of few microns are present in rim2 and 3. Late chlorite fractures dissect the entire garnet. Garnet in sample FG1420 is wrapped by the main external foliation and includes an internal foliation marked by quartz, epidote, and titanite (Fig. 3c).
- Phengite in IC samples displays a uniform composition except along grain boundaries, where lower Si and X_{Mg} contents are found, indicating retrograde overprinting (e.g. Fig. 3e; Group1 Si ~3.36 apfu, X_{Mg} ~0.83; Group2 Si ~3.3 apfu, X_{Mg} ~0.68 in sample FG1249; Table 2).

In the EC, two distinct generations of phengite are distinguished based on their microtextural position: the first one describes the main foliation and is characterized by high Si values (Fig. 3f; Group1 Si ~3.4 apfu, X_{Mg} ~0.61 in sample FG1420). The second phengite generation (Group2 Si ~3.32 apfu, X_{Mg} ~0.61) rims the first one and occurs in fold hinges that deform the main foliation.

5 Thermobarometry

5.1 Methods

25

5.1.1 Whole rock major element compositions

Major element compositions were analysed by X-ray fluorescence (XRF) spectrometry at the University of Lausanne (Switzerland). Representative quantities of samples were crushed and then pulverized in a tungsten carbide mill. The powder

was dried for two hours at 105°C. Loss of ignition was then determined by weight difference after heating to 1050°C for 3 hours.

5.1.2 Garnet thermobarometry using GrtMod

Garnet growth zones were carefully selected after detailed microstructural and compositional analysis of the high-resolution
X-ray maps to extract representative compositions that are used as input for modeling. Following the strategy proposed by Lanari et al. (2017), the composition of each growth zone was obtained directly from the quantitative maps by sampling chemically homogeneous areas amongst different garnet grains. Each composition was then assigned to a specific growth zone that is assumed to be uniform in the following. The regularity of the chemical zoning observed in garnet (Giuntoli et al., in review) supports the grain boundary equilibrium model adopted here (Lanari and Engi, 2017). Minor heterogeneities
observed in each growth zone (<0.01 in XAIm and XGrs; < 0.005 in XPrp, XSps) may be due to kinetic effects during growth, they do not affect the results of the equilibrium model.

To model the complex garnet textures adequately, fractionation and resorption processes must be taken into account in approximating the evolution of the reactive bulk composition. The latter is strongly affected by fractionation (i.e. the removal of refractive garnet, e.g. Evans, 2004;Konrad-Schmolke et al., 2008) and by resorption, which can shift the reactive

- 15 bulk composition back toward the garnet composition (Lanari and Engi, 2017). The program GRTMOD (Lanari et al., 2017) was specially developed to deal with samples in which garnet experienced a complex history involving several stages or growth, resorption and/or pseudomorphic replacement. In essence, the program refines the reactive bulk composition used in free energy minimization at each stage predicted if previously grown garnet was preserved or only partly dissolved. GRTMOD uses an iterative approach that refines the P-T conditions for successive garnet growth zones. For each inversion
- 20 (i.e. a single growth stage), a solution was deemed acceptable if the least squares residual value (the cost function C_0 used by Lanari et al., 2017) was <0.05, reflecting a sufficiently close match between the modeled and observed garnet compositions. In the IC samples, resorption and fractionation were constrained according to the volumetric proportion of each growth zone, as estimated from the thin section and the compositional maps. No resorption was permitted in the program for sample FG1420, as garnet textures in the compositional maps show no evidence of this process. To model the rims generation in the
- 25 IC samples the "go fast mode" function (Lanari et al., 2017) was used, with an initial starting P-T guess of 650 °C and 1.6 GPa. The initial starting guess is an essential technicality used for reasons discussed by Lanari et al. (2017, sections 5.2.3 and 6.3); the specific purpose is to avoid local minima in optimization. This function searches a solution around the starting guess and follows the gradient in the objective function; there cannot be local minima at high pressure for this range of bulk rock composition (see Fig. 8 in Lanari et al., 2017). The MnO component was used in the thermodynamic computations of
- 30 the relatively rich Mn-garnet in sample FG1420. In sample FG1249, MnO was used to model the garnet core but was ignored in the models of the following rims because the concentration found in garnet is low (< 1 wt% MnO; Table1). In the remaining samples, MnO was ignored (< 1 wt%), and the system considered in modelling was simplified to SiO₂-TiO₂-

 Al_2O_3 -FeO-MgO-CaO-Na₂O-K₂O-H₂O. The thermodynamic database used was the same as to compute the isochemical phase diagrams (see below).

5.1.3 Isochemical phase diagrams (pseudosections)

Isochemical equilibrium phase diagrams were computed using the Gibbs free energy minimization algorithm THERIAK/DOMINO (de Capitani and Petrakakis, 2010;De Capitani and Brown, 1987). The thermodynamic database of Berman (1988) with subsequent updates collected in JUN92.bs (distributed with Theriak-Domino 03.01.2012; Supplement S3) was used, together with the following solution models: Berman (1990) for garnet; Fuhrman and Lindsley (1988) for feldspar; Meyre et al. (1997) for omphacite; Keller et al. (2005) for white mica, and ideal mixing models for amphibole (Mäder and Berman, 1992;Mäder et al., 1994), epidote, and chlorite (Hunziker, 2003). All Gibbs free energy minimizations were carried out assuming an excess in pure H₂O fluid. The amount of H₂O component predicted at high-pressure is in line with the measured LOI (1.4-2.7 wt-%) in the present-day samples. Note that for the pre-Alpine HT computations no melt model was used. Fe³⁺ was ignored because of the lack of analytical data and suitable ferric end-members in solid solution

5.1.4 Chlorite and white mica multi-equilibrium

- To constrain the P-T conditions of retrograde stages, multi-equilibrium computations of the high-variance assemblages involving chlorite and white mica were carried out, using the standard state properties and solid solution models of Vidal et al. (2005; 2006) for chlorite, Dubacq et al. (2010) for phengite, and the program CHLMICAEQUI (Lanari, 2012). The activity of H₂O was set to unity. Three methods were successively employed:
 - (1) *Chlorite+Quartz+H*₂*O* thermometry: The chlorite formation temperature and XFe³⁺ were estimated at a fixed pressure of 1 GPa from the combination of four equilibria involving five chlorite end-members, quartz and H₂O (Lanari et al., 2012; Vidal et al., 2016).
 - (2) White-Mica+Quartz+H₂O barometry: A divariant P-T equilibrium line was estimated for each white mica analysis (assuming $XFe^{3+} = 0$) from the convergence of three equilibria involving five phengite end-members, quartz and H₂O (Dubacq et al., 2010).
- 25 (3) Chlorite+White-Mica+Quartz+H₂O thermobarometry: P and T of formation for each chlorite and white mica couple, as well as their respective XFe^{3+} , were estimated by minimizing the square root of the sum of $(\Delta G_{reaction})^2$ for 6 equilibria (see Supplement S4).

For the sake of clarity, only 64 equilibria (excluding the Pyrophyllite-1H₂O end-member) are shown in the P-T diagrams. The starting guess for T and P was taken from the result of Chlorite+Quartz+H₂O thermometry and White-Mica+Quartz+H₂O barometry. This multi-equilibrium approach relies on the assumption of local thermodynamic equilibrium between the selected chlorite and white mica at the P-T conditions of convergence. Chlorite and white mica

30

models.

couples were chosen where microtextural evidence suggested equilibrium, notably where sharp contacts were observed between these sheet silicates.

5.2 Bulk rock and reactive bulk composition

For samples FG1324 and FG1420 the original bulk rock compositions obtained by XRF were used to compute isochemical phase diagrams. In samples FG1315, FG12157, FG1347 and FG1249 however, the unmodified bulk rock composition cannot be used for modeling because a significantly high volume fraction of garnet is present (5 – 10 vol%), including a pre-Alpine core and Alpine rims generations. To compute equilibrium diagrams properly, the reactive bulk rock composition was approximated using the program GRTMOD (see section 5.1.2). Each isochemical phase diagram is thus valid for a single P-T stage only. To select the reactive bulk composition of this specific stage, a link must be established between the particular garnet generation that formed in equilibrium with the mineral phases present in the matrix. We established this link using petrographic observations, including textural equilibrium criteria, compositional zoning (visible in compositional maps) and inclusions relationships. Specifically, we determined that the garnet growth zones that coexisted with the mineral matrix are as follows: garnet rim3 for sample FG1315, rim1 for sample FG12157, rim3 for sample FG1347, and rim2 for

15 **5.3 Results:** Garnet thermobarometry and phase diagrams

20

separately below.

Fig. 4 shows isochemical P-T phase diagrams for each sample. The plots show the results of garnet thermobarometry (GRTMOD), garnet isopleths (X_{Grs} , X_{Alm} , and X_{Prp} in sample FG1324), and X_{Mg} and Si (in apfu) isopleths for phengite. For sample FG1420, results of chlorite – white mica thermobarometry are also displayed. A summary of mineral compositional data for the main phases, the modelling method used, the XRF analyses of major elements of each sample, and details of the GrtMod results are available as Supplement S5, S6, S7 and S8 respectively. Each sample from the IC and EC is presented

sample FG1249. The corresponding reactive bulk compositions used for the modeling are provided in the Supplement S5.

5.3.1 IC - FG1324 Omphacite, garnet, glaucophane and rutile micaschist

Garnet isopleths (Alm, Grs, Prp) intersect in two areas of the P-T diagram: at 1.65-1.75 GPa and 600-650 °C and at 1.9-2 GPa and ~550 °C. Assuming equilibrium between garnet and phengite, these conditions can be further constrained by matching this result with the Si and X_{Mg} isopleths of phengite (measured values: 3.38-3.46 and 0.76-0.81 respectively). Phengite isopleths match the garnet composition at 1.65-1.75 GPa and 600-650 °C (Fig. 4a). At these conditions, some discrepancies between observations and modelled assemblages are noted for this sample: paragonite instead of glaucophane is predicted to be stable in the model (probably because no solid solution model for sodic amphiboles is available in the thermodynamic database used). Also, 6 vol% biotite is predicted to be stable, whereas none was observed in thin section.

5.3.2 IC - FG1315 Garnet, epidote and rutile quartz-micaschist

5

The garnet core is found to be stable at 0.82 GPa and 750 °C, with a crystallization of ~14 vol% (details available in Supplement S5 and S8). The first garnet rim is predicted stable at ~1.5 GPa and 650 °C, with a resorption of ~3.7 vol% garnet core and crystallization of ~5.5 vol% rim1. The second and third garnet rims are found to be stable at similar conditions: 1.9 GPa, 650°C and 1.8 GPa (resorption of 2.3, 2.5 vol% core and rim1, crystallization of 5.6 vol% rim2), and at 670°C (resorption of 5, 0.4, and 3.1 vol% core, rim1, and rim2, plus crystallization of 8.6 vol% rim3). P-T conditions of rim2 and 3 match with the intersection of Si (3.37-3.40) and X_{Mg} (0.80-0.82; Fig. 4b) isopleths of phengite. The predicted assemblage conforms to the minerals observed in thin section (Supplement S1, S5).

5.3.3 IC - FG12157 Garnet, glaucophane, epidote and rutile micaschist

In this model, ~7 vol% of garnet is modelled stable at 0.6 GPa and 900 °C; its composition corresponds to the garnet core. The first garnet rim is predicted stable at 1.6 GPa - 650 °C (resorption of ~1.4 vol% garnet core and crystallization of ~10.8 vol% rim1), in agreement with the intersection of phengite isopleths for the observed values of Si (3.34-3.38) and X_{Mg} (0.76-0.82; Fig. 4c). Crossite rims around glaucophane in this sample mark a decompression stage ; this may correlate with the second rim in garnet, which is found to be stable at 1.4 GPa and 650 °C (resorption of 1.5 vol% core, 7.3 vol% rim1, and crystallization of 10.6 vol% rim2). 4-6% biotite is predicted to be stable, but none is observed in thin section.

5.3.4 IC - FG1347 Chloritoid, garnet and rutile micaschist

The garnet core is modelled stable at 0.8 GPa and 780 °C, with a crystallization of ~17.7 vol%. The three garnet rims show similar P-T conditions: 1.9 GPa, 590 °C (resorption of ~1.6 vol% garnet core and crystallization of ~2.8 vol% rim1); 1.8 GPa, 600 °C (resorption of 3.2 vol% core, 1.4 vol% rim1, and crystallization of 5 vol% rim2); then at 2.0 GPa , 600°C (crystallization of 2 vol% rim3). P-T estimates for the garnet rims are in perfect agreement with the intersection of phengite isopleths at Si apfu (3.29-3.33) and X_{Mg} (0.78-0.82; Fig. 4d). The predicted assemblage matches the minerals observed, except that kyanite (3 vol% predicted) was not detected in thin section.

5.3.5 IC - FG1249 Garnet, epidote and rutile micaschist

25

- The garnet core is modelled stable at 0.6 GPa, 730 °C, with crystallization of ~10 vol% garnet. rim1 is modelled stable at 0.6 GPa, 620 °C (resorption of ~2 vol% garnet core and crystallization of ~2 vol% rim1). The second rim is related to the peak pressure recorded by this sample (1.63 GPa, 615 °C with resorption of 0.2 vol% core and 0.05 vol% rim1, plus crystallization of 10 vol% rim2), conditions are in agreement with the Si apfu (3.35-3.4) and X_{Mg} (0.77-0.83; Fig. 4e) compositions of phengite. The last rim recorded may mark the thermal peak at lower pressure (1.56 GPa, 660 °C) with garnet resorption (6.9 vol% rim2) and growth (8.6 vol% rim3).
 - 11

5.3.6 EC - FG1420 Garnet orthogneiss

The garnet core is found stable at 0.48 GPa and 490 °C, rim1 at 0.67 GPa and 500 °C, rim2 at 0.73 GPa and 510 °C and rim3 at 0.8 GPa and 520 °C, with a total amout of 4.8 vol% garnet.

- Two phengite generations are presents (chapter 4.3; Fig. 3f): phengite describing the main foliation, with higher silica 5 content, displays Si apfu and X_{Mg} isoplets itersection at ~1.4 GPa, 550 °C. These conditions are not substantiated by the mineral assemblage, which is predicted to contain omphacite and rutile, but neither phase was observed in thin section. Also, at these P-T conditions further garnet growth is predicted, with a modal increase from 6.5 to >7.5 vol%, but is no garnet is observed with a composition compatible with these P-T conditions. We suspect that phengite grew at lower P-T conditions, as the appropriate Si apfu values intersect the P-T results derived from white mica + quartz + H_2O barometry at 0.6-0.8 GPa, 10
 - The second generation of phengite, post-dating the main foliation, shows Si values for which the isopleths intersect with the results derived from white mica + quartz + H_2O barometry at 0.55-0.75 GPa, 300-350 °C.

5.4 Results: Multi-equilibrium thermobarometry

350-400 °C (Fig. 4f; more details in the next chapter).

5.4.1 IC - FG1315 Garnet, epidote and rutile guartz-micaschist

Chlorite in this sample is retrograde and records formation temperatures decreasing from 450 °C to 300 °C (Fig. 5a). White 15 mica + quartz + H₂O barometry suggest pressures comprised between 1.5 and 0.4 GPa for the temperature range of chlorite. Chlorite and white mica grains in textural equilibrium were used to constrain the equilibrium conditions at 0.8 ± 0.2 GPa and 340 ± 50 °C for the retrograde stage (Fig. 5b; Tables 3 and 4).

5.4.2 IC - FG12157 Garnet, glaucophane, epidote and rutile micaschist

20 Chlorite records formation temperatures decreasing from 430 °C to 310 °C (Fig. 5c). White mica + quartz + H_2O barometry finds pressures between 0.02 and 1 GPa, for the temperature range shown by chlorite. Chlorite and white mica grains in textural equilibrium are used to approximate equilibrium conditions at 0.54 ± 0.2 GPa and 394 ± 50 °C for the retrograde stage (Fig. 5d).

5.4.3 IC - FG1347 Chloritoid, garnet and rutile micaschist

25 Chlorite registers temperatures from 370 °C to 250 °C (Fig. 5e). White mica + quartz + H_2O barometry indicates pressures between 0.02 and 1 GPa for the temperature range of chlorite. Chlorite and white mica grains record 0.78 ± 0.2 GPa and 341 \pm 50 °C for the retrograde stage (Fig. 5f).

5.4.4 EC - FG1420 Garnet orthogneiss

The chlorite and white mica multi-equilibrium technique was used to constrain the equilibrium conditions of three successive stages of retrogression using couples linked to different microstructural positions that developed after the main foliation. These show equilibrium conditions at 0.87 ± 0.2 GPa, 354 ± 50 °C in fold hinges; in pressure shadows at 0.6 ± 0.2 GPa, 331 ± 50 °C; and in static overgrowths over the main foliation at 0.42 ± 0.2 GPa, 231 ± 50 °C (Fig. 6).

6 Texture and geochronology of allanite and zircon

6.1 Methods

6.1.1 Allanite geochronology

- Allanite was first investigated in detail in thin section (see chapter 7), but for more efficient dating purposes, allanite grains
 were separated using high voltage pulsed power disintegration (Selfrag device at University of Bern; e.g. Rudashevsky et al., 1995) followed by magnetic separation and heavy liquids, hand-picked, mounted in acryl/epoxy and polished to equatorial section. The grains were imaged using SEM (BSE), to document the internal texture and compositional zoning. Allanite dating was performed using a LA-ICP-MS Geolas Pro 193 nm ArF excimer laser coupled to an Elan DRC-e ICPMS (Institute of Geological Sciences, University of Bern). The analytical procedure followed is described in detail in Burn et al.
 (2017). In particular, pre-ablation was performed for 4-5 pulses using an energy density of 2.5 J/cm², a repetition rate of 1 Hz and spot sizes of 40 and 32 µm. Ablation was performed using an energy density of 2.5 J/cm², a repetition rate of 9 Hz and
- spot sizes of 32 and 24 μ m. He (1 l/min) and H2 (0.08 l/min) were used as aerosol transport. Instrument setting was optimised for heavy masses and oxide production (ThO⁺/Th⁺) was decreased to be lower than 0.5 %. NIST SRM 610 measurements were performed for quantification of U- and Th-concentrations. Plešovice (Sláma et al., 2008) zircon was
- 20 used as primary standard. Cima d'Asta Pluton (CAP) allanite was used as secondary standard. The acquisition series were approximately 1h to minimise instrumental drift including between 8 and 12 unknowns analyses bracketed by 8 analyses of the primary standard Plešovice used for U-Th-Pb ratio calibration; SRM610 for trace element calibration, as well as 3 analyses of the secondary reference material CAP^b. Data reduction was performed with the in-house program TRINITY (Burn et al., 2017). Data for the secondary standards are reported in the Supplement S9. Uncertainties given in the text and figures
- 25 do not include the systematic uncertainties as defined by Horstwood et al. (2016), because the value appear to be quite variable depending the reference material used (see Burn et al., 2017). In theory, the total uncertainties may be slightly underestimated, whereas the low mean square weighted deviation (MSWD) values rather suggests that the total uncertainties are slightly overestimated.

6.1.2 Zircon geochronology

Individual grains of zircon were separated as described for allanite and investigated by SEM, using cathode luminescence (CL) imaging to document their internal textures. Zircon dates were obtained using the same LA-ICP-MS instrument, following the measurement procedures described by Kunz et al. (2017). Ablation was conducted with an energy density of

- 2.5 J/cm², a repetition rate 9 Hz and spot sizes of 32 or 16 µm. Samples were bracketed by zircon standard GJ-1 (Jackson et 5 al. 2004) and NIST SRM 612 measurements for quantification of U, Th and other trace element concentrations, using ²⁹Si as an internal standard. For accuracy and long-term reproducibility, Plešovice was measured as secondary zircon standard giving a ${}^{206}\text{Pb}/{}^{238}\text{U}$ weighted mean age of 339.2 \pm 1.6 Ma (n=34; error 2σ). Acquisition series took approximately one hour to minimise instrumental drift and were composed of the following: 2 SRM glasses, 3 GJ-1 zircon, 6-8 zircon unknowns, 3-4
- 10 Plešovice zircon, 6-8 zircon unknowns, 3 GJ-1 zircon, 2 SRM glasses. Data reduction was performed with Iolite 2.5 (Paton et al., 2011;Paton et al., 2010) with DRS 'Visual age' (Petrus and Kamber, 2012). All dates reported in this study are concordant within their uncertainty; no common Pb correction was applied.

6.2 Results: Allanite textures and their microstructural relations

In the samples of the IC, allanite prisms are elongate in the eclogite foliation, showing mutual intergrowth relations with

- 15 phengite, garnet, paragonite and rutile, which define this main foliation (Fig. 7a). In samples FG1324, FG1315, and FG1347 allanite crystals are characterized by one main growth zone. Some thin (< 20-30 μ m) allanite rims are observed as well as epidote or clinozoisite rims that appear dark in BSE photos (Fig. 7a, b); where present, these mark a retrograde greenschist overprint. In sample FG12157 and FG1249, BSE pictures show one or more allanite rims characterized by lower brightness (Fig. 7b, c). These rims may reflect minor retrograde stages that weakly altered the eclogite facies assemblage as well.
- 20 Again, a peripheral epidote rim is present. Monazite is occasionally observed as a relic in allanite cores in samples FG1324, FG1315, FG1347, and FG1249. Monazite shows lobate edges and is surrounded by symplectites (µm in size) of allanite and apatite or by discrete crystals of apatite and allanite (Fig. 7d). These features suggest prograde growth of allanite and apatite at the expense of monazite, a common allanite-forming reaction (e.g. Janots et al., 2008).
- Various mineral inclusions are found in allanite grains, as summarized in Table 5. In detail, allanite in sample FG1324 shows 25 intergrowths with garnet (Fig. 7e) suggesting synchronous growth of the two minerals. Phengite inclusions analysed in allanite show the same chemical composition as those marking the main foliation (representative mineral analyses are available in Supplement S⁵). Based on these features and the alignment in the foliation, allanite is interpreted to have grown syn-kinematically in the foliation and at the same time as garnet. In sample FG1315 allanite includes phengite, paragonite, and garnet (Fig. 7a, f); the latter is similar in composition to the Alpine HP-rims and atoll garnet. Phengite inclusions have
- 30

the same composition as phengite marking the main foliation and as phengite included in atoll garnet. Due to the relationships of these mutual inclusions in this sample, allanite growth again appears to be related to the development of the main foliation. In samples FG1347 and FG1249 allanite includes phengite and paragonite, in FG12157 only phengite; these micas have the same composition as those defining the eclogite foliation (Fig. 7b, c, d). In the case of FG1249 and FG1347, allanite also shows intergrowths with both white micas (Fig. 7c).

In the EC, allanite is rare and, where present, is typically magmatic; it appears dark brown and pleochroic in the optical microscope, with a grain size of some millimetres (Fig. 2h; Giuntoli and Engi, 2016). In only two samples metamorphic 5 allanite was found, preserved in the core of epidote crystals (FG1420, FG12107). The metamorphic allanite has a grain size less than 50 µm, is colourless or pale yellow in polarized light, with low interference colour and undulose extinction in crossed polarized light. Sample FG1420 shows both magmatic and metamorphic allanites (Fig. 7g, h). Allanite includes paragonite, with phengite and titanite occurring both in the epidote rim and at the allanite-epidote boundary (Fig. 7g). Very few tiny monazite grains (few µm) are found in the core of metamorphic allanite. Sample FG12107 also shows similar

10

20

epidote crystals preserving metamorphic allanite in their core, as in sample FG1420. Magmatic allanite preserved in sample FG1420 occurs as mm-size grains that are fractured and appear much brighter in BSE pictures than metamorphic allanite (Figs. 2h, 7h). Epidote crystals form satellites around magmatic allanite, suggesting partial breakdown (Fig. 7h). Note that these epidote crystals retain a BSE-bright core of newly grown (Alpine) metamorphic allanite.

6.3 Results: Allanite U-Th-Pb dating

15 For both IC and EC, only allanite cores were successfully dated. The rims showed too high common lead (Pb_c) contents, its correction (Gregory et al., 2007;Burn et al., 2017) would lead to large uncertainties in the age calculation.

In the IC and EC, the Tera-Wasserburg and 232 Th/ 206 Pb_c - 208 Pb/ 206 Pb_c isochron diagrams are concordant, within the range of uncertainty, in all the analyzed samples (Figs. 8. 9; every ellipse is a single spot measurement). Ages range between 77 and 56 Ma for the allanite cores of the IC. In the EC, the magmatic allanite cores in sample FG1420 yield ages of ~290 Ma, the metamorphic cores of 73.7±8.2 Ma. In sample FG12107, the metamorphic allanite cores were dated to 62.8±3.3 Ma. A summary of the Alpine allanite age data from each sample is listed in Table 5.

In detail, the IC Tera-Wasserburg diagrams show 207 Pb/ 206 Pb y-intercepts of 0.84 ± 0.01, 0.823 ± 0.018 and 0.83 ± 0.004 for samples FG1324, FG1315 and FG12157 respectively, with MSWD on the regression comprised between 1.2 and 2.1 (Fig. 8). 206 Pb_c-isochron diagrams display a 208 Pb/ 206 Pb y-intercept of 2.081 ± 0.027, 2.077 ± 0.066 and 2.085 ± 0.028 for samples

25 FG1324, FG1315 and FG12157 respectively, with MSWD on the regression comprised between 0.46 and 0.98. These values are close to the predicted values of Stacey and Kramers (1975) for model lead evolution of this time range (Fig.1 in Burn, 2016). The exception is sample FG1347, in which the Tera-Wasserburg diagram shows a 207 Pb/ 206 Pb y-intercept at 0.787 ± 0.04 (MSWD on the regression of 2.5) and displays a 208 Pb/ 206 Pb y-intercept on the 206 Pb_c-isochron diagram at 1.98 ± 0.082 (MSWD on the regression of 0.4). These values deviate from the values predicted by Stacey and Kramers (1975), probably 30 indicating local inheritance.

In the EC, data for magmatic allanite in sample FG1420 shows 207 Pb/ 206 Pb y-intercept at 0.85 \pm 0.11 in a Tera-Wasserburg diagram and displays a 208 Pb/ 206 Pb y-intercept at 2.09 ± 2.7 in a 206 Pb_c-isochron diagram (Fig. 9). These large uncertainties reflect few (8) spots analyses. Metamorphic allanites show 207 Pb/ 206 Pb y-intercepts of 0.825 ± 0.01 and 0.811 ± 0.014 in

Tera-Wasserburg diagrams (MSWD on the regression of 0.7 and 1.2) and displays 208 Pb/ 206 Pb y-intercepts of 2.061 ± 0.029 and 2.03 ± 0.031 in 206 Pb_c-isochron diagrams (MSWD on the regression of 0.4) for sample FG1420 and FG12107, respectively. These values are close to the predicted values of Stacey and Kramers (1975) for these time ranges.

6.4 Results: Zircon textures

- 5 Internal textures of zircon from the IC reveal complex zoning in CL-images (Fig. 10), showing detrital cores and several phases of resorption and (metamorphic) overgrowth. Zircon cores commonly preserve a variety of internal textures, most commonly oscillatory zoning (Fig. 10e, f) which is typical of zircon grown form melt. Many cores are affected by resorption, obliterating earlier features, but in some grains show sharp boundaries between core and rims, and these preserve features of sediment transport such as broken, rounded or pitted surfaces (Fig. 10b). The number of rims varies between and within
- 10 samples, from two rims in sample FG1257 to a maximum of five in FG1315. Most zircon grains show a first metamorphic rim with a light grey to bright CL-response, followed by a rim with dark CL-appearance. The third rim typically is again medium grey to bright white in CL. In sample FG1315, a forth (dark CL) and fifth (light grey) rim follow, and FG1347 occasionally shows a CL-dark forth rim. The internal textures of the different rims are not always clearly distinguishable, either because of limited width or, in case of very dark or bright CL-response, limited contrast. The second metamorphic rim
- in FG12157 is either uniform or shows fir-tree zoning (Vavra et al., 1996;Root et al., 2004), in sample FG1347 metamorphic rims are often too thin to distinguish or no texture is recognizable in the bright-CL third rim. The first two rims in sample FG1315 have cloudy textures, the third rim with the bright-CL shows no further internal textures but sometimes has inclusions and the two outermost rims commonly are uniform or cloudy in texture. The innermost and outermost metamorphic rims in sample FG1249 and FG1324 are bright and featureless, the second dark rim in sample FG1249 shows sector zoning and many inclusions. The third (medium grey) rim in FG1249 has a cloudy texture.
- Zircon in the External Complex displays pre-Alpine magmatic oscillatory zoning; rarely a bright CL rim, 5-10 µm in width, surrounds the core.

6.5 Results: Zircon U-Pb dating

- The range of Alpine ²⁰⁶Pb/²³⁸U zircon dates for each sample is summarized in Table 5. Details on the pre-Alpine dates are
 available in Kunz et al. (2017). Supplement S10 gives an overview of pre-Alpine dates as well as individual analyses of
 Alpine dates. Concordia plots of individual spots analyses of Alpine zircon are available in Supplement S11. For the five
 samples we obtained 27 analyses of detrital cores, 53 of Permian rims, and 34 of Alpine rims. Detrital zircon cores in
 samples from the IC range from ~793 to 353 Ma and partially overlap with the first rim in sample FG1324 and FG1249 (Fig. 10), for which the dates range from ~450–420 Ma. In most samples, at least two rims are found yielding Carboniferous to
 Triassic dates from ~313 to 222 Ma, as discussed in detail by Kunz et al. (2017); only one sample (FG1324) shows none of
- these. Alpine rims, with a range between 77 and 56 Ma, were measured in all samples except FG1249, where the third rim was too thin $(10\mu m)$ to be analysed. No correlation was found between rim types and ages within samples or amongst them,

and it was not possible to make absolute age distinctions among Alpine rim generations. This is due to relatively large uncertainty in the ages, as the narrow rims only allowed small spots for LA-ICP-MS measurements, decreasing their analytical precision.

Th/U ratios in detrital zircon cores range between 0.15 and >3, the rim generation with dates between 450–420 Ma have low Th/U ratios in sample FG1324 (0.006–0.14) whereas those in sample FG 1249 are between 0.1 and 0.5. The Carboniferous to Triassic rims show Th/U ratios between 0.002–0.4. The Alpine metamorphic rims show low Th/U ratios between 0.001–0.01.

7 Discussion

5

7.1 Linking time constraints with equilibrium conditions (Petrochronology)

- 10 Preliminary analysis of allanite for U-Th-Pb by LA-ICP-MS was performed *in situ* (i.e. using LA-ICPMS spot analyses), on polished thin sections. However, to obtain more material for dating, allanite grains were subsequently separated, mounted on resin mounts and polished. The main reasons why allanite grains were separated for dating are the following:
 - Grain mounts are more efficient as they enable selecting more spots to analyze a single growth zone (at least one order of magnitude more spot analyses per sample than in thin sections).
 - Grains mounts produce more likely equatorial cuts because polishing is optimized to obtain the largest sections possible at a given grain size. This is essential for LA-ICP-MS analysis of grains with several growth zones: if sectioning is not near-equatorial, one is more likely to drill through different growth zones.
 - We do consider petrographic control before dating to be essential: all mounts were imaged at the SEM with the BSE detector to document allanite textures. For each sample, both in the thin section and in the mounts, the allanite grains displayed the
 - 20 same textural patterns, growth zones and minerals inclusions at the sample scale (in separates and thin section). The microtextural features observed in thin sections were readily linked to those seen in grain mounts (compare Figs. 7a and 7f). Furthermore, we note that U-Th-Pb ages from grain mounts analyses agree within uncertainty with the spot analyses made in thin sections.

The fundamental task to reconstruct the P-T-t paths is to establish a strong link between P-T conditions, deriving from

25 thermodynamic modelling, and t data, deriving from age dating. Several criteria were used to achieve this goal, notably: (i) textural evidence, (ii) the presence and composition of distinctive minerals as inclusions, and (iii) the presence of intergrowths of allanite with other phases.

As described in section 6.2, allanite grains of the IC are intergrown with the main mineral phases that describe the eclogite facies foliation (Fig. 7; Table 5). Furthermore, the compositions of mineral inclusions in allanite matched to those found in

30 the matrix and to those predicted by thermodynamic modeling (representative chemical analyses in Supplement S5). Based on these observations (and details given in section 6.2 for each sample), we infer growth of allanite coeval with the minerals marking the eclogite facies foliation in all of the samples analyzed from the IC. This link between age data and P-T

conditions is shown by the red ellipses on the equilibrium phase diagrams shown in Fig. 4 and on the P-T-t paths shown in Fig. 11b and c (colour-coded for each sample). Additionally, in four samples of the IC, the ages obtained for allanite cores and Alpine zircon rims overlap within uncertainty, suggesting coeval growth of these two accessory minerals.

In the EC, metamorphic allanite is rare and preserved only in the core of epidote (section 6.2; Fig. 7 g, h). Epidote is 5 intergrown with white mica, chlorite and titanite, marking the retrograde greenschist overprint, which is constrained by Chlorite+White-Mica+Quartz+H₂O thermobarometry (~0.9 GPa, 350 °C to 0.4 GPa, 230 °C, see Fig. 6). According to these results, metamorphic allanite growth in the EC predates the retrograde greenschist overprint (red ellipse in Fig. 4f and purple ellipse in Fig. 11b-c).

7.2 P-T-t paths of samples

10 7.2.1 Pre-Alpine conditions in the IC

The history of the studied samples is summarized in P-T-t diagrams (Fig. 11) and compared with relevant data from the literature. The first metamorphic stage recorded in our samples from the IC is of Permian age, it reflects granulite facies conditions. Porphyroclastic garnet cores (present in samples FG1315, FG12157, FG1347 and FG1249) are the only major mineral relics of this HT stage. Among accessory minerals, late Carboniferous to Upper Triassic zircon rims (between 313 and 222 Ma) are preserved, and monazite relics in allanite most probably are Paleozoic as well. The ages of the Permian zircon cores fit well with those reflecting HT metamorphism in the Ivrea Zone and Adria-derived units in the Western Alps (Vavra et al., 1999;Vavra et al., 1996;Kunz et al., 2017;Ewing et al., 2013), thus supporting the long held view that the Sesia

Zone is closely related to the Ivrea Zone (e.g. Compagnoni et al., 1977). The single Upper Triassic date (222 ± 13 Ma) in

- sample FG1315 is similar to ages reported from the Ivrea Zone by Vavra et al. (1999), where they have been related to fluid 20 alteration. The P-T conditions for the pre-Alpine HT stage are constrained between 0.6-0.8 GPa and 700-900 °C from the garnet cores of the IC samples. These results confirm earlier data for the IC (0.6-0.9 GPa, ~850 °C; Lardeaux and Spalla, 1991; Rebay and Spalla, 2001; Fig. 11). A retrograde pre-Alpine path through amphibolite facies conditions is evident in sample FG1249, where a first garnet rim generation yields ~0.6 GPa and 600 °C (Fig. 11b), again in fair agreement with previous data (0.3-0.5 GPa, 570-670 °C; Lardeaux and Spalla, 1991; Rebay and Spalla, 2001). A sketch summarizing the chronology and thermodynamic modelling of garnet in sample FG1249 is shown in Giuntoli et al. (in review).
- 25

15

7.2.2 Pre-Alpine conditions in the EC

In the EC, sample FG1420 shows pre-Alpine ages of ~290 Ma for the magmatic allanite grains. These age data probably reflect magmatic crystallization ages of the granitoids; they are in good agreement with several age data of magmatic zircon and monazite for the Sesia-Dent Blanche nappe (Fig. 1 in Bussy et al., 1998; Table 2 and Fig. 5 in Kunz et al., 2017).

7.2.3 Alpine conditions in the IC

For the Alpine history, our samples from the IC indicate higher P and T than in the EC, based on detailed garnet compositional modelling (Lanari et al. 2017). The IC recorded a range of maximum pressures between 1.6 and 2 GPa at temperatures of 600-670 °C. Age constraints based on allanite dating of five samples from the IC fall into three groups:

- Group 1 with ages of ~73 Ma (from 77.2 ± 7.3 Ma to 72.4 ± 1.4 Ma, based on samples FG1324 and FG1249 respectively)
 - Group 2 at ~65 Ma (from 65.4 ± 3.5 Ma to 64.5 ± 4.3 Ma, based on samples FG1315 and FG12157)
 - Group 3 at 55.7 ± 4.5 Ma (sample FG1347).
- Metamorphic conditions for Group 1 show 1.6-1.75 GPa and 580-650 °C for the earliest stage. In Group 2, samples FG1315
 and FG12157 yield not only the same age, within analytical uncertainty, but also similar metamorphic conditions; the pressure difference of 0.1-0.2 GPa is within the uncertainty of the model, as indicated by error bars in Fig. 11b. Notably, the same P-T conditions were derived from garnet rim1 of FG12157 in Group 2 and from garnet rim3 of sample FG1249 in Group 1. Summarizing, Groups 1 and 2 experienced similar P-T conditions but Group 2 attained these some 5-10 Ma later. Furthermore, the outermost garnet rim in sample FG12157 preserves evidence of a retrograde stage at 1.4 GPa and 650 °C.
- Sample FG1347 (Group 3) shows the youngest allanite ages for a similar pressure range as Groups 1 and 2, but lower temperature conditions (~580-600°C). It thus appears that the samples from the IC reflect several stages of allanite growth, probably because rocks of slightly different bulk composition produced allanite by different metamorphic reactions (Engi, 2017). Three growth stages captured in our samples are at ~73 Ma, ~65 Ma, and ~56 Ma. The different P-T-t paths of Groups 1, 2 and 3 are interpreted to represent different tectonic sheets (Giuntoli and Engi, 2016) that experienced similar P-T conditions but at different times (further discussed in section 7.3).
- Regis et al. (2014; Druer slice) and Konrad-Schmolke and Halama (2014) suggested similar P-T paths for different parts of the IC (Fig. 11c), with the highest pressure reached at ~85 Ma. Halama et al. (2014) constrained the development of the retrograde blueschist facies Tallorno Shear Zone (Konrad-Schmolke et al., 2011) at 65.0 \pm 3.0 Ma using Ar-Ar data on phengite. This shear zone was related to external fluid influx occurring approximately at 1.35 GPa and 550 °C (Konrad-
- 25 Schmolke and Halama, 2014). These two P-T paths are similar to Group 3 of the present study, but the latter are up to 30 Ma younger. Groups 1 and 2 consistently display similar P-T paths, but temperatures are 50 100°C higher than those determined by Regis et al. (2014) and Konrad-Schmolke and Halama (2014). In our samples from the IC, we detected no evidence of pressure cycling such as documented by Rubatto et al. (2011) further south, in the Fondo slice (Regis et al., 2014).

30

5

7.2.4 Alpine conditions in the EC

5

In sample FG1420, metamorphic allanites yield ages of 73.7 ± 8.2 Ma associated with metamorphic conditions of 0.8 GPa and $350 - 500^{\circ}$ C, a stage constrained by the growth of garnet and phengite, the mica marking the main foliation. Sample FG12107 yields an allanite growth age of 62.8 ± 3.3 Ma (Fig. 11c highlighted by the star). Chlorite and white mica data from this sample give thermobarometric results in agreement with FG1420. The proximity of the two samples in the field, their similarity in P-T conditions and in textural features of allanite lead us to suggest that allanite grew at the same metamorphic stage in these two samples. Their (nominal) ages seem discrepant, but considering the overlapping (2 σ) uncertainties, we view them as a single age group (Group 4: ~63±3 Ma).

7.2.5 Comparison of P-T-t data for the IC and EC

10 Comparing P-T-t data for the IC and EC, we note that Group 2 (in the IC) and Group 4 (in the EC) recorded the same age data of ca. 65 Ma, but very different metamorphic conditions. This implies that the IC and EC at that time were at completely different structural position in the subduction zone: A difference of 1 GPa translates to a ~30 km vertical offset between the two complexes, assuming lithostatic pressures. Tectonic overpressure cannot be ruled out, but it cannot explain the observed difference in pressures, since Group 4 (at 0.8 GPa) also displays ~150 °C lower temperatures compared to Group 2 (at 1.6-1.8 GPa). We conclude that the IC resided in a substantially deeper and hotter part of the subduction system

than the EC ~65 Ma ago.

The retrograde P-T trajectories in the IC and EC are similar, based on our data for chlorite – white mica equilibria: ~0.8 GPa and 340 °C for samples FG1315, FG1347 (IC) and sample FG1420 (EC; Fig. 11b). Sample FG12157 recorded a low-pressure stage at slightly higher temperature (0.54 GPa and 393 °C) compared to the two other samples of the IC. The EC

- 20 records two further retrograde stages at 0.60 GPa, 330 °C and at 0.42 GPa, 220°C. However, these retrograde stages have yet to be precisely dated. Inger et al. (1996) and Reddy et al. (1999) constrained the greenschist facies overprint of the EC between 46 and 38 Ma (Rb-Sr using phengite). In the light of our data, we interpret these ages to reflect the greenschist facies conditions documented in both complexes (constrained between ~0.8 GPa and 400 °C to 0.4 GPa and 300 °C). The allanite core ages in Group 4 would then be related to an earlier deformation event, as the epidote rim marks the greenschist
- 25 overprint (Figs. 7g, 11c). Zircon fission track ages for the Sesia Zone range from 42 to 26 Ma (Hurford and Hunziker, 1985;Hurford et al., 1989;Hurford et al., 1991;Berger et al., 2012;Wagner and Reimer, 1972;Kapferer, 2010), suggesting that the terrain crossed the ~250°C isotherm in this age range during exhumation. For the most internal section of the Sesia terrain, the overlying Biella Volcanic Suite dated at 32.5 Ma (Kapferer et al., 2012;Kapferer, 2010) provide an age constraint for the final exhumation to the surface.

7.3 Tectonic assembly and exhumation of the Sesia Zone

The IC shows several tectonic sheets, from several hundred meters to a few kilometres in thickness (Giuntoli and Engi, 2016), some of which moved independently in the subduction channel (Rubatto et al., 2011;Regis et al., 2014) at some stages of the evolution. Some of the samples we studied, while taken at most a few kilometres apart in the field, recorded similar P-

- 5 T paths at different times, as reflected by the three age groups identified (Fig 1b-c). This age difference may reflect relative tectonic mobility at eclogite facies conditions between such sheets of basement-derived rocks, which are notoriously difficult to delimit in this terrain (Giuntoli and Engi, 2016). In addition to tectonic mobility, the combined effects of deformation and repeated hydration must be considered as triggers of recrystallization that promoted chemical equilibration to eclogite facies assemblages (including allanite and zircon, as discussed below).
- 10 The differences in the P-T-conditions (at eclogite facies) documented above for the IC samples do not indicate trajectories as tortuous as those suggested by numerical models for ablative subduction (e.g. Stöckhert and Gerya, 2005;Roda et al., 2012). Our P-T-t data as well as the nature and geometry of tectonic boundaries mapped by Giuntoli and Engi (2016) rather indicate progressive accretion at depths of 50-60 km, probably during final subduction-early exhumation stages, juxtaposing a series of continental sheets and forming a coherent complex (the IC Complex; Vitale Brovarone et al., 2013;Regis et al., 2014).
- 15 Eclogite facies conditions in the IC evidently prevailed for an extended period, at least ~15 Ma. In the IC, the main deformation stages and mineral reactions were related to pulses of external fluid passing through the rocks, as reported by Engi et al. (in review) and Giuntoli et al. (in review). Repeated fluid influx occurred at eclogite facies conditions, as shown by resorption and growth features in garnet and zircon, with hydrous fluid pervasively rehydrating rocks that had previously been almost completely dehydrated by Upper Paleozoic metamorphism (Engi et al., in review).
- 20 Thus Permian kinzigites (granulite facies metapelites) were transformed back to micaschists during Alpine subduction. As allanite and zircon ages from these samples are identical within analytical uncertainties, it appears that this fluid influx also triggered the coeval growth of accessory phases at eclogite facies conditions. Later and more localized fluid influx has also been documented at blueschist facies conditions (Konrad-Schmolke et al., 2011), and this probably continued to greenschist facies conditions, as reflected by the partial overprint of the main eclogite assemblages, which is frequently observed in the
- 25 IC. Our samples in the IC indicate higher temperatures than have been previously reported (Fig. 11). This discrepancy may reflect a regional thermal gradient, simply higher temperatures in more external areas (NW) of the IC, but we suspect it to be a direct effect of the (re)hydration of pre-Alpine HT rocks, which is an exothermic process (e.g. Walther and Orville, 1982;Peacock, 1987;Lyubetskaya and Ague, 2009). Walther and Orville (1982) analysed the thermal effects of (de-)hydration reactions during regional metamorphism and found them to be substantial. When applied to the present situation
- 30 of (re-)hydration, heat capacity data indicate that heating the HT Permian protolith requires ~1.0 kJ/K per kg of leucogneiss. The enthalpy released upon partial hydration of this protolith (producing the water content typical of these micaschists, 1.5 wt-% H₂O) adds ~77 kJ/kg in enthalpy. Such hydration should thus result in a temperature increase of some 80 °. This estimate lends credibility to the P-T estimates obtained here, which indeed are 20-80 °C higher than some maximum

temperatures recently reported from other parts of the Sesia Zone, e.g. 575°C (Konrad-Schmolke and Halama, 2014), 570-630°C for the Druer Slice (Regis et al., 2014), or >600°C for the Ivozio Complex (Zucali and Spalla, 2011). In addition, Zrin-rutile data reported by Kunz et al. (2017, their Table 3) gave 640 °C for one of the present samples (FG1249), confirming our results by an entirely independent method.

- 5 A counterclockwise P-T path is proposed here for the EC. This trajectory is based on the results of garnet and chlorite – white mica modeling. The path as shown in Fig. 11 implies that, at nearly isobaric conditions, this area of the EC experienced cooling by 100-150 degrees, possibly related to the entry of cold material into the subduction channel, as already proposed by Pognante (1989) for the southern Sesia Zone. Piemonte-Liguria oceanic units are a likely source of such material, which would be in line with age data of 58-40 Ma for the HP metamorphism in the Zermatt-Saas Zone (e.g.
- 10 Rubatto et al., 1998;de Meyer et al., 2014;Weber et al., 2015) and with tectonic-kinematic models for the evolution of the Sesia–Dent Blanche nappes (Manzotti et al., 2014; Angiboust et al., 2014). Summarizing, the major differences between the eclogite facies conditions recorded in the IC and the epidote blueschist facies condition in the EC are now quantitatively constrained by P-T-t data presented in this study (Fig. 11). Tectonic juxtaposition of these two complexes appears to have occurred during exhumation probably at 0.7-0.9 GPa and 350-400 °C,
- 15 since these are the first joint P-T conditions found in both complexes, around 46-38 Ma (age data from Inger et al., 1996 and Reddy et al., 1999). These data confirm the proposition by Williams and Compagnoni (1983) of a first order tectonic contact (Barmet Shear Zone in Giuntoli and Engi, 2016; Fig. 1b-c) between the two complexes. Our data are the first to quantify the P-T discontinuity at that contact, i.e. ~1 GPa and 100-180 °C. This age range is younger compared to age range proposed in the tectonic model by Angiboust et al. (2014) for the juxtaposition of the two complexes (70-57 Ma, based on data of Babist
- 20 et al., 2006, Konrad-Schmolke et al., 2011, and Halama et al., 2014). Whereas Angiboust et al. (2014) considered the Tallorno-Chiusella Shear Zone (Babist et al., 2006;Konrad-Schmolke et al., 2011) to represent the contact between the IC and EC, more recent mapping (Giuntoli and Engi, 2016) shows that this contact is located several kilometres further NW, in the Barmet Shear Zone.
- The data shown in Fig. 11 allow us to derive average exhumation rates for the IC (Table 6). Using Group 1 and Group 3 as 25 the two anchors, i.e. the oldest and the youngest groups, a first calculation considers Stage1, the interval from the highest pressures recorded in these two groups (~1.6 GPa at ~73 Ma and ~2 GPa at ~55 Ma, respectively) to the greenschist conditions upon juxtaposition with the EC (~0.6 GPa at ~38 Ma, age from Inger et al., 1996). The mean vertical exhumation velocity obtained ranges from 0.9 to 2.7 mm/year respectively (Table 6). If we take the juxtaposition to be at ~46 Ma (Reddy et al., 1999) for the same P conditions, values for the mean vertical exhumation velocity increase to 1.2 and 5.1 mm/year, 30 respectively. Plate convergence between Adria and Europe was ~15 mm/year in the time span of 68-38 Ma (Handy et al.,
- 2010). In the above calculations we considered subduction angles of 90° , 60° and 45° , a thickness of 20 km for the upper crust, 10 km for the lower crust; their densities were taken as 2.7 and 3.0, respectively, and 3.3 g/cm³ for the upper mantle. For Stage2, the final stage of exhumation of the IC and EC, we obtained mean vertical exhumation velocities using the juxtaposition conditions (~0.6 GPa) and ages quoted for the end of Stage1, and 32.5 Ma for the arrival at the surface. That

age of the Biella Volcanic Suite (Kapferer, 2010;Kapferer et al., 2012) is supported by zircon fission track ages (Berger et al., 2012). Depending on the time of juxtaposition adopted, at 38 or 46 Ma (Table 6), a mean vertical exhumation velocity of 4 or 1.6 mm/year results. For Stage2, only the vertical exhumation velocity was computed, and an average density of 2.7 g/cm³ was assumed. The rate of convergence in the 35-20 Ma period was ~13 mm/year (Handy et al., 2010). Data for Stage2

- 5 are taken as an approximate range of exhumation rates, since the last part of the exhumation was not uniform throughout the Sesia Zone; owing to brittle structures local differences may be important (e.g. Berger et al., 2012;Malusà et al., 2006). Furthermore, the values given are minimum exhumation velocities, as the emplacement of the Biella Volcanic Suite may have postdated the arrival of the Sesia Zone at the surface.
- The Stage1 exhumation rates we obtained for the IC agree with exhumation velocities proposed for the same complex by
 Zucali et al. (2002), Regis et al. (2014) and the mean exhumation rates for the Sesia Zone proposed by Roda et al. (2012). However, our rates are up to an order of magnitude lower than those proposed by Rubatto et al. (2011) and the maximum values of Roda et al. (2012). Mean exhumation velocities for the final exhumation stage of the Sesia Zone (Stage2) are in the same range (1.6-4 mm/year) as those for Stage1 (0.9-5.1 mm/year). Our data show no decrease in exhumation velocity from early exhumation (i.e. mantle to deep crustal positions) to late stages (i.e. crustal levels to the surface), such as has been proposed by Rubatto and Hermann (2001) for ultra-high pressure terranes.

7.4 Implication for the subduction, assembly and exhumation of continental fragments

Our work highlights that subducted continental terranes can be composed of several complexes that experienced major differences in their subduction histories (i.e. ~1 GPa in pressure and 100-180 °C in temperature for the Sesia Zone). These complexes can have been tectonically juxtaposed during late metamorphic stages (i.e. greenschist facies conditions) before

- 20 being jointly exhumed to the surface. Complexes can be lithologically heterogeneous and may comprise several tectonometamorphic subunits (from a few hundred metres to several kilometres in thickness). These may experience similar P-T paths but at different times (5-10 Ma apart). Differences in the P-T-t trajectories would thus reflect different tectonic sheets and attest to tectonic mobility in the subduction zone and/or several stages of internal deformation plus hydration at eclogite facies that triggered a pervasive HP-fabric and -assemblage (including datable accessory minerals).
- 25 To unravel such complex histories in subducted continental terranes, carefully established field relations, followed by microstructural and petrochronological analysis need to be combined to
 - map and identify the major and secondary tectono-metamorphic contacts;
 - characterize the different fabrics and the mineral phases defining them;
 - quantify the differences existing in P, T and t for each complex or subunit.
- 30 Finally, the heterogeneity of subducted continental terranes highlighted by this study should be considered when comparing results to numerical and analogue models that aim to investigate the mechanisms responsible for the subduction and exhumation of the continental crust.
 - 23

8 Conclusions

5

15

20

The present paper provides P-T-t data for the central the Sesia Zone, documenting in particular when and at what conditions the two main complexes of this HP-terrain were juxtaposed during the Alpine orogenic cycle.

In particular our data indicate that:

- In the Internal Complex the main stages of mineral growth, and probably the attendant deformation are related to • eclogite facies condition occurring between 77 and 55 Ma and triggered by repeated influx of external fluid. It is during this subduction phase that most of the Permian HT-assemblages were replaced, leaving only sparse relics of essentially dry granulites.
- The Internal Complex encompasses three groups of the samples studied, probably reflecting different tectonic 10 sheets. These experienced similar internal deformation and P-T paths, but at different times, reflecting minor (kmscale) adjustments in the subduction channel at 50-60 km depth and were juxtaposed probably during final subduction-early exhumation stages.
 - Comparing the two main complexes, diverse metamorphic evolutions emerge between 77 and 55 Ma, with conditions of 1.6-2 GPa and 600-670 °C in the Internal Complex, whereas only 0.7-0.9 GPa and ~500°C appear to have been reached in the External Complex.
 - The two complexes were juxtaposed between 46 and 38 Ma (Inger et al., 1996; Reddy et al., 1999) at ~0.8GPa and 350°C, so at mid-crustal levels during exhumation.
 - The different tectonic sheets of the Internal Complex were initially exhumed with mean vertical velocities of 0.9 -5.1 mm/year during Stage1, from the highest pressures recorded to the greenschist facies conditions attained upon juxtaposition with the External Complex.
 - During Stage2, the final exhumation of the entire Sesia Zone occurred with mean vertical velocities between 1.6 and 4 mm/year, but the late stages of exhumation in the area can display local differences (Malusà et al., 2006;Berger et al., 2012).
- This case study shows that subducted HP continental terranes can be composed of various complexes, characterized by 25 major differences in their tectonometamorphic histories, juxtaposed during exhumation by tectonic contacts operating at greenshist facies conditions. Such complexes include several subunits, which in the present case were some hundred metres to few kilometres in thickness, that acted as fragments in the subduction zone, recording slightly different P-T-t paths. The development of the main fabrics at eclogite facies conditions appears to have been triggered by external fluid influx. These processes could be dated based on the coeval growth of allanite and zircon. Petrochronology proved to be a powerful tool to quantify processes and unravel the metamorphic evolution in a complex geological setting, essentially because the detailed

analytical work at the microscale could be linked to solid field evidence.

30

Data availability

Original data underlying the material presented are available by contacting the authors.

Competing interests.

The authors declare that they have no conflict of interest.

5 Acknowledgments

This work was financially supported by the Swiss National Science Foundation (Project 200020-146175). We thank Roberto Compagnoni, Daniele Regis, and Jörg Hermann for fruitful discussions. We acknowledge constructive comments and suggestions from Samuel Angiboust and an anonymous referee, and we are grateful to Patrice Rey for editorial handling.

References

- Angiboust, S., Glodny, J., Oncken, O., and Chopin, C.: In search of transient subduction interfaces in the Dent Blanche–Sesia Tectonic System (W. Alps), Lithos, 205, 298-321, 2014.
 Angiboust, S., Yamato, P., Hertgen, S., Hyppolito, T., Bebout, G., and Morales, L.: Fluid pathways and high pressure metasomatism in a subducted continental slice (Mt. Emilius klippe, W. Alps), J. Metamorph. Geol., 2016.
 Austrheim, H.: Eclogitization of lower crustal granulites by fluid migration through shear zones, Earth Planet. Sci.
- Lett., 81, 221-232, 10.1016/0012-821X(87)90158-0, 1987.
 Babist, J., Handy, M. R., Hammerschmidt, K., and Konrad-Schmolke, M.: Precollisional, multistage exhumation of subducted continental crust: The Sesia Zone, western Alps, Tectonics, 25, <u>10.1029/2005TC001927</u>, 2006.
 Beltrando, M., Compagnoni, R., and Lombardo, B.: (Ultra-) High-pressure metamorphism and orogenesis: An Alpine perspective, Gondwana Res., 18, 147-166, 2010.
- 20 Berger, A., Mercolli, I., Kapferer, N., and Fügenschuh, B.: Single and double exhumation of fault blocks in the internal Sesia-Lanzo Zone and the Ivrea-Verbano Zone (Biella, Italy), Int. J. Earth Sci., 101, 1877-1894, 2012. Berman, R. G.: Internally consistent thermodynamic data for minerals in the system Na₂O-K₂O-CaO-MgO-FeO-Fe₂O₃-Al₂O₃-SiO₂-TiO₂-H₂O-CO₂, J. Petrol., 29, 445--522, 1988. Berman, R. G.: Mixing properties of Ca-Mg-Fe-Mn garnets, Am. Mineral., 75, 328-344, 1990.
- 25 Bertolani, M.: Contributo allo studio petrografico della cosiddetta "formazione dioritico-kinzigitica"; Ricerche in Val Sabbiola (Valsesia), Rendiconti della Società Mineralogica Italiana, 10, 92-207, 1954. Bertolani, M.: La formazione basica "Ivrea-Verbano" e la sua posizione nel quadro geologico-petrografico della bassa Val Sesia e del Biellese, Periodico di Mineralogia, 28, 151-209, 1959. Brun, J.-P., and Faccenna, C.: Exhumation of high-pressure rocks driven by slab rollback, Earth Planet. Sci. Lett.,
- 30 272, 1-7, 2008. Bucher, K., Fazis, Y., Capitani, C. D., and Grapes, R.: Blueschists, eclogites, and decompression assemblages of the Zermatt-Saas ophiolite: High-pressure metamorphism of subducted Tethys lithosphere, Am. Mineral., 90, 821-835, 2005.

Buick, I., and Holland, T.: The PTt path associated with crustal extension, Naxos, Cyclades, Greece, Geological Society, London, Special Publications, 43, 365-369, 1989.

Burn, M.: LA-ICP-QMS Th-U/Pb allanite dating: methods and applications, PhD thesis, University of Bern, Switzerland, 2016.

Burn, M., Lanari, P., Pettke, T., and Engi, M.: Non-matrix-matched standardisation in LA-ICP-MS analysis: General approach and application to allanite Th-U-Pb age-dating., J. Anal. At. Spectrom., 32, 1359-1377, 2017.

5 Bussy, F., Venturini, G., Hunziker, J., and Martinotti, G.: U-Pb ages of magmatic rocks of the western Austroalpine Dent-Blanche-Sesia unit, Schweiz. Mineral. Petrogr. Mitt., 78, 163-168, 1998.

Carraro, F., Dal Piaz, G. V., and Sacchi, R.: Serie di Valpelline e II Zona Diorito-Kinzigitica sono i relitti di un ricoprimento proveniente dalla Zona Ivrea-Verbano, Memorie della Societa Geologica Italiana, 9, 197–224, 1970.

Compagnoni, R., Dal Piaz, G., Hunziker, J., Gosso, G., Lombardo, B., and Williams, P.: The Sesia-Lanzo Zone, a slice
 of continental crust with Alpine high pressure-low temperature assemblages in the Western Italian Alps, Rend. Soc.
 Ital. Mineral. Petrol, 33, 281-334, 1977.

Compagnoni, R., Engi, M., and Regis, D.: Valle d'Aosta section of the Sesia Zone: multi-stage HP metamorphism and assembly of a rifted continental margin, ISPRA, Soc. Geol. Italiana, Geol. Field Trips, 6, 1-44, <u>http://dx.doi.org/10.3301/GFT.2014.02</u>, 2014.

- Dal Piaz, G. V., Gosso, G., and Martinotti, G.: La II Zona Diorito-Kinzigitica tra la Valle Sesia e la Valle d'Ayas (Alpi Occidentali), Memorie della Società Geologica Italiana, 11, 433–460, 1971.
 Dal Piaz, G. V., Hunziker, J. C., and Martinotti, G.: La Zona Sesia-Lanzo e l'evoluzione tettonico-metamorfica delle Alpi nordoccidentali interne, Memorie della Societa Geologica Italiana, 11, 433–466, 1972.
 Dal Piaz, G. V.: The Austroalpine-Piedmont nappe stack and the puzzle of the Alpine Tethys, Mem. Sci. Geol., 51,
- Dai 1122, G. V.. The Austroaphie-Fleamont happe stack and the puzzle of the Alphie Fethys, Went. Sci. Geol., 31,
 155-176, 1999.
 De Capitani, C., and Brown, T. H.: The computation of chemical equilibrium in complex systems containing non-

ideal solutions, Geochim. Cosmochim. Acta, 51, 2639-2652, 1987. de Capitani, C., and Petrakakis, K.: The computation of equilibrium assemblage diagrams with Theriak/Domino software, Am. Mineral., 95, 1006-1016, 10.2138/am.2010.3354, 2010.

de Meyer, C. M., Baumgartner, L. P., Beard, B. L., and Johnson, C. M.: Rb–Sr ages from phengite inclusions in garnets from high pressure rocks of the Swiss Western Alps, Earth Planet. Sci. Lett., 395, 205-216, 2014.
 Dewey, J., Helman, M., Knott, S., Turco, E., and Hutton, D.: Kinematics of the western Mediterranean, Geological Society, London, Special Publications, 45, 265-283, 1989.

Donaldson, D. G., Webb, A. A. G., Menold, C. A., Kylander-Clark, A. R., and Hacker, B. R.: Petrochronology of Himalayan ultrahigh-pressure eclogite, Geology, 41, 835-838, 2013.

- Dubacq, B., Vidal, O., and De Andrade, V.: Dehydration of dioctahedral aluminous phyllosilicates: thermodynamic modelling and implications for thermobarometric estimates, Contrib. Mineral. Petrol., 159, 159, 2010. Engi, M.: Petrochronology based on REE-minerals: monazite, allanite, xenotime, apatite, Rev. Mineral. Geochem., 83, 365-418, 2017.
- Engi, M., Lanari, P., and Kohn, M. J.: Significant Ages—An Introduction to Petrochronology Reviews in Mineralogy & Geochemistry, 83, 1-12, http://dx.doi.org/10.2138/rmg.2017.83.1, 2017.
 Engi, M., Giuntoli, F., Lanari, P., Burn, M., Kunz, B. E., and Bouvier, A.-S.: Pervasive eclogitization due to brittle deformation and rehydration of subducted basement. Effects on continental recycling?, Geochemistry, Geophysics, Geosystems (G³), in review.
- Erambert, M., and Austrheim, H.: The effect of fluid and deformation on zoning and inclusion patterns in polymetamorphic garnets, Contrib. Mineral. Petrol., 115, 204-214, 1993.
 Etheridge, M., Wall, V., and Vernon, R.: The role of the fluid phase during regional metamorphism and deformation, J. Metamorph. Geol., 1, 205-226, 1983.
- Evans, T. P.: A method for calculating effective bulk composition modification due to crystal fractionation in garnetbearing schist; implications for isopleth thermobarometry J. Metamorph. Geol., 22, 547-557, 2004.
- Ewing, T. A., Hermann, J., and Rubatto, D.: The robustness of the Zr-in-rutile and Ti-in-zircon thermometers during high-temperature metamorphism (Ivrea-Verbano Zone, northern Italy), Contr. Mineral. Petrol., 165, 757-779, 2013.
 Faccenda, M., Minelli, G., and Gerya, T.: Coupled and decoupled regimes of continental collision: numerical modeling, Earth Planet. Sci. Lett., 278, 337-349, 2009.

Foster, G., Parrish, R. R., Horstwood, M. S., Chenery, S., Pyle, J., and Gibson, H.: The generation of prograde P–T–t points and paths; a textural, compositional, and chronological study of metamorphic monazite, Earth Planet. Sci. Lett., 228, 125-142, 2004.

- Fuhrman, M. L., and Lindsley, D. H.: Ternary-feldspar modeling and thermometry, Am. Mineral., 73, 201--216, 1988.
 - Gasser, D., Bruand, E., Rubatto, D., and Stüwe, K.: The behaviour of monazite from greenschist facies phyllites to anatectic gneisses: An example from the Chugach Metamorphic Complex, southern Alaska, Lithos, 134, 108-122, 2012.
- Giuntoli, F., and Engi, M.: Internal geometry of the central Sesia Zone (Aosta Valley, Italy): HP tectonic assembly of continental slices, Swiss J. Geosci., 109, 445-471, 2016.
- Giuntoli, F., Lanari, P., and Engi, M.: Deeply subducted continental fragments: I. Fracturing, dissolutionprecipitation and diffusion processes recorded by garnet textures of the central Sesia Zone (Western Italian Alps), Solid Earth, in review.

Gregory, C. J., Rubatto, D., Allen, C. M., Williams, I. S., Hermann, J., and Ireland, T.: Allanite microgeochronology: a LA-ICP-MS and SHRIMP U–Th–Pb study, Chem. Geol., 245, 162-182, 2007.

- Halama, R., Konrad-Schmolke, M., Sudo, M., Marschall, H. R., and Wiedenbeck, M.: Effects of fluid-rock interaction on 40 Ar/39 Ar geochronology in high-pressure rocks (Sesia-Lanzo Zone, Western Alps), Geochim. Cosmochim. Acta, 126, 475-494, 2014.
- Handy, M. R., Schmid, S. M., Bousquet, R., Kissling, E., and Bernoulli, D.: Reconciling plate-tectonic reconstructions
 of Alpine Tethys with the geological-geophysical record of spreading and subduction in the Alps, Earth-Science Reviews, 102, 121-158, 2010.

Horstwood, M. S., Košler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., Pearson, N. J., Sircombe, K., Sylvester, P., and Vermeesch, P.: Community-Derived Standards for LA-ICP-MS U-(Th-) Pb Geochronology– Uncertainty Propagation, Age Interpretation and Data Reporting, Geostand. Geoanal. Res., 40, 311-332, 2016.

- Hunziker, P.: The stability of tri-octahedral Fe2+-Mg-Al chlorite: A combined experimental and theoretical study, Mineralogisch-Petrographisches Institut der Universität, 2003.
 Hurford, A., Flisch, M., and Jäger, E.: Unravelling the thermo-tectonic evolution of the Alps: a contribution from fission track analysis and mica dating, Geological Society, London, Special Publications, 45, 369-398, 1989.
 Hurford, A. J., and Hunziker, J.: Alpine cooling history of the Monte Mucrone eclogites (Sesia-Lanzo zone): fission
- Hurford, A. J., and Hunziker, J.: Alpine cooling history of the Monte Mucrone eclogites (Sesia-Lanzo zone): fission track evidence, Schweiz. Mineral. Petrogr. Mitt., 65, 325-334, 1985.

Hurford, A. J., Hunziker, J. C., and Stöckhert, B.: Constraints on the late thermotectonic evolution of the western Alps: evidence for episodic rapid uplift, Tectonics, 10, 758-769, 1991.

Inger, S., Ramsbotham, W., Cliff, R. A., and Rex, D. C.: Metamorphic evolution of the Sesia-Lanzo Zone, Western Alps: Time constraints from multi-system geochronology, Contrib. Mineral. Petrol., 126, 152-168, 1996.

35 Janots, E., Engi, M., Berger, A., Allaz, J., Schwarz, J.-O., and Spandler, C.: Prograde metamorphic sequence of REEminerals in pelitic rocks of the Central Alps: implications for allanite – monazite – xenotime phase relations from 250 to 610 °C, J. Metamorph. Geol., 26, 509-526, 10.1111/j.1525-1314.2008.00774.x, 2008. Janots, E., Engi, M., Rubatto, D., Berger, A., and Gregory, C.: Metamorphic rates in collisional orogeny from in situ

Janots, E., Engi, M., Rubatto, D., Berger, A., and Gregory, C.: Metamorphic rates in collisional orogeny from in situ allanite and monazite dating, Geology, 37, 11-14, 10.1130/G25192A.1, 2009.

40 Kapferer, N.: The evolution of the surface of the Sesia-Lanzo zone in Oligocene-Miocene times (Biella, NW Italy), PhD thesis, University of Bern, Switzerland, 2010.

Kapferer, N., Mercolli, I., Berger, A., Ovtcharova, M., and Fügenschuh, B.: Dating emplacement and evolution of the orogenic magmatism in the internal Western Alps: 2. The Biella Volcanic Suite, Swiss J. Geosci., 105, 67-84, 2012.

Keller, L. M., De Capitani, C., and Abart, R.: A quaternary solution model for white micas based on natural coexisting phengite-paragonite pairs J. Petrol., 46, doi:10.1093/petrology/egi050, 2005.

- Keppie, D. F., Currie, C. A., and Warren, C.: Subduction erosion modes: comparing finite element numerical models with the geological record, Earth Planet. Sci. Lett., 287, 241-254, 2009.
- Konrad-Schmolke, M., O'Brien, P. J., De Capitani, C., and Carswell, D. A.: Garnet growth at high- and ultra-high pressure conditions and the effect of element fractionation on mineral modes and composition, Lithos, 103, 309-332, 2008.

Konrad-Schmolke, M., O'Brien, P. J., and Zack, T.: Fluid migration above a subducted slab – Constraints on amount, pathways and major element mobility from partially overprinted eclogite-facies rocks (Sesia Zone, Western Alps), J. Petrol., 52, 457-486, 10.1093/petrology/egq087, 2011.

Konrad-Schmolke, M., and Halama, R.: Combined thermodynamic–geochemical modeling in metamorphic geology:
Boron as tracer of fluid–rock interaction, Lithos, 208, 393-414, 2014.

- Kunz, B. E., Manzotti, P., von Niederhäusern, B., Engi, M., Darling, J. R., Giuntoli, F., and Lanari, P.: Permian high-temperature metamorphism in the Western Alps (NW Italy), Int. J. Earth Sci., 1-27, 10.1007/s00531-017-1485-6, 2017.
- Kylander-Clark, A., Hacker, B., and Mattinson, J.: Slow exhumation of UHP terranes: titanite and rutile ages of the
 Western Gneiss Region, Norway, Earth Planet. Sci. Lett., 272, 531-540, 2008.
- Lanari, P.: Micro-cartographie P-T-e dans les roches métamorphiques. Applications aux Alpes et à l'Himalaya, Doctorate, Université de Grenoble, Grenoble, 534 pp., 2012.
- Lanari, P., Guillot, S., Schwartz, S., Vidal, O., Tricart, P., Riel, N., and Beyssac, O.: Diachronous evolution of the alpine continental subduction wedge: evidence from P-T estimates in the Briançonnais Zone houillere (France Western Alps), J. Geodyn., 56-57, 39-54, 2012.
- Lanari, P., Riel, N., Guillot, S., Vidal, O., Schwartz, S., Pêcher, A., and Hattori, K. H.: Deciphering high-pressure metamorphism in collisional context using microprobe mapping methods: Application to the Stak eclogitic massif (northwest Himalaya), Geology, 41, 111-114, 2013.
- Lanari, P., Vidal, O., Lewin, E., Dubacq, B., De Andrade, V., and Schwartz, S.: XMapTools a Matlab[©]-based graphic
 user interface for microprobe quantified image processing, Computers and Geosciences, 62, 227-240, 10.1016/j.cageo.2013.08.010, 2014.
 - Lanari, P., and Engi, M.: Local Bulk Composition Effects on Metamorphic Mineral Assemblages, Reviews in Mineralogy & Geochemistry, 83, 55–102, <u>http://dx.doi.org/10.2138/rmg.2017.83.1</u>, 2017.
- Lanari, P., Giuntoli, F., Loury, C., Burn, M., and Engi, M.: An inverse modeling approach to obtain P-T conditions of
 metamorphic stages involving garnet growth and resorption, Eur. J. Mineral., 29, 181-199, 10.1127/ejm/2017/0029-2597, 2017.

Lardeaux, J. M., and Spalla, M. I.: From granulites to eclogites in the Sesia zone (Italian Western Alps): a record of the opening and closure of the Piedmont ocean, J. metamorphic Geol., 9, 35-59, 1991.

Liati, A., and Gebauer, D.: Constraining the prograde and retrograde PTt path of Eocene H P rocks by SHRIMP 30 dating of different zircon domains: inferred rates of heating, burial, cooling and exhumation for central Rhodope, northern Greece, Contrib. Mineral. Petrol., 135, 340-354, 1999.

Lippitsch, R., Kissling, E., and Ansorge, J.: Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography, Journal of Geophysical Research: Solid Earth, 108, 2003.

- Lyubetskaya, T., and Ague, J.: Modeling the magnitudes and directions of regional metamorphic fluid flow in collisional orogens, J. Petrol., 50, 1505-1531, 2009.
 - Mäder, U. K., and Berman, R. G.: Amphibole thermobarometry: a thermodynamic approach, Geological Survey of Canada, Ottawa, Paper 92-1E, 393-400, 1992.

Mäder, U. K., Percival, J. A., and Berman, R. G.: Thermobarometry of garnet-clinopyroxene-hornblende granulites from the Kapuskasing structural zone, Can. J. Earth Sci., 31, 1134-1145, 1994.

40 Malusà, M., Philippot, P., Zattin, M., and Martin, S.: Late stages of exhumation constrained by structural, fluid inclusion and fission track analyses (Sesia–Lanzo unit, Western European Alps), Earth Planet. Sci. Lett., 243, 565-580, 2006.

Malusà, M., Faccenna, C., Garzanti, E., and Polino, R.: Divergence in subduction zones and exhumation of high pressure rocks (Eocene Western Alps), Earth Planet. Sci. Lett., 310, 21-32, 10.1016/j.epsl.2011.08.002, 2011.

- 45 Manzotti, P., Ballevre, M., Zucali, M., Robyr, M., and Engi, M.: The tectonometamorphic evolution of the Sesia–Dent Blanche nappes (internal Western Alps): review and synthesis, Swiss J. Geosci., 107, 309-336, 2014.
- Meyre, C., De Capitani, C., and Partzsch, J. H.: A ternary solid solution model for omphacite and its application to geothermobarometry of eclogites from the middle Adula Nappe (Central Alps, Switzerland), J. Metamorph. Geol., 15, 687-700, 1997.

Mottram, C. M., Warren, C. J., Regis, D., Roberts, N. M., Harris, N. B., Argles, T. W., and Parrish, R. R.: Developing an inverted Barrovian sequence; insights from monazite petrochronology, Earth Planet. Sci. Lett., 403, 418-431, 2014. Negro, F., Bousquet, R., Vils, F., Pellet, C.-M., and Hänggi-Schaub, J.: Thermal structure and metamorphic evolution of the Piemont-Ligurian metasediments in the northern Western Alps, Swiss J. Geosci., 106, 63-78, 2013.

5 Oliver, N.: Review and classification of structural controls on fluid flow during regional metamorphism, J. Metamorph. Geol., 14, 477-492, 1996.

Passchier, C. W., Urai, J. L., van Loon, J., and Williams, P. F.: Structural geology of the central Sesia Lanzo Zone, Geol. Mijnbouw, 60, 497–507, 1981.

Paton, C., Woodhead, J. D., Hellstrom, J. C., Hergt, J. M., Greig, A., and Maas, R.: Improved laser ablation U-Pb zircon geochronology through robust downhole fractionation correction, Geochem. Geophys. Geosyst., 11, 2010.

Paton, C., Hellstrom, J., Paul, B., Woodhead, J., and Hergt, J.: Iolite: Freeware for the visualisation and processing of mass spectrometric data, J. Anal. At. Spectrom., 26, 2508-2518, 2011.

Peacock, S. M.: Thermal effects of metamorphic fluids in subduction zones, Geology, 15, 1057-1060, 1987.

- Pennacchioni, G.: Progressive eclogitization under fluid-present conditions of pre-Alpine mafic granulites in the Austroalpine Mt Emilius Klippe (Italian western Alps), J. Struct. Geol., 18, 549-561, 1996.
- Petrus, J. A., and Kamber, B. S.: VizualAge: A novel approach to laser ablation ICP-MS U-Pb geochronology data reduction, Geostand. Geoanal. Res., 36, 247-270, 2012. Pognante, U.: Tectonic implications of lawsonite formation in the Sesia zone (western Alps), Tectonophysics, 162, 219-
- Pognante, U.: Tectonic implications of lawsonite formation in the Sesia zone (western Alps), Tectonophysics, 162, 219-227, 1989.
- Quick, J. E., Sinigoi, S., Snoke, A. W., Kalakay, T. J., Mayer, A., and Peressini, G.: Geologic map of the southern Ivrea-Verbano Zone, northwestern Italy, US Geological Survey, 2003.
 Rebay, G., and Spalla, M. I.: Emplacement at granulite facies conditions of the Sesia–Lanzo metagabbros: an early record of Permian riftig?, Lithos, 58, 85–104, 2001.
 Reddy, S. Wheeler, L. and Cliff, B.: The geometry and timing of orogenic extension: an example from the Western
- Reddy, S., Wheeler, J., and Cliff, R.: The geometry and timing of orogenic extension: an example from the Western
 Italian Alps, J. Metamorph. Geol., 17, 573-590, 1999.
- Regis, D., Rubatto, D., Darling, J., Cenki-Tok, B., Zucali, M., and Engi, M.: Multiple metamorphic stages within an eclogite-facies terrane (Sesia Zone, Western Alps) revealed by Th-U-Pb petrochronology, J. Petrol., 55, 1429-1456, 10.1093/petrology/egu029, 2014.
- Replumaz, A., Negredo, A. M., Guillot, S., and Villaseñor, A.: Multiple episodes of continental subduction during
 India/Asia convergence: Insight from seismic tomography and tectonic reconstruction, Tectonophysics, 483, 125-134, 2010.

Rivalenti, G., Garuti, G., and Rossi, A.: The origin of the Ivrea-Verbano basic formation (western Italian Alps); whole rock geochemistry, Boll. Soc. Geol. Ital., 94, 1149-1186, 1975.

Roda, M., Spalla, M. I., and Marotta, A. M.: Integration of natural data within a numerical model of ablative
 subduction: a possible interpretation for the Alpine dynamics of the Austroalpine crust, J. Metamorph. Geol., 30, 973-996, 2012.

Root, D. B., Hacker, B., Mattinson, J., and Wooden, J. L.: Zircon geochronology and ca. 400 Ma exhumation of Norwegian ultrahigh-pressure rocks: an ion microprobe and chemical abrasion study, Earth Planet. Sci. Lett., 228, 325-341, 2004.

- Rosenbaum, G., Lister, G. S., and Duboz, C.: Relative motions of Africa, Iberia and Europe during Alpine orogeny, Tectonophysics, 359, 117-129, 2002.
 Rubatto, D., Gebauer, D., and Fanning, M.: Jurassic formation and Eocene subduction of the Zermatt-Saas-Fee
 - ophiolites: implications for the geodynamic evolution of the Central and Western Alps, Contrib. Mineral. Petrol., 132, 269-287, 1998.
- 45 Rubatto, D., and Hermann, J.: Exhumation as fast as subduction?, Geology, 29, 3-6, 2001. Rubatto, D., Regis, D., Hermann, J., Boston, K., Engi, M., Beltrando, M., and McAlpine, S.: Yo-Yo subduction recorded by accessory minerals (Sesia Zone, Western Alps) Nat. Geosci., 4, 338-342, 10.1038/ngeo1124, 2011. Rubie, D. C.: The catalysis of mineral reactions by water and restrictions on the presence of aqueous fluid during metamorphism. Mineral. Mag., 50, 399-415, 1986.

Rudashevsky, N., Burakov, B., Lupal, S., and Thalhammer, O.: Liberation of accessory minerals from various rock types by electric-pulse disintegration-method and application, Transactions of the Institution of Mining and Metallurgy. Section C. Mineral Processing and Extractive Metallurgy, 104, 1995.

Schenk, V.: PTt path of the lower crust in the Hercynian fold belt of southern Calabria, Geological Society, London,
Special Publications, 43, 337-342, 1989.

Schmid, S. M., Kissling, E., Diehl, T., van Hinsbergen, D. J., and Molli, G.: Ivrea mantle wedge, arc of the Western Alps, and kinematic evolution of the Alps–Apennines orogenic system, Swiss J. Geosci., 1-32, 2017.

Scott, D. J., and St-Onge, M. R.: Constraints on Pb closure temperature in titanite based on rocks from the Ungava orogen, Canada: Implications for U-Pb geochronology and PTt path determinations, Geology, 23, 1123-1126, 1995.

10 Sláma, J., Košler, J., Condon, D. J., Crowley, J. L., Gerdes, A., Hanchar, J. M., Horstwood, M. S., Morris, G. A., Nasdala, L., and Norberg, N.: Plešovice zircon—a new natural reference material for U–Pb and Hf isotopic microanalysis, Chem. Geol., 249, 1-35, 2008. Stacey, J. t., and Kramers, J.: Approximation of terrestrial lead isotope evolution by a two-stage model, Earth Planet.

Statey, J. t., and Kramers, J.: Approximation of terrestrial lead isotope evolution by a two-stage model, Earth France.
 Sci. Lett., 26, 207-221, 1975.
 Stöckhert, B., and Gerya, T. V.: Pre-collisional high pressure metamorphism and nappe tectonics at active

Stockhert, B., and Gerya, T. V.: Fre-consional high pressure metanorphism and happe tectomes at active continental margins: a numerical simulation, Terra Nova, 17, 102-110, 2005.
 Tao, W. C., and O'Connell, R. J.: Ablative subduction: A two-sided alternative to the conventional subduction model, Journal of Geophysical Research: Solid Earth, 97, 8877-8904, 1992.
 Voure, C., Cabouer, D., Schmid, B., and Commuter, W.: Multiple gingen growth and non-stabilization during

- Vavra, G., Gebauer, D., Schmid, R., and Compston, W.: Multiple zircon growth and recrystallization during
 polyphase Late Carboniferous to Triassic metamorphism in granulites of the Ivrea Zone (Southern Alps): An ion
 microprobe (SHRIMP) study, Contrib. Mineral. Petrol., 122, 337-358, 1996.
 Vavra, G., Schmid, R., and Gebauer, D.: Internal morphology, habit and U-Th-Pb microanalysis of amphibolite-to granulite facies zircons: geochronology of the Ivrea Zone (Southern Alps), Contrib. Mineral. Petrol., 134, 380-404,
- 1999.
 Vidal, O., Parra, T., and Vieillard, P.: Thermodynamic properties of the Tschermak solid solution in Fe-chlorite: Application to natural examples and possible role of oxidation, Am. Mineral., 90, 347-358, 2005. Vidal, O., De Andrade, V., Lewin, E., Munoz, M., Parra, T., and Pascarelli, S.: P–T-deformation-Fe3+/Fe2+ mapping

at the thin section scale and comparison with XANES mapping: application to a garnet-bearing metapelite from the Sambagawa metamorphic belt (Japan), J. Metamorph. Geol., 24, 669-683, 2006.

- 30 Vidal, O., Lanari, P., Munoz, M., Bourdelle, F., and De Andrade, V.: Deciphering temperature, pressure and oxygenactivity conditions of chlorite formation, Clay Minerals, 51, 615-633, 2016. Vitale Brovarone, A., Beyssac, O., Malavieille, J., Molli, G., Beltrando, M., and Compagnoni, R.: Stacking and metamorphism of continuous segments of subducted lithosphere in a high-pressure wedge: the example of Alpine Corsica (France), Earth-Science Reviews, 116, 35-56, 2013.
- Wagner, G., and Reimer, G.: Fission track tectonics: the tectonic interpretation of fission track apatite ages, Earth Planet. Sci. Lett., 14, 263-268, 1972.
 Walther, J. V., and Orville, P. M.: Volatile production and transport in regional metamorphism, Contrib. Mineral. Petrol., 79, 252-257, 1982.
- Warren, C. J., Beaumont, C., and Jamieson, R. A.: Formation and exhumation of ultra-high-pressure rocks during
 continental collision: Role of detachment in the subduction channel, Geochem. Geophys. Geosyst., 9,
 doi:10.1029/2007GC001839, 2008.

Weber, S., Sandmann, S., Miladinova, I., Fonseca, R. O., Froitzheim, N., Münker, C., and Bucher, K.: Dating the initiation of Piemonte-Liguria Ocean subduction: Lu–Hf garnet chronometry of eclogites from the Theodul Glacier Unit (Zermatt-Saas zone, Switzerland), Swiss J. Geosci., 108, 183-199, 2015.

45 Wheeler, J., and Butler, R. W. H.: Evidence for extension in the western Alpine Orogen: the contact between the oceanic Piemonte and overlying continental Sesia units, Earth Planet. Sci. Lett., 117, 457-474, 1993. Whitney, D. L., and Evans, B. W.: Abbreviations for names of rock-forming minerals, Am. Mineral., 95, 185-187, 2010.

Williams, P., and Compagnoni, R.: Deformation and metamorphism in the Bard area of the Sesia Lanzo Zone,
Western Alps, during subduction and uplift, J. Metamorph. Geol., 1, 117-140, 1983.

Yamato, P., Burov, E., Agard, P., Le Pourhiet, L., and Jolivet, L.: HP-UHP exhumation during slow continental subduction: Self-consistent thermodynamically and thermomechanically coupled model with application to the Western Alps, Earth Planet. Sci. Lett., 271, 63-74, 2008.

Zhao, L., Paul, A., Guillot, S., Solarino, S., Malusà, M. G., Zheng, T., Aubert, C., Salimbeni, S., Dumont, T., and
Schwartz, S.: First seismic evidence for continental subduction beneath the Western Alps, Geology, 43, 815-818, 2015.
Zingg, A.: The Ivrea and Strona Ceneri Zones (Southern Alps, Ticino and Northern Italy): A review., Schweiz.
Mineral. Petrogr. Mitt., 63, 361-392, 1983.

Zucali, M., Spalla, M. I., and Gosso, G.: Strain partitioning and fabric evolution as a correlation tool: the example of the Eclogitic Micaschists Complex in the Sesia–Lanzo Zone (Monte Mucrone–Monte Mars, Western Alps, Italy), Schweiz. Mineral. Petrogr. Mitt., 82, 429–454, 2002.

Zucali, M., and Spalla, M. I.: Prograde lawsonite during the flow of continental crust in the Alpine subduction: Strain vs. metamorphism partitioning, a field-analysis approach to infer tectonometamorphic evolutions (Sesia-Lanzo Zone, Western Italian Alps), J. Struct. Geol., 33, 381-398, 2011.

10



Figure 1: (a) Simplified tectonic map of the Western Alps (modified from Manzotti et al., 2014). (b) Tectonic sketch of the study area (modified from Giuntoli and Engi, 2016) with sample locations and P-T-t data (this study). (c) Cross section through the study area (location show in a) with projection of the studied samples. Foliation traces: violet indicates the eclogite facies foliation (S2) of the IC, dark green indicates the composite epidote blueschist-HP greenschist facies foliation (S2+S3) of the EC, dark green indicates the greenschist facies mylonitic foliation (S4) at the contact IC-EC; BSZ Barmet Shear Zone, PLO Piemonte-Liguria Oceanic unit (modified from Giuntoli and Engi, 2016).



Figure 2: Thin section photos illustrating the metamorphic evolution of the studied samples. End-member mineral abbreviations used throughout text and figures are from Whitney and Evans (2010). (a) Eclogite foliation (S_{ecl}) marked by subparallel phengite, omphacite and glaucophane (plane-polarized light). (b) Eclogite foliation marked by the preferred orientation of chloritoid and phengite. Note the chloritoid crystal (left) is oriented perpendicular to S_{ecl} and overgrows it (plane-polarized light). (c) Garnet

5

phengite. Note the chloritoid crystal (left) is oriented perpendicular to S_{ecl} and overgrows it (plane-polarized light). (c) Garnet crystal with a bright pre-Alpine core and darker Alpine rims visible due to fine rutile inclusions (plane-polarized light). (d) Glaucophane crystal with a core displaying pale blue absorption colour, rimmed by darker blue crossite that marks a retrograde blueschist stage (plane-polarized light). (e) Colorless to pale green amphibole growing at the expense of omphacite during a retrograde greenschist stage (plane-polarized light). (f) Open folds with chlorite crystallizing in the hinge zone, marking a retrograde greenschist stage (plane-polarized light). (g) Garnet porphyroclast wrapped by the main greenschist foliation (S_{gr}); note the inner foliation inside the garnet (S_{int}; crossed-polarized light). (h) Brown magmatic allanite surrounded by finer grain epidote

10

crystals (plane-polarized light).



Figure 3: (a) Mineral phases in sample FG1249 (compare with Fig. 2c) . (b) X_{Grs} map highlights the porphyroclastic cores showing fractures sealed by higher X_{Grs} garnet and three rims (Further garnet end-members maps for the samples are available in Giuntoli et al. submitted). (c) Mineral phases in sample FG1420 (compare with Fig. 2g). (d) Garnet displaying concentric zoning related to a decrease in X_{Sps} . (Further garnet end-members maps for the samples are available in the Supplement S2). (e) Phengite groups in sample FG1249 (details in text). (f) Phengite groups in sample FG1420 (details in text). Note larger phengite flakes of Group2 localized in fold hinges.





Figure 4: Equilibrium phase diagrams of the studied samples computed with Theriak/Domino, assuming a free H₂O fluid, between 1.3-2 GPa and 400-700°C for (a), (b), (c), (d), (e) and between 0.3-1.5 GPa and 200-600°C for (f). The error bars departing from filled ellipses show the P-T uncertainty related to the analytical error of the garnet composition.



Figure 5: Multi-equilibrium thermobarometry results for the IC samples. (a), (c), and (e): intersection of the results deriving from *Chlorite+Quartz+H*₂O thermometry with White-Mica+Quartz+H₂O barometry. (b), (d), and (f): results of *Chlorite+White-Mica+Quartz+H*₂O thermobarometry.



Figure 6: Chlorite+White-Mica+Quartz+H₂O thermobarometry results for the EC sample.



Figure 7: Back-scattered electron photos illustrating some of the allanite crystals in the studied samples (a-h: thin sections; b-g: grain mounts). (a) Allanite grain elongate in the eclogite foliation with phengite and paragonite. It is rimmed by epidote and a simplectite of paragonite and chlorite. (b) Allanite displaying several growth zones and a dark outermost rim of epidote. Note the phengite inclusion and laser ablation pits (32 µm). (c) Allanite displaying several growth zones, the innermost of which are intergrown with paragonite. The tiny dark inclusions are fine grained phengite and paragonite. (d) Monazite with lobate edges preserved at the core of an allanite grain. (e) Allanite intergrown with garnet and wrapped by a darker epidote rim. (f) Allanite including garnet, rutile and phengite. (g) Allanite preserved at the core of epidote. Inclusions of paragonite and phengite are present at the boundary allanite-epidote rim. (h) Magmatic allanite with epidote crystals as satellites; some of the latter display cores of metamorphic allanite.





Figure 8: Results of the IC samples plotted in the Tera-Wasserburg and 232 Th/ 206 Pb_c - 208 Pb/ 206 Pb_c isochron diagram with intercept and isochron age respectively and initial common lead composition estimations.



Figure 9: Results of the EC samples plotted in the Tera-Wasserburg and 232 Th/ 206 Pb_c - 208 Pb/ 206 Pb_c isochron diagram with intercept and isochron age respectively and estimated initial common lead composition.



Figure 10: CL-images of zircon textures in the samples from the Internal Complex. Typically detrital cores show more or less resorption and are followed up by one to five metamorphic rim generations of different CL responses. The scale bar in all images is 50 µm. Solid circles correspond to 32µm LA-ICP-MS spots, while dashed circles are 16µm spot sizes. The dates are individual ²⁰⁶Pb/²³⁸U spots analysis given in Ma.







Figure 11: P-T-t paths of the study area (Fig. 1). (a) Data and paths from previous studies: proposed pre-Alpine evolution from the data of Lardeaux and Spalla (1991), and Rebay and Spalla (2001); Alpine evolution of the Internal Complex from Rubatto et al. (2011), Regis et al. (2014), Konrad-Schmolke and Halama (2014), and Halama et al. (2014); ages for the External Complex from Inger et al. (1996) and Reddy et al. (1999), related to greenschist facies conditions. (b) Data presented in this study. Ellipses show

5

P-T constraints for the age data, rectangles refer to retrograde stages. Error bars show the P-T uncertainty related to the analytical error of the garnet composition. (c) Interpreted P-T-t paths. Full lines = Internal Complex; dashed lines = External Complex. The star indicates the age data from sample FG12107 (see text for explanation).

Sample	e IC-FG1324		4	IC-FG1315			IC-FG12157			IC-FG1347			IC-FG1249				EC-FG1420						
Grt				CORE	RIM 1	RIM 2	RIM 3	CORE	RIM 1	RIM 2	CORE	RIM 1	RIM 2	RIM 3	RIM 4	CORE	RIM 1	RIM 2	RIM 3	CORE	RIM 1	RIM 2	RIM 3
	Spot a	nalyses (wt%)	Ave	rage comp	position (v	wt%)	Average composition (wt%)				Average composition (wt%)			Average composition (wt%)			Average composition (wt%)					
SiO2	37.75	37.93	37.74	37.70	37.96	37.81	37.44	38.17	38.43	38.74	37.75	37.88	38.18	37.75	37.33	36.91	36.62	37.80	37.98	38.29	37.57	37.77	37.61
TiO2	0	0.01	0.04	0.01	0.01	0.01	0.01	0.07	0.07	0.17	0.02	0.02	0.02	0.03	0.03	0.05	0.16	0.27	0.04	0.1	0.1	0.1	0.09
A12O3	21.59	21.62	21.53	21.27	21.44	21.46	21.32	21.07	21.20	21.24	21.25	21.28	21.09	21.21	21.33	20.83	21.36	21.18	21.77	21.04	21.3	21.11	21.22
FeO	31.16	31.12	31.77	31.46	27.94	30.03	32.07	31.23	29.05	26.83	31.30	30.19	30.82	31.96	32.07	33.06	34.23	28.79	27.12	23.98	25.57	27.04	28.47
MnO	0.65	0.63	0.60	0.61	0.41	0.42	0.31	1.00	0.57	0.56	0.92	0.37	0.39	0.35	0.32	2.31	0.37	0.66	0.56	3.52	1.93	0.76	0.37
MgO	4.5	4.73	4.35	7.19	5.33	6.18	6.30	6.41	5.08	4.22	7.05	5.76	6.60	6.62	7.30	4.65	3.86	5.18	4.86	0.58	0.63	0.73	0.94
CaO	4.1	3.94	3.87	1.31	6.72	4.17	2.08	1.38	5.59	8.39	1.14	3.96	2.13	1.49	0.96	1.77	3.27	6.35	8.38	12.45	12.83	12.47	11.36
Cr2O3	0	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.04	0.04	0.00	0.00	0.00	0.00	0.00	0.03	0.04	0.03	0.03	0	0	0	0
Total	99.75	99.97	99.90	99.56	99.81	100.08	99.52	99.37	100.03	100.18	99.41	99.47	99.24	99.41	99.33	99.62	99.90	100.27	100.73	99.95	99.93	99.98	100.06
Formulae l	based on 1	2 oxygei	ns																				
Si	2.99	3.00	2.99	2.97	2.97	2.96	2.96	3.02	3.02	3.03	2.98	2.99	3.02	2.98	2.94	2.95	2.93	2.96	2.95	3.05	2.99	3.01	3.00
Ti	0	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.02	0.00	0.01	0.01	0.01	0.01
Al	2.02	2.01	2.01	1.97	1.98	1.98	1.99	1.97	1.96	1.96	1.97	1.98	1.96	1.98	1.98	1.96	2.01	1.96	1.99	1.98	2.00	1.98	1.99
Fe	2.07	2.06	2.12	2.07	1.83	1.96	2.12	2.07	1.91	1.76	2.06	1.99	2.04	2.11	2.11	2.21	2.29	1.89	1.76	1.60	1.70	1.80	1.90
Mn	0.04	0.04	0.05	0.04	0.03	0.03	0.02	0.07	0.04	0.04	0.06	0.03	0.03	0.02	0.02	0.16	0.03	0.04	0.04	0.24	0.13	0.05	0.03
Mg	0.53	0.56	0.49	0.84	0.62	0.72	0.74	0.76	0.59	0.49	0.83	0.68	0.78	0.78	0.86	0.56	0.46	0.61	0.56	0.07	0.08	0.09	0.11
Ca	0.35	0.33	0.33	0.11	0.56	0.35	0.18	0.12	0.47	0.70	0.10	0.34	0.18	0.13	0.08	0.15	0.28	0.53	0.70	1.06	1.09	1.06	0.97
\sum cations	8.00	8.00	7.99	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00	8.00
Molecular proportions of garnet end members																							
Alm	0.69	0.69	0.70	0.68	0.60	0.64	0.69	0.69	0.63	0.59	0.68	0.66	0.67	0.69	0.69	0.72	0.75	0.62	0.58	0.54	0.57	0.60	0.63
Prp	0.18	0.19	0.17	0.28	0.20	0.24	0.24	0.25	0.20	0.16	0.27	0.22	0.26	0.26	0.28	0.18	0.15	0.20	0.18	0.02	0.03	0.03	0.04
Grs	0.12	0.11	0.11	0.04	0.19	0.11	0.06	0.04	0.16	0.24	0.03	0.11	0.06	0.04	0.03	0.05	0.09	0.17	0.23	0.36	0.37	0.35	0.32
Sps	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.05	0.01	0.01	0.01	0.08	0.04	0.02	0.01

Table 1: Representative analyses of garnet of the samples

Sample	IC-FG1324		IC-FG1315		IC-FG12157		IC-FG1347		IC-FG1249		EC-FG1420				
Ph												Average		Average	
	Spot a	nalysis	(wt%)	Average composition		Average composition		Average composition		Average composition		comp. (wt%)		(wt%)	
wt%				(wt%)	σ	(wt%)	σ	(wt%)	σ	(wt%)	σ	High Si	σ	Low Si	σ
SiO2	51.34	49.24	51.99	50.89	0.55	49.61	0.58	49.43	0.62	49.57	0.79	50.82	0.87	48.54	0.86
TiO2	0.41	0.53	0.53	0.26	0.15	0.31	0.06	0.23	0.05	0.24	0.08	0.15	0.04	0.13	0.03
Al2O3	26.67	26.82	26.83	28.15	0.40	27.58	0.35	29.78	0.50	27.92	0.52	27.45	0.69	30.17	0.76
FeO	1.67	1.39	1.64	1.49	0.11	2.09	0.43	1.40	0.14	1.30	0.42	3.21	0.25	2.63	0.28
MnO	0.00	0.00	0.00	0.06	0.01	0.19	0.04	0.06	0.01	0.05	0.01	0.08	0.01	0.08	0.01
MgO	3.00	3.37	3.45	3.68	0.14	3.65	0.15	3.29	0.16	3.45	0.18	2.85	0.25	2.30	0.21
CaO	0.00	0.00	0.00	0.00	0.00	0.19	0.04	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.01
Na2O	0.62	0.86	0.74	0.77	0.09	0.66	0.08	0.71	0.11	0.87	0.14	0.44	0.11	0.32	0.08
K2O	10.16	10.14	10.33	10.24	0.17	9.97	0.19	10.35	0.21	9.97	0.26	10.56	0.26	10.77	0.25
Total	93.87	92.35	95.51	95.54	-	94.26	-	95.25	-	93.37	-	95.56	-	94.93	-
Formulae	based o	on 11 ox	vygens												
Si	3.46	3.38	3.44	3.37	0.02	3.35	0.02	3.29	0.03	3.36	0.03	3.40	0.04	3.27	0.04
Ti	0.02	0.03	0.03	0.01	0.00	0.02	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00
Al	2.12	2.17	2.09	2.20	0.01	2.19	0.03	2.34	0.03	2.23	0.04	2.16	0.05	2.39	0.05
Fe	0.09	0.08	0.09	0.08	0.00	0.12	0.02	0.08	0.01	0.07	0.02	0.18	0.01	0.15	0.02
Mn	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00
Mg	0.30	0.35	0.34	0.36	0.01	0.37	0.02	0.33	0.02	0.35	0.02	0.28	0.03	0.23	0.02
Ca	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Na	0.08	0.11	0.10	0.10	0.01	0.09	0.01	0.09	0.01	0.11	0.02	0.06	0.01	0.04	0.01
Κ	0.87	0.89	0.87	0.87	0.01	0.86	0.02	0.88	0.02	0.86	0.02	0.90	0.02	0.93	0.02
\sum cations	6.94	7.01	6.97	7.00	-	7.01	-	7.02	-	7.00	-	6.99	-	7.02	-
XMg	0.76	0.81	0.79	0.82	0.012	0.76	0.04	0.81	0.02	0.83	0.05	0.61	0.03	0.61	0.03

12 Table 2: Representative analyses of phengite of the samples. σ is the standard deviation associated to the average composition analysis

Sample	ample IC-FG1315		IC-FG	F12157	IC-FO	G 1347	EC-FG1420			
Chl							Grp1	Grp2	Grp3	
	Chl-35093	Chl-35092	Chl-204162	Chl-204794	Chl-214479	Chl-215870	Chl-910650	Chl-547806	Chl-881538	
SiO2	25.64	26.22	25.81	25.91	24.91	25.62	25.11	25.91	25.45	
A12O3	19.62	19.71	20.64	20.1	21.7	20.97	20.09	21.63	20.34	
FeO	26.29	26.68	24.53	27.34	24.96	26.45	26.87	28.98	25.53	
MnO	0.05	0.05	0.21	0.33	0.11	0.19	0.28	0.31	0.27	
MgO	15.32	15.82	15.87	14.32	13.51	13.49	13.01	12.61	12.61	
CaO	0.05	0.06	0.03	0.03	0.13	0.09	0.08	0.09	0.17	
Na2O	0.05	0.03	0.04	0.03	0	0.03	0.02	0.03	0.03	
K2O	0.07	0.06	0.02	0.03	0.15	0.22	0.03	0.03	0.06	
Total	87.09	88.63	87.15	88.09	85.47	87.06	85.49	89.59	84.46	
Atom site dis	tribution (14	anhydrous-o	xygen basis iı	ncluding Fe3-	+)					
Si(T1+T2)	2.68	2.71	2.70	2.69	2.65	2.72	2.73	2.70	2.78	
Al(T2)	1.32	1.29	1.30	1.31	1.34	1.28	1.27	1.30	1.22	
Al(M1)	0.32	0.29	0.30	0.31	0.34	0.28	0.27	0.30	0.22	
Mg(M1)	0.28	0.33	0.35	0.27	0.24	0.31	0.30	0.27	0.30	
Fe2+(M1)	0.20	0.26	0.27	0.21	0.21	0.32	0.33	0.32	0.32	
V(M1)	0.20	0.12	0.09	0.21	0.21	0.10	0.10	0.11	0.16	
Mg(M2+M3)	2.10	2.10	2.13	1.95	1.90	1.83	1.80	1.69	1.75	
Fe(M2+M3)	1.46	1.61	1.64	1.57	1.62	1.87	1.94	2.03	1.85	
Al(M2+M3)	0.41	0.26	0.19	0.44	0.43	0.24	0.21	0.24	0.34	
Al(M4)	0.36	0.56	0.76	0.41	0.60	0.84	0.83	0.82	0.84	
XMg	0.59	0.57	0.56	0.55	0.54	0.49	0.48	0.45	0.49	
Fe3+(M4)	0.64	0.44	0.24	0.59	0.40	0.16	0.17	0.18	0.16	

13 Table 3: Representative analyses of chlorite used in the multi-equilibrium calculations

Sample	Sample IC-FG1315		IC-FO	G12157	IC-F	G1347	EC-FG1420			
Wm							Grp1	Grp2	Grp3	
	Wm-	Wm-	Wm-	Wm-	Wm-	Wm-	Wm-	Wm-	Wm-	
	20328	20330	232431	231796	245987	245295	883652	558805	851517	
SiO2	48.06	48.48	48.23	47.06	49.56	49.29	50.45	50.56	49.03	
A12O3	28.22	28.12	30.09	29.98	31.01	29.62	28.75	28.95	27.97	
FeO	1.86	1.77	2.5	2.32	2.64	2.83	2.48	3.13	3.28	
MnO	0.04	0.06	0.27	0.2	0.06	0.06	0.09	0.1	0.13	
MgO	3.15	3.03	2.67	2.71	2.89	2.98	2.27	2.52	2.44	
CaO	0.02	0.03	0.27	0.23	0.01	0.01	0	0	0	
Na2O	0.52	0.58	0.47	0.6	0.49	0.58	0.37	0.28	0.26	
K2O	9.49	9.36	10.04	10.17	10.14	10.5	10.17	11.01	11.04	
Total	91.36	91.43	94.54	93.27	96.8	95.87	94.58	96.55	94.15	
Atom site dis	tribution (12	1 anhydrous	-oxygen bas	is including	Fe3+)					
Si(T1+T2)	3.32	3.35	3.24	3.22	3.25	3.28	3.37	3.34	3.33	
Al(T2)	0.68	0.65	0.76	0.78	0.75	0.72	0.63	0.66	0.67	
V(M1)	0.94	0.95	0.98	0.97	0.94	0.95	1.00	0.99	1.00	
Mg(M1)	0.04	0.04	0.03	0.03	0.04	0.03	0.00	0.01	0.01	
Fe2+(M1)	0.01	0.01	0.01	0.01	0.02	0.02	0.00	0.01	0.00	
Al(M2+M3)	1.63	1.64	1.63	1.64	1.64	1.60	1.64	1.60	1.58	
Mg(M2+M3)	0.28	0.27	0.24	0.25	0.24	0.26	0.22	0.24	0.24	
Fe(M2+M3)	0.09	0.09	0.11	0.12	0.12	0.14	0.12	0.14	0.14	
XMg	0.75	0.75	0.68	0.68	0.66	0.65	0.65	0.64	0.64	
K(A)	0.84	0.82	0.86	0.89	0.85	0.89	0.87	0.93	0.96	
Na(A)	0.07	0.08	0.06	0.08	0.06	0.07	0.05	0.04	0.03	
V(A)	0.09	0.10	0.06	0.02	0.09	0.03	0.08	0.04	0.01	

15 Table 4: Representative analyses of white mica used in the multi-equilibrium calculations

	SAMPLE	GROUP	AGE Aln (2σ)	AGE Zrn	Р	т	CONSTRAINTS	NAME
	FG1324	1	77.2±7.3	77.2±7.3 77-63 1.65-1.75 GPa 600-650		Inclusion of Ph, Grt	Omp, Grt, Gln and Rt micaschist	
	FG1315	2	65.4±3.5	68-58	1.75-1.9 GPa	650-670	Inclusion of Ph, Grt (Rim1-2-3), Pg	Grt, Ep and Rt Qz- micaschist
C	FG12157 2		64.5±4.3	65-60	1.55-1.65 GPa	630-670	Inclusion of Ph	Grt, Gln, Ep and Rt micaschist
	FG1347 3		55.7±4.5	56±4	1.8-2 GPa	580-610	Inclusion of Ph, Pg, Rt	Cld, Grt and Rt micaschist
	FG1249	1	72.5±1.4	-	1.55-1.65 GPa	580-630	Inclusion of Ph, Pg	Grt, Ep and Rt micaschist
EC	FG1420	4	73.7±8.2	-	0.5-0.8 GPa	400-500	Inclusion of Ph, Pg,Ttn	Grt orthogneiss
	FG12107	4	62.8±3.3	-	no P	no T	Inclusion of Ph, Pg	Ep, Ph and Ab leucogneiss

16 Table 5: Constraints linking age data to pressure and temperature based on mineral inclusions in allanite; uncertainties

17 of age data are 2σ. Allanite age data based on Tera-Wasserburg intercepts. Zircon ²⁰⁶Pb/²³⁸U dates are given as the

18 minimum and maximum range of individual analysis. Groups are discussed in section 7.

		Age at ~0.6GPa	vertical velocity mm/year	along sub. inter. 60° mm/year	along sub. inter. 45° mm/year
	Group 1	~38 Ma	0.9	1.1	1.3
ge1	Group 1	~46 Ma	1.2	1.4	1.7
Sta	Group 3	~38 Ma	2.7	3.1	3.8
	Group 3	~46 Ma	5.1	5.9	7.2
ge2	INT-EXT	~38 Ma	4	-	-
Sta	INT-EXT	~46 Ma	1.6	-	-

21 Table 6: Exhumation velocities for Groups 1 and 3 (IC) from the highest pressures recorded to the greenschist

22 juxtaposition with the EC (Stage1) and for the late exhumation to the surface of the Sesia Zone (INT-EXT; Stage 2).