

# Permian plume beneath Tarim from receiver functions

Lev Vinnik<sup>1</sup>, Yangfan Deng<sup>2</sup>, Grigoriy Kosarev<sup>1</sup>, Sergey Oreshin<sup>1</sup>, Larissa Makeyeva<sup>1</sup>

1. Institute of physics of the Earth, Russian Academy of Sciences, Moscow, Russia
2. State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry,  
5 Chinese Academy of Sciences, Guangzhou 510640, China

Correspondence to: Yangfan Deng (Yangfandeng@gig.ac.cn) and Lev Vinnik ([vinnik@ifz.ru](mailto:vinnik@ifz.ru))

## Abstract

Receiver functions for the central Tien Shan and northern Tarim in central Asia reveal a pronounced depression on the 410-km discontinuity beneath the Permian basalts in Tarim. The 10 depression may most likely be caused by elevated temperature. The striking spatial correlation between the anomaly of the MTZ and the Permian basalts suggests that both may be effects of the same plume. This relation can be reconciled with reconstructed positions of paleo-continents since the Permian by assuming that the mantle layer which translated coherently with the Tarim plate extended to a depth of 410 km or more. Alternatively, lithosphere and the underlying 15 mantle are decoupled at a depth of ~ 200 km, but a cumulative effect of the Tarim plate motions since the Permian is by an order of magnitude less than predicted by the paleo-reconstructions. A similar explanation is applicable to the Siberian traps.

## 1. Introduction.

Theoretical considerations predict decoupling of the rigid lithosphere and the underlying 20 ductile upper mantle (asthenosphere) at the lithosphere-asthenosphere boundary (Eaton et al., 2009). The depth to the lithosphere-asthenosphere boundary (LAB) ranges from a few tens kilometers for a young lithosphere to about 300 km for Precambrian cratons (e.g. Artemieva and Mooney, 2001). Another idea postulates that the layer which translates coherently with the continental plate (tectosphere) may extend to a depth of at least 400 km (Jordan, 1978).

25 Examples of successful application of the concept of tectosphere to geophysical data are few. We test this idea by comparing the locations of possible remnants of extinct mantle plumes in the mantle transition zone (MTZ) and the related basaltic outcrops at the Earth's surface.

Recently this test was applied to the Siberian Large Igneous Province (LIP). The Siberian traps present the result of gigantic basalt eruptions which took place near the Permo-Triassic 30 boundary at about 250 Ma (Fedorenko et al., 1996). The analysis of structure of the mantle beneath the Siberian LIP was conducted with the aid of receiver function techniques that were applied to the recordings of seismograph station Norilsk (NRIL) within the Siberian LIP (Vinnik et al., 2017). This analysis has shown that the seismic boundary at the top of the MTZ with a standard depth of 410 km is depressed in the vicinity of NRIL by 10 km. The diagram of olivine 35 - wadsleyite phase transition may account for this depression by assuming about 100 K increase of the temperature.

In the depth range from 350 to 410 km, the S velocity beneath the Siberian LIP is reduced by 4 – 5% (Vinnik and Farra, 2007) relative to IASP91 model (Kennett and Engdahl, 1991). This is a likely effect of about 1 vol % melt (Hier-Majumder and Courtier, 2011) which is unusual for 40 cratons. Another low-velocity layer is found in the depth interval from 460 to 500 km. Previously this layer was found in the vicinities of several hot-spots (e.g., Vinnik et al., 2012). The low S-wave velocity coincides in depth with the abrupt decrease of the solidus temperature of carbonated mantle (Keshav et al., 2011) and may also be related to melting. The Siberian Craton shifted in the last 250 Myr by about 2000 km to the east (Torsvik et al., 2008). The 45 anomalies of the MTZ might preserve their position beneath the Siberian LIP in spite of the plate motion if they translated coherently with the Siberian plate.

A similar conclusion is obtained for Greenland by Kraft et al. (2018). Arrival times of P660s and P410s mode converted phases in P receiver functions (PRFs) were measured at 24 seismograph stations in central-eastern Greenland. In two regions corresponding to basaltic 50 outcrops about 55 Myr old, the differential time between P660s and P410s seismic phases is

reduced by more than 2 s relative to IASP91 reference model. The 410-km discontinuity in these regions is depressed by more than 20 km. The depression can be explained by elevated temperature. The basaltic outcrops and the related temperature anomalies are likely related to the passage of Greenland over the Iceland hot-spot. This explanation is consistent with the 55 concept of tectosphere and implies that the upper mantle beneath Greenland to a depth of at least 430 km translated coherently with the Greenland plate.

Here we describe a similar analysis for the central Tien Shan and Tarim in central Asia and discuss possible implications of these observations.

## 2. Seismic structure of the MTZ beneath the central Tien Shan and Tarim.

60 This section presents in condensed form the results of the recent seismic study (Kosarev et al., 2018) of the MTZ beneath the central Tien Shan and northern Tarim (Fig.1). The ongoing orogenesis in central Asia is a likely far-field effect of the India-Eurasia collision (Molnar and Tapponnier, 1975). Previous mountain-building episodes in the region of the Tien Shan took place in the Paleozoic (e.g., Windley et al., 1990), but for about 100 Myr prior to the onset of the 65 present-day mountain building the lithosphere of the Tien-Shan was quiet. Tectonic activity resumed at about 25-20 Ma in the southern Tien Shan (Sobel and Dumitru, 1997) and at 11 Ma in the north (Bullen et al., 2001). The lithosphere of Tarim underthrusts the relatively weak lithosphere of the Tien Shan at a rate of about 20 mm/yr (Reigber et al., 2001).

70 Teleseismic recordings of 64 broad-band stations in Fig. 1 were low-pass filtered with a corner at 6s and transformed into PRFs. The PRFs were calculated by using the LQ coordinate system, where L is parallel to the principal motion direction of the P wave and Q is normal to L in the wave propagation plane. The Q components were deconvolved by the L components in time domain and stacked to reduce noise. In the context of our study the most important elements 75 of the PRFs are P660s and P410s mode converted seismic phases. The 410-km and 660-km discontinuities mark the top and bottom of the MTZ and their depths are sensitive to the temperature and composition.

The times of P660s and P410s seismic phases depend not only on topography of the 660-km and 410-km discontinuities but also on volumetric velocity heterogeneities above the 410-km boundary. Separation of these two effects is the main problem of interpreting the observations of 80 P660s and P410s phases. This problem is solved by calculating the time difference (differential time) between the arrivals of P660s and P410s phases. The ray paths of P660s and P410s phases in the crust and upper mantle are close to each other for the same seismic recording, and, as a result, the differential time is insensitive to the properties of the Earth's medium above the MTZ.

To detect P660s and P410s phases and to map the differential time, a large number of the 85 PRFs should be stacked. One possibility is to apply a version of CCP (Common Conversion Point) stacking: to divide the Earth's surface into cells and to stack with appropriate move-out time corrections the PRFs, the projections of the conversion points of which fall into the same cell. However, the surface projections of the conversion points of P410s and P660s phases for the same recording are at different distances (around  $1^\circ$  and  $2^\circ$ , respectively) from the 90 seismograph station, and the set of PRFs thus selected for the detection of P410s phase may differ from that for P660s phase. Then the differential time of stacked P660s and P410s phases can be affected by lateral heterogeneity of the crust and mantle above the MTZ. This can be avoided by locating the conversion points in the middle of the MTZ (at a depth of 535 km) and stacking those PRFs, the projections of the conversion points of which are located within the 95 same cell. Then P410s and P660s phases for each cell are detected in the same set of PRFs and the effect of lateral heterogeneity above the 410-km discontinuity is minimized.

Epicenters of seismic events of sufficient magnitude in a distance range from  $35^\circ$  to  $90^\circ$  are abundant in a broad azimuth range. Surface projections of the conversion points at a depth of 535 km cover the area between  $38^\circ\text{N}$  and  $44^\circ\text{N}$  and between  $72^\circ\text{E}$  and  $82^\circ\text{E}$ . The cells were chosen 100 in the form of a rectangular box. The size of the box affects lateral resolution and, through the number of stacked PRFs, accuracy of the estimates of the differential time. The individual PRFs were visually inspected and only those of the best quality were stacked. The optimum size of the

box ( $2^{\circ}$  for NS and EW or 220 km and 160 km, respectively) was found by trial and error. The largest number of the stacked PRFs exceeds 1750, the smallest is 48 (See Fig. 3Table 4). These numbers are sufficient for a robust detection of P660s and P410s phases (see example in Fig. 2).  
105 The accuracy of the estimates of the differential time (confidence interval of 66%) which was determined by bootstrap resampling (Efron, Tibshirani, 1991) is typically 0.2 s. For most boxes the residuals of the differential time with respect to the IASP91 value (23.9 s) are on the order of a fraction of a second (Fig. 3). Large residuals (more than 1.0 s) are obtained for three boxes:  
110 ( $40^{\circ}$  -  $42^{\circ}$ N,  $76^{\circ}$  -  $78^{\circ}$ E, +1.5 s), ( $40^{\circ}$  -  $42^{\circ}$ N,  $72^{\circ}$  -  $74^{\circ}$ E, -1.1 s) and ( $38^{\circ}$  -  $40^{\circ}$ N,  $80^{\circ}$  -  $82^{\circ}$ E, -1.5 s). Further on these boxes are referred as a, b and c. The resulting anomalies of thickness of the MTZ for a, b and c are +15 km, -11 km, and -15 km, respectively. These anomalies are located beneath the south-central Tien Shan, Fergana Basin and Tarim. We note that while the number of stacked PRFs for c is minimal (48), quality of the PRFs (signal-noise ratio) in this box  
115 is very high and the accuracy of the differential time is comparable with the other boxes.

The further analysis (Kosarev et al., 2018) demonstrates that the increased thickness of the MTZ in a is the effect of an uplift of the 410-km discontinuity and a depression of the 660-km discontinuity. The MTZ might be cooled by a detached and sinking mantle lithosphere. The thinned MTZ in b and c is the effect of a depressed 410-km discontinuity and a stable 660-km discontinuity. The depressed 410-km discontinuity beneath b and c is likely a result of a temperature anomaly of about +100°C. The elevated temperature in b may be related to a plume which is responsible for small-scale basaltic volcanism in the Tien Shan from 72 Ma to 60 Ma.  
120 Possible origin of the anomaly in c (Tarim) is discussed in next Section.

### **3. Possible origin of the anomalous MTZ beneath Tarim.**

125 Tarim can be characterized as an Archean craton (Yuan et al., 2004) with a complex evolutional history (Zhang et al., 2013; Deng et al., 2017). In the Permian, basalts with the areal extent of about 200000 km<sup>2</sup> erupted in the west of the Tarim basin (Fig. 4). The thickness of basalt reaches 800 m. The age span of the magmatism extends from about 292 Ma to 272 Ma

with two peaks at 279 Ma and 289 Ma (Wei et al., 2014). The magmatism is interpreted as  
130 plume-induced (Zhang et al., 2010; Xu et al., 2014). Evidence for the mantle plume beneath  
Tarim includes the large volume of the Permian mafic rocks, OIB-like trace element signatures,  
Permian crustal doming and the high zircon saturation temperatures (Zhang et al., 2008, 2010).  
No magmatic activity is known after the Permian (Zhang et al., 2013; Deng et al., 2017).

Fig. 4 demonstrates a striking spatial correlation of the depressed 410-km discontinuity and  
135 the Permian magmatic province in Tarim, with implication of a causal relation between them. An  
alternative interpretation suggests that the topography on the 410-km discontinuity, though  
spatially correlated with the Permian basalts is caused by another, relatively young plume.  
However, this seems unlikely, as recently erupted (post-Permian) basalts are unknown in Tarim.  
The depressed 410-km discontinuity and the stable 660-km discontinuity are typical for hotspots  
140 and plumes (e.g. Du et al., 2006), though there are some exceptions (e.g. Vinnik et al., 2012). The  
stable depth of the 660-km discontinuity is either the result of a zero temperature anomaly at the  
base of the MTZ or an effect of two phase transitions at nearly the same depth but with opposite  
Clapeyron slopes (Hirose, 2002).

The assumed causal relation between the Permian basalts and the present-day anomaly  
145 implies that the anomaly at a depth of ~400 km may exist for ~300 Myr. To check this  
possibility we calculated the temperature for 1D conductive medium by using the well known  
expression (e.g., Zharkov et al., 1969)  $T(r,t) = Q \exp(-r^2/4\alpha t)/2\sqrt{\pi\alpha t}$ , where  $T$  is temperature,  $t$  is  
time,  $r$  is distance,  $\alpha$  is diffusivity,  $Q$  is constant, and the initial temperature anomaly distribution  
is taken in the form of  $\delta$ -function at  $r = 0$  and  $t = 0$ . The diffusivity  $\alpha$  is taken equal to 32  
150  $\text{km}^2/\text{m.y.}$  (e.g., Morgan and Sass, 1984). The results (Fig.5a) demonstrate that the temperature  
anomaly in the time interval of 300 m.y. (between 100 m.y. and 400 m.y.) is halved. The  
maximum temperature anomaly in plumes is  $\sim 300 \pm 100^\circ\text{C}$  (Campbell, 2005), which means that  
the temperature anomaly after 300 m.y. may be around  $150^\circ\text{C}$ , close to the seismic estimate. A

comparable result is obtained for 2D conductive medium (Fig. 5b). These calculations suggest  
155 that the thermal anomaly at a depth of 400 km may survive for a few hundred million years.

It is also possible that the anomalous depth of the 410-km discontinuity is an effect of  
anomalous composition. The pressure of the phase transition in  $(\text{Mg},\text{Fe})_2\text{SiO}_4$  depends on the  
Mg content (Mg#) relative to Fe (Fei and Bertka, 1999). Increasing Mg# from 89 to 92 results in  
up to 10-km deepening of the 410-km discontinuity (Schmerr and Garnero, 2007). The depleted  
160 composition and increased Mg# are commonly interpreted as effects of melting (e.g., Boyd,  
1989).

Relative positions of the present-day thermal anomaly in the MTZ and the Permian basalt  
eruptions depend on plate motions in the last ~300 Myr. Reconstruction of positions of old  
continents is difficult for the time exceeding the age of the oldest hot-spot trails (130 Ma). There  
165 are abundant paleomagnetic data for the earlier times, but they do not constrain paleo-longitudes.  
The motions of Tarim are constrained by paleomagnetic data. According to Zhao et al. (1996)  
Tarim might be attached to Eurasia since the Late Paleozoic time, and there are paleomagnetic  
indications of displacements of Tarim relative to Eurasia after the Cretaceous, apparently owing  
to the India-Eurasia collision (Zhao et al., 1996). The uncertainty of the paleoreconstructions for  
170 the Mesozoic can be minimized by selecting Africa as a reference continent that was most stable  
longitudinally (Torsvik et al. 2008). In this reference frame the Siberian traps shifted to the east  
by nearly 2000 km since they were erupted at 250 Ma (Torsvik et al., 2008). On the assumption  
that Tarim and the Siberian craton were parts of the same continental plate in the past 300 Myr,  
2000 km can be used as a rough estimate of the shift of Tarim.

175 The spatial correlation between the anomaly in the MTZ and the basalt eruptions in Tarim  
(Fig. 4) in spite of the shift of the Tarim craton to the east and north-east by a few thousand  
kilometers is possible if the layer which translates coherently with the plate includes the top of  
the MTZ. Alternatively this is possible without the recourse to the deep tectosphere, if the

available paleo-reconstructions for Asia are too rough and the actual shift of Tarim is by an order

180 of magnitude less than predicted. This would also be true for the Siberian traps.

#### 4. Conclusions

The striking spatial coincidence of the Permian basalts and a depression on the 410-km discontinuity beneath Tarim (Fig. 4) suggests that both may be related to the same mantle plume.

This relation allows a dual interpretation. Recent reconstructions (Torsvik et al., 2008)

185 demonstrate a shift of Tarim of about 2000 km in the past 300 Myr. Then the observed relation

between the deep and shallow features can be explained by a coherent translation of the crust and

mantle to a depth of 430 km. Alternatively the spatial coincidence of the deep and shallow

features is possible without the recourse to the deep tectosphere if the actual shift of Tarim is by

an order of magnitude less than predicted by the reconstructions. Practically similar conclusions

190 would apply to the Permo-Triassic traps of the Siberian Craton.

It would be useful in the future to find other evidence that may indicate whether tectosphere  
extends to over 400-km depth. For example, it would be interesting to look at the strength and  
direction of azimuthal anisotropy in the 100-400 km depth range to investigate if there is  
evidence for flow related to absolute plate motion shallower than the MTZ. It is difficult to do  
this with the available seismic data. The Tarim basin has the largest desert area in China,  
practically devoid of seismogram stations, but more data will be available in future.

#### Acknowledgment

This study was supported by the joint project between Russian Foundation for Basic Research

(RFBR, grant 17-55-53-117) and National Natural Science Foundation of China (NSFC, grant

200 41611530695), the Strategic Priority Research Program (B) of the Chinese Academy of Sciences

(grant XDB18000000). The authors appreciate comments from Dr. R. Porritt and an anonymous reviewer.

#### References

Artemieva, I. M., and Mooney, W. D.: Thermal thickness and evolution of Precambrian  
205 lithosphere: a global study, *Journal of Geophysical Research: Solid Earth*, 106(B8), 16387-16414, 2001

Boyd, F.R.: Compositional distinction between oceanic and cratonic lithosphere, *Earth Planet. Sci. Lett.*, 96, 15-26, 1989.

Bullen, M. E., Burbank, D. W., Garver, J. I., and Abdurakhmatov, K. Y.: Late Cenozoic tectonic  
210 evolution of the northwestern Tien Shan: New age estimates for the initiation of mountain building, *Geological Society of America Bulletin*, 113(12), 1544-1559, 2001.

Campbell, I. H.: Large igneous provinces and the mantle plume hypothesis, *Elements*, 1(5), 265-269, 2005.

Deng, Y., Levandowski, W., and Kusky, T.: Lithospheric density structure beneath the Tarim  
215 basin and surroundings, northwestern China, from the joint inversion of gravity and topography, *Earth and Planetary Science Letters*, 460, 244-254, 2017.

Du, Z., Vinnik, L. P., and Foulger, G. R.: Evidence from P-to-S mantle converted waves for a flat “660-km” discontinuity beneath Iceland, *Earth and Planetary Science Letters*, 241(1-2), 271-280, 2006.

220 Eaton, D. W., Darbyshire, F., Evans, R. L., Grütter, H., Jones, A. G., and Yuan, X.: The elusive lithosphere–asthenosphere boundary (LAB) beneath cratons, *Lithos*, 109(1-2), 1-22, 2009.

Efron, B., and Tibshirani, R.: Statistical data analysis in the computer age, *Science*, 253, 390-395, 1991.

Fedorenko, V. A., Lightfoot, P. C., Naldrett, A. J., Czamanske, G. K., Hawkesworth, C. J.,  
225 Wooden, J. L., and Ebel, D. S.: Petrogenesis of the flood-basalt sequence at Noril'sk, north central Siberia, *International Geology Review*, 38(2), 99-135, 1996.

Fei, Y., and Bertka, C. M.: Phase transitions in the Earth's mantle and mantle mineralogy, *Mantle petrology: field observations and high pressure experimentation*, 6, 189-207, 1999.

Jordan, T. H.: Composition and development of the continental tectosphere, *Nature*, 274(5671), 544, 1978.

Hier-Majumder, S., and Courtier, A.: Seismic signature of small melt fraction atop the transition zone, *Earth and Planetary Science Letters*, 308(3-4), 334-342, 2011.

Hirose, K.: Phase transitions in pyrolytic mantle around 670 - km depth: Implications for upwelling of plumes from the lower mantle, *Journal of Geophysical Research: Solid Earth*, 107(B4), 2002.

Kennett, B.L.N. and Engdahl, E.R.: traveltimes for global earthquake location and phase identification. *Geophysical Journal International*, 105(2), 429-465, 1991.

Keshav, S., Gudfinnsson, G.H., and Presnall, D.C.: Melting phase relations of simplified carbonated peridotite at 12 - 26 GPa in the systems CaO - MgO - SiO<sub>2</sub> - CO<sub>2</sub> and CaO - MgO - Al<sub>2</sub>O<sub>3</sub> - SiO<sub>2</sub> - CO<sub>2</sub>: highly calcic magmas in the transition zone of the Earth. *Journal of Petrology*, 52, 2265-2291, 2011.

Kosarev, G., Oreshin, S., Vinnik, L., and Makeyeva, L.: Mantle transition zone beneath the central Tien Shan: Lithospheric delamination and mantle plumes, *Tectonophysics*, 723, 172-177, 2018.

Kraft, H. A., Vinnik, L., and Thybo, H.: Mantle transition zone beneath central-eastern Greenland: Possible evidence for a deep tectosphere from receiver functions, *Tectonophysics*, 728, 34-40, 2018.

Molnar P., and Tapponnier P.: Cenozoic tectonics of Asia: effects of a continental collision, *Science*, 189, 419-426, 1975.

Morgan, P. and Sass, S.: Thermal regime of the continental lithosphere, *Journal of Geodynamics* 1, 143-166, 1984.

Reigber, C., Michel, G., Galas, R., Angermann, D., Klotz, J., Chen, J., Papschev, A., Arslanov, R., Tzurkov, V. and Ishanov, M.: New space geodetic constraints on the distribution of deformation in Central Asia. *Earth and Planetary Science Letters*, 191(1-2), 157-165, 2001.

255 Schmerr, N., and Garnero, E. J.: Upper mantle discontinuity topography from thermal and chemical heterogeneity, *Science*, 318(5850), 623-626, 2007.

Sobel, E. R., and Dumitru, T. A.: Thrusting and exhumation around the margins of the western Tarim basin during the India–Asia collision, *Journal of Geophysical Research: Solid Earth*, 102(B3), 5043-5063, 1997.

260 Torsvik, T. H., Steinberger, B., Cocks, L. R. M., and Burke, K.: Longitude: linking Earth's ancient surface to its deep interior, *Earth and Planetary Science Letters*, 276(3-4), 273-282, 2008.

Vinnik, L., and Farra, V.: Low S velocity atop the 410-km discontinuity and mantle plumes, *Earth Planet. Sci. Lett.*, 262, 398-412, 2007.

265 Vinnik, L., Silveira, G., Kiselev, S., Farra, V., Weber, M., and Stutzmann, E.: Cape Verde hotspot from the upper crust to the top of the lower mantle, *Earth and Planetary Science Letters*, 319, 259-268, 2012.

Vinnik, L. P., Oreshin, S. I., and Makeyeva, L. I.: Siberian traps: Hypotheses and seismology data, *Izvestiya, Physics of the Solid Earth*, 53(3), 332-340, 2017.

270 Wei, X., Xu, Y.-G., Feng, Y.-X., Zhao, J.-X.: Plume-lithosphere interaction in the generation of the Tarim large igneous province, NW China: geochronological and geochemical constraints. *American Journal of Science*, 314, 314-356, 2014.

Windley, B. F., Allen, M. B., Zhang, C., Zhao, Z. Y., and Wang, G. R.: Paleozoic accretion and Cenozoic redeformation of the Chinese Tien Shan range, central Asia. *Geology*, 18(2), 128-131, 1990.

275 Xu, Y.-G., Wei, X., Luo, Z.-Y., Liu, H.-Q., and Cao, J.: The Early Permian Tarim Large Igneous Province: main characteristics and a plume incubation model. *Lithos*, 204, 20–35, 2014.

Yuan, C., Sun, M., Yang, J., Zhou, H., and Zhou, M.-F.: Nb-depleted, continental rift-related Akaz metavolcanic rocks (West Kunlun): implication for the rifting of the Tarim Craton from Gondwana. *Geological Society, London, Special Publications*, 226(1), 131-143, 2004.

Zhang, C., Li, X., Li, Z., Ye, H., and Li, C.: A permian layered intrusive complex in the western tarim block, northwestern China: Product of a Ca. 275-ma mantle plume? *The Journal of Geology*, 116, 269-287, 2008.

285 Zhang, C.-L., Xu, Y.-G., Li, Z.-X., Wang, H.-Y., and Ye, H.-M.: Diverse Permian magmatism in the Tarim Block, NW China: Genetically linked to the Permian Tarim mantle plume? *Lithos*, 119, 537-552, 2010.

Zhang, C.-L., Zou, H.-B., Li, H.-K., and Wang, H.-Y.: Tectonic framework and evolution of the Tarim Block in NW China. *Gondwana Research*, 23(4), 1306-1315, 2013.

290 Zhao, X., Coe, R.S., Gilder, S.A. and Frost, G.M.: Paleomagnetic constraints on the palaeogeography of China: implications for Gondwanaland, *Australian Journal of Earth Sciences*, 43, 643-672, 1996.

Zharkov, V.N., Pan'kov, V.L., Kalachnikov, A.A. and Osnach, A.I. Introduction in the physics of the Moon , “Nauka”, Moscow, 1969. (in Russian).

295 **Figure captions**

**Figure 1.** Topographic map of the study region and the seismograph network.

**Figure 2.** Stacked PRFs for the box with the coordinates: 38 N, 40N, 80 E and 82 E. Trial conversion depths in kilometers for each trace are shown on the left-hand side. The detected P410s and P660s phases are marked by arrows. Note that the largest amplitudes of P410s and 300 P660s phases are observed at appropriate trial depths (around 400-500 km and 600=700 km, respectively).

**Figure 3.** Residuals of the differential time between P660s and P410s phases in seconds relative to IASP91. Strongly anomalous boxes are in south-central Tien Shan (1.5 s, blue, a), Fegana basin (-1.1 s, red, b) and Tarim (-1.5 s, red, c). Light shading indicates elevations greater than 305 1500 m, intermediate shading elevations greater than 3000 m. The number of stacked receiver functions in each box is shown by italics.

**Figure 4.** Superimposed Permian basalts in Tarim (orange) and the anomalous region on the 410-km discontinuity.

**Figure 5.** Temperature anomaly distributions in 1D (a) and 2D (b) conductive media with an

310 interval of 300 million years.

**Table 1. Numbers of piercing points in the boxes in Fig. 3.**

lat1	lat2	lon1	lon2	number of pp
38	40	72	74	132
38	40	74	76	267
38	40	76	78	402
38	40	78	80	96
38	40	80	82	48
40	42	72	74	414
40	42	74	76	1130
40	42	76	78	1512
40	42	78	80	616
40	42	80	82	322
42	44	72	74	317
42	44	74	76	757
42	44	76	78	1766
42	44	78	80	533
42	44	80	82	332
44	46	74	76	120