

1 **STABILITY OF SOIL ORGANIC MATTER IN CRYOSOLS OF MARITIME**  
2 **ANTARCTIC: INSIGHTS FROM <sup>13</sup>C NMR AND ELECTRON SPIN RESONANCE**  
3 **SPECTROSCOPY**

4  
5 **Evgeny Abakumov, Ivan Alekseev**

6 *Department of Applied Ecology, Saint-Petersburg State University, 199178, 16-line 2, Vasilyevskiy*  
7 *Island, Russian*

8  
9 *e-mail: [e.abakumov@spbu.ru](mailto:e.abakumov@spbu.ru), [e\\_abakumov@mail.ru](mailto:e_abakumov@mail.ru), [st014661@student.spbu.ru](mailto:st014661@student.spbu.ru)*

10 **Key words**

11 Antarctica, soil organic matter, stabilization, humic acids

12  
13 **Key points**

14 Investigation of Antarctic soil organic matter stability

15 Humic acids of superficial horizons contain more aromatic carbon

16 Humic acids of isolated layers contain more free radicals

17  
18  
19 **Abstract**

20  
21 Previously, the structure and molecular composition of the Antarctic SOM has been investigated  
22 using <sup>13</sup>C-NMR methods, which showed that in typical organo-mineral soils the aliphatic carbon  
23 prevails over the aromatic one, owing to the non-ligniferous nature of its precursor material. In  
24 this study, the soil organic matter (SOM) was analyzed from different sample areas (surface level  
25 and partially isolated supra-permafrost layer) of the tundra-barren landscape of the Fildes  
26 Peninsula, King George Island, Western Antarctica. We found that the humic acids (HAs) of the  
27 cryoturbated, buried areas had lower amounts of alkylaromatic and protonized aromatic  
28 compounds. In contrast, the HAs from the surface layers contain less alkyl carbon components.  
29 The free radical content was higher in the surface layers than in the buried layers due to the  
30 presence of fresh organic remnants in superficial soil samples. New data on SOM quality from  
31 these two representative Cryosols will enable more precise assessment of SOM stabilization rate  
32 in sub-Antarctic tundras. Comparison of the <sup>13</sup>C-NMR spectra of the HAs and the bulk SOM  
33 revealed that humification occurs in the Antarctic and results in accumulation of aromatic and  
34 carboxylic compounds and reductions in alkylic ones. This indicates that humification is one of  
35 the ways of soil organic matter stabilization.

36  
37  
38 **Highlights**

39 Stabilization of soil organic matter studied

40 Humic acids of superficial horizons contain more aromatic carbon

41 Humification is one of the ways of soil organic carbon stabilization

42

## 43 1. Introduction

44 Polar soils play a key role in global carbon circulation and stabilization as they contain  
45 maximum stocks of soil organic matter (SOM) within the whole pedosphere (Schoor et al, 2015).  
46 Cold climate and continuous and discontinuous permafrost result in the stabilization of essential  
47 amounts of organic matter in soils, biosediments, and grounds of the polar biome (Zubrzycki et al,  
48 2014). Global climate changes and permafrost degradation have led to the exposure of huge pools  
49 of organic matter to microbial degradation (Schoor et al, 2015) and other environmental risks.  
50 Polar SOM represents a vulnerable carbon source, susceptible to remobilization under increasing  
51 temperatures (Schoor et al, 2015, Ejarque, Abakumov, 2016). In order to better understand the  
52 implications of permafrost SOM for greenhouse gas emissions, accurate knowledge of its spatial  
53 distribution, both in terms of quantity and quality (e.g. biodegradability, chemical composition,  
54 and humification stage) is needed in addition to effective evaluation of SOM's temporal  
55 dynamics (Fritz et al, 2014, Vasilevitch et al, 2018 ).

56 Current estimations of soil organic carbon (SOC) stocks are around 1307 Pg throughout  
57 the northern circumpolar region (Hugelius et al, 2014). These amounts surpass previous  
58 estimates (Tarnocai et al, 2009) and grossly exceed the total carbon contained in the world's  
59 vegetation biomass (460 - 650 Pg) or in the atmosphere (589 Pg) (Tarnocai et al, 2009).  
60 However, the aforementioned SOM/SOC stock estimations are still poorly constrained (Hugelius  
61 et al, 2014). This uncertainty is largely caused by the estimates having been calculated from  
62 observations that are highly spatially clustered (Hugelius et al, 2014) while extensive land areas  
63 remain uncharacterized due to the logistic difficulties of reaching these sites. Additionally, the  
64 calculation of these stocks are based on estimated data on soil bulk density and carbon values  
65 derived from dichromate oxidation methods (Abakumov, Popov, 2005, Polyakov et al, 2017).

66 The stocks of SOM in the Antarctic are underestimated compared to the Arctic because  
67 of the lack of the data for many parts of this continent, due to the high content of stones in the  
68 soils and the high variability in the carbon content of the fine earth. Stocks of organic carbon in  
69 the Antarctic soil have been reported as 0.5 kg/m<sup>2</sup> in its polar deserts, about 1.0 kg/m<sup>2</sup> in its  
70 barrens, up to 3 - 5 kg/m<sup>2</sup> in the sub-Antarctic tundra, and up to 30 kg/m<sup>2</sup> in the penguin  
71 rookeries of the maritime islands (Abakumov, 2010, Abakumov, Mukhametova, 2014,  
72 Abakumov et al, 2016). To date, investigation on structural composition of the SOM from both  
73 superficial and partially isolated areas has only been performed on Cryosols of the Kolyma  
74 lowland (Lupachev et al, 2017), where the organic matter of modern and buried soils vary  
75 greatly in terms of their molecular composition and quality.

76 Stability and biodegradability are the key features of SOM that should be taken into  
77 account when estimating current and future carbon stocks and organic matter quality and  
78 dynamics. Stability is related to humification degree, as more advanced stages in the  
79 humification process involve depletion of the labile molecules, as well as an increase in the bulk  
80 aromaticity, which confers higher stability to the SOM. A number of proxies have been used to  
81 trace humification rate and SOM stability, including aromaticity level (Vasilevitch et al, 2018,  
82 Kniker, 2007). Also the ratio of C-Alkyl : C-Aryl and C-Alkyl : O-N-alkyl have been  
83 successfully used to assess humification degree (Kinker, 2007). C/H ratio from humic acids  
84 (HAs) has been used as an index of molecular complexity, as more degrees of conjugation imply  
85 less hydrogenation of the carbon chains (Zaccone et al, 2007) and C/N has been used as a  
86 measure of histic material degradation (Lodygin et al, 2014). <sup>13</sup>C-NMR spectrometry provides  
87 information on the diversity in carbon functional structures (carbon species) and has been used to

88 evaluate changes in SOM during decomposition and humification. More specifically, high  
89 phenolic (150 ppm), carboxyl-C (175 ppm) and alkyl-C (30 ppm) groups, combined with low O-  
90 alkyl carbons, have been associated with advanced humification stages (Zech et al, 1997). So far,  
91 studies of SOM quality from polar environments have revealed generally lowly-decomposed  
92 organic molecules (Dziadowiec, 1994, Lupachev et al, 2017), which preserve much of the  
93 chemical character of their precursor material due to slow progress of humification (Davidson  
94 and Jansens, 2006). This is very important because polar soils are characterized by the specific  
95 composition of the humification precursors.

96 The structure and molecular composition of the Antarctic SOM has been investigated  
97 using  $^{13}\text{C}$ -NMR methods (Beyer et al, 1997, Abakumov, 2017) and it was shown that in typical  
98 organo-mineral soils the aliphatic carbon prevails over the aromatic one, owing to the non-  
99 ligniferous nature of its precursor material (Calace et al, 1995). Also, analyses of cryptogam  
100 extracts were conducted towards identification of individual organic precursors (Chapman et al,  
101 1994). This feature was then shown to be typical for soils from different regions of the Antarctic  
102 (Abakumov, 2010), including soil formed on the penguin rockeries (Abakumov, Fattakhova,  
103 2015). The northern most soil of Arctic polar biome shows the same trend in organic molecules  
104 organization: higher prevalence of aliphatic structures over aromatic ones. The diversity of the  
105 individual components in aromatic and aliphatic areas is usually higher in Arctic soil because of  
106 the increased diversity of humification precursors (Ejarque, Abakumov, 2016, Abakumov,  
107 2010). A selective preservation of the alkyl moieties in the deeper soil layers has been suggested,  
108 and little transformation processes of the SOM are detectable because soil temperatures are not  
109 high enough to stimulate further microbial break-down, even in the summer (Beyer et al, 1997).  
110 It has been shown that ornithochoria play an essential role in redistribution of plant remnants in  
111 the Antarctic (Parnikoza et al, 2016) as birds transport considerable amounts of variably  
112 composed organic material within its inland landscapes. However, published data on SOM  
113 composition for the Antarctic are rare, and further studies that detail its structural compounds  
114 and their distribution are needed. Recently,  $^{13}\text{C}$ -NMR was successfully used to detail the soils  
115 found in endolithic communities in Eastern Antarctica and revealed that endolithic organic matter  
116 is characterized by a low prevalence of alkyl aromatic compounds (Mergelov et al, 2018).

117 This study aimed to compare the structural composition of the SOM from both superficial  
118 and partially isolated (i.e. buried spots on the border with permafrost) areas and to evaluated the  
119 stabilization rate of Antarctic Cryosols. The objectives of the study were: (1) to evaluate the  
120 alterations in the elemental compositions of the HAs under partial isolation (2) to assess the  
121 ratios of aromatic and aliphatic carbon species in the topsoil and isolated areas; (3) to  
122 characterize the biochemical activity of the HAs (e.g. free radical concentration).

## 123 **2. Materials and Methods**

### 124 **2.1. Study sites**

125 King George Island is the largest in the South Shetland archipelago and only around 5%  
126 of its 1400 km<sup>2</sup> area is free of ice (Fig. 1) (Rakusa-Suszczewski, 2002). The Fildes Peninsula and  
127 Ardley Island, together around 33 km<sup>2</sup>, comprise the largest ice-free area on King George Island  
128 and the second largest of the South Shetland Islands. It has a gentle landscape consisting of old  
129 coastal landforms with numerous rocky ridges and an average height of 30 m above main sea  
130 level (AMSL) (Michel et al, 2014). According to Smellie (Smellie et al, 2014), this area mainly  
131 consists of lava with small exposures of tuffs, volcanic sandstones, and agglomerates. The  
132 climate is cold and humid with a mean annual air temperature of -2.2°C and mean summer air  
133 temperatures above 0°C for only up to four months (Wen et al, 1994). The mean annual  
134 precipitation is 350 - 500 mm/year. The Fildes Peninsula and Ardley Island are among the first

135 areas in maritime Antarctica to become ice-free after the last glacial maximum (Birkenmajer,  
136 1989). The Fildes Peninsula was covered by glaciers from 8000 to 5000 BP (Mausbacher et al,  
137 1989, Haus et al, 2014). The patterned ground in this region dates from 720 to 2640 BP. In the  
138 South Shetland Islands, permafrost is sporadic or non-existent at altitudes below 20 m AMSL  
139 and occurs discontinuously in altitudes from 30 to 150 m AMSL (Bockheim et al, 2013).  
140 Mosses, lichens, and algae are common to this area along with two vascular plants (*Deschampsia*  
141 *antarctica* and *Colobanthus quitensis*). Penguins, seals, and seabirds inhabit the coastal areas and  
142 greatly impact the soil development. Major cryogenic surface-forming processes in this region  
143 include frost creep, cryoturbation, frost heaving and sorting, gravity, and gelifluction (Michel et  
144 al, 2014). Eight separate sites on the Fildes Peninsula have been collectively designated an  
145 Antarctic Specially Protected Area (ASP 125) largely due to their paleontological properties  
146 (Management plan, 2009). The average thickness of the soil is about 15 - 25 cm. Soils from King  
147 George Island have been divided into six groups (WRB, 2014): Leptosols, Cryosols, Fluvisols,  
148 Regosols, Histosols, and Technosols; this corresponds well with previously published data  
149 (Navas et al, 2008, Abakumov, 2017).

150 Three soils were selected for humic substance isolation and further investigation in this  
151 study. All soils have top humus layers with a high carbon content and distinguishable layers of  
152 suprapermafrost accumulation of organic matter. All three soils are classified as Turbic Cryosols  
153 (Histic, Stagnic) (WRB, 2014). Soil profiles 1, 2, and 3 (SP1, SP2, SP3) were collected from  
154 locations described by the following coordinates: 62,14,391 S, 58,58,549 W; 62,13,140 S,  
155 58,46,067 W; and 62,10,578 S, 58, 51,446 W respectively. Sampling depth was 0 - 10 cm for the  
156 superficial layers and 50 - 55, 15 - 20, 20 - 25 for SP1, SP2, and SP3 respectively. Images of the  
157 soil profiles are presented in Fig. 2. SP1 is from under the mixed lichen-bryophyta cover, SP2  
158 and SP3 are formed under species of *Bryophyta* and *Deschampsia antarctica* correspondingly.

## 159 **2.2. Sampling and laboratory analysis**

160 Soil samples were air-dried (24 hours, 20°C), ground, and passed through 2-mm sieve.  
161 Routine chemical analyses were performed using classical methods: C and N content were  
162 determined using an element analyzer (Euro EA3028-HT Analyser) and pH in water and in salt  
163 suspensions using a pH-meter (pH-150 M).

164 Humic acids (Has) were extracted from each sample according to a published protocol  
165 (Shnitzer, 1982), <http://humic-substances.org/isolation-of-ihss-samples/>). Briefly, the soil  
166 samples were treated with 0.1 M NaOH (soil/solution mass ratio of 1:10) under nitrogen gas.  
167 After 24 hours of shaking, the alkaline supernatant was separated from the soil residue by  
168 centrifugation at  $1,516 \times g$  for 20 minutes and then acidified to pH 1 with 6 M HCl to precipitate  
169 the HAs. The supernatant, which contained fulvic acids, was separated from the precipitate by  
170 centrifugation at  $1,516 \times g$  for 15 minutes. The HAs were then dissolved in 0.1 M NaOH and  
171 shaken for four hours under nitrogen gas before the suspended solids were removed by  
172 centrifugation. The resulting supernatant was acidified again with 6 M HCl to pH 1 and the HAs  
173 were again isolated by centrifugation and demineralized by shaking overnight in 0.1 M HCl/0.3  
174 M HF (soil/solution ratio of 1:1). Next, the samples were repeatedly washed with deionized  
175 water until pH 3 was reached and then finally freeze-dried. HA extraction yields were calculated  
176 as the percentage of carbon recovered from the original soil sample (Vasilevitch et al, 2018,  
177 Abakumov et al, 2018).

178 Isolated HAs were characterized for their elemental composition (C, N, H, and S) using the  
179 Euro EA3028-HT analyzer. Data were corrected for water and ash content. Oxygen content was  
180 calculated by difference. The elemental ratios reported in this paper are based on weight. Solid-  
181 state  $^{13}\text{C}$ -NMR spectra of HAs were measured with a Bruker Avance 500 NMR spectrometer in

182 a 3,2-mm ZrO<sub>2</sub> rotor. The magic angle spinning speed was 20 kHz in all cases and the nutation  
183 frequency for cross polarization was  $u/2p\ 1/4\ 62.5$  kHz. Repetition delay and number of scans  
184 were 3 seconds. Groups of structural compounds were identified by their chemical shifts values:  
185 alkyl C (-10 to 45 ppm), O/N-alkyl C (45 to 110 ppm), aryl/olefine C (110 to 160 ppm), and  
186 carbonyl/carboxyl/amide C (160 to 220 ppm) (Knicker, 2007). The <sup>13</sup>C-NMR study was also  
187 conducted in bulk soil samples towards characterizing changes in the initial soil material during  
188 humification.

189 The ESR spectra (only for HAs due to low ash content) were recorded on a JES FA 300  
190 spectrometer (JEOL, Japan) in X-diapason with a free-radical modulation amplitude of 0.06 mT  
191 and a microwave power in the cavity of 1 mW. Magnesium powder with fixed radical  
192 concentration was used as an external standard. The concentration of the paramagnetic centers in  
193 powdered samples was determined by comparison to relative signal intensities of the external  
194 standard using the program JES-FA swESR v. 3.0.0.1 (JEOL, Japan). (Chukov et al, 2017).

### 195 **2.3. Statistical analysis**

196 Statistical data analysis was performed using the STATISTICA 10.0 software (TX, USA).  
197 One-way analysis of variance (ANOVA) was applied to test the statistical significance of the  
198 differences between the data, based on estimation of the significance of the average differences  
199 between three or more independent groups of data combined by one feature (factor). Fisher's  
200 Least Significance Test (LST) was used for post-hoc analysis to provide a detailed evaluation of  
201 the average differences between groups. A feature of this post-hoc test is inclusion of intra-group  
202 mean squares when assessing any pair of averages. Differences were considered significant at  
203 the 95% confidence level. Concentrations of organic and inorganic contaminants were  
204 determined with at least three replicates. The calculated average concentrations are provided as  
205 mean  $\pm$  standard deviation (SD).  
206

## 207 **3. Results and Discussion**

208 Total organic carbon (TOC) content was high in both the superficial and buried soil  
209 layers. This is indicative of the low degree of decomposition and transformation of the precursor  
210 material and is comparable to the data on soils from the Yamal tundra (Ejarque, Abakumov,  
211 2016) and the Argentinian islands (Parnikoza et al, 2016). High TOC content is typical for the  
212 Antarctic Peninsula compared to soils of the Eastern Antarctic (Beyer et al, 1997, Mergelov et al,  
213 2017). While both were elevated, the TOC was higher in the superficial levels relative to the  
214 lower ones. Previous studies describe high variability in the TOC content from the soils of King  
215 George and Galindez Islands, mainly depending on the diversity of the ecotopes and the sources  
216 of organic matter (Abakumov, 2010, Parnikoza et al, 2016). Isolated (buried) soil spots are not  
217 connected with fresh sources of organic matter, explaining why the TOC content in these layers  
218 is lower. The carbon to nitrogen ratio was narrowest in SP1, which was affected by the scuas'  
219 activity (evidenced by remnants of nests). This is in line with previous studies that documented  
220 the well-pronounced ornithogenic effects on soil's nitrogen content (Simas et al, 2007, Parnikoza  
221 et al, 2016). Fine earth of soils investigated characterized by acid reaction, which is expected for  
222 soils of this region.

223 In terms of elemental composition, soil HAs are comparable with those previously  
224 reported for the Arctic and Antarctic soil. Current exposed organic layers contain HAs with  
225 higher carbon and nitrogen and lower oxygen content. Conversely, the HAs of isolated soil

226 patches show increased levels of oxidation. In comparison to soils of the tundra in the Komi  
227 Republic (Vasilevitch et al, 2018), HAs found in this study were more oxidized, comparable to  
228 those of the Kolyma Lowland (Lupachev et al, 2017) and previously published data from the  
229 Fildes Peninsula (Abakumov, 2017).

230 Data on the distribution of carbon species in HAs (fig. 3) and in bulk soil (fig. 4) samples  
231 indicated that aromatic compound content is generally lower than the alkyl components. This is a  
232 well-known peculiarity of the soils of the polar biome (McKniht et al, 1994, Beyer et al, 1997).  
233 At the same time, the degree of aromaticity of the isolated HAs is three fold higher than in the  
234 bulk organic matter. This suggests the presences of the humification process in the soils of  
235 Antarctica since humification involves increasing the aromatic compound content in  
236 macromolecules. This supports the classical humification hypothesis instead of new arguments,  
237 which are critical for this approach (Lehman, Kleber, 2015). Our data shows that SOM is on a  
238 continuum and HAs are the main acting constituent of this continuum; thereby confirming that  
239 this model is applicable even in Antarctica. The degree of aromaticity was higher in both isolated  
240 HAs and bulk soil samples from superficial levels compared to samples from isolated patches.  
241 Carbonyl/carboxyl/amide area (160 - 220 ppm) was more prevalent in the HAs of topsoils and  
242 less abundant in the organic matter of bulk samples (this region was presented mainly by  
243 carboxylic and amid carbon in the interval between 160 - 185 ppm) (Kniker, 2007). HAs  
244 extracted from SP1, located under the *Deshampsia antarctica*, exhibited wide peaks around 110 -  
245 140 ppm (H-aryl, C-aryl, olefinic-C) and at 140 - 160 ppm (O-aryl and N-aryl-C), while  
246 aromatic components of SP2 and SP3 were mainly represented by peaks between 110 - 140 ppm.  
247 This difference can be explained by the organic remnants of *Deshampsia antarctica* serving as  
248 the precursor for humification. All HA samples showed intensive areas of alkylic carbon (0 - 45  
249 ppm), aliphatic C and N, and methoxyl C (45 - 110 ppm), O-alkyl of carbohydrates and alcohols  
250 (60 - 95 ppm), and acetal and ketal carbon of carbohydrates (95 - 110 ppm). Carbon composition  
251 of the bulk samples was different from isolated HAs as evidenced mainly by the presence of  
252 alkyl carbon (0 - 45 ppm) and O- and N-alkyl carbon (45 - 110 ppm). Characteristic features of  
253 the bulk organic matter include carboxylic carbon and aryl compound content was low relative to  
254 isolated HAs. Only soils with prior ornithogenic interactions showed increases in carboxylic  
255 peaks, which corresponds well to data on relic ornithogenic soil (Beyer et al, 1997).

256 The C-alkyl : O-N-alkyl ratio used to indicate the degree of organic matter transformation  
257 was quite variable in all samples investigated. This can be caused by diversity in the origin and  
258 composition of the humification precursors. In case of comparisons with humic and fulvic acids  
259 of tundra soils (Vasilevitch et al, 2018), HAs of soils investigated are intermediated between  
260 HAs and fulvic acids of tundra Histosols with partially decomposed organic matter. These data  
261 are in line with a previous report (Hopkins et al, 2006) that showed soils of the Antarctic Dry  
262 Valleys have low alkyl-C : O-alkyl-C ratio using solid-state <sup>13</sup>C-NMR spectroscopy) and,  
263 therefore can serve as a labile, high-quality resource for micro-organisms. Beyer et al (1997)  
264 showed that both the CPMAS <sup>13</sup>C-NMR and the Py-FIMS spectra of the Terri-Gelic Histosol  
265 were dominated by signals from carbohydrates and alkylic compounds, which is corroborated by  
266 our findings. They also suggest that the <sup>13</sup>C-NMR data reflected decomposition of carbohydrates  
267 and enrichment of alkyl-C in deeper soil layers. In regards to the bulk SOM, this was true for  
268 SP2 and SP3 but not for SP1 that formed under the vascular plant *Deshampsia Antarctica*.

269 A representative electron spin resonance ESR spectrum of HAs is presented in fig 5 and the  
270 ESR parameters are similar to HAs and FAs of temperate soils (Senesi, 1990, Senesi et al, 2003).  
271 The spectra show a single, wide line with a g-factor ranging from 1,98890 to 1,99999,  
272 attributable to the presence of stable semiquinone free radicals in the HA-containing  
273 macromolecules (Table 5). The free radical content was higher in the superficial levels than in  
274 the isolated ones. This corresponds well with previous reports (Chukov et al, 2017, Abakumov et  
275 al, 2015) that connect the isolation of buried organic matter in the supra-permafrost with  
276 declining free radical content. This reveals the increased biochemical activity of HAs in topsoil.  
277 Compared to data from Lupachev (2017), the differences between exposed and isolated areas are  
278 less pronounced but, in general, the HAs of the Antarctic soils contain more unstable free  
279 radicals on average than the tundra soils of the Kolyma Lowland (Lupachev et al, 2017) and are  
280 comparable to the soils from the Yamal tundra (Chukov et al, 2017). Taken together, the free  
281 radical content found in our study was lower than in anthropogenically affected boreal and forest  
282 steppe soils of the East-European plains (Abakumov et al, 2018).

#### 283 4. Conclusions

284 High TOC content was fixed for the three studies representatives of Turbic Cryosols on  
285 King George Island, Northwest of the Antarctic Peninsula, Western Antarctic. High amounts of  
286 TOC are characteristic for both superficial and partially isolated soil materials. HAs contained  
287 three fold more aromatic carbon than bulk SOM, which indicates that humification appears and  
288 is active in soils of the Antarctic. Moreover, the amounts of aromatic carbon and carboxyl groups  
289 were higher in the HAs of the superficial layer, which is likely caused by the greater diversity of  
290 their organic precursors and more active humification than in sub-aerial conditions. The HAs of  
291 the superficial sample layers contained lower concentrations of free radicals, an indicator of  
292 active transformation in the topsoil. In general, the organic matter from partially isolated areas is  
293 less stable in terms of carbon species and free radical content. This likely results from the  
294 relative lack of fresh organic precursors and the different aeration and hydration conditions of  
295 stagnification bordering the permafrost table.

296 **Acknowledgments:** This work was supported by the Russian Foundation for Basic Research,  
297 project No 16-34-60010 and 18-04-00900 and the Saint Petersburg State University Internal  
298 Grant for the Modernization of Scientific Equipment No. 1.40.541.2017. Analyses were carried  
299 out at the Center for Magnetic Resonance and at the Center for Chemical Analysis and Materials  
300 Research of Research Park of St. Petersburg State University, Russia.

301  
302 The authors would like to thank Dr. A. Lupachev for assistance with field research and providing  
303 images for Figure 2 (2-1, 2-2 and 2-3).

304

305

307 **References**

- 308 Abakumov E. Vertical electric resistivity sounding of natural and anthropogenically affected  
309 cryosols of Fildes Peninsula, Western Antarctica. *Czech Polar Reports*, 7 (2), pp. 109-122, 2017  
310
- 311 Abakumov, E.V. The sources and composition of humus in some soils of West Antarctica  
312 *Eurasian Soil Science*, 43 (5), pp. 499-508, 2010
- 313 Abakumov, E.. Characterisation of humic acids, isolated from selected subantarctic soils by <sup>13</sup>C-  
314 NMR spectroscopy, *Czech Polar Reports*, 7 (1). 1-10, 2017
- 315 Abakumov, E., Maksimova, E., Tsibart, A. Assessment of postfire soils degradation dynamics:  
316 Stability and molecular composition of humic acids with use of spectroscopy methods *Land*  
317 *Degradation and Development* . Article in Press, 2018
- 318 Abakumov, E.V., Lodygin, E.D., Tomashunas V.M. <sup>13</sup>-C NMR and ESR characterization of  
319 Humic Substances, Isolates from Soils of two Siberian Arctic Islands, *International Journal of*  
320 *Ecology*. ID 390591, 2015
- 321 Abakumov, E , Mukhametova, N. Microbial biomass and basal respiration of selected Sub-  
322 Antarctic and Antarctic soils in the areas of some Russian polar stations, *Solid Earth*, 5. 705-712,  
323 2014
- 324 Abakumov, E.V., Parnikoza, I.Y., Vlasov, D.Y., Lupachev, A.V. (2016). Biogenic–abiogenic  
325 interaction in Antarctic ornithogenic soils. *Lecture Notes in Earth System Sciences*,  
326 (9783319249858), pp. 237-248.
- 327 Abakumov, E. V., Popov, A. I. Determination of the carbon and nitrogen contents and  
328 oxidizability of organic matter and the carbon of carbonates content in one soil sample,  
329 *Eurasian Soil Science*, 2:165–172, 2005
- 330 Abakumov, E.V., Fattakhova Yu.M. Structural composition of humic acids of ornitogenic soils  
331 on the data of <sup>13</sup>-C NMR resonance, *Russian Ornithology*, 1165: 2463-2466, 2014
- 332 Beyer, L; Knicker, H; Blume, H-P; Bölter, M; Vogt, B; Schneider. Soil organic matter of  
333 suggested spodic horiozns in relic ornitogenic soils of coastal Continental Antarctica (Casey  
334 Station, Wilkes Land) in comparison with that of spodic soil horizons in Germany, *Soil Science*,  
335 7. 518-527, 1997
- 336 Birkenmajer K. Geology and climatostratigraphy of Tertiary glacial and interglacial Successions  
337 on King George Island, South Shetland Islands (West Antarctica), *Ztrbl. Geol. Palaont.*, 1: 141–  
338 151, 1989
- 339 Bockheim J., Vieira G., Ramos M., López-Martínez, J., Serrano E., Guglielmin M., Wihelm K.,  
340 Nieuwendam A., . Climate warming and permafrost dynamics on the Antarctic Peninsula region,  
341 *Glob. Planet. Change*, 100: 215–223, 2013
- 342 Calace, N., Campanella, L., Paolis, F., and De Petronio, B. M. Characterization of Humic Acids  
343 Isolated from Antarctic Soils: *Int. J. Environ. Anal. Chem.*, 60: 71–78, 1995
- 344 Chapman, B., Roser, D., & Seppelt, R. <sup>13</sup>C NMR analysis of Antarctic cryptogam extracts.  
345 *Antarctic Science*, 6(3), 295-305, 1994



- 346 Chukov, S.N., Ejarque, E., Abakumov, E.V. Characterization of humic acids from tundra soils of  
347 northern Western Siberia by electron paramagnetic resonance spectroscopy, *Eurasian Soil*  
348 *Science*, 50 (1): 30-33, 2017
- 349 Davidson, E. A. and Janssens, I. A. Temperature sensitivity of soil carbon decomposition and  
350 feedbacks to climate change, *Nature*, 440: 165–173, 2006
- 351 Dziadowiec, H., Gonet, S., and Plichta, W. Properties of humic acids of Arctic tundra soils in  
352 Spitsbergen, *Polish Polar Res.*, (15), 71–81., 1994
- 353 Ejarque, E., Abakumov, E. Stability and biodegradability of organic matter from Arctic soils of  
354 Western Siberia: Insights from <sup>13</sup>C-NMR spectroscopy and elemental analysis (2016) *Solid*  
355 *Earth*, 7 (1), pp. 153-165, 2016
- 356 Fritz, M., Deshpande, B. N., Bouchard, F., Högström, E., Malenfant-Lepage, J., Morgenstern,  
357 A., Nieuwendam, A., Oliva, M., Paquette, M., Rudy, A. C. A., Siewert, M. B., Sjöberg, Y., and  
358 Weege, S. Brief Communication: Future avenues for permafrost science from the perspective of  
359 early career researchers, *The Cryosphere*, 9: 1715–1720, 2014
- 360 Haus, W., Serrano, E., Bockheim, J.G.. Soils and landforms from Fildes Peninsula and Ardley  
361 Island, Maritime Antarctica, *Geomorphology*, 225, 76–86, 2014
- 362 Hopkins D.W., Sparrow A.D., Elberling B., Gregorich E.G., Novis P.M., Greenfield L.G.,  
363 Tilston E.L., Carbon, nitrogen and temperature controls on microbial activity in soils from an  
364 Antarctic dry valley, *Soil Biology and Biochemistry*, 38: 3130-3140, 2006
- 365 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C.-L.,  
366 Schirrmeyer, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B.,  
367 Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P. Estimated stocks of circumpolar  
368 permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*,  
369 11: 6573–6593, 2014
- 370 ICCP – Intergovernmental panel on Climate Change: *Climate Change 2007.*, Cambridge,  
371 University Press, United Kingdom and New York, NY, USA, 2007
- 372 Kniker H. How does fire affect the nature and stability of soil organic nitrogen and carbon ?  
373 *Biogeosciences*, 2007, 86: 91-118, DOI 10.1007/s10533-007-9104-4, 2006
- 374 Lechman J., Kleber M. The contentious nature of soil organic matter. *Nature*, 528. 60-68, 2015
- 375 Lodygin, E.D., Beznosikov, V.A. and Vasilevich, R.S. Molecular composition of humic  
376 substances in tundra soils (<sup>13</sup>C-NMR spectroscopic study). *Eurasian Soil Sc.* (47) (5), 400–406,  
377 2014
- 378  
379 Lupachev, A., Abakumov, E., Gubin, S. The influence of cryogenic mass exchange on the  
380 composition and stabilization rate of soil organic matter in cryosols of the kolyma lowland  
381 (North Yakutia, Russia), *Geosciences (Switzerland)*, 7 (2), paper № 24, DOI:  
382 10.3390/geosciences7020024, 2017
- 383 Management Plan for Antarctic Specially Protected Area No. 125, Measure 6, Annex., Antarctic  
384 Treaty Secretariat, 2009
- 385 Mausbacher R., Muller J., Munnich M., Schmidt R. Evolution of postglacial sedimentation in  
386 Antarctic lakes (King Georges Island), *Z. Geomorphologie*, 33: 219–234, 1989

- 387 McKnight Diane M. , Andrews Edmund D. , Spaulding Sarah A. , Aiken George R. , Aquatic  
388 fulvic acids in algal-rich antarctic ponds, *Limnology and Oceanography*, 39, 1994
- 389 Mergelov, N., Mueller, C.W., Prater, I., Shorkunov, I., Dolgikh, A., Zazovskaya, E., Shishkov,  
390 V., Krupskaya, V., Abrosimov, K., Cherkinsky, A., Goryachkin, S. Alteration of rocks by  
391 endolithic organisms is one of the pathways for the beginning of soils on Earth. *Scientific*  
392 *Reports*, 8 (1), paper № 3367, 2018
- 393 Michel, R.F.M., Schaefer, C.E.G.R., López-Martínez, J., Simas, F.N.B., Haus, N.W., Serrano,  
394 E., Bockheim, J.G.. Soils and landforms from Fildes Peninsula and Ardley Island, Maritime  
395 Antarctica. *Geomorphology*. 225: 76-86. doi:10.1016/j.geomorph.2014.03.041, 2014
- 396 Navas, A., López-Martínez, J., Casas, J., Machín, J., Durán, J.J., Serrano, E., Cuchi, J.A., Mink,  
397 S.. Soil characteristics on varying lithological substrates in the South Shetland Islands, maritime  
398 Antarctica. *Geoderma* 144 :123–139. doi:10.116/j.geoderma.2007.10.01, 2008
- 399 Parnikoza, I., Abakumov, E., Korsun, S., Klymenko, I., Netsyk, M., Kudinova, A., Kozeretska, I.  
400 Soils of the argentine islands, antarctica: Diversity and characteristics *Polarforschung*, 86 (2), pp.  
401 83-96. 2016
- 402 Rakusa-Suszczewski S. King George Island — South Shetland Islands, Maritime Antarctic. In:  
403 Beyer L., Bölder M. (Eds.), *Geoecology of Antarctic Ice Free Coastal Landscapes*. Springer  
404 Verlag, Berlín, 23–40, 2002.
- 405 Polyakov V., Orlova K., Abakumov, E. Evaluation of carbon stocks in the soils of Lena River  
406 Delta on the basis of application of “dry combustion” and Tyurin’s methods of carbon  
407 determination. *Biological Communications*. 62(2): 67–72, 2017.
- 408 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S. Soil  
409 organic carbon pools in the northern circumpolar permafrost region, *Global Biogeochem.* , 23,  
410 1–11, 2009  
411
- 412 Schnitzer, M. Organic matter characterization, in: *Methods of soil analysis, Part 2, chemical and*  
413 *microbiological properties*, Agronomy monograph no. 9, edited by: Page, B., Miller, R., and  
414 Keeney, D., Soil Science Society of America, Madison, 581–594, 1982.
- 415 Simas, FNB., Schaefer, CEG.R., Melo, V.F., Albuquerque-Filho, MR., Michel, RFM., Pereira,  
416 VV., Gomes, MRM., da Costa, L.M. Ornithogenic cryosols from Maritime Antarctica:  
417 Phosphatization as a soil forming process, *Geoderma*. 138: 191-203, 2007
- 418 Senesi N, D’Orazio V, Ricca G. Humic acids in the first generation of eurosoils, *Geoderma*,  
419 116, 3-4: 325-344, 2003
- 420 Senesi N. Molecular and quantitative aspects of the chemistry of fulvic acids and its interactions  
421 with mineral ions and organic chemicals. *Analytica Chimica Acta*. 232: 51-75, 1990
- 422 Schuur, E.A.G., McGuire, A.D., Grosse, G., Harden, J.W., Hayes, D.J., Hugelius, G., Koven,  
423 C.D. and Kuhry, P. Climate change and the permafrost carbon feedback. *Nature*, 520, 171–179,  
424 2015
- 425 Smellie J.L., Pankhurst R.J., Thomson M.R.A., Davies R.E.S. The Geology of the South  
426 Shetland Islands: VI. Stratigraphy, Geochemistry and Evolution, *British Antarctic Survey*  
427 *Scientific Reports* 50: 87, 85, 1984

- 428 Vasilevich, R., Lodygin, E., Beznosikov, V., Abakumov, E. Molecular composition of raw peat  
429 and humic substances from permafrost peat soils of European Northeast Russia as climate  
430 change markers. *Science of the Total Environment*, 615, 1229-1238, 2018
- 431 Wen, J., Xie, Z., Han J., Lluberas A.. Climate, mass balance and glacial changes on small dome of  
432 Collins Ice Cap, King George Island, Antarctica, *Antarct. Res.*, 5, 52–61, 1994
- 433 WRB. World Reference Base for Soil Resources. International Soil Classification System for  
434 Naming Soils and Creating Legends for Soil Maps; World Soil Resources Reports No. 106;  
435 FAO: Rome, Italy, 2014.
- 436 Zaccone, C., Miano, T. M., and Shotyk, W. Qualitative comparison between raw peat and related  
437 humic acids in an ombrotrophic bog profile: *Org. Geochem.*, 38: 151–160, 2007
- 438 Zech, W., Senesi, N., Guggenberger, G., Kaiser, K., Lehmann, J., Miano, T. M., Miltner, A., and  
439 Schroth, G. Factors controlling humification and mineralization of soil organic matter in the  
440 tropics: *Geoderma*, 79, 117–161, 1997.
- 441 Zubrzycki, S., Kutzbach, L. and Pfeiffer, E.-M. Permafrost affected soils and their carbon pools  
442 with a focus on the Russian Arctic. *Solid Earth*, 5, 595–609. 2014
- 443
- 444
- 445
- 446
- 447

448 Table. 1. Basic characteristics of soils

Sample	TOC, %	N, %	C/N	pH <sub>H2O</sub>	pH <sub>CaCl2</sub>	Color
1 O	27.63±0.23	5.18±0.42	5.33	6.35	5.30	10 YR 4/7
2 [CRH]	19.05±0.15	2.20±0.05	8.66	5.67	4.89	2.5 YR 4/4
3 O	20.04±0.17	1.16±0.09	17.13	4.80	4.80	10 YR 4/4
5 [CRH]	12.33±0.24	0.78±0.09	15.80	4.70	4.50	2.5 YR 4/3
4 O	10.16±0.09	0.84±0.07	11.98	4.90	4.21	10 YR 5/3
6 [CRH]	6.66±0.07	0.81±0.09	8.20	4.70	4.35	2.5 YR 5/3

449

450

451

452 Table 2. Elemental composition (%) and atomic ratios in HAs. Data presented in atomic values.

Sample №	C	N	H	O	C/N	H/C	O/C
1	49.53±0.56	5.55±0.07	6.90±0.11	38.02±0.64	8.92	0.13	0.76
2	47.14±0.45	4.30±0.06	6.79±0.09	41.77±0.21	10.96	0.14	0.88
3	45.55±0.32	5.14±0.09	5.80±0.09	43.51±0.35	8.86	0.12	0.95
4	43.77±0.24	4.72±0.11	6.90±0.08	44.61±0.21	9.27	0.15	1.01
5	49.99±0.41	4.78±0.08	6.56±0.08	38.67±0.34	10.45	0.13	0.77
6	44.45±0.034	3.99±0.07	6.77±0.10	44.79±0.25	11.14	0.15	1.01
P, One way Anova, superficial/buried	0.14	<b>0.05</b>	0.29	<b>0.05</b>	n.d.	n.d.	n.d.

453

454

455

456 Table 3. Carbon species integration in molecules of the HAs, %

Sample №	Carbonyl/ carboxyl/ amide	Aryl- olefine	O-N alkyl	Calkyl	Calkyl/O-N alkyl	Caryl/Calkyl
Chemical shift, ppm	220-160	160-110	110-45	45-0		
1	11,38	33,59	39,86	14,18	0.35	2.36
2	10,75	30,45	31,86	26,05	0.81	1.16
3	19,24	23,34	29,54	27,85	0.94	0.83
4	16,48	21,42	34,23	27,87	0.81	0.77
5	16,75	33,40	29,12	20,71	0.71	1.61
6	14.39	26.86	40.07	18.68	0.46	1.43
P, One way Anova, superficial/buried	<b>0.02</b>	<b>0.03</b>	<b>0.02</b>	0.73	n.d.	n.d.

457

458

459

460 Table 4. Carbon species integration in molecules of the bulk organic matter, %

Sample №	Carbonyl/ carboxyl/ amide	Aryl- olefine	O-N alkyl	Calkyl	Calkyl/O -N alkyl	Caryl/Calkyl
Chemical shift, ppm	220-160	160-110	110-45	45-0		
1 =113=O	7.24	11.37	46.20	35.19	0.76	0.32
2 113-Ch	18.23	10.29	40.59	30.89	0.76	0.33
3 123 O	7.34	20.48	55.12	17.06	0.31	1.20
4 123 Ch	9.34	11.27	49.50	29.90	0.60	0.37
6 149 O	5.72	13.84	62.22	18.22	0.29	0.75
6 149 Ch	22.95	9.89	46.92	20.24	0.43	0.48
P, One way Anova, superficial/buried	0.53	<b>0.01</b>	<b>0.05</b>	0.56	n.d.	n.d.

461

462

463

464 Table. 5. Mass concentration of free radical in humic acids

Soil horizon	Mass concentration of free radical, $10^{15}$ spin*g <sup>-1</sup>	g-factor
1	3.67	2.0314
2	3.04	2.3150
3	3.51	2.0314
4	2.13	2.0303
5	6.10	2.0310
6	5.86	2.0314

465

466


467

468

469



470



471

472 Fig. 1.

473



474


1

2

3

475 Fig. 2

476






477

478

479 Fig. 3

480




481

482

483 Fig. 4

484



485

486 Fig. 5.

487

488

489 Fig. 1. Location of the Fildes peninsula

490 Fig.2. Soil morphology

491 Figure 3. <sup>13</sup>C NMR spectras of the HAs, isolated from soils (1-6 – according table 1)

492 Figure 4. <sup>13</sup>C NMR spectras of bulk organic matter of soils ((1-6 – according table 1)

493 Figure 5. Typical ESR spectrum of humic substances investigated

494 Table. 1. Basic characteristics of soils

495 Table 2. Elemental composition (%) and atomic ratios in HAs

496 Table 3. Carbon species integration in molecules of the HAs, %

497 Table 4. Carbon species integration in molecules of the bulk organic matter, %

498 Table. 5. Mass concentration of free radical in humic acids

499