

Myrmekite and strain weakening in granitoid mylonites

Alberto Ceccato^{1*}, Luca Menegon², Giorgio Pennacchioni¹, Luiz Fernando Grafulha Morales³

¹ Department of Geosciences, University of Padova, 35131 Padova, Italy

² School of Geography, Earth and Environmental Sciences, University of Plymouth, PL48AA Plymouth, UK

³ Scientific Centre for Optical and Electron Microscopy (ScopeM) - ETH Zürich, Switzerland

Correspondence to: Alberto Ceccato (alberto.ceccato.2@phd.unipd.it)

* Now at: School of Geography, Earth and Environmental Sciences, University of Plymouth, PL48AA Plymouth, UK

Abstract. At mid-crustal conditions, deformation of feldspar is mainly accomplished by a combination of fracturing, dissolution/precipitation and reaction-weakening mechanisms. In particular, K-feldspar

10 is reaction-weakened by formation of strain-induced myrmekite - a fine-grained symplectite of plagioclase and quartz. Here we investigate with EBSD the microstructure of a granodiorite mylonite, developed at 420-460 °C during cooling of the Rieserferner pluton (Eastern Alps), to assess the microstructural processes and the role of weakening associated with myrmekite development. Our analysis shows that the crystallographic orientation of the plagioclase of pristine myrmekite was 15 controlled by that of the replaced K-feldspar. Myrmekite nucleation resulted in both grain size reduction and ordered phase mixing by heterogeneous nucleation of quartz and plagioclase. The fine grain size of sheared myrmekite promoted grain size-sensitive creep mechanisms including fluid-assisted grain boundary sliding in plagioclase, coupled with heterogeneous nucleation of quartz within 20 creep cavitation pores. Flow laws calculated for monomineralic quartz, feldspar, and quartz + plagioclase aggregates (sheared myrmekite), show that during mylonitization at 450 °C grain-size-sensitive creep in sheared myrmekite accommodated strain rates several orders of magnitude higher than monomineralic quartz layers deforming by dislocation creep. Therefore, diffusion creep and grain size-sensitive processes contributed significantly to bulk rock weakening during mylonitization. Our results have implications for modelling the rheology of the mid-upper continental (felsic) crust.

25 1. Introduction

Localization of ductile strain within rocks arises from weakening associated with grain size refinement processes by dynamic recrystallization, metamorphic reactions, and microfracturing (e.g. de Bresser et al., 2001 and reference therein). Grain size reduction, accompanied by phase mixing in polymimetic rocks at high strains, commonly results in a switch of deformation mechanism from 30 grain-size-insensitive (GSI) to grain-size-sensitive (GSS) creep – one of the most effective strain weakening mechanisms within shear zones (Kruse and Stünitz, 1999; Kilian et al., 2011; Menegon et

al., 2013). Feldspars locally form the load-bearing framework of continental crust rocks (Handy, 1994). At mid-crustal conditions, feldspar deformation mainly occurs by microfracturing and dissolution/precipitation processes, typically associated with metamorphic reactions (Behrmann and Mainprice, 1987; Ree et al., 2005; Viegas et al., 2016). K-feldspar is commonly replaced by myrmekite – a fine-grained symplectic aggregate of quartz and plagioclase (Becke, 1908; Vernon, 1991). Myrmekite replacement is related either to K-feldspar chemical instability ([Cesare et al., 2002](#)), in some cases involving the presence of local metasomatic fluids ([Phillips, 1980](#)[Cesare et al., 2002](#)), or triggered by intracrystalline strain in K-feldspar during deformation (Simpson and Wintsch, 1989; Menegon et al., 2006). This replacement is acknowledged as a weakening mechanism during ductile deformation of granitoid rocks (Simpson and Wintsch, 1989; Tsurumi et al., 2003; Menegon et al., 2006; De Toni et al., 2016) and is particularly remarkable when it affects coarse-grained pegmatite (LaTour and Barnett, 1987; Pennacchioni, 2005; Pennacchioni and Zucchi, 2013). Deformation and shearing of myrmekite result into a fine-grained plagioclase + quartz aggregate, that is manifestly weaker than original coarse K-feldspar (Tsurumi et al. 2003; Ree et al., 2005; Ciancaleoni and Marquer, 2006; Viegas et al. 2016). Fine grain size and the local occurrence of fluid promote phase mixing processes and the development of ultramylonites (Vernon, 1991; Kilian et al., 2011, Czaplińska et al., 2015). The recognition of the key role of myrmekite in strain localization has not been accompanied with a quantitative analysis of the deformation mechanisms within myrmekite-derived fine-grained plagioclase + quartz aggregates. Here we present the detailed analysis of myrmekite evolution from breakdown of K-feldspar into plagioclase and quartz, grain size reduction and strain localization in the mylonites of the Rieserferner granitoid pluton (Eastern Alps). In this pluton, ductile shear zones nucleated along joints that were locally filled with quartz- and epidote- veins, during post-magmatic cooling (Ceccato et al., 2017; Ceccato and Pennacchioni, 2018). The progressive development of granodiorite mylonite included consumption of K-feldspar by myrmekite leading to increasingly interconnected, fine-grained plagioclase + quartz layers. Selected microstructures of granodiorite mylonite have been analysed to characterize: (i) the process of myrmekite nucleation; (ii) the deformation mechanisms during myrmekite shearing and transition to plagioclase + quartz aggregates; (iii) the deformation mechanisms of pure-quartz layers; (v) the deformation mechanisms of K-feldspar porphyroclasts and of K-feldspar neoblasts during mylonitization. Furthermore, the application of mixed flow-laws of the aforementioned deformation mechanisms for polymineralic aggregates allowed the quantification of the extent of rheological weakening resulting from the deformation of myrmekite.

2. Geological setting and field description

The tonalitic-granodioritic Rieserferner pluton (Eastern Alps) (Bellieni, 1978) emplaced at ~15 km depth (0.4 GPa; Cesare et al., 2010) into the Austroalpine nappe system at 32 Ma (Romer and Siegesmund, 2003). During post-magmatic cooling, a main set of ductile shear zones developed 5 exploiting precursor shallowly ESE-dipping joints and the locally associated quartz and epidote veins filling the joints (Ceccato, 2018; Ceccato and Pennacchioni, 2018). Nucleation of ductile shear zones on precursor tabular layers (e.g. dykes, veins) or “surface” discontinuities (joints) has been observed in many granitoid plutons (Adamello: Pennacchioni, 2005; Sierra Nevada: Segall and Simpson, 1986; Pennacchioni and Zucchi, 2013) and in meta-granitoid units (Pennacchioni and Mancktelow, 2007, 10 2018, and reference therein). The temperature of ductile shearing in the Rieserferner has been estimated at 420-460 °C based on thermodynamic modelling (Ceccato, 2018). Where associated with precursor joints filled with epidote, ductile shearing resulted in cm-thick heterogeneous shear zones with a sigmoidal-shaped foliation in the host granodiorite (Ceccato and Pennacchioni, 2018).

3. Sample description and microstructure

15 Polished thin sections of granodiorite mylonite were prepared, from rock chips cut parallel to the lineation and perpendicular to the shear plane (XZ plane of finite strain ellipsoid), for the study of the microstructure and of crystallographic preferred orientations (CPO). The microstructural and CPO analysis conducted via EBSD were complemented with microchemical analyses performed with an electron microprobe. Mineral abbreviations after Kretz (1983). Description of methods and analytical 20 conditions are reported in Appendix A.

The Rieserferner granodiorite consists of quartz, plagioclase, K-feldspar, biotite, allanite/epidote, hornblende, apatite and titanite. The magmatic plagioclase displays oscillatory zoning with a range in composition between An₅₈ (core) to An₃₂ (rim), and is arranged in glomeroclasts, included in K-feldspar (Or₉₃ – Ab₇). Various grain-size reduction mechanisms accompanied the development of a 25 mylonitic foliation in the granodiorite: (i) recrystallization of quartz and biotite (Fig. 1a,b); (ii) formation of myrmekite after K-Feldspar (Fig. 1c,d) and (iii) microfracturing of feldspar and (iv) formation of plagioclase (An₂₉Ab₇₁Or_{<1}) – titanite – muscovite symplectite at biotite-plagioclase boundaries (Pennacchioni et al., 2006; Johnson et al., 2008). Pristine myrmekite makes transition to fine grained aggregates of dominant plagioclase + quartz extended into the foliation (Fig. 1b,e). The 30 mylonitic foliation is defined by alternating layers of: (i) monomineralic quartz; (ii) plagioclase (An₂₆Ab₇₄Or_{<1}), quartz, K-feldspar; and (iii) biotite – recrystallized biotite/plagioclase (Fig. 1a).

Syn-kinematic K-feldspar neoblasts ($Or_{96} - Ab_4$) are found in strain shadows around porphyroclasts or dilatant fractures, and are in turn locally replaced by myrmekite (Fig. 1d). With increasing strain, there is a decrease in volume percentage of K-feldspar from 19 vol% (undeformed rock and protomylonite), to 1-6 vol% (scattered porphyroclasts in mylonite and ultramylonite) (Figs. 2). As

5 counterbalance, there is an increase in volume percentage of fine-grained myrmekite and derived plagioclase + quartz aggregates, from 3% (undeformed rock and protomylonite) to as much as 13% (mylonite and ultramylonite) (Figs. 2b). Ultramylonites consist of a fine-grained (ca. 10 μm grain size) well-mixed matrix of quartz, plagioclase, biotite, epidote, K-feldspar, titanite, apatite \pm garnet \pm white mica.

10 4. EBSD and cathodoluminescence analysis

EBSD maps and crystallographic orientation data are reported in Figs. 3 and 4. Results of phase spatial distribution analysis are reported in Fig. 5. Results of image analysis of grain size and shape are reported in Figs. 6 and 7.

4.1 Pristine myrmekite

15 Characteristics of pristine myrmekite are: (i) the preferential development along grain boundaries parallel to the mylonitic foliation, despite locally mantling entirely the K-feldspar porphyroblast (Fig. 1b-e); (ii) the lobate shape protruding into the K-feldspar (Fig. 1c); (iii) the monoecrystalline-single crystal structure of plagioclase within each lobe (20 to 50 μm in size: Fig. 7a), embedding vermicular quartz (sections up to 3 μm in equivalent size: Fig. 6a); (iv) the rather constant spacing between the 20 quartz vermicules of about 3-5 μm across the entire lobe; (v) the preferential elongation of the quartz vermicules orthogonal to the myrmekite/K-feldspar boundary.

The EBSD analysis of the crystallographic relationships between the K-feldspar and the replacing myrmekitic plagioclase and quartz shows that: (i) K-feldspar and plagioclase commonly show similar crystallographic orientations ($(100)_{\text{Kfs}} \parallel [100]_{\text{Plg}}$, $(010)_{\text{Kfs}} \parallel (010)_{\text{Plg}}$, $[001]_{\text{Kfs}} \parallel (001)_{\text{Plg}}$; Fig. 3b, c); (ii) 25 the quartz vermicules do not share any crystallographic plane or direction with K-feldspar or myrmekitic plagioclase (Fig. 3b-c-d); (iii) quartz vermicules do not show any obvious CPO, but they within a myrmekite lobe usually have similar-a crystallographic preferred orientation within a myrmekite lobe (see encircled clusters in Fig. 3d) (Abart et al., 2014); and (iv) Dauphiné and Albite twins are occasionally observed in quartz and plagioclase, respectively.

The plagioclase of myrmekite lobes exhibits rare low angle boundaries (misorientations $>2^\circ$, $>5^\circ$) that abut against the quartz vermicules (Figs. 3a and 4a). However, The internal distortion of myrmekitic plagioclase is very small ($<1^\circ$; Fig. SOM1a).

4.2 Sheared myrmekite: plagioclase + quartz aggregates

5 Shearing of myrmekite gave rise to plagioclase + quartz aggregates (\pm rare K-feldspar and biotite) elongated into the foliation (Fig. 1e). These aggregates are referred hereafter as sheared myrmekite.

Quartz grains of sheared myrmekite occur either as isolated single grains at triple/quadruple junctions of plagioclase grains or, less commonly, as polycrystalline aggregates elongated normal to the foliation (Fig. 4a). The quartz grain size is around $3\text{ }\mu\text{m}$ (Area B in Fig. 4a; Fig. 6b), but locally increases to

10 $>10\text{ }\mu\text{m}$ (Area C in Fig. 4a; Fig. 6c). Individual grains show polygonal, equant shapes (aspect ratio AR, $1.5 < \text{AR} < 1.75$) or a weak shape preferred orientation (SPO) oriented at low angle to the local mylonitic foliation (Fig. 6e). Quartz grains within sheared myrmekite have no CPO (Fig. 4b), show little internal distortion and rarely show low angle boundaries with scattered misorientation axis distribution (Fig. 4c). Misorientation angle distribution for correlated pairs displays higher frequency 15 for misorientation $< 15^\circ$ and at 60° than a random-pair distribution (Fig. 4d). The uncorrelated misorientation angle distribution approaches the random-pair distribution.

Plagioclase grains (average grain size of about $7\text{ }\mu\text{m}$: Figs. 7b-c) are mainly polygonal and range in shape from almost equant to elongated ($1.75 < \text{AR} < 2$). Elongated grains define an SPO almost parallel to the local mylonitic foliation (Fig. 7d for Area B in Fig. 4a). Plagioclase grains do not show any 20 obvious CPO (Fig. 4e), they display little internal distortion and rare low angle boundaries. The low and high angle misorientation axes are almost uniformly distributed in crystal directions (Fig. 4f). Even though very close to random-pair distribution, correlated misorientation distribution exhibits two weak peaks at very low angles ($<5\text{--}10^\circ$) and close to 180° (Fig. 4g). Misorientations $<70^\circ$ occurs with slightly higher frequency than the random-pair distribution. Albite-twins, and related 180° misorientations are 25 rarely observed inside new grains (Figs. 3a-4a). In cathodoluminescence (CL) both myrmekitic plagioclase and quartz have a CL grey shade similar to the surrounding non-myrmekitic plagioclase and quartz (similar to Hopson and Ramseyer, 1990) (Fig. 1f,h).

4.3 K-feldspar aggregates in strain shadows

K-feldspar neoblasts occur in strain shadows around feldspar porphyroclasts, as well as dispersed 30 within the sheared myrmekite (Figs. 3a, SOM2a; Area E in Fig. 4a). In strain shadows, the orientation of (100), (010), [001] planes of the neoblasts is similar to that of the porphyroblast (Fig. SOM2). The

grain size of K-feldspar dispersed within sheared myrmekite is ca. 7 μm , comparable to that of the plagioclase in the surrounding sheared myrmekite (Fig. SOM2a-f). The analysed K-feldspar aggregate shows a CPO for (010) planes close to the Y kinematic axis (Fig. SOM2d), which is similar to the orientation of (010) in the adjacent porphyroblast. Misorientation axis/angle distributions show very

5 few scattered data without any clear clustering (Fig. SOM2e).

The CL imaging of K-feldspar grains and porphyroblasts highlights a complex microstructure, in which porphyroblasts exhibit a homogeneous bright shade overprinted by a complex low-grey CL pattern (Fig. SOM3). K-feldspar grains in sheared myrmekite and tails around porphyroblasts show a homogeneous low-grey CL signature (Fig. SOM3). K-feldspar aggregates elongated parallel to the

10 foliation and enveloped by sheared myrmekite are characterized by bright irregularly-shaped K-feldspar cores (porphyroblasts) surrounded by low-grey shade K-feldspar.

4.4 Quartz layers along foliation

Monomineralic quartz layers defining the mylonitic foliation (Figs. 3, 4 and SOM4) show a variable grain size and a shape preferred orientation (SPO), inclined to the foliation consistent with the sense

15 of shear. Dauphiné twin boundaries are widespread (red boundaries in Fig. SOM4a). The quartz c-axis

CPO defines an asymmetric Type-I girdle roughly normal to the local mylonitic foliation (Fig. SOM4b).

The pole figures of c-axis and $\langle a \rangle$ directions show maxima close to the Y and to the X kinematic directions, respectively. Misorientation axis distribution for low angle misorientation ($<10^\circ$)

exhibits a wide maximum close to c-axis and $\langle \pi - \pi' \rangle$ directions in crystal coordinates. They are

20 preferentially oriented slightly off-set from the Y-kinematic direction in sample coordinates (Fig. SOM4c).

High angle misorientation axis distributions do not show any clear systematic pattern, except for misorientations around 60° . Misorientation angle distribution (Fig. SOM4d) shows two peaks at very low angle misorientations ($<10^\circ$) and around 60° for correlated misorientations. Un-correlated misorientation angle distribution is close to the random-pair distribution.

25 Quartz layers along the foliation show a variable grain size, usually ranging between 10 μm and 120 μm , mimicking a bimodal grain size distribution with maxima centred respectively at 20-35 μm and 50-70 μm (Figs. 6f and SOM5). The coarser grain sizes ($>40 \mu\text{m}$) is observed close to the centre of

quartz layers. These grains are usually characterized by subgrains ranging in size between 20 and 35 μm . The smaller grain size ($<40 \mu\text{m}$) commonly envelope the coarser grains, in addition to prevail at

30 the boundary between monomineralic quartz layers and sheared myrmekite, or around feldspar porphyroblasts (Figs. 3, 4 and SOM5e). CPOs and misorientation data of coarser grains do not differ

from those of finer grains. In CL images the quartz layers display an overall homogeneous signature, with lower-grey shades close to inclusions and layer boundaries (Fig. SOM3).

5. Image Analyses

The results of image analysis of EBSD phase maps indicate that pristine and sheared myrmekite have 5 the same phase ratio with ca. 18 vol% of quartz. We have analysed the phase spatial distribution of plagioclase and quartz in both pristine and sheared myrmekite to define their deviation from random distribution, either towards a clustered or an anticlustered distribution (Heilbronner and Barrett, 2014). Phase spatial distribution in deformed bimodal aggregates in shear zones is interpreted to reflect the activity of specific deformation mechanisms (Kruse and Stünitz, 1999; Menegon et al., 2013). 10 Dislocation creep usually results in either monomineralic aggregates or clustered phase distributions (Heilbronner and Barrett, 2014). Diffusion creep in polymimetic aggregates is commonly accompanied by heterogeneous phase nucleation that promotes phase mixing and a high degree of anticlustering in phase distribution (Kilian et al., 2011; Menegon et al., 2013). Phase spatial distribution analysis of a two-phase aggregate compares the cumulative lengths of phase boundaries (boundaries 15 between grains of a different phase) and of grain boundaries (boundaries between grains of the same phase) with those expected for a random distribution of the two phases. We have considered three types of boundaries: (i) plagioclase – plagioclase grain boundaries; (ii) quartz – quartz grain boundaries; and (iii) plagioclase – quartz phase boundaries. The results show that (Fig. 5), in pristine and sheared myrmekite: (i) the surface area fraction of quartz ranges between 0.55-0.75 and 0.55-0.65, 20 respectively; (ii) quartz-quartz grain boundaries occur with a probability lower than for a random distribution indicative of an anticlustered distribution; (iii) plagioclase-plagioclase grain boundaries occur with a higher probability than for a random distribution indicative of a more clustered distribution; (iv) plagioclase + quartz aggregates display an ordered/anticlustered distribution, with plagioclase – quartz phase boundaries occurring with higher probability than for random distribution 25 of phases.

6. Discussion

6.1 Formation and shearing of myrmekite

6.1.1 Crystallographic relationship between K-feldspar and myrmekitic phases

The EBSD analysis indicates that the K-feldspar and the overgrowing myrmekitic plagioclase have a 30 similar crystallographic orientation, though with some scattering (Fig. 3b; Wirth and Voll, 1987). This

suggests the occurrence of a topotactic replacive process where $(100)_{\text{Kfs}} \parallel [100]_{\text{Plg}}$, $(010)_{\text{Kfs}} \parallel (010)_{\text{Plg}}$, $[001]_{\text{Kfs}} \parallel (001)_{\text{Plg}}$. The scatter in crystallographic orientation between K-feldspar and myrmekitic plagioclase is interpreted to have resulted from deformation during and after myrmekite formation (see section 6.1.2). The crystallographic orientation of myrmekitic plagioclase and quartz was not controlled by neighbour plagioclase or quartz grains previously in contact with the K-feldspar, differently from what reported by other authors (Stel and Breedveld, 1990; Abart et al., 2014). As observed by Abart et al. (2014), the different myrmekite quartz vermicules have a similar crystallographic orientation. The anticlustered phase spatial distribution of pristine myrmekite is related to the process of heterogeneous phase nucleation during myrmekite formation (Wirth and Voll, 10 1987).

6.1.2. Transition from pristine- to sheared myrmekite (plagioclase + quartz aggregates)

The transition from pristine to sheared myrmekite was a dynamic process and here we try to constrain the processes involved as inferred from microstructural changes. These microstructural changes include: (i) randomization of plagioclase CPOs observed in pristine myrmekite; (ii) refinement of 15 plagioclase grain size distribution from scattered in pristine (3 to 50 μm) to homogeneous and centred at 7 μm in sheared myrmekite (Figs. 3, 4 and 7a); (iii) coarsening of quartz grains from $<3 \mu\text{m}$ vermicules to as large as 10 μm rounded-to-polygonal grains in sheared myrmekite (Fig. 6). These processes are probably related to the minimization of interfacial energy in the vermicular microstructure of the pristine myrmekite (e.g. Odashima et al., 2007; Dégi et al., 2010). Quartz grain 20 coarsening occurred as a consequence of annealing of the pristine vermicular microstructure after the reaction front moved further into the K-feldspar (Fig. 3a). Quartz coarsening implies simultaneous grain size refinement of plagioclase, which probably involved both annealing and micro-fracturing, as suggested by misorientation analysis on the few low and high misorientation angle boundaries and CPO randomization (Figs. 3, SOM1a). Microfractures could have originated from stress 25 concentrations within the 3-D geometrically/mechanically composite structure of myrmekite (see figure 2 of Hopson and Ramseyer, 1990; Dell'Angelo and Tullis, 1996; Xiao et al., 2002). Annealed myrmekite are then sheared along the mylonitic foliation from the contractional sites around the K-feldspar porphyroblast.

6.2. Deformation mechanisms in the Rieserferner mylonites

6.2.1. Sheared myrmekite

Plagioclase and quartz of sheared myrmekite both display: (i) a weak CPO; (ii) rare low angle boundaries without systematic pattern of misorientation axis distribution; (iii) correlated and uncorrelated misorientation angle distributions close to the theoretical random-pair distribution. All these features suggest very limited deformation by dislocation creep in both minerals (Kruse et al., 2001; Miranda et al., 2016). In addition, the sheared myrmekite show: (i) fine-grained plagioclase and quartz with polygonal, equant to slightly elongated shape ($AR < 2$); (ii) aligned grain boundaries (over the scale of several grain diameters) and common triple/quadruple-junctions; (iii) anticlustered spatial distribution of plagioclase and quartz. These microstructural features are consistent with GSS creep, including fluid-assisted grain boundary sliding (GBS) (White, 1977; Stünitz and Fitz Gerald, 1993; Fliervoet et al., 1997; Jiang et al., 2000; Wheeler et al., 2001; Lapworth et al., 2002; Bestmann and Prior, 2003; Kilian et al., 2011; Menegon et al., 2013). The occurrence of quartz in triple-quadruple junctions and quartz aggregates elongated orthogonal to the foliation suggest the activity of creep cavitation and heterogeneous quartz nucleation during GSS creep of plagioclase (Fusseis et al., 2009; Herwegen et al., 2011; Kilian et al., 2011). Heterogeneous phase nucleation in creep cavities led to the anticlustered phase spatial distribution observed in sheared myrmekite (Fig. 5) (Hiraga et al., 2013; Menegon et al., 2015). The constant plagioclase grain size of sheared myrmekite may then result from the combination of initial spacing between quartz vermicules in pristine myrmekite, diffusion creep processes and second-phase grain-boundary pinning during shearing (Herwegen et al., 2011). GSS processes, phase mixing and second-phase grain-boundary pinning inhibit grain growth and stabilizes grain size, hindering the efficiency of dynamic recrystallization processes and self-sustaining the activity of GSS processes.

6.2.2. K-feldspar tails and neoblasts

K-feldspar is abundant in the low-strain portions of the analysed mylonite (Fig. 2). K-feldspar porphyroclasts and tails do not show any microstructure, CPO or misorientation axis distribution referable to dislocation creep processes (Fig. SOM2c; Menegon et al., 2008, and reference therein). The similar crystallographic orientation between feldspar(s) porphyroclasts and either K-feldspar tails or fine neoblast aggregates can be explained invoking epitaxial nucleation and growth during dissolution – precipitation (Fig. SOM2). Dissolution – precipitation would be consistent with the K-feldspar aggregate microstructure observed under CL, which probably reflect either the different chemistry, or the different intragranular strain, observed between magmatic ($Or_{93}Ab_7$) and

synkinematic K-feldspar ($\text{Or}_{96}\text{Ab}_4$) (Ramseyer et al., 1992; Götze et al., 1999; Słaby et al., 2008). The modification of the inherited CPO in fine-grained aggregates could be then related to the occurrence of anisotropic dissolution – precipitation processes and GBS/rigid body rotation during myrmekite shearing (Behrmann and Mainprice, 1987; Menegon et al., 2008, 2013).

5 6.2.3. Monomineralic quartz layers

The microstructures, CL signatures and strong crystallographic preferred orientation of monomineralic quartz layers indicate deformation by dominant dislocation creep aided by subgrain rotation (SGR) recrystallization (e.g. Fliervoet et al., 1997; Wheeler et al., 2001; Stipp et al., 2002; Bestmann and Pennacchioni, 2015). The misorientation axes distributions suggest the preferential activation of 10 $\{m\} < a >$ and $\{r-z\} < a >$ slip systems (e.g. Ceccato et al., 2017 and references therein).

The analysis of the grain orientation spread (GOS), useful to distinguish different generation of relict and/or recrystallized grains (Cross et al., 2017), suggests that there are no meaningful correlations between grain size and average grain distortion. This missing correlation may reflect a non-steady-state quartz microstructure during a prolonged deformation history or, more likely, the development

15 of the microstructures at different temperature conditions during pluton cooling. The bimodal grain size of recrystallized quartz includes coarser grains that we infer developed during the relatively high-temperature bulk solid-state deformation of the host granodiorite predating the development of localized shear zones at 450 °C dominated by SGR recrystallization (Ceccato et al., 2017; Ceccato and Pennacchioni, 2018). Coarser grains in quartz layers (grain sizes from >40) record differential stresses 20 < 40 MPa and strain rates of $10^{-13} - 10^{-14} \text{ s}^{-1}$. Subgrain and finer grains (20-35 μm in diameter) suggest that localized deformation and shearing occurred at differential stresses close to 40-70 MPa and strain rates of $10^{-11} - 10^{-12} \text{ s}^{-1}$ (Stipp and Tullis, 2003; Cross et al., 2017).

6.3. The rheology of the Rieserferner mylonites

The rheological effect of transformation of coarse K-feldspar to fine-grained sheared myrmekite and 25 the transition to an interconnected, weak, fine grained microstructure (Handy, 1990) is estimated here by investigating the deformational behaviour of different mixtures of plagioclase and quartz, deforming via different contribution of dislocation creep and diffusion creep. Our simplified model does not include biotite, white mica and biotite + plagioclase aggregates.

The bulk strain rate ($\dot{\varepsilon}_{bulk}$) of a mineral aggregate is given by:

30 (1)
$$\dot{\varepsilon}_{bulk} = \dot{\varepsilon}_{Disl} + \dot{\varepsilon}_{Diff}$$

where: $\dot{\varepsilon}_{Disl}$ and $\dot{\varepsilon}_{Diff}$ represents the strain rates of dislocation creep and diffusion creep of mineral components, respectively. The flow law of Hirth et al. (2001) has been used to calculate the contribution of dislocation creep of quartz:

$$(2) \quad \dot{\varepsilon}_{q-Disl} = A_q f_h \sigma^n e^{(-\frac{Q_q}{RT})}$$

5 where: A_q is the pre-exponential factor for quartz (MPa⁻ⁿ s⁻¹); f_h is the water fugacity coefficient; σ is the differential stress (MPa); n is the stress exponent; Q_q is the activation energy (J); R is the gas constant (J/K*mol); T is the temperature (K). Following Platt (2015), the flow law of den Brok (1998) for thin-film model of pressure-solution has been used to calculate the contribution of pressure-solution to quartz deformation:

$$10 \quad (3) \quad \dot{\varepsilon}_{q-Diff} = \dot{\varepsilon}_{qps} = C_2 \frac{\rho_f}{\rho_s} \frac{\sigma}{d^3} \frac{VcD_w}{RT}$$

where: C_2 is a shape constant; ρ_f and ρ_s are the fluid and solid densities (Kg m⁻³), respectively; d is the grain size (μm); V is the molar volume (μm³ mol⁻¹); c is the solubility of the solid in the fluid phase (molar fraction); D_w is the diffusivity of the solid in the grain-boundary fluid film (μm² s⁻¹). The flow laws of Rybacki et al. (2006) have been used to calculate the contribution of dislocation and diffusion

15 creep of feldspar:

$$(4) \quad \dot{\varepsilon} = A_f f_h \frac{\sigma^n}{d^m} e^{(-\frac{Q_f + pV^{act}}{RT})}$$

$$(5) \quad \dot{\varepsilon}_{f-Disl} = A_f f_h \sigma^3 e^{(-\frac{Q_f + pV^{act}}{RT})}$$

$$(6) \quad \dot{\varepsilon}_{f-Diff} = A_f f_h \frac{\sigma}{d^3} e^{(-\frac{Q_f + pV^{act}}{RT})}$$

where: A_f is the pre-exponential factor for feldspar (MPa⁻ⁿ μm^m s⁻¹); d is the grain size (μm); m is the grain-size exponent ($m=3$ for diffusion creep; $m=0$ for dislocation creep); p is the confining pressure (MPa); V^{act} is the activation volume (m³ mol⁻¹). Deformation mechanisms maps for sheared myrmekite were calculated following the self-consistent approach presented in Dimanov and Dresen (2005) and Platt (2015), in which flow law parameters for the poly-phase aggregate deforming via dislocation creep are recalculated as follows:

$$25 \quad (7) \quad \text{Log}_{10} n_a = \phi_1 \text{log}_{10} n_1 + \phi_2 \text{log}_{10} n_2$$

$$(8) \quad Q_a = [Q_2(n_a - n_I) - Q_I(n_a - n_2)] / (n_2 - n_I)$$

$$(9) \quad \log_{10} A_a = [\log_{10} A_2(n_a - n_I) - \log_{10} A_I(n_a - n_2)] / (n_2 - n_I)$$

where: n_a is the stress exponent for dislocation creep of the two-phase mixture; ϕ_i is the volume fraction of the phase i ; n_i is the stress exponent of the i phase; Q_a is the dislocation creep activation energy for

5 the aggregate (J); Q_i is the activation energy for the i phase (J); A_a is the pre-exponential factor for the aggregate ($\text{MPa}^{-n} \mu\text{m}^m \text{s}^{-1}$); A_i is the pre-exponential factor for the i phase ($\text{MPa}^{-n} \mu\text{m}^m \text{s}^{-1}$). In order to account for water fugacity and activation volumes parameters in feldspar flow laws (Rybacki and Dresen, 2006), we implemented the model of Platt (2015), which includes the activation volume for feldspars in the calculation of the aggregate activation energy Q_a , as follow:

$$10 \quad (10) \quad Q_a = [(Q_f + pV^{act})(n_a - n_Q) - Q_Q(n_a - n_f)] / (n_f - n_Q)$$

where: p is the confining pressure (MPa); V^{act} is the activation volume for feldspars (m^3/mol). Water fugacity coefficients were integrated in the resulting flow law of sheared myrmekite deforming by dislocation creep:

$$(11) \quad \dot{\varepsilon}_{Disl} = A_a f_h \sigma^{na} e^{(-\frac{Q_a}{RT})}$$

15 The flow law for sheared myrmekite deforming by diffusion creep (plagioclase) and thin-film pressure solution (quartz) has been calculated, following the approach of Dimanov and Dresen (2005) and Platt (2015), by considering pressure-solution (in quartz) and diffusion creep (in feldspar) to contribute linearly to the bulk viscosity of the aggregate, μ_a :

$$(12) \quad 3\mu_a^2 + [2(\mu_q + \mu_f) - 5(\phi_q \mu_q + \phi_f \mu_f)]\mu_a - 2\mu_q \mu_f = 0$$

20 where

$$(13) \quad \mu_q = \frac{\sigma}{2\dot{\varepsilon}_{qps}}$$

is the viscosity of quartz deforming via pressure-solution processes calculated following the thin-film model of den Brok (1998); and

$$(14) \quad \mu_f = \frac{\sigma}{2\dot{\varepsilon}_{f-diff}}$$

is the viscosity of feldspar deforming via diffusion creep. Therefore, the deformation mechanism maps of the sheared myrmekite have been calculated using the following flow law:

$$(15) \quad \dot{\varepsilon}_{bulk} = A_a f_h \sigma^{n_a} e^{(-\frac{Q_a}{RT})} + \frac{\sigma}{2\mu_a}$$

The rheological modelling of poly-phase aggregates containing more than two-rheological phases (e.g.

5 a granitoid composed of plagioclase + quartz + myrmekite) has been performed applying iteratively the calculation of Platt (2015). For example, for a mixture composed of ϕ_X , ϕ_Y and ϕ_Z volume fractions of the phases X (plagioclase), Y (quartz), and Z (sheared myrmekite):

10 (i) Firstly, n_{XY} , Q_{XY} , A_{XY} flow law parameters for the XY two-phase mixture are calculated following equations (7), (8) and (9), adopting the volume fractions ϕ_{X1} and ϕ_{Y1} defined as follow:

$$(16) \quad \phi_{X1} = \phi_X / (1 - \phi_Z); \quad \phi_{Y1} = \phi_Y / (1 - \phi_Z).$$

(ii) Then for the calculation of the three-phase mixture flow law parameters n_{XYZ} , Q_{XYZ} , A_{XYZ} , considering the three-phase mixture as the result of mixing between phases “XY” and Z, volume fractions are recalculated as follow:

$$(17) \quad \phi_{XY} = \phi_X + \phi_Y; \quad \phi_Z = \phi_Z.$$

and the parameters are calculated as follow:

$$(18) \quad n_{XYZ} = 10^{n_{XY} \log_{10} n_{XY} + n_Z \log_{10} n_Z}$$

$$(19) \quad Q_{XYZ} = [Q_Z(n_{XYZ} - n_{XY}) - Q_{XY}(n_{XYZ} - n_Z)] / (n_Z - n_{XY})$$

$$(20) \quad A_{XYZ} = 10^{[n_{XYZ} \log_{10} A_Z(n_{XYZ} - n_{XY}) - n_{XY} \log_{10} A_{XY}(n_{XYZ} - n_Z)] / (n_Z - n_{XY})}.$$

20 For the calculation of the rheology of an aggregate in which dislocation and diffusion creep contribute in different proportions to the total strain rate (granitoid rock including a variable amount of myrmekite, see Discussion below), a limiting factor θ has been introduced in equation (15). For example, for an aggregate in which diffusion creep is limited to a specific volume proportion of phases:

$$(21) \quad \dot{\varepsilon}_{bulk} = A_a f_h \sigma^{n_a} e^{(-\frac{Q_a}{RT})} + \theta_{diff} \frac{\sigma}{2\mu_a}$$

25 where θ_{diff} represents the volume fraction of the phases undergoing diffusion creep in the aggregate. To consider the progressive transformation of feldspar into myrmekite with increasing strain, differential stress vs. strain rate curves have been calculated for a “granitoid” aggregate with increasing

vol% of myrmekite substituting for feldspar (Fig. 8c). The progression of the reaction is quantified by the reaction progress factor χ . The maximum volume percentage of feldspar substitution has been limited to

$$(22) \quad \phi_{MAX} = 20 \text{ vol\%} ;$$

5 the average concentration of K-feldspar in granite and granodiorite. For $\chi=0$ (no myrmekite), the rock is composed by 60 vol% plagioclase An_{100} and 40 vol% quartz. For $\chi>0$, plagioclase An_{100} (representing K-feldspar) is increasingly replaced by myrmekite. Myrmekite, plagioclase and quartz volume proportion in the rock are then calculated respectively as follow:

$$(23) \quad \phi_{Myrm}^1 = \chi * \phi_{MAX};$$

$$10 \quad (24) \quad \phi_{Plg}^1 = \phi_{Plg} - \chi * \phi_{MAX};$$

$$(25) \quad \phi_{Qtz}^1 = \phi_{Qtz}$$

The calculations represent the case in which grain size sensitive creep occurs only in sheared myrmekite, whereas granitoid quartz and feldspars deform only by dislocation creep. Therefore, the contribution of diffusion creep to bulk viscosity in equation (21) is proportional to

$$15 \quad (26) \quad \theta_{diff} = \chi * \phi_{MAX}$$

Deformation mechanisms maps have been calculated and plotted on grain-size vs. differential stress and on differential-stress vs. strain rate diagrams for the following three compositions (Fig. 8):

- (i) monomineralic quartz layer deforming via both dislocation and diffusion creep (Fig. 8a);
- (ii) sheared myrmekite, modelled as 80 vol% plagioclase (An_{60}) + 20 vol% quartz deforming via both dislocation creep and grain size sensitive creep (Fig. 8b); the input grain size is 7 μm , identical for both minerals;
- (iii) a mixture of 60% plagioclase (An_{100}) + 40% quartz assumed as a simplified composition representative of a mica-free granitoid rock deforming only by dislocation creep (after referred as “granitoid”) (Fig. 8c);

25 The above general flow laws and flow law parameters are estimated for the pressure-temperature conditions of mylonitization of the Rieserferner (450°C and 0.35GPa; Ceccato, 2018). At these conditions, the calculated water fugacity is $f_h = 97 \text{ MPa}$ (Pitzer and Stern, 1994). Fluid density, quartz

solubility and diffusivity in the thin-film (grain boundary) fluid has been calculated following Fournier and Potter (1982) and Burnham et al. (1969). The flow law parameters defined for An_{100} and An_{60} by Rybacki and Dresen (2004) have been adopted for our calculations to simulate different compositions of “granitoid” and myrmekitic feldspars. In our calculations All the K-feldspar has been considered as 5 plagioclase, given the lack of flow law parameters for K-feldspar (see discussion in Platt, 2015; Viegas et al., 2016). Our calculation includes the contribution of GBS to the bulk strain rate of the feldspar aggregate, which is considered in the flow law parameters adopted here (see discussions in Xiao et al., 2002; Rybacki and Dresen, 2004).

6.3.1. Relative strength, rheological ranking and strain localization

10 The grain-size vs. differential stress and differential stress vs. strain rate diagrams for the three above defined aggregates are shown in Fig. 8. Flow law parameters are listed in Table 1. The results indicate that the three aggregates can be ranked, from the strongest to the weakest, as follows: (i) quartz-feldspar “granitoid” aggregate; (ii) monomineralic quartz aggregates (grain size of 4-10-20-100 μm); (iii) sheared myrmekite. This ranking is consistent and validated by several field and microstructural 15 observations, that highlight the strain localization capability of monomineralic quartz layers (i.e. quartz veins) and two-phase microstructural domains (i.e. sheared myrmekite) in granitoid rocks (Pennacchioni, 2005; Pennacchioni and Mancktelow, 2007; Menegon and Pennacchioni, 2010; Pennacchioni and Zucchi, 2013; Pennacchioni et al., 2010; Ceccato et al., 2017). The results of rheological calculation of plagioclase + quartz aggregates deforming via diffusion creep (sheared 20 myrmekite) are consistent and comparable with some of the experimental results of Xiao et al., (2002) extrapolated to geological temperatures (Fig. 8c). The experimental data that best fit our estimated rheological curve are those obtained from triaxial deformation experiments of synthetic very fine-grained wet aggregate of 80 vol% An_{100} (6 μm) + 20 vol% quartz (10 μm). In addition, this rheological ranking supports the interpretation that quartz grains inside pristine myrmekite deforming via 25 dissolution – precipitation creep may behave as strong inclusions compared to feldspar grains deforming via diffusion creep; around these strong inclusions stress can concentrate and trigger microfracturing in the surrounding plagioclase.

These results show that in the Rieserferner mylonites an effective strength contrast occur between the host rock, mono- and poly-mineralic aggregates occurs as a consequence of the different deformation 30 mechanisms. To quantify the effective strength contrast between the modelled compositions, we consider both constant stress and constant strain-rate conditions. The differential stress of 40-70 MPa estimated from the finer grain size of quartz (20-35 μm) can be considered representative of the bulk

flow stress of the mylonite. At constant stress conditions, such differential stress corresponds to a strain rate of 10^{-11} - 10^{-13} s $^{-1}$ of the quartz aggregates deforming by dislocation creep (Fig. 8a). At such differential stress, a sheared myrmekite deforming via diffusion creep would flow at strain rates $>10^{-12}$ s $^{-1}$, depending on the actual grain size of the aggregate (red transparent area in Fig. 8b). For the grain size range of sheared myrmekite (4-7 μ m), the observed strain rates are always $>10^{-11}$ s $^{-1}$, and for the above defined differential stress range the calculated strain rate is on the order of 10 $^{-9}$ s $^{-1}$ (intersection between red transparent area and black box in Fig. 8b). Therefore, assuming constant differential stress conditions, a strain-rate partitioning of 2-4 orders of magnitude is expected between monomineralic quartz and sheared myrmekite (similarly to Behrmann and Mainprice, 1987). Such strain-rate partitioning at constant stress could also explain the observed decrease in quartz grain size from the core of monomineralic layers toward neighbouring sheared myrmekite (Fig. 4). The differential stress, calculated for sheared myrmekite deforming via diffusion creep at constant strain rate conditions of 10 $^{-11}$ - 10 $^{-12}$ s $^{-1}$, is always <45 MPa. Under constant strain rate assumption, the strength contrast between monomineralic quartz and sheared myrmekite is not quantifiable; however, the sheared myrmekite are always weaker than monomineralic quartz deforming via dislocation creep. Strain rates in the order of 10 $^{-11}$ -10 $^{-13}$ s $^{-1}$ would require grain sizes in the range of 10-100 μ m in the sheared myrmekite deforming by diffusion creep only (grey shaded areas in Fig. 8b).

6.3.2. The effect of myrmekite reaction

Figure 9d shows the different curves describing the rheological behaviour of a simplified granitoid rock where K-feldspar is progressively replaced, up to 20 vol%, by sheared myrmekite. The flow behaviour of the derived granitoid mylonite is represented by the grey curves, and is linear viscous for most of the investigated conditions. The complete consumption of K-feldspar results in 3-4 orders of magnitude increase of strain rate, consistent with experimental observations (Xiao et al., 2002). A similar increase in strain rate is already observed for reaction progress factor of $\chi = 0.25$, i.e. for a 5 vol% of sheared myrmekite in the total rock volume. These results can be compared to the different degree of myrmekite substitution observed along the strain gradient in the shear zone and also justify the progressive increase in strain toward the ultramylonite with increasing myrmekite substitution (Fig. 2), suggesting positive feedback between strain-induced myrmekite and strain accommodation. Dissolution-precipitation creep of K-feldspar and associated GSS creep in K-feldspar + plagioclase + quartz aggregates have been already described by Behrmann and Mainprice (1987) as an efficient strain accommodation and weakening mechanism in quartz-feldspar mylonites. In the Rieserferner mylonites, GSS creep of K-feldspar seems to be dominant in protomylonite, but its role decreases with increasing myrmekite substitution (Fig. 2). The positive correlation between accommodated strain and

myrmekite substitution suggests that GSS creep processes in K-feldspar are however not capable of accommodating strain at rates comparable to those produced by GSS creep in sheared myrmekite.

The effective role of small volume fraction of myrmekite in rheological weakening of the Rieserferner granitoid during mylonitization might be overestimated by our calculation, for two main reasons: (i)

5 other weakening mechanisms may concur during homogeneous deformation of the granitoid rock that are not considered in our simplified model of granitoid (such as feldspar GSS creep, biotite recrystallization); (ii) myrmekite initially occurred into scattered non interconnected pockets (e.g. Handy, 1994); it is only with increasing strain and increasing volume fraction of sheared myrmekite, that these isolate pockets coalesced to form an interconnected weak layer microstructure. In the
10 Rieserferner mylonites effective interconnection of sheared myrmekite is already developed in presence of 5 to 7 vol% of myrmekite and is particularly well developed at 10-15 vol% (Fig. 2). Therefore, mylonites containing up to 15 vol% of sheared myrmekite ideally underwent deformation at strain rates of 10^{-10} - 10^{-11} s⁻¹ and at differential stresses in the range between 14 and 70 MPa. These
15 mylonites were synkinematic to mylonitic quartz veins described in Ceccato et al. (2017), for which quartz paleopiezometry retrieved comparable strain rates of 10^{-11} s⁻¹ developed under 117 MPa differential stress (10 μ m of grain size).

7. Conclusions

Metamorphic reactions contribute importantly to strain weakening within the Rieserferner granitoid mylonites. A primary grain size reduction mechanism is related to the development of myrmekite

20 evolving, with increasing strain, to weak aggregates of quartz and plagioclase. Topotactic replacement has been inferred from the coincidence between myrmekitic plagioclase and parent K-feldspar grain crystal lattices. Transition from pristine myrmekite to fine-grained sheared myrmekite involved micro-fracturing, annealing and shearing of the resulting granoblastic aggregate. Sheared myrmekite consists of fine grained plagioclase + quartz aggregates (7 μ m and 4 μ m in grain size, respectively) that show
25 ordered (anticlustered) spatial distribution and well-defined shape preferred orientation; quartz usually occurs at triple- and quadruple-junction between plagioclase grains. Both plagioclase and quartz show weak CPOs and almost uniform misorientation angle distributions. Sheared myrmekite microstructural features suggest that different deformation mechanisms occur in plagioclase and quartz: plagioclase deforms mainly by GSS creep, whereas dissolution - precipitation and nucleation processes are
30 dominant in quartz. Myrmekite formation promotes also phase mixing, as the pristine myrmekite microstructure predispose the development of an “anticlustered” spatial distribution of phases in the recrystallized aggregate. Strong grain size reduction and the nucleation of plagioclase + quartz

polymineralic aggregates lead to a switch in the dominant deformation mechanisms, activating GSS creep processes and triggered phase mixing. GSS processes and phase mixing inhibit grain growth and stabilizes grain size, hindering the efficiency of dynamic recrystallization processes and self-sustaining the activity of GSS processes. Therefore, the stress-induced formation of myrmekite lead to the 5 activation of self-sustaining weakening processes. Results of rheological calculations show that, at the conditions of Rieserferner mylonitization, sheared myrmekite are several orders of magnitude weaker than pure quartz layers and ideal granitoid rock deforming via dislocation creep. Strain-rate partitioning is therefore expected to occur between sheared myrmekite and monomineralic quartz layers. 5 vol% of myrmekite would lead to an increase in strain rate of 3-4 orders of magnitude. 10 However, the effective role of myrmekite in rock weakening is however dependent on the evolution of the rock microstructure. Effective weakening need for interconnection of sheared myrmekite layers that occur after the development of 10-15 vol% of myrmekite. This work once again highlights the importance of metamorphic reactions and micro-cataclastic processes as grain size reduction mechanisms in feldspar and their role in localization of ductile deformation. The microstructural 15 results and the rheological calculation presented here will be useful for further development of detailed rheological models of feldspar-rich rocks (continental crust rocks) at mid-crustal conditions.

Code availability

The MATLAB script used for rheological calculation is available on request from the first author.

Data availability

20 Supplementary data are available in Supplementary Online Material (SOM).

Appendix A: Methods

A.1 EBSD analysis

Electron backscattered diffraction analysis was carried out on a JEOL 7001 FEG SEM equipped with a Nordlys Max EBSD detector (AZTec acquisition software, Oxford Instruments) at the Electron 25 Microscope Centre of Plymouth University. The thin section was SYTON-polished for ca. 3 hours and carbon coated. EBSD patterns were acquired on rectangular grids with step sizes of 0.2, 0.3 and 0.35 μm . Working conditions during acquisition of EBSD patterns were 20 kV, 70° sample tilt, high vacuum, and working distance between 17 and 23 mm. All data have been processed and analysed using CHANNEL5 software of HKL Technology, Oxford Instruments. Noise reduction was applied

following Bestmann and Prior (2003). Local mis-indexing between plagioclase and K-feldspar was resolved by nullifying the subset of selected grains with area $<1\mu\text{m}^2$ in each map. Dauphiné twins smaller than $0.5\mu\text{m}$ have been interpreted as an error from mis-indexing and were replaced by the average orientation of the neighbouring pixels. The indexed phases and relative symmetry group used for the indexing are: quartz – Trigonal -3m; plagioclase (anorthite) – Triclinic -1; orthoclase – Monoclinic 2/m; clinozoisite, biotite and garnet have been indexed where present, but orientation data have not been analysed. Critical misorientation for the distinction between low- and high-angle boundaries have been chosen at 10° . Quartz grain boundaries with $60^\circ\pm5^\circ$ of misorientation were disregarded from grain detection procedure, to avoid any contribution from Dauphiné twinning. Plagioclase grain boundaries with $180^\circ\pm5^\circ$ of misorientation around [010] were disregarded from grain detection procedure, to avoid any contribution from Albite twinning. The pole figures (one-point-per-grain, where not differently specified) are plotted as equal area, lower hemisphere projections oriented with the general shear zone kinematics reference system (X = stretching lineation; Z = pole to general shear plane/vein boundary); whereas the misorientation axis distributions in sample coordinates are plotted as equal area, upper hemisphere projections. The inverse pole figures for misorientation axis distribution in crystal coordinates are upper hemisphere projections. Contoured projections have constant contouring parameters (Halfwidth: 10°). Contouring lines are given only for the 0.5-10 m.u.d. (multiple of uniform distributions) range.

A.2 SEM – Cathodoluminescence

Cathodoluminescence imaging was performed in a FEI Quanta 200 FEI equipped with Gatan monocle detector. Imaging was performed using an accelerating voltage of 20 kV, beam current of 8 nA and working distance of 20 mm in C-coated (15 nm) thin sections used for EBSD analysis. To avoid incorrect interpretation of potential artifacts in the sample, secondary (SE) and backscatter electron images were collected simultaneously with CL.

A.3 Grain size analysis

Grain sizes were obtained from the grain detection routine of the HKL Channel5 Tango software. The grain size was calculated as diameter of the circle with an equivalent area. The minimum cut-off area was set to $1\mu\text{m}^2$ which means that only grains of a size ≥ 4 or ≥ 9 pixels (depending on the map acquisition step-size) were considered. Grain size data were represented as area-weighted distributions by plotting frequency against the square-root grain-size-equivalent grain diameters (as in Herwagh and Berger, 2004; Berger et al., 2011). The grain size distribution approaches a Gaussian distribution when plotted in this way, allowing a good estimate of the mean grain size. The geometric mean grain size

(red thick line in grain size distribution diagrams) was obtained graphically as the maximum frequency grain size of the distribution curve. The distribution curve (blue line in grain size distribution diagrams) was obtained interpolating distribution data with a 6th degree polynomial equation in Excel-MS Office. Relative frequencies are normalized to 1.

5 **A.4 Image analysis**

Image analysis of grain shape was performed on both SEM-BSE images and phase maps obtained from EBSD. Quantification of phase amount (vol%) was performed through segmentation of SEM-BSE images of a whole thin section collected at the Electron Microscopy Centre of the University of Plymouth. Image processing and thresholding was done with the ImageJ software, and further 10 processing together with manual correction were applied to improve data quality and to ensure the correspondence of greyscale ranges with specific mineral phases. Grain boundary images and phase distribution images were obtained directly from EBSD phase maps and grain boundary maps elaborated by Channel5 (HKL technology). Before the analysis with ImageJ software, images were manually corrected in order to exclude mis-indexing and non-indexed orientation pixels. Grain 15 boundaries and phase amount have been quantified by pixel counting.

A.5 Electron microprobe analysis (EMPA)

Microchemical analyses were performed with EM wavelength-dispersive spectroscopy (WDS) at Electron Microprobe Laboratory at the Università degli Studi di Milano with a Jeol 8200 Super Probe; the operating conditions were: 15 kV accelerating voltage; 5 nA (feldspar, epidote and phyllosilicate) 20 beam current. PAP correction program was applied to convert X-ray counts into oxide weight percentages.

Author contributions

LM, GP and AC developed the initial idea of the study and performed initial exploratory SEM study. GP collected the samples of Rieserferner mylonites. LM acquired EBSD data. AC performed EBSD 25 data processing and analysis, and performed rheological calculations. LFGM performed cathodoluminescence analysis. AC prepared the figures and the manuscript with contributions from all the co-authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

Simone Papa, Francesco Giuntoli, Luca Pellegrino are thanked for fruitful discussions. Andrea Risplendente is thanked for his help during EMPA data collection at the Università degli Studi di Milano. The staff at University of Plymouth Electron Microscopy Centre is thanked for the assistance 5 during EBSD data acquisition. Luca Menegon acknowledges the financial support from a FP7 Marie Curie Career Integration Grant (grant agreement PCIG13-GA-2013-618289). Financial support from the University of Padova (“Progetto di Ateneo” CPDA140255) and from the Foundation “Ing. Aldo Gini” is acknowledged.

References

Abart, R., Heuser, D., and Habler, G.: Mechanisms of myrmekite formation: case study from the Weinsberg granite, Moldanubian zone, Upper Austria. *Contributions to Mineralogy and Petrology* 168, 1074, 2014.

Becke, F.: Über myrmekite. *Mineralogie und Pétrographie Mitteilungen* 27, 377-390, 1908.

Behrmann, J.H. and Mainprice, D.: Deformation mechanisms in a high-temperature quartz-feldspar mylonite: evidence for superplastic flow in the lower continental crust. *Tectonophysics* 140, 297-305, 1987.

Bellieni, G.: Caratteri geochimici del massiccio granodioritico tonalitico delle Vedrette di Ries (Rieserferner) - Alto Adige Orientale. *Rendiconti della Società Italiana di Mineralogia e Petrologia* 34, 527-548, 1978.

Bestmann, M. and Prior, D.J.: Intragranular dynamic recrystallization in naturally deformed calcite marble: diffusion accommodated grain boundary sliding as a result of subgrain rotation. *Journal of Structural Geology* 25, 1597-1613, 2003.

Bestmann, M. and Pennacchioni, G.: Ti distribution in quartz across a heterogeneous shear zone within a granodiorite: The effect of deformation mechanism and strain on Ti resetting. *Lithos*, 227, 37-56, 2015.

Burnham, C.W., Holloway, J.R., and Davis, N.F.: Thermodynamic Properties of Water to 1, 0000 °C and 10,000 Bars. *Geological Society of America*, 132, <https://doi.org/10.1130/SPE132>, 1969.

Ceccato, A., Pennacchioni, G., Menegon, L., and Bestmann, M.: Crystallographic control and texture inheritance during mylonitization of coarse grained quartz veins. *Lithos* 290-291, 210-227, doi:10.1016/j.lithos.2017.08.005, 2017.

Ceccato, A. and Pennacchioni, G.: Structural evolution of the Rieserferner Pluton in the framework of the Tertiary tectonics of the eastern Alps. *Journal of Structural Geology*. 2018.

Cesare, B., Marchesi, C., and Connolly, J.A.D.: Growth of myrmekite coronas by contact metamorphism of granitic mylonites in the aureole of Cima di Vila, Eastern Alps, Italy. *Journal of Metamorphic Geology* 20, 203-213, 2002.

Cesare, B., Mazzoli, C., Sassi, R., Spiess, R., and Sassi, F.P.: Beauty and complexity of metamorphism: case studies from the frontal part of the Adria microplate. *Rendiconti Lincei* 21, 73-94, 2010.

Ciancaleoni, L., and Marquer, D.: Syn-extension leucogranite deformation during convergence in the Eastern Central Alps: example of the Novate intrusion. *Terra Nova* 18, 170-180, 2006.

Cross, A. J., Prior, D. J., Stipp, M., and Kidder, S.: The recrystallized grain size piezometer for quartz: An EBSD-based calibration. *Geophysical Research Letters*, 44, 6667-6674, 2017.

5 Czaplińska, D., Piazolo, S., and Zibra, I.: The influence of phase and grain size distribution on the dynamics of strain localization in polymimetic rocks. *Journal of Structural Geology* 72, 15-32 2015.

De Bresser, J., Ter Heege, J., and Spiers, C.: Grain size reduction by dynamic recrystallization: can it result in major rheological weakening? *International Journal of Earth Sciences* 90, 28-45, 2001.

10 De Toni, G. B., Bitencourt, M. F., and Nardi, L. V. S.: Strain partitioning into dry and wet zones and the formation of Ca-rich myrmekite in syntectonic syenites: A case for melt-assisted dissolution-replacement creep under granulite facies conditions. *Journal of Structural Geology*, 91, 88-101, 2016.

Dégi, J., Abart, R., Török, K., Bali, E., Wirth, R., and Rhede, D.: Symplectite formation during 15 decompression induced garnet breakdown in lower crustal mafic granulite xenoliths: mechanisms and rates. *Contributions to Mineralogy and Petrology*, 159, 293-314, 2010.

Dell'Angelo, L.N. and Tullis, J.: Textural and mechanical evolution with progressive strain in experimentally deformed aplite. *Tectonophysics* 256, 57-82, 1996.

Den Brok, S.W.J.: Effect of microcracking on pressure-solution strain rate: The Gratz grain-boundary 20 model. *Geology* 26, 915-918, 1998.

Dimanov, A., and Dresen, G.: Rheology of synthetic anorthite-diopside aggregates: Implications for ductile shear zones. *Journal of Geophysical Research: Solid Earth* 110. <https://doi.org/10.1029/2004JB003431>, 2005.

Farver, J. R. and Yund, R. A.: Interphase boundary diffusion of oxygen and potassium in K-25 feldspar/quartz aggregates. *Geochimica et Cosmochimica Acta*, 59, 3697-3705, 1995

Fliervoet, T.F., White, S.H., and Drury, M.R.: Evidence for dominant grain-boundary sliding deformation in greenschist-and amphibolite-grade polymimetic ultramylonites from the Redbank Deformed Zone, Central Australia. *Journal of Structural Geology* 19, 1495-1520, 1997.

Fournier, R.O. and Potter II, R.W.: An equation correlating the solubility of quartz in water from 25 30 to 900 °C at pressures up to 10,000 bars. *Geochimica et Cosmochimica Acta* 46, 1969-1973, 1982.

Fusseis, F., Regenauer-Lieb, K., Liu, J., Hough, R.M., and De Carlo, F.: Creep cavitation can establish a dynamic granular fluid pump in ductile shear zones. *Nature* 459, 974, 2009.

Götze, J., Habermann, D., Neuser, R. D., and Richter, D. K.: High-resolution spectrometric analysis of rare earth elements-activated cathodoluminescence in feldspar minerals. *Chemical Geology*, 153, 81-91, 1999.

Handy, M.R.: The solid-state flow of polymineralic rocks. *Journal of Geophysical Research: Solid Earth* 95, 8647-8661, 1990.

Handy, M.R.: Flow laws for rocks containing two non-linear viscous phases: a phenomenological approach. *Journal of Structural Geology* 16, 287-301, 1994.

Herwegh, M. and Berger, A.: Deformation mechanisms in second-phase affected microstructures and their energy balance. *Journal of Structural Geology* 26, 1483-1498, 2004.

10 Herwegh, M., Linckens, J., Ebert, A., Berger, A., and Brodhag, S. H.: The role of second phases for controlling microstructural evolution in polymineralic rocks: A review. *Journal of Structural Geology* 33, 1728-1750, 2011.

Hiraga, T., Miyazaki, T., Yoshida, H., and Zimmerman, M.E.: Comparison of microstructures in superplastically deformed synthetic materials and natural mylonites: Mineral aggregation via grain boundary sliding. *Geology* 41, 959-962, 2013

15 Hirth, G., Teyssier, C., and Dunlap, J.W.: An evaluation of quartzite flow laws based on comparisons between experimentally and naturally deformed rocks. *International Journal of Earth Sciences* 90, 77-87, 2001.

Hopson, R.F. and Ramseyer, K.: Cathodoluminescence microscopy of myrmekite. *Geology* 18, 336-339, 1990.

Jiang, Z., Prior, D.J., and Wheeler, J.: Albite crystallographic preferred orientation and grain misorientation distribution in a low-grade mylonite: implications for granular flow. *Journal of Structural Geology* 22, 1663-1674, 2000.

20 Johnson, S.E., Marsh, J.H., and Vernon, R.H.: From tonalite to mylonite: coupled mechanical and chemical processes in foliation development and strain localization. *Journal of Virtual Explorer* 30, 2008.

Kilian, R., Heilbronner, R., and Stünitz, H.: Quartz grain size reduction in a granitoid rock and the transition from dislocation to diffusion creep. *Journal of Structural Geology* 33, 1265-1284, 2011.

Kretz, R.: Symbols for rock-forming minerals. *American mineralogist*, 68, 277-279, 1983.

25 Kruse, R. and Stünitz, H.: Deformation mechanisms and phase distribution in mafic high-temperature mylonites from the Jotun Nappe, southern Norway. *Tectonophysics* 303, 223-249, 1999.

Kruse, R., Stünitz, H., and Kunze, K.: Dynamic recrystallization processes in plagioclase porphyroclasts. *Journal of Structural Geology* 23, 1781-1802, 2001.

Lapworth, T., Wheeler, J., and Prior, D.J.: The deformation of plagioclase investigated using electron backscatter diffraction crystallographic preferred orientation data. *Journal of Structural Geology* 24, 387-399, 2002.

LaTour, T. E. and Barnett, R. L.: Mineralogical changes accompanying mylonitization in the Bitterroot dome of the Idaho batholith: Implications for timing of deformation. *Geological Society of America Bulletin*, 98, 356-363, 1987.

Menegon, L., Pennacchioni, G., and Stünitz, H.: Nucleation and growth of myrmekite during ductile shear deformation in metagranites. *Journal of Metamorphic Geology* 24, 553-568, 2006.

Menegon, L., Pennacchioni, G., and Spiess, R.: Dissolution-precipitation creep of K-feldspar in mid-crustal granite mylonites. *Journal of Structural Geology*, 30, 565-579, 2008.

Menegon, L. and Pennacchioni, G.: Local shear zone pattern and bulk deformation in the Gran Paradiso metagranite (NW Italian Alps). *International Journal of Earth Sciences* 99, 1805-1825, doi:10.1007/s00531-009-0485-6 , 2010.

Menegon, L., Stünitz, H., Nasipuri, P., Heilbronner, R., and Svahnberg, H.: Transition from fracturing to viscous flow in granulite facies perthitic feldspar (Lofoten, Norway). *Journal of Structural Geology* 48, 95-112, 2013.

Menegon, L., Fusseis, F., Stünitz, H., and Xiao, X.: Creep cavitation bands control porosity and fluid flow in lower crustal shear zones. *Geology* 43, 227-230, 2015.

Miranda, E.A., Hirth, G., and John, B.E.: Microstructural evidence for the transition from dislocation creep to dislocation-accommodated grain boundary sliding in naturally deformed plagioclase. *Journal of Structural Geology* 92, 30-45, 2016.

Odashima, N., Morishita, T., Ozawa, K., Nagahara, H., Tsuchiyama, A., and Nagashima, R.: Formation and deformation mechanisms of pyroxene-spinel symplectite in an ascending mantle, the Horoman peridotite complex, Japan: An EBSD (electron backscatter diffraction) study. *Journal of Mineralogical and Petrological Sciences*, 103, 1-15, 2007.

Pennacchioni, G.: Control of the geometry of precursor brittle structures on the type of ductile shear zone in the Adamello tonalites, Southern Alps (Italy). *Journal of Structural Geology* 27, 627-644, 2005.

Pennacchioni, G., Di Toro, G., Brack, P., Menegon, L., and Villa, I.M.: Brittle–ductile–brittle deformation during cooling of tonalite (Adamello, Southern Italian Alps). *Tectonophysics* 427, 171-197, 2006.

Pennacchioni G., and Mancktelow, N.S.: Nucleation and initial growth of a shear zone network within compositionally and structurally heterogeneous granitoids under amphibolite facies conditions. *Journal of Structural Geology* 29, 1757-1780, 2007.

Pennacchioni, G. and Mancktelow, N.S.: Small-scale ductile shear zones: neither extending, nor thickening, nor narrowing. *Earth-Science Reviews*, doi:10.1016/j.earscirev.2018.06.004 , 2018.

Pennacchioni, G. and Zucchi, E.: High temperature fracturing and ductile deformation during cooling of a pluton: The Lake Edison granodiorite (Sierra Nevada batholith, California). *Journal of Structural Geology* 50, 54-81, 2013.

Phillips, E. R.: On polygenetic myrmekite. *Geological Magazine*, 117, 29-36, 1980.

Pitzer, K. S. and Stern, S.M.: Equations of state valid continuously from zero to extreme pressures for H₂O and CO₂. *The Journal of Chemical Physics* 101, 3111-3116, 1994.

Platt, J.P.: Rheology of two-phase systems: A microphysical and observational approach. *Journal of Structural Geology* 77, 213-227, 2015.

Ramseyer, K., AlDahan, A. A., Collini, B., and Landström, O.: Petrological modifications in granitic rocks from the Siljan impact structure: evidence from cathodoluminescence. *Tectonophysics*, 216, 195-204, 1992.

Ree, J.H., Kim, H.S., Han, R., and Jung, H.: Grain-size reduction of feldspars by fracturing and neocrystallization in a low-grade granitic mylonite and its rheological effect. *Tectonophysics* 407, 227-237, 2005.

Rybacki, E. and Dresen, G.: Deformation mechanism maps for feldspar rocks. *Tectonophysics* 382, 173-187, 2004.

Rybacki, E., Gottschalk, M., Wirth, R., and Dresen, G.: Influence of water fugacity and activation volume on the flow properties of fine-grained anorthite aggregates. *Journal of Geophysical Research: Solid Earth* 111. <https://doi.org/10.1029/2005JB003663> , 2006.

Romer, R.L. and Siegesmund, S.: Why allanite may swindle about its true age. *Contributions to Mineralogy and Petrology* 146, 297-307, 2003.

Segall, P. and Simpson, C.: Nucleation of ductile shear zones on dilatant fractures. *Geology*, 14, 56-59, 1986.

Simpson, C. and Wintsch, R.P.: Evidence for deformation-induced K-feldspar replacement by myrmekite. *Journal of Metamorphic Geology* 7, 261-275, 1989.

Ślaby, E., Götze, J., Wörner, G., Simon, K., Wrzalik, R., and Śmigielski, M.: K-feldspar phenocrysts in microgranular magmatic enclaves: a cathodoluminescence and geochemical study of crystal growth as a marker of magma mingling dynamics. *Lithos*, 105, 85-97, 2008.

Stel, H. and Breedveld, M.: Crystallographic orientation patterns of myrmekitic quartz: a fabric memory in quartz ribbon-bearing gneisses. *Journal of Structural Geology* 12, 19-28, 1990.

Stipp, M., Stünitz, H., Heilbronner, R., Schmid, S.M.: The eastern Tonale fault zone: a ‘natural laboratory’ for crystal plastic deformation of quartz over a temperature range from 250 to 700 °C. *Journal of Structural Geology* 24, 1861-1884, 2002.

Stipp, M. and Tullis, J.: The recrystallized grain size piezometer for quartz. *Geophysical Research Letters*, 30, 2003.

Stipp, M., Tullis, J., Scherwath, M., and Behrmann, J. H. A new perspective on paleopiezometry: Dynamically recrystallized grain size distributions indicate mechanism changes. *Geology*, 38, 759-762, 2010.

Stünitz, H. and Fitz Gerald, J.F.: Deformation of granitoids at low metamorphic grade. II: Granular flow in albite-rich mylonites. *Tectonophysics* 221, 299-324, 1993.

Tsurumi, J., Hosonuma, H., and Kanagawa, K.: Strain localization due to a positive feedback of deformation and myrmekite-forming reaction in granite and aplite mylonites along the Hatagawa Shear Zone of NE Japan. *Journal of Structural Geology* 25, 557-574, 2003.

Vernon, R.H.: Questions about myrmekite in deformed rocks. *Journal of Structural Geology* 13, 979-985, 1991.

Viegas, G., Menegon, L., and Archanjo, C.: Brittle grain-size reduction of feldspar, phase mixing and strain localization in granitoids at mid-crustal conditions (Pernambuco shear zone, NE Brazil). *Solid Earth* 7, 375-396, 2016.

Wheeler, J., Prior, D., Jiang, Z., Spiess, R., and Trimby, P.: The petrological significance of misorientations between grains. *Contributions to Mineralogy and Petrology* 141, 109-124, 2001.

White, S.: Geological significance of recovery and recrystallization processes in quartz. *Tectonophysics* 39, 143-170, 1977.

Wirth, R. and Voll, G.: Cellular intergrowth between quartz and sodium-rich plagioclase (myrmekite)—an analogue of discontinuous precipitation in metal alloys. *Journal of Materials Science* 22, 1913-1918, 1987.

Xiao, X., Wirth, R., and Dresen, G.: Diffusion creep of anorthite-quartz aggregates. *Journal of Geophysical Research: Solid Earth* 107(B11), 2002.

Figure and Tables Captions

Table 1. Parameters adopted in the rheological calculations. (a) List of the general parameters adopted in the rheological calculations. (b) Values of flow law parameters adopted in the rheological calculations according to mineral phase and deformation mechanism.

5 **Figure 1.** Microstructures of Rieserferner granodiorite mylonites. (a) Microphotograph (crossed nichols) showing the layered structure of granodiorite mylonites, composed of alternating layers of recrystallized quartz, recrystallized biotite + plagioclase + quartz, and plagioclase + quartz. White arrows indicate layers of recrystallized quartz (upper) and biotite (lower). (b) SEM-BSE image of the area shown in (a). (c) SEM-BSE image of a pristine myrmekite replacing K-feldspar. (d) SEM-BSE
10 image of the K-feldspar + biotite tails in strain shadows between two plagioclase porphyroclasts. K-feldspar in the strain shadows is in turn replaced by myrmekite (white arrows). (e) SEM-BSE image of a K-feldspar porphyroblast and of sheared myrmekite. Pristine myrmekite developed on K-feldspar boundaries parallel to the mylonitic foliation are sheared to form plagioclase + quartz aggregates (sheared myrmekite). The white polygon encloses K-feldspar neoblasts in strain shadows and sheared
15 myrmekite. (f) CL image of (e). Note the alteration of the CL signal after the EBSD scan. (g) K-feldspar and sheared myrmekite aggregate (particular of the EBSD map of Fig. 3). (h) CL image of (g).

Figure 2. Phase distribution across a strain gradient in a granodiorite mylonite. (a) Mosaic of SEM-BSE images with the K-feldspar and the myrmekite + sheared myrmekite coloured in red and pale
20 blue, respectively. The yellow rectangles indicate the location of the EBSD maps of Figs. 3, 4 and SOM2. (b) Bar diagram showing the volume percentage of K-feldspar (red bars) and myrmekite (pale blue bars) across the microstructure: PM = protomylonite; M = mylonite; and UM = ultramylonite.

Figure 3. EBSD map and crystallographic orientation data of incipient myrmekite and parent K-feldspar. (a) EBSD-derived phase map. The area delimited by dashed polygons represents pristine
25 myrmekite. Pole figures for: (b) K-feldspar grains on which pristine myrmekite nucleated; (c) plagioclase and (d) quartz in pristine myrmekite.

Figure 4. EBSD map and crystallographic orientation data of pristine and sheared myrmekite of Fig. 1e. (a) EBSD phase map including areas (A, B, C, D) selected for grain size analysis and phase distribution analysis. (b) Pole figures for quartz from the sheared myrmekite of area B. Upper row: 30 scattered data. Lower row: contoured data. (c) Misorientation axis distributions for quartz in sample (upper row) and crystal (lower row) coordinate system. (d) Misorientation angle distribution for quartz.

(e) Pole figures for plagioclase from sheared myrmekite of Area B. Upper row: scattered data. Lower row: contoured data. In this case, the [100] plagioclase pole figure is reported in upper hemisphere, where the maximum has been observed. (f) Misorientation axis distributions for plagioclase in sample (upper row) and crystal (lower row) coordinate system. (g) Misorientation angle distribution for 5 plagioclase.

Figure 5. Image analysis on phase spatial distribution. The diagram reports phase- and grain-boundary fraction for sheared myrmekite. Continuous curves represent the theoretical probability of phase- and grain-boundary fraction as a function of quartz content expected for a random distribution in a two-phase aggregate. The small maps on the left hand side report one of the analysed areas (Area C. Fig. 10 4), showing from the top to the bottom the phase map, the related plagioclase, quartz grain boundaries and phase boundaries.

Figure 6. Area-weighted grain size distributions and SPO for quartz. (a) Grain size distribution for quartz in incipient myrmekite A in Fig. 4a. (b) Grain size distribution for quartz in sheared myrmekite B in Fig. 4a. (c) Grain size distribution for quartz in sheared myrmekite C in Fig. 4a. (d) Grain size 15 distribution for quartz in monomineralic layer in Fig. 4a. (e) Rose diagram showing the orientation of major axis of quartz grains, defining a weak SPO.

Figure 7. Area-weighted grain size distributions and SPO for plagioclase. (a) Grain size distribution for plagioclase in myrmekite of Fig. 3. (b) Grain size distribution for plagioclase in incipient myrmekite A in Fig. 4a. (c) Grain size distribution for plagioclase in sheared myrmekite B in Fig. 4a. 20 (d) Rose diagram showing the orientation of major axis of plagioclase grains, defining a weak SPO.

Figure 8. Diagrams obtained from the calculation of the rheological model explained in the text. Grains size vs. Differential stress map with contoured strain rate curves obtained for: (a) quartz, (b) 25 80% plagioclase (An60) + 20% quartz aggregates. (a) The piezometric curve from Stipp and Tullis (2003) (black curve) and Cross et al. (2017) (red curve) are reported. Red and black stars mark the differential stress/strain-rate conditions defined by the grain size observed in pure quartz layers: (A) 35 μm ; (B) 20 μm ; (C) 10 μm (Ceccato et al., 2017). (b) A and B marked red polygons represent the differential stress range obtained from piezometric calculations on pure quartz layers (red and black stars along respective piezometric curves). The black dashed line represents the boundary between dislocation and diffusion creep dominated conditions. The black rectangle represents the grain size 30 range (4-7 μm) observed in the sheared myrmekite. The grey semi-transparent polygon defines the field of possible grain-size and differential stress conditions for iso-strain-rate conditions defined from piezometric relations. (c) Log Differential stress vs. Log Strain rate diagram reporting the curve

calculated for pure quartz with different grain sizes, sheared myrmekite, ideal granitoid rock and the curves representing the rheology of pure feldspar aggregates. For comparison, one of the curve obtained from experimental data of Xiao et al., (2002) is reported (black dashed curve). Grey field represent the uncertainties on the experimentally defined rheological curve. (d) Log Differential stress
5 vs. Log Strain rate diagram reporting the curve calculated for pure quartz, sheared myrmekite and ideal granitoid rock and the curves representing the rheology of a granitoid (60% An100 + 40% Qtz) with variable amount of sheared myrmekite (80% plagioclase An60 + 20% Qtz). Maximum substitution is limited to 20% of initial feldspar (see text for explanation).