Author comments (AC) and manuscript changes (MC) on review by editor (Renée Heilbronner) and referee (Florian Fusseis)

1) The rheological evolution of the northern and southern sections of the Woodroffe Thrust is still not very well explained. Simply referring to a follow-up paper is not enough. If the shear zones are strain softening, should not the efficiency of the softening mechanism be the main controlling factor determining the ultimate width? This issue remains unexplained. Why not focus more on the strain softening mechanisms, as suggested? With the key message being that (hydration) reactions control the rheological evolution of a mylonite, the paper would be a clear step forward compared to previous studies by Camacho et al and Bell & Etheridge.

AC/MC: We fully understand the point being made here, which is why the planned third paper in the trilogy is devoted to examining the dependence of rheology and microstructure on varying metamorphic conditions and water activity. There is a logical progression to these papers. The first established the overall geometry, kinematics and metamorphic conditions (Wex et al., 2017). The current manuscript (second) documents the preferential distribution of mylonites in the footwall and the variation in water content from south to north. The third considers the variation in microstructure, CPO (EBSD and Texturegoniometer data) and grain size (EBSD) as a basis for establishing the deformation mechanisms and rheology. We could "cut-and-paste" aspects of this third paper but these parts could not be thoroughly developed in the current manuscript, would introduce repetition and overlap with the upcoming paper, and do not really fit the well-defined aim of the current manuscript. We consider that adding limited information on deformation mechanisms without a robust documentation would decrease rather than increase the quality of the current manuscript. We therefore continue to argue against mixing the clearly defined aims of the three planned papers. The reader is referred to a forthcoming paper for a full description of the deformation mechanisms.

2) With respect to the CO_2 - your answer to the second reviewers question is not satisfying. If the Woodroffe Thrust did not channelize these fluids, how did the fluids get to where they are now? The hypothesis proposed is not very well supported by the CO_2 data and requires a better discussion.

AC/MC: We can only report the observations that we have with regard to the irregular distribution of very minor accessory calcite in otherwise non-carbonaceous rocks. We have now expanded on the discussion to make it clear that the irregular spatial distribution of small amounts and the lack of correlation with indicators for associated introduction of aqueous fluid argue for a local redistribution rather than a pervasive influx. The data from the stable isotope analyses on the calcite are also inconclusive, with the δ^{13} C values being partially rock-buffered, reflecting the small amount of fluid involved and the high rock/fluid ratio. We do not insist on publishing this admittedly inconclusive data in the supplementary material but think it is correct to make it available to future workers in the area.

³⁾ You did not properly describe (and justify) the thresholding technique you used, error estimates are missing, methods for the separation of peaks are not explained. Statements such as "sophisticated segmentation tools [...] require careful checking of the output results" do not qualify. The reliability of your results needs to be demonstrated - your "believe" that the results are "quite reliable" is not convincing. Looking at the histogram in figure S3, I wonder what motivated the choice of positions of

the boundaries between these phases - as clearly, the peaks are overlapping, and it is virtually impossible to separate Qtz, Plag and Kfsp.

AC/MC: We now include the 2 x standard deviation for each of our samples. We also extended the description as to how we estimated the modal abundance of hydrous minerals in our thin sections. The method section in the appendix now provides additional information with regard to our thresholding technique, emphasizing that we did not attempt to separate every single mineral phase from one another, but only the hydrous from the anhydrous ones. Thresholds were manually defined between the individual grey scale values ranges defined by the hydrous and anhydrous each minerals. In order to achieve a proper separation, we adjusted brightness and contrast in a manner that the grey scale range defined by the hydrous minerals covered a significant portion of the entire spectrum. As a result, the histogram data of the individual anhydrous minerals strongly overlap with one another (as mentioned be the reviewer for the peaks of quartz, K-feldspar and plagioclase in Fig. S3). This approach minimalizes the overlap between the peaks defined by the hydrous minerals with those of the anhydrous minerals and thus provides a proper basis for the desired separation. We agree with the reviewer that in this case a proper separation of quartz, K-feldspar and plagioclase would not be possible. However, separating individual anhydrous minerals from one another was never our goal.

A few minor points

Line 22 - increase in the abundance of fluids

AC/MC: done.

Line 26 - difference between fault and shear zones (cf. Ramsay's definition of shear zones)?

AC/MC: Reference added.

Line 53 onward - I think this could be pitched better - reads a bit like "just another contribution" and doesn't really grab the reader's attention/interest. Why not emphasize what new data you are adding?

AC/MC: We added the fact that we utilize the radiogenic signature of footwall and hanging wall rocks to constrain their respective degree of reworking the ductile mylonite zone.

Line 377 - For the sake of whose simplicity?

AC/MC: We removed this statement.

Line 447 - *This is avoiding the important question as to where these fluids come from and how they travelled.*

AC/MC: We cannot answer these questions, as the stable isotope data is inconclusive.

Line 518 - and increasing displacement?

AC/MC: According to Hull (1988), shear zone thickness decreases with decreasing displacement.

Line 535 - Why is this a contradiction?

AC/MC: The sentence has been rewritten for greater clarity and the word "contradiction" no longer appears.

Line 539 - effective viscosity contrast?

AC/MC: Rephrased.

Line 545 - It's not exactly new that even small amounts of fluid will support strain localization.

AC/MC: True, but we do not claim that we were the first to come up with this conclusion. We simply use it as an argument why the shearing is preferentially developed in the footwall, because it has a slightly higher water content than the hanging wall in an overall relatively dry environment – and this explains the "inverted" or "unexpected" distribution of mylonites between hanging wall and footwall.

Inverted distribution of ductile deformation in the relatively "dry" middle crust across the Woodroffe Thrust, central Australia

Sebastian Wex¹, Neil S. Mancktelow¹, Friedrich Hawemann¹, Alfredo Camacho², Giorgio Pennacchioni³

¹Department of Earth Sciences, ETH Zurich, Sonneggstrasse 5, 8092 Zurich, Switzerland

⁵ ²Department of Geological Sciences, University of Manitoba, 125 Dysart Rd, Winnipeg, Manitoba, R3T 2N2, Canada ³Department of Geosciences, University of Padova, Via Gradenigo 6, 35131 Padova, Italy

Correspondence to: Neil S. Mancktelow (neil.mancktelow@erdw.ethz.ch)

Abstract. Thrust fault systems typically distribute shear strain preferentially into the hanging wall rather than the footwall. The Woodroffe Thrust in the Musgrave Block of central Australia is a regional-scale example that does not fit this model. It
developed due to intracontinental shortening during the Petermann Orogeny (ca. 560-520 Ma) and is interpreted to be at least 600 km long in its E-W strike direction, with an approximate top-to-north minimum displacement of 60-100 km. The associated mylonite zone is most broadly developed in the footwall. The immediate hanging wall was only marginally involved in the mylonitization process, as can be demonstrated from the contrasting thorium signatures of mylonites derived from the upper amphibolite facies footwall and the granulite facies hanging wall protoliths. Thermal weakening cannot account for such an

- 15 inverse deformation gradient, as syn-deformational P-T estimates for the Petermann Orogeny in the hanging wall and footwall from the same locality are very similar. The distribution of pseudotachylytes, which acted as preferred nucleation sites for shear deformation, also cannot provide an explanation, since these fault rocks are especially prevalent in the immediate hanging wall. The most likely reason for the inverted deformation gradient across the Woodroffe Thrust is water-assisted weakening due to the increased, but still limited, presence of aqueous fluids in the footwall. We also infer that the relative increaseestablish
- 20 <u>a qualitative increase in the abundance of fluids in the footwall along a ca. 60 km section in the direction of thrusting, together with a the slight decrease in the temperature of mylonitization (ca. 100 °C), was responsible. These changes in ambient conditions are accompanied by an for the observed 6-fold decrease in thickness (from ca. 600 m to 100 m) of the Woodroffe Thrust mylonitic zone.</u>

1 Introduction

- 25 Continental fault and shear zone systems (cf., Ramsay, 1980) with displacements on the order of several tens to hundreds of kilometres generally show an asymmetric mylonite distribution across the main fault horizon that is opposite for reverse faults or thrusts and normal faults or detachments. Fault zones are predicted to become more viscous and broaden with depth (e.g., Fossen and Cavalcante, 2017; Handy et al., 2007; Mancktelow, 1985; Passchier and Trouw, 2005; Platt and Behr, 2011b). Juxtaposition of initially different crustal levels should therefore result in a geometry that, for a thrust, preferentially preserves
- 30 the broader ductile mylonite zone in the hanging wall, whereas, for a detachment, it should be in the footwall (e.g., Mancktelow,

1985, his Fig. 11; Passchier, 1984, his Fig. 2). This model is valid for many large-scale fault and shear zone systems, for example: the Moine Thrust Zone, NW Scotland (Christie, 1963; Coward, 1980), the Alpine Fault, New Zealand (Cooper and Norris, 1994; Sibson et al., 1981), the Saint-Barthélemy Massif shear zone (Passchier, 1984), the Simplon Fault, central European Alps (Mancktelow, 1985), the Grizzly Creek shear zone, Colorado (Allen and Shaw, 2011) and the Whipple

- 35 mountains detachment fault, south-western U.S. (Davis, 1988; Davis and Lister, 1988). The mid- to lower crustal Woodroffe Thrust of central Australia (Major, 1970) is an example that does not fit this model and predominantly developed a broader mylonite zone in the footwall (Bell and Etheridge, 1976; Camacho et al., 1995; Flottmann et al., 2004). An interpretation of the Woodroffe Thrust as an original detachment that was later re-oriented and exploited as a thrust can be excluded, both because the metamorphic grade decreases in the direction of tectonic transport and because field mapping shows that the fault
- 40 zone steepens and ramps down towards the internal part of the orogen, against the transport direction (Wex et al., 2017). Passive transport and thermal weakening also cannot account for the inverse deformation gradient, as there is no evidence for late brittle movement on the thrust plane and syn-deformational P-T estimates in the hanging wall and footwall from the same locality are very similar (Wex et al., 2017). Bell and Etheridge (1976) and Camacho et al. (1995) proposed that the inverted distribution of ductile deformation is explained by the difference in bulk water content between the upper amphibolite facies
- 45 (1.0 wt%) footwall and the granulite facies (0.2 wt%) hanging wall, reflecting the metamorphic conditions in the protolith prior to thrusting. Similarly, the preferential formation of shear zones in regions where the host rock mineralogy had previously been modified by fluid-rock interaction has, for example, also been documented in the Whipple mountains detachment fault in SE California (Selverstone et al., 2012) and the Neves area of the Tauern Window in the <u>E</u>eastern Alps (Mancktelow and Pennacchioni, 2005). In this paper, we discuss the control of host rock lithology and, potentially, of fluid activity on the
- 50 distribution of ductile deformation across the Woodroffe Thrust, in an attempt to critically test the local findings of Bell and Etheridge (1976) and Camacho et al. (1995) on a more regional scale (Fig. 1). In particular, we utilize the radiogenic signature of footwall and hanging wall rocks to constrain their respective degree of reworking in the ductile mylonite zone. We also investigate the effect of varying metamorphic temperatures and fluid conditions on the change variation in mylonite thickness over a distance of ca. 60 km parallel to the direction of thrusting.

55 2 Geology

The mid- to lower crustal Woodroffe Thrust outcrops in the Musgrave Block of central Australia (Major, 1970), which is are located within the suture zone of the north, west, and south Australian cratons, which. These cratons were amalgamated into the Australian continent at ca. 1300 Ma in an early stage of development of the supercontinent Rodinia (Myers et al., 1996). The Woodroffe Thrust is developed over an E-W strike length generally interpreted to exceed 600 km and separates the Mulga

60 Park Subdomain in the footwall (to the north) from the Fregon Subdomain in the hanging wall (to the south) (Camacho and Fanning, 1995; Edgoose et al., 1993; Major and Conor, 1993). Exposure of the thrust is poor to inexistent in its proposed western (e.g., Stewart, 1995, 1997) and eastern (e.g., Edgoose et al., 2004) prolongations, but is generally excellent for ca. 150

km in the central Musgrave Block (Bell, 1978; Camacho et al., 1995; Collerson et al., 1972; Wex et al., 2017), where the current study was conducted. In this region, both footwall and hanging wall predominantly consist of granitoids (more common

65 in the footwall) and quartzo-







Figure 2: Photograph and schematic sketch of a cross section through the Woodroffe Thrust at Kelly Hills. Ductile deformation is almost entirely concentrated in the immediate footwall (Mulga Park Subdomain), developing a sequence of protomylonites, mylonites and ultramylonites, with the degree of mylonitization decreasing into the footwall. In contrast, the mostly unaffected or only weakly foliated hanging wall (Fregon Subdomain) is characterized by ubiquitous and voluminous pseudotachylyte (pst) veins and breccias. Further into the hanging wall, a dip slope, characterized by a second zone of highly abundant unsheared pseudotachylyte, has also been documented. Photograph coordinates: 131.45077, -25.89823.

feldspathic gneisses (more common in the hanging wall), with subordinate metadolerites, mafic gneisses and metapelites (Fig.

- Rare quartzites, amphibolites and schists are restricted to the footwall (Camacho and Fanning, 1995; Collerson et al., 1972; Major, 1973; Major and Conor, 1993; Scrimgeour and Close, 1999; Young et al., 2002). Protoliths are inferred to have been felsic volcanics, sediments and intrusives with depositional or emplacement ages around ca. 1550 Ma (Camacho, 1997; Camacho and Fanning, 1995; Gray, 1977, 1978; Gray and Compston, 1978; Maboko et al., 1991; Major and Conor, 1993; Sun and Sheraton, 1992). These protoliths were regionally deformed and metamorphosed at upper amphibolite facies (Mulga Park Subdomain) to granulite facies (Fregon Subdomain) conditions during the ca. 1200 Ma Musgravian Orogeny (Camacho, 1997;
- Camacho and Fanning, 1995; Gray, 1978; Maboko et al., 1991; Sun and Sheraton, 1992), and syn- to post-tectonically intruded by the Pitjantjatjara Supersuite granitoids between ca. 1170-1130 Ma (Camacho, 1997; Camacho and Fanning, 1995; Scrimgeour et al., 1999; Smithies et al., 2011). Peak temperature and pressure conditions during the Musgravian Orogeny are controversial. Earlier estimates were ca. 820-900 °C and 1.1-1.4 GPa (Ellis and Maboko, 1992; Maboko et al., 1989, 1991)
- 85 but more recent studies have given conditions of ca. 800-1000°C and 0.6-0.9 GPa



Figure 3: Sharp contact between the ultramylonites of the Woodroffe Thrust (below) and the largely undeformed felsic granulite in the hanging wall (above). Photograph looking perpendicular to the direction of thrusting. Outcrop SW13-135 (coordinates:
131.87939, -26.21188; location 12 in Fig. 1).

(Camacho, 1997; Tucker et al., 2015; Walsh et al., 2015). Subsequent to this regional-scale tectono-metamorphic phase, the area experienced bimodal magmatism (Giles Complex, including the Alcurra Dolerite swarm) between ca. 1080-1050 Ma and mafic magmatism (Amata Dolerite) at ca. 800 Ma (Ballhaus and Glikson, 1995; Camacho et al., 1991; Clarke et al., 1995; Edgoose et al., 1993; Evins et al., 2010; Schmidt et al., 2006; Sun et al., 1996; Zhao et al., 1994; Zhao and McCulloch, 1993).

- 95 In the area considered in the current study, gabbro-norite intrusions, correlated to the Giles Complex, are restricted to the Fregon Subdomain. Except for these periods of magmatic intrusion, the central Musgrave Block remained largely unaffected by tectonic events between the Musgravian Orogeny at ca. 1200 Ma and the Petermann Orogeny at ca. 560-520 Ma (Camacho and Fanning, 1995; Maboko et al., 1992). Earlier studies by Maboko et al. (1989; 1991) and Ellis and Maboko (1992) proposed that the Musgravian and Petermann Orogenies represented successive stages of a single counterclockwise P-T-t path, whereas
- 100 Camacho (1997) and Camacho et al. (1997; 2015) suggested that the Musgrave Block was partially exhumed prior to the onset of the Petermann Orogeny and then reburied.

The Petermann Orogeny produced a number of large-scale mylonitic shear zones, amongst which the Woodroffe Thrust is the most prominent. Ductile deformation during top-to-north thrusting along the Woodroffe Thrust was largely accommodated within the Mulga Park Subdomain footwall (Bell and Etheridge, 1976; Camacho et al., 1995; Flottmann et al., 2004) and is

- 105 characterized by mylonites with varying degrees of strain, ranging from protomylonites to ultramylonites (Fig. 2), anastomosing around low-strain domains on the metre- to kilometre-scale. The syn-kinematic conditions of deformation along the Woodroffe Thrust were constrained by conventional geothermobarometry to be within the range of 520-650 °C and 0.8 1.3 GPa (Wex et al., 2017) and thus lower temperaturegrade when compared to peak conditions during the earlier Musgravian
- Orogeny. The metamorphic assemblage in both mylonitic samples (where the metamorphic aggregate developed synkinematically) and in statically overprinted undeformed samples (e.g. some Alcurra and Amata Dolerite dykes) record similar metamorphic conditions, consistent with ambient mid- to lower crustal levels (Wex et al., 2017). Stable mineral assemblages during the Petermann Orogeny in felsic units of the Woodroffe Thrust comprise (decreasing modal abundance from left to right) Qz + Pl + Kfs + Bt + Ilm ± Grt ± Ep ± Ms ± Ky ± Cpx ± Hbl ± Rt ± Ttn ± Mag ± Cal, whereas mafic units consist of Pl + Ilm ± Cpx ± Grt ± Rt ± Opx ± Bt ± Hbl ± Ky ± Mag ± Qz ± Kfs ± Cal (Wex et al., 2017). Mineral abbreviations are after
- 115 Whitney and Evans (2010). The degree of mylonitization progressively decreases into the footwall but shows a very abrupt transition into the immediate, dominantly brecciated hanging wall (Figs. 2, 3), which is characterized by ubiquitous and voluminous pseudotachylyte veins and breccias (Camacho et al., 1995; Lin et al., 2005). Even though this upper boundary of the mylonites is discrete or rapidly transitional in the field (Fig. 3), it does not necessarily represent the original boundary between the Mulga Park (footwall) and Fregon (hanging wall) Subdomains. In the Amata area (western edge of Fig. 1), Bell
- 120 (1978) reported up to 250 m of marginal hanging wall reworking into the mylonite zone. However, it remains uncertain how this value was exactly determined, since hanging wall and footwall mylonites are very similar in their field appearance.

3 Methods and general approach

The hanging wall, footwall and numerous transects across the Woodroffe Thrust have been studied and sampled along a N-S traverse, parallel to the direction of thrusting. The sampling locations are reported in Fig. 1. Thin sections of the sampled

- 125 mylonites were prepared from rock chips cut perpendicular to the foliation and parallel to the stretching lineation, and analysed using standard polarized light and scanning electron microscopy. Firstly, the distribution of ductile deformation along and across the Woodroffe Thrust was characterized by quantifying (1) the regional variation in the maximum thickness of the mylonitic zone and (2) the associated degree of hanging wall and footwall reworking. Secondly, field and thin section observations were compiled to assess in a qualitative manner (3) the presence/absence of fluids during deformation, and (4)
- 130 the regional variability in modal abundance of hydrous minerals in felsic units. Potential correlations between parameters (1) to (4) are then discussed. Sample/outcrop coordinates are given in the world geodetic system (WGS) 1984. Orientation measurements of structural elements are corrected for magnetic declination. A detailed description of all utilized methods is given in the Supporting Information S1.

4 Mylonite thickness

135 Field observations indicate that the thickness of the Woodroffe Thrust mylonites is variable. In a section perpendicular to strike, the thickness (T) of the mylonitic zone across the Woodroffe Thrust was calculated by trigonometry (Fig. 4) from: (1i)the angle of dip (α) of the thrust (measured in the field and averaged for each transect); (2ii) the respective difference in elevation (p) (derived from the 30x30 m digital elevation model ASTER) between the lower and upper structural boundaries of the mylonites to the unsheared country rocks (determined from field observations and remote sensing); and (3iii) the 140 apparent thickness (q) (derived from remote sensing). The upper boundary of the mylonitic zone is easily recognized (Fig. 3). The lower boundary is determined by the first appearance of Petermann mylonitic foliation, well characterized by the approximate N-S trend of the stretching lineation and by the top-to-north kinematic indicators. Errors for parameters α and q are considered negligible, whereas the 30x30 m resolution of the digital elevation model is prone to introduce an uncertainty on the order of 10-20 m. The geometrical arrangement for the estimate of T was the same for locations 2-8 and 12-14 (Fig. 145 4a), but slightly different for location 11 (Fig. 1), where the lower structural boundary of the mylonites was at a higher elevation than the upper boundary (Fig. 4b). The thickness of the mylonitic zone at location 1 (Fig. 1) was not calculated, because the Woodroffe Thrust is only exposed along strike. The results for all other studied transects are summarized in Table 1 and, independent of local variability, indicate a gradual increase in mylonite thickness from ca. 100 m in the north to ca. 600 m in the south.



155



Figure 4: Schematic illustration of the trigonometry applied to quantify the true thickness of the Woodroffe Thrust mylonitic zone in a section perpendicular to strike. The parameters are defined in the main text. (a) Geometry applicable to locations 2-8 and 12-14 of Fig. 1. (b) Geometry applicable to location 11 of Fig. 1.

location	coordinates	(WGS 1984)	trigonometrical parameters						
(Fig. 1)	longitude	Latitude	α (°)	<i>p</i> (m)	<i>q</i> (m)	<i>T</i> (m)			
2	131.454	-25.845	27	22	150	88			
3	131.452	-25.855	43	54	190	169			
4	131.449	-25.873	36	23	160	113			
5	131.442	-25.904	21	44	130	88			
6	132.143	-25.992	29	62	160	132			
7	131.644	-25.963	17	110	250	178			
8	131.663	-25.999	10	60	130	82			
11	131.926	-26.177	28	61	800	322			
12	131.879	-26.212	24	153	430	315			
13	131.844	-26.253	37	124	850	611			
14	131.774	-26.308	25	128	1200	623			

Table 1: Angle of dip (α), elevation difference (p), apparent thickness (q) and true thickness (T) of the Woodroffe Thrust mylonites

160 in the central Musgrave Block.



Figure 5: Airborne thorium (ppm) map of the central Musgrave Block. Data derived from the Australian Geophysical Archive Data Delivery System (GADDS) under www.ga.gov.au/gadds (Percival, 2010). The hanging wall Fregon Subdomain is generally depleted compared to the footwall Mulga Park Subdomain. Granitoid intrusives of the Pitjantjatjara Supersuite often have higher-thorium content, particularly in the footwall (one example surrounded by the blue dashed line), but also in the southern part of the hanging wall. Lower-thorium signatures forming diffuse tongues extending northward into the footwall are attributed to alluvial wash derived from the hanging wall. The distinction in thorium content is no longer evident for the northernmost klippe of the hanging wall in the Kelly Hills area (locations 1-5) and the northernmost part of the Mt. Fraser klippe (location 7).

5 Degree of hanging wall and footwall mylonitization

165

- 170 Mylonites derived from the hanging wall and footwall of the Woodroffe Thrust are very similar in both their field appearance and microstructural-petrographical characteristics. The high degree of recrystallization and general lack of porphyroclasts in the uppermost mylonites at locations 4, 6-8 and 12-14 (Fig. 1) suggest that these samples were potentially derived from the pseudotachylyte-rich Fregon Subdomain, hence indicating local hanging wall reworking. To establish whether mylonites were derived from either the Mulga Park or the Fregon Subdomains, we utilized their thorium concentrations, based on the general
- 175 observation that hanging wall rocks are depleted in thorium compared to footwall rocks (Fig. 5). This contrast is due to (1)

dehydration and melting reactions during the earlier ca. 1200 Ma Musgravian Orogeny, which depleted the granulite facies hanging wall to a greater degree than the upper amphibolite facies footwall (Heier and Adams, 1965; Lambert and Heier, 1967, 1968), (2) the predominance of granitoids rather than gneisses in the footwall, and (3) the fact that low-thorium Giles Complex gabbro-noritic intrusions are commonly exposed in the immediate hanging wall of the Woodroffe Thrust. Deformation and

- 180 metamorphism during mylonitization did not significantly alter the original thorium content of the rocks, since the thoriumbearing phases, such as zircon, allanite, monazite and apatite did not break down during the Petermann Orogeny. Consequently, the original variation in thorium between the granulite and amphibolite facies rocks was preserved (Fig. 5). Anomalies in this broad pattern are present locally and can be attributed to hanging wall-derived alluvial sediments in the immediate footwall (lower-thorium anomaly) and to granitoid intrusions of the Pitjantjatjara Supersuite (higher-thorium anomaly), which are syn-
- to post-Musgravian upper amphibolite to granulite facies metamorphism (Scrimgeour and Close, 1999; Young et al., 2002) (Fig. 5). The contrast between a lower-thorium hanging wall and a higher-thorium footwall is well-defined on the airborne thorium map for the central and southern locations, but less evident in the north (Fig. 5). Thorium measurements were carried out on thin section chips of felsic gneisses and granitoids via γ -ray spectrometry, with the measurements run until the error (2 x standard deviation) was below 10 %. The method is outlined in detail in the Supporting Information S1.

190 5.1 Thorium concentration in felsic units

Thorium contents of felsic gneisses and granitoids (Table 2) have been compiled from the current study as well as from previous studies of nearby areas (Camacho, 1997; Young et al., 2002). These data have been grouped into (1) the central and southern locations and (2) the northern locations, based on the airborne thorium map, as introduced above.

In group (1), samples unequivocally attributed to either the hanging wall or footwall of the Woodroffe Thrust were grouped together. This was done on the basis of their geographical position, with respective samples originating either from far into the non-mylonitized hanging wall and footwall or from the lowermost protomylonitic part of the Woodroffe Thrust. In the hanging wall, thorium concentrations range from 1 to 28 ppm in felsic gneisses and up to 63 ppm in granitic intrusions (higher-thorium anomalies), but are in both cases usually lower than 8 ppm. In the footwall, concentrations vary between 2 and 195 ppm and are typically higher than 10 ppm (Table 2). These concentrations are in accord with the regional-scale contrast in thorium

- 200 concentrations across the Woodroffe Thrust in the central and southern locations (Fig. 5), as well as with the results of Lambert and Heier (1968), who determined concentrations of 2.1 ppm for the granulite facies hanging wall and 11 ppm for the upper amphibolite facies footwall. Based on the compilation in Table 2, samples with thorium content <8 ppm were assigned to the hanging wall and samples with higher values to the footwall of the Woodroffe Thrust. In general, samples taken close to the boundary between the mylonites and unsheared rocks of the Fregon Subdomain (typically the uppermost few tens to one
- 205 hundred metres) were investigated. For the majority of samples, the assignment was straightforward since the inferred hanging wall samples are extremely low in thorium (<3 ppm), whereas most inferred footwall samples have values >8 ppm. Exceptions are samples SW14-243 and SW13-159, which have intermediate concentrations of 8 ppm and 6 ppm, respectively. Both

samples were assigned to the footwall since subsequent samples further towards the hanging wall, respectively SW14-244 and SW13-161, could clearly still be attributed to the footwall. Alternatively, samples SW14-243 and SW13-159 could reflect imbrication of the Mulga Park and Fregon Subdomains, but this is not supported by any field observation.

210

In group (2), thorium concentrations vary between 11 and 49 ppm (Table 2) but do not allow a clear distinction between samples derived from the footwall and hanging wall in a manner similar to the central and southern locations. This result is in accord with the airborne thorium concentrations, which also do not indicate a significant jump across the Woodroffe Thrust in these more northerly locations (Fig. 5).

	1	nanging wall	of the Woodr	offe Thrust			
location	مسساه	lith al a my	coordinates	(WGS 1984)) data	Th	
(Fig. 1)	sampie	ntnology	longitude	latitude	source	(ppm)	
9	SW14-029A	granite	131,74496	-26,00093	1	6	
9	SW14-025	granite	131,73269	-26,01569	1	3	
9	SW14-030B	granite	131,74295	-26,00222	1	5	
17	FW13-173	granite	131,54138	-26,34104	1	3	
18	FW13-228	granite	131,56005	-26,37413	1	28	
-	MP-2	granite	131,74845	-25,99871	3	10	
-	MP94/500	granite	131,73549	-25,99535	3	4	
-	MP94/501	granite	131,74105	-25,99389	3	11	
-	W-12	granite	131,55670	-26,38583	2	19	
-	W-34d	granite	131,52310	-26,34528	2	6	
-	W-70	granite	131,63910	-26,40186	2	4	
-	W-96	granite	131,69440	-26,38511	2	8	
-	W-104	granite	131,70010	-26,38119	2	4	on
-	W-127	granite	131,90190	-26,43750	2	7	cati
-	W-146	granite	131,87670	-26,35469	2	6	lo
-	W-148	granite	131,88030	-26,34653	2	10	cal
-	W-199a	granite	131,77080	-26.47550	2	54	phi
-	W-199b	granite	131,77080	-26,47550	2	63	gra
-	E-37	charnockite	132,27078	-26,19010	2	6	seo
-	E-38	charnockite	132,26899	-26,18286	2	3	u g
-	E-39	charnockite	132,14968	-26.22028	2	2	g
-	W-32	charnockite	131,46100	-26,29419	2	26	Dase
-	W-129	charnockite	131,89100	-26,38531	2	8	n h
16	NW13-043	felsic gneiss	131 49727	-26 28058	1	4	atic
16	SW13-032	felsic gneiss	131,49771	-26,28135	1	2	lei
19	NW13-016B	felsic gneiss	131,71053	-26.38530	1	28	Ö
20	NW13-026	felsic gneiss	131,66087	-26,41038	1	20	
20	NW13-030	felsic gneiss	131,66258	-26,40675	1	1	
-	F-27	felsic gneiss	132,14091	-26,18857	2	3	
-	E-47	felsic gneiss	132.04583	-26,29111	2	2	
-	MP97/513	felsic gneiss	131,66461	-25,97300	3	12	
-	W-16dg	felsic gneiss	131,52810	-26.26806	2	6	
-	W-16ug	felsic gneiss	131,52810	-26.26806	2	8	
-	W-18	felsic gneiss	131,53810	-26,27472	2	21	
_	W-122ada	felsic gneiss	131 69830	-26 28992	2	3	
_	W-122aug	felsic gneiss	131,69830	-26,28992	2	8	
	W-140b	felsic gneiss	131,0000	-26,26772	2	5	
-	W-1400	felsic gneiss	131,84010	-26,43042	2	N/D	
8	SW13_101	felsic gneise	131,66264	-25,00011	1	$\frac{10D}{N/D < 2}$	\vdash
11	SW14_181A	felsic gneiss	131 02505	-26,33311	1	N/D < 2	lc.
11	SW14-181A	felsic gneiss	131 02564	-26,17604	1	1	C01
12	NW12_197D	felsic gneiss	131,92504	-26,17081	1	1	Η
12	SW/13_135A	felsic gneiss	131,87917	-26,21191	1	2	on
12	SW13-135A	felsic gneiss	131,87939	-20,21108	1	2	ed
12	NW13-135D	felsic gneiss	131,87939	-20,21100	1	2	bas
12	SW14-2464	felsic gneiss	131 84444	-26,21191	1	2	E.
1/	SW13_164	felsic gneiss	131 77/62	-26 30772	1	2	1 S
14	SW15-104	icisic gueiss	151,77405	-20,50772	1	5	1

footwall of the Woodroffe Thrust										
location (Fig. 1)	sample	lithology	coordinates longitude	(WGS 1984) latitude	data source	Th (ppm)				
10 GW12 1004										
12	SW13-122A	granite	131,87206	-26,20364	1	73	E			
12	SW13-123	granite	131,87229	-26,20379	1	7	atio			
12	NW13-184	granite	131,87823	-26,20914	1	16	loc			
12	NW13-185B	granite	131,87830	-26,20946	1	14	cal			
14	SW13-151	granite	131,77438	-26,30117	1	20	hic			
14	SW13-153	granite	131,77411	-26,30173	1	N/D < 2	grap			
-	MP94/502	granite	131,78418	-25,95562	3	55	eog			
-	MP94/503	granite	131,62926	-25,86096	3	30	00 00			
-	BJ96/178	granite	131,48641	-25,78867	3*	18	op			
-	BJ96/201	granite	131,39771	-25,79945	3*	195	ase			
-	BJ96/276	granite	131,44734	-25,97263	3*	18	n p			
-	MP96/505	granite	131,64751	-25,95006	3*	35	tio			
-	MP96/509	granite	131,64996	-25,94452	3	12	rela			
12	SW13-125	felsic gneiss	131,87338	-26,20470	1	27	con			
-	MP97/43	felsic gneiss	131,81710	-25,98836	3	10	-			
8	GW13-415	felsic gneiss	131,66256	-25,99928	1	15				
8	SW13-193	felsic gneiss	131,66256	-25,99928	1	16	onc			
8	SW13-192	felsic gneiss	131,66258	-25,99915	1	23	1 C			
11	SW14-179	felsic gneiss	131,92603	-26,17661	1	15	E			
12	SW13-134	felsic gneiss	131,87913	-26,21151	1	9	lor			
13	SW14-243	felsic gneiss	131,83942	-26,25476	1	8	isec			
13	SW14-244	felsic gneiss	131,83949	-26,25472	1	13	; p			
14	SW13-159	felsic gneiss	131,77375	-26,30666	1	6	our			
14	SW13-161	felsic gneiss	131,77445	-26,30759	1	14				
		-								

data source:

location

(Fig. 1)

6

6

3

2

2

3

4

6

6

1 this study

2 Camacho (1997)

3 Young et al. (2002)

 values erroneously published in Young et al. (2002), but correctly presented here (personal communication A. Camacho)

northern studied locations

N/D below detection limit

SW13-322

NW13-134A

SW13-173A

SW13-173C

FW13-093

SW14-006

SW13-323A

SW14-034A

granite

samplelithologycoordinates (WGS 1984)
longitudedata
sourceSW13-321granite132,14333-25,991781

felsic gneiss 131,45284

felsic gneiss 131,45334

felsic gneiss 131,45334

felsic gneiss 131,45213

felsic gneiss 131,45047

felsic gneiss 132,14344

felsic gneiss 132,14337

132,14340

-25,99191

-25,85477

-25,84543

-25,84543

-25,85455

-25,87174

-25,99211

-25,99224

Th

(ppm) 23

49

34

17

31

11

20 19

12

1

1

1

1

1

1

1

1

2	1	5
4	T	\mathcal{I}

Table 2: Thorium concentrations in felsic gneisses and granitoids from the hanging wall and footwall of the Woodroffe Thrust, central Musgrave Block.



220 Figure 6: Sample-specific thorium (Th) concentrations (measured by γ-ray spectrometry) plotted across the Woodroffe Thrust (WT) mylonites at locations 8 and 11-14 (Fig. 1). The original boundary between the lower-thorium hanging wall and higher-thorium footwall is inferred (red line).

5.2 Determination of the hanging wall-footwall boundary

- 225 Measured thorium concentrations indicate that in locations 8 and 11-14 (Fig. 5) the uppermost mylonites of the Woodroffe Thrust developed in the lower-thorium Fregon Subdomain (Table 2). These results are in agreement with the proposed identification of hanging wall reworking based on field appearance. Similar field relationships, such as progressive downwards mylonitization of units clearly forming part of the hanging wall, also indicate limited reworking of the Fregon Subdomain at locations 4 and 6 (Fig. 5). However, the lack of a clear contrast in thorium concentrations between footwall and hanging wall
- 230 in these locations precludes any verification of these field observations, as well as any independent determination of the thickness of hanging wall-derived mylonites in the thrust zone. We therefore excluded these northern transects when using thorium concentration as a proxy to map the original boundary between the hanging wall and footwall (Fig. 6). It was also not possible to precisely determine the boundary at location 13 (Fig. 1), due to a large lateral gap between the last sample assigned to the footwall and the first sample assigned to the hanging wall. In an attempt to quantitatively calculate the degree of hanging
- 235 wall/footwall reworking into the total mylonite zone, we applied trigonometry based on the geometrical arrangement sketched in Fig. 4a. The only modification was that the lower mylonite boundary was now defined by the newly reconstructed boundary between hanging wall and footwall. Results are summarized in Table 3. In contrast to the 250 m proposed by Bell (1978) for the Amata area (western edge of Fig. 1), our results indicate that only the lowermost 3 m of the Fregon Subdomain were reworked into the Woodroffe Thrust mylonites at location 8, increasing up to 18-40 m at locations 11-14 (Fig. 1). These values
- represent 3-6 % of the entire thickness of the Woodroffe Thrust mylonites (Table 1) at locations 8, 11 and 14 and up to 13 % at location 12 (Fig. 1). However, the difference in elevation (*p*) is not well defined over short distances (*q*) given the limited (30x30 m) resolution of the digital elevation model (ASTER). This can introduce a significant uncertainty into the calculation of the degree of hanging wall reworking. Nevertheless, our analysis clearly shows that the majority of mylonites developed in the Mulga Park Subdomain (footwall) rather than the Fregon Subdomain (hanging wall).

location	coordinates	(WGS 1984)	trigonometrical parameters						
(Fig. 1)	longitude	latitude	a (°)	<i>p</i> (m)	<i>q</i> (m)	<i>T</i> (m)			
8	132.143	-25.992	10	1	10	3			
11	131.926	-26.177	28	5	30	18			
12	131.879	-26.212	21	22	55	40			
13	131.844	-26.253	-	-	-	>0			
14	131.774	-26.308	22	9	30	20			

245

Table 3: Angle of dip (α), elevation difference (p), apparent thickness (q) and true thickness (T) of the Woodroffe Thrust mylonites derived from the hanging wall.

6 Presence or absence of fluids during mylonitization

250 The syn-deformational presence or absence of fluids in the study area is established from a series of field and thin section observations. These include the regional variation in: (1) syntectonic quartz veins, (2) introduction of carbon and (3) plagioclase stability and breakdown.

6.1 Quartz veins

Syntectonic quartz veins (Fig. 7a) and associated quartz-rich pegmatite dykes are uncommon throughout the field area, being generally absent in the southern locations and only locally present in the northern locations (Fig. 8a). These quartz veins crosscut the mylonitic fabric but were themselves variably deformed during subsequent shearing, providing direct field evidence that they were broadly coeval with the Woodroffe Thrust, and thus associated with the Petermann Orogeny. Sense of shear is both top-to-north and top-to-south, which is contrary to the dominant top-to-north shear sense associated with the Woodroffe Thrust, but has also been documented by Bell and Johnson (1992) in the Amata region (western edge of Fig. 1).
Quartz veins are boudinaged within the mylonitic foliation and, although deformed, did not preferentially localize strain (Figs.

6.2 Introduction of carbon

7a,b).

Finely dispersed calcite is locally found with very low modal abundance (typically <1 %) in the otherwise non-carbonaceous rocks of the central Musgrave Block (Fig. 7c). Calcite-bearing samples are present throughout the study area, but are generally more common in the north (Fig. 8b). Microstructures indicate that the fine-grained (<10 µm) calcite nucleated during shearing (Fig. 7c). Calcium was made available from dynamic "recrystallization" of plagioclase to new grains with lower anorthite content (Wex et al., 2017), but carbon cannot have an immediately local source, because the studied rocks were initially entirely non-carbonaceous. In the attempt to place further constraints on the origin of this fluid, carbon and oxygen isotopes of calcite were measured (Supporting Information S2), yielding mean average values of -4.1 ‰ for δ¹³C_{Cal} (V PDB) and +10.1 ‰ for δ¹⁴O_{Cal} (SMOW). Within the same samples, the whole rock isotopic signature δ¹³C_{whole rock} (V PDB) is always lower (on average by 2.7 ‰) than the corresponding δ¹³C_{Cal} values.



Figure 7: Field and thin section images providing evidence for the presence or absence of fluids during deformation. (a) Quartz veins (encircled) crosscut the mylonitic foliation of the Woodroffe Thrust showing a sigmoidal shape consistent with rotation during

- 275 overall top-to-north shearing. The quartz vein at the bottom of the picture is boudinaged and did not localize deformation. Outcrop NW15-264 (coordinates: 131.46883, -25.83040; location 1 in Fig. 1). (b) Quartz vein adjacent to a sinistral shear zone that is not reactivated, even though shear zone and quartz vein are almost parallel to each other. Abbreviation: S_m = mylonitic foliation. Outcrop SW13-200 (coordinates: 131.73810, -25.99564; location 9 in Fig. 1). (c) Finely dispersed calcite (high birefringence) between recrystallized feldspar grains under crossed-polarized light. Thin section is not oriented. Sample NW14-423A (coordinates:
- 131.84368, -26.11423; location 10 in Fig. 1). (d) Plagioclase clast with muscovite (higher birefringence) and epidote (lower birefringence) inclusions under crossed-polarized light. Thin section is oriented N-S (left-right). Sample SW14-214A (coordinates: 131.44416, -25.90284; location 5 in Fig. 1). (e) Plagioclase clast with epidote inclusions under crossed-polarized light. Thin section is oriented N-S (left-right). Sample GW13-415 (coordinates: 131.66256, -25.99928; location 8 in Fig. 1). (f) Plagioclase clast with kyanite inclusions (greenish needles) and neo-crystallized garnet under plane-polarized light. Thin section is not oriented. Sample SW13-
- 285 167 (coordinates: 131.77475, -26.30845; location 14 in Fig. 1). (g) Inclusion-free plagioclase clast under plane-polarized light. Thin section is oriented NNE-SSW (left-right). Sample SW13-318 (coordinates: 132.14311, -25.99142; location 6 in Fig. 1).

6.3 Plagioclase stability and breakdown

Plagioclase dynamically recrystallized in the Woodroffe Thrust mylonites and associated shear zones (Bell and Johnson, 1989), forming typical "core-and-mantle structures". The composition of the newly recrystallized grains ranges from albite/oligoclase in felsic rocks to oligoclase/andesine in mafic rocks (Wex et al., 2017). Mineral inclusions within plagioclase clasts are common and allow the distinction of four different types of clasts, respectively termed microstructures 1 to 4:

- (1) Plagioclase studded with abundant inclusions of epidote and muscovite (Fig. 7d). This type is, with one exception, restricted to the northern locations (Fig. 8c).
- (2) Plagioclase containing only epidote inclusions (Fig. 7e), with a modal abundance far lower than that of epidote + muscovite inclusions of microstructure 1. This microstructure is restricted to the central locations (Fig. 8c).
- (3) Plagioclase crowded with kyanite needle inclusions (Fig. 7f) (see Supporting Information B of Wex et al. (2017) for identification techniques). This microstructure is restricted to the southernmost locations (Fig. 8c).
- (4) Plagioclase free of inclusions (Fig. 7g). This microstructure is found in almost all locations (Fig. 8c).

Figure 8c shows that the type of inclusions in plagioclase varies in a N-S direction, i.e. parallel to the tectonic transport direction 300 of the Woodroffe Thrust. From north to south, inclusions progressively change from muscovite + epidote (microstructure 1), to epidote (microstructure 2), and to kyanite (microstructure 3), with inclusion-free plagioclase clasts (microstructure 4) occurring throughout. There is no apparent variation of the type of plagioclase inclusions along strike of the Woodroffe Thrust (i.e. E-W).

290

(a) development of syntectonic quartz veins



305

Figure 8: Regional variation in the development of (a) syntectonic quartz veins, (b) the introduction of carbon and (c) plagioclase stability and breakdown, each plotted against sample latitude. The regional temperature gradient (Wex et al., 2017) and the position of the studied locations (Fig. 1) are indicated.

7 Abundance of hydrous minerals

- 310 Pseudotachylytes have been identified as preferred discontinuities for nucleation of ductile shear zones and strain localization under mid- to lower crustal conditions (Andersen and Austrheim, 2006; Austrheim and Andersen, 2004; Hawemann et al., 2014, 2018; Lund and Austrheim, 2003; Menegon et al., 2017; Passchier, 1982; Pennacchioni and Cesare, 1997; Pittarello et al., 2012; Wex et al., 2014, 2017). In order to study the control of host rock mineralogy on shear initiation, the modal abundance of hydrous minerals in deformed and undeformed pseudotachylytes (representative compilation in Supporting Information S3)
- 315 from felsic footwall and hanging wall samples was determined by image analysis of backscattered electron (BSE) images (method outlined in the Supporting Information S1). The different amounts of hydrous minerals between in the pseudotachylytes should reflect compositional variations between in the host rocks from which they formed. The results, summarized in Table 4, indicate that matrix mineral assemblages of felsic pseudotachylytes from the hanging wall and footwall are similar to each other. These assemblages dominantly comprise Pl + Kfs + Qz + Bt + Mag + Ilm with individual samples
- also containing Ep, Grt, Cpx, Opx, Ky, Ms, Rt or Hbl. However, there is a strong contrast with regard to the modal abundance of hydrous minerals in the studied pseudotachylytes. Samples have been grouped according to their latitude coordinates, revealing that: (i1) the abundance of hydrous minerals decreases from north to south in both the hanging wall and the footwall; and (ii2) at similar latitude, the footwall rocks are more hydrous than the hanging wall rocks. Two samples (SW14-029A and SW14-179) do not fit this regional trend and these outliers have been excluded in calculating the mean values, as they mask
- 325 what are otherwise clear trends.

1			hang	ging wa	dl of the Woo	droffe Thrust						ę	otwall of the Woo	odroffe Thrust		
sample		ənitsine	E. spesteq	BSE (coordinates WGS 1984)	mineral assemblage (pseudotachylyte)	hydroi minerals	us (%)	location (Fig. 1)	sample	oniteirq Sheared	in B	SE coordinates ges (WGS 1984)	mineral assemblage (pseudotachylyte)	hydroi minerals	sn (%)
SW13	-174A	×		3	131,45304 -25,84532	$\begin{array}{l} Pl + Kfs + \underline{Bt} + Cpx \\ + Qz + Mag + llm \end{array}$	10 ± 2		7	SW13-171	x		131,45397 -25,84535	$Qz + \underline{Bt} + \underline{Ms} + \underline{Ep}$ + Mag + P1 ± Kfs	40 ± 8	
NW 13	-139	×		ŝ	131,44826 -25,85366	Kfs + PI + Mag + <u>Bi</u>	4±2	/0.01	2	SW13-171	~		131,45397 -25,84535	$\begin{array}{l} Qz + \underline{Bt} + \underline{Ms} + \overline{Ep} \\ + Mag + Pl \pm Kls \end{array}$	4 0±2	
FW13	-096	×		5	131,45034 -25,85414	$PI + Kfs + \underline{Bt} + \underline{HbI}$ $+ \underline{Lp} + Mag$	7±1	<u>av. 10 %</u>	ŝ	NW13-134A	~		131,45284 -25,85477	PI + Qz + KGs + Lip + Lip + Lit	20 ± 3	av. 30%
SW13	-102	×		4	131,45164 -25,85433	$PI + \underline{Bt} + Opx + Qz + Mag$	17±4		m	FW13-093	~		131,45213 -25,85455	$\begin{array}{l} PI+Qz+Kfs+\underline{HbI}\\ +\underline{Bi}+Mag+Ilm \end{array}$	29±1	
									S	SW14-228A	x		131,44158 -25,90365	PI + Kfs + Ep + Bt + Mag ± Ilm	20 ± 1	
SW13	161-	×		4	131,66264 -25,99911	$\begin{array}{l} Pl+Cpx+\underline{Bl}+Qz\\ +Opx+Mag \end{array}$	16±1		9	SW13-321			132,14333 -25,99178	$\begin{array}{l} Pl + Kls + Qz + \underline{Bl} + \overline{Ep} \\ + Git + Mag + Ilm \end{array}$	13±2	
SW 14	H029B		х	ŝ	131,74496 -26,00093	P1 + Kfs + <u>bt</u> + Qz + Grt + <u>Fp</u> + Mag	13 ± 2		9	SW13-323A	х	4	132,14344 -25,99211	PI + Kfs + <u>13t</u> + Qz + Mag + Tlm	10 ± 2	
SW14	H029A	×		3	131,74496 -26,00093	$\frac{Hbl}{+Qz + \underline{Bt} + Ilm}$	$33 \pm 5^{*}$	av. 8 %	9	SW13-323A	x		132,14344 -25,99211	$\begin{array}{l} Pl+Kfs+\underline{Bt}+Qz\\ +Mag+Ilm \end{array}$	10 ± 4	av. 10%
SW1	H-029A	x		5	131,74496 -26,00093	KIs + PI + Qz + $\underline{BI} + Mag$	3 ± 0	[12 %]	9	SW13-323A	x		132,14344 -25,99211	$\begin{array}{l} PI+Kfs+\underline{Bt}+Qz\\ +Mag+Ilm \end{array}$	8 ± 1	
SW14	H025		×	9	131,73269 -26,01569	$\begin{array}{l} Pl+Kfs+Opx+Qz\\ +\underline{Bt}+Mag+Ilm+Cal \end{array}$	2±1		8	SW13-192	~		131,66258 -25,99915	Kfis + PI + Qz+ <u>Bt</u> + <u>En</u> + <u>Hbl</u> + Grt + Mag	8 ± 1	
IWN	3-203		x	3	131,74318 -26,00442	$\begin{array}{l} Kfs + PI + Grt \\ + \underline{Bt} + Qz + Mag \end{array}$	5 ± 1									
SW1	4-181A		×	5	131,92595 -26,17664	$\begin{array}{l} KIs + PI + Qz + Cpz \\ + Ght + Mag + \underline{Bi} \end{array}$	1±2		=	SW14-179	Ŷ		131,92603 -26,17661	$PI + Q_{X} + KI_{S} + H\overline{H}I + G_{H}I + O_{P}X + IIm + Mag \pm BI$	$21\pm6^{*}$	
IWS	4-181B		×		131,92595 -26,17664	$\begin{array}{l} Pl + Kfs + Qz + Cpx \\ + Grt + Ilm \pm \underline{Bt} \end{array}$	∇	1 02	12	SW13-134	^		131,87913 -26,21151	PI + Kfs + Qz + Bt + Opx + Grt + IIm	3 ± 1	
SWI	4-181B		Х	2	131,92595 -26,17664	PI + KIs + Grt + Qz + Cpz + Bt + TIm	3 ± 0	17 T - XP	13	SW14-237C	х	4	131,83544 -26,25303	Kfs + Pl + <u>Bi</u> + Mag	5±1	<u>av. 4 %</u> [7 %]
SW14	H181B	×		a,	131,92595 -26,17664	$PI + Kfs + Grt + Opx + Cpx + Qz + IIm \pm \underline{B}t$	∇		13	SW14-244	~		131,83949 -26,25472	PI + Kfs + Qz + Bt + IIbI + IIbI + IIbI + Ot	7±2	
									14	SW13-159	~		131,77375	$\begin{array}{l} Pl + KIs + Cpx + Git \\ + Qz + Ky + Hm + Rt \end{array}$	0	



^a Extreme outliers (*) are not considered. Mineral assemblages are listed in order of decreasing modal abundance (from left to right) with hydrous minerals underlined. A representative compilation of pseudotachylytes is given in the Supporting Information S3. Errors given are 2 x standard deviation.

330 8 Discussion

8.1 Plagioclase breakdown reactions

Within the range of mid- to lower crustal conditions, as estimated for the Woodroffe Thrust (Wex et al., 2017), the following plagioclase breakdown reactions are relevant:

(1) $An + Or + H_2O = Ms + Ep + Qz$ (Kretz, 1963; Ramberg, 1949)

335

- (2) $An + Grs + H_2O = Ep + Qz$ (Kretz, 1963)
- (3) An = Grs + Ky + Qz (Boyd and England, 1961; Hariya and Kennedy, 1968)
- (4) $An + H_2O = Ep + Ky + Qz$ (Goldsmith, 1982; Kretz, 1963; Ramberg, 1949)

For the sake of simplicity, t<u>T</u>hese reactions proposed in the cited publications are restricted to the system Ca-K-Al-Si-OH, and in particular do not consider the presence of iron. However, iron is certainly necessary to account for the crystallization of
epidote and was potentially derived from relict iron-bearing minerals (e.g., magnetite, ilmenite, biotite or garnet). Microstructure 1 is unequivocally correlated with hydration reaction 1. Reaction 2 is capable of producing microstructure 2, but garnet has been interpreted to serve as a calcium-sink rather than source during the high-pressure Petermann Orogeny (Camacho et al., 2009). Alternatively, we propose that the decrease in anorthite content during plagioclase recrystallization (Wex et al., 2017) provided the necessary source of calcium for producing microstructure 2. Microstructure 3 is correlated with reaction 3. Reactions 3 and 4 represent the high-pressure breakdown of plagioclase under anhydrous and hydrous conditions, respectively. According to Wayte et al. (1989), the transition between the two competing reactions occurs at a

water activity of ca. 0.004 for pressure and temperature conditions similar to those in the more southerly footwall locations. In samples with microstructure 3, epidote is never observed as a secondary inclusion phase together with kyanite (Supporting Information S4), indicating that reaction 3 always prevailed over reaction 4.

350 8.2 Fluid activity

In the southern locations (Fig. 1), plagioclase breakdown by reaction 3 indicates that water was not sufficiently available to drive reaction 4 (Goldsmith, 1980, 1982; Wayte et al., 1989). Consequently, reaction 3 is a good indicator for very low water activities (<0.004; Wayte et al., 1989) and the absence of a free fluid phase during deformation, since only very small amounts of water (ca. 20 ppm) are required for mineral reactions in a solid silicate system (Milke et al., 2013). Similarly, relict

355 plagioclase clasts without any inclusions (microstructure 4) also indicate the absence of a free fluid phase, as the studied rocks were all metastable with respect to hydration reactions 1, 2 and 4. Consequently, any of these breakdown reactions would have rapidly consumed any available free fluid, since all other reactants were present in the studied samples. Vice versa, plagioclase breakdown by reaction 1 clearly indicates the presence of a free fluid phase during deformation. Reaction 2 also involves hydration, as indicated by the marginal presence of epidote in plagioclase clasts (microstructure 2). However, free fluids were

not sufficiently available to facilitate reaction 1. Hence, we favour the interpretation that microstructure 2 indicates largely

360

anhydrous conditions, with only very minor fluid introduction.

Based on the classification above, the regional availability of a free fluid phase during deformation, respectively termed "wet" and "dry", can be determined from the mineral inclusions in plagioclase (Fig. 8c) and the corresponding inferred breakdown reactions (Sect. 8.1). The distinction is purely qualitative and refers only to whether or not free fluid was sufficiently available

- to facilitate the hydrous breakdown reaction 1. Consequently, the studied field area is characterized as dominantly "dry" (microstructures 2, 3 and 4) with a progression towards locally "wet" conditions (microstructure 1) in most of the northernmost exposures (Fig. 9). The regional variation in (1) the development of syntectonic quartz veins (Fig. 8a) and (2) the introduction of carbon (Fig. 8b) are each-both consistent with this interpretation (Fig. 9). However, within a single location, individual samples can have a "wet" microstructure 1 while others still preserve a "dry" microstructure 4 (Fig. 8c), indicating that the
- 370 <u>availability of fluids were only presentvaried</u> on a very local scale. Generally "dry" conditions are also in agreement with the fact that the metamorphic overprint during the Petermann Orogeny occurred under lower grade conditions compared to the earlier Musgravian Orogeny, and thus facilitated water consumption rather than water release.

8.3 Fluid source

Based on the above qualitative estimate of "fluid activity" in the study area, it is evident that aqueous and CO₂-bearing fluids
 were introduced in the northern exposures of the Woodroffe Thrust (Fig. 9). However, t<u>T</u>here does not seem to be any obvious link between the infiltration of local the distribution of aqueous and the CO₂-bearing fluids, since samples that preserve a "wet" microstructure 1 are not necessarily calcite-bearing and vice versa (Figs. 8b,c).

8.3.1 Aqueous fluids

Aqueous fluids infiltrating into the footwall of the Woodroffe Thrust were unlikely to have been derived from the "drier" hanging wall (Table 4) and consequently must originate from units within or underlying the current level of exposure of the footwall. Gneisses and granitoids in the "wet" northern part of the study area are clearly interleaved with and juxtaposed onto the basal units of the Amadeus Basin (Wells et al., 1970), as evident from outcrops of Dean Quartzite (Forman, 1965; Young et al., 2002) immediately west of the Kelly Hills klippe. These Neoproterozoic sedimentary rocks represent an ideal source for aqueous and CO₂-bearing fluids since they were metamorphosed and dehydrated during the Petermann Orogeny (Wells et al.,

385 1970). However, based on regional-scale geometric reconstructions, it has been argued that the Woodroffe Thrust and potentially underlying thrust planes developed in-sequence (Wex et al., 2017). If this is true, then the Dean Quartzite was only imbricated below the northernmost studied locations after movement and deformation on the Woodroffe Thrust had largely

ceased and the dynamic microstructures had already been frozen in. Therefore, we prefer a model in which the aqueous fluids were released from the



390

Figure 9: Projected schematic cross section through the central Musgrave Block. The horizontal scale is compressed by a factor of 3.5 with respect to the vertical scale. The regional temperature gradient is taken from Wex et al. (2017).

granitoids and upper amphibolite facies gneisses within the footwall. Such a model is in agreement with the fact that the regional trend towards higher abundance of hydrous minerals in the north parallels the shift towards "wet" conditions, as indicated by the plagioclase breakdown reactions (Fig. 9). An internal source is also consistent with the conclusion that "wet" conditions were only present on a very local scale and that "wet" and "dry" samples are often preserved in close proximity.

<u>The same is also true of calcite-bearing and calcite-absent samples.</u> <u>Consequently, wWe consider therefore conclude that the</u> studied field area was not <u>pervasively</u> infiltrated by externally derived aqueous<u>-rich</u> fluids and remained a relatively closed system. <u>The source of carbon remains unexplained</u>, even if the quantities involved are very small. As far as currently known,

units of the footwall Mulga Park Subdomain are non-carbonaceous, although small amounts of CO₂ can be present within

400

415

fluid-inclusions. Calcite

8.3.2 CO2 bearing fluids

The fact that the gneisses and granitoids of the central Musgrave Block are entirely non-carbonaceous (Collerson et al., 1972; Major, 1973; Major and Conor, 1993; Scrimgeour and Close, 1999) argues for an external source of the CO₂-bearing fluids.

405 <u>c</u>Carbon and oxygen ($\delta^{13}C_{Cal}$, $\delta^{18}O_{Cal}$) and carbon whole rock ($\delta^{13}C_{whole rock}$) isotropic signatures were measured in order to provide constraints on its nature with the aim of providing better constraints on the source but the results are inconclusive (see <u>Supporting Information S2 for more details</u>). The resultsValues for $\delta^{18}O$ are in agreement with calcite crystallization temperatures of 500-600 °C, while $\delta^{13}C_{Cal}\delta^{13}C$ values are strongly partially rock-buffered (see Supporting Information S2 for more details), and do not allow an unequivocal interpretation of the potential source (Supporting Information S2).

410 **8.4 Distribution of ductile deformation**

The distribution of thorium establishes that the Woodroffe Thrust mylonites preferentially developed in the Mulga Park rather than the Fregon Subdomain (Fig. 6). This conclusion is in agreement with our field observations and with those of Bell and Etheridge (1976), Camacho et al. (1995) and Flottmann et al. (2004). However, it is evident from our results that in the southern locations the lowermost hanging wall was also involved in the mylonitization process (Table 3). We discuss below the potential processes guiding the large-scale distribution of ductile deformation in the Woodroffe Thrust.

8.4.1 Footwall/hanging wall reworking

The proportion of the total shear strain accommodated in the narrow mylonitic to ultramylonitic zones developed in the lowermost hanging wall and uppermost footwall cannot be determined. However, what could be determined in this study is

- the relative thickness of the mylonite zones, using thorium concentrations to distinguish the original hanging wall and footwall protoliths. From this it is established that the hanging wall-derived parts of the Woodroffe Thrust generally make up <10% of the entire width of the mylonitic zone. This preferential development of a broader mylonite zone in the Mulga Park Subdomain footwall rather than in the Fregon Subdomain hanging wall is contrary to the expected simple model of a thrust or reverse fault system (e.g., Mancktelow, 1985; Passchier, 1984). Such a model would predict an asymmetric strain profile where the mylonite zone occurs in the initially "hotter" hanging wall rather than the "colder" footwall. An inverse distribution of ductile</p>
- 425 deformation, as a consequence of asymmetric thermal weakening, would be in agreement with the flower-like structure model proposed by Camacho and McDougall (2000) for the central Musgrave Block. Based on the preservation of pre-Petermann K-

Ar, 40 Ar- 39 Ar and Rb-Sr ages in hornblende, muscovite, biotite and K-feldspar in the undeformed gneissic country rocks, Camacho and McDougall (2000) argued that the Fregon Subdomain was rapidly buried and exhumed in less than 40 Myr. Consequently, these rocks failed to thermally equilibrate to temperatures above 350 °C at a pressure of ~1.2 GPa. With this

430 tectonic model, the hanging wall should have been at temperatures <350 °C (Camacho and McDougall, 2000) and thus should have been colder than the footwall (>500 °C; Wex et al., 2017). In such a model, thermal weakening could indeed account for preferential mylonitization of the Mulga Park Subdomain. However, Wex et al. (2017) and Hawemann et al. (2018) demonstrated that Petermann Orogeny metamorphic assemblages throughout the study area are directly comparable in dynamically and statically recrystallized units, arguing that the estimated metamorphic conditions were ambient and differed

435 little between shear zones and country rock. In addition, Wex et al. (2017) found that the syntectonic metamorphic conditions are similar in the hanging wall and footwall at the same locality, arguing against thermal weakening as a cause of the preferential mylonitization of the Mulga Park Subdomain (Camacho and McDougall, 2000).

Pseudotachylytes have been identified as preferred nucleation discontinuities for shearing under mid- to lower crustal conditions (Andersen and Austrheim, 2006; Austrheim and Andersen, 2004; Hawemann et al., 2014; Lund and Austrheim, 2003; Menegon et al., 2017; Passchier, 1982; Pennacchioni and Cesare, 1997; Pittarello et al., 2012; Wex et al., 2014, 2017). In the Musgrave Block however, the presence of precursor pseudotachylyte cannot account for the observed inverse distribution of ductile deformation, as they are significantly more abundant in the hanging wall compared to the footwall (Camacho et al., 1995; Lin et al., 2005).

A potential explanation for the observed inverse gradient of ductile deformation along the Woodroffe Thrust is a rheological
contrast between hanging wall and footwall assemblages, resulting in different degrees, and potentially different processes, of
water-assisted weakening in quartz and feldspar (e.g., Fitz Gerald et al., 2006; Griggs, 1967, 1974; Griggs and Blacic, 1965;
Hobbs, 1985; Kronenberg et al., 1990; Mancktelow and Pennacchioni, 2004; Stünitz et al., 2017; Tullis and Yund, 1989). As
discussed in Sect. 8.3.4, the rocks in the study area represent a relatively closed system with respect to the presence or absence
of aqueous fluids, although some CO₂-introduction from an external source is necessary to account for the syntectonic growth
of very minor amounts of calcite in previously non-carbonaceous rocks. The availability of aqueous fluids is thus directly
linked to the abundance of hydrous minerals in these rocks, which is observed to decrease from north to south (Fig. 9).
Similarly, the contrasting abundance of hydrous minerals in hanging wall and footwall (Table 4) indicates that there is

- consistently a higher potential for generating aqueous fluids in the footwall than in the hanging wall. If sufficient water was available, this could facilitate a preferential rheological weakening of the footwall due to new growth of micas and associated
- 455 phyllonite development. However, this is not observed because the majority of the studied rocks are still relatively "dry", so that only slight differences in the abundance of water could facilitate an asymmetric distribution of ductile deformation as a consequence of water-assisted weakening. The preferential mylonitization of the Mulga Park relative to the Fregon Subdomain (Fig. 6) therefore seems to be largely controlled by the precursor mineralogy established as the result of the earlier (ca. 1200)

Ma) Musgravian Orogeny metamorphism, with peak conditions of upper amphibolite facies in the footwall but granulite facies
 in the hanging wall. This supports the initial hypothesis of Bell and Etheridge (1976) and Camacho et al. (1995) that a more
 hydrous footwall underlying an anhydrous hanging wall can facilitate an inverted gradient of ductile deformation.

These arguments may explain the overall preferential distribution of deformation in the Mulga Park Subdomain, but fail to provide an adequate explanation for why the lowermost Fregon Subdomain was also incorporated into the mylonites (Figs. 2, 6, 9). This reworking of the hanging wall may potentially have been guided by the presence of pseudotachylytes, which are ubiquitous and voluminous in the immediate hanging wall of the Woodroffe Thrust (Camacho et al., 1995; Lin et al., 2005).

465

8.4.2 Variation in mylonite thickness

Mylonite thickness in the study area does not significantly vary parallel to strike (i.e. E-W). However, from south to north, i.e. parallel to the direction of tectonic transport, the Woodroffe Thrust mylonitic zone gradually decreases in thickness from over 600 m to less than 100 m (Table 1). This potentially indicates a 6-fold increase in shear strain (and therefore average strain 470 rate) within the mylonites, assuming a homogeneous distribution of strain and constant relative displacement across the Woodroffe Thrust thickness along the entire ca. 60 km transect. This trend from south to north is accompanied by a slight decrease in metamorphic temperature of ca. 100 °C (Wex et al., 2017) and the shift from "dry" to "wet" conditions, as reflected in the increasing abundance of hydrous minerals (Fig. 9). The thickness of large-scale shear zones is considered to decrease with decreasing temperature and depth (Platt and Behr, 2011a, 2011b) and decreasing displacement (Hull, 1988). A point in 475 the footwall only enters the shear zone when it passes the toe of the thrust (that is where the thrust meets the surface). This could potentially lead to a variation in the finite shear strain experienced in the footwall. However, the whole of the exposed ca. 60 km N-S section was formerly at mid- to lower crustal level (Wex et al., 2017) and thus nowhere near the thrust toe. There is also currently no evidence for major splays of the Woodroffe Thrust into the hanging wall. Distributed folding related to the Peterman Orogeny is observable on the regional scale in the footwall (Wex et al., 2017) but the amount of shortening 480 involved cannot be quantified. It follows that there is presently no indication for a significant change in relative displacement along the studied section and that variation in this parameter therefore cannot be used to account for the difference in mylonite thickness. It was argued above that the greater abundance of hydrous minerals in the footwall Mulga Park Subdomain compared to the hanging wall Fregon Subdomain, reflecting peak metamorphic conditions of upper amphibolite facies and granulite facies respectively during the earlier Musgravian Orogeny, resulted in slightly "wetter" conditions in the footwall during 485 activity of the Woodroffe Thrust. This was proposed as an explanation for the broader zone of mylonites in the footwall. However, it ereates the apparent contradiction the current study also establishes that the thickness of the mylonitic zone decreases with increasing water activity toward to the north and that this is associated with increasing water activity and decreasing temperature. Such an observation might reflect the possibility that even a small amount of fluid was able to weaken the footwall relative to the strong "drier" hanging wall, allowing it to preferentially take up deformation because of its lower 490 effective viscosity. An increase in the Increasing water content in the footwall to the north would could result in an increase of

27

in the effective viscosity rheological contrastratio between footwall and hanging wall, yielding a stronger localization toward the interface and a narrower mylonite zone that extends less into the stronger material. This hypothesis is supported by the current observations in this paper, but there is not yet a theoretical basis to provide a full explanation. Decreasing temperature to the north could also lead to increased localization and decreased shear zone thickness, both directly (Platt and Behr, 2011a, 2011b) and as a second order effect that accentuatesing the strength rheological contrast as both footwall and hanging wall become stronger with decreasing temperature. Water activity thus seems to be the crucial factor controlling both the preferential development of mylonites in the footwall and the decrease in thickness with increasing water activity towards the north.

9 Conclusions

495

- 500 Field and thin section observations establish that the rocks of the central Musgrave Block were predominantly "dry" during development of the midcrustal Woodroffe Thrust during the ca. 560-520 Ma Petermann Orogeny, but with a progression in the thrust direction towards locally "wet" conditions in some of the northernmost exposures. This is indicated by: (1) rare occurrence of syntectonic quartz veins and quartz-rich pegmatites (locally found only in the north), (2) metastability of plagioclase in the presence of K-feldspar, which rarely shows significant alteration via the reaction An + Or + H₂O = Ms + Ep + Qz (more common towards the north) and (3) preferential high-pressure breakdown of plagioclase via the reaction An = Grs + Ky + Qz (common in the southerly exposures), rather than An + H₂O = Ep + Ky + Qz. Aqueous fluids were most likely derived internally from hydrous minerals within the footwall gneisses and granitoids, implying that the rocks in the study area were a relatively closed system.
- The thickness of the Woodroffe Thrust mylonites generally increases decreases with increasing decreasing metamorphic grade and decreasing increasing availability of aqueous fluids. Atypical of a thrust, ductile deformation is more extensively developed in the footwall rocks and only marginally involved several tens of metres of the lowermost hanging wall. The inverse gradient of ductile deformation cannot be explained by thermal weakening or the distributed presence of pseudotachylyte (acting as preferred nucleation sites for shearing), but rather by preferential rheological weakening of the "wetter" footwall compared to the "drier" hanging wall. This reflects the earlier (Musgravian Orogeny) peak metamorphic 515 conditions (granulite facies in the hanging wall and upper amphibolite facies in the footwall) and the contrasting availability of aqueous fluids derived from relict hydrous minerals in the footwall and hanging wall.

Data availability

Supplementary data are available in the Supporting Information "S1" to "S4" and further information can be obtained on request from the corresponding author.

520 Competing interests

The authors declare that they have no conflict of interest.

Author contribution

Each of the listed authors took part in at least two of three field seasons, which formed the basis of this study. Alfredo Camacho's previous knowledge of the Musgrave Ranges and the local communities was essential for the success of the campaign. Neil Mancktelow and Giorgio Pennacchioni developed the initial idea of the study. Sebastian Wex prepared the

525 campaign. Neil Mancktelow and Giorgio Pennacchioni developed the initial idea of the study. Sebastian Wex j manuscript with contributions from all co-authors.

Acknowledgments

We thank the communities of the Anangu Pitjantjatjara Yankunytjatjara Lands (APY) for granting us access to the Musgrave Ranges. Logistical support from the Northern Territory Geological Survey (NTGS) of Australia, Basil Tikoff (Univ.

- 530 Wisconsin, Madison) and Shane and Alethea Nicolle are gratefully recognized. We further acknowledge the support of Karsten Kunze from the Scientific Center for Optical and Electron Microscopy (ScopeM) at the ETH, Zurich. Jost Eikenberg is thanked for supervising and conducting the thorium measurements at the Paul Scherrer Institute (PSI) in Villigen, Switzerland. We further acknowledge the support of Madalina Jaggi who carried out the stable isotope analyses at the Geological Institute at ETH, Zurich. This project was financed by the Swiss National Science Foundation (SNF) Grant 200021_146745 awarded to
- 535 Neil Mancktelow, with additional funding from the University of Padova (BIRD175145/17: The geological record of deep earthquakes: the association pseudotachylyte-mylonite) awarded to Giorgio Pennacchioni.

References

Allen, J. L. and Shaw, C. A.: Seismogenic structure of a crystalline thrust fault: fabric anisotropy and coeval pseudotachylytemylonitic pseudotachylyte in the Grizzly Creek Shear Zone, Colorado, in Geology of the Earthquake Source: A Volume in

540 Honour of Rick Sibson, edited by A. Fagereng, V. G. Toy, and J. V. Rowland, pp. 135–151, Geol. Soc. Spec. Publ., 359, London., 2011.

Andersen, T. B. and Austrheim, H.: Fossil earthquakes recorded by pseudotachylytes in mantle peridotite from the Alpine subduction complex of Corsica, Earth Planet. Sci. Lett., 242(1–2), 58–72, doi:10.1016/j.epsl.2005.11.058, 2006. Austrheim, H. and Andersen, T. B.: Pseudotachylytes from Corsica: fossil earthquakes from a subduction complex, Terra

- 545 Nov., 16(4), 193–197, doi:10.1111/j.1365-3121.2004.00551.x, 2004.
 Ballhaus, C. and Glikson, A. Y.: The petrology of layered mafic-ultramafic intrusions of the Giles complex, western Musgrave
 - Balinaus, C. and Glikson, A. Y.: The petrology of layered maric-ultramatic intrusions of the Glies complex, western Musgrave Block, Western Australia, AGSO J. Aust. Geol. Geophys., 16(1/2), 69–90, 1995.

Bell, T. H.: Progressive deformation and reorientation of fold axes in a ductile mylonite zone: The Woodroffe Thrust, Tectonophysics, 44(1–4), 285–320, doi:10.1016/0040-1951(78)90074-4, 1978.

- Bell, T. H. and Etheridge, M. A.: The deformation and recrystallization of quartz in a mylonite zone, central Australia, Tectonophysics, 32(3–4), 235–267, doi:10.1016/0040-1951(76)90064-0, 1976.
 Bell, T. H. and Johnson, S. E.: The role of deformation partitioning in the deformation and recrystallization of plagioclase and K-feldspar in the Woodroffe Thrust mylonite zone, central Australia, J. Metamorph. Geol., 7(2), 151–168, doi:10.1111/j.1525-1314.1989.tb00582.x, 1989.
- Bell, T. H. and Johnson, S. E.: Shear sense: a new approach that resolves conflicts between criteria in metamorphic rocks, J. Metamorph. Geol., 10(1), 99–124, doi:10.1111/j.1525-1314.1992.tb00074.x, 1992.
 Berglund, M. and Wieser, M. E.: Isotopic compositions of the elements 2009 (IUPAC Technical Report), Pure Appl. Chem., 83(2), 397–410, doi:10.1351/PAC-REP-10-06-02, 2011.
 Boyd, F. R. and England, J. L.: Melting of silicates at high pressures, in Carnegie Inst. Wash. Yr. Book 60, pp. 113–125, Port
- 560 City Press, Baltimore., 1961. Breitenbach, S. F. M. and Bernasconi, S. M.: Carbon and oxygen isotope analysis of small carbonate samples (20 to 100 μg) with a GasBench II preparation device, Rapid Commun. Mass Spectrom., 25(13), 1910–1914, doi:10.1002/rcm.5052, 2011. Camacho, A.: An Isotopic Study of Deep-Crustal Orogenic Processes: Musgrave Block, Central Australia, PhD Thesis. The Australian National University, Canberra., 1997.
- 565 Camacho, A. and Fanning, C. M.: Some isotopic constraints on the evolution of the granulite and upper amphibolite facies terranes in the eastern Musgrave Block, central Australia, Precambrian Res., 71(1–4), 155–181, doi:10.1016/0301-9268(94)00060-5, 1995.

Camacho, A. and McDougall, I.: Intracratonic, strike-slip partitioned transpression and the formation and exhumation of eclogite facies rocks: An example from the Musgrave Block, central Australia, Tectonics, 19(5), 978–996,

570 doi:10.1029/1999TC001151, 2000.

Camacho, A., Simons, B. and Schmidt, P. W.: Geological and palaeomagnetic significance of the Kulgera Dyke Swarm, Musgrave Block, NT, Australia, Geophys. J. Int., 107(1), 37–45, doi:10.1111/j.1365-246X.1991.tb01154.x, 1991. Camacho, A., Vernon, R. H. and Fitz Gerald, J. D.: Large volumes of anhydrous pseudotachylyte in the Woodroffe Thrust,

eastern Musgrave Ranges, Australia, J. Struct. Geol., 17(3), 371–383, doi:10.1016/0191-8141(94)00069-C, 1995.
575 Camacho, A., Compston, W., McCulloch, M. and McDougall, I.: Timing and exhumation of eclogite facies shear zones, Musgrave Block, central Australia, J. Metamorph. Geol., 15(6), 735–751, doi:10.1111/j.1525-1314.1997.00053.x, 1997.

Camacho, A., Yang, P. and Frederiksen, A.: Constraints from diffusion profiles on the duration of high-strain deformation in thickened crust, Geology, 37(8), 755–758, doi:10.1130/G25753A.1, 2009.

Camacho, A., Armstrong, R., Davis, D. W. and Bekker, A.: Early history of the Amadeus Basin: Implications for the existence
and geometry of the Centralian Superbasin, Precambrian Res., 259, 232–242, doi:10.1016/j.precamres.2014.12.004, 2015.

Chacko, T., Mayeda, T. K., Clayton, R. N. and Goldsmith, J. R.: Oxygen and carbon isotope fractionations between CO2 and

calcite, Geochim. Cosmochim. Acta, 55(10), 2867–2882, doi:10.1016/0016-7037(91)90452-B, 1991.

Christie, J. M.: Moine Thrust Zone in the Assynt region of northwest Scotland, Univ. Calif. Publ. Geol. Sci., 40(6), 345–440, 1963.

585 Clarke, G. L., Buick, I. S., Glikson, A. Y. and Stewart, A. J.: Structural and pressure-temperature evolution of host rocks of the Giles Complex, western Musgrave Block, central Australia: evidence for multiple high-pressure events, AGSO J. Aust. Geol. Geophys., 16(1/2), 127–146, 1995.

Collerson, K. D., Oliver, R. L. and Rutland, R. W. R.: An example of structural and metamorphic relationships in the Musgrave orogenic belt, central Australia, J. Geol. Soc. Aust., 18(4), 379–393, doi:10.1080/00167617208728776, 1972.

590 Cooper, A. F. and Norris, R. J.: Anatomy, structural evolution, and slip rate of a plate boundary thrust: The Alpine Fault at Gaunt Creek, Westland, New Zealand, Geol. Soc. Am. Bull., 106(5), 627–633, doi:10.1130/0016-7606(1994)106<0627:ASEASR>2.3.CO;2, 1994.

Coplen, T. B., Böhlke, J. K., De Bièvre, P., Ding, T., Holden, N. E., Hopple, J. A., Krouse, H. R., Lamberty, A., Peiser, H. S., Révész, K., Rieder, S. E., Rosman, K. J. R., Roth, E., Taylor, P. D. P., Vocke, R. D. and Xiao, Y. K.: Isotope-abundance

595 variations of selected elements (IUPAC Technical Report), Pure Appl. Chem., 74(10), 1987–2017, doi:10.1351/pac200274101987, 2002.
Coward, M. P.: The Caledonian thrust and shear zones of N.W. Scotland, J. Struct. Geol., 2(1–2), 11–17, doi:10.1016/0191-

8141(80)90029-2, 1980.

Davis, G. A.: Rapid upward transport of mid-crustal mylonitic gneisses in the footwall of a Miocene detachment fault, Whipple

- Mountains, southeastern California, Geol. Rundschau, 77(1), 191–209, doi:10.1007/BF01848684, 1988.
 Davis, G. A. and Lister, G. S.: Detachment faulting in continental extension; Perspectives from the Southwestern U.S. Cordillera, Geol. Soc. Am. Spec. Pap., 218, 133–159, doi:10.1130/SPE218-p133, 1988.
 Deines, P.: The carbon isotope geochemistry of mantle xenoliths, Earth-Science Rev., 58(3–4), 247–278, doi:10.1016/S0012-8252(02)00064-8, 2002.
- Edgoose, C. J., Camacho, A., Wakelin-King, G. A. and Simons, B. A.: 1:250 000 Geological Map Series Explanatory Notes.
 Kulgera SG 53-5, 2nd ed., North. Territ. Geol. Surv., Darwin., 1993.
 Edgoose, C. J., Scrimgeour, I. R. and Close, D. F.: Report 15: Geology of the Musgrave Block, Northern Territory, North.
 Territ. Geol. Surv., Darwin., 2004.

Eikenberg, J.: Radium Isotope Systematics in Nature: Applications in Geochronology and Hydrogeochemistry, Habilitation

- Thesis. Devision for Radiation Safety and Security, Paul Scherrer Institute (PSI), Villigen, Switzerland and Earth Science Department, Swiss Federal Institute of Technology (ETH), Zurich., 2002.
 Ellis, D. J. and Maboko, M. A. H.: Precambrian tectonics and the physicochemical evolution of the continental crust. I. The gabbro-eclogite transition revisited, Precambrian Res., 55(1–4), 491–506, doi:10.1016/0301-9268(92)90041-L, 1992.
 Evins, P. M., Smithies, R. H., Howard, H. M., Kirkland, C. L., Wingate, M. T. D. and Bodorkos, S.: Record 2010/6: Redefining
- 615 the Giles Event within the setting of the 1120-1020 Ma Ngaanyatjarra Rift, west Musgrave Province, central Australia, Geol.

Surv. West. Aust., Perth., 2010.

645

Fitz Gerald, J. D., Mancktelow, N. S., Pennacchioni, G. and Kunze, K.: Ultra-fine grained quartz mylonites from high-grade shear zones: Evidence for strong dry middle to lower crust, Geology, 34(5), 369–372, doi:10.1130/G22099.1, 2006.

Flottmann, T., Hand, M., Close, D., Edgoose, C. and Scrimgeour, I.: Thrust Tectonic Styles of the Intracratonic Alice Springs
and Petermann Orogenies, Central Australia, in AAPG Memoir 82: Thrust tectonics and hydrocarbon systems, edited by K.
R. McClay, pp. 538–557, Tulsa., 2004.

Forman, D. J.: 1:250,000 Geological Series Explanatory Notes. Ayers Rock, N.T. SG/52-8, 1st ed., Bur. Miner. Resour. Geol. Geophys., Canberra., 1965.

Fossen, H. and Cavalcante, G. C. G.: Shear zones – A review, Earth-Science Rev., 171, 434–455, doi:10.1016/j.earscirev.2017.05.002, 2017.

Goldsmith, J. R.: Plagioclase stability at elevated temperatures and water pressures., Am. Mineral., 67(7–8), 653–675, 1982. Gray, C. M.: The Geochemistry of Central Australian Granulites in Relation to the Chemical and Isotopic Effects of Granulite

Facies Metamorphism, Contrib. to Mineral. Petrol., 65(1), 79–89, doi:10.1007/BF00373573, 1977.
Gray, C. M.: Geochronology of granulite-facies gneisses in the western Musgrave Block, central Australia, J. Geol. Soc. Aust., 25(7), 403–414, doi:10.1080/00167617808729050, 1978.

Gray, C. M. and Compston, W.: A rubidium-strontium chronology of the metamorphism and prehistory of central Australian granulites, Geochim. Cosmochim. Acta, 42(11), 1735–1747, doi:10.1016/0016-7037(78)90259-4, 1978.

635 Griggs, D.: Hydrolytic Weakening of Quartz and Other Silicates, Geophys. J. R. Astron. Soc., 14(1–4), 19–31, doi:10.1111/j.1365-246X.1967.tb06218.x, 1967.

Griggs, D.: A Model of Hydrolytic Weakening in Quartz, J. Geophys. Res., 79(11), 1653–1661, doi:10.1029/JB079i011p01653, 1974.

Griggs, D. T. and Blacic, J. D.: Quartz: Anomalous Weakness of Synthetic Crystals, Science, 147(3655), 292–295, doi:10.1126/science.147.3655.292, 1965.

Handy, M. R., Hirth, G. and Bürgmann, R.: Continental fault structure and rheology from the frictional-to-viscous transition downward, in Tectonic Faults: Agents of Change on a Dynamic Earth, edited by M. R. Handy, G. Hirth, and N. Hovius, pp. 139–181, MIT Press, Cambridge., 2007.

Hariya, Y. and Kennedy, G. C.: Equilibrium study of anorthite under high pressure and high temperature, Am. J. Sci., 266(3), 193–203, doi:10.2475/ajs.266.3.193, 1968.

Hawemann, F., Mancktelow, N., Wex, S., Camacho, A. and Pennacchioni, G.: Strain localization on different scales and the importance of brittle precursors during deformation in the lower crust (Davenport Shear Zone, Central Australia), Geophys. Res. Abstr., 16, 5009, 2014.

Hawemann, F., Mancktelow, N. S., Wex, S., Camacho, A. and Pennacchioni, G.: Pseudotachylyte as field evidence for lower-

Goldsmith, J. R.: The melting and breakdown reactions of anorthite at high pressures and temperatures, Am. Mineral., 65(3–4), 272–284, 1980.

crustal earthquakes during the intracontinental Petermann Orogeny, Solid Earth, 9, 629–648, doi:10.5194/se-9-629-2018, 2018.

Heier, K. S. and Adams, J. A. S.: Concentration of radioactive elements in deep crustal material, Geochim. Cosmochim. Acta, 29(1), 53–61, doi:doi.org/10.1016/0016-7037(65)90078-5, 1965.

Hill, A. C., Arouri, K., Gorjan, P. and Walter, M. R.: Geochemistry of marine and nonmarine environments of a Neoproterozoic

655 cratonic carbonate/evaporite: the Bitter Springs Formation, central Australia, in Carbonate Sedimentation and Diagenesis in the Evolving Precambrian World, edited by J. P. Grotzinger and N. P. James, pp. 327–344, SEPM (Soc. Sediment. Geol.) Spec. Publ., 67, Tulsa., 2000.

Hobbs, B. E.: The hydrolytic weakening effect in quartz, in Point Defects in Minerals, edited by R. N. Schock, pp. 151–170, Am. Geophys. Union, Geophys. Monogr. Ser., 31, Washington D.C., 1985.

660 Hull, J.: Thickness-displacement relationships for deformation zones, J. Struct. Geol., 10(4), 431–435, doi:10.1016/0191-8141(88)90020-X, 1988.

Javoy, M., Pineau, F. and Delorme, H.: Carbon and nitrogen isotopes in the mantle, Chem. Geol., 57(1–2), 41–62, doi:10.1016/0009-2541(86)90093-8, 1986.

Kretz, R.: Note on some equilibria in which plagioclase and epidote participate, Am. J. Sci., 261(10), 973–982, doi:10.2475/ajs.261.10.973, 1963.

665

Kronenberg, A. K., Segall, P. and Wolf, G. H.: Hydrolytic Weakening and Penetrative Deformation Within a Natural Shear
Zone, in The Brittle-Ductile Transition in Rocks, edited by A. G. Duba, W. B. Durham, J. W. Handin, and H. F. Wang, pp. 21–36, Am. Geophys. Union, Geophys. Monogr. Ser., 56, Washington D.C., 1990.

Lambert, I. B. and Heier, K. S.: The vertical distribution of uranium, thorium and potassium in the Continental Crust, Geochim. 670 Cosmochim. Acta, 31(3), 377–390, doi:10.1016/0016-7037(67)90048-8, 1967.

Lambert, I. B. and Heier, K. S.: Geochemical investigations of deep-seated rocks in the Australian shield, Lithos, 1(1), 30–53, doi:10.1016/S0024-4937(68)80033-7, 1968.

Lin, A., Maruyama, T., Aaron, S., Michibayashi, K., Camacho, A. and Kano, K.: Propagation of seismic slip from brittle to ductile crust: Evidence from pseudotachylyte of the Woodroffe thrust, central Australia, Tectonophysics, 402(1–4), 21–35,

doi:10.1016/j.tecto.2004.10.016, 2005. Lund, M. G. and Austrheim, H.: High-pressure metamorphism and deep-crustal seismicity: evidence from contemporaneous formation of pseudotachylytes and eclogite facies coronas, Tectonophysics, 372(1–2), 59–83, doi:10.1016/S0040-1951(03)00232-4, 2003.

Maboko, M. A. H., McDougall, I. and Zeitler, P. K.: Metamorphic P-T path of granulites in the Musgrave Ranges, central

680 Australia, in Evolution of Metamorphic Belts, edited by J. S. Daly, R. A. Cliff, and B. W. D. Yardley, pp. 303–307, Geol. Soc. Spec. Publ., 43, London., 1989.

Maboko, M. A. H., Williams, I. S. and Compston, W.: Zircon U-Pb Chronometry of the Pressure and Temperature History of Granulites in the Musgrave Ranges, Central Australia, J. Geol., 99(5), 675–697, 1991.

Maboko, M. A. H., McDougall, I., Zeitler, P. K. and Williams, I. S.: Geochronological evidence for ~ 530-550 Ma juxtaposition of two Proterozoic metamorphic terranes in the Musgrave Ranges, central Australia, Aust. J. Earth Sci., 39(4), 457–471,

doi:10.1080/08120099208728038, 1992. Major, R. B.: Woodroffe Thrust Zone in the Musgrave Ranges, Q. Geol. Notes, 35, 9–11, 1970. Major, R. B.: Explanatory Notes for the Woodroffe 1:250 000 Geological Map SG/52-12, 1st ed., Geol. Surv. South Aust., Adelaide., 1973.

685

- Major, R. B. and Conor, C. H. H.: Musgrave Block, in Bulletin 54: The geology of South Australia. Volume 1. The Precambrian, edited by J. F. Drexel, W. V. Preiss, and A. J. Parker, pp. 156–167, Geol. Surv. South Aust., Adelaide., 1993. Major, R. B., Johnson, J. E., Leeson, B., Mirams, R. C. and Thomson, B. P.: 1:250 000 S. A. Geological Atlas Series Sheet. Woodroffe SG 52-12 Zone 4., 1st ed., Geol. Surv. South Aust., Adelaide., 1967. Mancktelow, N.: The Simplon Line: a major displacement zone in the western Lepontine Alps, Eclogae Geol. Helv., 78(1),
- 695 73–96, doi:10.5169/seals-165644, 1985.
 Mancktelow, N. S. and Pennacchioni, G.: The influence of grain boundary fluids on the microstructure of quartz-feldspar mylonites, J. Struct. Geol., 26(1), 47–69, doi:10.1016/S0191-8141(03)00081-6, 2004.
 Mancktelow, N. S. and Pennacchioni, G.: The control of precursor brittle fracture and fluid-rock interaction on the development of single and paired ductile shear zones, J. Struct. Geol., 27(4), 645–661, doi:10.1016/j.jsg.2004.12.001, 2005.
- Menegon, L., Pennacchioni, G., Malaspina, N., Harris, K. and Wood, E.: Earthquakes as Precursors of Ductile Shear Zones in the Dry and Strong Lower Crust, Geochemistry, Geophys. Geosystems, 18, 4356–4374, doi:10.1002/2017GC007189, 2017. Milke, R., Neusser, G., Kolzer, K. and Wunder, B.: Very little water is necessary to make a dry solid silicate system wet, Geology, 41(2), 247–250, doi:10.1130/G33674.1, 2013.

Myers, J. S., Shaw, R. D. and Tyler, I. M.: Tectonic evolution of Proterozoic Australia, Tectonics, 15(6), 1431–1446, doi:10.1029/96TC02356, 1996.

Passchier, C. W.: Pseudotachylyte and the development of ultramylonite bands in the Saint-Barthélemy Massif, French Pyrenees, J. Struct. Geol., 4(1), 69–79, doi:10.1016/0191-8141(82)90008-6, 1982.

Passchier, C. W.: Mylonite-dominated footwall geometry in a shear zone, central Pyrenees, Geol. Mag., 121(5), 429–436, doi:10.1017/S0016756800029964, 1984.

Passchier, C. W. and Trouw, R. A. J.: Microtectonics, 2nd ed., Springer, Berlin, Heidelberg., 2005.
 Pennacchioni, G. and Cesare, B.: Ductile-brittle transition in pre-Alpine amphibolite facies mylonites during evolution from water-present to water-deficient conditions (Mont Mary nappe, Italian Western Alps), J. Metamorph. Geol., 15(6), 777–791, doi:10.1111/j.1525-1314.1997.00055.x, 1997.

Percival, P. J.: Record 2010/13: Index of Airborne Geophysical Surveys, 11th ed., Geoscience Australia, Canberra., 2010.

Pittarello, L., Pennacchioni, G. and Di Toro, G.: Amphibolite-facies pseudotachylytes in Premosello metagabbro and felsic mylonites (Ivrea Zone, Italy), Tectonophysics, 580, 43–57, doi:10.1016/j.tecto.2012.08.001, 2012.
Platt, J. P. and Behr, W. M.: Deep structure of lithospheric fault zones, Geophys. Res. Lett., 38(24), 1–6,

doi:10.1029/2011GL049719, 2011a.

Platt, J. P. and Behr, W. M.: Lithospheric shear zones as constant stress experiments, Geology, 39(2), 127-130,

720 doi:10.1130/G31561.1, 2011b.

Ramberg, H.: The Facies Classification of Rocks: A Clue to the Origin of Quartzo-Feldspathic Massifs and Veins, J. Geol., 57(1), 18–54, doi:10.1086/625573, 1949.

Ramsay, J. G.: Shear zone geometry: A review, J. Struct. Geol., 2(1–2), 83–99, doi:10.1016/0191-8141(80)90038-3, 1980. Schmidt, P. W., Williams, G. E., Camacho, A. and Lee, J. K. W.: Assembly of Proterozoic Australia: Implications of a revised

725 pole for the ~1070 Ma Alcurra Dyke Swarm, central Australia, Geophys. J. Int., 167(2), 626–634, doi:10.1111/j.1365-246X.2006.03192.x, 2006.

Scrimgeour, I. and Close, D.: Regional high-pressure metamorphism during intracratonic deformation: The Petermann Orogeny, central Australia, J. Metamorph. Geol., 17(5), 557–572, doi:10.1046/j.1525-1314.1999.00217.x, 1999.

Scrimgeour, I. R., Close, D. F. and Edgoose, C. J.: 1:250,000 Geological Map Series and Explanatary Notes. Petermann Ranges
SG52-7, 2nd ed., North. Territ. Geol. Surv., Darwin., 1999.

Selverstone, J., Axen, G. J. and Luther, A.: Fault localization controlled by fluid infiltration into mylonites: Formation and strength of low-angle normal faults in the midcrustal brittle-plastic transition, J. Geophys. Res., 117, B06210, doi:10.1029/2012JB009171, 2012.

Sibson, R. H., White, S. H. and Atkinson, B. K.: Structure and distribution of fault rocks in the Alpine Fault Zone, New

735 Zealand, in Thrust and Nappe Tectonics, edited by K. R. McClay and N. J. Price, pp. 197–210, Geol. Soc. Spec. Publ., 9, London., 1981.

Smithies, R. H., Howard, H. M., Evins, P. M., Kirkland, C. L., Kelsey, D. E., Hand, M., Wingate, M. T. D., Collins, A. S. and Belousova, E.: High-Temperature Granite Magmatism, Crust-Mantle Interaction and the Mesoproterozoic Intracontinental Evolution of the Musgrave Province, Central Australia, J. Petrol., 52(5), 931–958, doi:10.1093/petrology/egr010, 2011.

Sprigg, R. C., Wilson, B., Coats, R. P., Webb, B. P. and O'Driscoll, E. S.: 4 Mile Geological Series Sheet. Alberga G 53-9
 Zone 4, 1st ed., Geol. Surv. South Aust., Adelaide., 1959.
 Stewart, A. J.: Western extension of the Woodroffe Thrust, Musgrave Block, central Australia, AGSO J. Aust. Geol. Geophys.,

16(1/2), 147–153, 1995.

Stewart, A. J.: Record 1997/5: Geology of the Bates 1:100 000 Sheet Area (4646), Musgrave Block, Western Australia, 1st 745 ed., Aust. Geol. Surv. Organ., Canberra., 1997.

Stünitz, H., Thust, A., Heilbronner, R., Behrens, H., Kilian, R., Tarantola, A. and Fitz Gerald, J. D.: Water redistribution in experimentally deformed natural milky quartz single crystals - Implications for H2O-weakening processes, J. Geophys. Res. Solid Earth, 122(2), 866–894, doi:10.1002/2016JB013533, 2017.

Sun, S., Sheraton, J. W., Glikson, A. Y. and Stewart, A. J.: A major magmatic event during 1050-1080 Ma in central Australia,

<sup>Sun, S. and Sheraton, J.: Zircon U/Pb chronology, tectono-thermal and crust-forming events in the Tomkinson Ranges,
Musgrave Block, Central Australia, AGSO Res. Newsl., 17, 9–11, 1992.</sup>

and an emplacement age for the Giles Complex, AGSO Res. Newsl., 24, 13-15, 1996.

Taylor, H. P.: Oxygen and Hydrogen Isotope Relationships in Hydrothermal Mineral Deposits, in Geochemistry of Hydrothermal Ore Deposits, edited by H. L. Barnes, pp. 229–302, John Wiley and Sons, New York., 1997.

Tucker, N. M. T., Hand, M., Kelsey, D. E. and Dutch, R. A.: A duality of timescales: Short-lived ultrahigh temperature metamorphism preserving a long-lived monazite growth history in the Grenvillian Musgrave–Albany–Fraser Orogen, Precambrian Res., 264, 204–234, doi:10.1016/j.precamres.2015.04.015, 2015.
 Tullis, J. and Yund, R. A.: Hydrolytic weakening of quartz aggregates: The effects of water and pressure on recovery, Geophys.

Res. Lett., 16(11), 1343–1346, doi:10.1029/GL016i011p01343, 1989.

775

760 Walsh, A. K., Kelsey, D. ., Kirklan, C. L., Hand, M., Smithies, R. H., Clark, C. and Howard, H. M.: P–T–t evolution of a large, long-lived, ultrahigh-temperature Grenvillian belt in central Australia, Gondwana Res., 28(2), 531–564, doi:10.1016/j.gr.2014.05.012, 2015.

Wayte, G. J., Worden, R. H., Rubie, D. C. and Droop, G. T. R.: A TEM study of disequilibrium plagioclase breakdown at high pressure: the role of infiltrating fluid, Contrib. to Mineral. Petrol., 101(4), 426–437, doi:10.1007/BF00372216, 1989.

- Wells, A. T., Forman, D. J., Ranford, L. C. and Cook, P. J.: Bulletin 100: Geology of the Amadeus Basin, Central Australia, Bur. Miner. Resour. Geol. Geophys., Canberra., 1970.
 Wex, S., Mancktelow, N., Hawemann, F., Camacho, A. and Pennacchioni, G.: Pseudotachylyte formation vs. mylonitization repeated cycles of seismic fracture and aseismic creep in the middle crust (Woodroffe Thrust, Central Australia), Geophys. Res. Abstr., 16, 5071, 2014.
- Wex, S., Mancktelow, N. S., Hawemann, F., Camacho, A. and Pennacchioni, G.: Geometry of a large-scale, low-angle, midcrustal thrust (Woodroffe Thrust, central Australia), Tectonics, 36(11), 2447–2476, doi:10.1002/2017TC004681, 2017. Whitney, D. L. and Evans, B. W.: Abbreviations for names of rock-forming minerals, Am. Mineral., 95(1), 185–187, doi:10.2138/am.2010.3371, 2010.

Young, D. N., Duncan, N., Camacho, A., Ferenczi, P. A. and Madigan, T. L. A.: 1:250 000 Geological Map Series and Explanatory Notes. Ayers Rock SG 52-8, 2nd ed., North. Territ. Geol. Surv., Darwin., 2002.

Zhao, J. and McCulloch, M. T.: Sm-Nd mineral isochron ages of Late Proterozoic dyke swarms in Australia: evidence for two distinctive events of mafic magmatism and crustal extension, Chem. Geol., 109(1–4), 341–354, doi:10.1016/0009-2541(93)90079-X, 1993.

Zhao, J., McCulloch, M. T. and Korsch, R. J.: Characterisation of a plume-related ~ 800 Ma magmatic event and its

780 implications for basin formation in central-southern Australia, Earth Planet. Sci. Lett., 121(3–4), 349–367, doi:10.1016/0012-821X(94)90077-9, 1994.

S1 Methodology

5

10

S1.1 Scanning electron microscopy (SEM)

Backscattered electron (BSE, atomic Z-contrast) analysis was performed using an FEI Quanta 200F scanning electron microscope equipped with a field emission gun at the Scientific Center for Optical and Electron Microscopy (ScopeM) at ETH, Zurich. BSE settings were: acceleration voltage 10-20 kV, working distance 10 mm, aperture 3-4, spot size 4 and high vacuum. Thin sections were carbon-coated.

y-ray spectrometry (²³²Th) S1.2

Trace concentrations of thorium in felsic gneisses and granitoids have been measured using γ -ray spectrometry. This method, rather than laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS), was applied, because it does not require extensive sample preparation and represents a quick and efficient method of acquiring the desired element concentration.

Thorium-232 accounts for 100 % of the element's natural abundance (Berglund and Wieser, 2011). Therefore, a sample's thorium content was assessed by quantifying the activity of this particular radioactive thorium isotope. Thorium-232 was not measured directly because it is a pure α -emitter without photon emission (Fig. S1).

- 15 However, due to the Mesoproterozoic age (ca. 1200 Ma) of the deformation event (Musgravian Orogeny) which produced the contrasting thorium concentrations in the hanging wall and footwall of the Woodroffe Thrust, all products of the thorium-232 decay series (Fig. S1) can be assumed to be in equilibrium (secular equilibrium in the thorium series is obtained after only 30 years). As a consequence, the activity of thorium-232 and all daughter isotopes is the same. Thus, the thorium concentration can be indirectly measured via the γ -decay of the shorter-
- 20 lived radium-228. The five most important peaks related to the γ -decay of radium-228 are located at 77.1 keV, 238.6 keV, 338.4 keV, 583.2 keV, 911.1 keV in the energy spectrum (Fig. S2). Measurements were conducted on thin section chips (weights ranging between 14-42 g) using γ -ray spectrometers at the Paul Scherrer Institute (PSI) in Villigen, Switzerland with a default measurement time of two hours. If the aspired error (2 x standard deviation) of 10 % was achieved earlier (thorium-rich samples), the measurement was stopped, otherwise it was run
- 25 overnight for up to 15 hours (thorium-poor samples). Results were weight-averaged, with the highest significance being attributed to the strongest peak at 238.6 keV, in order to obtain the activity of radium-228 normalized to sample weight (Bq/kg). Following up on the assumption that the activity of thorium-232 and radium-228 are the same, this value was divided by the specific activity of thorium-232 (4060 Bq/g; Eikenberg, 2002) in order to obtain the concentration of thorium-232. The detection limit was determined from the 238.6 keV peak of lead-
- 30 212. All measurements were done assuming a three centimetre plane source as sample geometry. Due to the dependence of the detector efficiency on the geometry and especially the thickness of each thin section chip, a correction factor was applied. This correction factor is equal to the relative difference between the detector efficiency of each sample with respect to the three centimetre plane source. All thin section chips were grouped according to their geometry and thickness and for each group a representative sample was chosen and the 35 correction factor determined using a Monte Carlo approach.



40 Figure S1: Schematic illustration of the thorium-232 decay series members. For simplicity of the scheme, in comparison to standard isotope tables, neutrons are not considered. From Eikenberg (2002).



Figure S2: γ-spectrum of radium-228 with the most relevant energy peaks highlighted. Sample SW13-323A (coordinates: 132.14344, -25.99211; location 6 in Fig. 1 of the main paper).

45 S1.3 Modal abundance of hydrous minerals

The modal abundance of hydrous minerals in thin section was quantified by running an image analysis script in Matlab R2013a (Fig. S3). Utilizing BSE images, thresholds were manually defined between the individual grey scale values ranges defined by the hydrous and anhydrous each minerals. In order to achieve a proper separation, we adjusted brightness and contrast in a manner such that the grey scale range defined by the hydrous minerals

50 covered a significant portion of the entire spectrum. As a result, the histogram data of the individual anhydrous minerals strongly overlap with one another (see peaks of quartz, K-feldspar and plagioclase in Fig. S3). This approach minimalizes the overlap between the peaks defined by the hydrous minerals with those of the anhydrous minerals and thus provides an optimal basis for the desired separation. The modal abundance of each phase was calculated after subtracting all Ppixels that were not part of the stable mineral assemblage (e.g. holes and clasts in pseudotachylytes) were subtracted. The magnification was kept roughly constant at an image width of between 60-80 µm. Depending on the grain size, a variable number of images (2-5) were used to assure that the derived value was representative of the pseudotachylyte matrix.



60

Figure S3: Calculation of the modal abundance of hydrous minerals from a BSE image. Thresholds (red lines) were set between the individual grey scale values ranges defined by each mineral. Thin section is not oriented. Sample FW13-093 (coordinates: 131.45213, -25.85455; location 3 in Fig. 1 of the main paper).

S1.4 Calcite isotope analysis

Carbon (δ¹³C_{Cal}) and oxygen (δ¹⁸O_{Cal}) isotope analysis of fine-grained, syntectonically crystallized calcite has been performed using a GasBench II preparation device connected to a Delta V mass spectrometer (both ThermoFisher Scientific, Bremen) at the Geological Institute at ETH, Zurich. The analytical procedures follow Breitenbach and Bernasconi (2011). Due to the low concentrations of calcite, slight modifications to these procedures include an increase in the amount of analysed sample material (9-13 mg) and utilized-amount of orthophosphoric acid (eight droplets)_a as well as the use of larger vials (12 ml). The data are reported in the conventional delta notation with respect to the V-PDB (Vienna Pee Dee Belemnite) and V-SMOW (Vienna Standard Mean Ocean Water) standard. For a reliable signal the benchmark Amplitude 44 (Ampl 44) should be above 2000.

S1.5 Whole rock carbon isotope analysis

Whole rock carbon isotopy (δ¹³C_{whole rock}) was determined using a ThermoFisher Flash-EA 1112 coupled with a Conflo IV interface to a ThermoFisher Delta V isotope ratio mass spectrometer (IRMS). Samples were combusted in the presence of oxygen in an oxidation column at 1020 °C. Combustion gases are passed through a reduction column (650 °C), and produced nitrogen and carbon dioxide gases separated chromatographically and transferred to the IRMS via an open split for on-line isotope measurements. Isotope ratios are reported in the conventional delta notation with respect to the V-PDB (Vienna Pee Dee Belemnite) standard. The system is calibrated with NBS22 (δ¹³C = -30.03 ‰) and IAEA CH-6 (δ¹³C = -10.46 ‰). Reproducibility of the measurements is better than 0.2 ‰. For a reliable signal the benchmark Amplitude 44 (Ampl 44) should be above 2000.

S2 Calcite and whole rock stable isotopic signature

- In an attempt to constrain the origin of the introduced carbon (Fig. 7c of the main paper), felsic and mafic samples, with and without fine-grained, syntectonically crystallized calcite, have been isotopically analysed (for details on the methods see Supporting Information S1), with the results for δ¹³C_{Cal}, δ¹⁸O_{Cal}, and δ¹³C_{whole rock} summarized in Table S1. With the exception of SW13-173A, calcite isotopic signatures were obtained for all calcite-bearing samples, yielding δ¹³C_{Cal} (V-PDB) values between -2.7 ‰ to -5.9 ‰ (mean average -4.1 ‰) and δ¹⁸O_{Cal} (SMOW)
 values between +5.7 ‰ and +14.5 ‰ (mean average +10.1 ‰). Within the same samples, the δ¹³C_{whole rock} (V-PDB)
- PDB) isotopic signature is always lower (on average by 2.7 ‰) than the corresponding δ¹³C_{cal} values. Calcite-free samples yield significantly lower δ¹³C_{whole rock} values between -10.7 ‰ and -22.6 ‰ (Table S1), with this carbon presumably coming from CO₂ contained in fluid inclusions. The amplitude (Ampl 44) of each measurement functions as a benchmark for the data's reliability. This parameter should typically be above 2000
 25 Second S18C and S18C
- for a measurement to be considered trustworthy. Consequently, $\delta^{13}C_{Cal}$ and $\delta^{18}O_{Cal}$ of SW13-159 and NW14-408 are well constrained, whereas the results for NW13-203, SW14-025 and SW13-124 are less reliable. The low benchmark values almost certainly reflect the very low modal abundance of calcite in the studied rocks (typically <1 %). Nevertheless, the determined values for each of the five samples are reproducible in different runs and are very similar to each other, which argues for their reliability.

100

					GasBench II							Combustion		
	location	sample	lithology	coordinates		d ¹³ C _{cal} (‰)		d ¹⁸ Oc	_{al} (‰)	_	d ¹³ C _{whole rock} (‰)		
	(Fig. 1)			(WGS 1984)	run	(V-PDB ±	±2σ)	(V-PI	B ± 2	5) (SMOW)	Ampl 44	(V-PDB)	Ampl 44	
	9 (HW)	NW13-203	granite	131,74318 -26,00442	1 2 3 4	$\begin{array}{rrrr} -4,12 & \pm \\ -4,12 & \pm \\ -4,47 & \pm \\ -4,08 & \pm \end{array}$	0,15 0,08 0,10 0,12	-21,19 -21,34 -23,30 -23,56	$\begin{array}{cccc} \pm & 0 \\ \pm & 0 \\ \pm & 0 \\ \pm & 0 \\ \pm & 0 \end{array}$	23 9,02 25 8,86 13 6,84 11 6,57	686 936 872 852	-9,36	4235	
ng	9 (HW)	SW14-025	granite	131,73269 -26,01569	1 2	-4,11 ± -3,92 ±	0,34 0,05	-24,37 -24,25	$ \begin{array}{cccc} \pm & 0, \\ \pm & 0, \end{array} $	25 5,74 23 5,86	432 667	-6,54	7500	
te-beari	10 (FW)	NW14-408	dolerite	131,85345 -26,11036	1 2	-5,83 ± -5,85 ±	0,05 0,10	-19,90 -19,90	$\begin{array}{ccc} \pm & 0 \\ \pm & 0 \end{array}$	15 10,34 08 10,35	2120 3463	-6,83	8069	
calci	12 (FW)	SW13-124	granite	131,87272 -26,20402	1 2	-3,72 ± -3,77 ±	0,14 0,12	-18,33 -18,16	$\begin{array}{ccc} \pm & 0 \\ \pm & 0 \end{array}$	29 11,97 04 12,14	625 1099	-6,50	6110	
	14 (FW)	SW13-159	felsic gneiss	131,77375 -26,30666	1 2	-2,76 ± -2,71 ±	0,07 0,07	-16,01 -15,88	$\begin{array}{c} \pm & 0 \\ \pm & 0 \end{array}$	16 14,36 04 14,49	1569 2305	-4,88	5743	
	2 (-)	SW13-173A	felsic gneiss	131,45334 -25,84543	1 2	-	-	-			5313 5312	-10,68	2353	
se	7 (FW)	SW13-104	granite	131,63509 -25,94122	1 2	-	-	-			7153 7156	-22,56	983	
lcite-fr	13 (FW)	SW14-194	dolerite	131,82577 -26,24931	1 2	-	-	-			5297 5292	-22,61	838	
са	14 (HW)	SW13-166	dolerite	131,77451 -26,30822	1 2	-	-	-		 	5304 5294	-17,29	960	

Table S1: Calcite ($\delta^{13}C_{Cal}$, $\delta^{18}O_{Cal}$) and whole rock ($\delta^{13}C_{whole rock}$) stable isotope results for the central Musgrave Block. Abbreviations: hanging wall (HW), footwall (FW).

- 105 $\delta^{13}C_{Cal}$ should provide information on the fluid source, since there are no other carbon-bearing phases in the studied rocks (Collerson et al., 1972; Major, 1973; Major and Conor, 1993; Scrimgeour and Close, 1999). $\delta^{18}O_{Cal}$, on the other hand, should reflect the calcite crystallization temperatures. Our results are in agreement with the range of $\delta^{18}O$ values for metamorphic waters (Taylor, 1997, his Fig. 6.4) and carbonates (Coplen et al., 2002, their Fig. 6) at temperatures of 500-600 °C, considering that the corresponding fluid phase should isotopically have
- 110 been ca. 5-6 ‰ heavier than the crystallizing calcite (Chacko et al., 1991). Our $\delta^{13}C_{Cal}$ results strongly overlap with the typical range of values for a rock-buffered system within igneous and metamorphic rocks (Coplen et al., 2002, their Fig. 4). However, from Table S1 it is evident that $\delta^{13}C_{Cal}$ is never identical to $\delta^{13}C_{whole rock}$. Consequently, full rock-buffering has not been achieved and it should be possible to place some constraints on the potential carbon source. The range of determined $\delta^{13}C_{Cal}$ values excludes a marine origin (Coplen et al., 2002,
- their Fig. 4), but is in agreement with a mantle (Deines, 2002; Javoy et al., 1986) or sedimentary source (Bitter Springs Formation; Hill et al., 2000). Within the underlying Amadeus Basin sediments, the most likely source would have been either the carbonate-bearing Bitter Springs Formation or its lateral equivalent the Pinyinna Beds (Wells et al., 1970; Young et al., 2002). However, both units do not crop out in the study area and, as outlined in Sect. 8.3.1 of the main paper, the Amadeus Basin sediments were potentially only imbricated below the Kelly
- 120 Hills klippe after shearing on the Woodroffe Thrust had largely ceased.

S3 Hydrous minerals in felsic pseudotachylytes



Figure S4: Representative BSE images of felsic pseudotachylytes from the hanging wall and footwall of the Woodroffe
Thrust arranged into northern, central and southern location groups (see Table 4 of the main paper). Overview thin section images are roughly 4 cm in width. (top left) Unoriented sample FW13-096 (coordinates: 131.45034, -25.85414; location 3 in Fig. 1 of the main paper). (top right) Unoriented sample FW13-093 (coordinates: 131.45213, -25.85455; location 3 in Fig. 1 of the main paper). (centre left) Unoriented sample SW14-029A (coordinates: 131.74496, -26.00093; location 9 in Fig. 1 of the main paper). (centre right) Sample SW13-321 (coordinates: 132.14333, -25.99178; location 6
in Fig. 1 of the main paper). Thin section is oriented N-S (left-right). (bottom left) Loose sample SW14-181B (coordinates: 131.92595, -26.17664; location 11 in Fig. 1 of the main paper). (bottom right) Unoriented sample SW14-237C (coordinates: 131.83544, -26.25303; location 13 in Fig. 1 of the main paper).

S4 Plagioclase stability microstructure 3



135 Figure S5: BSE image of plagioclase with kyanite inclusions (microstructure 3). Elemental maps (Al, Ca, Fe) do not indicate the presence of epidote. The host rock is a statically overprinted dolerite dyke with a stable mineral assemblage of Pl + Cpx + Grt + Ilm + Ky + Rt + Qz. Relict orthopyroxene is locally present (top right corner). Thin section is not oriented. Sample SW13-167 (coordinates: 131.77475, -26.30845; location 14 in Fig. 1 of the main paper).