

# Fluid-mediated, brittle-ductile deformation at seismogenic depth: Part II – Stress history and fluid pressure variations in a shear zone in a nuclear waste repository (Olkiluoto Island, Finland)

5 Francesca Prando<sup>1</sup>, Luca Menegon<sup>1,2</sup>, Mark. W. Anderson<sup>1</sup>, Barbara Marchesini<sup>3</sup>, Jussi Mattila<sup>4,5</sup> and  
Giulio Viola<sup>3</sup>

<sup>1</sup> School of Geography, Earth and Environmental Sciences, University of Plymouth, PL48AA Plymouth, UK

<sup>2</sup>The Njord Centre, Department of Geoscience, University of Oslo, P.O. Box 1048 Blindern, Norway

10 <sup>3</sup>Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università di Bologna, Italy

<sup>4</sup> Geological Survey of Finland, Espoo, Finland

<sup>5</sup> Currently at: Rock Mechanics Consulting Finland Oy (RMCF), Vantaa, Finland

Correspondence to: Francesca Prando (francesca.prando@plymouth.ac.uk) and Luca Menegon  
15 (luca.menegon@plymouth.ac.uk)

**Abstract.** Microstructural record of fault rocks active at the brittle-ductile transition zone (BDTZ) may retain information on the rheological parameters driving the switch in deformation mode, and on the role of stress and fluid pressure in controlling different fault slip behaviours. In this study we analysed the deformation microstructures of the strike-slip fault zone BFZ045 in Olkiluoto (SW Finland), located in the site of a deep geological repository for nuclear waste. We combined microstructural analysis, electron backscatter diffraction (EBSD), and mineral chemistry data to reconstruct the variations in pressure, temperature, fluid pressure and differential stress that mediated deformation and strain localisation along BFZ045 across the BDTZ. BFZ045 exhibits a mixed ductile-brittle deformation, with a narrow (< 20 cm thick) brittle fault core with cataclasites and pseudotachylytes that overprint a wider (60-100 cm thick) quartz-rich mylonite. Mylonitic deformation took place at 400-25 500° C and 3-4 kbar, typical of the greenschist facies metamorphism at the base of the seismogenic crust. We used the recrystallized grain size piezometry for quartz to document a progressive increase of differential stress, from ca. 50 MPa to ca. 120 MPa, towards the shear zone centre during mylonitization and strain localisation. Synkinematic quartz veins formed along the mylonitic foliation due to transiently high pore fluid pressure, up to lithostatic value. The overprint of the veins by dynamic recrystallization and mylonitic creep is further evidence of the occurrence of brittle events under overall ductile conditions.  
30 We propose a conceptual model in which the ductile-brittle deformation cycle was controlled by transient oscillations in fluid pressure and progressively higher differential stress, possibly occurring in a narrowing shear zone deforming towards the peak strength of the crust at the BDTZ.

## 1 Introduction

35 The change from fracturing and frictional sliding to dominant thermally activated creep processes accommodating viscous  
flow in mylonitic rocks occurs at the brittle-ductile transition zone (BDTZ; e.g. Kohlstedt et al., 1995; Handy et al., 2007).  
Strength envelopes predict that the BDTZ coincides with a peak strength in the crust at an approximate depth of 10-15 km  
between the brittle upper crust and the ductile middle- and lower crust. Naturally constrained stress profiles through exhumed  
mid-crustal rocks are consistent with this picture (Behr and Platt, 2011). However, cyclical switches in deformation style  
40 during the evolution of mid-crustal shear zones (e.g. Pennacchioni and Mancktelow, 2007; Fousseis and Handy, 2008; Wehrens  
et al., 2016; Melosh et al., 2018) demonstrate that the BDTZ occupies a depth interval that can vary transiently, reflecting  
changes in, e.g., bulk strength of the shear zones (Hirth and Tullis, 1994; Scholz, 1998; Fossen and Cavalcante, 2017; Melosh  
et al., 2018) and fluid pressure (Cox, 2010; Kjøl et al., 2015; Sibson and Rowland, 2003; Yardley and Baumgartner, 2007;  
Hirth and Beeler, 2015; Marchesini et al., 2019). Lithology, P-T conditions, as well as variations in stress, strain rate and fluid  
45 pressure are important factors controlling the occurrence of different deformation mechanisms (dislocation creep, diffusion  
creep, fluid-assisted veining, dissolution-precipitation creep, fracturing and cataclasis) that overlap in space and time at the  
BDTZ. It is important to assess whether evidence of cyclical fluctuations of those parameters is preserved in the geological  
record and whether the extent of such variations can be estimated by examining natural fault rocks.

Microstructures can record crucial information on the parameters steering deformation at the BDTZ, and are an invaluable tool  
50 that enables derivation of rheological parameters of shear zones (e.g. Stipp et al., 2002; Behr and Platt, 2011; Ceccato et al.,  
2018). However, the mutual overprinting relationships between brittle and ductile deformation and associated fault rocks at  
the BDTZ typically result in only partial microstructural records, in which the youngest deformation event might have  
completely overprinted the evidence of earlier deformation episodes. Recent deformation experiments have opened up new  
avenues for the investigations of natural deformation microstructures in quartz-rich rocks that result from stress variations  
55 during brittle-ductile deformation. The ‘kick and cook’ experiments, for example, have documented quartz microstructures  
formed during transient high stress deformation followed by stress relaxation (Trepmann et al., 2007). Deformation  
experiments conducted by Kidder et al. (2016) show that the microstructure associated with a stress increase in quartzite is a  
bimodal distribution of recrystallized grain size. The smaller grains accurately record the stress increase, whereas the surviving  
coarser grains formed during earlier, lower stress deformation. The smaller grains can be used to constrain differential stresses  
60 during the most recent (high stress) deformation event using a recrystallized grain size palaeopiezometer (Stipp and Tullis,  
2003; Cross et al., 2017).

Fluids can also play a fundamental role in triggering a transient switch from dominantly ductile to brittle deformation, as  
demonstrated, for example, by the synkinematic emplacement of quartz veins subsequently overprinted by crystal-plastic  
deformation (Handy et al., 2007; Kjøl et al., 2015; Trepmann and Seybold 2019; Marchesini et al., 2019). Cyclical ductile-

65 brittle-ductile deformation associated with high fluid fluxes involving a fault-valve behaviour (Sibson, 1990) implies cycles of fluid pressure build-up followed by fluid venting and pressure drop, and has been related to seismic fault behaviour (Sibson, 1992; Cox, 1995; Nguyen et al., 1998; Viola et al., 2006).

Fault modelling and field studies must attempt to quantify the thermal and structural history of fault rocks, as well as the fluid activity in faults to identify the parameters controlling the dominant deformation mode and mechanisms active at seismogenic  
70 depths. This study investigates the microstructural record of the deformation behaviour at the BDTZ of a subvertical sinistral strike-slip fault hosted in the basement of the Paleoproterozoic Baltic Shield in Finland. The fault occurs within the deep ONKALO spent nuclear fuel repository that is currently being built on the island of Olkiluoto in SW Finland (Fig. 1a). The present-day structure of the fault consists of a narrow (< 20 cm thick) brittle fault zone core that is contained inside a thicker (max 1 m thick) ductile, mylonitic shear zone. We constrain the deformation history of the fault zone and use quartz  
75 microstructure to estimate the stress history of the mylonitic precursor. We propose a conceptual model of the evolution of fault slip behaviour that incorporates the constraints on differential stress and fluid pressure derived from our microstructural analysis, and that applies to a narrowing shear zone that progressively localises strain when deforming across the BDTZ.

## 2 Geological Setting

The island of Olkiluoto in SW Finland (Fig. 1a) is part of the Baltic Shield, which was consolidated during the latest  
80 accretionary stages of the Svecofennian Orogeny between ca.1.89 and 1.80 Ga (Lahtinen et al 1994, Nironen et al. 1997, Lahtinen et al 2005, Pajunen et al. 2008). In the study area, the dominant rock types consist of amphibolite facies migmatitic metasediments formed between ca.1.84-1.82 Ga, at 3.7-4.2 kbar and 660-700° C (Tuisku and Karki, 2010, Aaltonen et al. 2016), ca. 1.89 and 1.85 Ma old calc-alkaline synorogenic TTG-type granitoids that predated the migmatisation (Mänttari et al., 2006), and syn- to late-orogenic leucogranites and pegmatites intruded ca. 1.85 to 1.79 Ga (Mänttari et al., 2010). Following  
85 the peak metamorphism leading to migmatisation, retrograde metamorphism under greenschist facies conditions affected the area and continued throughout the subsequent orogenic collapse, dated at ca.1.79-1.77 Ga for SW Finland (Lahtinen et al., 2005).

Polyphase ductile deformation affected Olkiluoto between ~1.86 and 1.79 Ga (Aaltonen et al., 2016), followed by a polyphase brittle deformation history between ~1.75 to 0.8 (Mattila and Viola, 2014). Field studies identified characteristic structures for  
90 three (D<sub>2</sub>-D<sub>4</sub>) deformation stages, which overprint a pre-migmatite, poorly preserved deformation stage (D<sub>1</sub>; Aaltonen et al. 2010, and references therein). Deformation during D<sub>2</sub> (~ 1.86-1.83 Ga) and D<sub>3</sub> (~1.83 to 1.81 Ga) occurred under amphibolite facies condition; D<sub>2</sub> developed a penetrative NE-SW striking high-grade (locally migmatitic) foliation dipping moderately towards SE, as well as NE-SW striking mesoscopic shear zones. D<sub>3</sub> developed NNE-SSW striking foliations, and E-W to NE-SW trending, S to SE dipping shear zones (Aaltonen et al. 2010). The latest stage, D<sub>4</sub>, developed under greenschist facies  
95 retrograde metamorphism around ~1.81–1.79 Ga according to U/Pb dating of syn-kinematic pegmatites (Mänttari et al., 2010).

D<sub>4</sub> structures consist of NNE-SSW and N-S striking subvertical ductile shear zones, varying in thickness from ~ 0.5 m to 200 m (Fig.1a).

100 Brittle deformation in Olkiluoto was characterized both by the reactivation of optimally oriented pre-existing ductile structures, and by the formation of new Andersonian-type faults and joints. The dominant brittle structures can be grouped into two main sets: (1) an E-W to NE-SW trending set of low angle faults exploiting the D<sub>2</sub> regional migmatitic foliation, and (2) a set of subvertical faults striking N-S to NW-SE (Fig. 1; Aaltonen et al., 2016). Paleostress inversion of fault slip data permitted to identify seven distinct brittle stages during the Paleoproterozoic-Mesoproterozoic structural history of SW Finland (Mattila and Viola, 2014). Of interest for this study is the first brittle deformation stage that developed conjugate systems of NNW–SSE sinistral and NW–SE dextral strike-slip faults, in accordance with the NW–SE to NNW–SSE compression proposed for 105 the late- to post- Svecofennian orogeny (Viola et al., 2009; Torvela and Ehlers, 2010; Saintot et al., 2011). Field observations indicate that faults optimally oriented for the first stage of brittle deformation commonly exploit NNE-SSW and N-S precursor shear zones characteristic of the D<sub>4</sub> stage of ductile deformation (Aaltonen et al, 2016; Nordbäck et al., 2018; Skyttä and Torvela, 2018).

110 A network of vertical N-S and NW-SE faults has been mapped and investigated at the repository scale in Onkalo with underground surveys and boreholes (Aaltonen et al. 2016, Fig 1b). N-S faults are typically localised on mica-rich precursor ductile shear zones (Pere, 2009). The role of subvertical, N-S striking ductile precursor zones in controlling the localisation of faults at the disposal site was investigated by Skyttä and Torvela (2018), who identified the ductile precursor structures as short limbs of D<sub>4</sub> asymmetric folds and as anastomosing networks of discrete retrograde (greenschist facies) ductile shear zones. Skyttä and Torvela (2018) proposed that the subvertical N-S faults formed as a result of progressive strain localisation 115 during the late stage of D<sub>4</sub>, which culminated in the development of discrete faults through linkage of individual fault segments that preferentially exploited optimally oriented branches of the anastomosing network of localised ductile shear zones. Therefore, the faults exploiting D<sub>4</sub> shear zones represent ideal targets to investigate the deformation processes and mechanisms at the brittle-ductile transition in Olkiluoto, and the associated rheological parameters recorded in the fault rock microstructures. This study uses the N-S sinistral strike-slip fault BFZ045 as a natural laboratory to investigate the stress 120 history of the ductile precursor and the possible role of fluids on the deformation processes active at the brittle-ductile transition. The companion paper by Marchesini et al. (2019) has thoroughly described the deformation history of the conjugate dextral BFZ300 fault.

### 3 Methods

#### 3.1 Sampling

125 Samples were obtained from two sub-horizontal drill cores from the underground facilities that intersect the BFZ045 fault (Fig. 2a). The analysed samples were selected from (i) a 2 m continuous section along drill core PH28, which was drilled at a depth of 433 m b.s.l. and is oriented ESE-WNW, and (ii) drill core PH16 from the Demonstration Facilities tunnels at 420 m

depth, described in Aaltonen et al. (2016). The PH28 samples were selected from the 99-101 m interval of the drill core, with core distance measured from ESE to WNW, which includes the fault core and the proximal damage zone. The selected interval of the drill core was cut in half parallel to the stretching lineation and perpendicular to the mylonitic foliation and subsampled at regular intervals of 2 cm. Nineteen polished thin sections were added to three samples from PH16 previously described in Aaltonen et al. (2016). An additional thin section from drill core PH21 (drilled at the same depth of PH16) was used to estimate the T of the graphite formation in the host rock using Raman spectroscopy of carbonaceous material (see 3.2 and 4.5).

### 135 **3.2 Microstructural observations and Electron Backscatter Diffraction (EBSD) analysis**

Deformation microstructures were studied using petrographic and scanning electron microscopy (SEM). SEM and EBSD analysis were performed at the Plymouth University Electron Microscopy Centre using a JEOL LV6610 SEM and a JEOL 7001 FEG-SEM. Thin sections used for Electron Backscatter Diffraction (EBSD) analysis were polished with colloidal silica before being carbon coated. Data were acquired on a NordlysNano and a NordlysMax EBSD detector (Oxford Instruments). Working conditions during acquisition of the EBSD patterns were 20 kV, 20 mm working distance, 70° sample tilt and high vacuum. AZtec software was used for pattern indexing on rectangular grids with step size of 0.7 µm, 1 µm and 1.8 µm. EBSD patterns were processed with the Channel 5 software (Oxford Instruments), and noise reduction was performed following the procedure suggested in Bestmann and Prior (2003).

Differential stresses during mylonitic creep were estimated using the recrystallized grain size piezometer for quartz of Cross et al. (2017). The method relies on the separation between relict and recrystallized grains based on the Grain Orientation Spread (GOS), which is a measure of the internal strain of a grain defined as the average misorientation angle between each pixel in a grain and that grain's mean orientation (Wright et al., 2011). Further details on the presentation of the EBSD data and on the recrystallized grain size piezometer for quartz of Cross et al. (2017) are presented in the supplementary material (S1).

### **3.3 Mineral chemistry and Raman spectroscopy**

150 Major element mineral chemistry of chlorite and white mica was measured with Electron Microprobe Analysis (EMPA), at the Department of Earth Sciences, University of Milan, Italy. Carbon coated thin sections were probed with a JEOL 8200 Super Probe equipped with 5 wavelength-dispersive spectrometer (WDS). Working conditions were set to 15 kV of probe current, 5nA current on sample, 1 µm beam diameter. Natural minerals were used for standardization, measurement times were 30s on peaks and 10s on backgrounds of the X-ray lines.

155 Raman spectroscopy was applied for feldspar and opaque phase identification and for carbonaceous material (CM) characterization. Data acquisition was conducted at the Department of Chemistry, University of Padua (Italy), using a Thermo Scientific DXR MicroRaman spectrometer, equipped with a 532 nm depolarised laser. Raman analysis was also carried out at the Department of Mathematical, Physical and Computer Sciences of the University of Parma (Italy) using a Jobin-Yvon Horiba LabRam spectrometer equipped with He-Ne laser (emission line 473.1 nm) and motorized XY stage. Spectra were

160 acquired from polished thin section, using a laser power of 5 mW, spectrograph aperture 25  $\mu\text{m}$  pinhole, and 50X or 50X low  
distance objective. The estimate spot size was 1-2  $\mu\text{m}$  in diameter and spectral resolution of 2 to 4.4  $\text{cm}^{-1}$ , with acquisition  
time of 30–90 s. Feldspars composition was classified on the basis of the acquired Raman spectra, as suggested in Freeman et  
al. (2008), using a comparison with standard Raman spectra from the RRUFF Project database (Lafuente et al. 2015). To  
165 assure a good statistical analysis of the CM structural heterogeneity, only samples with > 10 CM spectra were taken in  
consideration. Omnic software (Thermo Fisher Scientific) was used for Raman spectrum decomposition, using the software  
Lorentian/Gaussian function, following the procedure described in Koeketsu et al. (2009). Raman analytical conditions are  
detailed reported in the supplementary material (S1). A discussion about the analytical limitation of the chosen methods for  
temperature estimation is also presented in S1.

## 4 Results

### 170 4.1 BFZ045 fault zone structure

A schematic representation of BFZ045 fault geometry obtained from underground field mapping and detailed characterization  
along the horizontal PH28 drill core is represented in Fig. 2a. A 10-100 cm wide mylonite with a sinistral sense of shear,  
oriented 61/075 (dip/dip direction), hosts a sub-vertical network of cohesive cataclasites with an average orientation of 87/095  
(Fig. 2b). Rodding of quartz and feldspar in the mylonite defines a stretching lineation oriented 10/168 (plunge/trend; Fig.  
175 2b,c). Slickensides with chlorite mineral striations are abundant throughout BFZ045 and the average orientation of the  
striations is 07/169; stepped slickensides indicate a dominant sinistral sense of shear, although striations associated with dextral  
kinematics have also been observed (Aaltonen et al. 2016; Nordbäck et al., 2018).

A network of veins typically filled with chlorite, quartz and calcite, and chlorite filled fractures overprint the mylonite and  
host rock (Fig 2a; Aaltonen et al, 2016). The damage zone is localised between 0.5 to 1 m from the fault core, and is  
180 characterized by an increased fracture density towards the fault core, with the average spacing between fractures decreasing  
from 3 cm to ca. 0.5 cm. With the term “fault core”, we refer here to the brittle core of BFZ045 defined by cohesive cataclasites  
that overprint the mylonite. This means that the damage zone of BFZ045 affects both the migmatitic host rock and the BFZ045  
mylonite.

In figure 2d, representative samples of core PH28 are shown from left to right in a sequence from the damage zone in the host  
185 rock to the fault core. The host rock is a coarse-grained veined migmatite (Fig. 2d, I). The extent of damage in the host rock  
was defined by the farthest occurrence of chlorite filled fractures, identified microscopically as deformation bands associated  
with visible slip. A total of 12 fractures longer than 5 cm can be observed along the core, of which 7 within a distance of 10  
cm from the fault core (west side of the core). The contact between the host rock and the mylonite is sharp. The millimetre-  
spaced mylonitic foliation is defined by a compositional layering of alternating quartzofeldspathic domains and mica-rich  
190 domains (Fig. 2d, II). Multiple slip surfaces marked by 0.5 – 10 cm thick cataclastic domains overprint the mylonite along the  
foliation (Fig. 2d, III). Locally, phyllosilicates and trails of opaque minerals define thin (<1 mm thick) anastomosing foliation

planes within the cataclasites, which wrap around sub-angular fragments of the mylonitic precursor. Along a thin (<5 mm thick) slip surface, two pseudotachylyte injection veins intruding the mylonite at a high angle have been observed, which demonstrates the transient seismogenic behaviour of BFZ045. Calcite veins (1-3 mm thick) locally overprint the fault core both at high angle to the foliation and along the slip surfaces.

## 4.2 Petrography and microstructures

### 4.2.1 Host rock

The migmatitic host rock mineral assemblage observed in the PH28 samples consists of 50 % of quartz, 40 % K-feldspar, and plagioclase, and the remaining 10% of white mica, graphite, alteration minerals of plagioclase (sericite) and secondary chlorite, and calcite veins. Coarse-grained (1-2 mm) quartz and feldspars are homogeneously distributed in an equigranular texture and show mostly straight grain boundaries (Fig. 3a). K-feldspar occurs as orthoclase, with a well-developed veined perthitic texture. Plagioclase, albitic in composition as determined from Raman analysis, has lamellar twinning and is commonly altered to sericite. Both types of feldspar locally show bent twin lamellae and undulose extinction. Microfractures in feldspars are preferentially oriented parallel and at low angle ( $\leq 20^\circ$ ) to the mylonitic foliation. White mica was observed occasionally as millimetric sub-euhedral grains. Quartz displays intracrystalline deformation features such as undulatory extinction, wide extinction bands (WEBs, following the terminology of Derez et al., 2015; Fig. 3b), and bulges resulting in sutured grain boundaries (Fig. 3c; Stipp and Kunze, 2008). WEBs are locally bounded by fluid inclusion trails with different orientations, which give them a blocky or slightly elongated aspect (Figs. 3b-c). Two main sets of intracrystalline fluid inclusion trails are observed, one at a low angle with respect to the mylonitic foliation and the other perpendicular to the foliation. Fine bulges (10-20  $\mu\text{m}$  in size) occur along grain boundaries and intercrystalline fractures (Fig. 3b). Quartz grains in the proximity of the mylonite (sample PH28-2, Fig. 2b) develop intracrystalline bands of recrystallized grains sub-parallel to the foliation, with grain size of  $\sim 30\text{-}60 \mu\text{m}$  (Fig. 3c). Shear bands in the deformed host rock appear as cohesive micro-cataclasites, with fine grained ( $< 10 \mu\text{m}$ ) chlorite surrounding angular feldspar and quartz clasts.

### 4.2.2 Mylonite

The relative mineral abundances in the mylonite are slightly different from those in the host rock, and consist of 50% quartz, 20% white mica + chlorite, and 30% K-feldspar + plagioclase. Accessory phases are graphite, rutile, anatase, and apatite, which are typically found associated with chlorite to form black seams. The spaced mylonitic foliation is defined by a compositional layering between alternating millimetre-thick quartz bands and narrower (0.2 to 1 mm thick) mica- and feldspars-rich bands (Fig. 4a). Porphyroclasts of K-feldspar are up to 7 mm in size and show asymmetric pressure shadows filled with chlorite + muscovite  $\pm$  feldspars (albite and K-feldspar), with a geometry indicative of a sinistral sense of shear. Feldspars in the mylonite present brittle behaviour, with domino-type fragmented porphyroclasts. Neoblasts of K-feldspar are

commonly observed in the pressure shadows of porphyroclasts or together with fine-grained white micas along the mylonitic foliation.

225 Quartz in the mylonite presents various degrees of recrystallization (Figs. 4b, c). In zones where ductile deformation is less prominent (e.g. samples PH28\_3, PH16\_1, Fig. 2), quartz retains a coarser grain size (1-5 mm), and forms slightly asymmetric (sigmoidal-shaped) grains. Internally, the grains display patchy undulatory extinction, well-developed blocky to elongated WEBS (100-300  $\mu\text{m}$  wide), and discrete intracrystalline bands (< 200  $\mu\text{m}$  wide) of bulges and recrystallized grains preferentially oriented sub-parallel to the foliation and at ca. 45° from the foliation, measured anti-clockwise (Fig. 4b). The  
230 host grains contain small subgrains (< 60  $\mu\text{m}$ ), which, towards the boundaries of the host grain, make transition to aggregates of recrystallized grains of size comparable to the subgrains, forming typical core-and-mantle microstructures (Fig. 4b).

In zones of complete recrystallization, quartz forms highly elongated polycrystalline ribbons (up to 0.5 – 1 mm thick, and up to 2 cm long) parallel to the foliation (Fig. 4c). The recrystallized grains locally define a shape preferred orientation (SPO in Fig. 4c) inclined with 10° to 30° with respect to the trace of the foliation, consistently with the bulk sinistral sense of shear.

235 Adjacent to the cataclastic fault core (sample PH16\_3, Fig. 2), ataxial/unitaxial (i.e. with no visible median line; Bons, 2012) quartz veins occur parallel to the mylonitic foliation (and to the layers of recrystallized quartz) and contain grains elongated normal to the vein boundary (i.e. normal to the foliation; Fig. 4d). The vein crystals range in length from 200-400  $\mu\text{m}$  and have a maximum thickness of 150  $\mu\text{m}$  measured parallel to the vein. Quartz in the veins shows undulatory extinction and bulges at the grain boundaries (Fig. 4e) indicative of crystal plastic deformation. The recrystallized quartz in the mylonite surrounding  
240 the vein has a finer grain size than the one in the mylonite described in Figs. 4a-c, which is located farther away from the cataclastic core. Veins of radiate chlorite are observed cutting the mylonitic foliation at a high angle (~60°).

#### 4.2.3 Cataclasite

The brittle overprint in BFZ045 occurs mostly as 3 to 10 cm thick protocataclasites, that transition to 0.5 - 2 cm thick cataclasite bands in the fault core. The cataclasite is rich in chlorite and opaque minerals, which occur as fine-grained (2-10  $\mu\text{m}$ ) flaky  
245 aggregates within the fine grained (< 50  $\mu\text{m}$ ) quartz + feldspars + muscovite rich matrix. Locally, the cataclasite matrix contains a foliation defined by aligned phyllosilicates and anastomosing dark seams of opaque minerals. Clasts are predominantly angular fragments of the mylonite, ranging in size from 100  $\mu\text{m}$  to 5 mm, and surrounded by a variable proportion of matrix (Figs. 3a-c). Quartz in the protocataclasite occurs in almost entirely recrystallized ribbons with a finer grain size (ca. 10  $\mu\text{m}$ ) than the one observed at higher distances from the brittle fault core (Figs. 5b-c vs Figs. 4b-c). The quartz clasts in the cataclasite  
250 (Fig. 5c) preserve the deformation and recrystallization microstructures observed in the mylonite in close proximity to the cataclasite (Fig. 5b).

A pseudotachylyte generation surface is observed subparallel to a cataclastic band (Fig. 5a) and is identified from characteristic centimetric injection veins, branching in the mylonite at high angle to the foliation. The pseudotachylyte main generation surface is less than 1 mm thick and is parallel to the mylonitic foliation (Figs. 5a, d). The matrix of the pseudotachylyte is  
255 completely altered to a fine-grained, < 2  $\mu\text{m}$ , chlorite and muscovite rich matrix that surrounds survivor clasts of quartz and

rutile (Fig. 5d). Chlorite- and quartz aggregates commonly fill fractures within feldspar porphyroclasts in the mylonite. Similar fractured feldspar porphyroclasts with quartz + chlorite fillings are observed inside slightly rotated clasts of mylonite in the protocataclasites (Fig. 5e).

### 4.3 EBSD and grain size analysis of quartz

#### 260 4.3.1 Mylonite

EBSD analysis of the mylonite was conducted on sample PH16\_1, which is located at the mylonitic shear zone boundary at a distance of 4 cm from the brittle fault core (Fig. 2a). EBSD maps were acquired from intracrystalline bands of recrystallized grains within an elongated mm- sized quartz grain (Fig. 6a), and from a highly recrystallized quartz layer along the mylonitic foliation (Fig. 7a).

265 The recrystallized grain size within the intracrystalline bands ranges from 5 to 60  $\mu\text{m}$  (Fig. 6b, c). The Grain Orientation Spread (GOS) within the recrystallized bands varies between  $0^\circ$  and  $8.4^\circ$ , with a threshold value of  $1^\circ$  between the recrystallized grains and the relict grains when analysing the trade-off curve proposed by Cross et al. (2017). The average grain size of recrystallized grains ( $\text{GOS} < 1^\circ$ ) is  $16 \pm 7 \mu\text{m}$  whereas relict grains ( $\text{GOS} > 1^\circ$ ) have an average grain size of  $25 \pm 9 \mu\text{m}$ . Relict grains contain subgrains of an average size of  $17 \pm 7 \mu\text{m}$  (Fig. 6f).

270 Quartz grain hosting the intracrystalline band shows subgrains of approximately 25-50  $\mu\text{m}$  in size, which is comparable to the size of the coarser recrystallized grains observed in the intracrystalline bands (Fig. 6c). The size of the subgrains in the host quartz was estimated visually with the aid of Grain Relative Orientation Distribution maps (GROD, Fig. 6g).

In the recrystallized quartz layer (Fig. 7a), quartz grain shape ranges from equant to elongate parallel to the foliation, with grain size ranging from 5 to 87  $\mu\text{m}$  (Figs. 7b, c). GOS analysis identified a threshold value of  $1.56^\circ$  to separate recrystallized- and relict grains (Fig. 7d). Average grain size of the recrystallized grains is  $18 \pm 8 \mu\text{m}$ , while relict grains have an average size of  $28 \pm 11 \mu\text{m}$  (Fig. 7e).

275 The relict grains contain subgrains of an average size of  $17 \pm 7 \mu\text{m}$  (Figs. 7c, f). The crystallographic preferred orientation (CPO) of the c-axis of the relict grains and recrystallized grains forms a single girdle consistently inclined with the sinistral sense of shear of the sample (Fig. 7g). The EBSD-calibrated recrystallized grain size piezometer for quartz of Cross et al. (2017) was used to estimate the differential stresses during plastic flow in the mylonite. The estimated differential stress is 73- 80 MPa for the average recrystallized grain size of 16-18  $\mu\text{m}$ .

#### 4.3.2 Veins parallel to the mylonitic foliation

285 The quartz vein parallel to the foliation shown in Fig. 5c and 8a was analysed to identify possible evidence of crystal-plastic deformation and dynamic recrystallization. Grain shape varies from fibrous with elongation perpendicular to the vein wall, to more equant/less elongate. Grain boundaries of vein crystals are straight to lobate, the latter most commonly observed in association with fine recrystallized grains and bulges ( $< 15 \mu\text{m}$  in size) (Figs. 8 b, c). Irrespective of their shape, most of the

grains contain low-angle boundaries and Dauphiné twins. The low-angle boundaries are typically arranged to define polygonal to slightly elongated domains of  $\sim 10 \mu\text{m}$  in size, comparable to that of the surrounding recrystallized grains in the mylonite (Fig 8c).

290 Quartz in the mylonite flanking the vein shows fine grain size ( $< 20 \mu\text{m}$ ), with only a small fraction of coarser grains ( $30\text{--}60 \mu\text{m}$ ). GOS analysis indicates that dynamic recrystallization is pervasive. The mean recrystallized grain size is  $10 \pm 3 \mu\text{m}$ , and the relict average grain size is  $20 \pm 9 \mu\text{m}$ , with a GOS threshold value of  $1.94^\circ$  (Figs. 8d, e). Selected relict grains (size  $> 40 \mu\text{m}$ ) in the mylonite contain subgrains with size between  $5$  to  $25 \mu\text{m}$  range (Fig. 8f). Finer grains observed within the vein also present a GOS value below the threshold, which suggests they represent the recrystallized fraction in the quartz vein (Figs. 8c, 295 f). In the vein quartz, GOS analysis indicates that the average subgrain size is  $24 \pm 7 \mu\text{m}$ , although the largest fraction of subgrains is smaller than  $15 \mu\text{m}$ , i.e., similar in size to the recrystallized grains in the flanking mylonite and in the vein itself. The c-axis CPO of the recrystallized grains in the mylonite forms a single girdle synthetically inclined with the sinistral sense of shear (Fig. 8h). The c-axis CPO of recrystallized grains in the veins overlaps with the one of the relict grains (Fig. 8i). The differential stress estimated from the average recrystallized grain size in the mylonite ( $10 \mu\text{m}$ ) is  $106 \text{ MPa}$ .

### 300 4.3.3 Cataclasite

We analysed a largely recrystallized quartz clast in the cataclasite from sample PH28\_10 (Fig. 9a). The selected clast is rotated of less than  $10^\circ$  with respect to the adjacent mylonitic foliation. Quartz grain size in the clast ranges from  $4$  to  $60 \mu\text{m}$ . The coarser grains are elongated parallel to the foliation, show bulges and fine recrystallized grains at their boundaries, and contain a high density of low-angle boundaries (Fig. 9b). The low-angle boundaries define small polygonal domains of a size 305 comparable to the one of the recrystallized grains found at the grain boundaries (Fig. 9c). The GOS map in figure 9d identifies two grain size distributions, separated by a GOS threshold value of  $3.23^\circ$ . The recrystallized grains (average grain size:  $8 \pm 4 \mu\text{m}$ ) form equigranular aggregates at the boundaries of the coarser (average grain size:  $17 \pm 10 \mu\text{m}$ ) elongated relict grains. The c-axis CPO of the recrystallized grains and of the relict grains is the same, showing two maxima at an intermediate position between the centre of the pole figure and its periphery, and consistently inclined with the sinistral sense of shear of the sample 310 (Fig. 9f). The differential stress estimated from the average recrystallized grain size in the clast ( $8 \mu\text{m}$ ) is  $123 \text{ MPa}$ . Although the map has been acquired from a clast, these microstructures and recrystallized grain size are similar to those of the mylonite in the immediate vicinity ( $< 2 \text{ cm}$ ) of the brittle fault core (Figs. 5b, c).

## 4.4. Mineral chemistry, Raman spectroscopy, and pressure-temperature (P-T) conditions of deformation

We estimated the P-T conditions of mylonitic deformation using Raman spectroscopy of carbonaceous material (RSCM), 315 chlorite thermometry, and phengite barometry. Carbonaceous material was observed as grains and aggregates ranging in size from  $\sim 50$  to  $\sim 200 \mu\text{m}$  in the host rock (sample PH21\_1, Fig. 10a) and as smaller grains ( $20\text{--}50 \mu\text{m}$ ) along the mylonitic foliation (sample PH16\_1-2, Fig. 10b), along chlorite and muscovite rich layers. Data of Raman peaks deconvolution are reported in the Supplementary Material (S1). We estimated a maximum temperature of  $530^\circ \text{C} \pm 50^\circ \text{C}$  for the carbonaceous

material in the host rock (Fig. 11a) (using the thermometer calibration for a laser wavelength of 514 nm, Beyssac et al. 2002),  
320 and a lower T of  $440^{\circ}\text{C} \pm 50^{\circ}\text{C}$  for the mylonite (Aoya et al., 2010, using the thermometer calibration for a laser wavelength  
of 532 nm; Fig. 11a).

The pressure during mylonitization was estimated using the Si-in-phengite geobarometer (Massonne and Schreyer, 1987).  
Representative compositions of white mica are listed in Table 1. The full dataset of chemical compositions of white micas is  
reported in the Supplementary Material (S1). White mica composition was measured for grains parallel to the foliation  
325 associated with neoblasts of K-feldspar, as the application of the Si-in-phengite geobarometer requires stability of K-feldspar  
(Figs. 10c, d). Structural formulae were calculated assuming 11 oxygens. The range of Si apfu in the probed white mica grains  
is 3.12 –3.16. This compositional range indicates a pressure of 2-4 kbar for the mylonitization (Fig. 11b), considering the  
average temperature of  $440^{\circ}\text{C}$  derived for the mylonite with the graphite thermometry.

Chlorite composition was determined for i) chlorite grains intergrown with quartz and muscovite in the strain shadows around  
330 feldspar porphyroclasts in the mylonite (Fig. 10c, d), ii) chlorites flakes aggregate in the cataclasite quartz matrix (Fig. 10e),  
and iii) radiate chlorites filling veins at high angle to the mylonitic foliation (Fig. 10f). The structural formula of chlorite was  
calculated based on 14 oxygens, and representative composition are shown in Table 2. The full dataset of chemical  
compositions for chlorite is reported in the Supplementary Material (S2).

Chlorites along the mylonitic foliation have Si content ( $\sim 2.60 - 2.70$  apfu), Al between 2.74-2.88 apfu, and are moderately Fe-  
335 rich with a XFe ( $\text{XFe} = \text{Fe}/(\text{Fe} + \text{Mg})$ ) between 0.62 and 0.82. Chlorites in the mylonitic sample from PH\_16 have a more narrow  
range of XFe, between 0.57 and 0.73( see supplementary material S1) .In the cataclasite chlorites Si content range between  
2.53-2.772 apfu, and Al have a wider range of 2.61-2.94 apfu (S1). The cataclastic chlorite is richer in Fe compared to the  
mylonite, with a XFe range between 0.71-0.85. The radial chlorite filling the veins cutting the mylonite has Si content between  
2.51 and 2.780 apfu, Al between 2.74 and 3.00 apfu, and XFe vary between 0.69 and 0.81. In general, BFZ045 chlorites have  
340 a aphaesiderite-ripidolite composition and the microprobe results show that the composition of distinct chlorite generations is  
similar (Fig. 11c). The CHL(2) semi-empirical thermometer of Lanari et al. (2014) was applied to each EMPA analysis of  
chlorite with  $\text{Si} < 3$  apfu and  $(\text{Na} + \text{K} + \text{Ca}) < 0.1$  apfu. FeO was used as Fe total, and  $a_{\text{H}_2\text{O}} = 1$  and  $a_{\text{SiO}_2} = 1$  were assumed. The  
estimated temperature for the mylonite ranges from  $\sim 380^{\circ}\text{C}$  to  $500^{\circ}\text{C}$  (limit of the used thermometer) for an assumed P of  
3.5 kbar (Fig. 11d), with a temperature variation of  $10^{\circ}\text{C}$  every 0.5 kbar increment. Temperatures estimated for the cataclasite  
345 matrix and for the radiated chlorite in the veins have a larger range ( $\sim 300$ - $500^{\circ}\text{C}$ , Fig 11d). Compositions yielding T lower  
than  $400^{\circ}\text{C}$  are more frequent, especially in the cataclasite, however no clear relationship could be discerned between  
temperature and microstructural position of the chlorites.

## 5. Discussion

Our observations constrain the details of the structure and the deformation history of BFZ045. In particular, the microstructures  
350 of fault rocks indicate a sequence of deformation events where ductile deformation (mylonitisation) was punctuated by brittle

deformation (veining), and eventually culminated in the formation of the brittle, cataclastic fault core. We interpret this sequence to result from the evolving stress history and possible fluid pressure variations during the overall ductile-to-brittle deformation history of the strike-slip BFZ045 fault.

### **5.1 The sequence of deformation events in BFZ045: ductile-brittle deformation history**

355 Our microstructural observations are consistent with the general conclusion that brittle deformation along BFZ045 exploited a ductile (mylonitic) precursor (Nordbäck and Mattila al., 2018; Skytta and Torvela, 2018). Veins, cataclasites and pseudotachylytes are localised along the mylonitic fabric of BFZ045, and only minor evidence of brittle deformation (mostly in the form of fractures filled by chlorite) is present outside of the mylonitic fault core (Figs. 2a, 3a). The first stage of deformation of BFZ045 is represented by the development of a localised N-S trending mylonitic foliation (Figs. 2, 4a-c).  
360 Mylonitic creep was punctuated by transient brittle events, with the opening of extensional fractures along the mylonitic foliation filled by quartz veins (Figs. 4 d,e). Veining was again followed by mylonitic deformation, as indicated by dislocation creep and dynamic recrystallization microstructures of quartz in the veins. Mylonite and veins were then overprinted by brittle deformation that formed cm- thick cataclasites (Figs. 5a-c) and a < 0.5 cm thick pseudotachylyte (Fig. 5d) that, together, form the brittle fault core of BFZ045. Cataclasites and pseudotachylytes were not overprinted by mylonitic creep, which might  
365 indicate that they formed under condition favourable to predominantly brittle deformation along BFZ045.

The observed parallelism between stretching lineation in the mylonite and chlorite slickenlines in the cataclasites suggest that the ductile-brittle deformation history of BFZ045 occurred under a prolonged strike-slip regime with sinistral kinematics (Figs. 2b,c). This conclusion is consistent with the model of the brittle evolution of SW Finland proposed by Mattila and Viola (2014), which attributes the sinistral kinematics of NNW-SSE trending subvertical faults to the ductile-brittle transition stage  
370 of the basement at 1.75 Ga (Stage 1 of deformation in Mattila and Viola, 2014). BFZ045 experienced later reactivations during the prolonged brittle history of the SW Finland basement, as indicated by calcite veins cutting the brittle fault core and by (rare) slickenlines with dextral kinematics observed in the underground exposures (Aaltonen et al. 2016). Local dextral kinematics along BFZ045 is potentially consistent with the Stage 2 of deformation of Mattila and Viola (2014) at 1.7-1.6 Ga. These later features, however, are not discussed further in this paper as they are subordinate to, and did not obliterate the earlier  
375 history.

### **5.2 Deformation mechanisms and stress history during mylonitic creep of BFZ045**

Microstructural observations show that ductile deformation of the mylonite BFZ045 was accommodated in quartz by dislocation creep, while feldspars experienced dominantly brittle behaviour with only limited neocrystallization in pressure  
380 shadows (Figs. 10c). Given the lack of crystal plastic deformation and recovery features in the interior of feldspar porphyroclasts, we interpret the neocrystallization in pressure shadows as a possible indication of dissolution-precipitation,

which has been commonly reported in feldspars deforming at mid-crustal conditions (Fitz Gerald and Stünitz 1993; Menegon et al. 2008; Eberlei et al., 2014; Torgersen et al., 2015; Giuntoli et al., 2018; Hentschel et al., 2019).

385 Quartz recrystallization microstructures (bulges at grain boundaries and within intracrystalline bands, and core and mantle microstructures with subgrains of comparable size to that of the recrystallized grains; Figs. 4, 5) suggest that bulging and subgrain rotation were the dominant recrystallization mechanisms (Hirth and Tullis, 1992; Stipp and Kunze, 2008). The average recrystallized grain size of quartz ranges between 8 and 18  $\mu\text{m}$ , which falls within the  $< 40 \mu\text{m}$  size value representative of bulging as dominant recrystallization mechanism (defined as slow grain boundary migration coupled to localised subgrain rotation at the mantle of the host grain; Stipp et al., 2010). In all the studied samples, GOS analysis distinguished two quartz  
390 grain size populations, in which the coarser (relict) grain size contains subgrains of size comparable to the average size of the finer recrystallized grains (Figs. 7-9). The CPO of relict and recrystallized grains is the same, and this is consistent with the host-controlled development of a CPO during subgrain rotation recrystallization (e.g. Stünitz et al. 2003).

A key observation in the BFZ045 mylonite is the decrease in recrystallized grain size of quartz from the shear zone boundary (16-18  $\mu\text{m}$ ) towards the centre (8-12  $\mu\text{m}$ ; Fig. 7-9). The coarser (16-18  $\mu\text{m}$ ) grain size is the most representative of the partially-  
395 (Fig. 7) and of the nearly completely (Fig. 8) recrystallized quartz ribbons in the BFZ045 mylonite at distances  $\geq 4 \text{ cm}$  from the brittle fault core, and it also occurs as relict grain size in quartz clasts embedded in the cataclasite, where it is overprinted by the finer (8-12  $\mu\text{m}$ ) recrystallized grains (Fig. 10). We interpret this overprint and the overall decrease in recrystallized grain size to reflect a progressive increase in stress and strain rate towards the shear zone centre during mylonitic creep (Kidder et al. 2016). Throughout the samples the CPO of recrystallized quartz is consistently inclined with a sinistral sense of shear (  
400 Fig. 6-9); this is interpreted as evidence that they developed under constant kinematic conditions in BFZ045.

Relict quartz grains of 25-28  $\mu\text{m}$  in size are associated with the coarser (16-18  $\mu\text{m}$ ) grain size (Figs. 7-8), and un-recrystallized portions of quartz contain subgrains of similar size ( $\geq 25 \mu\text{m}$ , Figs. 7b-d). We consider that the 25-28  $\mu\text{m}$  grain size population might represent an early, dynamic recrystallization event within BFZ045, which was later overprinted by recrystallization occurring under progressively increasing differential stress.

405 Deformation and stress history of BFZ045 can be summarised in different stages. After the regional metamorphic peak and migmatite formation (i.e. 660-700° C, 3.7-4.2 kbar; Tuisku and Lauri, 2009), the basement of Olkiluoto was affected by different stages of ductile deformation (D<sub>2</sub>-D<sub>4</sub>, Aaltonen et al., 2010) under a metamorphic retrograde path toward greenschist facies conditions. The progressive change in P-T condition was favourable to the mobilization and subsequent recrystallization of carbonaceous material (e.g. Kirilova et al., 2017). We thus interpret the estimated metamorphic temperature of 530° C  $\pm$   
410 50° C in the host rock as the temperature of remobilization of carbonaceous material during retrograde metamorphism that culminated in the D<sub>4</sub> deformation stage. During the final stage of D<sub>4</sub> a localised N-S trending mylonitic foliation developed in the migmatites (Fig. 12a). Mylonitic creep took place under differential stresses increasing from  $\sim 54$  to  $\sim 80$  MPa, as determined from the recrystallized grain size of 16-18  $\mu\text{m}$  overprinting grains of 25-28  $\mu\text{m}$ . Foliation-parallel quartz veins crystallized transiently along the mylonitic foliation (Fig. 12b) and were overprinted by crystal plastic deformation and dynamic  
415 recrystallization under progressively higher stress (80-120 MPa, Figs. 8, 12c) during strain localisation towards the centre of

the shear zone. We are aware of the uncertainties and limitations of the palaeopiezometric calibrations, and our estimated flow stresses must be taken with care. However, the similarity between subgrain- and grain size, as well as the consistent sinistral asymmetry of the quartz c-axis CPO of relict and recrystallized grains in all the maps support that dynamic recrystallization occurred during the sinistral strike-slip movement of BFZ045 under progressively increasing differential stresses.

420 We estimate that mylonitization of BFZ045 (Fig. 12a-c) occurred at  $\sim 450^\circ\text{C}$  and 3.5 kbar, consistent with the retrograde greenschist facies conditions attributed to the final stages of  $D_4$  deformation in the SW Finland basement at 1.81-1.77 Ga (Lahtinen et al., 2005; Mänttari et al., 2010; Skytta and Torvela, 2018). Considering an average crustal density of  $2.7\text{ g/cm}^3$ , the depth of mylonitic deformation and of transient veining in BFZ045 corresponds to approximately 13 km.

The mylonite and quartz veins were eventually overprinted by more pervasive brittle deformation that formed chlorite veins, 425 cm-thick cataclasites and a  $< 0.5$  cm thick pseudotachylyte that, together, form the brittle fault core of BFZ045 (Fig. 12 d). We attempted to estimate the temperature of chlorite veins cutting the mylonitic foliation and of chlorite in the cataclastic matrix (Fig. 11), but we obtained a wide temperature range of  $\sim 300\text{-}500^\circ\text{C}$  that is unable to constrain the precise T of the final brittle deformation event(s). Although the conditions of formation of cataclasites and pseudotachylytes cannot be tightly constrained, it is interesting to note that the T estimates from the chlorite in the veins are generally higher than  $400^\circ\text{C}$ . This suggests that 430 chlorite veining occurred early in the deformation history of BFZ045 (i.e. at  $T \geq 400^\circ\text{C}$ ) under overall ductile conditions, as also supported by the precipitation of chlorite + quartz aggregates in microveins and in strain shadows (Fig. 5e). This is consistent with the results of Marchesini et al. (2019), who estimated a temperature of at least  $350^\circ\text{C}$  for the early stages of deformation and emplacement of quartz-chlorite veins along the dextral fault zone BFZ300 (conjugate to BFZ045).

### 435 **5.3 Conceptual model of the fault slip behaviour of BFZ045 at the base of the seismogenic zone**

Using a  $\lambda$ - $\sigma$  failure mode diagram (Cox, 2010), we propose a conceptual model of the structural evolution of BFZ045 (Fig. 13). Strain rates during mylonitic creep were estimated using the dislocation creep flow laws of quartz of Hirth et al. (2001). Parameters and assumptions used for the calculation of the strain rates and of the  $\lambda$ - $\sigma$  failure envelopes are listed in Table 3.

440 According to the failure envelope, differential stresses  $< 80$  MPa are necessary for extensional- and hybrid fractures to occur. In our model, this stage of fault evolution is represented by the  $\sim 54$  MPa creep recorded by quartz grain- and subgrain size of  $25\text{-}28\ \mu\text{m}$  (Figs. 7b-d). The failure envelope also shows that extensional failure and vein formation required transient high fluid pressure reaching lithostatic values ( $\lambda = 1$ ; Fig. 13b, stage 2). Although the overall geometric stress conditions during ductile deformation of BFZ045 are expected to generate en-echelon vein systems oblique to the mylonitic foliation, the 445 foliation-parallel veins are consistent with mode I opening mode due to hydrofracturing. A regional rotation of the stress field appears unlikely, given the constant orientation of the stretching lineation and slickenlines in the core samples, and the consistent asymmetry of the pre- and post-vein quartz c-axis CPO (Figs. 7-9). Thus, a transient reorientation of the stress field in the fault zone must be invoked to explain the switch from viscous creep to mode I fracturing along the mylonite foliation.

Veining was then followed by a decrease in fluid pressure (e.g. Sibson, 1989, 1993; Cox 1995) and a switch back to mylonitic  
450 creep under progressively higher stress and strain rate conditions (Fig. 13c, stage 3). The failure mode diagram indicates that  
these higher stresses are expected to result in brittle shear failure mode of the fault for a pore fluid pressure factor  $\lambda > 0.8$ . The  
cataclasite and pseudotachylytes in the brittle fault core likely represent the final product of BFZ045 deformation under  
progressively higher differential stresses and fluid pressure across the BDTZ. However, the time span between the high strain  
rate mylonitic creep and the formation of the brittle fault core is unknown, and so are the exact P-T conditions of cataclasites  
455 and pseudotachylytes. Thus, our model for stage 3 remains somewhat speculative.

The increase in stress towards the shear zone centre (Fig. 12c) may reflect the rheological evolution of a shear zone that is  
narrowing with exhumation under progressively decreasing T from the ductile to the brittle crust (e.g. Kohlstedt et al., 1995;  
Behr and Platt, 2011). Alternatively, it might be attributed to external stress loading from seismic faulting in the overlying  
upper crust (i.e. seismic loading; e.g. Küster and Stöckhert, 1998; Trepmann and Stöckhert, 2003; Trepmann et al., 2017;  
460 Trepmann and Seybold, 2019). The two models are not mutually exclusive and are both equally valid to explain our  
observations. We are unable to discern between the two options, due to the limitations of our P-T estimates. However, the  
presence of pseudotachylytes along the BFZ045 fault core indicates that the fault was capable of generating earthquakes, and  
the seismogenic behaviour of other faults in the Olkiluoto basement has been previously discussed (Marchesini et al., 2019  
and references therein). This suggests that transient seismic loading might have triggered the localised increase in creep rate  
465 during mylonitisation of BFZ045.

## Conclusions

This study shows that deformation microstructures can be used to evaluate the stress history of a shear zone deforming across  
the brittle-ductile transition in the continental crust, and to reconstruct the cyclical brittle-ductile deformation history of fault  
zones. The fault zone BFZ045 exploited a mylonitic precursor in the Paleoproterozoic basement in SW Finland and records  
470 transient brittle deformation in the form of syn-kinematic quartz veins emplaced during ongoing mylonitic creep at  $\sim 450^\circ \text{C}$   
and 3.5 kbar, in response to transiently high fluid pressure. Mylonitic deformation continued after vein emplacement, as  
evidenced by the dynamic recrystallization of the vein quartz. Mylonitic creep occurred under progressively increasing  
differential stress towards the shear zone centre in an overall narrowing shear zone. Mylonitic deformation at the shear zone  
centre records peak stress conditions of around 120 MPa, and was followed by brittle deformation that generated cataclasites  
475 and minor pseudotachylytes in the fault core.

The constraints derived from microstructural analysis shaped the proposed conceptual model of the evolution of BFZ045 slip  
behaviour, which highlights the important role of transiently sub-lithostatic fluid pressure in triggering vein emplacement  
during ongoing mylonitic deformation, as well as of the progressive increase in stress and strain rate during viscous creep  
towards peak conditions reached at the BDTZ in the Fennoscandian Shield. This study shows that microstructural studies  
480 leading to the acquisition of independent constraints offer the potential to reconstruct in detail the evolutionary history of fault

zones that experienced a transition in deformation mode at the BDTZ. In addition to deriving a conceptual model of varying fault slip behaviours at the BDTZ, the methods and the results of this work complement and expand thorough site characterization studies of deep geological disposal facilities.

## 485 **Acknowledgements**

This study has been funded by Posiva Oy (project no. 2104442) and by the University of Plymouth. Stephen F. Cox, Michael Stipp and Francesco Giuntoli are thanked for fruitful discussions in the course of the project. Glenn Harper is thanked for support during SEM and EBSD analysis at the Electron Microscopy Centre of the University of Plymouth. Andrea Risplendente is thanked for support with EMPA analysis at the University of Milan. Many thanks go to Fabrizio Nestola for  
490 giving us access to the microRaman spectrometer of the University of Padova. Danilo Bersani is thanked for support with Raman analysis at the University of Parma. Alfredo Camacho and Georg Dresen are thanked for their reviews, which helped us prepare a much improved version of the manuscript.

## **References**

- Aaltonen, I., Lahti, M., Engström, J., Mattila, J., Paananen, M., Paulamäki, S., Gehör, S., Kärki, A., Ahokas, T., Torvela, T.  
495 and Front, K.: Geological Model of the Olkiluoto Site. Version 2.0., Posiva Working Report 2010-70, Posiva Oy, Eurajoki, 2010.
- Aaltonen, I., Engström, J., Gehör, S., Kosunen, P. and Kärki, A.: Geology of Olkiluoto, Posiva Working Report 2016-16, Posiva Oy, Eurajoki, 2016.
- Aoya, M., Kouketsu, Y., Endo, S., Shimizu, H., Mizukami, T., Nakamura, D. and Wallis, S.: Extending the applicability of  
500 the Raman carbonaceous-material geothermometer using data from contact metamorphic rocks, *Journal of Metamorphic Geology*, 28(9), 895–914, doi:10.1111/j.1525-1314.2010.00896.x, 2010.
- Behr, W. M. and Platt, J. P.: A naturally constrained stress profile through the middle crust in an extensional terrane, *Earth and Planetary Science Letters*, 303(3), 181–192, doi:10.1016/j.epsl.2010.11.044, 2011.
- Bestmann, M. and Prior, D. J.: Intragranular dynamic recrystallization in naturally deformed calcite marble: diffusion  
505 accommodated grain boundary sliding as a result of subgrain rotation recrystallization, *Journal of Structural Geology*, 25(10), 1597–1613, doi:10.1016/S0191-8141(03)00006-3, 2003.
- Beyssac, O., Goffé, B., Chopin, C. and Rouzaud, J. N.: Raman spectra of carbonaceous material in metasediments: a new geothermometer, *Journal of Metamorphic Geology*, 20(9), 859–871, doi:10.1046/j.1525-1314.2002.00408.x, 2002.
- Bons, P. D., Elburg, M. A. and Gomez-Rivas, E.: A review of the formation of tectonic veins and their microstructures, *Journal*  
510 *of Structural Geology*, 43, 33–62, doi:10.1016/j.jsg.2012.07.005, 2012.

- Ceccato, A., Menegon, L., Pennacchioni, G. and Morales, L. F. G.: Myrmekite and strain weakening in granitoid mylonites, *Solid Earth*, 9(6), 1399–1419, doi:<https://doi.org/10.5194/se-9-1399-2018>, 2018.
- Cox, S. F.: Faulting processes at high fluid pressures: An example of fault valve behavior from the Wattle Gully Fault, Victoria, Australia, *Journal of Geophysical Research: Solid Earth*, 100(B7), 12841–12859, doi:10.1029/95JB00915, 1995.
- 515 Cox, S. F.: The application of failure mode diagrams for exploring the roles of fluid pressure and stress states in controlling styles of fracture-controlled permeability enhancement in faults and shear zones, *Geofluids*, doi:10.1111/j.1468-8123.2010.00281.x, 2010.
- Cross, A. J., Prior, D. J., Stipp, M. and Kidder, S.: The recrystallized grain size piezometer for quartz: An EBSD-based calibration: EBSD-Based Quartz Grain Size Piezometer, *Geophysical Research Letters*, 44(13), 6667–6674, doi:10.1002/2017GL073836, 2017.
- 520 Derez, T., Pennock, G., Drury, M. and Sintubin, M.: Low-temperature intracrystalline deformation microstructures in quartz, *Journal of Structural Geology*, 71, 3–23, doi:10.1016/j.jsg.2014.07.015, 2015.
- Eberlei, T., Habler, G., Grasemann, B. and Abart, R.: Upper–greenschist facies intragrain deformation of albite in mylonitic meta–pegmatite and the influence of crystallographic anisotropy on microstructure formation, *Journal of Structural Geology*, 69, 47–58, doi:10.1016/j.jsg.2014.10.001, 2014.
- 525 Fossen, H. and Cavalcante, G. C. G.: Shear zones – A review, *Earth-Science Reviews*, 171, 434–455, doi:10.1016/j.earscirev.2017.05.002, 2017.
- Freeman, J. J., Wang, A., Kuebler, K. E., Jolliff, B. L. and Haskin, L. A.: Characterization of natural feldspars by Raman spectroscopy for future planetary exploration, *The Canadian Mineralogist*, 46(6), 1477–1500, 2008.
- 530 Fusseis, F. and Handy, M. R.: Micromechanisms of shear zone propagation at the brittle–viscous transition, *Journal of Structural Geology*, 30(10), 1242–1253, doi:10.1016/j.jsg.2008.06.005, 2008.
- Giuntoli, F., Menegon, L. and Warren, C. J.: Replacement reactions and deformation by dissolution and precipitation processes in amphibolites, *Journal of Metamorphic Geology*, 36(9), 1263–1286, doi:10.1111/jmg.12445, 2018.
- 535 Handy, M. R., Hirth, G. and Hovius, N.: Continental Fault Structure and Rheology from the Frictional-to-Viscous Transition Downward, in *Tectonic Faults: Agents of Change on a Dynamic Earth*, MITP. [online] Available from: <https://ieeexplore.ieee.org/document/6285583> (Accessed 13 July 2019), 2007.
- Hey, M. H.: A new review of the chlorites, *Mineral. Mag. J. M. Soc.*, 224, 277–292, <https://doi.org/10.1180/minmag.1954.030.224.01>, 1954.
- 540 Hentschel, F., Trepmann, C. A. and Janots, E.: Deformation of feldspar at greenschist facies conditions – the record of mylonitic pegmatites from the Pfunderer Mountains, Eastern Alps, *Solid Earth*, 10(1), 95–116, doi:<https://doi.org/10.5194/se-10-95-2019>, 2019.
- Hirth, G. and Beeler, N. M.: The role of fluid pressure on frictional behavior at the base of the seismogenic zone, *Geology*, 43(3), 223–226, doi:10.1130/G36361.1, 2015.

- 545 Hirth, G. and Tullis, J.: Dislocation creep regimes in quartz aggregates, *Journal of Structural Geology*, 14(2), 145–159, doi:10.1016/0191-8141(92)90053-Y, 1992.
- Hirth, G. and Tullis, J.: The brittle-plastic transition in experimentally deformed quartz aggregates, *Journal of Geophysical Research: Solid Earth*, 99(B6), 11731–11747, doi:10.1029/93JB02873, 1994.
- Hirth, G., Teysier, C. and Dunlap, J. W.: An evaluation of quartzite flow laws based on comparisons between experimentally and naturally deformed rocks, *Int J Earth Sci*, 90(1), 77–87, doi:10.1007/s005310000152, 2001.
- 550 Kärki, A. and Paulamäki, S.: *Petrology of Olkiluoto*, 2006.
- Kidder, S., Hirth, G., Avouac, J.-P. and Behr, W.: The influence of stress history on the grain size and microstructure of experimentally deformed quartzite, *Journal of Structural Geology*, 83, 194–206, doi:10.1016/j.jsg.2015.12.004, 2016.
- Kirilova, M., Toy, V., Rooney, J. S., Giorgetti, C., Gordon, K. C., Collettini, C. and Takeshita, T.: Structural disorder of graphite and implications for graphite thermometry., *Solid Earth*, 9(1), 2018.
- 555 Kjöll, H. J., Viola, G., Menegon, L. and Sørensen, B. E.: Brittle–viscous deformation of vein quartz under fluid-rich lower greenschist facies conditions, *Solid Earth*, 6(2), 681–699, doi:https://doi.org/10.5194/se-6-681-2015, 2015.
- Kohlstedt, D. L., Evans, B. and Mackwell, S. J.: Strength of the lithosphere: Constraints imposed by laboratory experiments, *Journal of Geophysical Research: Solid Earth*, 100(B9), 17587–17602, doi:10.1029/95JB01460, 1995.
- 560 Kouketsu, Y., Mizukami, T., Mori, H., Endo, S., Aoya, M., Hara, H., Nakamura, D. and Wallis, S.: A new approach to develop the Raman carbonaceous material geothermometer for low-grade metamorphism using peak width, *Island Arc*, 23(1), 33–50, doi:10.1111/iar.12057, 2014.
- Lafuente, B., Downs, R. T., Yang, H. and Stone, N.: The power of databases: the RRUFF project, in *Highlights in mineralogical crystallography*, pp. 1–29, Walter de Gruyter GmbH., 2016.
- 565 Lahtinen, R.: *Crustal evolution of the Svecofennian and Karelian domains during 2.1 - 1.79 Ga: with special emphasis on the geochemistry and origin of 1.93 - 1.91 Ga gneissic tonalites and associated supracrustal rocks in the Rautalampi area, central Finland ; with 7 tables*, Geologian Tutkimuskeskus, Espoo., 1994.
- Lahtinen, R., Korja, A. and Nironen, M.: Chapter 11 Paleoproterozoic tectonic evolution, in *Developments in Precambrian Geology*, vol. 14, edited by M. Lehtinen, P. A. Nurmi, and O. T. Rämö, pp. 481–531, Elsevier., 2005.
- 570 Lanari, P., Wagner, T. and Vidal, O.: A thermodynamic model for di-trioctahedral chlorite from experimental and natural data in the system MgO–FeO–Al<sub>2</sub>O<sub>3</sub>–SiO<sub>2</sub>–H<sub>2</sub>O: applications to P–T sections and geothermometry, *Contributions to Mineralogy and Petrology*, 167(2), 968, 2014.
- Mänttari, I., Talikka, M., Paulamäki, S. and Mattila, J.: U-Pb ages for tonalitic gneiss, pegmatitic granite, and diabase dyke, Olkiluoto study site, Eurajoki, SW Finland, Posiva Oy., 2006.
- 575 Mänttari, I., Engstroem, J., Lahaye, Y. and Pere, T.: U-Pb ages for PGR dykes, KFP, and adjacent older leucosomic PGRs from ONKALO underground research facility, Olkiluoto, Eurajoki, SW Finland, Posiva Oy., 2010.

- Marchesini, B., Garofalo, P. S., Menegon, L., Mattila, J. and Viola, G.: Fluid-mediated, brittle–ductile deformation at seismogenic depth – Part 1: Fluid record and deformation history of fault veins in a nuclear waste repository (Olkiluoto Island, Finland), *Solid Earth*, 10(3), 809–838, doi:https://doi.org/10.5194/se-10-809-2019, 2019.
- 580 Massonne, H.-J. and Schreyer, W.: Phengite geobarometry based on the limiting assemblage with K-feldspar, phlogopite, and quartz, *Contributions to Mineralogy and Petrology*, 96(2), 212–224, 1987.
- Mattila, J. and Viola, G.: New constraints on 1.7 Gyr of brittle tectonic evolution in southwestern Finland derived from a structural study at the site of a potential nuclear waste repository (Olkiluoto Island), *Journal of Structural Geology*, 67, 50–74, 2014.
- 585 Melosh, B. L., Rowe, C. D., Gerbi, C., Smit, L. and Macey, P.: Seismic cycle feedbacks in a mid-crustal shear zone, *Journal of Structural Geology*, 112, 95–111, doi:10.1016/j.jsg.2018.04.004, 2018.
- Menegon, L., Pennacchioni, G. and Spiess, R.: Dissolution-precipitation creep of K-feldspar in mid-crustal granite mylonites, *Journal of Structural Geology*, 30(5), 565–579, doi:10.1016/j.jsg.2008.02.001, 2008.
- Nguyen, P. T., Harris, L. B., Powell, C. M. and Cox, S. F.: Fault-valve behaviour in optimally oriented shear zones: an example  
590 at the Revenge gold mine, Kambalda, Western Australia, *Journal of Structural Geology*, 20(12), 1625–1640, doi:10.1016/S0191-8141(98)00054-6, 1998.
- Nironen, M.: The Svecofennian Orogen: a tectonic model, *Precambrian Research*, 86(1–2), 21–44, 1997.
- Nordbäck, N. and Mattila, J.: *Brittle Fault Systems of the ONKALO Underground Research Facility*, 2018.
- Pajunen, M., Airo, M., Elminen, T., Mänttari, I., Niemelä, R., Vaarma, M., Wasenius, P. and Wennerström, M.: Tectonic  
595 evolution of the Svecofennian crust in southern Finland, *Geological Survey of Finland, Special Paper*, 47, 15–160, 2008.
- Pennacchioni, G. and Mancktelow, N. S.: Nucleation and initial growth of a shear zone network within compositionally and structurally heterogeneous granitoids under amphibolite facies conditions, *Journal of Structural Geology*, 29(11), 1757–1780, doi:10.1016/j.jsg.2007.06.002, 2007.
- Pere, T.: Fault-related local phenomena in the bedrock of Olkiluoto with particular reference to fault zone OL-BFZ100, Posiva  
600 Oy., 2009.
- Saintot, A., Stephens, M., Viola, G. and Nordgulen, Ø.: Brittle tectonic evolution and paleostress field reconstruction in the southwestern part of the Fennoscandian Shield, Forsmark, Sweden, *Tectonics*, 30(4), 2011.
- Scholz, C. H.: Earthquakes and friction laws, *Nature*, 391(6662), 37, doi:10.1038/34097, 1998.
- Sibson, R. H.: Earthquake faulting as a structural process, *Journal of Structural Geology*, 11(1), 1–14, doi:10.1016/0191-  
605 8141(89)90032-1, 1989.
- Sibson, R. H.: Conditions for fault-valve behaviour, *Geological Society, London, Special Publications*, 54(1), 15–28, doi:10.1144/GSL.SP.1990.054.01.02, 1990.
- Sibson, R. H.: Implications of fault-valve behaviour for rupture nucleation and recurrence, *Tectonophysics*, 211(1), 283–293, doi:10.1016/0040-1951(92)90065-E, 1992.

- 610 Sibson, R. H.: Load-strengthening versus load-weakening faulting, *Journal of Structural Geology*, 15(2), 123–128, doi:10.1016/0191-8141(93)90090-W, 1993.
- Sibson, R. H. and Rowland, J. V.: Stress, fluid pressure and structural permeability in seismogenic crust, North Island, New Zealand, *Geophys J Int*, 154(2), 584–594, doi:10.1046/j.1365-246X.2003.01965.x, 2003.
- Skyttä, P. and Torvela, T.: Brittle reactivation of ductile precursor structures: The role of incomplete structural transposition  
615 at a nuclear waste disposal site, Olkiluoto, Finland, *Journal of Structural Geology*, 116, 253–259, 2018.
- Stipp, M. and Kunze, K.: Dynamic recrystallization near the brittle-plastic transition in naturally and experimentally deformed quartz aggregates, *Tectonophysics*, 448(1), 77–97, doi:10.1016/j.tecto.2007.11.041, 2008.
- Stipp, M. and Tullis, J.: The recrystallized grain size piezometer for quartz, *Geophysical Research Letters*, 30(21), doi:10.1029/2003GL018444, 2003.
- 620 Stipp, M., Stünitz, H., Heilbronner, R. and Schmid, S. M.: The eastern Tonale fault zone: a ‘natural laboratory’ for crystal plastic deformation of quartz over a temperature range from 250 to 700°C, *Journal of Structural Geology*, 24(12), 1861–1884, doi:10.1016/S0191-8141(02)00035-4, 2002.
- Stipp, M., Tullis, J., Scherwath, M. and Behrmann, J. H.: A new perspective on paleopiezometry: Dynamically recrystallized grain size distributions indicate mechanism changes, *Geology*, 38(8), 759–762, doi:10.1130/G31162.1, 2010.
- 625 Stöckhert, B., Brix, M. R., Kleinschrodt, R., Hurford, A. J. and Wirth, R.: Thermochronometry and microstructures of quartz—a comparison with experimental flow laws and predictions on the temperature of the brittle–plastic transition, *Journal of Structural Geology*, 21(3), 351–369, doi:10.1016/S0191-8141(98)00114-X, 1999.
- Stünitz, H. and Gerald, J. D. F.: Deformation of granitoids at low metamorphic grade. II: Granular flow in albite-rich mylonites, *Tectonophysics*, 221(3), 299–324, doi:10.1016/0040-1951(93)90164-F, 1993.
- 630 Stünitz, H., Fitz Gerald, J. D. and Tullis, J.: Dislocation generation, slip systems, and dynamic recrystallization in experimentally deformed plagioclase single crystals, *Tectonophysics*, 372(3), 215–233, doi:10.1016/S0040-1951(03)00241-5, 2003.
- Torgersen, E., Viola, G., Zwingmann, H. and Harris, C.: Structural and temporal evolution of a reactivated brittle–ductile fault – Part II: Timing of fault initiation and reactivation by K–Ar dating of synkinematic illite/muscovite, *Earth and Planetary  
635 Science Letters*, 410, 212–224, doi:10.1016/j.epsl.2014.09.051, 2015.
- Torvela, T. and Ehlers, C.: From ductile to brittle deformation: structural development and strain distribution along a crustal-scale shear zone in SW Finland, *International Journal of Earth Sciences*, 99(5), 1133–1152, 2010.
- Trepmann, C. A. and Seybold, L.: Deformation at low and high stress-loading rates, *Geoscience Frontiers*, 10(1), 43–54, doi:10.1016/j.gsf.2018.05.002, 2019.
- 640 Trepmann, C. A., Stöckhert, B., Dorner, D., Moghadam, R. H., Küster, M. and Röller, K.: Simulating coseismic deformation of quartz in the middle crust and fabric evolution during postseismic stress relaxation — An experimental study, *Tectonophysics*, 442(1), 83–104, doi:10.1016/j.tecto.2007.05.005, 2007.
- Tuisku, P. and Kärki, A.: Metamorphic petrology of Olkiluoto, Posiva Oy., 2010.

- 645 Viola, G., Mancktelow, N. S. and Miller, J. A.: Cyclic frictional-viscous slip oscillations along the base of an advancing nappe complex: Insights into brittle-ductile nappe emplacement mechanisms from the Naukluft Nappe Complex, central Namibia, *Tectonics*, 25(3), doi:10.1029/2005TC001939, 2006.
- Viola, G., Venvik Ganerød, G. and Wahlgren, C.-H.: Unraveling 1.5 Ga of brittle deformation history in the Laxemar-Simpevarp area, southeast Sweden: A contribution to the Swedish site investigation study for the disposal of highly radioactive nuclear waste, *Tectonics*, 28(5), doi:10.1029/2009TC002461, 2009.
- 650 Wehrens, P., Berger, A., Peters, M., Spillmann, T. and Herwegh, M.: Deformation at the frictional-viscous transition: Evidence for cycles of fluid-assisted embrittlement and ductile deformation in the granitoid crust, *Tectonophysics*, 693, 66–84, doi:10.1016/j.tecto.2016.10.022, 2016.
- Wright, S. I., Nowell, M. M. and Field, D. P.: A Review of Strain Analysis Using Electron Backscatter Diffraction, *Microscopy and Microanalysis*, 17(3), 316–329, doi:10.1017/S1431927611000055, 2011.
- 655 Yardley, B. and Baumgartner, L.: Fluid processes in deep crustal fault zones, *Tectonic Faults—Agents of Change on a Dynamic Earth*, 295–318, 2007.

Sample name	PH28_7a	PH28_7b-1	PH28_5a	PH28_5b-1	PH28_5c	PH28_5d	PH28_9e	PH28_10-6
N. measurement	n=5	n=5	n=15	n=18	n=12	n=16	n=18	n=10
Chemical composition (wt%)								
SiO2	48.280	48.080	48.210	48.410	48.330	48.770	47.970	47.660
TiO2	0.093	0.098	0.124	0.099	0.104	0.074	0.043	0.136
Al2O3	33.910	33.900	33.860	34.520	34.590	34.680	35.690	34.910
Cr2O3	0.011	0.000	0.000	0.017	0.000	0.000	0.037	0.000
Fe2O3	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
FeO	2.540	2.430	2.710	2.080	2.060	1.940	3.010	2.730
MnO	0.060	0.001	0.052	0.000	0.000	0.005	0.078	0.028
MgO	1.563	1.383	1.446	1.411	1.384	1.360	0.546	1.415
CaO	0.017	0.005	0.000	0.032	0.010	0.033	0.015	0.054
Na2O	0.091	0.136	0.132	0.163	0.138	0.116	0.126	0.201
K2O	11.080	11.120	11.090	11.180	11.030	11.230	9.430	10.690
NiO	0.000	0.036	0.006	0.000	0.000	0.013	0.000	0.036
Total	97.645	97.189	97.630	97.912	97.646	98.221	96.944	97.859

Structural formula on the basis of 11 O

SiO2	3.15	3.16	3.15	3.15	3.15	3.16	3.13	3.11
TiO2	0.00	0.00	0.01	0.00	0.01	0.00	0.00	0.01
Al2O3	2.61	2.62	2.61	2.65	2.65	2.65	2.74	2.68
Cr2O3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Fe2O3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
FeO	0.14	0.13	0.15	0.11	0.11	0.11	0.16	0.15
MnO	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
MgO	0.15	0.14	0.14	0.14	0.13	0.13	0.05	0.14
CaO	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Na2O	0.01	0.02	0.02	0.02	0.02	0.01	0.02	0.03
K2O	0.92	0.93	0.93	0.93	0.92	0.93	0.78	0.89
NiO	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
H2O	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

655

**Table 1. Representative compositions of muscovite from the BFZ045 mylonite**


Fault rock	Mylonite				Cataclasite				Veins			
Sample name	PH16-2_2	PH16-2_5	PH28-10_3	PH28-10_15	PH28-9_5	PH28-9_7	PH28-9_10	PH28-9_19	PH28-6_6	PH28-6_9	PH28-6_11	PH28-6_13
Chemical composition (wt%)												
SiO <sub>2</sub>	25.040	25.820	24.790	24.790	23.600	23.800	23.290	24.170	25.470	23.930	23.290	24.070
TiO <sub>2</sub>	0.034	0.034	0.078	0.060	0.031	0.115	0.050	0.010	0.059	0.049	0.004	0.000
Al <sub>2</sub> O <sub>3</sub>	21.600	22.290	21.890	22.140	22.150	22.420	22.440	21.990	20.620	21.860	22.690	22.120
FeO	28.700	29.930	35.830	34.600	38.530	39.650	39.150	34.360	37.260	37.940	39.670	37.650
MnO	0.173	0.349	0.918	0.858	0.846	0.810	0.793	0.675	1.201	0.596	0.937	1.430
MgO	12.010	11.740	7.570	8.430	4.810	3.780	3.900	7.730	6.980	5.810	3.910	5.730
CaO	0.047	0.061	0.121	0.056	0.008	0.001	0.003	0.000	0.021	0.006	0.005	0.044
Na <sub>2</sub> O	0.014	0.055	0.010	0.000	0.000	0.046	0.036	0.011	0.000	0.000	0.015	0.000
K <sub>2</sub> O	0.064	0.047	0.020	0.040	0.021	0.000	0.080	0.087	0.008	0.000	0.005	0.008
Total	87.682	90.364	91.236	90.973	90.049	90.655	89.742	89.033	91.620	90.190	90.525	91.081
Structural formulae on the basis of 14 O												
Si	2.687	2.695	2.658	2.647	2.610	2.624	2.595	2.641	2.741	2.628	2.577	2.621
Ti	0.003	0.003	0.006	0.005	0.003	0.010	0.004	0.001	0.005	0.004	0.000	0.000
Al	2.731	2.742	2.766	2.786	2.887	2.913	2.947	2.832	2.615	2.829	2.959	2.839
Fe(3+)	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Fe(2+)	2.575	2.613	3.213	3.089	3.563	3.656	3.648	3.140	3.353	3.485	3.671	3.429
Mn	0.016	0.031	0.083	0.078	0.079	0.076	0.075	0.063	0.110	0.055	0.088	0.132
Mg	1.921	1.827	1.210	1.342	0.793	0.621	0.648	1.259	1.120	0.951	0.645	0.930
Ca	0.005	0.007	0.014	0.006	0.001	0.000	0.000	0.000	0.003	0.001	0.001	0.005
Na	0.003	0.011	0.002	0.000	0.000	0.010	0.008	0.002	0.000	0.000	0.003	0.000
K	0.009	0.006	0.003	0.005	0.003	0.000	0.011	0.012	0.001	0.000	0.001	0.001
X <sub>Fe</sub> = Fe/(Fe+Mg)	0.57	0.59	0.73	0.70	0.82	0.85	0.85	0.71	0.75	0.79	0.85	0.79

**Table 2. Representative compositions of chlorite from different domains of BFZ045.**

Parameter	Value	Reference/source	Notes
vertical stress $\sigma_2$	350 MPa	P estimate	lithostatic load during deformation for Andersonian type fault in a strike-slip regime
strain rate	stage 1 (50MPa) $4.3 \times 10^{-13} \text{ s}^{-1}$  stage 3 (80MPa) $1.1 \times 10^{-12} \text{ s}^{-1}$  stage 3 (120 MPa) $6.9 \times 10^{-12} \text{ s}^{-1}$	Hirth et al. (2001) Wither's fugacity calculator (Pitzer and Sterner, 1994)	$\dot{\epsilon} = A\sigma^n f_{\text{H}_2\text{O}}^m \exp(-Q/RT)$ A an empirical constant, $\sigma$ the differential stress, n the stress exponent, $f_{\text{H}_2\text{O}}$ the water fugacity (calculated at 350 MPa using Wither's fugacity calculator), m the water fugacity exponent, Q the activation energy, R the gas constant, and T the temperature
fluid pressure conditions	> 210 MPa	Marchesini et al. (2019)	
friction coefficient ( $\mu$ )	0.6	Sibson (1985)	common value for granitoids
cohesive strength	26 MPa	Aaltonen et al.(2010)	data from tensile strength measurements of granitic gneisses
tensile strength	13 MPa	Aaltonen et al.(2010)	


660

**Table 3. Parameters used for  $\lambda$ - $\sigma$  failure mode diagrams**



665 **Figure 1. Geological setting of Olkiluoto, SW Finland (inset on top-right). (a) Schematic geological and structural map, showing surface intersection of modelled brittle fault zone (BFZ) and ductile deformation zone, modified from Aaltonen et al. (2016) and Skytta and Torvela (2018). Coordinates for zone 34N in UTM coordinate system. The white line indicates the location of the**


underground Onkalo facility. A-B is the trace of the cross section shown in (b). (b) East-west cross section across the underground infrastructure, with the tunnel shown as white line. Sub vertical fault BFZ045 described in this study and its conjugate BFZ300 are shown as orange lines.



670

675


**Figure 2. BFZ045 fault geometry.** (a) Schematic representation of fault architecture from core logs, vertical axis not to scale. Grey rectangles locate the studied drill cores PH28 and PH16, black squares show sample location. The red rectangle indicates the samples shown in (d). (b) Stereoplot of BFZ045 fault orientation and mylonitic foliation observed at different drill hole intersection in the Onkalo facility (Aaltonen et al., 2016). (c) Core sample along PH28 drill core within the fault core unit. The core sample exposes the mylonitic foliation, where the blue line indicates the stretching lineation, which is parallel to chlorite striae. (d) Samples from PH28 drill core representative of different domains of the fault: damaged coarse-grained host rock (I), and fault core with mylonites and chlorite rich cataclasites (II-III). Dashed lines outline the area of petrographic thin sections. White arrow points to pseudotachylyte injection veins.




680

Figure 3. Polarized light microscope images of the damaged host rock (a), and representative quartz microstructures (b-c). Cross polarized light. (a) Host rock showing the original magmatic textures and mineral assemblage. White arrow shows a chlorite-rich shear band oriented at a low angle to the mylonitic foliation of BFZ045. (b) Quartz with wide extinction bands (WEBs) and undulatory extinction. WEBs are bounded by sets of fluid inclusions trails (FI trail, dashed lines). Intercrystalline deformation bands and well-developed FI trails developed sub-parallel to the mylonitic foliation. (c) Polygonal recrystallized quartz grains, with small grain size (~20 μm), forming bands oriented sub-parallel to the mylonitic foliation. The white arrow shows sutured grain boundaries between magmatic quartz grains, indicative of bulging. Foliation trace is projected as a white line on to the images from the adjacent foliated host rock.

685



690 Figure 4. Microstructure of the BFZ045 mylonite. Cross-polarized light. (a) The mylonitic foliation is defined by the alternation of  
 quartz-rich and mica-rich bands. Fractured porphyroclasts of feldspars are preferentially located in mica-rich domains. (b) Quartz  
 ribbons in the shear zone boundary are stretched along the foliation and show typical core-and-mantle microstructure, with  
 recrystallization localized at the grain boundaries. Ribbons contain also well-developed WEBS. (c) Completely recrystallized quartz  
 ribbon. The recrystallized grains show a shape preferred orientation indicating a sinistral sense of shear. Thin muscovite (Ms)  
 695 define the mylonitic foliation, together with the elongated and recrystallized quartz domains. (d) Quartz veins along the foliation,  
 filling a mode I fracture at a distance of ~ 1cm from the cataclasite fault core. Quartz grains elongation in the veins is normal to the  
 vein wall and to the foliation. (e) High magnification view of the quartz vein. Bulges along the grain boundaries, and subgrains  
 within the grains are visible.



705 **Figure 5.** . Microstructure of the BFZ045 brittle fault core. Cross-polarized light (a-c), plane-polarized light (a), and scanning electron microscope backscatter electron images (SEM-BSE) (d-e). (a) Cataclasite with characteristic pseudotachylyte injection veins (arrow). The cataclasite matrix is enriched in chlorite and Ti-oxides. (b) Contact between the mylonite (upper left corner) and the cataclasite. Quartz forms almost entirely recrystallized polycrystalline ribbons. (c) Detail of a sub-angular polycrystalline clast of quartz in the cataclasite. The trace of the mylonitic foliation is still visible in the clast, and is only slightly rotated with respect to the trace of the foliation in the mylonite. The surrounding matrix is a fine mixture of white mica and plagioclase. (d) Pseudotachylyte injection vein. Rounded quartz clast (dark grey) and Ti-oxides (white) are surrounded by a chlorite and mica rich ultrafine-grained

matrix. (e) Quartz + chlorite aggregates filling fractures within and strain shadows around plagioclase porphyroclast in the mylonite. The clast is slightly rotated within the protocataclasite. The white line in the top right corner indicates the trace of the mylonitic foliation.






Figure 6. EBSD data of quartz from the mylonitic fault core. In all the EBSD maps, black lines correspond to high-angle boundaries (misorientation > 10°), white lines to low-angle boundaries (misorientation between 2° and 10°), and red lines to Dauphiné twin boundaries (misorientation of 60° around the c-axis). (a) Quartz ribbons and intracrystalline bands of recrystallized grains (Fig. 5a). Cross polarized light. The box locates the EBSD map shown in (b-d, g). (b-c) Grain size map (diameter of the equivalent circle, μm), the higher magnification in (c) highlights the presence of subgrains in the large relict quartz grains and in the recrystallized grains. (d) Grain orientation spread (GOS) for each grain, coloured relative to the GOS threshold (black line) between recrystallized (blue) and relict (red) grains. (e) Histogram of grains size distribution of grains in the intercrystalline bands. (f) Histogram of subgrains

720 size distribution of subgrains in the relict quartz from the intercrystalline bands (light red in (d)). (g) Grain Orientation Distribution  
 725 maps (GROD) was used to estimate visually the subgrains size in the quartz ribbon.




725 Figure 7. EBSD data of quartz from a recrystallized ribbon in the mylonite. (a) Polycrystalline ribbons of recrystallized quartz  
 730 grains elongated parallel to the mylonitic foliation (Fig. 5b). Cross polarized light. The box indicates the EBSD maps shown in (b-  
 d). Colour coding of the boundaries like in Figure 6. (b, c) Grain size map (diameter of the equivalent circle, μm) and detail (c)  
 showing that the larger grains contain subgrains of the same size as the surrounding finer grains. (d) GOS map showing that the  
 GOS values are mostly under the threshold, indicative of high degree of recrystallization. (e) Histogram of the grain size distribution  
 for recrystallized and relict grains. (f) Histogram of the subgrain size distribution in the relict quartz grains identified in (d) and  
 (e). (g) Pole figure of the c-axis orientation of recrystallized and relict grains, colour coded like the GOS map in (d). Equal area, lower  
 hemisphere projection.




735 **Figure 8.** EBSD data of quartz from a foliation-parallel vein in the mylonite near the contact to the cataclasite. Colour coding of the boundaries like in Figure 6. (a) Quartz vein along the foliation. Quartz in the mylonite show a strong SPO consistent with the sinistral sense of shear of BFZ045. Cross polarized light. (b) Grain size map (diameter of the equivalent circle,  $\mu\text{m}$ ). (c) Details of previous maps. Bulges and subgrains (white arrows) of similar size of the bulges are evident within the vein quartz. (d) GOS map of quartz in the vein and of the surrounding mylonite. The GOS threshold value of  $1.94^\circ$  separates relict grains (red) from recrystallized grains (blue). Grey stars indicate relict grains plotted in (f) for subgrain size estimates. (e) Histogram of the grain size distribution in the mylonite, with relict and recrystallized grains separated with the GOS method. (f) Histogram of the subgrain size distribution (red) in relict quartz of the mylonite (grey). (g) Histogram of the recrystallized grain size (red) and subgrains size (blue) distribution in the vein. (h-i) Pole figures of the c-axis orientation of the recrystallized and relict quartz in the mylonite (h) and in the veins (i). Equal area, lower hemisphere projection.

740




745 **Figure 9.** EBSD data of quartz from a clast in the cataclasite. Colour coding of the boundaries like in Figure 6. (a) The analysed quartz clast (Fig. 5f). The white rectangle locates the EBSD map shown in (b-d). (b-c) Grain size map (diameter of the equivalent circle,  $\mu\text{m}$ ). The map highlights the presence of subgrains in the coarser elongated quartz grains with size comparable to the surrounding finer quartz. (d) GOS map. (e) Histogram of the grain size distribution of the recrystallized (blue) and relict (red) grains. (f) Pole figure of the c-axis orientation of recrystallized and relict grains. Equal area, lower hemisphere projection, color coded as GOS map.




750 **Figure 10. Light microscopy and SEM backscattered electron (BSE) images of characteristic microstructural domains and mineral assemblages used for geothermobarometry estimates. (a) Graphite flakes in association with radiate chlorite in the host rock. Plane polar light. (b) Detail of graphite grains along the mylonitic foliation of BFZ045. Plane polar light. (c) Chlorite in pressure shadows around a K-feldspar porphyroblast in the mylonite. White mica and a fine grained recrystallized K-feldspar assemblage is common along the foliation. (d) Light microscope image of a microstructure similar to (c), cross polar light. (e-f) SEM BSE images of radiate chlorite used for chlorite thermometry in the cataclasite matrix (e) and in a vein cutting the mylonitic foliation (f). The trace of the mylonitic foliation in (f) is oriented ca. NW-SE.**

755




**Figure 11. Results of P-T estimates. (a) Carbonaceous material Raman geothermometer. Average R2 ratio (refer to text for explanation) for graphite rich mylonitic and host rock samples was measured to derive a T estimate using the method of Beyssac et al. (2002) and Aoya et al. (2010). (b) Estimated P of mylonitization using the Si-in-phengite barometer (Massonne and Schreyer, 1987) for the average T of 440° C obtained with the carbonaceous material Raman thermometry. Red line show the total spread of the Si values obtained. Black square show the (c) Chlorite compositional diagram based on Hey (1954). (d) Chlorite formation temperature estimated for mylonitic foliation, veins and cataclasite using the method of Lanari et al. (2014).**

760



765 **Figure 12. Schematic representation of the microstructural evolution of BFZ045. (a-d) Grey lines: traces of metamorphic foliation in the host rock. In the schematic fault scheme evolution, red displays the active deformation process, grey represents the inactive deformation processes, and orange suggests a transition in the deformation. (a) The development of mylonite was punctuated by the emplace-**  
 770 **ment of quartz vein (b). (c) Ductile deformation localised toward the centre of the mylonitic fault core in an overall narrowing shear zone, and was followed by formation of cataclasite, chlorite veins and pseudotachylyte (d). (e) Schematic summary of the quartz recrystallized grain size, differential stresses, and P-T conditions of deformation for BFZ045 derived in the present study, in relationship to the fault core geometry. Each sample is coloured differently to indicate the spatial position of the results described in section 4 of this paper.**



775 **Figure 13.** Conceptual model of the temporal and mechanical evolution of the BFZ045 fault zone (see text for more details). BFZ045 was characterized by (a) the development of mylonite creeping at ca.  $10^{-13} \text{ s}^{-1}$ , followed by (b) a transient increase in fluid pressure responsible for the emplacement of quartz veins. Ductile deformation then continued under increasing differential stress and strain rates (c).