

Concerning the paper entitled ‘Quantifying the buffering of oceanic oxygen isotopes at ancient midocean ridges’  
(se-2019-169)

June 29, 2020

Dear Editors Drs. Patrice Rey and Joachim Gottsmann,

I express my gratitude to you for considering and accepting my paper for publication in SE. According to Topical Editor’s comments, I made the model codes available to public and have assigned a DOI to the version used for this paper. Corresponding revision was made in Code availability section. Please see just below for the list of all relevant changes in the revised manuscript and further below for the marked-up manuscript.

List of all relevant changes

Place	Previous manuscript	Revised manuscript
Lines 376–378 on the page 13	The source codes of the model are available upon request to the author.	The source codes of the model are available on GitHub ( <a href="https://github.com/kanzakiy/HT-Oiso">https://github.com/kanzakiy/HT-Oiso</a> ) under the GNU General Public License v3.0. The specific version used of the model in this paper is tagged as `v0.1` and has been assigned a DOI: 10.5281/zenodo.3922317 ( <a href="https://doi.org/10.5281/zenodo.3922317">https://doi.org/10.5281/zenodo.3922317</a> ). A readme file on the web provides the instructions for executing the simulations.

Yours sincerely,

Yoshiki Kanzaki

Department of Earth and Planetary Sciences  
University of California – Riverside  
University Ave. 900, Riverside, CA 92521, USA  
(phone: +1 (814) 424 4332;  
e-mail: [kanzakiy@ucr.edu](mailto:kanzakiy@ucr.edu))

# Quantifying the buffering of oceanic oxygen isotopes at ancient midocean ridges

Yoshiki Kanzaki<sup>1</sup>

<sup>1</sup>Department of Earth and Planetary Sciences, University of California - Riverside, Riverside, CA 92521, USA

**Correspondence:** Yoshiki Kanzaki (kanzakiy@ucr.edu )

**Abstract.** To quantify the intensity of oceanic oxygen-isotope buffering through hydrothermal alteration of oceanic crust, a two-dimensional hydrothermal circulation model was coupled with a two-dimensional reactive transport model of oxygen isotopes. The coupled model calculates steady-state distributions of temperature, water flow and oxygen isotopes of solid rock and porewater given physicochemical conditions of oceanic crust alteration and seawater  $\delta^{18}\text{O}$ . Using the present-day seawater  $\delta^{18}\text{O}$  under plausible modern alteration conditions, the model yields  $\delta^{18}\text{O}$  profiles for solid rock and porewater and fluxes of heat, water and  $^{18}\text{O}$  that are consistent with modern observations, confirming the model's validity. The model was then run with different assumed seawater  $\delta^{18}\text{O}$  values to evaluate oxygen isotopic buffering at the midocean ridges. The buffering intensity shown by the model is significantly weaker than previously assumed and, consistently, calculated  $\delta^{18}\text{O}$  profiles of oceanic crust are relatively insensitive to seawater  $\delta^{18}\text{O}$ . These results are attributed to the fact that isotope exchange at shallow depths does not reach equilibrium due to the relatively low temperatures, and  $^{18}\text{O}$  supply via spreading solid rocks overwhelms that through water flow at deeper depths. Further model simulations under plausible alteration conditions during the Precambrian showed essentially the same results. Therefore,  $\delta^{18}\text{O}$  records of ophiolites that are invariant at different Earth's ages can be explained by the relative insensitivity of oceanic rocks to seawater  $\delta^{18}\text{O}$  and do not require constant seawater  $\delta^{18}\text{O}$  through time.

15 *Copyright statement.* Author(s) 2019. CC BY 4.0 License.

## 1 Introduction

Hydrothermal alteration of oceanic crust at midocean ridges works as the dominant source/sink of several elements/isotopes in the ocean (e.g., Wolery and Sleep, 1976; Elderfield and Schultz, 1996). Notably, oxygen isotopes have been considered to be primarily controlled by isotope exchange at midocean ridges. The observation of oceanic crustal  $\delta^{18}\text{O}$  has revealed that oceanic rocks gain and lose  $^{18}\text{O}$  through low- and high-temperature hydrothermal alteration at shallow and deeper depths, respectively. The close balance between the addition and removal of the heavy isotope, together with the huge oxygen supply from the mantle, has led to a hypothesis that the water-rock interactions at midocean ridges have buffered oceanic  $\delta^{18}\text{O}$  at the present-day value (0 ‰ relative to standard mean ocean water, SMOW) throughout Earth's history (e.g., Muehlenbachs

and Clayton, 1976; Gregory and Taylor, 1981; Holland, 1984; Muehlenbachs, 1998). Age-invariant  $\delta^{18}\text{O}$  records of ophiolites  
25 (ancient oceanic crust) have been argued to support the hypothesis (e.g., Holmden and Muehlenbachs, 1993).

In contrast to ophiolites, sedimentary rocks (e.g., carbonates, cherts, phosphorites and shales) have shown secular  $\delta^{18}\text{O}$   
increases with the Earth's age; sedimentary rocks in the Precambrian are depleted in  $^{18}\text{O}$  by as much as 10 ‰ compared  
to those in the modern (e.g., Shemesh et al., 1983; Karhu and Epstein, 1986; Bindeman et al., 2016), and the exception to  
this has rarely been reported (e.g., Blake et al., 2010). In theory, if constituting minerals of sedimentary rocks were formed  
30 in equilibrium with seawater and if the later diagenetic/metamorphic modification of  $\delta^{18}\text{O}$  was negligible,  $\delta^{18}\text{O}$  records of  
sedimentary rocks can be utilized to infer surface temperatures in the past, using temperature-dependent isotope fractionation  
factors and assuming the  $\delta^{18}\text{O}$  value of seawater from which the constituting minerals formed. Indeed, several authors have  
suggested hot climates in the Precambrian from  $^{18}\text{O}$ -depleted sedimentary records assuming the present-day seawater  $\delta^{18}\text{O}$   
(e.g.,  $70 \pm 15$  °C at 3.2–3.5 Ga and 50–60 °C through the later Precambrian; Knauth and Lowe, 2003; Knauth, 2005). Such high  
35 temperatures are apparently at odds with glacial records observed through the Precambrian (e.g., Catling and Kasting, 2017, Ch.  
11). If one excludes hot climates based on the glacial records, one must conclude that sedimentary  $\delta^{18}\text{O}$  records do not reflect  
seawater  $\delta^{18}\text{O}$  (e.g., instead reflecting diagenetic/metamorphic overprints) and/or that seawater  $\delta^{18}\text{O}$  has changed through time  
(e.g., Walker and Lohmann, 1989). The latter conclusion has recently been supported by Proterozoic-Phanerozoic marine iron  
(oxyhydr)oxides, whose  $\delta^{18}\text{O}$  shows a similar secular evolution to those in sedimentary rocks despite its weak temperature  
40 dependence (Galili et al., 2019). On the other hand, the secular change in seawater  $\delta^{18}\text{O}$  apparently conflicts with the result  
derived from ophiolites: an invariant seawater  $\delta^{18}\text{O}$  resulting from strong oceanic- $\delta^{18}\text{O}$  buffering at midocean ridges. Overall,  
one cannot rely on sedimentary  $\delta^{18}\text{O}$  to reconstruct ancient surface temperatures and discuss associated climate forcings unless  
 $\delta^{18}\text{O}$  records of sedimentary rocks and ophiolites can be explained at the same time (cf. Godderis and Veizer, 2000; V  rard  
and Veizer, 2019).

Several hypotheses have been put forward to reconcile apparently conflicting invariant and variant  $\delta^{18}\text{O}$  records of respective  
45 ophiolites and sedimentary rocks. Perry et al. (1978) pointed out that oceanic rocks in the Precambrian were more mafic than  
those in the modern and this could have resulted in more intense low-temperature alteration of oceanic crust, removing more  
 $^{18}\text{O}$  from seawater than today. Walker and Lohmann (1989) argued that the midocean ridges could have been above sea level  
in shallower Precambrian oceans. Resultant subaerial water-rock interactions at low temperatures then could have removed  
50 a large amount of  $^{18}\text{O}$  from the Precambrian oceans. Even if midocean ridges were below the sea level, lower pressures  
on the seafloor in the shallow oceans could have brought less water into the oceanic crust, and water-rock interactions at  
midocean ridges could have been dominated by those at low temperatures at shallower depths, again resulting in lower oceanic  
 $\delta^{18}\text{O}$  in the Precambrian (Kasting et al., 2006). Absence of biogenic sediment cover during the Precambrian could have also  
contributed to more significant low-temperature seafloor alteration and a lower seawater  $\delta^{18}\text{O}$  (e.g., Jaffr  s et al., 2007). The  
55 above arguments, however, give possible explanations only for low oceanic  $\delta^{18}\text{O}$  during the Precambrian; they do not explain  
age-invariant ophiolite  $\delta^{18}\text{O}$  records. Wallmann and colleagues have indicated that the invariant  $\delta^{18}\text{O}$  records of ophiolites can  
be explained by decoupling ancient oceanic crust from the contemporaneous seawater  $\delta^{18}\text{O}$  (e.g., Wallmann, 2001; Jaffr  s et

al., 2007). However, the simple box models considered by these authors cannot explain whether and/or how this decoupling could have been made possible (Jaffrés et al., 2007).

60 To resolve the issue, a process-based approach that simulates oxygen isotope behavior during oceanic crust alteration is indispensable. Lécuyer and Allemand (1999) have developed an oxygen-isotope exchange model which utilizes prescribed distributions of temperature and water/rock ratio within oceanic crust and an equation for half-closed systems (e.g., Gregory et al., 1989), and thus is not entirely process-based. Using this model, Lécuyer and Allemand (1999) concluded that oceanic  $\delta^{18}\text{O}$  cannot change from the present-day value by more than 2 ‰ due to the strong buffering exhibited by their modelled  
65 isotope-exchange at midocean ridges. This conclusion contrasts with that by Wallmann (2001), who adopted a box model to argue that the buffering must have been weak and that the oxygen-isotopic composition of the ocean has evolved through the Phanerozoic. These conflicting conclusions from different models emphasize that a mechanistic understanding of oxygen isotope behavior during oceanic crust alteration is lacking.

The present study develops a process-based model to simulate oxygen isotope behavior during hydrothermal alteration of  
70 oceanic crust and presents its application to the Precambrian. A two-dimensional (2D) hydrothermal circulation model presented in the literature is combined with a 2D reactive transport model of oxygen isotopes. The coupled model can thus yield 2D distributions of solid-rock and porewater  $\delta^{18}\text{O}$  reflecting alteration conditions including seawater  $\delta^{18}\text{O}$  and hydrothermal fluid circulation. After confirming the model's validity by comparing the model results that assume the present-day seawater  $\delta^{18}\text{O}$  with modern observations, we examine the intensity of oceanic- $\delta^{18}\text{O}$  buffering by hydrothermal alteration of oceanic  
75 crust at midocean ridges by changing seawater  $\delta^{18}\text{O}$ . The buffering quantification is then conducted under different physico-chemical conditions that could have been the case during the Precambrian, to give insights into how oceanic  $\delta^{18}\text{O}$  could have been affected by water-rock interactions at midocean ridges during the Earth's early eons.

## 2 Methods

### 2.1 Hydrothermal circulation model

80 Hydrothermal flow circulating around midocean ridges is simulated two-dimensionally based on conservations of energy, mass and momentum (Steeffel and Lasaga, 1994; Cherkaoui et al., 2003; Iyer et al., 2010). Assuming steady state, the mass conservation of fluid is represented by

$$\nabla \cdot \mathbf{q} = 0, \quad (1)$$

where  $\nabla$  is the vector differential operator ( $\nabla = (\partial/\partial x, \partial/\partial y)$ ) and  $\mathbf{q}$  is the mass flux vector ( $\text{kg m}^{-2} \text{yr}^{-1}$ ). Conservation of  
85 momentum is realized by adopting Darcy's law (e.g., Steeffel and Lasaga, 1994):

$$\mathbf{q} = -\frac{k}{\nu} (\nabla P - \rho_f \mathbf{g}), \quad (2)$$

where  $k$  is the permeability of oceanic bulk rock ( $\text{m}^2$ ),  $\nu$  is the kinematic viscosity of water ( $\text{m}^2 \text{yr}^{-1}$ ),  $P$  is the fluid pressure (Pa),  $\rho_f$  is the fluid density ( $\text{kg m}^{-3}$ ) and  $\mathbf{g}$  is the gravity vector given by  $\mathbf{g} = (0, -g)$  where  $g$  represents the acceleration by

gravity ( $\text{m yr}^{-2}$ ) (e.g., Steefel and Lasaga, 1994). The energy conservation is represented by

$$90 \quad \{\phi \rho_f c_p^f + (1 - \phi) \rho_m c_p^m\} \frac{\partial T}{\partial t} = \nabla \cdot (-\mathbf{q} c_p^f T + \kappa \nabla T), \quad (3)$$

where  $t$  is time (yr),  $\phi$  is the porosity,  $\rho_m$  is the density of oceanic rock ( $\text{kg m}^{-3}$ ),  $c_p^f$  and  $c_p^m$  are the specific heat capacity of water and oceanic rock, respectively ( $\text{J kg}^{-1} \text{K}^{-1}$ ),  $T$  is the temperature (K) and  $\kappa$  is the thermal conductivity of oceanic rock ( $\text{J yr}^{-1} \text{m}^{-1} \text{K}^{-1}$ ). The thermodynamic and transport properties of water ( $c_p^f$ ,  $\rho_f$  and  $\nu$ ) are obtained through a FORTRAN90 library STEAM which is based on the NBS steam table package (Meyer et al., 1983; Haar et al., 1984). As in Iyer et al. (2010),  
 95 we assume pure water for hydrothermal fluid. The petrophysical parameters except for the permeability (i.e.,  $\kappa$ ,  $\phi$ ,  $\rho_m$  and  $c_p^m$ ) are assumed to be the same as those in Iyer et al. (2010). We assume that  $\log k$  exponentially decreases from  $-11.8$  to  $-16.8$  with a length scale of 300 m for an e-fold decrease, which is consistent with observations (Fisher, 1998; Supplementary material). Following Cherkaoui et al. (2003), oceanic rocks below the crust/mantle interface (6 km depth from the ocean/crust interface) and with temperatures above the critical temperature for rock cracking (600 °C) are impermeable. As long as the  
 100 model is consistent with observations, changing the assumptions about permeability does not affect the general conclusions given in the present study (Supplementary material).

The boundary conditions adopted in the present study follow those by Iyer et al. (2010) and/or Cherkaoui et al. (2003). Pressure at the ocean/crust interface is assumed as constant at 25 MPa, corresponding to the assumption of 2.5 km water depth. This assumption is modified when we explore the Precambrian hydrothermal circulation in Section 3.3. The right and  
 105 bottom sides of the calculation domain are assumed to be insulating. The ocean above the top boundary is assumed to have a constant temperature of 2 °C. The ridge axis on the left-hand boundary is assumed to supply a boundary heat flux  $J_b$  ( $\text{J m}^{-2} \text{yr}^{-1}$ ):

$$J_b = \rho_m w (T_m - T) c_p^m, \quad (4)$$

where  $w$  is the spreading rate ( $\text{m yr}^{-1}$ ) and  $T_m$  is the temperature of the intrusion (1200 °C). As a standard value, we assume  
 110  $3 \times 10^{-2} \text{ m yr}^{-1}$  for  $w$ , and this assumption is again changed in Section 3.3 where we consider the Precambrian hydrothermal circulation. Free flow is allowed at the top boundary, and the other boundaries are assumed to be impermeable (Iyer et al., 2010). The calculation domain is wide enough (see below) that making the right-hand boundary permeable with respect to heat and water will change the results only negligibly (cf. Supplementary material; Cherkaoui et al., 2003). See Table 1 for the definitions and values of the parameters used in the present study.

115 A finite difference approach is taken to solve Eqs. (1)–(3) for  $\mathbf{q}$ ,  $P$  and  $T$  using the second-order central differencing scheme for the second-order differential terms and the first-order upwind and forward differencing schemes for the first-order spatial and temporal differential terms, respectively (e.g., Steefel and Lasaga, 1994). The calculation procedure follows that by Iyer et al. (2010). First, we obtain  $P$  by solving Eqs. (1) and (2). Then,  $\mathbf{q}$  is obtained from the calculated  $P$  and Eq. (2). Finally, the calculated  $\mathbf{q}$  is used in Eq. (3) which is solved to obtain  $T$  for the same time step. The calculated  $P$  and  $T$  are used to update  
 120 the thermodynamic and transport properties of water for the next calculation step. The above procedure is repeated with a time step of  $3 \times 10^4 \text{ yr}$  until steady states are reached, which is accomplished within  $10^3$  time steps or  $3 \times 10^7$  model years (cf.

Cherkaoui et al., 2003). Note that only steady states are considered in this study because of the long time scale over which oceanic oxygen-isotopic composition can change ( $> \sim 5 \times 10^7$  yr; e.g., Holland, 1984; Walker and Lohmann, 1989; Wallmann, 2001). The model grid extends as deep as 12 km depth from the ocean/crust interface allowing changes in the location of the crust/mantle interface (though not varied in this study); and it extends to 30 km distance from the ridge axis, wide enough to simulate major changes in oxygen isotopic composition of oceanic crust which have been observed to be completed within less than 10 million years from the rock formation at midocean ridges (Muehlenbachs, 1979) (see Supplementary material for more details). The calculation domain ( $12 \times 30$  km<sup>2</sup>) is divided into a  $320 \times 200$  irregular grid, with the grid-cell size horizontally increasing from the ridge axis (1.1 to 330 m) and vertically from the ocean/crust interface (0.17 to 82 m) (cf. Cathles, 1983; Cherkaoui et al., 2003).

## 2.2 Reactive transport model of oxygen isotopes

Oxygen isotopes of solid rock are assumed to be transported by the spreading of oceanic rocks, while those of porewater are transported by water flow and molecular diffusion (cf. Norton and Taylor, 1979; Lécuyer and Allemand, 1999). At the same time, isotope exchange reactions transfer <sup>18</sup>O between the two phases (e.g., Norton and Taylor, 1979; Lécuyer and Allemand, 1999). Then, from the mass conservation of <sup>18</sup>O in the two phases, the time rates of change of the <sup>18</sup>O/total O mole ratios of solid rock and porewater ( $F_r$  and  $F_p$ , respectively) can be represented by

$$(1 - \phi) \rho_m m_s \frac{\partial F_r}{\partial t} = - (1 - \phi) \rho_m m_s w \frac{\partial F_r}{\partial x} - \rho_b m_s m_f k_{\text{ex}} \{ F_r (1 - F_p) - \alpha (1 - F_r) F_p \}, \text{ and} \quad (5)$$

$$\phi \rho_f m_f \frac{\partial F_p}{\partial t} = \nabla \cdot (-m_f \mathbf{q} F_p + \phi \rho_f m_f D \nabla F_p) + \rho_b m_s m_f k_{\text{ex}} \{ F_r (1 - F_p) - \alpha (1 - F_r) F_p \}, \quad (6)$$

where  $m_s$  and  $m_f$  are the oxygen concentrations of solid rock and porewater, respectively (mol kg<sup>-1</sup>),  $\rho_b$  is the density of bulk rock given by  $\rho_b = \phi \rho_f + (1 - \phi) \rho_m$ ,  $k_{\text{ex}}$  is the rate constant for oxygen isotope exchange (mol<sup>-1</sup> kg yr<sup>-1</sup>),  $\alpha$  is the oxygen isotope fractionation factor and  $D$  is the effective diffusion coefficient for <sup>18</sup>O (m<sup>2</sup> yr<sup>-1</sup>), determined by molecular diffusion and hydrodynamic dispersion. The first and second terms on the right-hand sides of Eqs. (5) and (6) represent oxygen isotope transport and oxygen isotope exchange, respectively. The kinetic expression of oxygen isotope exchange in Eqs. (5) and (6) is formulated based on Cole et al. (1983, 1987) and the rate constant is given by an Arrhenius equation:

$$k_{\text{ex}} = 10^{-8.5} \exp \left\{ - \frac{E}{R_g} \left( \frac{1}{T} - \frac{1}{278} \right) \right\}, \quad (7)$$

where  $E$  is the apparent activation energy ( $5 \times 10^4$  J mol<sup>-1</sup>) and  $R_g$  is the gas constant (8.314 J mol<sup>-1</sup> K<sup>-1</sup>). The value of  $10^{-8.5}$  mol<sup>-1</sup> kg yr<sup>-1</sup> at reference temperature 278 K (5 °C) is obtained by reducing the range  $10^{-7.2}$ – $10^{-6.6}$  mol<sup>-1</sup> kg yr<sup>-1</sup> at 5 °C extrapolated from  $10^{-2.1}$ – $10^{-1.5}$  mol m<sup>-2</sup> yr<sup>-1</sup> at 300 °C (Cole et al., 1987) with  $E = 5 \times 10^4$  J mol<sup>-1</sup> (cf. Cole et al., 1987) and  $10^3$  m<sup>2</sup> kg<sup>-1</sup> specific surface area of marine basalt (cf. Nielsen and Fisk, 2010) by a factor of  $10^{1.3}$ – $10^{1.9}$ . The reducing factor ( $10^{1.3}$ – $10^{1.9}$ ) accounts for the fact that the reaction rate in the field is generally slower than in the laboratory by a factor of up to  $10^3$  (e.g., Wallmann et al., 2008) (see Supplementary material for the details). A similar factor (up to  $10^4$ ) has been adopted by Cathles (1983) for the formulation of oxygen isotope exchange kinetics. Note that the general results and

conclusions in the present study are not affected by changes in the reference  $k_{\text{ex}}$  value within a plausible range (Supplementary material). The oxygen isotope fractionation factor for andesite by Zhao and Zheng (2003) is adopted in the present study:

$$10^3 \ln \alpha = \beta \left( \frac{6.673 \times 10^6}{T^2} + \frac{10.398 \times 10^3}{T} - 4.78 \right) \exp \left( \frac{1 - \beta}{R_g T} \right) - \frac{2.194 \times 10^6}{T^2} - \frac{15.163 \times 10^3}{T} + 4.72 + 1.767(2\beta - 1), \quad (8)$$

155

where  $\beta = 0.876$  (Zhao and Zheng, 2003), because Eq. (8) with  $\beta = 0.876$  yields comparable  $\alpha$  values to experimental results for basalt by Cole et al. (1987) at 300–500 °C while applicable over the wider range of temperature considered for the present study (2–1200 °C). The effective diffusion coefficient  $D$  considers both molecular diffusion and hydrodynamic dispersion; the former is obtained from the modified Stokes-Einstein relation for  $\text{H}_2^{16}\text{O}$  diffusion by Krynicki et al. (1978) and the isotope  
160 effect from Harris and Woolf (1980) and a homogeneous dispersivity of 10 m is assumed for the latter (Frind, 1982; Gelhar et al., 1992):

$$D = \tau 6.9 \times 10^{-15} \frac{T}{\rho_f \nu} \sqrt{\frac{18}{20}} + 10 \frac{|\mathbf{q}|}{\phi \rho_f}. \quad (9)$$

The first term on the right-hand side of Eq. (9) represents the molecular diffusion including the tortuosity factor  $\tau = \phi^{1.4}$  (Aachib et al., 2004), while the second term represents the hydrodynamic dispersion. The petrophysical parameters are as-  
165 sumed to be the same as those in the hydrothermal circulation model (Section 2.1, Table 1). The thermodynamic and transport properties of water are obtained through the hydrothermal circulation model (Section 2.1).

The steady-state values of  $F_r$  and  $F_p$  are obtained by solving  $\partial F_r / \partial t = \partial F_p / \partial t = 0$  in Eqs. (5) and (6). The intrusion on the left-hand boundary is assumed to have a  $^{18}\text{O}$ /total O mole ratio of fresh crust ( $F_m$ ) that corresponds to 5.7‰ relative to SMOW (e.g., Holmden and Muehlenbachs, 1993), and the ocean above the top boundary is assumed to have a constant  $^{18}\text{O}$ /total O mole  
170 ratio of seawater ( $F_{\text{sw}}$ ). The other boundaries are impermeable for  $^{18}\text{O}$  fluxes via water (cf. Section 2.1). A finite difference method is used for equation discretization (first-order upwind and second-order central differencing schemes for the first-order and second-order spatial differential terms, respectively) and Newton’s method is adopted to solve the difference equations. The calculation is conducted on the grid described earlier for the hydrothermal circulation model (Section 2.1). The  $^{18}\text{O}$  mole ratios are all reported in  $\delta$  notation relative to SMOW, using  $2.0052 \times 10^{-3}$  as the  $^{18}\text{O}/^{16}\text{O}$  mole ratio of SMOW (Fry, 2006).

## 175 3 Results

### 3.1 Application to the present day and model validation

The calculated flow geometry and temperature distribution (Fig. 1; note that Figs. 1c and d are plots of Figs. 1a and b, respectively, on logarithmic scales) are similar to those in previous studies (e.g., Cherkaoui et al., 2003). The 2D model results can be converted to associated mass and heat fluxes, assuming  $10^8$  m ridge length (cf. Wolery and Sleep, 1976). The total heat flux  
180 from the system is  $0.74 \times 10^{12}$  W, comparable to the observed cumulative heat flux within 1 Myr (corresponding to 30 km with  $3 \times 10^{-2}$  m yr $^{-1}$  spreading rate) from the ridge axis,  $0.4(\pm 0.3) \times 10^{12}$  W (Stein and Stein, 1994). The total mass of water

exchange is  $1.2 \times 10^{13}$  kg yr<sup>-1</sup>, falling within the constrained range ( $3 \times 10^{12}$  to  $1.6 \times 10^{14}$  kg yr<sup>-1</sup>) and close to the recommended value ( $3(\pm 1.5) \times 10^{13}$  kg yr<sup>-1</sup>) by Elderfield and Schultz (1996), and also comparable to the prediction from other hydrothermal circulation models (e.g., Norton and Knight, 1977; Fehn et al., 1983; Cherkaoui et al., 2003). The calculated 2D  
185 distributions of solid-rock and porewater  $\delta^{18}\text{O}$  (Figs. 2a, 2d, 3a and 3d; note that Fig 3 is a plot of Fig. 2 on logarithmic scales) are consistent with observations, especially at 30 km (1 Myr) from the ridge axis (black solid curves in Fig. 4; see Section 3.2 for paler curves). Where flow rate and temperature are highest near the ridge axis and close to the ocean/crust interface (at  $\sim 200$  m depths), porewater  $\delta^{18}\text{O}$  is in the range between 0 and 2 ‰ (Fig. 3d), which is comparable to the observed  $\delta^{18}\text{O}$  ranges for high-temperature hydrothermal fluids, e.g., 0.49–2.3 ‰ by Jean-Baptiste et al. (1997) and 0.3–1.4 ‰ by James et al. (2014). Away from the ridge axis, porewater  $\delta^{18}\text{O}$  is equivalent to the seawater value at the ocean/crust interface, but drops  
190 down to  $-4$  to  $-14$  ‰ within  $\sim 200$  m depths, again not inconsistent with observed porewater  $\delta^{18}\text{O}$  values (down to  $-8$  ‰) at  $< 500$  m depths where water exchange is limited (e.g., Lawrence and Gieskes, 1981). Solid rock  $\delta^{18}\text{O}$  at 1 Myr (i.e., 30 km from the ridge axis) is characterized by high ( $< 16$  ‰) and low ( $> 3$  ‰) values at shallow ( $\leq \sim 2$  km) and deeper ( $\geq \sim 2$  km) depths, respectively, consistent with observations of modern oceanic crust (e.g., Alt et al., 1986) and ophiolites (e.g.,  
195 Gregory and Taylor, 1981; crosses in Fig. 4) (Fig. S3 in Supplementary material). The  $^{18}\text{O}$  fluxes to the ocean from high- and low-temperature alteration are  $3.0 \times 10^9$  and  $-2.8 \times 10^9$  mol yr<sup>-1</sup>, respectively, well balanced, resulting in a net  $^{18}\text{O}$  flux of  $0.2 \times 10^9$  mol yr<sup>-1</sup>, consistent with the suggestion of zero net-flux by Gregory and Taylor (1981). Individual  $^{18}\text{O}$ -flux values are also comparable to those suggested in the previous studies. As examples,  $^{18}\text{O}$  fluxes through high-temperature alteration have been suggested to be  $4.5 \times 10^9$  (Holland, 1984),  $3.2 \times 10^9$  (Muehlenbachs, 1998) and  $2.8 \times 10^9$  mol yr<sup>-1</sup> (Wallmann,  
200 2001), and those through low-temperature alteration  $< -2.3 \times 10^9$  mol yr<sup>-1</sup> (Lawrence and Gieskes, 1981),  $-1.1 \times 10^9$  (Holland, 1984),  $-1.0 \times 10^9$  (Muehlenbachs, 1998) and  $-0.9 \times 10^9$  mol yr<sup>-1</sup> (Wallmann, 2001). The consistency between the model calculation and observations described above supports the validity of the model.

### 3.2 Evaluation of oceanic- $\delta^{18}\text{O}$ buffering capacity

As both the source and sink of oceanic  $^{18}\text{O}$ , hydrothermal systems can buffer oceanic  $\delta^{18}\text{O}$  but the strength of the buffering  
205 depends on the sensitivity of isotope exchange between oceanic crust and porewater to seawater  $\delta^{18}\text{O}$  (cf. Wallmann, 2001). As an extreme example, if the oxygen isotope fractionation between solid rock and porewater is independent of seawater  $\delta^{18}\text{O}$ , there should not be any feedbacks from the hydrothermal systems on changes in seawater  $\delta^{18}\text{O}$ , i.e., no oceanic- $\delta^{18}\text{O}$  buffering. The previous studies that support the strong oceanic- $\delta^{18}\text{O}$  buffering at midocean ridges assume linear relationships between oxygen isotope fractionation and seawater  $\delta^{18}\text{O}$  (e.g., Gregory and Taylor, 1981; Holland, 1984; Muehlenbachs,  
210 1998). This assumption regarding the sensitivity to seawater  $\delta^{18}\text{O}$  is difficult to confirm through observations of geological records because seawater  $\delta^{18}\text{O}$  is not known. We can examine the response of rocks to seawater  $\delta^{18}\text{O}$  with the present model by adopting different values for seawater  $\delta^{18}\text{O}$ . We can then measure the buffering capacity of the system by plotting net  $^{18}\text{O}$  flux against seawater  $\delta^{18}\text{O}$  and calculating the slope value, i.e.,  $\partial(\text{net } ^{18}\text{O flux})/\partial(\text{seawater } \delta^{18}\text{O})$ , as in, e.g., Muehlenbachs and Clayton (1976). For example, a hydrothermal system with a large negative slope value should exhibit a strong buffering



215 of oceanic  $\delta^{18}\text{O}$ , because a slight change in seawater  $\delta^{18}\text{O}$  makes the system introduce a large net  $^{18}\text{O}$  flux to the ocean that restores the change in seawater  $\delta^{18}\text{O}$  (e.g., Muehlenbachs and Clayton, 1976; Muehlenbachs, 1998).

Most features of 2D distributions of solid-rock and porewater  $\delta^{18}\text{O}$  are not significantly affected by changing seawater  $\delta^{18}\text{O}$  from 6 to  $-12\text{‰}$  (Figs. 2–4). Although the porewaters close to the ocean have  $\delta^{18}\text{O}$  compositions close to those of seawater (e.g., Fig. 3),  $\delta^{18}\text{O}$  signatures of solid rocks are not linearly proportional to seawater  $\delta^{18}\text{O}$  (e.g., Fig. 4). The relative insensitivity at shallow depths ( $\leq \sim 2\text{ km}$ ; Figs. 2–4) can be explained by the kinetics of oxygen isotope exchange. The distance from isotope exchange equilibrium can be measured by  $\Omega = F_r(1 - F_p)/\{\alpha(1 - F_r)F_p\}$  (Eqs. (5) and (6); Figs. 5b and d; note that Figs. 5c and d are plots of Figs. 5a and b, respectively, on logarithmic scales), and the shallow regions show non-equilibrium states ( $\Omega \neq 1$ ) because of their relatively low temperatures (Figs. 1b and d). Therefore, despite the approximate isotopic equivalence between porewater and seawater (Figs. 2 and 3), solid rocks do not directly reflect seawater  $\delta^{18}\text{O}$  in their oxygen-isotopic compositions. In deeper sections of oceanic crust, on the other hand, solid rocks attain isotope exchange equilibrium with porewater because of the relatively high temperatures and the correspondingly fast rates of isotope exchange (e.g., Figs. 5b and d). These deep regions, however, receive less water exchange than the shallow regions, as can be seen from the distribution of local water/rock oxygen-mole ratio  $\eta (\equiv m_f|\mathbf{q}|/\{(1 - \phi)\rho_m m_{sw}\})$  (Figs. 5a and c). Accordingly, deep oceanic rocks and porewaters are oxygen-isotopically buffered by spreading solid-rocks rather than by seawater, and  $\delta^{18}\text{O}$  values of deep porewaters are almost independent of seawater  $\delta^{18}\text{O}$  (Figs. 2d–f and 3d–f). Consequently, despite the isotope exchange equilibrium, deep oceanic rocks are almost completely insensitive to seawater  $\delta^{18}\text{O}$ . The combination of the above two factors (isotope exchange kinetics and  $^{18}\text{O}$  supply from spreading solid-rocks) explains the insensitivity of midocean ridge systems to seawater  $\delta^{18}\text{O}$  (Figs. 2–4). Note that the kinetics of oxygen isotope exchange and  $^{18}\text{O}$  supply from rocks have been invoked to explain the positive and negative  $\delta^{18}\text{O}$  anomalies, respectively, relative to the pristine crustal  $\delta^{18}\text{O}$  value by Cathles (1983), not inconsistent with the two factors causing the partial decoupling between oceanic crust and seawater  $\delta^{18}\text{O}$  in the present model.

Consistent with the insensitivity of oceanic rocks to seawater  $\delta^{18}\text{O}$  (Figs. 2–4), the net  $^{18}\text{O}$  flux from the midocean ridge systems is a very weak function of seawater  $\delta^{18}\text{O}$  (Fig. 6). Compared to the large buffering capacity suggested in the previous studies, which can be recognized from large negative slopes in Fig. 6, the intensity of oceanic- $\delta^{18}\text{O}$  buffering exhibited by the present model is weak; the magnitude of slope ( $-0.4 \times 10^9 \text{ mol yr}^{-1} \text{‰}^{-1}$ ) is smaller than those previously assumed by a factor of up to  $>7$ . The reason why the previous studies suggested strong buffering is because these studies assume isotope exchange equilibrium between porewaters and oceanic rocks and equivalence between porewater and seawater at the same time. These assumptions are not satisfied simultaneously in midocean ridge systems in general, as can be anticipated from Figs. 2–5. The exception to the previous studies is Wallmann (2001), which shows relatively small negative slope value (Fig. 6). Note, however, that Wallmann (2001) simulated weak buffering in a different way, i.e., by adopting low and temperature-independent rate constants for oxygen isotope exchange.

### 3.3 Application to the Precambrian

As the tectonics of the Earth likely evolved through time, the midocean ridge systems in the Precambrian could have been quite different from those in the present day. Indeed, differences in ocean volume and the spreading rate of oceanic crust have  
250 been discussed (e.g., Kasting et al., 2006; Korenaga et al., 2017). Here, we examine the oceanic- $\delta^{18}\text{O}$  buffering during the Precambrian by utilizing possible ranges of the spreading rate ( $1 \times 10^{-2}$  to  $30 \times 10^{-2}$  m yr $^{-1}$ ) and ocean volume (1 to 5 km of water depth at the ocean/crust interface) as well as seawater  $\delta^{18}\text{O}$  ( $-12$  to  $6$  ‰) suggested in the literature (e.g., Phipps Morgan, 1998; Kasting et al., 2006; Jaffrés et al., 2007; Korenaga et al., 2017; Galili et al., 2019; Johnson and Wing, 2020).

Increasing the spreading rate results in increases in the heat flux and associated water exchange between the crust and ocean,  
255 consistent with modern observations (e.g., Baker et al., 1996; Bach and Humphris, 1999):  $0.50 \times 10^{12}$ ,  $0.99 \times 10^{12}$  and  $1.25 \times 10^{12}$  W of heat flux and  $2.4 \times 10^{12}$ ,  $4.9 \times 10^{13}$  and  $8.5 \times 10^{13}$  kg yr $^{-1}$  of water exchange at spreading rates of  $1 \times 10^{-2}$ ,  $9 \times 10^{-2}$  and  $30 \times 10^{-2}$  m yr $^{-1}$ , respectively (Figs. 7a–f). On the other hand, differences in temperature and water-flow distributions caused by those in ocean depth (from 1 to 5 km) are relatively minor ( $0.74 \times 10^{12}$  W and  $(1.2\text{--}1.3) \times 10^{13}$  kg yr $^{-1}$ ; Figs. 7g–j). The distributions of solid-rock and porewater  $\delta^{18}\text{O}$  predicted under these different alteration conditions are correspondingly  
260 modified but show features common to those shown in the previous sections with the standard parameterization. Porewater  $\delta^{18}\text{O}$  is close to seawater  $\delta^{18}\text{O}$  only along the ocean/crust interface and becomes quite different from seawater  $\delta^{18}\text{O}$  at deep depths (Supplementary material). Solid rock  $\delta^{18}\text{O}$  is relatively insensitive to seawater  $\delta^{18}\text{O}$  in general, showing relative addition and removal of  $^{18}\text{O}$  at shallow and deeper depths, respectively (Fig. 8). The buffering capacity exhibited by models that reflect changes in the spreading rate and ocean depth is different from that in the standard case (Fig. 9), but still weaker than any  
265 of those assumed in the previous studies (cf. Fig. 6). Note that in the previous studies, the buffering capacity is assumed to increase linearly with the spreading rate (Gregory and Taylor, 1981; Holland, 1984; Muehlenbachs, 1998; Wallmann, 2001) and thus the slope values in Fig. 9 need be compared with those in the literature (Fig. 6) multiplied by a factor that accounts for changes in the spreading rate. As an example, the slope value of  $-1.9 \times 10^9$  mol yr $^{-1}$  ‰ $^{-1}$  with a spreading rate  $30 \times 10^{-2}$  m yr $^{-1}$  in Fig. 9 should be compared with  $-6 \times 10^9$  (Wallmann, 2001),  $-(14\text{--}27) \times 10^9$  (Gregory and Taylor, 1981),  $-24 \times 10^9$   
270 (Muehlenbachs, 1998) and  $-31 \times 10^9$  mol yr $^{-1}$  ‰ $^{-1}$  (Holland, 1984) (cf. Fig. 6); thus the buffering with the present model assuming  $30 \times 10^{-2}$  m yr $^{-1}$  of spreading rate is much weaker than those assumed in the previous studies. More details of changes in the system behavior are described below.

When spreading is slower, seawater penetrates deeper into the crust because of a weaker boundary heat flux (Eq. (4)) and thus hydrothermal flows are more homogenized but less intense, compared to those in the standard case (Fig. 7a). Although  
275 the changes of crustal  $\delta^{18}\text{O}$  from the pristine  $\delta^{18}\text{O}$  value (5.7 ‰) become larger over the whole crust (Fig. 8a), the sensitivity of solid-rock to seawater  $\delta^{18}\text{O}$  remains small (Fig. 8a). The small sensitivity is caused because oxygen isotope exchange at shallow depths of oceanic crust is still kinetically limited despite the elongated time duration for oxygen isotope exchange (3 Myr; cf. Figs. S13 and S16 in Supplementary material) and because the deeper section is buffered by spreading rocks rather than circulating seawater, given the simultaneous decreases in the amount of water exchange and the spreading rate. The lower  
280 O supply from the mantle combined with the small sensitivity makes the buffering intensity weaker than that with the standard

spreading rate (Fig. 9). With the spreading rate high, on the other hand, the hydrothermal penetration depth becomes shallower as a result of enhanced boundary heat flux (Eq. (4)) and hydrothermal flows are more intense and more localized close to the ocean and the ridge axis (Figs. 7c and e). Accordingly,  $^{18}\text{O}$  depletion from solid rocks near the ridge axis at relatively high temperatures and water/rock ratios is not recovered even after continued reactions at low temperatures. Nonetheless, the sensitivity of solid-rock to seawater  $\delta^{18}\text{O}$  does not significantly differ from that in the other spreading-rate cases (Figs. 8b and c) because the kinetics of oxygen isotope exchange and  $^{18}\text{O}$  buffering by spreading oceanic rocks remain effective to decouple the shallow and deeper oceanic crust from seawater  $\delta^{18}\text{O}$ , respectively. Note that the time duration for oxygen isotope exchange is reduced with the higher spreading rate ( $\geq 0.1$  Myr) but can still be long enough to complete major changes in the oxygen-isotopic composition of oceanic crust (cf. Fig. S15 in Supplementary material). Despite the weak rock sensitivity to seawater  $\delta^{18}\text{O}$ , the larger O supply from the mantle makes the buffering capacity larger than that with the standard spreading rate (Fig. 9). Overall, the system as a whole exhibits non-linear and reduced sensitivity to changes in the spreading rate as recognized from the relationship between the spreading rate and buffering capacity in Fig. 9 (cf. Gregory and Taylor, 1981; Holland, 1984; Muehlenbachs, 1998; Wallmann, 2001).

When assuming a shallow ocean with the ocean/crust interface at 1 km water depth, hydrothermal circulation becomes slightly stronger and occurs slightly closer to the ocean (Fig. 7g) (cf. Kasting et al., 2006). In contrast, in a deep ocean with the ocean/crust interface at 5 km depth from the sea surface, the mixing of water becomes slightly weaker and occurs slightly deeper (Fig. 7i). In both cases, the above changes in the flow geometry and intensity are relatively minor. The resultant oxygen-isotope behaviors are thus similar to the standard case (Figs. 4, 8d and 8e) and the oceanic- $\delta^{18}\text{O}$  buffering becomes only slightly weaker in the shallow- and deep-ocean cases than in the standard case (Fig. 9). Notably, oxygen isotope exchange through low-temperature hydrothermal alteration is relatively enhanced in the shallow ocean case (Fig. 9), which is not inconsistent with the prediction by Kasting et al. (2006). However, the magnitude of the ocean-depth effect is much smaller than anticipated by Kasting et al. (2006), probably because of the settings in the hydrothermal circulation model (Section 2.1) where two-dimensional distributions of permeability as well as boundary heat flux associated with the spreading rate (Eq. (4)) may exert a more significant control over the hydrothermal penetration depth (cf. Fig. 7 and Supplementary material).

## 305 4 Discussion

### 4.1 Interpretation of ophiolites

Simulations conducted in the present study suggest that oceanic rocks are not significantly affected by changes in seawater  $\delta^{18}\text{O}$  under any plausible alteration conditions (Figs. 4 and 8). Reported  $\delta^{18}\text{O}$  values of ophiolites and/or oceanic crust range from  $\sim 1$  to  $16$  ‰ (dashed lines in Fig. 8). By comparison, the simulated solid-rock  $\delta^{18}\text{O}$  values fall within this range for seawater  $\delta^{18}\text{O}$  values  $\leq \sim -10$ ,  $\sim -8$  to  $0$ ,  $\geq \sim -2$  and  $\geq \sim -2$  ‰ at a spreading rate of  $1 \times 10^{-2}$ ,  $3 \times 10^{-2}$ ,  $9 \times 10^{-2}$  and  $30 \times 10^{-2}$   $\text{m yr}^{-1}$ , respectively (Figs. 4 and 8). Accordingly, we conclude that the constant seawater  $\delta^{18}\text{O}$  at  $0$  ‰ is neither a necessary nor a sufficient condition for explaining the relatively invariant  $\delta^{18}\text{O}$  records of ophiolites. As ophiolite  $\delta^{18}\text{O}$  profiles can be affected more by alteration conditions (e.g., the spreading rate and permeability; Section 3.3 and Supplementary material)

with the control by seawater  $\delta^{18}\text{O}$  remaining relatively weak, feedbacks between the alteration parameters (e.g., spreading  
315 rate and permeability) could be more important in reproducing ophiolite records. Therefore, ophiolites may be interpreted to  
indicate the insensitivity of oceanic rocks to seawater  $\delta^{18}\text{O}$ , realized by the feedbacks between alteration parameters, and not  
necessarily a constant seawater  $\delta^{18}\text{O}$ . The weak buffering (Figs. 6 and 9) accompanying the partial decoupling between the  
oceanic crust and seawater  $\delta^{18}\text{O}$  (e.g., Fig. 8) shows that this interpretation is more plausible than the constant seawater  $\delta^{18}\text{O}$ .

#### 4.2 Controls of $\delta^{18}\text{O}$ in the Precambrian oceans

320 Although the buffering of oxygen isotopes through hydrothermal alteration of oceanic rocks is weaker than previously assumed  
under any alteration conditions (Section 3.3), it could have been relatively strong when the spreading rate is  $\geq \sim 30 \times 10^{-2}$   
 $\text{m yr}^{-1}$  (Figs. 6 and 9). Such high-spreading rate conditions could have been possible only during the earliest period of  
Earth's history ( $> 3.5$  Ga; Phipps Morgan, 1998) or even impossible according to Korenaga (2006). Excluding this earliest  
325 period, the buffering must have been weak despite the uncertainties in alteration conditions (Figs. 6 and 9). The weak buffering  
must have allowed variations of seawater  $\delta^{18}\text{O}$  through other surficial processes that exchange  $^{18}\text{O}$  with seawater, most likely  
through continental weathering (e.g., Walker and Lohmann, 1989). As surface environments likely have significantly changed  
through the eons (cf. Introduction),  $^{18}\text{O}$  fluxes from continental weathering could have correspondingly varied and might have  
contributed to the oceanic  $^{18}\text{O}$  budget more significantly than those from hydrothermal alteration of oceanic crust during certain  
periods in Earth's history.

330 Previous studies examining oceanic- $\delta^{18}\text{O}$  evolution on geological time scales have utilized box-modeling approaches to  
account for oxygen isotope exchange from both continental weathering and hydrothermal alteration of oceanic crust (Godderis  
and Veizer, 2000; Godd ris et al., 2001; Wallmann, 2001; Kasting et al., 2006; Jaffr s et al., 2007). Among them, the studies  
that assume strong buffering at midocean ridges have had difficulty in modifying oceanic  $^{18}\text{O}$  budget through other surficial  
processes including continental weathering (Godderis and Veizer, 2000; Godd ris et al., 2001). Other studies that assume  
335 weak buffering instead have shown the possibility of significant oceanic- $\delta^{18}\text{O}$  changes through modification of simplified  
continental-weathering parameters with the Earth's age (Wallmann, 2001; Kasting et al., 2006; Jaffr s et al., 2007). Revisiting  
theses previous box-model studies with the weak buffering suggested here (e.g., Figs. 6 and 9) and constructing a process-  
based model for continental weathering to be coupled with the present model will lead to a better understanding of controls on  
oceanic  $\delta^{18}\text{O}$  during the Precambrian.

#### 340 4.3 Comparison with other models

The results in the present study differ significantly from those by L cuyer and Allemand (1999), who utilized prescribed  
distributions of temperature and water/rock ratio along with an isotope-fractionation equation for a half-closed system (a  
system that is open to the water reservoir, but not to the rock reservoir; Gregory et al., 1989) (Introduction). The assumed range  
for water/rock ratio ( $\leq 10$ ) by these authors is comparable to the simulated range in the present study ( $\eta \leq 164$ ; Figs. 5a and  
345 c). States of equilibrium and non-equilibrium in the bulk-rock-based formulation of isotopes in the present study correspond to  
transport- and reaction-limited alteration states, respectively, in the mineral-based isotope formulation in L cuyer and Allemand

(1999); the difference in the formulation of isotopic composition of bulk rock should not cause any significant differences in oxygen isotope behavior of bulk rock (cf. Bindeman et al., 2019). The temperature dependence of kinetic constant for oxygen isotope exchange is described with the Arrhenius equation, based on the datasets by Cole and Ohmoto (1986) in Lécuyer and Allemand (1999) and by Cole et al. (1983, 1987) in the present study, and thus there should be little difference in the temperature dependence of isotope exchange kinetics between the two studies. The size difference in the calculation domain cannot explain the difference between the present study and Lécuyer and Allemand (1999) as additional numerical experiments on a wider calculation domain (300 km) with artificially imposing off-axis water exchange (up to  $5.3 \times 10^{15}$  kg yr<sup>-1</sup>) yield essentially the same results as those with the standard calculation domain width (30 km) (Supplementary material). Although a thorough comparison cannot be made between the two studies because Lécuyer and Allemand (1999) have not provided detailed model results, it is likely that the isotopic effect of spreading crust has not been explicitly considered by Lécuyer and Allemand (1999). Should this be the case, the sensitivity of oceanic rocks to seawater  $\delta^{18}\text{O}$  and the buffering capacity at midocean ridges could be overestimated by the model of Lécuyer and Allemand (1999). More generally, the models that do not explicitly account for the transport of solid rocks (e.g., Taylor, 1977; Norton and Taylor, 1979; Criss et al., 1987; Gregory et al., 1989; DePaolo, 2006) could overestimate the coupling strength between oceanic crust and seawater  $\delta^{18}\text{O}$  and the oceanic-<sup>18</sup>O buffering capacity of hydrothermal systems. Application of these models to ophiolites may also underestimate the uncertainty in reconstructed ancient seawater  $\delta^{18}\text{O}$  that is caused by the partial decoupling between oceanic crust and seawater  $\delta^{18}\text{O}$  (cf. Section 4.1 and Supplementary material).

## 5 Conclusions

The 2D reactive transport model of oxygen isotopes combined with 2D hydrothermal circulation simulations enables us to predict distributions of temperature, water flow and oxygen isotopes of solid rocks and porewaters within oceanic crust based on mass, momentum and energy conservations. The model assuming the present-day seawater  $\delta^{18}\text{O}$  reproduced those distributions consistent with modern observations, supporting the model's validity. The intensity of seawater- $\delta^{18}\text{O}$  buffering at midocean ridges was evaluated by calculating the net <sup>18</sup>O flux as a function of seawater  $\delta^{18}\text{O}$ . The buffering intensity predicted by the model is significantly weaker than those previously assumed under any plausible alteration conditions during the Precambrian. The weak buffering is realized because isotope exchange equilibrium is not reached in shallow sections of oceanic crust with low temperatures, and because <sup>18</sup>O supply via spreading solid rocks exceeds that through hydrothermal circulation in deeper high-temperature sections. Consistently with the weak buffering, oceanic rocks are insensitive to seawater with respect to oxygen isotopes. Thus, ophiolites can alternatively be explained by the insensitivity of oceanic rocks to seawater  $\delta^{18}\text{O}$  that could have evolved through the Precambrian because of the weak buffering at midocean ridges.

*Code availability.* ~~The source codes of the model are available upon request to the author.~~ The source codes of the model are available on GitHub (<https://github.com/kanzakiy/HT-Oiso>) under the GNU General Public License v3.0. The specific version used of the model in this

paper is tagged as 'v0.1' and has been assigned a DOI: 10.5281/zenodo.3922317 (<https://doi.org/10.5281/zenodo.3922317>). A readme file on the web provides the instructions for executing the simulations.

## 380 **Appendix A: Supplementary material**

Supplementary material related to this article can be found online at <https://doi.org/xxxxxx>.

*Author contributions.* YK conducted the whole research.

*Competing interests.* I declare no competing interests.

385 *Acknowledgements.* I thank Jim Kasting and Lee Kump for helpful discussions and Laurence Coogan and an anonymous reviewer for their useful comments on earlier version of manuscript. I am grateful to Benjamin Johnson and Itay Halevy for their useful comments on the current manuscript and to Patrice Rey for editorial handling.

## References

- Aachib, M., Mbonimpa, M. and Aubertin, M.: Measurement and prediction of the oxygen diffusion coefficient in unsaturated media, with applications to soil covers, *Water Air Soil Pollut.*, 156, 163–193, doi:10.1023/B:WATE.0000036803.84061.e5, 2004.
- 390 Agrinier, P., Javoy, M. and Girardeau, J.: Hydrothermal activity in a peculiar oceanic ridge: oxygen and hydrogen isotope evidence in the Xigaze ophiolite (Tibet, China), *Chem. Geol.*, 71, 313–335, doi:10.1016/0009-2541(88)90057-5, 1988.
- Alt, J. C. and Bach, W.: Oxygen isotope composition of a section of lower oceanic crust, ODP Hole 735B, *Geochem. Geophys. Geosyst.*, 7, Q12008, doi:10.1029/2006GC001385, 2006.
- Alt, J. C., Muehlenbachs, K. and Honnorez, J.: An oxygen isotope profile through the upper kilometer of the oceanic crust, DSDP Hole 395 504B, *Earth Planet. Sci. Lett.*, 80, 217–229, doi:10.1016/0012-821X(86)90106-8, 1986.
- Alt, J. C., Zuleger, E. and Erzinger, J.: Mineralogy and stable isotopic compositions of the hydrothermally altered lower sheeted dike complex, Hole 504B, Leg 140, *Proc. ODP Sci. Res.*, 137/140, 155–166, doi:10.2973/odp.proc.sr.137140.013.1995, 1995.
- Bach, W. and Humphris, S. E.: Relationship between the Sr and O isotope compositions of hydrothermal fluids and the spreading and magma-supply rates at oceanic spreading centers, *Geology*, 27, 1067–1070, doi:10.1130/0091-7613(1999)027<1067:RBTSAO>2.3.CO;2, 1999.
- 400 Baker, E. T., Chen, Y. J. and Morgan, J. P.: The relationship between near-axis hydrothermal cooling and the spreading rate of mid-ocean ridges, *Earth Planet. Sci. Lett.*, 142, 137–145, doi:10.1016/0012-821X(96)00097-0, 1996.
- Barrett, T. J. and Friedrichsen, H.: Strontium and oxygen isotopic composition of some basalts from Hole 504B, Costa Rica Rift, DSDP Legs 69 and 70, *Earth Planet. Sci. Lett.*, 60, 27–38, doi:10.1016/0012-821X(82)90017-6, 1982.
- Bickle, M. J. and Teagle, D. A. H.: Strontium alteration in the Troodos ophiolite: implications for fluid fluxes and geochemical transport in 405 mid-ocean ridge hydrothermal systems, *Earth Planet. Sci. Lett.*, 113, 219–237, doi:10.1016/0012-821X(92)90221-G, 1992.
- Bindeman, I. N., Bekker, A. and Zakharov, D. O.: Oxygen isotope perspective on crustal evolution on early Earth: A record of Precambrian shales with emphasis on Paleoproterozoic glaciations and Great Oxygenation Event, *Earth Planet. Sci. Lett.*, 437, 101–113, doi:10.1016/j.epsl.2015.12.029, 2016.
- Bindeman, I. N., Bayon, G. and Palandri, J.: Triple oxygen isotope investigation of fine-grained sediments from major world's rivers: Insights 410 into weathering processes and global fluxes into the hydrosphere *Earth Planet. Sci. Lett.*, 528, 115851, doi:10.1016/j.epsl.2019.115851, 2019.
- Blake, R. E., Chang, S. J. and Lepland, A.: Phosphate oxygen isotopic evidence for a temperate and biologically active Archaean ocean, *Nature*, 464, 1029–1033, doi:10.1038/nature08952, 2010.
- Cathles, L. M.: An analysis of the hydrothermal system responsible for massive sulfide deposition in the Hokuroku basin of Japan, In: *The Kuroko and Related Volcanogenic Massive Sulfide Deposits* (eds. H. Ohmoto and B. J. Skinner), Society of Economic Geologists, pp. 439–487, doi:10.5382/Mono.05.27, 1983.
- Catling, D. C. and Kasting, J. F.: *Atmospheric Evolution on Inhabited and Lifeless Worlds*, Cambridge University Press, doi:10.1017/9781139020558, 2017.
- Cherkaoui, A. S. M., Wilcock, W. S. D., Dunn, R. A. and Toomey, D. R.: A numerical model of hydrothermal cooling and crustal accretion 420 at a fast spreading mid-ocean ridge, *Geochem. Geophys. Geosyst.*, 4, 8616, doi:10.1029/2001GC000215, 2003.
- Cocker, J. D., Griffin, B. J. and Muehlenbachs, K.: Oxygen and carbon isotope evidence for seawater-hydrothermal alteration of the Macquarie Island ophiolite, *Earth Planet. Sci. Lett.*, 61, 112–122, doi:10.1016/0012-821X(82)90043-7, 1982.
- Cole, D. R. and Ohmoto, H.: Kinetics of isotopic exchange at elevated temperatures and pressures, *Rev. Mineral.*, 16, 41–90, 1986.

- Cole, D. R., Ohmoto, H. and Lasaga, A. C.: Isotopic exchange in mineral-fluid systems. I. Theoretical evaluation of oxygen isotopic exchange accompanying surface reactions and diffusion, *Geochim. Cosmochim. Acta*, 47, 1681–1693, doi:10.1016/0016-7037(83)90018-2, 1983.
- Cole, D. R., Mottl, M. J. and Ohmoto, H.: Isotopic exchange in mineral-fluid systems. II. Oxygen and hydrogen isotopic investigation of the experimental basalt-seawater system, *Geochim. Cosmochim. Acta*, 51, 1523–1538, doi:10.1016/0016-7037(87)90334-6, 1987.
- Criss, R. E., Gregory, R. T. and Taylor, Jr., H. P.: Kinetic theory of oxygen isotopic exchange between minerals and water, *Geochim. Cosmochim. Acta*, 51, 1099–1108, doi:10.1016/0016-7037(87)90203-1, 1987.
- 430 DePaolo, D. J.: Isotopic effects in fracture-dominated reactive fluid-rock systems, *Geochim. Cosmochim. Acta*, 70, 1077–1096, doi:10.1016/j.gca.2005.11.022, 2006.
- Driesner, T., Seward, T. M. and Tironi, I. G.: Molecular dynamics simulation study of ionic hydration and ion association in dilute and 1 molal aqueous sodium chloride solutions from ambient to supercritical conditions, *Geochim. Cosmochim. Acta*, 62, 3095–3107, doi:10.1016/S0016-7037(98)00207-5, 1998.
- 435 Elderfield, H. and Schultz, A.: Mid-ocean ridge hydrothermal fluxes and the chemical composition of the ocean, *Annu. Rev. Earth Planet. Sci.*, 24, 191–224, doi:10.1146/annurev.earth.24.1.191, 1996.
- Elthon, D., Lawrence, J. R., Hanson, R. E. and Stern, C.: Modelling of oxygen-isotope data from the Sarmiento ophiolite complex, Chile, *Geol. Soc. Spec. Publ.*, 13, 185–197, doi:10.1144/GSL.SP.1984.013.01.16, 1984.
- Gelhar, L. W., Welty, C. and Rehfeldt, K. R.: A critical review of data on field-scale dispersion in aquifers, *Water Resour. Res.*, 28, 1955–1974, doi:10.1029/92WR00607, 1992.
- 440 Fehn, U., Green, K. E., Von Herzen, R. P. and Cathles, L. M.: Numerical models for the hydrothermal field at the Galapagos spreading center, *J. Geophys. Res.*, 88, 1033–1048, doi:10.1029/JB088iB02p01033, 1983.
- Fisher, A. T.: Permeability within basaltic oceanic crust, *Rev. Geophys.*, 32, 143–182, doi:10.1029/97RG02916, 1998.
- Frind, E. O.: Simulation of long-term transient density-dependent transport in groundwater, *Adv. Water Resour.*, 5, 73–88, doi:10.1016/0309-1708(82)90049-5, 1982.
- 445 Fry, B.: *Stable Isotope Ecology*, Springer Science & Business Media, doi:10.1007/0-387-33745-8, 2006.
- Furnes, H., de Wit, M., Staudigel, H., Rosing, M. and Muehlenbachs, K.: A vestige of Earth's oldest ophiolite, *Science*, 315, 1704–1707, doi:10.1126/science.1139170, 2007.
- Galili, N., Shemesh, A., Yam, R., Brailovsky, I., Sela-Adler, M., Schuster, E. M., Collom, C., Bekker, A., Planavsky, N., Macdonald, F. A., Préat, A., Rudmin, M., Trela, W., Sturesson, U., Heikoop, J. M., Aurell, M., Ramajo, J. and Halevy, I.: The geologic history of seawater oxygen isotopes from marine iron oxides, *Science*, 365, 469–473, doi:10.1126/science.aaw9247, 2019.
- 450 Gao, Y., Vils, F., Cooper, K. M., Banerjee, N., Harris, M., Hoefs, J., Teagle, D. A. H., Casey, J. F., Elliott, T., Laverne, C., Alt, J. C. and Muehlenbachs, K.: Downhole variation of lithium and oxygen isotopic compositions of oceanic crust at East Pacific Rise, ODP Site 1256, *Geochem. Geophys. Geosyst.*, 13, Q10001, doi:10.1029/2012GC004207, 2012.
- 455 Godderis, Y. and Veizer, J.: Tectonic control of chemical and isotopic composition of ancient oceans: the impact of continental growth, *Am. J. Sci.*, 300, 434–461, doi:10.2475/ajs.300.5.434, 2000.
- Goddéris, Y., François, L. M. and Veizer, J.: The early Paleozoic carbon cycle, *Earth Planet. Sci. Lett.*, 190, 181–196, doi:10.1016/S0012-821X(01)00377-6, 2001.
- Gregory, R. T. and Taylor, Jr., H. P.: An oxygen isotope profile in a section of Cretaceous oceanic crust, Samail ophiolite, Oman: evidence for  $\delta^{18}\text{O}$  buffering of the oceans by deep (>5km) seawater-hydrothermal circulation at mid-ocean-ridges, *J. Geophys. Res.*, 86, 2737–2755, doi:10.1029/JB086iB04p02737, 1981.
- 460



- Gregory, R. T., Criss, R. E. and Taylor, Jr., H. P.: Oxygen isotope exchange kinetics of mineral pairs in closed and open systems: applications to problems of hydrothermal alteration of igneous rocks and Precambrian iron formations, *Chem. Geol.*, 75, 1–42, doi:10.1016/0009-2541(89)90019-3, 1989.
- 465 Haar, L., Gallagher, J. and Kell, G.: NBS/NRC Steam Tables: Thermodynamic and Transport Properties and Computer Programs for Vapor and Liquid States of Water in SI Units, Hemisphere Publishing Corporation, Washington, 1984.
- Harris, K. R. and Woolf, L. A.: Pressure and temperature dependence of the self diffusion coefficient of water and oxygen-18 water, *J. Chem. Soc., Faraday Trans. 1*, 76, 377–385, doi:10.1039/F19807600377, 1980.
- Holland, H. D.: *The Chemical Evolution of the Atmosphere and Oceans*, Princeton University Press, 1984.
- 470 Holmden, C. and Muehlenbachs, K.: The  $^{18}\text{O}/^{16}\text{O}$  ratio of 2-billion-year-old seawater inferred from ancient oceanic crust, *Science*, 259, 1733–1736, doi:10.1126/science.259.5102.1733, 1993.
- Iyer, K., Rüpke, L. H. and Morgan, J. P.: Feedbacks between mantle hydration and hydrothermal convection at ocean spreading centers, *Earth Planet. Sci. Lett.*, 296, 34–44, doi:10.1016/j.epsl.2010.04.037, 2010.
- Jaffrés, J. B. D., Shields, G. A. and Wallmann, K.: The oxygen isotope evolution of seawater: A critical review of a long-  
475 standing controversy and an improved geological water cycle model for the past 3.4 billion years, *Earth-Sci. Rev.*, 83, 83–122, doi:10.1016/j.earscirev.2007.04.002, 2007.
- James, R. H., Green, D. R. H., Stock, M. J., Alker, B. J., Banerjee, N. R., Cole, C., German, C. R., Huvenne, V. A. I., Powell, A. M. and Connelly, D. P.: Composition of hydrothermal fluids and mineralogy of associated chimney material on the East Scotia Ridge back-arc spreading centre, *Geochim. Cosmochim. Acta*, 139, 47–71, doi:10.1016/j.gca.2014.04.024, 2014.
- 480 Jean-Baptiste, P., Charlou, J. L. and Stievenard, M.: Oxygen isotope study of mid-ocean ridge hydrothermal fluids: Implication for the oxygen-18 budget of the oceans, *Geochim. Cosmochim. Acta*, 61, 2669–2677, doi:10.1016/S0016-7037(97)00090-2, 1997.
- Johnson, B. W. and Wing, B. A.: Limited Archaean continental emergence reflected in an early Archaean  $^{18}\text{O}$ -enriched ocean, *Nat. Geosci.*, 13, 243–248, doi:10.1038/s41561-020-0538-9, 2020.
- Karhu, J. and Epstein, S.: The implication of the oxygen isotope records in coexisting cherts and phosphates, *Geochim. Cosmochim. Acta*,  
485 50, 1745–1756, doi:10.1016/0016-7037(86)90136-5, 1986.
- Kasting, J. F., Howard, M. T., Wallmann, K., Veizer, J., Shields, G. and Jaffrés, J.: Paleoclimates, ocean depth, and the oxygen isotopic composition of seawater, *Earth Planet. Sci. Lett.*, 252, 82–93, doi:10.1016/j.epsl.2006.09.029, 2006.
- Knauth, L. P.: Temperature and salinity history of the Precambrian ocean: implications for the course of microbial evolution, In: *Geobiology: Objectives, Concepts, Perspectives* (ed. N. Noffke), pp. 53–69, doi:10.1016/B978-0-444-52019-7.50007-3, 2005.
- 490 Knauth, L. P. and Lowe, D. R.: High Archean climatic temperature inferred from oxygen isotope geochemistry of cherts in the 3.5 Ga Swaziland Supergroup, South Africa, *Geol. Soc. Am. Bull.*, 115, 566–580, doi:10.1130/0016-7606(2003)115<0566:HACTIF>2.0.CO;2, 2003.
- Korenaga, J.: Global water cycle and the coevolution of the Earth's interior and surface environment, In: *Archean Geodynamics and Environments* (eds. K. Benn, J.-C. Mareschal and K. C. Condie), pp.7-32, doi:10.1029/164GM03, 2006.
- 495 Korenaga, J., Planavsky, N. J. and Evans, D. A. D.: Archean geodynamics and the thermal evolution of Earth, *Phil. Trans. R. Soc. A*, 375, 20150393, doi:10.1098/rsta.2015.0393, 2017.
- Krynicky, K., Green, C. D. and Sawyer, D. W.: Pressure and temperature dependence of self-diffusion in water, *Faraday Discuss. Chem. Soc.*, 66, 199–208, doi:10.1039/DC9786600199, 1978.

- Lawrence, J. R. and Gieskes, J. M.: Constraints on water transport and alteration in the oceanic crust from the isotopic composition of porewater, *J. Geophys. Res.*, 86, 7924–7934, doi:10.1029/JB086iB09p07924, 1981.
- Lécuyer, C. and Allemand, P.: Modelling of the oxygen isotope evolution of seawater: Implication for the climate interpretation of the  $\delta^{18}\text{O}$  of marine sediments, *Geochim. Cosmochim. Acta*, 63, 351–361, doi:10.1016/S0016-7037(98)00277-4, 1999.
- Lécuyer, C. and Fourcade, S.: Oxygen isotope evidence for multi-stage hydrothermal alteration at a fossil slow-spreading center: the Silurian Trinity ophiolite (California, U.S.A.), *Chem. Geol.*, 87, 231–246, doi:10.1016/0168-9622(91)90023-P, 1991.
- 505 Meyer, C. A., McClintock, R. B., Silvestri, G. J. and Spencer, R. C.: AME Steam Tables: Thermodynamic and Transport Properties of Steam, Fifth Edition, American Society of Mechanical Engineers, 1983.
- Muehlenbachs, K.: The alteration and aging of the basaltic layer of the seafloor: Oxygen isotope evidence from DSDP/IPOD Legs 51, 52, and 53, *Int. Repts. DSDP*, 51, 1159–1167, doi:10.2973/dsdp.proc.515253.142, 1979.
- Muehlenbachs, K.: The oxygen isotopic composition of the oceans, sediments and the seafloor, *Chem. Geol.*, 145, 263–273, doi:10.1016/S0009-2541(97)00147-2, 1998.
- 510 Muehlenbachs, K. and Clayton, R. N.: Oxygen isotope composition of the oceanic crust and its bearing on seawater, *J. Geophys. Res.*, 81, 4365–4369, doi:10.1029/JB081i023p04365, 1976.
- Muehlenbachs, K., Furnes, H., Fonneland, H. C. and Hellevang, B.: Ophiolites as faithful records of the oxygen isotope ratio of ancient seawater: the Solund–Stavfjord Ophiolite Complex as a Late Ordovician example, *Geol. Soc. Spec. Publ.*, 218, 401–414, doi:10.1144/GSL.SP.2003.218.01.20, 2003.
- 515 Nielsen, M. E. and Fisk, M. R.: Surface area measurements of marine basalts: Implications for the subseafloor microbial biomass, *Geophys. Res. Lett.*, 37, L15604, doi:10.1029/2010GL044074, 2010.
- Norton, D. and Knight, J.: Transport phenomena in hydrothermal systems: cooling plutons, *Am. J. Sci.*, 277, 937–981, doi:10.2475/ajs.277.8.937, 1977.
- 520 Norton, D. and Taylor, Jr., H. P.: Quantitative simulation of the hydrothermal systems of crystallizing magmas on the basis of transport theory and oxygen isotope data: An analysis of the Skaergaard intrusion, *J. Petrol.*, 20, 421–486, doi:10.1093/petrology/20.3.421, 1979.
- Perry, Jr., E. C., Ahmad, S. N. and Swulius, T. M.: The oxygen isotope composition of 3,800 m.y. old metamorphosed chert and iron formation from Isukasia, West Greenland, *J. Geol.*, 86, 223–239, doi:10.1086/649676, 1978.
- Phipps Morgan, J.: Thermal and rare gas evolution of the mantle, *Chem. Geol.*, 145, 431–445, doi:10.1016/S0009-2541(97)00153-8, 1998.
- 525 Schiffman, P. and Smith, B. M.: Petrology and oxygen isotope geochemistry of a fossil seawater hydrothermal system within the Solea graben, northern Troodos ophiolite, Cyprus, *J. Geophys. Res.*, 93, 4612–4624, doi:10.1029/JB093iB05p04612, 1988.
- Shemesh, A., Kolodny, Y. and Luz, B.: Oxygen isotope variations in phosphate of biogenic apatites, II. Phosphorite rocks, *Earth Planet. Sci. Lett.*, 64, 405–416, doi:10.1016/0012-821X(83)90101-2, 1983.
- Stakes, D. S.: Oxygen and hydrogen isotope compositions of oceanic plutonic rocks: High-temperature deformation and metamorphism of oceanic layer 3, In: *Stable Isotope Geochemistry: A Tribute to Samuel Epstein* (eds. H. P. Taylor, J. R. O’Neil and I. R. Kaplan), The Geochemical Society, pp. 77–90, 1991.
- 530 Steefel, C. I. and Lasaga, A. C.: A coupled model for transport of multiple chemical species and kinetic precipitation/dissolution reactions with application to reactive flow in single phase hydrothermal systems, *Am. J. Sci.*, 294, 529–592, doi:10.2475/ajs.294.5.529, 1994.
- Stein, C. A. and Stein, S.: A model for the global variation in oceanic depth and heat flux with lithospheric age, *Nature*, 359, 123–129, doi:10.1038/359123a0, 1992.
- 535

- Stein, C. A. and Stein, S.: Constraints on hydrothermal heat flux through the oceanic lithosphere from global heat flow, *J. Geophys. Res.*, 99, 3081–3095, doi:10.1029/93JB02222, 1994.
- Taylor, Jr., H. P.: Water/rock interactions and the origin of H<sub>2</sub>O in granitic batholiths, *J. Geol. Soc. London*, 133, 509–558, doi:10.1144/gsjgs.133.6.0509, 1977.
- 540 V  rard, C. and Veizer, J.: On plate tectonics and ocean temperatures, *Geology*, 47, 881–885, doi:10.1130/G46376.1, 2019.
- Vibetti, N. J., Kerrich, R. and Fyfe, W. S.: Oxygen and carbon isotope studies of hydrothermal alteration in the Troodos ophiolite complex, Cyprus, *Geol. Surv. Canada Spec. Pap.*, 88-9, 221–228, 1989.
- Walker, J. C. G. and Lohmann, K. C.: Why the oxygen isotopic composition of sea water changes with time, *Geophys. Res. Lett.*, 16, 323–326, doi:10.1029/GL016i004p00323, 1989.
- 545 Wallmann, K.: The geological water cycle and the evolution of marine  $\delta^{18}\text{O}$  values, *Geochim. Cosmochim. Acta*, 65, 2469–2485, doi:10.1016/S0016-7037(01)00603-2, 2001.
- Wallmann, K., Aloisi, G., Haeckel, M., Tishchenko, P., Pavlova, G., Greinert, J., Kutterolf, S. and Eisenhauer, A.: Silicate weathering in anoxic marine sediments, *Geochim. Cosmochim. Acta*, 72, 3067–3090, doi:10.1016/j.gca.2008.03.026, 2008.
- Wolery, T. J. and Sleep, N. H.: Hydrothermal circulation and geochemical flux at mid-ocean ridges, *J. Geol.*, 84, 249–275, doi:10.1086/628195, 1976.
- 550 Zhao, Z. -F. and Zheng, Y. -F.: Calculation of oxygen isotope fractionation in magmatic rocks, *Chem. Geol.*, 193, 59–80, doi:10.1016/S0009-2541(02)00226-7, 2003.

Table 1. Symbols and their definitions and values.

Symbol	Definition	Value
$c_p^f$	Specific heat capacity of water ( $\text{J kg}^{-1} \text{K}^{-1}$ )	
$c_p^m$	Specific heat capacity of oceanic rock ( $\text{J kg}^{-1} \text{K}^{-1}$ )	$10^3$
$D$	Effective diffusion coefficient for $^{18}\text{O}$ ( $\text{m}^2 \text{yr}^{-1}$ )	
$E$	Apparent activation energy of oxygen isotope exchange ( $\text{J mol}^{-1}$ )	$5 \times 10^4$
$F_m$	Mole ratio of $^{18}\text{O}$ to total O ( $^{18}\text{O}$ plus $^{16}\text{O}$ ) of pristine crust (dimensionless)	$2.0126 \times 10^{-3}$
$F_p$	Mole ratio of $^{18}\text{O}$ to total O ( $^{18}\text{O}$ plus $^{16}\text{O}$ ) of porewater (dimensionless)	
$F_r$	Mole ratio of $^{18}\text{O}$ to total O ( $^{18}\text{O}$ plus $^{16}\text{O}$ ) of solid rock (dimensionless)	
$F_{sw}$	Mole ratio of $^{18}\text{O}$ to total O ( $^{18}\text{O}$ plus $^{16}\text{O}$ ) of seawater (dimensionless)	
$g$	Gravity acceleration ( $\text{m yr}^{-2}$ )	$9.76 \times 10^{15}$
$\mathbf{g}$	Gravity vector ( $\text{m yr}^{-2}$ )	
$J_b$	Boundary heat flux from the intrusion ( $\text{J m}^{-2} \text{yr}^{-1}$ )	
$k$	Permeability of oceanic rock ( $\text{m}^2$ )	
$k_{ex}$	Rate constant for oxygen isotope exchange between solid rock and porewater ( $\text{mol}^{-1} \text{kg yr}^{-1}$ )	
$m_f$	Mole concentration of oxygen per unit water mass ( $\text{mol kg}^{-1}$ )	55.56
$m_s$	Mole concentration of oxygen per unit solid mass ( $\text{mol kg}^{-1}$ )	31.25
$P$	Fluid pressure (Pa)	
$\mathbf{q}$	Water mass flux vector ( $\text{kg m}^{-2} \text{yr}^{-1}$ )	
$R_g$	Gas constant ( $\text{J mol}^{-1} \text{K}^{-1}$ )	8.314
$t$	Time (yr)	
$T$	Temperature (K)	
$T_m$	Temperature of the intrusion (K)	$1.473 \times 10^3$
$w$	Spreading rate of oceanic crust ( $\text{m yr}^{-1}$ )	
$x$	Distance from ridge axis (m)	
$y$	Depth from ocean/crust interface (m)	
$\alpha$	Oxygen isotope fractionation factor (dimensionless)	
$\beta$	Parameter relevant to the temperature dependence of $\alpha$ (dimensionless)	0.876
$\eta$	Water/rock oxygen-mole ratio (dimensionless)	
$\kappa$	Thermal conductivity of oceanic rock ( $\text{J yr}^{-1} \text{m}^{-1} \text{K}^{-1}$ )	$9.47 \times 10^7$
$\mu$	Kinematic viscosity of water ( $\text{m}^2 \text{yr}^{-1}$ )	
$\rho_b$	Density of bulk rock ( $\text{kg m}^{-3}$ )	
$\rho_f$	Density of water ( $\text{kg m}^{-3}$ )	
$\rho_m$	Density of solid rock ( $\text{kg m}^{-3}$ )	$3 \times 10^3$
$\tau$	Tortuosity factor (dimensionless)	
$\phi$	Porosity of crust (dimensionless)	$5 \times 10^{-2}$
$\Omega$	Degree of oxygen isotope exchange (dimensionless)	
$\nabla$	Vector differential operator ( $\text{m}^{-1}$ )	




Figure 1. Two-dimensional distributions of hydrothermal fluid flow (a and c) and temperature (b and d). Shown in a and c are logarithms of fluid velocity ( $\text{m yr}^{-1}$ ) together with mass-based stream lines that are depicted with white curves. The same data as in a and b are plotted on logarithmic scales in c and d, respectively. Gray zones in a and c represent where rocks are impermeable below 6 km depth and/or with temperatures above the rock-cracking threshold ( $600\text{ }^{\circ}\text{C}$ ).




Figure 2. Two-dimensional distributions of solid-rock and porewater  $\delta^{18}\text{O}$  (a–c and d–f, respectively) at seawater  $\delta^{18}\text{O}$  values of 0, -6 and -12 ‰ (a and d, b and e, and c and f, respectively).




Figure 3. As for Fig. 2, except plotted on logarithmic scales.




Figure 4. Solid rock  $\delta^{18}\text{O}$  as function of depth at 1 Myr (30 km) from the ridge axis at seawater  $\delta^{18}\text{O}$  values of 6, 4, ..., -12 ‰. Also plotted are Oman ophiolite data from Gregory and Taylor (1981) (crosses) converting their reported depth  $y'$  (km) to  $(7.4 - y')$  (km) to facilitate comparison.






Figure 5. Two-dimensional distributions of local water/rock oxygen-mole ratio (a and c) and degree of oxygen isotope exchange at a seawater  $\delta^{18}\text{O}$  value of 0‰ (b and d). The same data as in a and b are plotted on logarithmic scales in c and d, respectively. See the caption of Fig. 1 for the explanation of gray zones in a and c.




Figure 6. Net  $^{18}\text{O}$  flux to the ocean from hydrothermal systems as function of seawater  $\delta^{18}\text{O}$ . Slope values are denoted near the lines from previous studies and the standard simulation in this study.




Figure 7. Two-dimensional distributions of hydrothermal fluid flow (a, c, e, g and i) and temperature (b, d, f, h and j) from simulations with different spreading rates and ocean depths. Logarithms of fluid velocity and mass-based stream lines are depicted in a, c, e, g and i. The panels in the first three rows assume different spreading rates ( $1 \times 10^{-2} \text{ m yr}^{-1}$  in a and b,  $9 \times 10^{-2} \text{ m yr}^{-1}$  in c and d, and  $30 \times 10^{-2} \text{ m yr}^{-1}$  in e and f), while those in the last two rows assume different ocean depths (1 km in g and h and 5 km in i and j). Values of other parameters are the same as those in the standard simulation (e.g., Fig. 1). See the caption of Fig. 1 for the explanation of gray zones in a, c, e, g and i.




Figure 8. Solid rock  $\delta^{18}\text{O}$  as function of depth at 30 km from the ridge axis at seawater  $\delta^{18}\text{O}$  values of 6, 4, ..., -12 ‰ of from simulations with different spreading rates and ocean depths. Dashed lines denote 1 and 16 ‰, between which observed  $\delta^{18}\text{O}$  of ophiolites and/or oceanic crust ranges (Gregory and Taylor, 1981; Barrett and Friedrichsen, 1982; Cocker et al., 1982; Elthon et al., 1984; Alt et al., 1986, 1995; Agrinier et al., 1988; Schiffman and Smith, 1988; Vibetti et al., 1989; Lécuyer and Fourcade, 1991; Stakes, 1991; Bickle and Teagle, 1992; Holmden and Muehlenbachs, 1993; Muehlenbachs et al., 2003; Alt and Bach, 2006; Furnes et al., 2007; Gao et al., 2012). Standard parameter values are assumed except for the parameterization denoted at each panel.




Figure 9. Net  $^{18}\text{O}$  flux to the ocean from hydrothermal systems as function of seawater  $\delta^{18}\text{O}$  from simulations with different spreading rates and ocean depths. See the legend for the types of symbol and line for individual simulations. Slope values are denoted near the lines.