



Slab Break-offs in the Alpine Subduction Zone

Emanuel D. Kästle¹, Claudio Rosenberg², Lapo Boschi^{2,3}, Nicolas Bellahsen², Thomas Meier⁴, and Amr El-Sharkawy^{4,5}

¹Institute of Geological Sciences, Freie Universität Berlin, Germany

²Institut des Sciences de la Terre, Paris (iSTeP), Sorbonne Université, Paris, France

³Dipartimento di Geoscienze, Università degli Studi di Padova, Italy

⁴Christian Albrechts Universität Kiel, Germany

⁵National Research Institute of Astronomy and Geophysics (NRIAG), 11421, Helwan, Cairo, Egypt

Correspondence: E.D. Kästle (emanuel.kaestle@fu-berlin.de)

Abstract. After the onset of plate collision in the Alps, tectonic processes are inferred to have changed dramatically in the Alps: European plate break-offs in various places of the Alpine arc, as well as a subduction polarity reversal in the eastern Alps have been proposed. We review body-wave tomographic studies, compare them to our surface-wave-derived model, and interpret them in terms of slab geometries. We infer that the shallow subducting portion of the European plate is likely detached under both the western and eastern (but not the central) Alps. The Alps-Dinarides transition may be explained by a combination of European and Adriatic subduction. This implies that the deep high-velocity anomaly (>200 km depth) mapped by tomographers under the eastern Alps is a detached segment of the European plate. The shallower fast anomaly (100–200 km depth) can be ascribed to European or Adriatic subduction, or both. These findings are compared to previously proposed models for the eastern Alps in terms of slab geometry, but also integrated in a new, alternative geodynamic scenario that best fits both tomographic images and geological constraints.

Copyright statement.

1 Introduction

The Alps and the adjacent Apenninic, Dinaric and Carpathian mountain chains were formed by a complex history of subductions and collisions. During N-S convergence of Africa and Europe (e.g., Stampfli and Borel, 2002), between 84 Ma and 35 Ma, Adria-Africa acted as upper plate along most of the subduction front and the Alpine Tethys (Europe) was subducted towards the south (e.g., Schmid et al., 2004). An opposite subduction direction only affected what would later become the Dinaric-Hellenic arc (Laubscher, 1971; Pamić et al., 2000; Schmid et al., 2008; Ustaszewski et al., 2008, Fig. 1). During the transition between subduction and collision, slab break-offs are inferred to have occurred in the Alpine and Dinaric collision zones (e.g., Blanckenburg and Davies, 1995; Lippitsch et al., 2003; Harangi et al., 2006; Kissling et al., 2006), partly preceding reversals of subduction polarity, as in the Apenninic collision (Vignaroli et al., 2008; Handy et al., 2010; Molli and Malavieille, 2011) and possibly in the eastern Alps (Schmid et al., 2013). In this area, tomography results (Lippitsch et al., 2003) triggered a



discussion on the apparent northward dip of the slab between 50 and 250 km depth, suggesting that it may result from subduction of Adria under Europe (Schmid et al., 2004; Ustaszewski et al., 2008; Schmid et al., 2013; Handy et al., 2015), hence to a change in subduction polarity, after the onset of collision. This change would have been possible after the proposed break-off of the European slab, giving way to a northward indentation of the Adriatic crust and a subduction of its mantle (e.g., Schmid et al., 2013; Handy et al., 2015).

At present, the high-resolution regional tomographic models of the Alpine upper mantle disagree on important structures, such as the suggested present-day detachment of the European slab under the western Alps and the subduction polarity under the eastern Alps (Adriatic vs. European subduction) (Lippitsch et al., 2003; Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). An important limiting factor in the discussion on the structure of the Alpine slabs is the poor resolution in the uppermost mantle layer of teleseismic body-wave models, due to almost vertical ray paths and lower data coverage (e.g., Boschi et al., 2010). In the light of a new model which does not suffer from this limitation (Kästle et al., 2018), we discuss the Alpine slab geometries and their possible detachments. The latter model is reliable down to a depth of approximately 200 km with teleseismic body-wave models for the deeper parts of the sections. We apply this approach to a suite of published body-wave tomographic models of the Alpine area and illustrate the results along four orogen-perpendicular cross sections that allow one to systematically compare the results of different tomographic investigations. Such a thorough, quantitative comparison cannot currently be found in the literature; yet, it is a necessary step to establish what really is known of mantle structure under the Alpine belt. We interpret and discuss surface-wave and body-wave models, while also taking into account the constraints provided by estimates of crustal shortening.

2 Surface-wave Tomography

Surface-wave phase velocities are particularly sensitive to the vertical changes in shear-wave velocities, therefore they are well suited for probing the lithosphere and the asthenosphere. In a recent study, Kästle et al. (2018) combined phase-velocity-dispersion measurements from ambient-noise and earthquake data which resulted in a very broad frequency range (4 -300 seconds) of measurements, and an unprecedentedly dense ray-path coverage of the region. From the inversion of the phase-velocity measurements, we constructed a high resolution 3D shear-wave-velocity model underneath the Alpine region, showing that a 100 km thick anomaly (as from a lithospheric slab) can be identified down to 200 km depth, although the strength of the anomaly may weaken at depth. The shear-velocity model of Kästle et al. (2018), shown here in Fig. 1, thus provides an unique opportunity to study the crust-mantle transition and the approximate position and geometry of subduction slabs, under the Alps and adjacent orogens, despite the absence of deep earthquakes in this region.

The clearest velocity anomaly is obtained from the thick, vertical European slab under the central Alps (Fig. 1). In contrast, the Adriatic slab underneath Apennines and Dinarides is associated with a weaker high-velocity anomaly, which may be due to a thinner slab, or lower resolution, because the station distribution is not perfect. The intensity of the anomaly is different in the western, central and eastern Alps, which suggests that these three domains experienced a different evolution since the onset of collision around 35–30 Ma.




Figure 1. Map: Shear-velocity model from Kästle et al. (2018), showing deviations with respect to PREM (Dziewonski and Anderson, 1981). AF: Alpine frontal thrust, PF: Periadriatic fault, GF: Giudicarie fault. Faults and tectonic limits (red lines) simplified from Schmid et al. (2004, 2008); Handy et al. (2010). The white area marks the low-resolution region of the model. Cross-sections: Absolute shear velocity in the crust and PREM-deviations in the mantle. Moho boundaries from Spada et al. (2013). EU: Europe, AD: Adria, LPT: Liguro-Provençal and Tyrrhenian basins. Mantle anomalies: ApA: Apenninic Anomaly, CA: Central Alpine Anomaly, DA: Dinaric Anomaly, EAA: Eastern Alpine Anomaly, EANA: Eastern Alpine Northern Anomaly, WAA: Western Alpine Anomaly.

Central Alps: The strongest and most continuous high-velocity anomaly is found under the central Alps. It extends from the bottom of the crust down to the lower boundary of the mapped region at 200 km depth, suggesting a continuous slab of at least 150 km length (Fig. 1B). The slab anomaly extends in E–W direction from 7° to around 12.5°, with a thickness of up to 100 km and a (sub-) vertical dip.

- 5 **Western Alps:** In contrast to the central Alps, there is no clear, continuous high-velocity anomaly under the western Alps in our shear-velocity model (Fig. 1A). The fast, lithospheric velocity anomaly under the western Alps (European) vanishes at depths below 90 – 120 km. It is separated from the slab under the Apennines (Adriatic) by a low-velocity anomaly (Fig. 1A), suggesting the presence of a shallow European slab break-off.

- 10 **Eastern Alps:** The central Alpine anomaly extends eastward to 12–13°E, coinciding approximately with the Giudicarie fault (Fig. 1). At this longitude, we observe a sharp change in the amplitude of the high-velocity anomaly: further east, the strong, fast anomaly disappears below about 80 km in our surface-wave model (Fig. 1C,D). Below this depth, a region of more



diffuse, slightly elevated velocities (Fig. 1C) and no clear separation between eastern Alpine and Dinaric domain is observed. In addition, a separate high-velocity anomaly (EANA, profile 1C) is found more to the north beneath the European foreland at depth of about 100–150 km. The European slab under the eastern Alps must be dramatically thinned, or even broken off to explain the latter observations. Alternatively, the high-velocity anomaly under the eastern Alps at 60 – 200 km depth may correspond to an Adriatic slab, which generally shows a lower velocity anomaly strength underneath the Apennines in our model. The anomaly strength shows a small increase towards the Dinarides. Under the northern Dinarides, we interpret the anomaly attaining approx. 150 km depth to be of Adriatic origin (map view and DA in section 1D).

3 Body-wave Tomography

Using the tomographic models kindly provided by their authors (Lippitsch et al., 2003; Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017), we define 4 section traces and plot these sections from each model, applying the same velocity color map to each model (Fig. 2), hence allowing for an easy visual comparison. For the sake of clarity, we selected only three models for Figure 2 and present all other models in the supplementary material (Figs. S2–S10). The early model of Lippitsch et al. (2003) was selected because it has been and still is the reference (e.g., Handy et al., 2015); the model of Koulakov et al. (2009), because it covers much of Europe, allowing interpretations that go beyond the Alpine region; the model of Hua et al. (2017) is the most recent one which covers the entire Alps and also includes local-earthquake data. These models also illustrate the differences between different data sets and methods.

Western Alps: In the western Alps, Lippitsch et al. (2003) propose a slab break-off at lithospheric depth (Fig. 2A). Such a break-off is also proposed by the model of Beller et al. (2017), using full-waveform inversion along a dense seismic profile. However, others observe a continuity between the deeper anomaly and the lithospheric one (Koulakov et al., 2009; Zhao et al., 2016; Hua et al., 2017). At depths greater than 150 km, all models agree on a broad, fast anomaly (Fig. 2), which may represent a combination of European and Adriatic (from the Apenninic subduction) slabs (Zhao et al., 2016).

Central Alps: All body-wave models show a continuous slab to at least 200 km depth under the central Alps (Lippitsch et al., 2003; Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). Below this depth, the slab is thinned (Lippitsch et al., 2003; Mitterbauer et al., 2011; Hua et al., 2017), possibly indicating a break-off. It may also be merged with the Apenninic slab (Fig. 2B), which would be indistinguishable from a continuous slab to ≥ 400 km depth as proposed by Zhao et al. (2016). Towards the east, the slab anomaly is limited in some models by a slab gap (Lippitsch et al., 2003; Koulakov et al., 2009) or a northward step of the anomaly (maps in Fig. 2, Fig. S1, Zhao et al., 2016).

Eastern Alps: An eastward increase in the northward slab dip by Lippitsch et al. (2003), while the other models (Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017) show a sub-vertical, though slightly northdipping slab (Fig. 2C,D, S8–S10). Lippitsch et al. (2003) image the slab down to a depth of about 250 km (Fig. 2D). Dando et al. (2011) indicate that the slab may be more continuous (Fig. S8–S10) and linked vertically to a deep anomaly (>410 km) under the Pannonian Basin. A continuation of the fast anomaly down to at least 350 km, is also visible in models of Koulakov et al. (2009), Mitterbauer et al. (2011), Zhao et al. (2016) and Hua et al. (2017) (Figs. 2C,D, S8–S10). The latter




Figure 2. Comparison of teleseismic body-wave tomographic models along four cross-sections. The models show compressional velocity anomalies with respect to PREM (Dziewonski and Anderson, 1981). AF: Alpine frontal thrust, PF: Periadriatic fault, GF: Giudicarie fault. Faults and tectonic limits (red lines) simplified from Schmid et al. (2004, 2008); Handy et al. (2010). The crustal segments are attributed to the different domains according to Spada et al. (2013). EU: European plate; AD: Adriatic plate; LPT: Liguro-Provençal and Tyrrhenian domains. The white band in the Lippitsch et al. (2003) sections marks the shallow region which is not shown in the original model.



shows almost no fast anomaly in the upper 200 km, but a clear anomaly below, opposite to the model of Lippitsch et al. (2003) (Fig. 2D). Horizontal sections at 150 km (Fig. 2, S1) show that the eastern Alpine anomaly is separated from the central Alpine one by a slab gap (Lippitsch et al., 2003) or a northward step, spatially coinciding with the Giudicarie fault (Mitterbauer et al., 2011; Zhao et al., 2016).

- 5 An additional fast anomaly, whose maximum strength is at approximately 150 km depth is present north of the eastern Alps (Lippitsch et al., 2003; Zhao et al., 2016), similar to the EANA in the surface-wave model (Fig. 1).

At the transition between eastern Alps and Dinarides (Fig. 2D), Lippitsch et al. (2003) indicate a shallow anomaly (~ 100 km) under the northern Dinarides and a deeper one (~ 200 km) under the eastern Alps. These two anomalies seem to be laterally linked, indicating that the Adriatic slab from the northern Dinarides might extend into the area underneath the eastern Alps. A

- 10 similar link between northern Dinaric and eastern Alpine slabs is imaged by Dando et al. (2011) (Fig. S10). The other models do not show such a connection, but a deep anomaly, located below 300 km and detached from the lithospheric anomalies (Figs. 2D, S10 at 200 km distance; Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017).

4 Discussion

4.1 Western Alps

- 15 Based on our surface-wave model (Fig. 1A), the European slab under the western Alps shows a shallow break-off at approximately 100 km depth. This agrees with the models of Beller et al. (2017) (full-waveform modeling) and Lippitsch et al. (2003, body-wave tomography), who locate the gap between 80 and 150 km depth. In contrast, other tomographic models image a continuous slab down to at least 250 km depth (Koulakov et al., 2009; Zhao et al., 2016; Hua et al., 2017; Lyu et al., 2017). Zhao et al. (2016) use a synthetic test to argue that the thin (~ 50 km thick) link between European lithosphere and subducting
- 20 slab is a robust feature. Errors in crustal corrections or smearing through typical ray path geometries could, however, still prevent the detection of a gap in the downgoing slab.

At greater depth, Lippitsch et al. (2003) interpret the fast anomaly as the detached European slab, dipping towards the east. Zhao et al. (2016) argue that the western European Alpine slab and the Adriatic slab from the northern Apenninic subduction are too close to be distinguishable in tomographic imaging. This would create the apparent northward dip of the combined

- 25 slab under the northernmost part of the Apennines. Also in other models, there is no visible separation between European and Adriatic slab (Koulakov et al., 2009; Hua et al., 2017, Fig 2). Our surface-wave model, below approx. 100 km, can only resolve a single anomaly under the Apennines that we attribute to the Adriatic slab, because of its location and continuity with the lithosphere under the Apennines.

- 30 We favor the existence of a shallow slab gap in the western Alps which could also explain the observed surface uplift in the western Alps (Escher and Beaumont, 1997; Nocquet et al., 2016), due to the cease of slab pull. If the slab broke off 35 Ma ago, as inferred on the base of geochemical interpretations (Blanckenburg and Davies, 1995), it should now be at least at 350 km depth, taking low sinking velocities of 1 cm/yr into account (Capitanio et al., 2007; Replumaz et al., 2010). However, the geometry of the fast anomaly in some body-wave models between 150–300 km depth (Fig. 2A, S2–S4) suggest that it could



be related to European subduction which would be associated to a more recent (≤ 10 Ma) break-off. Although the resolution is limited, it is clear that the shallow attached European lithosphere and the subduction Adriatic lithosphere beneath the northern Apennines are located very close to each other and may have already collided.

4.2 Central Alps

- 5 The central Alpine slab appears as a thick and almost vertical anomaly in all models down to at least 200 km depth (Fig. 1B, 2B). The vertical termination of the slab around 200 km depth in most models (Lippitsch et al., 2003; Koulakov et al., 2009; Mitterbauer et al., 2011; Hua et al., 2017), points to a slab length of approximately 150 km, closely matching interpreted amounts of convergence since 35 Ma (Schmid et al., 2004). The portion of inferred European shortening (63 km: Schmid et al. (1996), 30–95 km Rosenberg et al. (2015)), suggests that break-off must have happened at a depth greater than 100 km to
- 10 explain the slab depth of ≥ 200 km. Recent results propose, however, that the shortening in the European plate may be in the range of 150 km (Rosenberg et al., 2019). We conclude that the fast anomaly in the top 200 km is caused by syn-collisional subduction of the European plate. Below this depth, a break-off is likely, given the lack of a clear signal in the body-wave images. The deeper anomalies shown by Koulakov et al. (2009), Zhao et al. (2016) and Hua et al. (2017) are roughly located under the Po-basin and could represent the broken-off European slab, or alternatively the Adriatic slab from the Apenninic
- 15 subduction.

4.3 Eastern Alps

- The classical interpretation (e.g. Hawkesworth et al., 1975; Lüschen et al., 2004) of a south-directed oceanic subduction followed by a south-directed continental subduction of Europe below Adria was questioned by Lippitsch et al. (2003), who
- 20 first argued in favor of a continental subduction of Adria below Europe, based on the imaged N-directed slab dip. This is also observed by other authors who interpret a vertically continuous Adriatic slab down to at least 400 km depth (Zhao et al., 2016; Hua et al., 2017). Mitterbauer et al. (2011) observe a more vertically oriented anomaly compared to the Lippitsch et al. (2003) model, hence arguing that the apparent dip direction is no reliable indication of Adriatic subduction. Dando et al. (2011) infer that the deepest part of the slab can be followed almost continuously into the deep seated high-velocity anomaly below the
- 25 Pannonian Basin that they interpret as remnant of the subducted European plate. Hence the imaged slab would rather be of European origin.

In order to assess the likelihood of these different scenarios we discuss below: (1) dip direction of the slab (2) continuity of the imaged slab with either the Adriatic or European plate and Moho (3) lateral discontinuity between central and eastern Alpine slabs, (4) vertical continuity of the slab, (5) geological constraints on the amounts of collisional shortening.

- 30 (1) Slab dip: the Alpine slabs are mostly sub-vertical, with dipping angles between 70 and 90° (Lippitsch et al., 2003; Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017; Kästle et al., 2018). The resolution limit and the geometry of the ray paths may furthermore introduce artifacts that bias the apparent dipping angle. If the entire slab is taken into account, a slight northward dip is visible in many sections through the eastern Alps (Lippitsch et al., 2003;



Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). Along the Dinarides-Alps transition, the apparent dip angle may be influenced by imaging northern Dinaric and eastern Alpine slabs at the same time, thus enhancing the northeastern dip direction of the anomaly (Fig. 2D).

(2) Moho structure: attributing the high-seismic-velocity, slab-like anomaly under the eastern Alps to either the European or Adriatic plate is difficult as teleseismic body-wave tomography suffers from smearing along ray paths in the uppermost mantle and is strongly dependent on crustal velocity corrections. Additionally, the Moho structure itself suffers from uncertainty in the eastern Alps and gives no clear hint on which plate is on top (e.g., Lüschen et al., 2004; Spada et al., 2013). The most recent work along the EASI dense-station profile (along 13° longitude) suggest that the Adriatic Moho goes below Europe, although the uncertainty is shown to be high (Hetényi et al., 2018b). Thus we do not find a reliable link between either European or Adriatic Moho and the slab anomalies below (Fig. 2C,D), and the surface-wave tomography does not provide any clear indications either (Fig. 1).

(3) Lateral continuity: the absence of a lateral continuity between central and eastern Alpine slabs is observed in all models (Lippitsch et al., 2003; Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017; Kästle et al., 2018). There is either a horizontal slab gap (Fig. 2 Lippitsch et al., 2003) or a northward step of the anomaly (Fig. 2 Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016). This discontinuity may be linked to the sinistral Giudicarie fault (e.g., Zhao et al., 2016).

(4) Vertical continuity: the eastern Alpine slab terminates sharply at 250 km depth in the model of Lippitsch et al. (2003, Fig. 2C,D), but all other models image also deeper anomalies (Fig. 2C,D, S8–S10; Koulakov et al., 2009; Mitterbauer et al., 2011; Dando et al., 2011), possibly in continuity to the shallow slab (Zhao et al., 2016; Hua et al., 2017). East of 14° longitude, several models indicate that the shallow part of the slab disappears and only a deep slab below 250 km is visible (Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). Our surface-wave model does not cover the deeper structures, but shows a weak but continuous fast anomaly in the top 200 km (Fig. 1).

(5) Shortening constraints: the total convergence since 25 Ma is estimated in the order of 190 km in the eastern Alps by assuming a 20° counter-clockwise rigid-body rotation of Adria (Ustaszewski et al., 2008). However, collisional shortening estimates provide minimum estimates of 75 km in the European plate (Rosenberg et al., 2017) and 50 km in the Adriatic plate (Schönborn, 1999; Nussbaum, 2000).

Based on the presented arguments we discuss four scenarios for the eastern Alps, three of which are taken from the literature and a fourth one in order to reconcile the presented findings (Fig. 3):

30 **Scenario (I): Continuous European slab down to at least 250 km**

This scenario represents a situation analogue to the one inferred for the central Alps and was the accepted one before the work of Lippitsch et al. (2003) started a discussion on a possible subduction of Adria under the eastern Alps. This scenario can explain the apparent vertical continuity of the slab, reaching down to depths ≥ 350 km as shown in several models (Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016). Only Lippitsch et al. (2003) show a shorter slab down to 250 km, which might suggest that the slab is broken-off at this depth.




Figure 3. Schematic illustration of the four discussed scenarios. The sections cross the eastern Alps approximately along profile C in Figs. 1 and 2.

However, scenario (I) cannot explain the above discussed lateral discontinuity. The surface-wave model shows a significant difference in anomaly strength when comparing central and eastern Alps, suggesting that the European slab is strongly thinned or detached in the uppermost mantle layer of the eastern Alps. This becomes also evident from the body-wave models, east of 14° longitude, where there is no high-velocity anomaly in the top 250 km (Koulakov et al., 2009; Dando et al., 2011; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). The slight northward dip of the slab in almost all models is another argument against European subduction, although it remains ambiguous and the slab could be overturned.

Scenario (II): Continuous Adriatic slab down to 400 km

Zhao et al. (2016), Hua et al. (2017) and (Hetényi et al., 2018b) propose that the vertically continuous and slightly northward dipping fast anomaly under the eastern Alps could represent a long Adriatic slab. We discard this possibility, because of the lack of geologic evidence of a long-lasting Adriatic subduction in the eastern Alps (Handy et al., 2010, and references therein) and the aforementioned estimates of collisional shortening that are by far too small to explain a continuous Adriatic slab down to the mantle transition zone.

Scenario (III): Adriatic subduction down to 250 km

This scenario (Lippitsch et al., 2003) requires a European slab break off and the onset of Adriatic subduction, or an extension of the Adriatic subduction from the Dinarides into the eastern Alps. Adriatic subduction would have been preceded by European slab tear and eastwards retreat, linked to the opening of the Pannonian Basin and the lateral movement along the Giudicarie Fault, which opened up the necessary space for the Adriatic slab (Schmid et al., 2004; Handy et al., 2015). This scenario can thus explain the lateral slab discontinuity from central to eastern Alps, evidenced in all tomographic models, approximately coinciding spatially with the Giudicarie Fault. It remains unclear whether such a European slab tear is related to the 32-30 Ma magmatism (Rosenberg, 2004) or whether a younger break-off is necessary to enable post-20 Ma Adriatic subduction (Handy et al., 2015).




Figure 4. Schematic model of a possible configuration of European and Adriatic plates as presented in Scenario (IV). Question marks indicate very speculative locations of the European broken-off slabs. Most of the Apenninic slab is cut out from the sketch to improve visual clarity.

A major problem with this scenario is the insufficient shortening on the Adriatic plate in the eastern Alps of only 50 km (Schönborn, 1999), as inferred from balancing of cross sections (Nussbaum, 2000). In order to explain a 200 km long Adriatic slab from the Lippitsch et al. (2003) model, Handy et al. (2015) propose that the slab may have deformed and stretched, driven by suction from the broken-off European slab, or that the tomographic image shows an amalgamation of European and Adriatic slab. The insufficient shortening of the Adriatic crust is even more difficult to explain with respect to the other models that show a slab length of 300 km and more (Fig. 2, S8–S10).

Scenario (IV): Detached European slab, shallow subduction of Europe and Adria

The discrepancy between minimum shortening estimates in the southern Alps and imaged slab lengths under the eastern Alps corresponds to ≥ 100 km for the model of Lippitsch et al. (2003) and ≥ 200 km for the other models. This can hardly be explained by Adriatic slab stretching or smearing, but could be explained by subduction of both European and Adriatic slabs into the upper mantle (Fig. 4). Considering the inferred, very shallow level of detachment of the European slab, as derived from the surface-wave model (Fig. 1) and the apparent continuity of the slab in many models, a young break-off age, significantly younger than the one inferred to cause the Periadriatic magmatism at 32–30 Ma, is required. Such a detachment would propagate in a wedge-like fashion from the east, possibly related to the eastward escape of the Carpathians (Schmid et al., 2004) and similar to the model of Qorbani et al. (2015) based on their SKS-splitting results. This model is in agreement with the vertical slab gap in the top 250 km, east of 14° longitude (Koulakov et al., 2009; Mitterbauer et al., 2011; Zhao et al., 2016; Hua et al., 2017). A small amount of northward directed subduction of Adria filling the gap left by the broken-off European



slab could explain why many tomographic models show a continuous vertical anomaly in the uppermost mantle layer between 12° and 14° longitude. Hence, the combination of remnants of Europe and onsetting Adriatic subduction could explain the weak, high-velocity anomaly in the surface-wave model.

A slight northward dip of the eastern Alpine slab is observed in most models, both in the shallow and in the deep parts of the slab. This may be caused by a retreat of the European slab (Kissling and Schlunegger, 2018) or northward directed asthenospheric flow under the northern Adriatic plate (e.g., Vignaroli et al., 2008), or by imaging a combination of shallow Adriatic and deep European slab (Fig. 3). A northward retreat of Europe would also explain the northward step of the anomaly from central to eastern Alps as observed in some models (Zhao et al., 2016; Hua et al., 2017). The deep, broken-off part of the European slab may be linked to the slab remnants in the mantle transition zone below the Pannonian basin (Dando et al., 2011, Fig. 4).

We assume that there is a structural transition along strike of the eastern Alps, related to the proposed wedge-like detachment: Adriatic subduction may play no or only a very minor role in the westernmost part (Tauern window) where the Adriatic slab should not be significantly longer than the inferred amounts of ~50 km (Fig. 3). Further east, parts of the Dinaric slab may have propagated laterally – along with the counter-clockwise rotation of Adria – into the eastern Alps, thus showing a longer Adriatic slab there and approaching Scenario (III) towards the eastern end of the eastern Alps. This would explain why the northward dip is clearest at the Alps-Dinarides transition, but less obvious close to the TRANSALP cross-section along approx. 12° (Figs. 2, 4, S8–S10).

The fast anomaly under the German Molasse, at around 150 km depth in the surface-wave model (Fig. 1C) and the one of Lip-pitsch et al. (2003) (Fig. 2) may be related to a European slab retreat, comparable to the Vrancea slab which is located under the Carpathian foreland (Knapp et al., 2005). The direction is compatible with the sinistral offset along the Giudicarie fault, the location at 200 km north of the eastern Alps is, however, quite far compared to the estimated 70 km of lateral movement along the fault (Scharf et al., 2013). The relation of the foreland anomaly to the eastern Alpine subduction may improve our understanding of the processes in the eastern Alps.

Similar to the western Alps, it will be necessary to improve the tomographic images in the eastern Alps and the northern Dinarides especially in the top 150 km, thus showing the crust-mantle transition and giving the possibility of attributing the deeper anomalies to either the European or Adriatic plate. This is linked to having better spatial and temporal constraints for the slab break-off(s) and testing the hypothesis of northward-directed European slab retreat. We expect that further insight will be provided by the AlpArray experiment (Hetényi et al., 2018a).

5 Conclusions

Our comparison of regional high-resolution tomography models shows that some of the body-wave models differ significantly in shape and length of the imaged slabs. As a consequence there is no consistent model of the Alpine subduction and slab geometries, yet. In order to understand detachments at lithospheric level, good resolution in the uppermost 200 km is required.



Teleseismic body-wave models are limited in this region due to steep incident angles of teleseismic rays, whereas the shear-velocity model from surface-wave recordings provides a better resolution.

In the western Alps, we conclude that the European slab is detached at the bottom of the lithosphere, hence the absence of slab pull may cause the recent uplift despite the absence of convergence. At greater depth, tomographic models cannot yet clearly distinguish between the detached European slab and Adriatic subduction under the northern Apennines that may be merged into one slab.

Concerning the eastern Alps, we discuss three scenarios from the literature for the subduction processes in the eastern Alps. We object the interpretation of a continuous Adriatic slab in the entire upper mantle, however, shallow Adriatic subduction cannot be ruled out. We suggest that the amount of subduction should be consistent with southern Alpine shortening estimates of ≥ 50 km and may only increase towards the northern Dinarides where the Adriatic slab is known to be of 150 km length. A European slab break-off could have propagated from the east related to the eastward migration of the Carpathians. Adria may have partially filled this gap. We summarize these findings in a new, fourth scenario, which will stimulate the discussion on the eastern Alpine slab geometries and provides a more evolved hypothesis to be further tested with the AlpArray (Hetényi et al., 2018a) experiment, ideally with a combination of methods to combine the strength of both body- and surface-wave tomography.

Data availability. All data used in the surface-wave study is stored in the EIDA archive (<http://www.orfeus-eu.org/eida>). The final surface-wave model is freely accessible, including Python routines for plotting (<http://hestia.lgs.jussieu.fr/boschil/downloads.html>). The body-wave tomographic models are available on request via the original authors.

Author contributions. EK created the model comparison and prepared the MS with contributions from all co-authors. CR contributed to the interpretation and discussions on the Alpine structures. EK and LB created the surface-wave tomographic model. NB contributed to the discussion on geological and tectonic context of the Alpine subduction. TM and AE contributed with an earthquake surface-wave dataset and assisted both the technical and interpretative discussions.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We want to thank A. Paul, B. Dando, Y. Hua, I. Koulakov, U. Mitterbauer and R. Lippitsch for providing the tomographic models. We are grateful to all the network operators providing data to the EIDA archive. Graphics were created with Python Matplotlib, Blender and Inkscape. EK has received funding from the German Science Foundation (SPP-2017, Project Ha 2403/21-1). LB has received



funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 641943.



References

- Beller, S., Monteiller, V., Operto, S., Nolet, G., Paul, A., and Zhao, L.: Lithospheric architecture of the South-Western Alps revealed by multiparameter teleseismic full-waveform inversion, *Geophysical Journal International*, 212, 1369–1388, 2017.
- Blanckenburg, F. and Davies, J. H.: Slab breakoff: a model for syncollisional magmatism and tectonics in the Alps, *Tectonics*, 14, 120–131, 1995.
- Boschi, L., Faccenna, C., and Becker, T.: Mantle structure and dynamic topography in the Mediterranean Basin, *Geophysical Research Letters*, 37, 2010.
- Capitanio, F., Morra, G., and Goes, S.: Dynamic models of downgoing plate-buoyancy driven subduction: Subduction motions and energy dissipation, *Earth and Planetary Science Letters*, 262, 284–297, 2007.
- 10 Dando, B., Stuart, G., Houseman, G., Hegedüs, E., Brückl, E., and Radovanović, S.: Teleseismic tomography of the mantle in the Carpathian-Pannonian region of central Europe, *Geophysical Journal International*, 186, 11–31, 2011.
- Dziewonski, A. M. and Anderson, D. L.: Preliminary reference Earth model, *Physics of the earth and planetary interiors*, 25, 297–356, 1981.
- Escher, A. and Beaumont, C.: Formation, burial and exhumation of basement nappes at crustal scale: a geometric model based on the Western Swiss-Italian Alps, *Journal of Structural Geology*, 19, 955–974, 1997.
- 15 Handy, M. R., Schmid, S. M., Bousquet, R., Kissling, E., and Bernoulli, D.: Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological–geophysical record of spreading and subduction in the Alps, *Earth-Science Reviews*, 102, 121–158, 2010.
- Handy, M. R., Ustaszewski, K., and Kissling, E.: Reconstructing the Alps–Carpathians–Dinarides as a key to understanding switches in subduction polarity, slab gaps and surface motion, *International Journal of Earth Sciences*, 104, 1–26, 2015.
- Harangi, S., Downes, H., and Seghedi, I.: Tertiary-Quaternary subduction processes and related magmatism in the Alpine-Mediterranean region, *Geological Society, London, Memoirs*, 32, 167–190, 2006.
- Hawkesworth, C., Waters, D., and Bickle, M.: Plate tectonics in the Eastern Alps, *Earth and Planetary Science Letters*, 24, 405–413, 1975.
- Hetényi, G., Molinari, I., Clinton, J., Bokelmann, G., Bondár, I., Crawford, W. C., Dessa, J.-X., Doubre, C., Friederich, W., Fuchs, F., et al.: The AlpArray Seismic Network: A large-scale European experiment to image the Alpine orogen, *Surveys in Geophysics*, pp. 1–25, 2018a.
- Hetényi, G., Plomerová, J., Bianchi, I., Exnerová, H. K., Bokelmann, G., Handy, M. R., Babuška, V., Group, A.-E. W., et al.: From mountain 25 summits to roots: Crustal structure of the Eastern Alps and Bohemian Massif along longitude 13.3° E, *Tectonophysics*, 744, 239–255, 2018b.
- Hua, Y., Zhao, D., and Xu, Y.: P-wave anisotropic tomography of the Alps, *Journal of Geophysical Research: Solid Earth*, 2017.
- Kästle, E. D., El-Sharkawy, A., Boschi, L., Meier, T., Rosenberg, C., Bellahsen, N., Cristiano, L., and Weidle, C.: Surface Wave Tomography of the Alps Using Ambient-Noise and Earthquake Phase Velocity Measurements, *Journal of Geophysical Research: Solid Earth*, 123, 1770–1792, 2018.
- 30 Kissling, E. and Schlunegger, F.: Rollback orogeny model for the evolution of the Swiss Alps, *Tectonics*, 37, 1097–1115, 2018.
- Kissling, E., Schmid, S. M., Lippitsch, R., Ansorge, J., and Fügenschuh, B.: Lithosphere structure and tectonic evolution of the Alpine arc: new evidence from high-resolution teleseismic tomography, *Geological Society, London, Memoirs*, 32, 129–145, 2006.
- Knapp, J. H., Knapp, C. C., Raileanu, V., Matenco, L., Mocanu, V., and Dinu, C.: Crustal constraints on the origin of mantle seismicity in the Vrancea Zone, Romania: The case for active continental lithospheric delamination, *Tectonophysics*, 410, 311–323, 2005.
- Koulakov, I., Kaban, M., Tesauero, M., and Cloetingh, S.: P- and S-velocity anomalies in the upper mantle beneath Europe from tomographic inversion of ISC data, *Geophysical Journal International*, 179, 345–366, 2009.



- Laubscher, H.: Das Alpen-Dinariden-Problem und die Palinspastik der südlichen Tethys, *Geologische Rundschau*, 60, 813–833, 1971.
- Lippitsch, R., Kissling, E., and Ansorge, J.: Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography, *Journal of Geophysical Research: Solid Earth*, 108, 2003.
- Lüschen, E., Lammerer, B., Gebrande, H., Millahn, K., Nicolich, R., Group, T. W., et al.: Orogenic structure of the Eastern Alps, Europe, from TRANSALP deep seismic reflection profiling, *Tectonophysics*, 388, 85–102, 2004.
- 5 Lyu, C., Pedersen, H. A., Paul, A., Zhao, L., and Solarino, S.: Shear wave velocities in the upper mantle of the Western Alps: new constraints using array analysis of seismic surface waves, *Geophysical Journal International*, 210, 321–331, 2017.
- Mitterbauer, U., Behm, M., Brückl, E., Lippitsch, R., Guterch, A., Keller, G. R., Koslovskaya, E., Rumpfhuber, E.-M., and Šumanovac, F.: Shape and origin of the East-Alpine slab constrained by the ALPASS teleseismic model, *Tectonophysics*, 510, 195–206, 2011.
- 10 Molli, G. and Malavieille, J.: Orogenic processes and the Corsica/Apennines geodynamic evolution: insights from Taiwan, *International Journal of Earth Sciences*, 100, 1207–1224, 2011.
- Nocquet, J.-M., Sue, C., Walpersdorf, A., Tran, T., Lenôtre, N., Vernant, P., Cushing, M., Jouanne, F., Masson, F., Baize, S., et al.: Present-day uplift of the western Alps, *Scientific reports*, 6, 28 404, 2016.
- Nussbaum, C.: Neogene tectonics and thermal maturity of sediments of the easternmost southern Alps (Friuli area, Italy), PhD thesis, 2000.
- 15 Pamić, J., Belak, M., Bullen, T., Lanphere, M., and McKee, E.: Geochemistry and geodynamics of a Late Cretaceous bimodal volcanic association from the southern part of the Pannonian Basin in Slavonija (Northern Croatia), *Mineralogy and Petrology*, 68, 271–296, 2000.
- Qorbani, E., Bianchi, I., and Bokelmann, G.: Slab detachment under the Eastern Alps seen by seismic anisotropy, *Earth and planetary science letters*, 409, 96–108, 2015.
- Replumaz, A., Negredo, A. M., Villasenor, A., and Guillot, S.: Indian continental subduction and slab break-off during Tertiary collision, *Terra Nova*, 22, 290–296, 2010.
- 20 Rosenberg, C.: Shear zones and magma ascent: a model based on a review of the Tertiary magmatism in the Alps, *Tectonics*, 23, 2004.
- Rosenberg, C. L., Berger, A., Bellahsen, N., and Bousquet, R.: Relating orogen width to shortening, erosion, and exhumation during Alpine collision, *Tectonics*, 34, 1306–1328, 2015.
- Rosenberg, C. L., Schneider, S., Scharf, A., Bertrand, A., Hammerschmidt, K., Rabaute, A., and Brun, J.-P.: Relating collisional kinematics to exhumation processes in the Eastern Alps, *Earth-Science Reviews*, 2017.
- 25 Rosenberg, C. L., Bellahsen, N., Kästle, E., Rabaute, A., and Boschi, L.: Amount and distribution of collisional shortening in the Alpine Chain: implications for slab geometries at depth, *EGU 2019 general assembly abstracts*, 2019.
- Scharf, A., Handy, M., Favaro, S., Schmid, S. M., and Bertrand, A.: Modes of orogen-parallel stretching and extensional exhumation in response to microplate indentation and roll-back subduction (Tauern Window, Eastern Alps), *International Journal of Earth Sciences*, 102, 1627–1654, 2013.
- 30 Schmid, S. M., Pfiffner, O.-A., Froitzheim, N., Schönborn, G., and Kissling, E.: Geophysical-geological transect and tectonic evolution of the Swiss-Italian Alps, *Tectonics*, 15, 1036–1064, 1996.
- Schmid, S. M., Fügenschuh, B., Kissling, E., and Schuster, R.: Tectonic map and overall architecture of the Alpine orogen, *Eclogae Geologicae Helveticae*, 97, 93–117, 2004.
- 35 Schmid, S. M., Bernoulli, D., Fügenschuh, B., Matenco, L., Schefer, S., Schuster, R., Tischler, M., and Ustaszewski, K.: The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units, *Swiss Journal of Geosciences*, 101, 139–183, 2008.
- Schmid, S. M., Scharf, A., Handy, M. R., and Rosenberg, C. L.: The Tauern Window (Eastern Alps, Austria): a new tectonic map, with cross-sections and a tectonometamorphic synthesis, *Swiss Journal of Geosciences*, 106, 1–32, 2013.



- Schönborn, G.: Balancing cross sections with kinematic constraints: the Dolomites (northern Italy), *Tectonics*, 18, 527–545, 1999.
- Spada, M., Bianchi, I., Kissling, E., Agostinetti, N. P., and Wiemer, S.: Combining controlled-source seismology and receiver function information to derive 3-D Moho topography for Italy, *Geophysical Journal International*, 194, 1050–1068, 2013.
- Stampfli, G. M. and Borel, G.: A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons, *Earth and Planetary Science Letters*, 196, 17–33, 2002.
- 5 Ustaszewski, K., Schmid, S. M., Fügenschuh, B., Tischler, M., Kissling, E., and Spakman, W.: A map-view restoration of the Alpine-Carpathian-Dinaridic system for the Early Miocene, in: *Orogenic Processes in the Alpine Collision Zone*, pp. S273–S294, Springer, 2008.
- Vignaroli, G., Faccenna, C., Jolivet, L., Piromallo, C., and Rossetti, F.: Subduction polarity reversal at the junction between the Western Alps and the Northern Apennines, Italy, *Tectonophysics*, 450, 34–50, 2008.
- 10 Zhao, L., Paul, A., Malusà, M. G., Xu, X., Zheng, T., Solarino, S., Guillot, S., Schwartz, S., Dumont, T., Salimbeni, S., et al.: Continuity of the Alpine slab unraveled by high-resolution P wave tomography, *Journal of Geophysical Research: Solid Earth*, 2016.