

1 **Seismicity characterization of oceanic earthquakes in the Mexican**
2 **territory**

3
4 **Quetzalcoatl Rodríguez-Pérez^{1,2,*}, Víctor Hugo Márquez-Ramírez², and Francisco Ramón**
5 **Zúñiga²**
6

7 ¹ Consejo Nacional de Ciencia y Tecnología, Dirección Adjunta de Desarrollo Científico, Mexico

8 ² Centro de Geociencias, Universidad Nacional Autónoma de México, Juriquilla, Querétaro, Mexico

9

10 **Correspondence:** Quetzalcoatl Rodríguez-Pérez (quetza@geociencias.unam.mx)

11 Víctor H. Márquez-Ramírez (marvh@geociencias.unam.mx)

12 F. Ramón Zúñiga (ramon@geociencias.unam.mx)

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

1 **Abstract.** We analyzed the seismicity of oceanic earthquakes in the Pacific oceanic regime of Mexico.
2 We used data from the earthquake catalogues of the Mexican National Service (SSN), and the
3 International Seismological Center (ISC) from 1967 to 2017. Events were classified into two different
4 categories: intraplate oceanic (INT), and transform faults zone and mid-ocean ridges events (TF-MOR),
5 respectively. For each category, we determined statistical characteristics such as magnitude frequency
6 distributions, the aftershocks decay rate, the non-extensivity parameters, and the regional stress field.
7 We obtained b -values of 1.17, and 0.82 for the INT, and TF-MOR events, respectively. TF-MOR events
8 also exhibit local b -value variations in the range of 0.72 – 1.30. TF-MOR events follow a tapered
9 Gutenberg-Richter distribution. We also obtained a p -value of 0.67 for the 1 May 1997 ($M_w = 6.9$)
10 earthquake. By analyzing the non-extensivity parameters, we obtained similar q -values in the range of
11 1.39-1.60 for both types of earthquakes. On the other hand, the parameter a showed a clear
12 differentiation, being higher for TF-MOR events than for INT events. An important implication is that
13 more energy is released for TF-MOR events than for INT events. Stress orientations are in agreement
14 with geodynamical models for transform faults zone and mid-ocean ridges zones. In the case of
15 intraplate seismicity, stresses are mostly related to a normal fault regime.

16

17 **1 Introduction**

18

19 Mid-ocean ridges and transform faults zones are two of the main morphological features of oceanic
20 environments. Most of the oceanic earthquakes take place in areas close to the active spreading ridges
21 where the seismogenic zone is narrow. For this reason, large aspect ratios are often required to generate
22 moderate-size strike-slip oceanic earthquakes. Nevertheless, the rupture process of oceanic events is
23 still poorly understood. Previous studies showed that these types of events have peculiar characteristics.
24 For example, estimates of seismic coupling for oceanic transform faults indicate that about three-

1 fourths of the accumulated moment are released aseismically (Abercrombie and Ekström, 2003;
2 Boettcher and Jordan, 2004) and some oceanic events exhibit slow slip ruptures (Kanamori and
3 Stewart, 1976; Okal and Stewart, 1992; McGuire et al., 1996). Earthquakes that have longer durations
4 than those predicted by scaling relationships are considered as slow (Abercrombie and Ekström, 2003).
5 These “slow” ruptures are mainly interpreted as having low rupture velocities. On the other hand,
6 others proposed that the slow ruptures may be explained as numerical artifacts generated by the
7 inversion procedures (e.g., Abercrombie and Ekström 2001; 2003). Several oceanic strike-slip events
8 were reported as being energy deficient at high-frequencies (Beroza and Jordan, 1990; Stein and
9 Pelayo, 1991; Ihmlé and Jordan, 1994), or having high apparent stresses (Choy and Boatwright, 1995;
10 Choy and McGarr, 2002). On another front, oceanic earthquakes also occur as intraplate events, but to
11 a lesser extent. The reason is that the oceanic plate interiors do not experience significant strain over
12 long periods of time (Bergman and Solomon 1980; Bergman, 1986). Oceanic intraplate earthquakes
13 originate from the following processes: stresses of the oceanic crust, in regions that concentrate
14 significant deformation, reactivation of faults, or thermoelastic stresses (Bergman, 1986).

15

16 From the statistical perspective, previous studies showed that the magnitudes of the major events in the
17 mid-oceanic ridges and transform faults zones are relatively smaller ($6.0 \leq M_w \leq 7.2$) compared to
18 continental events. The b -value in oceanic environments showed significant variability. For example,
19 Tolstoy et al. (2001) reported high b -values ($b \sim 1.5$) in the Gakkel Ridge associated with volcanic
20 activity. In the Southwest Indian Ridge, Läderach (2011) found b -values of about 1.28. Bohnenstiehl et
21 al. (2008) quantified the b -value in the East Pacific Rise, obtaining estimations in the range of $1.10 < b$
22 < 2.50 . Global studies have also shown that the mid-ocean ridge transform seismicity follows a tapered
23 frequency-moment distribution (Kagan and Jackson, 2000; Boettcher and McGuire, 2009). Cowie et al.
24 (1993) studied the seismic coupling on mid-ocean ridges. They found that fast-spreading ridges (≥ 9.0

1 cm/yr) are weakly coupled. On the contrary, slow-spreading ridges (≤ 4.0 cm/yr) are strongly coupled
2 (Cowie et al., 1993). In Mexico, oceanic earthquakes have been poorly studied. There are no systematic
3 studies on their statistical characteristics. In this article, we characterized the oceanic seismicity in
4 Mexico. We determined the orientation of the principal stresses, the b - and p -values, and the non-
5 extensivity parameters. The results may help to understand the ocean tectonics, particularly in Mexico.

6

7 **2 Tectonic Setting**

8

9 The Pacific oceanic regime of Mexico is an active area exhibiting ongoing tectonic plate interactions.
10 These interactions involve the Cocos (CO), the Pacific (PA), the Rivera (RI), and the North American
11 (NA) plates. The Gulf of California and the Middle America Trench (MAT) are separated by the
12 Tamayo Fracture Zone (TFZ). The convergence rate between the RI, and NA plates decreases
13 northward along the MAT (averaging about 2–3 cm/yr in the RI plate, which is slower than the adjacent
14 CO plate, about 5–7 cm/yr) (NUVEL-1a model, DeMets et al., 1994). Sea-floor spreading takes place
15 along the northernmost segment of the East Pacific Rise in the Cocos, and Rivera segments (EPR-CS,
16 and EPR-RS, respectively). In the EPR-RS, the spreading rates range from 5.3 cm/year at the northern
17 to 7.3 cm/year at the southern end of the rise (Bandy, 1992). The spreading rates at the EPR-CS are: 7.0
18 cm/yr near the Rivera Fracture Zone (RFZ); 8.2 cm/yr near the Orozco Fracture Zone (OFZ); 10.1
19 cm/yr near the Clipperton Fracture Zone (CFZ); and 10.7 cm/yr near the Siqueiros Fracture Zone (SFZ)
20 based on the NUVEL-1a model (DeMets et al., 1994; Pockalny et al., 1997). The Rivera Transform
21 (RT) is a left transform fault with fast slipping (~ 7.0 cm/year) (Bandy et al., 2011) (Fig. 1). Due to
22 these differences in subduction, and spreading rates, and convergence direction of the RI and CO
23 plates, complex seismicity patterns are generated in this region. In the last century, some intermediate-
24 size earthquakes ($6.8 < M < 7.1$) have taken place in the Pacific oceanic regime of Mexico (Table 1 and

1 Fig. 2).

2

3 **3 Data and Methods**

4

5 **3.1 Data**

6 We used earthquake catalogues of the Mexican National Service (SSN), and the International
7 Seismological Center (ISC) from 1967 to 2017. Events with **no reported** magnitude were excluded
8 from our analysis. Reported magnitudes (based on **surface**, M_s ; body, m_b ; and coda, M_c ; waves) were
9 converted to moment magnitude (M_w). The SSN reports M_w for events in Mexico. For the case of the
10 ISC events, M_s , and m_b were converted to M_w using the scaling relationships of Scordilis (2006). We
11 classified the seismic events into two different categories: 1) intraplate oceanic events (INT, red dots in
12 Fig. 2), and 2) transform faults zone and mid-ocean ridges events (TF-MOR, green dots in Fig. 2). The
13 INT catalogue consists of 177 events with magnitudes in the range of 2.9 - 6.0. The TF-MOR catalogue
14 **includes 2074 earthquakes with magnitudes between 2.7 and 6.9.** We also used the Global CMT focal
15 mechanism catalogue (Dziewonski et al., 1981; Ekström et al., 2012) with solutions from 1976 to 2017.
16 For the stress analysis, the focal mechanism catalog was divided into 6 sub-catalogues shown in Fig. 8
17 (R1 to R6).

18

19 **3.2 Methods**

20 **3.2.1 Moment/magnitude earthquake distributions**

21 The Gutenberg-Richter law describes the **earthquake magnitude distribution** (Ishimoto and Iida, 1939;
22 Gutenberg and Richter, 1944). Mathematically, this law is expressed by the following equation: \log_{10}
23 $N(M) = a - bM$, where $N(M)$ is the cumulative number of earthquakes with a magnitude larger than a
24 given magnitude limit (M), the constant b (or b -value) describes the slope of the **magnitude** distribution

1 and the constant a is proportional to the seismic productivity. The b -value describes the distribution of
2 small to large earthquakes in a sample, and it is considered to be specific for a given tectonic
3 environment (e.g., Scholz, 1968; Wyss, 1973; Smith, 1981; Wiemer and Benoit, 1996; Wiemer and
4 Wyss, 2002). In several tectonic environments, b is close to 1 (Utsu, 1961), with deviations affected by
5 many factors. Among them, high thermal gradients and rock heterogeneity (Mogi, 1962; Warren and
6 Latham, 1970) increases the b -values. On the contrary, increments in effective and shear stresses
7 (Scholz, 1968; Wyss, 1973; Urbancic et al., 1992) reduce the b -value. The b -value differs between
8 unrelated fault zones (Wesnousky, 1994; Schorlemmer et al., 2005), but also for specific space and time
9 periods (Nuannin et al., 2012). Schorlemmer et al. (2005) found a global dependence of the b -value on
10 focal mechanism, which was corroborated at a regional level by Rodríguez-Pérez and Zúñiga (2018).
11 According to those authors, the highest b -values correspond to normal-faulting events, followed by
12 strike-slip, and thrust earthquakes, respectively. To characterize the b -value of oceanic earthquakes and
13 compare the results with other tectonic environments, we calculated the b -value with a robust method
14 which has proven its validity in many studies. We estimated the b -value by means of the maximum
15 likelihood formulation of Aki (1965), and the completeness magnitude (M_c) employing the maximum
16 curvature method (Wiemer and Wyss, 2000) with the aid of the ZMAP software package (Wiemer,
17 2001).

18

19 As reported by previous authors, seismicity on the mid-ocean transform faults is better represented by a
20 tapered frequency moment distribution (e.g., Boettcher and McGuire, 2009). This distribution has the
21 following form (Kagan, 1997, 1999; Kagan and Jackson, 2000; Kagan and Schoenberg, 2001; Vere-
22 Jones et al., 2001):

23

1
$$N(M) = N_o \left(\frac{M_o}{M} \right)^\beta \exp\left(-\frac{M_o - M}{M_m} \right), \quad (1)$$

2

3 where β is one of the parameters to determine ($\beta = (2/3)b$, where b is the b -value), N_o is the cumulative
4 earthquake number over a completeness threshold seismic moment (M_o), and M_m is the maximum
5 expected moment. We analyzed if this frequency distribution is suitable for describing the seismicity of
6 oceanic events in Mexico. In order to calculate the tapered Gutenberg-Richter distribution, we used the
7 Matlab function `Get_GR_parameters.m` developed by Olive (2016). The tapered Gutenberg-Richter
8 moment distribution is fitted by means of a least-squares inversion following Frohlich (2007).

9

10 **3.2.2 Temporal distribution of aftershocks**

11 The frequency distribution of the decrement of earthquake aftershocks is described by the modified
12 Omori's law (Utsu, 1961; Utsu et al., 1995) as:

13

14
$$R(t) = \frac{k}{(t+c)^p}, \quad (2)$$

15

16 where $R(t)$ is the rate of occurrence of aftershocks within a given magnitude range, t is the time interval
17 from the mainshock, k is the productivity of the aftershock sequence, p is the power-law exponent (p -
18 value), and c is the time delay before the onset of the power-law aftershock decay rate. Variations of p -
19 values exist for different tectonic regimes and each aftershock sequence. Many authors have related the
20 p -value with crustal temperature, heat-flow, or rock heterogeneity in the fault zone. Thus, relevant
21 information can be extracted from these aftershock parameters in order to have a better understating of
22 the rupture process of oceanic earthquakes. As before, we used the ZMAP software package (Wiemer,

1 2001) for estimating the p -value of the aftershock sequence of the 1 May 1997 earthquake ($M_w = 6.9$).

2

3 3.2.3 Fragment-asperity model

4 Alternative statistical models that relate the earthquake magnitude distribution with the rheology of the
5 fault have been proposed. Among them, we have the fragment-asperity model. This model was
6 introduced by Sotolongo-Costa and Posadas (2004) to describe the earthquake dynamics in a Tsallis
7 entropy non-extensive framework (Tsallis, 1988). This model takes into consideration the irregular
8 surfaces of two fault planes in contact and the rock fragments of different shape and sizes that fill the
9 space between them. According to this model, earthquakes are triggered by the interaction along the
10 fault planes of these rock fragments. Considering that large fragments are more difficult to release than
11 small ones, the resulting energy is assumed to be proportional to the volume of the fragment (Telesca,
12 2010). Silva et al. (2006) improved the model and found a scaling law between the released energy (ε),
13 and the size of asperity fragments (r) by the following proportional factor: $\varepsilon \propto r^3$. The non-extensive
14 statistics is used to describe the volumetric distribution function of the fragments. A parameter that
15 represents the proportion between ε and r is introduced. This parameter is known as the a -value or
16 parameter a (Silva et al., 2006; Telesca, 2010). The parameter a is defined using a volumetric
17 distribution function of the fragments applying the maximum entropy principle for the Tsallis entropy
18 (for details in the mathematical expressions see Silva et al., 2006; Telesca, 2010). The magnitude
19 cumulative distribution function becomes:

20

$$21 \log_{10}(N > M) = \log_{10}(N) + \left(\frac{2-q}{1-q} \right) \log_{10} \left[1 - \left(\frac{1-q}{2-q} \right) \left(\frac{10^K}{a^{2/3}} \right) \right], \quad (3)$$

22

23 where N is the total number of earthquakes; $N (>M)$ represents the number of events with magnitude

1 larger than M ; a is a proportionality parameter between ε and r , and; q is the non-extensivity parameter.
2 K is defined as $K = 2M$ (Silva et al., 2006), or $K = M$ (Telesca, 2011). The magnitude (M) is related to
3 ε by the following relation: $M = 1/3 \log(\varepsilon)$ (Silva et al., 2006). Telesca (2011) considered that the
4 relation between ε and M is given by $M = 2/3 \log(\varepsilon)$ (Telesca, 2011). None of both models are preferred
5 over the other. We used both models in order to quantify the variability of the non-extensive
6 parameters. According to Telesca (2010), the physical meaning of the q -parameter consists in that it
7 provides information about the scale of interactions. It means that if q is close to 1, the physical state is
8 close to the equilibrium. As a result, few earthquakes are expected. On the other hand, as q rises, the
9 physical state goes away from the equilibrium state, this implies that the fault planes are able to
10 generate more earthquakes, thus resulting in an increment in the seismic activity (Telesca, 2009; 2011).
11 The physical meaning of the a -value lies in the fact that it provides a measure of the energy density. It
12 means that the a -value is large if the energy released is large (Telesca, 2011). For example, high a -
13 values are expected when the events with the highest magnitude take place. Previous studies have
14 shown that the q -value ranges mainly from 1.50 to 1.70 (Vilar et al., 2007; Vallianatos, 2009;
15 Rodríguez-Pérez and Zúñiga, 2017; among others). We obtained the a and q parameters by minimizing
16 the root mean square error (RMS) with the Nelder-Mead method (Nelder and Mead, 1965).

17

18 **3.2.5 Stress Inversion**

19 **Focal mechanisms are reliable indicators of the state of stress in a tectonic region.** In order to study the
20 regional stress field for oceanic earthquakes, we performed stress tensor inversion from focal
21 mechanisms reported in the Global CMT catalogue (Dziewonski et al., 1981; Ekström et al., 2012) with
22 the iterative joint inversion developed by Vavryčuk (2014). From the stress inversion, we obtained the
23 orientation of the principal stress axes σ_1 , σ_2 , and σ_3 (where $\sigma_1 \geq \sigma_2 \geq \sigma_3$), and the stress ratio R . We now
24 briefly explain each method. The first method (the iterative joint inversion), provides an accurate

1 estimation of R and stress orientations (Vavryčuk, 2014). In this method, the ratio is defined as $R = (\sigma_1$
2 $- \sigma_2) / (\sigma_1 - \sigma_3)$ (Gephart and Forsyth, 1984). A fault instability constraint is applied, and the fault is
3 identified with that nodal plane which is more unstable, and thus more susceptible to faulting
4 (Vavryčuk, 2014). By incorporating a fault instability constraint into the inversion, an iterative
5 procedure is imposed. The uncertainties are determined as the differences between the inverted results
6 considering noisy data (Vavryčuk, 2014). The stress inversion was carried out with the
7 STRESSINVERSE software developed by Vavryčuk (2014). The maximum horizontal stress (SH_{max})
8 was calculated using the formulation of Lund and Townend (2007). The stress inversion was performed
9 for each of the six different regions shown in Fig. 7.

10

11 **4 Results**

12

13 There is a large span of b -values (Table 2) which nevertheless sheds light on the seismicity
14 characteristics of oceanic earthquakes in Mexico. INT events exhibit higher b -values and M_c than TF-
15 MOR events (Fig. 3, Fig. 4a and Table 2). In particular, TF-MOR events also show local b -value
16 variations in the range of 0.72 – 1.30 (Fig. 4b) for each of the subregions R1 to R5 (Table 2). Previous
17 studies had shown large fluctuations in b -values of oceanic events. For example, Tolstoy et al. (2001)
18 reported b -values of about 1.5 associated with volcanic activity in the Gakkel Ridge. Läderach (2011)
19 reported b -values of 1.28 in the Southwest Indian Ridge. In a global study, Molchan et al. (1997)
20 estimated the b -value for mid-ocean, and transform zones, obtaining values of the following interval
21 0.97 – 1.47. In general, our b -value estimates agree with reported b -values in previous studies. On the
22 other hand, our results showed that M_c for oceanic events is higher than reported M_c for the subduction
23 zone, and continental regions of Mexico, which reflects on the capability of the global and regional
24 networks to appropriately register events in that region. The magnitude completeness for oceanic

1 earthquakes differs for different parts of the World, but in most cases, it is in the range of 4.0 – 5.0 on
2 average considering most of the global catalogues.

3
4 Our results also showed that transform faults zone and mid-ocean ridges events follow a tapered
5 Gutenberg-Richter distribution, as suggested in previous studies (Boettcher and McGuire, 2009). The
6 tapered Gutenberg-Richter distribution was fitted with the following parameters: $\beta = 0.64$, and an
7 estimated corner magnitude of $M_m = 6.7$ (Fig. 5a). These results are in agreement with previous studies
8 such as that of Bird et al. (2002) who studied the tapered Gutenberg-Richter distribution for spreading
9 ridges and oceanic transform faults based on global data obtaining a β -value of about 0.67 for both
10 types of events. They reported that M_m varies from 5.8 to 6.6 – 7.1 for mid-ocean ridge and transform
11 faults, respectively. The results for the non-extensive parameters are shown in Table 2. We found higher
12 q -values for TF-MOR events than for INT events (Fig. 5), meaning that TF-MOR events are farther
13 from the equilibrium than INT events. The results showed a better fitting for cumulative distribution
14 functions using the Telesca model for TF-MOR and each of the regions (Fig. 6). In regions R1-R5, our
15 results showed that q varies from 1.31 to 1.52, and from 1.57 to 1.63 using the Telesca's and Silva's
16 models, respectively. In the case of subduction zones, the q -value can vary from 1.35 to 1.70. For
17 example, in the Hellenic Subduction Zone, q is in the range of 1.35 - 1.55 (Papadakis et al., 2013); in
18 the Mexican subduction zone, Valverde-Esparza et al. (2012) found that q varies from 1.63 to 1.70.
19 Thus, our results conform to values obtained in regional studies.

20
21 The analysis of the aftershock sequence of the 1 May 1997 earthquake ($M_w = 6.9$), yielded a p -value of
22 0.67 ± 0.33 (Table 3). The magnitude of the largest aftershock of the 1997 event was $M_w = 5.3$ (Table
23 3). Oceanic strike-slip events seem to have lower p -values than mid-ocean ridges events. For example,
24 Bohnenstiehl et al. (2004) found a p -value of 0.95 for the 15 July 2003 ($M_w = 7.6$) central Indian Ridge

1 strike-slip event. For the Siqueiros, Discovery, and western Blanco transforms, the p -value varies from
2 0.94 to 1.29 (Bohnenstiehl et al., 2002). Davis and Frohlich (1991) determined a p -value of $0.928 \pm$
3 0.024 for the combined ridge and transform environments. Our results fall within the range of global
4 studies that showed that the p -value varies from 0.6 – 2.5 (Utsu et al., 1995). We also reported a c close
5 to 0 for the aftershock sequence of the 1 May 1997 ($M_w = 6.9$) (Table 3). Shcherbakov et al. (2004)
6 found that the parameter c of the Omori's law decreases as the magnitude of events considered
7 increases. According to them, this observation is due to the effect of an undercount of small aftershocks
8 in short time periods. This provides an explanation for our result of $c \sim 0$ because of the limited
9 magnitude detection reported in the regional and global catalogues used.

10

11 We classified the focal mechanisms used in the stress inversion into seven categories (1.- reverse, R; 2.-
12 reverse with lateral component, R-SS; 3.- strike-slip with reverse component, SS-R; 4.- strike-slip, SS;
13 5.- strike-slip with normal component, SS-N; 6.- normal with lateral component, N-SS; 7.- normal, N)
14 (Fig. 7). This classification was performed to identify the dominant type of faulting for each subregion.
15 Region R1 is composed of strike-slip (70.3%), strike-slip with normal and reverse components (21.6%,
16 and 5.4%, respectively), and normal-faulting (2.7%) focal mechanisms (Fig. 7b). Region R2 exhibits
17 the following focal mechanism distribution: strike-slip (82.4%), and strike-slip with normal and reverse
18 components (9.5 %, and 8.1 %, respectively) (Fig. 7b). In region R3, the focal mechanism classification
19 shows the following distribution: strike-slip (62.5%), strike-slip with normal component (25%),
20 normal-faulting with strike-slip component (6.3%), and reverse (6.3%)(Fig. 7b). Region R3 consists of
21 strike-slip (70.8%), strike-slip with normal and reverse components (8.3%, and 16.7 %, respectively),
22 and reverse earthquakes (4.2%)(Fig. 7b). Region R5 exhibits the following focal mechanism
23 distribution: strike-slip (53%), strike-slip with normal and reverse components (23.5%, and 17.6%,
24 respectively), and reverse (5.9%). For the case of earthquakes in R6, the classification shows the

1 following distribution: normal (83.3%) and normal-faulting with strike-slip component (16.7%) (Fig.
2 7b).

3
4 Table 4 summarizes the results from the stress inversion. Based on the orientation of stress axes, a
5 dynamical description of the tectonics of the oceanic earthquakes in Mexico can be carried out. A
6 quantitative comparison with other oceanic regions is discussed in what follows. Region R6 is only
7 dominated by N and N-SS earthquakes (Fig. 8). In regions R4 and R5, stress results showed moderate
8 similarities. The differences in these regions may also be related to the variability of focal mechanisms
9 (here we have SS, SS-N, SS-R, and to lesser extent R events) (Fig. 8). Variations are very significant in
10 regions R1 to R3 (particularly in σ_2) (Table 4). These regions also showed different types of events: SS,
11 SS-N, SS-R for R1; SS, SS-N, SS-R for R2; and SS, SS-N, N-SS, R for R3 (Fig. 8). In these regions,
12 strike-slip earthquakes are the dominant type of faulting. Events with unusual mechanisms have also
13 been reported in other oceanic regions. According to Wolfe et al. (1993), most of the anomalous
14 seismic activity is associated with mislocations, complex fault geometry, or large structural features
15 with an influence on the slip of the fault. DeMets and Stein (1990) showed that the strike direction and
16 earthquake slip vectors in the Rivera transform are rotated clockwise from the expected direction of the
17 Pacific-Rivera Euler vector. This deviation can be the result of morphologic features resulting in
18 unusual patterns of epicenters and focal mechanisms.

19
20 In the case of the East Pacific Rise Rivera segment (region R1), σ_2 is almost vertical, and SH_{\max} is ~
21 170° suggesting a strike-slip regime (Table 4). The main orientations of the P -axes are in the N-S, NW-
22 SE, and E-W directions. The orientation of the P -axes is NW-SE and, to a lesser extent, E-W directions
23 (Fig. 9). For the case of the Rivera Transform (region R2), σ_2 is quasi vertical, and the SH_{\max} is 157°
24 suggesting a strike-slip regime. The orientation of the P -axes is in the NW-SE direction and in the NE-

1 SW direction for the T -axis. In region R3, σ_2 is almost vertical, and the SH_{\max} is also 157° suggesting a
2 strike-slip regime. The orientation of the P -axis is in the NW-SE direction. The main orientation of the
3 T -axes is NE-SW, but E-W directions occur as well. For the region R4, σ_2 is 76° , and the SH_{\max} is 22°
4 suggesting a strike-slip regime. The predominant orientations of the P - and T -axes are NE-SW and
5 NW-SE, respectively. In R5, σ_2 is from 69° , and the SH_{\max} is 120° suggesting a strike-slip regime. The
6 main orientation of the P -axes is NW-SE while that of the T -axis is NE-SW. In R6, the principal axes
7 are related to a normal fault regime. σ_1 is almost vertical, and the SH_{\max} is $\sim 45^\circ$. The orientation of the
8 T -axes is in the NW-SE direction. The Mohr's circle diagram showed that most of the studied events are
9 clustered along the outer Mohr's circle in the area of validity of the Mohr-Coulomb failure criterion
10 (Fig 9). Reported focal mechanisms confirm Sykes's model for mid-ocean ridges (Sykes, 1967), where
11 events in transform zones tend to have strike-slip mechanisms, while ridge crest events have mainly
12 normal faults. The obtained orientation of the principal axes supports this model.

13

14 **5 Discussion**

15

16 One of the main problems for studying oceanic seismicity is that the epicenters are located far from
17 most of the recording stations in mainland Mexico. This has a direct effect on the earthquake
18 magnitude distributions (M_c and b -value). We first discuss the magnitude completeness of oceanic
19 earthquakes. Global studies showed that the magnitude completeness for oceanic earthquakes is in the
20 range of 4.0 – 5.0. Our results are in agreement with these global studies. However, as expected,
21 several microseismic surveys which have been conducted in different oceanic environments (e.g.,
22 Smith et al., 2003; Simão et al., 2010; McGuire et al., 2012; among others) can yield lower magnitude
23 thresholds. As a result of these studies, precise hypocenter locations and earthquake distributions with a
24 broader magnitude range were obtained. Thus, lower M_c has been reported for studies based on

1 microseismic surveys. For example, in the Mid-Atlantic Ridge, $M_c \sim 3.0$ with several smaller events
2 ($M_w < 2.5$) were reported (Bohnenstiehl et al., 2002; Smith et al., 2002, and 2003).

3

4 **Another factor that has to be discussed is the accuracy in the location of the epicenters.** The location
5 uncertainty plays an important role when earthquakes are assigned to an intraplate or a mid-ocean
6 ridge/transform fault environment. For example, some studies reported that for faults located at 4S on
7 the EPR, teleseismic locations could be off as much as 50 km (McGuire, 2008; Wolfson-Schwehr,
8 2014). As a consequence, some TF-MOR events are probably classified as INT events, and vice-versa
9 (for example, epicenters in color in Fig. 2). Some events located in the Tamayo Fracture Zone close to
10 the Rivera subduction zone may also be **misidentified**. This mislocation effect introduces uncertainties
11 in the estimation of the statistical parameters useful for understanding the tectonics of the region. In
12 order to have precise locations and avoid mislocation, ocean-bottom seismometers **off** the Mexican coast
13 **would be** needed. Being aware of this, one should avoid over-interpretation of the results. Local
14 monitoring of oceanic events represents an improvement of more than an order of magnitude relative to
15 the regional, and teleseismic detection levels.

16

17 Previous studies also showed that the seismicity **near** oceanic transform faults that connect mid-ocean
18 ridges may be thermally controlled (Abercrombie and Ekström, 2001; Boettcher et al., 2007). **The**
19 **thermal effect is most evident in the seismogenic zone.** It is essential to mention that faults along the
20 middle and southern segments of the EPR are shorter and faster-slipping. The faster slip rates and
21 shorter fault lengths result in narrower seismogenic zones because the thermal structure is shallow. On
22 the other hand, the Rivera Transform is longer, and has a slower slip rate, resulting in a wider
23 seismogenic zone. However, heat is not the only factor that regulates seismicity because the largest
24 events break a small part of the rupture areas predicted by thermal models (Boettcher and Jordan, 2004;

1 Roland et al., 2010). Thus, most slip occurs without producing large earthquakes (Boettcher and
2 Jordan, 2004; Roland et al., 2010). This can explain the occurrence of a few events with $M > 6.5$ in the
3 Rivera Transform. According to McGuire et al. (2012), the apparent lack of large events on mid-ocean
4 ridge transform faults may also be related to the heterogeneity of materials on the fault plane. The
5 maximum magnitude for transform fault events on the East Pacific Rise (in the latitude interval of $3^\circ <$
6 $\text{Lat} < 5^\circ$) is about 6.5 (McGuire et al., 2005). On the other hand, earthquakes in the Rivera Transform
7 and on the northern segment of the East Pacific Rise (in Mexico) have relative larger magnitudes ($M >$
8 6.8) based on reported seismicity in different catalogues (Fig. 1). This highlights a differentiation
9 between the mid-and southern and northern segments of the East Pacific Rise.

10

11 A further aspect of the analysis of oceanic earthquakes is their capacity to generate aftershocks as well
12 as their characteristics. Earthquake statistical studies showed that large oceanic events in transform
13 faults, fracture zones, and intraplate regions release low energy levels in their aftershock sequences
14 (Houston et al., 1993; Boettcher and Jordan, 2001; Antolik et al., 2006). Boettcher et al. (2012) found
15 that earthquakes on transform faults have an order of magnitude fewer aftershocks than intraplate
16 events. According to some authors, a low aftershock-to-mainshock energy ratio indicates an efficient
17 rupture or complete stress drop in the mainshock presupposing a weak fault (Hwang and Kanamori,
18 1992; Velasco et al., 2000). Many factors can affect the aftershock productivity. For example, the age
19 of the lithosphere and the heat flux have a direct influence on the rock strength (Antolik et al., 2006),
20 thus, explaining the low energy release in the aftershock sequence of oceanic events. The observed low
21 aftershock energy seems to be a common feature of oceanic earthquakes (Antolik et al., 2006). In this
22 regard, we studied the 1 May 1997 ($M_w = 6.9$) strike-slip event in the Rivera Transform and its largest
23 aftershock ($M_w = 5.3$). By considering the energy magnitude as $\log E = 1.5 M_w + 11.8$, we obtain that
24 the energy of the mainshock is 1.41×10^{22} ergs, and the energy of the largest aftershock is 5.62×10^{19}

1 ergs resulting in an aftershock-to-mainshock energy ratio of 0.003. This value is considered as low and
2 representative of strike-slip events, as shown by the comparison with the results reported by Velasco et
3 al. (2000).

4

5 A similar analysis comes from Båth's law by considering the magnitude difference between the
6 mainshock, and the largest aftershock. We determined that the magnitude difference for the 1997 event
7 is 1.6, which is higher than the theoretical value of 1.2. Both magnitude difference and the aftershock-
8 to-mainshock energy ratio showed large scatter (e.g., Velasco et al., 2000; Utsu, 2002), and results
9 ought to be taken with caution. The aftershock decay rate is the product of the strain relaxation around
10 the rupture plane. Aftershock studies have shown that oceanic ridges are prone to having larger p -
11 values than those of subduction zone regimes due to the high temperature of the oceanic crust which
12 results in rapid strain release (Kisslinger, 1996; Rabinowitz and Steinberg, 1998; Klein et al., 2006).
13 According to previous studies, extremely high p -values ($p > 2$), and short aftershock durations are
14 related to high temperatures (Bohnenstiehl et al., 2002; Simão et al., 2010), and/or migration of
15 hydrothermal fluids (Goslin et al., 2005). We found a p -value of 0.67 ± 0.33 for the 1 May 1997 ($M_w =$
16 6.9) strike-slip event in the Rivera transform. This p -value is consistent with other oceanic regions, but
17 it does not seem to conform to a high-temperature regime.

18

19 Regarding the magnitude distribution of oceanic events, our b -value estimates are in agreement with
20 global oceanic studies but differ from local studies. For example, along the East Pacific Rise (in the
21 latitude interval of $5^\circ\text{N} < \text{Lat} < 9.90^\circ\text{N}$), b -value estimations fluctuate from 1.10 to 2.50 (Bohnenstiehl
22 et al., 2008). Bohnenstiehl et al. (2008) determined the b -value of 9000 microearthquakes with
23 magnitudes in the range of $-1.5 - 1.0$ located in the southern part of our study zone. Due to this overlap,
24 we compare their results with our results for region R5. For this region, we obtained a b -value of 0.94

1 with a M_c of 4.2, whilst they found that the b -value approaches 2.5 at very shallow depths (< 0.3 km)
2 (with $M_c = -1.3$). At depths of 0.5 to 1.5 km, the b -values drops to a value of 1.10 (with $M_c = -0.4$).
3 According to Bohnenstiehl et al. (2008) at very shallow depths, the uppermost oceanic crust is
4 structurally heterogeneous because of the extrusion of lava, and the repeated emplacement of sheeted
5 dikes. As a consequence, there is a large proportion of small versus large earthquakes resulting in high
6 b -values. The b -values decreases with depth due to the decreasing heterogeneity, and/or changes in
7 ambient stress levels. Considering that events in our catalogue for R5 occur at a different depth
8 interval, and assuming the decreasing heterogeneity, fewer magnitude events would be expected
9 (reducing the b -value). Another explanation for the differences between our results and the results of
10 Bohnenstiehl et al. (2008) is that the magnitude ranges of the earthquake catalogues are extremely
11 different. This highlights how the b -value is affected by magnitude completeness.

12
13 Statistical studies suggested that β -value mainly takes values between 0.60 and 0.70 for a global range
14 (Kagan, 2002). Our estimates of β agree with global oceanic studies. It is essential to discuss the
15 tectonic implications of this parameter. Bird et al. (2002) also found a dependence of β -value on the
16 relative plate velocity. According to them, the β -value is higher (with $M_m = 7.1$) when the velocity is $<$
17 36 mm/yr than when the velocity is > 67 mm/yr (with $M_m = 6.6$) for spreading ridges, and oceanic
18 transform faults, respectively. These observations are in agreement with our estimate of $\beta = 0.64$, and
19 M_m of 6.6 for oceanic earthquakes in Mexico (Figure 5). For intraplate events, we obtained a $\beta > 0.70$.
20 According to Kagan (2010), β -values > 0.70 may be related to the mix of earthquake populations with
21 different maximum magnitudes (M_m). In the case of intraplate events, we associated the somewhat high
22 β -values with the mix of some intraplate, and mid-ocean- transform events. This could be related to
23 incorrect hypocenter locations due to the difficulty of precisely locating oceanic events by the
24 landbased networks.

1

2 The seismicity models based on non-extensivity consider the interaction of two irregular fault surfaces
3 (asperities), and rock fragments filling them. However, these models differ in their assumption of how
4 energy is stored in the fragments, and the asperities. This difference is expressed through the constant
5 a , which represents the proportionality between the released energy E , and the fragment size r . This
6 explains the difference in a parameter between Telesca's and Silva's models (Fig. 5). Both models
7 showed that a for TF-MOR is higher than a for the INT events (Fig. 5). This implies that more energy
8 is released for TF-MOR earthquakes. On the other hand, the q -value indicates if the physical state of a
9 seismic area moves away from equilibrium. The physical state is at equilibrium when q is equal to 1,
10 and as q increases, the system is in an instability state in which a more significant amount of seismic
11 energy is released.

12

13 Finally, we discuss the focal mechanisms and the calculated state of stress for oceanic earthquakes in
14 Mexico. Focal mechanisms provide useful information about the structure, and settings of faults, and
15 can describe the crustal stress field in which earthquakes take place. Our analysis is limited because we
16 only used focal mechanisms based on teleseismic data. The teleseismic detection threshold for oceanic
17 events in the East Pacific Rise is dependent on the region of the EPR. For example, Riedesel et al.
18 (1982) report a magnitude detection threshold in the range of 4.0 – 5.0. For the Quebrada, Discovery,
19 and Gofar faults, the CMT catalogue is only complete to $M_w = 5.4$. (McGuire, 2008; Wolfson-Schwehr
20 et al., 2014). Another limitation of our study is that we combine different types of earthquakes into a
21 single region, resulting in inaccurate estimations of the stress state for that specific region. Under these
22 circumstances, our study provides information on the stress field of major structures or the stress
23 associated with the dominant types of earthquakes.

24

1 In oceanic environments, the largest magnitude events along transform fault or intraplate earthquakes
2 usually show strike-slip mechanisms (Wiens and Stein, 1984; Kawasaki et al., 1985). In the adjacent
3 areas to the oceanic ridges where the oceanic lithosphere is young, Wiens and Stein (1984) report a
4 large variety of focal mechanisms and stress orientations. For example, in the East Pacific Rise, in the
5 Mexican territory, Wiens and Stein (1984) reported thrust and normal mechanism solutions for near
6 ridge intraplate seismicity. This explains the strike-slip with normal components, as well as thrust
7 events in regions R3, R4, and R5 (Fig. 7). In R3, and R4 (Fig. 7), the maximum horizontal axes
8 (compression) of thrust events show a preferred orientation perpendicular to the spreading direction.
9 On the other hand, in region R5 (Fig. 7), the compression axes, showed a weak preferred alignment
10 with respect to the spreading direction. In the Rivera transform, focal mechanisms showed right lateral
11 strike-slip motion implying oblique horizontal stresses (Fig. 7). Although most of the events in the
12 Rivera transform (R2 in Fig. 7) are strike-slip events, some events with unusual mechanisms have been
13 reported (normal faulting events) (Wolfe et al., 1993). Normal faulting events may be related to
14 extensional offsets or internal deformation of the Rivera plate (Wolfe et al., 1993).

15

16 **6 Conclusions**

17

18 We analyzed the seismicity of oceanic events in the Pacific oceanic regime of Mexico. Oceanic
19 earthquakes were classified into two different categories: intraplate oceanic (INT), and transform faults
20 zone and mid-ocean ridges events (TF-MOR), respectively. We conducted a stress state estimation for
21 the different regions. Because of the combination of different types of earthquakes into the regions, our
22 results only provide information on the stress field of major structures or the stress associated with the
23 dominant types of earthquakes. It is important to be aware of this limitation in order to avoid an over-
24 interpretation of the results. TF-MOR events have strike-slip, strike-slip with normal and reverse

1 components, normal and normal-faulting with the strike-slip component, and reverse focal
2 mechanisms. On the other hand, INT events have only normal, and normal-faulting with strike-slip
3 component focal mechanisms. The stress field from INT, and TF-MOR events agree with global
4 studies. Regarding the aftershock productivity, we found that the aftershock decay rate of the 1 May
5 1997 ($M_w = 6.9$) strike-slip event in the Rivera transform is also consistent with oceanic p -value
6 estimations. Although the limitation of the catalogues used, our results provided a comprehensive
7 insight into the seismicity of oceanic environments. The main problem is the location uncertainty and
8 mislabelling of the earthquakes. The b -value for INT events (1.17) is higher than that for TF-MOR
9 events (0.82). Our b -values estimations are in agreement with other regional studies but differ from b -
10 value estimates based on microseismicity studies. Our b -value estimates for mid-ocean ridge/transform
11 fault environments are lower ($0.72 < b < 1.30$) than those derived from microseismicity studies ($1.1 < b$
12 < 2.5). Our results also showed that TF-MOR events mostly follow a tapered Gutenberg-Richter
13 distribution.

14

15 From the non-extensivity analysis, we observed that TF-MOR events are farther from the equilibrium
16 than INT events. Thus high q -values take place in mid-ocean ridges and transform faults zones. This
17 means that mid-ocean ridge and transform faults are able to produce more seismicity. Low q -values are
18 also reported during relatively quiet periods, characterized mainly by the occurrence of small
19 magnitude events. This can be an explanation for the low q -values of regions R1 and R5. Our results
20 also showed that a -values are higher for TF-MOR events than for INT events using both models. This
21 implies that more earthquakes with larger magnitude occur (or more energy is released) in mid-ocean
22 ridge/transform fault environments than in an oceanic continental environment. Telesca's model fits
23 better with the cumulative magnitude distribution functions making a better option to study the oceanic
24 seismicity in Mexico.




Figure 1

Main tectonic features in the oceanic environment off the Pacific coast of Mexico discussed in the text.

CO is the Cocos plate, NA is the North American plate, PA is the Pacific plate, RI is the Rivera microplate, TMVB is the Trans-Mexican Volcanic Belt, TFZ is the Tamayo fracture zone, EPR-RS is the East Pacific Rise Rivera segment, EPS-CS is the East Pacific Rise Cocos segment, and RT is the Rivera Transform. Blue triangles are volcanoes. Dashed lines show contour lines of the subducted slab. Arrows indicate the motion of the PA, CO, and RI plates. R1 to R6 are the regions in which the study area was divided for analyzing stress and seismicity characteristics. Red number indicates the slipping rates. Pink numbers indicate convergence rates, and black numbers indicate spreading rates.

11
12
13




Figure 2

Seismicity in the oceanic environment off the Pacific coast of Mexico from 1899 to 2017. The size of the circles represents magnitude. Brown circles are relevant historical earthquakes shown in Table 1 with $M > 6.8$. Red circles are intraplate oceanic events, and green circles are transform faults zone, and mid-ocean ridges earthquakes. Epicenters were compiled from the Mexican National Service (SSN), and the International Seismological Center (ISC) catalogues.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15





Figure 3

1
 2
 3
 4 Main statistical characteristics for intraplate oceanic events (INT). Magnitude earthquake histograms
 5 (upper left panel); frequency magnitude distributions with M_c , and b -values (upper right panel). The
 6 normalized cumulative number of events as function of magnitude for intraplate oceanic events (INT)
 7 (lower panels). Color curves show the best fit for the non-extensivity parameters q , and a for the
 8 Telesca's (red lines), and the Silva's (green lines) models, respectively.


9
 10

1 (a)



2

3 (b)




4

5

Figure 4


6 Main statistical characteristics for the transform faults zone, and mid-ocean ridges events (TF-MOR)
7 (regions R1 to R5) (upper panels). Magnitude earthquake histograms, and frequency magnitude
8 distributions with M_c , and b -values for each of the different subregions shown in Fig. 8 (lower panels).



1
2
3
4
5
6
7
8
9

Figure 5

The cumulative annual seismic moment frequency distribution for the transform faults zone, and mid-ocean ridges events (TF-MOR) (regions R1 to R5) (upper panels). The blue lines are the moment tapered Gutenberg Richter distributions. The red lines represent the ordinary moment Gutenberg Richter distributions. The subregions that do not follow an ordinary moment Gutenberg Richter distribution are subregions R1 and R2 (lower panels).




1

2

Figure 6

3 The normalized cumulative number of events as function of magnitude for the transform faults zone,
 4 and mid-ocean ridges events (TF-MOR). Blue triangles show the completeness magnitude (M_c). Red
 5 curves show the best fit for the non-extensivity parameters q , and a for the Telesca's model (red lines).
 6 Green curves show the best fit for the non-extensivity parameters q , and a for the Silva's model (green
 7 lines).

1 (a)



2

3

4

5

6

7


8

9

10

11

1 (b)




2

3

Figure 7


4 Focal mechanism solutions of oceanic earthquakes in Mexico reported by the Global CMT catalogue
5 from 1976 to 2017. a) Focal mechanisms are divided into 6 regions (R1 to R6) for the stress inversion
6 analysis. b) Focal mechanism classification based on the Kaverina et al. (1996) projection technique
7 implemented by Álvarez-Gómez (2015): reverse, reverse with lateral component, strike-slip with
8 reverse component, strike-slip, strike-slip with normal component, normal with lateral component, and
9 normal (R, R-SS, SS-R, SS, SS-N, N-SS, and N, respectively).

1 (a)



2
3
4
5
6
7

1 (b)



2

3


Figure 8

4 Orientation of horizontal axes. a) maximum horizontal stresses (S_H); b) minimum horizontal stresses

5 (S_h).


6

7



2
3
4
5
6
7
8
9
10
11
12
13

1



2
3
4
5

Figure 9

6 Mohr's circle diagrams for all the regions (left column). *P*- and *T*-axes distributions for all the regions
7 (right column). Red circles represent pressure, while blue crosses represent tension.

8
9
10

1 **Table 1**

2 Major oceanic earthquakes in Mexico ($M > 6.8$)

Event	Date dd/mm/yyyy	Time hh:mm:ss	Lon (°)	Lat (°)	M_s	M_w	M_0 [Nm]	Reference
1	14/01/1899	02:36:00	-110.00	20.00	7.0			1
2	17/12/1905	05:27:00	-113.00	17.00	7.0	7.0	4.40×10^{19}	2
3	10/04/1906	21:18:00	-110.00	20.00	7.1	7.1	6.20×10^{19}	2
4	31/10/1909	10:18:00	-105.00	8.00	6.9			3
5	31/05/1910	04:54:00	-105.00	10.00	7.0			3
6	29/10/1911	18:09:00	-101.00	11.00	6.8			3
7	16/11/1925	11:54:00	-107.00	18.00	7.0			4
8	28/05/1936	18:49:01	-103.60	10.10	6.8			3
9	30/06/1945	05:31:21	-115.80	16.70	6.8			3
10	04/12/1948	04:00:00	-106.50	22.00	6.9			3
11	29/09/1950	06:32:00	-107.00	19.00	7.0			4
12	01/05/1997	11:37:40	-107.15	18.96	6.8	6.9	2.77×10^{19}	5

3 1 Data from the Decade of North American Geology Project (DNA) of the National Geophysical Data
4 Center (NGDC), and the Geological Society of America.

5 2 Pacheco and Sykes (1992)

6 3 ISC earthquake catalogue

7 4 Abe (1981)

8 5 Global CMT catalogue

9

10

11

12 **Table 2**

13 Statistical parameters

Type	M_c	b -value	q_s -value	a_s -value	q_T -value	a_T -value
INT	4.4	0.89	1.60	6.69×10^{12}	1.39	2.27×10^6
TF-MOR (R1- R5)	4.1	0.64	1.60	3.22×10^{13}	1.41	3.55×10^6
R1	4.1	0.72	1.62	3.22×10^{13}	1.43	2.53×10^6
R2	4.0	0.77	1.62	1.24×10^{13}	1.44	3.11×10^6
R3	4.4	1.30	1.57	6.81×10^{12}	1.31	2.98×10^6
R4	4.4	0.75	1.70	1.12×10^{13}	1.52	2.94×10^6
R5	4.3	0.94	1.63	5.79×10^{12}	1.38	3.15×10^6

14 INT are intraplate oceanic events; TF-MOR are transform faults zone, and mid-ocean ridges events; M_c
15 is the completeness magnitude; b is the slope of the Gutenberg-Richter distribution; q_s , a_s , q_T , and a_T
16 are the non-extensive parameters based on Silva et al. (2006), and Telesca (2011), respectively.

17

18

19

20

21

22

1 **Table 3**

2 Aftershocks characteristics of 1 May 1997 event

Date	M_m	M_a	D	p -value	c	k
01/05/1997	6.9	5.3	1.6	0.67 ± 0.33	0.00 ± 0.53	2.12 ± 1.53

3 M_m is the magnitude of the mainshock; M_a is the magnitude of the largest aftershock; D is the
 4 difference in magnitudes of the mainshock, and its largest aftershock; p , c , and k are the coefficients of
 5 the Omori's law.

6
7
8 **Table 4**

9 Stress inversion results

σ_1 Azimuth/plunge	σ_2 Azimuth/plunge	σ_3 Azimuth/plunge	SH_{max}	R	Region
169°/16°	2°/73°	260°/4°	169°	0.37	1 ^a
156°/0°	62°/83°	246°/7°	157°	0.58	2 ^a
157°/4°	31°/84°	247°/5°	157°	0.63	3 ^a
197°/3°	302°/76°	106°/13°	22°	0.84	4 ^a
299°/6°	44°/69°	207°/20°	120°	0.73	5 ^a
247°/80°	39°/9°	130°/5°	45°	0.73	6 ^a

10 Stress ratio is defined by $R = (\sigma_1 - \sigma_2) / (\sigma_1 - \sigma_3)$; ^a, stress inversion based on Vavryčuk (2014), and
 11 Lund and Townend (2007). Location of the regions are shown in Fig. 1.

12

13

14

15

16

17

18

19

20

21

22

23

24

1 **Code availability**

2

3 Generic Mapping Tools (GMT5), Available at: <http://gmt.soest.hawaii.edu/>, last access: 13 January
4 2020.

5

6 Get_GR_parameters.m, Available at: <https://jaolive.weebly.com/codes.html>, last access: 23 December
7 2019.

8

9 FMC, Available at: [https://josealvarezgomez.wordpress.com/2014/04/22/fmc-a-python-program-to-
10 manage-classify-and-plot-focal-mechanism-data/](https://josealvarezgomez.wordpress.com/2014/04/22/fmc-a-python-program-to-manage-classify-and-plot-focal-mechanism-data/), last access: 13 January 2020.

11

12 Stressinverse_1.1, Available at: <https://www.ig.cas.cz/en/stress-inverse/>, last access: 13 January 2020.

13

14 ZMAP, Available at: [http://www.seismo.ethz.ch/en/research-and-teaching/products-
15 software/software/ZMAP/](http://www.seismo.ethz.ch/en/research-and-teaching/products-software/software/ZMAP/), last access: 13 January 2020.

16

17 **Data availability**

18 Earthquake catalogues data available at:

19

20 Earthquake catalogue of the Servicio Sismológico Nacional, <http://www.ssn.unam.mx/>, last access: 13
21 January 2020.

22

23 Earthquake catalogue of the International Earthquake Center:

24 <http://www.isc.ac.uk/iscbulletin/search/catalogue/>, last access: 13 January 2020.

1

2 **Team list**

3 Q. Rodríguez-Pérez *E-mail:* quetza@geociencias.unam.mx

4 V.H. Márquez-Ramírez *E-mail:* marvh@geociencias.unam.mx

5 F.R. Zúñiga *E-mail:* ramon@geociencias.unam.mx

6

7 **Author contribution**

8

9 Quetzalcoatl Rodríguez-Pérez, Víctor H. Márquez, and F. Ramón Zúñiga designed the idea and
10 discussed the results. Quetzalcoatl Rodríguez-Pérez developed the methodology and performed the
11 analyses. Quetzalcoatl Rodríguez-Pérez prepared the manuscript with contributions from all co-authors.

12

13 **Competing interests**

14

15 The authors declare that they have no conflict of interest.

16

17

18 *Acknowledgments* We thank the Mexican National Seismological Service (SSN) for providing us with
19 the earthquake catalogue. Station maintenance, data acquisition, and distribution is thanks to its
20 personnel. Quetzalcoatl Rodríguez-Pérez was supported by the Mexican National Council for Science
21 and Technology (CONACYT) (Catedras program- project 1126).

22

23

24

1 **References**

2

3 Abe, K.: Magnitudes of large shallow earthquakes from 1904 to 1980, *Phys. Earth Planet. Int.*, 27, 72-
4 92, 1981.

5

6 Abercrombie, R.E., and Ekström, G.: Earthquake slip on oceanic transform faults, *Nature*, 410, 74-77,
7 2001.

8

9 Abercrombie, R.E., and Ekström, G.: A reassessment of the rupture characteristics of oceanic transform
10 earthquakes, *J. Geophys. Res.*, 108, B5, 2003.

11

12 Aki, K.: Maximum likelihood estimate of b in the formula $\log(N) = a - bM$ and its confidence limits, *B.*
13 *Earthq. Res. I. Tokyo*, 43, 237-239, 1965.

14

15 Álvarez-Gómez, J.A.: FMC: A program to manage, classify and plot focal mechanism data. Version
16 1.01, 2015.

17

18 Antolik, M., Abercrombie, R., Pan J., and Ekström, G: Rupture characteristics of the 2003 M_w 7.6 mid-
19 Indian Ocean earthquake: implications for seismic properties of young oceanic lithosphere, *J. Geophys.*
20 *Res.*, 111, B04302, 2006.

21

22 Bandy, W.L.: Geological and geophysical investigation of the Rivera-Cocos plate boundary:
23 implications for plate fragmentation, Ph.D. thesis, Texas A&M University, College Station, 195pp.,
24 1992.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Bandy, W.L., Michaud, F., Mortera Gutierrez, C.A., Dymant, J., Bourgois, J., Royer, J.Y., Calmus, T., Sosson, M., and Ortega-Ramirez, J.: The Mid-Rivera-Transform discordance: morphology and tectonic development, *Pure Appl. Geophys.*, 168, 1391-1413, 2011.

Bergman, E.A., and Solomon, S.C.: Oceanic intraplate earthquakes: implications for local and regional intraplate stress, *J. Geophys. Res.*, 85, B10, 5389-5410, 1980.

Bergman, E.A.: Intraplate earthquakes and the state of stress in oceanic lithosphere, *Tectonophysics*, 132, 1-35, 1986.

Beroza, G.C., and Jordan, T.: Searching for slow and silent earthquakes using free oscillations, *J. Geophys. Res.*, 95, B3, 2485-2510, 1990.

Bird, P., Kagan, Y. Y., and Jackson, D. D.: Plate tectonics and earthquake potential of spreading ridges and oceanic transform faults. In S. Stein and J.T. Freymueller (Eds.), *Plate Boundary Zones*, Geodynamics Series, American Geophysical Union, 203-218, 2002.

Boettcher, M.S., and Jordan, T.H.: Seismic behavior of oceanic transform faults, Fall Meeting, American Geophysical Union (AGU), San Francisco, California, December 10-14, S32E-07, 2001.

Boettcher, M.S., and Jordan, T.H.: Earthquake scaling relations for mid-ocean ridge transform faults, *J. Geophys. Res.*, 109, B12302, 2004.

- 1 Boettcher, M.S., Hirth, G., and Evans, B.: Olivine friction at the base of oceanic seismogenic zones, J.
2 Geophys. Res., 112, B01205, 2007.
- 3
- 4 Boettcher, M.S., and McGuire, J.J.: Scaling relations for seismic cycles on mid-ocean ridge transform
5 faults, Geophys. Res. Lett., 36, L21301, 2009.
- 6
- 7 Boettcher, M.S., Wolfson-Schwehr, M.L., Forestall, M., and Jordan, T.H.: Characteristics of oceanic
8 strike-slip earthquakes differ between plate boundary and intraplate settings, Fall Meeting, American
9 Geophysical Union (AGU), San Francisco, California, December 3-7, 7245 Seismology, 2012.
- 10
- 11 Bohnenstiehl, D.R., Tolstoy, M., Dziak, R.P., Fox, C.G., and Smith, D.K.: Aftershock sequences in the
12 mid-ocean ridge environment: an analysis using hydroacoustic data, Tectonophysics, 354, 49-70, 2002.
- 13
- 14 Bohnenstiehl, D.R., Tolstoy, M., and Chapp, E.: Breaking into the plate: A 7.6 M_w fracture-zone
15 earthquake adjacent to the central Indian Ridge, Geophys. Res. Lett., 31, L02615, 2004.
- 16
- 17 Bohnenstiehl, D.R., Waldhauser, F., and Tolstoy, M.: Frequency-magnitude distribution of
18 microearthquakes beneath the 9°50'N region of the East Pacific Rise, October 2003 through April
19 2004, Geochem. Geophys. Geosyst., 9, Q10T03, 2008.
- 20
- 21 Choy, G.L., and Boatwright, J.: Global patterns of radiated seismic energy and apparent stress, J.
22 Geophys. Res., 100, 18205-18226, 1995.
- 23
- 24 Choy, G.L., and McGarr, A.: Strike-slip earthquakes in the oceanic lithosphere: Observations of

1 exceptionally high apparent stress, *Geophys. J. Int.*, 100, 18205-18226, 2002.

2

3 Cowie, P.A., Scholz, C.H., Edwards, M., and Malinverno, A.: Fault strain and seismic coupling on Mid-
4 Ocean Ridges, *J. Geophys. Res.*, 98, 17911-17920, 1993.

5

6 Davis, S.D., and Frohlich, C.: Single-link cluster analysis, synthetic earthquake catalogues and
7 aftershock identification, *Geophys. J. Int.*, 104, 289-306, 1991.

8

9 DeMets , C., Gordon, R.G., Argus, D.F., and Stein, S.: Effect of recent revisions to the geomagnetic
10 reversal time scale on estimate of current plate motions, *Geophys. Res. Lett.*, 21, 2191-2194, 1994.

11

12 Dziewonski, A.M., Chou, T.A., and Woodhouse, J.H.: Determination of earthquake source parameters
13 from waveform data for studies of global and regional seismicity, *J. Geophys. Res.*, 86, 2825-2852,
14 1981.

15

16 Ekström, G., Nettles, M., and Dziewonski, A.M.: The global CMT project 2004-2010: centroid-
17 moment tensors for 13,017 earthquakes, *Phys. Earth Planet. Int.*, 200-201, 1-9, 2012.

18

19 Frohlich, C.: Practical suggestions for assessing rates of seismic-moment release, *B. Seismol. Soc.*
20 *Am.*, 97, 1158-1166, 2007.

21

22 Gephart, J.W., and Forsyth, D.W.: An improved method for determining the regional stress tensor using
23 earthquake focal mechanism data: application to the San Fernando earthquake sequence, *J. Geophys.*
24 *Res.*, 89, 9305-9320, 1984.

1

2 Goslin, J., et al.: Extent of Azores plume influence on the Mid-Atlantic Ridge north of the hotspot,
3 *Geology*, 27(11), 991-994, 1999.

4

5 Goslin, J., Lourenço, N., Dziak, R.P., Bohnenstiehl, D.R., Haxel, J., and Luis, J.: Long-term seismicity
6 of the Reykjanes Ridge (North Atlantic) recorded by a regional hydrophone array, *Geophys. J. Int.*,
7 162, 516-524, 2005.

8

9 Gutenberg, B., and Richter, C.F.: Frequency of earthquakes in California, *B. Seismol. Soc. Am.*, 34,
10 185-188, 1944.

11

12 Houston, H., Anderson, H., Beck., S. L., Zhang, J., and Schwartz, S.: The 1986 Kermadec earthquake
13 and its relation to plate segmentation, *Pure Appl. Geophys.*, 140, 331-364, 1993.

14

15 Hwang, L.J., and Kanamori, H.: Rupture process of the 1987-1988 Gulf of Alaska earthquake
16 sequence, *J. Geophys. Res.*, 97, 19881-19908, 1992.

17

18 Ihmlé, P.F., and Jordan, T.H.: Teleseismic search for slow precursors to large earthquakes, *Science*, 266,
19 1547-1551, 1994.

20

21 Ishimoto, M., and Iida, K.: Observations of earthquakes registered with the microseismograph
22 constructed recently, *B. Earthq. Res. I. Tokyo*, 17, 443-478, 1939.

23

24 Kagan, Y.Y.: Seismic moment-frequency relation for shallow earthquakes: regional comparisons, *J.*

- 1 Geophys. Res., 102, 2835-2852, 1997.
- 2
- 3 Kagan, Y.Y.: Universality of the seismic moment-frequency relation, Pure Appl. Geophys., 155, 537-
4 573, 1999.
- 5
- 6 Kagan, Y.Y., and Jackson, D.D.: Probabilistic forecasting of earthquakes, Geophys. J. Int., 143, 438-
7 453, 2000.
- 8
- 9 Kagan, Y.Y., and Schoenberg F.: Estimation of the upper cutoff parameter for the tapered pareto
10 distribution, J. Appl. Probab., 38A, 158-175, 2001.
- 11
- 12 Kagan, Y.Y.: Seismic moment distribution revisited: I. Statistical results, Geophys. J. Int., 148, 520-
13 541, 2002.
- 14
- 15 Kagan, Y.Y.: Earthquake size distribution: power-law with exponent $\beta \equiv 1/2?$, Tectonophysics, 490,
16 103-114, 2010.
- 17
- 18 Kanamori, H., and Stewart, G.S.: Mode of the strain release along the Gibbs fracture zone, Mid-
19 Atlantic Ridge, Phys. Earth Planet. Int., 11, 312-332, 1976.
- 20
- 21 Kaverina, A.N., Lander, A.V., and Prozorov, A.G.: Global creepex distribution and its relation to
22 earthquake-source geometry and tectonic origin, Geophys. J. Int., 125, 249-265, 1996.
- 23
- 24 Kawasaki, I., Kawahara, Y., Takata, I., and Kosugi, I.: Mode of seismic moment release at transform

1 faults, *Tectonophysics*, 118, 313-327, 1985.

2

3 Kisslinger, C.: Aftershocks and fault-zone properties, *Adv. Geophys.*, 38, 1-36, 1996.

4

5 Klein, F.W., Wright, T., and Nakata, J.: Aftershock decay, productivity, and stress rates in Hawaii:
6 indicators of temperature and stress from magma sources, *J. Geophys. Res.*, 111(B7), B07307, 2006.

7

8 Läderach, Ch.: Seismicity of ultraslow spreading mid-ocean ridges at local, regional and teleseismic
9 scales: A case study of contrasting segments, Ph.D thesis, University of Bremen, 116 pp., 2011.

10

11 Lund, B., and Townend, J.: Calculating horizontal stress orientations with full or partial knowledge of
12 the tectonic stress tensor, *Geophys. J. Int.*, 270, 1328-1335, 2007.

13

14 McGuire, J.J., Ihmlé, P.F., and Jordan, T.H.: Time-domain observations of a slow precursor to the 1994
15 Romanche transform earthquake, *Science*, 274, 82-85, 1996.

16

17 McGuire, J.J., Boettcher M.S., and Jordan, T.H.: Foreshock sequences and short-term earthquake
18 predictability on East Pacific Rise transform faults, *Nature*, 434, 457-461, 2005.

19

20 McGuire, J.J.: Seismic cycles and earthquake predictability on East Pacific Rise transform faults, *B.*
21 *Seismol. Soc. Am.*, 98, 1067-1084, 2008.

22

23 McGuire, J.J., Collins, J.A., Gouédard, P., Roland, E., and Lizarralde, D.: Variations in earthquake
24 rupture properties along the Gofar transform fault, East Pacific Rise, *Nat. Geosci.*, 5, 336-341, 2012.

1

2 Mogi, K.: Magnitude-frequency relation for elastic shocks accompanying fractures of various materials
3 and some related problems in earthquakes, B. Earthq. Res. I. Tokyo, 40, 831-853, 1962.

4

5 Molchan, G., Kronrod, T., and Panza, G.F.: Multi-scale seismicity model for seismic risk, B. Seismol.
6 Soc. Am., 87, 1220-1229, 1997.

7

8 Nelder, J.A., and Mead, R.: A simplex method for function minimization, Comput. J., 7, 308-313, 1965.

9

10 Nuannin, P., Kulhanek, O., and Persson, L.: Variations of b -value preceding large earthquakes in the
11 Andaman-Sumatra subduction zone, J. Asian Earth Sci., 61, 237-242, 2012.

12

13 Okal, E.A., and Stewart, L.M.: Slow earthquakes along oceanic fracture zones: evidence for
14 asthenospheric flow away from hotspots?, Earth Planet. Sci. Lett., 57, 75-87, 1992.

15

16 Olive, J-A.: Get_GR_parameters.m Matlab function for analysis of earthquake catalogs. Available at:
17 <https://jaolive.weebly.com/codes.html>, 2016.

18

19 Pacheco, J.F., and Sykes, L.R.: Seismic moment catalog of large shallow earthquakes, 1900 to 1989, B.
20 Seismol. Soc. Am., 82, 1306-1349, 1992.

21

22 Papadakis, G., Vallianatos, F., and Sammonds, P.: Evidence of nonextensive statistical physics behavior
23 of the Hellenic subduction zone seismicity, Tectonophysics, 608, 1037-1048, 2013.

24

1 Pockalny, R.A., Fox, P.J., Fornari, D.J., McDonald, K., and Perfit, M.R.: Tectonic reconstruction of the
2 Clipperton and Siqueiros Fracture zones: evidence and consequences of plate motion change for the
3 last 3Myr, *J. Geophys. Res.*, 102, 3167-3181, 1997.
4
5 Rabinowitz, N., and Steinberg, D.M.: Aftershock decay of the three recent strong earthquakes in the
6 Levant, *B. Seismol. Soc. Am.*, 88, 1580-1587, 1998.
7
8 Riedesel, M., Orcutt, J.A., McDonald, K.C., and McClain, J.S.: Microearthquakes in the Black Smoker
9 Hydrothermal Field, east Pacific Rise at 21° N, *J. Geophys. Res.*, 87, 10613-10623, 1982.
10
11 Rodríguez-Pérez, Q, and Zúñiga, F. R.: Seismicity characterization of the Maravatio-Acambay and
12 Actopan regions, central Mexico, *J. S. Am. Earth Sci.*, 76, 264-275, 2017.
13
14 Rodríguez-Pérez, Q, and Zúñiga, F. R.: Imaging *b*-value depth variations within the Cocos and Rivera
15 plates at the Mexican subduction zone, *Tectonophysics*, 734, 33-43, 2018.
16
17 Roland, E., Behn, M.D., and Hirth, G.: Thermal-mechanical behavior of oceanic transform faults:
18 Implications for the spatial distribution of seismicity, *Geochem. Geophys. Geosyst.*, 11, Q07001, 2010.
19
20 Scholz, C.H.: The frequency-magnitude relation of micro fracturing in rock and its relation to
21 earthquakes, *B. Seismol. Soc. Am.*, 58, 388-415, 1968.
22
23 Schorlemmer, D.S., Wiemer, S., and Wyss, M.: Variations in earthquake-size distribution across
24 different stress regimes, *Nature*, 437, 539-542, 2005.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Scordilis, E.M.: Empirical global converting M_s and m_b to moment magnitude, *J. Seismol.*, 10, 225-236, 2006.

Shcherbakov, R., Turcotte, D.L., and Rundle, J.B.: A generalized Omori's law for earthquakes aftershocks decay, *Geophys. Res. Lett.*, 31(11), L11613, 2004.

Silva, R., Franca, G., Vilar, C., and Alcaniz, J.: Nonextensive models for earthquakes, *Phys. Rev. E*, 73, 026102, 2006.

Smith, W.D.: The b -value as an earthquake precursor, *Nature*, 289, 136-139, 1981.

Smith, D.K., Tolstoy, M., Fox, C.G., Bohnenstiehl, D.R., Matsumoto, H., and Fowler, M.J.: Hydroacoustic monitoring of seismicity at the slow-spreading Mid-Atlantic Ridge, *Geophys. Res. Lett.*, 29(11), 2002.

Smith, D.K., Escartin, J., Cannat, M., Tolstoy, M., Fox, C.G., Bohnenstiehl, D.R., and Bazin, S.: Spatial and temporal distribution of seismicity along the northern Mid-Atlantic Ridge (15° - 35°), *J. Geophys. Res.*, 108(B3), 2003.

Simão, N., Escartín, J., Goslin, J., Haxel, J., Cannat, M., and Dziak, R.: Regional seismicity of the Mid-Atlantic Ridge: observations from autonomous hydrophone arrays, *Geophys. J. Int.*, 183, 1559-1578, 2010.

1 Sotolongo-Costa, O., and Posadas, M.A.: Fragment-asperity interaction model for earthquakes, Phys.
2 Rev. Lett., 92, 048501, 2004.
3
4 Stein, S., and Pelayo, A.: Seismological constraints on stress in the oceanic lithosphere, Philosophical
5 Transactions of the Royal Society of London A, 337, 53-72, 1991.
6
7 Sykes, L.R.: Mechanism of earthquakes and nature of faulting on the mid-oceanic ridges, J. Geophys.
8 Res., 72, 2131-2153, 1967.
9
10 Telesca, L.: Nonextensive analysis of seismic sequences, *Physica A*, 389, 1911-1914, 2009.
11
12 Telesca, L.: A non-extensive approach in investigating the seismicity of L'Aquila area (central Italy),
13 struck by the 6 April 2009 earthquake ($M_L = 5.8$), *Terra Nova*, 22, 87-93, 2010.
14
15 Telesca, L.: Tsallis-based nonextensive analysis of the Southern California seismicity, *Entropy*, 13,
16 1267-1280, 2011.
17
18 Tolstoy, M., Bohnenstiehl, D.R., and Edwards, M.H.: Seismic character of volcanic activity at the
19 ultraslow-spreading Gakkel Ridge, *Geology*, 29, 1139-1142, 2001.
20
21 Tsallis, C.: Possible generalization of Boltzmann-Gibbs statistics, *J. Stat. Phys.*, 52, 479-487, 1988.
22
23 Urbancic, T.I., Trifu, C.I., Long, J.M., and Young, R.P.: Space-time correlation of b -values with stress
24 release, *Pure Appl. Geophys.*, 139, 449-462, 1992.

1
2 Utsu, T.: A statistical study on the occurrence of aftershocks, *Geophysical Magazine*, 30, 521-605,
3 1961.
4
5 Utsu, T., Ogata, Y., and Mutsaers, R.S.: The centenary of the Omori formula for a decay law of
6 aftershock activity, *J. Phys. Earth.*, 43, 1-33, 1995.
7
8 Utsu, T.: Statistical features of seismicity, *International Handbook of Earthquake and Engineering*
9 *Seismology, Part A*. Academic Press, pp. 719-732, 2002.
10
11 Valverde-Esparza, S.M., Ramirez-Rojas, A., Flores-Marquez, E.L., and Telesca, L.: Non-extensivity
12 analysis of seismicity within four subduction regions in Mexico, *Acta Geophys.*, 60, 833–845, 2012.
13
14 Vavryčuk, V.: Iterative joint inversion for stress and fault orientations from focal mechanisms,
15 *Geophys. J. Int.*, 199, 69-77, 2014.
16
17 Velasco, A.A., Ammon, C.J., and Beck, S.L.: Broadband source modeling of the November 8, 1997,
18 Tibet ($M_w = 7.5$) earthquake and its tectonic implications, *J. Geophys. Res.*, 105(B12), 28065-28080,
19 2000.
20
21 Vere-Jones, D., Robinson, R., and Yang, W.Z.: Remarks on the accelerated moment release model:
22 problems of model formulation, simulation and estimation, *Geophys. J. Int.*, 144, 517-531, 2001.
23
24 Vilar, C.S., Franca, G., Silva, R., and Alcaniz, J.S.: Nonextensivity in geological faults?, *Physica A*,

1 377, 285-290, 2007.

2

3 Vallianatos, F.: A non-extensive approach to risk assessment, *Nat. Hazards Earth Syst. Sci.*, 9, 211-216,

4 2009.

5

6 Warren, N.W., and Latham, G.V.: An experimental study of the thermally induced microfracturing and

7 its relation to volcanic seismicity, *J. Geophys. Res.*, 75, 4455-4464, 1970.

8

9 Wesnousky, S.G.: The Gutenberg-Richter or characteristic earthquake distribution, which is it?, *B.*

10 *Seismol. Soc. Am.*, 84, 1940-1959, 1994.

11

12 Wiens, D.A., and Stein, S.: Intraplate seismicity and stresses in young oceanic lithosphere, *J. Geophys.*

13 *Res.*, 89, 11442-11464., 1984.

14

15 Wiemer, S., and Benoit, J.P.: Mapping the *b*-value anomaly at 100 km depth in the Alaska and New

16 Zealand subduction zones, *Geophys. Res. Lett.*, 23, 1557-1560, 1996.

17

18 Wiemer, S., and Wyss, M.: Minimum magnitude of completeness in earthquake catalogues: Examples

19 from Alaska, the western United States, and Japan, *B. Seismol. Soc. Am.*, 90, 859-869, 2000.

20

21 Wiemer, S.: A software package to analyze seismicity: ZMAP, *Seismol. Res. Lett.*, 72, 373-382, 2001.

22

23 Wiemer, S., Wyss, M.: Mapping spatial variability of the frequency-magnitude distribution of

24 earthquakes. In *Advances in geophysics* (Vol. 45, pp. 259-V). Elsevier, 2002.

1

2 Wolfe, C.J., Bergman, E.A., and Solomon, S.C.: Oceanic transform earthquakes with unusual
3 mechanisms or locations: relation to fault geometry and state of stress in the adjacent lithosphere, J.
4 Geophys. Res., 98, B9, 16187-16211, 1993.

5

6 Wolfson-Schwehr, M., Boettcher, M.S., McGuire, J.J., and Collins, J.A.: The relationship between
7 seismicity and fault structure on the Discovery transform fault, East Pacific Rise, *Geochem. Geophys.*
8 *Geosyst.*, 15, 3698-3712, 2014.

9

10 Wyss, M.: Towards a physical understanding of the earthquake frequency distribution, *Geophys. J. Roy.*
11 *Astron. Soc.*, 31, 341-359, 1973.