I acknowledge the important work realized by the authors. Based on my own reading of the manuscript and my experience in the field, I found that this contribution is of interest to the journal and may be suitable for publication. However, important revisions are required before its publication. The authors need to consider the very constructive suggestions provided by both

5 reviewers, and the manuscript has to be revised accordingly

10

Response: Thanks a lot for your affirmation. This is a very important paper for us that we really take much effort on it. We discussed and considered a lot when we did every step in this project. We take all the valuable comments by topical editor and reviewers very seriously. After the carefully and detailed revision, we believe that the paper has the remarkable progress in quality. And we really want to show

our gratitude to you with the real name in the acknowledgement, we have added this part in acknowledgement in the revised version.

1. In particular, the authors need to strengthen the discussions about the data quality and 15 robustness of the results, by quantifying the impact of applied corrections on data.

Response: This is very good questions we also considered when we did the gravity gradient data processing. We made three revisions in the paper for this comments.

- (1) The amplitude of the topography effects and the underground interface undulation effects is obvious,
 but topographic correction and underground interface undulation corrections balance each other out to a certain degree (Szwillus et al., 2016). In consideration of the gravity gradient effect caused by the far zones (the extended area outside the study area). We extended geographic area within E90°-E130° and N23°-N53° as the calculation area for corrections (Figure 13 e-f). And the obtained results in the study area are cut out from the calculation area. We have added the gravity gradient effects
 (Topographic and Interface undulation) with extended area in Fig. 13(e-f). The effect of the
- counteraction can be more clearly seen in the extended area.

(2) We have added gravity gradient corrections T_{zz} at two cross sections (Latitude 35° and Latitude 37.5°) in Fig. 13. The effect of the counteraction can be also observed at these two cross sections.

(3) We have added a table about the statistics for each gravity gradient effects in Table 3.

30

35

Revision: (Page 24, Paragraph 2)

The amplitude of the topography effects and the underground interface undulation effects is obvious, but the topographic correction and underground interface undulation correction balance each other to a certain degree (Szwillus et al., 2016). This counteraction is obvious in the extended study area, as shown in Fig. 13e (topographic correction) and Fig. 13f (underground interface undulation correction). The detailed statistical amplitude for each gravity gradient correction are summarized in Table 3. To present the different corrections more intuitively, several corrections T_{zz} at different cross sections of

latitude 35° and latitude 37.5° are presented in Fig. 13 (g-h).



40 Figure 13 Gravity gradient effects with extended area (e) Topographic effects T_{zz} , (f) Interface undulation T_{zz} (Sedimentary and Moho) effects. Several corrections T_{zz} at different cross sections (g) Latitude 35°, (h) Latitude 37.5°.

Gravity Gradient Effects (E)	Downward Continuation Maximum (E)	Downward Continuation Minimum (E)	Topographic Effects Maximum (E)	Topographic Effects Minimum (E)	Interface Undulation Effects Maximum (E)	Interface Undulation Effects Minimum (E)
T _{xx}	5.3	-5.7	6.3	-9.4	10.1	-4.9
$T_{\rm xz}$	7.2	-5.6	11.9	-5.6	5.2	-8.9
$T_{_{ m yy}}$	4.8	-3.6	7.2	-6.8	6.3	-6.7
T _{zz}	7.5	-6.5	13.8	-6.3	7.7	-14.1
Gravity Gradient Effects (E)	Long Wavelength Effects Maximum (E)	Long Wavelength Effects Minimum (E)	Corrected Effects Maximum (E)	Corrected Effects Minimum (E)		
T _{xx}	1.1	-0.8	8.7	-6.8		
$T_{\rm xz}$	1.2	-0.7	8.5	-4.7		
$T_{ m yy}$	1.4	-0.7	6.7	-6.6		
T ₇₇	1.5	-1.5	11.5	-12.3		

	Table 3	Statistics	for each	gravity	gradient	correction
--	---------	------------	----------	---------	----------	------------

50

55

45

2. More details are needed on the used methodology, and on provided information (e.g. Tables 1 and 2).

60 **Response:** (1) We have added the section "2.1 Kernel function calculation" in the section "2 Methodology". (Page 5)

(2) We have added the real densities of the true model, and RMS in both Table 1 and Table 2 as followings.

65 Revision: (Page 10)

Table 1 Conditions and results for initial density model one.

	True density at each layer (g/cm ³) (1 st to 4 th layer)	Initial density at each layer (g/cm ³) (1 st to 4 th layer)	Iterations (k)	RMS (E)	Maximum density at each layer (g/cm ³) (1 st to 4 th layer)
Test 1	(0.4, 0.7, 0.7, 1.0)	(0, 0, 0, 0)	10	7.35×10 ⁻¹¹	(0.34, 0.42, 0.53, 0.55)
Test 2	(0.4, 0.7, 0.7, 1.0)	(0.4, 0.4, 0.4, 0.4)	7	6.94×10 ⁻¹²	(0.51, 0.55, 0.66, 0.58)
Test 3	(0.4, 0.7, 0.7, 1.0)	(0.1, 0.2, 0.2, 0.3)	6	3.50×10 ⁻¹²	(0.36, 0.64, 0.67, 0.64)
Test 4	(0.4, 0.7, 0.7, 1.0)	(0.2, 0.4, 0.4, 0.7)	3	2.31×10 ⁻¹³	(0.38, 0.72, 0.76, 0.95)
	Minimum density	Average density			
	at each layer	at each layer			
	(g/cm^3)	(g/cm^3)			
	$(1^{st} to 4^{th} layer)$	$(1^{st} to 4^{th} layer)$			
Test 1	(0.23, 0.27, 0.34, 0.29)	(0.31, 0.34, 0.41, 0.35)			
Test 2	(0.42, 0.43, 0.51, 0.48)	(0.47, 0.50, 0.58, 0.55)			
Test 3	(0.32, 0.46, 0.33, 0.55)	(0.34, 0.55, 0.59, 0.58)			
Test 4	(0.33, 0.52, 0.57, 0.83)	(0.35, 0.59, 0.65, 0.91)			

70 (Page 12)

Table 2 Conditions and results for initial density model two.

	True density (Trapezoid, Cuboid) (g/cm ³)	Initial density (Trapezoid, Cuboid) (g/cm ³)	Iterations (k)	RMS (E)	Maximum density (g/cm ³)	Minimum density (g/cm ³)	Average density (g/cm ³)
Test 5	(0.8, 1.0)	(0, 0)	8	4.37×10 ⁻¹²	0.6887	0.1328	0.4225
Test 6	(0.8, 1.0)	(0.2, 0.2)	7	5.63×10 ⁻¹³	0.7121	0.1856	0.4886
Test 7	(0.8, 1.0)	(0.5, 0.5)	5	1.06×10 ⁻¹³	0.7556	0.4762	0.6256
Test 8	(0.8, 1.0)	(-0.2, -0.2)	13	8.49×10 ⁻¹¹	0.6399	0.0399	0.3276

3. An important issue is to appropriately address the interpretation of results in terms of geodynamical structure of North China Craton.

75

80

Response: As we have put forward the fundamental questions at the beginning of the abstract, the destruction mechanism and geodynamics of the NCC are the fundamental questions, we have added the section "5.4 The destruction mechanism of NCC" at the end of the section "5 Discussion" as followings. And we also interpreted results in terms of geodynamical structure in every section 5.1, 5.2 and 5.3, which in concert with the final discussion section 5.4.

Revision: (Page 34)

5.4 The destruction mechanism of NCC

- 85 The low-density anomalies in the Qilian block and northern Taihang Orogenic belt are affected by the high heat flux environments. Low-density anomalies in the central Taihang Orogenic belt exist but are not accompanied by continuous high heat flux environments at mantle depth. However, without mantle plumes, the low-density anomalies in Bohai Bay are affected by the extension of the Tancheng-Lujiang fault belt, and mechanical extension destroys the lithosphere in this area. Based on our studies, one 90 theory is that it is hard to explain the destruction phenomena and modes in the whole NCC. The destruction of NCC is not only affected by physical tension but also caused by thermal erosion and delamination. Previous dynamic studies have shown that (Zhu et al., 2012; Zhu, 2018) since the
- Mesozoic, the Pacific Plate subducted westward to the Taihang Orogenic belt in the central part of the NCC. The residual dehydration of the subducted plate in the mantle transition zone promoted an increase in the molten fluid content in the upper mantle beneath the NCC. The delamination and thermal erosion of the lithosphere in the NCC reflect different forms of mantle convection instability. Therefore, through this study, it is believed that the destruction in the NCC is caused by several forms. Several destruction modes of the NCC coexist in different geological structural backgrounds.

100 4. It is important to provide a point by point response to the reviewers' comments and a new manuscript with marked changes.

Response: We have provided a point by point response to the reviewers' comments, we also attached the new manuscript with marked changes in red after the response.

105

Sequential inversion of GOCE satellite gravity gradient data and terrestrial gravity data for the lithospheric density structure in the North China Craton

Yu Tian^{1,2,3}, Yong Wang^{2,3}

- ¹Ocean College, Minjiang University, 350108 Fuzhou, China
 ²State Key Laboratory of Geodesy and Earth's Dynamics, Institute of Geodesy and Geophysics, Chinese Academy of Sciences, 430077 Wuhan, China
 ³University of Chinese Academy of Sciences, 100049 Beijing, China
 Correspondence to: Yu Tian (tybgys455429145@163.com)
- 120 Abstract. The North China Craton (NCC) is one of the oldest cratons in the world. Currently, the destruction mechanism and geodynamics of the NCC remain controversial. All of the proposed views regarding the issues involve studying the internal density structure of the NCC lithosphere. Gravity field data are among the most important data in regard to investigating the lithospheric density structure, and gravity gradient data and gravity data each possess their own advantages.

Given the different observational plane heights between the on-orbit GOCE satellite gravity gradient and terrestrial

- 125 gravity and the effects of the initial density model on the inversion results, sequential inversion of the gravity gradient and gravity are divided into two integrated processes. By using the preconditioned conjugate gradient (PCG) inversion algorithm, the density data are calculated using the preprocessed corrected gravity anomaly data. Then, the newly obtained high-resolution density data are used as the initial density model, which can serve as constraints for the subsequent gravity gradient inversion. Several essential corrections are applied to the four gravity gradient tensors (T_{xx} , T_{yy} , T_{zz}) of the
- 130 GOCE satellite, after which the corrected gravity gradient anomalies (T'_{xx} , T'_{xz} , T'_{yy} , T'_{zz}) are used as observations. The lithospheric density distribution result within the depth range of 0-180 km in the NCC is obtained.

The inversion results are described as follows. In the crust, the eastern NCC is affected by lithospheric thinning with obvious local features. In the mantle, the presented obvious negative density areas are mainly affected by the high heat flux environment (An and Shi, 2007). In the eastern NCC, the density anomaly in the Bohai Bay area is mostly attributed to the

135 extension of the Tancheng-Lujiang major fault at the eastern boundary. In the western NCC, the crustal density anomaly distribution of the Qilian block is consistent with the northwest-southeast strike of the surface fault belt, whereas such an anomaly distribution experiences a clockwise rotation to a nearly north-south direction upon entering the mantle.

140

1 Introduction

The North China Craton (NCC, Fig. 1) is an outstanding example of a craton that has undergone both reconstruction and destruction. Studies on the NCC have enabled us to understand the formation, evolution, stabilization and destruction of the ancient continent. Two dominant NCC destruction mechanisms have been proposed, namely, delamination (Gao et al., 2009)

- 145 and thermal erosion (Zhu et al., 2012). However, both of these mechanisms involve internal tectonic deformation and substance distribution in the lithosphere of the NCC, which highlights the importance of obtaining the high-resolution density structure of the lithosphere. Gravity field data play an important role in determining and interpreting the lithospheric density structure and state of motion. For studies on the density structure of the NCC, Fang (1996) managed to invert the density distribution of the lithosphere in the North China area, with the constrained least square method using the Bouguer
- 150 anomaly. Additionally, using the Bouguer gravity anomaly, Wang et al. (2014) obtained the three-dimensional density structure of the NCC lithosphere through the algebraic reconstruction inversion method, which was constrained by seismic travel time. Based on gravity, geoidal surface and topography data, Xu et al. (2016) calculated the crustal density and depth of the interface between the lithosphere and asthenosphere using the rapid integrated inversion method. Using the preprocessed data of the GOCE satellite gravity gradient anomaly, Tian and Wang (2018) constructed a three-dimensional
- 155 density structure within the depth range of 0-120 km in the NCC lithosphere, during which the density variation induced by the temperature differences was incorporated. Previous studies on the NCC only adopted gravity or gravity gradient measurements, both of which have their own unique advantages as first-order and second-order derivatives of the gravity potential, respectively. In the frequency domain, the high frequencies of the deep structures are strongly attenuated due to the distance, which is masked by the signals of closer masses. The gravity data are mainly used to provide mid-low
- 160 frequency information about the deep structure. The amplitude of the gravity gradient data declines rapidly with increasing depth in the field source, which demonstrates that the gravity gradient anomaly is applicable to the high frequency signal information of the shallow structure, characterized by short wavelengths. By using both the gravity and gravity gradient data as the observation quantities, both the low frequency signal information and high frequency signal information of the gravity anomaly data can expand the frequency of the gravity field data. Sequential inversions of the gravity and gravity gradient
- 165 data are able to achieve mutual supplementation, which is favorable to enhance the reliability of the inversion result and obtain more reasonable analyses of the inversion solutions.

Currently, most studies are based on the joint inversion of gravity and gravity gradient data instead of sequential inversions. Zhdanov et al. (2004) introduced the concept of the curvature of gravity to carry out a joint inversion of gravity and gravity

170 gradient data and applied the proposed method to the existing models. Wu et al. (2013) inverted gravity and gravity gradients by transforming the formulas of the gravity and gravity gradients with the target body treated as a mass point. Given the varied decline rates of the kernel functions of the gravity and gravity gradient data, Capriotti and Li (2014) balanced the two decline rates using the density matrix to conduct a joint inversion of the gravity and gravity gradient data, after which the validity of the proposed method was confirmed with the published SEG model. Qin et al. (2016) developed an integrated

- 175 focusing inversion algorithm for gravity and gravity gradient data and applied this method to gravity and gravity gradient data from aerial surveys in the Vinton salt dome area. Li et al. (2017) inverted the lithospheric three-dimensional density structure of the Qinghai-Tibet Plateau and its adjacent area within a depth range of 0-120 km using the gravity gradient data measured by GOCE L2 together with the vertical gravity calculated by EGM2008. For small research areas, studies involving the gravity and gravity gradient joint inversion focused on applications of existing models or aerial gravity and
- 180 gravity gradient data at the same height. With respect to previous studies covering a large area, researchers often adopted gravity or gravity gradient data that were directly computed by the gravity field model. Although the gravity field model is able to rapidly calculate the gravity and gravity gradient data for large research areas worldwide, the resulting data are based on the spherical harmonic coefficients rather than actual measurements. In addition, directly using observations rather than the gravity field models is advantageous because it avoids the global average effect during gravity field modeling (Pavlis et
- 185 al., 2012). Thus, compared with the calculated data based on the gravity field model, the high precision gravity and gravity gradient data obtained from measurements with high resolution possess the same importance (Liu et al., 2003; Li et al., 2011).
- We selected GOCE satellite gravity gradient data along the orbit and gravity data of the terrestrial survey as the 190 measurements. The GOCE satellite data at the mean orbital height only reflect large structures of the earth. To highlight the high frequency information of shallow abnormal bodies and the detailed information of structural features, it is necessary to move the GOCE satellite data from the mean orbital height downward to the near surface in the NCC area. However, given the feasibility of the downward continuation result of the gravity gradient data from the GOCE satellite, the height of the observation plane after the downward continuation should be located outside the topographic mass unit (Sebera et al., 2014;
- 195 Li et al., 2017; Tian and Wang, 2018), while each observation location for gravity data acquired through the survey is always located on the topographic surface. Therefore, an inconsistency problem in the observation plane height is anticipated between the processed GOCE satellite gravity gradient data and the gravity data obtained by the terrestrial survey. Furthermore, although gravity and gravity gradient data are favored by higher resolutions and sensitivity to density, the inversion method is characterized by a strong nonuniqueness of solutions. To constrain the gravity and gravity gradient
- 200 inversion, seismic data are often input into the inversion after a transformation based on empirical formulas to suppress the nonuniqueness of solutions. In this regard, the effects of the initial model on the inversion results should be considered. For the two aforementioned aspects, the sequential inversions of gravity and gravity gradient are divided into two processes. First, the density data converted from the seismic wave velocity are used as the initial density model. The preprocessed corrected gravity anomaly data of the NCC terrestrial survey are collected as the observation quantity, and then, the density
- anomaly within a depth range of 0-180 km is calculated using the preconditioned conjugate gradient (PCG) algorithm in the first process. The obtained inversion results are used as the new initial density model and serve as constraints. The four high accuracy (Rummel et al., 2011; Yi et al., 2013) GOCE satellite gravity gradient anomaly tensors (T_{xx} , T_{xz} , T_{yy} , T_{zz}) in the

NCC are collected as the original observation data for downward continuation, topographic effect correction, underground interface undulation effect correction and long wavelength correction. The preprocessed corrected gravity gradient anomaly

210 data (T'_{xx} , T'_{xz} , T'_{yy} , T'_{zz}) are used as the new observation quantity, and the resulting density anomaly distribution within the depth range of 0-180 km in this area is obtained using the same PCG algorithm in the second process.

By considering the features of the gravity anomaly and gravity gradient anomaly data, we used the corrected gravity data as the initial measurements rather than gravity gradient data, which can be used first to determine the major and deep structures of the lithosphere based on the inversion results. Then, the inversion results of the corrected gravity gradient data are applied to identify the fine structures of the lithosphere. This presented method can be exempt from the limitation imposed by the observational plane height, which is able to sufficiently exploit and utilize the available actual gravity and gravity gradient measurements. The corrected initial model with the effect of the gravity inversion results can provide more reliable and effective initial density models for the following gravity gradient inversion. Compared with the inversion results based on either the gravity or gravity gradients in the NCC, the integrated inversion results offer both regional detailed information and a density structure model penetrating deeper underground, which is favorable for discussions and analyses of data regarding the crust and mantle in the same area; consequently, the phenomenon and origin of the NCC destruction



225 Figure 1 Topography and main tectonics of the study area. NCC - North China Craton.

2 Methods

mechanism can be investigated.

2.1 Kernel function calculation

230 Under the Cartesian coordinate system, according to the geological body with the residual density of ρ and the volume of V, the first and second derivatives of the gravitational potential at any point P₁ (x_0 , y_0 , z_0) in the outer space can be expressed as follows (Nagy et al. 2000):

$$T_{z}(\mathbf{P}_{1}) = \mathbf{G}\rho [[] \frac{(z - z_{0})}{[(x - x_{0})^{2} + (y - y_{0})^{2} + (z - z_{0})^{2}]^{3/2}} dx dy dz$$

$$T_{xx}(\mathbf{P}_{1}) = \mathbf{G}\rho [[] \frac{2(x - x_{0})^{2} - (y - y_{0})^{2} - (z - z_{0})^{2}}{[(x - x_{0})^{2} + (y - y_{0})^{2} + (z - z_{0})^{2}]^{5/2}} dx dy dz$$

$$T_{xz}(\mathbf{P}_{1}) = \mathbf{G}\rho [[] \frac{3(x - x_{0})(z - z_{0})}{[(x - x_{0})^{2} + (y - y_{0})^{2} + (z - z_{0})^{2}]^{5/2}} dx dy dz$$

$$T_{yy}(\mathbf{P}_{1}) = \mathbf{G}\rho [[] \frac{2(y - y_{0})^{2} - (x - x_{0})^{2} - (z - z_{0})^{2}}{[(x - x_{0})^{2} + (y - y_{0})^{2} + (z - z_{0})^{2}]^{5/2}} dx dy dz ,$$

$$T_{zz}(\mathbf{P}_{1}) = \mathbf{G}\rho [[] \frac{2(z - z_{0})^{2} - (x - x_{0})^{2} - (y - y_{0})^{2}}{[(x - x_{0})^{2} + (y - y_{0})^{2} + (z - z_{0})^{2}]^{5/2}} dx dy dz ,$$

$$(1)$$

235 where G is the Newtonian gravitational constant. By dividing the underground space of the detection area into k prisms with the same size, each cube is given a particular residual density, and the analytical expression of Eq. (1) can be expressed as follows:

$$T_{z}(P_{1}) = \sum_{i=1}^{k} G\rho \left\{ (x - x_{0}) ln[r + (y - y_{0})] + (y - y_{0}) ln[r + (x - x_{0}) - (z - z_{0}) arctan \frac{(x - x_{0})(y - y_{0})}{(z - z_{0})r}] \right\} \Big|_{x_{i}}^{x_{i+1}} \Big|_{y_{i}}^{y_{i+1}} \Big|_{z_{i}}^{z_{i+1}} = G_{z}^{(P_{1})}\rho$$

$$T_{xx}(P_{1}) = \sum_{i=1}^{k} G\rho_{i} [-tan^{-1} \frac{(y - y_{0})(z - z_{0})}{(x - x_{0})r}] \Big|_{x_{i}}^{x_{i+1}} \Big|_{y_{i}}^{y_{i+1}} \Big|_{z_{i}}^{z_{i+1}} = G_{xz}^{(P_{1})}\rho$$

$$T_{xz}(P_{1}) = \sum_{i=1}^{k} G\rho_{i} ln(y - y_{0} + r) \Big|_{x_{i}}^{x_{i+1}} \Big|_{y_{i}}^{y_{i+1}} \Big|_{z_{i}}^{z_{i+1}} = G_{xz}^{(P_{1})}\rho$$

$$T_{yy}(P_{1}) = \sum_{i=1}^{k} G\rho_{i} [-tan^{-1} \frac{(z - z_{0})(x - x_{0})}{(y - y_{0})r}] \Big|_{x_{i}}^{x_{i+1}} \Big|_{y_{i}}^{y_{i+1}} \Big|_{z_{i}}^{z_{i+1}} = G_{yy}^{(P_{1})}\rho$$

$$T_{zz}(P_{1}) = \sum_{i=1}^{k} G\rho_{i} [-tan^{-1} \frac{(x - x_{0})(y - y_{0})}{(z - z_{0})r}] \Big|_{x_{i}}^{x_{i+1}} \Big|_{y_{i}}^{y_{i+1}} \Big|_{z_{i}}^{z_{i+1}} = G_{yy}^{(P_{1})}\rho$$

$$(2)$$

240 where $G_z^{(P_1)}, G_{xx}^{(P_1)}, G_{xz}^{(P_1)}, G_{zz}^{(P_1)}$ represents the corresponding kernel function matrix at point P₁ and $\rho = (\rho_1, \rho_2, \dots, \rho_k)^T$ represents the density of each cube. The gravity gradient anomalies generated by all n points on the observation surface meet the linear relationship from the k discrete prisms in the underground space. The relationship can be expressed as the following equation:

$$T_{ij} = G_{ij}\rho ,$$

$$T_{ij} = [T_{ij}^{(P_1)}, T_{ij}^{(P_2)}, \dots, T_{ij}^{(P_n)}]^T ,$$

$$G_{ij} = [G_{ij}^{(P_1)}, G_{ij}^{(P_2)}, \dots, G_{ij}^{(P_n)}]^T , \quad ij = z, \, xx, \, xz, \, yy, \, zz$$
(3)

245 where T_{ij} represents the observation and G_{ij} represents the corresponding kernel function matrix for the observation surface.

2.2 Inversion method

Essentially, the inversion of the gravity and gravity gradient tensors is a process of solving a system of linear equations. The quantity of the unknown m greatly exceeds the acquired data vector, and the solutions of the equations are nonunique. Moreover, inversion is an ill-conditioned problem, and appropriate constraints on the objective function are required to narrow the range of solutions. Therefore, in the linear inversion theory, the objective function mostly consists of the data fitting function and the model objective function (Constable et al., 1987). Under such circumstances, solving the inversion problem is equivalent to finding a model vector *m* that can minimize the objective function while satisfying the data fitting function.

The objective function can be expressed as follows:

$$minimize: \phi = \phi_d + \mu \phi_m, \tag{4}$$

260 where μ represents the regularization parameter, which represents the weight factor that balances the data fitting function ϕ_d and the model objective function ϕ_m .

The data fitting function is defined as follows:

270

$$\phi_d = \sum_{i=1}^{N} \left(\frac{\Delta d_i - G_i \Delta m}{\sigma_i} \right)^2 = \left\| W_d (\Delta d - G \Delta m) \right\|^2, \tag{5}$$

265
$$W_d = diag\{1/\sigma_1, 1/\sigma_2, ..., 1/\sigma_N\},$$
 (6)

where $\Delta m = m - m_0$ is the correction between the model parameter vector m and the initial model m_0 ; G is the kernel function, namely, the linear projection operator from the model element to the observation (for gravity data, $G = G_z$, while for gravity gradient components, $G = [G_{xx}, G_{xz}, G_{yy}, G_{zz}]^T$); Δd is the correction of the corresponding measurement; and W_d is a diagonal matrix, with σ_i representing the standard deviation of the *i*-th data. The objective function of the model is constructed according to the minimization model function. The Lagrangian multiplier is used as the regularization parameter in the PCG inversion algorithm. In the process of solving a large-scale matrix, the calculation of the Lagrangian multiplier is complex; therefore, an empirical value is often adopted as the relative optimal value for the regularization parameter. Since the regularization parameter serves to balance the data fitting function and the model fitting function, an excessively large value will result in substantial differences between the inverse results response and the observation, while an overwhelmingly small value leads to ineffectiveness of the model

Given these problems, the L-curve method was developed for the selection of regularization parameters in the solution for

280

275

fitting function.

ill-posed problems (Hansen, 1992). The L-curve is a criterion that is based on a comparison between the actual data fitting function and the model objective function, which is applicable to solving large-scale problems. The value corresponding to the inflection point of the L-curve is assigned to the regularization parameter. The effectiveness of this method has been validated in previous studies (Tian et al., 2018; 2019). The curvature of the L-curve can be expressed as follows (Hansen, 1992):

285
$$k = \frac{\hat{\rho}'\hat{\eta}'' - \hat{\rho}''\hat{\eta}'}{\left[\left(\hat{\rho}'\right)^2 + \left(\hat{\eta}'\right)^2\right]^{3/2}},$$
(7)

where $\hat{\rho} = \log(\phi_d)$ and $\hat{\eta} = \log(\phi_m)$; and the superscripts ' and " represent the first-order and second-order derivatives of the function, respectively. Accordingly, during the inversion process, the algorithm seeks the maximum curvature of the L-curve based on the function constructed by actual data.

290 To constrain the spatial structure of the model and achieve a continuous variation in the inversion image along the three axis directions, a roughness matrix is introduced into the model objective function (Constable et al., 1987), with reference to the minimization model function. The three-dimensional model vector \mathbf{R} is the quadratic sum of the first-order partial difference of the model vector *m* along the x, y and z directions.

$$\boldsymbol{R} = \left\|\partial_{x}\Delta\boldsymbol{m}\right\|^{2} + \left\|\partial_{y}\Delta\boldsymbol{m}\right\|^{2} + \left\|\partial_{z}\Delta\boldsymbol{m}\right\|^{2} = \int \left(\frac{\partial\Delta\boldsymbol{m}}{\partial x}\right)^{2} dv + \int \left(\frac{\partial\Delta\boldsymbol{m}}{\partial y}\right)^{2} dv + \int \left(\frac{\partial\Delta\boldsymbol{m}}{\partial z}\right)^{2} dv, \qquad (8)$$

295

by meshing the model and replacing the partial differential form with the finite difference form, the above equation is converted into the following matrix:

$$\boldsymbol{R} = \Delta \boldsymbol{m} (\boldsymbol{R}_{x}^{\mathrm{T}} \boldsymbol{R}_{x} + \boldsymbol{R}_{y}^{\mathrm{T}} \boldsymbol{R}_{y} + \boldsymbol{R}_{z}^{\mathrm{T}} \boldsymbol{R}_{z}) \Delta \boldsymbol{m},$$

$$\tag{9}$$

where R_x , R_y and R_z are the roughness matrices of the model along the x, y and z directions, respectively.

300 Because the gravity data and gravity gradient data have no fixed depth resolution, the kernel function declines rapidly with increasing depth, and the inversion results are limited near the surface, which results in difficulties in capturing the true position of the anomaly. By introducing the depth weighting function into the model objective function, the kernel function is optimized to reflect the true weighted value of the anomaly element at each depth. The depth weighting function designed by Li and Oldenburg (1996), especially for the inversion of gravity data and gravity gradient data, is adopted as follows:

305
$$W(z) = \frac{1}{(Z + Z_0)^{\beta/2}},$$
 (10)

where Z is the burial depth of the center of the grid cell and Z_0 and β are constants. For gravity data, these values are used to counteract the decline in the kernel function G, with β often set to 2 and the function written as $(Z+Z_0)^{-1}$, while for the gravity gradient data, these values are used to compensate for the decline in the kernel function G, with β often set to 3 and the function written as $(Z+Z_0)^{-3/2}$.

310

320

In accordance with the minimization model function, the model objective function can be constructed as shown below, in reference to the roughness matrix and depth weighting function:

$$\phi_m(\boldsymbol{m}) = \alpha_s \int_V \left(\partial \boldsymbol{W}(z) \Delta \boldsymbol{m} \right)^2 dv + \alpha_s \int_V \left(\frac{\partial \boldsymbol{W}(z) \Delta \boldsymbol{m}}{\partial x} \right)^2 dv + \alpha_y \int_V \left(\frac{\partial \boldsymbol{W}(z) \Delta \boldsymbol{m}}{\partial y} \right)^2 dv + \alpha_z \int_V \left(\frac{\partial \boldsymbol{W}(z) \Delta \boldsymbol{m}}{\partial z} \right)^2 dv, \quad (11)$$

The model objective function can be converted into a matrix form by replacing the differential form with the finite difference method:

$$\phi_m(\boldsymbol{m}) = \Delta \boldsymbol{m}^{\mathrm{T}} (\boldsymbol{W}_S^{\mathrm{T}} \boldsymbol{W}_S + \boldsymbol{W}_x^{\mathrm{T}} \boldsymbol{W}_x + \boldsymbol{W}_y^{\mathrm{T}} \boldsymbol{W}_y + \boldsymbol{W}_z^{\mathrm{T}} \boldsymbol{W}_z) \Delta \boldsymbol{m} = \Delta \boldsymbol{m}^{\mathrm{T}} \boldsymbol{W}_i^{\mathrm{T}} \boldsymbol{W}_i \Delta \boldsymbol{m} , \qquad (12)$$

$$\boldsymbol{W}_{i} = \boldsymbol{\alpha}_{i} \boldsymbol{R}_{i} \boldsymbol{D}, \quad i = s, x, y, z,$$
(13)

where α_i is the weight coefficient for each term in the objective function; R_i is the difference operator for each component; and D is the discretized depth weighting function matrix. Substituting the model objective function into the objective function yields the following expression:

$$\phi = (\Delta d - G\Delta m)^{\mathrm{T}} (\Delta d - G\Delta m) + \mu \Delta m^{\mathrm{T}} W_{i}^{\mathrm{T}} W_{i} \Delta m \qquad (14)$$

Eq. (11) can be rearranged into the following matrix form:

$$\begin{bmatrix} \boldsymbol{G} \\ \sqrt{\mathbf{k}}\boldsymbol{W}_{\mathbf{i}} \end{bmatrix} \Delta \boldsymbol{m} = \begin{bmatrix} \Delta \boldsymbol{d} \\ \boldsymbol{0} \end{bmatrix}$$
(15)

by replacing the condition matrix of the objective function with $A = [G, \sqrt{k}W_i]^T$ and defining $b = [\Delta d, 0]^T$, Eq. (15) can be simplified as follows:

$A \Delta m = b$.

For the gravity data, we take T_z as the observation, where $A = \begin{bmatrix} G_z & \sqrt{\mu}W_i \end{bmatrix}^T$ and $b = \begin{bmatrix} T_z & 0 \end{bmatrix}^T$. For the gravity gradient data, we simultaneously selected four processed components T'_{xx} , T'_{xz} , T'_{yy} and T'_{zz} as the observations, which implies the Jacobian matrix, where $A = \begin{bmatrix} G_{xx} & G_{xz} & G_{yy} & G_{zz} & \sqrt{\mu}W_i \end{bmatrix}^T$ and $b = \begin{bmatrix} T'_{xx} & T'_{xz} & T'_{yy} & T'_{zz} & 0 \end{bmatrix}^T$. The contribution of each component and its differences from the joint inversion can be found in our previous studies (Tian et al., 2019).

Additionally, $A^{T}A = (G^{T}G + k^{-1}W_{i}^{T}W_{i})$. Because the condition number of the Jacobian matrix is normally very large, to increase the convergence speed and the solution stability, Eq. (16) is rewritten as follows:

$$335 \quad SA^{4}A\Delta m = SA^{4}b, \tag{17}$$

where S is the preconditioned factor, which is usually approximated as $(A^{T}A)^{-1}$. Thus, the eigenvalues of Eq. (17) will be concentrated along the diagonal, and the condition number will be improved so that the iteration efficiency is improved (Pilkington, 2009).

The PCG method is an algorithm for solving linear or nonlinear equations through an iterative calculation process and is mainly used to solve the quadratic equation $\phi = \frac{1}{2} \mathbf{m}^{T} A \mathbf{m} - \mathbf{m}^{T} \mathbf{b}$, with a solution of $\mathbf{m} = A^{-1} \mathbf{b}$, which is obviously equivalent to Eq. (16). The PCG method (Pilkington, 1997; 2009) is adopted for the inversion calculation.

2.3 Effects of the initial density model

345

350

330

Two models (Fig. 2 and Fig. 3) are designed to test the effects of the initial model on the inversion results. Model one (Fig. 2) consists of two independent cubes equipped with density values. The densities are different at each layer to study the role of the initial values in the inversion. Model two (Fig. 3) is composed of an upright cuboid and a trapezoid structure of multiple anomalies. This model test can be used to illustrate the initial model effects on the shape and magnitude of the inversion results. Since this study aims to capture the initial model effects upon the final inversion results, carrying out the integrated

inversion for the four components $(T_{xx}, T_{xz}, T_{yy}, T_{zz})$ is required.

As shown in Fig. 2 and Fig. 3, the inversion has been carried out for identical density anomaly bodies (Fig. 2a and Fig. 3a), during which the gravity gradient measurement, kernel function and algorithm parameters are all the same, except that the

- varied values are assigned to the initial model. In Model 1, the anomalies are set to 0 g/cm³ (Fig. 2b); 0.4 g/cm³ (Fig. 2c); 0.1 g/cm³ (1st layer), 0.2 g/cm³ (2nd and 3rd layers) and 0.3 g/cm³ (4th layer) in Fig. 2d; and 0.2 g/cm³ (1st layer), 0.4 g/cm³ (2nd and 3rd layers) and 0.6 g/cm³ (4th layer) in Fig. 2e. Then, in Model 2, the initial densities are set to 0 g/cm³ (Fig. 3b), 0.2 g/cm³ (Fig. 3c), 0.5 g/cm³ (Fig. 3d) and -0.2 g/cm³ (Fig. 3e).
- 360 In terms of the inversion of Model 1, the iterations are 10, 7, 6 and 3, and the inversion results are illustrated in Fig. 2b-e. The detailed results of every inversion test are summarized in Table 1.

	True density at each layer (g/cm ³) (1 st to 4 th layer)	Initial density at each layer (g/cm ³) (1 st to 4 th layer)	Iterations (k)	RMS (E)	Maximum density at each layer (g/cm ³) (1 st to 4 th layer)
Test 1	(0.4, 0.7, 0.7, 1.0)	(0, 0, 0, 0)	10	7.35×10 ⁻¹¹	(0.34, 0.42, 0.53, 0.55)
Test 2	(0.4, 0.7, 0.7, 1.0)	(0.4, 0.4, 0.4, 0.4)	7	6.94×10 ⁻¹²	(0.51, 0.55, 0.66, 0.58)
Test 3	(0.4, 0.7, 0.7, 1.0)	(0.1, 0.2, 0.2, 0.3)	6	3.50×10 ⁻¹²	(0.36, 0.64, 0.67, 0.64)
Test 4	(0.4, 0.7, 0.7, 1.0)	(0.2, 0.4, 0.4, 0.7)	3	2.31×10 ⁻¹³	(0.38, 0.72, 0.76, 0.95)
	Minimum density	Average density			
	at each layer	at each layer			
	(g/cm^3)	(g/cm^3)			
	$(1^{st} to 4^{th} layer)$	$(1^{st} to 4^{th} layer)$			
Test 1	(0.23, 0.27, 0.34, 0.29)	(0.31, 0.34, 0.41, 0.35)			
Test 2	(0.42, 0.43, 0.51, 0.48)	(0.47, 0.50, 0.58, 0.55)			
Test 3	(0.32, 0.46, 0.33, 0.55)	(0.34, 0.55, 0.59, 0.58)			
Test 4	(0.33, 0.52, 0.57, 0.83)	(0.35, 0.59, 0.65, 0.91)			

Table 1 Conditions and results for initial density model one.

- Fig. 2b-e indicates that as the initial model becomes more precise, the inversion results become more reliable, with the values of the anomalies becoming more similar to the true model. With no defined initial model (Fig. 2b), it is difficult to capture the layered structure of the anomaly, and the resulting average density is only 0.38 g/cm³, which is far different from the true model; however, in this case, the inversion results capture the position of the anomaly, its depth distribution, and its shape as two upright cuboids. The whole inversion process is relatively time consuming and finally reaches the convergence criterion after 10 iterations. We also compare the cases based on Model 1 with one another. Compared with Fig. 2d, although Fig. 2c presents smaller differences in the density anomaly values than the true model, the layered structure of the anomaly is difficult to capture from Fig. 2c. In Fig. 2d, it is feasible to conclude that certain differences exist between the density anomaly distributions of the 1st and 2nd layers, which is more similar to the true model. This finding suggests that the inversion results are not only related to the density value of the initial model but also to the distribution structure of the initial
- 375 density. Therefore, a more precise initial model with a layered density value distribution favors the generation of more reliable inversion results. Notably, the optimal inversion results are attributed to the case of Fig. 2e, which produces an inverted anomaly body shape that is almost identical to the true model. Apparently, the density distribution of the anomalous

body is laminated into 3 layers, with the 1st layer having a density anomaly of 0.3-0.4 g/cm³; the 2nd and 3rd layers generally having density anomalies of 0.6-0.7 g/cm³; and the 4th layer having a density anomaly of 0.8-1.0 g/cm³.



Figure 2 Initial density model experiments. (a) True model. Inversion results for (b) Test 1, (c) Test 2, (d) Test 3, and (e) Test 4.

For Model 2, the inversions for each case are calculated after 8, 7, 5 and 13 iterations. The inversion results are illustrated in

Fig. 3b-e, and the details of the results of each case are given in Table 2. With the aid of a more precise initial model and 385 more data, Fig. 3b-d demonstrates that it would be easier to determine the shape and dimensions of the density anomaly body 390

from the inversion results. In the inversion results of Fig. 3b, it is feasible to directly capture the upright cuboid of the density anomaly on the right, and for the complicated density anomaly body on the left, the interpretation based on the inversion results is limited to this anomaly body having an irregular shape and its size growing with increasing depth. In Fig. 3c, the anomaly body on the left reveals a stair-like shape, yet it is surrounded by multiple small anomaly bodies, and the density values of the upright cuboid on the right are closer to those of the true model. Next, the optimal inversion results are found in Fig. 3d, with anomalous body shapes that are completely consistent with those of the true model, and it is easy to discriminate the shape of the anomalous body. Meanwhile, it should also be noted that the density anomaly value of the upright cuboid on the right is higher than that of the stair-like body on the left. Finally, the inversion results in Fig. 3e make

it difficult to determine the shapes and sizes of both anomalous bodies. In addition, according to the inversion results 395 summary (Table 2), the average density anomaly is only 0.3276 g/cm³ in this case, which is very different from that of the true model.

2 Conditions and	f results for initial	density mod	del two.			
True density (Trapezoid, Cuboid) (g/cm ³)	Initial density (Trapezoid, Cuboid) (g/cm ³)	Iterations (k)	RMS (E)	Maximum density (g/cm ³)	Minimum density (g/cm ³)	Average density (g/cm ³)
(0.8, 1.0)	(0, 0)	8	4.37×10 ⁻¹²	0.6887	0.1328	0.4225
(0.8, 1.0)	(0.2, 0.2)	7	5.63×10 ⁻¹³	0.7121	0.1856	0.4886
(0.8, 1.0)	(0.5, 0.5)	5	1.06×10^{-13}	0.7556	0.4762	0.6256
(0.8, 1.0)	(-0.2, -0.2)	13	8.49×10 ⁻¹¹	0.6399	0.0399	0.3276
		(b)			(c)	
	0.9 0.8 0.7 0.6 0.5 0.5 0.9 0.4 0.6 0.5 0.5 0.4 0.4 0.2 0.1 x/m	0 -100 -200 -300 -400 -500 1000 800 60	0 400 200 y/m	0.9 0.8 0.7 % 5 0.6 3 0.5 1990 0.4 0.3 0.2 0.1 x/m	0 -100 -200 -400 -500 -500 -400 -500 -500 -400 -200 -400 -200 -400 -200 -400 -200 -400 -200 -400 -200 -400 -200 -400 -4	1 0.9 0.8 0.7 0.6 0.5 0.6 0.5 0.6 0.5 0.4 0.3 0.2 0.1 x/m
	1 0.9 0.8 0.7 0.6 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0 0.0	0 -100 -200 -300 -500 1000 800 60	0 400 200 0 200 0 200 200 200	1 0.9 0.8 0.7 (100) 0.6 Å) 0.5 Å) 0.4 0 0 0.0 0.2 0.1		
	True density (Trapezoid, Cuboid) (g/cm ³) (0.8, 1.0) (0.8, 1.0) (0.8, 1.0) (0.8, 1.0) (0.8, 1.0) $0^{00}_{200}_{0}_{0}_{0}_{0}_{0}_{0}_{0}_{0}_{0}^{0}_{0}$	True density Initial density (Trapezoid, Cuboid) (g/cm ³) (g/cm ³) (0.8, 1.0) (0, 0) (0.2, 0.2) (0.8, 1.0) (0.2, 0.2) (0.8, 1.0) (0.5, 0.5) (0.8, 1.0) (-0.2, -0.2) (0.8, 1.0) (-0.2, -0.2) (0.8, 1.0) (0.5, 0.5) (0.8, 0.5) (0.8, 0	Conditions and results for initial density True density Initial density Cuboid) Cuboid) Iterations Cuboid) Cuboid) Iterations (g/cm³) (g/cm³) (b) (0.8, 1.0) (0.2, 0.2) 7 (0.8, 1.0) (0.2, 0.2) 7 (0.8, 1.0) (0.2, 0.2) 7 (0.8, 1.0) (0.2, -0.2) 13 (b) (b) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c) (c)	$\begin{array}{c} \text{True density} & \text{Initial density} \\ (\text{Trapezoid,} \\ (\text{Cuboid}) \\ (g/cm^3) \\ \hline (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.2, 0.2) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.2, 0.2) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.2, 0.2) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.2, 0.2) \\ (0.8, 1.0) \\ (0.8, 1.0) \\ (0.2, 0.2) \\ (0.8, 1.0) \\ (0.8, $	True density Initial density (Trapezoid, Cuboid) (g/cm ³) (g/cm	True density (Trapezoid, Cuboid) (g/cm ³) (k) (k) (E) Maximum density (g/cm ³) (g/c

400 Figure 3 Initial density model experiment two. (a) True model. Inversion results for (b) Test 1, (c) Test 2, (d) Test 3, and (e) Test 4.

The model tests show that the inversion results are not only related to the observation but also affected by the physical and geometric parameters of the initial density model. Therefore, the density values for the initial model, as well as the general area where the density anomaly body is located, should be more accurately set for the initial density model.

x/m

405 **2.4** The process of the sequential inversion

The sequential inversion is realized in two steps, namely, gravity inversion and gravity gradient inversion. The detailed workflow of the whole inversion process is illustrated in Fig. 4. Given the constraint imposed on the inversion results by the initial density model, the high-precision results of the gravity inversion are used as the new initial density model for the subsequent steps, and the ultimate inversion results are somewhat constrained during the gravity gradient inversion

- 410 calculation. Specifically, the density data converted from the seismic wave velocity are first used as the initial density model, and the preprocessed corrected gravity anomaly data are collected to serve as the observation quantity. Then, the kernel function corresponding to the gravity data is calculated, and the density anomaly data within the depth range of 0-180 km in the study area are calculated through the PCG algorithm. Then, the obtained gravity inversion results are used as the new initial density model, and the gravity gradient components in this area that are measured by the GOCE satellite are processed,
- and then, the four corrected gravity gradient anomaly components (T'_{xx} , T'_{xz} , T'_{yy} , T'_{zz}) are simultaneously used as the new observations. The kernel functions corresponding to each gravity gradient component are calculated, and then, the ultimate density anomaly distribution within a depth range of 0-180 km in the study area is obtained through the same PCG algorithm.



420 Figure 4 The flowchart for the whole PCG inversion algorithm procedure.

3 Data processing

3.1 Inversion based on the corrected gravity anomaly data

The corrected gravity anomaly data (Wang et al., 2012) with depths of 0-180 km in the NCC have been directly collected. The corrected gravity anomaly data are based on the $5' \times 5'$ free air gravity anomaly data set, which comes from the China

425 Seismic Network. The corrected gravity data are first subjected to topographic correction and Bouguer correction, then to the underground interface undulation correction, which consists of sedimentary undulation correction and Moho undulation

correction and finally to the long wavelength correction, which corresponds to the gravity gradient effects of the 2nd-33rd order spherical harmonic coefficients. The gravity anomalies induced by the density heterogeneity of the crust and upper mantle are used as the observations for inversion. The gravity anomaly (Fig. 5a) after multiple corrections is shown in Fig.

- 430 5a. The figure shows that the gravity anomaly of the eastern NCC shows alternating distributions of positive and negative anomalies with local features. In the central NCC, most regions along the Taihang Orogenic belt and the Yinshan-Yanshan Orogenic belt are found to have negative gravity anomalies. For the western NCC, the whole Ordos block presents a prominent positive gravity anomaly distribution.
- 435 Although the gravity data are favored by their higher resolutions, the gravity inversion suffers from a strong nonuniqueness of solutions. It is effective to introduce seismic data into the gravity inversion to constrain the gravity inversion process and the nonuniqueness of the solution. Based on the 0.5°×0.5° resolution of the P-wave velocity structure obtained by seismic tomography (Tian et al., 2009), we constructed a three-dimensional initial density model for the NCC lithosphere using the empirical velocity to density conversion formula of the North China area (Eq. 16). The constructed models are divided into eight layers along the depth direction, with base depths of 10 km, 25 km, 42 km, 60 km, 80 km, 100 km, 140 km and 180 km.
- Along the horizontal direction, the model is meshed into grids of $0.5^{\circ} \times 0.5^{\circ}$. As shown in Fig. 5b, after 17 iterations, the curve of the defined RMS misfit tends to be horizontal, the variation is slight, and the calculated RMS misfit of 8.2×10^{-3} was obtained, at which point the inverse iterative calculation was complete.



445 Figure 5 (a) The corrected gravity anomalies after several corrections. (b) With the iterative calculation, residual mean square between the forward calculated theoretical gravity and the gravity measurements versus the iteration number in the PCG inversion algorithm.

The three-dimensional density anomaly distribution of the lithosphere within 0-180 km, which is meshed into $0.25^{\circ} \times 0.25^{\circ}$ 450 cells (Fig. 6), is inverted from the corrected gravity anomaly. Then, the results are used as the new initial model for gravity gradient inversion. In Sect. 5, the results of the initial and final inversion results (Fig. 6 and Fig. 15) are further analysed and discussed.





Figure 6 Lithospheric density distribution of the NCC based on gravity values at different depths.

455

460

3.2 Obtain GOCE satellite data for study area

The GOCE L2 (GOCE GO_CONS_EGG_TRF_2) gravity gradient measurement product provided for users includes the gravity gradient data along the orbit. This study directly downloads the preprocessed gravity gradient anomaly data with a spatial resolution of 10 arc-min, acquired over 48 months from November 2009 to October 2013 (Sebera et al., 2014) from the website GOCE+ Geoexplore II (http://http://goce.kma.zcu.cz/data.php). These data sets have undergone average orbit height correction (Sebera et al., 2014) and normal gravity gradient correction (Mortiz, 1980; Šprlák, 2012). The geographic area within E100°-E120° and N33°-N43° is defined as the study area (Fig. 7a-d), which is where the NCC is located.





Figure 7 Gravity gradient anomaly (T_{xx} , T_{xz} , T_{yy} , T_{zz}) after average orbital correction to an altitude of 250 km.3.3 Downward 465 continuation

The data of the GOCE satellite at the average orbit height only reflect the long wavelength information and the large structural features inside the earth. To highlight the high frequency information of the shallow anomaly body and detailed information of the structural features, such as depth and shape, the GOCE satellite data at the average orbit height for the 470 NCC have been downward continued to the near surface area (Martinec, 2014). The obtained results benefit the lithospheric structural analysis for local areas. The Tesseroids software developed by Uieda et al. (2016) is used in the topography calculation and the underground interface undulation effects. To ensure the accuracy of the calculation, the software recommends a distance of at least 1 km between the observation plane and the mass body, while the highest point of the topography in the study area is located at approximately 5 km. Thus, the gravity gradient components continued downward

475







Figure 8 Gravity gradient effect (T_{xx} , T_{xz} , T_{yy} , T_{zz}) after downward continuation to an altitude of 10 km.

3.4 Correction for the topographic effects

- 480 The gravity gradient anomaly components after downward continuation are a combination of the interface undulation and density heterogeneity. Thus, topographic correction, underground interface undulation correction and long wavelength correction should be carried out in accordance with the existing precise model. These three corrections are affected by the masses within and outside the study area, and far zone corrections outside the study area should be considered (Szwillus et al., 2016). Therefore, the correction radius in the study area is extended by 10° for all corrections. The extended geographic area within E00° E130° and N23° N53° is defined as the colculation area (Fig. 13 a.f.). The obtained results in the study area
- 485 area within E90°-E130° and N23°-N53° is defined as the calculation area (Fig. 13 e-f). The obtained results in the study area are removed from the calculation area.

First, the topographic effects are calculated using the Tesseroids software (https://tesseroids.readthedocs.io/en/latest) developed by Uieda et al. (2016), which is aimed at reducing the gravity anomaly induced by the topographic mass above the geoidal surface from the gravity gradient after continuation. The Tesseroids software can convert the spherical hexahedron (tesseroid) in spherical coordinates into the prism in Cartesian coordinates to calculate the corresponding gravity gradient effects for topography. The Tesseroids software provides a built-in module to calculate the topographic effects, for which the calculation is based on the topographic elevation data ETOPO1 with a spatial resolution of $1' \times 1'$ (Amante and Eakins

495

2009).

To directly reveal the correlation between the topography and calculated gravity gradient effects, the three-dimensional figure of the topographic gravity gradient effect T_{zz} is shown in Fig. 9e, where the interface undulation represents the topography (Fig. 1), and the color reflects the gravity gradient effects T_{zz} (Fig. 9d).



Figure 9 Gravity gradient effect caused by topographic masses: (a) T_{xx} , (b) T_{xz} , (c) T_{yy} , (d) T_{zz} , and (e) the 3D gravity 500 gradient effect T_{zz} of topography.

3.5 Correction for the underground interface undulation effects

First, the average depth and density of the sediment in this area should be calculated for the interface undulation effects. The average depth is calculated from the depth data of the sediment in NCC (Fig. 10a) provided by CRUST 1.0 with a spatial

- resolution of 1° × 1° (Laske et al., 2013), and then, the interface undulation data of the sedimentary layer (Fig. 10b) are obtained after correcting for the average sedimentary layer depth. With respect to the calculated average depth, all the density data corresponding to the average depth are extracted to calculate the average density corresponding to the average sedimentary layer depth. The average sedimentary layer depth in the NCC is calculated to be 2.3 km, and the corresponding average density is 2.45 g/cm³. Based on CRUST 1.0, the actual depth and density are adopted instead of the empirical value. Moreover, the differences in the actual depth and density at each point from the average depth and density are calculated 510 using the tesseroid forward modeling method based on the Tesseroids software. Consequently, the gravity gradient effects
- induced by the sedimentary layer undulation in the NCC are obtained (Fig. 10c-f).

515

To more intuitively illustrate the correlation between the sedimentary layer interface undulation and the calculated gravity gradient effects, in Fig. 10g, the interface undulation denotes the sedimentary layer interface undulation (Fig. 10b), and the contour lines represent the corresponding gravity gradient effects T_{zz} (Fig. 10f).





Figure 10 (a) Sedimentary layer interface of the NCC. (b) Sedimentary layer interface undulation of the NCC. The anomalous gravity gradient component caused by the relief at the sedimentary layer interface undulation, (c) T_{xx} , (d) T_{xz} , (e) T_{yy} , and (f) T_{zz} . (g) The 3D gravity gradient effect T_{zz} of the sedimentary layer interface undulation.

- 520 The depth of the Moho interface in the NCC provided by CRUST 1.0 is shown in Fig. 11a. The average depth calculated from the depth data of the Moho is 36.5 km, and the corresponding average density is 3.32 g/cm^3 . The Moho undulation after correction for the average depth is illustrated in Fig. 11b. Similarly, the gravity gradient effects induced by the Moho undulation can be calculated. The gravity gradient effects induced by the Moho interface undulation based on the Tesseroids software are presented in Fig. 11c-f. The three-dimensional gravity gradient effect T_{zz} induced by the Moho interface
- 525 undulation is shown in Fig. 11g, where the interface undulation reflects the Moho undulation (Fig. 11b) and the contour line represents the gravity gradient effects (Fig. 11f).















Figure 11 (a) Moho interface of the NCC. (b) Moho interface undulation of the NCC. (c) The anomalous gravity gradient effect caused by the Moho interface undulation. (c) T_{xx} , (d) T_{xz} , (e) T_{yy} , and (f) T_{zz} . (g) The 3D gravity gradient effect T_{zz} of the Moho interface 530 undulation.

3.6 Correction for the gravity gradient effects of the mantle under 180km

The long wavelength gravity gradient effects are mainly induced by substance density heterogeneity in the lithosphere at depths greater than 180 km. The correlation between the depth of the field source and the order of the spherical harmonic function of the gravity potential developed by Bowin et al. (1986) is adopted as follows:

535 Z=R/(n-1)

where Z is the depth of the field source; R is the radius of the Earth; and n is the order of the spherical harmonic function. The long wavelength gravity gradient anomaly effects corresponding to the 2nd-33rd order spherical harmonic coefficients (Wang et al., 2014) in EGM2008 are calculated. The gravity gradient effects induced by the long wavelength are presented in Fig. 12.

540







Figure 12 The gravity gradient effect (T_{xx} , T_{xz} , T_{yy} , T_{zz}) caused by the lithosphere under 180 km.

3.7 The corrected gravity gradient components

Since models with different spatial resolutions are used in several corrections, the spatial resolutions of the obtained results are different. Therefore, before calculating the corrected gravity gradient component, the resolutions of the results for each calculation are homogenized to the same spatial resolution of $0.5^{\circ} \times 0.5^{\circ}$. For the higher spatial resolution of the data (e.g., gravity gradient data after downward continuation and topographic correction), we extracted data from the calculated results. For the gravity gradient correction caused by the underground interface undulation, the common kriging interpolation method was adopted to obtain the data needed for the spatial resolution. The corrected gravity gradient anomaly components (Fig. 13) are obtained after removing the topography effects, the underground interface undulation effects and the long wavelength effects from the gravity gradient tensor data after downward continuation.

The amplitude of the topography effects and the underground interface undulation effects is obvious, but the topographic correction and underground interface undulation correction balance each other to a certain degree (Szwillus et al., 2016). This counteraction is obvious in the extended study area, as shown in Fig. 13e (topographic correction) and Fig. 13f (underground interface undulation correction). The detailed statistical amplitude for each gravity gradient correction are summarized in Table 3. To present the different corrections more intuitively, several corrections T_{zz} at different cross sections of latitude 35° and latitude 37.5° are presented in Fig. 13 (g-h).

560







8

6

O N F Gravity gradient (E)

-2

-4

12

10

8

6

4

a b b o b b Gravity gradient (E)

-6 -8

--10 --12

120°E

-40°N

-35°N

120°E

120°E

40°N

35°N

120°E



Figure 13 The corrected gravity gradient components (a) T_{xx} , (b) T_{xz} , (c) T_{yy} , and (d) T_{zz} . Gravity gradient effects with extended area (e) Topographic effects T_{zz} . (f) Interface undulation T_{zz} (Sedimentary and Moho) effects. Several corrections T_{zz} at different cross sections: (g) Latitude 35° and (h) Latitude 37.5°.

0,0										
	Gravity Gradient Effects (E)	Downward Continuation Maximum (E)	Downward Continuation Minimum (E)	Topographic Effects Maximum (E)	Topographic Effects Minimum (E)	Interface Undulation Effects Maximum (E)	Interface Undulation Effects Minimum (E)			
	T _{xx}	5.3	-5.7	6.3	-9.4	10.1	-4.9			
575	$T_{\rm xz}$	7.2	-5.6	11.9	-5.6	5.2	-8.9			
	$T_{_{ m yy}}$	4.8	-3.6	7.2	-6.8	6.3	-6.7			
	T_{zz}	7.5	-6.5	13.8	-6.3	7.7	-14.1			
580	Gravity Gradient Effects (E)	Long Wavelength Effects Maximum (E)	Long Wavelength Effects Minimum (E)	Corrected Effects Maximum (E)	Corrected Effects Minimum (E)					
	T _{xx}	1.1	-0.8	8.7	-6.8					
	$T_{\rm xz}$	1.2	-0.7	8.5	-4.7					
	$T_{_{ m yy}}$	1.4	-0.7	6.7	-6.6					
	T_{zz}	1.5	-1.5	11.5	-12.3					

Table 3 Statistics for each gravity gradient correction.

Results

In Fig. 14, after 8 iterations, the curve of the defined RMS misfit tends to be horizontal, and the variation is increasingly slighter between the values of the defined RMS misfit. A calculated RMS misfit of 9.0×10⁻³ was obtained, at which point the inverse iterative computation was complete. As shown in the inversion results (Fig. 15), obvious heterogeneity is present in

both the horizontal and vertical distributions of the lithospheric density within the NCC, and this heterogeneity is specifically featured by a segmented spatial distribution of the lithospheric density. In comparison (Fig. 6), the inversion based on the gravity gradient provides more local and detailed information about the density anomaly distribution within the entire NCC. The maximum and minimum values of the inversion results based on the gravity gradient have a larger range, and the

595 detailed data statistics are summarized in Table 4. The center of the anomalies is more concentrated with the regional anomaly features, which is more favorable for the discussion about the stability and destruction in different regions of the whole NCC area. In the eastern NCC, the boundary of density differences on both sides of the Tancheng-Lujiang fault belt zone is more obvious; the extreme value of low-density anomalies is continuously present in the Bohai Bay area at depths of

600

60 km-80 km. In the central NCC, it is easier to determine the center of the density anomaly distribution in the southern, middle and northern parts of the Taihang Orogenic belt at 42 km-80 km, as these areas have different regional block features of the density anomaly. In the western NCC, the gravity inversion results are connected overall; however, the result of gravity gradient inversion shows the southeastern trend of the Oilian block, which is more favorable for a geodynamic analysis in the western NCC.



Figure 14 For the iterative calculation, the residual mean square between the forward calculated theoretical gravity gradient and the 605 gravity gradient measurements versus the iteration number in the PCG inversion algorithm.

Depth Inversion Results	10km (g/cm ³)	25km (g/cm ³)	42km (g/cm ³)	60km (g/cm ³)	80km (g/cm ³)	100km (g/cm ³)	140km (g/cm ³)	180km (g/cm ³)
Maximum density of gravity inversion	0.023	0.021	0.011	0.035	0.053	0.056	0.058	0.079
Maximum density of gradient inversion	0.025	0.026	0.015	0.043	0.058	0.070	0.077	0.091
Minimum density of gravity inversion	-0.033	-0.026	-0.022	-0.020	-0.024	-0.061	-0.066	-0.064
Minimum density of gradient inversion	-0.035	-0.028	-0.025	-0.023	-0.028	-0.063	-0.078	-0.069

Table 4 Comparison of gravity and gravity gradient inversion results.

5 Discussion

- 610 In general, obvious density anomalies at a depth of 10 km are only seen in the eastern and central areas of the NCC. At depths of 25-40 km, significant negative density residuals are present in the Taihang Orogenic belt along the central NCC and in the Qilian and Qaidam blocks along the western NCC. Significant high-density residuals exist in the Ordos block. For depths of 60-80 km, low-density residuals are found in the Bohai Bay area in the eastern NCC, while widely distributed low-density residuals are seen in the Yinshan orogenic belt. At a depth of 100 km, the entire NCC is characterized by
- 615 spatially alternating positive and negative density residuals. In comparison, at depths of 140-180 km, an obvious regional differentiation in the density anomaly distribution is seen, with obvious low-density residuals in the Datong volcano at the junction of the Yinshan-Yanshan and Taihang Orogenic belts in the eastern NCC, as well as the Qilian block at the junction of the Qaidam and Ordos blocks in the western NCC.
- 620 According to the horizontal spatial distribution of the lithospheric density, the NCC is divided into three main areas, namely, the eastern NCC, which consists of the North China Basin and Bohai Bay area; the central NCC, which consists of the central transitional belts; and the western NCC, which consists of the Ordos Basin and its surrounding areas. Based on the three main local density anomaly areas in Fig. 15a-h, the density anomaly distribution in each area at different depths is discussed.







625 Figure 15 Lithospheric density distribution in the NCC at different depths: (a) 10 km, (b) 25 km, (c) 42 km, (d) 60 km, (e) 80 km, (f) 100 km, (g) 140 km, and (h) 180 km. Temperature differences at (i) 140 km and (j) 180 km. E represents the eastern NCC, N represents the central NCC, and W represents the western NCC; HFS is the Haiyuan fault system, RMF is the Riyue mountain fault, and SP is the Songliao plain.

630 5.1 Eastern NCC

deformation (Tian and Zhao, 2011).

635

The eastern NCC is characterized by obvious features with connections to Bohai Bay. In the eastern NCC, the distribution of positive and negative densities is always alternately characterized within depths 0-180 km. From the Tancheng-Lujiang fault belt (E2 and E3) in the south, to Bohai Bay (E5) and North China Basin (E1) in the middle, and the Songliao Basin (E6) in the north, the density always has the features of a regional distribution instead of large-scale distribution in the central NCC. The obvious spatial distributions of the density anomalies are consistent with the theory that this area has experienced strong

At the depth interval of 10-180 km in the eastern NCC, there are positive density anomalies in the E1 area of the North 640 China Basin, which are mainly distributed along the Tangshan-Xingtai earthquake belt. This result is consistent with the positive P-wave velocity along the earthquake belt (Huang and Zhao, 2004; 2009), which implies a positive density distribution affected by the Tangshan-Xingtai earthquake belt.

The Tancheng-Lujiang fault belt is one of the major north-northeast fault zones in the East Asia continent. This fault belt extends more than 2400 km in China and cuts through different structural units (Huang et al., 2011). The geological structure of the Tancheng-Lujiang fault belt is complicated. The Tancheng-Lujiang fault belt in the eastern NCC is extended in the north-northeast direction, and on both sides, (the E2 area on the right and the E3 area on the left), the density anomalies are significantly different. At a depth of 42 km, significant positive density residuals are present in the E3 area as the locally extreme value, while regional positive density residuals are observed in the E2 area (Fig. 15c). At a depth of 80 km,

- 650 significant positive density residuals are present in the E3 area, while no density anomalies are observed in the E2 area (Fig. 15e). At a depth range of 100-180 km, the density anomalies are obvious (Fig. 15h), including those in the E3 area with a direction consistent with the fault belt strike and those in the E2 area with alternating distributions of positive and negative density residuals. Moreover, these positive and negative density residuals are the local extremes, reaching 0.10 g/cm³ and -0.06 g/cm³, respectively. On both sides of the Tancheng-Lujiang fault belt, there are no connected large-scale density
- 655 anomalies, and the density anomalies are distributed along the Tancheng-Lujiang fault belt with different features. This distribution represents the density differences between different tectonic units. Therefore, the Tancheng-Luijang fault belt is obvious as the boundary of the tectonic units. In addition, based on the study of seismic receiver functions (Chen et al., 2006; Li et al., 2011), the Tancheng-Lujiang fault belt zone and its extension are the most obvious areas of lithospheric thinning in the eastern NCC, and the thickness of the lithosphere is only 60-70 km. The significantly different density distributions on

660 both sides of the Tancheng-Lujiang fault belt indicate that the fault belt may have penetrated the lithosphere.

In the E5 area of Bohai Bay, there are significant negative density anomalies at depths of 60-80 km. This area features a columnar shape density distribution at a depth of 80 km. However, as the depth increases from 100 km to 180 km, there are no continuous regional negative density residuals in the E5 area. In contrast, distinct low-density residuals are seen in the E4

- 665 area, which is in the extended region of the Tancheng-Lujiang fault belt. Meanwhile, density anomalies are seen in the extended region of the major fault belt in the area adjacent to the eastern boundary of the Tancheng-Lujiang fault belt. According to previous studies (Teng et al., 1997; Su et al., 2009), these density anomalies are ascribed to the extension of the Tancheng-Lujiang fault belt and the development of the mantle plume in the deep Bohai Bay area. However, the distribution of density anomalies in this region does not conform to the features of the mantle plume with continuous negative density
- 670 residuals in the mantle. Based on this study, these density anomalies are mainly ascribed to the extension of the Tancheng-Lujiang fault belt. Negative density residuals (E4) continuously exist at the extension of the Tancheng-Lujiang fault belt (100-180 km), which implies that the region at the extension of the fault zone has penetrated the crust to the mantle. The cutting effect of the Tancheng-Lujiang fault belt and lithospheric thinning result in asthenospheric upwelling with negative density residuals in Bohai Bay.

675

In the E6 area, which is a transitional area between the Yanshan block and Songliao Basin, significant low-density residuals are present at a depth of 10 km (Fig. 15a). As the depth increases, persistent high-density residuals are seen in the upper mantle, especially in the depth range of 100-140 km, with values reaching up to 0.07-0.08 g/cm³. Moreover, the density anomalies are aligned in the southwest direction at depths of 140-180 km relative to those at depths of 0-100 km. This

680

phenomenon is attributed to the southwest extension of the high-density mantle lithosphere below the Songliao Basin, reaching the region below the Yanshan Orogenic belt.

5.2 Central NCC

The central NCC, formed by the central transitional zone, is characterized by significant low-density residuals with obvious segmented distributions. The Taihang Orogenic belt is generally northeast-southwest oriented, and this belt can be divided

685

into three blocks, namely, southern block N1, middle block N2, and northern block N3. From 42 km to 100 km, N1 is connected to the high-density anomaly area of the western NCC (W4), while N3 is connected to the high-density anomaly area of the eastern NCC (E1) and the low-density anomaly area in the Yinshan-Yanshan blocks (N5). Blocks N3 and N5 are connected, forming low-density anomalies in large-scale areas. These features may indicate that the Taihang Orogenic belt has experienced different geological processes from south to north.

690

N4 in the northern part of the Linfen-Weihe Graben block is mainly distributed in the Datong volcanic area. As the depth increases, the density anomalies become significant high-density residuals at 25 km (Fig. 15b) and then become obvious low-density residuals from 42 km to 140 km. Although the Datong volcano is no longer active (Tian et al., 2009), its surrounding area still exhibits low-density residuals. Similarly, N5 is at the junction of Yinshan-Yanshan and north of the Taihang Orogenic belt, which features obvious low-density residuals at depths of 60-180 km. When the depth increases to 60 km (Fig. 15d), these anomalies become connected with N3 north of the Taihang Orogenic belt and with N4 at the junction of the Datong volcanic area, forming a large low-density anomaly zone. As the depth further increases, the low-density anomaly area covering N5 presents persistently negative density residuals that extend to a depth of 180 km (Fig. 15h). Moreover, the negative density anomaly value is more significant.

700

705

695

From the distribution of the temperature difference (An and Shi, 2007; Yang et al. 2013) in Fig. 15 i-j, N5 is located in an area with a continuous high heat environment. Based on previous studies using the magnetotelluric imaging method (Zhang et al., 2016), melting occurs in the mantle of the Datong volcanic area and north of the Taihang Orogenic belt. According to the seismic receiver function, the Poisson's ratio in the northern Taihang Orogenic belt is as high as 0.3, while in the southern Taihang Orogenic belt, the value is approximately 0.25-0.26 (Ge et al., 2011). The Poisson's ratio of the continent is generally between 0.25-0.27, and although the temperature and material composition seem to have a dominant influence on Poisson's ratio (Zandt and Ammon, 1995), it is difficult to increase the Poisson's ratio to 0.3 only by changing the material composition. Therefore, the obvious negative density residuals in this area are mainly affected by the high heat environment. The upwelling of the thermal materials from the deep asthenosphere formed the magma migration pathway, which

710 apparently transforms the lithosphere and upper mantle.

> In contrast, with the northern region, negative density residuals are observed in the central and southern parts of the Taihang Orogenic belt from 60 km to 180 km, with an extreme value of -0.048 g/cm³ at a depth of 180 km. However, at the depth of the mantle, the central southern part does not show features of a high continuous heat environment (Fig. 15i-j). At a depth of

- 715 140 km, the temperature in the southern block of the Taihang Orogenic (N1) is lower than the average temperature, and the temperature is higher than the average temperature only in the middle block of the Taihang Orogenic (N2). As the depth increases, the temperatures in the southern and middle blocks of the Taihang Orogenic (N1 and N2 areas) are all lower than the average temperature at a depth of 180 km. The Poisson's ratio of this area accords with the typical continental features (Ge et al., 2011). Thermal erosion is always accompanied by a high heat flow environment, which is not consistent with this
- 720 feature in the central and southern regions of the Taihang Orogenic belt. Therefore, it is inferred that the impact of temperature is limited, and the obvious negative density anomalies may be caused by delamination.

N6 in the central NCC exhibits significant positive density residuals at depths of 140-180 km, which are in contrast to the low-density residuals over a large area of the central NCC. The mantle part of N6 is connected to the positive density
anomaly area in the W1 area, Ordos block, and western NCC (Fig. 15e-f). In Fig. 15i-j, the temperature in N6 is normal at depths of 140-180 km, and the temperature boundary exists between the N6 and W1 areas. Based on previous studies (Ai et al., 2019), the Taihang Orogenic belt in the mantle part experienced a blocking effect from the rigid Ordos block during the expansion of the orogenic belt. The stable Ordos block area presents continuous large-scale high-density residuals. It is inferred that the Ordos block's blocking effect creates the positive density residuals in the central NCC (N6) and connects
with the Ordos block.

5.3 Western NCC

At a depth of 42 km, the W1 area in the central Ordos block is characterized by high-density residuals (Fig. 15c), whereas no obvious anomalies are seen in the northern part of the Ordos block. According to the receiver function (Tian and Zhao, 2011), the central part of the Ordos block is 41 km thick, which increases to 45 km in the northern part. Therefore, it can be inferred

- that the southern part of the Ordos block has entered the lithospheric mantle at a depth of 42 km. From 60 km to 100 km, the high-density residuals in the W1 area of the central Ordos block gradually change to low-density residuals. At the depth interval of 140-180 km, the high-density residuals still dominate the W1 area of the central Ordos block. Although Ordos is a relatively stable block with high-density features, a low-density block invasion appears at depths of 80-100 km. At a depth of 80 km, the low-density anomaly of the Ordos block (W1) on the east side is connected with the low-density anomaly in the
- 740 central NCC (N2), which indicates that Ordos is affected to some extent by the destruction of the central NCC. At a depth of 100 km, the low-density anomaly of the Ordos block (W1) is connected to the Qilian block (W3) and Qaidam block (W5), and the distributions of the density anomalies are consistent with the theories that the Ordos block is affected by the northeast compression of the Qinghai-Tibet Plateau and substantial deep expansion (Sheng et al., 2015).
- The whole Qilian block (W2 and W3 areas) features significant low-density residuals in the crustal part at 25-40 km, which falls in the range of the Haiyuan fault belt. The boundary of the negative density residuals is essentially consistent with the

strike of the fault belt shown on the surface. The W3 area at the junction of the Qilian and Ordos blocks is characterized by a sharp transition from low density to significant high density. At depths of 25-80 km, the density anomalies in the Oilian block present a dominant NW-SE strike; however, this strike turns clockwise to the NS direction as the depth further

750

increases to 180 km, especially after entering the mantle. This observation is consistent with previous research results in this region. The mantle convective stress field calculated based on gravity anomalies is demonstrated to be significantly inconsistent with the crustal movement pattern (Xiong and Teng, 2002; Wang et al., 2013), which indicates obvious decoupling between the crustal and mantle materials in the Oilian block. The Oilian block is located at the junction of the northeastern Qinghai-Tibet Plateau and the Alashan and Ordos blocks in the western NCC. The Qilian block has not only 755 been influenced by collisions and subduction of the Indian plate but also by the blocking effect of the Ordos Block, which results in a strong regional tectonic stress background in the deep part of the Qilian block with an associated clockwise rotation of the material movement.

As the depth increases in the range of 140-180 km, persistent obvious negative density residuals are increasingly dominant in 760 the Qilian block (W2 areas) of the western NCC. According to a previous study (Teng et al., 2010), the lithosphere in the western NCC is the thickest, with an average depth of 140-150 km. However, within this depth range, obvious low-density residuals are present in the orogenic belts of the western NCC, which eliminates the possibility that lithospheric thinning is the main cause of the density anomalies at depths of 140-180 km. According to the temperature differences (Figure 15 i-j), the Qilian block in the western NCC has a high temperature at depths of 140 km-180 km. Furthermore, according to research 765 on the terrestrial heat flux (An and Shi, 2007), the Qilian block has an average heat flux value of up to 68.340 mW/m². Therefore, it can be concluded that the high heat flux environments lead to negative density residuals in the Qilian block in the western NCC, accompanied by an upwelling of deep asthenosphere materials and the subsequent transformation of the mantle above the lithosphere.

770 The destruction mechanism of NCC 5.4

The low-density anomalies in the Oilian block and northern Taihang Orogenic belt are affected by the high heat flux environments. Low-density anomalies in the central Taihang Orogenic belt exist but are not accompanied by continuous high heat flux environments at mantle depth. However, without mantle plumes, the low-density anomalies in Bohai Bay are

- 775 affected by the extension of the Tancheng-Lujiang fault belt, and mechanical extension destroys the lithosphere in this area. Based on our studies, one theory is that it is hard to explain the destruction phenomena and modes in the whole NCC. The destruction of NCC is not only affected by physical tension but also caused by thermal erosion and delamination. Previous dynamic studies have shown that (Zhu et al., 2012; Zhu, 2018) since the Mesozoic, the Pacific Plate subducted westward to the Taihang Orogenic belt in the central part of the NCC. The residual dehydration of the subducted plate in the mantle 780 transition zone promoted an increase in the molten fluid content in the upper mantle beneath the NCC. The delamination and
 - 34

thermal erosion of the lithosphere in the NCC reflect different forms of mantle convection instability. Therefore, through this study, it is believed that the destruction in the NCC is caused by several forms. Several destruction modes of the NCC coexist in different geological structural backgrounds.

6 Conclusions

- The sequential inversion of the lithosphere density structure in the NCC is divided into two stages. The effects of the initial density model are considered. The inversion results obtained by the inversion of the corrected gravity anomaly are used as the initial model for the inversion of the GOCE satellite gravity gradient components. The GOCE satellite gravity gradient data were processed with several corrections to obtain the corrected gravity gradient components. The density distribution with a depth range of 0-180 km in the NCC is outlined as follows: (1) In the eastern NCC, affected by lithosphere thinning,
- 790 the eastern NCC has local features in the density anomaly distribution. Obvious differences in the density anomaly distribution are observed, and the Tancheng-Lujiang fault belt in the eastern NCC penetrates through the lithosphere. The density anomaly in Bohai Bay is mainly induced by the extension of the Tancheng-Lujiang major fault at the eastern boundary. (2) In the central NCC, the Taihang Orogenic belt located in the central NCC is characterized by a segmented density anomaly distribution. (3) In the western NCC, the Qilian block in the western NCC presents a clockwise rotation of
- 795 the density anomaly distribution with an increasing lithospheric depth, while the adjacent Ordos block remains continuously stable. (4) Across the Taihang Orogenic belt in the central NCC and the Qilian-Qaidam blocks in the western NCC, stronger impacts of the orogenic belt and a high heat flux environment are observed, which results in an upwelling of the deeply buried asthenospheric substances and consequently results in a reconstruction of the lithospheric density structure distribution.

800

Code and data availability. The on-orbit GOCE gravity gradient data can be viewed and downloaded from GOCE+ Geoexplore II (http://goce.kma.zcu.cz/data.php). The "Tesseroids" software for gravity gradient topographic correction and interface undulation can be viewed and downloaded from https://tesseroids.readthedocs.io/en/latest. The global crustal model CRUST1.0 can be downloaded from https://igppweb.ucsd.edu/~gabi/crust1.html. The processed gravity gradient data and inversion results can be viewed and downloaded from https://zenodo.org/record/3545809#.XdJ5H695vIU.

805

Author contributions. Conceptualization, YW; methodology, YT; validation, YT, writing original draft preparation, YT; writing review and editing, YT; supervision, YW; funding acquisition, YW.

Competing interests. The authors declare no conflict of interest.

- 810 Acknowledgements. We are grateful to editor Prof. Mioara Mandea and two reviewers for providing a lot of constructive comments and suggestions, which really help us a lot to make remarkable progress. We are grateful to Prof. Xinsheng Wang in Development Research Center of China Earthquake Administration for providing corrected gravity data. We are also grateful to Prof. Qi Lin and Prof. Bojie Yan in Minjiang University for providing a lot of help when we prepared this paper.
- 815 Financial support. This research was funded by the R&D of Key Instruments and Technologies for Deep Resources Prospecting (the National R&D Projects for Key Scientific Instruments), grant number No.ZDYZ2012-1-04.

References

820

835

- Amante, C., and Eakins, B.W.: ETOPO1 1 arc-minute global relief model: Procedures, data sources and analysis, NOAA Technical Memorandum NESDIS NGDC-24, Boulder, Colorado, USA, March 2009, 19, 2009.
- An, M.J., and Shi, Y.L.: Three-dimensional thermal structure of the Chinese continental crust and upper mantle, Sci. China Ser. D, Earth Sciences, 50 (10), 1441-1451, doi:10.1007/s11430-007-0071-3, 2007.
- Bowin, C., Scheer, E., and Smith, W.: Depth estimates from ratios of gravity, geoid, and gravity gradient anomalies, Geophysics, 51 (1), 123-136, doi:10.1190/1.1442025, 1986.
- 825 Capriotti, J., and Li, Y.: Gravity and gravity gradient data: Understanding their information content through joint inversions, Proceedings of the 84th SEG Annual International Meeting, Denver, Colorado, USA, October 2014, 1329-1333, 2014.
 - Chen, L., Zheng, T. Y., and Xu, W. W.: A thinned lithospheric image of the Tanlu Fault Zone, eastern China: Constructed from Wave equation based receiver function migration, J. Geophys. Res.: Sol. Ea., 111, B9, doi: 10.1029/2005JB003974, 2006.
- 830 Constable, S.C., Parker, R.L., and Constable, C.G.: Occam's inversion: A practical algorithm for generating smooth models from electromagnetic sounding data, Geophysics, 52 (3), 289-300, doi:10.1190/1.1442303, 1987.
 - Fang, J.: Three-dimensional density distribution of lithosphere in North China, Geophys. Geochem. Explor., 23 (3), 179-184, 1996.
 - Gao, S., Zhang, J.F., Xu, W.L., and Liu, Y.S.: Delamination and destruction of the North China Craton, Chinese Sci. Bull., 54, 3367-3378, doi:10.1007/s11434-009-0395-9, 2009.
 - Ge, C., Zheng, Y., and Xiong, X.: Study of crustal thickness and Poisson ratio of the North China Craton, Chinese J. Geophys., 54 (10): 2538-2548, doi:10.3969/j.issn.0001-5733.2011.10.011, 2011.

Hansen, P.C.: Analysis of discrete ill-posed problems by means of the L-curve, Siam Rev., 34 (4), 561-580, doi:10.1137/1034115, 1992.

- 840 Huang, J., and Zhao, D.: Seismic imaging of the crust and upper mantle under Beijing and surrounding regions, Phys. Earth Planet. In., 173 (3-4), 330-348, doi:10.1016/j.pepi.2009.01.015, 2009.
 - Laske, G., Masters, G., Ma, Z., and Pasyanos, M.: Update on CRUST1.0-a1-degree global model of Earth's crust, Geophysical Research Abstracts, 15, Abstract EGU2013-2658, doi:10.6092/1970-9870/128, 2013.

Li, H., Shen, C., Sun, S., Wang, X., Xiang, A., and Liu, S.: Recent gravity changes in China Mainland, Geodesy Geodyn., 2(1), 1-12, doi:10.3724/SP.J.1246.2011.00001, 2011.

845

860

875

- Li, H.L., Fang, J., Wang, X.S., Liu, J., Cui, R.H., and Chen, M.: Lithospheric 3-D density structure beneath the Tibetan plateau and adjacent areas derived from joint inversion of satellite gravity and gravity-gradient data, Chinese J. Geophys., 60 (6), 2469-2479, doi:10.6038/cjg20170634, 2017.
- Li, Y., and Oldenburg, D.W.: 3-D inversion of magnetic data, Geophysics, 61 (2), 394-408, doi:10.1190/1.1443968,1996.
- 850 Li, Z., Hao, T., and Xu, Y.: Uppermost mantle structure of the North China Craton: Constraints from interstation Pn travel time difference tomography, Chin. Sci. Bull., 56, 1691, doi.:10.1007/s11434-011-4487-y, 2011.
 - Liu, R., Wu, Z., Yin, C., Chen, Y., and Zhuang, C.: Development of China digital seismological observational systems, Acta Seismol. Sinica, 16(5), 568-573, doi:10.1007/BF02893477, 2003.

Moritz, H.: Geodetic reference system 1980, Bull. Geodesique, 54, 395-405, doi: 10.1007/BF02521480, 1980.

- Martinec, Z.: Mass-density Green's functions for the gravitational gradient tensor at different heights, Geophys. J. Int., 196 (3), 1455-1465, doi: 10.1093/gji/ggt495, 2014.
 - Nagy, D., Papp, G., and Benedek, J.: The gravitational potential and its derivatives for the prism, J. Geodesy, 74, 552-560, doi: 10.1007/s001900000116, 2000.
 - Pavlis, N.K., Holmes, S.A., Kenyon, S.C., and Factor, J.K. The development and evaluation of the Earth Gravitational Model 2008 (EGM2008), J. Geophys. Res.: Sol. Ea., 117, B04406, doi:10.1029/2011JB008916, 2012.
 - Pilkington, M.: 3-D magnetic imaging using conjugate gradients, Geophysics, 62 (4), 1132-1142, doi:10.1190/1.1444214, 1997.
 - Pilkington, M.: 3D magnetic data-space inversion with sparseness constraints, Geophysics, 74 (1), L7-L15, doi:10.1190/1.3026538, 2009.
- 865 Qin, P., Huang, D., Yuan, Y., Geng, M., and Liu, J.: Integrated gravity and gravity gradient 3D inversion using the non-linear conjugate gradient, J. Appl. Geophys., 126, 52-73, doi:10.1016/j.jappgeo.2016.01.013, 2016.
 - Rummel, R., Yi, W., and Stummer, C.: GOCE gravitational gradiometry, J. Geodesy, 85, 777-790, doi:10.1007/s00190-011-0500-0, 2011.
- Sebera, J., Šprlák, M., Novák, P., Bezděk, A., and Vaľko, M.: Iterative spherical downward continuation applied to magnetic
 and gravitational data from satellite, Surv. Geophys., 35 (4), 941-958, doi:10.1007/s10712-014-9285-z, 2014.
 - Sheng, S. Z., Wan, Y. G., Huang, J. C., Bu, Y. F., Li, X.: Present tectonics stress field in the Circum-Ordos region deducted from composite focal mechanism method, Chinese J. Geophys., 58 (2), 436-452, doi: 10.6038/cjg20150208, 2015.

Šprlák, M.: A graphical user interface application for evaluation of the gravitational tensor components generated by a level ellipsoid of revolution, Comput. Geosci., 46, 77-83, doi:10.1016/j.cageo.2012.04.013, 2012.

- Su, J., Zhu, W., Lu, H., Xu, M., Yang, W., and Zhang, Z.: Geometry styles and quantification of inversion structures in the Jiyang depression, Bohai Bay Basin, eastern China, Mar. Petrol. Geol., 26 (1), 25-38, doi:10.1016/j.marpetgeo.2007.08.003, 2009.
- Szwillus, W., Ebbing, J., Holzrichter, N.: Importance of far-field topographic and isostatic corrections for regional density modelling, Geophys. J. Int., 207(1), 274-287, doi:10.1093/gji/ggw270, 2016.
 - Teng, J., Wang, F., Zhao, W., Zhang, Y., Zhang, X., Yan, Y., Zhao, J., Li, M., Yang, H., Zhang, H., and Ruan, X.: Velocity structure of layered block and deep dynamic process in the lithosphere beneath the Yinshan orogenic belt and Ordos Basin, Chinese J. Geophys., 53 (1), 67-85, doi:10.3969/j.issn.0001-5733.2010.01.008, 2010.
- Teng, J., Zhang, Z., Zhang, B., Yang, D., Wan, Z., and Zhang, H.: Geophysical fields and background of exceptional
 structure for deep latent mantle plume in Bohai Sea, Chinese J. Geophys., 40, 468-480, 1997.
 - Tian, Y., Ke, X., and Wang, Y.: DenInv3D: A geophysical software for three-dimensional density inversion of gravity field data, J. Geophys. Eng., 15 (2), 354-365, doi:10.1088/1742-2140/aa8caf, 2018.
 - Tian, Y., Ke, X., and Wang, Y.: A folding calculation method based on the preconditioned conjugate gradient inversion algorithm of gravity gradient tensor, Pure Appl. Geophys., 176 (1), 215-234, doi:10.1007/s00024-018-1965-z, 2019a.
- 890 Tian, Y., Ke, X., and Wang, Y.: Inversion of three-dimensional density structure using airborne gradiometry data in Kauring test site, Geomatics and Information Science of Wuhan University, 44 (4), 501-509, doi: 10.13203/j.whugis20160503, 2019b.
 - Tian, Y., and Wang, Y.: Inversion of the density structure of the lithosphere in the North China Craton from GOCE satellite gravity gradient data, Earth Planets Space, 70, 173, doi:10.1186/s40623-018-0942-1, 2018.
- 895 Tian, Y., and Zhao, D.: Destruction mechanism of the North China Craton: Insight from P and S wave mantle tomography, J. Asian Earth Sci., 42 (6), 1132-1145, doi:10.1016/j.jseaes.2011.06.010, 2011.
 - Tian, Y., Zhao, D., Sun, R., and Teng, J.: Seismic imaging of the crust and upper mantle beneath the North China Craton, Phys. Earth Planet. In., 172 (3-4), 169-182, doi:10.1016/j.pepi.2008.09.002, 2009.
 - Uieda, L., Barbosa, V.C.F., and Braitenberg, C.: Tesseroids: Forward-modeling gravitational fields in spherical coordinates,
- 900 Geophysics, 81 (5), 41-48, doi:10.1190/geo2015-0204.1, 2016.
 - Wang, X.S., Fang, J., and Hsu, H.: Density structure of the lithosphere beneath North China Craton, Chinese J. Geophys., 55(4), 1154-1160, doi:10.6038/j.issn.0001-5733.2012.04.011, 2012.
 - Wang, X.S., Fang, J., and Hsu, H.: 3D density structure of lithosphere beneath northeastern margin of the Tibetan Plateau, Chinese J. Geophys., 56 (11), 3770-3778, doi:10.6038/cjg20131118, 10.1016/j.tecto.2013.11.002, 2013.
- 905 Wang, X., Fang, J., and Hsu, H.: Three-dimensional density structure of the lithosphere beneath the North China Craton and the mechanisms of its destruction, Tectonophysics, 610 (6), 150-158,doi:10.1016/j.tecto.2013.11.002, 2014.
 - Wu, L., Ke, X., Hsu, H., Fang, J., Xiong, C., and Wang, Y.: Joint gravity and gravity gradient inversion for subsurface object detection, IEEE Geosci. Remote S., 10(4), 865-869, doi:10.1109/LGRS.2012.2226427, 2013.

Xiong, X., and Teng, J.W.: Study on crustal movement and deep process in eastern Qinghai-Xizang Plateau, Chinese J.

- 910 Geophys., 45, 507-515, doi:10.1002/cjg2.266, 2002.
 - Xu, Y., Zeyen, H., Hao, T., Santosh, M., Li, Z., Huang, S., and Xing, J.: Lithospheric structure of the North China Craton: Integrated gravity, geoid and topography data, Gondwana Res., 34, doi:10.1016/j.gr.2015.03.010, 315-323, 2016.
 - Yang, S., Xiong, X., Zheng, Y., and Shan, B.: Upper-mantle temperature and lithospheric thickness of North China, Chinese J. Geophys., 56 (11), 3855-3867, doi:10.6038/cjg20131127, 2013.
- 915 Yi, W., Rummel, R., and Gruber, T.: Gravity field contribution analysis of GOCE gravitational gradient components, Stud. Geophys. Geod., 57 (2), 174-202, doi:10.1007/s11200-011-1178-8, 2013.
 - Zandt, G., and Ammon, C. J.: Continental crust composition constrained by measurements of crustal Poisson's ratio, Nature, 374 (6518), 152, doi:10.1038/374152a0, 1995.
 - Zhang, H. Q., Huang, Q. H., Zhao, G. Z., Guo, Z., and Chen, Y. J.: Three-dimensional conductivity model of crust and
- uppermost mantle at the northern trans North China Orogen: Evidence for a mantle source of Datong volcanoes,
 Earth Plant Sc. Lett., 453, 182–192, doi: 10.1016/j.epsl.2016.08.025, 2016.
 - Zhang, Y., Teng, J., Wang, F., Zhao, W., Li, M., and Wang, Q.: Structure of the seismic wave property and lithology deduction of the upper crust beneath the Yinshan orogenic belt and the northern Ordos block, Chinese J. Geophys., 54 (1), 87-97, doi: 10.3969/j.issn.0001-5733.2011.01.010, 2011.
- 925 Zhdanov, M.S., Ellis, R., and Mukherjee, S.: Three-dimensional regularized focusing inversion of gravity gradient tensor component data, Geophysics , 69 (4), 925-937, doi:10.1190/1.1778236, 2004.
 - Zhu, R.X., Yang, J.H., and Wu, F.Y.:Timing of destruction of the North China Craton, Lithos, 149 (15), 51-60, doi:10.1016/j.lithos.2012.05.013, 2012.

Zhu, R. X.: Review of the achievements of major research plan on "Destruction of North China Craton" (in Chinese), China

930 Sci Foundation, 32, 282–290.