

Dear editor and reviewer,

we sincerely thank Manel Prada for reading through the revised manuscript and point out some issues that need more clarification. We carefully worked through all the raised points. Please find below our answers for each comment in green coloured text. We used the revised manuscript to track the new changes. Line numbers given in our response correspond to

5 the manuscript without “track changes” visible.

On behalf of the authors,

with kind regards,

Anke Dannowski

## 10 **Reviewer report (Manel Prada)**

The authors have addressed most of the issues raised in the previous review, and the paper has improved considerably. However, as I mention in the attached PDF, there are minor to moderate issues that need to be explained/addressed to make the paper consistent.

15 The main moderate issue is that the authors should be more consistent with their main interpretation regarding the presence of exhumed mantle (i.e. whether Oligocene-Miocene extension led to extreme thinned continental crust or exhumed subcontinental mantle with a low grade of mantle serpentinisation remains enigmatic.) There are certain sections where the authors seem to favor the full exhumation of the mantle as opposed to the presence of hyperextended crust, which from the data they provide is not possible to discern.

Apart from this, I proposed some minor changes to the text and figures in the attached PDF file.

20

### **Point by point reply:**

Line 20 - perhaps the authors could mention here their hypothesis for the low grade of serpentinization (i.e. high sedimentation rate during syn-rift stage).

25 We included the hypothesis and re-phrased this sentence to: “... *thinned continental crust or exhumed subcontinental mantle remains enigmatic. A low grade of mantle serpentinisation indicates a high rate of syn-rift sedimentation.*”

Line 23 - this second sentence is a bit redundant with the first part of the sentence, please remove.

Deleted the second part of the sentence and re-phrased the two sentences to: “*However, rifting failed before oceanic spreading was initiated and continental crust thickens towards the NE within the northern Ligurian Basin.*”

Line 23 – Change “*is thickening*” to “*thickens*”

30 **Changed**

Line 45 – Change “*seismic refraction line*” to “*seismic refraction data*”

**Changed**

Line 182 – Change “*The starting models used in the analysis were 1D hanging below the seafloor (Fig. 4c).*” to “*The 1D starting models used were hanging below the seafloor (Fig. 4c).*”.

35 Changed

197 – Change “*crust, i.e. the crystalline basement.*” to “*(i.e. crystalline basement; CB in Fig. 4a)*”

Changed to “*crust, i.e. the crystalline basement (CB in Fig. 4a).*”

Line 211 – “*We interpret this section as syn-rift sediments, possibly ...*”

To be consistent with the abstract and the uncertainty of the nature of the rock at these depths the authors should also mention  
40 the possibility here that these velocities account for tilted fault blocks of continental crust.

We added the sentence: “*We will discuss the nature of these layers in a later section, since the observed seismic velocities also account for tilted fault blocks of stretched continental crust.*”

Line 244 – Change the second part of the sentence from “*that no typical oceanic crust and no thick layer of gabbroic rocks is present along the profile*” to “*the lack of a thick gabbroic layer, and thus, the lack of typical oceanic crust*”.

45 Changed.

246 – Change “*in the Tyrrhenian Sea (Prada ...)*.” to “*in the Central Tyrrhenian Sea (Prada ...)*.”.

Changed

Line 253 - As the authors mention later in this section (L. 282-284), they do not know whether exhumation or hyper-extension of the crust occurs in this region. I would suggest to remove this sentence to be consistent with the message of this section.

50 We would like to keep the second part of the sentence, since by this we open the discussion in the next section.

258 – Change “*high uncertainty*” to “*higher uncertainty*”

Changed

Line 270 – Change “*A continental crustal thickening*” to “*The thickening of the continental crust*”.

Changed

55 Line 271 – Change “*modelling of the free air anomaly (Fig. 5).*” to “*gravity modelling (Fig. 5)*”.

Changed

Lines 271 to 274 - This paragraph is not clear, and does not provide much as you already mention the lack of oceanic crust. I suggest to remove it.

We like to keep this sentence, because it supports our discussion that we observe continental crust. The observation of crustal  
60 thickening towards the NE contradicts an oceanic type of crust which we would expect to thin out towards the NE. We changed the last part of the sentence to make it more clearly to: “*... an abrupt change from oceanic to continental crust or oceanic crust to gradually thin out towards the NE.*”.

Line 285 – Delete “very well”

Deleted

65 Line 285 - I'm confused here. The authors interpret  $V_p$  between 4.5-5.7 km/s as syn-rift sediments (L. 224-225), then they suggest that mantle exhumation may have occurred (L. 267), but now it seems that their interpretation is unequivocally that hyper-extended crust is overlying the mantle. The authors need to be consistent along the manuscript and support a unique

interpretation (which may include both syn-rift or hyper-extended crust as they have way to discern between them), otherwise it turns to be rather confusing.

70 We try to clarify this and re-phrased this sentence at the end. This comparison is related to the northern end of our profile, where we indeed observe continental crust velocities. We changed this paragraph to: *“Our profile only provides information on the basin centre where the absolute velocities of Le Douaran et al. (1984) and Contrucci et al. (2001) fit continental crust velocities and support our interpretation for the northern end of the profile where we observe mantle material beneath thinned continental crust.”*

75 Line 289 – Remove the sentence: *“Seismic velocities of ~7.3 km/s and higher are too fast for magmatic crust (Grevemeyer et al., 2018; Christeson et al., 2019).”*.

Removed

Line 296 – Remove *“or a result of mafic intrusion”* - a mafic (e.g. gabbro) intrusion cannot have  $V_p > 7.8$  km/s, perhaps ultramafic rocks may reach to that velocities, and in this setting there is no evidence of crustal magmatic velocities as the authors say, so that I would remove this.

Removed

Lines 296 to 299 – Change *“that syn-rift sediments (nowadays showing high P-wave velocities) may have been directly deposited on top of the mantle or brittle continental crust (Pérez-Gussinyé, 2013).”* to *“that either continental crustal blocks are overlying the mantle, and thus it was not fully exposed during rifting or that syn-rift sediments (nowadays showing high P-wave velocities) may have been directly deposited on top of the mantle preventing its full serpentinisation (Perez-Gussinyé et al., 2013).”*.

Changed

Line 300 – This is an overstatement, as there are regions of the Atlantic margin that may not be as serpentinized as the Goban Spur and the Iberia margin. So perhaps the best here is to keep the comparison to these two regions rather than the entire Atlantic margin. In addition I would encourage the authors to include some values of serpentinisation from the Goban Spur and Iberia, so the reader can have a sense on how much different is the degree of serpentinization between the presented Atlantic margins and the Ligurian.

We re-phrased the sentence to *“However, the fast mantle in the Ligurian Sea would support a much lower degree of serpentinisation when compared to these two regions with a mantle serpentinisation of 100% at the top basement and <25% in 5-7 km depth at the Goban Spur and >75% at the top basement and <25% in 2 km depth at the Iberia margin.”*

Line 304 – Change *“In comparison to the Tyrrhenian Sea, ...”* to *“In comparison to the Central Tyrrhenian Sea, where exhumed mantle is inferred, ...”*

Changed

Line 305 – Change *“high (Fig. 4a).”* to *“faster (Fig. 4a).”*

100 Changed

Line 305 – Change “*Tyrrhenian Sea Domain #3*” to “*Magnaghi and Vavilov basins in the Tyrrhenian Sea (i.e. Domain #3 in Prada et al., 2014)*”.

#### Changed

105 Lines 310 to 314 – The Ligurian has a thicker syn (?) and post-rift layer of sediments than that of Domain #3 in the Tyrrhenian, that is a fact I agree. However, there is no quantitative evidence (at least is not provided here) that the Ligurian had a higher sedimentation rate than the Tyrrhenian.

The Ligurian is far older than the Domain #3 in the Tyrrhenian, which did not formed 9-10 Ma ago as the authors refer, but <5.9 Ma ago (Kastens and Mascle, et al., 1990 ODP results; Prada et al., 2016) and, consequently really fast (> 2-3 cm/yr ; Faccena et al., 2001). Thus, the Ligurian stopped opening 10Ma earlier than the opening of Domain #3 at 15-16Ma. 10 Ma  
110 gives a lot of time for sediments to cover up the basin, so there is no need to invoke a higher sediment rate to explain the difference in sediment thickness, but the difference in age of formation.

The authors use the presence of syn-rift sediments from the Gult of Lions to reinforce the hypothesis that a thicker syn-rift cover might be overlying the mantle in the Ligurian than in the Tyrrhenian. This could be a possibility, but the authors need to consider also the fact that the continental margin and thus the source of syn-rift sedimentation was closer to the center of  
115 the basin during the formation of the Ligurian than during the formation of Domain #3 in the Tyrrhenian, given that the Sardinia and Campanian margins were already formed, thus the center of the basin (where Domain #3 formed) in the Tyrrhenian was already at least 100 km away from emerged land (Prada et al., 2014).

Overall, once could would agree that perhaps the syn-rift sedimentary cover is thicker in the Ligurian than in Domain #3 in the Tyrrhenian because of the latter aspect (i.e. proximity of the margin to the center of the basin). In contrast, the hypothesis  
120 of a higher sedimentation rate looks rather weak and not supported by any quantitative observation.

We agree entirely with the reviewer’s comments and we re-phrased this section to make it more clear to the reader. We corrected the timing of the opening of the Tyrrhenian Sea and divided it to extension and spreading and provided the references. We changed the order of the observed differences between both basins and discuss them separately. While we used most of the phrasings we changed the order and added/changed 2 sentences. These changes are summarised in the one changed section:  
125 “(2) *The Ligurian Basin was stretched ~150 km during the ~16 million year opening phase, while the Tyrrhenian Sea was stretched ~170 km within ~5 million years prior the onset of oceanic spreading (Faccenna et al., 2001, Prada et al., 2014); thus, the stretching rate in the Tyrrhenian Sea was higher compared to the Ligurian Sea. Further, (3) the Ligurian Basin has a thick sedimentary cover of ~6-8 km, while the Tyrrhenian Sea Domain #3 shows a sedimentary cover of ~1-2 km (Prada et al., 2014). Syn-rift sedimentation was recorded in MCS data (Fig. 1, inlay profile 3) in the Gulf of Lion (Jolivet et al., 2015).*  
130 *The proximity of the two basins to the continental margin during their formation might result in a different syn-rift sedimentation rate that possibly was higher in the Ligurian Sea compared to the Tyrrhenian Sea.*”

Line 320 – This sentence needs to be rephrased, right now it does not make much sense.

Re-phrased to: “Also the interpretation of extremely thinned brittle continental crust requires syn-rift sedimentation since the stretching might open fluid pathways through the crust down to the mantle and would lead to a high degree of mantle serpentinitisation (Nagel and Buck, 2004).”

135 Line 336 – Change “The exhumation of” to “The nature of the COT along the Gulf of Lions is still debated”.

We ex-changed the entire sentence to: “The nature of the COT along the Gulf of Lion is still debated.”.

Line 343 – Change “... imaged in the travel time tomographic approach with the typical pattern observed at mid-ocean ridges (Gailler et al., 2009), with a high velocity gradient in the upper oceanic crust and a low velocity gradient in the lower crust.”

140 to “... interpreted on the basis of a 2D Vp model derived from travel time tomography as an anomalously thin oceanic crust (4–5 km) with the typical two-layer gradients clearly characteristic of oceanic layers 2 and 3 (Gailler et al. 2009).”

Changed to: “... interpreted on the basis of a 2D P-wave model derived from travel time tomography as an anomalously thin oceanic crust (4–5 km) with the typical two-layer gradients clearly characteristic of oceanic Layers 2 and 3 (Gailler et al., 2009).”

145 Line 362 – Change “..., which was indeed not mapped in MCS data so far.” to “..., absent in MCS data so far.”

We like to keep the original phrasing.

Line 365 - careful here, calc-alkaline signature is typical from subduction-related melts, not with post-rift magmatism. The fact that makes this seamount a potential post-rift feature is the age of basalts, which are younger than the age of the rifting (16Ma). The signature of the melt can be biased by the contamination of the subduction-related fluids.

150 Change to: “... 2012). The age is clearly indicating ...”.

Line 390 – Yet this phase is not certain from your observations. The authors should emphasize on that.

We included the sentence: “Yet, it is debated if we observe the latter phase in the central Ligurian Basin.”.

Line 403 – Delete the sentence and add the information to the previous one: “... pattern of magmatic domains, supporting again the lack of oceanic spreading during the formation of the Ligurian basin in the Oligocene-Miocene.”

155 Changed

Line 403 to 405

This is confusing towards the North where the Makris line is the crust thickens, thus the mantle won't exhume because there is not enough extension, not because it's not lasting enough. I would remove this sentence.

160 Changed to: “Continuing further north of our seismic line, the Oligocene-Miocene extension led to thinned continental crust, but the amount of extension was too small to extremely thin out the continental crust and exhume mantle.”

Line 412 – add “syn-rift” and change “thinned” to “thin”.

Changed

Line 416 - Again, from your data it is hard to conclude that the mantle is fully exposed. The authors should leave it as a they state above:

165 “it remains enigmatic if the mantle is overlain directly by sediments or by extremely thinned continental crust of up to 2.5 km”

We added this sentence: “*It remains enigmatic if the mantle is overlain directly by sediments or by extremely thinned continental crust of up to 2.5 km thickness.*”

Line 424 – Delete “*however, nearby*”

Removed

170 Figure 1 – yellow and orange triangles are mentioned in the figure caption 3 but not in this one. It should included here as it is the first figure where they appear.

Added

Figure 4 – the inset showing the 1D profiles in (d) needs to be larger as (c).

175 We keep it as it is, since the scale of the 1D profiles in Fig. 4d is horizontally the same and vertically larger than in 4c. It is an important background information on the modelling, however, we feel it is large enough to spot the tested model space.

**Marked-up manuscript:**

## 180 **Seismic evidence for failed rifting in the Ligurian Basin, Western Alpine Domain**

Anke Dannowski<sup>1</sup>, Heidrun Kopp<sup>1,2</sup>, Ingo Grevemeyer<sup>1</sup>, Dietrich Lange<sup>1</sup>, Martin Thorwart<sup>2</sup>, Jörg Bialas<sup>1</sup>, and Martin Wollatz-Vogt<sup>1</sup>

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany

<sup>2</sup>CAU, Christian-Albrechts-Universität zu Kiel, Germany

185 *Correspondence to:* Anke Dannowski (adannowski@geomar.de)

**Abstract.** The Ligurian Basin is located in the Mediterranean Sea to the north-west of Corsica at the transition from the western Alpine orogen to the Apennine system and was generated by the south-eastward trench retreat of the Apennines-Calabrian subduction zone. Late Oligocene to Miocene rifting caused continental extension and subsidence, leading to the opening of the basin. Yet, it remains enigmatic if rifting caused continental break-up and seafloor spreading. To reveal its lithospheric architecture, we acquired a 130-km long seismic refraction and wide-angle reflection profile in the Ligurian Basin. The seismic line was recorded in the framework of SPP2017 4D-MB, the German component of the European AlpArray initiative, and trends in a NE-SW direction at the centre of the Ligurian Basin, roughly parallel to the French coastline.

195 The seismic data were recorded on the newly developed GEOLOG recorder, designed at GEOMAR, and are dominated by sedimentary refractions and show mantle Pn arrivals at offsets of up to 70 km and a very prominent wide-angle Moho reflection. The main features share several characteristics (i.e. offset range, continuity), generally associated with continental settings rather than documenting oceanic crust emplaced by seafloor spreading. Seismic tomography results are complemented

by gravity data and yield a ~6-8 km thick sedimentary cover and the seismic Moho at 11-13 km depth below the sea surface. Our study reveals that the oceanic domain does not extend as far north as previously assumed. Whether Oligocene-Miocene extension led to extreme thinned continental crust or exhumed subcontinental mantle ~~remains enigmatic. A~~with a low grade of mantle serpentinisation ~~indicates a high rate of syn-rift sedimentation~~~~remains enigmatic~~. However, rifting failed before oceanic spreading was initiated, ~~accompanied by the formation of mantle derived oceanic crust, and Continental~~continental crust ~~is thickening~~~~thickens~~ towards the NE within the northern Ligurian Basin.

## 1 Introduction

The Ligurian Sea is situated in the north-western Mediterranean Sea at the transition from the western Alpine orogen to the Apennine system. The geodynamic setting of the area is controlled by the convergence of the African and Eurasian plates (e.g. Dercourt et al., 1986). Despite the existing large collection of seismic and other geophysical data, the present-day crustal architecture of the Ligurian Basin is still under discussion and the kinematic boundaries are poorly resolved, in particular, the continent-ocean transition (COT) along the margins as well as its termination to the north-northeast. Imaging clear fault structures within the crust has proven challenging due to the presence of thick Messinian salt layers and due to the masking effect of the first seafloor multiple which roughly coincides with the arrival of the reflection of the acoustic basement (Béthoux et al., 2008). Deep drilling data are lacking and the magnetic data are complex and anomalies discontinuous (Bayer et al., 1973). Based on integrated seismic and magnetic data, maps indicating the extent of the oceanic domain were created (i.e. Burrus, 1984; Gueguen et al., 1998; Rollet et al., 2002), however, no axial ridge was imaged near the centre of the basin (Rollet et al., 2002). To explain the mismatch between the expected oceanic domain and the observed seismic signal, the crust in the north-eastern basin was interpreted to be ‘atypical’ oceanic crust (Mauffret et al., 1995; Chamot-Rooke et al., 1997; Contrucci et al., 2001; Rollet et al., 2002). A clear change from continental to oceanic crust was only shown for the southern area of the Ligurian Basin, in the Gulf of Lion and offshore Sardinia (Gailler et al., 2009). It is proposed that the oceanic domain is separated from the continental margins by a transitional domain characterised by a high-velocity lower crust (Fig. 1). An overview of seismic experiments until 2002 is presented in Rollet et al. (2002). Furthermore, the area was revisited or data were re-analysed with modern seismic techniques including the CROP deep seismic profiles (Finetti et al., 2005), the TGS-NOPEC and the SARDINIA profiles (Gailler et al., 2009; Jolivet et al., 2015), as well as more recent studies along the French and Italian Riviera with the 3D seismic refraction GROSMarin project (Dessa et al., 2011) and an amphibious ambient noise study (Guerin et al., 2019).

In the frame of the LOBSTER project, we obtained a new state-of-the-art seismic refraction ~~line~~~~data~~ (Fig. 1, red line with orange and yellow triangles). Here, we present the analysis of the seismic refraction data from the central Ligurian Basin, which is the extension of a pre-existing seismic profile (Makris et al., 1999), which we call MAKRIS (Fig. 1, black line). We aim to unravel the present-day crustal structure and its nature in the centre of the Ligurian Basin, map the depth of the crust-mantle boundary (seismic Moho), and reveal the styles of deformation during the last extensional phase. We investigate the

hypothesis that Oligocene-Miocene rifting led to either extended continental crust or exhumation of sub-continental mantle  
230 below post-rift sediments in the north-eastern Ligurian Basin.

## 2 Geological structures and geodynamics of the Ligurian Sea and the Corsica-Sardinia block

The Ligurian Sea has a width of ~150 km, reaching from the northern tip of Corsica to the Ligurian coast near the city of Sanremo. It widens towards the southwest to ~175 km between Calvi (Corsica) and Cannes. South of an imaginary line between Ajaccio (Corsica) and Toulon, the Ligurian Sea is roughly 225 km wide and opens entirely towards the Balearic Sea. The  
235 Ligurian Basin itself is smaller with a width of 70 km, 120 km, and 170 km, respectively, along the three dashed grey lines in Figure 1 and the seafloor reaches a depth of ~2700 m. The Ligurian margin is characterised by a narrow and steep slope (10-20 km) with a few listric normal faults (Finetti et al., 2005). The Corsica slope is wider (20-50 km) and the margin is characterised by several listric faults extending over a wider area (-Contrucci et al., 2001; Rollet et al., 2002).

The Ligurian Sea formed as a back-arc basin at the transition from the western Alpine orogen to the Apennine system (e.g.  
240 Doglioni et al., 1997; Faccenna et al., 1997; Réhault et al., 1984). The Alpine transition is characterised by a change in subduction polarity between the two orogens (Jolivet and Faccenna, 2000; Handy et al., 2010). The Ligurian Basin is the oldest back-arc basin in the Western Mediterranean Sea and developed from Late Oligocene to Early Miocene (Réhault and Béthoux, 1984; Roca and Desegaulx, 1992; Fernández et al., 1995; Jolivet and Faccenna, 2000; Rosenbaum et al., 2002; Finetti et al., 2005; Advokaat et al., 2014). The extension is related to the south-east trench retreat of the Apennines-Calabrian subduction  
245 zone initiated in the Oligocene (Montigny et al., 1981; Réhault and Béthoux, 1984; Vigliotti and Langenheim, 1995; Gueguen et al., 1998; Rosenbaum et al., 2002; Faccenna et al., 2001).

Rifting has initiated ~30 Ma ago at a rate of ~1 cm/yr in the NE and ~2 cm/yr in the SW (Rollet et al., 2002). The initiation is associated with magmatism on land along the western Ligurian margin (Rollet et al., 2002). At roughly 21 Ma, rifting terminated while an anticlockwise rotation of the Corsica-Sardinia block was initiated (Rollet et al., 2002; Speranza et al.,  
250 2002). During this phase, the commencement of oceanic spreading was proposed (Pascal et al., 1993; Contrucci et al., 2001; Rollet et al., 2002; Finetti et al., 2005). These authors referred to tholeiitic volcanic edifices to solidify their interpretation and interpreted the pattern of magnetic data (Bayer et al., 1973) to be a result of two main discontinuous volcanic lineaments, sub-parallel to the basin axis related to oceanic spreading and unroofing of mantle material. The opening of the Ligurian Basin ended ~16-15 Ma ago and was associated with a second calc-alkaline volcanic phase along the Corsican margin (Rollet et al.,  
255 2002) that is linked to the migration of the subducting lithosphere towards the E-SE. The extension of the Ligurian Basin terminated and shifted to the Tyrrhenian Sea while the Apennines-Calabrian subduction zone continued to roll back further southeast until late the Messinian, ~6 Ma (Faccenna et al., 2001; Advokaat et al., 2014). The opening rate was calculated with 7.8-10.3 mm/yr (Moeller et al., 2013). In the north of the Tyrrhenian Sea, extension led to continental crustal thinning (Moeller et al., 2013), while further south in the centre of the Tyrrhenian Basin, the mantle was exhumed and serpentinitised and intruded  
260 by Mid-Ocean-Ridge type (MOR-type and intraplate basalts) (Prada et al., 2016). Similar to the Ligurian Basin, the Tyrrhenian



Sea shows distributed, non-linear magnetic anomalies (Cella et al., 2008). Anomalies often coincide with volcanic islands, seamounts or other morphological units of igneous composition. During the Ocean Drilling Project (ODP) Leg 107 at site 651, serpentinised mantle rocks were drilled forming the top of the basement (Bonatti et al., 1990).

265 Gueguen et al. (1998) and Rollet et al. (2002) suggest that the central Ligurian Basin is comprised of oceanic crust. These authors divided the basin into different zones of continental and oceanic domains based on seismic, magnetic and gravity data (Fig. 1): (1) atypical oceanic crust with (2) transitional zones to (3) continental crust. The location of the northeast-southwest trending continent-ocean transition is proposed to be situated in the vicinity of the volcanic Tristanites Massif (Fig. 1) (Makris et al., 1999) (yellow bar perpendicular to the MAKRIS profile in Figure 1). Based on re-analysed expanding spread profiles (ESP), Contrucci et al. (2001) proposed a 40 km wide area of oceanic crust near the Median Seamount (Fig. 1).

270

### 3 Data acquisition, processing, and modelling

Data at different scales resolving the subsurface structure were acquired in the Ligurian Sea in February of 2018 during the cruise MSM71 aboard the German research vessel Maria S. Merian (Kopp et al., 2018). Active seismic refraction data were obtained along the centre of the basin. Our NE-SW trending seismic refraction and wide-angle reflection line is situated in the 275 prolongation of an existing refraction profile in the northern Ligurian Basin (Makris et al., 1999) (Fig. 1).

#### 3.1 Data acquisition and processing

The active seismic data were simultaneously recorded on short period ocean bottom seismometers (OBS) and ocean bottom hydrophones (OBH) as well as on a short streamer (280 m long) that was towed behind the vessel at 5 m water depth. Additionally, Parasound sediment echo sounding data were recorded along the profiles. The 127.5 km long refraction seismic 280 profile consists of 15 OBH/OBS at a station spacing of ~ 8km (Fig. 1). A total of 1079 shots were fired by an ~89-liter (5420 inch<sup>3</sup>) G-gun array, consisting of 2 sub-arrays. Each sub-array with a cluster of 2x8.5 litres (520 inch<sup>3</sup>), followed by a cluster in the middle of 2x6.2 litres (2x380 inch<sup>3</sup>, port) and 2x4.1 litres (2x250 inch<sup>3</sup>, starboard), and the third cluster again of 2x8.5 litres for both sub-arrays. The array with a string distance of 12 m was towed at 8 m below the sea-surface and 40 m behind the vessel. A shot interval of 60 s resulted in a shot distance of ~123 m. The guns were shot at ~190 bar providing a dominant 285 frequency band of approximately 5-70 Hz. The location of the stations on the seafloor was determined using the symmetry of the direct water arrivals from the shots on both sides. For this purpose, the direct arrival was picked and the deviation between computed and real travel times was minimised by adjusting the OBS's position along the profile. Dislocation off-line cannot be corrected with this method. For 2D traveltimes modelling, the stations were projected on to the profile. The airgun shots were recorded using newly developed GEOLOG data loggers designed at GEOMAR. All recorders operated reliably during 290 the deployment of 2 days with a negligible absolute clock drift between -1.03 ms and +0.72 ms. The sampling frequency was 250 Hz. The data processing included the conversion of the continuous data from GEOLOG format into the standard

continuous SEG-Y format using the GEOLOG programming interface. Afterwards, the continuous SEG-Y data were converted into standard trace-based SEG-Y format (Fig. 2b). Simultaneously, the clock drift was corrected, a step important for OBS data, since the instruments cannot be continuously synchronized via GPS during deployment as commonly done onshore. A gated Wiener multi-trace deconvolution with an autocorrelation average of 51 traces was applied to the shot gathers to compress the basic wavelet, to leave the Earth's reflectivity in the seismic trace and to remove the source signature and the hydrophone and geophone responses.

### 3.2 The GEOLOG recorder

The GEOLOG is a 32-bit seismic data logger designed to digitise data from a three-component seismometer and a hydrophone. We recorded the hydrophone output on two channels (channels 1 and 5) at two different amplification levels providing well amplified long-range records (gain=16) and preventing clipped amplitudes from short-range airgun shots (gain=1) to minimise difficulties with amplitude restoration because no gain range was implemented. The gain for the three seismometer channels 2 to 4 was set to 16, which provided good signal to noise ratios for all record offsets without clipping of amplitudes. Two additional analogue pins can be used as general-purpose input/output (GPIO) for measuring power levels for example. 3.3 V and 5 V connectors can serve external devices. Sampling intervals between 50 Hz and 4 kHz are controlled either by an atomic clock or by a temperature compensated clock (SEASCAN). We used an external GPS receiver for synchronization of the internal clock prior and after deployment, which was driven by the GEOLOG itself. Our seismic data were stored on two micro SD cards with a volume of 32 GB each. The recorder has been tested and proved reliable for writing speeds and SD cards of up to 128 GB (larger capacities are possible). The low power consumption of 375 mW (average battery drain) allowed us to save batteries. We used only 8 alkaline batteries per station for our short-term deployment. Thus, using lithium batteries, long-term deployments of more than 9 months can be performed. Battery power can further be saved by a delayed start of recording up to 31 days after programming. We set the recording parameters, i.e. the number of channels, gain and sampling rate, using a graphical user interface. The recorders can be programmed through any terminal program on a Windows or Linux operating system. The programming device was connected via RS232 using an RS232-USB adapter. A second RS232 interface can be used to drive external sensors (e.g. levelling of broadband seismometers). The GPS system used for the internal clock time synchronisation was developed together with the recorder and can operate with GPS, GLONASS, GALLILEO and QZSS enabling operation worldwide and in polar regions. Besides stable output of NMEA data (defined by the National Marine Electronics Association) and a PPS (pulse-per-second) time signal, the German DCF-77 code is also available. Moreover, the GPS system is available to deliver time or distance based trigger with TTL output, NMEA sequence and records of time stamps on an SD card.

### 3.3 Data description and analysis

The airgun shots can be followed for offsets up to 60 km on all 15 stations (Fig. 2). In general, the sections look very similar with clear sedimentary arrivals and wide-angle Moho reflections (PmP) as well as mantle phases (Pn) at a critical distance

between 25 km and 35 km to the stations (Fig. 2a). Although phase arrivals show common features in all record sections (Fig. 2a), the characteristics of the seismic phases change slightly from south to north (Fig. 2b-2d).

As a result of decreasing water depth towards the northeast, the direct wave through water (Pw) arrive later at the southern stations than at the northern stations (Fig. 2a). Arrivals from a shallow sedimentary reflection phase (PsP) occur approximately 0.5 s to 1 s after the direct wave and result from the top of salts that become shallower towards the north (as imaged in the multichannel seismic data in Fig. 3a). The red picks (Ps1) and the orange picks (Ps2) (Fig. 2b-2d) are interpreted as refracted phases through Plio-Quaternary and older sediments, respectively. The apparent seismic velocity of the Ps2 is very constant at ~4.3 km/s to ~4.6 km/s. The phase shows many undulations and some shadow zones (Fig. 3b) caused by the salt unit that displays intense doming and is possibly disrupted by some volcanic structures that are imaged in the MCS (Fig. 3a) and the Parasound data (Fig. 3c). This phase continues as a secondary arrival (Ps3) with a similar apparent velocity of ~4.6 km/s at the southern stations but disappears at the northern stations. Based on the apparent velocity and forward modelling, we interpret phase Ps3 as a refracted phase through the sediments. Simultaneously, when phase Ps3 disappears from OBS208 towards the north (compare to OBS209 (Fig. 2c), where Ps3 only occurs on the southern branch), an additional refracted phase (Pg) (green picks in Fig. 2c-2d) occurs with an increasing range of offsets observed on the stations northwards. The phase has an apparent velocity of ~6.2 km/s. At an offset of about 25 km, an abrupt change in the apparent seismic velocity to ~8 km/s occurs for the first arrival, as typically observed in the oceanic upper mantle. The yellow picks (Fig. 2b-2d) are refracted mantle phases (Pn) that show a similar apparent seismic velocity of ~8 km/s at the northern stations. However, the critical distance at the northern stations moves to slightly larger offsets of up to ~30 km. Furthermore, an earlier very short reflection occurs at 20-25 km offset. Pn phases at the southern stations are very weak, while the PmP is relatively strong compared to typical oceanic crust characteristics. The observed slight changes in the seismic signal are accompanied by slight changes in the free-air gravity anomaly around profile KM 60 (approx. 20 km south of OBS209), as discussed below.

### 3.4 P-wave traveltimes tomography modelling strategy and parameters

A preliminary seismic velocity model was build using RAYINVR (Zelt, 1999) to (1) reveal the overall structure of the profile, (2) manually assign the picked phases to certain layers, and (3) serve as starting point for the travel time tomography. Travel times were picked on the hydrophone channels using the interactive analysis tool for wide-angle seismic data PASTEUP (Fujie et al., 2008). The overall quality of the hydrophone data was slightly better compared to the vertical geophone channel, however, the vertical component was used for picking to confirm and to complement the picks observed on the hydrophone channel. In addition, multiples were picked when above the noise level (because of constructive interference) and where primary waves are below the noise level (Meléndez et al., 2014). Picks of water layer multiple phases were used during the forward modelling approach to confirm the layer boundaries and seismic velocities. Thereafter, a travel time tomographic inversion (tomo2D from Korenaga et al., 2000) was applied to invert the seismic P-wave velocity model and yield model uncertainties. The picks were assigned pick uncertainties ranging from 20 ms for clear near offset phases (Ps1), 30 ms for intermediate offsets (Ps2 and Pg), and up to 50-70 ms for picks at larger offset (Pn and PmP) taking into account the decreased

resolution due to the increased wave length of the seismic signal and the decreased signal-noise-ratio. Subsequently, all first arrivals and the mantle reflections were inverted with a set of starting models that converged to  $\chi^2$  values of less than 1 within 5 iterations. To test the model space and its limits, starting models, ranging from velocities between 1.8 km/s and 2.5 km/s at the seafloor with different velocity gradients, and ranging from 4.5 km/s to 7.5 km/s at 12-13.5 km depth to mimic the different types of crust, were manually created using RAYINVR (Zelt, 1999). [The starting models used in the analysis were 1D hanging below the seafloor \(Fig. 4e\). The 1D starting models were hanging below the seafloor \(Fig. 4c\).](#) To carefully evaluate the resulting velocity models, we used three criteria: (1) travel times need to fit the data (Fig. 2a), (2) travel time residuals, RMS misfit and  $\chi^2$  had to be low (i.e.  $\chi^2 \sim 1$ ), and (3) the gravity response (calculated after a velocity-density conversion after Korenaga et al., 2001) of the resulting density model must yield comparable results to the satellite gravity data. Based on this evaluation, 17 models (Fig. 4c) were chosen to generate an average model for the crustal part (Fig. 4a, above the Moho) and the standard deviation was calculated (Fig. 4b). Overall, the standard deviation in the crust down to the acoustic basement is smaller than 0.15 km/s, indicating small differences between the inverted velocity models and hence an excellent resolution. Random Gaussian noise was not added, to the travel time picks, however, during modelling re-picking of phases (mainly fine adjustments to the picks) did not lead to major differences in the resulting velocity model. In a further step, the average model was edited by adding different 1D profiles with mantle velocities underneath the crust-mantle boundary (inlay in Fig. 4d). A set of 14 mantle velocity starting models was used to invert for refracted mantle phases, while the model above the seismic Moho was overdamped. Again, an average model and the standard deviation for the mantle were calculated (Fig. 4d). Standard deviations for the mantle P-wave velocities are small ( $<0.1$  km/s), indicating a good resolution of upper mantle velocities. Lastly, the very short reflected phases interpreted to result from the top of continental crust were calculated as a floating reflector without implementing a velocity discontinuity into the model to confirm the top of crust, i.e. the crystalline basement ([CB in Fig. 4a](#)).

## 4 Results

### 4.1 Seismic P-wave velocity distribution

In general, the average P-wave velocity along the profile (Fig. 4a) shows only minor lateral variations, mainly caused by the salt layers and the corresponding tectonic features at 4-6 km depth. The uppermost portion of the velocity model is characterised by a strong velocity gradient of  $\sim 1 \text{ s}^{-1}$  that is laterally constant. P-wave velocities increase from 2.2 km/s at the seafloor to 3.5 km/s approximately 1.3 km depth below the seafloor. We interpret this unit as Plio-Quaternary sediments mixed with the upper evaporite unit after Rollet et al. (2002), using their multi-channel seismic data profile MA24 (Fig. 1, inlay profile 6). The Plio-Quaternary sediments are imaged as horizontally layered strata in the multi-channel seismic data in Figure 3a. This high velocity-gradient layer thins towards the north, from 1.5 km to 1.2 km thickness and shows slightly slower velocities at the southern end (2.2 km/s) compared to the northern end (2.4 km/s) at the seafloor. Between  $\sim 4$  km and 6 km depth, the velocities range from 3.5 km/s to 4.5 km/s, and there are areas where minor velocity inversions are observed. These

low velocity units have a lateral extent of up to 10 km and a velocity contrast of up to ~0.2 km/s. We identify this section as the Messinian salt unit. From 6 km to ~10 km depth, the seismic velocities increase from ~4.5 km/s at the top to 5.7 km/s at the bottom. We interpret this section as syn-rift sediments, possibly Aquitanian according to Jolivet et al. (2015), to post-rift sediments, until Pre-Messinian. [We will discuss the nature of these layers in a later section, since the observed seismic velocities also account for tilted fault blocks of stretched continental crust.](#)

In the north-eastern half of the profile, starting roughly at profile KM 70, we determine the crystalline basement (CB) (red dashed line in Fig. 4a) at a depth of 10 km to 11.5 km below the sea surface. The basement velocities increase from 5.8 km/s to 6.6 km/s (marked with “Y” in Fig. 4a); they are interpreted, based on absolute velocities, as continental crust, thickening towards the north. The acoustic basement here is at a depth of ~10 km below the sea surface. At the opposite southern half of the profile, we could not identify the CB in the OBS data. However, a strong velocity jump occurs from 5.7 km/s to  $\geq 7.3$  km/s that we interpret as the crust-mantle boundary (Moho). The uppermost mantle is characterised by seismic velocities  $> 7.3$  km/s that increase to ~8 km/s over a depth interval of 2-3 km. The histogram (Fig. 4f) images a gap in seismic velocities between 6.6 km/s and 7.3 km/s, which suggests that no fresh oceanic crust material (gabbroic rocks) is present along the profile.

## 4.2 Gravity modelling

To constrain the crustal structure along the profile, we calculated the gravity response (Talwani et al., 1959) of the final seismic velocity model and compared it to the free-air gravity anomaly derived from satellite data (Sandwell et al., 2014). The fact that the profile is situated in the centre of the basin allows us to assume that only minor 3D side-effects occur in our 2D-modelling approach, caused by topography. The velocity-depth distribution was used to assign densities by applying different density-velocity relations. The water layer is assumed to have a density of 1.03 g/cm<sup>3</sup>. Gardeners rule,  $\rho = 1.74 * V_p^{0.25}$ , valid for sediments between 1.5 km/s  $< V_p < 6.1$  km/s (Gardner et al., 1974), was used for the sedimentary layers. For crystalline (non-volcanic) rocks the relation:  $\rho = 0.541 + 0.3601 * V_p$  (Christensen and Mooney, 1995) was used. A density of 3.3 g/cm<sup>3</sup> was assigned to the mantle. In areas with reduced seismic mantle velocities, the mantle density was reduced to 3.15 g/cm<sup>3</sup> (Carlson and Miller, 2003). The converted densities explain the observed free-air gravity anomaly for the part covered by our deployed instruments. We extended the profile further northeast over the marine part of the MAKRIS line (Fig. 1, inlay profile 4). From profile KM 127.5 northwards, we related the gross density model structure to the results of the MAKRIS line (Makris et al., 1999). However, we removed a large step of 10 km in Moho depth and replaced it by a more gradually deepening Moho, which closely follows the top of the layer of underplating in the MAKRIS line. The fit of observed and calculated gravity data reasonably well supports the interpretation of a thickening continental crust towards the northeast.

## 5 Discussion

### 5.1 Nature of the crust

420 The seismic velocity model along our refraction profile (Fig. 4a) shows no common features of oceanic crust. Oceanic crust typically consists of a high-velocity gradient in Layer 2 and a lower velocity gradient in Layer 3 (e.g. White et al., 1992; Grevenmeyer et al., 2018; Christeson et al., 2019). The absolute seismic velocities are highly variable, however, for a gabbroic crust, velocities are typically between 6.7 km/s and 7.2 km/s (Grevenmeyer et al., 2018; Christeson et al., 2019). The histogram in Figure 4f shows a gap for this range of velocities suggesting the lack of a thick gabbroic layer, and thus, the lack of typical  
425 oceanic crust~~that no typical oceanic crust and no thick layer of gabbroic rocks is present along the profile~~. In any case, the lack of seismic velocities expected for oceanic crust does not support the occurrence of larger units of oceanic crust as observed in the Central Tyrrhenian Sea (Prada et al., 2014).

Continental crust is characterised by a low seismic velocity gradient throughout the crystalline crustal layers and shows typical velocities of ~5.8 to ~6.6 km/s (Christensen and Mooney, 1995). We observe this velocity range in the northern half of the  
430 profile, starting from profile KM 70, at a depth of 10 km to 13 km (marked with “Y” in Fig. 4a). The observed seismic velocities provide only two possible interpretations: (1) hyper-extended continental crust or (2) a laterally isolated magmatic intrusion within the sedimentary units feeding the volcanic extrusion observed in the MCS and Parasound data (Fig. 3c). Based on the gravity model (Fig. 5), we favour the first scenario of extremely thinned continental crust, which is decreasing in thickness towards the SW and may even lead to exhumed mantle during the rifting phase in the south.

435 The velocity model for the southern half along our refraction profile is well constrained, however, the lower part (9 km to 11 km depth), above the Moho, shows higher uncertainty compared to the shallow, sedimentary units. The depth of the crust-mantle boundary is well constrained with an uncertainty range of  $\pm 0.25$  km along the southern profile half in contrast to  $\pm 0.75$  km along the northern profile half (Fig. 4b). We observe seismic velocities  $>5.5$  km/s that we interpret as fast syn-rift sediments due to a missing crystalline basement reflection. Alternatively, the change from sediments to the crystalline  
440 basement might not be characterised by a high impedance contrast, and thus, not imaged in our refraction seismic data as a strong in amplitude reflection event (compare to Fig. 2) and is expressed in a higher uncertainty of the determined CB at the northern profile end (Fig. 4). The MCS line MA24 (Rollet et al., 2002) (Fig. 1, inlay profile 6) records the acoustic basement at ~6.5 s two-way traveltime (stwt), while the seafloor occurs at 3.6 stwt (~2.7 km below sea surface). By means of a simple time to depth conversion using an average seismic velocity of 4.2 km/s, we estimate a minimum sedimentary thickness of  
445 ~6.1 km (2.9 stwt), resulting in an acoustic basement depth of ~8.8 km, as a most shallow approximation (drawn as a red dotted line in figure 4a). This line roughly fits the 5.5 km/s isoline accounting for a standard deviation of 0.2 km/s (Fig. 4b). For the southern half of our profile, this leaves a maximum continental crustal thickness of 2-2.5 km, thickening northwards.

Based on the refraction seismic data along our profile (southern half) we are not able to distinguish between sediments with high seismic velocities and extremely thinned continental crust. However, we can give a minimum and maximum continental  
450 crustal thickness, ranging from 0 km to 2.5 km. Based on the velocity model (Fig. 4a) it is not possible to distinguish whether

the crystalline basement is upper, middle, or lower continental crust. ~~The thickening of the continental crust A continental crustal thickening~~ towards the north-east is as well supported by the ~~modelling of the free air anomaly-gravity modelling~~ (Fig. 5). Additionally, a thickening crustal layer supports the interpretation as continental crust, since we would expect the COT to be manifested in an abrupt change from oceanic to continental crust or ~~oceanic crust~~ to gradually thin out towards the NE, towards the rotational pole (Rosenbaum et al., 2002), depending on the position of the profile with respect to the proposed spreading axis.

An expanding spread profile (ESP) (Le Douaran et al., 1984; Contrucci et al., 2001) crosses the northern end of our profile (Fig. 1, inlay profile 5). There the crust-mantle boundary was defined at a depth of 13-15 km while the acoustic basement was observed at ~9 km depth. Contrucci et al. (2001) retrieved crustal velocities of 6.3 - 6.9 km/s for the basin centre, which in general, is in good agreement with our results. Based on MCS data (LISA01) (Contrucci et al., 2001) with an observed major step in the basement on the Ligurian margin, they interpreted the central basin as an oceanic domain. On the Corsica margin, this major step was not observed; however, magnetic anomalies were used to constrain the interpretation. The MCS data resolve only the sedimentary portion of the crust and yield no information on the internal structures of the crystalline basement itself. Thus, a different explanation for the major step in the basement near the Ligurian margin could be that upper-crustal blocks sit on top of continental mantle similar to the Galicia margin (Nagel and Buck, 2004). Our profile only provides information on the basin centre where the absolute velocities of Le Douaran et al. (1984) and Contrucci et al. (2001) fit continental crust velocities ~~very well~~ and support our interpretation ~~for the northern end of the profile with here we observe of~~ mantle material beneath thinned continental crust ~~observations~~.

## 470 **5.2 Low degree of mantle serpentinisation**

~~Seismic velocities of ~7.3 km/s and higher are too fast for magmatic crust (Grevemeyer et al., 2018; Christeson et al., 2019).~~ Seismic velocities of unaltered mantle are >7.8 km/s (Carlson and Miller, 2003; Grevemeyer et al., 2018). Based on the seismic P-wave velocities, we interpret the uppermost mantle to be serpentinised at a low grade, which is supported by the Pn phases that are weak in amplitude at the southern stations (Fig. 2). P-wave velocities of ~7.5 km/s in the south-western half of the profile (Fig. 4a) are in-line with up to ~20% serpentinisation (Carlson and Miller, 2003). From OBS204 to OBS207, the PmP phase is extremely high in amplitude and unusually clearly visible over a wide distance of up to 20 km (in ~10 km to ~30 km offset to the station). This area (profile KM 40-KM 60) is marked by  $V_p > 7.8$  km/s directly underneath the basement ("Z" in Fig. 4a), possibly an area of unaltered mantle due to a left over (and possibly rotated) block of continental crust as observed in other magma-poor passive margins (Bayracki et al., 2016) ~~or a result of a mafic intrusion~~. The fact that the mantle is only partly serpentinised suggests ~~that either continental crustal blocks are overlying the mantle, and thus it was not fully exposed during rifting or that syn-rift sediments (nowadays showing high P-wave velocities) may have been directly deposited on top of the mantle preventing its full serpentinisation (Perez-Gussinyé et al., 2013) that syn-rift sediments (nowadays showing high P-wave velocities) may have been directly deposited on top of the mantle or brittle continental crust (Pérez-Gussinyé, 2013).~~

485 Thus, structurally, the Ligurian Sea is mimicking the Atlantic non-volcanic passive margins of Iberia (Minshull et al., 2014) and Goban Spur (Bullock and Minshull, 2005). However, the fast mantle in the Ligurian Sea would support a much lower degree of serpentinisation when compared to these two regions with a mantle serpentinisation of 100% at the top basement and less than <25% in 5-7 km depth at the Goban Spur and >75% at the top basement and <25% in 2 km depth at the Iberia margin.

490 In comparison to the Central Tyrrhenian Sea, where exhumed mantle is inferred~~Tyrrhenian Sea~~, the P-wave velocities of the upper mantle in the Ligurian Sea are high-faster (Fig. 4a). P-wave velocities of the upper mantle in the ~~Tyrrhenian Sea Domain #3~~ Magnaghi and Vavilov basins in the Tyrrhenian Sea (i.e. Domain #3 in Prada et al., 2014) are significantly lower with 4.5 km/s at the top of the mantle (Prada et al., 2016). We observe significant differences between both basins: (1) Along the southern half of our profile we observe strong PmP reflections indicating a high-velocity contrast at the crust-mantle boundary, while in Domain #3 in the Tyrrhenian Sea PmP reflections are absent. ~~(2) The Ligurian Basin has a thick sedimentary cover of ~6-8 km, while the Tyrrhenian Sea Domain #3 shows a sedimentary cover of ~1-2 km (Prada et al., 2014). Further, (3) the Ligurian Basin has a thick sedimentary cover of ~6-8 km, while the Tyrrhenian Sea Domain #3 shows a sedimentary cover of ~1-2 km (Prada et al., 2014) was stretched ~150 km during the ~16 million year opening phase, while the Tyrrhenian Sea was stretched ~300 km within 9-10 million years. Although the extension in the Ligurian Basin lasted longer and occurred earlier, the sedimentation rate is significantly higher compared to the Tyrrhenian Sea. Syn-rift sedimentation was recorded in MCS data (Fig. 1, inlay profile 3) in the Gulf of Lion (Jolivet et al., 2015).~~ (2) The Ligurian Basin was stretched ~150 km during the ~16 million year opening phase, while the Tyrrhenian Sea was stretched ~170 km within ~5 million years prior the onset of oceanic spreading (Faccenna et al., 2001; Prada et al., 2014). Thus, the stretching rate in the Tyrrhenian Sea was higher compared to the Ligurian Sea. Further, (3) the Ligurian Basin has a thick sedimentary cover of ~6-8 km, while the Tyrrhenian Sea Domain #3 shows a sedimentary cover of ~1-2 km (Prada et al., 2014). Syn-rift sedimentation was recorded in MCS data (Fig. 1, inlay profile 3) in the Gulf of Lion (Jolivet et al., 2015). The proximity of the two basins to the continental margin during their formation might result in a different syn-rift sedimentation rate that possibly was higher in the Ligurian Sea compared to the Tyrrhenian Sea. Sediments are known to reduce the permeability and thus, the amount of water that reaches the mantle rocks, necessary for serpentinisation (Ruepke et al., 2013). Two other factors can play a role for the degree of mantle serpentinisation in the Ligurian Basin: Ruepke et al. (2013) show in thermo-tectono-stratigraphic basin models the effects of sedimentary blanketing and low stretching factors on serpentinisation. Hence our seismic velocity model (Fig. 4a) can be well explained if mantle rocks have been partially exhumed from continental crust, without being directly exposed to sea water due to syn-rift sedimentation. ~~However, a~~ Also the interpretation of extremely thinned brittle continental crust requires syn-rift sedimentation since the stretching might open fluid pathways through the crust down to the mantle and would lead to a high degree of mantle serpentinisation (Nagel and Buck, 2004).

515



### 5.3 Continent-ocean transition and magmatic intrusions

The MCS line MA24 (Rollet et al., 2002) was shot along an ESP profile consisting of four measurements with a spacing of ~35 km (Le Douaran et al., 1984). The two transects are crossing our profile at the southern end (Fig. 1, inlay profile 6). The MCS data resolve sedimentary units, while the seismic velocities retrieved along the ESP profile show no absolute seismic velocities similar to oceanic crust. Both transects do not map a spreading axis. Further west along the Ligurian margin, a multichannel seismic study (Jolivet et al., 2015) and a wide-angle refraction seismic study (Gailler et al., 2009) of the Ligurian margin (Fig. 1, inlay profiles 2a and 3), in the Gulf of Lion, show a wide continent-ocean transition zone. The travel time tomography model along the OBS profiles (Gailler et al., 2009) images a succession of three domains: (1) continental, (2) transitional, and (3) oceanic towards the basin centre, following the zonation of Rollet et al. (2002). The same succession was found for both continental margins, though, the Corsica margin's transitional zone is much narrower. The transitional domain is interpreted to consist of a mixture of continental crust, exhumed mantle, and magmatic intrusions (Gailler et al., 2009; Rollet et al., 2002). In contrast, Jolivet et al. (2015) interpret the transitional zone as exhumed lower continental crust overlying the continental mantle which is in the distal part exhumed and serpentinitised. [The nature of the COT along the Gulf of Lion is still debated](#) ~~The exhumation of lower continental crust in the Gulf of Lion is still debated~~. For example, numerical modelling of continental rifting at the magma-poor Galicia margin showed that the lower crust is scarcely preserved or absent in the continental tip (Nagel and Buck, 2004). Our velocity model at the base of the continental crust is not well enough resolved (Fig. 4b) to distinguish between upper and lower continental crust, but we emphasize again, that we can exclude oceanic crust based on the seismic velocity structure (Fig. 4f) and the results of gravity modelling (Fig. 5) along our seismic profile. The oceanic domain on both conjugated margins in the Gulf of Lion (Fig. 1, inlay profile 2a) and offshore Sardinia (Fig. 1, inlay profile 2b) was [imaged in the travel-time tomographic approach with the typical pattern observed at mid-ocean ridges \(Gailler et al., 2009\), with a high velocity gradient in the upper oceanic crust and a low velocity gradient in the lower crust, interpreted on the basis of a 2D P-wave model derived from travel time tomography as an anomalously thin oceanic crust \(4–5 km\) with the typical two-layer gradients clearly characteristic of oceanic Layers 2 and 3 \(Gailler et al., 2009\)](#).

The extension process in the Ligurian Basin stopped roughly 16 Ma and was replaced by the extension and opening of the Tyrrhenian Sea as the Apennines-Calabrian subduction zone continued to roll back. The magnetic data (Bayer et al., 1973; Cella et al., 2008) in both basins show a similar anomaly distribution with discontinuous, partially isolated anomalies. Prada et al. (2014) analysed a seismic refraction profile crossing the Tyrrhenian Sea from Sardinia to Italy mainland. Similar to the Ligurian Basin, the western margin is more elongated than the eastern margin. They divide the analysed profile into 3 different domains from Sardinia to the central basin: In domain #1 continental crust thins from 22 km to 13 km over a distance of 80 km. Domain #2 is interpreted as magmatic back-arc crust with blocks of continental crust and stretches over a distance of ~80 km on the Corsican side of the basin (Prada et al., 2014). The change from continental to magmatic crust is marked by an abrupt increase of seismic velocities to >7 km/s in the lower crust, similar to the observation of Gailler et al. (2009) on the Ligurian Basin side of Sardinia. Prada et al. (2014) interpret the seismic velocities, which are slightly lower than found in 0-7

Ma old-oceanic crust, to be a result of back-arc spreading close to the active volcanic arc. Domain #3 is interpreted to be  
550 composed of serpentinised mantle to a depth of 5-6 km with basaltic intrusions and shows a width of ~140 km. Prada et al.,  
(2014) suggest that rifting in the Central Tyrrhenian Basin started with extension of continental crust, continued with back-arc  
spreading, followed by mantle exhumation. Later, the area underwent magmatic episodes with magmatic intrusions into the  
sedimentary layer or cropping out, forming volcanoes. These volcanoes and magmatic intrusions could be related to magnetic  
anomalies (Prada et al., 2016). Using the Tyrrhenian Sea as an analogy, we suggest that many of the isolated magnetic  
555 anomalies in the Ligurian Sea are caused by magmatic intrusions or extrusions manifested as volcanic edifices (Median  
Seamount, Tristanites Massif, Monte Doria; see Fig. 1) (Rollet et al., 2002), rather than related to a spreading axis, which was  
indeed not mapped in MCS data so far. However, in MCS data, intrusions of volcanic sills into younger sediments were  
observed (Finetti et al., 2005). At the Monte Doria Seamount, 11-12 Ma old basalts with a calc-alkaline signature were sampled  
by dredges and submersible dives (Rollet et al., 2002; Réhault et al., 2012). ~~The age is~~ clearly indicating post-rift magmatism.  
560 Further, volcanism related to the slab roll-back of the Apennines-Calabrian subduction zone was observed at the Ligurian  
continental margin and dated to the initiation of the rifting phase (Rollet et al., 2002). Volcanism was as well associated with  
the end of the opening of the Ligurian Basin and related to the trench retreat of the Apennines-Calabrian subduction zone  
(Rollet et al., 2002). This implies that volcanism also occurred during the rifting phase and could add to the discontinuous  
magnetic anomalies.

565

#### 5.4 Opening of the Ligurian Basin

The opening of the Ligurian Basin in a back-arc position during late Oligocene and early Miocene was driven by the south-  
east retreating Apennines-Calabria-Maghrebides subduction zone (e.g. Doglioni et al., 1997; Faccenna et al., 1997; Réhault et  
al., 1984; Carminati et al., 1998). The shift of active expansion from the Ligurian Basin to the Tyrrhenian Sea is considered a  
570 result of the Alpine collision that locked the Corsica-Sardinia drift towards the east and slab break-offs along the northern  
African margin and along the Apennines (Carminati et al., 1998). Thus, the opening of the Ligurian Basin was limited in time  
and space. Two different conceptual scenarios of rifting could explain our observations: (1) Rifting causing continental crust  
to thin until continental lower crust and mantle are exhumed and afterwards oceanic spreading is induced as observed in the  
Gulf of Lion (Gailler et al., 2009; Jolivet et al., 2015). (2) Rifting causing continental crust to thin until back-arc spreading is  
575 initiated and the continuation of extension leads to exhumation of mantle with magmatic intrusions (Prada et al., 2016).

Depending on the scenario, our profile is situated in the Ligurian transitional domain #2 or in the Tyrrhenian domain #3.  
Rifting scenario (2) would imply that well developed oceanic back-arc crust should occur southeast and northwest of the  
profile. The transect reaches into the area of a 3D seismic study of the Ligurian margin offshore Sanremo (Dessa et al., 2011).  
The authors state that they were surprised not to see a distinct change in the velocity field at the COT. Dessa et al. (2011) could  
580 not find clear evidence for a kind of back-arc crust as shown by Prada et al. (2014) or Gailler et al. (2009). However, continental  
crustal thinning is well imaged. Considering these aspects, we rather favour rifting scenario (1) which is also supported by the

conceptual model described by Decarlis et al. (2017) for the evolution of magma-poor rifted margins. The model includes three phases of extension: (1) An initial stretching phase forming widely distributed half-grabens in the upper crust. Afterwards (2) a thinning phase leads to hyper-extended crust and is followed by (3) an exhumation phase during which subcontinental mantle rocks were exhumed. [Yet, it is debated if we observe the latter phase in the central Ligurian Basin.](#)

Furthermore, the Ligurian Basin width in our study area (70 - 120 km) is much narrower than further south (~200 km) where domain #2 is cumulating to ~100 km in length for both conjugated margins together, which would entirely fill the basin in our study area, leaving little or no space for oceanic spreading. This is i.e. supported by petrological and geophysical observations at the West Iberia margin, that suggest that a COT zone can reach a width of up to 200 km (Pérez-Gussinyé, 2013).

Additionally, the opening rate becomes lower towards the north and the amount of stretching becomes less, which is probably caused by the anti-clockwise rotation of the Corsica-Sardinia block. Stretching of the crust as a result of the opening of the basin becomes less intense towards the north and thus controls the NE termination of the ultra-thin continental crust. Further, the extension of the basin decreases towards the north and assuming oceanic crust to be present, the crust should become less thick towards the proposed ridge axis tip. However, our seismic data and gravity data indicate a gradual thickening of the crystalline crust, at least a gradual deepening of the mantle, indicating thickening continental crust northwards. This is as well supported by the magnetic data (Bayer et al., 1973), which do not show the typical oceanic crust pattern of magnetisation stripes, but rather a lateral patchy pattern of magmatic domains. [supporting again the lack of oceanic spreading during the formation of the Ligurian basin in the Oligocene-Miocene. This could imply that oceanic spreading was not initiated during the Oligocene-Miocene extension in the northern Ligurian Basin, along our seismic line.](#) Continuing further north [of our seismic line](#), extension led to ~~extreme thinning~~ of continental crust, but ~~lasted not long enough to~~ [the amount of extension was too small to extremely thin out the continental crust and](#) exhume mantle.

## 6 Conclusion

The P-wave velocity model determined in this study images the uppermost lithospheric structure of the central Ligurian Basin. Syn- and post-rift sediments of ~6-8 km thickness filled the basin during and after the 15 Ma long lasting opening phase. Based on the image of the seismic velocity distribution along the southern half of the profile it remains enigmatic if the mantle is overlain directly by [syn-rift](#) sediments or by extremely thinned continental crust of up to 2.5 km. The degree of mantle serpentinisation with up to 20% is low. The northern half of the profile indicates a northward thickening of continental crust and a deepening crust-mantle boundary from 11 km to 13 km. Based on the retrieved velocity distribution, gravity modelling and results of surrounding studies, we conclude that the extension of the Ligurian Basin led to:

(1) Extended and very thin continental crust or exhumed, partially serpentinised mantle. [It remains enigmatic if the mantle is overlain directly by sediments or by extremely thinned continental crust of up to 2.5 km thickness.](#)

(2) Continental crustal thinning from north to south related to the increase of extension with increasing distance from the rotation pole of the anti-clockwise rotation of the Corsica-Sardinia block.

615 Furthermore, our study documents that:

(3) Seafloor spreading and formation of mantle-derived oceanic crust was not initiated during the extension of the Ligurian Basin.

Thus, we conclude that the oceanic domain does not extend as far north as previously stated and that the transition from the continental domain and the real oceanic domain with a potential spreading axis is situated south or south-west, ~~however, nearby~~  
620 [of](#) our seismic line.

*Data availability.* Seismic data are available on request from the first or second author and are available via the German marine data archive PANGAEA at <https://doi.org/10.1594/PANGAEA.910561>.

625 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* This project is funded by Deutsche Forschungsgemeinschaft (DFG), grant number KO\_2961/6-1. We thank captain and crew of RV Maria S. Merian cruise MSM71 for their support during the OBS work. We thank the MSM71 cruise participants for their effort. The LOBSTER project comprises the offshore component of the AlpArray seismic network  
630 (Hetényi et al., 2018) and is part of the German priority program SPP2017 4D-MB. Generic Mapping Tools (Wessel & Smith, 1998) were used to produce the figures. We thank the reviewers L. Jolivet, J-X. Dessa, and M. Prada for their fruitful comments.

## References

- Advokaat, E. L., van Hinsbergen, D. J. J., Maffione, M., Langereis, C. G., Vissers, R. L. M., Cherchi, A., Schroeder, R., Madani, H. and Columbu, S.: Eocene rotation of Sardinia, and the paleogeography of the western Mediterranean region, Earth  
635 Planet. Sci. Lett., 401, 183–195, doi:10.1016/j.epsl.2014.06.012, 2014.
- Bayrakci, G., Minshull, T.A., Sawyer, D.S., Reston, T. J., Klaeschen, D., Papenberg, C., Ranero, C., Bull, J. M., Davy, R. G., Shillington, D. J., Perez-Gussinye, M., and Morgan, J. K.: Fault-controlled hydration of the upper mantle during continental rifting, Nature Geoscience , doi: 10.1038/NGEO2671, 2016.
- Bayer, R., Le Mouel, J. L. and Le Pichon, X.: Magnetic anomaly pattern in the western mediterranean, Earth Planet. Sci. Lett.,  
640 19(2), 168–176, doi:10.1016/0012-821X(73)90111-8, 1973.
- Béthoux, N., Tric, E., Chery, J. and Beslier, M.-O.: Why is the Ligurian Basin (Mediterranean Sea) seismogenic? Thermomechanical modeling of a reactivated passive margin: MODELING OF THE LIGURIAN BASIN, Tectonics, 27(5), doi:10.1029/2007TC002232, 2008.

- Bonatti, E., Seyler, M., Channell, J., Giraudeau, J. and Mascle, G., Eds.: Peridotites drilled from the Tyrrhenian Sea, ODP Leg 107, Ocean Drilling Program 107, 1990.
- Bullock, A. and Minshull, T.: From continental extension to seafloor spreading: Crustal structure of the Goban Spur rifted margin, southwest of the UK, *Geophys. J. Int.*, 163, 527–546, doi: 10.1111/j.1365-246X.2005.02726.x, 2005.
- Burrus, J.: Contribution to a geodynamic synthesis of the Provençal Basin (NorthWestern Mediterranean), *Mar. Geol.*, 55(3–4), 247–269, [https://doi.org/10.1016/0025-3227\(84\)90071-9](https://doi.org/10.1016/0025-3227(84)90071-9), 1984.
- Carlson, R. L. and Miller, D. J.: Mantle wedge water contents estimated from seismic velocities in partially serpentinized peridotites: MANTLE WEDGE WATER CONTENTS, *Geophys. Res. Lett.*, 30(5), doi:10.1029/2002GL016600, 2003.
- Carminati et al., 1998, Carminati, E., Wortel, M.J.R., Meijer, P.T., and Sabadini, R.: The two-stage opening of the western-central mediterranean basins: a forward modeling test to a new evolutionary model, *Earth Planet. Sci. Lett.*, 160, pp. 667-679, doi:10.1016/S0012-821X(98)00118-6, 1998.
- Cella, F., Fedi, M., Florio, G., Paoletti, V. and Rapolla, A.: A review of the gravity and magnetic studies in the Tyrrhenian Basin and its volcanic districts, *Annals of Geophysics*, 51, doi:10.4401/ag-3035, 2008.
- Chamot-Rooke, N., Jestin, F. and Gaulier, J.-M.: Constraints on Moho Depth and Crustal Thickness in the Liguro-Provençal Basin from a 3d Gravity Inversion: Geodynamic Implications, *Rev. Inst. FRANCAIS Pet.*, 52, 557–583, doi:10.2516/ogst:1997060, 1997.
- Christensen, N. I. and Mooney, W. D.: Seismic velocity structure and composition of the continental crust: A global view, *J. Geophys. Res. Solid Earth*, 100(B6), 9761–9788, doi:10.1029/95JB00259, 1995.
- Christeson, G. L., Goff, J. A. and Reece, R. S.: Synthesis of Oceanic Crustal Structure From Two-Dimensional Seismic Profiles, *Rev. Geophys.*, 57(2), 504–529, doi:10.1029/2019RG000641, 2019.
- Contrucci, I., Nercessian, A., Béthoux, N., Mauffret, A. and Pascal, G.: A Ligurian (Western Mediterranean Sea) geophysical transect revisited, *Geophys. J. Int.*, 74–97, 2001.
- Decarlis, A., Beltrando, M., Manatschal, G., Ferrando, S. and Carosi, R.: Architecture of the Distal Piedmont-Ligurian Rifted Margin in NW Italy: Hints for a Flip of the Rift System Polarity, *Tectonics*, 36(11), 2388–2406, doi:10.1002/2017TC004561, 2017.
- Dercourt, J., Zonenshain, L. P., Ricou, L.-E., Kazmin, V. G., Le Pichon, X., Knipper, A. L., Grandjacquet, C., Sbertshikov, I. M., Geysant, J., Lepvrier, C., Pechersky, D. H., Boulin, J., Sibuet, J.-C., Savostin, L. A., Sorokhtin, O., Westphal, M., Bazhenov, M. L., Lauer, J. P. and Biju-Duval, B.: Geological evolution of the tethys belt from the atlantic to the pamirs since the LIAS, *Tectonophysics*, 123(1–4), 241–315, doi:10.1016/0040-1951(86)90199-X, 1986.
- Dessa, J.-X., Simon, S., Lelievre, M., Beslier, M.-O., Deschamps, A., Bethoux, N., Solarino, S., Sage, F., Eva, E., Ferretti, G., Bellier, O. and Eva, C.: The GROSMarin experiment: three dimensional crustal structure of the North Ligurian margin from refraction tomography and preliminary analysis of microseismic measurements, *Bull. Soc. Geol. Fr.*, 182(4), 305–321, doi:10.2113/gssgfbull.182.4.305, 2011.


- Doglioni, C., Gueguen, E., Sàbat, F. and Fernandez, M.: The Western Mediterranean extensional basins and the Alpine orogen, *Terra Nova*, 9(3), 109–112, doi:10.1046/j.1365-3121.1997.d01-18.x, 1997.
- Faccenna, C., Mattei, M., Funiciello, R. and Jolivet, L.: Styles of back-arc extension in the Central Mediterranean, *Terra Nova*, 680 9(3), 126–130, doi:10.1046/j.1365-3121.1997.d01-12.x, 1997.
- Faccenna, C., Funiciello, F., Giardini, D. and Lucente, P.: Episodic back-arc extension during restricted mantle convection in the Central Mediterranean, *Earth Planet. Sci. Lett.*, 187(1–2), 105–116, doi:10.1016/S0012-821X(01)00280-1, 2001.
- Fernàndez, M., Foucher, J. P. and Jurado, M. J.: Evidence for the multi-stage formation of the south-western Valencia Trough, *Mar. Pet. Geol.*, 12(1), 101–109, doi:10.1016/0264-8172(95)90390-6, 1995.
- 685 Finetti, I. R., Boccaletti, M., Bonini, M., Ben, A., Pipan, M., Prizzon, A. and Sani, F.: Lithospheric Tectono-Stratigraphic Setting of the Ligurian Sea–Northern Apennines–Adriatic Foreland from Integrated CROP Seismic Data, in *Deep Seismic Exploration of the Central Mediterranean and Italy, CROP PROJECT*, pp. 119–158., 2005.
- Fujie, G., Kasahara, J., Murase, K., Mochizuki, K. and Kaneda, Y.: Interactive analysis tools for the wide-angle seismic data for crustal structure study (Technical Report), *Explor. Geophys. - EXPLOR GEOPHYS*, 39, doi:10.1071/EG08006, 2008.
- 690 Gailler, A., Klingelhoefer, F., Olivet, J.-L. and Aslanian, D.: Crustal structure of a young margin pair: New results across the Liguro–Provencal Basin from wide-angle seismic tomography, *Earth Planet. Sci. Lett.*, 286(1–2), 333–345, doi:10.1016/j.epsl.2009.07.001, 2009.
- Gardner, G. H. F., Gardner, L. W. and Gregory, A. R.: FORMATION VELOCITY AND DENSITY—THE DIAGNOSTIC BASICS FOR STRATIGRAPHIC TRAPS, *GEOPHYSICS*, 39(6), 770–780, doi:10.1190/1.1440465, 1974.
- 695 Grevemeyer, I., Ranero, C. R. and Ivandic, M.: Structure of oceanic crust and serpentinization at subduction trenches, *Geosphere*, 14(2), 395–418, doi:10.1130/GES01537.1, 2018.
- Gueguen, E., Doglioni, C. and Fernandez, M.: On the post-25 Ma geodynamic evolution of the western Mediterranean, *Tectonophysics*, 298(1–3), 259–269, doi:10.1016/S0040-1951(98)00189-9, 1998.
- Guerin, G., Rivet, D., Deschamps, A., Larroque, C., Mordret, A., Dessa, J.-X. and Martin, X.: High resolution ambient noise 700 tomography of the South-Western Alps and the Ligurian margin, *Geophys. J. Int.*, doi:10.1093/gji/ggz477, 2019.
- Handy, M. R., M. Schmid, S., Bousquet, R., Kissling, E. and Bernoulli, D.: Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological–geophysical record of spreading and subduction in the Alps, *Earth-Sci. Rev.*, 102(3–4), 121–158, doi:10.1016/j.earscirev.2010.06.002, 2010.
- Hetényi, G., Molinari, I., Clinton, J., Bokelmann, G., Bondár, I., Crawford, W. C., Dessa, J.-X., Doubre, C., Friederich, W., 705 Fuchs, F., Giardini, D., Gráczner, Z., Handy, M. R., Herak, M., Jia, Y., Kissling, E., Kopp, H., Korn, M., Margheriti, L., Meier, T., Mucciarelli, M., Paul, A., Pesaresi, D., Piromallo, C., Plenefisch, T., Plomerová, J., Ritter, J., Rümpler, G., Šipka, V., Spallarossa, D., Thomas, C., Tilmann, F., Wassermann, J., Weber, M., Wéber, Z., Wesztergom, V., Živčić, M., Abreu, R., Allegretti, I., Apoloner, M.-T., Aubert, C., Besançon, S., Bès de Berc, M., Brunel, D., Capello, M., Čarman, M., Cavaliere, A., Chèze, J., Chiarabba, C., Cougoulat, G., Cristiano, L., Czifra, T., D’Alema, E., Danesi, S., Daniel, R., Dannowski, A., 710 Dasović, I., Deschamps, A., Egdorf, S., Fiket, T., Fischer, K., Funke, S., Govoni, A., Gröschl, G., Heimers, S., Heit, B., Herak,

- D., Huber, J., Jarić, D., Jedlička, P., Jund, H., Kligen, S., Klotz, B., Kolínský, P., Kotek, J., Kühne, L., Kuk, K., Lange, D., Loos, J., Lovati, S., Malengros, D., Maron, C., Martin, X., Massa, M., Mazzarini, F., Métral, L., Moretti, M., Munzarová, H., Nardi, A., Pahor, J., Péquegnat, C., Petersen, F., Piccinini, D., Pondrelli, S., Prevolnik, S., Racine, R., Régnier, M., Reiss, M., Salimbeni, S., et al.: The AlpArray Seismic Network: A Large-Scale European Experiment to Image the Alpine Orogen, *Surv. Geophys.*, 39(5), 1009–1033, doi:10.1007/s10712-018-9472-4, 2018.
- 715 Jolivet, L. and Faccenna, C.: Mediterranean extension and the Africa-Eurasia collision, *Tectonics*, 19(6), 1095–1106, doi:10.1029/2000TC900018, 2000.
- Jolivet, L., Gorini, C., Smit, J. and Leroy, S.: Continental breakup and the dynamics of rifting in back-arc basins: The Gulf of Lion margin: Backarc rift and lower crust extraction, *Tectonics*, 34(4), 662–679, doi:10.1002/2014TC003570, 2015.
- 720 Kopp, H., Lange, D., Thorwart, M., Paul, A., Dannowski, A., Petersen, F., Aubert, C., Beeck, F., Beniest, A., Besançon, S., Brotzer, A., Caielli, G., Crawford, W., Deen, M., Lehmann, C., Marquardt, K., Neckel, M., Papanagnou, L., Schramm, B., Schröder, P., Steffen, K.-P., Wolf, F., Xia, Y.: RV MARIA S. MERIAN Fahrtbericht / Cruise Report MSM71 LOBSTER: Ligurian Ocean Bottom Seismology and Tectonics Research, Las Palmas (Spain) – Heraklion (Greece) 07.02.-27.02.2018. GEOMAR Report, N. Ser. 041. GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Kiel, Germany, 47 pp, doi:10.3289/GEOMAR\_REP\_NS\_41\_2018, 2018.
- 725 Korenaga, J., Holbrook, W. S., Kent, G. M., Kelemen, P. B., Detrick, R. S., Larsen, H.-C., Hopper, J. R. and Dahl-Jensen, T.: Crustal structure of the southeast Greenland margin from joint refraction and reflection seismic tomography, *J. Geophys. Res. Solid Earth*, 105(B9), 21591–21614, doi:10.1029/2000JB900188, 2000.
- Korenaga, J., Holbrook, W. S., Detrick, R. S. and Kelemen, P. B.: Gravity anomalies and crustal structure at the southeast Greenland margin, *J. Geophys. Res. Solid Earth*, 106(B5), 8853–8870, doi:10.1029/2000JB900416, 2001.
- 730 Le Douaran, S., Burrus, J. and Avedik, F.: Deep structure of the north-western Mediterranean Basin: Results of a two-ship seismic survey, *Mar. Geol.*, 55(3–4), 325–345, doi:10.1016/0025-3227(84)90075-6, 1984.
- Makris, J., Egloff, F., Nicolich, R. and Rihm, R.: Crustal structure from the Ligurian Sea to the Northern Apennines — a wide angle seismic transect, *Tectonophysics*, 301(3–4), 305–319, doi:10.1016/S0040-1951(98)00225-X, 1999.
- 735 Mauffret, A., Pascal, G., Maillard, A. and Gorini, C.: Tectonics and deep structure of the north-western Mediterranean Basin, *Mar. Pet. Geol.*, 12(6), 645–666, doi:10.1016/0264-8172(95)98090-R, 1995.
- Meléndez, A., Sallarès, V., Ranero, C.R., and Kormann, J.: Origin of water layer multiple phases with anomalously high amplitude in near-seafloor wide-angle seismic recordings, *Geophys. J. Int.*, 196, 243–252, doi:10.1093/gji/ggt391, 2014.
- 740 Minshull, T. A., Dean, S. M. and Whitmarsh, R. B.: The peridotite ridge province in the southern Iberia Abyssal Plain: Seismic constraints revisited, *J. Geophys. Res. Solid Earth*, 119(3), 1580–1598, doi:10.1002/2014JB011011, 2014.
- Moeller, S., Grevemeyer, I., Ranero, C. R., Berndt, C., Klaeschen, D., Sallares, V., Zitellini, N. and de Franco, R.: Early-stage rifting of the northern Tyrrhenian Sea Basin: Results from a combined wide-angle and multichannel seismic study: Rifting Tyrrhenian, *Geochem. Geophys. Geosystems*, 14(8), 3032–3052, doi:10.1002/ggge.20180, 2013.

- Montigny, R., Edel, J. B. and Thuizat, R.: Oligo-Miocene rotation of Sardinia: KAr ages and paleomagnetic data of Tertiary volcanics, *Earth Planet. Sci. Lett.*, 54(2), 261–271, doi:10.1016/0012-821X(81)90009-1, 1981.
- Nagel, T. J. and Buck, W. R.: Symmetric alternative to asymmetric rifting models, *Geology*, 32(11), 937, doi:10.1130/G20785.1, 2004.
- Pascal, G. P., Mauffret, A. and Patriat, P.: The ocean-continent boundary in the Gulf of Lion from analysis of expanding spread profiles and gravity modelling, *Geophys. J. Int.*, 113(3), 701–726, doi:10.1111/j.1365-246X.1993.tb04662.x, 1993.
- 745 Pérez-Gussinyé, M.: A tectonic model for hyperextension at magma-poor rifted margins: an example from the West Iberia–Newfoundland conjugate margins, *Geol. Soc. Lond. Spec. Publ.*, 369(1), 403–427, doi:10.1144/SP369.19, 2013.
- Prada, M., Sallares, V., Ranero, C. R., Vendrell, M. G., Grevenmeyer, I., Zitellini, N. and de Franco, R.: Seismic structure of the Central Tyrrhenian basin: Geophysical constraints on the nature of the main crustal domains: CRUSTAL AFFINITY OF THE TYRRHENIAN BASIN, *J. Geophys. Res. Solid Earth*, 119(1), 52–70, doi:10.1002/2013JB010527, 2014.
- 755 Prada, M., Ranero, C. R., Sallarès, V., Zitellini, N. and Grevenmeyer, I.: Mantle exhumation and sequence of magmatic events in the Magnaghi–Vavilov Basin (Central Tyrrhenian, Italy): New constraints from geological and geophysical observations, *Tectonophysics*, 689, 133–142, doi:10.1016/j.tecto.2016.01.041, 2016.
- Réhault, J.-P. and Bethoux, N.: Earthquake relocation in the Ligurian Sea (western Mediterranean): Geological interpretation, *Mar. Geol.*, 55(3), 429–445, doi:10.1016/0025-3227(84)90080-X, 1984.
- 760 Réhault, J.-P., Boillot, G. and Mauffret, A.: The Western Mediterranean Basin geological evolution, *Mar. Geol.*, 55(3), 447–477, doi:10.1016/0025-3227(84)90081-1, 1984.
- Réhault, J.-P., Honthaas, C., Guennoc, P., Bellon, H., Ruffet, G., Cotten, J., Sosson, M. and Maury, R. C.: Offshore Oligo-Miocene volcanic fields within the Corsica-Liguria Basin: Magmatic diversity and slab evolution in the western Mediterranean Sea, *J. Geodyn.*, 58, 73–95, doi:10.1016/j.jog.2012.02.003, 2012.
- 765 Roca, E. and Desegaulx, P.: Analysis of the geological evolution and vertical movements in the València Trough area, western Mediterranean, *Mar. Pet. Geol.*, 9(2), 167–185, doi:10.1016/0264-8172(92)90089-W, 1992.
- Rollet, N., Déverchère, J., Beslier, M.-O., Guennoc, P., Réhault, J.-P., Sosson, M. and Truffert, C.: Back arc extension, tectonic inheritance, and volcanism in the Ligurian Sea, Western Mediterranean: LIGURIAN SEA BACK ARC STRUCTURE AND EVOLUTION, *Tectonics*, 21(3), 6-1-6–23, doi:10.1029/2001TC900027, 2002.
- 770 Rosenbaum, G., Lister, G. and Duboz, C.: Reconstruction of the tectonic evolution of the Western Mediterranean since the Oligocene, *J. Virtual Explor.*, 8, 107–130, doi:10.3809/jvirtex.2002.00053, 2002.
- Ruepke, L.H., Schmid, D.W., Perez-Gussinye, M., and Hartz, E.: Interrelation between rifting, faulting, sedimentation, and mantle serpentinization during continental margin formation—including examples from the Norwegian Sea, *Geochem. Geophys. Geosyst.* 14, doi: 10.1002/ggge.20268, 2013.
- 775 Ryan, W.B.F., Carbotte, S., Coplan, J.O., O’Hara, S., Melkonian, A., Arko, R., Weissel, A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., Zemsky, R.: Global Multi-Resolution Topography synthesis, *Geochem. Geophys. Geosyst.*, 10, doi:10.1029/2008GC002332, 2009.




- Sandwell, D. T., Müller, R. D., Smith, W. H. F., Garcia, E. and Francis, R.: New global marine gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure, *Science*, 346(6205), 65, doi:10.1126/science.1258213, 2014.
- 780 Speranza, F., Villa, I. M., Sagnotti, L., Florindo, F., Cosentino, D., Cipollari, P. and Mattei, M.: Age of the Corsica–Sardinia rotation and Liguro–Provençal Basin spreading: new paleomagnetic and Ar/Ar evidence, *Tectonophysics*, 347(4), 231–251, doi:10.1016/S0040-1951(02)00031-8, 2002.
- Talwani, M., Worzel, J. L. and Landisman, M.: Rapid gravity computations for two-dimensional bodies with application to the Mendocino submarine fracture zone, *J. Geophys. Res.* 1896-1977, 64(1), 49–59, doi:10.1029/JZ064i001p00049, 1959.
- 785 Vigliotti, L. and Langenheim, V. E.: When did Sardinia stop rotating? New palaeomagnetic results, *Terra Nova*, 7(4), 424–435, doi:10.1111/j.1365-3121.1995.tb00538.x, 1995.
- Wessel, P., & Smith, W. H. F.: New improved version of the generic mapping tools released. *Eos, Transactions American Geophysical Union*, 79(47), 579, doi: 10.1029/98EO00426, 1998.
- White, R. S., McKenzie, D. and O’Nions, R. K.: Oceanic crustal thickness from seismic measurements and rare earth element  
790 inversions, *J. Geophys. Res.*, 97(B13), 19683, doi:10.1029/92JB01749, 1992.
- Zelt, C. A.: Modelling strategies and model assessment for wide-angle seismic travelttime data, *Geophys. J. Int.*, 139(1), 183–204, doi:10.1046/j.1365-246X.1999.00934.x, 1999.




800 **Figure 1: Relief map (GMRT data, Ryan et al., 2009) of the study area with the seismic refraction line (thick red line) and OBH/OBS**  
**locations (yellow and orange triangles, respectively) that extend the MAKRIS profile (thick long black line) (Makris et al., 1999).**  
**Thin black polygons and grey shaded areas mark volcanic extrusion after Rollet et al. (2002). The different crustal domains (Rollet**  
**et al., 2002) are marked by thin orange and red lines and are labelled with: AOD - atypical oceanic domain, CCM - Corsica**  
**continental margin, LCM - Ligurian continental margin, TD - transitional domain. A thin yellow line marks the oceanic domain**  
805 **(ODG) after Gueguen et al. (1998). Thin red dashed lines show proposed fracture zones (Rollet et al., 2002). Short thick yellow bar**  
**perpendicular to the MAKRIS profile marks the continent-ocean transition (COT) (Makris et al., 1999). Green triangles and thin**  
**dotted black lines are the OBS locations and shot profiles of Dessa et al. (2011). The black and white inset in the lower left**  
**corner show previous seismic refraction and reflection lines: 1 - Prada et al. (2014), 2a/2b - Gailler et al. (2009), 3 - Jolivet et al. (2015), 4 -**  
**Makris et al. (1999), 5 - Contrucci et al. (2001), 6 - MA24 from Rollet et al. (2002).**

810




**Figure 2: (a) Stacked travel time picks of all 15 stations showing very similar arrivals suggesting an almost 1D structure along the profile. (b) Record section of station OBS205 (time reduced with a velocity of 8 km/s). The lower panel shows the calculated travel time picks from the final velocity model superimposed on the seismic data. (c) Record section and calculated travel times of station OBS209 (d) Record section and calculated travel times of station OBH212.**

815



820 **Figure 3:** (a) Multi-channel seismic data (MCS data) simultaneously shot with the refraction seismic line. The orange/yellow triangles mark the OBS/OBH positions along the profile. (b) Upper panel shows OBS205 from shot point 300 to shot point 600 with a reduction velocity of 4.5 km/s. The lower panel is a zoom into the MCS section (black box in a). The white lines show that the undulations in the sedimentary phases fit well with faults and salt diapirs. (c) Parasound sediment echo sounder data (orange box in 3a).



825 **Figure 4: (a) Final velocity model based on the averaged velocities from the plausible starting models. The red dashed line marks the crystalline basement (CB) as determined from the refraction seismic data. The red dotted line presents the CB inferred from**  
**MCS data (CB-MCS) crossing our profile (details given in the text). The solid red line marks the crust-mantle boundary (Moho);**  
**(b) Standard deviation for 17 inverted velocity models, covering the crustal part down to the Moho; (c) Starting models used in the inversion and to calculate the resulting average model in 4a. (d) Standard deviation for 14 inverted velocity models (starting models**  
**in the inlay), covering the upper mantle up to the Moho; (e) Ray coverage for the final average velocity model; (f) Histogram with the velocity distribution of the final average velocity model.**  
830




Figure 5: Density model (lower panel) converted from seismic velocities to densities (details given in the text) for the SW part covered by seismic stations. The profile was extended towards the NE using the marine part of the seismic refraction line of Makris et al. (1999). The upper panel shows the data fit between the satellite derived free-air anomaly data (Sandwell et al., 2014) (dashed blue) and the model response (solid red line).

835