

SED

Interactive comment

Interactive comment on "Evolution of a long-lived continental arc: a geochemical approach (Arequipa Batholith, Southern Peru)" by Sophie Demouy et al.

Sophie Demouy et al.

mathieu.benoit@get.omp.eu

Received and published: 16 May 2019

We have red carefully the anonymous referee #1 review by and we agree that the introduction and discussion should be improved according to the new insights coming mostly from the geochronologist community. You will find below a detailed answer to referee #1.

First, it exists a philosophical controversy enclose within his criticism. We are working at the batholith scale using whole rock (WR) chemical and isotopic data obtained on a large sample set collected regularly along this massif. Referee #1 argues that we don't take into account recent results coming from more and more precise geochronological

Printer-friendly version

Discussion paper



and isotopic measurements of zircons extracted in similar lithologies, using analytical apparatus (or methods, or skills) that we don't have. The philosophical question is: should we banish the naturalistic (or descriptive) approach at large scale of a single batholith in regards to what can be measured using the zircon/WR combined work on a less amount of samples? It is true to say that the latter allow to access cryptic chemical and geochronological variations, which are used to propose more and more complex models of magma petrogenesis, like those cited by referee #1. However, are they all directly transposable at the batholith spatial and temporal scale, like for Areguipa? The main criticism that we are facing is the same that the scientific community working on mantle processes had 15 years ago: the global, naturalistic approach should erase itself with regard to in-situ measurements of melt inclusions trapped in olivine extracted from mantle peridotites. At the end, the scientific community agrees to say that both may co-exist: as it is not possible to measure all melt inclusions coming from 100 samples collected in a 50x50 kms geological area, the same conclusions may apply to zircons. And even so, the data will provide so much detail that a de-magnifier may be necessary in order to transfer the signification of the observed heterogeneity at the batholith scale. Therefore, we maintain that the naturalistic approach of the isotopic signal enclosed in our 100 sample set is relevant, considering that it would have been a huge effort, maybe not scientifically relevant, to analyse all zircons in those samples. This is a first step, which will help scientist like referee #1 to define some targets where zirconology may be applied. This approach may be considered "outdated" by isotope geochemists (no fancy new isotopes) or "descriptive" by geochronologists (no

Secondly, and as the referee says, models and approaches become outdated. We have no doubt that modern, fractal, stochastic, ab-initio, thermodynamic modelization of processes happening 30 to 90 kms below the surface of the Earth will be

more details than all U/Pb ages available may do.

in-situ ID-TIMS TEA on zircon), we consider it as "naturalistic" (even if radiogenic isotopes/trace elements are measured) and we agree that it does not allow to "enhance our knowledge or arc magmatism" but it's a useful tool to explore an entire massif, in

SED

Interactive comment

Printer-friendly version

Discussion paper



SED

Interactive comment

Printer-friendly version

Discussion paper



genesis at zero ages.

Again, we do agree that our introduction and conclusion need a severe overhaul, but we still maintain that the descriptive approach is valuable and scientifically relevant, considering the number of samples processed for a single area.

Interactive comment on Solid Earth Discuss., https://doi.org/10.5194/se-2019-43, 2019.

SED

Interactive comment

Printer-friendly version

Discussion paper

