Responses

Dear Editor and reviewers,

Thank you very much for your comments and the positive assessment of our work. Your comments and suggestions have greatly improved our paper. We hope our revisions will meet with approval. The detailed responses to your comments are listed bellows:

Responses to Editor (Dr. Antonella Longo)

Comment 1: pag. 1, line 18 I expect that your investigation is carried out with passive seismic imagine, if so, it would be important to state that, considering the wide range of earthquakes that usually occurs in the region you are investigating.

Response 1: The seismic reflection data we use was collected during hydrocarbon exploration using an active source (i.e. an air-gun).

Comment 2: pag. 2, line 34 I think that a general reader would be interested to know the importance to give such attention to submarine eruptions. Just a quick summary of the potentiality of your study. Any implication for hazard? Just to ask.

Response 2: Because our study is focused on the 3D structure and growth of deepwater volcanoes, we have decided not to discuss, in too great a detail, the implications for hazards. By omitting this material we keep the abstract concise and to-the-point. However, we agree that briefly mentioning the potential geohazards associated with deep-water volcanoes could be of interest to the general reader. We have therefore added the following sentences (Lines 58-61) to the Introduction: "Without such information on the structure of deep-water volcanoes, we cannot assess how they grow or what hazard they may pose (e.g. tsunamis induced by flank collapse, seabed deformation and instability induced by highly explosive eruptions)".

Comment 3: pag. 3, line 60 I am curious to know why did you choose these two volcanoes and not other ones.

Response 3: Although several volcanoes are imaged in our study area, only two of them are physically isolated (i.e. the others are physically linked because their related lava flow fields merge). We can therefore confidently separate them, calculate the volume of eruptive material contained in their edifices and flanking lava flow fields, and thus calculate the ratio between material in the volcano edifice vs. the lava flow field. To clarify this, we have added: "*These two volcanoes are physically isolated and appear to have been fed by independent, sub-volcanic intrusive bodies (i.e. sills; see below); we can thus confidently characterize each individual volcano and its associated lava flows (Fig. 1b)." (Lines 66-68).*

Comment 4: pag. 6, line 115-116 Could you please explanate (I am not an expert on

the subject) the words\Bin spacing is 25 m". Furthermore, why the interval of interest of frequencies is 0-400 ms two-way time (twt) of ~40 Hz?

Response 4: Bin is a square and comprises two inlines and two crosslines in the 3D seismic reflection data. Here, 'Bin spacing is 25 m' means the inline and crossline are 25 m, respectively. To clarify this, we have changed this sentence to: "*The inline and crossline spacing are 25 m, respectively*" (Line 122).

The frequency of the seismic data is lies within a frequency band ranging from a few Hz to a few hundreds of Hz. The dominant frequency is the maximum frequency encountered in a frequency spectrum. The dominant (and peak) frequency typically decrease downwards from the seabed due to attentuation. Therefore, different depths are characterized by different dominant frequency spectrums (and dominant and peak frequencies). In this study, the dominant frequency in the interval of interest (i.e. 0–400 ms two-way time (twt)) is ~40 Hz.

Comment 5: Table 1 Unit of measure should be written as in Table 2. **Response 5**: We have revised the units of Table 1 to be the same as those in Table 2.

Comment 6: Table 2 How do you justify the computation of diameter from the area assuming it is a circle?

Response 6: The boundaries of lava flow apron are irregular and thus we cannot directly measure its diameter. In this study, we directly measured the lengths of the apron's irregular boundaries and calculated an approximate diameter by assuming it was a circle. We think this was the best estimate for estimating the crude diameter of an apron with irregular boundaries.

Comment 7: Fig. 2 (b) In caption D/T, DT, RHOB, and RC are mentioned but not indicated in Fig. 2(b).

Response 7: We have added these to Fig. 2a (Lines 704-705).

Comment 8: pag. 8, line 175 For a better understanding, I would replace the \-" with a \:", the same for the other Seismic Facies.

Response 8: We have revised this, in addition to those in Lines 181, 184 and 187.

Comment 9: pag. 10, line 206 Put the deg on 15: $15.0^{\circ} \pm 3.6^{\circ}$. **Response**: We have revised this in the text (Line 213).

Comment 10: pag. 11, line 226 Is it possible to assign an error to the $\sim 14 \text{ km}^2$ area? **Response 10**: It is difficult to assign a precise error to this area estimation because we do not know how much eruptive material is thinner than the vertical resolution of the seismic data, and thus lies outside of the high-amplitude area used to define the lava flow apron. We used '~' to represent the uncertainty here.

Comment 11: pag. 11, line 234 As for pag. 10, line 206. **Response 11**: We have revised this in the text (Line 241).

Comment 12: pag. 11, line 239 Every measurement has an error, or is \sim , why the 9.2 km long lava flow channel has been defined without uncertainty?

Response 12: The lava flow channel extend beyond the area imaged by the seismic reflection data, and thus is *at least* 9.2 km long. To clarify this, we have changed this sentence to: "V2 lacks a lava apron, instead being directly flanked by relatively straight, >9.2 km long lava flow channels extending beyond the seismic survey boundary on its south-eastern side (C4-C7) (Fig. 5a)" (Lines 245-247).

Comment 13: pag. 11, line 241 As for pag. 10, line 206. pag. 12, line 245 As for pag. 10, line 206. pag. 15, line 317 As for pag. 10, line 206. pag. 16, line 334 As for pag. 10, line 206. Response 13: We have revised all of these in the text.

Comment 14: pag. 16, line 337 How is lava viscosity of 9-38 Pa computed? **Response 14**: 9-38 Pa is an estimate of bulk fluid viscosity in the center of the lava flow calculated using the Giordano et al. (2008) melt viscosity model and an approximated MORB melt composition. This calculation assumes that the melt has equilibrium solubility of H_2O at a pressure pertaining to the eruption depth of 2 km (20 MPa) and at temperatures of 1100-1200°C. The purpose of this calculation is to provide a viscosity comparison with an equivalent basaltic lava flow erupted subaerially vs. that erupted in water depths of 2 km.

Comment 15: pag. 16, line 351 Is it \controls" or \control"? **Response 15**: We have revised the 'controls' to 'control' in the text (L361).

Comment 16: pag. 17, line 363 It would be interesting to give an estimate of the cooling rate of underwater lavas. If possible.

Response 16: The cooling rates of submarine lava flows are highly dependent on their thickness and effusion rate. While we have good volumetric constraints, effusion rates are much harder to determine for whole-scale lava flow cooling rates. There has been prior work by Gregg and Fornari (1998) that looks at the theoretical surficial cooling rate of lava flows. Compositional data of these lava flows would also be required to make informative estimates of cooling rate. It is an interesting area of study that certainly requires more attention but, we argue, outside of the scope of this paper.

Comment 17: pag. 19, line 417 I would rephrase into \can play a critical role in understanding".

Response 17: We have rephrase this sentence (Lines 422-424).

Responses to Dr. Alexander L. Peace

Comment 1: Figure 1a would benefit from a key (i.e. the red, green and blue symbols that are described in the caption).

Response 1: We have added the symbols in the figure caption to the image.

Comment 2: On Figure 2 some of the text for the different logs is very small and difficult to read, particularly the units. I suggest making these larger.

Response 2: We have made the font twice as large to make them easier to read.

Comment 3: In the caption for Figure 5, the mentions of '(a)' and '(b)' could be better placed to describe the figure. As it is they are both at the start of the caption which reads a little awkwardly. Also, Figure 5a might be better with a colour bar.

Response 3: We have changed the locations of (a) and (b) in the figure caption to make them easier to understand. We also added a colour bar to Figure 5(a).

Comment 4: There is a minor grammatical error in the acknowledgements (the 2nd "have" isn't necessary).

Response 4: We could not find 'have' in the acknowledgements; maybe you were referring to the 'Author Contribution'? Please note that we have carefully revised the main text to remove all grammatical errors.

Comment 5: Finally, another good example of a seismic reflection study on offshore volcanoes that may be of interest to the authors is by Keen et al. (2014) on the Charlie-Gibbs Volcanic Province.

References

Keen, C.E., Dafoe, L.T., and Dickie, K., 2014, A volcanic province near the Western termination of the Charlie-Gibbs Fracture Zone at the rifted margin, offshore northeast Newfoundland: Tectonics, v. 33, no. 6, p. 1133–1153, doi: 10.1002/2014TC003547. **Response 5**: We have read and now cite this paper.

Responses to Dr. William W. Chadwick,

Comment 1: Line 1: I think a better title for this paper would be something like "3D seismic imaging of Miocene volcanoes in the South China Sea" – something that is more informative to the reader about the real content of the paper. I don't think this paper is a general discussion about "extrusion dynamics of deep-water volcanoes".

Response 1: Considering a lot of contents were referred to the extrusion dynamics of deep-water volcanoes (See the detailed Responses 2, 4 and 7) and we used 3D seismic data in this paper, we changed the title to "Extrusion dynamics of deep-water volcanoes

revealed by 3D seismic data" to make this title more informative to the reader.

Comment 2: Line 17 and throughout: What does "extrusion dynamics" mean here and throughout the manuscript? The authors need to explain what this means to them somewhere early in the paper. How can 3D (static) seismic images tell you about "dynamics"?

Response 2: Here, 'extrusion dynamics' means how the erupted materials flow and accumulate. Yes, the seismic images are 'static' in the sense volcanism has long-since ceased. However, the present structure and distribution of the volcanoes and their lavas allow us to infer how they formed (i.e. their "dynamics").

Comment 3: Lines 24-25: I suggest taking out "shallow sub-surface depths" because it is unnecessary and potentially confusing with the "deep-water" emplacement of the volcano as a whole. (water depths vs. subsurface depths within sediment)

Response 3: We have deleted "shallow sub-surface depths" from the revised manuscript.

Comment 4: Line 26: In my experience high hydrostatic pressure has little effect on eruption processes (1000 m vs 4000 m depth), so I'm skeptical about this sentence.

Response 4: This has been modified in the revised manuscript (Lines 25-28: *Extrusion dynamics were likely controlled by low magma viscosities as a result of increased dissolved H2O due to high hydrostatic pressure, and soft, near-seabed sediments, which collectively are characteristic of deep-water environments). We attribute the long run out of the lava flows due to increased effusion rate and low lava viscosity. The control of hydrostatic pressure in this setting is on the solubility of the erupted melt, where up to (and over) 20 MPa of hydrostatic pressure may allow lavas to be very H₂O-rich, with viscosities up to an order of magnitude lower than their subaerial counterparts and/or submarine lavas erupted in drier tectonic settings (e.g. at Axial seamount). This is detailed more within the discussion (Lines 336-339).*

Comment 5: Line 49: It seems to me a distinction should be made here. With beforeand-after bathymetric surveys, the volumes of individual eruptions CAN be wellconstrained. It is only if you don't have information on the pre-existing topography or bathymetry – or you are estimating over longer periods of time (multiple eruptions or an entire volcano's history) that volume estimation is more difficult.

Response 5: Yes, we agree with you that the volumes of individual eruptions can be relatively well-constrained, if we carry out the before- and –after-eruption bathymetric surveys and meanwhile *less erosion occurs at the basal surface*. We have expressed this meaning in the sentence "*By collecting high-resolution, quantitative data on the morphology of modern volcanic edifices and surrounding lava flows from airborne/shuttle radar topography or time-lapse multi-beam bathymetry, we can estimate erupted volumes, at least for individual eruptive episodes (e.g. Holcomb et al., 1988; Walker, 1993; Goto and McPhie, 2004; Cocchi et al., 2016; Somoza et al., 2017; Allen et al., 2018; Chadwick et al., 2018; Grosse and Kervyn, 2018)" (Lines 41-46).*

In fact, a pseudo three-dimensional data with only the upper and lower surfaces has been made, if the before- and –after-eruption bathymetric data are available. This pseudo three-dimensional data can use it to characterize the distribution of erupted material and calculate the volume. However, it cannot image the *internal* or *basal* structures of the erupted materials, and thus cannot establish how volcanoes grow and lava is emplaced over multiple eruptive episodes. Moreover, *if large-scale erosion* occurs at the basal surface of lava flow, the volume estimate only based on the before-and –after-eruption bathymetric surveys will be incorrect. Therefore, we still need 'full 3D structure of these extrusive systems' to 'assess the accuracy of estimated volumes of total erupted materials, or test volcano growths and lava emplacement models'. Because we have expressed similar meaning as the reviewer suggested in the sentence mentioned above, we only made a minor revision to this sentence (added 'total'; Line 49) to highlight the volume represents that of entire volcano's history.

Comment 6: Line 51-58: The authors should mention these papers on seismic imaging of Axial Seamount (an active basaltic caldera with a summit depth of \sim 1400 m):

Arnulf, A. F., A. J. Harding, G. M. Kent, S. M. Carbotte, J. P. Canales, and M. R. Nedimovic (2014), Anatomy of an active submarine volcano, Geology, 42(8), 655-658, doi:10.1130/G35629.1.

Arnulf, A. F., A. J. Harding, G. M. Kent, and W. S. D. Wilcock (2018), Structure, seismicity, and accretionary processes at the hotspot-influenced Axial Seamount on the Juan de Fuca Ridge, J. Geophys. Res., 123, doi:10.1029/2017JB015131.

Response 6: We have read and cited Arnulf et al. (2014). However, Arnulf et al. (2018) focuses on a tomographic inversion of OBS data, and although 2D multi-channel seismic data are used, it is quite different from these we mention in Lines 51-58. We therefore choose not to cite Arnulf et al. (2018).

Comment 7: Line 68: Why would pressure have an effect on rheology? Observations from are recent eruption site at \sim 4000 m depth in the Mariana back-arc suggest that high hydrostatic pressure there had little or no effect on eruption dynamics and lava morphology, compared to submarine eruptions observed at shallower depths (for example Axial Seamount at \sim 1500 m):

Chadwick, W. W., Jr., S. G. Merle, E. T. Baker, S. L. Walker, J. A. Resing, D. A. Butterfield, M. O. Anderson, T. Baumberger, and A. M. Bobbitt (2018), A recent volcanic eruption discovered on the central Mariana back-arc spreading center, Front. Ear. Sci., 6:172, doi:10.3389/feart.2018.00172.

Response 7: As addressed in a previous comment (Response 4), the greater hydrostatic pressure will also control the solubility of H_2O in melt at the point of eruption (e.g. >20 MPa). Increased H_2O content in melt may lower bulk lava viscosity enough to modulate effusion rate and propagation dynamics. Dissolved H_2O content in magma will depend on the continental setting, so we may not expect H_2O -controlled viscosity effects to be as observable in tectonic settings where H_2O -undersaturated (even dry) magmas are being erupted on the seafloor e.g. mid ocean ridges and back arc basins. We have modified this sentence to: "*We suggest the high hydrostatic pressure of the deep-water*

environment controlled melt H2O content and internal lava viscosity, effusion rate and, consequently, volcano and lava flow morphology and run-out distance" (Lines 73-75).

Comment 8: Line 71: This statement is inaccurate. Before-and-after multibeam bathymetry calculates depth changes from the shape of the pre-eruption seafloor to the post-eruption seafloor, so does NOT assume a smooth base. You should re-phrase this to something like: "Any eruption volume estimates that do not include pre-eruption topography may be grossly underestimated."

Response 8: We have revised this sentence; "Our results also show that erupted volumes calculated from airborne/shuttle radar topography or time-lapse multi-beam bathymetry data, without knowledge of detailed geometry of the basal surfaces of the lava flows and the volcanoes themselves, may be grossly underestimated, particularly if extrusion was explosive and/or involved erosion of the seabed" (Lines 76-79).

Comment 9: Line 331: I question whether any of the referenced papers here support the statement that "extensive lava flows in deep water... occur primarily because of high hydrostatic pressure...". In fact, I question that conclusion at all.

Response 9: We have revised this sentence. The sentence now reads; "*Extensive lava flows have also been observed at other deep-water volcanoes (e.g. Chadwick et al., 2018; Embley and Rubin, 2018; Ikegami et al., 2018) where greater dissolved H2O contents in melt imply lower melt viscosity while the lavas were mobile." (Lines 336-338). See also response to comment 7. A detailed geochemical analyses of samples would be required to test this hypothesis; unfortunately this is not within the scope of the present study.*

Comment 10: Line 344: You need to explain why you interpret that there are lava tubes (vs. just channels).

Response 10: In this study, the lava tube is core channel or channelization. In the plan view, it looks like a channel with high-low sinuosity. Therefore, we are apt to call them as 'lava flow channel' in this study.

Comment 11: Line 385-386: These references do not support this statement (in the 2nd half of the sentence). The Caress et al. paper describes an eruption in which the largest volume was erupted after lateral intrusion (not transport on the surface), and the Carey et al paper describes an eruption for which the largest volume was erupted as a pumice raft that floated to the ocean surface.

Response 11: Here we simply consider the ratio between the volume of erupted materials contained within the main volcano edifice vs. volume of lava flows transported from the volcano edifice; we do not consider how the erupted materials are transported. To avoid confusion, we have deleted the latter part of this sentence.

Comment 12: Figure 1: If the contour lines are in ms what do they show? The twt to the seafloor? Or some sub-surface horizon? Why not just use depth contours?

Response 12: Figure 1b is the bathymetry of the study area and the contours are in ms

(twt). Our 3D seismic data are in a time- rather than depth domain, thus all maps and profiles are presented in time domain. A precise velocity model is needed to convert from time to depth.

Comment 13: Figure 5: "Lava" is misspelled in the figure 5b legend. **Response 13**: We have revised Figure 5b.

Responses to Dr. Weiwei Ding,

Comment 1: Line 45: change 'volume' to 'volumes'; **Response 1**: We have changed 'volume' to 'volumes' (Line 43).

Comment 2: Line 65: only the volcanoes that presently stand out the seabed can be imaged by the bathymetry and remote sensing data. Therefore, please add 'present' before 'deep-water volcanoes';

Response 2: We have added the word 'modern' before 'deep-water volcanoes' (Line 70).

Comment 3: Line 120: add 'edifice' after volcano; **Response 3**: We have added 'edifice' (Line 128).

Comment 4: Line 205: change 'volcano' to 'volcano edifice' or 'volcano construction'. V1 is a volcano edifice and don't include the lava flows; **Response 4**: We added 'edifice' after 'conical volcano' (Line 212).

Comment 5: Line 236: Please label the age of V1 here; it is easier for the author to compare the ages of V1 and V2; **Response 5**: We have added the age of V1 ('~6.5 Ma') (Line 244).

Comment 6: Line 259: Same to above; please label the age of latest Pliocene? 2.58 Ma? **Response 6**: Yes, it is about the same age or slightly older than T0. It is labeled in Line 267.

Comment 7: Lines 671-683: please carefully check the figure captions and make it clearer. For example, Ds is marked in the Figure 1a. However, it is not interpreted in the figure caption;

Response 7: We have improved labelling in Figure 1 and updated its figure caption (Lines 687-700).

Comment 8: Line 690: revise 'reflection' to 'reflection'; **Response 8**: We have revised the caption for Figure 2 (Line 750). **Comment 9**: Please mention A-A' and B-B' are the locations of Figure 6 and 7 in the caption.

Response 9: We have modified the caption to show the locations of Figure 6 and Figure 7 (Line 726).

1	Extrusion dynamics of deep-water volcanoes revealed by 3D seismic				
2	data				
3					
4	Qiliang Sun ^{1,2,3} , Christopher A-L. Jackson ⁴ , Craig Magee ^{4,5} , Samuel J. Mitchell ⁶ and Xinong Xie ^{1,3}				
5	¹ Key Laboratory of Tectonics and Petroleum Resources, China University of Geosciences (Wuhan),				
6	Ministry of Education, Wuhan 430074, China;				
7	² Laboratory for Marine Mineral Resources, Qingdao National Laboratory for Marine Science and				
8	Technology, Qingdao 266061, China;				
9	³ College of Marine Science and Technology, China University of Geosciences (Wuhan), Wuhan,				
10	Hubei 430074, PR China;				
11	⁴ Basins Research Group (BRG), Department of Earth Science & Engineering, Imperial College,				
12	London, SW7 2BP, UK				
13	⁵ School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK				
14	⁶ School of Earth Sciences, University of Bristol, Bristol, BS8 1RJ UK				
15					
16	Abstract				
17	Submarine volcanism accounts for c. 75% of the Earth's volcanic activity. Yet difficulties with				
18	imaging their exteriors and interiors mean the extrusion dynamicsand erupted volumes of deep-				
19	water volcanoes remain poorly understood. Here, we use high-resolution 3-D seismic reflection data				
20	to examine the external and internal geometry, and extrusion dynamics of two Late Miocene-				
21	Quaternary, deep-water (>2 km emplacement depth) volcanoes buried beneath 55-330 m of				
22	sedimentary strata in the South China Sea. The volcanoes have crater-like basal-basescontacts,				

23	which truncate underlying strata and suggest extrusion was initially explosive, and erupted lava
24	flows that feed lobate lava fans. The lava flows are >9 km long and contain lava tubes that have
25	rugged basal contacts defined by ~90 \pm 23 m high erosional ramps. We suggest the lava flows eroded
26	down into and were emplaced at shallow sub-surface depths within wet, unconsolidated, near-
27	seafloor sediments. Extrusion dynamics were likely controlled by low magma viscosities as a result
28	of,-increased dissolved H ₂ O due to high hydrostatic pressure,s, and soft, near-seabed sediments,
29	which collectively are characteristic of deep-water environments. Because the lava flows and
30	volcanic edifices are imaged in 3D, wWe calculate thate long run-out lava flows account for 50-
31	97% of the total erupted volume-based on the 3D seismic data, and the remaining. Our results
32	indicate deep-water volcanic edifices may thus formwith a surprisingly - a minor component (~3-
33	50%) of being preserved in the extrusive system deposits main volcanic edifice, A and that
34	aAccurate estimates of erupted volumes therefore requires knowledge of the volcano and lava basal
35	surface <u>morphology</u> -of genetically related lava flows. We conclude that 3D seismic reflection data
36	is a powerful tool for constraining the geometry, volumes, and extrusion dynamics volume of
37	buried, ancient or active deep-water volcanoes and lava flowsic features; such data should be used
38	to image and quantify <u>unravel</u> extrusion dynamics of modern deep-water volcanoes.
39	

40 Keywords

41 Volcano, deep-water, lava flow, seismic reflection, South China Sea

42

I

43 **1. Introduction**

44 The external morphology of volcanoes and their eruptive products reflect, and provide insights

45	into, the processes controlling magma extrusion and volcano construction (e.g. Walker, 1993; Planke
46	et al., 2000; Grosse and Kervyn, 2018). By extracting collecting high-resolution, quantitative data
47	on the morphology of modern and, in some cases, still active volcanic edifices and surrounding lava
48	flows from airborne/shuttle radar topography or time-lapse multi-beam bathymetry, we can estimate
49	erupted volumes, at least for individual eruptive episodes and reconstruct volcano growth
50	mechanisms histories (e.g. Holcomb et al., 1988; Walker, 1993; Goto and McPhie, 2004; Cocchi et
51	al., 2016; Somoza et al., 2017; Allen et al., 2018; <u>Chadwick et al., 2018;</u> Grosse and Kervyn, 2018).
52	Whilst remote sensing data capture the external morphology of volcanoes and lava flows, both
53	before, during, and after eruptions, they do not image their basal surface or internal architecture.
54	Without access to the full 3D structure of these extrusive systems, it is difficult to assess the accuracy
55	of estimated volumes of total erupted materials over multiple eruptive episodes, and to thus test or
56	long-term test-volcano growths and lava emplacement models.
57	Several studies demonstrate that seismic reflection data can be used to map the external
58	morphology and internal architecture of buried volcanoes in 3D (e.g. Planke et al., 2000; Calvès et
59	al., 2011; Jackson, 2012; Magee et al., 2013; <u>Arnulf et al., 2014;</u> Reynolds et al., 2017 <u>; Arnulf et al.</u> ,
60	2018). To-date, most seismic-based studies have focused on volcanoes formed in sub-aerial or
61	shallow-marine environments (e.g. Planke et al., 2000; Jackson, 2012; Magee et al., 2013; Keen et
62	al., 2014; Reynolds et al., 2018), although seismic reflection surveys have been used to image the
63	shallowly buried flanks of deep-water volcanoes (e.g. Funck et al. 1996). The 3D geometry, internal
64	structure, extrusion dynamics, and volume of deep-water volcanoes thus remain poorly documented.
65	Without such information on the structure of deep-water volcanoes, we cannot assess how they grow
66	or what potential-hazards and they may pose (e.g. tsunamis induced by flank collapse, seabed

67 <u>deformation and instability induced by highly explosive eruptions</u>).

68	We use high-resolution 3D seismic reflection data to examine the external morphology and
69	internal architecture of two, Late Miocene-Quaternary submarine volcanoes that were emplaced in
70	deep-water (>2.0 km) on highly stretched continental crust in the northern South China Sea (Fig. 1).
71	The volcanoes and associated lava flows are now buried by a ~55-330 m thick sedimentary
72	succession (Fig. 1). Moreover, tThese two volcanoes are physically isolated and appear to have been
73	fed by independent, sub-volcanic intrusive bodies (i.e. sills; see below); we can thus confidently
74	characterize each individual volcano and its associated lava flows (Fig. 1b). By interpreting volcano
75	and lava flow 3D structure, distribution, and size, we aim to determine extrusion dynamics, calculate
76	accurate erupted volumes, and relate our findings to presentmodern deep-water volcanoes studied
77	using bathymetry and remote sensing data. We show <u>that the</u> basal surfaces of <u>these</u> volcanic edifices
78	and lava flows are <i>ruggederosive</i> , with 50–97% of the total erupted material hosted within the latter;
79	i.e. the volcano edifices only comprise only a small portion of the total erupted magma-volume. We
80	suggest the high hydrostatic pressure of the deep-water environment controlled erupting lava
81	rheologymelt H ₂ O content and internal lava viscosity, effusion rate and, consequently, volcano and
82	lava flow morphology and run-out distance. Our results also show that erupted volumes calculated
83	from airborne/shuttle radar topography or time-lapse multi-beam bathymetry data, without the
84	knowledge of detailed geometry of the basal surfaces of the lava flows and the volcanoes
85	themselves,, which typically assume imaged volcanoes and lava flows have a smooth base, may be
86	grossly underestimated, particularly if extrusion was explosive and/or involved erosion of the
87	seabed.

89 2. Geological setting

90	The study area is located in the south of Pearl River Mouth Basin, on the northern, highly
91	stretched continental crust of the South China Sea (Franke, 2013; Zhao et al., 2016) (Fig. 1a). The
92	South China Sea was an area of subduction in the late Mesozoic, before the onset of continental
93	rifting and subsequent seafloor spreading (~33-15 Myr) in the Cenozoic (e.g. Taylor and Hayes,
94	1983; Briais et al., 1993; Franke et al., 2014; Li et al., 2014; Sun et al., 2014a; Ding and Li, 2016).
95	A lack of seaward-dipping reflections (SDRs), and low volumes of rift-related igneous rocks,
96	suggest the northern part of the South China Sea is a magma-poor margin (e.g. Clift et al., 2001;
97	Yan et al., 2006; Cameselle et al., 2017). Seafloor spreading ceased at ~15 Ma (Li et al., 2014), with
98	post-rift thermal cooling driving subsidence of the northern South China Sea margin since the Early
99	Miocene (Ru and Pigott, 1986; Yu, 1994). During this phase of thermal subsidence ₂ the Dongsha
100	Event (~5.3 Ma) occurred, which involved widespread uplift and normal faulting (e.g. Lüdmann et
101	al., 2001). Several mechanisms may have triggered the Dongsha Event, including the collision
102	between Taiwan and the East Asian continent (Lüdmann et al., 2001; Hall, 2002), isostatic rebound
103	(Zhao et al., 2012), post-rift magmatism (Franke, 2013), lithospheric bending (Wu et al., 2014),
104	and/or subduction of the South China Sea beneath the Philippine Sea plate (Xie et al., 2017).
105	Post-spreading magmatism in the South China Sea may reflect ascent of magma triggered by
106	subduction of the South China Sea along the Manila trench and collision with Taiwan Island
107	(Lüdmann et al., 2001), convective removal of continental lithosphere by warm asthenosphere
108	(Lester et al., 2014), or magma ascent from a high-velocity layer in the lower crust fed by the Hainan
109	mantle plume (Xia et al., 2016; Fan et al., 2017). Volcanoes generated by post-rift magmatism in
110	the early Miocene and Quaternary were emplaced both onshore and offshore (e.g. Zou et al., 1995;

111	Yan et al., 2006; Franke, 2013; Li et al., 2014; Sun et al., 2014b; Zhao et al., 2014, 2016; Fan et al.,
112	2017), with the latter typically extruded onto the continental slope in relatively shallow water depths
113	(<300 m; Yan et al., 2006; Zhao et al., 2016). Boreholes reveal these shallow-water volcanoes are
114	composed of basalt, dacite, and rhyolitic tuff (Li and Liang, 1994; Yan et al., 2006; Zhao et al.,
115	2016). In addition to the onshore and shallow-water volcanoes, several volcanoes were emplaced
116	further basinwards on the continental slope in deeper water, close to the Continent-Ocean Boundary
117	(COB; Fig. 1) (Clift et al., 2001; Wang et al., 2006; Cameselle et al., 2017) (Fig. 1). We examine
118	two of these deep-water volcanoes, which are situated in an area currently characterized by water
119	depths of 1850–2680 m and that are now buried by overlyingbeneath sedimentary strata up to 330
120	m thick (Fig. 1). Micropalaeontological data from the Pearl River Mouth Basin (Xu et al., 1995;
121	Qin, 1996), and microfauna data from ODP borehole sites 1146 and 1148, indicate the Middle
122	Miocene (16.5 Ma) to Recent, nanofossil-bearing clays encasing overlying the volcanoes were
123	deposited in a deep-water setting (1.0-3.0 km; Wang et al., 2000).

125 **3. Data and Methods**

We use a time-migrated 3D seismic reflection survey acquired in 2012 and covering an area of ~350 km² (Fig. 1b). The seismic data are zero-phase processed and displayed with SEG (Society of Exploration Geophysicists) normal polarity, whereby a downward increase in acoustic impedance (a function of rock velocity and density) corresponds to a positive reflection event (red on seismic profiles) (e.g. Brown, 2004). Bin The inline and crossline spacingspacing is are 25 m, respectively.; and tThe seismic data have a dominant frequency in the interval of interest (i.e. 0–400 ms two-way time (twt)) of ~40 Hz.

133	Stacking velocities are not available for the survey and no wells intersect the studied Late
134	Miocene-Quaternary, buried, deep-water volcanic features. We thus have no direct control
135	onobservations of the composition or velocities of the seismically imaged volcanic materials. Depth-
136	conversion of volcano edifice and lava flow thickness measurements in milliseconds (twt) to meters
137	is therefore based on velocity estimates, which introduces some uncertainty into our erupted volume
138	calculations. To derive a reasonable velocity estimate, we use velocity data for submarine volcanoes
139	obtained from boreholes (i.e. BY7-1 and U1431) (Li et al., 2015; Zhao et al., 2016) and OBS (Ocean
140	Bottom Seismometer) profiles (Yan et al., 2001; Wang et al., 2006; Chiu, 2010; Wei et al., 2011) in
141	the South China Sea. The boreholes, which are situated >300 km away from our study area, intersect
142	buried basaltic volcanoes with p-wave velocities of ~4.5 km/s (BY7-1; Zhao et al., 2016) and ~3.0-
143	5.0 km/s (IODP U1431; Li et al., 2015). OBS profiles reveal that submarine volcanoes located 140
144	km from the study area (Fig. 1a) typically have p-wave velocities of >3.0 km/s, and occasionally up
145	to ~5.5 km/s (Yan et al., 2001; Wang et al., 2006; Chiu, 2010; Wei et al., 2011). The <u>se</u> basaltic
146	composition and p-wave velocities of ~3.0-5.5 km/s for volcanoes intersected by boreholes and
147	studied using OBS data-are consistent with p-wave velocity data for shallow-water, mafic volcanoes
148	located offshore western India (~3.3-5.5 km/s; Calvès et al., 2011), and southern Australia in the
149	Bight (~2.4–6.7 km/s, with an average velocity of 4.0 km/s; Magee et al. 2013) and Bass (~2.2–4.0
150	km/s with an average of 3.0 km/s; Reynolds et al. 2018) basins. Based on these velocity data, we
151	assume the imaged volcanic material studied deposits here have mafic compositions and p-wave
152	velocities of 4.0 (\pm 1.0) km/s. It is important to note that using a range of estimated velocities does
153	not affect our calculation of the <i>relative</i> amount of material contained within volcanic edifices versus
154	the flanking lava flows.

155	We calculate a vertical resolution ($\lambda/4$) of ~10 m for the sedimentary strata encasing overlying
156	the volcanic materials, given a dominant frequency of 40 Hz and assuming a seismic velocity of 2.2
157	km/s for the nanofossil-bearing clay (based on seismic refraction profiles OBS1993, Yan et al., 2001;
158	OBS2001, Wang et al., 2006; OBS2006-3, Wei et al., 2011). The calculated vertical resolution for
159	the volcanic materials is 19-31 m, based on a dominant frequency of 40 Hz and estimated seismic
160	velocities of 4.0 (\pm 1.0) km/s. The top and base of volcanic structures can be distinguished in seismic
161	reflection data when their thickness is greater than the estimated vertical resolution of these data (i.e.
162	19-31 m) (Brown, 2004). Volcanic structures with thicknesses below the vertical resolution, but
163	above the detection limit (i.e. $\lambda/8 = 10-16$ m), are imaged as tuned reflection packages whereby
164	reflections from their top and base contacts interfere on their return to the surface and cannot be
165	distinguished (Brown, 2004). The lava flows we image are typically >2 seismic reflection thick
166	(>41 \pm 10 m), suggesting they too are thicker than the tuning thickness and are represented by discrete
167	top and basal reflections (Tables 1-3).
168	We used a regional 2D seismic profile and interpreted four seismic surfaces tied to ODP Site 1146,
169	which is located ~65 km west of the study area (Figs. 1a, 2), and two horizons locally mappable
170	around the volcanoes: T0 (~2.58 Ma), T1 (~5.3 Ma), TRa (~6.5 Ma), and TRb (~8.2 Ma), TM (top
171	of the volcanic material) and BM (base of the volcanic material). The youngest age of the volcanoes
172	and associated lava flows are determined using the first seismic reflection that onlaps or overlies
173	them (Fig. 3). After mapping TM and BM, we calculated the volumes of the volcanic features
174	(Tables 1-4), with errors largely arising from uncertainties in the velocities $(4.0\pm1.0 \text{ km/s})$ used to
175	undertake the depth conversion (see above).

176 Root mean square (RMS) amplitude extractions and slices through a variance volume were used

to constrain the geometry, scale, and distribution of the submarine volcanoes (Figs. 3-8). The RMS 177 amplitude attribute computes the square root of the sum of squared amplitudes, divided by the 178 179 number of samples within the specified window used; put simply, the RMS attribute measures the 180 reflectivity of a given thickness of seismic data (Fig. 4a) (Brown, 2004). The variance attribute is 181 free of interpreter bias because it is directly derived from the processed data (Fig. 5). Variance measures the variability in shape between seismic traces; this can be done in a specified window 182 along a picked horizon or within a full 3D seismic volume. Variance is typically used to map 183 184 structural and stratigraphic discontinuities related to, for example, faults and channels (Brown, 185 2004).

186

4. Seismic expression and interpretation of igneous features

188

189 **4.1. Observations**

190 We identify three main types of seismic structures and associated facies related to these buried deep-water volcanoesvolcanic deposits: (1) Seismic Facies 1 (SF1): -two (V1 and V2) conical-191 192 shaped features up to ~ 202 ms twt ($\sim 404 \pm 101$ m) thick, which internally are weakly-to-moderately 193 reflective or chaotic with distinguished reflections downlapping onto BM, capped by a positive polarity, high-amplitude reflection (TM) onlapped by overlying strata (Figs. 3a, 7); (2) Seismic 194 195 Facies 2 (SF2): -ribbon-like, broadly strata-concordant, high-amplitude, positive polarity 196 reflections, which emanate from the conical structures (SF1) and extend up to ~9.2 km downslope 197 (Figs. 3a-b, 6-7); and (3) Seismic Facies 3 (SF3):- saucer-shaped, strata-discordant, high-amplitude reflections situated beneath SF1 and SF2 (Fig. 6). 198

200 **4.2. Interpretations**

201 The conical shape of SF1 and downlap of its internal reflections (where developed) onto BM, 202 coupled with onlap of overlying reflections onto TM, suggest SF1 is an extrusive rather than 203 intrusive feature. SF1 is similar in terms of its conical shape, highly reflective top, and internally chaotic reflections to mud volcanoes documented elsewhere in the northern South China Sea (Sun 204 205 et al., 2012; Yan et al., 2017). It is therefore plausible SF1 could represent a mud volcano that fed 206 long run-out mud flows (i.e. SF2). Alternatively, the highly reflective, ribbon-like geometry of SF2 207 is similar to that associated with shallow/free gas accumulations (Sun et al., 2012). We consider 208 these two interpretations unlikely because: (i) the limited supply and high viscosity of mud means 209 mud volcanoes are rarely associated with long run-out flows, although we note that one mud flow in the Indus Fan was ~5.0 km long (Calvès et al. 2009); and (ii) the top of SF2 is defined by a 210 211 positive polarity reflection (downward increase in acoustic impedance), which is opposite to that typically associated with shallow/free gas accumulations (e.g. Judd and Hovland, 2007; Sun et al., 212 213 2012). Based on their geometric and geophysical characteristics, spatial relationships, and similarity 214 to structures observed on other rifted continental margins, we interpret these features as volcanic 215 edifices (SF1), genetically related lava flows (SF2), and saucer-shaped shallow sills (SF3) (e.g. 216 Berndt et al., 2000; Planke et al., 2000; Thomson and Hutton, 2004; Calvès et al., 2011; Jackson, 217 2012; Magee et al., 2013; Keen et al., 2014; Reynolds et al., 2018). We now focus on the detailed 218 external morphology and internal architecture of the two deep-water volcanoes that are shallowly 219 buried (<330 m) and thus well-imaged.

4.3. Volcano edifice 1 (V1) and associated lava flows

222 V1 is a prominent, ~ 202 ms twt high (404±101 m) and ~ 3.0 km diameter conical velcanovolcanic <u>edifice</u> covering ~7.2 km², with a volume of ~0.94 \pm 0.24 km³ and an average flank dip of ~15.0° \pm 3.6° 223 224 (Figs. 3-4; Table 1). V1 is onlapped by overlying reflections, with the oldest onlapping reflection 225 correlating to TRa (~6.5 Ma); this suggests V1 was emplaced in the latest Miocene-earliest Pliocene 226 (>6.5 Ma) (Fig. 3a). V1 is underlain by a downward-tapering, >1.1 km deep, up to 2.0 km wide, 227 sub-vertical zone of chaotic reflections (Fig. 3a). We attribute the poor imaging within this chaotic 228 sub-vertical zone to: (1) the presence of sub-vertical feeder intrusions that disrupt background 229 reflections and scatter energy (cf. Thomson, 2007); (2) increased fluid flow and hydrothermal 230 alteration in fractured and deformed host rock adjacent to the magma plumbing system; and/or (3) scattering of energy travelling through the volcano, leading to 'wash-out' of the underlying data (i.e. 231 232 a geophysical artefact; Magee et al. 2013). This reduction in imaging beneath the volcanoes partly 233 obscures their basal surface, but where visible it is clear BM undulates and truncates underlying stratal reflections (Fig. 3b). 234

235 Volcano V1 is surrounded by an asymmetric apron of moderate-to-high amplitude reflections 236 extending up to 1.5 km from the main edifice. The apron is up to ~ 115 ms twt thick ($\sim 230\pm 58$ m), and has a dip of <0.5° (Figs. 4a-b; Table 2). A package of moderate-to-very high-amplitude 237 238 reflections extending a further c. 1.5 km down-dip of this apron contains very high-amplitude, 239 channel-like geometries (marked with-C1-C3 in (Fig. 4a), which terminate down-dip into or are 240 flanked at prominent bends by, moderate-amplitude, fan-like geometries (marked with F1-F4 in Fig. 241 4a). We interpret these two features as lava flow channels and fans, respectively (Fig. 3-4). The lava 242 flow channels are sinuous, <340 m wide, and usually bisect the lava fans (Figs 4a-b). Lava flow-

related features (i.e. apron, channels, and fans) emanating from V1 cover an area of ~14 km² (Tables 3-4), have an average thickness of ~33 ms twt (~ 66 ± 17 m), and a volume of ~ 0.92 ± 0.23 km³; this volume is nearly equal to that of V1 (~ 0.94 ± 0.24 km³) and thus represents ~50% of the total erupted volume (~ 1.86 ± 0.47 km³).

247

248 4. 4. Volcano edifice 2 (V2) and associated lava flows

249 V2 covers ~ 0.44 km² and is elliptical in plan-view, with long and short axes of ~ 1.2 km and ~ 0.6 250 km, respectively (Figs. 5, 7). The volcano is ~ 100 ms twt high ($\sim 200\pm 50$ m), with an irregular base, 251 has flank dips of ~27.8°±5.9°, and a volume of 0.03±0.01 km³ (Figs. 5, 7; Table 1). The top of V2 is 252 of moderate amplitude and is irregular, with the oldest onlapping reflections correlating to Reflector 253 T1 (~5.3 Ma) suggesting V2 is latest Miocene-earliest Pliocene (>5.3 Ma), but probably younger 254 than V1 (>6.5 Ma) (Fig. 7). Reflections within V2 are chaotic and, similar to V1, V2 is underlain 255 by a vertical zone of disturbance (Fig. 7). V2 lacks a lava apron, instead being directly flanked by 256 relatively straight, up to ->9.2 km long lava flow channels extending beyond the seismic survey 257 boundary on its south-eastern side (C4-C7) (Fig. 5a). Lava flow C6 is unusual in that underlying 258 strata are truncated at the base of the flow, defining 'ramps' that are up to \sim 32.5 ms twt high (\sim 65 \pm 16 259 m) high and dip towards V2 at ~25.5°±5.8° (Fig. 8). Beyond the main ramp at the base of lava flow 260 C6 (Fig. 5b), the lava flows thickens to \sim 130 ms twt (\sim 260±65 m), where it is defined by stacked, 261 high-amplitude reflections that have a lobate geometry in plan-view (F5) (Figs. 5, 7, 8c-d). At its distal end, the pinch out of F5 occurs where it abuts a basal ramp that is ~90±23 m tall and that dips 262 263 ~9.3°±2.3° (Figs. 8c-d). F5 is capped by a younger lava fan (F6) (Figs. 8c-d). The V2-sourced lava flows (C4-C7 and F5) cover ~11.5 km²; ~4.20 km² of this comprises lava flow channels and ~7.32 264

265 km² lava fan. Given the average thickness of the lava flow channels ($\sim 61\pm16$ m) and fans ($\sim 109\pm27$ 266 m), we estimate the total volume of V2-sourced lava flows to be $\sim 1.05\pm0.27$ km³; this volume 267 estimate is ~ 35 times greater than that of the main V2 edifice (0.03 ± 0.01 km³), representing $\sim 97\%$ 268 of the total erupted volume.

269

4.5. Shallow sills and associated lava flows

South of V2, we map two areally extensive, partly merged lava flows emanating from the upper 271 272 tips of inclined sheets fringing saucer-shaped sills (i.e. S1 and S2) (Figs. 1b, 5-6). A narrow, vertical, 273 seismically chaotic/blanking zone occurs directly below the saucer-shaped sills (Fig. 6). Several 274 linear structures, rooted at the junction between sills, and feeding the overlying lava fan (F6), are 275 also observed (Fig. 6). F6 covers an area of \sim 49 km², with a diameter of \sim 7.9 km and thickness of 276 55±14 m (Table 4). F6 is directly onlapped by surface T0 (~2.58 Ma), suggesting it was emplaced 277 in the latest Pliocene (≥ 2.58 Ma) (Fig. 6). Similar to other lava fans, F6 is characterized by a single, 278 positive, high-amplitude seismic event (Fig. 6). F6 extends beyond the seismic coverage and is much 279 bigger than other lava fans imaged in the study area (Figs. 5-6; Table 4).

281 **5. Discussion**

280

282 5.1. Water depths during volcano emplacement

The different burial depths and onlap relationships of the volcano edifices and lava flows studied here suggest three phases of volcanism: i.e. ~6.5 Ma for V1, ~5.3 Ma for V2, and ~2.58 Ma for S1/S2 (Figs. 2-3, 6-7). According to the relative sea-level change curve of the Pearl River Mouth Basin acquired from nannofossils (Xu et al., 1995; Qin, 1996) and the dating of volcanic phases, the

287	water depths during V1 and V2 emplacement were likely \sim 75 m and \sim 150 m shallower than the
288	present depths of ~2.25 km and ~2.14 km, respectively. The water depth during the emplacement of
289	F6, fed by S1/S2, was probably ~150 m greater than the present depth of ~2.32 km (Xu et al., 1995;
290	Qin, 1996). To be conservative, we estimate that volcanism in the study area occurred in water
291	depths of a little over 2.0 km.

5.2. Origin of post-spreading volcanism in the SCS

294 The volcanoes documented here ($\sim 6.3-2.58$ Ma) have similar ages with those documented in 295 theat Hainan Island (e.g. Tu et al., 1991; Shi et al., 2011) and southwestern SCS (e.g. Li et al., 2013) 296 (Fig. 1a). However, our these volcanoes are substantially younger than those previously observed 297 in the central SCS (~13.8-7.0 Ma; Expedition 349 Scientists, 2014; Li et al., 2015) and on the 298 middle-lower slope of the northern SCS (~23.8-17.0 Ma; Yan et al., 2006; Zhao et al., 2016; Fan et 299 al., 2017). We note such that the small-scale, buried, post-spreading volcanic features studied here 300 have not been identified by lower-resolution techniques (e.g. gravity, magnetism, OBS and 2D 301 seismic data). These young volcanic features may be widespread and diagnostic of post-spreading 302 magmatism across the northern SCS (e.g. Briais et al., 1993; Yan et al., 2006).

303 Given that the volcanoes documented here were emplaced after SCS rifting (>32 Ma ago; e.g.

Taylor and Hayes, 1983; Franke et al., 2014; Li et al., 2015) and spreading (>15 Ma ago; Li et al.,

305 2014), it is clear they have a different origin to-than the breakup-related volcanoes described

- elsewhere (e.g. Yan et al., 2006; Expedition 349 Scientists, 2014; Li et al., 2015; Zhao et al., 2016;
- 307 Fan et al., 2017). The post-spreading age of volcanism may suggest that mantle melting (Clift et al.,
- 308 2001) and convective removal of continental lithosphere by warm asthenosphere (Lester et al.,

309	2014), processes typically associated with rifting and breakup, were not responsible for the
310	generation of this phase of igneous activity. Magmatism gets younger south-eastwards, from ~23.8-
311	17.0 Ma on the proximal continental slope (Yan et al., 2006; Zhao et al., 2016; Fan et al., 2017) to
312	\sim 6.30–2.58 Ma in the deeper water study area. This observation is seemingly in agreement with the
313	results of teleseismic imaging, which shows southeastward migration of the eastern branch of the
314	Hainan mantle plume (Xia et al., 2016). This suggests that plume melt (Xia et al., 2016; Fan et al.,
315	2017) may have supplied magma to the observedthese volcanoes. However, where the Hainan
316	mantle plume was located or even whether the Hainan mantle plume occurred or not are still
317	questioned controversial at present (e.g. Wheeler and White, 2000; He and Wen, 2011; Zhang and
318	Li, 2018). Another possibility for the origin of <u>the magma feeding these volcanoes</u> -is related to
319	the Dongsha Event, which may have that likely triggered the mantle upwelling of mantle materials
320	as well as transtensional faulting (Lüdmann et al., 1999). The Dongsha Event peaked at ~5.3 Ma
321	and 2.58 Ma (Lüdmann et al., 2001) and was broadly synchronous with the main period of eruptive
322	magmatism documented here. Faults generated during the Dongsha Event may have provided high-
323	permeability zones that promoted the vertical migration of magma that fed the eruptive centers.

325 5.3. Volcano construction

Both V1 and V2 are underlain by sub-vertical, pipe-like zones of chaotic reflections, which we suggest demarcate the limits of their magma plumbing systems. The basal surfaces of V1 and V2 truncate underlying strata (Figs. 3a, 7). Apparent erosion of the sub-volcanic substrate may indicate the initial eruptions were explosive, similar to eye-shaped hydrothermal vents documented by, for example, Hansen et al. 2006; Magee et al. 2016). Alternatively, subsidence of the volcano load into

- underlying, wet, unconsolidated sediments may have caused the strata to locally compact andthereby change the reflection configuration, making it appear that they are truncated.
- Internal reflections that lie sub-parallel to the flanks of V1 and V2 suggest the volcanoes grew by increasing both edifice height and diameter by the accretion of volcanic material (Magee et al. 2013). Flank dips of $\sim 15^{\circ}_{-}-28^{\circ}$ likely indicate that the volcanic material building the edifices constitutes coherent lava flows and/or a dome structure, rather than a pyroclastic cone of tephra (Francis and Thorpe, 1974; Griffiths and Fink, 1992). Construction via emplacement of coherent lava flows is consistent with the presence of internal reflections in V1 and V2; i.e. boundaries between blocky lava flows would be irregular and scatter seismic energy, meaning they would not likely be imaged.

341 5.4. Lava flow extrusion dynamics

342 In addition to the formation of volcanic edifices, both V1 and V2, as well as S1 and S2, are associated with extensive lava flows. In particular, we show V1 and V2 are flanked either by an 343 asymmetric lava apron, which is broader on their downslope (SE) side, or lava flow channels that 344 345 flowed south-eastwards for up to >9 km (Figs. 3a, 4a-b, 5a). At sub-aerial volcanoes (e.g. Walker, 346 1993; Cashman et al., 1999), high eruption rates and low magma viscosities are the dominant causes 347 of long run-out lava flows. Extensive lava flows have also been observed at other deep-water 348 volcanoes (e.g. Chadwick et al., 2018; Embley and Rubin, 2018; Ikegami et al., 2018) and they are 349 possibly related tooccur primarily because of the high hydrostatic pressure in deep-water 350 environments where greater dissolved H₂O contents in melt imply lower melt viscosity while the lavas were mobile. These low viscosity, highly mobile <u>and thus, lavas had the</u>could potentially 351 352 have longer run-out distances to propagate further from source. (e.g. Chadwiek et al., 2018; Embley 353 and Rubin, 2018; Ikegami et al., 2018). In particular, <u>Mh</u>igher ambient pressure can also affect bulk 354 lava rheology, (lower viscosity, e.g. decreased vesicularity and ,-crystal content.), Lower gas 355 fractions will also suppress magma decompression and ascent prior to eruption, and, thereby, 356 effusion rates and extrusion dynamics -(Bridges, 1997; Gregg and Fornari, 1998). For example, 357 upon eruption of a 1200°-1100°C basalt (MORB composition) at a confining pressure of 20 MPa (i.e. a hydrostatic-equivalent water depth of 2 km), lava the melt can contain up to 1.4 wt% H₂O at 358 equilibrium volatile solubility (Newman and Lowenstern, 2002). Using the viscosity model of 359 360 Giordano et al. (2008) and 1.4 wt% H₂O, T the resulting lava viscosity of 9–38 Pa s is significantly 361 lower than a dry (e.g. 0.1 wt% H₂O) sub-aerial basalt, having a viscosity range of 41–248 Pa s 362 (calculated using Giordano et al., 2008). Higher H_2O content in lavas erupted in deep-water, compared to those extruded in sub-aerial settings, will mean: (1) there are fewer bubbles from 363 364 suppressed degassing or brittle fragmentation to hinder flow (Gregg and Fornari, 1998); (2) crystallization may be inhibited, reducing the effect of crystal interactions on viscosity; and (3) the 365 366 glass transition temperature is suppressed (Giordano et al. 2008), allowing lavas to flow further as 367 the interiors cool.

From our seismic reflection data it is also clear <u>that</u> channelization in lava tubes, in addition to the water content effects described above, also facilitated long distance lava transport. We suggest these tubes formed by rapid cooling and <u>hardening-thickening</u> of a surficial crust that insulated and focused lava flow through a core channel (e.g. Cashman et al., 1999). Based on the long run-out lava distances, we consider our initial assumption that the imaged volcanic features have a mafic composition remains validis likely. Overall, whilst we do not know the composition of the lavas imaged in our seismic reflection data, pressure-related changes in lava rheology and channelization of any lava type (i.e. mafic to silicic) will allow it to flow hotter for longer. Given the downslope
topographic controls during eruption, a combination of rheology changes and channelization
allowed lavas to flow for >9 km from associated volcanic edifices.

378 The overall geometry and internal architecture of the imaged lava flows indicate substrate 379 rheology was a keyalso a controlled on emplacement dynamics. Our 3D seismic reflection data show that relatively long run-out lava flows (>9 km) erupted from deep-water volcanoes have a rugged 380 381 basal surface that is locally defined by erosional basal 'ramps'. Truncation of underlying strata 382 suggests the lavas were able to erode down into the seabed, perhaps because the pre-eruption 383 substrate was cold, wet, and unconsolidated. We suggest erosion of the lava substrate was promoted by: (1) the dense (bubble-poor) lava sinking down into or 'dredging' the soft sediments (Duffield et 384 al., 1986; Ikegami et al. 2018); (2) thermal erosion (Griffiths, 2000); and/or (3) more "turbulent" 385 386 flow dynamics of channelized lava, consistent with the inferred low viscosities (<10 Pa s).

387 Lava flows eventually ceased in distal areas due to gradual cooling and crystallization (Cashman et al., 1999). We suggest that, in the case of the straight lava flows (C5 and C6), lava transported 388 389 within the axial tube temporarily accumulated at the transient end of the flow, possibly forming a 390 lava pool (Greeley, 1987). Lava entering the tube from the ongoing or new volcanic eruption caused 391 an increase in pressure, with the cooled and crystallized material at the flow toe forming an 392 impermeable, albeit, transient barrier. High hydrostatic pressure (>26 MPa at C5 and C6) and thick 393 surficial crusts inhibited the release of pressure build up by significant lava inflation (Gregg and 394 Fornari, 1998). Eventually, pressure build-up was sufficient to rupture this frontal barrier, leading 395 to emplacement of a downdip fan downdip of the front-most base-lava rampbarrier (F5; Fig. 5a, 7-8) (Griffiths, 2000). However, in the case of fans (e.g. F1-4) fed by sinuous channels (Figs. 4a-b), 396

397	we suggest these were emplaced in a process similar to that documented by Miles and Cartwright
398	(2010), with lobate lava flows fed and bisected by a 'lava tube' through magma inflation and
399	increases in eruption rate. At the end of sinuous lava flow channels (e.g. C1), the main channel
400	bifurcated to form a lobate fan (F3, Figs. 4a-b), which was also probably caused by flow branching
401	triggered by magma cooling (Griffiths, 2000).

403 5.5. Volume balance of volcano edifice and lava flow

404 Inaccurate constraints on total erupted volumes compromises limits our understanding of volcano 405 construction, lava propagation, eruption rates, eruption durations, magma storage conditions, 406 melting processes, and risk assessment of volcanism in deep-water settings (Carey et al., 2018). 407 High-resolution 3D seismic reflection data allow us to calculate the volumes of material contained 408 within volcano edifices and in flanking lava flows, if they are thick enough to be imaged. In this 409 study, We we show that most (i.e. 50-97%) of the erupted material is was transported away from the 410 imaged edifices, an observation comparable to that made for deep ocean volcanic eruptions (Caress 411 et al., 2012; Carey et al., 2018). A critical outcome An important result of our work is that flanking 412 lava flows, and to a lesser extent the volcanic edifices, have rugged and discordant bases (Fig. 7); 413 accurately calculating the volume of deep-water volcanoes and lava flows therefore requires an 414 understanding of their pre-eruption basal morphology. Erupted volume estimates based solely on 415 remote sensing of the post-eruption seabed may be thus incorrect (e.g. Robinson and Eakins, 2006). Although we show the accuracy of total erupted volume estimates can be improved by constraining 416 417 basal volcano and lava morphologies, seismic images capturing the geological record of deep-water 418 volcanoes cannot determine how much, if any, <u>clastic</u> volcanic material was transported away from the eruption site as pumice rafts (e.g. Carey et al. 2018)or through ocean current clast suspension
and remobilizationsubsequent transport (e.g. Jutzeler et al., 2014; Carey et al. 2018). Nevertheless,
3D seismic imaging can significantly improve quantitative volume estimates of recent and ancient
volcanic features (e.g. volcanoic edifices and lava flows), either currently outcropping on the
seafloor or now buried by sedimentary successions.

424

425 **6.** Conclusions

426 High-resolution 3-D seismic data from the South China Sea allow us to image and map the 427 internal structure, calculate the volume of erupted material, and to better understand the extrusion 428 dynamics of buried deep-water volcanoes; such insights cannot readily be gained from analysis of 429 remote sensing data (e.g. airborne/shuttle radar topography). Volcanism occurred ~6.3-2.58 Ma, 430 after seafloor spreading had ceased in the area, and may be related to the Dongsha Event and/or a hypothesized Hainan mantle plume. High hydrostatic pressure, an inclined seabed ($\sim 1^{\circ}$), and low-431 strength, very fine-grained, near-seabed sediments, combined with formation of lava tubes and 432 433 extrusion of low-viscosity magmas, are likely responsible for observed long-distance lava run-outs 434 (>9 km) in this deep-water environment. We show the imaged volcanic edifices and associated lava 435 flows have rugged, erosional bases, meaning traditional remote sensing-based volume calculations 436 of deep-water volcanic features, which typically assume smooth bases, are underestimated. Because 437 seismic reflection data images the base of deep-water volcanoes and lava flows, weWe calculate a 438 large amount (as high as ~97%) of the erupted materials are were transported away from the volcano 439 edifices as lava flows, suggesting that volume of deep-water volcanic edifices may not faithfully archive eruption size or magma production. Considering deep-water conditions (e.g. high 440

441 hydrostatic pressure and unconsolidated sediments) in the study area are common elsewhere, the 442 conclusions derived from this study can likely be used in other deep-water sedimentary basins and 443 <u>perhaps sedimentary some-mid-ocean ridges.</u> Our study highlights that 3D seismic reflection data 444 can <u>play a critical contribute toto</u> understanding volcano morphology in 3D and accurately 445 estimating <u>the volumes of crupted materials</u>.

446

447 Author Contribution

Qiliang Sun, Christopher A-L. Jackson, Craig Magee and Xinong Xie have contributed to the
conceptualization, data analysis, writing and revising the original draft. Samuel J. Mitchell have
contributed to the conceptualization and revising the original draft.

451

452 **Competing interests**

453 The authors declare that they have no conflict of interest.

454

455 Acknowledgment

456 This work was supported by the National Scientific Foundation of China (Grant Nos. 91528301,

457 41676051 and 41372112), the Programme of Introducing Talents of Discipline to Universities (No.

458 B14031) and the Fundamental Research Funds for the Central Universities-the China University of

459 Geosciences (Wuhan) (No. CUG160604). We thank the China National Offshore Oil Company

- 460 (CNOOC) for permission to release the data; reflection seismic data may be requested from CNOOC
- 461 (http://www.cnooc.com.cn/en/). Dieter Franke, Gerome Calvès and Nick Schofield are thanked for
- their invaluable comments and suggestions on a previous version of this manuscript. We thank

- 463 Weiwei Ding, Alexander Peace, and William Chadwick for their constructive comments on this
- 464 version of the manuscript, and Antonella Longo for editorial handling. Rebecca Bell is thanked for
- 465 generously providing office space during the visit of Qiliang Sun to Imperial College.

467 **References**

- 468 Allen, R.W., Berry, C., Henstock, T.J., Collier, J.S., Dondin, F.J-Y., Rietbrock, A., Latchman, J.L., and Robertson,
- 469 R.E.A.: 30 Years in the Life of an Active Submarine Volcano: A Time Lapse Bathymetry Study of the Kick-'em-
- 470 Jenny Volcano, Lesser Antilles, Geochem. Geophy. Geosy., 19, 715-731, https://doi.org/10.1002/2017GC007270,
- **471** 2018.
- 472 <u>Arnulf, A. F., Harding, A.J., Kent, G.M., Carbotte, S.M., Canales, J.P., and Nedimovic, M.R.: Anatomy of an active</u>
 473 submarine volcano, Geology, 42, 655-658, https://doi.org/10.1130/G35629.1, 2014.
- 474 Berndt, C., Skogly, O.P., Planke, S., Eldholm, O., and Mjelde, R.: High-velocity break up-related sills in the Vøring
- 475 Basin, off Norway, J. Geophy. Res., 105, 28443-28454, https://doi.org/10.1029/2000JB900217, 2000.
- 476 Briais, A., Patriat, P., and Tapponnier, P.: Updated interpretation of magnetic anomalies and seafloor spreading stages
- 477 in the South China Sea: Implications for the Tertiary tectonics of Southeast Asia, J. Geophy. Res., 98, 6299-6328,
- 478 https://doi.org/10.1029/92JB02280, 1993.
- 479 Bridges, N.T.: Ambient effects on basalt and rhyolite lavas under Venusian, subaerial, and subaqueous conditions, J.
- 480 Geophy. Res., 102(E4), 9243-9255, https://doi.org/10.1029/97JE00390, 1997.
- 481 Brown, A.R.: Interpretation of three-dimensional seismic data: AAPG Memoir 42, 6thed. SEG Investigations in
- **482** Geophysics, 2004.
- 483 Calvès, G., Schwab, A.M., Huuse, M., Clift, P.D., Gaina, C., Jolley, D., Tabrez, A.R., and Inam, A.: Seismic
- 484 volcanostratigraphy of the western Indian rifted margin: The pre-Deccan igneous province, J. Geophy. Res., 116,

- 485 B01101, https://doi.org/10.1029/2010JB000862, 2011.
- 486 Calvès, G., Schwab, A.M., Huuse, M., van Rensbergen, P., Clift, P.D., Tabrez, A.R., and Inam, A.: Cenozoic mud
- 487 volcanoe activity along the Indus Fan: offshore Pakistan, Basin Res., 22, 398-413, https://doi.org/10.1111/j.1365-
- **488** 2117.2009.00448.x, 2009.
- 489 Cameselle, A.L., Ranero, C.R., Franke, D., and Barckhausen, U.: The continent-ocean transition on the northwestern
- 490 South China Sea, Basin Res., 29, 73-95, https://doi.org/10.1111/bre.12137, 2017.
- 491 Caress, D.W., Clague, D.A., Paduan, J.B., Martin, J.F., Dreyer, B.M., Chadwick Jr, W.W., Denny, A., and Kelley,
- 492 D.S.: Repeat bathymetric surveys at 1-metre resolution of lava flows erupted at Axial Seamount in April 2011,
- 493 Nat. Geosci., 5, 483-488, https://doi.org/10.1038/NGEO1496, 2012.
- 494 Carey, R., Soule, S.A., Manga, M., White, J.D.L., McPhie, J., Wysoczanski, R., Jutzeler, M., Tani, K., Yoerger, D.,
- 495 Fornari, D., Caratori-Tontini, F., Houghton, B., Mitchell, S., Ikegami, F., Conway, C., Murch, A., Fauria, K., Jones,
- 496 M., Cahalan, R., and McKenzie, W.: The largest deep-ocean silicic volcanic eruption of the past century, Sci. Adv.,
- 497 4, e1701121, https://doi.org/10.1126/sciadv.1701121, 2018.
- 498 Cashman, K.V., Thornber, C.R., and Kauahikaua, J.P.: Cooling and crystallization of lava in open channels, and the
- 499 transition of pahochoe lava to `a`a, B. Volcanol., 61, 306-323, https://doi.org/10.1007/s004450050, 1999.
- 500 Chadwick Jr, W.W., Merle, S.G., Baker, E.T., Walker, S.L., Resing, J.A., Butterfield, D.A., Anderson, M.O.,
- 501 Baumberger, T. and Bobbitt, A.M.: A recent volcanic eruption discovered on the central Mariana back-arc
- 502 spreading center: Front. Earth Sci., 6, 172, https://doi.org/10.3389/feart.2018.00172, 2018.
- 503 Chiu, M.: The p-wave velocity modeling of the transitional crust in northern South China Sea continental margin,
- 504 M.S. dissertation, National Taiwan Ocean University, Keelung, 112 pp., 2010.
- 505 Clift, P.D., Lin, J., and ODP Leg 184 Scientific Party: Patterns of extension and magmatism along the continent-
- 506 ocean boundary, South China margin, Geological Society, London, Special Publications, 187, 489-510,

- 507 https://doi.org/10.1144/GSL.SP.2001.187.01.24, 2001.
- 508 Cocchi, L., Masetti, G., Muccini, F., and Carmisciano, C.: Geophysical mapping of Vercelli Seamount: Implications
- 509 for Miocene evolution of the Tyrrhenian back arc basin, Geosci. Front., 7, 835-849,
- 510 https://doi.org/10.1016/j.gsf.2015.06.006, 2016.
- 511 Ding, W.W., and Li, J.B.: Propagated rifting in the Southwest Sub-basin, South China Sea: Insights from analogue
- 512 modelling, J. Geodyn., 100, 71-86, https://doi.org/10.1016/j.jog.2016.02.004, 2016
- 513 Duffield, W.A., Bacon, C.R., and Delaney, P.T.: Deformation of poorly consolidated sediment during shallow
- 514 emplacement of a basalt sill, Coso Range, California, B. Volcanol., 48, 97-107,
- 515 https://doi.org/10.1007/BF01046545, 1986.
- 516 Embley, R.W. and Rubin, K.H.: Extensive young silicic volcanism produces large deep submarine lava flows in the
- 517 NE Lau Basin, B. Volcanol., 80, 36, https://doi.org/10.1007/s00445-018-1211-7, 2018.
- 518 Expedition 349 Scientists: South China Sea tectonics: Opening of the South China Sea and its implications for
- 519 southeast Asian tectonics, climates, and deep mantle processes since the late Mesozoic, International Ocean
- 520 Discovery Program Preliminary Report, 349, https://doi.org/10.14379/iodp.pr.349.2014, 2014.
- 521 Fan, C.Y., Xia, S.H., Zhao, F., Sun, J.L., Cao, J.H., Xu, H.L., and Wan, K.Y.: New insights into the magmatism in
- 522 the northern margin of the South China Sea: Spatial features and volume of intraplate seamounts, Geochem.
- 523 Geophy. Geosy., 18, 2216-2239, https://doi.org/10.1002/2016GC006792, 2017.
- 524 Francis, P.W. and Thorpe, R.S.: Significance of lithologic and morphologic variations of pyroclastic cones, Geo. Soc.
- 525 Am. Bull., 85, 927-930, https://doi.org/10.1130/0016-7606(1974)85<927:SOLAMV>2.0.CO;2, 1974.
- 526 Franke, D.: Rifting, lithosphere breakup and volcanism: comparison of magma-poor and volcanic rifted margins,
- 527 Marine and Petroleum Geology, 43, 63-87, https://doi.org/10.1016/j.marpetgeo.2012.11.003, 2013.
- 528 Franke, D., Savva, D., Pubellier, M., Steuer, S., Mouly, B., Auxietre, J., Meresse, F., and Chamot-Rooke, N.: The

- 529 final rifting evolution in the South China Sea, Mar. Petrol. Geol., v. 58, p. 704-720,
 530 https://doi.org/10.1016/j.marpetgeo.2013.11.020, 2014.
- 531 Funck, T.: Structure of the volcanic apron north of Gran Canaria deduced from reflection seismic, bathymetric and
- borehole data, Ph.D. dissertation, University of Kiel, 156 pp., 1996.
- 533 Giordano, D., Russell, J.K., Dingwell, D.B.: Viscosity of magmatic liquids: a model, Earth Planet. Sci. Lett., 271,
- 534 123-134, https://doi.org/10.1016/j.epsl.2008.03.038, 2008.
- 535 Goto, Y., and McPhie, J.: Morphology and propagation styles of Miocene submarine basanite lavas at Stanley,
- 536 northwestern Tasmania, Australia, J. Volcanol. Geoth. Res., 130, 307-328, https://doi.org/10.1016/S0377-
- **537** 0273(03)00311-1, 2004.
- 538 Grosse, P., and Kervyn, M.: Morphometry of terrestrial shield volcanoes, Geomorphology, 304, 1-14,
- 539 https://doi.org/10.1016/j.geomorph.2017.12.017, 2018.
- 540 Greeley, R.: The role of lava tubes in Hawaiian volcanoes, U.S. Geological Survey Professional Paper 1350, 1589-

541 1602, 1987.

- 542 Gregg, T.K.P., and Fornari, D.J.: Long submarine lava flows: Observations and results from numerical modeling, J.
- 543 Geophy. Res., v. 103, p. 27517-27531, https://doi.org/10.1029/98JB02465, 1998.
- 544 Griffiths, R.W. and Fink, J.H.: Solidification and morphology of submarine lavas: A dependence on extrusion rate,
- 545 J. Geophy. Res., 97(B13), 19729-19737, https://doi.org/10.1029/92JB01594, 1992.
- 546 Griffiths, R.W.: The Dynamics of lava flows, Annu. Rev. Fluid Mech., 32, 477-518,
 547 https://doi.org/10.1146/annurev.fluid.32.1.477, 2000.
- 548 Hall, R.: Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: Computer-based
- reconstructions, model and animations, J. Asian Earth Sci., 20, 353-431, https://doi.org/10.1016/S1367-

550 9120(01)00069-4, 2002.

- 551 He, Y.M., and Wen, L.X.: Seismic velocity structures and detailed features of the D" discontinuity near the core-
- 552 mantle boundary beneath eastern Eurasia, Phys. Earth Planet. In., 189, 176-184,
 553 https://doi.org/10.1016/j.pepi.2011.09.002, 2011.
- 554 Holcomb, R.T., Moore, J.G., Lipman, P.W., and Belderson, R.H.: Voluminous submarine lava flows from Hawaiian
- volcanoes, Geology, 16, 400-404, https://doi.org/10.1130/0091-7613(1988)016<0400:VSlava flow
 fanH>2.3.CO;2, 1988.
- 557 Ikegami, F., McPhie, J., Carey, R., Mundana, R., Soule, S.A. and Jutzeler, M.: The eruption of submarine rhyolite
- 558 lavas and domes in the deep ocean-Havre 2012, Kermadec Arc, Front. Earth Sci., 6, 147,
- 559 https://doi.org/10.3389/feart.2018.00147, 2018.
- 560 Jackson, C.A.-L.: Seismic reflection imaging and controls on the preservation of ancient sill-fed magmatic vents, J.
- 561 Geol. Soc. London, 169, 503-506, https://doi.org/10.1144/0016-76492011-147, 2012.
- 562 Judd, A.G., and Hovland, M. (Eds.): Seabed Fluid Flow: The Impact on Geology, Biology and the Marine
- 563 Environment, Cambridge University Press, Cambridge, 2007.
- 564 Jutzeler, M., Marsh, R., Carey, R.J., White, J.D., Talling, P.J. and Karlstrom, L., 2014.: On the fate of pumice rafts
- 565 formed during the 2012 Havre submarine eruption. Nature communications, 5, <u>p.3660.</u>
- 566 <u>https://doi.org/10.1038/ncomms4660</u>https://doi.org/10.1038/ncomms4660, 2014.
- 567 Keen, C.E., Dafoe, L.T., and Dickie, K.: A volcanic province near the western termination of the Charlie-Gibbs
- 568 Fracture Zone at the rifted margin, offshore northeast Newfoundland: Tectonics, 33, 1133-1153,
- 569 <u>https://doi.org/10.1002/2014TC003547, 2014.</u>
- 570 Lester, R., Van Avendonk, H.J.A., McIntosh, K., Lavier, L., Liu, C.S., Wang, T.K., and Wu, F.: Rifting and
- 571 magmatism in the northeastern South China Sea from wide-angle tomography and seismic reflection imaging: J.
- 572 Geophy. Res., 119, 2305-2323, https://doi.org/10.1002/2013JB010639, 2014.

- 573 Li, C.F., Lin, J., Kulhanek, D.K., and the Expedition 349 Scientists: Proceedings of the International Ocean
- 574 Discovery Program, 349, https://doi.org/10.14379/iodp.proc.349.103.2015, 2015.
- 575 Li, C.F., Xu, X., Lin, J., Sun, Z., Zhu, J., Yao, Y.J., Zhao, X.X., Liu, Q.S., Kulhanek, D.K., Wang, J., Song, T.R.,
- 576 Zhao, J.F., Qiu, N., Guan, Y.X., Zhou, Z.Y., Williams, T., Bao, R., Briais, A., Brown, E.A., Chen, Y.F., Clift, P.D.,
- 577 Colwell, F.S., Dadd, K.A., Ding, W.W., Almeida, I.H., Huang, X.L., Hyun, S., Jiang, T., Koppers, A.A.P., Li, Q.Y.,
- 578 Liu, C.L., Liu, Z.F., Nagai, R.H., Peleo-Alampay, A., Su, X., Tejada, M.L.G., Trin, H.S., Yeh, Y.C., Zhang, C.L.,
- 579 Zhang, F., and Zhang, G.L.: Ages and magnetic structures of the South China Sea constrained by the deep tow
- 580 magnetic surveys and IODP Expedition 349: Geochem. Geophy. Geosy., 15, 4958-4983,
- 581 https://doi.org/10.1002/2014JB011686, 2014.
- 582 Li, L., Clift, P.D., and Nguyen, H.T.: The sedimentary, magmatic and tectonic evolution of the southwestern South
- 583 China Sea revealed by seismic stratigraphic analysis, Mar. Geophys. Res., 34, 341-365,
- 584 https://doi.org/10.1007/s11001-013-9171-y, 2013.
- 585 Li, P., and Liang, H.: Cenozoic magmatism in the Pearl River Mouth Basin and its relationship to the basin evolution
- and petroleum accumulation, Guangdong Geology, 9, 23-34, 1994.
- 587 Lüdmann, T., and Wong, H.K.: Neotectonic regime on the passive continental margin of the northern South China
- 588 Sea, Tectonophysics, 311, 113-138, https://doi.org/10.1016/S0040-1951(99)00155-9, 1999.
- 589 Lüdmann, T., Wong, H.K., and Wang, P.: Plio-Quaternary sedimentation processes and neotectonics of the northern
- 590 continental margin of the South China Sea, Mar. Geol., 172, 331-356, https://doi.org/10.1016/S0025591 3227(00)00129-8, 2001.
- 592 Magee, C., Hunt-Stewart, E., and Jackson, C.A.-L.: Volcano growth mechanisms and the role of sub-volcanic
- 593 intrusions: Insights from 2D seismic reflection data, Earth Planet. Sci. Lett., 373, 41-53,
- 594 https://doi.org/10.1016/j.epsl.2013.04.041, 2013.

- 595 Miles, A., and Cartwright, J.: Hybrid flow sills: A new mode of igneous sheet intrusion, Geology, 38, 343-346,
- 596 https://doi.org/10.1130/G30414.1, 2010.
- 597 Newman, S., and Lowenstern, J.B.: VolatileCalc: a silicate melt-H₂O-CO₂ solution model written in Visual Basic for
- 598 excel, Comput. Geosci., 28, 597-604, https://doi.org/10.1016/S0098-3004(01)00081-4, 2002,
- 599 Planke, S., Symonds, P., Alvestad, E., and Skogseid, J.: Seismic volcanostratigraphy of large-volume basaltic
- 600 extrusive complexes on rifted margins, J. Geophy. Res., 105, 19335-19351,
 601 https://doi.org/10.1029/1999JB900005, 2000.
- 602 Qin, G.Q.: Application of micropaleontology to the sequence stratigraphic studies of late Cenozoic in the Pearl River
- Mouth Basin, Marine Geology& Quaternary Geology, 16, 1-18, https://doi.org/10.16562/j.cnki.02561492.199.04.001, 1996.
- 605 Reynolds, P., Holford, S., Schofield, N., and Ross, A.: Three-dimensional seismic imaging of ancient submarine lava
- flows: an example from the southern Australian margin, Geochem. Geophy. Geosy., 18, 3840-3853,
- 607 https://doi.org/10.1002/2017GC007178, 2017.
- 608 Reynolds, P., Schofield, N., Brown, R.J. and Holford, S.P.: The architecture of submarine monogenetic volcanoes-
- 609 insights from 3D seismic data, Bas. Res., 30, 437-451, https://doi.org/10.1111/bre.12230, 2018.
- 610 Robinson, J.E., and Eakins, B.W.: Calculated volumes of individual shield volcanoes at the young end of the
- 611 Hawaiian Ridge, J. Volcanol. Geoth. Res., 151, 309-617, https://doi.org/10.1016/j.jvolgeores.2005.07.033, 2006.
- 612 Ru, K., and Pigott, J.D.: Episodic rifting and subsidence in the South China Sea, AAPG Bull., 9, 1136-1155, 1986.
- 613 Shi, X., Kohn, B., Spencer, S., Guo, X., Li, Y., Yang, X., Shi, H., Gleadow, A.: Cenozoic denudation history of
- 614 southern Hainan Island, South China Sea: constraints from low temperature thermochronology, Tectonophysics,
- 615 504, 100-115, https://doi.org/10.1016/j.tecto.2011.03.007, 2011.
- 616 Sibuet, J.-C., Yeh, Y.-C., and Lee, C.-S.: Geodynamics of the South China Sea, Tectonophysics, 692, 98-119,

- 617 https://doi.org/10.1016/j.tecto.2016.02.022, 2016.
- 618 Somoza, L., Gonzalez, F.J., Barker, S.J., Madureira, P., Medialdea, T., de Ignacio, C., Lourenco, N., Leon, R.,
- 619 Vazquez, J.T., and Palomino, D.: Evolution of submarine eruptive activity during the 2011-2012 El Hierro event
- 620 as documented by hydroacoustic images and remotely operated vehicle observations, Geochem. Geophy. Geosy.,
- 621 18, 3109-3137, https://doi.org/10.1002/2016GC006733, 2017.
- 622 Sun, Q.L., Xie, X.N., Piper, D.J.W., Wu, J., and Wu, S.G.: Three dimensional seismic anatomy of multi-stage mass
- 623 transport deposits in the Pearl River Mouth Basin, northern South China Sea: Their ages and kinematics, Mar.
- 624 Geol., 393, 93-108, https://doi.org/10.1016/j.margeo.2017.05.005, 2017.
- 625 Sun, Q.L., Wu, S.G., Cartwright, J., Wang, S.H., Lu, Y.T., Chen, D.X., and Dong, D.D.: Neogene igneous intrusions
- 626 in the northern South China Sea: evidence from high resolution three dimensional seismic data, Mar. Petrol. Geol.,
- 627 54, 83-95, https://doi.org/10.1016/j.marpetgeo.2014.02.014, 2014b.
- 628 Sun, Q.L., Wu, S.G., Cartwright, J., and Dong, D.D.: Shallow gas and focused fluid flow systems in the Pearl River
- 629 Mouth Basin, northern South China Sea, Mar. Geol., 315-318, 1-14, https://doi.org/10.1016/j.margeo.2012.05.003,
- **630** 2012.
- 631 Sun, Z., Xu, Z., Sun, L., Pang, X., Yan, C., Li, Y., Zhao, Z., Wang, Z., Zhang, C.: The mechanism of post-rift fault
- 632 activities in the Baiyun Sag, Pearl River Mouth Basin, J. Asian Earth Sci., 89, 76-87,
- 633 https://doi.org/10.1016/j.jseaes.2014.02.018, 2014a.
- fal Taylor, B., and Hayes, D.E.: Origin and history of the South China Sea Basin, in: The Tectonic and Geologic
- 635 Evolution of Southeast Asian Seas and Islands, edited by Hayes, D.E., AGU, Washington, DC, 23-56, 1983.
- 636 Thomson, K.: Determining magma flow in sills, dykes and laccoliths and their implications for sill emplacement
- 637 mechanisms, B. Volcanol., 70, 183-201, https://doi.org/10.1007/s00445-007-0131-8, 2007.
- 638 Thomson, K., and Hutton, D.: Geometry and growth of sill complexes: Insights using 3-Dseismic from the North

- 639 Rockall Trough, B. Volcanol., 66, 364–375, https://doi.org/10.1007/s00445-003-0320-z, 2004.
- 640 Tu, K., Flower, M.F.J., Carlson, R.W., Zhang, M., Xie, G.: Sr, Nd, and Pb isotopic compositions of Hainan basalts
- 641 (south China): implications for a subcontinental lithosphere Dupal source, Geology, 19, 567-569,
- 642 https://doi.org/10.1130/0091-7613(1991)019<0567:SNAPIC>2.3.CO;2, 1991.
- 643 Wang, T.K., Chen, M.K., Lee, C.S., and Xia, K.Y.: Seismic imaging of the transitional crust across the northeastern
- 644 margin of the South China Sea, Tectonophysics, 412, 237-254, https://doi.org/10.1016/j.tecto.2005.10.039, 2006.
- 645 Wang, P., Prell, W.L., and ODP 184 scientists.: Proceedings of the Ocean Drilling Program, Initial Reports, 184.
- 646 Ocean Drilling Program, College Station, TX 2000, 2000.
- 647 Walker, G.P.L.: Basaltic-volcano systems, in: Magmatic Processes and Plate Tectonics, edited by Pritchard, H.M.,
- Alabaster, T., Harris, N.B.W., and Neary, C.R., Geological Society Special Publication, 76, 3-38, 1993.
- 649 Wei, X.D., Ruan, A.G., Zhao, M.H., Qiu, X.L., Li, J.B., Zhu, J.J., Wu, Z.L., and Ding, W.W.: A wide-angle OBS
- profile across the Dongsha uplift and Chaoshan depression in the mid-northern South China Sea, Chinese J.
- 651 Geophys-CH., 54, 3325-3335, https://doi.org/10.3969/j.issn.0001-5733.2011.12.030, 2011.
- 652 Wu, S.G., Gao, J.W., Zhao, S.J., Lüdmann, T., Chen, D.X., and Spence, G.: Post-rift uplift and focused fluid flow in
- 653 the passive margin of Northern South China Sea, Tectonophysics, 615-616, 27-39,
- 654 https://doi.org/10.1016/j.tecto.2013.12.013, 2014.
- 55 Xia, S.H., Zhao, D.P., Sun, J.L., and Huang, H.B.: Teleseismic imaging of the mantle beneath southernmost China:
- new insights into the Hainan plume, Gondwana Res., 36, 33-43, https://doi.org/10.1016/j.gr.2016.05.003, 2016.
- 657 Xie, Z.Y., Sun, L.T., Pang, X., Zheng, J.Y., and Sun, Z.: Origin of the Dongsha Event in the South China Sea: Mar.
- 658 Geophys. Res., 38, 357-371, https://doi.org/10.1007/s11001-017-9321-8, 2017.
- 659 Xu, S.C., Yang, S.K., and Huang, L.F.: The application of sequence stratigraphy to stratigraphic correlation, Earth
- 660 Sci. Front., 2, 115-123, 1995.

- 661 Yan, P., Deng, H., Liu, H.L., Zhang, Z., and Jiang, Y.: The temporal and spatial distribution of volcanism in the South
- 662 China Sea region, J. Asian Earth Sci., 27, 647-659, https://doi.org/10.1016/j.jseaes.2005.06.005, 2006.
- 463 Yan, P., Wang, Y.L., Liu, J., Zhong, G.J., and Liu, X.J.: Discovery of the southwest Dongsha Island mud volcanoes
- amid the northern margin of the South China Sea, Mar. Petrol. Geol., 88, 858-870,
- 665 https://doi.org/10.1016/j.marpetgeo.2017.09.021, 2017.
- 666 Yan, P., Zhou, D., and Liu, Z.S.: A crustal structure profile across the northern continental margin of the South China
- 667 Sea, Tectonophysics, 338, 1-21, https://doi.org/10.1016/S0040-1951(01)00062-2, 2001.
- 468 Yang, S., Qiu, Y., and Zhu, B.: Atlas of Geology and Geophysics of the South China Sea, China Navigation
- 669 Publications, Tianjin, 2015.
- 670 Yu, H.S.: Structure, stratigraphy and basin subsidence of Tertiary basins along the Chinese southeastern continental
- 671 margin, Tectonophysics, 253, 63-76, 1994.
- 672 Zhang, N., and Li, Z.X.: Formation of mantle "lone plumes" in the global downwelling zone A multiscale modelling
- of subduction-controlled plume generation beneath the South China Sea, Tectonophysics, 723, 1-13,
- 674 https://doi.org/10.1016/j.tecto.2017.11.038, 2018.
- 675 Zhao, F., Alves, T.M., Wu, S.G., Li, W., Huuse, M., Mi, L.J., Sun, Q.L., and Ma, B.J.: Prolonged post-rift magmatism
- on highly extended crust of divergent continental margins (Baiyun Sag, South China Sea), Earth Planet. Sci. Lett.,
- 677 445, 79-91, https://doi.org/10.1016/j.epsl.2016.04.001, 2016.
- 678 Zhao, F., Wu, S.G., Sun, Q.L., Huuse, M., Li, W., and Wang, Z.J.: Submarine volcanic mounds in the Pearl River
- Mouth Basin, northern South China Sea, Mar. Geol., 355, 162-172, https://doi.org/10.1016./j.margeo.2014.05.018,
- **680** 2014.
- 681 Zhao, S.J., Wu, S.G., Shi, H.S., Dong, D.D., Chen, D.X., and Wang, Y.: Structures and dynamic mechanism related
- to the Dongsha Event at the northern margin of the South China Sea, Progress in Geophysics, 27, 1008-1019,

- 683 https://doi.org/10.6038/j.issn.1004-2903.2012.03.022, 2012.
- 684 Zou, H., Li, P., and Rao, C.: Geochemistry of Cenozoic volcanic rocks in Zhu Jiangkou Basin and its geodynamic
- 685 significance, Geochimica, 24, 33-45, 1995.
- 686
- 687

688 **Tables**

689

Table 1: Dimensions of volcano edifices. ^adiameter and dip are average values.

Volcano edifice	^a Diameter_	Height	Area	Volume_	ªDip <u>(</u> ⁰)∕⁰
	<u>(m)</u> /m	<u>(m)</u> /m	<u>(km²)</u> /km²	<u>(km³)/km³</u>	
Volcano edifice 1 (V1)	3018	404±101	7.15	0.940±0.235	15.0±3.6
Volcano edifice 1 (V2)	714	200±50	0.44	0.030 ± 0.008	27.8±5.9

691

Table 2: Dimensions of lava flow apron. ^aDiameter is calculated from the area as a circle. V =

693 Volcano edifice.

Lava flow apron	Diameter Area		Thickness	cness Volume		Shape
	(m)	(km ²)	(m)	(km ³)		
Lava flow apron	3182 ^a	7.95	80±20	0.637±0.159	V1	Ring

694

Table 3: Dimensions of lava flow channels (C). Please note that all the lengths of lava flow

696 channels are measured along their axes. ^aMaximum lengths (including the inferred part of lava

flow channels); ^bMinimum length (C3 extends beyond the 3D survey); ^cThickneses cannot be

698 measured, because of lava flow channels (C1 and C2) are only identified on the plan-view map

699 (RMS and variance slice map); ^dArea and volume don't include the inferred part of C5.

Lava flow channels		Length	Width	Thickness	Area	Volume
		(km)	(m)	(m)	(km ²)	(km ³)
Volcano edifices 1-related	C1	2.86 ^a	55-273	unknown ^c	0.31ª	unknown ^c
	C2	3.66 ^a	94-340	unknown ^c	0.56ª	unknown ^c
	C3	4.60 ^b	163-340	52±13	0.84 ^a	0.044±0.011
	C4	2.80	172-229	61±15	0.54	0.032±0.008
Volcano edifices 2-related	C5	9.15 ^a	185-267	64±16	1.52 ^d	$0.097{\pm}0.024^{d}$
	C6	6.39	203-285	60±15	1.47	0.088±0.022
	C7	1.93	236-427	57±14	0.67	0.037±0.009

700

Table 4: Dimensions of lava flow fans. ^aDiameter is calculated from the area as a circle.

Lava flow fans	flow fans Diameter		Thickness	Volume	Feeder	Shape
	(m)	(km ²)	(m)	(km ³)		
Lava flow fan 1	944 ^a	0.70	41±10	0.028±0.007	C1	Lobate
(F1)						
Lava flow fan 2	1050 ^a	0.87	41±10	0.035±0.009	C1	Lobate
(F2)						
Lava flow fan 3	997ª	0.78 ^b	41±10	0.031±0.008b	C1	Lobate
(F3)						
Lava flow fan 4	2171ª	3.70 ^b	41±10	0.148±0.037 ^b	C2	Lobate
(F4)						
Lava flow fan 5	3054ª	7.32	109±27	0.791±0.198	C5/C6	Lobate
(F5)						
Lava flow fan 6	7906 ^a	49.07 ^b	55±14	2.650 ± 0.662^{b}	S1/S2	Lobate
(F6)						

^bMinimum areas and volumes, because of limited data coverage. C = Lava flow channel; S = Sill.



Figure 1: Geological setting of the study area. (a) Bottom leftInset: regional setting of the South 709 710 China Sea that is bounded by the Red River Strike-slip faults (RRFs) to the west and by the 711 subduction trench (Manila Trench) to the east. Hainan Island (HN; Tu et al., 1991; Shi et al., 2011) and southwestern South China Sea (SWSCS; Li et al., 2013) in which the magmatism has the similar 712 713 ages with the studied volcanoes are labelled in blue. In Figure (a): The study area (marked with red square) is located to the south of Dongsha Islands (labeled Ds). The green dashed line outlines the 714 715 boundary of Pearl River Mouth Basin. Locations of boreholes (Red dots; Exploration well BY7-1 716 and ODP sites 1146 and 1148), crustal structure profiles (Purple solid lines; OBS1993 (Yan et al., 2001), OBS2001 (Wang et al., 2006), OBS2006-3 (Wei et al., 2011), and OBS2008 (Chiu, 2010)) 717

718	and mud volcanoes (Mvs) (Light blue dots; Mvs; Sun et al., 2012; Yan et al., 2017) are labeled. Ds
719	= Dongsha Islands; COB = Continent ocean boundary (Adopted from Sibuet et al., 2016). The base
720	map is modified from Yang et al. (2015); (b) Seabed morphologies of the study area. Distributions
721	of volcano edifices (red), sills (blue), lava flows (green) and locations of Figures 4a and 5a are
722	labeled. The contour lines are in 100 ms (twttwo way travel time).
723	







Figure 3: Seismic characteristics of deep-water volcano (V1) and associated lava flow channels/fans.
(a) Seismic profile crosscuts the volcano edifice and associated lava flow; (b) Seismic profile
crosscuts the lava flow (enhanced seismic anomalies). TM = top of volcano/lava flow; BM = base
of volcano/lava flow. <u>See locations of seismic profiles in Figure 4.</u>







Figure 4: (a) and (b) RMS amplitude map (± 30 ms along the surface BM) and its interpretations.
Volcanic apron, lava flow channels/fans are labeled. See map location in Figure 1b. Red lines in (a)







750	Figure 5: (a) Variance slice (extracted from the surface BM) and (b) its interpretations. Lava flows
751	are clearly identified by their texture and marked in (b). $C = lava$ flow channel; $S = shallow sill; F$
752	= lava fan. Red lines A-A' and B-B' in (a) are seismic profiles shown in Figures 6 and 7. (a) and (b)
753	Variance slice (extracted from the surface BM) and its interpretations. Lava flows are clearly
754	identified and marked. C = lava flow channel; S = shallow sill; F = lava fan.
755	







Figure 7: (a) Seismic profile crosscuts V2 and along lava flow channel (C6) and Lava fans (F5 and F6). The V2 has a sharp boundary to the upslope. Lava fan 6 (F6) is directly overlying the Lava fan 5 (F5). BM = base of volcano/lava flow; See location of seismic profile B-B' in Figure 5aSee location in Figure 5a.





Figure 8: (a) and (b) Enlargement of the end of lava flow channel (ramp structure) and its line
drawings; (c) and (d) Enlargement and its line drawings of the lava fans (F5 and F6). BM = base of
volcano/lava flow. See locations in Figure 7.