

# 1 **Insights from elastic thermobarometry into exhumation of high-** 2 **pressure metamorphic rocks from Syros, Greece**

3 Miguel Cisneros<sup>1,2\*</sup>, Jaime D. Barnes<sup>1</sup>, Whitney M. Behr<sup>1,2</sup>, Alissa J. Kotowski<sup>1,3\*</sup>, Daniel F. Stockli<sup>1</sup>, and  
4 Konstantinos Soukis<sup>4</sup>

5 <sup>1</sup>Department of Geological Sciences, Jackson School of Geosciences, University of Texas at Austin, Austin, TX, USA

6 <sup>2\*</sup>Current address: Geological Institute, ETH Zürich, Zürich, Switzerland

7 <sup>3\*</sup>Current address: Department of Earth and Planetary Sciences, McGill, Montreal, Canada

8 <sup>4</sup>Faculty of Geology and Geoenvironment, NKUA, Athens, Greece

9 *Correspondence to:* Miguel Cisneros (miguel.cisneros@erdw.ethz.ch)

10 **Abstract.** Retrograde metamorphic rocks provide key insights into the pressure-temperature (P-T) evolution of exhumed  
11 material, and resultant P-T constraints have direct implications for the mechanical and thermal conditions of subduction  
12 interfaces. However, constraining P-T conditions of retrograde metamorphic rocks has historically been challenging and has  
13 resulted in debate about the conditions experienced by these rocks. In this work, we combine elastic thermobarometry with  
14 oxygen isotope thermometry to quantify the P-T evolution of retrograde metamorphic rocks of the Cycladic Blueschist Unit  
15 (CBU), an exhumed subduction complex exposed on Syros, Greece. We employ quartz-in-garnet and quartz-in-epidote  
16 barometry to constrain pressures of garnet and epidote growth near peak subduction conditions and during exhumation,  
17 respectively. Oxygen isotope thermometry of quartz and calcite within boudin necks was used to estimate temperatures during  
18 exhumation and to refine pressure estimates. Three distinct pressure groups are related to different metamorphic events and  
19 fabrics: high-pressure garnet growth at ~1.4 - 1.7 GPa between 500 - 550 °C, retrograde epidote growth at ~1.3 - 1.5 GPa  
20 between 400 - 500 °C, and a second stage of retrograde epidote growth at ~1.0 GPa and 400 °C. These results are consistent  
21 with different stages of deformation inferred from field and microstructural observations, recording prograde subduction to  
22 blueschist-eclogite facies and subsequent retrogression under blueschist-greenschist facies conditions. Our new results indicate  
23 that the CBU experienced cooling during decompression after reaching maximum high-pressure/low-temperature conditions.  
24 These P-T conditions and structural observations are consistent with exhumation and cooling within the subduction channel  
25 in proximity to the refrigerating subducting plate, prior to Miocene core-complex formation. This study also illustrates the  
26 potential of using elastic thermobarometry in combination with structural and microstructural constraints, to better understand  
27 the P-T-deformation conditions of retrograde mineral growth in HP/LT metamorphic terranes.

## 28 **1 Introduction**

29           Constraining the pressure-temperature (P-T) evolution of metamorphic rocks is fundamental for understanding the  
30 mechanics, timescales, and thermal conditions of plate tectonic processes operating on Earth. Historically, one of the most  
31 challenging aspects of thermobarometry has been deciphering the P-T evolution of rocks during their exhumation from peak  
32 depths back to the surface (e.g., Essene, 1989; Kohn and Spear, 2000; Pattison et al., 2003; Schliestedt and Matthews, 1987;  
33 Spear and Pattison, 2017; Spear and Selverstone, 1983). Exhumation P-T paths are particularly challenging to reconstruct  
34 because during retrogression rocks are cooled, fluids are consumed by metamorphic reactions, and strain is progressively  
35 localized, all of which result in more sluggish reaction kinetics and lesser degrees of chemical equilibrium (e.g., Baxter, 2003;  
36 Carlson, 2002; Jamtveit et al., 2016; Rubie, 1998). These issues are especially pronounced in high-pressure/low-temperature  
37 (HP/LT) environments characteristic of subduction zones.

38           Elastic thermobarometry offers an alternative to conventional thermobarometry. Rather than relying on equilibrium  
39 metamorphic reactions, this approach constrains the P-T conditions at which a host crystal entraps an inclusion (e.g., Adams  
40 et al., 1975a, 1975b; Rosenfeld, 1969; Rosenfeld and Chase, 1961). Because inclusion-host-pair bulk moduli and thermal  
41 expansivities commonly differ, upon ascent, an inclusion develops residual strain(s) that can be determined from measurements  
42 of Raman shifts. A residual inclusion pressure can be calculated from strain(s) by using Grüneisen tensors (Angel et al., 2019;  
43 Murri et al., 2018, 2019) or experimental hydrostatic calibrations (e.g., Ashley et al., 2014; Enami et al., 2007; Thomas and  
44 Spear, 2018). Elastic modeling is then used to calculate the initial entrapment conditions of when the host grew around the  
45 inclusion, and thus can be used to determine the conditions at which individual host minerals grew during metamorphism (e.g.,  
46 Alvaro et al., 2020; Ashley et al., 2014; Enami et al., 2007).

47           The purpose of this study is to illustrate the potential of using elastic thermobarometry in combination with structural  
48 and microstructural observations, to better understand the P-T-deformation (D) conditions of prograde-to-peak and retrograde  
49 mineral growth in subduction-related HP/LT metamorphic rocks. We focus on a subduction complex exposed on Syros Island,  
50 Cyclades, Greece, where despite several decades of petrological study, the early exhumation history remains enigmatic. We  
51 combine the recently tested quartz-in-epidote (qtz-in-ep) barometer (Cisneros et al., 2020), quartz-in-garnet (qtz-in-grt)  
52 barometry (e.g., Ashley et al., 2014; Bonazzi et al., 2019; Thomas and Spear, 2018), and oxygen isotope thermometry (e.g.,  
53 Javoy, 1977; Urey, 1947), to constrain metamorphic growth pressures and temperatures near peak subduction depths and  
54 during early exhumation. The results demonstrate that combining qtz-in-ep barometry with careful structural and  
55 microstructural observations allows us to delineate a retrograde P-T-D path that is contextually constrained, and **provide new**  
56 **insights into the exhumation history of the CBU on Syros, Greece.**

## 57 **2. Geologic Setting**

58           Syros Island in the Cyclades of Greece consists of metamorphosed tectonic slices of oceanic and continental affinity  
59 that belong to the Cycladic Blueschist Unit (CBU), structurally below the Pelagonian Upper Unit (Fig. 1). CBU rocks on Syros

60 record Eocene subduction (~52 - 49 Ma) to peak blueschist-eclogite facies conditions (Bröcker et al., 2013; Cliff et al., 2017;  
61 Lagos et al., 2007; Laurent et al., 2017; Lister and Forster, 2016; Putlitz et al., 2005; Tomaschek et al., 2003; Uunk et al.,  
62 2018), followed by exhumation during Oligo-Miocene (~25 Ma) back-arc extension (e.g., Jolivet and Brun, 2010; Ring et al.,  
63 2010). A retrograde regional metamorphic event occurred between 25-18 Ma and caused greenschist- to amphibolite facies  
64 metamorphism in the Cycladic islands, but was most pervasive in the footwall adjacent to the large-scale extensional North  
65 and West Cycladic Detachment Systems (e.g., Bröcker et al., 1993; Bröcker and Franz, 2006; Gautier et al., 1993; Grasemann  
66 et al., 2012; Jolivet et al., 2010; Pe-Piper and Piper, 2002; Schneider et al., 2018). Despite these documented metamorphic  
67 events, the exhumation history of the CBU between ~52 and ~25 Ma remains enigmatic and poorly constrained; yet, this period  
68 spans exhumation of the CBU from maximum subduction to middle crust pressures (~0.3 - 0.7 GPa). Previous work has  
69 constrained some aspects of the early exhumation history of the CBU on Syros, including: the timing of peak and retrograde  
70 metamorphism (e.g., Bröcker et al., 2013; Cliff et al., 2017; Lagos et al., 2007; Laurent et al., 2017; Skelton et al., 2018;  
71 Tomaschek et al., 2003), prograde and exhumation-related kinematics (e.g., Behr et al., 2018; Keiter et al., 2011; Kotowski  
72 and Behr, 2019; Laurent et al., 2016; Philippon et al., 2011; Rosenbaum et al., 2002), and the retrograde P-T path (e.g., Laurent  
73 et al., 2018; Ring et al., 2020; Schumacher et al., 2008; Skelton et al., 2018; Trotet et al., 2001a, 2001b); however, debate  
74 remains about the relationship between deformation events and retrograde metamorphism, the maximum pressure reached by  
75 different CBU rock types, the retrograde P-T path, and the mechanisms and kinematics of CBU exhumation.

76 In this work, we focus on rocks within the CBU, which consist of intercalated metavolcanic and metasedimentary  
77 rocks, metabasites, and serpentinites (e.g., Keiter et al., 2011). The CBU has been separated into the “Upper Cycladic  
78 Blueschist Nappe” and the “Lower Cycladic Blueschist Nappe” on Milos Island; the Upper Nappe records peak pressure  
79 conditions above ~0.8 GPa (~2.0 GPa and 550 °C; Grasemann et al., 2018). **Previous studies have reported a wide range of  
80 maximum P-T conditions for rocks from the Upper Cycladic Blueschist Nappe on different Cycladic islands [Sifnos: ~1.4 –  
81 2.2 GPa and 450 – 550 °C (e.g., Schmädicke and Will, 2003; Groppo et al., 2009; Dragovic et al., 2012, 2015; Schliestedt and  
82 Matthews, 1987; Matthews and Schliestedt, 1984; Ashley et al., 2014; Spear et al., 2006); Tinos: ~1.4 – 2.6 GPa and ~450 –  
83 550 °C (e.g., Bröcker et al., 1993; Lamont et al., 2020; Parra et al., 2002); Naxos: ~1.2 – 2.0 GPa and ~450 – 600 °C (e.g.,  
84 Avigad, 1998; Peillod et al., 2017, 2021); Sikinos: ~1.1 – 1.7 GPa and ~ 500 °C (e.g., Augier et al., 2015; Gupta and Bickle,  
85 2004)].** Some conventional thermobarometry (i.e., thermobarometry techniques that rely on chemical equilibrium) suggests  
86 that the CBU on Syros reached peak P-T conditions of ~1.5 GPa and ~500 °C (Ridley, 1984). Trotet et al. (2001a) and Laurent  
87 et al. (2018) suggest higher peak P-T conditions of ~2.0 - 2.4 GPa and ~500 - 550 °C; however, multi-mineral phase equilibria  
88 of marbles (Schumacher et al., 2008) and elastic thermobarometry of metabasites from Kini beach (Behr et al., 2018) support  
89 the original P-T estimates of ~1.5 GPa and 500 °C. Published exhumation P-T paths for the CBU on Syros are also highly  
90 variable, ranging from cooling during decompression, near-isothermal decompression, to cooling during decompression  
91 followed by reheating at moderate pressures (Laurent et al., 2018; Schumacher et al., 2008; Skelton et al., 2018; Trotet et al.,  
92 2001a). Because of these conflicting P-T paths, several models have been proposed to explain the exhumation history of the  
93 CBU, including coaxial vertical thinning (Rosenbaum et al., 2002), extrusion wedge tectonics (Keiter et al., 2011; Ring et al.,

94 2020), multiple cycles of thrusting and extension (Lister and Forster, 2016), continuous accretion and syn-orogenic extension  
95 (Trotet et al., 2001a, b), and subduction channel exhumation (Laurent et al., 2016).

### 96 **3. Field and Microstructural Observations**

97 We studied four localities on Syros (Kalamisia, Delfini, Lotos, Megas Gialos; Fig. 1). Each locality exhibits multiple  
98 stages of mineral growth, and the same deformation and P-T progression. Kalamisia records blueschist facies metamorphism,  
99 and Delfini, Lotos, and Megas Gialos record blueschist-greenschist facies metamorphism. GPS coordinates of collected  
100 samples and their associated mineralogy are provided in the supplementary material (Supplementary Table S1). 1 – 4 samples  
101 from each locality were examined petrographically.

#### 102 **3.1 Kalamisia**

103 Mafic rocks from Kalamisia preserve retrograde blueschist facies metamorphism (Fig. 1). Protoliths of Kalamisia  
104 rocks are fine-grained basalts. They exhibit an early foliation ( $S_s$ ) characterized by relict blueschist and eclogite facies minerals.  
105 The early  $S_s$  fabric is re-folded by upright folds ( $F_{t1}$ ) with steeply dipping axial planes, NE-SW-oriented fold hinge lines, and  
106 NE-SW-oriented stretching lineations primarily defined by white mica, glaucophane, and epidote; this indicates syn-blueschist  
107 facies folding ( $D_{t1}$ ).

108 Garnets in Kalamisia mafic samples occur as ~1 - 4 mm porphyroblasts (KCS70A, Supplementary Fig. S1), lack a  
109 well-defined internal foliation, and the  $S_s$  foliation deflect around garnets. Glaucophane typically grows within pressure  
110 shadows and brittle fractures of garnet, and omphacite displays breakdown and alteration to glaucophane; this indicates  
111 retrograde glaucophane growth. Glaucophane inclusions within epidote are commonly oriented parallel to  $S_s$ , and no omphacite  
112 is observed as inclusions within epidote; these observations support epidote (ep1) growth during retrograde metamorphism.

#### 113 **3.2 Delfini Beach**

114 Metasedimentary rocks (quartz-rich lenses intermixed with metavolcanics) at Delfini Beach show retrogression from  
115 eclogite- and blueschist- to greenschist facies (Fig. 1). Protoliths of Delfini rocks remain enigmatic, but may be graywackes  
116 or sandstones variably intermixed with mafic tuffitic intercalations. The rocks at Delfini exhibit an early foliation (also  
117 considered  $S_s$ ) characterized by relict blueschist and eclogite facies minerals (garnet porphyroblasts, and foliation-parallel  
118 white mica, blue amphibole, and epidote) aligned in tight isoclinal folds ( $F_s$ ) with shallow axial planes. This early fabric was  
119 locally retrogressed and re-folded by upright folds (considered  $F_{t2}$ ) with steeply dipping axial planes, E-W-oriented fold hinge  
120 lines, and E-W-oriented stretching lineations primarily defined by white mica, chlorite, and actinolite (considered  $D_{t2}$ , Fig.  
121 2a,b); this indicates folding under greenschist facies conditions.  $D_{t2}$  folding was associated with boudinage of earlier-  
122 generation epidote parallel to the fold hinge lines, and simultaneous precipitation of new coarse-grained epidote (ep2), along

123 with quartz, calcite and iron oxides in boudin necks (Fig. 3). In some areas of tight  $D_{12}$  folding, a new generation of fine-  
124 grained epidote (also interpreted as ep2) grows within a newly developed crenulation cleavage ( $S_{12}$ , Fig. 2c,d,e).

125 Garnets in Delfini metasedimentary samples occur as ~1 - 4 mm, partially chloritized porphyroblasts (KCS34, Fig.  
126 2c), and as <1 mm garnets that are commonly found as inclusions within epidote (KCS1621, Supplementary Fig. S3). Foliation  
127 parallel epidotes (ep1) found within early blueschist-greenschist facies outcrops (KCS1621) range in size from ~0.5 – 5 mm  
128 (b-axis length), are strongly poikiloblastic, lack late greenschist facies inclusions such as chlorite, and commonly contain an  
129 internal foliation that is oblique to the external matrix  $S_s$  foliation (Fig. 2f,g; Supplementary Fig. S3). Late epidote (ep2)  
130 crystals are found within sample KCS34 from the core of an upright fold ( $F_{12}$ ). During upright folding, a predominant portion  
131 of the rock is recrystallized to late-stage greenschist facies minerals, and contains new epidote (ep2) that is oriented parallel to  
132 the  $S_{12}$  crenulation cleavage. Ep2 crystals range from ~50 - 300  $\mu\text{m}$  along the b-axis (Fig. 2c,d,e), tend to be euhedral (Fig.  
133 2d,e), sometimes contain titanite inclusions (Fig. 2d), and show textural equilibrium with white mica and titanite that also  
134 formed in the  $S_{12}$  cleavage (Fig. 2d,e). Ep2 crystals are not poikiloblastic and rarely preserve quartz inclusions, thus only a few  
135 analyses were possible.

### 136 3.3 Lotos Beach

137 The rocks from Lotos Beach exhibit the same structural and petrological progression as those from Delfini (Fig. 1),  
138 showing retrogression from eclogite- and blueschist- to greenschist facies. Protoliths of Lotos rocks are primarily fine-grained  
139 volcanics. An early  $S_s$  foliation was locally retrogressed and re-folded by upright  $F_{12}$  folds with steeply dipping axial planes,  
140 E-W-oriented fold hinge lines, and E-W-oriented stretching lineations primarily defined by white mica, chlorite, and actinolite  
141 ( $D_{12}$ ).  $D_{12}$  folding was associated with boudinage of earlier-generation epidote parallel to the fold hinge lines, and simultaneous  
142 precipitation of new coarse-grained epidote (ep2), along with quartz, calcite and iron oxides in boudin necks (Fig. 3).

143 Garnets in Lotos samples occur as ~1 - 3 mm chloritized porphyroblasts (e.g., KCS3), that deflect the external  $S_s$   
144 foliation (KCS3). Foliation parallel epidotes (ep1) found within early blueschist-greenschist facies outcrops (SY1402, SY1405,  
145 KCS2, KCS3) range in size from ~0.5 – 5 mm (b-axis length), are strongly poikiloblastic, and commonly contain an internal  
146 foliation that is oblique to the external matrix  $S_s$  foliation (Supplementary Fig. S4). Boudinage of ep1 parallel to stretching  
147 lineations is common in thin sections (Supplementary Fig. S4).

### 148 3.4 Megas Gialos

149 The rocks from Megas Gialos exhibit the same structural and petrological progression as those from Lotos and Delfini  
150 Beaches (Fig. 1). Protoliths of Megas Gialos rocks remain enigmatic, but may be sediments intermixed with volcanics. Rocks  
151 from Megas Gialos show retrogression from eclogite- and blueschist- to greenschist facies. An early  $S_s$  foliation was locally  
152 retrogressed and stretching lineations primarily defined by white mica, chlorite, and actinolite are E-W-oriented.

153 No garnets were found within the analyzed sample from Megas Gialos. Foliation parallel epidotes (ep1) found within  
154 early blueschist-greenschist facies outcrops range in size from ~0.5 – 3 mm (b-axis length), are strongly poikiloblastic, and

155 commonly contain an internal foliation that is oblique to the external matrix  $S_s$  foliation (Supplementary Fig. S5). Boudinage  
156 of ep1 parallel to stretching lineations is common in thin sections (Supplementary Fig. S5).

## 157 **4. Methods**

158 We determined P-T conditions using elastic thermobarometry and oxygen isotope thermometry. Raman spectroscopy  
159 was used to measure Raman shifts of strained quartz inclusions entrapped within epidote or garnet, and a laser fluorination  
160 line and a GasBench II coupled to a gas source mass spectrometer was used to measure oxygen isotope ratios of quartz and  
161 calcite separates, respectively.

### 162 **4.1 Raman Spectroscopy measurements**

163 Our Raman spectroscopy measurements are taken from  $\sim 30\ \mu\text{m}$ ,  $\sim 80\ \mu\text{m}$ , and  $\sim 150\ \mu\text{m}$  thin and thick sections, that  
164 consist of sections cut perpendicular to foliation ( $S_s$ ) and parallel to stretching lineations (e.g., KCS1621), and perpendicular  
165 to the  $F_2$  fold axial plane (KCS34). Quartz inclusions were measured from multiple epidotes and garnets within individual  
166 sections (Supplementary Table S3). Measured quartz inclusions were small in diameter relative to the host, and were two-to-  
167 three-times the inclusion radial distance from other inclusions, fractures, and the host exterior to avoid overpressures or stress  
168 relaxation (Fig. 4a,b; Campomenosi et al., 2018; Zhong et al., 2020). No geometric corrections were applied (Mazzucchelli et  
169 al., 2018).

170 Raman spectroscopy measurements of quartz inclusions within garnet and epidote were carried-out at Virginia Tech  
171 (VT) and ETH Zürich (ETHZ) by using JY Horiba LabRam HR800 and DILOR Labram Raman systems, respectively.  
172 Analyses at VT used an 1800 grooves  $\text{mm}^{-1}$  grating, 100x objective with a 0.9 numerical aperture (NA), 400  $\mu\text{m}$  confocal  
173 aperture, and a 150  $\mu\text{m}$  slit width. Raman spectra were centered at  $\sim 360\ \text{cm}^{-1}$ . We used a 514.57 nm wavelength Ar laser, and  
174 removed the laser interference filter for all analyses to apply a linear drift correction dependent on the position of the 116.04  
175  $\text{cm}^{-1}$ , 266.29  $\text{cm}^{-1}$ , and 520.30  $\text{cm}^{-1}$  Ar plasma lines (Fig. DR4). Measurements at ETHZ used a 532 nm laser, an 1800 grooves  
176  $\text{mm}^{-1}$  grating, a 100x objective with a 0.9 NA, a 200  $\mu\text{m}$  confocal aperture, and a 300  $\mu\text{m}$  slit width. Raman spectra were  
177 centered at  $\sim 850\ \text{cm}^{-1}$ .

178 All Raman spectra was reduced with a Bose-Einstein temperature-dependent population factor (Kuzmany, 2009). All  
179 Raman bands were fit by using PeakFit v4.12 from SYSTAT Software Inc. A Gaussian model was used to fit Ar plasma lines  
180 (only VT analyses), and a Voigt model was used to fit the quartz 128  $\text{cm}^{-1}$ , 206  $\text{cm}^{-1}$ , and 464  $\text{cm}^{-1}$  bands, epidote bands, and  
181 garnet bands. Raman bands of quartz, epidote, and garnet, and Ar plasma lines were fit simultaneously, and a linear background  
182 subtraction was applied during peak fitting. Baseline-to-baseline deconvolution of quartz and garnet bands was simple and  
183 generally required fitting quartz bands and a few shoulder garnet bands. Deconvolution of quartz and epidote bands required  
184 more complicated deconvolution; we followed a fitting approach similar to that described by Cisneros et al. (2020).

## 185 4.2 Inclusion and entrapment pressure calculations

186 The fully encapsulated inclusions preserve strain that causes the Raman active vibrational modes of inclusions to be  
187 shifted to higher or lower wavenumbers relative to minerals that are unstrained (fully exposed). We calculated the Raman  
188 shift(s) of inclusions ( $\omega_{\text{inc}}$ ) relative to Raman shift(s) of an unencapsulated Herkimer quartz standard ( $\omega_{\text{ref}}$ ) at ambient  
189 conditions ( $\Delta\omega = \omega_{\text{inc}} - \omega_{\text{ref}}$ ) (Fig. 4). For VT analyses,  $\omega_{\text{inc}}$  was measured relative to a Herkimer quartz standard that was  
190 analyzed 5 times prior to same day analyses. A drift correction was applied to  $\omega_{\text{inc}}$  by monitoring the position of Ar plasma  
191 lines (Supplementary Tables S2; S3). For ETHZ analyses, a Herkimer quartz standard was analyzed 3 times prior to and after  
192 quartz inclusion analyses. A time-dependent linear drift correction was applied to  $\omega_{\text{inc}}$  based on the drift shown by Herkimer  
193 quartz analyses that bracketed inclusion analyses ( $< 0.2 \text{ cm}^{-1}$ ).

194 We calculated residual inclusion pressures ( $P_{\text{inc}}$ ) by using hydrostatic calibrations and by accounting for quartz  
195 anisotropy. To calculate a  $P_{\text{inc}}$  from individual quartz Raman bands, we used pressure-dependent Raman shift(s) ( $P\text{-}\Delta\omega$ ) of the  
196 quartz  $128 \text{ cm}^{-1}$ ,  $206 \text{ cm}^{-1}$ , and  $464 \text{ cm}^{-1}$  bands, that have been experimentally calibrated under hydrostatic stress conditions  
197 by using diamond anvil cell experiments (Schmidt and Ziemann, 2000). To account for quartz anisotropy, we calculated  $P_{\text{inc}}$   
198 from strains. Calculating quartz strains requires that the Raman shift of at least 2 quartz vibrational modes can be measured.  
199 When we were able to measure the quartz  $128$ ,  $206$  and  $464 \text{ cm}^{-1}$  band positions of inclusions, we calculated strains from the  
200  $\Delta\omega$  of 3 bands. If only two bands were measured, we calculated strains from the  $\Delta\omega$  of 2 bands (Supplementary Table S3). For  
201 the remaining analyses with low  $128$  and  $206 \text{ cm}^{-1}$  intensities, we report  $P_{\text{inc}}$  calculated from the  $464 \text{ cm}^{-1}$  band hydrostatic  $P\text{-}$   
202  $\Delta\omega$  relationship (Supplementary Table S3). Strains were determined from the  $\Delta\omega$  of the  $128 \text{ cm}^{-1}$ ,  $206 \text{ cm}^{-1}$ , and  $464 \text{ cm}^{-1}$   
203 quartz bands by using Strainman (Angel et al., 2019; Murri et al., 2018, 2019), wherein a weighted fit was applied based on  
204 the  $\Delta\omega$  error associated with each quartz Raman band. Calculated strains were converted to a mean stress [ $P_{\text{inc}} = (2\sigma_1 + \sigma_3)/3$ ]  
205 using the matrix relationship  $\sigma_i = c_{ij}\epsilon_j$ , where  $\sigma_i$ ,  $c_{ij}$ , and  $\epsilon_j$ , are the stress, elastic modulus, and strain matrices, respectively. We  
206 used the  $\alpha$ -quartz trigonal symmetry constraints of Nye (1985) and quartz elastic constants of Wang et al. (2015).

207 We assumed constant mineral compositions for all modeling (epidote:  $X_{\text{ep}} = 0.5$  and  $X_{\text{cz}} = 0.5$ ; garnet:  $X_{\text{Alm}} = 0.7$ ,  
208  $X_{\text{Gr}} = 0.2$ , and  $X_{\text{Py}} = 0.1$ ). Garnet compositions have a negligible effect on entrapment pressures ( $P_{\text{trap}}$ ) because the  
209 thermodynamic and physical properties of garnet end-members are similar (e.g., Supplementary Table S8). Epidote  
210 composition has a greater effect on  $P_{\text{trap}}$ , but the compositional dependence is minor  $< 1.5 \text{ GPa}$  (Cisneros et al., 2020). To  
211 account for epidote and garnet solid solutions, we implemented linear mixing of shear moduli and molar volumes ( $V$ ). Ideal  
212 mixing of molar volumes has been shown to be an appropriate approximation for epidote-clinzoisite solid solutions (Cisneros  
213 et al., 2020; Franz and Liebscher, 2004). Garnet molar volumes were modeled using the thermodynamic properties of Holland  
214 and Powell (2011) (almandine and pyrope) and Milani et al. (2017) (grossular), and a Tait Equation of State (EoS) with a  
215 thermal pressure term. We used the shear moduli of Wang and Ji (2001) (almandine and pyrope) and Isaak et al. (1992)  
216 (grossular). Epidote molar volumes were modeled using the thermodynamic properties and shear moduli given by Cisneros et  
217 al. (2020), and a Tait EoS and thermal pressure term. Epidote and clinzoisite regressions are based on the  $P\text{-}V\text{-}T$  data of Gatta

218 et al. (2011) ( $X_{ep} = 0.74$ ), and T-V and P-V data of Pawley et al. (1996) ( $X_{ep} = 0$ ) and Qin et al. (2016) ( $X_{ep} = 0.39$ ), respectively.  
219 Clinzoisite and epidote have similar thermal expansivities but differing bulk moduli (Supplementary Table S4). To account  
220 for the composition of epidotes used in P-V-T experiments, we normalized the composition of our unknown epidotes across  
221 the compositional range of P-V experimental epidotes, i.e., the molar volume of our unknown epidote ( $X_{ep} = 0.5$ ) is estimated  
222 as 31 % ( $X_{ep} = 0.74$ ) and 69 % ( $X_{ep} = 0.39$ ) of each experimental epidote. Quartz molar volumes were modeled using the  
223 thermodynamic properties and approach of Angel et al. (2017a). Entrapment pressures were calculated from residual quartz  
224  $P_{inc}$  by using the Angel et al. (2017b) 1D elastic model equation, and a MATLAB program available in Cisneros and Befus  
225 (2020) that implements mixing of shear moduli and molar volumes. A comparison of entrapment pressures calculated from  
226 the Cisneros and Befus (2020) MATLAB program and EoSFit-Pinc (Angel et al., 2017b) is given in Supplementary Table S4;  
227 entrapment pressure calculations of mineral end-members accounts for the reproducibility of molar volume and elastic  
228 modeling calculations.

### 229 **4.3 Stable isotope measurements**

230 Samples were measured by using a ThermoElectron MAT 253 isotope ratio mass spectrometer (IRMS) at the  
231 University of Texas at Austin. Quartz  $\delta^{18}O$  values were measured by laser fluorination (Sharp, 1990), and ~2.0 mg of quartz  
232 were used in each analysis. Quartz from samples SY1613, SY1617, and SY1623 was duplicated to determine isotopic  
233 homogeneity and reproducibility. An internal quartz Lausanne-1 standard ( $\delta^{18}O = +18.1\text{‰}$ ) was analyzed with all samples to  
234 evaluate precision and accuracy. All  $\delta^{18}O$  values are reported relative to standard mean ocean water (SMOW), where the  $\delta^{18}O$   
235 value of NBS-28 is +9.65‰. Measurement precision based on the long-term reproducibility of standards is  $\pm 0.1 \text{‰}$  (1  $\sigma$ ).  
236 Precision of Lausanne-1 on the day of analysis was  $\pm 0.3 \text{‰}$  (1  $\sigma$ ), whereas samples reproduced with a precision of  $\pm 0.1 \text{‰}$   
237 (1  $\sigma$ ) or better (Supplementary Table S5). Calcite  $\delta^{18}O$  values were measured on a Thermo Gasbench II coupled to a  
238 ThermoElectron 253 mass spectrometer. Each analysis used 0.25 - 0.5 mg of calcite that was loaded into Exetainer vials,  
239 flushed with ultra-high purity helium, and reacted with 103 % phosphoric acid at 50 °C for ~2 hours. Headspace  $CO_2$  was then  
240 transferred to the mass spectrometer. Samples were calibrated to an in-house standard, NBS-18, and NBS-19. Measurement  
241 precision is  $\pm 0.04 \text{‰}$  (1  $\sigma$ ) based on the long-term reproducibility of standards.

### 242 **4.4 Stable isotope temperature calculations**

243 Temperatures derived from stable isotope measurements were calculated by using the Sharp and Kirschner (1994)  
244 quartz-calcite oxygen isotope fractionation calibration ( $A = 0.87 \pm 0.06$ ; equation A1; Supplementary Table S5). Isotopic  
245 equilibrium was assumed for all samples. Several observations support that this assumption is appropriate: 1) duplicate  $\delta^{18}O$   
246 analysis of quartz and calcite grains give the same isotopic value, suggesting grain isotopic homogeneity, 2) the stage of  
247 deformation that these mineral pairs are related to is not affected by further deformation in either outcrop or thin section, and  
248 3) all quartz-calcite pairs suggest a similar temperature of isotopic equilibration.



249 Temperature errors from quartz-calcite oxygen isotope measurements were calculated through the square-root of the  
250 summed quadratures of all sources of uncertainty (equations A2, A3). These uncertainties included  $\delta^{18}\text{O}$  value errors of quartz  
251 and calcite of  $\pm 0.1 \text{ ‰}$  ( $1 \sigma$ ) and  $\pm 0.04 \text{ ‰}$  ( $1 \sigma$ ), respectively, and errors associated with the Sharp and Kirschner (1994)  
252 quartz-calcite oxygen isotope fractionation calibration (A parameter).

#### 253 4.5 Electron probe measurements

254 Electron probe analyses were carried-out at ETHZ using a JEOL JXA-8230 Electron Probe Microanalyzer (EPMA).  
255 The EPMA is equipped with five wavelength-dispersive spectrometers. Epidote and pyroxene were analyzed for Si, Al, Na,  
256 Mg, Ca, Cr, K, Ti, Fe, and Mn on TAP (Si, Al), TAPH (Al, Ca), PETJ (Ca, Cr), PETL (K, Ti), and LIFH (Fe, Mn) crystals.  
257 Beam parameters included a 20 nA beam current, 10  $\mu\text{m}$  beam size, and a 15 keV accelerating voltage. All elements were  
258 measured for 30 s on peak and a mean atomic number background correction was applied. Primary calibration standards used  
259 included: albite (Si, Na), anorthite (Al, Ca), synthetic forsterite (Mg), chromite (Cr), microcline (K), synthetic rutile (Ti),  
260 synthetic fayalite (Fe), and synthetic pyrolusite (Mn). Mole fraction expressions from Franz and Liebscher (2004) were used  
261 to calculate epidote ( $X_{\text{ep}}$ ), clinozoisite ( $X_{\text{cz}}$ ), and tawmawite ( $X_{\text{taw}}$ ) compositions. Further information on mineral chemistry  
262 calculations is available in Supplementary Table S6. Garnets were analyzed for Al, Ca, Mn, Fe, Mg on TAP (Al), PETJ (Ca),  
263 LIFL (Mn), LIFH (Fe), and TAPH (Mg) crystals. Si was calculated stoichiometrically. X-ray maps were collected with a 50  
264 nA beam current, 15 keV accelerating voltage, 100 ms dwell time, and 5  $\mu\text{m}$  (KCS34 Garnet 1) and 4  $\mu\text{m}$  (KCS34 Garnet 3)  
265 step sizes. X-ray maps were reduced using CalcImage (Probe for EPMA).

#### 266 5. Thermobarometry Results

267 Determined pressures were categorized into three groups according to outcrop and microstructural context (Fig. 5;  
268 Fig. 7; Supplementary Table S3): garnet growth near peak metamorphic conditions (Group 1), growth of foliation-parallel  
269 epidote during blueschist-greenschist facies metamorphism (ep1, Group 2), and late-stage epidote growth in the new  
270 crenulation ( $S_{12}$ ) associated with  $F_{12}$  folds during greenschist facies metamorphism (ep2, Group 3). New ep2 growth is also  
271 supported by the mineral chemistry of different epidote generations within the  $S_{12}$  crenulation. Epidotes show a progressive  
272 chemical evolution that is recorded by an early generation epidote inclusion in titanite that occurs parallel to  $S_{12}$  ( $X_{\text{ep}} \cong 0.1$ ),  
273 the ep2 core ( $X_{\text{ep}} \cong 0.5$ ), and the ep2 rim ( $X_{\text{ep}} \cong 0.8$ ) (Fig. 2g; Supplementary Table S6).

274 The entrapment temperature ( $T_{\text{trap}}$ ) of quartz inclusions in garnet (garnet growth temperature) is estimated as 500 -  
275 550  $^{\circ}\text{C}$ ; this is based on good agreement between previous studies on the maximum temperature reached by CBU rocks from  
276 Syros (e.g., Laurent et al., 2018; Ridley, 1984; Schumacher et al., 2008; Skelton et al., 2018; Trotet et al., 2001a).  $T_{\text{trap}}$  for the  
277 ep2 population (Group 3) is deduced from oxygen isotope thermometry of quartz-calcite boudin-neck precipitates. The mean  
278 temperature from quartz-calcite pairs from boundin necks is  $411 \pm 23 \text{ }^{\circ}\text{C}$  ( $n = 4$ , Supplementary Table S5).  $T_{\text{trap}}$  for the ep1  
279 population (Group 2) is estimated as being intermediate between garnet and ep2 growth ( $\sim 400 - 500 \text{ }^{\circ}\text{C}$ ). As shown by qtz-in-

280 ep isomekes (constant  $P_{inc}$  lines along which fractional volume changes of an inclusion and host are equal), the assumed  $T_{trap}$   
281 has a minimal effect on  $P_{trap}$  (Fig. 7a; Cisneros et al., 2020).

## 282 **5.1 Kalamisia**

283 Group 1 quartz-inclusions-in-garnet record a mean  $P_{inc}$  of  $600 \pm 78$  MPa (Fig. 5; Supplementary Table S3). This  
284 corresponds to an entrapment pressure ( $P_{trap}$ ) of  $1.43 - 1.49 \pm 0.14$  GPa ( $n = 5$ ), at an estimated  $T_{trap}$  between  $500 - 550$  °C (Fig.  
285 7a, Supplementary Table S3). Group 2 quartz-inclusions-in-ep1 record a mean  $P_{inc}$  of  $544 \pm 57$  MPa, corresponding to a  $P_{trap}$   
286 of  $1.43 \pm 0.12$  GPa ( $n = 6$ ) at an estimated  $T_{trap}$  of  $450$  °C. No Group 3 epidotes are found within our analyzed section from  
287 Kalamisia.

## 288 **5.2 Delfini**

289 Group 1 records a mean  $P_{inc}$  of  $731 \pm 54$  MPa (Fig 5; Supplementary Table S3). This corresponds to a  $P_{trap}$  of  $1.66 -$   
290  $1.72 \pm 0.10$  GPa ( $n = 22$ ), at an estimated  $T_{trap}$  between  $500 - 550$  °C (Fig. 7a, Supplementary Table S3). Group 2 records a  
291 mean  $P_{inc}$  of  $518 \pm 52$  MPa, corresponding to a  $P_{trap}$  of  $1.38 \pm 0.11$  ( $n = 5$ ) at an estimated  $T_{trap}$  of  $450$  °C. Group 3 records a  
292 mean  $P_{inc}$  of  $343 \pm 23$  MPa, corresponding to a  $P_{trap}$  of  $0.98 \pm 0.05$  GPa ( $n = 3$ ) at  $411$  °C (Supplementary Table S3).

## 293 **5.3 Lotos**

294 Group 1 records a mean  $P_{inc}$  of  $751 \pm 76$  MPa (Fig 5; Supplementary Table S3). This corresponds to a  $P_{trap}$  of  $1.70 -$   
295  $1.76 \pm 0.14$  GPa ( $n = 2$ ), at an estimated  $T_{trap}$  between  $500 - 550$  °C (Fig. 7a; Supplementary Table S3). Group 2 records a mean  
296  $P_{inc}$  of  $531 \pm 78$  MPa, corresponding to a  $P_{trap}$  of  $1.41 \pm 0.17$  ( $n = 15$ ) at an estimated  $T_{trap}$  of  $450$  °C. No Group 3 epidotes were  
297 analyzed from Lotos.

## 298 **5.4 Megas Gialos**

299 Group 2 records an average  $P_{inc}$  of  $494 \pm 29$  MPa (Fig. 5), corresponding to a  $P_{trap}$  of  $1.33 \pm 0.03$  ( $n = 6$ ) at an estimated  
300  $T_{trap}$  of  $450$  °C (Fig. 7a; Supplementary Table S3). No Group 1 garnets or Group 3 epidotes were analyzed from Megas Gialos.

## 301 **6. Discussion**

### 302 **6.1 Elastic thermobarometry pressure groups**

303 Group 1 garnets either lack an internal foliation or contain a weak foliation that is defined by inclusions oblique to  
304 the  $S_s$  fabric, which indicates a previous stage of deformation (Fig. 2c; Supplementary Figs. S1, S2, S3). Garnets record similar  
305 pressures, regardless of the location of quartz inclusions (Fig. 6, Supplementary Table S3). Pyroxene inclusions within different  
306 garnet zones (core:  $X_{jd} \approx 0.84$ , rim:  $X_{jd} \approx 0.81$ ) also show no difference in composition, which is consistent with qtz-in-grt

307 barometry results (Delfini: KCS1621, Supplementary Table S6). Group 2 epidotes (ep1) overgrow garnets, are aligned parallel  
308 to the S<sub>s</sub> foliation but sometimes preserve an internal foliation that is oblique to S<sub>s</sub>, and lack late greenschist facies inclusions  
309 (Fig. 2f,g; Supplementary Figs. S1, S3, S4, S5). Group 3 epidotes (ep2, KCS34, Fig. 2c, d, e) are short in length, are aligned  
310 parallel to a late S<sub>t2</sub> crenulation, contain minimal quartz inclusions, and only record Group 3 pressures, independent of the  
311 position of quartz inclusions within epidotes.

312 Based on these observations, the Group 1 P<sub>trap</sub> estimates from the qtz-in-grt barometer record high-P conditions on  
313 Syros associated with prograde-to-peak garnet growth, and the Group 2 and 3 P<sub>trap</sub> estimates from the qtz-in-ep barometer  
314 record epidote growth during early blueschist-greenschist facies retrogression (ep1, D<sub>t1</sub>) and subsequent D<sub>t2</sub> deformation (ep2),  
315 respectively. We interpret the low-P epidote group (Group 3) to be associated with D<sub>t2</sub> folding, and best recorded in areas that  
316 experienced late greenschist facies mineral growth due to enhanced deformation and/or fluid influx during this stage of  
317 deformation (e.g., core of F<sub>t2</sub> fold).

## 318 6.2 Comparison of peak pressure constraints for the CBU on Syros and Sifnos

319 Based on qtz-in-grt measurements (Group 1), our P<sub>trap</sub> calculations suggest maximum P conditions of ~1.6 - 1.8 GPa  
320 were reached by the CBU on Syros. Garnets from metasedimentary and metavolcanic rocks record the statistically highest P<sub>trap</sub>  
321 (~1.5 - 1.8 GPa), whereas garnets from metamafic rocks (Kalamisia) record the lowest P<sub>trap</sub> (~1.3 - 1.6 GPa) (Fig. 7a). **Several**  
322 **observations support that the qtz-in-grt barometry results record max P conditions of the CBU on Syros: 1) quartz inclusion**  
323 **measurements across core-to-rims of garnets that show prograde growth (decreasing Mn), show no systematic change in P<sub>trap</sub>**  
324 **(Fig. 6), 2) max pressures from this study are equivalent to qtz-in-grt barometry results from prograde-to-peak eclogites and**  
325 **blueschists (non-retrogressed) from the CBU on Syros (Behr et al., 2018), 3) retrograde ep1 pressures, do not exceed those**  
326 **recorded by qtz-in-grt barometry, and 4) several studies from the CBU have used garnets to constrain max pressures, suggesting**  
327 **that garnets are suitable for constraining maximum pressures (e.g., Laurent et al., 2018; Dragovic et al., 2012, 2015; Groppo**  
328 **et al., 2009). We herein discuss our qtz-in-grt barometry results as max pressures constraints, but acknowledge that we may**  
329 **have missed high-P rims that have been found in other studies from the CBU on Syros (e.g., Laurent et al., 2018). We present**  
330 **a compilation of previous P-T constraints on CBU rocks from Syros and Sifnos, Greece, and discuss how our P<sub>trap</sub> constraints**  
331 **compare with previous studies.**

332 Elastic thermobarometry, mineral stability constraints, and multi-phase equilibrium modeling results from Sifnos  
333 CBU rocks suggest maximum P conditions of ~1.8 ± 0.1 GPa (Ashley et al., 2014), ~1.4 ± 0.2 GPa (Matthews and Schliestedt,  
334 1984), and ~2.0 - 2.2 GPa (Dragovic et al., 2012, 2015; Groppo et al., 2009; Trotet et al., 2001a), respectively. **Elastic**  
335 **thermobarometry (Ashley et al., 2014) and garnet modelling results (Dragovic et al., 2012, 2015; Groppo et al., 2009) from**  
336 **Sifnos, suggest near isobaric conditions during garnet growth.** The results of Ashley et al. (2014) are commonly cited as  
337 evidence that the CBU reached high pressure conditions (≥ 2.0 GPa, from elastic thermobarometry); however, their P<sub>trap</sub>  
338 calculations were carried out by using fits to quartz molar volume (P-T-V) data that have recently been re-evaluated (Angel et

339 al., 2017a) . Improved fits to quartz molar volume experiments “soften” quartz, and remodeling  $P_{inc}$  values from Ashley et al.  
340 (2014) reduces maximum mean  $P_{trap}$  conditions to  $\sim 1.6 \pm 0.1$  GPa (Fig. 7b, Supplementary Table S7).

341 Elastic thermobarometry, mineral stability constraints, glaucophane-bearing marble mineral equilibria, and multi-  
342 phase equilibria modeling results from Syros CBU rocks suggest peak pressure conditions of  $\sim 1.5 \pm 0.1$  GPa (Behr et al.,  
343 2018),  $\sim 1.4 - 1.9$  GPa (Ridley, 1984),  $\sim 1.5$  GPa (Schumacher et al., 2008), and  $\sim 1.9 - 2.4$  GPa (Laurent et al., 2018; Skelton  
344 et al., 2018; Trotet et al., 2001a), respectively. Elastic thermobarometry results from **prograde-to-peak eclogites and blueschists**  
345 **from** Syros, Greece were reduced using the approach outlined in Ashley et al. (2016), wherein a correction to  $P_{trap}$  is applied  
346 based on the assumed  $T_{trap}$ . Recent studies suggest that not using a temperature-dependent  $P_{trap}$  correction produces suitable  
347 results that accurately reproduce experimental conditions of quartz entrapment by garnet (Bonazzi et al., 2019; Thomas and  
348 Spear, 2018). Recalculation of the Behr et al. (2018)  $P_{inc}$  data (no temperature-dependent  $P_{trap}$  correction) results in a mean  $P_{trap}$   
349 of  $\sim 1.7 \pm 0.1$  GPa (Fig. 7b, Supplementary Table S8). The re-evaluation of data from Ashley et al. (2014) and Behr et al.  
350 (2018) suggests that our results are in good agreement with previous elastic thermobarometry constraints, and that to date, no  
351 qtz-in-grt elastic thermobarometry results suggest pressures  $\geq 2.0$  GPa.

352 Different methodologies applied to CBU rocks from Syros have resulted in a wide range of maximum P estimates.  
353 Schumacher et al. (2008) used mineral-equilibria modeling of glaucophane-bearing marbles to place constraints on maximum  
354 P-T conditions. Maximum P-T conditions are constrained by the presence of glaucophane +  $CaCO_3$  + dolomite + quartz,  
355 which suggests that the marbles exceeded the albite/Na-pyroxene + dolomite + quartz  $\rightarrow$  glaucophane +  $CaCO_3$  reaction, but  
356 did not cross the dolomite + quartz  $\rightarrow$  tremolite +  $CaCO_3$  or the glaucophane + aragonite-out reactions. The mineral reaction  
357 constraints suggest maximum P-T conditions of  $\sim 1.5 - 1.6$  GPa and  $500^\circ C$  for the CBU marbles. Ridley (1984) used the  
358 stability of paragonite and lack of kyanite to deduce max P constraints of  $\sim 1.4 - 1.9$  GPa. Trotet et al. (2001b, 2001a), Laurent  
359 et al. (2018), and Skelton et al. (2018) employed thermodynamic phase-equilibria modeling and supplementary methods to  
360 constrain P-T conditions for CBU rocks from Syros. Skelton et al. (2018) used the Powell and Holland (1994) Thermocalc  
361 database, Trotet et al. (2001b, 2001a) used the Berman (1991) thermodynamic database and the TWEEQC approach, and  
362 Laurent et al. (2018) used empirical thermobarometry, GrtMod (Lanari et al., 2017), and isochemical phase diagrams. Trotet  
363 et al. (2001b, 2001a), Laurent et al. (2018), and Skelton et al. (2018) found high-P conditions for the CBU ( $\geq 1.9$  GPa), and  
364 results from Laurent et al. (2018) suggest some rocks reached conditions as high as  $2.2 \pm 0.2$  GPa. Results from Laurent et al.  
365 (2018) suggest most garnet growth occurred at  $\sim 1.7$  GPa and  $450 \pm 50^\circ C$ ; however, some garnet modeling results suggest that  
366 garnet rims grew at  $\sim 2.4$  GPa and  $500 - 550^\circ C$ , albeit errors are increasingly large for these results ( $\pm 0.4 - 0.9$  GPa). These  
367 errors reflect the spacing between garnet isopleths (optimal P-T conditions), that result from uncertainties in chemical analyses.

368 Some GrtMod results suggest prograde core and rim garnet growth at  $\sim 1.8$  GPa and  $475^\circ C$ , and  $\sim 2.4$  GPa and  $475^\circ C$ ,  
369 respectively (sample SY1418 from; Laurent et al., 2018); however, the optimal P-T conditions for garnet rims have large  
370 errors and plot within uncertainty of garnet core conditions. Garnet results from another sample (SY1401) suggest core and  
371 rim garnet growth at  $\sim 1.8$  GPa and  $475^\circ C$ , and  $\sim 2.4$  GPa and  $550^\circ C$ , respectively. Sample SY1401 is collected from the same  
372 locality as ours (Kalamisia), but our qtz-in-grt results from this study suggest that garnets from this outcrop record the

373 statistically lowest  $P_{\text{trap}}$ . It is possible, however, that we did not sample the same rocks as Laurent et al. (2018), or that we have  
374 not found or analyzed garnets that record high pressures.

375 Previous studies have also suggested that pressures  $\geq 2.0$  GPa are unreasonable for Syros because paragonite is  
376 abundant in CBU rocks, but kyanite has not been reported. This suggests that CBU rocks did not cross the reaction paragonite  
377  $\rightarrow$  jadeite<sub>50</sub> + kyanite + H<sub>2</sub>O (~1.9 - 2.0 GPa); however, we recognize that the occurrence of kyanite may require high  
378 Al<sub>2</sub>O<sub>3</sub>:SiO<sub>2</sub> ratios for metabasites (e.g., Liati and Seidel, 1996), and that the pressure of this reaction is compositionally  
379 dependent. Pseudosections of eclogite CBU rocks show that kyanite would not be found in these bulk compositions below  
380 ~2.3 GPa (Skelton et al., 2018). It is possible that the high-P conditions found in previous studies may be real, but may only  
381 be recorded locally within some eclogite blocks.

382 In general, phase stability relationships (e.g., Matthews and Schliestedt, 1984; Ridley, 1984; Schumacher et al., 2008)  
383 and qtz-in-grt barometry results are in good agreement, but do not agree with high-pressure results ( $\geq 1.9$  GPa) deduced from  
384 thermodynamic modeling using approaches such as GrtMod and TWEEQC. The difference between our results and those of  
385 previous studies is important to reconcile, because the maximum P conditions reached by the CBU has considerable  
386 implications for the internal architecture of the CBU, its geodynamic evolution, and the mechanisms that can accommodate  
387 exhumation mechanisms of high-P subduction zone rocks from Syros. A comparison of qtz-in-grt barometry with  
388 thermodynamic modeling results from samples that record high pressures would be appropriate for further testing differences  
389 between the two techniques.

### 390 **6.3 Comparison of exhumation P-T conditions**

391 Previous studies have presented varying P-T paths and associated exhumation histories for Syros CBU rocks (Fig.  
392 7a; Laurent et al., 2018; Schumacher et al., 2008; Skelton et al., 2018; Trotet et al., 2001a). We present a compilation of  
393 previous P-T constraints and interpretations and discuss how our results compare with previous studies.

394 Schumacher et al. (2008) do not provide quantitative constraints for the retrograde P-T path (schematic), and samples  
395 do not have structural context; however, the authors suggest that a “cold” P-T path during exhumation is required for Syros  
396 CBU rocks based on the occurrence of lawsonite + epidote assemblages across Syros, and the P-T path required to avoid  
397 crossing the lawsonite  $\rightarrow$  kyanite + zoisite reaction (Fig. 7b). The authors suggest that exhumation of CBU packages occurred  
398 shortly after juxtaposition near peak metamorphic conditions.

399 Both Trotet et al. (2001a, 2001b) and Laurent et al. (2018) constrain high-P conditions for the CBU ( $> 2.0$  GPa),  
400 however, their proposed exhumation histories differ. Trotet et al. (2001b) suggested that CBU eclogites, blueschists and  
401 greenschists underwent different T-t histories during exhumation and were juxtaposed late along ductile shear zones. Laurent  
402 et al. (2018) suggested that the entire CBU reached peak metamorphic conditions of ~2.2 GPa, and that units that preserved  
403 blueschist facies assemblages underwent cooling during decompression, whereas rocks of southern Syros from lower structural  
404 levels experienced isobaric heating (~550 °C) at mid-crustal depths (~1.0 GPa) followed by subsequent cooling. Laurent et al.

405 (2018) interpreted reheating to indicate that CBU rocks on Syros reached high-P conditions, and then transitioned from a  
406 forearc to back-arc setting at  $\sim 1.0$  GPa, thus experiencing a period of increasing temperatures.

407 Skelton et al. (2018) also estimated peak and exhumation P-T conditions of rocks from Fabrikas (southern Syros),  
408 and interpreted exhumation of the CBU within an extrusion wedge (Ring et al., 2020). The authors constrained maximum P-  
409 T conditions of  $\sim 1.9$  GPa and  $540$  °C, and retrograde conditions of  $\sim 1.4 - 1.6$  GPa and  $510 - 520$  °C (blueschist facies) and  
410  $\sim 0.3$  GPa and  $450$  °C (greenschist facies) based on Thermocalc end-member activity modeling (Powell and Holland, 1994).  
411 Retrograde blueschist conditions (inferred from garnet growth) are similar between their estimates and ours, but greenschist  
412 facies conditions vastly differ. However, Skelton et al. (2018) focused on greenschist facies outcrops wherein metamorphism  
413 occurred locally over short length scales (e.g.  $\sim 10 - 100$  m), adjacent to late-stage brittle normal faults. We interpret our  $D_{12}$   
414 stage of greenschist facies metamorphism to pre-date late-stage normal faulting that has been attributed to Neogene block  
415 rotations (Cooperdock and Stockli, 2016) or possible coeval granitoid magmatism during Miocene back-arc extension (Keiter  
416 et al., 2011).

417 Gyomlai et al. (2021) estimate max and retrograde P-T conditions, but from metasomatic rocks from the Kampos  
418 belt in northern Syros. The authors estimated maximum T conditions of  $561 \pm 78$  °C, and two retrograde pressure-temperature  
419 conditions:  $1.02 \pm 0.15$  GPa and  $505 \pm 155$  °C, and  $1.03 \pm 0.11$  GPa and  $653 \pm 27$  °C. The retrograde pressures are reasonable  
420 ( $\sim 1.0 \pm 0.1 - 0.2$  GPa), but the max temperatures raise questions that the authors discuss. Specifically, temperatures above  
421  $\sim 600$  °C (at  $\sim 1.0$  GPa) would lead to serpentine breakdown (Guillot et al., 2015; Wunder and Schreyer, 1997); however,  
422 serpentine is abundant across Syros. The authors used the  $505 \pm 155$  °C temperature constraint, and a temperature below  $600$   
423 °C, to suggest their studied rocks reached temperatures between  $500 - 600$  °C at  $\sim 1.0$  GPa. Several other studies on retrograde  
424 metasomatic rocks from Kampos constrain P-T conditions:  $\sim 1.17 - 1.23$  GPa and  $500 - 550$  °C (Breeding et al., 2004),  $\sim 0.60$   
425  $- 0.75$  GPa and  $400 - 430$  °C (Marschall et al., 2006), and  $\sim 1.20$  GPa and  $430$  °C (Miller et al., 2009). Breeding et al. (2004)  
426 did not constrain a temperature, but used an estimated temperature from Trotet et al. (2001a), and constrained a pressure of  
427  $\sim 1.17 - 1.23$  GPa at the estimated T of  $\sim 500 - 550$  °C using Thermocalc V. 3.2. Marschall et al. (2006) used the garnet-  
428 clinopyroxene thermometer and Thermocalc V. 3.01 to calculate temperatures, and estimated a pressure based on jadeite +  
429  $\text{SiO}_2 \rightarrow \text{albite}$  reaction. Miller et al. (2009) used *Perple\_X* and the thermodynamic database of Holland and Powell (1998) to  
430 calculate P-T conditions from reaction zones. In general, most studies indicate cooling during decompression for metasomatic  
431 rocks from Kampos, with the exception of interpretations by Gyomlai et al. (2021); however, the large uncertainty of their  
432 temperature estimate ( $505 \pm 155$  °C) makes it difficult to differentiate between cooling during decompression, isothermal  
433 decompression, or re-heating.

434 Our results show that rocks from Kalamisia, Delfini, Lotos, and Megas Gialos, reached peak P-T conditions and  
435 underwent cooling during retrograde blueschist and greenschist facies metamorphism (Fig. 7a). Peak P-T conditions of the  
436 CBU are  $\sim 1.6 - 1.8$  GPa and  $500 - 550$  °C (Group 1 qtz-in-grt  $P_{\text{trap}}$  estimates), indicating a subduction zone geothermal gradient  
437 of  $\sim 9 - 10$  °C  $\text{km}^{-1}$  at  $\sim 55 - 60$  km (assuming  $30 \text{ MPa km}^{-1}$ ). Group 2 and 3 qtz-in-ep  $P_{\text{trap}}$  estimates indicate geothermal gradients  
438 of  $\sim 10$  °C  $\text{km}^{-1}$  and  $\sim 12$  °C  $\text{km}^{-1}$  at  $\sim 47$  and  $33$  km depths, respectively (Fig. 7a), demonstrating a similar P-T trajectory during

439 exhumation. We do not have a temperature constraint for the ep1 population; however, we consider cooling during  
440 decompression from garnet growth (~500 – 550 °C) to ep2 growth (~400 °C), to be the most likely P-T path for CBU rocks  
441 from Syros. Isothermal decompression from ~1.8 GPa and ~500 – 550 °C to ~ 1.0 GPa, would lead to terminal lawsonite  
442 breakdown above ~ 450 °C and produce kyanite + zoisite (Hamelin et al., 2018; Schumacher et al., 2008); however, kyanite  
443 has not been found on Syros, therefore requiring temperatures below ~450 °C at ~ 1.0 GPa. It is possible that sluggish kinetics  
444 did not lead to lawsonite breakdown, but given the prevalent evidence of retrograde deformation on Syros and the extensive  
445 presence of retrograde overprinting/mineral growth, we consider kinetic-limitations to be unlikely. Furthermore, the chemical  
446 evolution of amphiboles (magnesio-riebeckite → winchite → actinolite) suggests that CBU rocks from Syros followed a cold  
447 P-T path during decompression (c.f., Kotowski et al., 2020). Our P-T constraints are also inconsistent with reheating to ~550  
448 °C and 1.0 GPa, wherein amphibolite facies mineralogy may be stable. Our samples and the sample from which Laurent et al.  
449 (2018) determined reheating (SY1407), preserve no mineralogical evidence for having reached epidote-amphibolite facies  
450 (Fig. 7b; e.g., pargasite/hornblende, biotite/muscovite). Instead, the matrix mineralogy of sample SY1407 (glaucophane,  
451 phengite, rutile) suggests that these rocks formed under a cold geothermal gradient, rather than in a back-arc setting with an  
452 elevated geothermal gradient. Laurent et al. (2018) suggest that sample SY1407 records albite-epidote-blueschist conditions,  
453 a field metamorphic facies that can expand to higher T conditions; however, a pseudosection created for a similar bulk  
454 composition suggests that the determined P-T constraints (~1.0 GPa and 550 °C) are within epidote-amphibolite facies (Trotet  
455 et al., 2001a). Furthermore, results from sample SY1407 of Laurent et al. (2018) sometimes disagree when using local vs. bulk  
456 compositions for modeling. Models that use bulk compositions and consider Mn suggest that the core and mantle of the garnet  
457 record P-T conditions of ~1.8 GPa and 475 °C, whereas models that use local compositions or do not consider Mn suggest that  
458 the garnets do not record conditions above ~1.0 GPa (model residuals are lower using local bulk composition models).

459 Our results suggest that rocks from different Syros outcrops record similar peak and exhumation P-T conditions, but  
460 experienced different extents of deformation and thus recrystallization during exhumation. The similar peak pressures (> 0.8  
461 GPa) between different Syros outcrops suggests that these rocks belong to the Upper Cycladic Blueschist Nappe (Grasemann  
462 et al., 2018), even though in some cases significant retrogression overprinted indicators that would suggest these rocks reached  
463 P conditions above ~0.8 GPa. The observation of similar P-T conditions reached at different locations is inconsistent with  
464 results that suggest individual P-T paths for rocks that preserve different metamorphic facies (Trotet et al., 2001b, a), and  
465 different sections of the CBU (Laurent et al., 2018); however, we do not have T constraints for rocks from southern Syros. Our  
466 results are in better agreement with a P-T evolution resembling that of Schumacher et al. (2008), and a geothermal gradient of  
467 ~10 – 12 °C km<sup>-1</sup> that has also been proposed for CBU rocks from Sifnos, Greece (Schmädicke and Will, 2003).

#### 468 **6.4 Limitations of elastic thermobarometry**

469 Elastic thermobarometry has rapidly gained interest due to its limited dependence on mineral and fluid chemistry.  
470 Recent hydrostatic experiments that grow garnet around quartz have also shown the quartz-in-garnet barometer is accurate  
471 from ~ 0.8 – 3.0 GPa (± 0.1 - 0.2 GPa, Thomas and Spear, 2018; Bonazzi et al., 2019). The results suggest that the applied 1-

472 dimensional elastic model that assumes a spherical inclusion and isotropic inclusion-host pairs (Guiraud and Powell, 2006;  
473 Angel et al., 2017b), and the currently applied EoS' (Angel et al., 2017a; Holland and Powell, 2011; Milani et al., 2017),  
474 sufficiently replicate the elastic behaviour of an isotropic mineral (quartz) in a near isotropic host (garnet). Nonetheless,  
475 multiple secondary processes may affect quartz-in-garnet entrapment conditions: 1) mineral anisotropy (e.g., Murri et al.,  
476 2018), 2) inclusion shape effects (e.g., Cesare et al., 2021; Mazzucchelli et al., 2018), 3) relaxation adjacent to fractures or the  
477 host exterior, or overpressures adjacent to other inclusions (e.g., Zhong et al., 2020), 4) non-ideal tensile strain (e.g., Cisneros  
478 and Befus, 2020), or 5) non-elastic strain (i.e., viscous strain, e.g., Zhang, 1998). We propose that none of these processes have  
479 affected our quartz-in-garnet barometry results for the following reasons: 1)  $P_{inc}$  values calculated from different quartz bands  
480 and by accounting for anisotropy (strains) center around the hydrostatic stress lines (1:1 line, Fig. 5), and  $P_{inc}$  calculated from  
481 strains changes the final  $P_{trap}$  by  $< 0.2$  GPa (relative to  $P_{inc}$  calculated from the  $464\text{ cm}^{-1}$  band). 2) Near spherical quartz  
482 inclusions were analysed to minimize shape effects, and measurements were taken from the center of quartz inclusions to avoid  
483 stress effects at inclusion-host boundaries. 3) Quartz inclusions were a minimum two-to-three-times the radial distance away  
484 from fractures, cleavage, and the host exterior, or other inclusions to minimize under- or overpressures, respectively. 4) All  
485 quartz inclusions from this study exist under compression, thus tensile strain limits are not relevant. 5) The maximum estimated  
486 temperature of CBU rocks from Syros is  $\sim 500 - 550$  °C, and garnet flow laws predict that viscous creep of garnet occurs above  
487  $\sim 650$  °C at geologic strain rates (Wang and Ji, 2001; Ji and Martignole, 1994); therefore, viscous strain of garnet is unlikely  
488 to have occurred. Considering the current state-of-knowledge in elastic thermobarometry, we propose that our pressure results  
489 have been minimally influenced by secondary effects.

490 In contrast, the quartz-in-epidote barometer is less studied. Recent studies have explored the suitability of using an  
491 isotropic elastic model to model the elastic evolution of two anisotropic minerals (Cisneros et al., 2020). Results showed that  
492 an isotropic elastic model suitably simulates the pressure evolution of two anisotropic minerals during heating, and that the  
493 calculated entrapment pressures agree with independent thermobarometry constraints. However, it is unknown if isotropic  
494 elastic models correctly simulate the elastic evolution of anisocoric mineral pairs during compression, and additional processes  
495 may influence the entrapment pressures calculated from quartz-inclusions-in-epidote: 1) the orientation of quartz inclusions  
496 relative to the orientation of epidote, and 2) the material properties of epidote (i.e., at what conditions does viscous creep  
497 become important for epidote). 1) Cisneros et al. (2020) showed that the orientation of quartz inclusions relative to epidote  
498 may have had a minimal effect on the elastic evolution of quartz-epidote pairs, but the orientation of quartz and epidote were  
499 not determined. We hypothesize that in this study, the mutual orientation of quartz-epidote inclusion-host pairs had a minimal  
500 effect on the calculated entrapment pressures. If the mutual orientation of quartz-epidote pairs had a large effect, we expect  
501 that the  $P_{inc}$  calculated from different quartz-inclusions-in-epidote would exhibit significant scatter; however,  $P_{inc}$  values from  
502 different quartz-inclusions-in-epidote are similar, and  $P_{inc}$  values from different quartz bands and strains, center around the  
503 hydrostatic stress line (Fig. 5). The  $P_{inc}$  scatter from different quartz-inclusions-in-epidote (same ep population, e.g., ep2) and  
504 the  $P_{inc}$  variation from different quartz bands and strains, generally does not exceed that of quartz-inclusions-in-garnet. The  
505 minimal  $P_{inc}$  variation between quartz-inclusions-in-epidote from the same epidote population may result from the orientation



506 of quartz and epidote parallel to the primary foliation. The orientation of quartz-epidote pairs may lead to a bulk stress tensor  
507 that produces minimal orientation-dependent effects, or the lower bulk modulus of epidote (relative to garnet) may result in a  
508 small stress anisotropy. 2) No epidote flow law exists (to the best of our knowledge); therefore, the temperature at which  
509 viscous strain will be important for epidote is unknown. Nonetheless, in contrast to garnet (isotropic), evidence for viscous  
510 creep in epidote can be observed in thin section. In epidotes from this study, we have observed no thin-section scale evidence  
511 of dislocation creep; however,  $\mu\text{m}$ -scale viscous creep in epidote adjacent to quartz inclusions cannot be excluded.

## 512 **6.5 Implications for exhumation mechanisms**

513 Our results indicate that the CBU followed a “cooling during decompression” P-T trajectory that required a heat sink  
514 at depth to cool rocks during exhumation. Cooling could be achieved under a steady-state subduction zone thermal gradient  
515 with slab-top temperatures similar to those of warm subduction zones, such as in Cascadia (e.g., Syracuse et al., 2010;  
516 Walowski et al., 2015). This would suggest that exhumation was achieved parallel to the subducting plate, in a subduction  
517 channel geometry prior to core-complex formation. Results from this study cannot differentiate between extrusion wedge  
518 models (“extrusion” of a wedge of CBU rocks within a subduction channel) that require a kinematically necessary thrust fault  
519 at the base (the subducting slab) and a kinematically necessary normal fault at the top (upper plate), and other general  
520 subduction channel models (e.g., Ring et al., 2020). Subduction channel and extrusion models have slight differences, i.e., the  
521 extrusion wedge model calls for a specific geometry that should produce opposing shear sense indicators at distinct locations  
522 that define the base (subduction plate) and top (upper plate) of the wedge (within a subduction channel). A subduction channel  
523 model has a looser definition (without a specific geometric structure) that merely reflects the plate interface structure (discrete  
524 or broad interface), and does not require this deformation. Because we do not present sufficient kinematic information in this  
525 study to differentiate these models, we prefer to use a general “subduction channel” model nomenclature, to indicate that we  
526 interpret CBU rocks to have been exhumed parallel to the subducting plate, within a broad, viscous shear zone that defines the  
527 subduction interface.

528 During this phase of exhumation, CBU rocks remained within a cold forearc until they reached the mid-crust ( $\sim 1.0$   
529 GPa), and exhibit a progressive change in kinematics, from N-S stretching lineations during subduction (e.g., Behr et al., 2018;  
530 Laurent et al., 2016; Philippon et al., 2011), to lineations that swing towards the NE (this study, Roche et al., 2016; Sifnos)  
531 and E-W during exhumation (c.f., Kotowski and Behr, 2019; Laurent et al., 2016). We propose that N-S ( $D_s$ ) lineations  
532 (subduction-related) and exhumation-related upright folds that generate NE ( $D_{t1}$ ) and E-W ( $D_{t2}$ ) extension parallel to fold hinge  
533 lines, document the transition from subduction to exhumation as rocks turn the corner to be exhumed within the subduction  
534 channel. Stretching lineations in the footwall of the North and West Cycladic Detachment Systems have top-to-the- NE and  
535 SW orientations, respectively (e.g., Brichau et al., 2007; Grasemann et al., 2012; Jolivet et al., 2010; Mehl et al., 2005). The  
536 inferred P-T conditions and kinematics of our studied samples are consistent with Syros recording early deformation and  
537 metamorphism within a forearc setting, whereas adjacent Cycladic islands that border the North and West Cycladic  
538 Detachment Systems record late-stage kinematics and greenschist facies metamorphism that capture the CBU transition to a

539 warmer back-arc setting (e.g., Laurent et al., 2016; Ring et al., 2020; Roche et al., 2016; Schmädicke and Will, 2003). Our  
540 data suggests that during generation of exhumation-related upright folds ( $D_{11-12}$ ), rocks from the CBU on southern Syros (below  
541 the Kampos nappe) followed similar P-T conditions exhumation (Fig. 7). It is unclear if the upper Kampos nappe exhibited  
542 the same deformation because it preserves less structural coherency; however, rocks from Kampos and southern Syros seem  
543 to have experienced similar P-T conditions during exhumation. Rocks from different sections of the CBU may have reached  
544 peak P conditions at different times, and thus experienced the same exhumation-related deformation at different times  
545 (Kotowski et al., 2020); however, our data suggest that rocks from different sections of the southern CBU on Syros were  
546 exhumed within a forearc setting up to ~33 km depth. We propose that CBU on Syros may not record back-arc deformation  
547 until the Vari detachment accommodated exhumation of the CBU at ~ 10 – 8 Ma (~5 - 7 km depth, Soukis and Stockli, 2013).  
548 Back-arc related deformation occurs directly adjacent to the Vari detachment, as evidenced by semi-brittle to brittle cataclastic  
549 deformation (greenschist facies) that affects the Upper Unit and the underlying CBU (Soukis and Stockli, 2013).

## 550 7. Conclusions

551 This work highlights the potential of using elastic thermobarometry in combination with structural (macro and micro)  
552 and petrographic constraints, to better constrain P-T conditions of challenging rock assemblages. Our results allow us to place  
553 robust P-T constraints on distinct textural fabrics that are related to well-constrained outcrop scale structures. In particular, the  
554 work highlights how the qtz-in-ep barometer is well suited for constraining formation conditions of epidote, a common mineral  
555 that is found within a large range of geologic settings and P-T conditions. Combining the qtz-in-ep barometer with other elastic  
556 thermobarometers (e.g., qtz-in-grt) allows determination of protracted P-T histories from minerals that record different  
557 geologic stages within single rocks samples.

558 Our new results show that CBU rocks from Syros, Greece, experienced similar P-T conditions during subduction and  
559 exhumation, inconsistent with results that suggest different P-T histories for CBU rocks for Syros or increasing temperatures  
560 during exhumation. Our targeted stages of deformation and metamorphism suggest that CBU rocks from Syros record cooling  
561 during decompression, consistent with exhumation within a subduction channel and early deformation and metamorphism  
562 within a forearc (at least to ~33 km depth), prior to Miocene core-complex formation and transition to a warmer back-arc  
563 setting.

## 564 Appendix A: Stable isotope temperature error calculations

565 Temperature errors from oxygen isotope measurements were calculated through the square-root of the summed  
566 quadratures of all sources of uncertainty. These uncertainties included error of  $\delta^{18}\text{O}$  values of quartz (qtz) and calcite (cc) of  $\pm$   
567 0.1 ‰ (1  $\sigma$ ) and  $\pm$  0.04 ‰ (1  $\sigma$ ), respectively, and errors associated with the Sharp and Kirschner (1994) quartz-calcite oxygen

568 isotope fractionation calibration (A parameter). Errors from the sum of propagated analytical errors, were propagated through  
569 the empirical calibration of quartz-calcite oxygen isotope fraction that was used for temperature calculations:

$$570 \quad \Delta_{qtz-cc} = \frac{A \times 10^6}{T^2} \quad \#A1$$

571  
572 where  $A = 0.87 \pm 0.06$  (1  $\sigma$ ). The square-root of the summed quadratures is expressed as:

$$573 \quad \sigma_T = \sqrt{\sigma_A^2 \left(\frac{\partial T}{\partial A}\right)^2 + \sigma_{\Delta_{qtz-cc}}^2 \left(\frac{\partial T}{\partial \Delta_{qtz-cc}}\right)^2} \quad \#A2$$

$$574 \quad \sigma_T = \sqrt{\sigma_A^2 \left(\frac{0.5 * 10^3}{\sqrt{A} * \sqrt{\Delta_{qtz-cc}}}\right)^2 + \sigma_{\Delta_{qtz-cc}}^2 \left(-0.5 * \frac{\sqrt{A} * 10^3}{\Delta_{qtz-cc}^{1.5}}\right)^2} \quad \#A3$$

## 576 **Author Contribution**

577 All authors contributed to this manuscript. M. Cisneros developed the epidote barometer, collected the data, and wrote  
578 the manuscript. J. Barnes, W. Behr, A. Kotowski, D. Stockli, and K. Soukis helped with conceiving the project, field work,  
579 and writing.

## 580 **Acknowledgements**

581 We thank J. Schumacher and V. Laurent for constructive reviews that helped improve this manuscript, and F. Rossetti  
582 for editorial handling and additional comments that helped improve this manuscript. We thank N. Raia for field work  
583 assistance, J. Allaz for assistance on the microprobe at ETH Zürich, and C. Farley and R. Bobnar for access to the Raman  
584 Spectrometer at Virginia Tech. This work was supported by a GSA Student Research Grant and a Ford Foundation Fellowship  
585 awarded to M.C., an NSF Graduate Research Fellowship awarded to A.K., and NSF Grant (EAR-1725110) awarded to J.B.,  
586 W.B., and D.S.

## 587 **References**

588 Adams, H. G., Cohen, L. H., and Rosenfeld, J. L.: Solid inclusion piezothermometry I: comparison dilatometry, 60, 574–583,  
589 1975a.  
590 Adams, H. G., Cohen, L. H., and Rosenfeld, J. L.: Solid inclusion piezothermometry II: geometric basis, calibration for the  
591 association quartz-garnet, and application to some pelitic schists, 60, 584–598, 1975b.

592 Alvaro, M., Mazzucchelli, M. L., Angel, R. J., Murri, M., Campomenosi, N., Scambelluri, M., Nestola, F., Korsakov, A.,  
593 Tomilenko, A. A., Marone, F., and Morana, M.: Fossil subduction recorded by quartz from the coesite stability field, *Geology*,  
594 48, 24–28, <https://doi.org/10.1130/G46617.1>, 2020.

595 Angel, R. J., Alvaro, M., Miletich, R., and Nestola, F.: A simple and generalised P-T-V EoS for continuous phase transitions,  
596 implemented in EosFit and applied to quartz, *Contrib Mineral Petrol*, 172, 29, <https://doi.org/10.1007/s00410-017-1349-x>,  
597 2017a.

598 Angel, R. J., Mazzucchelli, M. L., Alvaro, M., and Nestola, F.: EosFit-Pinc: A simple GUI for host-inclusion elastic  
599 thermobarometry, 102, 1957–1960, <http://dx.doi.org/10.2138/am-2017-6190>, 2017b.

600 Angel, R. J., Murri, M., Mihailova, B., and Alvaro, M.: Stress, strain and Raman shifts, 234, 129–140,  
601 <https://doi.org/10.1515/zkri-2018-2112>, 2019.

602 Ashley, K. T., Caddick, M. J., Steele-MacInnis, M. J., Bodnar, R. J., and Dragovic, B.: Geothermobarometric history of  
603 subduction recorded by quartz inclusions in garnet, *Geochemistry, Geophysics, Geosystems*, 15, 350–360,  
604 <https://doi.org/10.1002/2013GC005106>, 2014.

605 Ashley, K. T., Steele-MacInnis, M., Bodnar, R. J., and Darling, R. S.: Quartz-in-garnet inclusion barometry under fire:  
606 Reducing uncertainty from model estimates, *Geology*, 44, 699–702, <https://doi.org/10.1130/G38211.1>, 2016.

607 Augier, R., Jolivet, L., Gadenne, L., Lahfid, A., and Driussi, O.: Exhumation kinematics of the Cycladic Blueschists unit and  
608 back-arc extension, insight from the Southern Cyclades (Sikinos and Folegandros Islands, Greece), 34, 152–185,  
609 <https://doi.org/10.1002/2014TC003664>, 2015.

610 Avigad, D.: High-pressure metamorphism and cooling on SE Naxos (Cyclades, Greece), *European Journal of Mineralogy*, 10,  
611 1309–1319, 1998.

612 Baxter, E. F.: Natural constraints on metamorphic reaction rates, 220, 183–202,  
613 <https://doi.org/10.1144/GSL.SP.2003.220.01.11>, 2003.

614 Behr, W. M., Kotowski, A. J., and Ashley, K. T.: Dehydration-induced rheological heterogeneity and the deep tremor source  
615 in warm subduction zones, *Geology*, 46, 475–478, <https://doi.org/10.1130/G40105.1>, 2018.

616 Berman, R. G.: Thermobarometry using multi-equilibrium calculations; a new technique, with petrological applications, *The*  
617 *Canadian Mineralogist*, 29, 833–855, 1991.

618 Bonazzi, M., Tumiati, S., Thomas, J., Angel, R. J., and Alvaro, M.: Assessment of the reliability of elastic geobarometry with  
619 quartz inclusions, *Lithos*, 105201, <https://doi.org/10.1016/j.lithos.2019.105201>, 2019.

620 Breeding, C. M., Ague, J. J., and Bröcker, M.: Fluid–metasedimentary rock interactions in subduction-zone mélangé:  
621 Implications for the chemical composition of arc magmas, *Geology*, 32, 1041–1044, <https://doi.org/10.1130/G20877.1>, 2004.

622 Brichau, S., Ring, U., Carter, A., Monié, P., Bolhar, R., Stockli, D., and Brunel, M.: Extensional faulting on Tinos Island,  
623 Aegean Sea, Greece: How many detachments?, 26, <https://doi.org/10.1029/2006TC001969>, 2007.

624 Bröcker, M. and Franz, L.: Dating metamorphism and tectonic juxtaposition on Andros Island (Cyclades, Greece): results of  
625 a Rb–Sr study, 143, 609–620, <https://doi.org/10.1017/S001675680600241X>, 2006.

626 Bröcker, M., Kreuzer, H., Matthews, A., and Okrusch, M.: 40Ar/39Ar and oxygen isotope studies of polymetamorphism from  
627 Tinos Island, Cycladic blueschist belt, Greece, 11, 223–240, <https://doi.org/10.1111/j.1525-1314.1993.tb00144.x>, 1993.

628 Bröcker, M., Baldwin, S., and Arkudas, R.: The geological significance of 40Ar/39Ar and Rb–Sr white mica ages from Syros  
629 and Sifnos, Greece: a record of continuous (re)crystallization during exhumation?, 31, 629–646,  
630 <https://doi.org/10.1111/jmg.12037>, 2013.

631 Campomenosi, N., Mazzucchelli, M. L., Mihailova, B., Scambelluri, M., Angel, R. J., Nestola, F., Reali, A., and Alvaro, M.:  
632 How geometry and anisotropy affect residual strain in host-inclusion systems: Coupling experimental and numerical  
633 approaches, 103, 2032–2035, <https://doi.org/10.2138/am-2018-6700CCBY>, 2018.

634 Carlson, W. D.: Scales of disequilibrium and rates of equilibration during metamorphism, *American Mineralogist*, 87, 185–  
635 204, <https://doi.org/10.2138/am-2002-2-301>, 2002.

636 Cesare, B., Parisatto, M., Mancini, L., Peruzzo, L., Franceschi, M., Tacchetto, T., Reddy, S., Spiess, R., Nestola, F., and  
637 Marone, F.: Mineral inclusions are not immutable: Evidence of post-entrapment thermally-induced shape change of quartz in  
638 garnet, *Earth and Planetary Science Letters*, 555, 116708, <https://doi.org/10.1016/j.epsl.2020.116708>, 2021.

639 Cisneros, M. and Befus, K. S.: Applications and Limitations of Elastic Thermobarometry: Insights From Elastic Modeling of  
640 Inclusion-Host Pairs and Example Case Studies, 21, e2020GC009231, <https://doi.org/10.1029/2020GC009231>, 2020.

641 Cisneros, M., Ashley, K. T., and Bodnar, R. J.: Evaluation and application of the quartz-inclusions-in-epidote mineral  
642 barometer, *American Mineralogist*, 105, 1140–1151, <https://doi.org/10.2138/am-2020-7379>, 2020.

643 Cliff, R. A., Bond, C. E., Butler, R. W. H., and Dixon, J. E.: Geochronological challenges posed by continuously developing  
644 tectonometamorphic systems: insights from Rb–Sr mica ages from the Cycladic Blueschist Belt, Syros (Greece), 35, 197–211,  
645 <https://doi.org/10.1111/jmg.12228>, 2017.

646 Cooperdock, E. H. G. and Stockli, D. F.: Unraveling alteration histories in serpentinites and associated ultramafic rocks with  
647 magnetite (U-Th)/He geochronology, *Geology*, 44, 967–970, <https://doi.org/10.1130/G38587.1>, 2016.

648 Dragovic, B., Samanta, L. M., Baxter, E. F., and Selverstone, J.: Using garnet to constrain the duration and rate of water-  
649 releasing metamorphic reactions during subduction: An example from Sifnos, Greece, 314–317, 9–22,  
650 <https://doi.org/10.1016/j.chemgeo.2012.04.016>, 2012.

651 Dragovic, B., Baxter, E. F., and Caddick, M. J.: Pulsed dehydration and garnet growth during subduction revealed by zoned  
652 garnet geochronology and thermodynamic modeling, Sifnos, Greece, *Earth and Planetary Science Letters*, 413, 111–122,  
653 <https://doi.org/10.1016/j.epsl.2014.12.024>, 2015.

654 Enami, M., Nishiyama, T., and Mouri, T.: Laser Raman microspectrometry of metamorphic quartz: A simple method for  
655 comparison of metamorphic pressures, *American Mineralogist*, 92, 1303–1315, <https://doi.org/10.2138/am.2007.2438>, 2007.

656 Essene, E. J.: The current status of thermobarometry in metamorphic rocks, 43, 1–44,  
657 <https://doi.org/10.1144/GSL.SP.1989.043.01.02>, 1989.

658 Franz, G. and Liebscher, A.: Physical and Chemical Properties of the Epidote Minerals—An Introduction—, *Reviews in  
659 Mineralogy and Geochemistry*, 56, 1–81, <https://doi.org/10.2138/gsrmg.56.1.1>, 2004.

660 Gatta, G. D., Merlini, M., Lee, Y., and Poli, S.: Behavior of epidote at high pressure and high temperature: a powder diffraction  
661 study up to 10 GPa and 1,200 K, *Phys Chem Minerals*, 38, 419–428, <https://doi.org/10.1007/s00269-010-0415-y>, 2011.

662 Gautier, P., Brun, J.-P., and Jolivet, L.: Structure and kinematics of Upper Cenozoic extensional detachment on Naxos and  
663 Paros (Cyclades Islands, Greece), 12, 1180–1194, <https://doi.org/10.1029/93TC01131>, 1993.

664 Grasemann, B., Schneider, D. A., Stöckli, D. F., and Iglseider, C.: Miocene bivergent crustal extension in the Aegean: Evidence  
665 from the western Cyclades (Greece), *Lithosphere*, 4, 23–39, <https://doi.org/10.1130/L164.1>, 2012.

666 Grasemann, B., Huet, B., Schneider, D. A., Rice, A. H. N., Lemonnier, N., and Tschegg, C.: Miocene postorogenic extension  
667 of the Eocene synorogenic imbricated Hellenic subduction channel: New constraints from Milos (Cyclades, Greece), *GSA*  
668 *Bulletin*, 130, 238–262, <https://doi.org/10.1130/B31731.1>, 2018.

669 Groppo, C., Forster, M., Lister, G., and Compagnoni, R.: Glaucofane schists and associated rocks from Sifnos (Cyclades,  
670 Greece): New constraints on the P–T evolution from oxidized systems, *Lithos*, 109, 254–273,  
671 <https://doi.org/10.1016/j.lithos.2008.10.005>, 2009.

672 Guillot, S., Schwartz, S., Reynard, B., Agard, P., and Prigent, C.: Tectonic significance of serpentinites, *Tectonophysics*, 646,  
673 1–19, <https://doi.org/10.1016/j.tecto.2015.01.020>, 2015.

674 Guiraud, M. and Powell, R.: P–V–T relationships and mineral equilibria in inclusions in minerals, *Earth and Planetary Science*  
675 *Letters*, 244, 683–694, <https://doi.org/10.1016/j.epsl.2006.02.021>, 2006.

676 Gupta, S. and Bickle, M. J.: Ductile shearing, hydrous fluid channelling and high-pressure metamorphism along the basement-  
677 cover contact on Sikinos, Cyclades, Greece, 224, 161–175, <https://doi.org/10.1144/GSL.SP.2004.224.01.11>, 2004.

678 Gyomlai, T., Agard, P., Marschall, H. R., Jolivet, L., and Gerdes, A.: Metasomatism and deformation of block-in-matrix  
679 structures in Syros: The role of inheritance and fluid-rock interactions along the subduction interface, *Lithos*, 386–387, 105996,  
680 <https://doi.org/10.1016/j.lithos.2021.105996>, 2021.

681 Hamelin, C., Brady, J. B., Cheney, J. T., Schumacher, J. C., Able, L. M., and Sperry, A. J.: Pseudomorphs after lawsonite from  
682 Syros, Greece, <https://doi.org/10.1093/petrology/egy099>, 2018.

683 Holland, T. and Powell, R.: An internally consistent thermodynamic data set for phases of petrological interest, 16, 309–343,  
684 <https://doi.org/10.1111/j.1525-1314.1998.00140.x>, 1998.

685 Holland, T. and Powell, R.: An improved and extended internally consistent thermodynamic dataset for phases of petrological  
686 interest, involving a new equation of state for solids, 29, 333–383, <https://doi.org/10.1111/j.1525-1314.2010.00923.x>, 2011.

687 Isaak, D. G., Anderson, O. L., and Oda, H.: High-temperature thermal expansion and elasticity of calcium-rich garnets, *Phys*  
688 *Chem Minerals*, 19, 106–120, <https://doi.org/10.1007/BF00198608>, 1992.

689 Jamtveit, B., Austrheim, H., and Putnis, A.: Disequilibrium metamorphism of stressed lithosphere, *Earth-Science Reviews*,  
690 154, 1–13, <https://doi.org/10.1016/j.earscirev.2015.12.002>, 2016.

691 Javoy, M.: Stable isotopes and geothermometry, 133, 609–636, <https://doi.org/10.1144/gsjgs.133.6.0609>, 1977.

692 Ji, S. and Martignole, J.: Ductility of garnet as an indicator of extremely high temperature deformation, *Journal of Structural*  
693 *Geology*, 16, 985–996, [https://doi.org/10.1016/0191-8141\(94\)90080-9](https://doi.org/10.1016/0191-8141(94)90080-9), 1994.

694 Jolivet, L. and Brun, J.-P.: Cenozoic geodynamic evolution of the Aegean, *Int J Earth Sci (Geol Rundsch)*, 99, 109–138,  
695 <https://doi.org/10.1007/s00531-008-0366-4>, 2010.

696 Jolivet, L., Lecomte, E., Huet, B., Denèle, Y., Lacombe, O., Labrousse, L., Le Pourhiet, L., and Mehl, C.: The North Cycladic  
697 Detachment System, *Earth and Planetary Science Letters*, 289, 87–104, <https://doi.org/10.1016/j.epsl.2009.10.032>, 2010.

698 Keiter, M., Ballhaus, C., and Tomaschek, F.: A new geological map of the Island of Syros (Aegean Sea, Greece): implications  
699 for lithostratigraphy and structural history of the Cycladic Blueschist Unit, Geological Society of America, 2011.

700 Kohn, M. J. and Spear, F.: Retrograde net transfer reaction insurance for pressure-temperature estimates, *Geology*, 28, 1127–  
701 1130, [https://doi.org/10.1130/0091-7613\(2000\)28<1127:RNTRIF>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<1127:RNTRIF>2.0.CO;2), 2000.

702 Kotowski, A. J. and Behr, W. M.: Length scales and types of heterogeneities along the deep subduction interface: Insights  
703 from exhumed rocks on Syros Island, Greece, *Geosphere*, 15, 1038–1065, <https://doi.org/10.1130/GES02037.1>, 2019.

704 Kotowski, A. J., Behr, W. M., Cisneros, M., Stockli, D. F., Soukis, K., Barnes, J. D., and Ortega-Arroyo, D.: Subduction,  
705 underplating, and return flow recorded in the Cycladic Blueschist Unit exposed on Syros Island, Greece,  
706 <http://dx.doi.org/10.1002/essoar.10504307.1>, 2020.

707 Lagos, M., Scherer, E. E., Tomaschek, F., Münker, C., Keiter, M., Berndt, J., and Ballhaus, C.: High precision Lu–Hf  
708 geochronology of Eocene eclogite-facies rocks from Syros, Cyclades, Greece, *Chemical Geology*, 243, 16–35,  
709 <https://doi.org/10.1016/j.chemgeo.2007.04.008>, 2007.

710 Lamont, T. N., Searle, M. P., Gopon, P., Roberts, N. M. W., Wade, J., Palin, R. M., and Waters, D. J.: The Cycladic Blueschist  
711 Unit on Tinos, Greece: Cold NE Subduction and SW Directed Extrusion of the Cycladic Continental Margin Under the  
712 Tsiknias Ophiolite, 39, e2019TC005890, <https://doi.org/10.1029/2019TC005890>, 2020.

713 Lanari, P., Giuntoli, F., Loury, C., Burn, M., and Engi, M.: An inverse modeling approach to obtain P–T conditions of  
714 metamorphic stages involving garnet growth and resorption, *European Journal of Mineralogy*, 29, 181–199,  
715 <https://doi.org/10.1127/ejm/2017/0029-2597>, 2017.

716 Laurent, V., Jolivet, L., Roche, V., Augier, R., Scaillet, S., and Cardello, G. L.: Strain localization in a fossilized subduction  
717 channel: Insights from the Cycladic Blueschist Unit (Syros, Greece), 672–673, 150–169,  
718 <https://doi.org/10.1016/j.tecto.2016.01.036>, 2016.

719 Laurent, V., Huet, B., Labrousse, L., Jolivet, L., Monié, P., and Augier, R.: Extraneous argon in high-pressure metamorphic  
720 rocks: Distribution, origin and transport in the Cycladic Blueschist Unit (Greece), *Lithos*, 272–273, 315–335,  
721 <https://doi.org/10.1016/j.lithos.2016.12.013>, 2017.

722 Laurent, V., Lanari, P., Naïr, I., Augier, R., Lahfid, A., and Jolivet, L.: Exhumation of eclogite and blueschist (Cyclades,  
723 Greece): Pressure–temperature evolution determined by thermobarometry and garnet equilibrium modelling, *Journal of*  
724 *Metamorphic Geology*, 36, 769–798, <https://doi.org/10.1111/jmg.12309>, 2018.

725 Liati, A. and Seidel, E.: Metamorphic evolution and geochemistry of kyanite eclogites in central Rhodope, northern Greece,  
726 *Contrib Mineral Petrol*, 123, 293–307, <https://doi.org/10.1007/s004100050157>, 1996.

727 Lister, G. S. and Forster, M. A.: White mica  $^{40}\text{Ar}/^{39}\text{Ar}$  age spectra and the timing of multiple episodes of high-P metamorphic  
728 mineral growth in the Cycladic eclogite–blueschist belt, Syros, Aegean Sea, Greece, *Journal of Metamorphic Geology*, 34,  
729 401–421, <https://doi.org/10.1111/jmg.12178>, 2016.

730 Marschall, H. R.: Syros Metasomatic Tourmaline: Evidence for Very High- 11B Fluids in Subduction Zones, 47, 1915–1942,  
731 <https://doi.org/10.1093/petrology/egl031>, 2006.

732 Marschall, H. R., Ludwig, T., Altherr, R., Kalt, A., and Tonarini, S.: Syros Metasomatic Tourmaline: Evidence for Very High-  
733  $\delta^{11}\text{B}$  Fluids in Subduction Zones, *Journal of Petrology*, 47, 1915–1942, <https://doi.org/10.1093/petrology/egl031>, 2006.

734 Matthews, A. and Schliestedt, M.: Evolution of the blueschist and greenschist facies rocks of Sifnos, Cyclades, Greece, *Contr.*  
735 *Mineral. and Petrol.*, 88, 150–163, <https://doi.org/10.1007/BF00371419>, 1984.

736 Mazzucchelli, M. L., Burnley, P., Angel, R. J., Morganti, S., Domeneghetti, M. C., Nestola, F., and Alvaro, M.: Elastic  
737 geothermobarometry: Corrections for the geometry of the host-inclusion system, *Geology*, 46, 231–234,  
738 <https://doi.org/10.1130/G39807.1>, 2018.

739 Mehl, C., Jolivet, L., and Lacombe, O.: From ductile to brittle: Evolution and localization of deformation below a crustal  
740 detachment (Tinos, Cyclades, Greece), 24, <https://doi.org/10.1029/2004TC001767>, 2005.

741 Milani, S., Angel, R. J., Scandolo, L., Mazzucchelli, M. L., Ballaran, T. B., Klemme, S., Domeneghetti, M. C., Miletich, R.,  
742 Scheidl, K. S., Derzsi, M., Tokár, K., Prencipe, M., Alvaro, M., and Nestola, F.: Thermo-elastic behavior of grossular garnet  
743 at high pressures and temperatures, *American Mineralogist*, 102, 851–859, <https://doi.org/10.2138/am-2017-5855>, 2017.

744 Miller, D. P., Marschall, H. R., and Schumacher, J. C.: Metasomatic formation and petrology of blueschist-facies hybrid rocks  
745 from Syros (Greece): Implications for reactions at the slab–mantle interface, *Lithos*, 107, 53–67,  
746 <https://doi.org/10.1016/j.lithos.2008.07.015>, 2009.

747 Murri, M., Mazzucchelli, M. L., Campomenosi, N., Korsakov, A. V., Prencipe, M., Mihailova, B. D., Scambelluri, M., Angel,  
748 R. J., and Alvaro, M.: Raman elastic geobarometry for anisotropic mineral inclusions, 103, 1869–1872,  
749 <https://doi.org/10.2138/am-2018-6625CCBY>, 2018.

750 Murri, M., Alvaro, M., Angel, R. J., Prencipe, M., and Mihailova, B. D.: The effects of non-hydrostatic stress on the structure  
751 and properties of alpha-quartz, *Phys Chem Minerals*, <https://doi.org/10.1007/s00269-018-01018-6>, 2019.

752 Nye, J. F.: *Physical properties of crystals: their representation by tensors and matrices*, Oxford university press, 1985.

753 Parra, T., Vidal, O., and Jolivet, L.: Relation between the intensity of deformation and retrogression in blueschist metapelites  
754 of Tinos Island (Greece) evidenced by chlorite–mica local equilibria, *Lithos*, 63, 41–66, [https://doi.org/10.1016/S0024-4937\(02\)00115-9](https://doi.org/10.1016/S0024-4937(02)00115-9), 2002.

756 Pattison, D. R. M., Chacko, T., Farquhar, J., and McFARLANE, C. R. M.: Temperatures of Granulite-facies Metamorphism:  
757 Constraints from Experimental Phase Equilibria and Thermobarometry Corrected for Retrograde Exchange, *J Petrology*, 44,  
758 867–900, <https://doi.org/10.1093/petrology/44.5.867>, 2003.



759 Pawley, A. R., Redfern, S. A. T., and Holland, T. J. B.: Volume behavior of hydrous minerals at high pressure and temperature:  
760 I. Thermal expansion of lawsonite, zoisite, clinozoisite, and diaspore, 81, 335–340, <https://doi.org/10.2138/am-1996-3-407>,  
761 1996.

762 Peacock, S. M.: The importance of blueschist → eclogite dehydration reactions in subducting oceanic crust, *GSA Bulletin*,  
763 105, 684–694, [https://doi.org/10.1130/0016-7606\(1993\)105<0684:TIOBED>2.3.CO;2](https://doi.org/10.1130/0016-7606(1993)105<0684:TIOBED>2.3.CO;2), 1993.

764 Peillod, A., Ring, U., Glodny, J., and Skelton, A.: An Eocene/Oligocene blueschist-/greenschist facies P–T loop from the  
765 Cycladic Blueschist Unit on Naxos Island, Greece: Deformation-related re-equilibration vs. thermal relaxation, 35, 805–830,  
766 <https://doi.org/10.1111/jmg.12256>, 2017.

767 Peillod, A., Majka, J., Ring, U., Drüppel, K., Patten, C., Karlsson, A., Włodek, A., and Tehler, E.: Differences in  
768 decompression of a high-pressure unit: A case study from the Cycladic Blueschist Unit on Naxos Island, Greece, *Lithos*, 386–  
769 387, 106043, <https://doi.org/10.1016/j.lithos.2021.106043>, 2021.

770 Pe-Piper, G. and Piper, D. J. W.: *The igneous rocks of Greece: The anatomy of an orogen*, 2002.

771 Philippon, M., Brun, J.-P., and Gueydan, F.: Tectonics of the Syros blueschists (Cyclades, Greece): From subduction to Aegean  
772 extension: TECTONICS OF THE SYROS BLUESCHISTS, 30, n/a-n/a, <https://doi.org/10.1029/2010TC002810>, 2011.

773 Powell, R. and Holland, T.: Optimal geothermometry and geobarometry, *American Mineralogist*, 79, 120–133, 1994.

774 Putlitz, B., Cosca, M. A., and Schumacher, J. C.: Prograde mica <sup>40</sup>Ar/<sup>39</sup>Ar growth ages recorded in high pressure rocks  
775 (Syros, Cyclades, Greece), *Chemical Geology*, 214, 79–98, <https://doi.org/10.1016/j.chemgeo.2004.08.056>, 2005.

776 Qin, F., Wu, X., Wang, Y., Fan, D., Qin, S., Yang, K., Townsend, J. P., and Jacobsen, S. D.: High-pressure behavior of natural  
777 single-crystal epidote and clinozoisite up to 40 GPa, *Phys Chem Minerals*, 43, 649–659, <https://doi.org/10.1007/s00269-016-0824-7>, 2016.

778

779 Ridley, J.: Evidence of a temperature-dependent ‘blueschist’ to ‘eclogite’ transformation in high-pressure metamorphism of  
780 metabasic rocks, 25, 852–870, 1984.

781 Ring, U., Glodny, J., Will, T., and Thomson, S.: The Hellenic Subduction System: High-Pressure Metamorphism, Exhumation,  
782 Normal Faulting, and Large-Scale Extension, 38, 45–76, <https://doi.org/10.1146/annurev.earth.050708.170910>, 2010.

783 Ring, U., Pantazides, H., Glodny, J., and Skelton, A.: Forced Return Flow Deep in the Subduction Channel, Syros, Greece,  
784 39, e2019TC005768, <https://doi.org/10.1029/2019TC005768>, 2020.

785 Roche, V., Laurent, V., Cardello, G. L., Jolivet, L., and Scaillet, S.: Anatomy of the Cycladic Blueschist Unit on Sifnos Island  
786 (Cyclades, Greece), *Journal of Geodynamics*, 97, 62–87, <https://doi.org/10.1016/j.jog.2016.03.008>, 2016.

787 Rosenbaum, G., Avigad, D., and Sánchez-Gómez, M.: Coaxial flattening at deep levels of orogenic belts: evidence from  
788 blueschists and eclogites on Syros and Sifnos (Cyclades, Greece), *Journal of Structural Geology*, 24, 1451–1462,  
789 [https://doi.org/10.1016/S0191-8141\(01\)00143-2](https://doi.org/10.1016/S0191-8141(01)00143-2), 2002.

790 Rosenfeld, J. L.: Stress effects around quartz inclusions in almandine and the piezothermometry of coexisting aluminum  
791 silicates, *Am J Sci*, 267, 317–351, <https://doi.org/10.2475/ajs.267.3.317>, 1969.

792 Rosenfeld, J. L. and Chase, A. B.: Pressure and temperature of crystallization from elastic effects around solid inclusions in  
793 minerals?, *Am J Sci*, 259, 519–541, <https://doi.org/10.2475/ajs.259.7.519>, 1961.

794 Rubie, D. C.: Disequilibrium during metamorphism: the role of nucleation kinetics, 138, 199–214,  
795 <https://doi.org/10.1144/GSL.SP.1996.138.01.12>, 1998.

796 Schliestedt, M. and Matthews, A.: Transformation of blueschist to greenschist facies rocks as a consequence of fluid  
797 infiltration, Sifnos (Cyclades), Greece, *Contr. Mineral. and Petrol.*, 97, 237–250, <https://doi.org/10.1007/BF00371243>, 1987.

798 Schmädicke, E. and Will, T. M.: Pressure–temperature evolution of blueschist facies rocks from Sifnos, Greece, and  
799 implications for the exhumation of high-pressure rocks in the Central Aegean, 21, 799–811, <https://doi.org/10.1046/j.1525->  
800 1314.2003.00482.x, 2003.

801 Schmidt, C. and Ziemann, M. A.: In-situ Raman spectroscopy of quartz: A pressure sensor for hydrothermal diamond-anvil  
802 cell experiments at elevated temperatures, *American Mineralogist*, 85, 1725–1734, <https://doi.org/10.2138/am-2000-11-1216>,  
803 2000.

804 Schneider, D. A., Grasemann, B., Lion, A., Soukis, K., and Draganits, E.: Geodynamic significance of the Santorini  
805 Detachment System (Cyclades, Greece), 30, 414–422, <https://doi.org/10.1111/ter.12357>, 2018.

806 Schumacher, J. C., Brady, J. B., Cheney, J. T., and Tonnsen, R. R.: Glaucofane-bearing Marbles on Syros, Greece, *J*  
807 *Petrology*, 49, 1667–1686, <https://doi.org/10.1093/petrology/egn042>, 2008.

808 Sharp, Z. D.: A laser-based microanalytical method for the in situ determination of oxygen isotope ratios of silicates and  
809 oxides, *Geochimica et Cosmochimica Acta*, 54, 1353–1357, [https://doi.org/10.1016/0016-7037\(90\)90160-M](https://doi.org/10.1016/0016-7037(90)90160-M), 1990.

810 Sharp, Z. D. and Kirschner, D. L.: Quartz-calcite oxygen isotope thermometry: A calibration based on natural isotopic  
811 variations, 58, 4491–4501, [https://doi.org/10.1016/0016-7037\(94\)90350-6](https://doi.org/10.1016/0016-7037(94)90350-6), 1994.

812 Skelton, A., Peillod, A., Glodny, J., Klonowska, I., Månbro, C., Lodin, K., and Ring, U.: Preservation of high-P rocks coupled  
813 to rock composition and the absence of metamorphic fluids, *Journal of Metamorphic Geology*, 0,  
814 <https://doi.org/10.1111/jmg.12466>, 2018.

815 Soukis, K. and Stockli, D. F.: Structural and thermochronometric evidence for multi-stage exhumation of southern Syros,  
816 Cycladic islands, Greece, *Tectonophysics*, 595–596, 148–164, <https://doi.org/10.1016/j.tecto.2012.05.017>, 2013.

817 Spear, F. S. and Pattison, D. R. M.: The implications of overstepping for metamorphic assemblage diagrams (MADs), 457,  
818 38–46, <https://doi.org/10.1016/j.chemgeo.2017.03.011>, 2017.

819 Spear, F. S. and Selverstone, J.: Quantitative P-T paths from zoned minerals: Theory and tectonic applications, *Contr. Mineral.*  
820 *and Petrol.*, 83, 348–357, <https://doi.org/10.1007/BF00371203>, 1983.

821 Spear, F. S., Wark, D. A., Cheney, J. T., Schumacher, J. C., and Watson, E. B.: Zr-in-rutile thermometry in blueschists from  
822 Sifnos, Greece, *Contrib Mineral Petrol*, 152, 375–385, <https://doi.org/10.1007/s00410-006-0113-4>, 2006.

823 Syracuse, E. M., van Keken, P. E., and Abers, G. A.: The global range of subduction zone thermal models, *Physics of the Earth*  
824 *and Planetary Interiors*, 183, 73–90, <https://doi.org/10.1016/j.pepi.2010.02.004>, 2010.

825 Thomas, J. B. and Spear, F. S.: Experimental study of quartz inclusions in garnet at pressures up to 3.0 GPa: evaluating validity  
826 of the quartz-in-garnet inclusion elastic thermobarometer, *Contrib Mineral Petrol*, 173, 42, [https://doi.org/10.1007/s00410-](https://doi.org/10.1007/s00410-018-1469-y)  
827 018-1469-y, 2018.

828 Tomaschek, F., Kennedy, A. K., Villa, I. M., Lagos, M., and Ballhaus, C.: Zircon from Syros, Cyclades, Greece—  
829 Recrystallization and Mobilization of Zircon During High-Pressure Metamorphism, *J Petrology*, 44, 1977–2002,  
830 <https://doi.org/10.1093/petrology/egg067>, 2003.

831 Trotet, F., Vidal, O., and Jolivet, L.: Exhumation of Syros and Sifnos metamorphic rocks (Cyclades, Greece). New constraints  
832 on the P-T paths, *European Journal of Mineralogy*, 13, 901–902, <https://doi.org/10.1127/0935-1221/2001/0013-0901>, 2001a.

833 Trotet, F., Jolivet, L., and Vidal, O.: Tectono-metamorphic evolution of Syros and Sifnos islands (Cyclades, Greece),  
834 *Tectonophysics*, 338, 179–206, [https://doi.org/10.1016/S0040-1951\(01\)00138-X](https://doi.org/10.1016/S0040-1951(01)00138-X), 2001b.

835 Urey, H. C.: The thermodynamic properties of isotopic substances, 562–581, 1947.

836 Uunk, B., Brouwer, F., ter Voorde, M., and Wijbrans, J.: Understanding phengite argon closure using single grain fusion age  
837 distributions in the Cycladic Blueschist Unit on Syros, Greece, *Earth and Planetary Science Letters*, 484, 192–203,  
838 <https://doi.org/10.1016/j.epsl.2017.12.031>, 2018.

839 Walowski, K. J., Wallace, P. J., Hauri, E. H., Wada, I., and Clynne, M. A.: Slab melting beneath the Cascade Arc driven by  
840 dehydration of altered oceanic peridotite, 8, 404–408, <https://doi.org/10.1038/ngeo2417>, 2015.

841 Wang, J., Mao, Z., Jiang, F., and Duffy, T. S.: Elasticity of single-crystal quartz to 10 GPa, *Phys Chem Minerals*, 42, 203–  
842 212, <https://doi.org/10.1007/s00269-014-0711-z>, 2015.


843 Wang, Z. and Ji, S.: Elasticity of six polycrystalline silicate garnets at pressure up to 3.0 GPa, *American Mineralogist*, 86,  
844 1209–1218, <https://doi.org/10.2138/am-2001-1009>, 2001.

845 Wunder, B. and Schreyer, W.: Antigorite: High-pressure stability in the system MgO–SiO<sub>2</sub>–H<sub>2</sub>O (MSH), *Lithos*, 41, 213–  
846 227, [https://doi.org/10.1016/S0024-4937\(97\)82013-0](https://doi.org/10.1016/S0024-4937(97)82013-0), 1997.

847 Zhang, Y.: Mechanical and phase equilibria in inclusion–host systems, *Earth and Planetary Science Letters*, 157, 209–222,  
848 [https://doi.org/10.1016/S0012-821X\(98\)00036-3](https://doi.org/10.1016/S0012-821X(98)00036-3), 1998.

849 Zhong, X., Moulas, E., and Tajčmanová, L.: Post-entrapment modification of residual inclusion pressure and its implications  
850 for Raman elastic thermobarometry, 11, 223–240, <https://doi.org/10.5194/se-11-223-2020>, 2020.

851



852


853

854

855

856

**Figure 1.** Simplified geologic map of Syros, Greece [modified from Keiter et al. (2011)]. Inset map shows Syros relative to the North and West Cycladic, and Naxos-Paros Detachment Systems (NCSD, WCDS, NPDS, modified from Grasemann et al., 2012). Shear zones within the CBU and the Vari detachment are after Laurent et al., 2016 and Soukis and Stockli (2013), respectively. Stereonets from each studied outcrop are shown, and arrows indicate the outcrop location.



858

859 **Figure 2. Outcrop, micrograph, and electron images showing stages of retrograde deformation present in southern Delfini. a)**  
 860 **Upright folds ( $F_{12}$ ) that refold the primary  $S_s$  foliation. b): Core of  $F_{12}$  folds (below Fig. 2a, KCS34). c): Plane light image of sample**  
 861 **KCS34; sample cut perpendicular to the  $F_{12}$  fold axial plane. Epidotes (ep2) from the upright fold exhibit recrystallization as**  
 862 **indicated by alignment with a late  $S_{12}$  crenulation, and a reduction in inclusions and grain size. d) Ep2 with late titanite (ttn)**  
 863 **inclusions. Ep2 is parallel to white mica (wm) that defines  $S_{12}$  (KCS34). e) Ep2 in textural equilibrium with tttn (KCS34). f) Ep1**  
 864 **parallel to  $S_s$ , with garnet (grt) and quartz (qtz) inclusions that do not define an internal foliation (KCS1621). g) Poikiloblastic ep1**  
 865 **parallel to  $S_s$ , with a weak internal foliation defined by qtz (KCS1621).**

866

867

868


869

870

871

872

873




875

876

877

**Figure 3. Outcrop photos of epidote boudins sampled for oxygen isotope thermometry. a) SY1613 (Lotos), b) SY1617 (Delfini), c) SY1618 (Delfini), d) SY1623 (Delfini).**




878

879

880

881

**Figure 4. Photomicrographs of measured quartz inclusions in garnet from Delfini (a) and Raman spectrums of unstrained Herkimer quartz and strained quartz inclusions (b). b) Shown for comparison are Herkimer quartz (red) and quartz inclusion (blue) measurements from Virginia Tech and ETH Zürich. Quartz bands and Ar plasma lines (only VT analyses) are numerically labelled.**



882

883 **Figure 5. Comparison of  $P_{inc}$  determined from different quartz bands using hydrostatic calibrations, and by using phonon-mode**  
 884 **Grüneisen tensors (strains). Red, blue, and yellow symbols indicate qtz-in-grt (Group 1), qtz-in-ep1 (Group 2), and qtz-in-ep2**  
 885 **(Group 3) results, respectively. Diamonds, squares, and circles indicate  $P_{inc}^{464}$  vs  $P_{inc}^{128}$ ,  $P_{inc}^{464}$  vs  $P_{inc}^{206}$ , and  $P_{inc}^{464}$  vs  $P_{inc}^{strains}$**   
 886 **results, respectively. No border, filled, and open symbols indicate analyses from Kalamisia, Delfini, and Lotos samples, respectively.**

887

888

889

890

891

892

893

894

895


896

897

898

899






900

901 **Figure 6. Compositional x-ray maps of two garnets from sample KCS34 (Delfini). Blue dots indicate the location of measured**  
 902 **inclusions; systematic  $P_{\text{trap}}$  differences are not observed across garnets ( $P_{\text{trap}}$  units are GPa, calculated at  $T_{\text{trap}} = 500$  °C.). Subscripts**  
 903 **indicate the inclusion number (see Supplementary Table S3).**

904



905

906 **Figure 7. (a) P-T conditions deduced from elastic thermobarometry and oxygen isotope thermometry superimposed on modeled**  
 907 **Cascadia slap-top geotherm (Syracuse et al., 2010) and b) reference P-T conditions. (a)  $P_{\text{trap}}$  from Groups 1, 2, and 3, that reflect**  
 908 **peak (qtz-in-garnet), retrograde blueschist-greenschist facies (qtz-in-ep1), and late greenschist facies (qtz-in-ep2) conditions. Solid**  
 909 **red, blue, and yellow lines and rectangles are the  $P_{\text{trap}}$  isomekes (calculated from the mean residual inclusion pressure of each group)**  
 910 **and our best-estimate entrapment conditions, respectively. Transparent lines are  $P_{\text{trap}}$  errors ( $1\sigma$  around the mean) for analyses from**  
 911 **Delfini samples. Grey box bounds the range of temperatures calculated from oxygen isotope thermometry of quartz-calcite boudin**  
 912 **neck precipitates. b) Recalculated  $P_{\text{trap}}$  values from Behr et al. (2018) (Syros) and Ashley et al. (2014) (Sifnos) and are shown in**  
 913 **purple (solid border) and green (dashed border) rectangles, respectively. Metamorphic facies are taken from (Peacock, 1993).**  
 914 **Metamorphic facies fields (Peacock, 1993): zeolite (ZE), prehnite-pumpellyite (PrP), prehnite-actinolite (PrAc), pumpellyite-**  
 915 **actinolite (PA), lawsonite-chlorite (LC), greenschist (GS), lawsonite-blueschist (LB), epidote-blueschist (EB), epidote-amphibolite**  
 916 **(EA), amphibolite (AM), eclogite (EC). RSCM = Raman Spectroscopy of Carbonaceous Material (data from Laurent et al., 2018).**

917

918