

1 Evolution of the Iberian Massif as deduced from its crustal thickness and 2 geometry of a mid-crustal (Conrad) discontinuity

3 Puy Ayarza¹, José Ramón Martínez Catalán¹, Ana Martínez García², Juan Alcalde², Juvenal Andrés^{1,2},
4 José Fernando Simancas³, Immaculada Palomeras¹, David Martí^{2,4}, Irene DeFelipe², Chris Juhlin⁵,
5 Ramón Carbonell²

- 6 1. Geology Department, Salamanca University. Pza de la Merced, s/n. Salamanca 37008. Spain
7 2. Geosciences Barcelona, CSIC. Lluís Solé i Sabaris, s/n. Barcelona 08028. Spain
8 3. Geodynamics Department, Granada University. Av/ de la Fuente Nueva, s/n. Granada 18071. Spain
9 4. Lithica SCCL. Avinguda Farners 16, Santa Coloma de Farners, Girona 17430. Spain
10 5. Department of Earth Sciences, Uppsala University. Villavägen 16, Uppsala, 75236. Sweden

11
12 Corresponding author: Puy Ayarza (puy@usal.es)

13 Abstract

14 Normal incidence seismic data provide the best images of the crust and lithosphere. When
15 properly designed and continuous, these sections greatly contribute to understanding the
16 geometry of orogens and, together with surface geology, to unravel their evolution. In this
17 paper we present ~~an~~the almost complete transect, to date, of the Iberian Massif, the
18 westernmost exposure of the European Variscides. Despite the heterogeneity of the dataset,
19 acquired during the last 30 years, the images resulting from reprocessing with a homogeneous
20 workflow allow us to clearly define the crustal thickness and its internal architecture. The
21 Iberian Massif crust, formed by the amalgamation of continental pieces belonging to
22 Gondwana and Laurussia (Avalonian margin) is well structured in upper and lower crust. A
23 conspicuous mid-crustal discontinuity is clearly defined by the top of the reflective lower crust
24 and by the asymptotic geometry of reflections that merge into it, suggesting that it has often
25 acted as a detachment. The geometry and position of this discontinuity can give us insights on
26 the evolution of the orogen, i.e. of the magnitude of compression and the effects and extent of
27 ~~the late~~ Variscan gravitational collapse. Also, ~~its position and~~ the limited thickness of the
28 lower crust in central and NW Iberia might have constrained the response of the Iberian
29 microplate to Alpine shortening. This discontinuity, featuring a Vp increase, is here observed as
30 an orogeny-scale ~~feature~~boundary with characteristics compatible with those of the
31 worldwide debated, Conrad discontinuity.

32 **Keywords:** Iberian Massif, vertical incidence seismic data, mid-crustal detachment, Conrad
33 discontinuity, geodynamic evolution

34 Glossary:

35 CIA: Central Iberian Arc

36 CIZ: Central Iberian Zone

37 CU: Central Unit

38 CZ: Cantabrian Zone

39 DB: Duero Basin

40 GTMZ: Galicia Tras-os-Montes Zone

41 JAA: Ibero-Armorican Arc

Con formato: Fuente: Negrita, Español (alfab. internacional)

Con formato: Español (alfab. internacional)

Con formato: Espacio Después: 0 pto

Con formato: Español (alfab. internacional)

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

42 [ICS: Iberian Central System](#)
43 [NI: Normal Incidence](#)
44 [OMA: Ossa-Morena Zone](#)
45 [SPZ: South Portuguese Zone](#)
46 [TB: Tajo Basin](#)
47 [WA: Wide Angle](#)
48 [WALZ: West Asturian-Leonese Zone](#)
49

50 1. Introduction

51 In the last 35 years, controlled source seismic data have greatly contributed to the
52 understanding of the European Variscides. National research programs like DEKORP (Bortfeld,
53 1985; DEKORP Research Group, 1987; Franke et al., 1990; Oncken, 1998), BIRPS and ECORS
54 (BIRPS and ECORS, 1986) have sampled this orogen providing a detailed picture of its
55 lithospheric architecture. In the Iberian Massif, normal incidence (NI) seismic reflection profiles
56 often acquired with coincident wide angle (WA) reflection/refraction seismic data have
57 allowed scientists to depict its crustal structure, infer its P and S waves velocity distribution,
58 place constraints on its geodynamic evolution, visualize the accommodation pattern of
59 shortening at different crustal levels and, sometimes, deduce the effect of Alpine reactivation
60 on this Paleozoic orogen.

61 In this regard, seismic datasets acquired in the Iberian Massif (DeFelipe et al., 2020)(~~De Felipe~~
62 ~~et al., 2020~~) from the programs ESCIN (Ayarza et al., 1998, 2004; Pérez-Estaún et al., 1991;
63 Pulgar et al., 1996), IBERSEIS (Flecha et al., 2009; Palomeras et al., 2009; Simancas et al., 2003),
64 ALCUDIA (e.g., Ehsan et al., 2014, 2015; Martínez Poyatos et al., 2012) and CIMDEF (Andrés et
65 al., 2019) have helped so far to identify several outstanding features such as, i) clear
66 differences in the intensity and geometry of reflectivity at upper and lower crustal levels, ii)
67 contrasting deformation patterns deduced from the good correlation between reflectivity and
68 upper crustal structures as, regardless of the many factors that trigger the concentration of
69 deformation along narrow thrust zones (Butler and Mazzoli, 2006), the latter often follow
70 lithological boundaries in the Iberian Massif (e.g., Alonso, 1987), thus being candidates to
71 appear as outstanding reflections in NI seismic sections, and iii) a very reflective and
72 sometimes thick lower crust, even in areas where the upper crust is weakly deformed. In order
73 to explain these features, decoupling of the upper and lower crust has been invoked along
74 large parts of the sampled area ~~-(Simancas et al., 2013)(Simancas et al., 2013a)~~. The existence
75 of a mid-crustal detachment has been addressed by these authors as the reason why different
76 shortening mechanisms exist at different crustal levels. However, coupled crustal deformation
77 has been inferred for NW Iberia, where large crustal thickening took place. Also, no inference
78 has been made on how this detachment acted during later Alpine deformation in the central
79 and northern Iberian Massif.

80 In this paper, we present a more complete new composite seismic section of the Iberian Massif
81 that integrates results of ~~We benefit from the existence of two~~ new datasets: (CIMDEF and
82 ALCUDIA WA ~~The former~~) ~~that~~ fill the gaps of areas previously unexplored, like Central Iberia,
83 altogether providing an almost complete transect. In addition, we include the N-S ESCIN-2

84 | seismic profile and new time-migrated sections of all datasets, some of them, as yet,
85 | unmigrated (e.g., ESCIN-1). Although the detailed interpretation of the upper crustal
86 | reflectivity and its correlation with the surface geology does not significantly change, later on,
87 | we revisit the deep reflectivity and redefine the extension and implications of a mid-crustal
88 | discontinuity that, in our view, exists below the entire Iberian Massif, affecting the two
89 | continents involved in the Variscan Orogeny and certainly playing a critical role in the
90 | decoupling between upper and lower crustal deformation. Finally, we infer the geometry and
91 | nature of this feature, discuss its tectonic significance and its role during the younger Alpine
92 | Orogeny, and relate it with the long-debated (e.g., Finlayson et al., 1984; Litak and Brown,
93 | 1989; Wever, 1989; Xiaobo and Tae Kyung, 2010) Conrad discontinuity (Conrad, 1925) of the
94 | classic continental seismology.

95 | 2. Geological setting

96 | The Iberian Massif represents the westernmost outcrop of the European Variscides, exposing
97 | an almost complete section of this Paleozoic oOrogen. It is divided into six zones (Fig. 1; Arenas
98 | et al., 1988; Farias et al., 1987; Julivert et al., 1972) that from N to S and E to W are: Cantabrian
99 | (CZ), West Asturian-Leonese (WALZ), Galicia-Trás-os-Montes (GTMZ), Central Iberian (CIZ),
100 | Ossa-Morena (OMZ) and South Portuguese (SPZ). The CZ and the SPZ represent the external
101 | zones whereas the rest represent the hinterland. The CZ, WALZ and CIZ belong to the northern
102 | margin of the Paleozoic Gondwana. The GTMZ represents the remnants of a large nappe stack
103 | formed by pieces of the outermost margin of the Gondwana margin, i.e., a pulled-apart peri-
104 | Gondwanan terrane and oceanic units derived from the oceanic realm separating them. They
105 | were emplaced above the autochthonous CIZ in NW Iberia and are preserved as a large klippen
106 | at the core of late Variscan synforms (Martínez Catalán et al., 1997, 2007; Ries and Shackleton,
107 | 1971). Its rootless suture (ophiolitic units) is thought to represent a branch of the Rheic
108 | oceanic realm (Martínez Catalán et al., 1997). The OMZ has been interpreted as a continental
109 | fragment that rifted and probably drifted away from Gondwana (Matte, 2001; Robardet,
110 | 2002), docking back later to the CIZ giving rise to a suture and thus representing another peri-
111 | Gondwanan terrane (Azor et al., 1994; Gómez-Pugnaire et al., 2003; Simancas et al., 2001).
112 | Finally, the SPZ is thought to be separated from the OMZ by the Rheic Ocean suture (Simancas
113 | et al., 2003) which was later overprinted by early Carboniferous extension accompanied by
114 | mafic intrusions (Azor et al., 2008) and by transpression (Pérez-Cáceres et al., 2015). In this
115 | context, the basement of the SPZ represents a fragment of Avalonia (Braid et al., 2011, 2012;
116 | Pereira et al., 2014; Rodrigues et al., 2015), and its ~~correlation~~ affinity with the Rheno-
117 | Hercynian Zone in Germany (Franke, 2000; Franke et al., 1990) further supports this
118 | affinity~~correlation~~.

119 | From a tectonic point of view, the Iberian Massif shows evidences of pre-Variscan activity. The
120 | Cadomian Neoproterozoic event is characterized by continental arc magmatism, deformation
121 | and metamorphism (Bandrés et al., 2004; Dallmeyer and Quesada, 1992; Ochsner, 1993). It
122 | developed above a previous non-outcropping continental crust, and formed the basement
123 | above which the Ediacaran and Paleozoic sedimentation took place, favored by Ediacaran-
124 | Cambrian and Cambro-Ordovician rifting which developed a wide continental platform
125 | (Linnemann et al., 2008; Sánchez-García et al., 2008, 2010). However, most of the outcropping
126 | tectonic features observed in the surface are the result of the Devonian and Carboniferous

127 collision between Gondwana, some peri-Gondwanan terranes and the Avalonian border of
128 Laurentia, which resulted in the Variscan Orogen (Matte, 2001). The deformation associated to
129 the latter was diachronous along the Iberian Massif. Next, we describe ~~– this part of the~~
130 [European Variscides](#), from N to S (in present coordinates), together with ~~the-its~~ most
131 important [tectonic and](#) stratigraphic features.

132 The CZ (Fig. 1) is an external zone located at the core of the Ibero-Armorican Arc (IAA; Dias and
133 Ribeiro, 1995; Lotze, 1929; Matte and Ribeiro, 1975; Stille, 1924). It is a thin-skinned thrust and
134 fold belt with a transport direction towards the foreland in the E, and overprinted by [the](#)
135 oroclinal folding giving rise to the IAA (Alonso et al., 2009; Pérez-Estaún et al., 1988).
136 Stratigraphically, it is characterized by a Precambrian sequence, [outcropping](#) ~~out~~ at its western
137 part and overlain by a Paleozoic stratigraphic succession that ranges from the Cambrian to a
138 well-developed Carboniferous: pre-orogenic up to early Carboniferous, and syn-orogenic in the
139 Upper Carboniferous (Sánchez de Posada et al., 1990; Truyols et al., 1990). Scarce tholeiitic
140 and alkaline magmatism is related to Cambro-Ordovician rifting (Corretgé and Suárez, 1990),
141 ~~and a~~ regional metamorphism accompanied deformation, indicating shallow crustal
142 conditions. In this area, deformation began at ~325-320 Ma, in the Late Mississippian
143 (Dallmeyer et al., 1997), and the emplacement of nappes that characterize the deformation
144 and the formation of folds within each sequence took place between the Westphalian B and
145 the Stephanian (313-300 Ma), in the Upper Carboniferous (Pérez-Estaún et al., 1988). An
146 extensional episode related to the end of the orogeny led to the formation of Permian Basins
147 (Martínez García, 1981). Later on, extension related to the opening of the Bay of Biscay
148 triggered the development of deep Cretaceous basins (Quintana et al., 2015; Rat, 1988). Alpine
149 tectonics uplifted the Pyrenean-Cantabrian range from the end of the Late Cretaceous to the
150 Miocene (DeFelipe et al., 2019; Teixell et al., 2018 [and references therein](#)) ~~(DeFelipe et al.,~~
151 [2019; Teixell et al., 2018 and references therein](#)) reactivating Variscan thrusts and Mesozoic
152 normal faults (Gallastegui et al., 2016).

153 The WALZ lies to the W of the CZ (Fig. 1). Stratigraphically it consists of a Neoproterozoic
154 terrigenous sequence ~~unconformably~~ overlain by a Paleozoic platform succession that
155 ranges from the Cambrian to the Lower Devonian (Pérez-Estaún et al., 1990) ~~(Martínez Catalán,~~
156 [1985](#)), much thicker than that of the CZ. These sediments were actively deformed along three
157 compressional (C1, C2, and C3) and two extensional (E1 and E2) phases during the Variscan
158 Orogeny (Martínez Catalán et al., 2014). Large E vergent folds witness the C1 related
159 compression (360-340 Ma). Those were later affected by E vergence thrusts resulting from
160 ongoing shortening (345-325 Ma). Crustal thickening followed by thermal relaxation led to syn-
161 orogenic extension during E1 (330-315 Ma). A last compressional episode (C3, 315-305 Ma)
162 produced upright folds associated with wrench shear zones while simultaneous extension (E2,
163 315-300 Ma) continued, characterizing the latest stages of the orogeny in this area. Crustal
164 melting triggered by compression and thickening led to extension and to the intrusion of
165 granitoids in the western part of the WALZ. Thermal models show that the crust could have
166 started to melt within 30 Ma after the ~~start-onset~~ of crustal thickening, which is then
167 constrained by the ages of Variscan granitoids (Alcock et al., 2009).

168 The CIZ is the largest of the Iberian Massif zones. Curvature of magnetic anomalies and that of
169 early (C1) Variscan folds depict the Central Iberian Arc (CIA; Martínez Catalán, 2011a, 2011b),

Código de campo cambiado

170 | partly explaining the width of this internal zone (Fig. 1). The stratigraphic sequence differs
171 | from N to S: Ordovician felsic metavolcanic, subvolcanic and intrusives (Diez-Montes et al.,
172 | 2010) represent the most ancient lithologies ~~outcropping-out~~ in the N, defining the 'Ollo de
173 | Sapo' domain whereas to the S, Upper-Proterozoic-Lower Cambrian metasediments outcrop
174 | (Díez Balda et al., 1995) (~~Díez Balda, 1986; Díez Balda et al., 1995~~), defining the 'Schist-
175 | Greywacke Complex' domain. The pre-orogenic sedimentary sequence continues to the
176 | Devonian, followed by a syn-orogenic Carboniferous sequence (Martínez Catalán et al., 2004;
177 | Robardet, 2002). This area represents a relatively stable Gondwana margin characterized by
178 | the Early Ordovician extension that opened the Rheic Ocean and allowed intrusion of
179 | essentially felsic magmas (Díez Montes et al., 2010). The deformation phases described for the
180 | WALZ affected most of the CIZ although C1, C2 and E1 are somewhat older, according to the
181 | propagation of deformation from the hinterland. Slight differences in the importance of
182 | phases can also be found to the center and S (Martínez Catalán et al., 2019), allowing the CIZ
183 | to be divided in two zones. In the NW, intense recumbent C1 folds and important C2 thrusts
184 | ~~exist,~~ related to the emplacement of the GTMZ ~~occur~~. Outcropping rocks show epi- to
185 | catazonal metamorphism and ductile detachments. Gneiss domes of both E1 and E2
186 | extensional phases ~~existexist~~, evidencing significant crustal thinning, and Variscan granitoids
187 | are abundant. To the S, C1 folds are upright, C2 deformation is limited to the southernmost
188 | part, and upright C3 folds are the most important structures (Martínez Catalán et al., 2012;
189 | Martínez Poyatos, 2002). Metamorphism is generally weak and the amount of granitoids
190 | decreases, except in the Iberian Central System (ICS). Here extension postdates C3 upright
191 | folding and thus, it is considered E2.

192 | Alpine tectonics in the CIZ reactivated previous Variscan fractures and triggered the
193 | development of the Iberian Central System (ICS) mountain belt (de Vicente et al., 1996),
194 | allowing the products of syn- and post-E2 crustal melting to outcrop in large areas.

195 | The GTMZ is represented by five klippen that are the remnants of the emplacement of a thick
196 | nappe stack on top of the CIZ. This includes, from bottom to top, a relatively distal part of the
197 | northern Gondwana margin (Parautochthon), the outermost edge of that margin (Lower
198 | Allochthon), a few oceanic units of Cambro-Ordovician and Lower Devonian age (Middle
199 | Allochthon) and a peri-Gondwanan terrane with magmatic evidences of Cambro-Ordovician
200 | rifting and a continental arc setting (Upper Allochthon). Several units show high-P
201 | metamorphism reflecting subduction of the ocean represented by the Middle Allochthon and
202 | involving also the Upper and Lower ones (Arenas et al., 2007; Gómez Barreiro et al., 2007;
203 | Martínez Catalán et al., 2007; Sánchez Martínez et al., 2007). Ongoing subduction during most
204 | of the Devonian (400-365 Ma) built an accretionary wedge that was subsequently emplaced on
205 | top of the CIZ during the early Carboniferous (C1-C2 events, c. 360-340 Ma).

206 | The boundary between the CIZ and the OMZ (the Badajoz Córdoba Shear Zone) has been
207 | largely interpreted as a suture (Gómez-Pugnaire et al., 2003; Simancas et al., 2001), although
208 | no true oceanic units have been identified. It includes amphibolites of oceanic affinity from the
209 | early Paleozoic, as well as eclogite relics. In SW Iberia, outcropping lithologies range from the
210 | Upper Precambrian to the Upper Carboniferous, with an angular unconformity at the Lower
211 | Carboniferous. In the OMZ, the Serie Negra (Black Series) is a thick Neoproterozoic sequence
212 | that includes graphitic quartzites and schists and underwent Cadomian arc-related magmatism

Con formato: Ajustar espacio entre texto latino y asiático, Ajustar espacio entre texto asiático y números

213 and regional metamorphism (Dallmeyer and Quesada, 1992; Ochsner, 1993; Quesada and
214 Dallmeyer, 1994). The pre-orogenic Paleozoic sequence is rather complete and was deposited
215 at the peri-Gondwanan platform, as for the CIZ, although differences in the faunal content and
216 in the Paleozoic facies, generally more pelitic in the OMZ, point to a more distal position
217 (Robardet, 2002; Robardet and Gutiérrez Marco, 1990). Ediacaran-Cambrian and Cambro-
218 Ordovician magmatism reflects two rifting events. The latter is the most important one, it
219 includes alkaline magmatism and is related ~~with-to~~ the opening of the Rheic Ocean (García
220 Casquero et al., 1985; Ochsner, 1993; Sánchez-García et al., 2008, 2010). The first deformation
221 event, of Devonian age, formed overturned and recumbent folds and thrust faults with SW
222 vergence (Expósito et al., 2002, 2003). Syn-orogenic, early Carboniferous basins developed in
223 an extensional context and are related to calc-alkaline volcanism and magmatism (Casquet et
224 al., 2001). These deposits unconformably overlay the early folds and thrusts. Later,
225 deformation continued with middle and upper Carboniferous sinistral transpression and
226 associated upright NW-SE folds.

227 A salient seismic reflector, the Iberseis Reflective Body (IRB, Carbonell et al., 2004; Simancas et
228 al., 2003) seems to be the result of a mantle-derived intrusion located along a mid-crustal
229 detachment ~~at~~ around 350-340 Ma. It was emplaced in the context of early Carboniferous
230 extension in the SW of the Iberian Massif, while the hinterland to the NW was undergoing the
231 first stages of compression (C1-C2). Magmatic activity in the SW triggered a high-T/low-P
232 metamorphism that, otherwise, has a low grade elsewhere in the OMZ (Díaz Azpiroz et al.,
233 2006; Pereira et al., 2009).

234 The boundary between the OMZ and the SPZ has been long understood as a suture on the
235 basis of geometric assumptions (e.g., Carvalho, 1972). Later evidences have reinforced this
236 point of view suggesting that the above mentioned boundary represents the remnants of the
237 Rheic Ocean, although Carboniferous transtension and transpression have largely obliterated it
238 (Pérez-Cáceres et al., 2015 and references therein). The SPZ is a Variscan foredeep basin
239 strongly deformed by thin-skinned thrust tectonics, and is usually correlated with the
240 Rhenohercynian Zone of Kossmat (1927) in the Bohemian Massif. It features wide outcrops of
241 low or very low grade Devonian phyllites, quartzites and sandstones overlain by a lower
242 Carboniferous (Early Mississippian) volcano-sedimentary sequence topped by middle and
243 upper Carboniferous flysch (Oliveira, 1990). From a tectonic point of view, it is characterized by
244 Carboniferous S vergent thrusts and folds, the latter featuring axial traces oblique to the
245 northern boundary of the zone, evidencing transpression (Simancas et al., 2003 and references
246 therein). Deformation propagated towards the S along the lower and upper Carboniferous
247 (Oliveira, 1990).

248 Although the start of the Variscan collision seems to have been frontal or maybe right-lateral
249 in most of Europe (Shelley and Bossière, 2000), surface geology and interpretation of seismic
250 data evidences the existence of relevant left lateral transpression and oblique-slip syn-
251 metamorphic shear zones in the OMZ, SPZ and their boundaries (Pérez-Cáceres et al., 2016;
252 Simancas et al., 2003 and references therein). In the OMZ, folds and thrusts witnessing
253 Devonian and early Carboniferous compression are oblique to the OMZ/SPZ boundary,
254 indicating a transpressional setting. These features are disrupted by later Mississippian
255 transtensional tectonics (Expósito et al., 2002) that gave way to the intrusion of the Beja-

Código de campo cambiado

256 Acebuches mafic and ultramafic rocks (Azor et al., 2008). Convergence resumed soon after,
257 leading to the emplacement of the Beja–Acebuches unit onto the OMZ (Pérez-Cáceres et al.,
258 2015). Inside the OMZ, Devonian and Carboniferous left lateral deformation accounts for ~400
259 km, higher than perpendicular shortening. Likewise, inside the SPZ, left-lateral displacement is
260 estimated to reach 90 km whereas the orthogonal one amounts ~60 km (Pérez-Cáceres et al.,
261 2016).

262 3. Geophysical setting: Existing datasets, their reprocessing and a brief description

263 3.1. Seismic datasets sampling the Iberian Massif

264 Since the early 1990's, the Iberian Massif has been sampled by different controlled source
265 seismic experiments (DeFelipe et al., 2020) (~~De Felipe et al., 2020~~): the ESCIN (1991-1992),
266 IBERSEIS (2000 and 2003), and ALCUDIA (2007-2012) experiments acquired normal incidence
267 (NI) and coincident wide-angle (WA) data. The latest project, carried out with the target of
268 understanding the structure and effect of the Alpine reactivation across the central part of the
269 Iberian Massif, is the CIMDEF experiment (2017-2019). It ~~acquired-recorded~~ densely spaced
270 controlled source WA reflection and natural source (earthquakes and noise) seismic data.
271 However, the acquisition of coincident NI data along this transect has not currently been
272 planned ~~along this transect~~, regardless of its potential quality and relevance, due to the
273 relatively high costs of this kind of experiments.

274 From N to S, and from E to W, the ESCIN project sampled the northern part of the Iberian
275 Massif (Fig. 1). Profile ESCIN-1 (1991) is an onshore E-W line crossing the CZ from its eastern,
276 most external part to its boundary with the WALZ to the W; Profile ESCIN-2 (1991) is an
277 onshore N-S profile crossing the most external and eastern part of the CZ and reaching the
278 northern end of the Duero Basin (DB) to the S, which represents the Cantabrian Mountains
279 foreland basin. The ESCIN-3 (1992) profiles sampled the WALZ and the CIZ along the northern
280 Iberia shelf. Although it consists of three parts (ESCIN-3.1, 3.2 and 3.3) only the easternmost
281 ones (3.2 and 3.3.) are relevant for the study of the Variscan crust and thus, included here.
282 ESCIN-3.3 crossed the entire WALZ to its western boundary with the CIZ, which in this area was
283 surveyed by the ESCIN-3.2. Geographically, the latter also sampled the alloct~~h~~thonous GTMZ.
284 But as this is an offshore profile, it shows no evidences of the presence of the GTMZ, and most
285 of the imaged crust corresponds to that of its relative autochthon, the CIZ.

286 A significant geographical and methodological gap exists between the ESCIN profiles to the N
287 and the location of the CIMDEF experiment (Fig. 1). The latter crosses central Iberia from the N
288 part of the CIZ, then samples the DB down to the ICS, and goes on S across the Tajo Basin (TB)
289 till it reaches again the CIZ metasediments to the S of the ICS.

290 In the southern part of the Iberian Massif, the onshore ALCUDIA seismic line (NI and WA),
291 striking NE-SW, was acquired across the CIZ, going from the S of the ICS to the boundary with
292 the OMZ. Finally, the NE-SW IBERSEIS dataset (NI and WA) is also an onshore profile that
293 partially overlaps the same structures as the SW end of the ALCUDIA line although with some
294 50 km of offset to the W. This seismic line samples the southern part of the CIZ, the OMZ and
295 the SPZ.

296 Altogether these seismic profiles account for a ~1500 km long seismic transect geared to
297 understand the crustal and, in places, lithospheric structure of the Iberian Massif and to
298 constrain its evolution.

299 3.2. Processing of datasets

300 The data used in this work have been acquired at different times, have different characteristics
301 (onshore and offshore) and accordingly exhibit very heterogeneous quality. Table 1 shows the
302 acquisition parameters of all these datasets. The most outstanding differences are: i) the
303 quality and characteristics of the offshore (ESCIN-3) vs the onshore data, ii) the difference
304 between the low fold (30) ESCIN-1, ESCIN-2 and ESCIN-3 data acquired with an explosive
305 source and airguns respectively and the high fold (>60) IBERSEIS and ALCUDIA datasets, which
306 used Vibroseis trucks as source of energy, and iii) the fact that the CIMDEF dataset lacks NI
307 data and only provides lower resolution noise and earthquake data, since WA profiles are, as
308 yet, ~~un-interpreted~~published. Thus, reprocessing ~~the~~all NI data was mandatory, at least at
309 stack and post-stack level. Figure 2 shows the processing flow followed to homogenize the
310 display of datasets while preserving the true amplitude (Martínez García, 2019). The software
311 package used for reprocessing was GLOBE Claritas (www.globeclaritas.com/) and the most
312 important steps were related ~~with~~to frequency filtering, amplitude weighting and
313 equalization, Kirchhoff time migration and coherency filtering (Fig. 2). In addition, up to 20
314 multi-trace attribute analysis were tested with the goal to enhance structural and lithological
315 impedance contrasts that allowed to improve the interpretation (Chopra and Alexeev, 2005;
316 Taner and Sheriff, 1977). Although this methodology has been mostly used in sedimentary
317 reservoirs, we have seen that the application of these techniques can enhance the continuity
318 of reflections and help to identify different types of crust, thus easing the interpretation. Some
319 of the boundaries ~~resulting~~of ~~from~~ this attribute analysis (e.g., variance and chaos attribute
320 filters, the former estimating the local variance in the signal and the latter measuring the lack
321 of organization in the reflectivity) are included in the interpretations.

Código de campo cambiado

Con formato: Inglés (Estados Unidos)

322 3.3. Description of the seismic sections

323 The NI datasets included in this paper have already been presented, so the reader will be
324 referred, in every sub-section, to previous publications ~~for that include~~ detailed descriptions of
325 pre-stack processing and interpretations. Here we will just focus on those features that are
326 essential to our interpretation.

327 ~~Geological cross-sections coincident with Reprocessed-reprocessed time~~-migrated sections and
328 their interpretations are presented in figure 3 (ESCIN-1), figure 4 (ESCIN-2), figure 5 (ESCIN-
329 3.3), figure 6 (ESCIN-3.2) figure 7 (ALCUDIA) and figure 8 (IBERSEIS). Migration velocities (Fig. 2)
330 are average crustal velocities~~values~~ as calculated from coincident or nearby wide-angle
331 data~~WA models~~. Depth conversion using migration velocities is also carried out. The
332 description of sections will be done from N to S and from E to W. The CIMDEF dataset will be
333 only described in the discussion (Figs. 9 and 10) as it ~~does not include NI data but~~ is key to
334 understanding the geometry of the mid-crustal discontinuity, its late Variscan reworking and
335 its Alpine reactivation.

336 3.3.1. Cantabrian Zone (ESCIN-1 section)

337 The ESCIN-1 section is a ~130 km long, E-W profile crossing the CZ from its most external part
338 to the Narcea Antiform to the W, in the boundary with the hinterland (WALZ, Figs. 1 and 3). It
339 consists of two slightly overlapping parts, 1.1 and 1.2, separated a few kilometers in the N-S
340 direction. The complete ESCIN-1 section migrated at $v=5600$ m/s (Fig. 2) and its interpretation
341 are presented in figure 3.

342 This section was first described and interpreted over an unmigrated image by Pérez-Estaún et
343 al. (1994). Later works revisited the interpretation, adding travel-time modeling to help on the
344 understanding of the unmigrated data (Gallastegui et al., 1997). The reader is referred to these
345 papers for further details than those provided here.

346 In the upper crust, the western part shows W-dipping reflections that represent the Variscan
347 imbrication, through a thrust ramp (t in Fig. 3), of the basement under the Paleozoic
348 succession sequence (a in Fig. 3), indicating the proximity of the hinterland (WALZ). In fact, a
349 Neoproterozoic, non-metamorphic sequence outcrops in this area, which is probably underlain
350 by an older crystalline basement. One Another prominent W-dipping reflection roughly parallel
351 to t- (ta') crosscuts subhorizontal ones, defining a pattern that might indicate its be providing
352 an slightly out of the plane provenance image of the above mentioned thrust ramp, as this
353 profile lies in the hinge of an arcuate structure, the IAA (Fig. 1b). To the E, the thin skinned
354 tectonics characteristic of this external zone can be interpreted from shallow subhorizontal to
355 W dipping reflections often coincident with outcropping thrusts (bot), as observed in figure
356 3a. The main one among these, running at around 5 s (TWT), is interpreted as the sole thrust
357 of the thin-skinned orogenic wedge (ste). To the W, it gets involved in the crustal ramp (t in Fig.
358 3) observed at the Narcea Antiform, suggesting that it ends down rooting into the upper part
359 of the lower crust (cd). A low reflectivity wedge of undifferentiated basement (be) located
360 between 4-5 and 8.5 s (TWT) exists underneath the easternmost reflections. This may image
361 some pre-Paleozoic basement that is interpreted as upper crust, since the pattern of
362 reflections changes below, suggesting that a significant boundary occurs underneath.

363 The lower crust shows little reflectivity but seems to be present in the interval between 8.5-14
364 s (TWT) in the E and between 8.5 and 12 s (TWT) in the W. It features subhorizontal (irf) and W
365 dipping internal reflectivity to the E, the latter (irf') crosscutting the former reflections. These
366 might represent the imprint of Alpine tectonics over a previously deformed/reflective lower
367 crust. To the W, reflectivity seems to be subhorizontal or dipping to the E (irg). Some of the
368 dipping and arcuated reflectivity observed at the edges of section ESCIN-1 (mig in Fig.3 and
369 thereafter) might be related to the migration effects over a little reflective
370 section discontinuous features and caution should be taken when interpreting it.

371 The Moho along this section (m) is located at nearly 14.5 s TWT (~45 km) in the eastern part,
372 and shallower (12 s TWT, ~36 km) to the W. The crustal thickening observed to the E (h) is
373 probably related with to an out of section image of the crustal Alpine root, better observed in
374 profile ESCIN-2, which is described next.

375 **3.3.2. Cantabrian Zone and Duero Basin (ESCIN-2 section)**

376 The ESCIN-2 seismic line is a 65 km long, N-S section that samples the transition between the
377 CZ and the DB (Fig. 1). Even though this profile was geared to study the Alpine structures, it

378 shows how the Variscan features have been inherited and reactivated during the Cenozoic
379 compression between the Iberian Peninsula and the European plate. The section was first
380 presented by Pulgar et al. (1996). Later on, some authors have used this image to constraint
381 the Alpine structure in the North Iberian Margin (e.g., Fernández-Viejo et al., 2000; Gallastegui
382 et al., 2016). However, only Teixell et al. (2018) used a migrated version (4000 m/s) of this
383 section. Here we present the results of a Kirchhoff time migration at $v=5600$ m/s (Fig. 4).

384 This seismic line shows, in places, a conspicuous reflectivity that allows a straightforward
385 interpretation. To the S end, the upper crust is characterized by high amplitude horizontal
386 reflectivity representing the DB sedimentary sequence (Fig. 4a). It occupies the interval from 0-
387 3.5 s TWT (sa in Fig. 4b, c and d) and appears to be offset by N dipping reflections (tb). The
388 latter have been interpreted as S vergent Alpine thrusts affecting the CZ basement and partly
389 the DB sediments. The rest of the crust is less reflective although N dipping reflectivity (ote),
390 also interpreted as imaging Alpine thrusts on the basis of the clearer stack image (Pulgar et al.,
391 1996), crosscuts shallow subhorizontal weak reflections that represent the Paleozoic
392 sedimentary sequence of the CZ (psd).

393 The lower crust presents higher amplitude reflectivity. In general, a thick band of horizontal
394 reflections located between 7.5 and 12 s (TWT) at the southern part of the profile, bends and
395 dips to the N in the northern part of the line (lce) in response to Alpine compression. Although
396 the stacked section shows that this N dipping reflectivity reaches 14.5 s TWT (Pulgar et al.,
397 1996) the migrated sections (Teixell et al., 2018 and Fig. 4) indicate that these reflections move
398 southward and upward to less than 14 s (TWT), while losing amplitude and coherence. In fact,
399 the geometry of the bottom of the lowermost crust (Moho, m) is deduced on the basis of the
400 geometry of its uppermost part (c), the lower crust internal reflectivity, the stack image (Pulgar
401 et al., 1996), and the amplitude contrasts observed in the attribute analysis (Fig. 4).
402 Furthermore, its depth is solely established on the basis of the position of the strongest
403 subhorizontal reflections to the S.

404 Even though this profile shows the imprint of recent Alpine shortening, no reflections are
405 observed to crosscut the entire crust. In contrast, reflectivity suggests that deformation is
406 decoupled between the upper and lower crust. However, this section is not long and/or
407 reflective enough as to image where the Alpine thrusts (ote) root. Possibly, they merge into
408 the roof of the underthrust CZ lower crust. ~~In addition, the migration effects on the edges of
409 the section produce misleading reflectivity than hinders more detailed interpretations.~~

410 A 1D Vp profile (Fig. 4e) extracted from the coincident WA model (Pulgar et al., 1996) shows a
411 conspicuous velocity increase in the lower crust, at a depth roughly coincident with (c). Depth
412 misfits are due to the effect of the low Vp of the DB sediments not being taken into account in
413 the depth conversion.

414 3.3.3. West Asturian-Leonese Zone (ESCIN-3.3 section)

415 The ESCIN-3.3 profile is part of a ~375 km long crooked offshore seismic line consisting in
416 ESCIN-3.1, 3.2 and 3.3. The latter is 137 km long, parallel to the coast and close to it across the
417 WALZ (Fig. 1). It was first presented by Martínez Catalán et al. (1995) and Ayarza et al. (1998,
418 2004). Later on, its image has been used to constrain the structure of the western North

419 Iberian Margin and that of the transition between the WALZ and the CIZ (Martínez Catalán et
420 al., 2012, 2014). The cross-section presented in figure 5a corresponds to the equivalent
421 onshore transect of this profile.

422 Reflectivity in the upper crust is characterized by the image of Mesozoic sedimentary basins
423 (sa in Fig. 5) related to the extension that led to the opening of the Bay of Biscay. Underneath,
424 W dipping reflections (t) are interpreted as the imprint of the first stages of Variscan
425 compressional deformation in the WALZ (C1 and C2), developing E-vergent thrust faults (Fig.
426 5ab). These affect the pre-Paleozoic basement and root in the upper part (c) of a thick
427 reflective band interpreted as the lower crust (lc) or in a sole thrust (st) that also reaches the
428 lower crust.

429 The lower crust (lc) is represented by a thick band of subhorizontal reflectivity (8-12 s TWT)
430 that thickens (6-12 s TWT) in the westernmost part of the WALZ (CDP 3000) underneath the
431 Lugo Dome, an extensional structure bounded to the E by the Viveiro normal fault (Fig. 5).
432 Then it thins towards the end of the line, when entering the CIZ, coinciding with an area of E
433 dipping sub-crustal reflections (sc) thus defining a Moho offset of ~2 s TWT altogether defining
434 a less clear Moho (m) in this area. The ESCIN-3.3 lower crust seems to feature an internal layer
435 with mantle P-wave velocities when modeled from coincident WA data. Accordingly it was
436 interpreted as consisting of the WALZ lower crust underthrust by the CZ lower crust (Ayarza
437 et al., 1998). This model would compensate the high shortening observed in the upper crust of
438 the CZ, a thin-skinned belt whose sole thrust roots at the contact with the WALZ. ~~However, the~~
439 ~~ESCIN-3.3 WA data need to be revisited as it is a fan profile difficult to model with old~~
440 ~~conventional 2D algorithms.~~ The internal reflectivity of the lower crust shows W dipping
441 reflectors (ire), similar to the ones observed in the upper crust and probably imaging Variscan
442 deformation in the lower crust, either compressional or extensional. They crosscut
443 subhorizontal reflectivity, thus postdating it. A 1D P-wave velocity profile (Fig. 5e) derived from
444 coincident WA data (Ayarza et al., 1998) shows again an important increase in relation to the
445 top of the lower crust (c).

446 Even though migration (v= 5200 m/s, Fig. 2) over discontinuous reflections blurs the seismic
447 section in the edges (mig in Fig. 5), reflectivity never seems to cross-cut the crust and/or the
448 Moho, indicating that deformation is decoupled at upper and lower crustal level. Subcrustal E
449 dipping reflections (sc) are interpreted as the out-of-the-plane image of the Alpine southward
450 subduction of the Bay of Biscay oceanic crust (Ayarza et al., 1998, 2004), which is out of the
451 scope of this paper.

452 The boundary between the WALZ and the CIZ is the Viveiro Fault, one of the most striking
453 surface expressions of Late Variscan extensional tectonics, featuring a decompression of ~4kb
454 or 14 km (Reche et al., 1998). To its W, gravitational collapse of a thickened crust and
455 associated crustal extension and melting have played a key role in the orogenic evolution of
456 the CIZ. However, to the E, crustal re-equilibration after C1 and C2 thickening was less
457 important and igneous activity decreases. Even though this fault itself is not identified in the
458 seismic section (Fig. 4), the reflectivity in general varies on both sides of it, featuring a thinner
459 (9 s TWT vs 12 s TWT) and more transparent crust to the W (Fig. 6). In fact the geometry of
460 some reflections (ef) in the boundary between the WALZ and the CIZ, above the thickest lower

461 | crust, and the subtractive way the sole thrust (~~ste~~) merges with the lower crust (~~ste'~~) seem to
462 | indicate the effect of extensional tectonics, sometimes reactivating compressional structures
463 | (~~ste-ste'~~). Such a reactivation has been described ~~for-in~~ the base of the main thrust sheet in
464 | the WALZ based on structural and metamorphic considerations (Alcock et al., 2009).
465 | Conversely, further to the E of the section, reflectivity probably represents the geometry of
466 | preserved compressional deformation.

467 | 3.3.4. Northern Central Iberian Zone (ESCIN-3.2 section)

468 | The seismic line ESCIN-3.2 is a 97 km long profile, also parallel and close to the coast, ~~and that~~
469 | ~~sampling~~ the relative autochthon to the GTMZ, i.e. the CIZ (Figs. 1 and 6). It was first
470 | described by Álvarez-Marrón et al. (1996) and later by Ayarza et al. (2004). ~~Here it is presented~~
471 | ~~migrated with a v=5200 m/s (Fig. 2). The cross-section presented in figure 6a corresponds to~~
472 | ~~the equivalent onshore transect of this profile and depicts allochthonous sequences not~~
473 | ~~imaged by the NI data.~~

474 | This profile shows, in the upper part, a band of high subhorizontal reflectivity ~~that coincides~~
475 | ~~related to with the location of~~ Mesozoic basins, as in profile ESCIN-3.3 (~~sa~~ in Fig. 6). The rest of
476 | the upper crust is not very reflective although a couple of W-dipping reflections (~~efb~~) rooting in
477 | a thin band of strong reflectivity are observed. These reflections, located in the E of the section
478 | from 4.5 s to 8 s TWT, define a sort of duplex, extensional or compressional, but later
479 | extended, indicating in any case boudinage and crustal thinning. To the W, the upper crust is
480 | very transparent and just a few weak reflections can be observed.

481 | The narrow reflective band at 8-9 s TWT represents the lower crust (~~lc~~), and is the most
482 | striking feature of this profile. ~~This-Its~~ 1 s TWT thick~~ness feature~~ contrasts with ~~that observed~~
483 | ~~in~~ the neighboring ESCIN-3.3 and ~~even~~ ESCIN-1 sections, which show a much thicker lower
484 | crust (4-5 s TWT). Reflectivity in this band is subhorizontal (~~e~~), although somewhat undulated,
485 | while the band itself is slightly inclined to the W. In the E, the Moho (~~m~~) is located ~~at~~
486 | ~~aroundabove~~ 9 s TWT (~~~27 km~~), the shallowest identified so far in the Iberian Massif.
487 | Subcrustal ~~E~~ dipping reflections (~~scd~~) are again associated to the 3D image of the southward
488 | subduction of the oceanic crust of the Bay of Biscay during the Alpine convergence (~~Ayarza et~~
489 | ~~al., 2004) whereas W dipping features might be related with the CZ lower crust (czlc)~~
490 | ~~underthrusted also underneath the easternmost part of the CIZ. They have been already~~
491 | ~~modeled by Ayarza et al. (2004) and will not be further discussed in this paper.~~

492 | This profile samples the northern CIZ, where Variscan crustal thickening during C1 and C2, was
493 | most important. Consequently, later gravitational collapse triggered extensional tectonics and
494 | crustal melting, allowing the intrusion of granites and the development of extensional
495 | detachments (with associated metamorphic offsets). The image of line ESCIN-3.2 shows a
496 | transparent upper crust to the W suggesting that granites occupy most of it, which is
497 | supported by onland geological mapping. Some thrust faults, as those imaged by W-dipping
498 | reflections in profile ESCIN-3.3, probably root ~~here along this section~~ and are represented by
499 | the W-dipping reflections (~~b~~) at the base of the upper crust. However, these were later
500 | flattened and/or reactivated as extensional detachments by crustal thinning (~~ef~~). The
501 | narrowness of the highly reflective lower crust here suggests that crustal thinning was largely
502 | accommodated at this level, as the upper crust has basically the same thickness as in the

503 ESCIN-3.3 line (up to 6-7 s TWT). In addition, crustal melting might have also affected the top
504 of the lower crust. But even though large parts of the crust were melted, reflectivity exists
505 deep in the upper crust, suggesting that crustal melting was not pervasive and/or reflectivity is
506 linked to syn- or late-tectonic features.

507 3.3.5. Southern Central Iberian Zone (ALCUDIA section)

508 The ALCUDIA seismic profile was first presented by Martínez Poyatos et al. (2012) and
509 reprocessed and further interpreted by Ehsan et al. (2014). It is a more than 220 km long, NE-
510 SW seismic profile sampling the CIZ to the S of the ICS down to the boundary with the OMZ
511 (Fig. 1). [Here we presented this section migrated and depth converted with using a v=6200 m/s](#)
512 [\(Figs. 2 and 7\).](#)

513 This profile presents a fairly transparent upper crust [when compared to other nearby sections](#)
514 [\(e.g., IBERSEIS, Fig. 8\) although where scarce reflectivity exists to the S coinciding with is related](#)
515 ~~to~~ the boundary (suture) between the CIZ and the OMZ, namely, the Central Unit (CU; [cu](#) in
516 Fig. 7) and to the presence of vertical folds ([vfb](#)). Some very transparent zones ([ge](#)) appear to
517 be in relation with the existence of granitic batholiths. To the N, the intrusion of these
518 granites, associated to the existence of normal faults ([efd](#)), is one of the evidences of
519 extensional tectonics affecting the southern part of the CIZ. The rest of the upper crust shows
520 weak and discontinuous reflectivity that responds to the existence of vertical folding affecting
521 lithologies with little impedance contrast. In fact, deformation in the upper crust of this part of
522 the CIZ is weak, with absence of low-dipping structures typical of tangential tectonics.

523 The lower crust shows a very different image to that of the upper crust. It is a thick band, of up
524 to 6 s TWT (from 4 s to 10 s), of mostly subhorizontal high amplitude reflectivity ([lce](#)) that at
525 some points appears to be cut across by N-dipping reflectors ([ccf](#)). S-dipping internal
526 reflectivity is also identified although more scarce ([irg](#)). The lower crust thins in the northern
527 end of the profile, near the ICS, where intrusion of granites and other evidences of crustal re-
528 equilibration suggest that extension played a key role. Accordingly, we suggest that the
529 mechanisms that triggered this lower crustal thinning are related ~~with to~~ melting and
530 extension and not with compression, as previously proposed (Ehsan et al., 2014; Martínez
531 Poyatos et al., 2012), and that the N dipping reflectivity observed ~~above in the top of~~ the lower
532 crust ([c](#)) in that area (~~h~~) is the expression of extensional tectonics.

533 One of the most striking features of this profile is the crocodile-like structure affecting the
534 lower crust at around CMP 10000 ([cc and cc'f](#)). This structure, most likely related to Variscan
535 shortening, accommodates an important part of the deformation at lower crustal level and
536 evidences that sub-horizontal reflectivity of the lower crust is pre-Variscan, thus raising the
537 question about its precise age and origin. Despite the presence of this feature in the depth
538 continuation of the suture between the CIZ and the OMZ ([Fig. 7a](#)), reflectivity does not
539 crosscut the whole crust, suggesting the existence of a detachment in the top of the lower
540 crust. This contrasts with the presence in the upper crust (CU) of retro-eclogites with peak
541 metamorphic conditions of 19 kbar and ~550°C (López Sánchez-Vizcaíno et al., 2003). Finally,
542 the Moho boundary ([m](#)) is located at a fairly constant depth (~10 s TWT, i.e. 30-33 km),
543 although the lower crust seems to be preserved and a local crustal imbrication into the mantle
544 is observed underneath the crocodile-like structure. [A 1D Vp profile \(inset in Fig. 7d\) from a](#)

545 coincident WA-data model (Palomeras et al, in press) shows a conspicuous increase of values
546 along more than 15 km starting in the top of the lower crust (c) thus supporting its important
547 thickness.

Con formato: Fuente:
(Predeterminado) +Cuerpo (Calibri)

548 3.3.6. Central Iberia, Ossa-Morena and South Portuguese Zones (IBERSEIS section)

549 The IBERSEIS seismic line was first presented by Simancas et al. (2003). A number of later
550 works added information and details to its interpretation (e.g., Carbonell et al., 2004;
551 Schmelzbach et al., 2007, 2008; Simancas et al., 2006). This section crosses the southernmost
552 CIZ, the whole OMZ, and most of the external SPZ (Fig. 1). It samples two major boundaries
553 interpreted as suture zones: that between the CIZ and the OMZ (CU, Azor et al., 1994) and the
554 one bounding the OMZ and the SPZ, which has been largely affected by younger Carboniferous
555 events (Pérez-Cáceres et al., 2015). The IBERSEIS profile structurally overlaps the ALCUDIA
556 profile along ~30 km, but it is displaced some 50 km to the W. A cross-section along this
557 transect together with its interpretation is shown in figure 8 after migration at $v=6000$ m/s
558 (Fig. 2) are shown in figure 8.

559 This section is ~300 km long and features an outstanding reflectivity at upper and lower crustal
560 levels. In the upper crust, a wealth of N dipping reflections (ta in Fig. 8) image a S verging
561 thrust and fold belt. In the SPZ, these are most reflective events and probably related with
562 to normal faults derived from the extension that led to the opening of the Rheic Ocean and
563 were later reactivated as thrusts during the Late Carboniferous compression. Some authors
564 link the highest reflective features to the middle Carboniferous volcano-sedimentary complex
565 (Schmelzbach et al., 2008), which might have used these fractures as a conduit, thus enhancing
566 reflectivity. In the OMZ, N dipping reflections probably image Variscan thrust faults (ot) as
567 some coincide with such mapped structures. Their lesser reflectivity might indicate the lack of
568 involvement in the thrusts of lithologies that increase the impedance contrast.

569 Upper crustal reflectivity in both, the ZOM and SPZ, does not cross to the lower crust, rooting
570 at a mid-crustal level that, in the SPZ is transparent and does not have any particular
571 expression itself but does coincide with the top of the lower crust (c). However, in the OMZ, a
572 reflective layer exists at this depth (irbb): it has been defined as the IBERSEIS reflective body
573 (IRB, Simancas et al., 2003), a 140 km long, high velocity conductive feature (Palomeras et al.,
574 2009) that is supposed to represent an early Carboniferous mantle-derived intrusion. Its origin
575 has been related to mantle plume activity that thinned the lithosphere and extracted mantle-
576 derived melts from the ascending asthenosphere (Carbonell et al., 2004). Its surface expression
577 are intraorogenic transtensional features (Rubio Pascual et al., 2013; Simancas et al., 2006).
578 Alternatively, Pin et al. (2008) have suggested, based on geochemical constraints, a tectonic
579 scenario of slab break-off for this feature. Internal reflectivity along the IRB is mostly
580 subhorizontal, probably due to the effect of the intrusion along a subhorizontal detachment,
581 and evidences little imprint from Variscan deformation. The body is slightly inclined to the S, at
582 odds with the detachment being the sole thrust of the OMZ upper crustal imbricates. Perhaps
583 it was, but its inclination changed during subsequent deformation, as later suggested.

584 The lower crust shows slightly different patterns in the CIZ and OMZ on one side and the SPZ
585 on the other. In the southernmost part of the CIZ and northern OMZ, N and S dipping
586 reflections define a wedge (cc) that might be the western continuation of the crocodile-like

587 | structure observed in the ALCUDIA seismic line in an equivalent structural position (~~ccf~~ and ~~g~~ in
588 | Fig. 7). In this section, the limited crustal imbrication into the mantle identified in the ALCUDIA
589 | line (~~cc'~~ in Fig. 7) is not observed, perhaps because it only occurs further to the N or E. This
590 | structure may be the reason why the IRB is shallower at this point, indicating that the latter is
591 | older than the crocodile compressional feature. The rest of the lower crust shows S dipping (~~d~~)
592 | and sub-horizontal (~~lce~~) reflectivity that does not exhibit clear crosscutting relationships, thus
593 | hindering their interpretation. However, near the boundary with the SPZ, this reflectivity
594 | seems to be affected by N dipping features (~~irf~~) overprinting them. In the SPZ, the lower crust
595 | shows a more homogeneous image, with subhorizontal reflectivity (~~lce~~) that is often cut across
596 | by longer scale S dipping features (~~lcth~~) that postdate them. The latter probably represent
597 | fractures that firstly accommodated the extension linked to the opening of the Rheic Ocean
598 | and were then reactivated as thrusts during the late Carboniferous compression and collision
599 | of the SPZ basement with the OMZ. The most conspicuous of these reflections (~~lcth'~~) cuts the
600 | IRB in its southern part and seems to offset the lower crustal upper boundary between the SPZ
601 | and the OMZ. Two 1D Vp profiles derived from coincident WA-data (Palomeras et al., 2009)
602 | and shown as insets in figure 8d indicate a velocity increase starting at the top of lower crust
603 | (c) and along the IRB.

604 | Even though the lower crust in the OMZ and SPZ shows dipping features, none of them crosses
605 | to the upper crust, thus rooting at a mid-crustal level as does the upper crustal reflectivity. This
606 | implies again the existence of a discontinuity (~~c~~) in the mid-crust.

607 | Despite of crossing two suture zones and imaging part of a crocodile-like structure, the
608 | IBERSEIS profile shows a fairly flat Moho (~~m~~) located at ~10 s TWT, the same apparent depth as
609 | in the ALCUDIA line (30-33 km). Its signature is very clear underneath the SPZ and a bit blurry
610 | below the IRB.

611 | 4. Discussion

612 | Simancas et al. (2013) already undertook an integrated interpretation of most of the seismic
613 | sections presented here focusing on, i) the accommodation of orogenic shortening at crustal
614 | scale, (ii) the relationships between convergence, crustal thickening and collisional granitic
615 | magmatism, and (iii) the development of the Iberian Variscan oroclinal. In this paper the same
616 | sections are presented, but they have been reprocessed at stack level and time migrated using
617 | a Kirchhoff algorithm. In addition, two extra sections that image the alleged mid-crustal
618 | discontinuity after the Alpine reactivation are taken into account. The first one is the N-S
619 | ESCIN-2 NI dataset (Fig. 4), in the CZ, where this discontinuity has remained untouched during
620 | late Variscan evolution but was reactivated during the Alpine Orogeny. The second one results
621 | from the CIMDEF experiment, carried out in the CIZ across the ICS, where the mid-crustal
622 | discontinuity has probably been affected by crustal melting during the Late-Variscan extension
623 | and by later Alpine reactivation. The latter sections somehow fill the gap existing in Simancas
624 | et al. (2013).

625 | Figures 3 to 8 represent an effort to show a homogeneous seismic image of the Iberian Massif
626 | crust that eased its integrated interpretation. Next, we discuss the main observed features,
627 | their implications and how they contribute to the understanding of the structure and evolution
628 | of the Iberian Massif, adding constraints to the origin of the elevation of the central Iberian

629 Peninsula. Figure 9 presents a simplified sketch of the crustal layers observed in the Iberian
630 Massif. Figure 10 shows a compendium of the position of the mid-crustal discontinuity and the
631 Moho depth (in TWT) along the entire Iberian Massif as deduced from seismic NI data together
632 with a map of the entire Iberian Peninsula Moho depth (Palomeras et al., 2017) that includes
633 the position of the seismic profiles for comparison. We will refer to these figures along most of
634 the discussion.

635 A particular feature of the SW Iberian Massif is the great importance of out-of-section, mainly
636 left-lateral shear zones associated to its suture boundaries. They displaced central and
637 northern Iberia to the NW with respect to southern Iberia (Simancas et al., 2013). The seismic
638 sections do not provide constraints about this movement, as it is perpendicular to their layout.
639 Thus, interpretations in these areas must be taken with caution.

640 **4.1. The upper crust in the Iberian Massif: a depth image of outcropping geology**

641 Most of the seismic sections display a moderate to thick upper crust (4 to 8 s TWT, Fig. 9), with
642 very variable reflectivity. Reflections ~~coincide with have been confidently related to~~
643 outcropping Variscan structures ~~and thus, a link has been established~~. As such, N dipping
644 reflectivity in the SPZ and the OMZ is related to S vergent folds and thrust faults mapped in the
645 surface. W dipping reflections in the CZ are related to mapped thin-skinned thrusts. The same
646 type of reflectivity observed in the WALZ, reaching deeper levels in the crust and rooting in the
647 lower crust, has been addressed as evidence of thick-skinned thrust tectonics, which in the
648 hinterland affects the pre-Paleozoic basement. Particularly interesting is the upper crustal SPZ
649 seismic image in contrast with that of the CZ, both representing external zones. While in the
650 latter thrusts are observed to root in a shallow sole detachment, in the former one
651 reflections/thrusts root in the lower crust. This feature will be discussed in the next section.

652 Only a few seismic profiles feature a transparent upper crust. Lack of reflectivity has been
653 ~~related-addressed~~ to low fold data (ESCIN-1 and ESCIN-2, Figs 3 and 4), and most importantly
654 to the existence of a re-equilibrated upper crust having recorded large amounts of partial
655 melting, as shown by voluminous outcropping granitoids (ESCIN-3.2 and N of ALCUDIA, Figs. 6
656 and 7). The existence of vertical folds affecting little reflective monotonous lithologies also
657 results in a fairly transparent upper crust in most of the ALCUDIA section (Fig. 7).

658 None of the upper crustal reflections observed and interpreted in the presented Iberian Massif
659 NI seismic sections seems to cut across the whole crust, always rooting in a sole thrust (parts
660 of ESCIN-1 and ESCIN-3.3, Figs. 3 and 5) or in the lower crust (the rest of them).

661 **4.2. The lower crust in the Iberian Massif: accommodation of shortening, extension and its** 662 **nature**

663 The Iberian Massif dataset presented here shows a very coherent image of the lower crust. Its
664 reflectivity is high and usually subhorizontal. However, cross cutting relationships with later
665 features of opposite dips evidence a multi-phase origin for this reflectivity.

666 The SPZ, OMZ, WALZ, CZ and the southern CIZ show that this part of the crust is also thick (4 to
667 6 s TWT). However, in NW Iberia and the northern part of the ALCUDIA section (Figs. 1, 6 and
668 7), the few existing NI profiles indicate that the in the northern CIZ, the lower crust is much

669 thinner (1 to 2 s TWT) and irregular (ESCIN-3.2, Figs. 6 and 9). This thin lower crust has been
670 observed in the area characterized by outcropping syn-collisional granitoids (zone II of
671 Simancas et al., 2013). These witness the onset of crustal re-equilibration processes triggered
672 by gravitational collapse, extension and crustal melting during the Late Carboniferous. The
673 straightforward conclusion is to attribute the architecture of this lower crust to late Variscan
674 orogenic extension, which features at the surface high metamorphic offsets (chlorite to
675 sillimanite zone, (Díez Balda et al., 1995) and melting, implying that crustal thinning has been
676 mostly accommodated by its lowermost part.

677 Nevertheless, a gap of crustal-scale NI data exists in most of the northern CIZ. The CIMDEF
678 noise autocorrelation profiles (Figs. 1 and 9 and) show a thick (~5 s TWT) lower crust in most
679 of this area, which essentially corresponds to the CIZ (Andrés et al., 2019, 2020)(~~Andrés et al.,~~
680 ~~2019 and this volume~~). This is in conflict with the NI sections ESCIN-3.2 and northernmost
681 ALCUDIA, where the highly reflective lower crust is less than half as thick. However, granitoids
682 are probably scarce in the Variscan basement hidden under the DB, which can then present a
683 thick lower crust. But in and near the ICS, a rather continuous internal reflection in the lower
684 crust could be interpreted as its top part (Figs. 9 and 10), thus indicating that crustal thinning
685 and melting, observed in the surface, has also affected the lower crust (Andrés et al.,
686 2020)(~~Andres et al., this volume~~).

687 Extension in the northern CIZ occurred simultaneously with shortening in the SW Iberian
688 Massif. According to Simancas et al. (2013) this suggests that the tectonic stresses would be
689 dominantly compressional, still induced by ongoing collision. In fact, gravitational instabilities
690 in a thickened crust should mostly be affecting the upper crust. In this context, theoretical
691 models (Royden, 1996; Seyferth and Henk, 2004) indicate that beneath the areas of extension
692 in the upper crust, shortening may prevail in the lower crust. This mechanism is an efficient
693 way for syn-convergent exhumation of deep rocks.

694 Indeed, from a regional tectonic perspective, compression was active till the end of the
695 Variscan orogeny, and at times, clearly simultaneous with extension (C3 and E2 overlapped in
696 the interval 315-305 Ma; Martínez Catalán et al., 2014). But it is clear that extension affected
697 the lower crust, as it appears thinned in areas of transparent, extended molten crust (ESCIN-
698 3.2 and ALCUDIA sections, Figs. 6 and 7). However, the irregular pattern observed in the ESCIN-
699 3.2 lower crust might indicate the existence of folds in this re-equilibrated layer, witnessing the
700 simultaneity of extension and compression even at lower crustal level (Fig. 6). In addition, we
701 cannot rule out that these undulations represent boudinage (i.e. extension) or Alpine folding,
702 although we consider the latter less likely.

703 In the ALCUDIA section, the imaged part of the CIZ underwent only moderate upper crustal
704 shortening (Martínez Poyatos et al., 2012). According to Simancas et al. (2013), the thick
705 laminated lower crust, representing pre-Pennsylvanian (most probably pre-Variscan) ductile
706 deformation, appears deformed in two sectors near both ends of the profile, concentrating
707 shortening in discrete structures that compensate the upper crustal deformation. The first of
708 them is the very conspicuous crocodile-like structure observed in the southern end, and also
709 imaged in the northern part of the IBERSEIS line (Fig. 9b). This structure mimics localized
710 crustal indentation of the OMZ into the CIZ, producing a local underthrusting of the latter to

711 the S that is still (partly?) preserved. Indentation generated tectonic inversion of the Los
712 Pedroches early Carboniferous basin (Simancas et al., 2013) and bending of the overlying
713 upper crust, as seen by the uplift of the IRB, both of which predate the imbrication. The Los
714 Pedroches batholith intruded above at 314-304 Ma in an extensional setting (Carracedo et al.,
715 2009), postdating the age of the wedge as no further deformation affected the batholith.
716 Indeed, the crocodile-like feature must represent early Carboniferous Variscan compressional
717 deformation and must account for part of the shortening observed at upper crustal level.

718 However, to the NE of this section, a ramp-and-flat geometry has been interpreted as a major
719 lower crustal thrust (Martínez Poyatos et al., 2012; Simancas et al., 2013) that helps to
720 compensate upper and lower crustal shortening. However, the highly reflective lower crust is
721 not repeated in the hanging wall to the structure, so that a subtractive character is a
722 reasonable alternative. As stated above, the thin lower crust to the N of the ramp seems to be
723 clear evidence of lower crustal thinning (Fig. 9b), supported by the fact that it underlies an
724 area of upper crustal extension, the Toledo gneiss dome, characterized by normal faulting and
725 pervasive partial melting (Barbero, 1995; Hernández Enrile, 1991). Regardless of how much
726 shortening that area accommodated during crustal thickening and even though the observed
727 ramp could be a former thrust fault reactivated during later extension, the present image of
728 the lower crust does not suggest compensation of upper crustal shortening. In fact, the lower
729 crust in the ALCUDIA section is anomalously thick elsewhere (up to 6 s TWT, 18 km) suggesting
730 the possibility of ductile thickening previous to the extension that triggered thinning at its
731 northern part.

732 In the IBERSEIS profile, lower crustal dominant reflectivity is also subhorizontal but disrupted
733 by N and S dipping features (Fig. 9b). Whereas in the OMZ these features usually dip to the N,
734 as do the upper crustal reflections representing Variscan thrusts, in the SPZ they surprisingly
735 mirror the upper crustal Variscan thrusts, dipping to the S. Furthermore, one of these features,
736 placed close to the boundary with the OMZ, affects almost the entire lower crust.

737 Orogenic orthogonal shortening in the OMZ upper crust has been estimated in 120 km (~57%)
738 and in the SPZ around 80 km (~45%) or even less (Pérez-Cáceres et al., 2016). According to
739 Simancas et al. (2013), the crocodile structure and a not observed associated northward
740 subduction of the OMZ might account for this shortening in the OMZ. Similarly, in the SPZ, the
741 lower crustal imbricated structures represent only ~ 20 km of shortening so that according to
742 these authors, detached lower crustal subduction along the OMZ/SPZ might have
743 accommodated the other 60 km.

744 | In this regard, we suggest that the present day SPZ crustal image represents its decoupled
745 early Carboniferous extension and later compression. This evolution would have erased any
746 evidences of previous (pre-Carboniferous) subduction, and forced the SPZ to thin during
747 extension. i.e., the lower crust had to decrease its thickness ductilely, perhaps first in a more
748 or less distributed way and later through localized shear zones (brittle or not depending on the
749 depth) as it became shallower. However, the upper crust could have preserved most of its
750 original thickness, as the developing basins associated to extension would have been
751 constantly fed by sediments and igneous extrusions and intrusions (like the IRB in the OMZ).
752 Later compression would have folded and thrust the upper crust, and also thickened the

753 lower crust. A few lower crustal normal shear zones might have developed during extension
754 and then be reactivated as ductile thrusts during compression. Those are today observed as S
755 dipping reflections that disrupt the subhorizontal previous reflectivity in the lower crust and
756 mirror thrusts in the upper crust. Accordingly, distributed ductile deformation and thrusting
757 might have thickened the lower crust back to its original (or simply stable) thickness in the SPZ
758 and elsewhere, something that cannot be measured but would need to be accounted for when
759 comparing shortening at upper and lower crustal level. The resulting seismic image of the SPZ
760 would then be that of an extended and then inverted margin, with mirroring reflectors in the
761 upper and lower crust merging in a mid-crustal discontinuity and providing a seismic image
762 different to that of a typical foreland thrust and fold belt (e.g., CZ; Fig. 3). This evolution differs
763 from that of a hyperextended magma-rich margin as stretching of the upper and lower crust is
764 not coupled and faults do not cut across the crust and penetrate down into the mantle. In any
765 case, the S dipping lower crustal reflections, active during the Late Carboniferous, postdate the
766 sub horizontal reflectivity of the lower crust. It is worth mentioning here that the SPZ seismic
767 image is identical to that of the Rhenohercynian Massif in Germany (Franke et al., 1990;
768 Oncken, 1998) suggesting a similar evolution.

769 The discussion above shows that the lower crust in the Iberian Massif is thick, except when it is
770 affected by late orogenic extension. The mechanisms that produced lower crustal thickening
771 are probably related ~~with to~~ compressional deformation, mostly ductile. Continental
772 underthrusting of the CZ underneath the WALZ (Ayarza et al., 1998, 2004) and part of the CIZ
773 (Fig. 6, 9 and 10), indentation of the OMZ in the CIZ (Figs. 9 and 10b) and Variscan thrust-like
774 structures probably played an important role. In addition, the latter help to constrain the age
775 of the subhorizontal reflectivity. Frequent disruption of subhorizontal lower crustal lamination
776 by Variscan (late Carboniferous) dipping features indicates that the lamination developed prior
777 to Variscan compressional deformation. What this lamination represents is still an open
778 question.

779 Many vertical incidence seismic reflection profiles worldwide have shown reflective lower
780 crusts (e.g., Meissner et al., 2006; Wever, 1989). Lower crust seismic lamination has been
781 often related to late orogenic extensional events (Meissner, 1989). In the Iberian Massif,
782 surface geology shows that late orogenic extension affects the upper crust, mainly in areas of
783 large previous thickening. ~~In~~ But in contrast to the ~~latter's~~ author's models, important thinning
784 of the lower crust takes place in those areas (ESCIN-3.2 and northern ALCUDIA, Figs. 6, 7 and
785 9). Certainly, lower crustal lamination might come from underplating eased by extension in
786 magma rich margins (Klempner et al., 1986). But also, ductile deformation is a very likely
787 source of lower crustal lamination. Dipping events observed in the lower crust crosscutting a
788 strong banded reflectivity represent the latest orogeny-related shortening, which will be
789 further flattened and horizontalized in the next orogeny. Continuous superposition of
790 deformational events at lower crustal level managed to decrease the dip of
791 structural/lithological markers and define a subhorizontal fabric. These deformation
792 mechanisms can generate structures with a strongly defined anisotropy, which result in a
793 strongly laminated lower crustal fabric (Carbonell and Smithson, 1991; Okaya et al., 2004).
794 Accordingly, a laminated lower crust may represent an overly reworked lower crust that has
795 been ductilely deformed over several orogenies. Opposite to the model by Meissner (1989),
796 such a horizontal reflectivity is observed along the Iberian Massif in areas where late orogenic

797 extension is absent or weak: the SPZ (Avalonia), the OMZ (peri-Gondwana), the not extended
798 CIZ, the WALZ and the CZ (Gondwana). Thus, we suggest that strong lamination in the deep
799 crust is probably a global characteristic of reworked lower crusts not affected by late orogenic
800 extension in the latest orogeny.

801 **4.3. The Moho and crustal thickness in the Iberian Massif**

802 The crust-mantle boundary, i.e., the Moho, is basically flat in the Iberian Massif except where
803 affected by the Alpine tectonics (Fig. 10). This is rather surprising as the lower crust seems to
804 be quite well preserved, suggesting that the Moho geometry has been flattened out through
805 slow, not invasive, readjustments.

806 Flat Mohos imply the existence of either isostatic and/or thermal, late to post-orogenic
807 processes that have managed to eliminate crustal roots (Cook, 2002). NW Iberia was affected
808 by late Carboniferous extension that heated and reworked the CIZ, possibly without significant
809 mantle involvement (Alcock et al., 2009, 2011), but producing crustal thinning (Palomeras et
810 al., 2017: see Moho depth map in Fig. 10). Thick and thermally mature crusts might experience
811 lateral flow of its low-viscosity deeper part that contributed to reduce crustal roots (Seyferth
812 and Henk, 2004). This process might have partly occurred in the CIZ sampled by the ESCIN-3.2
813 section (Fig. 6) where an outstanding change in lower crustal thickness and signature exist,
814 manifested by a thinner and very reflective lower crust in contrast to that to the E, in the WALZ
815 (1 vs 3 s TWT) or to the S, in the ALCUDIA profile (1 vs 5-6 s TWT). In the ~~latter~~former, the
816 Variscan crust is still thick even though it experienced late-Variscan extension in its western
817 part and the whole area was slightly affected offshore by the extension linked to the opening
818 of the Bay of Biscay. In fact, underthrusting of the CZ lower crust is still preserved in the
819 eastern CIZ (Figs. 6 and 9).

820 In the SW Iberian Massif, a thick laminated lower crust is still observable while the Moho depth
821 is fairly constant (~10s TWT). Carboniferous-to-Permian isostatic rebound in response to
822 tectonic thickening, erosion and localized Permian thermal readjustments must have
823 contributed to flatten the Moho. However, seismic reflections show that crustal imbrication
824 into the mantle has locally survived post-orogenic Moho resetting. This indicates that isostatic
825 equilibrium has been reached in a long wavelength scale, but that local features can still
826 remain if they ~~can beare~~ supported by the crustal strength and do not pose an isostatic
827 constraint.

828 **4.4. The (missing) middle crust in the Iberian Massif (and elsewhere?)**

829 One of the highlights of this work is the lack of a seismic layer that can be identified with the
830 middle crust. But, what is the middle crust?

831 From a metamorphic point of view, the middle crust could be ascribed to the mesozone, which
832 may be correlated ~~with-to~~ the amphibolite facies, whose temperature ranges between 400-
833 500 and 600-800°C, the precise limits depending on the pressure (Spear, 1993). In addition, the
834 epizone, between 200-250 and 400-500°C and typically represented by the greenschist facies,
835 is also a metamorphic entity which develops during metamorphism under several kilometers
836 of anchi- and no metamorphic rocks. The depths corresponding to these temperature intervals

837 vary with the geothermal gradient. For a Barrovian gradient, typical of a continental crust
838 undergoing collision, the depths for epizone and mesozone can be estimated around 10-20
839 and 20-30 (\pm 5) km respectively. ~~However~~Although, the boundaries of these metamorphic
840 zones might have a gravity, i.e. density signature, ~~they lack but not~~ a seismic one. Furthermore,
841 epi-, meso- and catazonal rocks outcrop everywhere in any eroded orogenic belt defining a
842 very complex pattern that contrasts with the simplicity of seismic images. This implies that,
843 implying that they do not represent a the metamorphic middle-crust does not need to coincide
844 with a hypothetic seismic middle crust, the former often ~~but actually occur in~~ being part the
845 upper crust in ancient orogens.

846 Seismic data are sensible to velocity and density contrasts and not to the absolute value of
847 velocity and density. If a sharp contrast exists, a reflection appears, but metamorphic zones
848 usually lack sharp boundaries. So fFrom a seismic point of view, a middle crust ~~could should~~ be
849 a crustal level bounded in its upper and lower parts by ~~characteristic~~ reflections ~~that~~
850 indicative of the existence of important impedance contrasts at its top and bottom. In this
851 regard, only the IRB, intruded between the upper and the lower crust (Carbonell et al., 2004;
852 Simancas et al., 2003), and providing conspicuous velocity contrasts (Palomeras et al., 2009,
853 2011) fulfill that requirement. However, it is most probably an intrusion emplaced at a mid-
854 crustal discontinuity and does not represent the middle crust.

855 WA reflection seismic data from the northern Iberian Massif have often resulted in
856 multilayered models despite weak evidences of continuous reflectivity at these levels (Ayarza
857 et al., 1998; Fernández-Viejo et al., 1998, 2000; Pedreira et al., 2003). Even though local
858 velocity contrasts capable of providing weak and patchy reflectivity contrasts exist at different
859 crustal depths (e.g., thrust faults and normal detachments may represent lithological
860 boundaries with a noticeable velocity contrast), these are not orogen-scale features but local
861 reflectors. ~~However, m~~Many of these reflections ~~observed, have been~~ extrapolated and
862 interpreted and/or extrapolated as middle crust in seismic WA datasets, despite of ~~belong in~~
863 fact to the being part of upper crust, when compared with NI data, i.e., lie above the mid-
864 crustal discontinuity (e.g., Vp increase at ~10 km in Fig. 5e). In this regard, the short
865 wavelength heterogeneities of the crust ~~can behave been often seen~~ considered by low
866 resolution WA datasets as laterally continuous features (Levander and Holliger, 1992),
867 something that has led us to wrong models.

868 According to the above we argue that, in the Iberian Massif, no seismic middle crust can be
869 identified. In the hinterland, reflectors imaging deformation in the upper crust root in the top
870 of the lower crust. Only in ESCIN-1, which depicts the thin-skinned deformation of the CZ,
871 thrust faults root in a sole thrust and one could argue that the basement underneath these
872 shallow reflections represents the middle crust. But in the shallower part, to the E, early
873 Paleozoic and Neoproterozoic sediments occur on both sides of the sole thrust. Also, in the
874 deeper parts, to the W, the previous crystalline basement is probably involved in imbrications
875 affecting the upper crust. Thus, in our opinion, that non-reflective basement represents the
876 seismic upper crust.

877 In the Iberian Massif, the Paleozoic was deposited unconformably above Neoproterozoic
878 sediments which could be considered as its basement, but these were not metamorphic then.

879 Only in the OMZ, greenschist to amphibolite facies Neoproterozoic represents the Cadomian
880 basement, but it cannot be distinguished from the overlying Paleozoic metasediments in the NI
881 profiles. An even older crystalline basement of felsic composition exists, as indicated by
882 inherited zircons of 830-2000 Ma found in Ediacaran orthogneisses, Lower Ordovician
883 volcanics and Variscan granitoids that resulted from partial melting of such a basement
884 (Fernández-Suárez et al., 1998; Montero et al., 2007; Villaseca et al., 2012). Again, its upper
885 boundary is not imaged on NI profiles. These data also suggest that, in the Iberian Massif,
886 there are no crustal intervals that can be related ~~with to~~ a seismic middle crust. Decoupling of
887 reflectivity, i.e. deformation, at a mid-crustal level ~~led us to~~ define just an upper and a lower
888 crust.

889 4.5. Significance of a mid-crustal discontinuity: the Conrad discontinuity?

890 Inspection of the Iberian Massif NI seismic dataset leads us to conclude that an orogenic-scale
891 mid-crustal discontinuity exists. This surface does not always provide a clear reflection, as in the
892 SPZ, but it is clearly defined by the geometry of the upper and lower crustal reflections,
893 asymptotically merging into it. The discontinuity coincides with the top of the lower crust,
894 which is often much more reflective than the upper crust ~~and features a Vp increase~~.
895 Furthermore, this discontinuity has probably acted as a detachment for Variscan deformation
896 in the hinterland of the orogen and in the SPZ. However, in the CZ, the transition between
897 upper and lower crust is poorly defined, ~~in accordance with the fact that~~ ~~as most of~~ its
898 basement was not affected by Variscan tectonics. ~~There, a detachment level interpreted as the~~
899 ~~sole thrust of the thin-skinned wedge occurs above the lower crust, and no deformation~~
900 ~~decoupling is identified above or below this feature.~~

901 Simancas et al. (2013) already described this discontinuity on the basis of the asymptotic
902 geometry of the SPZ faults towards the middle of the crust. These authors concluded that its
903 depth greatly varies when reaching suture boundaries, where the discontinuity roots. Although
904 we do not observe a subduction zone in the reworked elusive suture between the SPZ and the
905 OMZ (Pérez-Cáceres et al., 2015), and interpret the OMZ/CIZ suture as an indentation between
906 two continental crusts, triggering imbrication into the mantle of the latter (crocodile
907 structure), we agree that this discontinuity would have eased the decoupling of the Iberian
908 crust, allowing subduction of its lower part while the upper part was deformed by folds and
909 thrust faults. ~~In fact, t~~ This is clearly observed in the Alpine northward subduction of the Iberian
910 Massif lower crust underneath the CZ (ESCIN-2, Fig. 4) and also, in the Pyrenees, where a
911 detached Iberian lower crust subducts to the N (Teixell et al., 2018). ~~But i~~ in the Iberian Massif,
912 the complexity of Variscan tectonics and late-Variscan crustal re-equilibration has mostly
913 removed evidences of a such mechanisms, ~~but although~~ a comparable example has been
914 preserved in the NW: the thick lower crust imaged by ESCIN-3.3 is interpreted as
915 underthrusting of the CZ lower crust under that of the WALZ (Ayarza et al., 1998; Martínez
916 Catalán et al., 2003, 2012, 2014) ~~and even reaching the CIZ, as shown in profile ESCIN-3.2.-~~

917 ~~Some authors have interpreted the Iberian Massif~~ The mid crustal discontinuity ~~has been~~
918 ~~interpreted~~ as the brittle-ductile transition (e.g. Ehsan et al., 2014; Simancas et al., 2013).
919 Indeed, it bounds a lower crust, highly reflective and ductilely deformed from the upper crust.
920 However, Variscan ductile deformation occurs also above the discontinuity ~~in the entire and is~~

921 | ~~a general feature of the whole~~ Iberian Massif, ~~with the~~ exception of ~~for~~ the CZ. ~~On the other~~
922 | ~~side, if~~ we deal with present deformation mechanisms, it is unlikely that the brittle-ductile
923 | transition, which depends on the values of P and T, coincide with the described discontinuity,
924 | because, i) it does not necessarily imply an impedance contrast (Litak and Brown, 1989), and
925 | ii) according to figure 10, the depth of the discontinuity varies from 4 s TWT (~12 km, ALCUDIA
926 | section) to 8 s TWT (~~~2~~14 km, ESCIN-2 section) which would imply unrealistic variations on P
927 | and T in the present crust. Accordingly, a different interpretation must be sought.

928 | The Iberian laminated lower crust is probably very old. Granulites dredged in Mesozoic
929 | sediments of the Cantabrian margin have yielded ages of up to 1400 Ma (Capdevila et al., 1980
930 | and references therein). Even older values have been obtained for the Galicia Bank granulites
931 | ~~from the Galicia Bank~~ (Gardien et al., 2000), which featured Ar-Ar ages of up to 2500 Ma.
932 | These granulites have been deformed ductilely during several orogenies. ~~But r~~Rocks lying
933 | above the lower crust, whatever their nature ~~is~~, are separated from it by a discontinuity that
934 | fosters decoupled deformation between both crustal layers. Accordingly, ~~we think that~~ the
935 | observed mid-crustal discontinuity probably represents a rheological boundary that separates
936 | rocks that have been deformed differently. Thise boundary, located at the top of the lower
937 | crust, represents a velocity ~~contrast-increase~~ as the latter is probably composed of dense
938 | granulites and includes relatively abundant basic rocks, which makes it easily identifiable in NI
939 | and WA seismic sections.

940 | The geometry of this discontinuity and its depth, together with that of the Moho (Fig. 10),
941 | provide insights of the evolution of the Iberian Massif. Along the SW Iberian Massif, the mid-
942 | crustal discontinuity is sub-horizontal and lies at a depth between 4-6 s TWT. In the OMZ, the
943 | intrusion of the IRB allows to establish its depth in the top or the bottom of this feature, but in
944 | average, its location would fit the above given values. However, in the center and NW, the
945 | position of the discontinuity varies, deepening down to 8 s TWT (Figs. 9 and 10).

946 | The low resolution noise autocorrelation models obtained along the CIMDEF profile shows
947 | confusing results along the central Iberian Massif. ~~In central Iberia~~There, the mid-crustal
948 | discontinuity might lie at 5-6 s TWT, deepening around the ICS to 8 s TWT as it has been
949 | affected by pervasive extension and melting, thus defining a thin lower crust (2 s TWT, ~6 km,
950 | Figs. 9 and 10). Accordingly, this feature appears redefined in this area, and now follows the
951 | geometry of the ICS batholith. The change in the depth and geometry of this discontinuity and
952 | the thinning of the lower crust might have allowed coupled deformation, letting part of the
953 | upper crust to the S of the ICS to underthrust it (Andrés et al., 2020)~~(Andrés et al. this volume)~~.
954 | This would foster the 400-500 m topographic change between the N and S foreland basins of
955 | this Alpine mountain range (Fig. 9). In fact, Simancas et al. (2013) argues that coupled crustal
956 | deformation takes place when a relatively weak lower crust exists something that might well
957 | represent the context of the ICS. The resulting geometry of this Alpine reactivation and its
958 | topographic imprint is different to that observed to the N, in the CZ, where late orogenic
959 | extension and melting does not exist and the mid-crustal discontinuity has been preserved.

960 | On the other hand, the lower crust imaged along the CIMDEF transect presents a conspicuous
961 | internal reflection that could also be interpreted as the top of the lower crust (Andrés et al.,
962 | 2020). If this were the case, the lower crust would be even thinner along the entire section,

963 matching the characteristics observed to the N of the ALCUDIA section and in the ESCIN-3.2
964 | line. In any case, we argue that the mid-crustal discontinuity and the lower crust ~~we are seeing~~
965 | in the CIMDEF profile are both probably reworked by extension but not totally re-equilibrated
966 | and thus, its seismic image is confusing to the N and S of the ICS. Moho depth models (Fig. 10)
967 | derived from shear wave tomography (Palomeras et al., 2017) indicate that along the CIMDEF
968 | profile the crust is thin (except in the Alpine root) but not as much as in NW Iberia, so that
969 | lower crustal extension and re-equilibration may have not been as intense as in the GTMZ and
970 | CIZ of the NW Iberian Massif.

971 The most outstanding change in the mid-crustal discontinuity architecture appears in NW
972 | Iberia, along the ESCIN-3.2 profile. This section features the thinnest crust (9 s TWT)
973 | accompanied by the thinnest lower crust (~1 s TWT). The mid-crustal discontinuity lies at 8 s
974 | TWT in contrast to the depth where it appears in the neighbouring ESCIN-3-3 and ESCIN-1
975 | lines, ~~where it is located between~~ (-6 and 8 s TWT), suggesting that it has been redefined.
976 | Nevertheless, clear reflections root in its upper part indicating that it still acted as a
977 | discontinuity/detachment. The depth of this feature in ~~the NW corner of~~ Iberia is similar to
978 | that of the high amplitude lower crustal internal reflection near the ICS. Accordingly, we
979 | suggest that in NW Iberia, gravitational collapse followed by crustal melting and extension has
980 | thinned the crust (Fig. 10), and specially the lower crust, relocating the mid-crustal
981 | discontinuity.

982 NW Iberia was importantly thickened (up to 50-70 km) due to the emplacement of the GTMZ
983 | allochthonous complexes. Thermal models by Alcock et al. (2009, 2015) show that as a result,
984 | the upper mantle continued increasing its temperature 60-65 Ma after the start of
985 | compressional deformation at 360 Ma. This implies large thinning of the mantle lithosphere,
986 | from 70 to 25-30 km, due to the ascent of the 1300 °C isotherm. It is not surprising that the
987 | lower crust there became the most highly extended as a consequence of the heat increase, as
988 | in the models it reached 800 °C after 45 Ma and 900 °C after 55 Ma (315-305 Ma).

989 The idea of a mid-crustal velocity discontinuity was put forward in the 1920's (Conrad, 1925).
990 | Early analysis of natural source earthquake recordings and later images from controlled source
991 | seismic reflection data provided further evidences that supported a clear distinction between
992 | upper and lower crust. These evidences led to considering the Conrad discontinuity, a global
993 | scale feature present in the continental crust. However, this was later challenged as some
994 | results of deep seismic reflection profiling did not show a clear distinction between upper and
995 | lower crust (Litak and Brown, 1989).

996 | Mid-crustal discontinuities have, however, been observed ~~very often and~~ in different types of
997 | seismic data worldwide (e.g., Fianco et al., 2019; Hobbs et al., 2004; Melekhova et al., 2019;
998 | Oncken, 1998; Ross et al., 2004; Snelson et al., 2013). Important changes in the rheology of the
999 | crust have also been reported at those depths (Maggini and Caputo, 2020; Wever, 1989)
1000 | supporting the idea that a mechanical boundary must exist. Thus, we suggest that, even
1001 | though it is not observed everywhere (Litak and Brown, 1989), this feature is an orogen-scale,
1002 | world class ~~crustal~~ continental crustal discontinuity (Artemieva, 2009), often coinciding with
1003 | the top of the highly laminated lower crust (when there is one). Its existence might determine
1004 | the way the crust deforms, easing decoupled deformation. Orogenic evolution, i.e. rifting,

1005 extension, melting, etc. may modify it or even erase it, thus its existence and geometry might
1006 help us to understand the geologic history of continents. In this regard, and coming back to the
1007 long-forgotten discussion of the nature of the Conrad discontinuity (Conrad, 1925) and its
1008 position on top of the laminated lower crust (Wever, 1989), we suggest that, in the Iberian
1009 Massif, the observed mid-crustal feature fulfills the characteristics of this debated
1010 discontinuity. Its clear signature and regional extension contributes to unravel its nature and
1011 significance.

1012 **5. Conclusions**

1013 Normal incidence seismic data acquired across the Iberian Massif in the last 30 years have
1014 provided an entire section of a well exposed and almost complete part of the European
1015 Variscides. Existing gaps in the central part have been recently sampled by passive source
1016 seismic recordings (noise and earthquakes) that provide fairly good constraints on the crustal
1017 structure.

1018 Results show that crustal thickness varies from ~9 s TWT in late-Variscan extended areas (NW
1019 of the Central Iberian Zone), to ~10 s TWT (30-33 km) in the external South Portuguese Zone to
1020 ~12 s TWT (36-38 km) in the internal West Asturian-Leonese Zone. Alpine reactivation has
1021 managed to further thicken the crust to at least ~14 s TWT (42-45 km) in the external
1022 Cantabrian Zone and to 35-38 km in the Iberian Central System, a Tertiary orogenic belt
1023 developed in Central Spain. The top of an **often** thick (up to 6 s TWT) and very reflective lower
1024 crust helps to define a mid-crustal discontinuity across the entire Iberian Massif. This boundary
1025 represents a level where reflections from the upper and lower crust merge asymptotically,
1026 thus suggesting that it has often acted as a detachment or a decoupling level. Its position and
1027 geometry varies mostly in relation to the late Variscan evolution. Accordingly, it is deeper in
1028 NW and central Iberia (~8 s TWT), where Variscan crustal thickening was important and
1029 gravitational collapse melted and extended the crust, thus defining a very thin lower crust.
1030 However, it appears between 4-6 s TWT to the SW, where the crust did not thicken as much
1031 and its original structure is better preserved, being later re-equilibrated through slow isostasy
1032 and erosion.

1033 This discontinuity exists in all the Iberian Massif tectonic zones, regardless of their Gondwana
1034 or Avalonia affinity, thus suggesting it is an orogenic-scale discontinuity. We interpret it as the
1035 rheological boundary between an overly ductilely deformed old lower crust and a
1036 heterogeneous variably (often also ductilely) deformed upper crust that mostly (but not only)
1037 shows evidences of the latest orogenic event. Its geometry, position and extent match the
1038 characteristics defined for the long-forgotten Conrad discontinuity. The identification of similar
1039 features in normal incidence profiles worldwide supports its inclusion as a major crustal
1040 discontinuity.

1041 **Acknowledgements**

1042 The seismic data was reprocessed using the commercial seismic signal processing software
1043 Claritas. Funding for this research was provided by the Junta de Castilla y León (SA065P17), the
1044 Spanish Ministry of Science and Innovation (CGL2016-78560-P) and the Generalitat de

1045 Catalunya (grant 2017SGR1022). [The comments of Prof. R.W.H. Butler and an anonymous](#)
 1046 [reviewer have greatly contributed to improve this manuscript.](#)

1047 References

- 1048 Alcock, J. E., Martínez Catalán, J. R., Arenas, R. and Díez Montes, A.: Use of thermal modeling
 1049 to assess the tectono-metamorphic history of the Lugo and Sanabria gneiss domes, Northwest
 1050 Iberia, *Bull. la Soc. Geol. Fr.*, 180(3), 179–197, doi:10.2113/gssgfbull.180.3.179, 2009.
- 1051 Alcock, J. E., Martínez Catalán, J. R. and Arenas, R.: One- and two-dimensional models are
 1052 equally effective in monitoring the crust's thermal response to advection by large-scale
 1053 thrusting during orogenesis, *Comput. Geosci.*, 37(8), 1205–1207,
 1054 doi:10.1016/j.cageo.2011.02.012, 2011.
- 1055 Alcock, J. E., Martínez Catalán, J. R., Rubio Pascual, F. J., Díez Montes, A., Díez Fernández, R.,
 1056 Gómez Barreiro, J., Arenas, R., Dias da Silva, Í. and González Clavijo, E.: 2-D thermal modeling
 1057 of HT-LP metamorphism in NW and Central Iberia: Implications for Variscan magmatism,
 1058 rheology of the lithosphere and orogenic evolution, *Tectonophysics*, 657, 21–37,
 1059 doi:10.1016/j.tecto.2015.05.022, 2015.
- 1060 Alonso, J. L.: Sequences of thrusts and displacement transfer in the superposed duplexes of the
 1061 Esla Nappe Region (cantabrian zone, nw spain), *J. Struct. Geol.*, 9(8), 969–983,
 1062 doi:10.1016/0191-8141(87)90005-8, 1987.
- 1063 Alonso, J. L., Marcos, A. and Suárez, A.: Paleogeographic inversion resulting from large out of
 1064 sequence breaching thrusts: The León Fault (Cantabrian zone, NW Iberia). A new picture of the
 1065 external Variscan thrust belt in the Ibero-Armorican arc, *Geol. Acta*, 7(4), 451–473,
 1066 doi:10.1344/105.000001449, 2009.
- 1067 Álvarez-Marrón, J., Pérez-Estaún, A., Dañobeitia, J. J., Pulgar, J. A., Martínez Catalán, J. R.,
 1068 Marcos, A., Bastida, F., Ayarza, P., Aller, J., Gallart, A., González-Lodeiro, F., Banda, E., Comas,
 1069 M. C. and Córdoba, D.: Seismic structure of the northern continental margin of Spain from
 1070 ESCIN deep seismic profiles, *Tectonophysics*, 264, 153–174, doi:10.1016/S0040-
 1071 1951(96)00124-2, 1996.
- 1072 Andrés, J., Draganov, D., Schimmel, M., Ayarza, P., Palomeras, I., Ruiz, M. and Carbonell, R.:
 1073 Lithospheric image of the Central Iberian Zone (Iberian Massif) using global-phase seismic
 1074 interferometry, *Solid Earth*, 10(6), 1937–1950, doi:10.5194/se-10-1937-2019, 2019.
- 1075 Andrés, J., Ayarza, P., Schimmel, M., Palomeras, I., Ruiz, M. and Carbonell, R.: What can seismic
 1076 noise tell us about the Alpine reactivation of the Iberian Massif? An example in the Iberian
 1077 Central System, *Solid Earth*, 11(6), 2499–2513, doi:10.5194/se-11-2499-2020, 2020.
- 1078 Arenas, R., Farias, P., Gallastegui, G., Gil Iburguchi, I., González Lodeiro, F., Klein, E., Marquínez,
 1079 J., Martín Parra, L. M., Martínez Catalán, J. R., Ortega, E., de Pablo Maciá, J. G., Peinado, M. and
 1080 Rodríguez Fernández, L. R.: Características geológicas y significado de los dominios que
 1081 componen la Zona de Galicia-Trás-os- Montes. Simposio sobre Cinturones Orogénicos, in
 1082 Simposio sobre Cinturones Orogénicos. II congreso Geológico de España, pp. 75–84, SGE., 1988.
- 1083 Arenas, R., Martínez Catalán, J. R., Martínez Sánchez, S., Díaz García, F., Abati, J., Fernández-
 1084 Suárez, J., Andonaegui, P. and Gómez-Barreiro, J.: Paleozoic ophiolites in the Variscan suture of
 1085 Galicia (northwest Spain): Distribution, characteristics, and meaning, *Mem. Geol. Soc. Am.*,
 1086 200(22), 425–444, doi:10.1130/2007.1200(22), 2007.

- 1087 Artemieva, I.: Continental Crust, Geophysics and Geochemistry Vol.II -Encyclopedia of Earth
1088 and Atmospheric Sciences in the Global Encyclopedia of Life Support Systems, UNESCO., 2009.
- 1089 Ayarza, P., Martínez Catalán, J. R., Gallart, J., Pulgar, J. A. and Dañobeitia, J. J.: Estudio Sismico
1090 de la Corteza Ibérica Norte 3 . 3 : A seismic image of the Variscan crust in the hinterland of the
1091 NW Iberian Massif, *Tectonics*, 17(2), 171–186, 1998.
- 1092 Ayarza, P., Martínez Catalán, J. R., Alvarez-Marrón, J., Zeyen, H. and Juhlin, C.: Geophysical
1093 constraints on the deep structure of a limited ocean-continent subduction zone at the North
1094 Iberian Margin, *Tectonics*, 23(1), doi:10.1029/2002TC001487, 2004.
- 1095 Azor, A., Lodeiro, F. G. and Simancas, J. F.: Tectonic evolution of the boundary between the
1096 Central Iberian and Ossa-Morena zones (Variscan belt, southwest Spain), *Tectonics*, 13(1), 45–
1097 61, doi:10.1029/93TC02724, 1994.
- 1098 Azor, A., Rubatto, D., Simancas, J. F., González Lodeiro, F., Martínez Poyatos, D., Martín Parra,
1099 L. M. and Matas, J.: Rhenic Ocean ophiolitic remnants in southern Iberia questioned by SHRIMP
1100 U-Pb zircon ages on the Beja-Acebuches amphibolites, *Tectonics*, 27(5), 1–11,
1101 doi:10.1029/2008TC002306, 2008.
- 1102 Bandrés, A., Eguíluz, L., Pin, C., Paquette, J. L., Ordóñez, B., Le Fèvre, B., Ortega, L. A. and Gil
1103 Ibarra, J. I.: The northern Ossa-Morena Cadomian batholith (Iberian Massif): Magmatic arc
1104 origin and early evolution, *Int. J. Earth Sci.*, 93(5), 860–885, doi:10.1007/s00531-004-0423-6,
1105 2004.
- 1106 Barbero, L.: Granulite-facies metamorphism in the Anatectic Complex of Toledo, Spain: late
1107 Hercynian tectonic evolution by crustal extension, *J. Geol. Soc. London*, 152(2), 365–382,
1108 doi:10.1144/gsjgs.152.2.0365, 1995.
- 1109 BIRPS, B. I. R. P. S. and ECORS, E. de la C. C. et O. par R. et R. S.: Deep seismic reflection
1110 profiling between England, France and Ireland., *J. - Geol. Soc.*, 143(1), 45–52,
1111 doi:10.1144/gsjgs.143.1.0045, 1986.
- 1112 Bortfeld, R. K.: First results and preliminary interpretation of deep- reflection seismic
1113 recordings along profile DEKORP 2-South (FRG.), *J. Geophys. - Zeitschrift fur Geophys.*, 57(3),
1114 137–163, 1985.
- 1115 Braid, J. A., Brendan Murphy, J., Quesada, C. and Mortensen, J.: Tectonic escape of a crustal
1116 fragment during the closure of the Rhenic Ocean: U-Pb detrital zircon data from the late
1117 Palaeozoic Pulo do Lobo and South Portuguese zones, Southern Iberia, *J. Geol. Soc. London.*,
1118 168(2), 383–392, doi:10.1144/0016-76492010-104, 2011.
- 1119 Braid, J. A., Murphy, J. B., Quesada, C., Bickerton, L. and Mortensen, J. K.: Probing the
1120 composition of unexposed basement, South Portuguese Zone, southern Iberia: Implications for
1121 the connections between the Appalachian and Variscan orogens, *Can. J. Earth Sci.*, 49(4), 591–
1122 613, doi:10.1139/E11-071, 2012.
- 1123 Butler, R. W. H. and Mazzoli, S.: Styles of continental contraction: A review and introduction,
1124 *Spec. Pap. Geol. Soc. Am.*, 414(414), 1–10, doi:10.1130/2006.2414(01), 2006.
- 1125 Capdevila, R., Boillot, G., Lepvrier, C., Malod, J. A. and Mascle, G.: Les formations cristallines du
1126 Banc Le Danois (marge nord-ibérique), *CR Acad. Sci. Paris, D*, 291(January), 317–320 [online]
1127 Available from:
1128 [http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Les+formations+cristallines+du+Banc+Le+Danois+\(marge+nord+ibérique\)#0](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Les+formations+cristallines+du+Banc+Le+Danois+(marge+nord+ibérique)#0), 1980.
1129

- 1130 Carbonell, R. and Smithson, S. B.: Large-scale anisotropy within the crust in the Basin and
1131 Range province, *Geology*, 19(July), 698–701, doi:doi:10.1130/0091-7613(1991)019, 1991.
- 1132 Carbonell, R., Simancas, J. F., Juhlin, C., Pous, J., Pérez-Estaún, A., Gonzalez-Lodeiro, F., Munoz,
1133 G., Heise, W. and Ayarza, P.: Geophysical evidence of a mantle derived intrusion in SW Iberia,
1134 *Geophys. Res. Lett.*, 31(11), 1–4, 2004.
- 1135 Carracedo, M., Paquette, J. L., Alonso Olazabal, A., Santos Zalduegui, J. F., de García de
1136 Madinabeitia, S., Tiepolo, M. and Gil Ibarguchi, J. I.: U-Pb dating of granodiorite and granite
1137 units of the Los Pedroches batholith. Implications for geodynamic models of the southern
1138 Central Iberian Zone (Iberian Massif), *Int. J. Earth Sci.*, 98(7), 1609–1624, doi:10.1007/s00531-
1139 008-0317-0, 2009.
- 1140 Carvalho, D.: The metallogenetic consequences of plate tectonics and the upper Paleozoic
1141 evolution of southern Portugal, *Estud. Notas e Trab. S.F.M.*, 20(3–4), 297–320, 1972.
- 1142 Casquet, C., Galindo, C., Tornos, F., Velasco, F. and Canales, A.: The Aguablanca Cu-Ni ore
1143 deposit (Extremadura, Spain), a case of synorogenic orthomagmatic mineralization: Age and
1144 isotope composition of magmas (Sr, Nd) and ore (S), *Ore Geol. Rev.*, 18(3–4), 237–250,
1145 doi:10.1016/S0169-1368(01)00033-6, 2001.
- 1146 Chopra, S. and Alexeev, V.: Application of texture attribute analysis to 3D seismic data, *Soc.*
1147 *Explor. Geophys. - 75th SEG Int. Expo. Annu. Meet. SEG 2005*, 30(7), 767–770,
1148 doi:10.1190/1.2144439, 2005.
- 1149 Conrad, V.: Laufzeitkurven des Tauernbens vom 28, *Mitt. Erdb. Komm. Wlen Akad. Wiss.*, 59, 1,
1150 1925.
- 1151 Cook, F. A.: Fine structure of the continental reflection Moho, *Bull. Geol. Soc. Am.*, 114(1), 64–
1152 79, doi:10.1130/0016-7606(2002)114<0064:FSOTCR>2.0.CO;2, 2002.
- 1153 Corretgé, L. G. and Suárez, O.: Cantabrian and Palentian Zones. Igneous Rocks, in *Pre-Mesozoic*
1154 *Geology of Iberia*, edited by R. D. Dallmeyer and E. Martínez García, pp. 72–79, Springer-
1155 Verlag, Berlin., 1990.
- 1156 Dallmeyer, R. D. and Quesada, C.: Cadomian vs. Variscan evolution of the Ossa-Morena zone
1157 (SW Iberia): field and ⁴⁰Ar/³⁹Ar mineral age constraints, *Tectonophysics*, 216(3–4), 339–364,
1158 doi:10.1016/0040-1951(92)90405-U, 1992.
- 1159 Dallmeyer, R. D., Martínez Catalán, J. R., Arenas, R., Gil Ibarguchi, J. I., Gutiérrez-Alonso, G.,
1160 Farias, P., Bastida, F. and Aller, J.: Diachronous Variscan tectonothermal activity in the NW
1161 Iberian Massif: Evidence from ⁴⁰Ar/³⁹Ar dating of regional fabrics, *Tectonophysics*, 277(4),
1162 307–337, doi:10.1016/S0040-1951(97)00035-8, 1997.
- 1163 DeFelipe, I., Pedreira, D., Pulgar, J. A., van der Beek, P. A., Bernet, M. and Pik, R.: Unraveling
1164 the Mesozoic and Cenozoic Tectonothermal Evolution of the Eastern Basque-Cantabrian Zone–
1165 Western Pyrenees by Low-Temperature Thermochronology, *Tectonics*, 38(9), 3436–3461,
1166 doi:10.1029/2019TC005532, 2019.
- 1167 DeFelipe, I., Alcalde, J., Ivandic, M., Martí, D., Ruiz, M., Marzán, I., Diaz, J., Ayarza, P.,
1168 Palomeras, I., Fernandez-Turiel, J. L., Molina, C., Bernal, I., Brown, L., Roberts, R. and Carbonell,
1169 R.: Reassessing the lithosphere: SeisDARE, an open access seismic data repository, *Earth Syst.*
1170 *Sci. Data Discuss.*, (September), 1–32, doi:10.5194/essd-2020-208, 2020.
- 1171 DEKORP Research Group: Results of deep reflection seismic profiling in the Oberpfalz (Bavaria),

- 1172 Geophys. J. R. Astron. Soc., 89(1), 353–360, doi:10.1111/j.1365-246X.1987.tb04430.x, 1987.
- 1173 Dias, R. and Ribeiro, A.: The Ibero-Armorican Arc: A collision effect against an irregular
1174 continent?, *Tectonophysics*, 246(1–3), 113–128, doi:10.1016/0040-1951(94)00253-6, 1995.
- 1175 Díaz Azpiroz, M., Fernández, C., Castro, A. and El-Biad, M.: Tectonometamorphic evolution of
1176 the Aracena metamorphic belt (SW Spain) resulting from ridge-trench interaction during
1177 Variscan plate convergence, *Tectonics*, 25(1), 1–20, doi:10.1029/2004TC001742, 2006.
- 1178 Díez-Montes, A., Martínez Catalán, J. R. and Bellido Mulas, F.: Role of the Ollo de Sapo massive
1179 felsic volcanism of NW Iberia in the Early Ordovician dynamics of northern Gondwana,
1180 *Gondwana Res.*, 17(2–3), 363–376, doi:10.1016/j.gr.2009.09.001, 2010.
- 1181 Díez Balda, M. A., Martínez Catalán, J. R. and Ayarza, P.: Syn-collisional extensional collapse
1182 parallel to the orogenic trend in a domain of steep tectonics: the Salamanca Detachment Zone
1183 (Central Iberian Zone, Spain), *J. Struct. Geol.*, 17(2), 163–182, doi:10.1016/0191-
1184 8141(94)E0042-W, 1995.
- 1185 Díez Montes, A., Martínez Catalán, J. R. and Bellido Mulas, F.: Role of the Ollo de Sapo massive
1186 felsic volcanism of NW Iberia in the Early Ordovician dynamics of northern Gondwana,
1187 *Gondwana Res.*, 17(2–3), 363–376, doi:10.1016/j.gr.2009.09.001, 2010.
- 1188 Ehsan, S. A., Carbonell, R., Ayarza, P., Martí, D., Pérez-Estaún, A., Martínez-Poyatos, D. J.,
1189 Simancas, J. F., Azor, A. and Mansilla, L.: Crustal deformation styles along the reprocessed deep
1190 seismic reflection transect of the Central Iberian Zone (Iberian Peninsula), *Tectonophysics*, 621,
1191 159–174, doi:10.1016/j.tecto.2014.02.014, 2014.
- 1192 Ehsan, S. A., Carbonell, R., Ayarza, P., Martí, D., Martínez Poyatos, D., Simancas, J. F., Azor, A.,
1193 Ayala, C., Torné, M. and Pérez-Estaún, A.: Lithospheric velocity model across the Southern
1194 Central Iberian Zone (Variscan Iberian Massif): The ALCUDIA wide-angle seismic reflection
1195 transect, *Tectonics*, 34(3), 535–554, doi:10.1002/2014TC003661, 2015.
- 1196 Expósito, I., Simancas, J. F. and Lodeiro, F. G.: Estructura de la mitad septentrional de la zona
1197 de Ossa-Morena: Deformación en el bloque inferior de un cabalgamiento cortical de evolución
1198 compleja: Deformación en el bloque inferior de un cabalgamiento cortical de evolución
1199 compleja, *Rev. la Soc. Geológica España*, 15(1), 3–14, 2002.
- 1200 Expósito, I., Simancas, J. F., González Lodeiro, F., Bea, F., Montero, P. and Salman, K.:
1201 Metamorphic and deformational imprint of Cambrian - Lower Ordovician rifting in the Ossa-
1202 Morena Zone (Iberian Massif, Spain), *J. Struct. Geol.*, 25(12), 2077–2087, doi:10.1016/S0191-
1203 8141(03)00075-0, 2003.
- 1204 Farias, P., Gallastegui, G., González Lodeiro, F., Marquinez, J., Martín-Parra, L. M., Martínez
1205 Catalán, J. R. and Pablo-Maciá, J. G.: “Aportaciones al conocimiento de la litoestratigrafía y
1206 estructura de Galicia Central,” *Memórias da Fac. Ciências, Univ. do Porto*, 1(January 1987),
1207 411–431, 1987.
- 1208 Fernández-Suárez, J., Gutiérrez-Alonso, G., Jenner, G. A. and Jackson, S. E.: Geochronology and
1209 geochemistry of the Pola de Allande granitoids (northern Spain): their bearing on the
1210 Cadomian-Avalonian evolution of northwest Iberia, *Can. J. Earth Sci.*, 35(12), 1439–1453,
1211 doi:10.1139/cjes-35-12-1439, 1998.
- 1212 Fernández-Viejo, G., Gallart, J., Pulgar, J. A., Gallastegui, J., Dañobeitia, D. and Córdoba, D.:
1213 Crustal transition between continental and oceanic domains along the North Iberian margin
1214 from wide angle seismic and gravity data, *Geophys. Res. Lett.*, 25(23), 4249–4252, 1998.

- 1215 Fernández-Viejo, G., Gallart, J., Pulgar, A., Córdoba, D. and Dañoibeitia, J. J.: Seismic signature
1216 of Variscan and Alpine tectonics in NW Iberia: Crustal structure of the Cantabrian Mountains
1217 and Duero Basin, *J. Geophys. Res.*, 105(1999), 3001–3018, 2000.
- 1218 Fianco, C. B., França, G. S., Albuquerque, D. F., Vilar, C. da S. and Argollo, R. M.: Using the
1219 receiver function for studying earth deep structure in the Southern Borborema Province, *J.*
1220 *South Am. Earth Sci.*, 94(April), 102221, doi:10.1016/j.jsames.2019.102221, 2019.
- 1221 Finlayson, D. M., Collins, C. D. N. and Lock, J.: P-wave velocity features of the lithosphere under
1222 the Eromanga Basin, Eastern Australia, including a prominent MID-crustal (Conrad?)
1223 discontinuity, *Tectonophysics*, 101(3–4), 267–291, doi:10.1016/0040-1951(84)90117-3, 1984.
- 1224 Flecha, I., Palomeras, I., Carbonell, R., Simancas, J. F., Ayarza, P., Matas, J., González-Lodeiro, F.
1225 and Pérez-Estaún, A.: Seismic imaging and modelling of the lithosphere of SW-Iberia,
1226 *Tectonophysics*, 472(1–4), 148–157, doi:10.1016/j.tecto.2008.05.033, 2009.
- 1227 Franke, W.: The mid-European segment of the Variscides: Tectonostratigraphic units, terrane
1228 boundaries and plate tectonic evolution, *Geol. Soc. Spec. Publ.*, 179, 35–56,
1229 doi:10.1144/GSL.SP.2000.179.01.05, 2000.
- 1230 Franke, W., Bortfeld, R. K., Brix, M., Drozdowski, G., Dürbaum, H. J., Giese, P., Janoth, W.,
1231 Jödicke, H., Reichert, C., Scherp, A., Schmoll, J., Thomas, R., Thünker, M., Weber, K., Wiesner,
1232 M. G. and Wong, H. K.: Crustal structure of the Rhenish Massif: results of deep seismic
1233 reflection lines Dekorp 2-North and 2-North-Q, *Geol. Rundschau*, 79(3), 523–566,
1234 doi:10.1007/BF01879201, 1990.
- 1235 Gallastegui, J., Pulgar, J. A. and Alvarez-Marrón, J.: 2-D seismic modeling of the Variscan
1236 foreland thrust and fold belt crust in NW Spain from ESCIN-1 deep seismic reflection data,
1237 *Tectonophysics*, 269(1–2), 21–32, doi:10.1016/S0040-1951(96)00166-7, 1997.
- 1238 Gallastegui, J., Pulgar, J. A. and Gallart, J.: Alpine tectonic wedging and crustal delamination in
1239 the Cantabrian Mountains (NW Spain), *Solid Earth*, 7(4), 1043–1057, doi:10.5194/se-7-1043-
1240 2016, 2016.
- 1241 García Casquero, J. L., Boelrijk, N. A. I. M., Chacón, J. and Priem, H. N. A.: Rb-Sr evidence for the
1242 presence of Ordovician gsrinites in the deformed basement of the Badajoz-Córdoba belt, SW
1243 Spain, *Geol. Rundschau*, 74(2), 379–384, doi:10.1007/BF01824904, 1985.
- 1244 Gardien, V., Arnaud, N. and Desmurs, L.: Petrology and ar-ar dating of granulites from the
1245 galicia bank (spain): African craton relics in western europe, *Geodin. Acta*, 13(2–3), 103–117,
1246 doi:10.1080/09853111.2000.11105367, 2000.
- 1247 Gómez-Pugnaire, M. T., Azor, A., Fernández-Soler, J. M. and López Sánchez-Vizcaíno, V.: The
1248 amphibolites from the Ossa-Morena/Central Iberian Variscan suture (Southwestern Iberian
1249 Massif): Geochemistry and tectonic interpretation, *Lithos*, 68(1–2), 23–42, doi:10.1016/S0024-
1250 4937(03)00018-5, 2003.
- 1251 Gómez Barreiro, J., Martínez Catalán, J. R., Arenas, R., Castiñeiras, P., Abati, J., Díaz García, F.
1252 and Wijbrans, J. R.: Tectonic evolution of the upper allochthon of the Órdenes complex (
1253 northwestern Iberian Massif): Structural constraints to a polyorogenic peri-Gondwanan
1254 terrane, *Geol. Soc. Am. Spec. Pap.*, 423, 315–332, doi:10.1130/2007.2423(15)., 2007.
- 1255 Hernández Enrile, J. L.: Extensional tectonics of the toledo ductile-brittle shear zone, central
1256 Iberian Massif, *Tectonophysics*, 191(3–4), 311–324, doi:10.1016/0040-1951(91)90064-Y, 1991.

- 1257 Hobbs, B. E., Ord, A., Regenauer-Lieb, K. and Drummond, B.: Fluid reservoirs in the crust and
1258 mechanical coupling between the upper and lower crust, *Earth, Planets Sp.*, 56(12), 1151–
1259 1161, doi:10.1186/BF03353334, 2004.
- 1260 Julivert, M., Fontboté, M., Ribeiro, A. and Conde, L. E.: Mapa tectónico de la Península Ibérica
1261 y Baleares Notas Incluye mapa : Unidades estructurales de la Península Ibérica . Escala 1 :
1262 1.000.000., Instituto Geológico y Minero de España., 1972.
- 1263 Klemperer, S. L., Hauge, T. A., Hauser, E. C., Oliver, J. E. and Potter, C. J.: The Moho in the
1264 northern Basin and Range Province, Nevada, along the COCORP 40oN seismic- reflection
1265 transect. (USA), *Geol. Soc. Am. Bull.*, 97(5), 603–618, doi:10.1130/0016-
1266 7606(1986)97<603:TMITNB>2.0.CO;2, 1986.
- 1267 Kossmat, F.: Gliederung des varistischen Gebirgsbaues. Abhandlungen des Sächsischen, in
1268 Abhandlungen des Sächsischen Geologuschen Landesamts, edited by A. des S. G. Landesamts,
1269 p. 39, Leipzig, Heft 1., 1927.
- 1270 Levander, A. R. and Holliger, K.: Small-scale heterogeneity and large-scale velocity structure of
1271 the continental crust, *J. Geophys. Res.*, 97(B6), 8797–8804, doi:10.1029/92JB00659, 1992.
- 1272 Linnemann, U., Pereira, M. F., Jeffries, T. E., Drost, K. and Gerdes, A.: The Cadomian Orogeny
1273 and the opening of the Rheic Ocean: The diacrony of geotectonic processes constrained by LA-
1274 ICP-MS U-Pb zircon dating (Ossa-Morena and Saxo-Thuringian Zones, Iberian and Bohemian
1275 Massifs), *Tectonophysics*, 461(1–4), 21–43, doi:10.1016/j.tecto.2008.05.002, 2008.
- 1276 Litak, R. K. and Brown, L. D.: A modern perspective on the Conrad Discontinuity, *Eos, Trans.*
1277 *Am. Geophys. Union*, 70(29), 713–725, doi:10.1029/89EO00223, 1989.
- 1278 López Sánchez-Vizcaíno, V., Gómez-Pugnaire, M. T., Azor, A. and Fernández-Soler, J. M.: Phase
1279 diagram sections applied to amphibolites: A case study from the Ossa-Morena/Central Iberian
1280 Variscan suture (Southwestern Iberian Massif), *Lithos*, 68(1–2), 1–21, doi:10.1016/S0024-
1281 4937(03)00017-3, 2003.
- 1282 Lotze, F.: Stratigraphie und Tektonik des Keltiberischen Grundgebirges (Spanien),
1283 Abhandlungen Gesellschaft der Wissenschaften zu Göttingen, Math. (Abh. Ges. Wiss.
1284 Göttingen, Math.-Phys.), 14(2), 143–162, 1929.
- 1285 Maggini, M. and Caputo, R.: Sensitivity analysis for crustal rheological profiles: Examples from
1286 the Aegean region, *Ann. Geophys.*, 63(3), 1–29, doi:10.4401/ag-8244, 2020.
- 1287 Martínez Catalán, J. R.: Are the oroclines of the Variscan belt related to late Variscan strike-slip
1288 tectonics?, *Terra Nov.*, 23(4), 241–247, doi:10.1111/j.1365-3121.2011.01005.x, 2011.
- 1289 Martínez Catalán, J. R.: The Central Iberian arc, an orocline centered in the Iberian Massif and
1290 some implications for the Variscan belt, *Int. J. Earth Sci.*, 101(5), 1299–1314,
1291 doi:10.1007/s00531-011-0715-6, 2012.
- 1292 Martínez Catalán, J. R., Ayarza, P., Pulgar, J. A., Pérez-Estaún, A., Gallart, J., Marcos, A., Bastida,
1293 F., Álvarez-Marrón, J., González Lodeiro, F., Aller, J., Dañobeitia, J. J., Banda, E., Córdoba, D.
1294 and Comas, M. C.: Results from the ESCIN-N3.3 marine deep seismic profile along the
1295 Cantabrian continental margin, *Rev. Soc. Geol. España*, 8(4), 341-354., 1995.
- 1296 Martínez Catalán, J. R., Arenas, R., Díaz García, F. and Abati, J.: Variscan accretionary complex
1297 of northwest Iberia: Terrane correlation succession of tectonothermal events, *Geology*, 25(12),
1298 1103–1106, doi:10.1130/0091-7613(1997)025<1103:VACONI>2.3.CO;2, 1997.

- 1299 Martínez Catalán, J. R., Fernández-Suárez, J., Jenner, G. A., Belousova, E. and Díez Montes, A.:
 1300 Provenance constraints from detrital zircon U–Pb ages in the NW Iberian Massif: implications
 1301 for Palaeozoic plate configuration and Variscan evolution, *J. Geol. Soc. London.*, 161(3), 463–
 1302 476, doi:10.1144/0016-764903-054, 2004.
- 1303 Martínez Catalán, J. R., Arenas, R., Diágarcía, F., González Cuadra, P., Gómez-Barreiro, J., Abati,
 1304 J., Castiñeiras, P., Fernández-Suárez, J., Sánchez Martínez, S., Andonaegui, P., González Clavijo,
 1305 E., Díez Montes, A., Rubio Pascual, F. J. and Valle Aguado, B.: Space and time in the tectonic
 1306 evolution of the northwestern Iberian Massif: Implications for the Variscan belt, *Mem. Geol.*
 1307 *Soc. Am.*, 200(21), 403–423, doi:10.1130/2007.1200(21), 2007.
- 1308 Martínez Catalán, J. R., Álvarez Lobato, F., Pinto, V., Gómez Barreiro, J., Ayarza, P., Villalaín, J. J.
 1309 and Casas, A.: Gravity and magnetic anomalies in the allochthonous rdenes Complex (Variscan
 1310 belt, northwest Spain): Assessing its internal structure and thickness, *Tectonics*, 31(5), 1–18,
 1311 doi:10.1029/2011TC003093, 2012.
- 1312 Martínez Catalán, J. R., Rubio Pascual, F. J., Díez Montes, A., Díez Fernández, R., Gómez
 1313 Barreiro, J., Dias da Silva, Í., González Clavijo, E., Ayarza, P. and Alcock, J. E.: The late Variscan
 1314 HT/LP metamorphic event in NW and Central Iberia: relationships to crustal thickening,
 1315 extension, orocline development and crustal evolution, *Geol. Soc. London, Spec. Publ.*, 405(1),
 1316 225–247, doi:10.1144/SP405.1, 2014.
- 1317 Martínez Catalán, J. R., Díez Balda, M. A., Escuder Viruete, J., Villar Alonso, P., Ayarza, P.,
 1318 Gonzalez Clavijo, E. and Díez Montes, A.: Cizallamientos dúctiles de escala regional en la
 1319 provincia de Salamanca, in *Geo-Guías: Rutas Geológicas por la Península Ibérica, Canarias,*
 1320 *Sicilia y Marruecos*, edited by M. Diaz Azpiroz, I. Exposito Ramos, S. Llana Fúnez, and B. Bauluz
 1321 Lázaro, pp. 109–118, *Sociedad Geológica de España.*, 2019.
- 1322 Martínez García, A.: Seismic characterization and geodynamic significane of the mid-crustal
 1323 discontinuity across the Variscan Iberia, *Universidad de Barcelona.*, 2019.
- 1324 Martínez García, E.: El Paleozoico de la Zona Cantabrica Oriental (Noroeste de España), *Trab.*
 1325 *Geol.*, 11, 95–127, 1981.
- 1326 Martínez Poyatos, D., Carbonell, R., Palomeras, I., Simancas, J. F., Ayarza, P., Martí, D., Azor, A.,
 1327 Jabaloy, A., González Cuadra, P., Tejero, R., Martín Parra, L. M., Matas, J., González Lodeiro, F.,
 1328 Pérez-Estaún, A., García Lobón, J. L. and Mansilla, L.: Imaging the crustal structure of the
 1329 Central Iberian Zone (Variscan Belt): The ALCUDIA deep seismic reflection transect, *Tectonics*,
 1330 31(3), 1–21, doi:10.1029/2011TC002995, 2012.
- 1331 Martínez Poyatos, D. J.: Estructura del borde meridional de la Zona Centroibérica y su relación
 1332 con el contacto entre las Zonas Centroibérica y de Ossa-Morena, in *Serie Nova Terra*, 18,
 1333 edited by L. X. de Laxe, p. 295, *Instituto Universitario de Xeoloxía, A Coruña, Spain.*, 2002.
- 1334 Matte, P.: The Variscan collage and orogeny (480–290 Ma) and the tectonic definition of the
 1335 Armorica microplate: a review - Matte - 2003 - *Terra Nova - Wiley Online Library*, *Terra Nov.*,
 1336 13(1997), 122–128 [online] Available from: <http://onlinelibrary.wiley.com/doi/10.1046/j.1365-3121.2001.00327.x/full%5Cnpapers2://publication/uuid/85445613-EAD8-49D7-AE4E-704E3D282C36>, 2001.
- 1339 Matte, P. and Ribeiro, A.: Forme et orientation de l'ellipsoïde de déformation dans la virgation
 1340 hercynienne de Galice. Relations avec le plissement et hypothèses sur la genèse de l'arc ibéro-
 1341 armoricain, *Comptes Rendus l'Académie des Sci.*, 280, 2825–2828, 1975.

- 1342 Meissner, R.: Rupture, creep, lamellae and crocodiles: happenings in the continental crust,
1343 *Terra Nov.*, 1(1), 17–28, doi:10.1111/j.1365-3121.1989.tb00321.x, 1989.
- 1344 Meissner, R., Rabbel, W. and Kern, H.: Seismic lamination and anisotropy of the lower
1345 continental crust, *Tectonophysics*, 416(1–2), 81–99, doi:10.1016/j.tecto.2005.11.013, 2006.
- 1346 Melekhova, E., Schlaphorst, D., Blundy, J., Kendall, J. M., Connolly, C., McCarthy, A. and
1347 Arculus, R.: Lateral variation in crustal structure along the Lesser Antilles arc from petrology of
1348 crustal xenoliths and seismic receiver functions, *Earth Planet. Sci. Lett.*, 516, 12–24,
1349 doi:10.1016/j.epsl.2019.03.030, 2019.
- 1350 Montero, P., Bea, F., González-Lodeiro, F., Talavera, C. and Whitehouse, M. J.: Zircon ages of
1351 the metavolcanic rocks and metagranites of the Ollo de Sapo Domain in central Spain:
1352 Implications for the Neoproterozoic to Early Palaeozoic evolution of Iberia, *Geol. Mag.*, 144(6),
1353 963–976, doi:10.1017/S0016756807003858, 2007.
- 1354 Ochsner, A.: U-Pb geochronology of the Upper Proterozoic-Lower Paleozoic geodynamic
1355 evolution in the Ossa-Morena Zone (SW Iberia): constraints on the timing of the Cadomian
1356 orogeny, University of Zurich, Diss. ETH10, 192., 1993.
- 1357 Okaya, D., Rabbel, W., Beilecke, T. and Hasenclever, J.: P wave material anisotropy of a
1358 tectono-metamorphic terrane: An active source seismic experiment at the KTB super-deep drill
1359 hole, southeast Germany, *Geophys. Res. Lett.*, 31(24), 1–4, doi:10.1029/2004GL020855, 2004.
- 1360 Oliveira, J. T.: Part VI: South Portuguese Zone, stratigraphy and synsedimentary tectonism, in
1361 *Pre-Mesozoic Geology of Iberia*, edited by E. Dallmeyer, R. D., and Martínez García, pp. 334–
1362 347, Springer, Berlin, Germany., 1990.
- 1363 Oncken, O.: Orogenic mass transfer and reflection seismic patterns - Evidence from DEKORP
1364 sections across the European variscides (central Germany), *Tectonophysics*, 286(1–4), 47–61,
1365 doi:10.1016/S0040-1951(97)00254-0, 1998.
- 1366 Palomeras, I., Carbonell, R., Flecha, I., Simancas, J. F., Ayarza, P., Matas, J., Poyatos, D. M.,
1367 Azor, A., Lodeiro, F. G. and Pérez-Estaún, A.: Nature of the lithosphere across the Variscan
1368 orogen of SW Iberia: Dense wide-angle seismic reflection data, *J. Geophys. Res. Solid Earth*,
1369 114(2), 1–29, 2009.
- 1370 Palomeras, I., Carbonell, R., Ayarza, P., Martí, D., Brown, D. and Simancas, J. F.: Shear wave
1371 modeling and Poisson's ratio in the Variscan Belt of SW Iberia, *Geochemistry, Geophys.*
1372 *Geosystems*, 12(7), 1–23, doi:10.1029/2011GC003577, 2011.
- 1373 Palomeras, I., Villaseñor, A., Thurner, S., Levander, A., Gallart, J. and Harnafi, M.: Lithospheric
1374 structure of Iberia and Morocco using finite-frequency Rayleigh wave tomography from
1375 earthquakes and seismic ambient noise, *Geochemistry, Geophys. Geosystems*, 1–17,
1376 doi:10.1002/2016GC006657, 2017.
- 1377 Pedreira, D., Pulgar, J. A., Gallart, J. and Díaz, J.: Seismic evidence of Alpine crustal thickening
1378 and wedging from the western Pyrenees to the Cantabrian Mountains (north Iberia), *J.*
1379 *Geophys. Res.*, 108(B4), 1–21, doi:10.1029/2001JB001667, 2003.
- 1380 Pereira, M. F., Chichorro, M., Williams, I. S., Silva, J. B., Fernández, C., Diaz-Azpiroz, M., Apraiz,
1381 A. and Castro, A.: Variscan intra-orogenic extensional tectonics in the Ossa – Morena-Évora –
1382 Arcena – Lora del Río metamorphic belt, SW Iberian Zone (E Massif): SHRIMP zircon U – Th –
1383 Pb geochronology, in *Ancient Orogens and Modern Analogues*, edited by J. B. Murphy, J. D.
1384 Keppie, and A. J. Hynes, pp. 327, 215–237, Geological Society of London, Special Publications.,

- 1385 2009.
- 1386 Pereira, M. F., Ribeiro, C., Vilallonga, F., Chichorro, M., Drost, K., Silva, J. B., Albardeiro, L.,
1387 Hofmann, M. and Linnemann, U.: Variability over time in the sources of South Portuguese Zone
1388 turbidites: Evidence of denudation of different crustal blocks during the assembly of Pangaea,
1389 *Int. J. Earth Sci.*, 103(5), 1453–1470, doi:10.1007/s00531-013-0902-8, 2014.
- 1390 Pérez-Cáceres, I., Martínez Poyatos, D., Simancas, J. F. and Azor, A.: The elusive nature of the
1391 Rhenic Ocean suture in SW Iberia, *Tectonics*, 34(12), 2429–2450, doi:10.1002/2015TC003947,
1392 2015.
- 1393 Pérez-Cáceres, I., Simancas, J. F., Martínez Poyatos, D., Azor, A. and González Lodeiro, F.:
1394 Oblique collision and deformation partitioning in the SW Iberian Variscides, *Solid Earth*, 7(3),
1395 857–872, doi:10.5194/se-7-857-2016, 2016.
- 1396 Pérez-Estaún, A., Bastida, F., Alonso, J. L., Marquínez, J., Aller, J., Alvarez-Marrón, J., Marcos, A.
1397 and Pulgar, J. A.: A thin-skinned tectonics model for an arcuate fold and thrust belt: The
1398 Cantabrian Zone (Variscan Ibero-Armorican Arc), *Tectonics*, 7(3), 517–537,
1399 doi:https://doi.org/10.1029/TC007i003p00517, 1988.
- 1400 Pérez-Estaún, A., Bastida, F., Martínez Catalán, J.R. Gutierrez Marco, J. C., Marcos, A. and
1401 Pulgar, J. A.: West Asturian-Leonese Zone. Stratigraphy, in *Pre-Mesozoic Geology of Iberia*,
1402 edited by E. Dallmeyer, R.D. and Martínez Garcia, pp. 92–102, Springer-Verlag, Berlin., 1990.
- 1403 Pérez-Estaún, A., Martínez Catalán, J. R. and Bastida, F.: Crustal thickening and deformation
1404 sequence in the footwall to the suture of the Variscan belt of northwest Spain, *Tectonophysics*,
1405 191(3–4), 243–253, doi:10.1016/0040-1951(91)90060-6, 1991.
- 1406 Pérez-Estaún, A., Pulgar, J. A., Banda, E. and Alvarez-Marrón, J.: Crustal structure of the
1407 external variscides in northwest Spain from deep seismic reflection profiling, *Tectonophysics*,
1408 232(1–4), doi:10.1016/0040-1951(94)90078-7, 1994.
- 1409 Pin, C., Fonseca, P. E., Paquette, J. L., Castro, P. and Matte, P.: The ca. 350 Ma Beja Igneous
1410 Complex: A record of transcurrent slab break-off in the Southern Iberia Variscan Belt?,
1411 *Tectonophysics*, 461(1–4), 356–377, doi:10.1016/j.tecto.2008.06.001, 2008.
- 1412 Pulgar, J. A., Gallart, J., Fernández-Viejo, G., Pérez-Estaún, A. and Álvarez-Marrón, J.: Seismic
1413 image of the Cantabrian Mountains in the western extension of the Pyrenees from integrated
1414 ESCIN reflection and refraction data, *Tectonophysics*, 264(1–4), 1–19, doi:10.1016/S0040-
1415 1951(96)00114-X, 1996.
- 1416 Quesada, C. and Dallmeyer, R. D.: Tectonothermal evolution of the Badajoz-Córdoba shear
1417 zone (SW Iberia): characteristics and $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age constraints, *Tectonophysics*,
1418 231(1–3), 195–213, doi:10.1016/0040-1951(94)90130-9, 1994.
- 1419 Quintana, L., Pulgar, J. A. and Alonso, J. L.: Displacement transfer from borders to interior of a
1420 plate: A crustal transect of Iberia, *Tectonophysics*, 663, 378–398,
1421 doi:10.1016/j.tecto.2015.08.046, 2015.
- 1422 Rat, P.: The Basque-Cantabrian basin between the Iberian and European plates: Some facts but
1423 still many problems, *Rev. la Soc. Geológica España*, 1(3–4), 327–348, 1988.
- 1424 Reche, J., Martínez, F. J., Arboleya, M. L., Dietsch, C. and Briggs, W. D.: Evolution of a kyanite-
1425 bearing belt within a HT-LP orogen: the case of NW Variscan Iberia, *J. Metamorph. Geol.*, 16(3),
1426 379–394, doi:10.1111/j.1525-1314.1998.00142.x, 1998.

- 1427 Ries, A. C. and Shackleton, R. M.: Catazonal Complexes of North-West Spain and North
1428 Portugal, Remnants of a Hercynian Thrust Plate, *Nat. Phys. Sci.*, 234(47), 65–68,
1429 doi:10.1038/physci234065a0, 1971.
- 1430 Robardet, M.: Alternative approach to the Variscan Belt in southwestern Europe: Preorogenic
1431 paleobiogeographical constraints, *Spec. Pap. Geol. Soc. Am.*, 364, 1–15, doi:10.1130/0-8137-
1432 2364-7.1, 2002.
- 1433 Robardet, M. and Gutiérrez Marco, J. C.: Ossa-Morena Zone. Stratigraphy. Passive Margin
1434 Phase (Ordovician-Silurian-Devonian), in *Pre-Mesozoic Geology of Iberia*, edited by R. D.
1435 Dallmeyer and E. Martínez García, pp. 267–272, Springer-Verlag, Berlin., 1990.
- 1436 Rodrigues, B., Chew, D. M., Jorge, R. C. G. S., Fernandes, P., Veiga-Pires, C. and Oliveira, J. T.:
1437 Detrital zircon geochronology of the Carboniferous Baixo Alentejo Flysch Group (South
1438 Portugal); Constraints on the provenance and geodynamic evolution of the South Portuguese
1439 Zone, *J. Geol. Soc. London.*, 172, 294–308, doi:10.1144/jgs2013-084, 2015.
- 1440 Ross, A. R., Brown, L. D., Pananont, P., Nelson, K. D., Klemperer, S. L., Haines, S., Wenjin, Z. and
1441 Jingru, G.: Deep reflection surveying in central Tibet: Lower-crustal layering and crustal flow,
1442 *Geophys. J. Int.*, 156(1), 115–128, doi:10.1111/j.1365-246X.2004.02119.x, 2004.
- 1443 Royden, L.: Coupling and decoupling of crust and mantle in convergent orogens: Implications
1444 for strain partitioning in the crust, *J. Geophys. Res.*, 101(B8), 17679–17705, 1996.
- 1445 Rubio Pascual, F. J., Arenas, R., Martínez Catalán, J. R., Rodríguez Fernández, L. R. and
1446 Wijbrans, J. R.: Thickening and exhumation of the Variscan roots in the Iberian Central System:
1447 Tectonothermal processes and 40Ar/39Ar ages, *Tectonophysics*, 587, 207–221,
1448 doi:10.1016/j.tecto.2012.10.005, 2013.
- 1449 Sánchez-García, T., Quesada, C., Bellido, F., Dunning, G. R. and González del Tánago, J.: Two-
1450 step magma flooding of the upper crust during rifting: The Early Paleozoic of the Ossa Morena
1451 Zone (SW Iberia), *Tectonophysics*, 461(1–4), 72–90, doi:10.1016/j.tecto.2008.03.006, 2008.
- 1452 Sánchez-García, T., Bellido, F., Pereira, M. F., Chichorro, M., Quesada, C., Pin, C. and Silva, J. B.:
1453 Rift-related volcanism predating the birth of the Rheic Ocean (Ossa-Morena zone, SW Iberia),
1454 *Gondwana Res.*, 17(2–3), 392–407, doi:10.1016/j.gr.2009.10.005, 2010.
- 1455 Sánchez de Posada, L. C., Martínez Chacón, M. L., Méndez Fernández, C., Menéndez Alvarez,
1456 J.R. Truyols, J. and Villa, E.: Carboniferous Pre-Stephanian rocks of the Asturian-Leonese
1457 Domain (Cantabrian Zone), in *Pre-Mesozoic Geology of Iberia*, edited by R. D. Dallmeyer and E.
1458 Martínez García, pp. 24–33, Springer-Verlag., 1990.
- 1459 Sánchez Martínez, S., Arenas, R., Andonaegui, P., Martínez Catalán, J. R. and Pearce, J. A.:
1460 Geochemistry of two associated ophiolites from the Cabo Ortegal Complex (Variscan belt of
1461 NW Spain), *Mem. Geol. Soc. Am.*, 200(23), 445–467, doi:10.1130/2007.1200(23), 2007.
- 1462 Schmelzbach, C., Juhlin, C., Carbonell, R. and Simancas, J. F.: Prestack and poststack migration
1463 of crooked-line seismic reflection data: A case study from the South Portuguese Zone fold belt,
1464 southwestern Iberia, *Geophysics*, 72(2), 9–18, doi:10.1190/1.2407267, 2007.
- 1465 Schmelzbach, C., Simancas, J. F., Juhlin, C. and Carbonell, R.: Seismic reflection imaging over
1466 the South Portuguese Zone fold-and-thrust belt, SW Iberia, *J. Geophys. Res. Solid Earth*, 113(8),
1467 1–16, doi:10.1029/2007JB005341, 2008.
- 1468 Seyferth, M. and Henk, A.: Syn-convergent exhumation and lateral extrusion in continental

- 1469 collision zones - Insights from three-dimensional numerical models, *Tectonophysics*, 382(1–2),
1470 1–29, doi:10.1016/j.tecto.2003.12.004, 2004.
- 1471 Shelley, D. and Bossière, G.: A new model for the Hercynian Orogen of Gondwanan France and
1472 Iberia, *J. Struct. Geol.*, 22(6), 757–776, doi:10.1016/S0191-8141(00)00007-9, 2000.
- 1473 Simancas, J. F., Martínez Poyatos, D., Expósito, I., Azor, A. and González Lodeiro, F.: The
1474 structure of a major suture zone in the SW Iberian massif: The Ossa-Morena/central Iberian
1475 contact, *Tectonophysics*, 332(1–2), 295–308, doi:10.1016/S0040-1951(00)00262-6, 2001.
- 1476 Simancas, J. F., Carbonell, R., González Lodeiro, F., Pérez-Estaún, A., Juhlin, C., Ayarza, P.,
1477 Kashubin, A., Azor, A., Martínez Poyatos, D., Almodóvar, G. R., Pascual, E., Sáez, R. and
1478 Expósito, I.: Crustal structure of the transpressional Variscan orogen of SW Iberia: SW Iberia
1479 deep seismic reflection profile (IBERSEIS), *Tectonics*, 22(6), doi:10.1029/2002TC001479, 2003.
- 1480 Simancas, J. F., Carbonell, R., Gonzalez Lodeiro, F., Pérez-Estaún, A., Juhlin, C., Ayarza, P.,
1481 Kashubin, A., Azor, A., Martínez Poyatos, D., Saez, R., Almodovar, G. R., Pascual, E., Flecha, I.
1482 and Marti, D.: Transpressional collision tectonics and mantle plume dynamics: the Variscides of
1483 southwestern Iberia, *Geol. Soc. London, Mem.*, 32(1), 345–354, 2006.
- 1484 Simancas, J. F., Ayarza, P., Azor, A., Carbonell, R., Martínez Poyatos, D., Pérez-Estaún, A. and
1485 González Lodeiro, F.: A seismic geotraverse across the Iberian Variscides: Orogenic shortening,
1486 collisional magmatism, and orocline development, *Tectonics*, 32(3), 417–432,
1487 doi:10.1002/tect.20035, 2013.
- 1488 Snelson, C. M., Randy Keller, G., Miller, K. C., Rumpel, H. M. and Prodehl, C.: Regional Crustal
1489 Structure Derived from the CD-ROM 99 Seismic Refraction/Wide-Angle Reflection Profile: The
1490 Lower Crust and Upper Mantle, in *The Rocky Mountain Region: An Evolving Lithosphere:*
1491 *Tectonics, Geochemistry, and Geophysics*, pp. 271–291., 2013.
- 1492 Spear, F. S.: *Metamorphic Phase Equilibria and pressure-temperature-time paths*, Monograph.,
1493 edited by M. Mineralogical Society of America, Chelsea., 1993.
- 1494 Stille, H.: *Grundfragen der Vergleichenden. Tectonik*, Gebrueder Borntragen, Berlin., 1924.
- 1495 Taner, M. T. and Sheriff, R. E.: Application of Amplitude, Frequency, and Other Attributes to
1496 Stratigraphic and Hydrocarbon Determination, in *AAPG Memoir, Seismic Stratigraphy —*
1497 *Applications to Hydrocarbon Exploration*, vol. 26, pp. 301–327., 1977.
- 1498 Teixell, A., Labaume, P., Ayarza, P., Espurt, N., de Saint Blanquat, M. and Lagabrielle, Y.: Crustal
1499 structure and evolution of the Pyrenean-Cantabrian belt: A review and new interpretations
1500 from recent concepts and data, *Tectonophysics*, 724–725(July 2017), 146–170,
1501 doi:10.1016/j.tecto.2018.01.009, 2018.
- 1502 Truyols, J., Arbizu, M., Garcia-Alcalde, J. L., Garcia-López, S., Mendez-Bendia, I., Soto, F. and
1503 Truyols, M.: The Asturian-Leonese Domain (Cantabrian Zone), in *Pre-Mesozoic Geology of*
1504 *Iberia*, edited by R. D. Dallmeyer and E. Martínez García, pp. 10–19, Springer-Verlag., 1990.
- 1505 de Vicente, G., Giner, J. L., Muñoz-Martí, A., González-Casado, J. M. and Lindo, R.:
1506 Determination of present-day stress tensor and neotectonic interval in the Spanish Central
1507 System and Madrid Basin, central Spain, *Tectonophysics*, 266(1–4), 405–424,
1508 doi:10.1016/S0040-1951(96)00200-4, 1996.
- 1509 Villaseca, C., Orejana, D. and Belousova, E. A.: Recycled metaigneous crustal sources for S- and
1510 I-type Variscan granitoids from the Spanish Central System batholith: Constraints from Hf

1511 isotope zircon composition, *Lithos*, 153, 84–93, doi:10.1016/j.lithos.2012.03.024, 2012.

1512 Wever, T.: The Conrad discontinuity and the top of the reflective lower crust-do they
 1513 coincide?, *Tectonophysics*, 157(1–3), 39–58, doi:10.1016/0040-1951(89)90339-9, 1989.

1514 Xiaobo, H. and Tae Kyung, H.: Evidence for strong ground motion by waves refracted from the
 1515 Conrad discontinuity, *Bull. Seismol. Soc. Am.*, 100(3), 1370–1374, doi:10.1785/0120090159,
 1516 2010.

1517

1518

1519 Table 1: Acquisition parameters of the NI seismic profiles shown from figures 3 to 8 and
 1520 former processing flows. These are grouped according to their similarities.

1521	Acquisition parameters	ESCIN-1 (onshore)	ESCIN-2 (onshore)
1522	Source	Dynamite-single-hole	Dynamite-single-hole
1523	Charge	20 kg at 24 m depth	20 kg at 24 m depth
1524	Trace Interval	60 m	60 m
1525	# of traces	240	240
1526	Spread configuration	Symmetrical Split-spread	Symmetrical Split-spread
1527	Fold	30	30
1528	Geophones per group	18	18
1529	Spread length	14.5 km	14.5 km
1530	Sample interval rate	4 ms	4 ms
1531	<u>Kirchoff time migration</u>	<u>5600 m/s</u>	<u>5600 m/s</u>
1532			
1533		ESCIN-3.2 and ESCIN-3.3 (offshore)	
1534			
1535	Source	Airgun (5490 cu.in)	
1536	Shot spacing	75 m	
1537	Receiver interval	12.5 m	
1538	Spread length	4500 m	
1539	Fold	30	
1540	Internal offset	240 m	
1541	Sample rate	4 ms	
1542	Record length	20 s	
1543	<u>Kirchoff time migration</u>	<u>5200 m/s</u>	
1544			
1545		IBERSEIS (onshore)	ALCUDIA (onshore)
1546			
1547	Source	4, 22T vibrators	4 (+1), 22T vibrators
1548	Recording instrument	SERCEL 388, 10 Hz	SERCEL 388, 10 Hz
1549	# active channels	240 minimum	240 minimum
1550	Station spacing	35 m	35 m
1551	Station configuration	12 geophones	12 geophones
1552	Source spacing	70 m	70 m
1553	Sweep frequencies	non-linear 8-80 Hz	non-linear 8-80 Hz
1554	Sweep length	20 s	20 s
1555	Listening time	40 s	40 s
1556	Sample rate	2 ms	4 ms

1557	Spread type	Asymmetric split-spread	Asymmetric split-spread
1558	Nominal fold	60 (minimum)	60 (minimum)
1559	<u>Kirchoff time migration</u>	<u>6000 m/s</u>	<u>6200 m/s</u>

1560
1561
1562 Figure captions
1563

1564 Figure 1: (a) Map of the Iberian Peninsula showing the outcrops of the Variscan basement and
1565 the subdivision in zones of the Iberian Massif. The main strike-slip shear zones, traces of
1566 Variscan folds and gneiss domes are also included. Blue lines show the position of normal
1567 incidence seismic reflection profiles and that of the CIMDEF transect. ~~Abbreviations:~~
1568 ~~Allochthonous complexes of NW Iberia: B: Bragança; CO: Cabo Ortegal; M: Morais; MT:~~
1569 ~~Malpica-Tui; O: Órdenes. Strike-slip shear zones: BCSZ: Badajoz-Córdoba; DBSZ: Douro-Beira;~~
1570 ~~JPSZ: Juzbado-Penalva; PTSZ: Porto Tomar; SISZ: Southern Iberian. See legend for other~~
1571 ~~abbreviations. Traces of the main Variscan folds and the Variscan granitoids are also included.~~
1572 ~~See legend for abbreviations. (b) Map of the outcropping granitoids in the Iberian Peninsula~~
1573 ~~together with the main structures.~~

1574 Figure 2: Processing flow carried over the SEG-Y original stack sections. This task was
1575 geared to improve and homogenize the resolution of the seismic images while creating
1576 new migrated sections. See Martínez García, (2019) for further details.

1577 Figure 3: Cross-section (a) and depth converted time migrated section (v=5600 m/s) of
1578 along the NI seismic profile ESCIN-1 (Fig. 1), without (ba) and with interpretation (cb). A
1579 sketch of the most important features is presented in (de). CDP: Common Depth Point.
1580 TWT: Two-way travel time. WALZ: West Asturian-Leonese Zone. CZ: Cantabrian Zone. The
1581 position of the Narcea Antiform is indicated. Red dashed lines represent the boundaries
1582 provided by chaos and variance attribute analyses. Nomenclature for reflections goes as
1583 follows: (ot), outcropping thrusts; (t), thrusts affecting the basement; (b), indifferiated
1584 basement; (st), sole thrust; (c), top of the lower crust; (ir), lower crust internal reflectivity;
1585 (m) is the Moho; (mig) are curved features resulting from the migration of discontinuous
1586 reflections. Depth conversion is based on migration velocities.

1587 Figure 4: Cross-section ((a) modified from Gallastegui et al., 2016) and depth converted
1588 time migrated section (v=5600 m/s) of the NI seismic profile ESCIN-2 (Fig. 1), without (ba)
1589 and with interpretation (cb). A sketch of the most important features is presented in (de).
1590 A 1D velocity profile as modeled from wide-angle data (Pulgar et al., 1996) appears in (e).
1591 CDP: Common Depth Point. TWT: Two-way travel time. ~~WALZ: West Asturian-Leonese~~
1592 ~~Zone.~~ CZ: Cantabrian Zone. ~~DB: Duero Basin.~~ Some discontinuous reflections are traced on
1593 the basis of their geometry on the stack image (Pulgar et al., 1996). Red dashed lines
1594 represent the boundaries provided by chaos and variance attribute analyses. (s),
1595 sediments; (t), thrusts; (ot), outcropping thrusts; (ps), Paleozoic sediments; (c), top of the
1596 lower crust; (lc), lower crust; (m), Moho. Depth conversion is based on migration
1597 velocities.

1598 Figure 5: Cross-section (a) and depth converted time migrated section (v=5200 m/s) of
1599 the NI seismic profile ESCIN-3.3 (Fig. 1), without (ba) and with interpretation (cb). A sketch
1600 of the most important features is presented in (de). A 1D velocity profile as modeled from

1601 wide-angle data (Ayarza et al., 1998) in presented in (e). CDP: Common Depth Point. TWT:
1602 Two-way travel time. CZ: Cantabrian Zone. WALZ: West Asturian-Leonese Zone. CIZ:
1603 Central Iberian Zone. The offshore projection of the Viveiro Fault is indicated. Red dashed
1604 lines represent the boundaries provided by chaos and variance attribute analyses. (s),
1605 sediments; (t), thrusts; (st), sole thrust; (ef), extensional features; (c), top of the lower
1606 crust; (lc), lower crust; (ir) lower crust internal reflectivity; (m), Moho; (sc), subcrustal
1607 reflections; (mig) curved features resulting from the migration of discontinuous reflections.
1608 Depth conversion is based on migration velocities.

1609 Figure 6: Cross-section (a) and depth converted time mMigrated section (v=5200 m/s) of
1610 the NI seismic profile ESCIN-3.2 (Fig. 1), without (ba) and with interpretation (cb). A sketch
1611 of the most important features is presented in (de). CDP: Common Depth Point. TWT: Two-
1612 way travel time. GTMZ: Galicia-Trás-os-Montes Zone. CIZ: Central Iberian Zone. (s),
1613 sediments; (ef), extensional features; (lc), lower crust; (m), Moho; (sc), subcrustal
1614 reflections; (czlc) CZ underthrust lower crust. Depth conversion is based on migration
1615 velocities.

1616 Figure 7: Cross-section ((a) modified from Martínez Poyatos et al., 2012) and depth
1617 converted time mMigrated section (v=6200 m/s) of the NI seismic profile ALCUDIA (Fig. 1),
1618 without (ba) and with interpretation (cb). A sketch of the most important features is
1619 presented in (de). A 1D velocity profile as modeled from wide-angle data (Palomeras et al.,
1620 in press) is overlapped in (d). CDP: Common Depth Point. TWT: Two-way travel time. CIZ:
1621 Central Iberian Zone. OMZ: Ossa-Morena Zone. (cu): Central Unit; (vf): vertical folds; (g):
1622 granites; (ef): extensional features; (c): top of the lower crust; (lc): lower crust; (cc) and
1623 (cc'): crocodile structure; (m): Moho. ~~CU: Central Unit.~~ Red dashed lines represent the
1624 boundaries provided by chaos and variance attribute analyses. Depth conversion is based
1625 on migration velocities.

1626 Figure 8: Cross-section ((a), modified from Simancas et al., 2003) and depth converted time
1627 mMigrated section (v=6000 m/s) of the NI seismic profile IBERSEIS (Fig. 1), without (ba) and
1628 with interpretation (cb). A sketch of the most important features is presented in (de). Two
1629 1D velocity profiles as modeled from wide-angle data (Palomeras et al., 2009)are
1630 overlapped in (d). CDP: Common Depth Point. TWT: Two-way travel time. CIZ: Central
1631 Iberian Zone. CU: Central Unit. OMZ: Ossa-Morena Zone. RORS: Rheic ocean reworked
1632 suture. SPZ: South Portuguese Zone. (t), thrusts; (ot), outcropping thrusts; (irb), Iberseis
1633 Reflective Body; (c), top of the lower crust; (lc), lower crust; (ir) lower crust internal
1634 reflectivity; (lct), lower crust thrusts; (m), Moho. Depth conversion is based on migration
1635 velocities.

1636 Figure 9: Joint geological interpretation of all the seismic sections (normal incidence and
1637 seismic noise) whose location is shown in figure 1. (a): ESCIN-1, ESCIN3-3 and ESCIN-3.2.
1638 (b): ALCUDIA and IBERSEIS. (c): CIMDEF (Andrés et al., 2020). Special attention should be
1639 paid to the depth and geometry of the Moho and mid-crustal discontinuity. Alpine
1640 structures (i.e. crustal thickening) appear in ESCIN-1 and in CIMDEF. The rest are Variscan
1641 features.

1642 Figure 10: Map of the Moho depth as derived from tomography of shear waves (seismic
1643 noise and earthquakes, Palomeras et al., 2017) with the projection of the seismic profiles

Código de campo cambiado

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

1644 already shown in figure 1 and described along the text. A sketch of the geometry of the
1645 main discontinuities (Moho and Conrad) is also shown.