

# Early Cenozoic Eurekan strain partitioning and decoupling in central Spitsbergen, Svalbard

Jean-Baptiste P. Koehl<sup>1,2,3,4</sup>

<sup>1</sup>Centre for Earth Evolution and Dynamics (CEED), University of Oslo, PO Box 1028 Blindern, N-0315 Oslo Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, P.O. Box 5003 NMBU, NO-1432 Ås, Norway.

<sup>2</sup>Department of Geosciences, UiT The Arctic University of Norway in Tromsø, NO-9037 Tromsø, Norway.

<sup>3</sup>Research Centre for Arctic Petroleum Exploration (ARCEX), University of Tromsø, NO-9037 Tromsø, Norway.

<sup>4</sup>CAGE – Centre for Arctic Gas Hydrate, Environment and Climate, NO-9037 Tromsø, Norway.

**Correspondence:** Jean-Baptiste P. Koehl (jean-baptiste.koehl@uit.no)

## Abstract

The present study of field, petrological, exploration well and seismic data ~~shows that~~ describes backward-dipping duplexes comprised of phyllitic coal and bedding-parallel décollements and thrusts, ~~which~~ localized along lithological transitions in tectonically thickened Lower–lowermost Upper Devonian, uppermost Devonian–Mississippian and uppermost Pennsylvanian–lowermost Permian sedimentary strata of the Wood Bay and/or Widje Bay and/or Grey Hoek formations, of the Billefjorden Group and of the Wordiekammen Formation respectively. ~~The study shows that~~ these structures partially decoupled uppermost Devonian–Permian sedimentary rocks of the Billefjorden and Gipsdalen groups from Lower–lowermost Upper Devonian rocks of the André Land Group and Mimerdalen Subgroup during early Cenozoic Eurekan deformation in central Spitsbergen. Eurekan strain decoupling along these structures explains differential deformation between Lower–lowermost Upper Devonian rocks of the André Land Group/Mimerdalen Subgroup and overlying uppermost Devonian–Permian sedimentary strata of the Billefjorden and Gipsdalen groups in central–northern Spitsbergen without requiring an episode of (Ellesmerian) contraction in the Late Devonian. Potential formation mechanisms for bedding-parallel décollements and thrusts include shortcut faulting, and/or formation as a roof décollement in a fault-bend hanging wall (or ramp) anticline, as an imbricate fan, as an antiformal thrust stack, and/or as fault-propagation folds over reactivated/overprinted basement-seated faults. The

interpretation of seismic data in Reindalspasset indicates that Devonian sedimentary rocks of the Andrée Land Group and Mimerdalen Subgroup might be preserved east of the Billefjorden Fault Zone, suggesting that the Billefjorden Fault Zone did not accommodate reverse movement in the  
35 Late Devonian. Hence, the thrusting of Proterozoic basement rocks over Lower Devonian sedimentary rocks along the Balliolbreen Fault and fold structures within strata of the Andrée Land Group and Mimerdalen Subgroup in central Spitsbergen may be explained by a combination of down-east Carboniferous normal faulting with associated footwall rotation and exhumation, and subsequent top-west early Cenozoic Eurekan thrusting along the Billefjorden Fault Zone. Finally,  
40 the study shows that major east-dipping faults, like the Billefjorden Fault Zone, may consists of several, discrete, unconnected (soft-linked and/or stepping) or, most probably, offset fault segments that were reactivated/overprinted with varying degree during Eurekan deformation due to strain partitioning and/or decoupling along sub-orthogonal NNE-dipping reverse faults.

## 45 1. Introduction

The main goal of this contribution is to examine the influence of strain decoupling and partitioning on deformation patterns within Devonian–Permian sedimentary successions in central Spitsbergen during the early Cenozoic Eurekan tectonic event. The impact of this event, though well studied in western Spitsbergen where it resulted in the formation of the West Spitsbergen  
50 Fold-and-Thrust Belt (Dallmann et al., 1988, 1993; Braathen et al., 1999) with multiple levels of detachment and décollement (Maher, 1984; Maher et al., 1986; Bergh et al., 2000), lacks detailed characterization in central Spitsbergen (~~Figure 1~~~~Figure 1a~~; see DataverseNO for high-resolution versions of all figures and supplements <https://doi.org/10.18710/MXKQPE>).

The study ~~has potential implications for strain partitioning in rift systems and distal parts of fold-and-thrust belts. Notably, the study discusses the presence of~~~~describes~~ bedding-parallel décollement levels and imbricate link thrusts (McClay and Insley, 1986) arranged into gently dipping duplexes within weak sedimentary beds of the Andrée Land Group, Billefjorden Group and Wordiekammen Formation, ~~which were, thus far, not described~~, and ~~discusses~~ their role in partially decoupling Eurekan deformation in late Paleozoic sedimentary successions. Potential  
60 formation mechanisms, such as shortcut faulting (Buitter and Pfiffner, 2003), and the influence of preexisting inherited structures (e.g., Billefjorden Fault Zone) are reviewed.

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

The identification of structures showing comparable geometries and kinematics (e.g., bedding-parallel décollements) within discrete stratigraphic units (e.g., coals and coaly shales of the Billefjorden Group) both on nearshore seismic data and onshore during structural fieldwork further validates the use of seismic interpretation in areas where extensive (glacial) erosion resulted in partial destruction and covering of outcrop transects with loose material, and where large portions of the outcrops available for field mapping are hardly accessible for detailed inspection because located on steep slopes and cliffs. The study also illustrates the complementarity between fieldwork, which provide detailed lithological and structural data, and seismic transects providing continuous transects through deformation belts and fault zones.

The study also briefly discusses implications for the Ellesmerian Orogeny, a poorly constrained short-lived episode of contractional deformation in the Late Devonian that presumably explains the juxtaposition of Proterozoic basement against Lower–lowermost Upper Devonian sedimentary strata of the Andrée Land Group and Mimerdalen Subgroup (Vogt, 1938; Harland et al., 1974; McCann, 2000; Piepjohn, 2000; Piepjohn et al., 2000; Piepjohn and Dallmann, 2014), and differential deformation between folded Devonian rocks of the Andrée Land Group and Mimerdalen Subgroup and poorly deformed rocks of the uppermost Devonian–Permian Billefjorden and Gipsdalen groups in Dickson Land in central Spitsbergen. Notably, the contribution shows that Eurekan deformation localized in weak, intensely deformed sedimentary units of the uppermost Devonian–Permian sedimentary rocks of the Billefjorden and Gipsdalen rocks, that Devonian sedimentary rocks of the Andrée Land Group are possibly preserved east of the Billefjorden Fault Zone and, thus, that the Billefjorden Fault Zone most likely did not act as a reverse fault in the Late Devonian, and that juxtaposition of Proterozoic basement against Lower Devonian rocks in central Spitsbergen may be achieved through Carboniferous normal faulting and early Cenozoic Eurekan top-west thrusting. Hence, the study contributes to our understanding of deformation partitioning in fold-and-thrust belts consisting of thick sedimentary successions, and for the extent of the Ellesmerian Orogeny in the Arctic, which presumably extends from Arctic Canada and northern Greenland to Spitsbergen.

Finally, the study has implication for the segmentation and linkage of rift-bounding fault with long-lived tectonic histories. Thus far, although segmentation of the Billefjorden Fault Zone was described (e.g., Bælum and Braathen, 2012), along-strike geometrical and kinematics variations along the Billefjorden Fault Zone have been poorly addressed and tentatively attributed

to the complex tectonic history of this fault. The present study further discusses ~~considers~~ the significant along-strike variations in geometry and kinematics ~~of the Billefjorden Fault Zone, and discusses the, the~~ extent, and potential segmentation of ~~the~~ Billefjorden Fault Zone ~~fault~~ in conjunction with a new trend of NNE-dipping faults striking suborthogonal to the main N–S-trending structural grain in the study area. The role of these suborthogonal faults in Eurekan strain partitioning is briefly discussed.

## 2. Geological setting

### 2.1. Caledonian Orogeny

Spitsbergen is composed of three terranes that started assembling during the late Cambrian–Silurian Caledonian Orogeny and were juxtaposed against one another by N–S-striking crustal faults like the Billefjorden Fault Zone (Harland and Wright, 1979; Ohta et al., 1989, 1995; Gee and Page, 1994). Caledonian deformation was accompanied by tectonothermal events with high-grade (eclogite and blueschist) metamorphism from mid-Cambrian to late Silurian times that occurred during subduction and closure of the Iapetus Ocean and that are partly preserved in northwestern (Ohta et al., 1989) and western Spitsbergen (Horsfield, 1972; Kościńska et al., 2014).

Caledonian grain in western, northwestern, central and eastern Spitsbergen forms major, gently plunging, N–S-trending folds and thrust stacks with well-developed foliation. An example ~~is e.g.~~ the Atomfjella Antiform in Ny-Friesland (Figure 1 ~~Figure 1b~~), an antiformal thrust stack that consists of a succession of nappes composed of Proterozoic granite and metasedimentary rocks separated by west-verging (Flood et al., 1969; Balashov et al., 1993; Witt-Nilsson et al., 1998; Johansson and Gee, 1999; Johansson et al., 2004, 2005) and/or top-east thrusts (Manby and Michalski, 2014).

### 2.2. Devonian late–post-orogenic collapse

In the Early Devonian, late–post-Caledonian gravitational collapse initiated (Chorowicz, 1992; Roy, 2007, 2009; Roy et al., unpublished) leading to the deposition of several km-thick (Old Red Sandstone) basins throughout Spitsbergen (Birkenmajer and Turnau, 1962; Harland et al., 1974; Manby and Lyberis, 1992; Manby et al., 1994; Dallmann and Piepjohn, 2020) and emplacement of late-orogenic plutons in northwestern, central and eastern Spitsbergen (Hamilton et al., 1962; Gayer et al., 1966; Ohta et al., 2002; Myhre et al, 2008).

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

125 In northern Spitsbergen, Devonian sedimentary rocks of the Siktefjellet, Red Bay and  
Andrée Land groups (Gee and Moody-Stuart, 1966; Friend et al., 1966; Friend and Moody-Stuart,  
1972; Murascov and Mokin, 1979; Friend et al., 1997; Figure 2) deposited during extension and  
subsidence along N–S-striking normal faults, forming west-tilted (half-) grabens, e.g., in  
Raudfjorden, Bockfjorden (Manby and Lyberis, 1992; Manby et al., 1994), Andrée Land and Kota  
(Roy, 2007, 2009; Roy et al., unpublished; Figure 1~~Figure 1a~~). However, other works argue that  
130 Devonian sedimentary deposits of the Andrée Land Group and Mimerdalen Subgroup (Figure 2)  
in Svalbard deposited along low-angle, post-Caledonian detachments that accommodated large  
amounts of top-east, normal movement (e.g., the Woodfjorden detachment) and are associated with  
syn-kinematic east-verging folds (Roy, 2007, 2009; Roy et al., unpublished). In addition, recent  
studies show that basement ridges, e.g., the Bockfjorden Anticline in northwestern Spitsbergen,  
135 may have exhumed as core complexes along low-angle extensional detachments (e.g., the  
Keisarhjelmen detachment), and K–Ar geochronology suggests that exhumation occurred from late  
Silurian to Late Devonian times (Braathen et al., 2018).

### 2.3. Ellesmerian Orogeny

140 Ellesmerian deformation is thought to have initiated in the Late Devonian–Early  
Mississippian, possibly in the Late Frasnian–Famennian (Vigran, 1964; Allen, 1965, 1973; Pcelina  
et al., 1986; Brinkmann, 1997; Schweitzer, 1999; Piepjohn et al., 2000) and was presumably  
recorded by the deposition of coarse-grained sedimentary rocks of the Mimerdalen Subgroup  
(Planteryggen and Plantekløfta formations; Piepjohn and Dallmann, 2014). However, recent fossil  
145 and spore analysis suggest an early Frasnian (ca. 380 Ma) age for these stratigraphic units (Berry  
and Marshall, 2015). Deformation is believed to have stopped prior to the deposition of middle–  
late Famennian–Mississippian (Scheibner et al., 2012; Lindemann et al., 2013; Marshall et al.,  
2015; Würtzen et al., 2019; Lopes, pers. comm. 2019) sedimentary rocks of the Billefjorden Group  
(Vogt, 1938; Piepjohn, 2000). Previous works also suggested that hundreds–thousands of  
150 kilometer-scale strike-slip movement along N–S-striking faults, e.g., Billefjorden Fault Zone,  
finalized the accretion of basement terranes constituting the Svalbard Archipelago (Harland et al.,  
1974; Harland and Wright, 1979; Ohta et al., 1989), while more recent studies argue for limited  
amounts of strike-slip movement (McCann, 2000; Piepjohn, 2000).

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

Field Code Changed

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

In Pyramiden, in Dickson Land (northern–central Spitsbergen; ~~Figure 1~~~~Figure 1b~~),  
155 Proterozoic basement rocks were thrust top-west onto Lower Devonian sedimentary rocks of the  
Wood Bay Formation along the Balliolbreen Fault (Harland et al., 1974; Piepjohn, 2000; Bergh et  
al., 2011) in Late Devonian times, and presumably undeformed uppermost Devonian–  
Mississippian clastic and coal-bearing sedimentary deposits of the Billefjorden Group overlie  
160 folded Lower–lowermost Upper Devonian metasedimentary rocks that were involved in  
Ellesmerian deformation. In Triungen (~~Figure 1~~~~Figure 1a–b~~), folded–gently dipping Lower  
Devonian rocks of the Wood Bay Formation are juxtaposed against flat-lying, undeformed,  
uppermost Devonian–Permian strata of the Billefjorden Group and Wordiekammen Formation  
along the Triungen–Grønhordalen Fault Zone (McCann and Dallmann, 1996). In Sentinelfjellet  
and Odellfjellet (~~Figure 1~~~~Figure 1b~~), the Balliolbreen Fault thrust Proterozoic basement rocks in  
165 the hanging wall over Devonian sedimentary rocks of the Andrée Land Group and Mimerdalen  
Subgroup in the footwall and is thought to be unconformably overlain by undeformed, uppermost  
Devonian–Mississippian sedimentary rocks of the Billefjorden Group, thus suggesting Late  
Devonian top-west thrusting (Friend and Moody-Stuart, 1972; Harland et al., 1974; Lamar et al.,  
1986).

170

#### 2.4. Carboniferous basins

In Carboniferous times, ENE–WSW extension formed narrow, kilometer- to tens of  
kilometer-wide, N–S- to NW–SE-trending troughs, e.g., Billefjorden Trough (Maher Jr., 1996;  
McCann and Dallmann, 1996; Braathen et al., 2011), bounded by major faults such as the  
175 Billefjorden Fault Zone (Harland et al., 1974), which was reactivated as a normal fault from  
Odellfjellet in the north to Reindalspasset in the south (Bælum and Braathen, 2012; ~~Figure 1~~~~Figure  
1a–b~~).

Shortly after the end of Ellesmerian deformation, partly eroded Devonian sedimentary  
rocks of the Andrée Land Group and Mimerdalen Subgroup were covered by uppermost Devonian–  
180 Mississippian (Marshall et al., 2015), fluvial, coal- and clastic-rich deposits of the Billefjorden  
Group (Cutbill and Challinor, 1965; Cutbill et al., 1976; Aakvik, 1981; Gjelberg, 1981, 1984;  
~~Figure 2~~). These are divided into the Hørbye breen and Mumien formations, which are composed  
of the Triungen and Hoelbreen, and Sporehøgda and Birger Johnsonfjellet members respectively.  
The Triungen and Sporehøgda members dominantly consist of clastics whereas the Hoelbreen and

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

185 Birger Johnsonfjellet members are composed of coal seams and coaly shales (Cutbill and Challinor, 1965; Cutbill et al., 1976; Aakvik, 1981; Gjelberg and Steel, 1981; Gjelberg, 1984; [Figure 2](#)).

**Formatted:** Font: (Default) Times New Roman, 12 pt, Not Italic

**Formatted:** Font: Not Bold, Not Italic

190 These deposits are found in Arctic areas stretching from the Barents Sea (Bugge et al., 1995; Larssen et al., 2002) to Arctic Canada (Emma Fiord Formation; Davies and Nassichuck, 1988) and were presumably deposited during a period of tectonic quiescence (Johannessen and Steel, 1992; Braathen et al., 2011; Smyrak-Sikora et al., 2018), though a syn-tectonic deposition was also proposed for these rocks in Arctic Canada (Beauchamp et al., 2018), the Barents Sea (Koehl et al., 2018), Bjørnøya (Gjelberg, 1981), and in Spitsbergen in the northern part of the Billefjorden Trough (Koehl and Muñoz-Barrera, 2018).

195 In the Pennsylvanian, fluvial to shallow marine sedimentary strata of the Gipsdalen Group were deposited in subsiding basins. These are divided into the Hultberget, Ebbadalen, Minkinfjellet, Wordiekammen and Gipshuken formations in central Spitsbergen (Cutbill and Challinor, 1965; Johannessen, 1980; Gjelberg and Steel, 1981; Johannessen and Steel, 1992; Braathen et al., 2011; Smyrak-Sikora et al., 2018), all of which range from late Serpukhovian to earliest Permian in age ([Figure 2](#)).

**Formatted:** Font: (Default) Times New Roman, 12 pt, Not Italic

**Formatted:** Font: Not Bold, Not Italic

200 Sedimentary strata of the Gipsdalen Group are mostly composed of clastic, carbonate and evaporitic deposits and karst breccia, and represent the thickest sedimentary succession in the Billefjorden Trough (McWhae, 1953; Cutbill and Challinor, 1965; Holliday and Cutbill, 1972; Johannessen, 1980; Lønøy, 1995; [Figure 2](#)). The deposition of sedimentary strata of the Hultberget, Ebbadalen, and Minkinfjellet formations was accompanied by kilometer scale normal displacement along N–S-striking faults like the Billefjorden Fault Zone, whereas the Wordiekammen and Gipshuken formations were deposited during minor tectonic activity (Gjelberg and Steel, 1981; Fedorowski, 1982; Braathen et al., 2011; Smyrak-Sikora et al., 2018).

**Formatted:** Font: (Default) Times New Roman, 12 pt, Not Italic

**Formatted:** Font: Not Bold, Not Italic

### 2.5. Eureka deformation

210 In the Paleocene (ca. 62 Ma), Eureka deformation initiated in western Spitsbergen due to the opening of the Labrador Sea and Baffin Bay between Canada and Greenland (Chalmers and Pulvertaft, 2001; Oakey and Chalmers, 2012) and resulted in the formation of the West Spitsbergen Fold-and-Thrust Belt between Kongsfjorden and Sørkapp (Harland, 1969; Lowell, 1972; Harland and Horsfield, 1974; Maher et al., 1986; Dallmann et al., 1988, 1993; Andresen et al., 1994; Bergh and Grogan, 2003) and formation of a foreland basin, the Tertiary Central Basin, in central

Spitsbergen (Larsen, 1988; Petersen et al., 2016). Eurekan thrusts and folds in Spitsbergen dominantly strike and trend NNW–SSE (Harland and Horsfield, 1974; Bergh and Andresen, 1990; Dallmann et al., 1993; Bergh et al., 2011; Blinova et al., 2012) except in Kongsfjorden (Figure 1, Figure 1a) where they strike and trend WNW–ESE (Bergh and Andresen, 1990; Bergh et al., 2000; Saalman and Thiedig, 2000, 2001; Piepjohn et al., 2001). Early Cenozoic thrusts in western Spitsbergen commonly form décollements in shaly beds, e.g., in Triassic shales in Midterhuken (Maher, 1984; Maher et al., 1986; Figure 1, Figure 1a). In central–eastern Spitsbergen, major N–S-striking brittle faults like the Billefjorden Fault Zone were partly reactivated by Eurekan deformation in Flowerdalen (Harland et al., 1974; Haremo et al., 1990; Haremo and Andresen, 1992; Figure 1, Figure 1b) and Reindalspasset (Bælum and Braathen, 2012) in the south, but were apparently unaffected in northern areas like Sentinelfjellet (see location in Figure 1, Figure 1b) where uppermost Devonian–Mississippian strata seem to unconformably lie over the fault (Harland et al., 1974).

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

### 3. Methods

#### *3.1. Seismic, field and petrological data, and satellite images*

The present contribution uses ~~structural strike and dip~~ measurements of bedding and fracture surfaces in Devonian–Mississippian ~~sandstone, coals, and coaly shales of the Billefjorden Group~~ sedimentary strata collected in summer 2016 in Pyramiden (Figure 1, Figure 1b). ~~These were used to determine the state of deformation of the various lithological units of the Billefjorden Group, to infer the presence of major faults, and to assess fault kinematics. Unfortunately only few slickensides of poor quality were recorded and these are not presented in the present study.~~

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

-The study also uses microscopic analysis of fault rocks and sedimentary rocks adjacent to brittle faults as a confirmation tool (included in supplement 1). ~~Thin sections were cut perpendicular to potential brittle faults in the field to better observed brittle deformation and offset. Cohesive fault rock was exclusively encountered in a gully below the mine entrance in Pyramiden, along the potential field occurrence of the Balliolbreen Fault.~~

Seismic data in nearshore fjords in central Spitsbergen are from the Norwegian Petroleum Directorate and uninterpreted seismic lines are provided in supplement 2. Seismic interpretation was tied to data from exploration well 7816/12-1 in Reindalspasset (Figure 1, Figure 1a–b; Eide et al., 1991) and time–depth conversion of well data is based on checkshots from Equinor and Store

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Norske Spitsbergen Kulkompani. The well penetrated late Paleozoic–Mesozoic sedimentary rocks and ends at a depth of 2261 m with 54 m of uppermost Devonian–Mississippian strata of the Billefjorden Group.

250

#### 4. Results and interpretations

##### 4.1. Field and petrological data

###### 4.1.1. Pyramiden

255 In Pyramiden, a steeply east-dipping, N–S-striking brittle fault crops out in a gully below the entrance of the Russian coal mine (Figure 3Figure 2). This fault is located half-way to the mine in the gully and crosscuts steeply east-dipping Lower Devonian sedimentary rocks of the Wood Bay Formation (Figure 2), which are involved into a large fold structure with Devonian bedding surfaces locally overturned to the east (Figure 3Figure 2 and Figure 4Figure 3a, and supplement 3). The fault shows meter-thick lenses of cataclastic fault rock (supplement 1). Devonian 260 sedimentary rocks are dominated by poorly deformed quartz crystals showing undulose extinction and limited recrystallization (supplement 1), whereas cataclastic fault rock shows distributed fractures with little (centimeter-scale) to no displacement.

There is no trace of Proterozoic basement rocks in this area although field studies and geological maps suggest that Proterozoic basement was thrust over Lower Devonian strata along 265 the Balliolbreen Fault (McCann, 1993; McCann and Dallmann, 1996; Piepjohn et al., 1997; Dallmann et al., 1999, 2004; Bergh et al., 2011; svalbardkartet.npolar.no). Sample preparation for thin sectioning actually proved problematic for Devonian sedimentary rocks (quartzitic sandstone) located in the hanging wall of the presumed fault, which resulted in misleading thick sections showing quartz crystals resembling pyroxenes (supplement 1). Thus, it is more likely that earlier 270 maps showing exclusively Devonian–Mississippian sedimentary rocks of the Wood Bay Formation and Billefjorden Group (Figure 2) below the mine entrance by Harland et al. (1974), Aakvik (1981), Lamar et al. (1986), and Trust Arktikugol (1988; Sirotkin, pers. comm. 2019) are correct.

Farther up the gully, a one–two meter-thick succession of interbedded sandstone and coal is juxtaposed against steeply east-dipping Lower Devonian strata to the west and overlain by a (at 275 least three meter) thick layer of uppermost Devonian–Mississippian coals of the Billefjorden Group that shows phyllitic shear fabrics (Figure 3Figure 2 and Figure 4Figure 3b and supplement 4). The presence of abundant coal suggests that this one–two meters thick unit is part of the Billefjorden

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Group as well (Figure 2). Bedding surfaces within the one–two meter-thick succession dip gently–steeply to the east (Figure 4Figure 3a), display sigmoidal geometries with Z-like shapes, and terminate abruptly against the three meter-thick layer of uppermost Devonian–Mississippian phyllitic coal upwards and against Lower Devonian rocks downwards (dashed yellow lines in Figure 4Figure 3b). In addition, coaly shales within this succession display phyllitic fabrics similar to those observed within overlying coals, and seem to form repeated successions of alternating beds of sandstone and coaly shale truncated by steeply east-dipping sigmoidal fault surfaces (thin dashed red lines in Figure 4Figure 3b). The Z-like sigmoidal shape of bedding surfaces, phyllitic shear fabrics of the coaly shales, and possible repetitions of the succession suggest that the steeply east-dipping, sigmoidal faults crosscutting the succession are imbricate thrust faults (stereonet 3 in Figure 3Figure 2), i.e., possible link thrusts (McClay and Insley, 1986), which accommodated top-west to top-WNW movements. The truncation of sandstone–coaly shale beds upwards and downwards, the abrupt transition (partly covered by screens) with underlying Lower Devonian rocks and overlying uppermost Devonian–Mississippian coals, and Z-shaped phyllitic shear fabrics within overlying coals suggest that the sandstone–coaly shale succession is bounded by moderate–low-angle, east-dipping floor- and roof-thrusts (McClay, 1992) with top-west to top-WNW sense of shear. In cross-section, the interaction of intra-succession, steeply east-dipping link thrusts and inter-succession, moderate–low-angle floor- and roof-thrusts defines an east-dipping duplex structure (Boyer and Elliott, 1982) of imbricate thrusts bounded upwards and downwards by potential décollements and/or detachments parallel to original (i.e., prior to deformation) bedding surfaces (e.g., thick red lines showing the transition from interbedded coaly shales and sandstone to coal, and from coal to sandstone in Figure 4Figure 3b). The nomenclature of hindward/forward-dipping duplexes of Boyer and Elliott (1982) does not apply here since the foreland of the West Spitsbergen Fold-and-Thrust Belt (Tertiary Central Basin) is located southeast of Pyramiden. Thus, the term “backward” is used to describe the east-dipping character of the duplexes, i.e., oppositely to the inferred transport direction.

Above the mine entrance, sedimentary rocks of the Billefjorden Group are dominated by yellow sandstone that are crosscut by dominant WNW–ESE-striking fractures and subsidiary N–S- and ENE–WSW-striking fractures (stereonets 1 and 2 in Figure 3Figure 2) showing oblique-slip kinematics. Poorly preserved slickenside lineations did not yield any information on relative displacement between footwall and hanging wall. In the west, dark sandstone and quartzite crop

**Formatted:** Font: (Default) Times New Roman, 12 pt, Not Italic

**Formatted:** Font: Not Bold, Not Italic

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

out and contain fossil wood, which are probably Lower Devonian in age. The contact between the  
310 Lower (–lowermost Upper?) Devonian dark sandstone and uppermost Devonian–Mississippian  
yellow sandstone of the Billefjorden Group, and intra-Devonian lithological contacts (e.g., between  
Devonian quartzite and dark sandstone; [Figure 4Figure 3a](#)), although partly covered by screens  
and/or mostly made of loose blocks, do not appear to be faulted or tectonized and trend c. WNW–  
ESE to NW–SE as bedding surfaces appear to change from moderately–steeply east-dipping below  
315 the mine entrance to gently NNE-dipping above the mine entrance ([Figure 3Figure 2](#) and [Figure  
4Figure 3a](#)), i.e., parallel to the dominant fault trend in both uppermost Devonian–Mississippian  
(stereonet 1 in [Figure 3Figure 2](#)) and Lower (–lowermost Upper?) Devonian rocks (stereonet 4 in  
[Figure 3Figure 2](#)).

Noteworthy, most outcrops of uppermost Devonian–Mississippian strata in this part of the  
320 study area trend E–W to WNW–ESE. Thus, the dominance of WNW–ESE-striking faults is  
unlikely the result of measurements flawed by a preferential outcrop trend, since E–W- to WNW–  
ESE-trending outcrops would rather favor identification and measurement of N–S-striking faults.

A possible interpretation of outcrops and structures in Pyramidén ([Figure 1Figure 1b](#)) is  
that the subvertical, N–S-striking brittle fault within steeply east-dipping Lower Devonian strata in  
325 the gully below the coal mine entrance ([Figure 3Figure 2](#) and [Figure 4Figure 3a](#)) represents the  
Balliolbreen Fault segment of the Billefjorden Fault Zone, and that low-angle roof/floor thrusts  
between Lower (–lowermost Upper?) Devonian rocks and the overlying succession of uppermost  
Devonian–Mississippian sandstone, coaly shale and coal ([Figure 4Figure 3b](#)) correspond to the  
upward-flattening continuation of this fault. However, no fault was observed between Lower (–  
330 lowermost Upper?) Devonian rocks of the André Land Group (and Mimerdalen Subgroup) and  
sandstones of the Billefjorden Group above the mine, and lithological and stratigraphic contacts there  
display significantly different trends (WNW–ESE to NW–SE; [Figure 4Figure 3a](#)).

#### 4.1.2. Triungen

335 Fieldwork in Triungen (see location in [Figure 1Figure 1a–b](#)) shows that the Triungen–  
Grønhorgdalen Fault Zone (McCann and Dallmann, 1996) and the contact between Lower  
Devonian of the Wood Bay Formation and overlying uppermost Devonian–Mississippian  
sedimentary rocks of the Billefjorden Group along the fault are largely covered by dark screens  
([Figure 4Figure 3c–e](#)). In the hanging wall though, Lower Devonian sedimentary strata are

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

340 moderately tilted to the south and define an angular unconformity with overlying, flat-lying strata  
of the Billefjorden Group (Figure 4Figure 3c). Based on the presence of thick, flat-lying, coal-rich  
strata in the lower part of the Billefjorden Group overlying Lower Devonian sedimentary strata in  
the hanging wall of the fault, the dark screens along the fault trace (Figure 4Figure 3d–e) are ~~believed~~  
interpreted to represent uppermost Devonian–Mississippian coals–coaly shales that might have  
345 been dragged along the Triungen–Grønhorgdalen Fault Zone during tectonic movements.

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

## 4.2. Seismic data

### 4.2.1. Seismic units and stratigraphy

350 In seismic sections, Precambrian–Caledonian basement rocks commonly show chaotic  
reflections, most likely arising from their complex tectonic history (e.g., Caledonian folding,  
shearing, thrusting and post-Caledonian extensional and contractional overprints), and subparallel  
reflections, possibly corresponding to seismic artifacts (e.g., multiples; Figure 5Figure 4a–g; see  
DataverseNO for high-resolution versions of all figures and supplements;  
<https://doi.org/10.18710/MXKQPE>).

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

355 Potential Devonian rocks of the André Land Group in Reindalspasset (Figure 1Figure 1a–  
b and Figure 2) are characterized by partly disrupted, semi-continuous, sub-parallel to chaotic,  
moderate- to low-amplitude seismic reflections (Figure 5Figure 4g). The moderate- to low-  
amplitude character of internal seismic reflections within this seismic unit suggests that it is made  
up with relatively homogeneous deposits with minor lithological variations. Thus, Devonian rocks  
360 in Reindalspasset are ~~interpreted to consist~~likely composed of thick successions of medium- to  
fine-grained sedimentary rocks such as siltstone and shales, possibly of the Lower– Devonian  
Wood Bay (or time-equivalent Marietoppen Formation in southern Spitsbergen) and/or Middle  
Devonian Grey Hoek and/or Wijde Bay formations.

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

365 Uppermost Devonian–Mississippian sedimentary rocks (Figure 2) are characterized by  
high-amplitude seismic reflections that are most likely the product of acoustic impedance contrast  
between low density coal seams interbedded with clastic deposits. Such seismic facies is relatively  
common for uppermost Devonian–Mississippian sedimentary rocks in the Norwegian Barents Sea  
(Koehl et al., 2018; Tonstad, 2018). In Reindalspasset, uppermost Devonian–Mississippian,  
phyllitic, coal-rich deposits of the Billefjorden Group were penetrated by exploration well 7816/12-

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

370 1 at a depth of 2261 m (Eide et al., 1991), which corresponds to a time of 0.96 s (TWT) when time-converted (Figure 5Figure 4g).

Pennsylvanian–Permian sedimentary strata of the Gipsdalen Group (Figure 2) are mostly composed of packages of subparallel low- to moderate-amplitude seismic reflections separated by discrete, moderate- to high-amplitude reflections. The Hultberget and Ebbadalen formations dominantly show partly disrupted, subparallel reflections possibly representing medium- to fine-grained sedimentary strata (e.g., of the Trikolorfjellet Member) that, in places, alternate with chaotic seismic facies probably characterizing coarse-grained sedimentary deposits (e.g., of the Odellfjellet and/or Ebbaelva members; Johannessen, 1980; Johannessen and Steel, 1992; Braathen et al., 2011; Smyrak-Sikora et al., 2018). The Minkinfjellet and Wordiekammen formations are dominated by a thick package of sub-parallel, moderate- to low-amplitude seismic reflections mostly representing carbonate and gypsum deposits (Figure 5Figure 4). The top reflection of the Wordiekammen Formation is characterized by high amplitude and is relatively easy to trace throughout the study area (Figure 5Figure 4). Finally, the Gipshuken Formation displays chaotic to subhorizontal and subparallel low-amplitude seismic reflections (Figure 5Figure 4). The Wordiekammen and Gipshuken formations are easily identified on seismic data because they crop out at sea level along the northern shore of Sassenfjorden and Tempelfjorden and, hence, can be directly tied to onshore geology (Dallmann et al., 2004, 2009; Dallmann, 2015). Mesozoic sedimentary rocks are not the focus of the present study and were therefore not described.

#### 390 4.2.2. Structures in Sassenfjorden–Tempelfjorden

Seismic data in Sassenfjorden–Tempelfjorden (Figure 1Figure 1a–b) show that basement rocks and overlying, uppermost Devonian–Permian sedimentary rocks of the Billefjorden and Gipsdalen groups are folded into two open, upright, NW–SE- to WNW–ESE-trending fold structures that coincide with similarly trending, several kilometer-wide, elongated ridges representing uplifted portion of the seafloor in Sassenfjorden and Billefjorden (Koehl, 2020; Koehl et al., submitted), and with steeply NNE-dipping, basement-seated faults mostly confined to basement (–Devonian?) rocks and uppermost Devonian–Mississippian coal-rich deposits of the Billefjorden Group, or that die out upwards in the lower part of the Gipsdalen Group (Figure 5Figure 4a). Based on the minor reverse, top-SSW offset of thickened uppermost Devonian–Mississippian sedimentary strata, it is probable that the two gentle fold structures formed in the

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

early Cenozoic as fault-propagation folds due to upward propagation and reverse reactivation/overprinting of NNE-dipping basement-seated faults.

Seismic data in Sassenfjorden and Tempelfjorden also show that high-amplitude seismic reflections characterizing uppermost Devonian–Mississippian sedimentary rocks significantly thicken (approximately twice thicker) towards the south-southwest, near the intersection of the east-dipping Billefjorden Fault Zone with NNE-dipping basement-seated faults, potentially suggesting that uppermost Devonian–Mississippian rocks represent early syn-rift sedimentary deposits (Prosser, 1993) and are part of the initiation stage (Gawthorpe and Leeder, 2000) of the Billefjorden Trough (Figure 5Figure 4a–b). There, high-amplitude seismic reflections representing coal-rich uppermost Devonian–Mississippian strata display laterally disrupted, (SSW-) tilted, Z-shaped geometries (Figure 5Figure 4b and e) that contrast with continuous, subparallel, subhorizontal geometries of the reflections in the northeast (Figure 5Figure 4a, and c–e). Since similar Z-shaped geometries interpreted as duplex structures comprised of bedding-parallel décollements (floor- and roof-thrusts) connected by bedding-oblique link-thrusts were encountered in locally thickened, coal-rich, uppermost Devonian–Mississippian sedimentary deposits in Pyramiden (Figure 4Figure 3b), it is conceivable that, in Sassenfjorden–Tempelfjorden too, significant rheological contrasts between uppermost Devonian–Mississippian coal-coaly shale and sandstone of the Billefjorden Group localized the formation of duplex-related décollements and thrust faults during early Cenozoic deformation.

Locally, moderate- to low-amplitude, subparallel seismic reflections of the Hultberget, Ebbadalen, Minkinfjellet and Wordiekammen formations are disrupted by and slightly bending along moderate to shallow dipping, bedding-oblique reflections, which are interpreted as minor early Cenozoic thrust faults (Figure 5Figure 4a, c, d and f). These minor thrusts appear to flatten downwards and die out within high-amplitude seismic reflections of the Billefjorden Group, thus supporting the presence of bedding-parallel décollements in uppermost Devonian–Mississippian sedimentary rocks (Figure 5Figure 4c).

Seismic reflections within the overlying Gipshuken Formation dip gently to moderately and display continuous to partly chaotic facies (Figure 5Figure 4f). These are disrupted by possible gently NE- to east- and SW- to west-dipping thrusts that seem to flatten downwards into the Top Wordiekammen Formation reflection, forming part of possible imbricate thrust systems (Figure 5Figure 4a, c, d and f) resembling thrusts within coals and coaly shales of the Billefjorden Group

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

(~~Figure 4~~~~Figure 3~~b). This interpretation is supported by onshore Eurekan thrust geometries on the northern shore of Sassenfjorden (supplement 5). This suggests the presence of (a) décollement level(s) within the Wordiekammen Formation and/or at the boundary between the Wordiekammen and Gipshuken formations. Internal seismic packages within the Gipshuken Formation display significant thickness variations, pinching out laterally and, in places, becoming as thick as the whole Gipshuken Formation (~~Figure 5~~~~Figure 4~~a, c, d and f). These thickness variations are tentatively related to tectonic thickening due to early Cenozoic thrusting and, potentially, to the presence of partially mobile evaporite within the Gipshuken Formation (Dallmann et al., 1999).

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

#### 4.2.3. Structures in Reindalspasset

Seismic data in Reindalspasset show a N–S-trending open fold structure (~~Figure 5~~~~Figure 4~~g). In Lower–Middle Devonian rocks, the lowermost part of the fold shows semi-continuous to chaotic, moderate- to low-amplitude, locally undulating seismic reflections that display intensive disruption and wedgesigmoid-shaped geometries (~~Figure 5~~~~Figure 4~~g). Moderate- to low-amplitude reflections within wedgesigmoid-shaped seismic packages display S- and Z-shaped geometries that are disrupted respectively by moderately west- and east-dipping reflections that appear to be responsible for the thickening of internal units and that flatten and die out upwards prior to or at the boundary with overlying uppermost Devonian–Mississippian rocks (~~Figure 5~~~~Figure 4~~g). These wedgesigmoid-shaped seismic packages are interpreted as thickened sheets crosscut by early Cenozoic thrust faults that, in places, form duplex structures comprised of floor- and roof-thrusts connected by link thrusts. Associated undulating reflection geometries are thought to represent folding. Based on the sub-continuous, low- to moderate-amplitude seismic facies and on the presence of folds and bedding-subparallel thrusts, it is probable that (at least the upper part of) this seismic unit is composed of shale-rich, Lower–Middle Devonian sedimentary strata of the Wood Bay and/or Grey Hoek and/or Wijde formations.

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

The core of the fold is partly composed of gently west-dipping to flat-lying, high-amplitude seismic reflections representing coal-rich sedimentary strata of the Billefjorden Group, which were penetrated by exploration well 7816/12-1 at a depth of 2261 m (Eide et al., 1991), i.e., 0.96 s (TWT; ~~Figure 5~~~~Figure 4~~g). In the east, sedimentary strata of the Billefjorden Group can be traced as continuous, gently west-dipping, sub-parallel reflections that thicken westwards against the eastern limb of the fold and that are locally folded and disrupted by a few gently west-dipping, bedding-

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

subparallel reflections that accommodate local thickening of the Billefjorden Group and, hence, may represent minor early Cenozoic thrust faults (Figure 5Figure 4g). High-amplitude reflections of the Billefjorden Group are thickest within the fold hinge, where they show undulating geometries and are intensively disrupted. These disruptions may be the result of early Cenozoic thrusting along low-angle, bedding-subparallel faults, which are probably responsible for the thickening of uppermost Devonian–Mississippian strata within the fold hinge and are possibly forming part of an antiformal stack or ramp anticline (Figure 5Figure 4g). The largest of these potential early Cenozoic thrusts localized along the boundary between uppermost Devonian–Mississippian and Pennsylvanian sedimentary strata, i.e., parallel to the eastern limb of the fold, and splays upwards into four faults. This fault and associated splays quickly die out upwards within the fold hinge in the upper part of the uppermost Devonian–Mississippian and in the lower part of the Pennsylvanian sedimentary succession, offset sediments of the Billefjorden and Gipsdalen groups in a reverse manner (possible repeated portion of the Billefjorden Group), and flatten into the base of the Billefjorden Group or uppermost part of the Lower–Middle Devonian succession (Figure 5Figure 4g). The lowermost splay of this thrust was most likely penetrated by exploration well 7816/12-1 and consists of phyllitic coal and sheared coaly shales of the Billefjorden Group (Eide et al., 1991; Figure 5Figure 4g). Bedding-parallel thrusts in uppermost Devonian–Mississippian strata are further supported by the presence of an analogous, sub-horizontal, bedding-parallel fault within the overlying Middle–Upper Triassic sedimentary rocks of the Barentsøya Formation, which was also penetrated by well 7816/12-1 and represents a possible early Cenozoic décollement (Eide et al., 1991; see uppermost sub-horizontal fault in Figure 5Figure 4g).

Continuous to semi-continuous, parallel, dominantly moderate- to high-amplitude seismic reflections representing Pennsylvanian–lower Permian sedimentary strata of the Hultberget, Ebbadalen, Minkinfjellet and Wordiekammen formations thicken eastwards and westwards away from the fold hinge, i.e., opposite to sedimentary rocks of the Billefjorden Group, and appear to be affected by much fewer disruptions and, therefore, to be only mildly deformed (Figure 5Figure 4g). Pennsylvanian–lower Permian strata are thickest along the eastern fold limb where they are crosscut by three splays of the early Cenozoic thrust localized along the boundary between the Billefjorden and Gipsdalen groups and by a steeply east-dipping brittle fault. This steeply east-dipping fault shows a planar geometry in cross-section, thickening of the Hultberget, Ebbadalen,

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Minkinfjellet and Wordiekammen formations in the hanging wall, minor normal offsets of seismic reflections within these stratigraphic units, and dies out within the lower part of the Wordiekammen Formation upwards and the upper part of the Lower–Middle Devonian succession downwards. Based on cross-section geometries, offset kinematics, and thickening of stratigraphic units, this steeply dipping normal fault is interpreted as a Carboniferous normal fault possibly representing the southwards continuation of the Billefjorden Fault Zone.

## 5. Discussion

### 5.1. Implications of contractional duplexes and décollements in Devonian–Mississippian sedimentary rocks for Ellesmerian and Eurekan deformation

Uppermost Devonian–Mississippian sedimentary rocks of the Billefjorden Group in Pyramiden (Figure 3b) and Sassenfjorden–Tempelfjorden (Figure 4a, b, d and e) are arranged in gently dipping duplexes comprised of interbedded coal–coaly shale and sandstone deposits with sigmoidal shear fabrics and (imbricate) link thrusts (McClay and Insley, 1986) connecting bedding-parallel décollements (roof and floor thrusts/detachments; McClay, 1992) localized along lithological boundaries. These structures and geometries are typical in coal deposits reworked by contractional deformation (Phillipson, 2003, 2005; Molinda, 2003; Elizalde et al., 2016). The interpretation of bedding-parallel décollements is supported by minor early Cenozoic thrusts crosscutting the Hultberget, Ebbadalen, Minkinfjellet and Wordiekammen formations in Sassenfjorden–Tempelfjorden (Figure 1a–b) that flatten downwards and die out within sedimentary strata of the Billefjorden Group (Figure 5c), and by the presence of analogous shallow-dipping, bedding-parallel décollements in uppermost Devonian–Mississippian coals and coaly shales sedimentary strata of the Billefjorden Group in Odellfjellet (Koehl and Muñoz-Barrera, 2018), in Robertsonbreen (between the uppermost Devonian–Mississippian Hørbyebreen Formation and Pennsylvanian–Permian Wordiekammen Formation; Dißmann and Grewing, 1997), in northeastern Bjørnøya (Koehl, in prep.), at Midterhuken and in St. Jonsfjorden (where the unconformity between uppermost Devonian–Mississippian and Pennsylvanian sedimentary rocks possibly acted as a décollement/subhorizontal thrust; Maher and Welbon, 1992; Gasser and Andresen, 2013; Figure 1a), in Nordenskiöld Land (Braathen and Bergh, 1995), and, potentially, in Oscar II Land (Bergh and Andresen, 1990) and Wedel Jarlsberg Land–Torell Land (Dallmann and Maher, 1989; Figure 1a). Imbrication within the duplexes in Pyramiden

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check spelling and grammar

525 indicates top-west thrusting, and most likely reflects Eurekan contraction–transpression since it is the only post-Mississippian episode of contraction–transpression recorded in Spitsbergen. Similar Eurekan duplex geometries with sigmoidal bedding surfaces and link thrusts were also observed in Triassic strata in Spitsbergen (Andresen et al., 1992; Haremo and Andresen, 1992; Andresen, 2009), thus further supporting an interpretation of early Cenozoic thrusting in Pyramiden.

530 In Reindalspasset, potential décollements and low-angle thrusts folded into a gentle upright anticline and possibly forming an antiformal thrust stack were identified on seismic data within Lower–Middle Devonian strata of the Wood Bay and/or Grey Hoek and/or Wijde Bay formations and uppermost Devonian–Mississippian rocks of the Billefjorden Group (Figure 5Figure 4g). In tectonically thickened and mildly folded uppermost Devonian–Mississippian rocks, low-angle brittle–ductile thrust faults are comprised of phyllitic (i.e., sheared) and brittle coals (penetrated by well 7816/12-1 at a depth of 2261–2280 meters; Eide et al., 1991) that are similar to sheared uppermost Devonian–Mississippian coals in Pyramiden, and are arranged into potential duplexes that are comparable to duplexes and thrust systems in uppermost Devonian–Mississippian sedimentary rocks in Pyramiden (Figure 1Figure 1 and Figure 4Figure 3b) and Sassenfjorden–Tempelfjorden (Figure 1Figure 1, and Figure 5Figure 4b and e). The geometries of these duplexes, thrusts and décollements on seismic data in Spitsbergen are similar to analogous structures on seismic data worldwide (e.g., Morley et al., 2017, their figure 8). Potential Lower–Middle Devonian rocks show ~~wedgesigmoid~~-shaped duplex structures, décollements, folding and thrusting comparable to deformation structures in analogous rocks in Andrée Land, e.g., Bråvallafjella Fold Zone (Piepjohn, 2000; Dallmann and Piepjohn, 2020), and in southern Spitsbergen (e.g., Røkensåta; Figure 1Figure 1a; Dallmann, 1992), thus potentially supporting the preservation of Devonian sedimentary rocks of the Andrée Land Group (and/or Mimerdalen Subgroup) east of the Billefjorden Fault Zone in Reindalspasset, pending that the observed normal fault does actually represent the southern continuation of the Billefjorden Fault Zone (Figure 5Figure 4g; see section 5.3). The presence of décollements within Lower–Middle (–lowermost Upper?) Devonian rocks is further supported by the observation of similar structures between shale and sandstone units of the Wood Bay and Grey Hoek formations in Andrée Land (Roy, 2007, 2009; Roy et al., unpublished).

545 Based on the significant differences in deformation styles, it is probable that the décollements and backward-dipping duplexes in sheared uppermost Devonian–Mississippian coals–coaly shales decoupled early Cenozoic Eurekan deformation between folded, shale-rich, 555

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Lower Devonian rocks and undeformed to poorly-deformed uppermost Devonian–Permian sedimentary strata in Pyramiden (Figure 3Figure 2 and Figure 4Figure 3b). Seismic data in Sassenfjorden–Tempelfjorden also show potential duplexes and décollements within uppermost Devonian–Mississippian coal-rich deposits (Figure 5Figure 4a, b, d and e). In these fjords, steeply dipping, basement-seated brittle faults seem to have propagated upwards during early Cenozoic Eureka deformation, resulting in fault-propagation folding and reverse offsets in uppermost Devonian–Permian sedimentary strata (Figure 5Figure 4a and c). These faults die out upwards within uppermost Devonian–Pennsylvanian sedimentary rocks, while minor early Cenozoic thrusts crosscutting Pennsylvanian–Permian sedimentary strata appear to flatten downwards and die out into high-amplitude seismic reflections interpreted as uppermost Devonian–Mississippian coals, thus, also suggesting decoupling of Eureka deformation by early Cenozoic décollements in uppermost Devonian–Mississippian coals of the Billefjorden Group.

In Reindalspasset, early Cenozoic duplexes and thrusts within potential Lower–Middle Devonian strata of the Wood Bay and/or Grey Hoek and/or Widje Bay formations die out upwards and minor thrusts within Pennsylvanian–Permian rocks die out downwards near or at the boundary with coal-rich sedimentary rocks of the Billefjorden Group (Figure 5Figure 4g), thus also supporting the presence of early Cenozoic décollements within uppermost Devonian–Mississippian coaly shales and coals and (partial) decoupling of Eureka deformation. Thickened coal-rich deposits are long known to be able to decouple deformation both in contractional (Frodsham and Gayer, 1999, their figures 1b, 2, 7 and 9) and extensional settings (Wilson and Wojtal, 1986, their figures 7 and 10). In Svalbard, recent field studies by Koehl and Muñoz-Barrera (2018) in the northern part of the Billefjorden Trough in Odellfjellet (Figure 1Figure 1b) showed that bedding-parallel duplex-shaped décollements in uppermost Devonian–Mississippian coaly shales may have partly inhibited early Cenozoic Eureka contraction–transpression in overlying Pennsylvanian strata, thus further supporting the presence of such décollements in Pyramiden (Figure 4Figure 3b), Sassenfjorden–Tempelfjorden (Figure 5Figure 4a–f) and Reindalspasset (Figure 5Figure 4g).

Uppermost Devonian–Mississippian coal-rich strata are locally thicker in Pyramiden, thus resulting in their exploitation by Russia until the early 90s (Livshitz, 1966; Cutbill et al., 1976). They are also thicker in Sassenfjorden in the hanging wall of the east-dipping Billefjorden Fault Zone near the intersection with a NNE-dipping basement-seated fault (Figure 1Figure 1, and Figure

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

~~5~~Figure 4a–d), and within the hinge zone of the anticline adjacent to the possible southward continuation of the Billefjorden Fault Zone in Reindalspasset (~~Figure 5~~Figure 4g). Recent studies of sedimentary rocks of the Billefjorden Group in the Ottar Basin (Tonstad, 2018), the Finnmark Platform (Koehl et al., 2018) in the SW Barents Sea, and the northern part of the Billefjorden Trough (Koehl and Muñoz-Barrera, 2018) show that uppermost Devonian–Mississippian sedimentary strata were deposited into subsiding basins bounded by normal faults. In addition, high-amplitude seismic reflections in the Ottar Basin representing thickened, coal-rich, uppermost Devonian–Mississippian sedimentary strata analog to those observed in Sassenfjorden–Tempelfjorden are thickest on basin edges where fluvial systems dominated in latest Devonian–Mississippian times (Tonstad, 2018). It is possible that, in Spitsbergen too, thick uppermost Devonian–Mississippian coal seams ~~we~~are restricted to the basin edges along boundary faults. ~~This would explain, thus explaining~~ the localization of contractional duplexes and décollements in areas such as Pyramiden, Sassenfjorden, Reindalspasset and (potentially) Triungen during early Cenozoic deformation. ~~These contractional duplexes partially decoupled~~ing deformation between Lower–lowermost Upper Devonian sedimentary rocks of the Andrée Land Group and Mimerdalen Subgroup and thick Pennsylvanian–Permian deposits of the Gipsdalen Group, and locally shielded~~ing~~ the latter from Eurekan deformation, while Pennsylvanian sedimentary rocks in basinal areas in the hanging wall of the Odellfjellet Fault were involved in Eurekan deformation, and Carboniferous normal faults were inverted, e.g., in Odellfjellet (Koehl and Muñoz-Barrera, 2018), Løvehovden–Hultberget (Dallmann, 1993; Maher and Braathen, 2011), Adolfbukta (Harland et al., 1988), Lykteneset (Koehl et al., ~~submitted~~2020), Anservika (Ringset and Andresen, 1988), and Sassenfjorden (~~Figure 5~~Figure 4a–f).

Based on field and seismic data in central Spitsbergen (present study; Koehl and Muñoz-Barrera, 2018; Koehl et al., ~~submitted~~2020) and on analog modelling (Bonini, 2001), it is possible that Lower–lowermost Upper Devonian sedimentary deposits of the Andrée Land Group and Mimerdalen Subgroup were folded exclusively in early Cenozoic times since the differences in deformation style and intensity between Devonian and Carboniferous–Permian deposits can be explained simply by decoupling of Eurekan deformation by weak, uppermost Devonian–Mississippian, coal- and shale-rich sedimentary deposits of the Billefjorden Group (Figs. ~~Figure 4~~Figure 3b, and ~~Figure 5~~Figure 4a–e and g; Koehl and Muñoz-Barrera, 2018). Hence, a short-lived episode of Late Devonian (Ellesmerian) deformation is not required to explain differential

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

620 deformation within Lower Devonian to Permian sedimentary successions in central Spitsbergen, thus potentially simplifying the late Paleozoic tectonic history of the area by reducing it to the Caledonian Orogeny and late–post-Caledonian extensional collapse–rifting. This is further supported by a field study in Robertsonbreen (central Spitsbergen; Figure 1Figure 1b), where Dißmann and Grewing (1997) noticed that sedimentary strata of the lowermost Upper Devonian Plantekløfta Formation and uppermost Devonian–Mississippian Hørbyebreen Formation are both similarly folded, i.e., suggesting that early Cenozoic deformation may be (at least partially) responsible for folding of Lower–lowermost Upper Devonian rocks of the Andrée Land Group and Mimerdalen Subgroup in central Spitsbergen.

630 Strain decoupling, décollements and contractional duplexes are common features in the West Spitsbergen Fold-and-Thrust Belt and were described at various locations and within varied rock types and stratigraphic units. Notably, Ringset and Andresen (1988) and Harland et al. (1988) discussed the presence of subhorizontal, bedding-parallel décollements within Pennsylvanian evaporites of the Ebbadalen and Minkinfjellet formations in eastern Billefjorden, from which early Cenozoic Eurekan thrusts may have ramped upwards into trailing imbricate fans (Boyer and Elliott, 1982) due to lateral lithological variations within Pennsylvanian formations (Ringset and Andresen, 1988). In addition, in western Spitsbergen, Maher (1988), Saalman and Thiedig (2000) and Bergh and Andresen (1990) described early Cenozoic décollements and gently hinterland-dipping duplexes in uppermost Pennsylvanian–Permian sedimentary deposits of the Wordiekammen, Gipshuken and Kapp Starostin formations, which may represent analogs to duplex structures and associated bedding-parallel décollements and low-angle thrusts within uppermost Devonian–Mississippian coals and coaly shales in Pyramiden, Sassenfjorden–Tempelfjorden and Reindalspasset (Figure 4Figure 3b and Figure 5Figure 4). Noteworthy, a model of critical wedge taper for the West Spitsbergen Fold-and-Thrust Belt predicted an increasing influence of decoupling (as observed in Pyramiden, Sassenfjorden–Tempelfjorden and Reindalspasset; Figure 4Figure 3b and Figure 5Figure 4) towards the foreland of the fold and thrust belt, i.e., near the study area in central Spitsbergen (Braathen et al., 1999). All these earlier models and observations are in agreement with the model of strain partitioning and decoupling along bedding-parallel décollements and thrusts proposed by the present study in Pyramiden.

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline

5.2. *Formation mechanism for duplexes and décollements in uppermost Devonian–Mississippian rocks in Pyramiden*

650 Backward-dipping duplexes in Pyramiden are juxtaposed against east-dipping (and locally overturned west-dipping) Devonian strata of the Andrée Land Group and Mimerdalen Subgroup (Figures 2 and 3a and b) adjacent to and showing similar attitude to major fold structures in Mimerdalen thus far ascribed to the Ellesmerian Orogeny (Vogt, 1938; Piepjohn, 2000; Bergh et al., 2011). It is possible that, during early Cenozoic folding, Lower–lowermost Upper Devonian  
655 rocks of the Andrée Land Group and Mimerdalen Subgroup in the west may have acted as a relatively rigid buttress, i.e., partly deforming but not as easily as overlying weak uppermost Devonian–Mississippian coals and coaly shales of the Billefjorden Group that localized the formation of duplexes and décollements, and, thus, allowing these structures to ramp upwards to the west. This is supported by field studies (Fard et al., 2006) and analog modelling (Bahroudi and  
660 Koyi, 2003) in the Zagros Fold-and-Thrust Belt showing buttressing, backward-dipping duplexes and décollements in the hanging wall of deep-seated faults, and by analog modelling of décollements in weak sedimentary layers with limited lateral extent (Costa and Vendeville, 2002, their model 3). Notably, Costa and Vendeville’s model shows that initially sub-horizontal sedimentary strata may have been tilted backwards (i.e., eastwards in Pyramiden) during  
665 contraction, and that décollement lithology (i.e., uppermost Devonian–Mississippian coal–coaly shale) may be incorporated and transported (top-west to top-WNW in Pyramiden; ~~Figure 4~~  
~~Figure 3a~~) as part of the hanging wall sequence during thrusting. In Pyramiden, this is supported by drill data from Trust Arktikugol showing that coal seams of the Billefjorden Group at the mine continue eastwards and preserve a gentle–moderate dip to the east (Aakvik, 1981, his figure 8.2.5). This  
670 interpretation implies the presence of the Balliolbreen Fault in Pyramiden, which is discussed in section 5.3.

Another possibility is that the Pyramiden outcrop represents a mildly inverted extensional fault-block that was gently folded due to upward propagation of the Balliolbreen Fault (if present at all in Pyramiden; see section 5.3) and Odelfjellet Fault (e.g., gentle tilt to the east-southeast of  
675 strata of the Minkinfjellet Formation in Pyramiden; Koehl et al., 2016). Fault-propagation folds (Schlische, 1995) were discussed along the Løvehovden Fault (Maher and Braathen, 2011) and Billefjorden Fault Zone (Braathen et al., 2011; Bælum and Braathen, 2012) in central Spitsbergen. However, this model implies the existence of the Balliolbreen Fault in Pyramiden as a steeply east-

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline,  
Check spelling and grammar

dipping fault, which is not obvious (see section 5.3), and, alone, does not explain the presence of bedding-parallel décollements and backward-dipping duplexes within uppermost Devonian–Mississippian coals and coaly shales of the Billefjorden Group in Pyramiden and Sassenfjorden–Tempelfjorden (Figure 3b, and Figure 4b and e). Moreover, seismic data in Reindalspasset show that a steeply east-dipping normal fault potentially representing the southwards continuation of the Billefjorden Fault Zone (Odellfjellet Fault?) is located along the eastern flank of a broad, gentle anticline (Figure 4g) and, hence, might be related to (or might have interacted with) the fold structure but is most likely not the cause of folding in this area.

Analog modelling of inversion in asymmetric half-graben basins shows features similar to those observed in Pyramiden, demonstrating a potential relationship between weak, early syn-rift sedimentary deposits and segments of basin-bounding faults (Buiter and Pfiffner, 2003, their figure 6a). Notably, in presence of weak, syn-rift sedimentary rocks in basin-edge fault-blocks, newly-formed shortcut shear zones or faults (McClay, 1989) may branch off preexisting inverted basin-bounding normal faults, and ramp up into the weak, syn-rift sedimentary strata, potentially using décollement levels to accommodate contraction. Buiter and Pfiffner (2003) further argue that basement blocks experience much less contraction-related rotation along preexisting normal faults. Thus, a possible scenario for the early Cenozoic tectonic history of the Billefjorden Fault Zone in Pyramiden might involve the formation of a shortcut shear zone or fault along an inverted portion of the Billefjorden Fault Zone at depth, branching off and ramping upwards into, weak, coal- and coaly shale-dominated syn-rift sedimentary rocks of the Billefjorden Group, forming bedding-parallel décollements (Phillipson, 2003, 2005; Molinda, 2003; Elizalde et al., 2016) and east-dipping, backward-dipping duplexes (Figure 3b).

Alternatively, early Cenozoic reverse reactivation/overprinting of the potentially upward-flattening Balliolbreen Fault (if present at all in Pyramiden; see section 5.3) might have triggered the development of a décollement within and of a fault-bend hanging wall anticline (e.g., the Kuqa Fold Belt in northwestern China; Wang et al., 2013; Izquierdo-Llavall et al., 2017) above uppermost Devonian–Mississippian coals, e.g., in Reindalspasset (Figure 4g). In this scenario, backward-dipping duplexes and décollements in uppermost Devonian–Mississippian coals–coaly shales may have acted as a roof décollement decoupling uppermost Devonian–Permian strata from (Lower–lowermost Upper) Devonian rocks, passively thrusting the former over the latter (Bonini, 2001). Through this process, the length of the roof sequence (uppermost Devonian–

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

710 Permian sedimentary strata) remains essentially the same, whereas the length of the floor sequence  
(Lower–lowermost Upper Devonian rocks of the Andrée Land Group and Mimerdalen Subgroup)  
decreases through intense folding (Bonini, 2001). This may (partially) explain the significant  
differences of deformation between folded Lower–lowermost Upper Devonian of the Andrée Land  
Group/Mimerdalen Subgroup (Vogt, 1938; Harland et al., 1974; Piepjohn et al., 1997; Michaelsen  
715 et al., 1997; Michaelsen, 1998; Piepjohn, 2000), strongly sheared uppermost Devonian–  
Mississippian strata of the Billefjorden Group (~~Figure 4~~Figure 3b), and poorly deformed to gently  
tilted uppermost Devonian–Permian strata of the Billefjorden and Gipsdalen groups in central  
Spitsbergen (e.g., Braathen et al., 2011) without a short-lived episode of Ellesmerian contraction  
in the Late Devonian. The lack of uppermost Devonian–Mississippian coals and coaly shales of  
720 the Billefjorden Group directly on top of folded Lower (–lowermost Upper?) Devonian  
sedimentary rocks above the mine entrance in Pyramiden may suggest that uppermost Devonian–  
Mississippian coals–coaly shales were too thin or too localized (syn-rift?) to allow décollements to  
ramp all the way up to the mine entrance or that early Cenozoic Eurekan contraction–transpression  
was too mild to form a complete ramp-anticline (assuming that the Balliolbreen Fault is present in  
725 Pyramiden) with roof décollement over Lower Devonian sedimentary rocks (e.g., Faisal and Dixon,  
2015).

Another plausible interpretation might be that of (a) west-directed imbricate fan(s) in  
Pennsylvanian evaporitic deposits and/or uppermost Devonian–Mississippian coals and coaly  
shales at depth in the Billefjorden Trough with east-dipping imbricate thrusts ramping upwards  
730 into coals and coaly shales of the Billefjorden Group in the footwall of the Odellfjellet Fault, in  
Pyramiden. This interpretation is supported by field studies of Ringset and Andresen (1988) who  
discussed imbricate (thrust) fans and associated basal décollement developed along lithological  
boundaries within the Ebbadalen Formation in Anservika–Gipshuken (see ~~Figure 1~~Figure 1b for  
location), Harland et al. (1988) who described sheared evaporites within the Ebbadalen and  
735 Gipshuken formations in eastern Billefjorden, and by recent field studies showing the presence of  
a potentially gently east-dipping, bedding-parallel thrust–décollement within the Billefjorden  
Group and Hultberget Formation in Anservika (Henningsen et al., pers. comm. 2019), and within  
the Hultberget Formation in Lykteneset (Koehl et al., submitted2020).

Based on field data, backward-dipping duplexes and bedding-parallel décollements in  
740 uppermost Devonian–Mississippian coals and coaly shales of the Billefjorden Group in Pyramiden

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline,  
Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check  
spelling and grammar

are believed to have formed through a combination of at least two or more mechanisms, including Devonian rocks of the André Land Group and Mimerdalen Subgroup acting as a relatively rigid buttress to the west (e.g., [Figure 5](#)[Figure 4g](#)), fault-propagation folding of (a) preexisting fault(s) like the Balliolbreen Fault and/or Odellfjellet Fault (although not very likely), shortcut faulting propagating upwards and westwards from the Billefjorden Fault Zone (e.g., Buitter and Pfiffner, 2003), ramp/fault-bend hanging wall anticline with roof décollement (e.g., Faisal and Dixon, 2015), and imbricate fan with basal décollement in the Billefjorden Trough (e.g., Ringset and Andresen, 1988; Henningsen et al., pers. comm. 2019).

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

### 750 5.3. Geometry and kinematics of the Balliolbreen Fault and implications for Ellesmerian and Eureka deformation events, and Carboniferous normal faulting

Structural field analysis in the gully below the entrance of the Russian coal mine in Pyramiden has shown the presence of a sub-vertical, steeply east-dipping brittle fault tentatively interpreted as the Balliolbreen Fault and comprised of cataclastic fault rock that, half-way to the mine, crosscuts steeply east-dipping, quartzitic, (Lower–lowermost Upper?) Devonian sedimentary rocks involved in a fold structure with bedding locally overturned to the east ([Figure 3](#)[Figure 2](#) and [Figure 4](#)[Figure 3a](#), and supplement 3). Thin section analysis on both sides of this fault (supplement 1) shows cataclased (Lower–lowermost Upper?) Devonian quartzitic sandstone both in the fault footwall and hanging wall, suggesting that there are no basement rocks at this locality, which is supported by geological maps of Harland et al. (1974), Aakvik (1981), Lamar et al. (1986), and geological maps and logs of Trust Arktikugol (1988; Sirotkin, pers. comm. 2019). In addition, the steeply east-dipping fault does not seem to extend upwards into overlying uppermost Devonian–Mississippian clastic deposits above phyllitic coal-rich sedimentary strata. It is possible that the décollements within uppermost Devonian–Mississippian coals–coaly shales represent the upward low-angle continuation of the steeply east-dipping fault, but the structural location of the décollements (almost directly over the fault) would require an abrupt change of geometry of the fault from subvertical to low-angle (c. 30°; [Figure 4](#)[Figure 3b](#)) within a narrow zone, which is unlikely. In addition, fault surfaces and lithological transitions switch from dominant N–S to NNW–SSE strikes and trends in uppermost Devonian–Mississippian coals–coaly shales below the coal-mine entrance ([Figure 3](#)[Figure 2](#) and [Figure 4](#)[Figure 3a](#), and stereonet 3 in [Figure 3](#)[Figure 2](#)) to dominantly WNW–ESE in Lower (–lowermost Upper?) Devonian rocks and uppermost

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Devonian–Mississippian sandstone above the mine entrance ([Figure 3Figure 2](#) and [Figure 4Figure 3a](#), and stereonet 2 in [Figure 3Figure 2](#)), i.e., parallel to most outcrops sections of uppermost Devonian–Mississippian strata in this part of the study area.

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

775 Above the coal mine in Pyramiden, the contact between Lower (–lowermost Upper?) Devonian sedimentary strata and uppermost Devonian–Mississippian sedimentary rocks is not clearly exposed (partly loose blocks) and its nature is relatively speculative. It may be (1) a (folded?) stratigraphic unconformity and/or (2) a bedding-parallel décollement. Based on the internal geometry of bedding surfaces and deformation state of uppermost Devonian–Mississippian sedimentary strata of the Billefjorden Group, which are arranged into contractional, west-verging duplexes separated by low-angle, bedding-parallel décollements ([Figure 4Figure 3b](#)), it is possible that the stratigraphic contact hosts a décollement, e.g., the potential prolongation of one of the décollements within coal- and coaly shale-rich deposits of the Billefjorden Group ([Figure 3Figure 2](#) and [Figure 4Figure 3b](#)). However, uppermost Devonian–Mississippian deposits above the coal mine appear to consist only of clastic deposits and, hence, lack weak coals–coaly shales into which décollements preferentially localize. Thus, the contact between Lower (–lowermost Upper?) Devonian and uppermost Devonian–Mississippian sedimentary rocks above the mine in Pyramiden most likely corresponds to a (folded?) unconformity.

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

790 Even if the décollements within uppermost Devonian–Mississippian coals and shales ([Figure 4Figure 3b](#)) were to represent the upwards continuation of the steeply east-dipping fault ([Figure 3Figure 2](#)), these most likely do not extend into Lower (–lowermost Upper?) Devonian and uppermost Devonian–Mississippian sandstone units above the mine entrance. Based on the similarity between the strike and dip of the steeply east-dipping fault and the trend and dip of (locally overturned) Lower Devonian bedding surfaces in Pyramiden ([Figure 3Figure 2](#) and [Figure 4Figure 3a](#)), it is possible that the steeply east-dipping fault formed as a minor, bedding-parallel (fold-limb parallel) fault related to post-Caledonian gravitational collapse processes and low-angle detachments (e.g., the Woodfjorden detachment in André Land; Roy, 2007, 2009; Roy et al., unpublished; [Figure 1Figure 4a](#)) in Lower–lowermost Upper Devonian sedimentary rocks of the André Land Group and Mimerdalen Subgroup in northern Spitsbergen (e.g., Chorowicz, 1992), or formed as a minor, bedding-parallel Eurekan accommodation thrust (e.g., Cosgrove, 2015) in the early Cenozoic. Since no major fault was identified in Pyramiden it is probable that the Balliolbreen Fault does not crop out or is not present there. This is supported by microstructures

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

along the steeply east-dipping fault in Pyramiden (Figure 3Figure 2), e.g., mild undulose extinction and limited recrystallization and low amounts of displacement along distributed brittle cracks in fault rock (supplement 1), which indicate mild deformation associated with low-grade pressure–temperature conditions (< 280°C; Stipp et al. 2002).

In Reindalspasset, the planar, east-dipping normal fault that offsets Pennsylvanian–lower Permian sedimentary rocks may represent the potential continuation of the basin-bounding Odelfjellet Fault (“BFZ?” in Figure 5Figure 4g), and the Eureka thrust (and associated splays) localized along the boundary between uppermost Devonian–Mississippian and Pennsylvanian sedimentary successions (Figure 5Figure 4g) the continuation of the (inverted?) Balliolbreen Fault. Fault relationships in cross section in Reindalspasset are comparable to what is proposed for the Balliolbreen and Odelfjellet faults in Pyramiden, e.g., possible merging at depth and hundreds of meter- to kilometer-scale lateral spacing between the faults (see previous section), assuming that the Balliolbreen Fault is present in Pyramiden. The preservation of Lower–lowermost Upper Devonian sedimentary rocks of the Andrée Land Group and/or Mimerdalen Subgroup east of the Billefjorden Fault Zone in Reindalspasset suggests that this fault did not accommodate top-west reverse movement in Late Devonian times as proposed by previous works in Dickson Land (Vogt, 1938; Friend, 1961; Piepjohn, 2000; Dallmann and Piepjohn, 2020). Would such movements have occurred, Devonian sedimentary rocks of the Andrée Land Group and/or Mimerdalen Subgroup in the upthrust block east of the Billefjorden Fault Zone would have been exposed and subjected to continental erosion. This is clearly not the case in Reindalspasset where potential Lower–lowermost Upper Devonian sedimentary rock units appear to thicken eastwards. The presence of Devonian sedimentary east of the Billefjorden Fault Zone in Reindalspasset is also supported by the interpretation of well bore data in the Raddedalen-1 well in Edgeøya (Harland and Kelly, 1997).

~~Based on field data in Pyramiden and seismic data in Sassenfjorden and Reindalspasset, and on previous work (Harland et al., 1974; Lamar et al., 1982, 1986; McCann, 1993; Lamar and Douglass, 1995),~~ The Balliolbreen Fault displays significant along-strike variations in geometry and kinematics. In the north, in Odelfjellet and Sentinelfjellet (Figure 1Figure 1b), the Balliolbreen Fault dips c. 60–65° to the east and juxtaposes Precambrian basement unconformably overlain by uppermost Devonian–Mississippian strata of the Billefjorden Group in the hanging wall against Lower Devonian strata of the Wood Bay Formation supposedly unconformably overlain by uppermost Devonian–Mississippian rocks of the Billefjorden Group (Harland et al., 1974; Lamar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

et al., 1986; Lamar and Douglass, 1995). Both in Odellfjellet and Sentinelfjellet, it is unclear  
835 whether the Balliolbreen Fault offsets uppermost Devonian–Mississippian strata, or if the fault is  
unconformably overlain by uppermost Devonian–Mississippian rocks (Lamar et al., 1982, 1986;  
Lamar and Douglass, 1995). Although Harland et al. (1974) argue that the Triungen Member of  
the Hørbyebeen Formation is unfaulted in Sentinelfjellet (thus potentially supporting Late  
Devonian top-west thrusting along the Balliolbreen Fault and no further reactivation), stratigraphic  
840 contacts in this area are covered by screens and poorly–not exposed (like in Triungen in [Figure  
~~4~~Figure 3c–e](#)) and inaccessible because located on very steep slopes–cliffs (see  
toposvalbard.npolar.no). The presence of newly evidenced décollements in the lower part of the  
Billefjorden Group in Pyramiden, Sassenfjorden–Tempelfjorden and Reindalspasset ([Figure  
~~4~~Figure 3b](#) and [Figure 5Figure 4](#)) suggests that the nature of the contact of the Billefjorden Group  
845 with underlying Devonian rock units must be interpreted with care, especially were covered by  
screens. If sedimentary strata of the Billefjorden Group are actually truncated by the Balliolbreen  
Fault in Sentinelfjellet (e.g., McCann, 1993, his figures 5.9 and 5.10), then early Cenozoic thrusting  
may, in conjunction with Carboniferous normal faulting, explain the observed juxtaposition of  
Precambrian basement and Lower Devonian sedimentary rocks ([Figure 6Figure 5](#)). In this scenario,  
850 basement rocks constituting the Caledonian Atomfjella Antiform were located close to the surface  
at the end of the Caledonian Orogeny, thus leaving no (or limited) accommodation space east of  
the Billefjorden Fault Zone in Ny-Friesland ([Figure 1Figure 1a](#)) during Devonian sedimentation  
sourced from the collapsing orogen and exhuming core complexes [in the west and east](#) (e.g.,  
Bockfjorden Anticline; Braathen et al., 2018; [Figure 6Figure 5a](#)). In the Carboniferous, normal  
855 faulting and footwall rotation along the Odellfjellet Fault possibly exhumed [or kept](#) a small portion  
of basement rocks [relatively close to the surface](#) in the footwall of the fault ([Figure 6Figure 5b–c](#));  
[and Later on](#), subsequent early Cenozoic deformation may have thrust part of the exposed  
basement rocks in the footwall as a kilometer-scale lens along a possibly inverted Carboniferous  
normal fault, the Balliolbreen Fault ([Figure 6Figure 5d](#)) potentially leading to extensive  
860 deformation of Lower–lowermost Upper Devonian rocks in Dickson Land, which acted as a  
buttress absorbing most of Eurekan deformation together with sheared uppermost Devonian–  
Mississippian coals and shales of the Billefjorden Group (note that deformation within Lower–  
lowermost Upper Devonian rocks is not detailed in [Figure 6Figure 5d](#)). In this model,  
Carboniferous normal and early Cenozoic reverse offsets along the Balliolbreen Fault have similar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold

**Formatted:** Font: Not Bold, Not Italic

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

865 magnitude, as shown in Mumien (juxtaposition of the Ebbadalen Formation and Billefjorden Group  
against the Wordiekammen Formation and the Billefjorden Group with no apparent offset at top  
Billefjorden Group level; Dallmann et al., 2004; Dallmann, 2015; [Figure 1Figure 1b](#)), and in  
Sentinelfjellet and Odellfjellet (top of Billefjorden Group offset by 0–40 m; Harland et al., 1974;  
Lamar et al., 1986; [Figure 6Figure 5e](#)). Thus, it is possible that the above mentioned localities  
870 reflect different structural levels of the same fault system ([Figure 6Figure 5e](#)). Eurekan inversion  
of Carboniferous normal faults in central Spitsbergen is also supported by reverse offset and thrust-  
related folding along the Overgangshytta fault in Odellfjellet (Koehl and Muñoz-Barrera, 2018),  
and by minor reverse offset of thickened, uppermost Devonian–Mississippian and Pennsylvanian  
sedimentary deposits in the hanging wall of the east-dipping Billefjorden Fault Zone, near the  
875 intersection with a steeply NNE-dipping basement-seated fault in Sassenfjorden ([Figure 5Figure  
4b](#)).

The high degree of uncertainty in the relationship (truncated or truncating) between the  
Balliolbreen Fault and uppermost Devonian–Mississippian sedimentary strata of the Billefjorden  
Group (especially in Odellfjellet and Sentinelfjellet; Harland et al., 1974; Lamar et al., 1986; Lamar  
and Douglass, 1995), and the uncertainty regarding the nature of the contact (unconformity or  
880 bedding-parallel décollements–thrusts) between Lower Devonian and uppermost Devonian–  
Mississippian sedimentary strata shed by the presence of bedding-parallel Eurekan décollements  
and thrusts in Pyramiden ([Figure 4Figure 3b](#)), Sassenfjorden–Tempelfjorden ([Figure 5Figure 4b, c  
and e](#)) and Reindalspasset ([Figure 5Figure 4g](#)) call for caution and further (re-) examination of  
885 outcrops of uppermost Devonian–Mississippian rocks along the Balliolbreen Fault in central  
Spitsbergen.

-Notably, the significant along strike differences in cross-section geometry from  
subvertical, e.g., in Pyramiden (if present at all; [Figure 3Figure 2](#) and [Figure 4Figure 3b](#)) to shallow  
dipping, e.g., in Reindalspasset (Eurekan thrust localized along the Billefjorden–Gipsdalen groups  
890 boundary; [Figure 5Figure 4g](#)), together with the strong contrasts in offset stratigraphic units, e.g.,  
Pennsylvanian rocks of the Ebbadalen Formation overlain by carbonates of the Wordiekammen  
Formation in the hanging wall against Lower Devonian rocks of the Wood Bay Formation  
unconformably overlain by strata of the Wordiekammen Formation in the footwall in  
Ygdrasilkampen (Dallmann et al., 2004; [Figure 1Figure 1b](#)), Pennsylvanian Ebbadalen Formation  
895 against uppermost Pennsylvanian–lower Permian Wordiekammen Formation in Mumien

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, No underline, Do not check spelling or grammar

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Not Italic

Formatted: Font: Not Bold, No underline

Formatted: Font: Not Bold, Not Italic, No underline, Check spelling and grammar

Formatted: Font: Not Bold, Not Italic, No underline

Formatted: Font: Not Bold, No underline, Check spelling and grammar

(Dallmann et al., 2004; Dallmann, 2015), Lower Devonian rocks overlain by uppermost Devonian–Mississippian Billefjorden Group in the hanging wall against Lower Devonian rocks in the footwall in Pyramiden (if present at all; ~~Figure 3~~~~Figure 2~~ and ~~Figure 4~~~~Figure 3a~~), Precambrian basement rocks in the hanging wall against Lower Devonian rocks in the footwall in Odellfjellet and Sentinelfjellet (Harland et al., 1974; Lamar et al., 1986), and in inferred timing and kinematics, e.g., Carboniferous normal faulting in Yggdrasilkampen (Dallmann et al., 2004), early Cenozoic reverse movement in Pyramiden (if present at all; ~~Figure 4~~~~Figure 3b~~) and possibly in Reindalspasset (if present at all; ~~Figure 5~~~~Figure 4g~~) and Flowerdalen (Harland et al., 1974; Haremo et al., 1990; Haremo and Andresen, 1992), Carboniferous normal and early Cenozoic reverse faulting in Mumien (Dallmann, 2015), and potential Late Devonian (e.g., Harland et al., 1974; Piepjohn, 2000; Dallmann and Piepjohn, 2020) or early Cenozoic thrusting (this study; Koehl and Muñoz-Barrera, 2018) in Odellfjellet and Sentinelfjellet, suggest that the Balliolbreen Fault might consist of several, discrete, disconnected (soft-linked and/or stepping?) or possibly offset fault segments crosscut by suborthogonal faults (McCann, 1993, his figure 5.11; Koehl, 2020).

For example, a basement-seated reverse fault in Sassenfjorden–Tempelfjorden accommodated top-SSW thrusting during Eurekan deformation (~~Figure 5~~~~Figure 4a–b~~) and seem to have limited the amount of Eurekan reactivation/overprinting (strain partitioning) along east-dipping segments of the Billefjorden Fault Zone in this area, which shows mainly down-east Carboniferous normal offset with limited amount of early Cenozoic reworking along the main east-dipping fault (e.g., ~~Figure 5~~~~Figure 4d~~), and may be responsible for restricting sediment deposition/preservation to the southwest of Sassenfjorden during Eurekan tectonism in the early Cenozoic, thus, explaining sediment provenance from the northeast (e.g., Petersen et al., 2016). Another example where strain partitioning may have occurred along suborthogonal faults is Yggdrasilkampen, where the possible continuation of the Balliolbreen Fault juxtaposes Pennsylvanian (hanging wall) against Lower Devonian (footwall) sedimentary rocks suggesting that Carboniferous normal faulting was followed by limited early Cenozoic reactivation/overprinting if any at all. The character of the Billefjorden Fault Zone in Sassenfjorden and Yggdrasilkampen contrasts sharply with areas farther north (e.g., in Sentinelfjellet and Odellfjellet; Harland et al., 1974; Lamar et al., 1986; Lamar and Douglass, 1995; Dallmann et al., 2004; Dallmann, 2015) and farther south (e.g., in Flowerdalen; Harland et al., 1974; Haremo et al., 1990; Haremo and Andresen, 1992; ~~Figure 1~~~~Figure 4b~~) where the east-dipping Billefjorden Fault

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold

**Formatted:** Font: Not Bold, Not Italic

**Formatted:** Font: Not Bold, No underline, Do not check spelling or grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, No underline

**Formatted:** Font: Not Bold, Not Italic, No underline, Check spelling and grammar

**Formatted:** Font: Not Bold, Not Italic, No underline

**Formatted:** Font: Not Bold, No underline, Check spelling and grammar

Zone displays clear evidence of top-west Eurekan movements. More of these (WNW–ESE-striking) suborthogonal faults are described and their impact on Eurekan strain partitioning further discussed in Koehl et al. ([submitted2020](#)).

930

## 6. Conclusion

1. Thickened uppermost Devonian–Mississippian sedimentary deposits of the Billefjorden Group in central Spitsbergen are arranged in duplexes comprised of phyllitic coal–coaly shale interbedded with sandstone showing sigmoidal shear fabrics separated by imbricate thrusts linking an upper (roof thrust) and a lower (floor thrust) décollements that localized along lithological transitions.
2. Early Cenozoic bedding-parallel décollements and thrusts in tectonically thickened, coal-rich sedimentary rocks of the Billefjorden Group, in the Wordiekammen Formation, and in Lower–lowermost Upper Devonian sedimentary rocks partially decoupled Eurekan deformation, resulting in intense folding in Devonian sedimentary rocks of the Andrée Land Group and Mimerdalen Subgroup and uppermost Devonian–Mississippian coals of the Billefjorden Group, and mild to no deformation in Carboniferous–Permian strata in central Spitsbergen, thus suggesting that Late Devonian Ellesmerian contraction is not required to explain differential deformation within Lower Devonian to Permian sedimentary successions in central Spitsbergen.
3. Early Cenozoic backward-dipping duplexes and bedding-parallel décollements in the Billefjorden Group in Pyramiden formed through shortcut faulting propagating upwards and westwards from the Odellfjellet Fault, and/or as roof décollements of a ramp/fault-bend hanging wall anticline, and/or as part of an imbricate fan with basal décollement in the Billefjorden Trough. Early Cenozoic contractional structures in uppermost Devonian–Mississippian coals–coaly shales also include fault-propagation folds over preexisting basement-seated faults in Sassenfjorden, and a possible antiformal thrust stack (or ramp anticline) in Reindalspasset.
4. Lower–lowermost Upper Devonian sedimentary rocks might be preserved east of the Billefjorden Fault Zone in Reindalspasset, thus suggesting that the Billefjorden Fault Zone did not act as a reverse fault in Late Devonian times.

955

5. Thrusting of Proterozoic basement rocks over Lower Devonian sedimentary rocks along the Balliolbreen Fault and fold structures within strata of the Andrée Land Group and Mimerdalen Subgroup in central Spitsbergen may be explained by a combination of down-east Carboniferous normal faulting with associated footwall rotation and exhumation, and subsequent top-west early Cenozoic Eurekan thrusting along the Billefjorden Fault Zone
- 960
6. The uncertain relationship of the Balliolbreen Fault with uppermost Devonian–Mississippian sedimentary strata and the poorly constrained nature of the contact (unconformity or bedding-parallel décollements–thrusts) between Devonian of the Andrée Land Group and Mimerdalen Subgroup and uppermost Devonian–Mississippian sedimentary strata, as well as significant along strike variations in cross-section geometry, offset stratigraphy, and inferred timing and kinematics suggest that the Balliolbreen Fault consists of several, discrete, unconnected (soft-linked and/or stepping) or most likely offset
- 965
- 970 fault segments that were reactivated/overprinted with varying degree during Eurekan deformation due to strain partitioning along suborthogonal Eurekan reverse faults.

#### **Data availability**

High-resolution versions of the figures and supplements of the manuscript necessary to identify individual reflections and structures can be found at DataverseNO (<https://doi.org/10.18710/MXKQPE>).

975

#### **Competing interests**

The author declares that he has no conflict of interest.

980

#### **Acknowledgements**

The present study is part of the ARCEX (Research Centre for Arctic Petroleum Exploration), which is funded by the Research Council of Norway (grant number 228107) together with 10 academic and eight industry partners, and of the SEAMSTRESS project supported by a starting grant of the Tromsø Research Foundation and the Research Council of Norway (grant number 287865). The author would like to thank all the persons from these institutions that are involved in this project. Most grateful thanks to Prof. John Marshall (University of Southampton), Dr. Christopher Berry (University of Cardiff), Dr. Gilda Lopes (University of Algarve), and Prof.

985

Gunn Mangerud (University of Bergen) for their help with the palynology of Devonian–  
990 Mississippian rocks in Spitsbergen. Many thanks to Alexander Andreas for field collaborations, to  
Prof. Steffen Bergh, Dr. Winfried Dallmann, Prof. Holger Stunitz and Dr. Mélanie Forien  
(University of Tromsø), and Dr. Karsten Piepjohn (BGR) and Assoc. Prof. Jaroslaw Majka  
(Uppsala University) for fruitful discussions. Sebastian Sikora and the University Centre in  
Svalbard (UNIS) are thanked for boat-transportation to Pyramiden in summer 2016. The  
995 Norwegian Petroleum Directorate, Equinor and Store Norske Spitsbergen Kulkompani are thanked  
for granting access to seismic and well data. Many thanks to Ivar Stokkeland from the Norwegian  
Polar Institute Library in Tromsø, Norway, for his help with finding old, non-digitized publications  
about the geology of Svalbard (available at the Norwegian Polar Institute Library in Tromsø,  
Norway; list of publications included in Appendix A). The Ph.D. Thesis of John G. Gjelberg (1984)  
1000 was also digitized and, thanks to the University of Bergen and to John G. Gjelberg’s family, is now  
available from the University of Bergen Library at <http://bora.uib.no/handle/1956/20981>.

## References

- Aakvik, R.: Fasies analyse av Undre Karbonske kullførende sedimenter, Billefjorden, Spitsbergen,  
1005 Ph.D. Thesis, University of Bergen, Bergen, Norway, 219 pp., 1981.
- Allen, K. C.: Lower and Middle Devonian spores of north and central Vestspitsbergen,  
Palaeontology, 8, 678–748, 1965.
- Allen, K. C.: Further information on the Lower and Middle Devonian spores from Dickson Land,  
Spitsbergen, Norsk Polarinstitutt Årbok 1971, 7–15, 1973.
- 1010 Andresen, A.: Geology of Svalbard – A Window into the Barents Sea Hydrocarbon Province,  
Svalex 2009 cruise report, Statoil, 2009.
- Andresen, A., Haremo, P., Swensson, E. and Bergh, S. G.: Structural geology around the southern  
termination of the Lomfjorden Fault Complex, Agardhdalen, east Spitsbergen, Norsk Geol.  
Tidsskr., 72, 83–91, 1992.
- 1015 Andresen, A., Bergh, S. G. and Haremo, P.: Basin inversion and thin-skinned deformation  
associated with the Tertiary transpressional west Spitsbergen Orogen, in: Proceedings of  
the International Conference on Arctic Margins, edited by: Thurston, D. K. and Fujita, K.,  
Anchorage, Alaska, USA, September 1992, 161–166, 1994.

- 1020 Bahroudi, A. and Koyi, H. A.: Effect of spatial distribution of Hormuz salt on deformation style in the Zagros fold and thrust belt: an analogue modelling approach, *J. Geol. Soc. London*, 160, 719–733, 2003.
- Balashov, Yu. A., Larionov, A. N., Gannibal, L. F., Sirotkin, A. N., Tebenkov, A. M., Ryüngen, G. I. and Ohta, Y.: An Early Proterozoic U–Pb zircon age from an Eskolabreen Formation gneiss in southern Ny Friesland, Spitsbergen, *Polar Res.*, 12:2, 147–152, 1993.
- 1025 Beauchamp, B., Alonso-Torres, D., Piepjohn, K., Thériault, P. and Grasby, S. E.: Early Carboniferous syn-rift sedimentation in the Sverdrup Basin (Yelverton Pass area, northern Ellesmere Island, Arctic Canada): A solution to the Okse Bay problem, in: *Circum-Arctic Structural Events: Tectonic Evolution of the Arctic Margins and Trans-Arctic Links with Adjacent Orogens*, edited by: Piepjohn, K., Strauss, J. V., Reinhardt, L. and McClelland, W. C., *GSA Spec. Paper*, 541, 255–284, 2018.
- 1030 Bergh, S. G. and Andresen, A.: Structural development of the Tertiary fold-and-thrust belt in east Oscar II Land, Spitsbergen, *Polar Res.*, 8, 217–236, 1990.
- Bergh, S. G. and Grogan, P.: Tertiary structure of the Sørkapp–Hornsund Region, South Spitsbergen, and implications for the offshore southern continuation of the fold-thrust Belt, *Norsk Geol. Tidsskr.*, 83, 43–60, 2003.
- 1035 Bergh, S. G., Maher Jr., H. D. and Braathen, A.: Tertiary divergent thrust directions from partitioned transpression, Brøggerhalvøya, Spitsbergen, *Norsk Geol. Tidsskr.*, 80, 63–82, 2000.
- Bergh, S. G., Maher Jr., H. D. and Braathen, A.: Late Devonian transpressional tectonics in Spitsbergen, Svalbard, and implications for basement uplift of the Sørkapp–Hornsund High, *J. Geol. Soc. London*, 168, 441–456, 2011.
- 1040 Berry, C. M. and Marshall, J. E. A.: Lycopsid forests in the early Late Devonian paleoequatorial zone of Svalbard, *Geology*, 43, 1043–1046, 2015.
- Birkenmayer, K. and Turnau, E.: Lower Carboniferous age of the so-called Wijde Bay Series in Hornsund, Vestspitsbergen, *Nor. Polarinst. Årb.* 1961, 41–61, 1962.
- 1045 Blinova, M., Faleide, J. I., Gabrielsen, R. H. and Mjelde, R.: Seafloor expression and shallow structure of a fold-and-thrust system, Isfjorden, west Spitsbergen, *Polar Res.*, 31, 11209, 2012.

- Bonini, M.: Passive roof thrusting and forelandward fold propagation in scaled brittle–ductile  
1050 physical models of thrust wedges, *J. of Geophys. Res.*, 106, 2291–2311, 2001.
- Boyer, S. E. and Elliott, D.: Thrust Systems, *AAPG Bulletin*, 66, 9, 1196–1230, 1982.
- Braathen, A. and Bergh, S. G.: Kinematics of Tertiary deformation in the basement-involved fold-  
thrust complex, western Nordenskiöld Land, Svalbard: tectonic implications based on fault-  
slip data analysis, *Tectonophys.*, 249, 1–29, 1995.
- 1055 Braathen, A., Bergh, S. G. and Maher Jr., H. D.: Application of a critical wedge taper model to the  
Tertiary transpressional fold-thrust belt on Spitsbergen, Svalbard, *GSA Bull.*, 111, 10,  
1468–1485, 1999.
- Braathen, A., Bælum, K., Maher Jr., H. D. and Buckley, S. J.: Growth of extensional faults and  
folds during deposition of an evaporite-dominated half-graben basin; the Carboniferous  
1060 Billefjorden Trough, Svalbard, *Norsk Geol. Tidsskr.*, 91, 137–160, 2011.
- Braathen, A., Osmundsen, P. T., Maher Jr., H. D. and Ganerød, M.: The Keisarhjelmen detachment  
records Silurian–Devonian extensional collapse in Northern Svalbard, *Terra Nova*, 30, 34–  
39, 2018.
- Brinkmann, L.: Geologie des östlichen zentralen Dickson Landes und Palynologie der Mimerdalen  
1065 Formation (Devon), Spitzbergen. (Geology of eastern–central Dickson Land and  
palynology of the Mimerdalen Formation [Devonian], Spitsbergen.), unpublished Master’s  
Thesis, University of Münster, Münster, Germany, 94 pp., 1997.
- Bugge, T., Mangerud, G., Elvebakk, G., Mørk, A., Nilsson, I., Fanavoll, S. and Vigran, J. O.: The  
Upper Palaeozoic succession on the Finnmark Platform, Barents Sea, *Norsk Geol. Tidsskr.*,  
1070 75, 3–30, 1995.
- Buiter, S. J. H. and Pfiffner, O. A.: Numerical models of the inversion of half-graben basins,  
*Tectonics*, 5, 1057, doi:10.1029/2002TC001417, 2003.
- Bælum, K. and Braathen, A.: Alongstrike changes in fault array and rift basin geometry of the  
Carboniferous Billefjorden Trough, Svalbard, Norway, *Tectonophys.*, 546–547, 38–55,  
1075 2012.
- Chalmers, J. A. and Pulvertaft, T. C. R.: Development of the continental margins of the Labrador  
Sea: a review, in: *Non-Volcanic Rifting of Continental Margins: A Comparison of  
Evidence from Land and Sea*, edited by: Wilson, R. C. L., Taylor, R. B. and Froitzheim,  
N., *Geol. Soc. London, Spec. Publi.*, 187, 77–105, 2001.

- 1080 Chorowicz, J.: Gravity-induced detachment of Devonian basin sediments in northern Svalbard, Norsk Geol. Tidsskr., 72, 21–25 1992.
- Cosgrove, J. W.: The association of folds and fractures and the link between folding, fracturing and fluid flow during the evolution of a fold–thrust belt: a brief review, in: Industrial Structural Geology: Principles, Techniques and Integration, edited by: Richards, F. L.,  
1085 Richardson, N. J., Rippington, S. J., Wilson, R. W. and Bond, C. E., Geol. Soc. London, Spec. Publi., 421, 41–68, 2015.
- Costa, E. and Vendeville, B. C.: Experimental insights on the geometry and kinematics of fold-and-thrust belts above weak, viscous evaporitic décollement, Journal of Structural Geology, 24, 1729–1739, 2002.
- 1090 Cutbill, J. L. and Challinor, A.: Revision of the Stratigraphical Scheme for the Carboniferous and Permian of Spitsbergen and Bjørnøya, Geol. Mag., 102, 418–439, 1965.
- Cutbill, J. L., Henderson, W. G. and Wright, N. J. R.: The Billefjorden Group (Early Carboniferous) of central Spitsbergen, Norsk Polarinst. Skr., 164, 57–89, 1976.
- Dallmann, W. K.: Multiphase tectonic evolution of the Sørkapp–Hornsund mobile zone (Devonian, Carboniferous, Tertiary), Svalbard, Norsk Geol. Tidsskr., 72, 49–66, 1992.
- 1095 Dallmann, W. K.: Notes on the stratigraphy, extent and tectonic implications of the Minkinfjellet Basin, Middle Carboniferous of central Spitsbergen, Polar Res., 12:2, 153–160, 1993.
- Dallmann, W. K.: Geoscience Atlas of Svalbard, Norsk Polarinstitut, Tromsø, Norway, Rapportserie nr. 148, 2015.
- 1100 Dallmann, W. K. and Maher Jr., H. D.: The Supanberget area – basement imbrication and detached foreland thrusting in the Tertiary fold-and-thrust belt, Svalbard, Polar Res., 7:2, 95–107, 1989.
- Dallmann, W. K. and Piepjohn, K.: The structure of the Old Red Sandstone and the Svalbardian Orogenic Event (Ellesmerian Orogeny) in Svalbard, Norg. Geol. Unders. B., 15, 106 pp.,  
1105 2020.
- Dallmann, W. K., Ohta, Y. and Andresen, A.: Tertiary Tectonics of Svalbard, Norsk Polarinstitut Rapportserie, 46, 112 pp., 1988.
- Dallmann, W. K., Andresen, A., Bergh, S. G., Maher Jr., H. D. and Ohta, Y.: Tertiary fold-and-thrust belt of Spitsbergen Svalbard, Norsk Polarinstitut Meddelelser 128, 51 pp., 1993.

- 1110 Dallmann, W. K., Dypvik, H., Gjelberg, J. G., Harland, W. B., Johannessen, E. P., Keilen, H. B., Larssen, G. B., Lønøy, A., Midbøe, P. S., Mørk, A., Nagy, J., Nilsson, I., Nøttvedt, A., Olaussen, S., Pcelina, T. M., Steel, R. J. and Worsley, D.: Lithostratigraphic Lexicon of Svalbard, edited by: Dallmann, W. K., Norwegian Polar Institute, Polar Environmental Centre, Tromsø, Norway, 1999.
- 1115 Dallmann, W. K., Piepjohn, K. and Blomeier, D.: Geological map of Billefjorden, Central Spitsbergen, Svalbard, with geological excursion guide, Scale 1 : 50,000, Temakart No. 36, Norsk Polarinstitutt, 2004.
- Davies, G. R. and Nassichuck, W. W.: An Early Carboniferous (Visean) Lacustrine Oil Shale in Canadian Arctic Archipelago, AAPG Bulletin, 72, 8–20, 1988.
- 1120 Dißmann, B. and Grewing, A.: Post-svalbardische kompressive Strukturen im westlichen Dickson Land (Hugindalen), Zentral-Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 235–242, 1997.
- Eide, J. R., Ree, R. and Rockman, P. O.: Final well report – 7816/12-1 July 1991, Norsk Hydro A.S., Harstad, Norway, 1991.
- 1125 Elizalde, C., Griffith, W. A. and Miller, T.: Thrust fault nucleation due to heterogeneous bedding plane slip: evidence from an Ohio coal mine, Engineering Geology, 206, 1–17, 2016.
- Faisal, S. and Dixon, J. M.: Physical analog (centrifuge) model investigation of contrasting structural styles in the Salt Range and Potwar Plateau, northern Pakistan, Journal of Structural Geology, 77, 277–292, 2015.
- 1130 Fard, I. A., Braathen, A., Mokhtari, M. and Alavi, S. A.: Interaction of the Zagros Fold–Thrust Belt and the Arabian-type, deep-seated folds in the Abadan Plain and the Dezful Embayment, SW Iran, Petroleum Geoscience, 12, 347–362, 2006.
- Fedorowski, J.: Coral thanatocoenoses and depositional environment in the upper Treskelodden beds of the Hornsund area, Spitsbergen, Palaeontologica Polonica, 43, 17–68, 1982.
- 1135 Flood, B., Gee, D. G., Hjelle, A., Siggerud, T. and Winsnes, T. S.: The geology of Nordaustlandet, northern and central parts, Norsk Polarinst. Skr., 146, 145 pp., 1969.
- Friend, P. F.: The Devonian stratigraphy of north and central Vestspitsbergen, Proceedings of the Yorkshire Geological Society, 33, 77–118, 1961.

- 1140 Friend, P. F. and Moody-Stuart, M.: Sedimentation of the Wood Bay Formation (Devonian) of Spitsbergen: Regional analysis of a late orogenic basin, *Norsk Polarinst. Skr.*, 157, 80 pp., 1972.
- Friend, P. F., Heintz, N. and Moody-Stuart, M.: New unit terms for the Devonian of Spitsbergen and a new stratigraphical scheme for the Wood Bay Formation, *Polarinst. Årbok*, 1965, 59–64, 1966.
- 1145 Friend, P. F., Harland, W. B., Rogers, D. A., Snape, I. and Thornley, R. S.: Late Silurian and Early Devonian stratigraphy and probable strike-slip tectonics in northwestern Spitsbergen, *Geol. Mag.*, 134, 4, 459–479, 1997.
- Frodsham, K. and Gayer, R. A.: The impact of tectonic deformation upon coal seams in the South Wales coalfield, UK, *Int. J. Coal Geol.*, 38, 297–332, 1999.
- 1150 Gasser, D. and Andresen, A.: Caledonian terrane amalgamation of Svalbard: detrital zircon provenance of Mesoproterozoic to Carboniferous strata from Oscar II Land, western Spitsbergen, *Geol. Mag.*, 150, 6, 1103–1126, 2013.
- Gawthorpe, R. L. and Leeder, M. R.: Tectono-sedimentary evolution of active extensional basins, *Basin Res.*, 12, 195–218, 2000.
- 1155 Gayer, R. A., Gee, D. G., Harland, W. B., Miller, J. A., Spall, H. R., Wallis, R. H. and Winsnes, T. S.: Radiometric age determinations on rocks from Spitsbergen, *Norsk Polarinstitutt Skrifter*, 137, 43 pp., 1966.
- Gee, D. G. and Moody-Stuart, M.: The base of the Old Red Sandstone in central north Haakon VII Land, Vestspitsbergen, *Polarinst. Årbok*, 1964, 57–68, 1966.
- 1160 Gee, D. G. and Page, L. M.: Caledonian terrane assembly on Svalbard: New evidence from  $^{40}\text{Ar}/^{39}\text{Ar}$  dating in Ny Friesland, *American Journal of Science*, 294, 1166–1186, 1994.
- Gjelberg, J. G.: Upper Devonian (Famennian) – Middle Carboniferous succession of Bjørnøya, a study of ancient alluvial and coastal marine sedimentation, *Norsk Polarinst. Skr.*, 174, 67 pp., 1981.
- 1165 Gjelberg, J. G.: Early–Middle Carboniferous sedimentation on Svalbard. A study of ancient alluvial and coastal marine sedimentation in rift- and strike-slip basins, Ph.D. Thesis, University of Bergen, Bergen, Norway, 306 pp., 1984.
- Gjelberg, J. G. and Steel, R. J.: An outline of Lower–Middle Carboniferous sedimentation on Svalbard: Effects of tectonic, climatic and sea level changes in rift basin sequences, in:

- 1170 Geology of the North Atlantic Borderlands, edited by: Kerr, J.W. and Ferguson, A. J., Can. Soc. Of Petrol. Geol. Mem., 7, 543–561, 1981.
- Hamilton, E., Harland, W. B. and Miller, J. A.: Isotopic Ages from Some Spitsbergen Rocks, Nature, 195, 1191–1192, 1962.
- Haremo, P. and Andresen, A.: Tertiary décollements thrusting and inversion structures along Billefjorden and Lomfjorden Fault Zones, East Central Spitsbergen, in: Structural and Tectonic Modelling and its Application to Petroleum Geology, edited by: Larsen, R. M., Brekke, H., Larsen, B. T. and Talleraas, E., Norwegian Petroleum Society (NPF) Special Publications, 1, 481–494, 1992.
- 1175 Haremo, P., Andresen, A., Dypvik, H., Nagy, J., Elverøi, A., Eikeland, T. A. and Johansen, H.: Structural development along the Billefjorden Fault Zone in the area between Kjellströmdalen and Adventdalen/Sassendalen, central Spitsbergen, Polar Res., 8, 195–216, 1990.
- 1180 Harland, W. B.: Contribution of Spitsbergen to understanding of tectonic evolution of North Atlantic region, AAPG Memoirs, 12, 817–851, 1969.
- 1185 Harland, W. B. and Horsfield, W. T.: West Spitsbergen Orogen, in: Mesozoic–Cenozoic orogenic belts, edited by: Spencer, A. M., Geol. Soc. London Spec. Publ., 4, 747–755, 1974.
- Harland, W. B. and Kelly, S. R. A.: Eastern Svalbard Platform, in: Geology of Svalbard, edited by: Harland, W. B., Geol. Soc. London Mem., 17, 75–95, 1997.
- Harland, W. B. and Wright, N. J. R.: Alternative hypothesis for the pre-Carboniferous evolution of Svalbard, Norsk Polarinst. Skr., 167, 89–117, 1979.
- 1190 Harland, W. B., Cutbill, L. J., Friend, P. F., Gobbett, D. J., Holliday, D. W., Maton, P. I., Parker, J. R. and Wallis, R. H.: The Billefjorden Fault Zone, Spitsbergen – the long history of a major tectonic lineament, Norsk Polarinst. Skr., 161, 1–72, 1974.
- Harland, W. B., Mann, A. and Townsend, C.: Deformation of anhydrite-gypsum rocks in central Spitsbergen, Geol. Mag., 125, 103–116, 1988.
- 1195 Holliday, D. W. and Cutbill, J. L.: The Ebbadalen Formation (Carboniferous), Spitsbergen, Proceedings of the Yorkshire Geological Society, 39, 1, 1, 1–32, 1972.
- Horsfield, W. T.: Glauconite schists of Caledonian age from Spitsbergen, Geol. Mag., 109, 1, 29–36, 1972.

- 1200 Izquierdo-Llavall, E., Roca, E., Xie, H., Pla, O., Muñoz, J.A., Rowan, M. G., Yuan, N. and Huang, S.: Structural style of a fold-and-thrust belt involving laterally-changing, multiple décollements: the Kuqa fold-and-thrust belt (NW China), in: *Fold and Thrust Belts: structural style, evolution and exploration*, Petroleum Group of the Geological Society of London, Burlington House, London, UK, 31<sup>st</sup> October – 02<sup>nd</sup> November, 2017.
- 1205 Johannessen, E.: *Facies analysis of the Ebbadalen Formation, Middle Carboniferous, Billefjorden Trough, Spitsbergen*, unpublished Master's Thesis, University of Bergen, Bergen, Norway, 314 pp., 1980.
- Johannessen, E. P. and Steel, R. J.: Mid-Carboniferous extension and rift-infill sequences in the Billefjorden Trough, Svalbard, *Norsk Geol. Tidsskr.*, 72, 35–48, 1992.
- 1210 Johansson, Å. and Gee, D. G.: The late Palaeoproterozoic Eskolabreen granitoids of southern Ny Friesland, Svalbard Caledonides – geochemistry, age, and origin, *GFF*, 121:2, 113–126, 1999.
- Johansson, Å., Larionov, A. N., Gee, D. G., Ohta, Y., Tebenkov, A. M. and Sandelin, S.: Greenvillian and Caledonian tectono-magmatic activity in northeasternmost Svalbard, in: *The Neoproterozoic Timanide Orogen of Eastern Baltica*, edited by: Gee, D. G. and Pease, V., *Geol. Soc. London Memoirs*, 30, 207–232, 2004.
- 1215 Johansson, Å., Gee, D. G., Larionov, A. N., Ohta, Y. and Tebenkov, A. M.: Greenvillian and Caledonian evolution of eastern Svalbard – a tale of two orogenies, *Terra Nova*, 17, 317–325, 2005.
- 1220 Koehl, J.-B. P.: Impact of Timanian thrusts on the Phanerozoic tectonic history of Svalbard, Keynote lecture, EGU General Assembly, May 3<sup>rd</sup>–8<sup>th</sup> 2020, Vienna, Austria, 2020.
- Koehl, J.-B. P.: Latest Devonian–Mississippian tectonic history of Bjørnøya, *Solid Earth*, in prep.
- Koehl, J.-B. P. and Muñoz–Barrera, J. M.: From widespread Mississippian to localized Pennsylvanian extension in central Spitsbergen, Svalbard, *Solid Earth*, 9, 1535–1558, 2018.
- 1225 Koehl, J.-B. P., Tveranger, J., Osmundsen, P. T., Braathen, A., Taule, C. and Collombin, M.: Fault-growth deposit in a Carboniferous rift-basin: the Billefjorden Trough, Svalbard, *Geophys. Res. Abstracts*, 18, EGU General Assembly 2016, Vienna, Austria, 17<sup>th</sup>–22<sup>nd</sup> April, 2016.
- Koehl, J.-B. P., Bergh, S. G., Henningsen, T. and Faleide, J. I.: Middle to Late Devonian–Carboniferous collapse basins on the Finnmark Platform and in the southwesternmost Nordkapp basin, SW Barents Sea, *Solid Earth*, 9, 341–372, 2018.
- 1230

- Koehl, J.-B. P., Collombin, M., Taule, C., Christophersen, G., Allaart, L., Tveranger, J. and Noormets, R.: Influence of WNW–ESE-striking faults on Devonian–Permian sedimentary rocks in Billefjorden and implications for Ellesmerian and Eurekan tectonic events, submitted.
- 1235 Kościńska, K., Majka, J., Mazur, S., Krumbholz, M., Klonowska, I., Manecki, M., Czerny, J. and Dwornik, M.: Blueschist facies metamorphism in Nordenskiöld Land of west-central Svalbard, *Terra Nova*, 26, 377–386, 2014.
- Lamar, D. L. and Douglass, D. N.: Geology of an area astride the Billefjorden Fault Zone, Northern Dicksonland, Spitsbergen, Svalbard, *Polarist. Skr.*, 197, 46 pp., 1995.
- 1240 Lamar, D. L., Reed, W. E. and Douglass, D. N.: Structures bearing on the sense and magnitude of displacement and tectonic significance of Billefjorden Fault Zone, Dicksonland, Spitsbergen, Svalbard: Progress report, 1982 field season, Lamar-Merifield, Geologists, Technical report 82-6, 48 pp., 1982.
- Lamar, D. L., Reed, W. E. and Douglass, D. N.: Billefjorden fault zone, Spitsbergen: Is it part of a major Late Devonian transform?, *GSA*, 97, 1083–1088, 1986.
- 1245 Larsen, B. T.: Tertiary thrust tectonics in the east of Spitsbergen, and implications for the plate-tectonic development of the North-Atlantic, in: *Tertiary Tectonics of Svalbard*, edited by: Dallmann, W. K., Ohta, Y. and Andresen, A., *Norsk Polarinstitutt Rapportserie*, 46, 85–88, 1988.
- 1250 Larssen, G. B., Elvebakk, G., Henriksen, S. E., Nilsson, I., Samuelsen, T. J., Svånå, T. A., Stemmerik, L. and Worsley, D.: Upper Palaeozoic lithostratigraphy of the Southern Norwegian Barents Sea, *Norwegian Petroleum Directorate Bulletin*, 9, 1–76, 2002.
- Lindemann, F.-J., Volohonsky, E. and Marshall, J. E.: A bonebed in the Hørbyreen Formation (Famennian-Viséan) on Spitsbergen, *NGF Abstracts and Proceedings*, 1, Winter Meeting, 1255 Oslo, 8–10<sup>th</sup> January, 2013.
- Livshitz, Ju. Ja.: New data on the geological structure of the Pyramiden area (Vestspitsbergen), *Uchenie Zapiski (Inst. Geol. Arctic) Regional'naya geologiya (NIIGA)*, 9, 36-56, 1966.
- Lowell, J. D.: Spitsbergen Tertiary Orogenic Belt and the Spitsbergen Fracture Zone, *Geol. Soc. Am. Bul.*, 83, 3091–3102, 1972.
- 1260 Lønøy, A.: A Mid-Carboniferous, carbonate-dominated platform, Central Spitsbergen, *Norsk Geol. Tidsskr.*, 75, 48–63, 1995.

- Maher, H. D.: Structure and stratigraphy of the Midterhuken peninsula, Bellsund, west Spitsbergen, PhD Thesis, University of Wisconsin–Madison, Madison, USA, 437, 1984.
- 1265 Maher Jr., H. D.: Photointerpretation of Tertiary structures in platform cover strata of interior Oscar II Land, Spitsbergen, *Polar Res.*, 6, 155–172, 1988.
- Maher Jr., H. D.: Atypical rifting during the Carboniferous of the NW Barents Shelf, Report for Saga Petroleum, November 1996, 1996.
- Maher Jr., H. D. and Braathen, A.: Løvehovden fault and Billefjorden rift basin segmentation and development, Spitsbergen, Norway, *Geol. Mag.*, 148, 1, 154–170, 2011.
- 1270 Maher Jr., H. D. and Welbon, A. I.: Influence of Carboniferous structures on Tertiary tectonism at St. Jonsfjorden and Bellsund, Western Svalbard, *Norsk Geol. Tidsskr.*, 72, 67–75, 1992.
- Maher Jr., H. D., Craddock, C. and Maher, K.: Kinematics of Tertiary structures in upper Paleozoic and Mesozoic strata on Midterhuken, west Spitsbergen, *GSA Bulletin*, 97, 1411–1421, 1986.
- 1275 Manby, G. M. and Lyberis, N.: Tectonic evolution of the Devonian Basin of northern Spitsbergen, *Norsk Geol. Tidsskr.*, 72, 7–19, 1992.
- Manby, G. M. and Michalski, K.: Contrasting metamorphic terranes of Ny-Friesland and their place in the Arctic Caledonides, in: *SvalGeoBase: Proterozoic and Lower Palaeozoic basement of Svalbard – state of knowledge and new perspectives of investigations*, Workshop report, edited by: Dallmann, W. K., Manecki, M., Michalski, K. and Glowacki, P., Tromsø, January 2014, 2014.
- 1280 Manby, G. M., Lyberis, N., Chorowicz, J. and Thiedig, F.: Post-Caledonian tectonics along the Billefjorden fault zone, Svalbard, and implications for the Arctic region, *Geol. Soc. Am. Bul.*, 105, 201–216, 1994.
- 1285 Marshall, J., Lindemann, F. J., Finney, S. and Berry, C.: A Mid Fammenian (Late Devonian) spore assemblage from Svalbard and its significance, CIMP Meeting, Bergen, Norway, 17–18<sup>th</sup> September, 2015.
- 1290 McCann, A. J.: The Billefjorden Fault Zone, Dickson Land, Svalbard: Basement fault control on cover deformation, Ph.D. Thesis, Imperial College, London, UK, 1993.

- McCann, A. J.: Deformation of the Old Red Sandstone of NW Spitsbergen; links to the Ellesmerian and Caledonian orogenies, in: *New Perspectives on the Old Red Sandstone*, edited by: Friends, P. F. and Williams, B. P. J., Geol. Soc. London, 180, 567–584, 2000.
- 1295 McCann, A. J. and Dallmann, W. K.: Reactivation of the long-lived Billefjorden Fault Zone in north central Spitsbergen, Svalbard, *Geol. Mag.*, 133, 63–84, 1996.
- McClay, K. R.: Analogue models of inversion tectonics, *Geol. Soc. London Spec. Publ.*, 44, 41–59, 1989.
- McClay, K. R.: Glossary of thrust tectonics terms, in: *Thrust tectonics*, edited by: McClay, K. R., London, Chapman and Hall, 419–433, 1992.
- 1300 McClay, K. R. and Insley, M. W.: Duplex structures in the Lewis thrust sheet, Crowsnest Pass, Rocky Mountains, Alberta, Canada, *Journal of Structural Geology*, 8, 8, 911–922, 1986.
- McWhae, J. R. H.: The major fault zone of central Vestspitsbergen, *Quart. J. Geol. Soc. Lond.*, 108, 209–232, 1953.
- 1305 Michaelsen, B.: *Strukturgeologie des svalbardischen Überschiebungs- und Faltengürtels im zentralen, östlichen Dickson Land, Spitzbergen (Structural geology of the Svalbardian fold-and-thrust belt in central–eastern Dickson Land, Spitsbergen)*, Master’s Thesis, University of Münster, Münster, Germany, 134 pp., 1998.
- 1310 Michaelsen, B., Piepjohn, K. and Brinkmann, L.: *Struktur und Entwicklung der svalbardischen Mimerelva Synklina im zentralen Dickson Land, Spitzbergen*, Münster. *Forsch. Geol. Paläont.*, 82, 203–214, 1997.
- Molinda, G. M.: *Geologic Hazards and Roof Stability in Coal Mines*, Information Circular, 9466, U.S. Department of Health and Human Services, 2003.
- Murascov, L. G. and Mokin, Ju. I.: Stratigraphic subdivision of the Devonian deposits of Spitsbergen, *Polarinst. Skr.*, 167, 249–261, 1979.
- 1315 Myhre, P. I., Corfu, F. and Andresen, A.: Caledonian anatexis of Grenvillian crust: a U/Pb study of Albert I Land, NW Svalbard, *Norw. J. Geol.*, 89, 173–191, 2008.
- Oakey, G. N. and Chalmers, J. A.: A new model for the Paleogene motion of Greenland relative to North America: Plate reconstructions of the Davis Strait and Nares Strait regions between Canada and Greenland, *J. of Geophys. Res.*, 117, B10401, 2012.
- 1320 Ohta, Y., Dallmeyer, R. D. and Peucat, J. J.: Caledonian terranes in Svalbard, *GSA Spec. Paper*, 230, 1–15, 1989.

- Ohta, Y., Krasil'shchikov, A. A., Lepvrier, C. and Teben'kov, A. M.: Northern continuation of Caledonian high-pressure metamorphic rocks in central-western Spitsbergen, *Polar Res.*, 14, 3, 303–315, 1995.
- 1325 Ohta, Y., Larionov, A. N., Tebenkov, A. M., Lepvrier, C., Maluski, H., Lange, M. and Hellebrandt, B.: Single-zircon Pb-evaporation and  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of the metamorphic and granitic rocks in north-west Spitsbergen, *Polar Res.*, 21, 1, 73–89, 2002.
- Petersen, T. G., Thomsen, T. B., Olausen, S. and Stemmerik, L.: Provenance shifts in an evolving Eurekan foreland basin: the Tertiary Central Basin, Spitsbergen, *J. of Geol. Soc.*, 173, 634–648, 2016.
- 1330 Phillipson, S. E.: The control of coal bed decollement-related slickensides on roof falls in North American Late Paleozoic coal basins, *International Journal of Coal Geology*, 53, 181–195, 2003.
- Phillipson, S. E.: Effects of late Paleozoic foreland deformation on underground coal mine ground instability, Illinois and Appalachians Basins, *International Journal of Coal Geology*, 64, 3–19, 2005.
- 1335 Piepjohn, K.: The Svalbardian–Ellesmerian deformation of the Old Red Sandstone and the pre-Devonian basement in NW Spitsbergen (Svalbard), in: *New Perspectives on the Old Red Sandstone*, edited by: Friend, P. F. and Williams, B. P. J., *Geol. Soc. London Spec. Publi.*, 180, 585–601, 2000.
- 1340 Piepjohn, K., and Dallmann, W. K.: Stratigraphy of the uppermost Old Red Sandstone of Svalbard (Mimerdalen Subgroup), *Polar Res.*, 33:1, 19998, 2014.
- Piepjohn, K., Brinkmann, L., Dißmann, B., Grewing, A., Michaelsen, B. and Kerp, H.: Geologische und strukturelle Entwicklung des Devon im zentralen Dickson Land, Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 175–202, 1997.
- 1345 Piepjohn, K., Brinkmann, L., Grewing, A. and Kerp, H.: New data on the age of the uppermost ORS and the lowermost post-ORS strata in Dickson Land (Spitsbergen) and implications for the age of the Svalbardian deformation, in: *New Perspectives on the Old Red Sandstone*, edited by: Friend, P. F. and Williams, B. P. J., *Geol. Soc. London Spec. Publi.*, 180, 603–609, 2000.
- 1350 Piepjohn, K., Thiedig, F. and Manby, G. M.: Nappe Stacking on Brøggerhalvøya, NW Spitsbergen, *Geol. Jb*, B 91, 55–79, 2001.

- Prosser, S.: Rift-related linked depositional systems and their seismic expression, in: *Tectonics and Seismic Sequence Stratigraphy*, edited by: Williams, G. D. and Dobb, A., Geol. Soc., London, Spec. Publi., 71, 35–66, 1993.
- 1355 Pčelina, T. M., Bogač, S. I. and Gavrilov, B. P.: Novye dannye po litostratigrafii devonskih otloženji rajona Mimerdalen arhipelaga Špicbergen (New data on the lithostratigraphy of the Devonian deposits of the region of Mimerdalen of the Svalbard Archipelago), in: *Geologija osadocnogo cehla arhipelaga Špicbergen (Geology of the sedimentary blanket of the archipelago of Spitsbergen)*, edited by: Krasil'ščikov, A. A. and Mirzaev, M. N., Leningrad: Sevmorgeologija, 7–19, 1986.
- 1360 Ringset, N. and Andresen, A.: The Gipshuken Fault System – Evidence for Tertiary thrusting along the Billefjorden Fault Zone, in: *Tertiary Tectonics of Svalbard*, edited by: Dallmann, W. K., Ohta, Y. and Andresen, A., Norsk Polarinstitut Rapportserie, 46, 67–70, 1988.
- 1365 Roy, J.-C., L. G.: *La géologie du fossé des Vieux Grès Rouges du Spitzberg (archipel du Svalbard, territoire de l'Arctique) – Synthèse stratigraphique, conséquences paléoenvironnementales et tectoniques synsédimentaires*, Mémoires des sciences de la Terre de l'Université Pierre et Marie Curie, Ph.D. Thesis, Pierre and Marie Curie University, Paris, France, 2007-15, 242 pp., 2007.
- 1370 Roy, J.-C.: *La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle*, Charenton-le-pont: Auto-Edition Roy-Poulain, 290 pp., 2009.
- 1375 Roy, J.-C., Chorowicz, J., Deffontaines, B., Lepvrier, C. and Tardy, M.: Clues of gravity sliding tectonics at the Eifelian–Givetian boundary in the Old Red Sandstone of the [late Silurian?]-Devonian trough of Andrée Land (Spitsbergen), in: *La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle*, edited by: Charenton-le-pont: Auto-Edition Roy-Poulain, Norw. J. Geol., unpublished.
- Saalman, K. and Thiedig, F.: Structural Evolution of the Tertiary West Spitsbergen Fold-and-Thrust Belt on Brøggerhalvøya, NW-Spitsbergen, *Polarforschung*, 68, 111–119, 2000.
- 1380 Saalman, K. and Thiedig, F.: Tertiary West Spitsbergen fold and thrust belt on Brøggerhalvøya, Svalbard: Structural evolution and kinematics, *Tectonics*, 20, 6, 976–988, 2001.

- Scheibner, C., Hartkopf-Fröder, C., Blomeier, D. and Forke, H.: The Mississippian (Lower Carboniferous) in northeast Spitsbergen (Svalbard) and a re-evaluation of the Billefjorden Group, *Z. Dt. Ges. Geowiss.*, 163/3, 293–308, 2012.
- 1385 Schlische, R. W.: Geometry and Origin of Fault-Related Folds in Extensional Settings, *AAPG Bull.*, 79, 11, 1661–1678, 1995.
- Schweitzer, H.-J.: Die Devonflora Spitzbergens (The Devonian flora of Spitsbergen), *Palaeontographica Abteilung B Band*, 252, Stuttgart, Schweizerbart Science Publishers, 1999.
- 1390 Smyrak-Sikora, A. A., Johannessen, E. P., Olaussen, S., Sandal, G. and Braathen, A.: Sedimentary architecture during Carboniferous rift initiation – the arid Billefjorden Trough, Svalbard, *J. Geol. Soc. Lond.*, 176, 2, 225–252, 2018.
- Stipp, M., Stünitz, H., Heilbronner, R. and Schmid, S. M.: The eastern Tonale fault zone: a ‘natural laboratory’ for crystal plastic deformation of quartz over a temperature range from 250 to 700 °C, *Journal of Structural Geology*, 24, 1861–1884, 2002.
- 1395 Tonstad, S. A.: The Late Paleozoic development of the Ottar basin from seismic 3D interpretation, Master’s Thesis, University of Tromsø, Tromsø, Norway, 135 pp., 2018.
- Vigran, J. O.: Spores from Devonian deposits, Mimerdalen, Spitsbergen, *Norsk Polarinstitutt Skrifter*, 132, 49 pp., 1964.
- 1400 Vogt, T.: The stratigraphy and tectonics of the Old Red formations of Spitsbergen, *Abstracts of the Proceedings of the Geological Society London*, 1343, 88, 1938.
- Wang, G., Liu, Y., Gu, Y., Xiong, X., Yuan, K., Li, L., Zhou, H., Liu, J. and Shang, G.: Multi-Décollement Structure Modelling of Kuqa Fold-Thrust Belt in Tarim Basin, in: *International Petroleum Technology Conference*, Beijing, China, 26–28 March, 2013.
- 1405 Wilson, R. L. and Wojtal, S. F.: Cumberland Plateau décollement zone at Dunlap, Tennessee, *GSA Centennial Field Guide—Southeastern Section*, 143–148, 1986.
- Witt-Nilsson, P., Gee, D. G. and Hellman, F. J.: Tectonostratigraphy of the Caledonian Atomfjella Antiform of northern Ny Friesland, Svalbard, *Norsk Geol. Tidsskr.*, 78, 67–80, 1998.
- 1410 Würtzen, C. L., Stemmerik, L., Olaussen, S. and Ahokas, J.: Facies analysis of the uppermost Devonian to Lower Carboniferous Billefjorden Group, central Spitsbergen, Svalbard, *NGF Winter conference*, Bergen, Norway, 7–9<sup>th</sup> January, 2019.



**Legend**

Lower Cenozoic	Permian (Tempelfjorden Group)
Cretaceous dolerite	Pennsylvanian–Permian (Gipsdalen Group)
Lower Cretaceous	Uppermost Devonian–Mississippian (Billefjorden Group)
Upper Jurassic	Lower–lowermost Upper Devonian
Lower Triassic–Middle Jurassic	Precambrian basement
Brittle faults	Folds
Thrusts	Coastline

1415

**Figure 14:** (a) Topographic–bathymetric map around Spitsbergen modified after Jakobsson

et al. (2012). Abbreviations: Bi: Billefjorden; Bk: Bockfjorden; Kg: Kongsfjorden; Mi: Midterhuken; Ra: Raudfjorden; Re: Reindalspasset; Rø: Røkensåta; Ss: Sassenfjorden; SJ: St-Jonsfjorden; Tp: Tempelfjorden; Tr: Triungen; (b) Geological map modified from

1420

svabardkartet.npolar.no showing the main tectono-stratigraphic units and structures in the study area in central Spitsbergen. Abbreviations: AA: Atomfjella Antiform; Af: Adolfbukta; An: Anservika; BF: Balliolbreen Fault; Fw: Flowerdalen; Gh: Gipshuken; Ly: Lykteneset; Lø: Løvehovden–Hultberget; Mu: Mumien; Od: Odelfjellet; OF: Odelfjellet Fault; Py: Pyramiden; Re: Reindalspasset; Rs: Robertsonbreen; RT: Robertsonbreen thrust; Se: Sentinelfjellet; TGFZ: Triungen–Grønhorgdalen Fault Zone; Tr: Triungen; Yg: Yggdrasilkampen.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic





|



**Formatted:** Font: Bold, Not Italic








**Figure 332:** Aerial photograph of the Pyramiden coal mine locality showing intensely folded (dotted orange lines) Lower Devonian rocks in the west juxtaposed against clastic- and coal-rich, uppermost Devonian–Mississippian sedimentary rocks of the Billefjorden Group, which are overlain by Pennsylvanian–lower Permian strata of the Gipsdalen Group in the east. Dashed lines represent lithostratigraphic transitions. Dotted lines represent bedding surfaces as seen on the photograph, whereas white symbols indicate bedding trend and dip in map view (see [Figure 4Figure 3a](#)). Note the Z-shaped fabrics of uppermost Devonian–Mississippian sedimentary strata (yellow lines) along potential bedding-parallel décollements (red lines) near the boundary between Lower Devonian and uppermost Devonian–Mississippian sedimentary rocks. The white frame shows the location of [Figure 4Figure 3b](#). Lower hemisphere Schmidt stereonet show (1) contoured poles of fracture surfaces in the uppermost Devonian–Mississippian Billefjorden Group (red indicates high and blue low density), (2) contoured poles of fracture surfaces within sandstone units of the Billefjorden Group, (3) great circles of fracture surfaces within coaly shale- and coal-bearing units of the Billefjorden Group, and (4) contoured poles of fracture surfaces in Lower Devonian rocks. Photo by Åsle Strøm.

Formatted: No underline, Do not check spelling or grammar

Formatted: No underline



1450 **Figure 443:** (a) Satellite photograph of the Pyramiden locality (Figure 3Figure 2) from  
 toposvalbard.npolar.no. See legend in Figure 3Figure 2. Bedding surface measurements are  
 shown in white. The lower hemisphere Schmidt stereonet shows bedding surface  
 measurements in the Billefjorden Group as contoured poles (red indicates high and blue low  
 density); (b) Field photograph of the base of uppermost Devonian–Mississippian, coaly shale-  
 and coal-rich sedimentary rocks of the Billefjorden Group below the mine entrance in  
 1455 Pyramiden. The photo shows gently east-dipping stratigraphic unit boundaries that localized  
 the formation of bedding-parallel décollements (thick red and thick dashed red lines). Within

Formatted: No underline

Formatted: No underline

1460

individual units, coal displays phyllitic, Z-shaped shear fabrics (yellow lines) parallel-subparallel to steeply east-dipping, intra-unit bedding surfaces (dashed yellow lines) that are truncated by subparallel, steeply east-dipping thrusts (thin dashed red lines). See blue hammer (c. 40 cm) on the foreground and person (c. 1.75 m) in the background for scales.

1465

See supplement 4 for uninterpreted photograph. Location is shown in Figure 3~~Figure 2~~; (c) Field photograph showing gently south-dipping Lower Devonian rocks of the Wood Bay Formation unconformably overlain by flat-lying strata of the Billefjorden Group (dashed green bedding surfaces), Hultberget Formation (dashed red bedding surfaces) and Wordiekammen Formation (dashed blue bedding surfaces) in the hanging wall of the Triungen-Grønhorgdalen Fault Zone in Triungen (see Figure 1~~Figure 1b~~ for location). The upper right inset displays the angular unconformity (dotted yellow line) between gently south-dipping (tilted?) Lower Devonian sedimentary rocks of the Wood Bay Formation


1470

(bedding surfaces in dashed orange) and overlying flat-lying strata of the Billefjorden Group; (d) Field photograph of the inferred location of the Triungen-Grønhorgdalen Fault Zone in Triungen showing that the fault trace is not exposed and is covered by local black scree probably belonging to uppermost Devonian-Mississippian coals of the Billefjorden Group. View is towards the west-northwest; (e) Same as (d) with view towards the north.


Formatted: No underline

Formatted: No underline, Font color: Auto

Formatted: Font: Not Italic, No underline, Font color: Auto, Check spelling and grammar



1475




**Figure 554:** Seismic sections in Two-Way Time (TWT) and associated zoomed-in portions in Saassenfjorden–Tempelfjorden (a–f) and Reindalspasset (g). See **Figure 1** **Figure 1b** for locations. (a) NE–SW-trending section showing minor reverse offset and fault-propagation folding in thickened uppermost Devonian–Mississippian sedimentary rocks along WNW–ESE- to NW–SE-striking, deep-seated basement faults, and Eurekan thrusts in overlying Pennsylvanian–Permian strata; (b) Zoom in SW-verging, coal-bearing duplexes acting as top-SW Eurekan décollements in thickened, uppermost Devonian–Mississippian sedimentary deposits; (c) Zoom in NW–SE-striking Eurekan thrusts that flatten into décollements within uppermost Devonian–Mississippian coals and at the top of the Wordiekammen Formation; (d) NE–west-trending, arch-shaped section showing the potential continuation of the Billefjorden Fault Zone bounding thick uppermost Devonian–Permian sedimentary deposits and top-west Eurekan thrusts within lower Permian rocks; (e)

1480


1485

**Formatted:** No underline, Font color: Auto  
**Formatted:** Font: Not Italic, No underline, Font color: Auto, Check spelling and grammar


1490 **Zoom in coal-bearing duplexes in uppermost Devonian–Mississippian sedimentary strata**  
indicating top-west early Cenozoic movement; (f) **Zoom in Eurekan thrusts flattening into a**  
décollement near the top of the Wordiekammen Formation; (g) **West–east-trending section**  
showing a top-east Eurekan detachment in Mesozoic sedimentary rocks and a broad anticline  
in Devonian–Permian strata. Shale-rich Devonian–Mississippian sedimentary strata thicken  
1495 into the anticline whereas Pennsylvanian–Permian sedimentary rocks thicken away from the  
anticline. The former are truncated by numerous early Cenozoic Eurekanthrusts arranged  
into duplex-like structures that flatten into intra Devonian–Mississippian décollements. Note  
that the thick vertical black line represents the location of exploration well 7816/12-1 ([total  
depth: 2261 meters](#); Eide et al., 1991). Abbreviations: BFZ: Billefjorden Fault Zone.




**(A) Devonian normal faulting**




**(B) Latest Devonian–Mississippian normal faulting**




**(C) Latest Mississippian–Pennsylvanian normal faulting**



**(D) Cenozoic thrusting**



**(E) Analog localities**



**Legend**

- uppermost Pennsylvanian–lowermost Permian (Wordiekammen Fm)
- uppermost Mississippian–Pennsylvanian (Hultberget, Ebbadalen, and Minkinjellet Fms)
- uppermost Devonian–Mississippian (Billefjord Group)
- Devonian
- Basement
- Faults
- Bedding-parallel décollements and thrusts in coal-rich deposits
- Coal-rich deposits

1505 | **Figure 665:** Schematic cross-sections showing the possible evolution of the Billejorden Fault Zone and its relationship with deformation within the Billejorden Group in Devonian–early Cenozoic times. (a) Relatively high pre-Devonian basement in the east and collapse basin in the west with erosion-related exhumation of basement rocks in the footwall of the Odellfjellet Fault, (b) widespread latest Devonian–Mississippian normal faulting and localization of thick coal deposits on basin edges, (c) latest Mississippian–Pennsylvanian normal faulting localized along the Billejorden Fault Zone (Balliolbreen and Odellfjellet faults), (d) inversion of Devonian–Carboniferous normal faults and basins during early Cenozoic Eurekan deformation, including top-west thrusting of Proterozoic basement onto Lower–lowermost Upper Devonian rocks of the Andrée Land Group and Mimerdalen Subgroup along the Balliolbreen Fault, the formation of bedding-parallel décollements and thrusts in uppermost Devonian–Mississippian coal-rich deposits, and intense internal deformation of Lower–lowermost Upper Devonian sedimentary rocks (not detailed here) that acted as a buttress, and (e) parts of the cross-section in (d) that fit field observations (from the present contribution and by previous workers) in key localities discussed in the text (**Figure 1** **Figure 4a–b**). **The location of the schematics in (e) is shown as black frames in (d).** Abbreviations: BaF: Balliolbreen Fault; OF: Odellfjellet Fault; TGFZ: Triungen–Grønhordalen Fault Zone.

**Formatted:** No underline, Font color: Auto

**Formatted:** Font: Not Italic, No underline, Font color: Auto, Check spelling and grammar

**Appendix A: List of digitized publications from the Norwegian Polar Institute's Library.**

- 1520 Abakumov, S. A.: The Lower Hecla Hoeck of the Ny Friesland Peninsula, in: *Geology of Spitsbergen*, Vol. 1, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 98–115, 1965.
- Andresen, A., Bergh, S. G., Haremo, P., Maher Jr., H. and Welbon, A.: Extrem strain partitioning during evolution of a transform plate boundary, Spitsbergen, North Atlantic, 1992, unpublished.
- 1525 Birkenmajer, K.: Course of the geological investigations of the Hornsund area, Vestspitsbergen, in 1957–1958, *Studia Geologica Polonica*, 4, 7–35, 1960.
- Birkenmajer, K.: Course of the geological investigations of the Hornsund area, Vestspitsbergen, in 1530 1959 and 1960, *Studia Geologica Polonica*, 11, 7–33, 1964a.
- Birkenmajer, K.: Devonian, Carboniferous and Permian formations of Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 11, 47–123, 1964b.
- Birkenmajer, K.: Cambrian succession in South Spitsbergen, *Studia Geologica Polonica*, 59, 7–46, 1978a.
- 1535 Birkenmajer, K.: Ordovician succession in South Spitsbergen, *Studia Geologica Polonica*, 59, 47–81, 1978b.
- Birkenmajer, K.: Palaeotransport and source of Early Carboniferous fresh-water clastics of South Spitsbergen, *Studia Geologica Polonica*, 60, 39–43, 1979.
- Birkenmajer, K.: Tertiary tectonic deformation of Lower Cretaceous dolerite dykes in a 1540 Precambrian terrane, South-West Spitsbergen, *Studia Geologica Polonica*, 59, 31–44, 1986.
- Birkenmajer, K.: Precambrian succession at Hornsund, South Spitsbergen: A lithostratigraphic guide, *Studia Geologica Polonica*, 98, 7–66, 1992.
- Birkenmajer, K. and Morawski, T.: Dolerite intrusions of Wedel-Jarlsberg Land Vestspitsbergen, *Studia Geologica Polonica*, 4, 103–123, 1960.
- 1545 Birkenmajer, K. and Narebski, W.: Precambrian amphibolite complex and granitization phenomena in Wedel-Jarlsberg Land, Vestspitsbergen, *Studia Geologica Polonica*, 4, 37–82, 1960.
- Birkenmajer, K. and Wojciechowski, J.: On the age of ore-bearing veins of the Hornsund area, Vestspitsbergen, *Studia Geologica Polonica*, 11, 179–184, 1964.

- 1550 Burov, Yu. P.: Peridotite inclusions and bombs in the trachybasalts of Sverre Volcano in Vestspitsbergen, in: *Geology of Spitsbergen, Vol. 2*, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 267–279, 1965.
- Burov, Yu. P. and Livshits, Yu. Ya.: Poorly differentiated dolerite intrusions in Spitsbergen, in: *Geology of Spitsbergen, Vol. 2*, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E.
- 1555 in 1970, 255–266, 1965.
- Burov, Yu. P. and Murashov, L. G.: Some results of the lithological and stratigraphic study of the Kapp Kjeldsen series in the Bockfjorden area, in: *Material on the geology of Spitsbergen*, edited by: Sokolov, V.N., NIIGA, Leningrad (English translation: The British Library, Lending Division, 1977), 89–97, 1977.
- 1560 Cerny, J., Lipien, G., Manecki, A. and Piestrzynski, A.: Geology and ore-mineralization of the Hecla Hoek succession (Precambrian) in front of Werenskioldbreen, South Spitsbergen, *Studia Geologica Polonica*, 98, 67–113, 1992a.
- Cerny, J., Plywacz, I. and Szubala, L.: Siderite mineralization in the Hecla Hoek succession (Precambrian) at Strypegga, South Spitsbergen, *Studia Geologica Polonica*, 98, 153–169,
- 1565 1992b.
- Dißmann, B. and Grewing, A.: Post-svalbardische kompressive Strukturen im westlichen Dickson Land (Hugindalen), Zentral-Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 235–242, 1997.
- Firsov, L. V. and Livshits, Yu. Ya.: Potassium–Argon dating of dolerites from the region of
- 1570 Sassenfjorden, Vestspitsbergen, in: *Material on the geology of Spitsbergen*, edited by: Sokolov, V.N., NIIGA, Leningrad (English translation: The British Library, Lending Division, 1977), 228–234, 1965.
- Greving, S., Werner, S. and Thiedig, F.: Post-keldonische Ganggesteine auf der nordöstlichen Mitrahavøya (Albert I Land, Nordwest-Spitzbergen), *Münster. Forsch. Geol. Paläont.*, 82,
- 1575 73–78, 1997.
- Guddingsmo, J.: *Strukturgeologisk analyse av tertiært deformerte karbon/perm-bergater ved Svartfjella, nordvestlige Oscar II Land, Spitsbergen*, Master's Thesis, University of Tromsø, Tromsø, Norway, 150 pp.
- Haczewski, G.: Lower Carboniferous alluvial sandy deposits (Hornsundneset Formation) of South
- 1580 Spitsbergen, *Studia Geologica Polonica*, 80, 91–97, 1984.

- Haremo, P.: Geological map of the area between Kjellstrømdalen and Adventdalen/Sassendalen, central Spitsbergen, in: Post – Paleozoic tectonics along the southern part of the Billefjorden and Lomfjorden fault zones and their relation to the west Spitsbergen foldbelt, edited by: Haremo, P. (1992), 1989.
- 1585 Haremo, P.: Post – Paleozoic tectonics along the southern part of the Billefjorden and Lomfjorden fault zones and their relation to the west Spitsbergen foldbelt, Ph.D. Thesis, University of Oslo, Oslo, Norway, 135 pp., 1992.
- Haremo, P., Andresen, A. and Dypvik, H.: Mesozoic extension versus Tertiary compression along the Billefjorden Fault Zone south of Isfjorden, central Spitsbergen, 1993, unpublished.
- 1590 Kempe, M., Niehoff, U., Piepjohn, K. and Thiedig, F.: Kaledonische und Svalbardische Entwicklung im Grundgebirge auf der Blomstrandhalvøya, NW-Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 121–128, 1997.
- Kieres, A. and Piestrzynski, A.: Ore-mineralization of the Hecla Hoek succession (Precambrian) around Werenskioldbreen, South Spitsbergen, *Studia Geologica Polonica*, 98, 115–151, 1992.
- 1595 Klubov, B. A.: The main features of the geological structure of Barentsøya, in: *Geology of Spitsbergen*, Vol. 1, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 89–97, 1965.
- Klubov, B. A., Alekseeva, A. B. and Drozdova, I. N.: On the Triassic coals of Spitsbergen, in: *Material on the geology of Spitsbergen*, edited by: Sokolov, V.N., NIIGA, Leningrad (English translation: The British Library, Lending Division, 1977), 219–227, 1977.
- 1600 Krasil'shchikov, A. A.: Some aspects of the geological history of North Spitsbergen, in: *Geology of Spitsbergen*, Vol. 2, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 32–48, 1965.
- 1605 Lamar, D. L., Reed, W. E. and Douglass, D. N.: Structures bearing on the sense and magnitude of displacement and tectonic significance of Billefjorden Fault Zone, Dicksonland, Spitsbergen, Svalbard: Progress report, 1982 field season, Lamar-Merifield, Geologists, Technical report 82-6, 48 pp., 1982.
- Lange, M. and Hellebrandt, B.: *Geologie, Petrographie und Tektonik des südwestlichen Haakon VII Landes, Nordwest-Spitzbergen*, Münster. Forsch. Geol. Paläont., 82, 99–119, 1997.
- 1610

- Lange, M., Hellebrandt, B., Piepjohn, K., Saalman, K. and Donath, H.-J.: Münstersche Forschungen zur Geologie und Paläontologie, Beiträge zur geologischen Evolution Nordwest-Spitzbergen, 82, 242 pp., 1997.
- 1615 Laptas, A.: Sedimentary evolution of Lower Ordovician carbonate sequence in South Spitsbergen, *Studia Geologica Polonica*, 89, 7–30, 1986.
- Litjes, B. and Thiedig, F.: Geologie und Petrographie des kristallinen Basements und des paläozoischen Bulltinden Konglomerats am Südufer des St. Jonsfjords (Oscar II Land, NW-Spitzbergen), *Münster. Forsch. Geol. Paläont.*, 82, 165–174, 1997.
- 1620 Livshits, Yu. Ya.: Tectonic of central Vestspitsbergen, in: *Geology of Spitsbergen*, Vol. 1, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 59–75, 1965a.
- Livshits, Yu. Ya.: Paleogene deposits of Nordenskiöldbreen Land, Vestspitsbergen, in: *Geology of Spitsbergen*, Vol. 2, edited by: Sokolov, V. N., translated by Bradley, Dr. J. E. in 1970, 193–215, 1965b.
- 1625 McCann, A. J.: The Billefjorden Fault Zone, Dickson Land, Svalbard: Basement fault control on cover deformation, Ph.D. Thesis, Imperial College, London, UK, 1993.
- Michaelsen, B.: Strukturgeologie des svalbardischen Überschiebungs- und Faltengürtels im zentralen, östlichen Dickson Land, Spitzbergen (Structural geology of the Svalbardian fold-and-thrust belt in central–eastern Dickson Land, Spitsbergen), Master’s Thesis, University of Münster, Münster, Germany, 134 pp., 1998.
- 1630 Michaelsen, B., Piepjohn, K. and Brinkmann, L.: Struktur und Entwicklung der svalbardischen Mimerelva Syncline im zentralen Dickson Land, Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 203–214, 1997.
- 1635 Peletz, G., Greving, S. and Thiedig, F.: Der tektonische Bau des überschiebungsgürtels auf der Mitrahelvøya, Albert I Land, NW-Spitzbergen, *Münster. Forsch. Geol. Paläont.*, 82, 79–86, 1997.
- Piepjohn, K.: Geological Map of Woodfjorden Area (Haakon VII Land, Andrée Land), NW-Spitsbergen, Svalbard, Scale 1 : 150 000, Fachhochschule Karlsruhe, Department of Surveying and Cartography, 1992.
- 1640 Piepjohn, K.: Geologische Karte Germaniahelvøya, Haakon VII Land Spitzbergen (Svalbard), Scale 1 : 50 000, Fachhochschule Karlsruhe, 1993.

- Piepjoh, K.: Überblick über die Arktis-Expeditionen der Spitzbergen-Arbeitsgruppe von Prof. Dr. F. Thiedig, Geologisch-Paläontologisches Institut der Universität Münster, Münster. Forsch. Geol. Paläont., 82, 1–14, 1997a.
- 1645 Piepjoh, K.: Erläuterungen zur Geologischen Karte 1:150.000 des Woodfjorden-Gebietes (Haakon VII Land, Andrée Land), NW-Spitzbergen, Svalbard, Münster. Forsch. Geol. Paläont., 82, 15–37, 1997b.
- Piepjoh, K. and Thiedig, F.: Erläuterungen zur Geologischen Karte 1:50.000 der Germaniahelvøya, Haakon VII Land, Spitzbergen (Svalbard), Münster. Forsch. Geol. Paläont., 82, 39–52, 1997a.
- 1650 Piepjoh, K. and Thiedig, F.: Geologisch-tektonische Evolution NW-Spitzbergens im Paläozoikum, Münster. Forsch. Geol. Paläont., 82, 215–233, 1997b.
- Piepjoh, K., Greving, S., Peletz, G., Thielemann, T., Werner, S. and Thiedig, F.: Kaledonische und svalbardische Entwicklung im kristallinen Basement auf der Mitrahelvøya, Albert I Land, NW-Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 53–72, 1997a.
- 1655 Piepjoh, K., Brinkmann, L., Dißmann, B., Grewing, A., Michaelsen, B. and Kerp, H.: Geologische und strukturelle Entwicklung des Devon im zentralen Dickson Land, Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 175–202, 1997b.
- Roy, J.-C.: La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle, Charenton-le-pont: Auto-Edition Roy-Poulain, 290 pp., 2009.
- 1660 Roy, J.-C., Chorowicz, J., Deffontaines, B., Lepvrier, C. and Tardy, M.: Clues of gravity sliding tectonics at the Eifelian–Givetian boundary in the Old Red Sandstone of the [late Silurian?]-Devonian trough of Andrée Land (Spitzbergen), in: La saga des vieux grès rouges du Spitzberg (archipel du Svalbard, Arctique): Une histoire géologique et naturelle, edited by: Charenton-le-pont: Auto-Edition Roy-Poulain, Norw. J. Geol., unpublished.
- 1665 Rozycki, S. Z.: Geology of the north-western part of Torrell Land, Vestspitsbergen, *Studia Geologica Polonica*, 2, 4–98, 1959a.
- Rozycki, S. Z.: Geological cross-sections of the north-western part of Torrell Land, Vestspitsbergen, 1 : 25000, *Studia Geologica Polonica*, 2, 1959b.
- 1670 Rozycki, S. Z.: Geological map of the north-western part of Torrell Land, Vestspitsbergen, 1 : 50000, *Studia Geologica Polonica*, 2, 1959c.

- Saalmann, K. and Brommer, A.: Stratigraphy and structural evolution of eastern Brøggerhalvøya, NW-Spitsbergen, Münster. Forsch. Geol. Paläont., 82, 147–164, 1997.
- 1675 Saalmann, K., Piepjohn, K. and Thiedig, F.: Involvierung des Tertiärs von Ny-Ålesund in den alpidischen Deckenbau der Brøggerhalvøya, NW-Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 129–145, 1997.
- Siedlecki, S.: Culm beds of the SW. coast of Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 4, 93–102, 1960.
- 1680 Siedlecki, S. and Turnau, E.: Palynological investigations of culm in the area SW of Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 11, 125–140, 1964.
- Smulikowski, W.: Petrology and Some Structural Data of Lower Metamorphic Formations of the Hecla Hoek Succession in Hornsund, Vestspitsbergen, *Studia Geologica Polonica*, 18, 3–107, 1965a.
- 1685 Smulikowski, W.: Geological sketch-map of upper Revdalen, 1 : 2500, *Studia Geologica Polonica*, 18, 1965b.
- Smulikowski, W.: Geological map of Kvartsittsletta and of SW margin of Werenskioldbreen, 1 : 5000, *Studia Geologica Polonica*, 18, 1965c.
- Smulikowski, W.: Directions of linear structures in the Hecla Hoek succession of the SW Wedel-Jarlsberg Land, 1 : 25000, *Studia Geologica Polonica*, 18, 1965d.
- 1690 Smulikowski, W.: Sketch-map to show areas of detailed petrological and structural investigations in the Hecla Hoek succession of the SW Wedel-Jarlsberg Land, 1 : 25000, *Studia Geologica Polonica*, 18, 1965e.
- 1695 Smulikowski, W.: Some petrological and structural observations in the Hecla Hoek succession between Werenskioldbreen and Torellbreen, Vestspitsbergen, *Studia Geologica Polonica*, 21, 97–161, 1968a.
- Smulikowski, W.: Geological map of the environs of Werenskioldbreen, 1 : 25000, *Studia Geologica Polonica*, 21, 1968b.
- Thielmann, T. and Thiedig, F.: Paläozoisch-postkaledonische Sedimente auf Mitrahålvøya, NW-Spitzbergen, Münster. Forsch. Geol. Paläont., 82, 87–98, 1997.
- 1700 Ustritskii, V. I.: Main features of the stratigraphy and palaeogeography of the upper Palaeozoic of Spitsbergen, in: *Material on the geology of Spitsbergen*, edited by: Sokolov, V.N., NIIGA,

- Leningrad (English translation: The British Library, Lending Division, 1977), 98–124, 1967.
- 1705 Wojciechowski, J.: Ore-bearing veins of the Hornsund area, Vestspitsbergen, *Studia Geologica Polonica*, 11, 173–177, 1964.
- Witt-Nilsson, P. W.: The West Ny Friesland Terrane: An Exhumed Mid-Crustal Obliquely Convergent Orogen, Ph.D. Thesis, Uppsala University, Uppsala, Sweden, 121 pp., 1998.
- Witt-Nilsson, P. W.: Caledonian mid-crustal oblique convergence in eastern Svalbard, 34 pp., 1998, unpublished.
- 1710 Witt-Nilsson, P. W., Hellmann, F. J., Johansson, Å, Larionov, A. N. and Tebenkov, A. M.: Structural and geochronological studies of mylonites along a major Caledonian fault zone, northeastern Spitsbergen, 36 pp., 1998, unpublished.
- Wright, N. J. R.: The Billefjorden Group Central and Eastern Spitsbergen, Cambridge Arctic Shelf Program Report, 6, 79 pp., 1975a.
- 1715 Wright, N. J. R.: The Billefjorden Group of Western Spitsbergen, Cambridge Arctic Shelf Program Report, 9, 37 pp., 1975b.
- Wright, N. J. R.: The Carboniferous and Permian evolution of Svalbard, Cambridge Arctic Shelf Program Report, 25, 51 pp., 1976.