

Interactive comment on “A reconstruction of Iberia accounting for W-Tethys/N-Atlantic kinematics since the late Permian-Triassic” by Paul Angrand et al.

Paul Angrand et al.

paul.angrand@get.omp.eu

Received and published: 14 April 2020

1. Response to the major point

We acknowledge the comment done by an anonymous referee about our manuscript. This comment raises a major issue about a geological implication of our model that needs to be clarified. This is about the role of the Ventaniella Fault as a good candidate to accommodate the Iberia-Ebro relative motion.

This point is critical as most of the recent models accounting for a segmented Iberian plate invoke the Ventaniella Fault as a main boundary fault accommodating a large

Printer-friendly version

Discussion paper



amount of Iberia-Ebro movement (Tugend et al., 2015; Nirrengarten et al., 2018). As pointed out by the referee, the estimated left-lateral displacement along the Ventaniella Fault (after correction of the right-lateral Cenozoic displacement) is expected to be in the order of a few kilometers during late Paleozoic-middle Cretaceous (Tavani et al., 2011).

We actually do not propose that the Ventaniella Fault accommodates in our reconstruction the large displacement between the Ebro block and Europe. We made this point clearer by reorganizing the discussion (please note that the shape of the final version in the revised manuscript may differ a little bit from this one). The structure of the discussion is now:

"Discussion: Implications for strike-slip movements and the Europe-Iberia boundary

1. Amount of strike-slip displacement # former discussion section
2. Strike-slip structures in the intra-Iberian basins # new section

Despite the requirement of 245 km left-lateral strike-slip displacement along the Iberia-Ebro boundary from 160 to 100 Ma, there are no simple geological evidence in support of a unique major crustal-scale fault in the Iberian Range-Basque Cantabrian Basin system.

Several studies have suggested that a left-lateral shear zone can be recognized along the Iberian Range and the Basque-Cantabrian rifts system. Geological evidence includes the High Tagus Fault in the Iberian Range (Aldega et al., 2019; Aurell et al., 2019) and the Ventaniella Fault in the Basque-Cantabrian region (e.g., Tavani et al., 2011). The latter fault is often considered in recent reconstructions to accommodate alone the Iberia-Ebro movement (Tugend et al., 2015; Nirrengarten et al., 2018). However, the estimated left-lateral displacement along the Ventaniella Fault is only in the order of magnitude of a few kilometers (Tavani et al., 2011) and therefore cannot be used as a North Pyrenean Fault equivalent.

[Printer-friendly version](#)[Discussion paper](#)

In the Basque Cantabrian Basin, the Ventaniella Fault is part of a NW-SE fault system that acted as left-lateral shear zone during the Late Jurassic-Early Cretaceous and has been subsequently inverted with a right-lateral kinematic during the Cenozoic (De Vicente et al., 2011; Tavani et al., 2011; Cámara, 2017). These faults have a Triassic origin (Tavani and Granado, 2015). Tectonic activity along these faults gets younger NE-ward (Ubierna fault: Late Jurassic-Early Cretaceous; Zamanza-Oña fault: Early-Middle Cretaceous; salt tectonics in the center of the basin, Cámara, 2017). The required Iberia-Ebro displacement could have been distributed along these structures.

The role of the weak Triassic evaporites in efficiently decoupling deformation in the pre-salt basement from the thin-skinned extension in sedimentary cover has been emphasized largely in the Pyrenees (Lagabriele et al., 2020; Grool et al., 2019; Duretz et al., 2019; Jourdon et al., 2020). Salt tectonics has also been suggested to have been particularly significant from the Jurassic through the Early Cretaceous in Mesozoic basins that shaped NW-directed boundary between Ebro and Iberia, including the Basque-Cantabrian Basin (Cámara, 2017), Parentis Basin (Ferrer et al., 2012), Cameros Basin (Rat et al., 2019) and Maestrat Basin (Vergés et al., 2020). The surface expression of the crustal strike-slip movements is inferred to have been limited in supra-salt layers."

2. Responses to minor points

L2: done

L3: done

L4: we shortened this sentence: "The Late Permian-Triassic Iberian rift basins have accommodated extension, but. . ."

L8: done

L19-21: done

L42: "An alternative scenario has recently emerged (Tugend et al., 2015; Nirrengarten et al., 2018, Tavani et al., 2018), proposing a spatiotemporal partitioning of the defor-

[Printer-friendly version](#)

[Discussion paper](#)



mation in a wider deformation corridor than the single Pyrenean belt. It suggests that the major strike-slip movement required to accommodate the eastwards movement of Iberia first occurred during the Late Jurassic-Early Cretaceous in Northern Iberia along the NW-SE-trending Iberian Massifs. Indeed, along these massifs, several extensional basins registered major subsidence and strike-slip deformation during the Late Permian to middle Cretaceous time interval (Álvaro et al., 1978; Salas and Casas, 1993; Salas et al., 2001; Aldega et al., 2019; Aurell et al., 2019, Soto et al., 2019)."

L 46: merged with L42.

L 59: We add a sentence about the stratigraphy. "This late Permian-Lower Triassic phase is associated with the deposition of thick detrital non-marine deposits in intra-continental basins. Sedimentation became carbonaceous during the middle Triassic. Finally, the Late Triassic is characterized by a thick evaporitic (mainly salt) sequence (e.g., Orti et al., 2017)."

L 71: We keep this as we added informations about the stratigraphy.

L 76-81: This paragraph is needed to introduce the following paragraph.

L 82-83: We reorganized this paragraph according to RC2's comments.

L 90-93 and L 93-94: "The persistence of shallow-marine to non-marine deposition during this period contrasts with the large accommodation space that is required at larger scale to sediment the giant evaporitic province in the Late Permian (Jackson et al., 2019) and in the Late Triassic (Stolfova and Shannon, 2009; Leleu et al., 2016; Orti et al., 2017). Therefore subsidence associated with crustal thinning expected for this period does not fit with increase of tectonic subsidence as predicted by modeling (McKenzie, 1978)."

L 94-104: We prefer keeping this paragraph in that position as it supports the extension phase we discuss later in the text.

L 96: done

Printer-friendly version

Discussion paper



L112-115: done

L119: We agree that it would be interesting to add more subsidence data but we chose on purpose to limit the number of curves to not overload the figure. In addition, and as reported by the referee, subsidence curves in Gomez et al. (2002) are not covering the all Mesozoic and do not show the Triassic phase.

L 139: we removed "strike slip deformation".

L 174: done.

3. References

Aldega, L., Viola, G., Casas-Sainz, A., Marcén, M., Román-Berdiel, T., and van der Lelij, R.: Unraveling Multiple Thermotectonic Events Accommodated by Crustal-Scale Faults in Northern Iberia, Spain: Insights From K-Ar Dating of Clay Gouges, *Tectonics*, 38, 3629–3651, <https://doi.org/10.1029/2019TC005585>, 2019.

Alvaro, M. (1979). Modelo De Evolución Geotectónica Para. *Acta Geològica Hispànica*, 14, 172–177.

Aurell, M., Fregenal-Martínez, M., Bádenas, B., Muñoz-García, M. B., Élez, J., Meléndez, N., and de Santisteban, C.: Middle Jurassic–Early Cretaceous tectono-sedimentary evolution of the southwestern Iberian Basin (central Spain): Major palaeogeographical changes in the geotectonic framework of the Western Tethys, *Earth-Science Reviews*, 199, 102–198, <https://doi.org/10.1016/j.earscirev.2019.102983>, <https://doi.org/10.1016/j.earscirev.2019.102983>, 2019.

Cámara, P. (2017). Salt and Strike-Slip Tectonics as Main Drivers in the Structural Evolution of the Basque-Cantabrian Basin, Spain. In *Permo-Triassic Salt Provinces of Europe, North Africa and the Atlantic Margins* (pp. 371-393). Elsevier.

De Vicente, G., Cloetingh, S. A. P. L., Van Wees, J. D., & Cunha, P. P. (2011). Tectonic classification of Cenozoic Iberian foreland basins. *Tectonophysics*, 502(1-2), 38-61.

Printer-friendly version

Discussion paper



Duretz, T., Asti, R., Lagabrielle, Y., Brun, J. P., Jourdon, A., Clerc, C., and Corre, B.: Numerical modelling of Cretaceous Pyrenean Rifting: The interaction between mantle exhumation and syn-rift salt tectonics, *Basin Research*, pp. 1–16, <https://doi.org/10.1111/bre.12389>, 2019.

Ferrer, O., Jackson, M. P. A., Roca, E., & Rubinat, M. (2012). Evolution of salt structures during extension and inversion of the Offshore Parentis Basin (Eastern Bay of Biscay). *Geological Society, London, Special Publications*, 363(1), 361-380.

Gómez, M., Vergés, J., & Riaza, C. (2002). Inversion tectonics of the northern margin of the Basque Cantabrian Basin. *Bulletin de La Societe Geologique de France*, 173(5), 449–459. <https://doi.org/10.2113/173.5.449>

Grool, A. R., Huismans, R. S., and Ford, M.: Salt décollement and rift inheritance controls on crustal deformation in orogens, *Terra Nova*, 31, 562–568, <https://doi.org/10.1111/ter.12428>, 2019.

Jackson, C. A. L., Gawthorpe, R. L., Elliott, G. M., & Rogers, E. R. (2019). Salt thickness and composition influence rift structural style , northern North Sea , offshore Norway. July 2018, 514–538. <https://doi.org/10.1111/bre.12332>

Jourdon, A., Mouthereau, F., Pourhiet, L. L., and Callot, J. P.: Topographic and Tectonic Evolution of Mountain Belts Controlled by Salt Thickness and Rift Architecture, pp. 1–14, <https://doi.org/10.1029/2019TC005903>, 2020.

Lagabrielle, Y., Asti, R., Duretz, T., Clerc, C., Fourcade, S., Teixell, A., Labaume, P., Corre, B., and Saspiturry, N.: A review of cretaceous smooth-slopes extensional basins along the Iberia-Eurasia plate boundary: How pre-rift salt controls the modes of continental rifting and mantle exhumation, *Earth-Science Reviews*, 201, 103–171, <https://doi.org/10.1016/j.earscirev.2019.103071>, <https://doi.org/10.1016/j.earscirev.2019.103071>, 2020.

Leleu, S., Hartley, A. J., van Oosterhout, C., Kennan, L., Ruckwied, K., & Gerdes,

[Printer-friendly version](#)[Discussion paper](#)

K. (2016). Structural, stratigraphic and sedimentological characterisation of a wide rift system: The Triassic rift system of the Central Atlantic Domain. *Earth-Science Reviews*, 158, 89–124. <https://doi.org/10.1016/j.earscirev.2016.03.008>

McKenzie, D. (1978). Some remarks on the development of sedimentary basins. *Earth and Planetary Science Letters*, 40(1), 25–32. [https://doi.org/10.1016/0012-821X\(78\)90071-7](https://doi.org/10.1016/0012-821X(78)90071-7)

Nirrengarten, M., Manatschal, G., Tugend, J., Kuszniir, N. J., and Sauter, D.: Kinematic Evolution of the Southern North Atlantic: Implications for the Formation of Hyperextended Rift Systems, *Tectonics*, 37, 89–118, <https://doi.org/10.1002/2017TC004495>, 2018.

Ortí, F., Pérez-López, A., & Salvany, J. M. (2017). Triassic evaporites of Iberia: Sedimentological and palaeogeographical implications for the western Neotethys evolution during the Middle Triassic–Earliest Jurassic. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 471, 157–180. <https://doi.org/10.1016/j.palaeo.2017.01.025>

Rat, J., Mouthereau, F., Bricchau, S., Crémades, A., Bernet, M., Balvay, M., Ganne, J., Lahfid, A., and Gautheron, C.: Tectonothermal Evolution of the Cameros Basin: Implications for Tectonics of North Iberia, *Tectonics*, 38, 440–469, <https://doi.org/10.1029/2018TC005294>, 2019.

Salas, R., & Casas, A. (1993). Mesozoic extensional tectonics, stratigraphy and crustal evolution during the Alpine cycle of the eastern Iberian basin. *Tectonophysics*, 228(1–2), 33–55. [https://doi.org/10.1016/0040-1951\(93\)90213-4](https://doi.org/10.1016/0040-1951(93)90213-4)

Salas, R., Guimerà, J., Mas, R., Martín-Closas, C., Melendez, A., & Alonso, A. (2001). Evolution of the Mesozoic Central Iberian Rift System and its Cainozoic inversion (Iberian chain). In *Mémoires du Muséum national d'histoire naturelle* (Vol. 186, Issue 2).

Soto, R., Casas-Sainz, A. M., Oliva-Urcia, B., García-Lasanta, C., Izquierdo-Llavall,

[Printer-friendly version](#)[Discussion paper](#)

E., Moussaid, B., Kullberg, J. C., Román-Berdiel, T., Sánchez-Moya, Y., Sopeña, A., Torres-López, S., Villalaín, J. J., El-Ouardi, H., Gil-Peña, I., & Hirt, A. M. (2019). Triassic stretching directions in Iberia and North Africa inferred from magnetic fabrics. *Terra Nova*, 31(5), 465–478. <https://doi.org/10.1111/ter.12416>

Štolfová, K., & Shannon, P. M. (2009). Permo-Triassic development from Ireland to Norway: basin architecture and regional controls. *Geological Journal*, 44(6), 652–676.

Tugend, J., Manatschal, G., and Kusznir, N. J.: Spatial and temporal evolution of hyper-extended rift systems: Implication for the nature, kinematics, and timing of the Iberian-European plate boundary, *Geology*, 43, 15–18, <https://doi.org/10.1130/G36072.1>, 2015.

Tavani, S., Bertok, C., Granado, P., Piana, F., Salas, R., Vigna, B., & Muñoz, J. A. (2018). The Iberia-Eurasia plate boundary east of the Pyrenees. *Earth-Science Reviews*, 187(August), 314–337. <https://doi.org/10.1016/j.earscirev.2018.10.008>

Tavani, S., & Granado, P. (2015). Along-strike evolution of folding, stretching and breaching of supra-salt strata in the Plataforma Burgalesa extensional forced fold system (northern Spain). *Basin Research*, 27(4), 573-585.

Tavani, S., Quintà, A., & Granado, P. (2011). Cenozoic right-lateral wrench tectonics in the Western Pyrenees (Spain): the Ubierna Fault System. *Tectonophysics*, 509(3-4), 238-253.

Vergés, J., Poprawski, Y., Almar, Y., Drzewiecki, P.A., Moragas, M., Bover-Arnal, T., Macchiavelli, C., Wright, W., Messenger, G., Embry, J.-C., Hunt, D., 2020. Tectono-Sedimentary Evolution of Jurassic-Cretaceous diapiric structures: Miravete anticline, Maestrat Basin, Spain. *Basin Research* n/a. doi:10.1111/br.12447

Interactive comment on *Solid Earth Discuss.*, <https://doi.org/10.5194/se-2020-24>, 2020.

Printer-friendly version

Discussion paper

