



April 22, 2020

To  
Prof. Soliva Roger  
The guest editor  
Special Issue in Solid Earth

***Subject: Submission of the REVISED MANUSCRIPT for the Special Issue in Solid Earth entitled "Faults, fractures, and fluid flow in the shallow crust".***

Dear Editor,

We are grateful to you as well as Prof. Francesco Mazzarini (Reviewer#1) and Prof. Christophe Pascal (Reviewer#2) for a detailed review and constructive suggestions on the manuscript **se-2020-30**, and for giving us an opportunity to revise the same. We believe that the comments have helped us to think more deeply and have helped in revising the manuscript considerably, which we hope will be to your satisfaction. *In the revised manuscript, we have taken into account all the recommendations/suggestions provided by the two Reviewers and have tried to clarify our viewpoint for the benefit of the reader.*

Please note that in the marked (revised) manuscript, the changes made on basis of comments of Reviewer#1 appear in **RED**, while changes based on comments of Reviewer#2 appear in **BLUE** color.

At first we outlined the main issues that we have addressed in the revision, followed by comment wise response to the specific points mentioned in the comments section. It may be noted that we have also responded the comments of Reviewer#1 and Reviewer#2 separately, below, after the section “response to Editor”.

### **Response to the editor**

#### **Response to main issues:**

##### ***Issue 1: Better justification of considering BTS and minimum principal stress***

We would like to thank you as well as reviewer#2 for raising this issue. The tensile strength of metabasalt was obtained from **Brazilian Tensile Strength (BTS)** studies which was further used to denote the magnitude of the minimum principle stress ( $\sigma_3$ ). Here we explain the rationale behind this assumption. It is interpreted that the tectonic stress was more dominant during the initial fracturing mechanism which was followed by episodic fluid pressure pulses leading to fracture reactivation and vein emplacement. The 3D Mohr circle diagrams in Fig.7 and Fig.8 (of the revised manuscript), represents the relative fluid pressure conditions only. We have provided the effective normal stress conditions separately in Fig. 11 (Schematic model in the revised manuscript). However, in order to address the Reviewers concern for  $\sigma_3=12$  MPa, we have added a few sentences in the main manuscript justifying the reason behind such a consideration. It definitely enhanced the quality of the manuscript. The magnitude of  $\sigma_3$  can also be estimated using the stress intensity factor (fracture toughness),  $K_{IC}$ .

for mode-I fractures, can be determined using the following equation (Lawn and Wilshaw, 1975; Atkinson, 1989).

$$K_{IC} = \sigma_3 Y (\pi a)^{1/2},$$

where  $\sigma_3$  is the minimum far field stress acting normal to the crack,  $Y$  is a numerical modification factor to account for the crack geometry and edge effect, and 'a' is the crack half-length. Microstructural investigation suggests that the metabasalts are generally fine to medium grained, massive and consists of plagioclase laths and altered pyroxene with minor chlorite, ankerite and quartz (Gupta et al., 2014). Within the metabasalts, variably oriented larger phenocryst grains range from 2 mm to 8 mm in length. It is evident that the large phenocrysts were more susceptible to generate the initial fractures as compared to the medium to fine grained groundmass in these types of rocks. Therefore, we have considered that the length of the initial crack ranges from 2 mm to 8 mm (Brace, 1964).  $K_{IC}$  value for the metabasalts are found to be 1.069 (Donovan, 2003). Considering this  $K_{IC}$  value and the range for fracture half-length (a), the  $\sigma_3 Y$  value for the metabasalt ranges between 9-19 MPa. From this strength range we considered that 12 MPa is a reasonable estimate for the in situ tensile strength (similar to the laboratory strength) of the corresponding lithology. It may also be noted that, previous studies by Mondal and Acharyya, 2018, conducted in Chitradurga Granite, in close vicinity of the study area also regarded the magnitude of  $\sigma_3 \sim 10$  MPa, to be a good estimation. Combining these estimations with the results obtained from the present studies, we constrained the value for  $\sigma_3 \sim 12$  MPa.

## **Issue 2: Estimation of $P_f$ condition**

You have raised your concern regarding the estimation of  $P_f$  condition, which was also raised by the Reviewer#2. We would again thank the Reviewer for pointing out this problem. In Fig.7, we tried to quantify multiple fluid pressure pulses from the vein pole distribution data. Although, it is critical to differentiate various episodes of fluid ingression from a single large event. Initially we considered the girdle distribution of data points indicating,  $P_f > \sigma_2$  (Jolly and Sanderson, 1997; McKeagney et al., 2004; Mazzarini and Isola, 2007; Martinez-Poza et al., 2016). However, as the distribution of data points show three prominent clusters, we decided to extend our analysis to all the respective clusters and not only the WSW cluster (with highest cluster density). We have incorporated the analysis for the other two clusters as well (NE and SE clusters) in the revised manuscript, as also suggested by Reviewer-1 (see Fig. 8 in the revised manuscript). The number of obtained clusters can also be testified through mixed Bingham analysis using  $K$  vs BIC (i.e., the number of Bingham component of a mixed Bingham distribution vs Bayesian information criterion; Yamaji and Sato, 2011). We have found that the lowest BIC values are obtained when  $K=3$  (number of possible clusters for the given data set), thereby, justifying the selection of the three clusters for the analysis. In each case,  $P_f$  values for the respective clusters were obtained, which is most likely to differ, as mentioned by the Reviewer. It is absolutely true that if we select a tighter cluster, thereby reducing the contour interval, the  $P_f$  value will be further reduced. However, field evidences suggest that most of the vein orientations representing the WSW cluster show a NW-SE to NNW-SSE trend. Also, veins along this orientation attained maximum thickness along with multiple median lines. Thus, we decided to extend the contour interval beyond the data points in order to incorporate the maximum range of vein orientations ( $\theta$ ) lying parallel/sub-parallel to the internal anisotropy (as evident from the anisotropy of magnetic susceptibility study) of the host rock. Also, the contour interval and significance level for each of the clusters were selected in such a way that maximum number of data points are included, in order to obtain a statistically viable data cluster. It is however difficult to quantify the lowest  $P_f$  value; we therefore intend to use the obtained  $P_f$  values from

the respective clusters as examples of low  $P_f$  conditions denoting  $P_f$  fluctuation rather than quantifying the lowest  $P_f$  condition of the study area. We have obtained different shape values ( $\phi$ ) for the inversions, this is because of the variation in the magnitude of  $\sigma_2$ . Both magnitude and orientation of  $\sigma_2$  changes from high to low  $P_f$  conditions as explained by the  $k_2/k_1$  ratio, regarded as the stress ratio ( $\phi$ ), which is expressed as  $k_1 \approx k_2$ , for clustered distribution and  $k_1 \ll k_2$ , for girdle distribution ( $k_1, k_2$  are the concentration parameters of a Bingham distribution; Yamaji, 2016). We hope this provides a better explanation for the interpretation of episodic fluid pressure condition prevailing in the region. In the revised manuscript we have added a number of lines in section 4.3, which increases the clarity of the manuscript.

You have also suggested to include the new figure of the first revision (in accordance with reviewer#1) into the manuscript. In the revised manuscript we have incorporated all the new figures related to the first revision.

### **Issue 3: Fault valve.**

*You have mentioned that “ I consider your revision, especially the new photograph suggesting crack seal with two episodes. Please, better introduce and discuss the new proofs of fault valve mechanism both in the introduction and your result section. Also include the photographs into the new version of the manuscript (not as a supplementary material) and if possible, replace the close view with another photograph showing better evidences of more cycles into a same vein (here we can really see two episodes). It could be more convincing for the fault valve behaviour. I think to see more convincing cases with multiple adjacent branches in the part (a) of you new figure”*

We would like to thank you for your kind words related to the first revision.

We have included a few lines justifying the fault-valve mechanism both in the introduction and in the result section as you have suggested. We have also included the photographs into the main manuscript. We appended Figure 3g, with multiple median lines, where more than two cycles of fluid ingression are observed.

### ***Response to the specific points:***

Revise the sense of shear marked with all the arrows in the 3 bloc diagrams of Figure 10b, c and d, which are kinematically inconsistent with the wing cracks drawn. Also better align the yellow arrow with the T criteria in the riedel plane analysis. **CORRECTED**

Provide more justification in the discussion about the deformation mechanism related to the magnetic fabric. Can we interpret the magnetic fabric as non-coaxial (simple shear) or multi episodic deformation (2 poles on the stereogram), and then having a shortening oblique to the foliation ? **CORRECTED and INCORPORATED the suggestion**

Apart from the above we have also modified the manuscript in accordance with the comments mentioned in the annotated pdf.

With the above revisions we hope that all the questions raised by the Reviewers have been addressed. We hope the revised version is to your satisfaction.

Thanking you  
Yours sincerely  
Tridib Kumar Mondal  
(Corresponding author)

## Response to Reviewer#1 (Prof. Francesco Mazzarini)

We thank Prof. Francesco Mazzarini for his extensive review, and many constructive comments. Although it is suggested for a moderate revision, we believe that the comments have made us to think more deeply and have helped in revising the manuscript considerably. We have implemented, most of the changes suggested and addressed some of his points in details, below. In other cases, we clarified our arguments. It may be noted that the changes are marked in **RED** color.

### **Response to main Issues**

#### **Issue-1: Fault-valve action.**

The reviewer has raised his first concern on *fault-valve* action and commented “*Fault valving: the authors propose that fault valving was the driving mechanism to pump fluids. I think this may be possible but both “fragments of host rock” occurrence within veins and mutually cross-cutting relationships are not exclusively indications of fault-valving. These may also indicate a very high fluid pressure surge and “hydrofracturing” of the host rock (fragments) and coeval emplacement of all the veins (mutual crosscutting veins). Probably a fabric analysis of vein may disentangle this point, the occurrence of crack and seal textures in the veins may clearly indicate cyclic fluid ingressions and fault valve mechanism.*”

Yes, we do agree with the reviewer that the presence of host rock fragments in quartz veins and mutually cross-cutting relationships does not exclusively indicate *fault-valve* action. Therefore, as suggested by the Reviewer, in the revised manuscript we have incorporated a new figure 3g, consisting field photographs of metabasalt hosting quartz veins with multiple median lines. This exemplifies crack seal texture in the veins, thereby clearly indicating cyclic fluid ingression and *fault-valve* action that led to the formation of veins in the metabasalts of the Chitradurga region.

#### **Issue-2: Cluster analysis**

The reviewer has pointed out a number of clustering of veins in the lower hemisphere equal area projection and mentioned that “*In analysing the veins distribution the authors define a clear vein cluster (NNW-SSE veins). If the Jolly and Sanderson method is applied to a cluster of vein clearly it will indicate a cluster distribution. So it is not possible to differentiate different episodes of fluid injection from a large single event where most of the fluid injected the well-oriented NNW-SSE trend- ing fractures (they attain also the maximum vein thicknesses) and also reactivated the other fractures. Moreover, other clusters are apparent in the distribution (for example that around the s2 pole). I suggest to expand the cluster analysis of the data sets and to apply the Jolly and Sanderson method to all the clusters and also to the data set minus the NNW-SSE cluster.*”

Yes, there are a number of clusters, however the WSW cluster is quite prominent with a very high cluster density as compared to the rest. This was the motivation behind primarily considering the WSW cluster for analyzing the fluid pressure condition. Although, it is critical to differentiate various episodes of fluid ingression from a single large event. Therefore, in accordance with the suggestion made by the reviewer we have expanded the analysis and incorporated the other clusters (such as the NE and the SE clusters) in the lower hemisphere equal area projection of pole to veins. The figure 8, provides all the cluster analysis in this regard. In the revised manuscript we have incorporated the same and changes have been made accordingly which definitely increases the clarity of the manuscript.

### **Response to minor comments**

Apart from the above comments the Reviewer has commented few more in the specific line positions of the manuscript, which are listed below.

Lines 47-48. *This is not completely true. Fault valve action develops when fluid pressure shows a cyclic increase. All veins forms because of dilation of fracture's walls. Please reword the sentence.*

**Reply:** We have reframed the entire sentence in the revised manuscript as suggested by the Reviewer.

Lines 55-58. *not clear, please reword.*

**Reply:** The sentences have been reframed in the revised manuscript.

Line 124. *angular enclaves suggest high fluid pressure at the time of vein formation with rupture of wall rocks (hydrofracture). Fault valving is testified by crack and seal fabric of veins. The occurrence of striated veins suggests active stress field after the vein formation.*

**Reply:** This is one of the major issues suggested by the Reviewer. This has been previously addressed in Issue-1: Fault-valve action, and accordingly a new figure 3g has been added with the main manuscript as a response to the comment. The process of fracturing, fracture reactivation/faulting along with vein emplacement prevailed in an active stress field as pointed out by the Reviewer.

Lines 147-148. *this is not clear. Why mutually cross-cutting relationships do imply several cycles of emplacement? It is possible that this pattern of veins indicates a coeval formation of veins?*

**Reply:** At high fluid pressure condition (supra-lithostatic levels),  $P_f$  exceeds the stresses acting on the fracture/fault plane, thus causing failure and forming mesh of fractures. Subsequently, the fluid flows into this mesh of fractures, a phenomenon analogous to “burping” (Sibson and Scott, 1998), and the fluid pressure drops. The veins seal the fractures/faults, and the system is once again ready for the next cycle of fluid pressure build-up, failure, fluid flow (*burping*) and subsequent vein formation. This phenomenon involving repeated cycles leads to crosscutting veins and fractures (mesh-like structure; Mondal and Mamtani, 2013; Sibson, 2004 and the references therein). Under successive low  $P_f$  conditions, the orientations represented by the individual clusters (WSW, NE and SE) were reactivated selectively depending on their respective orientation to the tectonic stress field and the fluid pressure magnitude.

Lines 256-259. *This is an important point. Looking at the veins' pole distribution also a cluster around the intermediate stress ( $s_2$ ) is apparent. Construction of data set from clustered data clearly will lead to a cluster distribution. I suggest here to provide a clear cluster analysis of the data and to extend the analysis to all the resulting clusters. Once these results are obtained, the question to face is: are these different pluses of fluid injections? Is clustering inherited by the fracture network exploited by fluids?*

**Reply:** All the respective clusters have been separately evaluated as suggested by the Reviewer (Fig. 8). Fluid pressure conditions ( $P_f$ ) are determined discretely for each of these clusters. The variation in  $P_f$  clearly indicates the existence of different pulses of fluid injection. All such orientations represented in the cluster have been exploited by fluids as evident from vein emplacement along these orientations. However, the maximum vein thickness is recorded along NNW-SSE direction.

Lines 348-356. *These sentences are repetitive. You may expand the geological back- ground and cut off here these sentences.*

**Reply:** The sentences that are repetitive have been omitted as suggested by the Reviewer.

Line 387. *This is ok from a tectonic point of view. Now rocks are exposed. When did rocks' exposure occur? Unloading from a few kilometres to the surface did not generate any jointing? This point here should be discussed.*

**Reply:** The host metabasalts of the study area have an island arc affinity (Chakrabarti et al., 2006) and can be related to the accretion of EDC (Eastern Dharwar craton) and WDC (Western Dharwar craton) along the CSZ (Chitradurga Shear Zone). Unloading from a few kilometers to the surface might generate jointing in host rocks, however, we did not find any such field evidence to support this.

Line 427. *please expand this point to clearly state that these fracture systems are coeval.*

**Reply:** Necessary change has been made in the manuscript as suggested by the Reviewer.

Line 470. *as in the above comment. How do you ascribe to vein formation the cluster? The cluster may be inherited by the existing fracture network at the time of vein emplacement.*

**Reply:** We have separately analyzed each of the clusters as suggested by the Reviewer. In each case the respective fluid pressure conditions ( $P_f$ ) have been evaluated. The WSW (highest density cluster, Fig. 7c) ascribes to the NNW-SSE trending vein orientations (attaining the maximum thickness). The NE and the SE clusters (Fig. 8) can be related to the vein orientations emplaced along the R and X shear fractures respectively (see figure 11 of the revised manuscript).

Lines 474-476. *As discussed above. Angular fragments of hosting rock within veins indicate hydrofracture with rupture of wall rocks. This is not directly associated with fault-valve action. Moreover, it is not clear if indications of hydrofracturing are observed in all vein sets or only in the NNW-SSE set. To clearly define a fault-valve action a look at the vein microfabric (crack and seal textures) will help.*

**Reply:** This point has been addressed in Issue-1 (response to the main issues) discussed above.

Figure 1. *Add definition of TTG and Supracrustals in the key legend.* Figure 6. *label (c) is poorly visible.* Figure 10. *The stress field in (c) has  $s_1$  vertical while the cartoon has ESE-WNW contraction (yellow arrows). If fluid overpressure acted this should be put into the figure caption.*

**Reply:** All the suggested corrections in Figures. 1, 6 and 10 have been incorporated in the main manuscript.

*Apart from the above, the reviewer suggested a few minor corrections in the annotated manuscript (pdf), which has been addressed in the revised manuscript.*

## Response to Reviewer#2 (Prof. Christophe Pascal)

We thank the Prof. Christophe Pascal for the valuable and constructive comments on the manuscript. The comments have made us think more deeply and have helped in revising the manuscript considerably. The reviewer has mentioned two issues regarding the manuscript. It may be noted that the changes are made in [BLUE](#) color

Issue: 1

The reviewer has raised his first concern regarding the quantification of  $\sigma_3$  and mentioned “*In page 14 (lines 273-274), it is mentioned: “Tensile strength of metabasalt ( $\sim 12$  MPa; obtained from BTS studies) indicates that the minimum principal stress ( $\sigma_3$ ) has to be  $\sigma_3 \sim 12$  MPa.” Thereafter, this value is selected as magnitude for the minimum principal stress for the stress reconstructions (e.g. Fig. 7). I am particularly puzzled by the reasoning. The condition for tensile fracturing is  $\sigma_3 - \text{pore pressure}$  less or equal to the negative value of tensile strength. That is if  $\sigma_3 = 12$  MPa then  $P_f$  has to be equal or higher than 24 MPa... Starting from there could the authors explain how they constrain  $\sigma_3$ ?”*”

Reply:

We would like to thank the reviewer for addressing this issue. The tensile strength of metabasalt was obtained from BTS studies which was further used to denote the magnitude of the minimum principle stress ( $\sigma_3$ ). Here we explain the rationale behind this assumption. It is interpreted that the tectonic stress was more dominant during the initial fracturing, which was followed by episodic fluid pressure pulses leading to fracture reactivation and vein emplacement. The 3D Mohr circle diagrams in Fig.7 and Fig.8 (of the revised manuscript), represents the relative fluid pressure conditions only. We have provided the effective normal stress conditions separately in Fig. 11 (Schematic model in the revised manuscript). However, in order to address the Reviewers concern for  $\sigma_3=12$  MPa, we have added a few sentences in the main manuscript justifying the reason behind such a consideration. It definitely enhanced the quality of the manuscript. The magnitude of  $\sigma_3$  can also be estimated using the stress intensity factor (fracture toughness),  $K_{IC}$ .  $K_{IC}$  for mode-I fractures, can be determined using the following equation (Lawn and Wilshaw, 1975; Atkinson, 1989).

$$K_{IC} = \sigma_3 Y (\pi a)^{1/2},$$

where  $\sigma_3$  is the minimum far field stress acting normal to the crack, Y is a numerical modification factor to account for the crack geometry and edge effect, and ‘a’ is the crack half-length. Microstructural investigation suggests that the metabasalts are generally fine to medium grained, massive and consists of plagioclase laths and altered pyroxene with minor chlorite, ankerite and quartz (Gupta et al., 2014). Within the metabasalts, variably oriented larger phenocryst grains range from 2 mm to 8 mm in length. It is evident that the large phenocrysts were more susceptible to generate the initial fractures as compared to the medium to fine grained groundmass in these types of rocks. Therefore, we have considered that the length of the initial crack ranges from 2 mm to 8 mm (Brace, 1964).  $K_{IC}$  value for the metabasalts are found to be 1.069 (Donovan, 2003). Considering this  $K_{IC}$  value and the range for fracture half-length (a), the  $\sigma_3 Y$  value for the metabasalt ranges between 9-19 MPa. From this strength range we considered that 12 MPa is a reasonable estimate for the in situ tensile strength (similar to the laboratory strength) of the corresponding lithology. It may also be noted that, previous studies by Mondal and Acharyya, 2018, conducted in Chitradurga Granite, in close vicinity of the study area also regarded the magnitude of  $\sigma_3 \sim 10$  MPa, to be a good estimation. Combining these estimations with the results obtained from the present studies, we constrained the value for  $\sigma_3 \sim 12$  MPa.

Issue: 2

The reviewer has raised his second concern regarding the  $P_f$  condition and mentioned “*It is proposed that the veins formed in response to various fluid pressure pulses (see pages 13 and 14, and Fig. 7), which is reasonable and supported by numerous studies (e.g. discussion in Yamaji et al. 2010, p. 1139). The analysis suggests two extreme values for the fluid pressure: a maximum  $P_f$  derived from the Bingham distribution of all the measured veins and a minimum  $P_f$ , corresponding to the Bingham distribution of a selected subset of veins forming an elliptical cluster around  $\sigma_3$  (see Fig. 7). It is totally unclear how the authors have selected the subset indicating, presumably, the actual minimum pore pressure. For example if one decides to select another subset of poles forming a tighter cluster around  $\sigma_3$ ,  $P_f$  will be further reduced. It is just a mathematical consequence (see Jolly and Sanderson 1997)... In addition, stress anisotropy is theoretically insensitive to pore pressure variations. Surprisingly, the two inversions shown in Fig. 7 indicate rather different shape values, though pore pressure is the only quantity that is expected to change. In conclusion, the results themselves suggest that the selection of the subset of poles is merely subjective and the advanced minimum pore pressure value badly supported by the analysis”.*

Reply:

We would again thank the Reviewer for pointing out this problem. In Fig.7, we tried to quantify multiple fluid pressure pulses from the vein pole distribution data. Although, it is critical to differentiate various episodes of fluid ingression from a single large event. Initially we considered the girdle distribution of data points indicating,  $P_f > \sigma_2$  (Jolly and Sanderson, 1997; McKeagney et al., 2004; Mazzarini and Isola, 2007; Martinez-Poza et al., 2016). However, as the distribution of data points show three prominent clusters, we decided to extend our analysis to all the respective clusters and not only the WSW cluster (with highest cluster density). We have incorporated the analysis for the other two clusters as well (NE and SE clusters) in the revised manuscript, as also suggested by Reviewer-1 (see fig. 8 in the revised manuscript). The number of obtained clusters can also be testified through mixed Bingham analysis using K vs BIC (i.e., the number of Bingham component of a mixed Bingham distribution vs Bayesian information criterion; Yamaji and Sato, 2011). We have found that the lowest BIC values are obtained when  $K=3$  (number of possible clusters for the given data set), thereby, justifying the selection of the three clusters for the analysis. In each case,  $P_f$  values for the respective clusters were obtained, which is most likely to differ, as mentioned by the Reviewer. It is absolutely true that if we select a tighter cluster, thereby reducing the contour interval, the  $P_f$  value will be further reduced. However, field evidences suggest that most of the vein orientations representing the WSW cluster show a NW-SE to NNW-SSE trend. Also, veins along this orientation attained maximum thickness along with multiple median lines. Thus, we decided to extend the contour interval beyond the data points in order to incorporate the maximum range of vein orientations ( $\theta$ ) lying parallel/sub-parallel to the internal anisotropy (as evident from the anisotropy of magnetic susceptibility study) of the host rock. Also, the contour interval and significance level for each of the clusters were selected in such a way that maximum number of data points are included, in order to obtain a statistically viable data cluster. It is however difficult to quantify the lowest  $P_f$  value; we therefore intend to use the obtained  $P_f$  values from the respective clusters as examples of low  $P_f$  conditions denoting  $P_f$  fluctuation rather than quantifying the lowest  $P_f$  condition of the study area. We have obtained different shape values ( $\phi$ ) for the inversions, this is because of the

variation in the magnitude of  $\sigma_2$ . Both magnitude and orientation of  $\sigma_2$  changes from high to low  $P_f$  conditions as explained by the  $k_2/k_1$  ratio, regarded as the stress ratio ( $\phi$ ), which is expressed as  $k_1 \approx k_2$ , for clustered distribution and  $k_1 \ll k_2$ , for girdle distribution ( $k_1, k_2$  are the concentration parameters of a Bingham distribution; Yamaji, 2016). We hope this provides a better explanation for the interpretation of episodic fluid pressure condition prevailing in the region. In the revised manuscript we have added a number of lines in section 4.3, which increases the clarity of the manuscript.

With the above revisions I hope that all the questions raised by the Reviewer #2 has been addressed.

**Control of pre-existing fabric in fracture formation, reactivation and vein emplacement under variable fluid pressure conditions: An example from Archean Greenstone belt, India.**

Sreyashi Bhowmick and Tridib Kumar Mondal

Department of Geological Sciences, Jadavpur University

5 Kolkata-700032, West Bengal, India

\*Corresponding author: **Tridib Kumar Mondal** ([tridibkumarmondal@gmail.com](mailto:tridibkumarmondal@gmail.com))

Alternate E-mail: [tridibk.mondal@jadavpuruniversity.in](mailto:tridibk.mondal@jadavpuruniversity.in)

**Abstract**

10 Most of the upper crustal fluid flows are strongly influenced by the pre-existing fractures/foliations in the rocks under a certain state of tectonic stress and fluid pressure condition. In the present study, we analysed a wide range of crosscutting fractures that are filled with quartz veins of variable orientations and thicknesses, from the gold bearing massive metabasalts (supracrustal) of the Chitradurga Schist Belt adjacent to the Chitradurga Shear Zone (CSZ), western Dharwar craton, south India. The study involves  
15 the following steps: 1) analysing the internal magnetic fabric using anisotropy of magnetic susceptibility (AMS) studies, and strength of the host metabasalts, 2) quantifying the fluid pressure condition through lower hemisphere equal area projection of pole to veins by determining the driving pressure ratio ( $R'$ ), stress ratio ( $\phi$ ), and susceptibility to fracturing, and 3) deciphering the paleostress condition using fault  
20 slip analysis. We interpret that the NNW-SSE to NW-SE (mean  $337^\circ/69^\circ$  NE) oriented magnetic fabric in the rocks of the region developed during regional D1/D2 deformation on account of NE-SW shortening. However, D3 deformation manifested by NW-SE to E-W shortening led to the sinistral movement along

CSZ. As a consequence of this sinistral shearing, fractures with prominent orientations formed riedel shear components, with CSZ as the shear boundary. Subsequently, all the pre-existing fabrics along with the riedel shear components were reactivated and vein emplacement took place through episodic fluid pressure fluctuation from high to low  $P_f$  at shallow depth (~2.4 Km). However, NNW-SSE orientations were susceptible to reactivate under both high and low  $P_f$  conditions thereby attaining maximum vein thickness along these orientations. The deduced paleostress from fault-slip analysis, along with the kinematics of the fractures and veins are in good agreement with previously estimated regional tectonics. Thus, integrating multiple domains of studies, help in the logical interpretation of fluid flow condition and vein emplacement mechanism in the study area that has not been ventured before.

## 1. Introduction

The upper crust is replete with fractures/faults, which act as pathways for fluid flow and vein emplacement. Fracture formation and vein emplacement mechanisms are closely interrelated, and require a detailed study for finding out potential hydrothermal deposits. Fracture formation and reactivation involves a combination of regional stress field (far field stress), stress ratio ( $\Phi$ ) and driving pressure ratio ( $R'$ ) that helps to determine the prevailing fluid pressure condition (Delaney et al., 1986; Jolly and Sanderson, 1997; McKeagney et al., 2004; Mazzarini and Isola, 2007; Martinez-Poza et al., 2016; Cucci et al., 2017). However, previous studies suggest that pre-existing anisotropy in host rocks play a significant role in formation and propagation of fractures provided the anisotropy is favorably oriented to the far field stresses (Ikari et al., 2015; Donath, 1961; Hoek, 1964; Attwell and Sandford, 1974). Presence

of such favorably oriented anisotropy lowers the shear strength of the host rocks, enabling failure/slip along them at minimum compressive stress, prior to/during vein emplacement. Such conduits are reactivated at both high and low fluid pressures forming pathways for fluid flow (Mondal and Mamtani 2013, Lahiri and Mamtani, 2016). Thus, vein emplacement mechanism requires favorably oriented fractures that can be reactivated, when fluid pressure gradually builds and exceeds the normal stresses acting on the fracture wall (Gudmundsson, 2011). This enables the fractures to open up, under cyclic increase of fluid pressure along such pre-existing conduits, a mechanism known as *fault-valve action* (Sibson et al., 1988; Sibson, 1992, 1996, 2000; Boullier and Robert, 1992; Sibson and Scott, 1998; Petit et al., 1999; Cox et al., 2001 among others). Subsequently, fluid flows into the fractures, a phenomenon analogous to *burping*, triggering an immediate drop in fluid pressure. This sudden drop in fluid pressure is responsible for mineral deposition and vein formation (Cox et al., 1991, Cox, 1995; 2001). Vein materials thus deposited seals the fracture/fault planes, preparing the system for the next cycle of fluid pressure build up, rupture, fluid flow and vein formation (Mondal and Mamtani, 2013; Lahiri and Mamtani, 2016; Marchesini et al., 2019). Thus, repeated cycles of elevated and depleted fluid pressure generate widespread networking of veins in host rocks. Studies conducted in this approach help to understand the mechanism of fabric development and fracture formation vis-à-vis vein emplacement. This study also aims to provide a detailed insight about the development of brittle structures and their role in understanding the tectonic evolution of Archean cratons.

60 The Chitradurga Schist belt (western Dharwar craton, south India), is a NW-SE trending Archean greenstone belt, known to harbor a widespread network of veins with potential epigenetic gold bearing lodes (Gupta et al., 2014; Gopalakrishna et al., 2018). We have conducted this study in the meta-volcanic

(metabasalt), hosting quartz veins of variable orientations and thicknesses, in and around Chitradurga region. We emphasize on understanding the mechanism of vein emplacement under a tectonic environment where propensity of fracture reactivation for vein emplacement is mutually dependent on both fluid pressure condition and the regional far field stresses. The present paper is a comprehensive work which quantifies the fabric in the visually isotropic metabasalts of the Chitradurga Greenstone belt and its role in fracture **formation** and in channelizing upper crustal fluids. We also found the **potential reactivation ability** of the fractures/faults at variable fluid pressure conditions to investigate the role of fractures/faults which are not favorably oriented to the pre-existing anisotropy and regional stress field, and their contribution **for** vein emplacement in the study area.

## 2. Geology of the study area

The study has been conducted in the Chitradurga Greenstone Belt of Dharwar craton (southern India; Fig. 1a), which represents a complex geological history. Dharwar craton exposes >3.0 Ga, Archean continental crusts, represented by the TTG (trondjemite-tonalite-granodiorite gneiss) also known as the peninsular gneiss (Jayananda et al., 2006). The craton stabilized during the accretion of Eastern Dharwar Craton (EDC) and Western Dharwar Craton (WDC) at 2.75-2.51 Ga. The zone of accretion of the two tectonic blocks is marked by a shear zone, referred to as Chitradurga Shear Zone (e.g., Naqvi and Rogers, 1987; Chadwick et al., 2003; Jayananda et al. 2006). The eastern part of WDC marked by Chitradurga Schist Belt (CSB) (Fig. 1b), comprises of peninsular gneiss (3.4-3.0 Ga), and younger supracrustal rocks (Beckinsale et al., 1980; Sarma et al., 2011; Taylor et al., 1984). The latter comprises metavolcanics/metabasalts (greenstone belt; greenschist/lower amphibolite facies metamorphism),

metamorphosed greywacke-argillite (interbanded with ferruginous chert and banded iron formation),  
85 polymict conglomerate, and ferruginous chert. **The basaltic rocks** of the schist belt have an island arc  
affinity (Chakrabarti et al., 2006). Regionally, the belt is surrounded by older peninsular gneisses and  
younger granitoids (Ramakrishna and Vaidyanadhan, 2010). Previous geological investigations, revealed  
the presence of various younger granites ~2.6 Ga; (Jayananda et al., 2006; Chardon et al., 2002) which  
are associated with the schist belt, such as Chitradurga granite, J. N. Kote granite. The structural  
90 investigations of the adjacent meta-sedimentary rocks of the region show that the area has undergone  
three phases of deformation: D1, D2 and D3 respectively (Chadwick et al., 1989; Jayananda et al., 2006;  
Mondal and Mamtani, 2014). The D1/D2 deformations are coaxial with NW-SE striking axial plane. D1  
folds are tight to isoclinal and asymmetric while the D2 folds are open to tight and upright. Earlier studies  
revealed that the folds (regional open; NE-SW striking vertical axial plane), related to early D3  
95 deformation superposed the D1/D2 structures thus resulting in dome-basin geometry in the meta-  
sedimentary rocks of that region (Chakrabarti et al., 2006; Mondal and Mamtani, 2014). However, later  
phase of D3 deformation led to the formation of brittle structures in the younger granites of CSB (Mondal  
and Mamtani, 2016). The NE-SW shortening during D1/D2 deformation was responsible for NW-SE  
oriented structural elements in the CSB (Chadwick et al., 2003), while NW-SE to E-W shortening  
100 direction prevailed during D3 deformation (Jayananda et al., 2006; Mondal, 2018; Mondal and Acharyya,  
2018). A number of petrological and geochemical investigations have been carried out on the area in the  
past. These studies suggest the presence of actinolite, albite, chlorite, epidote, quartz and calcite in the  
basaltic rocks (Chakrabarti et al., 2006). The CSB holds evidence of progressive metamorphism with a  
gradual increase in P-T conditions (approximately 3-4 Kb of pressure), from greenschist facies **in the**

105 north to amphibolite facies in the south suggesting the depth of deformation to be 10-12 Km (during D1/D2). However, recent studies by Acharyya and Mondal, (2019) shows that the brittle deformation during late D3 took place at a shallow depth of ~2-4 km. The dashed rectangular area in figure 1b demarcates the study area. The present study focuses on the dark greyish to blackish, massive to fine grained, altered metabasalt hosting quartz veins, of the region surrounded by meta-sedimentary sequences  
110 (Fig. 2). The metabasalts of the study area are devoid of any well-defined mesoscopic foliation. The foliation planes in the adjacent meta-sedimentary rocks of the region are found to be NW-SE oriented (mean strike/dip is 323°/71° NE). Since, the metabasalts are devoid of mesoscopic field fabric, the anisotropy of magnetic susceptibility (AMS) study has been conducted in order to quantify their internal magnetic fabric.

115

### 3. Overview of brittle structures

Metabasalts of the study area are replete with fractures and faults of multiple orientations. Some of these fractures and faults are found to host quartz veins forming criss-cross pattern (Fig. 3a and c) and some of them are devoid of any vein material. Growth of vein materials (quartz crystals) are often found to be  
120 perpendicular to the vein wall suggesting that dilation was significant in these veins (Fig.3b). The maximum width and length of the quartz veins are ~1 meter and ~130 meters respectively. Some of the quartz veins are found to be displaced by other ones forming crisscross network of veins in the study area (Fig. 3c). Wing cracks filled up with quartz veins are also observed (Fig. 3d). At few places, the thicker quartz veins show series of successive fault planes with slickenside lineations on them (Fig. 3e and in-  
125 set). Those veins are often found to enclose angular metabasalt enclaves (host material, Fig. 3f),

suggesting that fault valve action was predominant in the region. Quartz veins are recorded with multiple median lines as an evidence of crack-seal mechanism due to cyclic fluid ingress (Fig. 3g). Some of the fault planes recorded in the study area have shallow plunging slickenside lineations while others have moderate to high plunge. Both Left Lateral Faults (LLF's) and Right Lateral Faults (RLF's) are recorded based on the movement of the hanging wall with respect to the footwall. Presence of congruous steps helps in the discrete identification of the fault planes (Fig. 3h). Although, quartz veins of variable orientations and thicknesses are found, however, most of the veins are predominantly NNW-SSE trending (Fig. 4a). Most of the fractures and faults show NNW-SSE trend (maxima) whereas some others form a WNW-ESE to NE-SW sub maxima respectively (Fig. 4b & 4c). Some WNW-ESE trending Mode-I (tensional) cracks with prominent tips are also recorded from the study area which are often filled up with quartz veins. It may be noted that the veins with maximum thicknesses are oriented along NNW-SSE direction (Fig. 4d). In this study, 992 fracture data (strike/dip), 378 vein data (strike, dip and thickness) and 73 fault data (strike, dip and slip) have been measured from the metabasalt exposures at about 13 locations in the entire study area. All 73 shallow to moderately plunging fault planes are used to decipher the paleostress condition. Thus, the strict spatial and temporal relationships between brittle structures and veins provided by field observations help to quantify the fluid pressure conditions that prevailed during fracture reactivation and vein emplacement. It also provides information for reconstructing the tectonic stress conditions of the craton.

#### 145 4. Methods of analysis and results

The quartz veins have been emplaced in the massive metabasalts of the region that are devoid of any prominent mesoscopic foliation as mentioned above. At places, veins of one orientation are dissected and sometimes displaced by others that led to the formation of mesh like structures (Fig. 3a, c). Sibson (1992) has mentioned that such a mesh is formed when a rock contains fractures of variable orientations that may get reactivated due to rise in fluid pressure. It is mentioned that in the Chitradurga region, veins of various orientations show mutually cross-cutting relationships, which implies repeated cycles of vein emplacement (see section 3; Mondal and Mamtani, 2013; Sibson, 1992). Although veins have various orientations, NNW-SSE striking veins are the most common (Fig. 4a). It may be noted that NW-SE to NNW-SSE direction, which defines the maximum strike orientation of quartz veins, fractures and faults, is also the orientation of the adjacent Chitradurga shear zone, the overall trend of the schist belt. This implies that there is a strong structural control on the formation of these quartz veins. It has been shown earlier that the pre-existing anisotropy plays a critical role in propagating fractures and channelizing fluid in rocks (Sanderson and Zhang, 1999, Cox et al., 2001 and Ikari et al., 2015). It is also known from rock mechanics investigations that the rock strength variation controls the strain partitioning and influences fluid flow (e.g., Tsidzi, 1990; Vishnu et al 2018). Therefore, it is crucial to determine the orientation of anisotropy in host rocks and the state of stresses governing the upper crustal fluid flow vis-à-vis vein emplacement.

3-D Mohr circle analysis has been performed by Jolly and Sanderson (1997) using dyke orientation data to examine the magma pressure condition that was responsible for the opening of pre-existing fractures during dyke emplacement. Further the work has been extended to understand the fluid pressure condition and vein emplacement mechanism by McKeagney et al., 2004 (also see Yamaji et al., 2010). We present

vein orientation data from the Chitradurga region (southern India), which is a province of epigenetic gold deposit (Gupta et al., 2014;Gopalakrishna et al., 2018).

We conduct anisotropy of magnetic susceptibility (AMS) study, followed by Brazilian Tensile strength (BTS) determination in order to quantify the internal magnetic fabric and tensile strength of the rocks within the study area. The 3-D Mohr circle construction using quartz vein orientation data helps to recognise the fluid pressure conditions under which they were emplaced. Further, these fluid pressure conditions were integrated with dilation tendency, slip tendency and fracture susceptibility in order to understand the mechanism of vein emplacement in the Chitradurga region. A combination of data obtained from these methods along with paleostress analysis using fault-slip data recorded from the study area provides a comprehensive evaluation of vein forming conditions in Chitradurga greenstone belt.

#### ***4.1 Anisotropy of Magnetic Susceptibility (AMS)***

The quartz veins occur in massive metabasalt, which does not show any visible field foliation. However, such visibly massive rocks may preserve an internal fabric, which can be recognized through anisotropy of magnetic susceptibility (AMS) studies (e.g., Tarling and Hrouda, 1993; Maffione et al., 2015; Mamtani and Greiling, 2005; Raposo et al., 2007; Looock et al., 2008; Mondal and Mamtani, 2014; Mondal, 2018). The study involves preparation of cylindrical core samples (25.4 mm diameter × 22 mm height) from oriented metabasalt blocks. These metabasalt block samples have been collected from 13 different locations (Fig. 5) in the study area. The prepared core samples are subjected to an external magnetic field and the induced magnetization for each core sample is measured in different directions. AMS is considered to be a symmetric second-rank tensor, represented by an ellipsoid with three mutually

perpendicular **principal** axes,  $K_1$ ,  $K_2$  and  $K_3$  respectively where ( $K_1 \geq K_2 \geq K_3$ ). The orientation and magnitude of each of these principal axes are determined in this analysis, where  $K_1$  represents the magnetic lineation,  $K_3$  is the pole to the magnetic foliation ( $K_1K_2$ ). Using the magnitudes of  $K_1$ ,  $K_2$  and  $K_3$ , several AMS parameters are calculated such as magnetic susceptibility ( $K_m$ ), magnitude of the magnetic foliation (F) and magnetic lineation (L), degree of magnetic anisotropy ( $P_j$  or  $P'$ ) and shape parameter (T). The formulae for the parameters are given below (after Tarling and Hrouda, 1993; Jelínek, 1981):

$$K_m = (K_1 + K_2 + K_3) / 3, \quad (1)$$

$$F = (K_2 - K_3) / K_m, \quad (2)$$

$$L = (K_1 - K_2) / K_m, \quad (3)$$

$$P_j = \exp \sqrt{2 [( \eta_1 - \eta_m )^2 + ( \eta_2 - \eta_m )^2 + ( \eta_3 - \eta_m )^2 ]}, \quad (4)$$

$$T = (2\eta_2 - \eta_1 - \eta_3) / (\eta_1 - \eta_3), \quad (5)$$

Here,  $\eta_1 = \ln K_1$ ,  $\eta_2 = \ln K_2$ ,  $\eta_3 = \ln K_3$  and  $\eta_m = (\eta_1 \cdot \eta_2 \cdot \eta_3)^{1/3}$ . In the above equations,  $P_j$  and T give the measure of the eccentricity and shape of the AMS ellipsoid respectively. The value of T ranges between -1 to +1. Positive and negative values represent oblate and prolate shapes of the AMS ellipsoid (Tarling and Hrouda, 1993).

AMS measurements have been conducted in the spinner mode (field intensity of  $300 \text{ Am}^{-1}$ ) using the KLY-4S Kappabridge (AGICO, Czech Republic); see Hrouda et al. (2006) for instrument details. The SUFAR program has been used to calculate the required AMS parameters described above for each sample. A total of 65 cores are prepared and analysed. The mean value of the AMS parameters at each

location is calculated using the program Anisoft (version 4.2, AGICO; Jelinek statistics, Jelinek, 1981). It is noted that the  $K_m$  varies between 37 and  $1280 \times 10^{-6}$  SI units, with most of the samples having  $K_m$  below 210  $1000 \times 10^{-6}$  SI units (see supplementary sheet-1). This indicates that paramagnetic and ferromagnetic minerals contribute significantly to the AMS. Petrographic studies reveal the presence of actinolite, hornblende, chlorite, albite, epidote, pyrite which are inferred to contribute to the susceptibilities recorded in the samples. The  $P_j$  lies between 1.003 and 1.539, and the shape of the AMS ellipsoid is dominantly oblate (positive  $T$  values; see supplementary sheet-1). It is noted that the magnetic foliation ( $K_1K_2$  plane) 215 is consistently NW-SE striking (mean orientation:  $337^\circ/69^\circ$  towards NE; Fig. 5a). The magnetic lineations plunge variably from NNW, through sub-vertical to SSE (Fig. 5b).

#### ***4.2 Tensile strength determination***

It is known that the dilation occurs in a direction parallel to the minimum compressive principal 220 stress ( $\sigma_3$ ), when the fluid pressure ( $P_f$ ) exceeds the normal stress acting on the fracture wall. Therefore, the fractures may occur at any depth when the effective stress ( $\sigma_3 - P_f$ ) is sufficient to counteract the tensile strength ( $T$ ) of the rocks (Gudmundsson, 2011). We measure the tensile strength of the host metabasalts to quantify the  $P_f$  that prevailed during vein emplacement in the region. The measurement of direct tensile strength requires machined specimens and also involves 225 difficulty in applying tensile load on the cylindrical specimen during analysis. Therefore, tensile strength measurement of rocks using Brazilian test has become imperative in rock mechanics. Compression-induced extensional fractures are generated in the test which essentially involves line-loading on a circular disk placed between two platens (Aydin and Basu, 2006; Basu et al., 2013; see

Fig.6). This tensile strength (T) is estimated from the elastic theory (ISRM, 1978; ASTM D3967,  
230 2001):

$$T = \frac{2P}{\pi LD} \quad (6)$$

Here, P is peak/failure load, and L and D, are the length and diameter of the disk respectively. For this  
analysis, 18 core samples were drilled from metabasalt blocks which were later resized to obtain the  
desirable cores for the analysis (length: diameter = 1: 2). The maximum tensile strength of each specimen  
235 at the instance of failure is recorded. The maximum tensile strength for 18 samples are averaged out to  
obtain the approximate tensile strength of metabasalts, which is ~12 MPa. This tensile strength value is  
further used for quantifying the  $P_f$  in 3D Mohr circle using vein orientation data.

#### ***4.3 Fluid pressure determination***

240 Here, we have used the method proposed by Jolly and Sanderson (1997) to quantify the  $P_f$  conditions that  
led to the vein emplacement in metabasalts of Chitradurga region (southern India). We have used the  
lower hemisphere equal area projection of the poles to quartz vein data. According to Jolly and Sanderson  
(1997), girdle distribution of vein pole data implies  $P_f > \sigma_2$ , described as a condition where large number  
of fracture orientations are susceptible to reactivate, while  $P_f < \sigma_2$ , represents clustered distribution of vein  
245 pole data, where only limited range of fracture orientations reactivate. Depending on the type of  
distribution (girdle/cluster), parameters such as stress ratio ( $\phi$ ) and driving pressure ratio ( $R'$ ) are

calculated using ranges of fracture orientations ( $\theta_1$ ,  $\theta_2$  and  $\theta_3$ ) from the following equations provided by Jolly and Sanderson, 1997 and Baer et al., 1994.

$$R' = \frac{P_f - \sigma_3}{\sigma_1 - \sigma_3} = \frac{1 + \cos 2\theta_2}{2}. \quad (7)$$

250 For  $P_f > \sigma_2$ ,

$$\Phi = \frac{\sigma_2 - \sigma_3}{\sigma_1 - \sigma_3} = 1 - \frac{1 - \cos 2\theta_2}{1 - \cos 2\theta_3}. \quad (8)$$

For  $P_f < \sigma_2$ ,

$$\Phi = \frac{\sigma_2 - \sigma_3}{\sigma_1 - \sigma_3} = \frac{1 + \cos 2\theta_2}{1 + \cos 2\theta_1}. \quad (9)$$

In figure 7a, the lower hemisphere equal area projection of pole to vein data shows girdle distribution, implying high fluid pressure condition ( $P_f > \sigma_2$ ). From this distribution the orientations of the principal stress axes ( $\sigma_1$ ,  $\sigma_2$  and  $\sigma_3$ ) are determined using the Bingham statistics of the Stereonet 9 software (<http://www.geo.cornell.edu/geology/faculty/RWA/programs/stereonet.html>).  $\sigma_1$  is sub-vertical lying in the empty space devoid of any vein pole data. Subsequently, following Jolly and Sanderson (1997), the planes  $\sigma_1\sigma_2$ ,  $\sigma_1\sigma_3$  and  $\sigma_2\sigma_3$  are constructed and the range of fracture orientations,  $\theta_2$  and  $\theta_3$  are determined along the  $\sigma_1\sigma_3$  and  $\sigma_1\sigma_2$  planes respectively. For this high  $P_f$  condition,  $\theta_2 = 27^\circ$ ,  $\theta_3 = 59^\circ$  from which  $\phi = 0.72$  and  $R' = 0.8$  are calculated. Thus, such a  $P_f$  condition enhances the chances of vein emplacement along various orientations. Although pole to vein data represents a girdle distribution pattern, however the distribution of the data points shows three prominent clusters. The WSW cluster forms the highest density cluster around the  $\sigma_3$  axis indicating a number of veins with similar orientations. The cluster primarily represents the NNW-SSE to NW-SE trending veins, attaining the maximum thickness. It is to

be noted that majority of the veins are oriented along this direction (see Fig. 4a), which also suggests that the vein forming fluid must have been channelized through a pre-existing anisotropy (trending NW-SE to NNW-SSE, see section 4.1). These data are segregated and plotted separately in the lower hemisphere equal area projection and thus the obtained contour defines the WSW cluster (Fig. 7c). We decided to  
270 extend the contour interval beyond the data points in order to incorporate the maximum range of vein orientations ( $\theta$ ) lying parallel/sub-parallel to the internal anisotropy (as evident from the anisotropy of magnetic susceptibility study) of the host rock. Also, the contour interval and significance level for each of the clusters were selected in such a way that maximum number of data points are included, in order to obtain a statistically viable data cluster. **Apart from the WSW cluster (with a high cluster density), the SE  
275 and the NE clusters (with higher data spreading) are also evaluated discretely (Fig. 8).** The number of obtained clusters can also be testified through mixed Bingham analysis using  $K$  vs BIC (i.e., the number of Bingham component of a mixed Bingham distribution vs Bayesian information criterion; Yamaji and Sato, 2011). We have found that the lowest BIC values are obtained when  $K=3$  (number of possible clusters for the given data set), thereby, justifying the selection of the three clusters for the analysis. It is  
280 however difficult to quantify the lowest  $P_f$  value; we therefore intend to use the obtained  $P_f$  values from the respective clusters as examples of low  $P_f$  conditions denoting  $P_f$  fluctuation rather than quantifying the lowest  $P_f$  condition of the study area. The orientations of the principal stress axes ( $\sigma_1$ ,  $\sigma_2$  and  $\sigma_3$ ) are determined using the Bingham statistics of the Stereonet 9 software. Similarly, the  $\sigma_1\sigma_2$ ,  $\sigma_1\sigma_3$  and  $\sigma_2\sigma_3$  planes are constructed and the range of fracture orientations  $\theta_1$  and  $\theta_2$ , are calculated along the  $\sigma_2\sigma_3$  and  
285  $\sigma_1\sigma_3$  planes. Again, for low  $P_f$  conditions,  $\theta_1 = 38^\circ$ ,  $\theta_2 = 43.2^\circ$ ,  $\phi = 0.85$  and  $R' = 0.53$  (WSW cluster, Fig. 7c);  $\theta_1 = 32^\circ$ ,  $\theta_2 = 35^\circ$ ,  $\phi = 0.93$  and  $R' = 0.67$  (NE cluster, Fig. 8a);  $\theta_1 = 30^\circ$ ,  $\theta_2 = 44^\circ$ ,  $\phi = 0.69$  and  $R' =$

0.52 (SE cluster, Fig. 8c) This implies that under low fluid pressure condition ( $P_f < \sigma_2$ ) only limited range of fracture orientations are susceptible to reactivate. In each case, for determining the absolute  $P_f$  magnitude, pole to vein data are plotted in 3D Mohr circles. Recent studies by Mondal and Acharyya (2018), documented the depth of faulting and fracturing in close vicinity of the study area (Chitradurga granite) to be ~2.4km. Similar depth of fracturing has also been reported by Acharyya and Mondal (2019), from the elliptical clasts of conglomerate bed within the Chitradurga schist belt. We have considered this depth for determining the magnitude of the maximum compressive stress ( $\sigma_1$ ) during fracture formation, using  $\sigma_1 = h\rho g$ , where  $h$  = depth of fracturing in metabasalts (~2.4km),  $\rho$  = approximate bulk density of crust (2700 kg/m<sup>3</sup>),  $g$  = 9.8 m/sec<sup>2</sup>. Therefore,  $\sigma_1 \geq 63.5$  MPa, when depth of fracturing in metabasalt ~2.4km. The magnitude of the minimum compressive stress ( $\sigma_3$ ) can be estimated using the stress intensity factor (fracture toughness),  $K_{IC}$ .  $K_{IC}$  for mode-I fractures, can be determined using the following equation (Lawn and Wilshaw, 1975; Atkinson, 1989).

$$K_{IC} = \sigma_3 Y(\pi a)^{1/2},$$

where  $\sigma_3$  is the minimum far field stress acting normal to the crack,  $Y$  is a numerical modification factor to account for the crack geometry and edge effect, and 'a' is the crack half-length. Microstructural investigation suggests that the metabasalts are generally fine to medium grained, massive and consists of plagioclase laths and altered pyroxene with minor chlorite, ankerite and quartz (Gupta et al., 2014). Within the metabasalts, variably oriented larger phenocryst grains range from 2 mm to 8 mm in length. It is evident that the large phenocrysts were more susceptible to generate the initial fractures as compared to the medium to fine grained groundmass in these types of rocks. Therefore, we have considered that the length of the initial crack ranges from 2 mm to 8 mm (Brace, 1964).  $K_{IC}$  value for the metabasalts are

found to be 1.069 (Donovan, 2003). Considering this  $K_{IC}$  value and the range for fracture half-length ( $a$ ), the  $\sigma_3 Y$  value for the metabasalt ranges between 9-19 MPa. From this strength range we considered that  
310 12 MPa is a reasonable estimate for the in situ tensile strength (similar to the laboratory strength) of the corresponding lithology. It may also be noted that, previous studies by Mondal and Acharyya, 2018, conducted in Chitradurga Granite, in close vicinity of the study area also regarded the magnitude of  $\sigma_3$  ~10 MPa, to be a good estimation. Combining these estimations with the results obtained from the present studies, we constrained the value for  $\sigma_3$  ~12MPa.

315 In each case we consider the limiting values for both maximum and minimum compressive stresses. Magnitude for the intermediate compressive stress ( $\sigma_2$ ) is determined using the respective stress ratios ( $\phi$ ) for both high and low  $P_f$  respectively. Following Jolly and Sanderson (1997), the 3D Mohr circles are constructed using the above magnitudes of principal stresses and the angles determining the range of fracture orientations. Thus, well-defined  $P_f$  lines for both high  $P_f$  ( $P_f=53.2$  MPa; Fig. 7b) and low  $P_f$   
320 conditions ( $P_f=39.3$  MPa; Fig. 7d,  $P_f=46.5$ , Fig. 8b and  $P_f=38.8$ , Fig. 8d) are obtained using the Fractend code (Github, 2017).

#### 4.4 Dilation tendency, slip tendency and fracture susceptibility

Dilation tendency ( $T_d$ ) and slip tendency ( $T_s$ ) determine the propensity of any fracture orientation to  
325 reactivate through dilation or shearing, under a certain state of stress condition (Mazzarini et al., 2019). While, high dilation tendency ensures reactivation through dilation, high slip tendency elevates chances of opening through shearing (Ferrill et al., 1999). According to, Stephens et al., 2017, fracture planes suffer dilation when the difference between  $\sigma_1$  and the normal stress acting on

the plane is close enough to the magnitude of differential stress ( $\sigma_D = \sigma_1 - \sigma_3$ ) and  $T_d = (\sigma_1 - \sigma_n) / \sigma_D$ . Slip  
330 tendency is denoted by the ratio of shear stress ( $\sigma_s$ ) to normal stress ( $\sigma_n$ ); ( $T_s = \sigma_s / \sigma_n$ ) and also depends on  
the frictional characteristics of the rock (Morris et al., 1996), along with the fracture plane orientation.  
Under a particular state of stress condition if the ratio of shear stress to normal stress is significantly large,  
then that particular fracture orientation is susceptible to reactivate. Fracture susceptibility ( $S_f$ ), is defined  
as the variation of fluid pressure ( $\Delta P_f$ ) within a fracture plane that can lead to fluid induced shear  
335 reactivation (Mildren et al., 2002; Stephen et al., 2017). Such reactivations depend on the shear and  
normal stresses acting on the fracture plane, along with the cohesion ( $= 0$  in this case) and the static  
coefficient of friction ( $\mu_s$ );  $S_f = \sigma_n - (\sigma_s / \mu_s)$ .

Lower hemisphere equal area projections (see Fig. 9) of the poles to fracture (vein-filled) data help us to  
understand the variation in dilation tendency, slip tendency and susceptibility of fractures with respect to  
340 their orientations, under both high and low  $P_f$  conditions. The diagrams are prepared using Fractend code  
(Github, 2017). It is evident from figure 9a, that the dilation tendency is high for the fracture orientations  
which are at a high angle to the  $\sigma_3$  axis, i.e., pole to these fractures form a well-defined cluster around  $\sigma_3$ .  
These fractures show greater tendency towards dilational opening for both high and low  $P_f$  (Fig. 9a and  
b). For fracture orientations having higher slip tendency, i.e., susceptible to shear opening, pole to the  
345 fractures are at a low angle to the  $\sigma_3$  axis. The fracture planes are therefore oriented at an angle to the  
maximum compressive stress axis  $\sigma_1$ , condition favorable for shear reactivation for both, high and low  $P_f$   
(Fig. 9c and d). Fracture susceptibility, which involves variation in fluid pressure ( $\Delta P_f$ ) is low for the

fracture orientations having high dilation and slip tendencies, which indicates fluid-induced fracture reactivation in metabasalts (Fig. 9e and f) in the study area.

350

#### ***4.5 Paleostress analysis***

The shallow to moderately plunging normal faults of the study area with prominent slip directions are used to determine the stress regime under which these fractures and faults were formed and reactivated. Fault-slip data (orientations of fault planes and slip directions) recorded from the field were used for paleostress determination. Several methods are proposed for paleostress analyses using fault-slip data (e.g. Angelier, 1994; Dupin et al., 1993; Etchecopar et al., 1981; Gapais et al., 2000; Marrett and Allmendinger, 1990; Ramsay and Lisle, 2000; Twiss and Unruh, 1998; Yamaji, 2000; Žalohar and Vrabec, 2007 and the references therein). Since, the fault-slip analysis methods are well established, here we prefer to represent only the salient aspects. The fault-slip analysis can be divided into two categories based on whether the fault-slip data are viewed as representing kinematic or dynamic information (Blenkinsop 2006; Gapais et al. 2000; Twiss & Unruh, 1998) based on the following assumptions: (1) bulk state of stress is uniform and movement on the fault planes are independent of each other; (2) slip on the fault plane occurs along the direction of the maximum resolved shear stress under a given stress state (Wallace–Bott Hypothesis); (3) faults are homogeneous and a part of the same tectonic event (Angelier, 1994; Gapais et al., 2000; Gephart & Forsyth, 1984; Twiss & Unruh, 1998). In this study, we have determined the paleostress direction, using fault-slip data measured from 73 shallow to moderately **dipping** normal faults (spatially distributed) in the metabasalt by Right Dihedron method. Since, some of the fault planes show variation in their strike orientations, the small amount of inhomogeneity in the data set is reduced in this process.

360

365

Thus, the data sets are segregated methodically into homogeneous data subsets using the ‘Win\_Tensor’  
370 software program (version 5.8.6; Delvaux and Sperner, 2003; Delvaux, 2011). In the present analysis all  
the collected data are represented in a single set and separation is done by using the Right Dihedron  
method without any sort of manual intervention. Following Delvaux and Sperner, 2003, the data is filtered  
on the basis of stress ratio (R), orientation of the stress axes and symmetry of the measured sets. Out of  
73 data, 30 data are accepted with a low value of counting deviation and nominal counting values of 0  
375 and 100 for  $\sigma_1$  and  $\sigma_3$  respectively. Thus, the best fitted reduced stress tensor is obtained for the accepted  
data subset (30 out of 73 fault data; see Fig. 10) at a “C” quality ranking. It also provides the relative  
orientations of the principal stress axes, stress ratio (R= 0.72) and stress regime index (R’=1.25). The  
NNE-SSW directed extension direction obtained from this paleostress analysis (see Fig. 10) coincides  
well with the regional D3 extension direction. Data rejected in this process to obtain the best fit stress  
380 tensor when treated separately yields, NNE-SSW oriented extension direction with small variations in the  
R and R’ values. According to Delvaux and Sperner (2003), the obtained stress regime index indicates a  
pure-strike slip domain which is in a good agreement with sinistral shearing along CSZ.

## 5 Discussions

### 385 *5.1 Fabric development vs. regional tectonics*

It has been mentioned earlier that the metabasalts of the study area lack any distinct visible foliation.  
However, the AMS analysis suggests, a prominent NNW-SSE to NW-SE oriented magnetic fabric in  
metabasalts. This magnetic fabric also matches well with the field foliation of the meta-sedimentary  
sequences, surrounding the metabasalts and is also parallel to the regional trend of CSZ (see Fig. 2). This

390 implies that the fabric in metabasalts of the study area must have been controlled by the regional D1/D2  
deformation under NE-SW directed shortening that generated the field foliation in the meta-sedimentary  
sequences. The magnetic lineations in the metabasalts plunges due NNW through sub-vertical to SSE  
(Fig. 5b). In the previous studies, these variations in the plunge of the magnetic lineations had been  
interpreted as a consequence of superposed deformation (e.g., Mamtani and Sengupta, 2010; Mondal and  
395 Mamtani, 2013). Therefore, in the light of regional structural information and above discussions, it is  
inferred that these variations in magnetic lineations of the metabasalts are the manifestation of dome-  
basin geometry that were produced due to the superposition of D3 over D1/D2 regional deformation.  
Recently, Mondal (2018) documented similar results from the magnetic fabric analysis in adjacent  
Chitradurga granite. The studies by Bhatt et al., (2017); Mondal and Acharyya (2018), suggest that the  
400 D1/D2 deformation lasted between 2614 and 2555 Ma while the D3 deformation is approximated around  
~2537 Ma.

### ***5.2 Control of regional far-field stress on developing the brittle structures in Chitradurga region.***

The above discussions suggest that (a) NNW-SSE oriented magnetic foliation developed during D1/D2  
405 deformation under NE-SW directed shortening, (b) variation in plunge of magnetic lineation is a  
manifestation of dome-basin geometry on account of D3 deformation. It is argued that during D3  
deformation under NW-SE to E-W directed shortening the CSZ evolved as a sinistral shear zone (Mondal  
and Acharyya, 2018). It may be noted that the angle between the mean orientation of the schist belt and  
the compression direction for D3 deformation are found to be  $\sim 45^\circ$  which also supports the sinistral  
410 movement along Chitradurga shear boundary.

It is mentioned in sections 2 and 3 that, the study area is replete with a number of brittle structures such as fractures and faults. At places, these fractures are filled-up with quartz veins. Therefore, it is now essential to evaluate whether and how these brittle structures and their kinematics can be fitted to the regional far-field stresses responsible for deformation in the Chitradurga region. The quartz vein orientation data from northern part of the Chitradurga schist belt reveals that the vein emplacement took place during regional D3 deformation (Mondal and Mamtani, 2014). Recently, Mondal and Acharyya (2018), and Acharyya and Mondal (2019) suggested that the brittle structures (fractures/faults) in the Chitradurga granite (in close vicinity to the study area) are related to the D3 deformation. Moreover, paleostress investigation using fault-slip data also reveals NNE-SSW directed extension was dominant during D3 deformation. Apart from this, evidences of any later deformation, i.e., post D3 deformation, have not been recorded from the study area (from previous studies and as per our field observations). Thus, based on the present studies and above discussions, it is logically inferred that the formation of brittle structures (fractures/faults) as well as vein emplacement must be explained as a consequence of regional D3 deformation at a shallow depth. Mondal and Mamtani (2014) suggested that the Mulgund granite ( $2555 \pm 6$  Ma; Sarma et al., 2011), which lies in close vicinity to the metabasalts of the Chitradurga region, was emplaced syn-tectonically with CSZ. It is shown that the Mulgund granite underwent ductile deformation as it cooled and crystallized syn-tectonically with D3 deformation. The ductile deformation features in the granite were then superimposed by brittle structures during late D3, when the granite was fully solidified and had achieved a shallow depth (Mondal and Mamtani, 2016). Since both the lithologies (metabasalts and granite) have undergone same deformation and the fractures in them developed syntectonically with adjacent CSZ, it is favourable to interpret that the brittle structures, such as fractures

and faults in metabasalts developed on account of late D3 deformation under NW-SE to E-W directed compression at a shallow depth of about ~ 2.4 Km. For evaluating the nature (mode) of fracturing in the metabasalts of the study area, we used tensile strength (T) to be 12 MPa (from BTS studies). We estimated  $\Delta\sigma = 51.5$  MPa, which is greater than 4T, suggesting that the fractures in the metabasalts are not purely tensile, except for the cracks parallel to D3 shortening direction. However, the calculated value of  $\Delta\sigma$  is less than 5.7T, indicating that the normal stress on the fractures are not purely compressive also. Therefore, the value of  $\Delta\sigma$  satisfies  $4T < \sigma_1 - \sigma_3 < 5.7 T$  (Sibson, 2000), indicating that these fractures in the study area are extensional shear mode fractures.

440

### ***5.3 Regional tectonics and the mechanism of fracturing, faulting***

We discussed earlier in sect. -3, that the fractures and faults recorded in the metabasalts show a wide range of orientations with NNW-SSE maxima and WNW-ESE to NE-SW sub-maxima respectively (Fig. 4b). Among which, fractures trending along WNW-ESE to E-W are sub-parallel to the D3 (late phase) shortening direction. As previously mentioned in sect. -3, these WNW-ESE trending fractures have been regarded as tensional fractures. Similar orientations have also been recorded and interpreted as tensile fractures from the micro-granitoid enclaves in Chitradurga granite as evident from the studies of Mondal and Acharyya (2018). However, it is essential to explain the predominance of fractures and faults along the NNW-SSE orientation (forming the maxima). In order to explain this, we refer to the pre-existing fabric in the metabasalts of the study area, i.e., the NNW-SSE to NW-SE oriented magnetic fabric developed during D1/D2 regional deformation (Fig. 5a). Earlier studies suggest, that fractures are more likely to propagate along a pre-existing anisotropy, if and when the anisotropy is favorably oriented with

respect to the regional stress field (Ikari et al., 2015). The CSZ being a sinistral shear zone exhibits a pure strike slip stress regime, coeval with D3 deformation (NW-SE to E-W directed shortening).

455 In the later phase of D3 deformation, the favorably oriented NNW-SSE fabrics were reactivated under a **compatible** stress field, thereby, causing reactivation of the pre-existing fabrics in the metabasalts. This however, fails to justify the occurrence of the ~NW-SE and ~NE-SW oriented fractures within the metabasalts. From field investigations, the NE-SW oriented fractures show dextral movements, while the NNW-SSE and NW-SE oriented fractures are recorded with sinistral movements respectively (see Fig. 460 3c and d). However, the fracture disposition and consistency in their respective orientations indicate that all of these fractures **are coeval**, related to the same deformational event and have been reactivated under similar stress conditions. Moreover, any other brittle deformational event post D3, have not been recorded from the study area as mentioned earlier. Therefore, we need to explain the occurrence of such variably oriented fractures/faults within a single kinematic framework. The NNW-SSE and NW-SE orientations 465 are most likely to be sinistral, whereas NE-SW orientations form the dextral shear components respectively, considering CSZ to be the sinistral shear boundary. (See Fig. 11b and 4b, c). Thus, the NNW-SSE to NW-SE (P, Y and R) and the NE-SW to ENE-WSW (X and R') fractures coincide with the shear components of a riedel shear system considering the angle of internal friction ( $\Phi$ ) in metabasalts of the study area to be  $\sim 30^\circ$ . It may be noted that the value of  $\Phi$  is approximated from the Uniaxial 470 Compressive Strength (UCS) studies of the metabasalts core samples following Sivakugan et al., 2014.

These fracture planes (P, Y, R, X and R') acted as pathways for fluid flow and vein emplacement during the late D3 deformation.

#### *5.4 Understanding the mechanism of fluid flow and vein emplacement in Chitradurga region*

475 In order to explain the vein emplacement mechanism along these weak zones we need to consider the fluid pressure conditions **that prevailed at the time of vein emplacement**. In sect. 4.4, we mentioned earlier that the lower hemisphere equal area projection of pole to vein data shows girdle distribution, indicating high  $P_f$  condition ( $\sim 53.2$  MPa;  $> \sigma_2$ ) in the study area. Under such high  $P_f$  conditions, veins were emplaced along all possible orientations, including NNW-SSE, NW-SE, WNW-ESE, NE-SW and ENE-WSW  
480 trending fractures. The building fluid pressure surpassed the normal stresses acting on the fracture wall, and hence, fluid burped into the weak planes leading to fluid-induced reactivation of the fractures, promoting vein emplacement along them (Fig. 11c). However, the NNW-SSE trending veins show greater thickness and abundance with respect to other orientations (see Fig. 4d and 3e). We found that pole to these veins (NNW-SSE trending orientations) lie within the warm zones of the stereoplots obtained from  
485 the Fractend code (Github, 2017), indicating higher dilation tendencies (see Fig. 9a and 9b). In Fig. 9a and 9b, it is perceived that pole to these orientations form a cluster around the  $\sigma_3$  stress axis. Similarly, pole to the orientations trending NW-SE and NE-SW respectively, have higher slip tendencies, indicating shear reactivation along them (Fig. 9c and d). Most of the poles to the NNW-SSE trending orientations  
490 lie within the overlapping warm zones, indicating high potential to reactivate in both shear and dilation mode (high slip and dilation tendencies). These orientations are susceptible for dilation as well as shear reactivation, depending on the predominance of fluid pressure and availability of fluid for inducing

dilation or shear reactivation. Availability of fluid significantly reduces the effective normal stress ( $\sigma_n' = \sigma_n - P_f$ ); where  $\sigma_n'$  and  $\sigma_n$  are the effective normal stress and normal stress respectively) acting on the fracture planes (see Fig. 11c and d). Under this condition, fluid burps into the fractures leading to a significant drop in the fluid pressure. This sudden drop in the fluid pressure reduces the solubility of fluid materials promoting deposition of vein followed by sealing of the pathways (Cox et al., 1991, 2001). Pathways thus remains sealed till the next cycle of fluid pressure build up, fracture reactivation and vein deposition. However, in between two such cycles of high fluid pressure, intermediate low fluid pressure cycles persist, during which fluid pressure might not be substantially high to reactivate all the pre-existing pathways. Thus, a high fluid pressure cycle is followed by subsequent pulses of low fluid pressure causing selective reactivation of some of the pre-existing orientations. The orientations represented by the individual clusters (WSW, NE and SE) can be reactivated selectively depending on their respective orientation to the tectonic stress field and the fluid pressure magnitude. Hence, it is envisaged that, these cycles of high and low  $P_f$  conditions might have been repeated multiple times (n-times) until the fluid source was completely exhausted. In the present case, the low  $P_f$  conditions are interpreted by discretely analyzing the clusters within the girdle distribution of pole to vein data in the lower hemisphere equal area projection. (see sect. 4.4). Such low  $P_f$  could not surpass the normal stresses of all the pre-existing pathways. And only the NNW-SSE oriented ones (related to the magnetic fabric of the metabasalts) having high dilation and slip tendencies were favorably oriented for reactivation and fluid flow. The above mechanism of *fault-valve action* is supported with field evidences showing occurrences of thick quartz veins enclosing angular chunks of metabasalt hosts with multiple sub-parallel fault planes (identified from the slickenside lineations), see Fig. 3e & 3f. Quartz veins with multiple median lines indicate

predominance of crack-seal mechanism due to cyclic fluid ingression (Fig. 3g). This suggests that the process of fracturing, fracture reactivation through faulting and vein emplacement (*fault-valve* action) prevailed throughout the D3 deformation. Thus, the process of new fracture formation and reactivation were continuously persistent during the entire late D3 phase, with intermittent episodes of vein emplacement under both high and low  $P_f$  conditions.

## 7 Conclusions

In the present study, we commented on the vein emplacement mechanism of the Chitradurga greenstone belt (Dharwar craton, south India). We analysed the magnetic fabric data recorded from AMS analysis of the metabasalt that hosts the quartz veins. 3D Mohr circle and paleostress analysis have been used to evaluate the vein emplacement vs. regional deformation. Following are the main findings and conclusions from the study:

1. The NW-SE oriented magnetic fabric recorded in the metabasalts (as evident from the AMS analysis) is a product of the D1/D2 regional deformation on account of NE-SW directed shortening. This fabric was also favourably oriented and therefore, suitable for fracture propagation in relation to the prevailing stress field.
2. D3 deformation manifested by NW-SE to E-W directed shortening was coeval with the sinistral movement along CSZ. It is concluded that during late D3 deformation all pre-existing fabrics and the ones constituting the riedel shear system were reactivated, with CSZ acting as a shear boundary.

3. The variably oriented fractures and faults i.e., the NNW-SSE, NW-SE, WNW-ESE, NE-SW and ENE-WSW oriented ones are identified as the P, Y, R, T, X and R' shear components of the riedel shear system. The NNW-SSE oriented shears (P, Y and R) are abundant owing to their favourable correspondence with the pre-existing AMS fabric.
- 535
4. Paleostress analysis using fault-slip data recorded from field studies reveals NNE-SSW directed extension, asserting reactivation under a WNW-ESE directed compression (related to D3 deformation).
- 540
5. Vein emplacement took place under both high and low  $P_f$  conditions. It is envisaged that multiple number of such alternating high and low  $P_f$  cycles prevailed in the process of fluid flow and vein emplacement in the region.
6. Vein emplacement took place along all possible orientations. Evidences of *fault-valve behaviour* and fluid induced fracture/fault reactivation have been recorded from field studies. However, the NNW-SSE trending orientations having higher values of slip and dilation tendencies, channelized fluid during both high and low  $P_f$  conditions, thereby attaining maximum vein thickness.
- 545
7. The process of fracture formation, reactivation and faulting prevailed under a far field compression related to late D3 deformation. However, intermittent episodes of fluid pressure build up led to fluid induced faulting, rupturing and vein emplacement in the region.

550

**Author contribution:** SB: conceptualization, methodology, data curation, formal analysis, writing.  
TKM: supervision, conceptualization, methodology, data curation, funding acquisition, writing.

**Competing interests:** The authors declare that they have no conflict of interest.

555

### **Acknowledgments**

The study is funded by DST-SERB (File No. ECR/2015/000079) and RUSA 2.0 to TKM. This study is a part of SB's doctoral research; being funded by DST Inspire (IF170912). Geological Survey of India (Bangalore) is acknowledged for helping with logistic support and discussions during fieldwork. Prof. Manish A. Mamtani is thanked for allowing the authors to use KLY-4S Kappabridge system (IIT Kharagpur, India) for AMS analysis. Prof. Arindam Basu and Dr. Bikash K. Ram are thanked for helping with tensile strength measurements using GCTS<sup>©</sup> at IIT Kharagpur, India. **Detailed reviews by Prof. Francesco Mazzarini and Prof. C. Pascal helped to improve the paper considerably. Editorial handling by Prof. Roger Soliva is gratefully acknowledged.** Assistance provided by Ayan Patsa, Subha Saha and Swarnasree Mondal are acknowledged. SB also acknowledges the wholehearted support of Sunil Kumar Bhowmick and Gouri Bhowmick.

## References

- Acharyya, S.S., Mondal, T.K.: Stress enhanced tensile fractures in elliptical clast in conglomerate. *Journal of Structural Geology* 122 (2019) 81–88, <https://doi.org/10.1016/j.jsg.2019.02.001>, 2019.
- 570 Angelier, J., 1994.: Fault slip analysis and palaeostress construction. In: Hancock, P.L. (Ed.), *Continental Deformation*. Pergamon Press, London.
- ASTM (2001) American Society for Testing and Materials. ASTM Standards on Disc, 04.08. West Conshohocken, PA, 2001.
- 575 Attewell, P.B., Sandford, M.R.: Intrinsic shear strength of a brittle, anisotropic Rock-I. Experimental and mechanical interpretation. *Int. J. Rock Mech. Min. Sci. Geomech. Abstract.* 11, 423-430, 1974.
- Atkinson, B.K., and Meredith, P. G., 1989. Experimental fracture mechanics data for rocks and minerals, in Atkinson, B. K., ed., *Fracture mechanics of rocks*: London, Academic Press. 477-525.
- 580 Aydin A, Basu A.: The use of Brazilian test as a quantitative measure of rock weathering. *Rock Mech Rock Eng* 39:77–85, <https://doi.org/10.1007/s00603-005-0069-0>, 2006.
- Baer, G., Beyth, M., Reches, Z.: Dikes emplaced into fractured basement, Timna Igneous Complex, Israel. *Journal of Geophysical Research* 99, 24039–24051, <https://doi.org/10.1029/94JB02161>, 1994.
- 585 Basu, A., Mishra, D.A., Roychowdhury, K.: Rock failure modes under uniaxial compression, Brazilian, and point load tests. *Bull EngGeol Environ* (2013) 72:457–475, <http://dx.doi.org/10.1007%2Fs10064-013-0505-4>, 2013.
- Beckinsale, R. D., Drury, S. A., Holt, R. W.: 3305–Myr old gneisses from south Indian craton. *Nature* 283, 469–470, 1980.
- 590 Bhatt, S., Rana, V., Mamtani, M.A.: Deciphering relative timing of fabric development in granitoids with similar absolute ages based on AMS study (Dharwar Craton, South India). *Journal of Structural Geology* 94 (2017) 32-46,

[https://ui.adsabs.harvard.edu/link\\_gateway/2017JSG...94...32B/doi:10.1016/j.jsg.2016.11.002](https://ui.adsabs.harvard.edu/link_gateway/2017JSG...94...32B/doi:10.1016/j.jsg.2016.11.002),  
2017.

595 Blenkinsop, G. T.: Kinematic and dynamic fault slip analyses: implications from the surface rupture of  
the 1999 Chi–Chi, Taiwan, earthquake, *Journal of Structural geology*, 28, 1040–1050,  
<https://doi.org/10.1016/j.jsg.2006.03.011>, 2006.

Boullier, A. M., Robert, F.: Palaeoseismic events recorded in Archaean gold quartz vein networks, Val  
d'Or, Abitibi, Quebec, Canada. *Journal of Structural Geology* 14, 161–179,  
600 [https://doi.org/10.1016/0191-8141\(92\)90054-Z](https://doi.org/10.1016/0191-8141(92)90054-Z), 1992.

Chadwick, B., Vasudev, V. N., Hedge, G. V.: The Chitradurga schist belt and its adjacent plutonic rocks  
NW of Tungabhadra, Karnataka: a duplex in the late Archean convergent setting of the Dharwar  
craton. *Journal of the Geological Society of India* 61, 611–613, 2003.

Chadwick, B., Ramakrishnan, M., Vasudev, V. N., Viswanatha, M. N.: Facies Distributions and Structure  
605 of a Dharwar Volcano sedimentary Basin: Evidence for Late Archaean transpression in Southern  
India? *Journal of the Geological Society of London* 146, 825–834, 1989.

Chakrabarti, C., Mallick, B. S., Pyne, T. K., Guha, D.: *A manual of the Geology of India*. Geological  
Survey of India, Kolkata, 2006.

Chardon, D., Peucat, J.–J., Jayananda, M., Choukroune, P., Fanning, C.M.: Archaean granite–greenstone  
610 tectonics at Kolar (south India): interplay of diapirism and bulk inhomogeneous contraction during  
juvenile magmatic accretion. *Tectonics* 21, 1–17, <https://doi.org/10.1029/2001TC901032>, 2002.

Cox, S. F., Knackstedt, M. A., Braun, J.: Principles of structural control on permeability and fluid flow  
in hydrothermal systems. *Society of Economic Geologists Reviews* 14, 1–24, 2001.

Cox, S. F.: Faulting processes at high fluid pressures: an example of fault–valve behaviour from the  
615 Wattle Gully Fault, Victoria, Australia. *Journal of Geophysical Research* 100, 12841–12860,  
<https://doi.org/10.1029/95JB00915>, 1995.

Cox, S. F., Wall, V. J., Etheridge, M. A., Potter, T. F.: Deformational and metamorphic processes in the  
formation of mesothermal vein–hosted gold deposits — examples from the Lachlan Fold Belt in

- central Victoria, Australia. *Ore Geology Reviews* 6, 391–423, [https://doi.org/10.1016/0169-1368\(91\)90038-9](https://doi.org/10.1016/0169-1368(91)90038-9), 1991.
- 620
- Cucci, L., Luccio, F.D., Esposito, A., Ventura, G.: Vein networks in hydrothermal systems provide constraints for the monitoring of active volcanoes. *Scientific reports* 7:146, <https://dx.doi.org/10.1038/s41598-017-00230-8>, 2017.
- Delaney, P. T., Pollard, D. D., Zioney, J. I., McKee, E. H.: Field relations between dikes and joints: emplacement processes and palaeostress analysis. *Journal of Geophysical Research* 91 (B5), 4920–4938, [doi.org/10.1029/JB091iB05p04920](https://doi.org/10.1029/JB091iB05p04920), 1986.
- 625
- Delvaux, D.: EGU General Assembly. Win-tensor, an Interactive Computer Program for Fracture Analysis and Crustal Stress Reconstruction, vol. 13 , *Geophysical Research Abstract*, Vienna, 2011
- 630 Delvaux, D., Sperner, B.: Stress tensor inversion from fault kinematic indicators and focal mechanism data: the TENSOR program. In: Nieuwland, D. (Ed.), *New Insights into Structural Interpretation and Modelling*: Geol. Soc. Lond. Special Publication 212, 75–100, 2003.
- Donath, F.A.: Experimental study of shear failure in anisotropic rocks. *Geol. Soc. Am. Bull.* 72, 985-990, 1961.
- 635 [Donovan, J. G. 2003. Fracture toughness based model for the prediction of power consumption, product size, and capacity of jaw crushers. Doctoral dissertation, Virginia Polytechnic Institute and State University.](#)
- Dupin, J. M., Sassi, W., Angelier, J.: Homogeneous stress hypothesis and actual fault slip: a distinct element analysis, *Journal of Structural Geology* 15, 1033–1043, [doi.org/10.1016/0191-8141\(93\)90175-A](https://doi.org/10.1016/0191-8141(93)90175-A), 1993.
- 640
- Etchecopar, A., Vasseur, G., Daigniers, M. : An inverse problem in microtectonics for the determination of stress tensor from fault striation analysis, *Journal of Structural Geology* 3, 51–65, DOI: 10.1016/0191-8141(81)90056-0, 1981.

- Ferrill, D.A., Winterle, J., Wittmeyer, G., Sims, D., Colton, S., Armstrong, A., Morris, A.P., 1999.  
645 Stressed rock strains groundwater at Yucca Mountain, Nevada. *GSA Today* 9, 1–8.
- Gapais, D., Cobbold, P. R., Bourgeois, O., Rouby, D., de Urreiztieta, M.: Tectonic  
Significance of fault–slip data, *Journal of structural Geology*. 22, 881–888,  
[doi.org/10.1016/S0191-8141\(00\)00015-8](https://doi.org/10.1016/S0191-8141(00)00015-8), 2000.
- Gephart, J. W., Forsyth, D. W.: An improved method for determining the regional stress tensor using  
650 earthquake focal mechanism data: application to the San Fernando earthquake sequence, *Journal  
Geophysical Research* 89(B11), 9305–9320, [doi.org/10.1029/JB089iB11p09305](https://doi.org/10.1029/JB089iB11p09305), 1984.
- GitHub, 2017. FracTend MATLAB code. <https://github.com/DaveHealy-Aberdeen/FracTend>
- Gopalakrishna, G., Shareef, M., Nagesh, P.C., 2018. Shear-Controlled Gold Mineralization of G. R.Halli  
Area of Chitradurga Schist Belt, DharwarCraton: Insights from Fluid Inclusion Study. *Open*  
655 *Journal of Geology*, 8, 662-673, DOI: [10.4236/ojg.2018.87039](https://doi.org/10.4236/ojg.2018.87039) , 2018.
- Gudmundsson, A., 2011. *Rock Fractures in Geological Processes*. Cambridge University Press.
- Gupta, S., Jayananda, M., Fareeduddin, 2014. Tourmaline from the Archean G.R. Halli Gold Deposit,  
Chitradurga Greenstone Belt, Dharwar Craton (India): Implications for the Gold Metallogeny.  
*Geoscience Frontiers*, 5, 877-892, [doi.org/10.1016/j.gsf.2013.12.004](https://doi.org/10.1016/j.gsf.2013.12.004), 2014.
- 660 Hoek, E.: Fracture of anisotropic rock. *J. South Afr. Inst. Min. Metal.* 64 (10),  
501-518, 1964.
- Hrouda, F., Chlupáčová, M., Pokorný, J.: Low–field variation of magnetic susceptibility measured by the  
KLY–4S Kappabridge and KLF–4A magnetic susceptibility meter: Accuracy and interpretational  
programme. *Studia Geophysica et Geodaetica* 50, 283–299, 2006.
- 665 Ikari, M.J., Neimeijer, A.R., Marone, C.: Experimental investigation of incipient shear failure in foliated  
rock. *Journal of Structural Geology* 77, 82-91, [doi.org/10.1016/j.jsg.2015.05.012](https://doi.org/10.1016/j.jsg.2015.05.012), 2015.
- ISRM, Suggested methods for determining tensile strength of rock materials. *Int. J. Rock Mech. Min. Sci.  
& Geomech. Abstr.*, Vol. 15, 99-103, 1978.

- Jayananda, M., Chardon, D., Peucat, J.-J., Capdevila, R.: 2.61 Ga potassic granites and crustal reworking  
670 in the western Dharwar craton, southern India: tectonic, geochronologic and geochemical  
constraints. *Precambrian Research* 150, 1–26, DOI: 10.1016/j.precamres.2006.05.004, 2006.
- Jelínek, V.: Characterization of magnetic fabric of rocks. *Tectonophysics* 79, T63–T67,  
DOI:10.1016/0040-1951(81)90110-4, 1981.
- Jolly, R.J.H., Sanderson, D.J. : A Mohr circle reconstruction for the opening of a pre-existing fracture.  
675 *Journal of Structural Geology* 19, 887-892, DOI. 10.1016/S0191-8141(97)00014-X, 1997
- Lahiri, S., Mamtani, M.A.: Scaling the 3-D Mohr circle and quantification of paleostress during fluid  
pressure fluctuation–Application to understand gold mineralization in quartz veins of Gadag  
(southern India). *Journal of Structural geology* 88, 63–72, [doi.org/10.1016/j.jsg.2016.05.003](https://doi.org/10.1016/j.jsg.2016.05.003), 2016.
- Lawn, B.R., and Wilshaw, T.R., 1975. *Fracture of brittle solids: Cambridge, England, Cambridge*  
680 *University Press*.
- Logan, J. M., Dengo, C. A., Higgs, N. G., Wang, Z. Z.: Fabrics of experimental fault zones: their  
development and relationship to mechanical behaviour. In: Evans, B. & Wong, T. (eds) *Fault  
Mechanics and Transport Properties of Rocks*. Academic Press, San Diego, CA, 33–67, 1992.
- Loock, S., Diot, H., Van Wyk de Vries, B., Launeau, P., Merle, O., Vadeboin, F., Petronis, M.S.: Lava  
685 flow internal structure found from AMS and textural data: An example in methodology from the  
Chaîne des Puys, France. *Journal of Volcanology and Geothermal Research* 177, 1092–1104,  
[doi.org/10.1016/j.jvolgeores.2008.08.017](https://doi.org/10.1016/j.jvolgeores.2008.08.017), 2008.
- Maffione, M., Hernandez-Moreno, C., Ghiglione, M.C., Speranza, F., van Hinsbergen, D.J.J., Lodolo, E.:  
690 Deformation of the Southern Andes since the Late Cretaceous: constraints from anisotropy of  
magnetic susceptibility (AMS). *Tectonophysics* 665, 236–250,  
[doi.org/10.1016/j.tecto.2015.10.008](https://doi.org/10.1016/j.tecto.2015.10.008), 2015.
- Mamtani, M.A., Greiling, R.O.: Granite emplacement and its relation with regional deformation in the  
Aravalli Mountain Belt–inferences from magnetic fabric. *Journal of Structural Geology* 27, 2008–  
2029, [doi.org/10.1016/j.jsg.2005.06.004](https://doi.org/10.1016/j.jsg.2005.06.004), 2005.

- 695 Mamtani, M.A., Sengupta, P.: Significance of AMS analysis in evaluating superposed folds in quartzites. Geological Magazine 147, 910–918, DOI: 10.1017/S0016756810000397, 2010
- Marchesini, B., Garofalo, P.S, Menegon, L., Mattila, J., Viola, G.: Fluid-mediated, brittle–ductile deformation at seismogenic depth –Part 1: Fluid record and deformation history of fault veins in a nuclear waste repository (Olkiluoto Island, Finland). Solid Earth, 10, 809–838, doi.org/10.5194/se-10-809-2019, 2019.
- 700
- Marrett, R., Allmendinger, R. W.: Kinematic analysis of fault–slip data , J.Struc.Geol. 12, 973–986. doi.org/10.1016/0191-8141(90)90093-E, 1990.
- Martínez-Poza, A. I., Druguet, E., Castaño, L. M., Carreras, J.: Dyke intrusion into a pre-existing joint network: The Aiguablava lamprophyre dyke swarm (Catalan Coastal Ranges). Tectonophysics 705 630, 75–90, doi.org/10.1016/j.tecto.2014.05.015, 2014.
- Mazzarini, F., Isola, I.: Hydraulic connection and fluid overpressure in upper crustal rocks: evidence from geometry and spatial distribution of veins at Botrona quarry, southern Tuscany, Italy. Journal of Structural Geology 29, 1386–1399, doi.org/10.1016/j.jsg.2007.02.016, 2007.
- Mazzarini, F., Musumeci, G., Viola, G., Garofalo, P.S., Mattila, J.: Structural and lithological control on fluid circulation, dilation and ore mineralization (Rio Albano mine, Island of Elba, Italy). Journal of Structural Geology 126 (2019) 210–230, doi.org/10.1016/j.jsg.2019.06.012, 2019.
- 710
- McKeagney, C.J., Boulter, C.A., Jolly, R.J.H., Foster, R.P.: 3-D Mohr Circle analysis of vein opening, Indarama lode-gold deposit, Zimbabwe: implications for exploration. Journal of Structural Geology 26, 1275–1291, doi.org/10.1016/j.jsg.2003.11.001, 2004.
- 715
- Mondal, T.K., Acharyya, S.S.: Fractured micro-granitoid enclaves: a stress marker. Journal of Structural Geology 113, 33–41, DOI: 10.1016/j.jsg.2018.05.011, 2018.
- Mondal, T.K.: Evolution of fabric in Chitradurga granite (south India)-A study based on microstructure, anisotropy of magnetic susceptibility (AMS) and vorticity analysis. Tectonophysics 723, 149–161, doi.org/10.1016/j.tecto.2017.12.013, 2018.

- 720 Mondal, T.K., Mamtani, M.A.: Palaeostress analysis of normal faults in granite implications for interpreting Riedel shearing related to regional deformation. *Journal of Geological Society* 173, 216–227, [doi.org/10.1144/jgs2014-136](https://doi.org/10.1144/jgs2014-136), 2016.
- Mondal, T.K., Mamtani, M.A.: Fabric analysis in rocks of the Gadag region (southern India)-implications for time relationship between regional deformation and gold mineralization. *Tectonophysics* 629, 725 238–249, DOI: 10.1016/j.tecto.2013.09.021, 2014.
- Mondal, T.K., Mamtani, M.A.: 3-D Mohr circle construction using vein orientation data from Gadag (southern India) e implications to recognize fluid pressure fluctuation. *J. Struct. Geol.* 56, 45-56, [doi.org/10.1016/j.jsg.2013.08.005](https://doi.org/10.1016/j.jsg.2013.08.005), 2013.
- Mildren, S.D., Hillis, R.R., Kaldi, J.: Calibrating predictions of fault seal reactivation in the Timor Sea. 730 *APPEA J.42* (1), 187–202, 2002
- Morris, A., Ferrill, D.A., Henderson, D.B., 1996. Slip-tendency analysis and fault reactivation. *Geology* 24, 275–278.
- Naqvi, S.M., Rogers, J.J.W.: *Precambrian geology of India: Oxford monographs on geology and geophysics No. 6.* Oxford University Press, New York. 1987
- 735 Petit, J. P., Wibberley, C. A. J., Ruiz, G.: 'Crack±seal', slip: a new fault valve mechanism? *Journal of Structural Geology* 21, 1199-1207, 1999, 10.1016/S0191-8141(99)00038-3.
- Ramakrishnan, M., Vaidyanadhan, R., 2010. *Geology of India. Vol. 1.* Geological Society of India, Bangalore.
- Ramadass, G., Himabindu, D., Srinivasulu, N., 2003. Structural appraisal of the Gadag schist belt from 740 gravity investigations. *Proceedings of the Indian Academy of Sciences (Earth and Planetary Science)* 112, 577–586.
- Ramsay, J.G. & Lisle, R.J.: *The Techniques of Modern Structural Geology. Vol. 3: Applications of Continuum Mechanics in Structural Geology.* Academic Press, London, 2000.
- Raposo, M.I.B., D'Agrella-Filho, M.S., Pinese, J.P.P.: Magnetic fabrics and rock magnetism of Archaean and Proterozoic dike swarms in the southern São Francisco Craton, Brazil. *Tectonophysics* 443, 745 53–71 [doi.org/10.1016/j.tecto.2007.08.001](https://doi.org/10.1016/j.tecto.2007.08.001), 2007.

- Sanderson, D. J., Zhang, X., 1999. Critical stress localization of flow associated with deformation of well-fractured rock masses, with implications for mineral deposits. In: McCaffrey, K. J. W., Lonergan, L. and Wilkinson, J. J. (Eds.) *Fractures, Fluid Flow and Mineralization*. Geological Society of London, Special Publications 155, 69–81.
- 750 Sarma, D. S., Fletcher, I. R., Rasmussen, B., McNaughton, N. J., Mohan, M. R., Groves, D. I.: Archean gold mineralization synchronous with late cratonization of the Western Dharwar Craton, India: 2.52 Ga U–Pb ages of hydrothermal monazite and xenotime in gold deposits. *Mineralium Deposita* 46, 273–288, DOI: 10.1007/s00126-010-0326-3, 2011.
- 755 Sibson, R. H.: A brittle failure mode plot defining conditions for high flux–flow. *Economic Geology* 95, 41–48, 10.2113/gsecongeo.95.1.41, 2000.
- Sibson, R. H., Robert, F., Poulsen, K. H.: High–angle reverse faults, fluid–pressure cycling, and mesothermal gold–quartz deposits. *Geology* 16, 551–555, doi.org/10.1130/0091-7613(1988)016<0551:HARFFP>2.3.CO;2, 1988
- 760 Sibson, R. H., Scott, J.: Stress/fault controls on the containment and release of over pressured fluids: examples from gold–quartz vein systems in Juneau, Alaska, Victoria, Australia, and Otago, New Zealand. *Ore Geology Reviews* 13, 293–306, 10.1016/S0169-1368(97)00023-1, 1998.
- Sibson, R. H.: Structural permeability of fluid–driven fault–fracture meshes. *Journal of Structural Geology* 18, 1031–1043, doi.org/10.1016/0191-8141(96)00032-6, 1996.
- 765 Sibson, R. H.: Implications of fault–valve behaviour for rupture nucleation and recurrence. *Tectonophysics* 211, 283–293, doi.org/10.1016/0040-1951(92)90065-E, 1992.
- Sivakugan, N., Das, B. M., Lovisa, J., Patra, C. R.: Determination of  $c$  and  $w$  of rocks from indirect tensile strength and uniaxial compression tests. *International Journal of Geotechnical Engineering* 8, 59–65, doi.org/10.1179/1938636213Z.00000000053, 2014.
- 770 Stephens, T.L., Walker, R.J., Healy, D., Bubeck, A., England, R.W., McCaffrey, K.J.: Igneous sills record far-field and near-field stress interactions during volcano construction: Isle of Mull, Scotland. *Earth Planet. Sci. Lett.* 478, 159–174. doi.org/10.1016/j.epsl.2017.09.003, 2017.
- Tarling, D. H., Hrouda, F., 1993. *The Magnetic Anisotropy of Rocks*. Chapman and Hall, London.

- 775 Taylor, P.N., Chadwick, B., Moorbath, S., Ramakrishnan, M., Viswanatha, M.N.: Petrography, chemistry and isotopic ages of Peninsular Gneiss, Dharwar acid volcanic rocks and the Chitradurga granite with special reference to the late Archaean evolution of the Karnataka craton. *Precambrian Research* 23, 349- 375. [10.1016/0301-9268\(84\)90050-0](https://doi.org/10.1016/0301-9268(84)90050-0), 1984.
- Thyng, K.M., Greene, C.A., Hetland, R.D., Zimmerle, H.M., DiMarco, S.F.; True Colors of Oceanography Guidelines for Effective and Accurate Colormap Selection. *Oceanography*, 9-13, [doi.org/10.5670/oceanog.2016.66](https://doi.org/10.5670/oceanog.2016.66), 2016
- 780 Tsidzi, K. E. N.: The influence of foliation on point load strength anisotropy of foliated rocks. *Engineering Geology* 29, 49–58. [10.1016/0013-7952\(90\)90081-B](https://doi.org/10.1016/0013-7952(90)90081-B), 1990.
- Twiss, R. J., Unruh, J. R.: Analysis of fault slip inversions; do they constrain stress or strain rate? , *Journal of Geophysical Research* 103 (B6), 12205–12222, [doi.org/10.1029/98JB00612](https://doi.org/10.1029/98JB00612).1998.
- 785 Vishnu, C. S., Lahiri, S., Mamtani, M. A.: The relationship between magnetic anisotropy, rock-strength anisotropy and vein emplacement in gold-bearing metabasalts of Gadag (South India). *Tectonophysics* 722, 286-298, [doi.org/10.1016/j.tecto.2017.09.011](https://doi.org/10.1016/j.tecto.2017.09.011). 2018.
- Yamaji, A.: Genetic algorithm for fitting a mixed Bingham distribution to 3D orientations: a tool for the statistical and paleostress analyses of fracture orientations. *Island Arc* 25, 72-83. DOI: [10.1111/iar.12135](https://doi.org/10.1111/iar.12135), 2016.
- 790 Yamaji, A., Sato, K.: Clustering of fracture orientations using a mixed Bingham distribution and its application to paleostress analysis from dike or vein orientations. *Journal of Structural Geology* 33, 1148-1157. DOI: [10.1016/j.jsg.2011.05.006](https://doi.org/10.1016/j.jsg.2011.05.006), 2011.
- Yamaji, A., Sato, K., Tonai, S.: Stochastic modeling for the stress inversion of vein orientations: Paleostress analysis of Pliocene epithermal veins in south western Kyushu, Japan. *Journal of Structural Geology* 32, 1137–1146. DOI: [10.1016/j.jsg.2010.07.001](https://doi.org/10.1016/j.jsg.2010.07.001), 2010.
- 795 Yamaji, A.: The multiple inverse methods: a new technique to separate stresses from heterogeneous fault-slip data, *Journal of Structural Geology* 22, 441–452. DOI: [10.1016/S0191-8141\(99\)00163-7](https://doi.org/10.1016/S0191-8141(99)00163-7) 2000.

800 Žalohar, J., Vrabec, M.: Paleostress analysis of heterogeneous fault–slip data: the Gauss method, *Journal  
of Structural Geology* 29, 1798–1810. DOI: 10.1016/j.jsg.2007.06.009, 2007.

805

810


**Figures:**

815

820


825

830



**Figure 01:** (a) Regional map of western Dharwar craton, South Indian Shield [after Chadwick et al., 2003]. Inset shows the map of India. EDC = Eastern Dharwar Craton; WDC = Western Dharwar Craton, **TTG=Tonalite-Trondhjemite-Granodiorite; Supracrustals=volcano-sedimentary assemblages.** (b) Regional geological map (DEM) of the Chitradurga Schist Belt [modified after Jayananda et al., 2013]. Dotted line (in b) marks the eastern boundary of the Chitradurga Schist Belt, representing the Chitradurga Shear Zone (CSZ). Rectangular box near Chitradurga demarcates the study area.

840



**Figure 02:** Geological map of the study area showing the foliation trends of the supracrustals. Lower hemisphere equal area projection shows the pole to foliation planes recorded from the region. Note that the mean foliation (dotted great circle) is 323°/71°NE which is parallel to the CSZ (n= number data). Color scheme of the legend indicates variation in the contour density.

865

870

875




880



885



890




895


**Figure 03.** Field photographs from the study area. (a) Criss-cross orientation of quartz veins in metabasalt.  
900 (b) Close-up of a quartz vein in metabasalt showing crystal growth direction perpendicular to the vein wall. (c) Cross-cutting nature in quartz veins showing dextral displacement (marked by yellow half-arrows). (d) Wing cracks filled up with quartz veins associated with sinistral shearing (marked by yellow arrow). (e) NE dipping quartz vein showing slickenside lineations (maximum width recorded=120 cm), inset showing close up of the fault plane found in e. Marker pen placed along the orientation of the slickenside lineations.  
905 (f) Angular chunks of metabasalt (enclaves) enclosed within faulted quartz vein. Dotted red line demarcates the enclave boundaries. Black arrow marks the slickenside lineation on the fault plane. (g) Field photograph of quartz vein (close view) showing multiple median lines (marked with red arrows) as an evidence of crack-seal mechanism. This clearly indicates cyclic fluid ingress and fault valve action that led to the formation of veins in the metabasalts of the Chitradurga region. (h) Fault  
910 planes in metabasalt showing congruous steps (marked by yellow arrow). Blue arrow in the respective figures point towards N.

915

920



940 **Figure 04.** Rose diagrams and graphs of structural data recorded from the study area. (a), (b) and (c) are  
 945 the rose diagrams of the strike orientation of the quartz veins, fractures and fault planes respectively. Note  
 that the veins are mostly NNW-SSE striking whereas the fractures and fault planes show variable  
 orientations, with a NNW-SSE maxima and a WNW-ESE to NE-SW sub maxima. (d) Graph of vein  
 orientation vs. thickness, showing maximum vein thickness along NNW-SSE (marked with orange box).  
 n= number of data.



975

**Figure 05.** Map showing the location points (marked with brown boxes) from which oriented metabasalt samples for AMS analysis have been collected. (a) Lower hemisphere equal area projection of poles ( $K_3$ ) to magnetic foliation ( $K_1K_2$ ) plane. Mean orientation of  $K_1K_2$  plane (dashed great circle)=  $337^\circ/69^\circ E$  (b) Lower hemisphere equal area projection showing the distribution of magnetic lineation ( $K_1$ ). Color scheme of the legend indicates variation in the contour density.

985

990

995



**Figure 06.** Tensile strength determination of metabasalt samples. (a) Sample cores for Brazilian tensile strength (BTS) determination. Diameter: length=2:1. (b) Instruments used for measuring BTS. (c) Sample cores obtained after failure.

1000

1005


1010

1015

1020

1025

1030



1035

1040

**Figure 07.** State of stresses and fluid pressure ( $P_f$ ) conditions determined from vein orientation data in the study area. (a) Lower hemisphere equal area projection of pole to veins shows girdle distribution, implying  $P_f > \sigma_2$  (following Jolly and Sanderson, 1997). The empty space devoid of any vein pole data helps to determine the position of  $\sigma_1$  (using Bingham statistics of the Stereonet 9 software) and thereby defines  $\sigma_1\sigma_2$  and  $\sigma_1\sigma_3$  planes. Angles  $\theta_2$  and  $\theta_3$  are measured which are used to determine the stress ratio ( $\phi$ ) and driving pressure ratio ( $R'$ ) respectively. Color scheme of the legends indicate variation in the contour density. Pink circle ( $\sigma_1$ ), pink triangle ( $\sigma_2$ ) and pink square ( $\sigma_3$ ) respectively. (b) 3D Mohr circle diagram for high  $P_f$  conditions. Red line in the Mohr circle represents the reactivation envelope for cohesionless fractures. Vein pole data lying within the blue zone, i.e., to the left of the  $P_f$  (black) line represent fractures filled up with veins that are susceptible to reactivate (Fractend code available via


Github, 2017). (c) Lower hemisphere equal area projection of poles to vein data forming the SW cluster in (a), this cluster distribution of vein pole data indicates  $P_f < \sigma_2$ . Cluster maxima defines  $\sigma_3$  axis. Angles  $\theta_1$  and  $\theta_2$  are measured and similarly  $\phi$  and  $R'$  are determined. (d) 3D Mohr circle diagram for low  $P_f$  condition. Only a limited range of fracture filled up with veins are susceptible to reactivate. Red dots represent pole to vein data, red line forms the reactivation envelope for cohesionless fractures.

1050


1055

1060

1065




**b**




1070

1075

1080



**d**



1085 **Figure 08.** State of stress and fluid pressure ( $P_f$ ) conditions determined from the vein orientation data in  
the study area. (a) Lower hemisphere equal area projection of poles to vein data forming the NE cluster.  
(b) 3D Mohr circle diagram for NE cluster. (c) Lower hemisphere equal area projection of poles to vein  
data forming the SE cluster. (d) 3D Mohr circle diagram for SE cluster. In the respective lower hemisphere  
1090 equal area projections, cluster distribution of vein pole data indicates  $P_f < \sigma_2$ . Cluster maxima defines  $\sigma_3$   
axis. The ranges of fracture orientations ( $\theta$ ) are measured. Subsequently,  $\phi$  and  $R'$  values are also  
determined similar to Fig 7. Pink circle ( $\sigma_1$ ), pink triangle ( $\sigma_2$ ) and pink square ( $\sigma_3$ ) respectively. In the  
3D Mohr circle diagrams, only a limited range of fracture filled up with veins are susceptible to reactivate.  
Red dots represent pole to vein data, red line forms the reactivation envelope for cohesionless fractures.  
1095 Vein pole data lying within the blue zone, i.e., to the left of the  $P_f$  (black) line represent fractures filled  
up with veins that are susceptible to reactivate.

1100


1105

1110


1115

1120

1125




1130 **Figure 09.** Lower hemisphere equal area projection for dilation tendency, slip tendency and fracture
 1135 susceptibility in a given fluid pressure condition and for a specific stress state (Fractend code available
 via Github, 2017). (a) and (b) represents dilation tendency for high and low  $P_f$ . (c) and (d) represents slip
 tendency for high and low  $P_f$ . (e) and (f) fracture susceptibility for high and low  $P_f$  conditions respectively.
 Warm color zones in (a) and (b), represents vein orientations with higher propensity for tensional opening.
 In (c) and (d) warm colors indicate vein attitudes that suffered shearing. In (e) and (f) the warm color
 zones stand for vein attitudes that are more susceptible to reactivate under low  $P_f$  variation. ‘Thermal’
 color scheme from Thyng et al., 2016. White square ( $\sigma_1$ ), white diamond ( $\sigma_2$ ) and white triangle ( $\sigma_3$ ).



1145

**Figure 10.** Paleostress analysis (using right dihedral method) of faults recorded in the metabasalts of the study area; for both left lateral and right lateral normal faults. Red arrow marks the extension direction ( $\sigma_3$ ) and pink arrow marks the principal compression direction. The histogram shows the minimum values for counting deviation.  $n$  and  $nt$  are the number of data accepted for tensor calculation and total number of data respectively.  $R$  (stress ratio) =  $(\sigma_2 - \sigma_3) / (\sigma_1 - \sigma_3)$ . Rose diagrams for fault plane strike, dip and plunge variations are also presented in the top right corner. 3D Mohr diagrams demonstrate possible areas of fault reactivation with variation in susceptibility, colored circles represent pole to the fault planes. The failure envelope for Mohr's circle is based on Byerlee (1978) initial friction angle of  $16.7^\circ$ . Note that the maximum extension is along NNE-SSW direction.



**Figure 11.** Schematic model showing formation and reactivation of fractures, along with vein emplacement in the metabasalt host rock. (a) Development of the NNW-SSE to NW-SE oriented magnetic fabrics in the rock unit under a NE-SW shortening related to D1/D2 deformation. Lower hemisphere equal area projection shows mean orientation of the magnetic fabric  $\sim 337^\circ/69^\circ\text{NE}$ . (b) Formation of the Y, P, R, R', X and T shears related to the riedel shear system under a NW-SE to E-W shortening related to the late D3 deformation (half arrows representing the shear zone boundary). The angles ( $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\lambda$  and  $\Delta$ ) between the shear components are given after Logan et al. 1992.  $\Phi=30^\circ$  (angle of internal friction) is measured from UCS studies of the samples from the study area. Corresponding riedel shear model, with CSZ as the shear boundary and rose diagram showing orientations of the respective shear components. (c) and (d) showing vein emplacement under high  $P_f$  and low  $P_f$  conditions respectively (**state of stresses along the fracture planes are given, in c,  $\sigma_1$  is vertical due to fluid overpressure**). Corresponding 3D-mohr circle diagrams quantifying the effective normal stresses ( $\sigma_n' = \sigma_n - P_f$ , in Mpa) under both high and low  $P_f$  conditions, difference in  $P_f$  ( $\Delta P_f \sim 13.9$  MPa), obtained from the dashed lines representing the mean stress in each case. A conceptual graph shows multiple cycles (n-times) of high and low  $P_f$  conditions in the study area justifying *fault-valve action* that led to the emplacement of vein in the Chitradurga region.