

Comparative geochemical study on Furongian–earliest Ordovician (Toledanian) and Ordovician (Sardic) felsic magmatic events in south-western Europe: underplating of hot mafic magmas linked to the opening of the Rheic Ocean

J. Javier Álvaro^{a*}, Teresa Sánchez-García^b, Claudia Puddu^c, Josep Maria Casas^d, Alejandro Díez-Montes^e, Montserrat Liesa^f & Giacomo Oggiano^g

^a *Instituto de Geociencias (CSIC-UCM), Dr. Severo Ochoa 7, 28040 Madrid, Spain, jj.alvaro@csic.es*

^b *Instituto Geológico y Minero de España, Ríos Rosas 23, 28003 Madrid, Spain, t.sanchez@igme.es*

^c *Dpto. Ciencias de la Tierra, Universidad de Zaragoza, 50009 Zaragoza, Spain, claudiapuddugeo@gmail.com*

^d *Dpt. de Dinàmica de la Terra i de l'Oceà, Universitat de Barcelona, Martí Franquès s/n, 08028 Barcelona, Spain, casas@ub.edu*

^e *Instituto Geológico y Minero de España, Plaza de la Constitución 1, 37001 Salamanca, Spain, al.diez@igme.es*

^f *Dept. de Mineralogia, Petrologia i Geologia aplicada, Universitat de Barcelona, Martí Franquès s/n, 08028 Barcelona, Spain, mliesa@ub.edu*

^g *Dipartimento di Scienze della Natura e del Territorio, 07100 Sassari, Italy, giacoggi@uniss.it*

* Corresponding author

ABSTRACT

A geochemical comparison of Early Palaeozoic felsic magmatic episodes throughout the south-western European margin of Gondwana is **made, and includes** (i) Furongian–Early Ordovician (Toledanian) activities recorded in the Central Iberian and Galicia-Trás-os-Montes Zones of the Iberian Massif, and (ii) Early–Late Ordovician (Sardic) activities in the eastern Pyrenees, Occitan Domain (Albigeois, Montagne Noire and Mouthoumet massifs) and Sardinia. Both phases are related to uplift and denudation of an inherited palaeorelief, and stratigraphically preserved as distinct angular discordances and paraconformities involving gaps of up to 22 m.y. The geochemical features of the Toledanian and Sardic, felsic-dominant activities point to a predominance of **magmatic** byproducts derived from the melting of metasedimentary rocks, rich in SiO₂ and K₂O and with peraluminous character. Zr/TiO₂, Zr/Nb, Nb/Y and Zr vs. Ga/Al ratios, and REE and $\epsilon\text{Nd}_{(t)}$ values suggest the contemporaneity, for both phases, of two geochemical scenarios characterized by arc and extensional features evolving to distinct extensional and rifting conditions associated with the final outpouring of mafic tholeiitic-dominant lava flows. The Toledanian and Sardic **magmatic** phases are linked to neither metamorphism nor penetrative deformation; on the contrary, their unconformities are associated with foliation-free open folds subsequently affected by the Variscan deformation. The geochemical and structural framework precludes subduction **generated melts** reaching the crust in a magmatic arc to back-arc setting, but favours partial melting of sediments and/or granitoids in a continental lower crust triggered by the underplating of hot mafic magmas related to the opening of the Rheic Ocean.

Keywords: granite, orthogneiss, geochemistry, Cambrian, Ordovician, Gondwana.

1. Introduction

A succession of Early–Palaeozoic magmatic episodes, ranging in age from Furongian (former “late Cambrian”) to Late Ordovician, is widespread along the south-western European margin of Gondwana. Magmatic pulses are characterized by preferential development in different palaeogeographic areas and linked to the development of stratigraphic unconformities, but they are related to neither metamorphism nor penetrative deformation (Gutiérrez Marco et al., 2002; Montero et al., 2007). In the Central Iberian Zone of the Iberian Massif (representing the western branch of the Ibero-Armorican Arc; Fig. 1A–B), this magmatism is mainly represented by the Ollo de Sapo Formation, which has long been recognized as a Furongian–Early Ordovician (495–470 Ma) assemblage of felsic-dominant volcanic, subvolcanic and plutonic igneous rocks. This magmatic activity is contemporaneous with the development of the Toledanian Phase, which places Lower Ordovician (upper Tremadocian–Floian) rocks onlapping an inherited palaeorelief formed by Ediacaran–Cambrian rocks and involving a sedimentary gap of ca. 22 m.y. This unconformity can be correlated with the “Furongian gap” identified in the Ossa-Morena Zone of the Iberian Massif and the Anti-Atlas Ranges of Morocco (Álvaro et al., 2007, 2018; Álvaro and Vizcaino, 2018; Sánchez-García et al., 2019), and with the “lacaune normande” in the central and North-Armorican Domains (Le Corre et al., 1991).

Another felsic-dominant magmatic event, although younger (Early–Late Ordovician) in age, has been recognized in some massifs situated along the eastern branch of the Variscan Ibero-Armorican Arc, such as the Pyrenees, the Occitan Domain and Sardinia (Fig. 1A, C–E). This magmatism is related to the Sardic unconformity, where Furongian–Lower Ordovician rocks are unconformably overlain by those attributed to the Sandbian–lower Katian (former Caradoc). The Sardic Phase is related to a sedimentary gap of ca. 16–20 m.y. and geometrically ranges from 90° (angular

discordance) to 0° (paraconformity) (Barca and Cherchi, 2004; Funneda and Oggiano, 2009; Álvaro et al., 2016, 2018; Casas et al., 2019).

Although a general consensus exists to associate this Furongian–Ordovician magmatism with the opening of the Rheic Ocean and the drift of Avalonia from northwestern Gondwana (Díez Montes et al., 2010; Nance et al., 2010; Thomson et al., 2010; Álvaro et al., 2014a), the origin of this magmatism has received different interpretations. In the Central Iberian Zone, for instance, **several geodynamic models have been proposed, such as:** (i) **subduction-related melts** reaching the crust in a magmatic arc to back-arc setting (Valverde-Vaquero and Dunning, 2000; Castro et al., 2009); (ii) partial melting of sediments or granitoids in a continental lower crust affected by the underplating of hot mafic magmas during an extensional regime (Bea et al., 2007; Montero et al., 2009; Díez Montes et al., 2010); and (iii) post-collisional decompression melting of an earlier thickened continental crust, and without significant mantle involvement (Villaseca et al., 2016). In the Occitan Domain (southern French Massif Central and Mouthoumet massifs) and the Pyrenees, Marini (1988), Pouclet et al. (2017) and Puddu et al. (2019) have suggested a link to mantle thermal anomalies. Navidad et al. (2018) proposed that the Pyrenean magmatism was induced by progressive crustal thinning and uplift of lithospheric mantle isotherms. In Sardinia, Oggiano et al. (2010), Carmignani et al. (2001), Gaggero et al. (2012) and Cruciani et al. (2018) have suggested that a subduction scenario, mirroring an Andean-type active margin, **caused** the main Mid–Ordovician magmatic activity. In the Alps, the Sardinian counterpart is also interpreted as a result of the collision of the so-called Qaidam Arc with the Gondwanan margin, subsequently followed by the accretion of the Qilian Block (Von Raumer and Stampfli, 2008; Von Raumer et al., 2013, 2015). This geodynamic interpretation is mainly suggested for the Alpine Briançonnais-Austroalpine basement, where the volcanosedimentary complexes postdating the Sardinian tectonic inversion and folding stage portray a younger arc-arc oblique collision (450 Ma) of the eastern tail of the internal Alpine margin with the Hun terrane, succeeded by conspicuous exhumation

in a transform margin setting (430 Ma) (Zurbriggen et al., 1997; Schaltegger et al., 2003; Franz and Romer, 2007; Von Raumer and Stampfli, 2008; Von Raumer et al., 2013; Zurbriggen, 2015, 2017).

Until now the Toledanian and Sardinic magmatic events had been studied on different areas and interpreted separately, without taking into account their similarities and differences. In this work, the geochemical affinities of the Furongian–Early Ordovician (Toledanian) and Early–Late Ordovician (Sardic) felsic magmatic activities recorded in the Central Iberian and Galicia-Trás-os-Montes Zones, Pyrenees, Occitan Domain and Sardinia are compared. The re-appraisal is based on 17 new samples from the Pyrenees, Montagne Noire and Sardinia, completing the absence of analysis in these areas and wide-ranging a dataset of 93 previously published geochemical analyses throughout the study region in south-western Europe. This comparison may contribute to a better understanding of the meaning and origin of this felsic magmatism, and thus, to discuss the geodynamic scenario of this Gondwana margin (Fig. 1A) during Cambrian–Ordovician times, bracketed between the Cadomian and Variscan orogenies.

2. Emplacement and age of magmatic events

This section documents the emplacement (summarized in Fig. 2) and age (Fig. 3) of the Toledanian and Sardinic magmatic events throughout a SW-NE palaeogeographic transect of the south-western European margin of Gondwana during Cambro–Ordovician times.

2.1. Iberian Massif

In the Ossa Morena and southern Central Iberian Zones of the Iberian Massif (Fig. 1A–B), the so-called Toledanian Phase is recognized as an angular discordance that

separates variably tilted Ediacaran–Cambrian Series 2 rifting volcanosedimentary packages from overlying passive-margin successions. The Toledanian gap comprises, at least, most of the Furongian and basal Ordovician, but the involved erosion can incise into the entire Cambrian and the upper Ediacaran Cadomian basement (Gutiérrez-Marco et al., 2019; Álvaro et al., 2019; Sánchez-García et al., 2019). Recently, Sánchez-García et al. (2019) have interpreted the Toledanian Phase as a break-up (or rift/drift) unconformity with the Armorican Quartzite (including the Purple Series and Los Montes Beds; McDougall et al., 1987; Gutiérrez-Alonso et al., 2007; Shaw et al., 2012, 2014) sealing an inherited Toledanian palaeorelief (Fig. 2).

The phase of uplift and denudation of an inherited palaeorelief composed of upper Ediacaran–Cambrian rocks is associated with the massive outpouring of felsic-dominant calc-alkaline magmatic episodes related to neither metamorphic nor cleavage features. This magmatic activity is widely distributed throughout several areas of the Iberian Massif, such as the Cantabrian Zone and the easternmost flank of the West Asturian-Leonese Zone, where sills and rhyolitic lava flows and volcanoclastics mark the base of the Armorican Quartzite (dated at ca. 477.5 Ma; Gutiérrez-Alonso et al., 2007, 2016), and the lower Tremadocian Borrachón Formation of the Iberian Chains (Álvaro et al., 2008). Similar ages have been reported from igneous rocks of the Basal Allochthonous Units and the Schistose Domain in the Galicia-Trás-os-Montes Zone (500–462 Ma; Valverde-Vaquero et al., 2005, 2007; Montero et al., 2009; Talavera et al., 2008, 2013; Dias da Silva et al., 2012, 2014; Díez Fernández et al., 2012; Farias et al., 2014) and different areas of the Central Iberian Zone, including the contact between the Central Iberian and Ossa-Morena Zones, where the Carrascal and Portalegre batoliths are intruded and the felsic volcanosedimentary Urra Formation marks the unconformity that separates Cambrian and Ordovician strata (494–470 Ma, Solá et al., 2008; Antunes et al., 2009; Neiva et al., 2009; Romão et al., 2010; Rubio-Ordóñez et al., 2012; Villaseca et al., 2013) (Fig. 1B).

The most voluminous Toledanian-related volcanic episode is represented by the Ollo de Sapo Formation, which **crops out throughout** the northeastern Central Iberian Zone. It mainly consists of felsic volcanosedimentary and volcanic rocks, interbedded at the base of the Lower Ordovician strata and plutonic bodies. The Ollo de Sapo volcanosedimentary Formation has long been recognized as an enigmatic Furongian–Early Ordovician (495–470 Ma) magmatic event exposed along the core of a 600 km-long antiform (labelled as 77 in [Fig. 1B](#)) (Valverde-Vaquero and Dunning, 2000; Bea et al., 2006; Montero et al., 2007, 2009; Zeck et al., 2007; Castiñeiras et al., 2008a; Díez Montes et al., 2010; Navidad and Castiñeiras, 2011; Talavera et al., 2013; López-Sánchez et al., 2015; Díaz-Alvarado et al., 2016; Villaseca et al., 2016; García-Arias et al., 2018). The peak of magmatic activity was reached at ca. 490–485 Ma and its most recognizable characteristic is the presence of abundant megacrysts of K-feldspar, plagioclase and blue quartz. There is no evident space-time relationship in its distribution (for a discussion, see López-Sánchez et al., 2015) and, collectively, the Ollo de Sapo Formation rocks **record** a major tectonothermal event whose expression can be found in most of the Variscan massifs of continental Europe including the Armorican and Bohemian massifs (e.g., von Quadt, 1997; Kröner and Willmer, 1998; Linnemann et al., 2000; Tichomirowa et al., 2001; Friedl et al., 2004; Mingram et al., 2004; Teipel et al., 2004; Ballèvre et al., 2012; El Korh et al., 2012; Tichomirowa et al., 2012; for a summary, see Casas and Murphy, 2018). The large **volume** of magmatic rocks located in the European Variscan Belt led some authors to propose the existence of a siliceous Large Igneous Province (LIP) (Díez Montes et al., 2010; Gutiérrez-Alonso et al., 2016), named Ibero-Armorican LIP by García-Arias et al. (2018).

2.2. Central and Eastern Pyrenees

In the **central and eastern** Pyrenees ([Fig. 1D](#)), earliest Ordovician volcanic-free passive-margin conditions, represented by the Jujols Group (Padel et al., 2018), were

succeeded by a late Early–Mid Ordovician phase of uplift and erosion that led to the onset of the Sardinian unconformity (Fig. 2). Uplift was associated with magmatic activity, which continued until Late Ordovician times. An extensional interval took place then developing normal faults that controlled the sedimentation of post–Sardinian siliciclastic deposits infilling palaeorelief depressions. Acritharchs recovered in the uppermost part of the Jujols Group suggest a broad Furongian–earliest Ordovician age (Casas and Palacios, 2012), conterminous with a maximum depositional age of ca. 475 Ma, based on the age of the youngest detrital zircon populations (Margalef et al., 2016). On the other hand, a ca. 459 Ma U–Pb age for the Upper Ordovician volcanic rocks overlying the Sardinian Unconformity has been proposed in the eastern Pyrenees (Martí et al., 2019), and ca. 452–455 Ma in the neighbouring Catalan Coastal Ranges, which represent the southern prolongation of the Pyrenees (Navidad et al., 2010; Martínez et al., 2011). Thus, a time gap of about 16–23 m.y. can be related to the Sardinian Phase in the eastern Pyrenees and the neighbouring Catalan Coastal Ranges.

Coeval with the late Early–Mid Ordovician phase of generalized uplift and denudation, a key magmatic activity led to the intrusion of voluminous granitoids, about 500 to 3000 m thick and encased in strata of the Ediacaran–Lower Cambrian Canaveilles Group (Fig. 2). These granitoids constitute the protoliths of the large orthogneissic laccoliths that punctuate the backbone of the central and eastern Pyrenees. These are, from west to east (Fig. 1D), the Aston (467–470 Ma ; Denèle et al., 2009; Mezger and Gerdes, 2016), Hospitalet (about 472 Ma, Denèle et al., 2009), Canigó (472–462 Ma, Cocherie et al., 2005; Navidad et al., 2018), Roc de Frausa (477–476 Ma; Cocherie et al., 2005; Castiñeiras et al., 2008b) and Albera (about 470 Ma; Liesa et al., 2011) massifs, which comprise a dominant Floian–Dapingian age. It is noticeable the fact that only a minor representation of coeval basic magmatic rocks are outcropped. The acidic volcanic equivalents have been documented in the Albera massif, where subvolcanic rhyolitic porphyroid rocks have yielded similar ages to those

of the main gneissic bodies at **about** 474–465 Ma (Liesa et al., 2011). Similar acidic byproducts are represented by the rhyolitic sills of Pierrefite (Calvet et al., 1988).

The late Early–Mid Ordovician (“Sardic”) phase of uplift was succeeded by a Late Ordovician extensional **interval** responsible for the opening of (half-)grabens infilled with the basal Upper Ordovician alluvial-to-fluvial conglomerates (La Rabassa **Conglomerate** Formation). At **map** scale, a set of NE-SW trending normal faults abruptly **controlling** the thickness of the basal Upper Ordovician formations can be recognized in the La Cerdanya area (Casas and Fernández, 2007; Casas, 2010). Sharp variations in the thickness of the Upper Ordovician strata have been documented by Hartevelt (1970) and Casas and Fernández (2007). Drastic variations in grain size and thickness can be attributed to the development of palaeotopographies controlled by faults and subsequent erosion of uplifted palaeoreliefs, with subsequent infill of depressed areas by alluvial fan and fluvial deposits, finally sealed by Silurian sediments (Puddu et al., 2019). A Late Ordovician magmatic pulse contemporaneously yielded a varied set of magmatic rocks. Small granitic bodies are encased in the Canaveilles strata of the Canigó massif. They constitute the protoliths of the Cadí (**about** 456 Ma; Casas et al., 2010), Casemí (446 **to** 452 Ma; Casas et al., 2010), Núria (**ca.** 457 Ma; Martínez et al., 2011) and Canigó G-1 type (**ca.** 457 Ma; Navidad et al., 2018) gneisses.

The lowermost part of the Canaveilles Group (the so-called Balaig Series) host metre-scale thick bodies of metadiorite **sills** related to an Upper Ordovician protolith, (**ca.** 453 Ma, SHRIMP U–Pb in zircon; Casas et al., 2010). Coeval calc-alkaline ignimbrites, andesites and volcanoclastic rocks are interbedded in the Upper Ordovician succession of the Bruguera and Ribes de Freser areas (Robert and Thiebaut, 1976; Ayora, 1980; Robert, 1980; Martí et al., 1986, 2019). In the Ribes area, a granitic body with granophyric texture, dated at **ca.** 458 Ma by Martínez et al. (2011), intruded at the base of the Upper Ordovician succession. In the La Pallaresa dome, some metre-scale rhyodacitic to dacitic subvolcanic sills, Late Ordovician in age (**ca.** 453 Ma, Clariana et

al., 2018), occur interbedded within the pre-unconformity strata and close to the base of the Upper Ordovician.

2.3. Occitan Domain: Albigeois, Montagne Noire and Mouthoumet massifs

The parautochthonous framework of the southern French Massif Central, named Occitan Domain by Pouclet et al. (2017), includes among others, from south to north, the Mouthoumet, Montagne Noire and Albigeois massifs. The domain represents the southeastern prolongation of the Variscan South Armorican Zone (including southwestern Bretagne and Vendée). Since Gèze (1949) and Arthaud (1970), the southern edge of the French Massif Central has been traditionally subdivided, from north to south, into the northern, axial and southern Montagne Noire (Fig. 1C). The Palaeozoic succession of the northern and southern sides includes sediments ranging from late Ediacaran to Silurian and from Terreneuvian (Cambrian) to Visean in age, respectively. These successions are affected by large scale, south-verging recumbent folds that display a low to moderate metamorphic grade. Their emplacement took place in Late Visean to Namurian times (Engel et al., 1980; Feist and Galtier, 1985; Echtler and Malavieille, 1990). The Axial Zone consists of plutonic, migmatitic and metamorphic rocks forming a regional ENE-WSW oriented dome (Fig. 1C), where four principal lithological units can be recognized (i) schists and micaschists, (ii) migmatitic orthogneisses, (iii) metapelitic metatexites, and (iv) diatexites and granites (Cocherie, 2003; Faure et al., 2004; Roger et al., 2004, 2015; Bé Mézème, 2005; Charles et al., 2009; Rabin et al., 2015). The Rosis micaschist synform subdivides the eastern Axial Zone into the Espinouse and Caroux sub-domes, whereas the southwestern edge of the Axial Zone comprises the Nore massif.

In the Occitan Domain, two main Cambro–Ordovician felsic events can be identified giving rise to the protoliths of (i) the Larroque metarhyolites in the northern Montagne

Noire and Albigeois, thrust **southward** from Rouergue; and (ii) the migmatitic orthogneisses **that form** the Axial Zone of the Montagne Noire (Fig. 2).

(i) The Larroque volcanosedimentary Complex is a thick (500–1000 m) package of porphyroclastic metarhyolites located on the northern Montagne Noire (Lacaune Mountains), Albigeois (St-Salvi-de-Carcavès and St-Sernin-sur-Rance nappes) and Rouergue; the Variscan setting of the formation is allochthonous in the Albigeois and parautochthonous in the rest. This volcanism emplaced above the Furongian strata and the so-called “Série schisto-gréseuse verte” (see Guérangé-Lozes et al., 1996; Guérangé-Lozes and Alabouvette, 1999), and is encased in the upper part of the Miaolingian La Gardie Formation (Pouclet et al., 2017) (Fig. 2). The Larroque volcanic rocks consist of deformed porphyroclastic rhyolites rich in largely fragmented, lacunous (rhyolitic) quartz and alkali feldspar phenocrysts. The metarhyolites occur as porphyritic lava flows, sills and other associated facies, such as aphyric lava flows, porphyritic and aphyric pyroclastic flows of welded or unwelded ignimbritic types, fine to coarse tephra deposits, and epiclastic and volcanoclastic deposits. **These** rocks are named “augen gneiss” or **augengneiss** and do not display a high-grade gneiss paragenesis but a general lower grade metamorphic mineralogy. The Occitan augengneisses mimic the Ollo de Sapo facies from the Central Iberian Zone because of their large bluish quartz phenocrysts. Based on geochemical similarities and contemporaneous emplacement, Pouclet et al. (2017) suggested that this event also supplied the Davejean acidic volcanic rocks in the Mouthoumet Massif, which represent the southern prolongation of the Montagne Noire (Fig. 2), and the Génis rhyolitic unit of the western Limousin sector.

(ii) Some migmatitic orthogneisses make up the southern Axial Zone, from the western Cabardès to the eastern Caroux domes. The orthogneisses, derived from Ordovician metagranites bearing large K-feldspar phenocrysts, were emplaced at **about** 471 Ma (Somail Orthogneiss, Cocherie et al., 2005), 456 **to** 450 Ma (Pont de Larn and Gorges d’Héric gneisses, Roger et al., 2004) and **ca.** 455 Ma (Sain Eutrope

gneiss, Pitra et al., 2012). They intruded a metasedimentary pile, traditionally known as “Schistes X” and formally named St. Pons-Cabardès Group (Fig. 2). The latter consists of schists, greywackes, quartzites and subsidiary volcanic tuffs and marbles (Demange et al., 1996; Demange, 1999; Alabouvette et al., 2003; Roger et al., 2004; Cocherie et al., 2005). The group is topped by the Sériès Tuff, dated at about 545 Ma (Lescuyer and Cocherie, 1992), which represents a contemporaneous equivalent of the Cadomian Rivernous rhyolitic tuff (542.5 to 537.1 Ma) from the Lodève inlier of the northern Montagne Noire (Álvaro et al., 2014b, 2018; Padel et al., 2017). Age of migmatization has been inferred from U–Pb dates on monacite from migmatites and anatectic granites at 333 to 327 Ma (Bé Mézème, 2005; Charles et al., 2008); as a result, the 330–325 Ma time interval can represent a Variscan crustal melting event in the Axial Zone.

As in the Pyrenees, the Middle Ordovician is absent in the Occitan Domain. Its gap allows distinction between a Lower Ordovician pre-unconformity sedimentary package para- to unconformably overlain by an Upper Ordovician–Silurian succession (Álvaro et al., 2016; Pouclet et al., 2017).

2.4. Sardinia

In Sardinia the Cambro–Ordovician magmatism is well represented in the external (southern) and internal (northern) nappe zones of the exposed Variscan Belt (Fig. 1E), and ranges in age from late Furongian to Late Ordovician. A Furongian–Tremadocian (ca. 491–480 Ma) magmatic activity, predating the Sardic phase, is mostly represented by felsic volcanic and subvolcanic rocks encased in the San Vito sandstone Formation. The Sardic-related volcanic products differ from one nappe to another: intermediate and basic (mostly metandesites and andesitic basalts) are common in the nappe stacking of the central part of the island (Barbagia and Goceano), whereas felsic metavolcanites prevail in the southeastern units. Their age is bracketed between 465

and 455 Ma (Giacomini et al., 2006; Oggiano et al., 2010; Pavanetto et al., 2012; Cruciani et al., 2018) and matches the Sardinian gap based on biostratigraphy (Barca et al., 1988).

Teichmüller (1931) and Stille (1939) were the first to recognize in southwestern Sardinia an intra-Ordovician stratigraphic hiatus. Its linked erosive unconformity is supported by a correlatable strong angular discordance in the Palaeozoic basement of the Iglesias-Sulcis area, External Zone (Carmignani et al., 2001). This major discontinuity separates the Cambrian–Lower Ordovician Nebida, Gonnese and Iglesias groups (Pillola et al., 1998) from the overlying coarse-grained (“Pudding”) Monte Argentu metasediments (Leone et al., 1991, 2002; Laske et al., 1994). The gap comprises a chronostratigraphically constrained minimum gap of about 18 m.y. that includes the Floian and Dapingian (Barca et al., 1987, 1988; Pillola et al., 1998; Barca and Cherchi, 2004) (Fig. 2). The hiatus is related to neither metamorphism nor cleavage, though some E–W folds have been documented in the Gonnese Anticline and the Iglesias Syncline (Cocco et al., 2018), which are overstepped by the “Pudding” metaconglomerates. Both the E–W folds and the overlying metaconglomerates were subsequently affected by Variscan N–S folds (Cocco and Funneda, 2011, 2017). Sardinian-related volcanic rocks are not involved in this area, but Sardinian-inherited palaeoreliefs are lined with breccia slides that include metre- to decametre-scale carbonate boulders (“Olistoliti”), some of them hosting synsedimentary faults contemporaneously mineralized with ore bodies (Boni and Koeppel, 1985; Boni, 1986; Barca, 1991; Caron et al., 1997). The lower part of the unconformably overlying Monte Argentu Formation deposited in alluvial to fluvial environments (Martini et al., 1991; Loi et al., 1992; Loi and Dabard, 1997).

A similar gap was reported by Calvino (1972) in the Sarrabus-Gerrei units of the External Nappe Zone. The so-called “Sarrabese Phase” is related to the onset of thick, up to 500 m thick, volcanosedimentary complexes and volcanites (Barca et al., 1986; Di Pisa et al., 1992) with a Darriwilian age for the protoliths of the metavolcanic rocks

(465.4 to 464 Ma; Giacomini et al., 2006; Oggiano et al., 2010). In the Iglesiente-Sulcis region (Fig. 1E), Carmignani et al. (1986, 1992, 1994, 2001) suggested that the “Sardic-Sarrabese phase” should be associated with the compression of a Cambro–Ordovician back-arc basin that originated the migration of the Ordovician volcanic arc toward the Gondwanan margin.

Some gneissic bodies, interpreted as the plutonic counterpart of metavolcanic rocks, are located in the Bithia unit (e.g., the Monte Filau area, 458 to 457 Ma, surrounded by a Mid–Ordovician andalusite thermal aureole; Pavanetto et al., 2012; Costamagna et al., 2016) and in the internal units (Lodè orthogneiss, ca. 456 Ma; Tanaunella orthogneiss, ca. 458 Ma, Helbing and Tiepolo, 2005; Golfo Aranci orthogneiss, ca. 469 Ma, Giacomini et al., 2006).

The Sardic palaeorelief is sealed by Upper Ordovician transgressive deposits. The sedimentary facies show high variability, but the –mostly terrigenous– sediments vary from grey fine- to medium-sized sandstones, to muddy sandstones and mudstones. They are referred to the Katian Punta Serpeddì and Orroledu formations (Pistis et al., 2016). This post–Sardic sedimentary succession is coeval with a new magmatic pulsation represented by alkaline to tholeiitic within-plate basalts (Di Pisa et al., 1992; Gaggero et al., 2012).

3. Geochemical data

3.1. Materials and methods

The rocks selected for geochemical analysis (231 samples; see tectonostratigraphic location in Fig. 1 and stratigraphic emplacement in Fig. 2) have recorded different degrees of hydrothermalism and metamorphism, as a result of which only the most immobile elements have been considered. The geochemical calculations, in which the

major elements take part, have been made from values recalculated to 100 in volatile free compositions; Fe is reported as FeO_t .

The geochemical dataset of the Central Iberian Zone includes 152 published geochemical data, from which 85 are plutonic and 67 volcanic and volcanoclastic rocks from the Ollo de Sapo Formation (Galicia, Sanabria and Guadarrama areas), and the contact between the Central Iberian and Ossa Morena Zones (Urza Formation and Portalegre and Carrascal granites). Other data were yielded from six volcanic rocks of the Galicia-Trás-os-Montes Zone (Saldanha area) ([Fig. 1B](#); [Repository Data](#)).

The dataset of the eastern Pyrenees consists of 38 samples, six of which are upper Lower Ordovician volcanic rocks, and seven upper Lower Ordovician plutonic rocks, together with nine Upper Ordovician volcanic and 14 Upper Ordovician plutonic rocks ([Repository Data](#)). New data reported below include two samples of subvolcanic sills intercalated in the pre-Sardic unconformity succession (Clariana et al., 2018; Margalef, unpubl.; [Table 1](#)).

The study samples from the Occitan Domain comprise six metavolcanic rocks, four from the Larroque volcanosedimentary Complex in the Albigeois and northern Montagne Noire and two from the Mouthoumet massif (Pouclet et al., 2017) ([Repository Data](#)), and four new samples for the Axial Zone gneisses ([Table 1](#)).

In the Sardinian dataset, 25 published analyses are selected: five correspond to the Golfo Aranci orthogneiss (Giacomini et al., 2006), six to metavolcanics from the central part of the island (Giacomini et al., 2006; Cruciani et al., 2013), and five to metavolcanics and one to gneisses from the Bithia unit (Cruciani et al., 2018) ([Repository Data](#)). Ten new analyses are added from the Monte Filau and Capo Spartivento gneisses of the Bithia unit, and from the Punta Bianca gneisses embedded within the migmatites of the High-grade Metamorphic complex of the Inner Zone ([Table 1](#)).

Whole-rock major and trace elements and rare earth element (REE) compositions were determined at ACME Laboratories, Vancouver, Canada. LiBO_2 fusion followed by

X-ray fluorescence spectroscopy (XRF) analysis was used to determine major elements. Rare earth and refractory elements were measured by ICP–MS following a lithium metaborate/tetraborate fusion and nitric acid digestion on 0.2 g of sample. For base metals, 0.5 g of sample was digested in Aqua Regia at 95 °C and analyzed by inductively coupled plasma - atomic emission spectrometry (ICP–AES). Analyses of standards and duplicate samples indicate precision to better than 1 % for major oxides, and 3–10 % for minor and trace elements.

Additional Sm–Nd isotopic analyses were performed at Centro de Geocronología y Geoquímica Isotópica from the Complutense University, Madrid. They were carried out in whole-rock powders using a ^{150}Nd – ^{149}Sm tracer by isotope dilution-thermal ionization mass spectrometry (ID–TIMS). The samples were first dissolved through oven digestion in sealed Teflon bombs with ultra pure reagents to perform two-stage conventional cation-exchange chromatography for separation of Sm and Nd (Strelow, 1960; Winchester, 1963), and subsequently analysed using a Sector 54 VG-Micromass multicollector spectrometer. The measured $^{143}\text{Nd}/^{144}\text{Nd}$ isotopic ratios were corrected for possible isobaric interferences from ^{142}Ce and ^{144}Sm (only for samples with $^{147}\text{Sm}/^{144}\text{Sm} < 0.0001$) and normalized to $^{146}\text{Nd}/^{144}\text{Nd} = 0.7219$ to correct for mass fractionation. The Lajolla Nd international isotopic standard was analysed during sample measurement, and gave an average value of $^{143}\text{Nd}/^{144}\text{Nd} = 0.5114840$ for 9 replicas, with an internal precision of ± 0.000032 (2σ). These values were used to correct the measured ratios for possible sample drift. The estimated error for the $^{147}\text{Sm}/^{144}\text{Nd}$ ratio is 0.1%.

A general classification of the analyzed samples, following Winchester and Floyd (1977), can be seen in Figure 4A–B, and the geographical coordinates of the new samples in Table 1. For geochemical comparison (summarized in Table 2), two large groups or suites are differentiated in order to check the similarities and differences between the magmatic rocks, and to infer a possible geochemical trend following a

palaeogeographic SW–NE transect. The description reported below follows the same palaeogeographic and chronological order.

3.2. Furongian–to–Mid Ordovician Suite

In the Central Iberian and Galicia-Trás-os-Montes Zones, the Furongian–to–Mid Ordovician magmatic activity is pervasive. Their main representative is the Ollo de Sapo Formation, which includes volcanic and subvolcanic rocks (67 samples) as well as plutonic rocks (85 samples) (data from Murphy et al., 2006; Díez-Montes, 2007; Montero et al., 2007, 2009; Solá, 2007; Solá et al., 2008; Talavera, 2009; Villaseca et al., 2016). From the Parautochthon Schistose Domain of the Galicia-Trás-os Montes Zone, six samples of rhyolite tuffs of the Saldanha Formation (Dias da Silva et al., 2014) are selected, which share geochemical features with the Ollo de Sapo Formation. In summary, five facies are differentiated in the Central Iberian and Galicia-Trás-os Montes Zones: the Ollo de Sapo orthogneisses, some leucogneisses, metagranites and volcanic rocks, and the San Sebastián orthogneiss (for a geochemical characterization, see Table 2).

In the central and eastern Pyrenees, an Early–Mid Ordovician magmatic activity gave rise to the intrusion of voluminous (about 500–3000 m in size) aluminous granitic bodies, encased into the Canaveilles beds (Álvaro et al., 2018; Casas et al., 2019). They constitute the protoliths of the large orthogneissic laccoliths that form the core of the domal massifs scattered throughout the backbone of the Pyrenees. Rocks of the Canigó, Roc de Frausa and Albera massifs have been taken into account in this work, in which volcanic rocks of the Pierrefite and Albera massifs, and the so-called G2 and G3 orthogneisses by Guitard (1970) are also included. All subgroups vary compositionally from subalkaline andesite to rhyolite, as illustrated in the Pearce's (1996) diagram of Figure 5 (data compiled from Vilà et al., 2005; Castiñeiras et al., 2008b; Liesa et al., 2011; Navidad et al., 2018).

Although most rocks in this area are acidic, it is remarkable the presence of minor mafic bodies (Cortalet and Marialles metabasites, not studied in this work), which could indicate a mantle connection with parental magmas during the Mid and Late Ordovician. As well, it should be noted that there are no andesitic rocks in the area.

In the Occitan Domain, six samples of the Larroque volcanosedimentary Complex (Early Tremadocian in age) represent basin floors and subaerial explosive and effusive rhyolites (Pouclet et al., 2017). The porphyroclastic rocks of the Larroque metarhyolites were sampled in the Saint-Géraud and Larroque areas from the Saint-Sernin-sur-Rance nappe and the Saint-André klippe above the Saint-Salvi-de-Carcavès nappe (Pouclet et al., 2017).

In the Middle Ordovician rocks of Sardinia, 11 samples are selected, five of which correspond to orthogneisses of the Aranci Gulf, in the Inner Zone of the NE island (Giacomini et al., 2006), completed with six volcanic rocks of the External Zone (Giacomini et al., 2006; Cruciani et al., 2018) (Table 2).

3.3 Upper Ordovician Suite

In the central and eastern Pyrenees, four Upper Ordovician subgroups are distinguished based on their field occurrence and geochemical and geochronological features: the G1-type orthogneisses *sensu* Guitard (1970); the Cadí and Casemí orthogneisses and the metavolcanic rocks that include the Ribes de Freser rhyolites; the Els Metges volcanic tuffs; and the rhyolites from Andorra and Pallaresa areas (the latter dated at ca. 453 Ma; Clariana et al., 2018) (Table 2). The suite is completed with the Smail orthogneisses of the Axial Montagne Noire (dated at ca. 450 Ma at Gorges d'Héric; Roger et al., 2004) and the orthogneisses from the Sardinian External Zone (dated at ca. 458–457 Ma at Monte Filau; Pavanetto et al., 2012) and the volcanic rocks from the Sardinian Nappe Zone (Table 2).

4. Geochemical framework

A geochemical comparison between the Furongian–Ordovician felsic rocks of all the above-reported groups offers the opportunity to characterize the successive sources of crustal-derived melts along the south-western European margin of Gondwana.

The geochemical features point to a predominance of materials derived from the melting of metasedimentary rocks, rich in SiO_2 and K_2O (average $\text{K}_2\text{O}/\text{Na}_2\text{O} = 2.25$) and peraluminous ($0.4 < C_{\text{norm}} < 4.5$ and $0.94 < A/\text{CNK} < 3.12$), with only three samples with $A/\text{CNK} < 1$ (samples 100786 of the Casemí subgroup, and T26 and T27 of the San Sebastián subgroup).

The result of plotting the REE content vs. average values of continental crust (Rudnick and Gao, 2004; Fig. 6) yields a flat spectra and a base level shared by most of the considered groups. The total content in REE is moderate to high (average REE = 176 ppm, ranging between 482.2 and 26.0 ppm; Fig. 7), with a maximum in the subgroup of the Middle Ordovician volcanic rocks from Sardinia (average REE = 335 ppm, *VOL-SMO*), and with LREE values more fractionated than HREE ones, and negative anomalies of Eu, which would indicate a characteristic process of magmatic evolution with plagioclase fractionation. These features are common in peraluminous granitoids.

All subgroups display similar chondritic normalized REE patterns (Fig. 7), with an enrichment in LREE relative to HREE, which should indicate the involvement of crustal materials in their parental magmas. Nevertheless, some variations can be highlighted, such as the lesser fractionation in REE content of some subgroups. These are the leucogneisses from the Iberian massif (*LG*, $\text{La}/\text{Yb}_n = 2.01$), the Upper Ordovician orthogneisses from Sardinia (*OG-SUO*, $\text{La}/\text{Yb}_n = 2.94$), the Casemí orthogneisses ($\text{La}/\text{Yb}_n = 4.42$) and the Middle Ordovician volcanic rocks from Sardinia (*OG-SUO*, $\text{La}/\text{Yb}_n = 2.94$). This may be interpreted as a greater degree of partial fusion in the origin of their parental magmas (Rollinson, 1993).

There are three geochemical groups displaying $(\text{Gd/Yb})_n$ values > 2 , and $(\text{La/Yb})_n$ values ≥ 9 . These groups are OSS (Central Iberian Zone), VOL-OD (Occitan Domain) and G1 (Pyrenees), and share higher alkalinity features.

Some V1 rocks from the Pyrenees (Pierrefite Formation) show no negative anomalies in Eu. Their parental magmas could have been derived from deeper origins and related to residual materials of the lower continental crust, in areas of production of K-rich granites (Taylor and McLennan, 1989).

The spider diagrams (Fig. 8), however, exhibit strong negative anomalies in Nb, Sr and Ti, which indicate a distinct crustal affiliation (Díez-Montes, 2007). Only the San Sebastián orthogneisses (OSS) show distinct discrepancies in respect of the remaining samples from the Ollo de Sapo Formation. They display lower negative anomalies in Nb and a more alkaline character by comparison with the rest of the Ollo de Sapo rocks, which point to alkaline affinities and greater negative anomalies in Nb.

Despite some small differences in the chemical ranges of some major elements, most felsic Ordovician rocks from the Iberian massif (Central Iberian and Galicia-Trás-os Montes Zones), eastern Pyrenees, Occitan Domain and Sardinia share a common chemical pattern. The Lower–Middle Ordovician rocks of the eastern Pyrenees show less variation in the content of Zr and Nb (Fig. 8B). The volcanic rocks of these groups show a different REE behaviour, which would indicate different sources. Two groups are distinguished in Figure 7, one with greater enrichment in REE and negative anomaly of Eu, and another with lesser content of HREE and without Eu negative anomalies.

Figure 9 illustrates how the average of all the considered groups approximates the mean values of the Rudnick and Gao's (2003) upper continental crust (UCC). In this figure, small deviations can be observed, some of them toward lower continental crust (LCC) values and others toward bulk continental crust (BCC), indicating variations in their parental magmas but with quite similar spectra. Overall chondrite-normalized

patterns are close to the values that represent the upper continental crust, with slight enrichments in the Th/Nb, Th/La and Th/Yb ratios.

Finally, in the Occitan volcanic rocks (*VOL-OD*) the rare earth elements are enriched and fractionated ($33.2 \text{ ppm} < \text{La} < 45.6 \text{ ppm}$; $11.2 < \text{La/Yb} < 14.5$). The upper continental crust normalized diagram exhibits negative anomalies of Ti, V, Cr, Mn and Fe associated with oxide fractionation, of Zr and Hf linked to zircon fractionation, and of Eu related to plagioclase fractionation. The profiles are comparable to the Vendean Saint-Gilles rhyolitic ones. The Th vs. Rb/Ba features are also similar to those of the Saint-Gilles rhyolites, and the Iberian Ollo de Sapo and Urre rhyolites (Solá et al., 2008; Díez Montes et al., 2010).

4. Discussion

4.1 Inferred tectonic settings

In order to clarify the evolution of geotectonic environments, the data have been represented in different **discrimination** diagrams. The Zr/TiO₂ ratio (Lentz, 1996; Syme, 1998) is a key index of compositional evolution for intermediate and felsic rocks. In the Syme diagram ([Fig. 10](#)), most rocks from the Central Iberian Zone represent a characteristic arc association, although there are some contemporaneous samples characterized by extensional-related values ($\text{Zr/Ti} = 0.10$, *LG*). The rocks of the Middle–Ordovician San Sebastián orthogneisses (*OSS*) show values of $\text{Zr/Ti} = 0.08$, intermediate between extensional and arc conditions. This could be interpreted as a sharp change in geotectonic conditions toward the Mid Ordovician ([Fig. 10A](#)). For a better comparison, the samples of the San Sebastián orthogneisses (*OSS*) and the granites (*GRA*) have been distinguished with a shaded area in all the diagrams, since they have slightly different characteristics to the rest of the samples from the Ollo de Sapo group. The samples *G1* (Pyrenees) and *VOL* (Central Iberian Zone) broadly

share similar values, as a result of which, the three latter groups (*OSS*, *G1* and *VOL*) arrange following a good correlation line. The same trend seems to be inferred in the eastern Pyrenees (Fig. 10B), where the Middle Ordovician subgroups display arc features, but half of the Upper Ordovician subgroups show extensional affinities (*G1* and Casemí orthogneisses). In the case of the Occitan orthogneisses (Fig. 10C), they show arc characters, which contrast with the contemporaneous volcanic rocks displaying extensional values with $Zr/Ti = 0.10$. This disparity between plutonic and volcanic rocks could be interpreted as different conditions for the origin of these magmas. In Sardinia (Fig. 10D), the same evolution from arc to extensional conditions is highlighted for the Upper Ordovician samples, although some Middle Ordovician volcanic rocks already shared extensional patterns ($Zr/Ti = 0.09$). In summary, there seems to be a geochemical evolution in the Ordovician magmas grading from arc to extensional environments.

In the Nb–Y tectonic discriminating diagram of Pearce et al. (1984) (Fig. 11), most samples plot in the volcanic arc-type, though some subgroups project in the within-plate and anomalous ORG. The majority of samples display very similar Zr/Nb and Nb/Y ratios, typical of island arc or active continental margin rhyolites (Díez-Montes et al., 2010). Only some samples plot separately: *OSS* samples with highest Nb contents (>20 ppm), and some volcanic rocks of the Occitan Domain (average Nb =16.87 ppm). In the eastern Pyrenees, the Middle Ordovician rocks plot in the volcanic arc field, whereas the Upper Ordovician ones point in the ORG type, except the Casemí samples. This progress of magmatic sources agrees with the evolution seen in Figure 10. In the Occitan Domain, *VOL-OD* samples share values with those of the San Sebastián orthogneiss, while *OG-OD* shares values with those of *OG* from the Central Iberian Zone.

The Zr vs. Nb diagram (Leat et al., 1986; modified by Piercey, 2011) (Fig. 12) illustrates how magmas evolved toward richer values in Zr and Nb, which is consistent with what it is observed in the Syme diagram (Fig. 10). Figure 12A documents how

most samples show a general positive **correlation**. These different groups correspond to the OSS and Portalegre granites, highlighted in the figure. The two groups indicate a tendency toward alkaline magmas. Some samples, such as the Pyrenean *G1*, some Occitan *VOL-OD* samples and some Sardinian *OG-UOS* samples share the same affinity, clearly distinguished from the general geochemical trend exhibited by the Central Iberian Zone.

On a Zr vs. Ga/Al diagram (Whalen et al., 1987) (Fig. 13), the samples depict an intermediate character between **anorogenic or alkaline (A-type) and orogenic (I&S-type)**. In the Central Iberian Zone, samples from the San Sebastián orthogneisses and Portalegre granites show characters of A-type granites, while the remaining samples display affinities of I&S-type granites. For the Central Iberian Zone, a clear magmatic shift toward more extensional geotectonic environments is characterized. For the eastern Pyrenees, we find the same situation **as in** the Central Iberian Zone, with a magmatic evolution toward A-granite type characteristics, indicating more extensional geotectonic environments. In the Occitan Domain, the samples show a clear I&S character. In the Sardinian case, the same seems to happen as in the Central Iberian Zone: the Upper Ordovician orthogneisses suggest a more extensional character.

In summary, all the reported diagrams point to a magmatic evolution through time, grading from arc to extensional geotectonic environments (with increased Zr/Ti ratios) and to granite type-A characters. This geotectonic framework is consistent with that illustrated in Figure 10. The geochemical characters of these rocks show a rhyodacite to dacite composition, peraluminous and calc-alkaline K-rich character, and an arc-volcanic affinity for most of samples, but without intermediate rocks associated with andesitic types. Hence a change in time is documented toward more alkaline magmas.

4.2 Interpretation of ϵ_{Nd} values

$\epsilon\text{Nd}_{(t)}$ values are useful to interpret the nature of magmatic sources. Most samples of the above-reported groups show no significant differences in isotopic $\epsilon\text{Nd}_{(t)}$ values, and Nd_{CHUR} model ages (Fig. 14). Some exceptions are related to granites from the southern Central Iberian Zone, which display positive values (from +2.6 to –2.4) and T_{DM} values from 0.90 to 3.46 Ga. These granites, space-related with calcalkaline diorites and gabbros, were interpreted by Solá et al. (2008) as the result of underplating and temporal storage of mantle-derived magmas as a potential source for the intrusive “orogenic melts” during Early Palaeozoic extension.

Some samples from (i) the Central Iberian Zone, such as VI-3 (Leucogneiss subgroup) and PORT2 and PORT15 (Granite subgroup); (ii) the eastern Pyrenees, such as 99338 (G1 subgroup) and 100786 samples (Casemí subgroup); and (iii) the Sardinian CS5, CS8 and CC5 samples (Upper Ordovician Orthogneiss subgroup) display anomalous T_{DM} values and $^{147}\text{Sm}/^{144}\text{Nd}$ ratios > 0.17 (Table 2; Fig. 14), a character relatively common in some felsic rocks (DePaolo, 1988; Martínez et al., 2011). According to Stern et al. (2012), these values should not be considered, but a possible explanation for these high ratios may be related to the M-type tetrad effect (e.g., Irber, 1999; Monecke et al., 2007; Ibrahim et al., 2015), which affects REE fractionation in highly evolved felsic rocks due to the interaction with hydrothermal fluids. This process can be reflected as an enrichment of Sm related to Nd. Other authors, however, explain this enrichment as a result of both magmatic evolution (e.g., McLennan, 1994; Pan, 1997) and weathering processes after exhumation (e.g., Masuda and Akagi, 1989; Takahashi et al., 2002).

In the granites of the southern Central Iberian Zone and the volcanic rocks of Sardinia, positive values in $\epsilon\text{Nd}_{(t)}$ could be interpreted as a more primitive nature of their parental magmas, even though the samples with highest T_{DM} values are those that display higher $^{147}\text{Sm}/^{144}\text{Nd}$ ratios (> 0.17 ; Table 2).

The volcanic rocks of the Central Iberian Zone display some differences following a N-S transect, being $\epsilon\text{Nd}_{(t)}$ values less variable in the north ($\epsilon\text{Nd}_{(t)}$: –4.0 to –5.0) than in

the south ($\epsilon\text{Nd}_{(t)}$: -1.6 to -5.5). The isotopic signature of the Urrea volcanoclastic rocks is compatible with magmas derived from young crustal rocks, with intermediate to felsic igneous compositions (Solá et al., 2008). The volcanic rocks of the northern Central Iberian Zone could be derived from old crustal rocks (Montero et al., 2007). The isotopic composition of the granitoids from the southern Central Iberian Zone has more primitive characters than those of the northern Central Iberian Zone, suggesting different sources for both sides (Talavera et al., 2013). OSS shows lower inheritance patterns, more primitive Sr–Nd isotopic composition than other rocks of the Ollo de Sapo suite, and an age some 15 m.y. younger than most meta-igneous rocks of the Sanabria region (Montero et al., 2009), likely reflecting a greater mantle involvement in its genesis (Díez-Montes et al., 2008).

According to Talavera et al. (2013), the Cambro–Ordovician rocks of the Galicia-Trás-os-Montes Zone schistose area and the magmatic rocks of the northern Central Iberian Zone are contemporary. Both metavolcanic and metagranitic rocks almost share the same isotopic compositions.

The Upper Ordovician orthogneisses from the Occitan Domain show very little variation in $\epsilon\text{Nd}_{(t)}$ values (-3.5 to -4.0), typical of magmas derived from young crustal rocks. The variation in TDM values is also small (1.4 to 1.8 Ga) indicating **similar** crustal residence times **to other rock groups**.

In Sardinia, $\epsilon\text{Nd}_{(t)}$ values present a greater variation (-1.6 to -3.3), but they are also included in the typical continental crustal range. As noted above, anormal TDM values (between 1.2 to 4.5 Ga) may be due to post-magmatic hydrothermal alteration processes.

5. Geodynamic setting

In the Iberian Massif, the Ediacaran–Cambrian transition was marked by paraconformities and angular discordances indicating the passage from Cadomian

volcanic arc to rifting conditions. The axis of the so-called Ossa-Morena Rift lies along the homonymous Zone (Quesada, 1991; Sánchez-García et al., 2003, 2008, 2010) close to the remains of the Cadomian suture (Murphy et al., 2006). Rifting conditions were accompanied by a voluminous magmatism that changed from peraluminous acid to bimodal (Sánchez-García et al., 2003, 2008, 2016, 2019). Some authors (Álvaro et al., 2014; Sánchez-García et al., 2019) propose that this rift resulted from a SW-to-NE inward migration, toward innermost parts of Gondwana, of rifting axes from the Anti-Atlas in Morocco to the Ossa-Morena Zone in the Iberian Massif. According to this proposal the rifting developed later (in Cambro–Ordovician times) in the Iberian, Armorican and Bohemian massifs.

The Furongian–Ordovician transition to drifting conditions is associated, in the Iberian Massif, Occitan Domain, Pyrenees and Sardinia, with a stepwise magmatic activity contemporaneous with the record of the Toledanian and Sardic unconformities. These, related to neither metamorphism nor penetrative deformations, are linked to uplift, erosion and irregularly distributed mesoscale deformation that gave rise to angular unconformities up to 90°. The time span involved in these gaps is similar (22 m.y. in the Iberian Massif, 16–23 m.y. in the Pyrenees and 18 m.y. in Sardinia). This contrasts with the greater time span displayed by the magmatic activity (30–45 m.y.), which started before the unconformity formation (early Furongian in the Central Iberian Zone vs. Floian in the Pyrenees, Occitan Domain and Sardinia), **continued** during the unconformity formation (Furongian and early Tremadocian in the Central Iberian Zone vs. Floian–Darriwilian in the Pyrenees, Occitan Domain and Sardinia), and ended during the sealing of the uplifted and eroded palaeorelief (Tremadocian–Floian volcaniclastic rocks at the base of the Armorican Quartzite in the Central Iberian Zone vs. Sandbian–Katian volcanic rocks at the lowermost part of the Upper Ordovician successions in the Pyrenees, Occitan Domain and Sardinia; Gutiérrez-Alonso et al., 2007, 2016; Navidad et al., 2010; Martínez et al., 2011; Álvaro et al., 2016; Martí et al., 2019). In the Pyrenees, Upper Ordovician magmatism and sedimentation coexist with

normal faults controlling marked thickness changes of the basal Upper Ordovician succession and cutting the lower part of this succession, the Sardinian unconformity and the underlying Cambro–Ordovician sequence (Puddu et al., 2018, 2019).

Although the Toledanian and Sardinian Phases reflect similar geodynamic conditions in two distinct palaeogeographic areas, at present forming the western and eastern branches of the Variscan Ibero-Armorican Arc, they display different peaks in magmatic activity with a minor chronological overlapping (Fig. 3). This may reflect a SW-to-NE “zip-like” propagation of the latest Ediacaran–Terreneuvian rifting axes in the so-called Atlas-Ossa Morena Rift.

Toledanian Phase

The Early Ordovician (Toledanian) magmatism of the Central Iberian Zone evolved to a typical passive-margin setting, with geochemical features dominated by acidic rocks, peraluminous and rich in K, and lacking any association with basic or intermediate rocks. Some of the orthogneisses of the Galicia-Trás-os-Montes Zone basal and allochthonous complex units share these same patterns. This fact has been interpreted by some authors as a basin environment subject to important episodes of crustal extension (Martínez-Catalán et al., 2007; Díez-Montes et al., 2010). In contrast, Villaseca et al. (2016) interpreted this absence as evidence against rifting conditions, though the absence of contemporary basic magmatism may be explained by the partial fusion of a thickened crust, through recycling of Neoproterozoic crustal materials. The thrust of a large metasedimentary sequence could generate dehydration and metasomatism of the rocks above this sequence, triggering partial fusion at different levels, although the increase in peraluminosity with the basicity of the orthogneisses is against any AFC process involving mantle materials. However, this increase in peraluminosity with the basicity has not been revealed in the samples studied above. Following Villaseca et al.’s (2016) model, a flat subduction of the southern part of the

Central Iberian Zone would have taken place under its northern prolongation, whereas the reflection of such a subduction is not evident in the field. The calc-alkaline signature of this magmatism has also been taken into account as proof of its relationship with volcanic-arc environments (Valverde-Vaquero and Dunning, 2000). However, calc-alkaline features may be also interpreted as a result of a variable degree of continental crustal contamination and/or previously enriched mantle source (Sánchez-García et al., 2003, 2008, 2016, 2019; Díez-Montes et al., 2010). Finally, other granites not considered here of Tremadocian age have been reported in the southern Central Iberian Zone, such as the Oledo massif and the Beira Baixa-Central Extremadura, which display a I-type affinity (Antunes et al., 2009; Rubio Ordóñez et al., 2012). These granites could represent different sources for the Ordovician magmatism in the Central Iberian Zone.

Sánchez-García et al. (2019) have proposed that the anomaly that produced the large magmatism throughout the Iberian Massif could have migrated from the rifting axis to inwards zones and the acid, peraluminous, K-rich rocks of Mid Ordovician in age should represent the initial stages of a new rifting pulse, resembling the peraluminous rocks of the Early Rift Event *sensu* Sánchez-García et al. (2003) from the Cambrian Epoch 2 of the Ossa-Morena Rift.

In the parautochthon of the Galicia-Trás-os-Montes Zone, the appearance of tholeiitic and alkaline-peralkaline magmatism in the Mid Ordovician would signal the first steps toward extensional conditions (Díez Fernández et al., 2012; Dias da Silva et al., 2016). In the Montagne Noire and the Mouthoumet massifs contemporaneous tholeiitic lavas indicate a similar change in the tectonic regimen (Álvaro et al., 2016). This gradual change in geodynamic conditions is also marked by the appearance of rocks with extensional characteristics in some of subgroups considered here, such as the Central Iberian Zone (San Sebastián orthogneisses), eastern Pyrenees (Casemí orthogneisses, and G1), volcanic rocks of the Occitan Domain, and the orthogneisses and volcanic rocks from Sardinia.

Sardic Phase

In the eastern Pyrenees, two peaks of **Ordovician** magmatic activity **are observed** (Casas et al., 2019). Large Lower–Middle Ordovician peraluminous granite bodies are known representing the protoliths of numerous gneissic bodies with laccolithic morphologies. In the Canigó massif, the Upper Ordovician granite bodies (protoliths of Cadí, Casemí, G1) are encased in sediments of the Canaveilles and Jujols groups. During this time span, there was generalized uplift and erosion that culminated with the onset of the Sardic unconformity. The Sardic Phase was succeeded by an extensional **interval** related to the formation of normal faults affecting the pre–unconformity strata (Puddu et al., 2018, 2019). The volcanic arc signature can be explain by crustal recycling (Navidad et al., 2010; Casas et al., 2010; Martínez et al., 2011), as in the case of the Toledanian Phase in the Central Iberian Zone, although, according to Casas et al. (2019), the Pyrenees and the Catalan Coastal Ranges were probably fringing the Gondwana margin in a different position than that occupied by the Iberian Massif. As a whole, the Ordovician magmatism in the Pyrenees lasted about 30 m.y., from ca 477 to 446 Ma, in a time span contemporaneous with the formation of the Sardic unconformity (Fig. 2). Recently, Puddu et al. (2019) proposed that a thermal doming, bracketted between 475 and 450 Ma, **could** have stretched the Ordovician lithosphere. The emersion and denudation of the inherited Cambrian–Ordovician palaeorelief would have given rise to the onset of the Sardic unconformity. According to these authors, thermal doming triggered by hot mafic magma underplating may also be responsible for the late Early–Late Ordovician coeval magmatic activity.

In the Occitan Domain, there was a dramatic volcanic event in early Tremadocian times, with the uprising of basin floors and the subsequent effusion of abundant rhyolitic activities under subaerial explosive conditions (Larroque volcanosedimentary Complex in the Montagne Noire, and Davejean acidic volcanic counterpart in the

Mouthoumet Massif). Pouclet et al., (2017) interpreted this as a delayed Ollo de Sapo-style outpouring where a massive crustal melting required a rather significant heat supply. Asthenospheric upwelling leading to the interplay of lithospheric doming, continental break-up, and a decompressionally driven mantle melting can explain such a great thermal anomaly. The magmatic products accumulated on the mantle-crust contact would provide enough heat transfer for crustal melting (Huppert and Sparks, 1988). Subsequently, a post-Sardic reactivation of rifting conditions is documented in the Cabrières klippen (southern Montagne Noire) and the Mouthoumet massif. There, a Late Ordovician fault-controlled subsidence linked to the record of rift-related tholeiites (Roque de Bandies and Villerouge formations) were contemporaneous with the record of the Hirnantian glaciation (Álvaro et al., 2016). Re-opening of rifting branches (Montagne Noire and Mouthoumet massifs) was geometrically recorded as overlapping patterns and final sealing of Sardic palaeoreliefs by Silurian and Lower Devonian strata.

Sardinia illustrates an almost complete record of the Variscan Belt (Carmignani et al., 1994; Rossi et al., 2009). Some plutonic orthogneisses of the Inner Zone belong to this cycle, such as the orthogneisses of Golfo Aranci (Giacomini et al., 2006). Gaggero et al. (2012) described three magmatic cycles. The first cycle is well represented in the Sarrabus unit by Furongian–Tremadocian volcanic and subvolcanic interbeds within a terrigenous succession (San Vito Formation) which is topped by the Sardic unconformity. Some plutonic orthogneisses of the Inner Zone belong to this cycle, such as the orthogneisses of Golfo Aranci (Giacomini et al., 2006) and the PB orthogneiss of Punta Bianca). The second Mid–Ordovician cycle, about 50 m.y. postdating the previous cycle, is of an arc-volcanic type with calc-alkaline affinity and acidic-to-intermediate composition. The acidic metavolcanites are referred in the literature as “porphyroids”, which crop out in the External Nappe Zone and some localities of the Inner Zone. The intermediate to basic derivatives are widespread in Central Sardinia (Serra Tonnai Formation). Some plutonic rocks (Mt. Filau orthogneisses and Capo

Spartivento) of the second cycle are discussed above. The third cycle consists of alkalic meta-epiclastites interbedded in post–Sandbian strata and metabasites marking the Ordovician/Silurian contact and reflecting rifting conditions. In this work only the first two cycles **are** considered. Giacomini et al. (2006) cite coeval mafic rocks of felsic magmatism of Mid Ordovician age (Cortesogno et al., 2004; Palmeri et al., 2004; Giacomini et al., 2005), although they interpret a subduction scenario of the Hun terrain below Corsica and Sardinia in the Mid Ordovician.

Origin of intracrustal siliceous melts

In this scenario, the key to generate large volumes of acidic rocks in an intraplate context would be the existence of a lower-middle crust, highly hydrated, in addition to a high heat flow, possibly caused by mafic **melts** (Bryan et al., 2002; Díez-Montes, 2007). This could be the scenario **initiated** by the arrival of a thermal anomaly in a subduction-free area (Sánchez-García et al., 2003, 2008, 2019; Álvaro et al., 2016). The formation of large volumes of intracrustal siliceous melts could act as a viscous barrier, preventing the rise of mafic magmas within volcanic environments, and causing the underplating of these magmas at the contact between the lower crust and the mantle (Huppert and Sparks, 1988; Pankhurst et al., 1998; Bindeman and Valley, 2003). The cooling of these magmas could lead to crustal thickening and in this case, the volcanic arc signature can be explained by crustal recycling (Navidad et al., 2010; Díez-Montes et al., 2010; Martínez et al., 2011).

Sánchez-García et al. (2019) have proposed that the anomaly that produced the large magmatism throughout the Iberian Massif could have migrated from the rifting axis to inwards zones and the acid, peraluminous, K-rich rocks of Mid Ordovician in age should represent the initial stages of a new rifting pulse, resembling the peraluminous rocks of the Early Rift Event *sensu* Sánchez-García et al. (2003) from the Cambrian Epoch 2 of the Ossa-Morena Rift. In the parautochthon of the Galicia-Trás-

os-Montes Zone, the appearance of tholeiitic and alkaline-peralkaline magmatism in the Mid Ordovician would signal the first steps toward extensional conditions (Díez Fernández et al., 2012; Dias da Silva et al., 2016). In the Montagne Noire and the Mouthoumet massifs contemporaneous tholeiitic lavas indicate a similar change in the tectonic regimen (Álvaro et al., 2016). This change in geodynamic conditions is also marked by the appearance of rocks with extensional characteristics in some of subgroups considered here, such as the Central Iberian Zone (San Sebastián orthogneisses), eastern Pyrenees (Casemí orthogneisses, and G1), volcanic rocks of the Occitan Domain, and the orthogneisses and volcanic rocks from Sardinia. In the Pyrenees, Puddu et al. (2019) proposed that a thermal doming, between 475 and 450 Ma, should have stretched the Ordovician lithosphere leading to emersion and denudation of a Cambrian–Ordovician palaeorelief, and giving rise to the onset of the Sardinic unconformity. According to these authors, thermal doming triggered by hot mafic magma underplating may also be responsible for the late Early–Late Ordovician coeval magmatic activity

A major continental break-up, leading to the so-called Tremadocian Tectonic Belt, was suggested by Pouclet et al. (2017), which initiated by upwelling of the asthenosphere and tectonic thinning of the lithosphere. Mantle-derived mafic magmas were underplated at the mantle-crust transition zone and intruded the crust. These magmas provided heat for crustal melting, which supplied the rhyolitic volcanism. After emptying the rhyolitic crustal reservoirs, the underlying mafic magmas finally rose and reached the surface. According to Pouclet et al. (2017), the acidic magmatic output associated with the onset of the Larroque metarhyolites resulted in massive crustal melting requiring a rather important heat supply. Asthenospheric upwelling leading to lithospheric doming, continental break-up, and a decompressionally driven mantle melting can explain such a great thermal anomaly. Magmatic products accumulated on the mantle-crust contact providing enough heat transfer for crustal melting.

6. Conclusions

A geochemical comparison of 231 plutonic and volcanic samples of two major suites, Furongian–Mid Ordovician and Late Ordovician in age, from the Central Iberian and Galicia-Trás-os-Montes Zones of the Iberian Massif and in the eastern Pyrenees, Occitan Domain (Albigeois, Montagne Noire and Mouthoumet massifs) and Sardinia points to a predominance of materials derived from the melting of metasedimentary rocks, peraluminous and rich in SiO_2 and K_2O . The total content in REE is moderate to high. Most felsic rocks display similar chondritic normalized REE patterns, with an enrichment of LREE relative to HREE, which should indicate the involvement of crustal materials in their parental magmas.

Zr/ TiO_2 , Zr/Nb, Nb/Y and Zr vs. Ga/Al ratios, and REE and ϵ_{Nd} values reflect contemporaneous arc and extensional scenarios, which progressed to distinct extensional conditions finally associated with outpouring of mafic tholeiitic-dominant rifting lava flows. Magmatic events are contemporaneous with the formation of the Toledanian (Furongian–Early Ordovician) and Sardic (Early–Late Ordovician) unconformities, related to neither metamorphism nor penetrative deformation. The geochemical and structural framework precludes subduction generated melts reaching the crust in a magmatic arc to back-arc setting. On the contrary, it favours partial melting of sediments and/or granitoids in a continental lower crust triggered by the underplating of hot mafic magmas related to the opening of the Rheic Ocean as a result of asthenospheric upwelling.

7. Acknowledgements

The authors thank the constructive and useful revisions made by Laura Gaggero (Genoa, Italy) and Jochen Mezger (Fairbanks, USA). This paper is a contribution to projects CGL2017-87631-P and PGC2018-093903-B-C22 from Spanish Ministry of

Science and Innovation. We acknowledge support of the publication fee by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

Data availability - All data included in the paper and the Repository Data.

Author contributions - JJA, TSG and JMC: Methodology (Lead), Supervision (Lead), Writing – Original Draft (Lead), Writing – Review & Editing (Lead); CP, ADM, ML & GO: Methodology (Supporting), Supervision (Supporting), Writing – Original Draft (Supporting), Writing – Review & Editing (Supporting).

Competing interests - No competing interests

References

- Alabouvette, B., Demange, M., Guérangé-Lozes, J., Ambert, P., 2003. Notice explicative de la carte géologique de Montpellier au 1/250 000. BRGM, Orléans.
- Álvaro, J.J., Vizcaíno, D., 2018. The Furongian break-up (rift-drift) transition in the Anti-Atlas, Morocco. *J. Iberian Geol.* 44, 567–587.
- Álvaro, J.J., Ferretti, F., González-Gómez, C., Serpagli, E., Tortello, M. F., Vecoli, M., Vizcaíno, D., 2007. A review of the Late Cambrian (Furongian) palaeogeography in the western Mediterranean region, NW Gondwana. *Earth-Sci. Rev.* 85, 47–81.
- Álvaro, J.J., Ezzouhairi, H., Ribeiro, M.L., Ramos, J.F., Solá, A.R., 2008. Early Ordovician volcanism of the Iberian Chains (NE Spain) and its influence on preservation of shell concentrations. *Bull. Soc. géol. France* 179(6), 569–581.
- Álvaro, J.J., Bellido, F., Gasquet, D., Pereira, F., Quesada, C., Sánchez-García, T., 2014a. Diachronism of late Neoproterozoic–Cambrian arc-rift transition of North

- 939 Gondwana: a comparison of Morocco and the Iberian Ossa-Morena Zone. *J. Afr.*
 940 *Earth Sci.* 98, 113–132.
- 941 Álvaro, J.J., Bauluz, B., Clausen, S., Devaere, L., Gil Imaz, A., Monceret, E., Vizcaïno,
 942 D., 2014b. Stratigraphy of the Cambrian–Lower Ordovician volcanosedimentary
 943 complexes, northern Montagne Noire, France. *Stratigraphy* 11, 83–96.
- 944 Álvaro, J.J., Colmenar, J., Monceret, E., Pouclet, A., Vizcaïno, D., 2016. Late
 945 Ordovician (post–Sardic) rifting branches in the North Gondwanan Montagne Noire
 946 and Mouthoumet massifs of southern France. *Tectonophysics* 681, 111–123.
- 947 Álvaro, J.J., Casas, J.M., Clausen, S., Quesada, C., 2018. Early Palaeozoic
 948 geodynamics in NW Gondwana. *J. Iberian Geol.* 44, 551–565.
- 949 Álvaro, J.J., Cortijo, I., Jensen, S., Lorenzo, S., Palacios, T., Pieren, A., 2019. Updated
 950 stratigraphic framework and biota of the Ediacaran and Terreneuvian in the Alcudia-
 951 Toledo Mountains of the Central Iberian Zone, Spain. *Est. Geol.* 75(2), e093.
- 952 Antunes, I.M.H.R., Neiva, A.M.R., Silva, M.M.V.G., Corfu, F., 2009. The genesis of I-
 953 and S-type granitoid rocks of the Early Ordovician Oledo pluton, Central Iberian
 954 Zone (central Portugal). *Lithos* 111, 168–185.
- 955 Arthaud, F., 1970. Etude tectonique et microtectonique comparée de deux domaines
 956 hercyniens: les nappes de la Montagne Noire (France) et l'anticlinorium de
 957 l'Iglesiente (Sardaigne). PhD, Univ. Montpellier.
- 958 Ayora, C., 1980. Les concentrations métal·liques de la Vall de Ribes. PhD, Univ.
 959 Barcelona.
- 960 Ballèvre, M., Fourcade S., Capdevila, R., Peucat, J.J., Cocherie, A., Mark Fanning, C.,
 961 2012. Geochronology and geochemistry of Ordovician felsic volcanism in the
 962 Southern Armorican Massif (Variscan belt, France): Implications for the breakup of
 963 Gondwana. *Gondwana Res.* 21, 1019–1036.
- 964 Barca, S., 1991. Phénomènes de resédimentation et flysch hercynien à faciès Culm
 965 dans le “synclinal du Sarrabus” (SE de la Sardaigne, Italie). *C. R. Acad. Sci., Paris*
 966 313(2), 1051–1057.

- 967 Barca, S., Cherchi, A., 2004. Regional geological setting. In: Barca, S., Cherchi, A.
 968 (eds.), Sardinian Palaeozoic Basement and its Meso–Cainozoic Cover (Italy). 32nd
 969 Int. Geol. Congress. Field Trip Guide Book–P39 (5), 3–8.
- 970 Barca, S., Carmignani, L., Maxia, M., Oggiano, G., Pertusati, P.C., 1986. The Geology
 971 of Sarrabus. In: Guide-Book to the Excursion on the Paleozoic Basement of Sardinia
 972 (Carmignani, L., Coccozza, T., Ghezzi, C., Pertusati, P.C., Ricci, C.A., eds.). IGCP
 973 Newsl., Spec. Iss. 5, 51–60.
- 974 Barca, S., Coccozza, T., Del Rio, M., Pillola, G.L., Pittau Demelia, P., 1987. Datation de
 975 l'Ordovicien inférieur par *Dictyonema flabelliforme* et acritarches dans la partie
 976 supérieure de la formation “cambrienne” de Cabitza (SW de la Sardaigne, Italie):
 977 conséquences géodynamiques. C. R. Acad. Sci., Paris 305(2), 1109–1113.
- 978 Barca, S., Del Rio, M., Pittau Demelia, P., 1988. New geological and stratigraphical
 979 data and discovery of Lower Ordovician acritarchs in the San Vito Sandstone of the
 980 Genn'Argiolas Unit (Sarrabus, Southeastern Sardinia). Riv. It. Paleontol. Stratigr. 94,
 981 339–360.
- 982 Bea, F., Montero, P., Talavera, C., Zinger, T., 2006. A revised Ordovician age for the
 983 Miranda do Douro orthogneiss, Portugal. Zircon U–Pb ion-microprobe and LA–
 984 ICPMS dating. Geol. Acta 4, 395–401.
- 985 Bea, F., Montero, P., González Lodeiro, F., Talavera, C., 2007. Zircon inheritance
 986 reveals exceptionally fast crustal magma generation processes in Central Iberia
 987 during the Cambro–Ordovician. J. Petrol. 48, 2327–2339
- 988 Bé Mézème, E., 2005. Contribution de la géochronologie U–Th–Pb sur monazite à la
 989 compréhension de la fusion crustale dans la chaîne Hercynienne française et
 990 implication géodynamique. PhD, Univ. Orléans.
- 991 Bindeman, I.N., Valley, J.W., 2003. Rapid generation of both high- and low- $\delta^{18}\text{O}$, large-
 992 volume silicic magmas at the Timber Mountain/Oasis Valley caldera complex,
 993 Nevada. Geol. Soc. Am. Bull. 115(5), 581–595.

- 994 Boni, M., 1986. The Permo–Triassic vein and paleokarst ores in southwest Sardinia:
 995 contribution of fluid inclusion studies to their genesis and paleoenvironment. *Mineral.*
 996 *Deposita* 21, 53–62.
- 997 Boni, M., Koeppel, V., 1985. Ore-lead isotope pattern from the Iglesiente-Sulcis area
 998 (SW Sardinia) and the problem of remobilization of metals. *Mineral. Deposita* 20,
 999 185–193.
- 1000 Bryan, S.E., Riley, T.R., Jerram, D.A., Stephens, Ch.J., Leat, Ph.T., 2002. Silicic
 1001 volcanism: An undervalued component of large igneous provinces and volcanic
 1002 rifted margins. In: *Volcanic Rifted Margins* (Menzies, M.A., Klemperer, S.L., Ebinger,
 1003 C.J., Baker, J., eds.). *Geol. Soc. Am., Spec. Pap.* 362, 99–120.
- 1004 Calvet, P., Lapierre, H., Chavet, J., 1988. Diversité du volcanisme Ordovicien dans la
 1005 région de Pierrefitte (Hautes Pyrénées): rhyolites calco-alkalines et basaltes
 1006 alcalins. *C. R. Acad. Sci., Paris* 307, 805–812.
- 1007 Calvino, F., 1972. Note Illustrative della Carta Geologica d'Italia, Foglio 227 –
 1008 Muravera. Servizio Geologico d'Italia, Roma, 60 p.
- 1009 Carmignani, L., Cocozza, T., Ghezzi, C., Pertusati, P.C., Ricci, C.A., 1986. Guide-
 1010 book to the Excursion on the Palaeozoic Basement of Sardinia. IGCP Project no. 5,
 1011 *Newsl. Spec. Iss.*, 1–102.
- 1012 Carmignani, L., Pertusati, P.C., Barca, S., Carosi, R., Di Pisa, A., Gattiglio, M., et al.,
 1013 1992. Struttura della Catena Ercinica in Sardegna. Guida all'escursione del Gruppo
 1014 Informali di Geologia Strutturale in Sardegna, 24–29 (Maggio), 1–177.
- 1015 Carmignani, L., Carosi, R., Di Pisa, A., Gattiglio, M., Musumeci, G., Oggiano, G. et al.,
 1016 1994. The Hercynian chain in Sardinia (Italy). *Geodin. Acta* 7, 31–47.
- 1017 Carmignani, L., Oggiano, G., Barca, S., Conti, P., Salvadori, I., Eltrudis, A. et al. 2001.
 1018 Geologia della Sardegna. Note illustrative della Carta Geologica della Sardegna a
 1019 scala 1:200.000. *Memorie Descrittive della Carta Geologica d'Italia, Servizio*
 1020 *Geologico* 60, 1–283. Istituto Poligrafico e Zecca dello Stato, Roma.

- 1021 Caron, C., Lancelot, J., Omenetto, P., Orgeval, J.J., 1997. Role of the Sardic tectonic
1022 phase in the metallogenesis of SW Sardinia (Iglesiente): lead isotope evidence.
1023 *European J. Miner.* 9, 1005–1016.
- 1024 Casas, J.M., 2010. Ordovician deformations in the Pyrenees: new insights into the
1025 significance of pre–Variscan ('sardic') tectonics. *Geol. Mag.* 147, 674–689.
- 1026 Casas, J.M., Fernández, O., 2007. On the Upper Ordovician unconformity in the
1027 Pyrenees: New evidence from the La Cerdanya area. *Geol. Acta* 5, 193–198.
- 1028 Casas, J.M., Murphy, J.B. 2018. Unfolding the arc: the use of pre-orogenic constraints
1029 to assess the evolution of the Variscan belt in Western Europe. *Tectonophysics* 736,
1030 47–61.
- 1031 Casas, J.M., Palacios, T., 2012. First biostratigraphical constraints on the pre–Upper
1032 Ordovician sequences of the Pyrenees based on organic-walled microfossils. *C. R.*
1033 *Geosci.* 344, 50–56.
- 1034 Casas, J.M., Castiñeiras, P., Navidad, M., Liesa, M., Carreras, J., 2010. New insights
1035 into the Late Ordovician magmatism in the Eastern Pyrenees: U–Pb SHRIMP zircon
1036 data from the Canigó massif. *Gondwana Res.* 17, 317–324.
- 1037 Casas, J.M., Álvaro, J.J., Clausen, S., Padel, M., Puddu, C., Sanz-López, J., Sánchez-
1038 García, T., Navidad, M., Castiñeiras, P., Liesa, M., 2019. Palaeozoic basement of
1039 the Pyrenees. In: *The Geology of Iberia: A Geodynamic Approach* (Quesada, C.,
1040 Oliveira, J.T., eds.). *Regional Geology Reviews*, vol. 2, 229–259. Springer,
1041 Heidelberg.
- 1042 Castiñeiras, P., Villaseca, C., Barbero, L., Martín-Romera, C., 2008a. SHRIMP U–Pb
1043 zircon dating of anatexis in high-grade migmatite complexes of Central Spain:
1044 implications in the Hercynian evolution of Central Iberia. *Int. J. Earth Sci.* 97, 35–50.
- 1045 Castiñeiras, P., Navidad, M., Liesa, M., Carreras, J., Casas, J.M., 2008b. U–Pb zircon
1046 ages (SHRIMP) for Cadomian and Lower Ordovician magmatism in the Eastern
1047 Pyrenees: new insights in the pre–Variscan evolution of the northern Gondwana
1048 margin. *Tectonophysics* 46, 228–239.

- 1049 Castro, A., García-Casco, A., Fernández, C., Corretgé, L.G., Moreno-Ventas, I., Gerya,
1050 T., Löw, I., 2009. Ordovician ferrosilicic magmas: experimental evidence for
1051 ultrahigh temperatures affecting a metagreywacke source. *Gondwana Res.* 16, 622–
1052 632
- 1053 Charles, N., Faure, M., Chen, Y., 2008. The emplacement of the Montagne Noire axial
1054 zone (French Massif Central): New insights from petro-textural, geochronological
1055 and AMS studies. *22ème Réunion des Sciences de la Terre, Nancy*, 155.
- 1056 Charles, N., Faure, M., Chen, Y., 2009. The Montagne Noire migmatitic dome
1057 emplacement (French Massif Central): New insights from petrofabric and AMS
1058 studies. *J. Struct. Geol.* 31, 1423–1440.
- 1059 Clariana, P., Valverde-Vaquero, P., Rubio-Ordóñez, A., Beranoaguirre, A., García-
1060 Sansegundo, J., 2018. Pre–Variscan tectonic events and Late Ordovician
1061 magmatism in the Central Pyrenees: U–Pb age and Hf in zircon isotopic signature
1062 from subvolcanic sills in the Pallaresa massif. *J. Iberian Geol.* 44, 589–601.
- 1063 Cocco, F., Funedda, A., 2011. New data on the pre–Middle Ordovician deformation in
1064 SE Sardinia: a preliminary note. *Rend. online Soc. Geol. It.* 15, 34–36.
- 1065 Cocco, F., Funedda, A., 2017. The Sardinic Phase: field evidence of Ordovician tectonics
1066 in SE Sardinia. *Italy. Geol. Mag.* 156, 25–38.
- 1067 Cocco, F., Oggiano, G., Funedda, A., Loi, A., Casini, L., 2018. Stratigraphic, magmatic
1068 and structural features of Ordovician tectonics in Sardinia (Italy): a review. *J. Iberian*
1069 *Geol.* 44, 619–639.
- 1070 Cocherie, A., 2003. Datation avec le SHRIMP II du métagranite oeilé du Somail-
1071 Montagne Noire. *C. R. technique ANA–ISO/NT, BRGM.*
- 1072 Cocherie, A., Baudin, T., Guerrot, C., Autran, A., Fanning, M.C., Laumonier, B., 2005.
1073 U–Pb zircon (ID–TIMS and SHRIMP) evidence for the early Ordovician intrusion of
1074 metagranites in the late Proterozoic Canaveilles Group of the Pyrenees and the
1075 Montagne Noire (France). *Bull. Soc. géol. France* 176, 269–282.

- 1076 Cortesogno, L., Gaggero, L., Oggiano, G., Paquette, J.L., 2004. Different tectono-
1077 thermal evolution paths in eclogitic rocks from the Axial Zone of the Variscan Chain
1078 in Sardinia (Italy) compared with the Ligurian Alps. *Ophioliti* 29, 125–144.
- 1079 Costamagna, L.G., Elter, F.M., Gaggero, L., Mantovani, F., 2016. Contact
1080 metamorphism in Middle Ordovician arc rocks (SW Sardinia, Italy): New
1081 paleogeographic constraints. *Lithos* 264, 577–593.
- 1082 Cruciani, G., Franceschelli, M., Musumeci, G., Spano, M.E., Tiepolo, M., 2013. U–Pb
1083 zircon dating and nature of metavolcanics and metarkoses from the Monte Grighini
1084 Unit: new insights on Late Ordovician magmatism in the Variscan belt in Sardinia,
1085 Italy. *Int. J. Earth Sci.* 102, 2077–2096.
- 1086 Cruciani, G., Franceschelli, M., Puxeddu, M., Tiepolo, M., 2018. Metavolcanics from
1087 Capo Malfatano, SW Sardinia, Italy: New insight on the age and nature of
1088 Ordovician volcanism in the Variscan foreland zone. *Geol. J.* 53(4), 1573–1585.
- 1089 Demange, M., 1999. Evolution tectonique de la Montagne noire: un modèle en
1090 transpression. *C. R. Acad. Sci., Paris* 329, 823–829.
- 1091 Demange, M., Guérangé-Lozes, J., Guérangé, B., 1996. Notice explicative de la feuille
1092 de Lacauene (987) au 1:50 000. BRGM, Orléans.
- 1093 Denèle, Y., Barbey, P., Deloule, E., Pelleter, E., Olivier, Ph., Gleizes, G., 2009. Middle
1094 Ordovician U–Pb age of the Aston and Hospitalet orthogneissic laccoliths: their role
1095 in the Variscan evolution of the Pyrenees. *Bull. Soc. géol. France* 180, 209–221.
- 1096 DePaolo, D.J., 1981. Neodymium isotopes in the Colorado Front Range and crust-
1097 mantle evolution in the Proterozoic. *Nature* 291, 193–196.
- 1098 DePaolo, D.J., 1988. Neodymium isotope geochemistry. An introduction. *Minerals and*
1099 *Rocks Series* 20, 1–187. Springer-Verlag, Berlin.
- 1100 DePaolo, D.J., Wasserburg, G.J., 1976. Nd isotopic variations and petrogenetic
1101 models. *Geophys. Res. Lett.* 3(5), 249–252.

- 1102 Dias da Silva Í., 2014. Geología de las Zonas Centro Ibérica y Galicia-Tras-os-Montes
1103 en la parte oriental del Complejo de Morais, Portugal/España. Serie Nova Terra 45,
1104 1–424.
- 1105 Dias da Silva, I., Valverde-Vaquero, P., González-Clavijo, E., Díez-Montes, A.,
1106 Martínez-Catalán, J.R., 2012. Structural and stratigraphical significance of U–Pb
1107 ages from the Saldanha and Mora volcanic complexes (NE Portugal, Iberian
1108 Variscides). Géol. France 1, 105–106.
- 1109 Dias da Silva Í., Valverde-Vaquero, P., González Clavijo E., Díez-Montes A., Martínez
1110 Catalán J.R., 2014. Structural and stratigraphical significance of U–Pb ages from the
1111 Mora and Saldanha volcanic complexes (NE Portugal, Iberian Variscides). In:
1112 Schulmann, K., Martínez Catalán, J. R., Lardeaux, J. M., Janousek, V., Oggiano, G.,
1113 (eds.), The Variscan Orogeny: Extent, Timescale and the Formation of the
1114 European Crust. Geol. Soc., London, Spec. Publ. 405, 115–135.
- 1115 Dias da Silva, I., Díez Fernández, R., Díez Montes, A., González Clavijo, E., Foster,
1116 D.A., 2016. Magmatic evolution in the N-Gondwana margin related to the opening of
1117 the Rheic Ocean – evidence from the Upper Parautochthon of the Galicia-Trás-os-
1118 Montes Zone and from the Central Iberian Zone (NW Iberian Massif). Int. J. Earth
1119 Sci. 105, 1127–1151.
- 1120 Díaz-Alvarado, J., Fernández, C., Chichorro, M., Castro, A., Pereira, M.F., 2016.
1121 Tracing the Cambro–Ordovician ferrosilicic to calc-alkaline magmatic association in
1122 Iberia by *in situ* U–Pb SHRIMP zircon geochronology (Gredos massif, Spanish
1123 Central System batholith). Tectonophysics 681, 95–110.
- 1124 Díez Fernández, R., Castiñeiras, P., Gómez Barreiro, J., 2012. Age constraints on
1125 Lower Paleozoic convection system: Magmatic events in the NW Iberian Gondwana
1126 margin. Gondwana Res. 21, 1066–1079.
- 1127 Díez-Montes, A., 2007. La Geología del Dominio “Ollo de Sapo” en las Comarcas de
1128 Sanabria y Terra do Bolo. PhD, Univ. Salamanca. Laboratorio Xelóxico de Laxe,
1129 Serie Nova Terra no. 34, A Coruña.

- 1130 Díez Montes, A., Martínez Catalán, J.R., Bellido Mulas, F., 2010. Role of the Ollo de
1131 Sapo massive felsic volcanism of NW Iberia in the Early Ordovician dynamics of
1132 northern Gondwana. *Gondwana Res.* 17, 363–376.
- 1133 Di Pisa, A., Gattiglio, M., Oggiano, G., 1992. Pre–Hercynian magmatic activity in the
1134 nappe zone (internal and external) of Sardinia: evidence of two within plate basaltic
1135 cycles. In: *Contributions to the Geology of Italy with Special Regard to the Paleozoic*
1136 *Basements* (Carmingnani, L., Sassi, F.P., eds.). *Newsl. IGCP* 276, 107–116.
- 1137 Echtler, H., Malavieille, J., 1990. Extensional tectonics, basement uplift and Stephano–
1138 Permian collapse basin in a late Variscan metamorphic core complex (Montagne
1139 Noire, southern Massif Central). *Tectonophysics* 177, 125–138.
- 1140 El Korh, A., Schmidt, S.Th., Ballèvre, M., Ulianov, A., Bruguier, O., 2012. Discovery of
1141 an albite gneiss from the Ile de Groix (Armorican Massif, France): geochemistry and
1142 LA–ICP–MS U–Pb geochronology of its Ordovician protolith. *Int. J. Earth Sci.* 101,
1143 1169–1190.
- 1144 Engel, W., Feist, R., Franke, W., 1980. Le Carbonifère anté–stéphanien de la Montagne
1145 Noire: rapports entre mise en place des nappes et sédimentation. *Bull. BRGM* 2,
1146 341–389.
- 1147 Farias, P., Ordoñez-Casado, B., Marcos, A., Rubio-Ordóñez, A., Fanning, C.M., 2014.
1148 U–Pb zircon SHRIMP evidence for Cambrian volcanism in the Schistose Domain
1149 within the Galicia-Trás-os-Montes Zone (Variscan Orogen, NW Iberian Peninsula).
1150 *Geol. Acta* 12(3), 209–218.
- 1151 Faure, M., Ledru, P., Lardeaux, J.M., Matte, P., 2004. Paleozoic orogenies in the
1152 French Massif Central. A cross section from Béziers to Lyon. 32nd Int. Geol.
1153 Congress Florence (Italy), Field-trip guide book, 40 p.
- 1154 Feist, R., Galtier, J., 1985. Découverte de flores d'âge namurien probable dans le
1155 flysch à olistolithes de Cabrières (Hérault). Implications sur la durée de la
1156 sédimentation synorogénique dans la Montagne Noire (France Méridionale). *C. R.*
1157 *Acad. Sci., Paris* 300, 207–212.

- 1158 Franz, L., Romer, R.L., 2007. Caledonian high-pressure metamorphism in the Strona-
1159 Ceneri-Zone (Southern Alps of southern Switzerland and northern Italy). *Swiss J.*
1160 *Geosci.* 100, 457–467.
- 1161 Friedl, G., Finger, F., Paquette, J.L., von Quadt, A., McNaughton, N.J., Fletcher, I.R.,
1162 2004. Pre-Variscan geological events in the Austrian part of the Bohemian Massif
1163 deduced from U–Pb zircon ages. *Int. J. Earth Sci.* 93, 802–823
- 1164 Funedda, A., Oggiano, G., 2009. Outline of the Variscan basement of Sardinia. In: *The*
1165 *Silurian of Sardinia. Volume in Honour of Enrico Serpagli (Corradini, C., Ferretti, A.,*
1166 *Štorch, P., eds.). Rend. Soc. Paleontol. It.* 3, 23–35.
- 1167 Gaggero L., Oggiano G., Funedda A., Buzzi L., 2012. Rifting and arc-related Early
1168 Paleozoic volcanism along the North Gondwana margin: Geochemical and
1169 geological evidence from Sardinia (Italy). *J. Geol.* 120, 273–292.
- 1170 García-Arias, M., Díez-Montes, A., Villaseca, C., Blanco-Quintero, I.F. 2018. The
1171 Cambro–Ordovician Ollo de Sapo magmatism in the Iberian Massif and its Variscan
1172 evolution: A review. *Earth-Sci. Rev.* 176, 345–372.
- 1173 Gèze, B., 1949. Etude géologique de la Montagne Noire et des Cévennes
1174 méridionales. *Mém. Soc. géol. France* 62, 1–215.
- 1175 Giacomini, F, Bomparola, R.M., Ghezzo, C., 2005. Petrology and geochronology of
1176 metabasites with eclogite facies relics from NE Sardinia: constraints for the
1177 Palaeozoic evolution of Southern Europe. *Lithos* 82, 221–248
- 1178 Giacomini, F., Bomparola, R.M., Ghezzo, C., Guldbransen, H., 2006. The geodynamic
1179 evolution of the Southern European Variscides: constraints from the U/Pb
1180 geochronology and geochemistry of the lower Palaeozoic magmatic-sedimentary
1181 sequences of Sardinia (Italy). *Contrib. Miner. Petr.* 152, 19–42.
- 1182 Guérangé-Lozes, J., Alabouvette, B., 1999. Notice explicative, Carte géol. France (1/50
1183 000), feuille Saint-Sernin-sur-Rance (960). BRGM, Orléans, 84 p.

- 1184 Guérangé-Lozes, J., Guérangé, B., Mouline, M.P., Delsahut, B., 1996. Notice
1185 explicative, Carte géol. France (1/50 000), feuille Réalmont (959). BRGM, Orléans,
1186 78 p.
- 1187 Guitard, G., 1970. Le métamorphisme hercynien mésozonal et les gneiss ocellés du
1188 massif du Canigou (Pyrénées orientales). Mém. BRGM 63, 1–353.
- 1189 Gutiérrez-Alonso, G., Fernández-Suárez, J., Gutiérrez-Marco, J.C., Corfu, F., Murphy,
1190 J.B., Suárez Martínez, S., 2007. U–Pb depositional age for the upper Barrios
1191 Formation (Armorican Quartzite facies) in the Cantabrian zone of Iberia: implications
1192 for stratigraphic correlation and paleogeography. In: The Evolution of the Rheic
1193 Ocean: from Avalonian–Cadomian Active Margin to Alleghenian–Variscan Collision
1194 (Linnemann, R.D., Nance, P., Kraft, G.Z., eds.). Geol. Soc. Am., London, 287–296.
- 1195 Gutiérrez-Alonso, G., Gutiérrez-Marco, J.C., Fernández-Suárez, J., Bernárdez, E.,
1196 Corfu, F., 2016. Was there a super-eruption on the Gondwanan coast 477 Ma ago?
1197 Tectonophysics 681, 85–94.
- 1198 Gutiérrez-Marco, J.C., Robardet, M., Rábano, I., Sarmiento, G., San José Lancha,
1199 M.A., Herranz Araújo, P., Pieren Vidal, A., 2002. Ordovician. In: The Geology of
1200 Spain, (Gibbons, W., Moreno, T., eds.), 31–49, Geological Society, London
- 1201 Gutiérrez-Marco, J.C., Piçarra, J.M., Meireles, C.A., Cózar, P., García-Bellido, D.C.,
1202 Pereira, Z. et al., 2019. Early Ordovician–Devonian Passive margin stage in the
1203 Gondwanan units of the Iberian massif. In: The Geology of Iberia: A Geodynamic
1204 Approach (Quesada, C., Oliveira, J.T., eds.). Regional Geology Reviews, vol. 2, 75–
1205 98. Springer, Heidelberg.
- 1206 Hartevelt, J.J.A., 1970. Geology of the upper Segre and Valira valleys, central
1207 Pyrenees, Andorra/Spain. Leid. Geol. Meded. 45, 167–236.
- 1208 Helbing, H., Tiepolo, M., 2005. Age determination of Ordovician magmatism in NE
1209 Sardinia and its bearing on Variscan basement evolution. J. Geol. Soc. 162, 689–
1210 700.

- 1211 Huppert, H.E., Sparks, R.S.J., 1988. The generation of granitic magmas by intrusion of
1212 basalt into continental crust. *J. Petrol.* 29, 599–624.
- 1213 Ibrahim, M.E., El-Kalioby, B.A., Aly, G.M., El-Tohamy, A.M., Watanabe, K., 2015.
1214 Altered granitic rocks, Nusab El Balgum Area, Southwestern Desert, Egypt.
1215 Mineralogical and geochemical aspects of REEs. *Ore Geol. Rev.* 70, 252–261.
- 1216 Irber, W., 1999. The lanthanide tetrad effect and its correlation with K/Rb, Eu/Eu*,
1217 Sr/Eu, Y/Ho, and Zr/Hf of evolving peraluminous granite suites. *Geochim.*
1218 *Cosmochim. Acta* 63, 489–508.
- 1219 Jégouzo, P., Peucat, J.J., Audren, C., 1986. Caractérisation et signification
1220 géodynamique des orthogneiss calco-alcalins d'âge ordovicien de Bretagne
1221 méridionale. *Bull. Soc. géol. France* 2, 839–848.
- 1222 Kröner, A., Willner, A.P., 1998. Time of formation and peak of Variscan HP–HT
1223 metamorphism of quartz-feldspar rocks in the central Erzgebirge, Saxony, Germany.
1224 *Contrib. Mineral. Petrol.* 132, 1–20
- 1225 Lancelot, J., Allegret, A., Iglesias Ponce de León, M., 1985. Outline of Upper
1226 Precambrian and Lower Paleozoic evolution of the Iberian Peninsula according to
1227 U–Pb dating of zircons. *Earth Planet. Sci. Lett.* 74, 325–337.
- 1228 Laske, R., Bechstädt, T., Boni, M., 1994. The post–Sardic Ordovician series. In:
1229 Sedimentological, stratigraphical and ore deposits field guide of the autochthonous
1230 Cambro–Ordovician of Southeastern Sardinia (Bechstädt, T., Boni, M., eds.).
1231 *Memorie descrittive della carta geologica d'Italia* 48, 115–146.
- 1232 Leat, P.T., Jackson, S.E., Thorpe, R.S., Stillman, C.J., 1986. Geochemistry of bimodal
1233 basalt-subalkaline/peralkaline rhyolite provinces within the Southern British
1234 Caledonides. *J. Geol. Soc.* 143, 259–273.
- 1235 Le Corre, C., Auvray, B., Ballèvre, M., Robardet, M., 1991. Le Massif Armoricaïn. *Sci.*
1236 *Géol., Bull.* 44, 31–103.
- 1237 Lentz, D., 1996. U, Mo and REE mineralization in late-tectonic granite pegmatites,
1238 south-west Grenville Province, Canada. *Ore Geol. Rev.* 11, 197–227.

- 1239 Leone, F., Hamman, W., Laske, R., Serpagli, E., Villas, E., 1991. Lithostratigraphic
1240 units and biostratigraphy of the post–Sardic Ordovician sequence in south-west
1241 Sardinia. *Boll. Soc. Paleontol. It.* 30, 201–235.
- 1242 Leone, F., Ferretti, A., Hammann, W., Loi, A., Pillola, G.L., Serpagli, E., 2002. A
1243 general view of the post–Sardic Ordovician sequence from SW Sardinia. *Rend. Soc.*
1244 *Paleontol. It.* 1, 51–68.
- 1245 Lescuyer, J.L., Cocherie, A., 1992. Datation sur monozircons des métadacites de
1246 Sériès: arguments pour un âge protérozoïque terminal des “schistes X” de la
1247 Montagne Noire (Massif central français). *C. R. Acad. Sci., Paris (sér. 2)* 314, 1071–
1248 1077.
- 1249 Liesa, M., Carreras, J., Castiñeiras, P., Casas, J.M., Navidad, M., Vilà, M., 2011. U–Pb
1250 zircon age of Ordovician magmatism in the Albera Massif (Eastern Pyrenees). *Geol.*
1251 *Acta* 9, 1–9.
- 1252 Linnemann, U., Gehmlich, M., Tichomirowa, M., Buschmann, B., Nasdala, L., Jonas, P.
1253 et al. 2000. From Cadomian subduction to early Palaeozoic rifting: the evolution of
1254 Saxo-Thuringia at the margin of Gondwana in the light of single zircon
1255 geochronology and basin development (Central European Variscides, Germany). In:
1256 *Orogenic Processes: Quantification and Modelling in the Variscan Belt* (Franke, W.,
1257 Haak, V., Oncken, O., Tanner, D., eds.). *Geol. Soc., London, Spec. Publ.* 179, 131–
1258 153.
- 1259 Loi, A., Dabard, M.P., 1997. Zircon typology and geochemistry in the palaeogeographic
1260 reconstruction of the Late Ordovician of Sardinia (Italy). *Sediment. Geol.* 112, 263–
1261 279.
- 1262 Loi, A., Barca, S., Chauvel, J.J., Dabard, M.P., Leone, F., 1992. Analyse de la
1263 sédimentation post-phase sarde les dépôts initiaux à placers du SE de la Sardaigne.
1264 *C. R. Soc. géol. France (sér. 2)* 315, 1357–1364.
- 1265 López-Sánchez, M.A., Iriondo, A., Marcos, A., Martínez, F.J., 2015. A U–Pb zircon age
1266 (479 ± 5 Ma) from the uppermost layers of the Ollo de Sapo Formation near Viveiro

- 1267 (NW Spain): implications for the duration of rifting-related Cambro–Ordovician
1268 volcanism in Iberia. *Geol. Mag.* 152, 341–350.
- 1269 Ludwig, K.R., Turi, B., 1989. Paleozoic age of the Capo Spartivento Orthogneiss,
1270 Sardinia, Italy. *Chem. Geol.* 79, 147–153.
- 1271 Margalef, A., Castiñeiras, P., Casas, J.M., Navidad, M., Liesa, M., Linnemann, U.,
1272 Hofmann, M., Gärtner, A., 2016. Detrital zircons from the Ordovician rocks of the
1273 Pyrenees: Geochronological constraints and provenance. *Tectonophysics* 681, 124–
1274 134.
- 1275 Marini, F., 1988. “Phase” sarde et distension ordovicienne du domaine sud-varisque,
1276 effets de point chaud? Une hypothèse fondée sur les données nouvelles du
1277 volcanisme albigeois. *C. R. Acad. Sci., Paris (sér. 2)* 306, 443–450.
- 1278 Martí, J., Muñoz, J.A., Vaquer, R., 1986. Les roches volcaniques de l’Ordovicien
1279 supérieur de la région de Ribes de Freser-Rocabruna (Pyrénées catalanes):
1280 caractères et signification. *C. R. Acad. Sci., Paris* 302, 1237–1242.
- 1281 Martí, J., Solari, L., Casas, J.M., Chichorro, M., 2019. New late Middle to early Late
1282 Ordovician U–Pb zircon ages of extension-related felsic volcanic rocks in the
1283 Eastern Pyrenees (NE Iberia): tectonic implications. *Geol. Mag.* 156(10), 1783–
1284 1792.
- 1285 Martínez, F., Iriondo, A., Dietsch, C., Aleinikoff, J.N., Peucat, J.J., Cirès, J., Reche, J.,
1286 Capdevila, R., 2011. U–Pb SHRIMP–RG zircon ages and Nd signature of lower
1287 Paleozoic rifting-related magmatism in the Variscan basement of the Eastern
1288 Pyrenees. *Lithos* 127, 10–23.
- 1289 Martínez Catalán, J.R., Arenas, R., Díaz García, F., Gómez Barreiro, J., González
1290 Cuadra, P., Abati, J. et al. 2007. Space and time in the tectonic evolution of the
1291 northwestern Iberian Massif. Implications for the comprehension of the Variscan
1292 belt. In: 4–D Framework of Continental Crust (Hatcher, R.D.Jr., Carlson, M.P.,
1293 McBride, J.H., Martínez Catalán, J.R., eds.). *Geol. Soc. Am., Mem.* 200, 403–423.

- 1294 Martini, I.P., Tongiorgi, M., Oggiano, G., Coccozza, T., 1991. Ordovician alluvial fan to
1295 marine shelf transition in SW Sardinia, Western Mediterranean Sea: tectonically
1296 (“Sardic” phase”) influenced clastic sedimentation. *Sediment. Geol.* 72, 97–115.
- 1297 Masuda, A., Akagi, T., 1989. Lanthanide tetrad effect observed in leucogranites from
1298 China. *Geochem. J.* 23, 245–253.
- 1299 McDougall, N., Brenchley, P.J., Rebelo, J.A., Romano, M., 1987. Fans and fan deltas –
1300 precursors to the Armorican Quartzite (Ordovician) in western Iberia. *Geol. Mag.*
1301 124, 347–359.
- 1302 McLennan, S.M., 1994. Rare earth element geochemistry and the “tetrad” effect.
1303 *Geochim. Cosmochim. Acta* 58, 2025–2033.
- 1304 Mezger, J., Gerdes, A., 2016. Early Variscan (Visean) granites in the core of central
1305 Pyrenean gneiss domes: implications from laser ablation U–Pb and Th–Pb studies.
1306 *Gondwana Res.* 29, 181–198.
- 1307 Mingram, B., Kröner, A., Hegner, E., Krentz, O., 2004. Zircon ages, geochemistry, and
1308 Nd isotopic systematics of pre–Variscan orthogneisses from the Erzgebirge, Saxony
1309 (Germany), and geodynamic interpretation. *Int. J. Earth Sci.* 93, 706–727.
- 1310 Monecke, T., Dulski, P., Kempe, U., 2007. Origin of convex tetrads in rare earth
1311 element patterns of hydrothermally altered siliceous igneous rocks from the
1312 Zinnwald Sn–W deposit, Germany. *Geochim. Cosmochim. Acta* 71, 335–353.
- 1313 Montero, P., Bea, F., González-Lodeiro, F., Talavera, C., Whitehouse, M.J., 2007.
1314 Zircon ages of the metavolcanic rocks and metagranites of the Ollo de Sapo Domain
1315 in central Spain: implications for the Neoproterozoic to Early Palaeozoic evolution of
1316 Iberia. *Geol. Mag.* 144, 963–976.
- 1317 Montero, P., Talavera, C., Bea, F., Lodeiro, F.G., Whitehouse, M.J., 2009. Zircon
1318 geochronology of the Ollo de Sapo Formation and the age of the Cambro–
1319 Ordovician rifting in Iberia. *J. Geol.* 117, 174–191.

- 1320 Murphy, J.B., Gutiérrez-Alonso, G., Nance, R.D., Fernández-Suárez, J., Keppie, J.D.,
 1321 Quesada, C. et al., 2006. Origin of the Rheic Ocean: rifting along a Neoproterozoic
 1322 suture? *Geology* 34, 325–328.
- 1323 Nance, R.D., Gutiérrez-Alonso, G., Keppie, J.D., Linnemann, U., Murphy, J.B.,
 1324 Quesada, C. et al., 2010. Evolution of the Rheic Ocean. *Gondwana Res.* 17, 194–
 1325 222.
- 1326 Navidad, M., Castiñeiras, P., 2011. Early Ordovician magmatism in the northern
 1327 Central Iberian Zone (Iberian Massif): new U–Pb (SHRIMP) ages and isotopic Sr–
 1328 Nd data. 11th ISOS, Alcalá de Henares, May 2011.
- 1329 Navidad, M., Castiñeiras, P., Casas, J.M., Liesa, M., Fernández-Suárez, J., Barnolas,
 1330 A., Carreras, J., Gil-Peña, I., 2010. Geochemical characterization and isotopic ages
 1331 of Caradocian magmatism in the northeastern Iberia: insights into the Late
 1332 Ordovician evolution of the northern Gondwana margin. *Gondwana Res.* 17, 325–
 1333 337.
- 1334 Navidad, M., Castiñeiras, P., Casas, J.M., Liesa, M., Belousova, E., Proenza, J.,
 1335 Aiglsperger, T., 2018. Ordovician magmatism in the Eastern Pyrenees: Implications
 1336 for the geodynamic evolution of northern Gondwana. *Lithos* 314–315, 479–496.
- 1337 Neiva, A.M.R., Williams, I.S., Ramos, J.M.F., Gomes, M.E.P., Silva, M.M.V.G.,
 1338 Antunes, I.M.H.R., 2009. Geochemical and isotopic constraints on the petrogenesis
 1339 of Early Ordovician granodiorite and Variscan two-mica granites from the Gouveia
 1340 area, central Portugal. *Lithos* 111, 186–202.
- 1341 Oggiano, G., Gaggero, L., Funedda, A., Buzzi, L., Tiepolo, M., 2010. Multiple early
 1342 Paleozoic volcanic events at the northern Gondwana margin: U–Pb age evidence
 1343 from the southern Variscan branch (Sardinia, Italy). *Gondwana Res.* 17, 44–58.
- 1344 Padel, M., Álvaro, J.J., Clausen, S., Guillot, F., Poujol, M., Chichorro, M., Monceret, E.,
 1345 Pereira, M.F., Vizcaíno, D., 2017. U–Pb laser ablation ICP–MS zircon dating across
 1346 the Ediacaran–Cambrian transition of the Montagne Noire, southern France. *C. R.*
 1347 *Geosci.* 349, 380–390.

- 1348 Padel, M., Clausen, S., Álvaro, J.J., Casas, J.M., 2018. Review of the Ediacaran–
1349 Lower Ordovician (pre-Sardic) stratigraphic framework of the Eastern Pyrenees,
1350 southwestern Europe. *Geol. Acta* 16, 339–355
- 1351 Palme, H., O'Neill, H.S.C., 2004. Cosmochemical estimates of mantle composition. In:
1352 Treatise on Geochemistry 2 (Holland, H.D., Turekian, K.K., eds.), 1–38. Elsevier-
1353 Pergamon, Oxford.
- 1354 Palmeri, R., Fanning, M., Franceschelli, M., Memmi, I., Ricci, C.A., 2004. SHRIMP
1355 dating of zircons in eclogite from the Variscan basement in northeastern Sardinia
1356 (Italy). *N. Jb. Miner., Mh.* 6, 275–288.
- 1357 Pan, Y., 1997. Controls on the fractionation of isovalent trace elements in magmatic
1358 and aqueous systems: evidence from Y/Ho, Zr/Hf, and lanthanide tetrad effect – a
1359 discussion of the article by M. Bau, 1996. *Contrib. Mineral. Petrol.* 128, 405–408.
- 1360 Pankhurst, R.J., Rapela, C.W., Saavedra, J., Baldo, E., Dahlquist, J., Pascua, I.,
1361 Fanning, C.M., 1998. The Famatinian magmatic arc in the central Sierras
1362 Pampeanas, and Early to Middle Ordovician continental arc on the Gondwana
1363 margin. In: *The Proto-Andean Margin of Gondwana* (Pankhurst, R.J., Rapela, C.E.,
1364 eds.). *Geol. Soc., London, Spec. Publ.* 142, 343–367.
- 1365 Pavanetto, P., Funedda, A., Northrup, C. J., Schmitz, M., Crowley, J., Loi, A., 2012.
1366 Structure and U–Pb zircon geochronology in the Variscan foreland of SW Sardinia,
1367 Italy. *Geol. J.* 47, 426–445.
- 1368 Pearce, J.A., 1996. Sources and settings of granitic rocks. *Episodes* 19, 120–125.
- 1369 Pearce, J.A., Harris, N.B.W., Tindle, A.G., 1984. Trace element discrimination
1370 diagrams for the tectonic interpretation of granitic rocks. *J. Petrol.* 25, 956–983.
- 1371 Pereira, M.F., Solá, A.R., Chichorro, M., Lopes, L., Gerdes, A., Silva, J.B., 2012. North-
1372 Gondwana assembly, break-up and paleogeography: U–Pb isotope evidence from
1373 detrital and igneous zircons of Ediacaran and Cambrian rocks of SW Iberia.
1374 *Gondwana Res.* 22(3–4), 866–881.

- 1375 Piercey, S.J., 2011. The setting, style, and role of magmatism in the formation of
1376 volcanogenic massive sulphide deposits. *Miner. Deposita* 46, 449–471.
- 1377 Pistis, M., Loi, A., Dabard, M.P., 2016. Influence of relative sea-level variations on the
1378 genesis of palaeoplacers, the examples of Sarrabus (Sardinia, Italy) and the
1379 Armorican Massif (western France). *C. R. Geosci.* 348(2), 150–157.
- 1380 Pillola, G.L., Leone, F., Loi, A., 1998. The Cambrian and Early Ordovician of SW
1381 Sardinia. *Gior. Geol. (ser. 3), Spec. Iss.* 60, 25–38.
- 1382 Pitra, P., Poujol, M., Den Driessche, J.V., Poilvet, J. C., Paquette, J.L., 2012. Early
1383 Permian extensional shearing of an Ordovician granite: the Saint-Eutrope “C/S-like”
1384 orthogneiss (Montagne Noire, French Massif Central). *C. R. Geosci.* 344, 377–384.
- 1385 Pouclet, A., Álvaro, J.J., Bardintzeff, J.M., Gil Imaz, A., Monceret, E., Vizcaíno D.,
1386 2017. Cambrian–Early Ordovician volcanism across the South Armorican and
1387 Occitan Domains of the Variscan Belt in France: Continental break-up and rifting of
1388 the northern Gondwana margin. *Geosci. Frontiers* 8, 25–64.
- 1389 Puddu, C., Álvaro J.J., Casas, J.M., 2018. The Sardic unconformity and the Upper
1390 Ordovician successions of the Ribes de Freser area, Eastern Pyrenees. *J. Iberian*
1391 *Geol.* 44, 603–617.
- 1392 Puddu, C., Álvaro, J.J., Carrera, N., Casas, J.M., 2019. Deciphering the Sardic
1393 (Ordovician) and Variscan deformations in the Eastern Pyrenees. *J. Geol. Soc.*
1394 176(6), 1191–1206.
- 1395 Quesada, C., 1991. Geological constraints on the Paleozoic tectonic evolution of
1396 tectonostratigraphic terranes in the Iberian Massif. *Tectonophysics* 185, 225–245.
- 1397 Rabin, M., Trap, P., Carry, N. Fréville, K. Cenki-Tok, B. Lobjoie, C. Gonçalves, P.,
1398 Marquer, D., 2015. Strain partitioning along the anatectic front in the Variscan
1399 Montagne Noire massif (southern French Massif Central). *Tectonics* 34, 1709–1735.
- 1400 Robert, J. F. 1980. Étude géologique et métallogénique du val de Ribas sur le versant
1401 espagnol des Pyrénées catalanes. PhD, Univ. Franche-Comté.

- 1402 Robert, J.F., Thiebaut, J., 1976. Découverte d'un volcanisme acide dans le Caradoc de
1403 la région de Ribes de Freser (Prov. de Gérone). C. R. Acad. Sci., Paris 282, 2050–
1404 2079.
- 1405 Roger, F., Respaut, J.P., Brunel, M., Matte, Ph., Paquette, J.L., 2004. Première
1406 datation U–Pb des orthogneiss ocellés de la zone axiale de la Montagne Noire (Sud
1407 du Massif central): nouveaux témoins du magmatisme ordovicien dans la chaîne
1408 varisque. C. R. Geosci. 336, 19–28
- 1409 Roger, F., Teyssier, C., Respaut, J.P., Rey, P.F., Jolivet, M., Whitney, D.L., Paquette,
1410 J.L., Brunel, M., 2015. Timing of formation and exhumation of the Montagne Noire
1411 double dome, French massif Central. Tectonophysics 640–641, 53–69.
- 1412 Rollison, H.R., 1993. Using Geochemical Data: Evaluation, Presentation, Interpretation.
1413 Longman Group, London, 352 p.
- 1414 Romão, J., Dunning, G., Marcos, A., Dias, R., Ribeiro, A., 2010. O lacólito granítico de
1415 Mação-Penhascoso: idade e as suas implicações (SW da Zona Centro-Ibérica). e-
1416 Terra 16, 1–4.
- 1417 Rossi, P., Oggiano, G., Cocherie, A., 2009. A restored section of the “southern
1418 Variscan realm” across the Corsica-Sardinia microcontinent. C. R. Geosci. 34, 224–
1419 238.
- 1420 Rubio-Ordóñez, A., Valverde-Vaquero, P., Corretgé, L. G., Cuesta-Fernández, A.,
1421 Gallastegui, G., Fernández-González, M., Gerdes, A., 2012. An Early Ordovician
1422 tonalitic-granodioritic belt along the Schistose-Greywacke Domain of the Central
1423 Iberian Zone (Iberian Massif, Variscan Belt). Geol. Mag. 149, 927–939.
- 1424 Rudnick, R.L., Gao, S., 2003. Composition of the Continental Crust. In: Treatise on
1425 Geochemistry (Holland, H.D., Turekian, K.K., eds.), vol. 3, 1–64. Elsevier-
1426 Pergamon, Oxford.
- 1427 Sánchez-García, T., Bellido, F., Quesada, C., 2003. Geodynamic setting and
1428 geochemical signatures of Cambrian–Ordovician rift-related igneous rocks (Ossa-
1429 Morena Zone, SW Iberia). Tectonophysics 365, 233–255.

- 1430 Sánchez-García, T., Quesada, C., Bellido, F., Dunning, G., González de Tánago, J.,
 1431 2008. Two-step magma flooding of the upper crust during rifting: the Early Paleozoic
 1432 of the Ossa-Morena Zone (SW Iberia). *Tectonophysics* 461, 72–90.
- 1433 Sánchez-García, T., Bellido, F., Pereira, M.F., Chichorro, M., Quesada, C., Pin, Ch.,
 1434 Silva, J.B., 2010. Rift-related volcanism predating the birth of the Rheic Ocean
 1435 (Ossa-Morena zone, SW Iberia). *Gondwana Res.* 17, 392–407.
- 1436 Sánchez-García, T., Quesada, C., Bellido, F., Dunning, G.R., Pin, Ch., Moreno-Eiris,
 1437 E., Perejón, A., 2016. Age and characteristics of the Loma del Aire unit (SW Iberia):
 1438 Implications for the regional correlation of the Ossa-Morena Zone. *Tectonophysics*
 1439 681, 58–72.
- 1440 Sánchez-García, T., Chichorro, M., Solá, R., Álvaro, J.J., Díez Montes, A., Bellido, F. et
 1441 al. 2019. The Cambrian–Early Ordovician Rift Stage in the Gondwanan Units of the
 1442 Iberian Massif. In: *The Geology of Iberia: A Geodynamic Approach* (Quesada, C.,
 1443 Oliveira, J.T., eds.). *Regional Geol. Rev.* 1, 27–74.
- 1444 Schaltegger, U., Abrecht, J., Corfu, F., 2003. The Ordovician orogeny in the Alpine
 1445 basement: constraints from geochronology and geochemistry in the Aar Massif
 1446 (Central Alps). *Schweizerische Miner. Petrogr. Mitteil.* 83, 183–195.
- 1447 Shaw, J., Johnston, S., Gutiérrez-Alonso, G., Weil, A.B., 2012. Oroclines of the
 1448 Variscan orogen of Iberia: paleocurrent analysis and paleogeographic implications.
 1449 *Earth Planet. Sci. Lett.* 329–330, 60–70.
- 1450 Shaw, J., Gutiérrez-Alonso, G., Johnston, S., Pastor Galán, D., 2014. Provenance
 1451 variability along the early Ordovician north Gondwana margin: paleogeographic and
 1452 tectonic implications of U–Pb detrital zircon ages from the Armorican Quartzite of
 1453 the Iberian Variscan belt. *Geol. Soc. Am. Bull.* 126(5–6), 702–719.
- 1454 Solá, A.R., 2007. *Relações Petrogeoquímicas dos Maciços Graníticos do NE*
 1455 *Alentejano*. PhD, Univ. Coimbra.
- 1456 Solá, A.R., Pereira, M.F., Williams, I.S., Ribeiro, M.L., Neiva, A.M.R., Montero, P., Bea,
 1457 F., Zinger, T., 2008. New insights from U–Pb zircon dating of Early Ordovician

- 1458 magmatism on the northern Gondwana margin: the Urro formation (SW Iberian
1459 Massif, Portugal). *Tectonophysics* 461, 114–129.
- 1460 Stern, R.J., 2002. Crustal evolution in the East African Orogen: a neodymium isotopic
1461 perspective. *J. African Earth Sci.* 34, 109–117.
- 1462 Stille, H., 1939. Bemerkungen betreffend die “Sardische” Faultung und den Ausdruck
1463 “Ophiolitisch”. *Zeits. Deuts. Gess. Geowiss.* 91, 771–773.
- 1464 Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic
1465 basalts: implications for mantle composition and processes. In: *Magmatism in the*
1466 *Ocean Basins* (Saunders, A.D., Norry, M.J., eds.). *Geol. Soc., Spec. Publ.* 42, 13–
1467 345.
- 1468 Syme, E.C., 1998. Ore-Associated and Barren Rhyolites in the central Flin Flon Belt:
1469 Case Study of the Flin Flon Mine Sequence. Manitoba Energy and Mines, Open File
1470 Report OF98–9, 1–32
- 1471 Takahashi, Y., Yoshida, H., Sato, N., Hama, K., Yusa, Y., Shimizu, H., 2002. W- and
1472 M-type tetrad effects in REE patterns for water-rock systems in the Tono uranium
1473 deposit, central Japan. *Chem. Geol.* 184, 311–335.
- 1474 Talavera, C., 2009. Pre–Variscan magmatism of the Central Iberian Zone: chemical
1475 and isotope composition, geochronology and geodynamic significance. PhD, Univ.
1476 Granada.
- 1477 Talavera, C., Bea F, Montero P., Whitehouse, M., 2008. A revised Ordovician age for
1478 the Sisargas orthogneiss, Galicia (Spain). Zircon U–Pb ion-microprobe and LA–
1479 ICPMS dating. *Geol. Acta* 8, 313–317.
- 1480 Talavera, C., Montero, P., Bea, F., González Lodeiro, F., Whitehouse, M., 2013. U–Pb
1481 Zircon geochronology of the Cambro–Ordovician metagranites and metavolcanic
1482 rocks of central and NW Iberia. *Int. J. Earth. Sci.* 102, 1–23.
- 1483 Taylor, S.R., McLennan, S.M., 1985. *The Continental Crust: Its Composition and*
1484 *Evolution*. Blackwell, London, 312 pp.


- 1485 Teichmüller, R., 1931. Zur Geologie des Thyrrhenisgebietes, Teil 1: Alte und junge
1486 Krustenbewegungen im südlinchen Sardinien. Abh. Der wissen. Gess. Göttingen
1487 (Math.-Phys. Kl) 3, 857–950.
- 1488 Teipel, U., Eichhorn, R., Loth, G., Rohrmüller, J., Höll, R., Kennedy, A., 2004. U–Pb
1489 SHRIMP and Nd isotopic data from the western Bohemian Massif (Bayerischer
1490 Wald, Germany): Implications for Upper Vendian and Lower Ordovician magmatism.
1491 Int. J. Earth Sci. 93, 782–801.
- 1492 Thompsom, M.D., Grunow, A.M., Ramezani, J., 2010. Cambro–Ordovician
1493 paleogeography of the Southeastern New England Avalon Zone: Implications for
1494 Gondwana breakup. Geol. Soc. Am. Bull. 122, 76–88.
- 1495 Tichomirowa, M., Berger, H.J., Koch, E.A., Belyatski, B., Götze, J., Kempe, U.,
1496 Nasdala, L., Schaltegger, U., 2001. Zircon ages of high-grade gneisses in the
1497 Eastern Erzgebirge (Central European Variscides) – constraints on origin of the
1498 rocks and Precambrian to Ordovician magmatic events in the Variscan foldbelt.
1499 Lithos 56, 303–332.
- 1500 Tichomirowa, M., Sergeev, S., Berger, H.J., Leonhardt, D., 2012. Inferring protoliths of
1501 high-grade metamorphic gneisses of the Erzgebirge using zirconology,
1502 geochemistry and comparison with lower-grade rocks from Lusatia (Saxothuringia,
1503 Germany). Contrib. Mineral. Petrol. 164, 375–396.
- 1504 Valverde-Vaquero, P., Dunning, G.R., 2000. New U–Pb ages for Early Ordovician
1505 magmatism in Central Spain. J. Geol. Soc. London 157, 15–26.
- 1506 Valverde-Vaquero, P., Marcos, A., Farias, P., Gallastegui, G., 2005. U–Pb dating of
1507 Ordovician felsic volcanism in the Schistose Domain of the Galicia-Trás-os-Montes
1508 Zone near Cabo Ortegal (NW Spain). Geol. Acta 3, 27–37.
- 1509 Valverde-Vaquero, P., Farias, P., Marcos, A., Gallastegui, G., 2007. U–Pb dating of
1510 Siluro–Ordovician volcanism in the Verín synform (Orense, Schistose Domain,
1511 Galicia-Trás-os-montes Zone). Geogaceta 41, 247–250.

- 1512 Vilà, M., Pin, C., Enrique, P., Liesa, M., 2005. Telescoping of three distinct magmatic
1513 suites in an orogenic setting: Generation of Hercynian igneous rocks of the Albera
1514 Massif (Eastern Pyrenees). *Lithos* 83, 97–127.
- 1515 Villaseca, C., Castiñeiras, P., Orejana, D., 2013. Early Ordovician metabasites from the
1516 Spanish Central System: A remnant of intraplate HP rocks in the Central Iberian
1517 Zone. *Gondwana Res.* 27, 392–409.
- 1518 Villaseca, C., Merino Martínez, E., Orejana, D., Andersen, T., Belousova, E., 2016.
1519 Zircon Hf signatures from granitic orthogneisses of the Spanish Central System:
1520 Significance and sources of the Cambro–Ordovician magmatism in the Iberian
1521 Variscan Belt. *Gondwana Res.* 34, 60–83.
- 1522 Von Quadt, A., 1997. U–Pb zircon and Sr–Nd–Pb whole-rock investigations from the
1523 continental deep drilling (KTB). *Geol. Rundsch.* 86 (suppl.), S258–S271.
- 1524 Von Raumer, J.F., Stampfli, G.M., 2008. The birth of the Rheic Ocean – early
1525 Palaeozoic subsidence patterns and tectonic plate scenarios. *Tectonophysics* 461,
1526 9–20.
- 1527 Von Raumer, J.F., Stampfli, G.M., Borel, G., Bussy, F., 2002. The organization of pre–
1528 Variscan basement areas at the Gondwana margin. *Int. J. Earth Sci.* 91, 35–52.
- 1529 Von Raumer, J.F., Bussy, F., Schaltegger, U., Schulz, B., Stampfli, G., 2013. Pre–
1530 Mesozoic Alpine basements: their place in the European Paleozoic framework.
1531 *Geol. Soc. Am. Bull.* 125, 89–108.
- 1532 Von Raumer, J.F., Stampfli, G.M., Arenas, R., Sánchez Martínez, S., 2015. Ediacaran
1533 to Cambrian oceanic rocks of the Gondwanan margin and their tectonic
1534 interpretation. *Int. J. Earth Sci.* 104, 1107–1121.
- 1535 Whalen, J.B., Currie, K.L., Chappell, B.W., 1987. A-type granites: Geochemical
1536 characteristics, discrimination and petrogenesis. *Contr. Miner. Petrol.* 95, 407–419.
- 1537 Winchester, J.A., Floyd, P.A., 1977. Geochemical discrimination of different magma
1538 series and their differentiation products using immobile elements. *Chem. Geol.* 20,
1539 325–343.

- 1540 Zeck, H.P., Whitehouse, M.J., Ugidos, J.M., 2007. 496 ± 3 Ma zircon ion microprobe
1541 age for pre-Hercynian granite, Central Iberian Zone, NE Portugal (earlier claimed
1542 618 ± 9 Ma). *Geol. Mag.* 144, 21-31.
- 1543 Zurbruggen, R., 2015. Ordovician orogeny in the Alps – a reappraisal. *Int. J. Earth Sci.*
1544 104, 335–350.
- 1545 Zurbruggen, R., 2017. The Cenerian orogeny (early Paleozoic) from the perspective of
1546 the Alpine region. *Int. J. Earth Sci.* 106, 517–529.
- 1547 Zurbruggen, R., Franz, L., Handy, M.R., 1997. Pre-Variscan deformation,
1548 metamorphism and magmatism in the Strona-Ceneri Zone (southern Alps of
1549 northern Italy and southern Switzerland). *Schweiz. Miner. Petrograph. Mitteil.* 77,
1550 361–380.
- 1551

FIGURE CAPTIONS


Figure 1. A. Reconstruction of the south-western European margin of Gondwana in Late Carboniferous–Early Permian times; modified from Pouclet et al. (2017). B. Setting of samples in the Central Iberian and Galicia-Trás-os-Montes zones; 59- Carrascal, 68- Guadarrama, 70- Sanabria, 74- Miranda do Douro, 77- Olo de Sapo, 79- Portalegre, 82- Saldanha, 84- San Sebastián, 99- Urra, Sa Sanabria; modified from Sánchez-García et al. (2019). C. Setting of samples in the Montagne Noire and Mouthoumet massifs; Am1-2 Larroque hamlet (Ambialet), Stg- St.Géraud Sta- St. André, Mj- Montjoi, Qu- Quintillan, GL- Roque de Bandies, VLR- Villerouge-Termenès, VIN- Le Vintrou, HER- Gorges d'Héric (Caroux massif), Ax1- S Mazamet (Nore massif), Ax2 (Rou)- S Rouayroux (Agout massif); modified from Álvaro et al. (2016). D. Setting of Pyrenean samples; modified from Casas et al. (2019). E. Setting of Sardinian samples; CS 2,3,4,8- Spartivento Cap, T2- Tuerreda, CC5- Cuile Culurgioni, MF1- Monte Filau, MFS1-Monte Settiballas, PB- Punta Bianca; modified from Oggiano et al. (2010).



1568

1569

Figure 2. Stratigraphic comparison of the Cambro-Ordovician successions from the Central Iberian Zone, Galicia Trás-os-Montes Zone, Occitan Domain, Eastern Pyrenees and Sardinia; modified from Álvaro et al. (2014b, 2016, 2018), Pouclet et al. (2017) and Sánchez-García et al. (2019); abbreviations: **Ag Agualada**, **Ca** Campelles ignimbrites (ca. 455 Ma, Martí et al., 2014), **CD** Cadí gneiss (456 ± 5 Ma, Casas et al., 2010), **Cg** Canigó gneiss (472–462 Ma, Cocherie et al., 2005; Navidad et al., 2018), **Co** Cortalets metabasite (460 ± 3 Ma, Navidad et al., 2018), **Cs** Casemí gneiss (446 ± 5 and 452 ± 5 Ma, Casas et al., 2010), **Es** Estremoz rhyolites (499 Ma, Pereira et al., 2012), **Ga Galiñero**, **GA** Golfo Aranci orthogneiss (469 ± 3.7 Ma, Giacomini et al., 2006), **GH** Gorges d’Heric orthogneiss (450 ± 6 Ma, Roger et al., 2004), **La** Larroque Volcanic Complex, **Ma** Marialles microdiorite (453 ± 4 Ma, Casas et al., 2010), **Lo** Lodè orthogneiss (456 ± 14 Ma, Helbing and Tiepolo, 2005), **MF** Monte Filau-Capo Spartivento orthogneiss (449 ± 6 Ma, Ludwing and Turi, 1989; 457.5 ± 0.3 and 458.2 ± 0.3 Ma, Pavanetto et al., 2012), **Mo Mora** (493.5 ± 2 Ma, Dias Da Silva et al., 2014), **Nu** Núria gneiss (457 ± 4 Ma, Martínez et al., 2011), **OS** Ollo de Sapo rhyolites and ash-fall tuff beds (ca. 477 Ma., Gutiérrez-Alonso et al., 2016), **PL** Pont de Larn orthogneiss (456 ± 3 Ma, Roger et al., 2004), **Qb** Queralbs gneiss (457 ± 5 Ma, Martínez et al., 2011), **PB** Punta Bianca orthogneiss (broadly Furongian–Tremadocian in age), **PC** Porto Corallo dacites (465.4 ± 1.9 and 464 ± 1 Ma, Giacomini et al., 2006; Oggiano et al., 2010), **Ri** Ribes granophyre (458 ± 3 Ma, Martínez et al., 2011), **Rf** Roc de Frausa gneiss (477 ± 4, 476 ± 5 Ma, Cocherie et al., 2005; Castiñeiras et al., 2008), **So** Somaill orthogneiss (471 ± 4 Ma, Cocherie et al. 2005), **Sa Saldanha** (483.7 ± 1.5; Dias da Silva, 2014), **SE** Saint Eutrope gneiss (455 ± 2 Ma, Pitra et al., 2012), **Ta** Tanaunella orthogneiss 458 ± 7 Ma (Helbing and Tiepolo, 2005), **Tr** Turchas and **Ur** Urra rhyolites.



1594

1595

Figure 3. Relative probability plots of the age of the Cambrian–Ordovician magmatism for (A) the Ollo de Sapo domain from the Central Iberian Zone; and (B) Pyrenees (Guilleries and Gavarres massifs), French Central Massif (including Montagne Noire), Sardinia, Corsica and Sicily (n = number of analyses). Data obtained from references cited in the text.




Figure 4. SiO_2 vs. Zr/TiO_2 and Zr/TiO_2 vs. Nb/Y plots (Winchester and Floyd, 1977) showing the composition of new samples (purple diamonds) and those taken from the literature (green triangles).




Figure 5. Zr/Ti vs. Nb/Y discrimination diagram (after Winchester and Floyd, 1977; Pearce, 1996). A. Lower–Middle Ordovician rocks of Iberian Massif (Central Iberian and Galicia-Trás-os-Montes zones). B. Middle–Upper Ordovician rocks of the eastern Pyrenees. C) Middle Ordovician rocks of the Occitan Domain. C–D. Middle–Upper Ordovician rocks of Sardinia.





Figure 6. Upper Crustal-normalized REE patterns (Rudnick and Gao, 2003) with average values for all distinguished groups; symbols as in Figure 4.



CIZ -	Mid-Ord	○ OSS
GTmZ	Low-Ord	✕ OG ✦ LG ✧ VOL ✦ GRA

PYR	Upp-Ord	✧ G1 ▼ CASEMI □ CADI ◆ V2
	Mid-Ord	▼ V1 ● G2 ✕ G3

Occ.	Upp-Ord	◆ OG-OD
Dom.	Mid-Ord	▲ VOL-OD

Sar.	Upp-Ord	★ OG-SUO ✕ VOL-SUO
	Mid-Ord	✦ OG-SMO ■ VOL-SMO

Figure 7. Chondrite-normalized REE patterns (Sun and McDonough, 1989) for all study samples.






Figure 8. Multi-element diagram normalised to Primitive Mantle of Palme and O'Neill (2004) for all study samples.



1631 **Figure 9.** Chondrite-normalised isotope ratio patterns (Sun and McDonough, 1989) for
1632 standard comparison for all study samples. Blue area: limits of continental crustal
1633 values (Lower and Upper) of Rudnick and Gao (2003).
1634



References

- Continental Crust (limits: upper, UCC and lower, LCC) of Rudnick and Gao (2004)
- Ocean Island Basalts (OIB) of Sun and McDonough (1989)
- Enriched Mid-ocean ridge basalts (EMORB) of Sun and McDonough (1989)
- Chondrite of Sun & McDonough (1989)

Figure 10. Tectonic discriminating diagram of Zr vs. TiO_2 (Syme, 1998) for all study samples. Double-sided arrows indicate ranging of different fields: rhyolites in tholeiitic and calc-alkaline arc suites have Zr/TiO_2 ratios ranging from about 0.016 to 0.04, and extension-related rhyolites from about 0.13 to 0.28 (Syme, 1989).






Figure 11. Tectonic discriminating diagram of Y vs. Nb (Pearce et al., 1984) for all study samples.



1649 **Figure 12.** Zr vs. 10^4 Ga/Al discrimination diagram (Whalen et al., 1987).


1650



1651

1652

Figure 13. Zr–Nb plot diagram (Leat et al., 1986; modified by Piercey, 2011) for all study samples.






Table 1. Chemical analyses of magmatic rocks. ICP and ICP–MS methods at ACME–LABS in Canada.

Sample	PYRENEES			MONTAGNE NOIRE				SARDINIA										
	Albera	Pallaresa	Andorra	Axial		Zone		External					Zone		Inner		Zone	
	A-08-03	IC1803	BN 1	Ax - 1	Ax - 2	HER	VIN	CC 5	CS 2	CS 3	CS 5	CS 8	MF 1	MFS 1	T 2	PB50	PB100	
Long. (E)	3°7'39"	1°27'43"	1°33'29"	2°13'50"	2°33'58"	2°57'58"	2°13'50"	8°50'37"	8°50'35"	8°50'35"	8°50'40"	8°50'35"	8°50'47"	8°52'02"	8°48'54"	9°09'32"	9°09'32"	
Lat. (N)	42°25'2"	42°36'1"	42°32'30"	43°34'32"	43°29'3"	43°34'32"	43°17'45"	38°54'16"	38°52'38"	38°52'38"	38°52'36"	38°52'39"	38°54'58"	38°53'57"	38°53'57"	41°11'1"	41°11'04"	
SiO ₂	68.38	71.67	69.18	70.38	67.43	68.31	73.97	76.43	75.14	76.52	76.61	76.36	72.13	75.94	75.55	68.93	67.24	
TiO ₂	0.57	0.63	0.61	0.36	0.64	0.61	0.20	0.08	0.08	0.09	0.04	0.06	0.31	0.13	0.18	0.41	0.46	
Al ₂ O ₃	15.68	14.24	15.05	14.90	15.76	15.39	13.82	13.28	12.81	11.80	12.71	12.63	13.80	13.16	12.94	16.32	15.79	
Fe ₂ O ₃	4.09	4.54	4.20	3.04	4.11	4.19	2.05	0.69	1.39	1.44	1.28	1.35	2.96	1.55	1.62	3.19	4.78	
MnO	0.07	0.06	0.05	0.04	0.04	0.04	0.04	0.01	0.01	0.01	0.01	0.01	0.02	0.03	0.04	0.08	0.08	
MgO	1.35	0.78	1.16	0.78	1.33	1.34	0.43	0.08	0.15	0.16	0.06	0.05	0.36	0.19	0.08	1.15	1.58	
CaO	0.21	0.53	1.78	1.22	1.44	1.58	0.62	0.32	0.25	0.15	0.20	0.35	0.61	0.38	0.17	3.05	2.70	
Na ₂ O	4.07	1.67	3.40	3.33	2.78	2.93	2.87	3.04	1.71	1.58	2.91	3.35	2.89	2.57	2.53	3.85	3.43	
K ₂ O	2.84	2.91	2.71	4.35	4.68	4.03	4.55	4.79	7.84	7.43	5.16	4.91	5.47	4.94	5.36	2.26	2.96	
P ₂ O ₅	0.17	0.24	0.20	0.21	0.2	0.19	0.18	0.15	0.05	0.05	0.03	0.04	0.12	0.11	0.07	0.15	0.14	
L.O.I.	2.03	2.60	1.50	1.2	1.3	1.2	1.2	1.1	0.4	0.7	0.9	0.8	1.1	0.9	1.4	0.90	0.70	
Total	99.05	99.42	99.42	99.51	99.30	99.39	99.73	99.90	99.69	99.79	99.78	99.78	99.47	99.75	99.78	99.97	99.37	
As	77.20	1.70	6.80	2.50	6.00	1.80	1.90	0.70	1.00	0.50	2.80	1.10	1.80	101.10	4.00	5.00	5.00	
Ba	742.50	388.00	398.00	499	1050	767	256	60	467	109	21	27	784	194	192	689.00	600.00	
Be	2.44	3.00	2.00	4.00	2.00	5.00	3.00	6.00	3.00	1.00	9.00	2.00	7.00	3.00	7.00	3.00	5.00	
Bi	0.30	0.20	0.10	0.20	0.20	0.20	0.40	0.30	0.10	0.10	0.10	0.10	0.10	0.70	0.40	4.00	4.00	
Cd	0.18	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	
Co	5.84	4.60	6.20	5.20	5.20	5.40	2.70	0.50	1.60	1.00	0.80	0.60	2.30	1.50	1.20	5.00	14.00	
Cs	9.79	5.60	4.90	14.30	7.10	6.80	7.30	4.20	3.40	1.60	4.50	4.60	6.40	3.90	4.10	4.20	9.40	
Cu	16.34	13.20	10.30	7.20	7.40	10.10	8.70	4.70	4.60	8.20	26.80	2.50	5.00	5.50	5.00	10.00	60.00	
Ga	21.03	19.80	18.80	19.10	19.20	18.90	16.70	19.30	14.90	15.30	19.40	19.20	20.70	19.00	19.90	17.00	18.00	
Hf	6.40	7.30	6.40	5.00	6.90	5.70	3.10	3.10	4.10	4.30	3.50	3.80	8.80	3.70	5.80	5.90	5.30	
Mo	1.20	0.90	1.00	0.60	0.90	0.60	0.30	0.70	0.70	0.70	0.80	0.50	1.70	0.80	1.60	2.00	2.00	
Nb	10.49	11.30	11.30	9.60	12.40	11.90	7.90	10.30	7.70	12.10	13.20	13.30	20.20	9.10	20.60	9.00	11.00	
Ni	16.56	8.00	7.70	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	20.00	80.00	
Pb	7.94	9.80	22.90	3.50	4.60	5.10	3.60	2.90	7.40	8.60	4.50	5.50	5.10	6.30	5.50	21.00	24.00	
Rb	124.40	123.70	137.20	204.6	161.6	142.2	188.2	289.9	206.1	187.4	294.1	275.1	208.7	256.4	227.1	85.00	118.00	
Sb	2.27	0.10	0.30	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	5.00	5.00	
Sc		10.00	10.00	6.00	9.00	9.00	4.00	3.00	3.00	4.00	4.00	4.00	15.00	4.00	8.00	9.00	12.00	
Sn	2.11	5.00	5.00	9.00	3.00	3.00	7.00	9.00	4.00	3.00	13.00	15.00	7.00	15.00	12.00	3.00	3.00	
Sr	158.00	201.80	83.70	91.20	160.30	150.10	68.70	30.70	73.90	25.20	7.90	8.10	59.90	45.60	25.00	217.00	167.00	
Ta	1.07	1.10	1.10	0.80	1.00	0.80	0.70	2.10	0.90	1.10	3.40	1.70	1.60	1.70	2.30	1.00	1.20	
Th	11.90	15.70	13.50	11.10	14.40	14.30	5.90	9.10	14.10	17.00	13.50	13.10	22.80	10.20	26.90	13.30	11.50	
U	3.70	5.10	4.60	4.10	3.60	3.20	4.80	3.30	2.90	3.20	3.50	3.50	4.60	8.10	4.90	4.50	2.20	
V	44.49	49.00	36.00	36.00	63.00	68.00	22.00	8.00	8.00	8.00	8.00	8.00	15.00	8.00	10.00	62.00	53.00	
W	1.80	1.90	2.50	3.20	2.60	1.60	3.00	5.60	0.90	2.10	5.20	3.00	2.40	4.40	3.50	1.00	20.00	
Y	29.29	43.90	50.60	28.30	38.40	36.20	27.80	28.00	60.10	53.60	44.40	46.00	61.60	31.80	55.80	29.00	24.00	
Zn	63.71	52.00	70.00	55.00	71.00	78.00	46.00	7.00	35.00	39.00	15.00	24.00	37.00	30.00	22.00	70.00	70.00	
Zr	233.30	263.20	237.10	174.40	249.20	219.10	93.70	73.50	93.80	105.10	62.20	74.50	311.80	108.10	161.90	245.00	214.00	
La	27.90	45.30	38.00	29.60	39.50	38.70	13.60	10.50	22.70	19.50	12.10	13.40	54.20	17.90	31.30	26.90	34.30	
Ce	59.00	86.90	75.50	58.10	77.00	78.20	26.70	21.60	42.10	39.70	26.20	29.90	109.80	37.40	97.60	53.20	70.50	
Pr	7.26	9.80	8.47	6.99	9.41	9.55	3.36	2.36	4.73	4.85	3.00	3.24	11.94	4.07	6.86	5.88	8.20	
Nd	27.83	35.60	31.20	26.00	36.40	36.40	12.60	8.40	16.60	17.10	10.50	10.90	44.70	15.00	24.00	21.60	29.40	
Sm	5.80	7.69	7.16	5.70	7.55	7.63	3.15	2.43	4.10	4.41	3.28	3.44	9.37	3.88	4.93	4.70	6.00	
Eu	0.98	1.05	1.03	0.87	1.27	1.15	0.41	0.14	0.43	0.13	0.06	0.09	1.17	0.30	0.19	0.95	0.93	
Gd	5.22	8.32	7.89	5.59	7.28	7.05	3.38	3.20	5.60	5.50	4.42	4.69	10.60	4.50	6.34	4.00	5.10	
Tb	0.87	1.26	1.27	0.89	1.17	1.10	0.67	0.69	1.13	1.18	1.03	1.07	1.70	0.82	1.27	0.70	0.80	
Dy	5.30	6.68	8.00	5.09	6.89	6.39	4.59	4.30	7.69	8.23	7.31	7.66	10.28	5.24	9.00	3.70	4.30	
Ho	1.06	1.52	1.73	0.99	1.42	1.30	0.98	0.91	1.91	1.91	1.59	1.65	2.13	1.12	2.01	0.70	0.80	
Er	2.98	4.52	4.96	2.64	3.92	3.56	3.07	2.85	5.80	6.46	5.35	5.38	6.25	3.64	6.17	2.20	2.10	
Tm	0.46	0.60	0.73	0.38	0.57	0.50	0.44	0.43	0.91	1.00	0.85	0.85	0.89	0.52	0.92	0.35	0.32	
Yb	3.00	3.98	4.72	2.33	3.56	3.11	2.83	2.95	5.81	6.60	6.10	6.16	5.53	3.70	6.04	2.50	2.20	
Lu	0.44	0.58	0.69	0.33	0.53	0.45	0.39	0.44	0.90	0.94	0.92	0.94	0.86	0.56	0.90	0.41	0.36	

1669 **Table 2.** Summarized geochemical features of the Furongian and Ordovician felsic
1670 episodes described in the text; data from Lancelot et al. (1985), Calvet et al. (1988),
1671 Valverde-Vaquero and Dunning (2000), Roger et al. (2004), Vilà et al. (2005),
1672 Giacomini et al. (2006), Díez-Montes (2007), Montero et al. (2007, 2009), Solá (2007),
1673 Zeck et al. (2007), Castiñeiras et al. (2008b), Talavera (2009), Casas et al. (2010),
1674 Navidad et al. (2010, 2018), Liesa et al. (2011), Martínez et al. (2011, 2018), Navidad
1675 and Castiñeiras (2011), Gaggero et al. (2012), Talavera et al. (2013), Villaseca et al.
1676 (2016), Pouclet et al. (2017), Cruciani et al. (2018) and this work. Abbreviations: *CIZ*
1677 Central Iberian Zone, *GTOMZ* Galicia-Trás-os-Montes Zone, *OCC* Occitan Domain,
1678 *PYR* Pyrenees and *SAR* Sardinia; * *sensu* Guitard (1970); *A/CNK* ratio is always
1679 peraluminous.

ORTHOGNEISS FACIES		code	composition	SiO2 wt. %	Na2O wt. %	K2O wt. %	A/CNK ratio	εNd	TDM (Ga)	¹⁴⁷ Sm/ ¹⁴⁴ Nd	area
(1) <i>Furongian-Mid Ordovician Suite</i>											
CIZ - Olo de Sapo orthogneiss	OG	K-rich dacite to rhyolite	75-60.3	3.9-0.1	5.9-3.4	3.1-1.0	-5.1 to -1.8	1.8-1.1	0.15-0.09	Sanabria (ca. 472 Ma) and Guadarrama (ca. 488-473 Ma)	
CIZ - Leucogneiss	LG	K-rich dacite to rhyolite	75.9-73.6	3.1-2.7	5.3-4.2	1.3-1.1	-5.1 to -4.9	4.1	0.22-0.18	Guadarrama	
CIZ - Metagranite	GRA	K-rich dacite to rhyolite	77-64.6	4.8-0.5	6.3-2.5	1.8-1.0	-5.2 to +2.6	3.6-0.9	0.19-0.09	NE Central System, Sanabria, Miranda do Douro (ca. 496-473 Ma), CIZ (496-471 Ma for Carrascal, Femosselle, Ledesma, Portalegre and Vitgudino granites)	
CIZ/GTMZ - Volcanic rocks	VOL	andesite to rhyolite	79.3-64.6	3.2-0.1	6.3-2.2	2.7-1.1	-5.5 to -1.6	1.7-1.3	0.15-0.13	Saldanha Fm. in GTMZ, Olo de Sapo Fm. in Sanabria, and Urza Fm.	
CIZ - San Sebastián orthogneiss	OSS	rhyolite	75.4-73.8	3.1-2.5	5.4-4.9	1.2-1.1	-4.0 to 0	1.6-1.2	0.14-0.14	Sanabria (ca. 470-465 Ma)	
PYR - augengneiss	G2*	dacite to rhyolite	73.6-68.3	3.9-3.2	4.4-2.5	1.2-1.1	-4.4 to -3.0	1.4-1.2	0.14-0.13	ca. 476-462 Ma	
PYR - orthogneiss	G3*	K-rich dacite	73.5-68.4	2.9-2.4	4.4	1.2	-4.2	1.33	0.13	ca. 463 Ma	
PYR - volcanic rocks	V1	Na-rich rhyolite	73.5-68.4	7.8-2.4	3.2-1.3	2.0-1.1	-5.1 to -2.6	1.7-1.6	0.19-0.13	Pierrefite Fm. and Albera massif (ca. 472-465 Ma)	
OCC - volcanic rocks	VOL-OD	K-rich dacite to rhyolite	75.6-66.7	3.7-0.6	9.3-2.3	2.4-1.3				Saint-Sernin-sur-Rance and Saint-Salvi-de-Carcavès nappes	
SAR - orthogneiss	OG-SMO	K-rich rhyolite	74-67.2	3.8-2.6	5.8-2.3	1.3-1.1				ca. 469 Ma	
SAR - volcanic rocks	VOL-SMO	K-rich dacite to rhyolite	76.7-67.6	4.7-1.9	5.4-2.9	2.0-1.2			0.16	ca. 464-462 Ma	
(2) <i>Upper Ordovician Suite</i>											
PYR - orthogneiss	G1*	K-rich dacite to rhyodacite	76.4-73.4	3.1-2.6	5.3-4.7	1.2-1.1	-5.3 to -3.1	2.7-1.5	0.17-0.12	ca. 457 Ma	
PYR - orthogneiss	CADi	K-rich dacite to rhyodacite	69.4	3	4	1.2	-4.1	1.5	0.13	Cadí massif (ca. 456 Ma)	
PYR - orthogneiss	CASEMi	K-rich dacite to rhyodacite	76-71.9	4-1.8	6.3-3.2	1.2-0.9	-3.6 to -1.3	2.6-1.3	0.17-0.13	Casemí massif (ca. 451-446)	
PYR - volcanic rocks	V2	andesite to rhyodacite	86.1-63	6-0	4.3-0.6	3.6-1	-5.1 to -2.6	1.7-1.6	0.14-0.14	Ribes de Freser, Andorra (ca. 457 Ma), Pallaresa (ca. 453 Ma), Els Maïges (ca. 455.2 Ma)	
OCC - orthogneiss	OG-OD	K-rich dacite to rhyolite	73.9-67.4	3.3-2.8	4.7-4	1.3-1.2	-4 to -3.5	1.8-1.4	0.15-0.13	Gorges d'Héric (ca. 450 Ma; Caroux), S Mazamet (Nore), S Rouairoux (Agout), Le Vintrou	
SAR - External Zone orthogneiss	OG-SUD	K-rich dacite to rhyolite	76.6-72.1	3.3-1.6	7.8-4.8	1.3-1.1	-3.3 to -1.6	4.2-1.2	0.19-0.12	Capo Spartivento, Culla Culurgioni, Tuerredda, Monte Filau, Monte Setiballas (ca. 458-457 Ma)	
SAR - Nappe Zone volcanic rocks	VOL-SUD	K-rich dacite to rhyodacite	76.7-70.7	3.3-1.6	7.8-4.8	1.3-1.1				Truzzulla Fm. at Monte Grighini	