Editors of Solid Earth

Dear Editors,

Please find enclosed the reviewed version of the paper entitled: "The enigmatic curvature of Central Iberia and its puzzling kinematics.", co-authored by Daniel Pastor-Galán, Gabriel Gutiérrez-Alonso and Arlo B. Weil to be considered in Solid Earth.

Following this letter a point by point answered to the reviewers is attached.

Sincerely,

Daniel Pastor Galán on behalf of all co-authors

Response to Anonymous Referee #1 (R1 hereafter)

We want to thank R1 the time invested in the review and the nice words about our manuscript.

We have followed all the suggestions and corrected all typos marked in the annotated PDF. We have also reviewed the text as indicated by R1 and identified some other mistakes.

R1 points out that the available structural information is not so relevant as the paleomagnetic one to rule out the Central Iberian curve as a secondary orocline. Following R1 recommendation, we highlighted the milestones and limitations of the structural analyses in the Central Iberian curve in section **6 Get Back: Ideas flowing out and endless questions:**

"In addition are the curved traces of C1 fold-axes, whose geometry and kinematics are reasonably well constrained around Galicia Tras-os-Montes (e.g. Dias da Silva et al., in press), but in many other areas their strong curvature remain largely unstudied (Fig. 2) and therefore, to date we can only speculate on their origin."

Finally, R1 is right on the previous figure 11. We agree it is not a fully necessary figure and their content is properly described and referred in the text. We have removed this figure from the revised version of the paper.

Response to Referee #2 (Dominique Jacques)

We want to thank Dominique Jacques for the thorough review and insights provided.

We have followed all the suggestions and corrected all typos, spelling mistakes and suggestions annotated in the attached PDF.

Apart from stylistic and orthographic corrections Dominique Jacques has some in detail comments:

"the most significant of these revisions concerns the supposed existence and nature of the post-Variscan C4 stage. The authors suggest a subdivision between a C3 (315-300Ma) and C4 (300-280Ma deformation stage, although both stages are characterized by N-S shortening and the widespread development of shear zones. The main distinction is the contrast of mainly ductile C3 folding and C4 brittle faulting. I would argue that a gradual evolution from ductile to brittle deformation as the orogeny is uplifted and cools does not warrant a separate deformation stage if the dynamics remain similar. Figure 13E even suggests continued tightening of the orocline, prior to E2."

Here Dominique is right, C3 and C4 are roughly N-S shortening events (in present day coordinates), that likely respond to the same strain field. The same could be said in the Iberian sector of the Variscan belt about C1 and C2. This paper is a review and we followed the most recent terminology for the Central Iberian Zone for consistency (Martínez Catalán et al., 2014). In addition, we think that phase naming does not need to reflect only dynamic or kinematic separate events, but also structural styles or metamorphic conditions changing progressively in accordance with the orogenic evolution. In this sense, we think C1, C2, E1 C3, C4 and E2 is a really useful terminology for the Central Iberian Zone. We have slightly changed the description of C4 to clarify this point:

"The N-S shortening (in present day coordinates) of C3 deformation continued through the Early Permian under brittle conditions (so-called C4 event) (e.g. Dias da Silva et al., in press) and overlapped with the formation of E2 extensional faults (Fig. 2A; Dias and Ribeiro 1991; Dias et al. 2003; Rubio Pascual et al., 2013; Arango et al., 2013; Fernández-Lozano et al. 2019; Dias da Silva et al., in press). C4 N-S compression produced a series of NNE-SSW and NNW-SSE brittle faults (Gil Toja et al. 1985; Dias and Ribeiro 1991; Dias et al. 2003; Fernández-Lozano et al., 2019) and associated sub-vertical and sub-horizontal widespread kink-bands (e.g. Aller et al., 2020; Dias da Silva et al., in press) that are today exposed in Northern Iberia. E2 developed core complex-like structures with extensional shear zones that further telescoped M2 metamorphic isograds between the anatectic cores of gneiss domes and the associated hanging wall units. This event favored subhorizontal folding, and crenulation cleavage development in the footwall together with kink-band generation in the upper low-grade structural levels."

In addition, I also disagree with the notion that the early Permian (i.e. posterior to oroclinal buckling) was characterized by N-S compression. Instead, literature generally agrees that Gondwana-Laurussia convergence during this time period was NW-SE to W-E directed. In the southern Variscan orogeny this led to NW-SE transpression and strikeslip deformation, especially observed in Stephanian-Permian intramontane basins. Simultaneously, to the west of the Variscan orogeny, frontal collision of Gondwana and Laurussia led tothe formation of the Alleghenides, Mauritanides and the Ouachita-Marathon-Sonora orogenic belts. The authors do not take into account the related literature, which are not in accordance with their geodynamic model in Fig. 13E. Hence, I would suggest that the authors either (i) limit themselves to the geodynamics of oroclinal buckling (C3 - 315-295Ma) in their interpretation, or (ii) strongly extend their literature study on the kinematics of post-Variscan deformation (295-280Ma).

Dominique Jacques is right about the west- and south-wards evolution of the Variscan Alleghanian orogen. In our paper, however we are reviewing exclusively the structural and kinematic evolution of Iberia, where all studied C4 structures (developed in Early to Mid Permian) indicate N-S shortening in present coordinates (which would be WSW-WNW in paleogeographic coordinates). We have added extra references supporting the C4 N-S compression in Iberia. Our kinematic model is restricted to Iberia and it is out of the scope of the paper to fit it within a global or large scale plate reconstruction. We want to remark that strain patterns in Iberia do not necessarily indicate or contradict any particular far field stress. Instead, these strain patterns should be taken into account for regional paleogeographic and global tectonic models, and not the opposite.

"What about the stratigraphic successions of the Southern Central Iberian Zone and the West Asturian-Leonese zones, which are connected in the Shaw et al. (2012) model (Fig. 2C)? Can these indeed be connected from this viewpoint, because literature appears to suggest that their lithological succession is different. As your Fig. 3A demonstrates, the lithological succession in the WALZ is much more proximal to the Gondwana shelf than in the CIZ, which consists of more shallow marine lithologies.In a similar fashion, is the metamophic grade of regional metamorphism (mainly active during D1-D2) in both zones comparative?"

This is a good question that is still unsolved. The stratigraphy of the southermost part of the Central Iberian Zone has been studied in lesser detail, probably because of its poorer exposure.

Best stratigraphical sections coming from Almadén area are quite similar to the Tamames area. However, the area covered is small and the correlations are not yet certain. This is another exciting problem yet to be solved. Nevertheless, there is consensus that the stratigraphical architecture is controlled by the presence of structural highs and troughs that caused differences in the shallowness of the deposits and, if so, it cannot be correlated directly with proximity to the Gondwanan shelf. From this point of view, the deeper facies rocks in the WALZ only indicates the presence of a marked through, despite its location within the Gondwanan margin.

The enigmatic curvature of Central Iberia and its puzzling kinematics

Daniel Pastor-Galán^{1,2,3}, Gabriel Gutiérrez-Alonso^{4, 5}, Arlo B. Weil⁶

¹Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

²Department of Earth Science, Tohoku University

³Center for Northeast Asian Studies, Tohoku University, 41 Kawauchi, Aoba-ku, Sendai, 980-8576, Japan. pastor.galan.daniel.a8@tohoku.ac.jp

⁴Dept. of Geology. Faculty of Sciences. University of Salamanca. Plaza de la Merced s/n. 38007, Salamanca (Spain). gabi@usal.es

⁵Geology and Geography Department, Tomsk State University, Lenin Street, 36, Tomsk 634050 Russia

⁶Department of Geology, Bryn Mawr College, PA, USA 19010

Abstract

The collision between Gondwana and Laurussia that formed the latest supercontinent, Pangea, occurred during Devonian to Early Permian times and resulted in <u>a</u> large-scale orogeny that today transects Europe, northwest Africa and eastern North America. This orogen is characterized by an 'S' shaped corrugated geometry in Iberia. The northern curve of the corrugation is the well-known and studied Cantabrian (or Ibero-Armorican) Orocline and is convex to the east and towards the hinterland. Largely ignored for decades, the geometry and kinematics of the southern curvature, known as the Central Iberian curve, are still ambiguous and hotly debated. Despite the paucity of data, the enigmatic Central Iberian curvature has inspired a variety of kinematic models that attempt to explain its formation with little consensus. This paper presents the advances and milestones in our understanding of the geometry and kinematics of the Central Iberian curve from the last decade, with particular attention to structural and paleomagnetic studies.

When combined, the currently available datasets suggest that the Central Iberian curve did not undergo regional differential vertical-axis rotations during or after the latest stages of the Variscan orogeny, and did not form as the consequence of a single process. Instead, its core is likely a primary curve (i.e. inherited from previous physiographic features of the <u>Iberian</u> crust) whereas the curvature in areas outside the core are dominated by folding interference during the Variscan orogeny or more recent Cenozoic (Alpine) tectonic<u>event</u>s.

Keywords

Central Iberia Curve, Variscan orogen, Iberia, Cantabrian Orocline, Curved orogens,

Pangea

1 Introduction

Mountain belt systems are the most striking product of plate tectonics. In addition to their astonishing visual effectimpact, marking the locations where ancient and modern plates collided, orogenic belts often preserve a variety of rocks that have the potential to illuminate the entirety of the systems pre- and syn-orogenic historiesy. One of the most striking characteristics of the majority of most of Earth's orogens are their curvature in plan-view (e.g. van der Voo, 2004; Marshak, 2004; Rosenbaum, 2014). The degree of orogenic curvature may range from a few degrees of deflection in structural trend (e.g. Kopet Dag, Iran), to 180° of arc curvature (e.g. Kazakhstan arc and the Carpathians). The kinematics, structural and geodynamic implications of these systems are as varied as their geometries (Marshak, 2004; Weil and Sussman, 2004; Johnston et al., 2013). For example, some orogenic curvatures are hypothesized to be the consequence of physiographic features of the basement that pre-date orogen formation, such as irregular basin architectures or plate margin salients and recesses that formed from rift-todrift processes (e.g. Jura mountains, Hindle et al., 2000), which then control the growth geometry of the ensuing orogen. These systems are known as primary arcs, and reflect preorogenic geometries and show no significant or systematic vertical-axis rotations along their structural length. On the other hand, oroclines, as classically defined by Carey in 1956, involve systematic differential vertical-axis rotations subsequent to initial orogenic shortening: different sectors of an orogen rotate with variable magnitudes or in opposite directions (e.g. Li et al., 2012). Rotations in <u>o</u>-Proclines may occur at a range of scales, from thrust emplacement at upper crustal levels (e.g. Izquierdo-Llavall et al., 2018), up to a-lithospheric-scale vertical-axis folding (e.g. Li et al., 2018). They can occur as single curves (e.g. Maffione et al., 2009), coupled curves (Johnston, 2001), or in trains of curves (Li and Rosenbaum, 2014). Oroclines can form during the main orogenic building event, known as progressive oroclines (Johnston et al., 2013; e.g., the Wyoming salient, Yonkee and Weil, 2010, and Weil et al., 2010), or in-during a subsequent tectonic pulse, so-called secondary oroclines (Weil and Sussman, 2004). UnderstandingUnraveling the kinematics and mechanisms of curvature formation in mountain belts is a critical step to understanding orogenesis in 4D and to evaluate their geodynamic consequences and paleogeographic implications.

The Variscan-Alleghanian orogeny resulted in the suturing of Gondwana and Laurussia during Devonian-Carboniferous times, and ultimately led to the formation of the supercontinent, Pangea. This long and sinuous orogen runs for >8000 km along strike and is ca. 1000 km wide, transecting across Europe, to northwest Africa and into eastern North America. The final stages

of Pangea amalgamation (e.g. Nance et al., 2010) modified the Western Europe sector of the belt into its characteristic sinuous shape. <u>Today</u>, this tectonic belts, which today_tracess at least one, and perhaps four arcs from Poland to Brittany, and then across the Bay of Biscay (Cantabrian Sea) into Iberia, where the system is today-truncated by the <u>younger</u> Betic Alpine orogeny in southeast Iberia (Fig. 1; e.g. Weil et al., 2013). The southern truncation of the Variscan in Europe hinders a precise correlation with equivalent age outcrops in NW Africa.

Within the Iberian Peninsula, the orogen is characterized by two large-scale curves (Fig. 2): (1) to the north is the well-studied and nearly 180° secondary orocline, the Cantabrian (a.k.a. Ibero-Armorican) Orocline, which buckled a segment of the Variscan belt from ~315 to ~290 Ma (e.g. Weil et al., 2019 and references therein); and (2) to the south is a curve with disputed magnitude and kinematics that, and is usually referred to as the Central Iberian curve/orocline or Castillian bend (Martínez-Catalán et al., 2015). Though there remains tremendous uncertainty on the geometry and kinematics of the Central Iberian curve, multiple hypotheses exist as to its nature, and disagreements continue on its importance in the tectonic evolution of Europe during the waning stages of Paleozoic global supercontinent construction. The diversity of author's interpretations of the Central Iberian curve range from a nonexistent structure (Dias et al., 2016), to being one of the most important pieces toin our understanding of the late Carboniferous and Permian geodynamics of the Iberian Variscan system (e.g. Martínez-Catalán et al., 2011; 2014).

This paper reviews the most recent advances on the geometry and kinematics of the Central Iberian curve, synthesizing what we know and what we don't, and ending with a discussion of the main unsolved issues. We hope that this paper fosters novel studies that will lead to a better understanding of when and which mechanisms acted in the aftermath of the Variscan-Alleghanian orogeny.

2 The long and winding orogen

The Variscan (Europe-NW Africa)-Alleghanian (North America) orogeny is a continentalscale tectonic system (1000 km wide and ≥8000 km long) that sutured Gondwana and Laurussia together, forming the supercontinent Pangea (e.g. Domeier and Torsvik, 2014; Edel et al., 2018; Pastor-Galán et al., 2019a). The fragments of this system are now dispersed over three continents, Europe, Africa and North America due to the Mesozoic break-up of Pangea (Buiter and Torsvik, 2014; Keppie, 2015). This orogen formed as a consequence of a long and protracted tectonic history that involved several different events, from initial convergence (ca. 420 Ma; e.g. Franke et al., 2017), to the consumption of multiple putative oceanic tracts and/or basins that existed between Gondwana and Laurussia (ca. 280 Ma; e.g. Kirsch et al., 2012). The Variscan-Alleghanian orogen itself represents the closing of at least one major ocean, the Rheic (e.g. Nance et al., 2010), whose axial ridge likely failed or subducted at ca. 395 Ma along its paleo-northern margin (e.g. Woodcock et al., 2007; Gutiérrez-Alonso et al., 2008a). Perhaps the orogeny involved other large oceans (Stampfli and Borel, 2002; Franke et al., 2017; 2019), but most surely involved several minor seaways and basins that existed between Gondwana, Laurussia, and several intervening micro-continents (e.g. Azor et al., 2008, Dallmeyer et al., 1997; Kroner and Romer, 2013; Díez-Fernández et al., 2016; Pérez-Cáceres et al., 2017). The final continent-continent collision began after closure of all oceans and intervening seaways. The commencement of this deformation was diachronistic and became progressively younger westwards (in present-day coordinates): with Devonian continent-continent collisions along the eastern boundary, progressing to earliest Permian ages in the westernmost sector (McWilliams et al., 2013; Chopin et al., 2014; López-Carmona et al., 2014; Franke et al., 2017).

The present-day geometry of the Variscan-Alleghanian systems has a contorted trace (Fig. 1). In Europe, from east to west, the trend starts with a prominent curve around the Bohemian massif (e.g. Tait et al., 1996), followed by a deflection in the Ardennes-Bravant (e.g. Zegers et al., 2003). In Brittany the outer curvature of the Cantabrian or Ibero-Armorican orocline begins (e.g. van der Voo et al., 1997), and wraps nearly 180° around and across the Bay of Biscay as it turns in NW Iberia. The Central Iberian curve marks the final concave to the west curve (in present-day coordinates) and is the focus of this paper (e.g. Aerden, 2004; Martínez Catalán, 2011; Shaw et al., 2012). The orogen continues in North America where, from north to south, it has salients and recesses that undulate back and forth from Atlantic Maritime Canada (e.g. O'Brien, 2012) down along the Pennsylvanian and <u>into the</u> Alabama curves (e.g. Thomas, 1977).

Interpretation on the origin of these curvatures varies widely. The curvatures in North America are argued to be the result of a preexisting irregular margin of Laurentia due to the break-up of the Rodinia supercontinent, which resulted in the formation of orogenic salients and recesses during subsequent Appalachian collision (e.g. Rankin, 1976; Thomas, 1977, 2004). In this case, vertical-axis rotations affected only mainly the upper crustal levels during orogenesis (e.g. Marshak, 1988; Bayona et al., 2003; Hnat and van der Pluijm, 2011). In Europe, the Bohemian and Ardennes-Bravant massif curvatures have poor kinematic constraints. In the Bohemian Massif, some suggest secondary rotations that formed an orocline (Tait et al., 1996), while others suggest little to no vertical-axis rotations and a primary arc (Chopin et al., 2012). The Ardennes-Bravant Massif recorded some vertical-axis-rotations (e.g. Molina-Garza and

Zeijderveld, 1996), but it is unclear if these are a response to progressive or secondary oroclinal bending, or whether rotations only affected the upper crust. The most outstanding example of Variscan-Alleghanian orogen curvature is exposed in the Iberian Massif, with the Cantabrian Orocline and the coupled curvature of Central Iberia.

2.1 Two of us: The Variscan orogen in Iberia

The western half of the Iberian Peninsula constitutes the Iberian massif, one of the largest exposures of the Variscan orogen and the only place that contains an almost continuous cross section of the orogen (Fig. 2; e.g. Lotze 1945, Julivert 1974, Pérez-Estaún et al., 1991; Ayarza et al., 1998; Simancas et al., 2003; Ribeiro et al. 2007, Martínez Catalán et al., 2014, 2019). The majority of the Iberian Massif contains Gondwanan affinity rocks (e.g. Murphy et al., 2008; Pastor-Galán et al., 2013a; Gutiérrez-Marco et al., 2017) and likely represents a proximal piece of the Gondwana margin until its final amalgamation with Pangea (e.g. Pastor-Galán et al., 2013b). Owing to the stratigraphic, structural and petrological styles, the Iberian Massif has been traditionally divided into six tectonostratigraphic zones (Fig. 2; Lozte, 1945; Julivert, 1971): (1) The Cantabrian Zone represents a Gondwanan thin-skinned foreland fold-and-thrust belt. It has overall low-grade internal deformation and metamorphism, and represents shortening that occurred during Mississippian times (e.g. Marcos and Pulgar, 1982; Pérez Estaún et al., 1988; Gutiérrez-Alonso 1996; Alonso et al., 2009; Pastor-Galán et al., 2009; 2013b). (2) The West-Asturian Leonese Zone represents a metamorphic fold-and-thrust belt with Bbarrovian metamorphism that collapsed coevally with thrust emplacement onto the Cantabrian Zone (e.g. Martínez-Catalán et al., 1992; Alcock et al., 2009; Martínez-Catalán et al., 2014). (3) The Central Iberian Zone represents the Gondwanan hinterland with Barrovian and Buchan metamorphism and is intruded by igneous rocks of various ages (e.g. Macaya et al., 1991; Díez Balda, 1995; Gutiérrez-Alonso et al., 2018). (4) The Ossa-Morena Zone represents the most distal zone of the Gondwana platform, and is characterized by a metamorphic fold-and-thrust belt with dominantly sinistral displacement (e.g. Robardet and Gutiérrez-Marco, 2004; Quesada, 2006). (5) The Galicia-Tras-os-Montes Zone represents a far-travelled allochthonous terrane that contains high-pressure units and relicts of oceanic-like crust (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). (6) The South Portuguese Zone represents a foreland fold-and-thrust belt with little internal deformation and metamorphism with Avalonian affinity and a strong sinistral left-lateral component of shear (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2016; Oliveira et al., 2019). Geographically, the external zones of the Gondwana margin are nested to the north into the core of the Cantabrian Orocline, whereas the hinterland zones are to the west and center of the massif (Fig. 2; e.g. Díaz Balda, 1995; Azor et al., 2019). The southwestern-most extent of Iberia contains a putative suture of the Rheic Oecean, as well as a piece of the Laurussian margin fold-and thrust belt, today preserved in the South Portuguese Zone (e.g. Pereira et al., 2012, 2017; Oliveira et al., 2019).

The Gondwanan authochton stratigraphy (Cantabrian, West Asturian-Leonese, Central Iberian and Ossa Morena zzones) consist of a Neoproterozoic arc and back-arc basin (e.g. Fernández-Seuárez et al., 2014), which evolved to a rift-to-drift Cambrian to Early Ordovician sequence and then to an Ordovician to Late Devonian passive margin basin sequence (e.g. Sánchez-García et al., 2019; Gutiérrez-Marco et al., 2019; Gutiérrez-Alonso et al., submitted). Overall the system transitioned from a relatively isolated Early Cambrian continental basin, to a restricted marine basin, to development of an open marine platform that was locally punctuated by magmatism (e.g. Gutiérrez-Alonso et al., 2008b; Palero-Fernández, 2015). The Ossa Morena $\mathbb{Z}_{\mathbb{Z}}$ one represents the outermost platform, followed by an intermediate platform characterized by an asymmetric horst (Central Iberian Zone) and graben (West-Asturian Leonese Zone), which ends in the innermost shelf environment of the Cantabrian $Z_{\overline{z}}$ one (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The differences between the West Asturian-Leonese and Central Iberian Zone are mainly deeper vs. shallower sedimentary facies (respectively) and a local Lower Ordovician unconformity in the Central Iberian Zone (Toledanian, e.g. Alvaro et al., 2018), that which places Lower Ordovician strata atop pre-Cambrian to Cambrian rocks (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The Central Iberian Zone is divided into two domains: (1) The Ollo de Sapo domain, which contains abundant Lower Ordovician calc-alkaline-magmatism of calc-alkaline affinity (e.g. Díez Montes, 2006; Gutiérrez-Marco et al., 2019); and (2) the 'Schistose–<u>G</u>ereywacke <u>d</u>-Domain' characterized by a predominance of outcrops of Neoproterozoic to Lower Cambrian sedimentary rocks (e.g. Gutiérrez-Marco et al., 2019 and references therein).

The Galicia Tras-os-Montes Zone (Farias et al., 1987) is a complex structural stack including a basal schistose unit (Parautochthon; Dias da Silva et al., in press Dias da Silva et al., 2020) structurally overlain by mafic rocks with an oceanic-like signature and other allochthonous far-traveled rocks under high-pressure metamorphism (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). The oceanic rocks of this zone are classically interpreted as a Rheic Ocean suture (e.g. Martínez Catalán et al., 2009). Recent interpretations support its origin as a minor oceanic basin or seaway within the realm of Gondwana (e.g. Pin et al., 2002; Arenas et al., 2016).

The South Portuguese Zone constitutes the Laurussian foreland fold-and-thrust belt in

the Iberian Variscides (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2017). It contains three units: (1) the Pulo de Lobo, a low grade metamorphic accretionary prism with clastic sedimentary rocks and basalts with MORB signature (e.g. Azor et al., 2019; Pérez-Cáceres et al., this volume) <u>some times considered an independent zone;</u> (2) The Iberian Pyrite Belt, which is a world class volcanogenic massive sulfide deposit formed between 390 and 330 Ma (e.g. Oliveira et al., 2019a; 2019b); and (3) the Baixo Alentejo Flysch, which is located to the southwest and is a syn-orogenic composite turbiditic sequence with ages from ~330 to ~310 Ma (Oliveira et al., 2019b). The boundary between the South Portuguese and Ossa Morena zones is a sinistral shear zone (so-called the so-called ftsSouthern Iberian shear zone in the Beja-Acebuches <u>oceanic-like unit</u>, <u>Crespo-Blanc and Orozco, 1988;</u> Quesada and Dallmeyer., 1994; Pérez-Cáceres et al., 2016) that contains a strongly deformed amphibolitic belt with oceanic affinity (Munha et al., 1986; Munha, 1989; Quesada et al., 2019). This belt potentially represents dismembered relics of the Rheic <u>Q</u>ecean and/or a subsidiary seaway that opened during a Variscan transtension event in SW Iberia (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019).

Finally, Paleozoic rocks occur sporadically within the Alpine Betic chain. Their lithological monotony, paucity of fossils, and the intensity of deformation and metamorphism during Alpine orogeny, make recognizing the original features of the different successions challenging (e.g. Martín-Algarra et al., 2019). Some faunal and detrital zircon studies suggest that the Paleozoic outcrops in the Betics may be similar to the most <u>continentalsea</u>ward realms of the Gondwanan platform (i.e., the Cantabrian Zone; e.g. Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018). Following the latest plate reconstructions of the Mediterranean during Meso-Cenozoic times, the Paleozoic units of the Betic-Rif chain may have been located proximal to the present-day position of the Balearic Islands (van Hinsbergen et al., 2020).

The Variscan orogen in Iberia shows multiple deformation, metamorphic, and magmatic events (e.g. Martínez-Catalán et al., 2014; Azor et al., 2019; Fig. 2) that evolved diachronously from the suture towards the external zones (Dalmeyer et al., 1997): (1) An initial continent-continent collision began ca. 370-365 Ma, which produced high_pressure metamorphism (e.g. Lopez-Carmona et al. 2014). (2) Between 360 and 330 Ma a protracted shortening phase occurred, frequently divided into main phases C1 and C2, that were accompanied by Barrovian type metamorphism (e.g. Dias da Silva et al., in pressDias da Silva et al., 2020) and plutonism at ~340 Ma (e.g. Gutiérrez-Alonso et al., 2018). (3) An extensional collapse event, so-called E1, occurred at ~333-317 Ma, which formed core-complexes and granitic domes in the Central Iberian and West Asturian-Leonese zones (Fig. 2C; e.g. Alcock et al., 2009; Díez-Fernández

and Pereira, 2016; López-Moro et al., 2018). This event is coeval and genetically linked to the formation of the foreland fold-and-thrust-belt of the Cantabrian Zone (e.g. Gutiérrez-Alonso, 1996). (4) A late Carboniferous shortening event (C3) occurred ca. 315-290 Ma and is arguedinterpreted to have resulted in the formation of the Cantabrian Orocline and was accompanied by the intrusion of mantle derived granitoids (Fing.Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a, 2011b; Pastor-Galán et al., 2012a). (5) A final early Permian extensional event (E2), mostly found in the Central Iberian Zone, resulted in the formation of core complexes and regional doming (Dias da Silva et al., in pressDias da Silva et al., 2020). (6) A final shortening event (C4), possibly coeval with E2, resulted in widespread brittle deformation (e.g. Azor et al., 2019; Fernández-Lozano et al., 2019).

In SW Iberia, the aforementioned Variscan deformation events are characterized by a dominant sinistral component, which contrasts with the general dextral component recognized in most other regions of the orogen (e.g. Martínez Catalán et al., 2011; Gutiérrez-Alonso et al., 2015). Early collisional structures (C1) formed NE-vergent recumbent folds in the southernmost Central Iberian Zone and SW-vergent folds and thrusts in the Ossa Morena and South Portuguese zones. This phase continued with a transtensional event that heterogeneously extended the continental lithosphere (e.g. Pérez-Cáceres et al. 2015). Coevally, an important extension-related magmatic event happened, perhaps assisted by a plume-type mantle (Simancas et al. 2006) or <u>a-because of</u> slab break-off (Pin et al. 2008). After this transtensional event, significant <u>left-lateralsinistral</u> transpression occurred forming the extensive shear zones to the north and south of <u>the</u>Ossa Morena Zone (Fig. 2B), which accommodated the majority of the transcurrent motion. However, <u>left-lateralsinistral</u> displacements are observed all along the Ossa Morena and South Portuguese zones. Pérez-Cáceres et al. (2016) estimated over 1000 km of collisional convergence in SW Iberia, most of which corresponds with <u>left-lateralsinistral</u> displacements parallel to terrane boundaries.

3 Synthesis on the Geometry and Kinematics of the Cantabrian Orocline

Understanding the geometry, kinematic evolution and mechanics of curved mountain systems is crucial to developing paleogeographic and tectonic reconstructions<u>and</u> <u>understanding past geodynamics</u> (e.g. Marshak, 2004; Van der Voo, 2004; Li et al., 2012; van Hinsbergen et al., 2020). Introduced by Carey (1955 p.257), an orocline (from Greek opoç, mountain, and $\kappa\lambda$ ivo, bend) is "...an orogenic system, which has been flexed in plan to a horseshoe or elbow shape." Although sometimes used in the literature as a geometric description of any orogenic curvature, herein orocline is strictly used as a the term for map-scale bends that underwent vertical-axis rotations (Weil and Sussman, 2004; Johnston et al., 2013; Pastor-Galán et al., 2017a). The kinematic classification of curved mountain belts (Weil and Sussman, 2004; Johnston et al, 2013) distinguishes two end members: (1) Primary orogenic curves, which describe those systems in which curvature is a primary feature of the orogen and formed without significant or systematic vertical-axis rotations, and (2) Secondary oroclines, where orogenic curvature was acquired due to vertical-axis rotations subsequent to primary orogenic building. Those systems whose curvature is the product of vertical-axis rotation during the primary orogenic pulse and/or only a portion of the observed curvature is secondary are termed progressive oroclines.

The orocline test (or strike test), evaluates the relationship between changes in regional structural trend (relative to a reference trend for an orogen) and the orientations of a given geologic fabric element or magnetization (relative to a reference direction). In terms of evaluating developmental kinematics, the most relevant geologic marker is paleomagnetic declination, which can be used to quantitatively evaluate total and systematic rotations as a function of along-strike variability. Once acquired, data is plotted on Cartesian coordinate axes with the strike (S) of the orogen (relative to a reference) along the horizontal axis, and the fabric azimuth (F, relative to a reference) along the vertical axis. The test originally used a basic leastsquares (OLS) regression (Schwartz and Van der Voo, 1983) to estimate the slope (coded m in formulas), ideally between 0 and 1, which then is interpreted with respect to vertical-axis kinematics. More recently, Yonkee and Weil (2010b) and Pastor-Galán et al. (2017a) introduced more robust statistics to estimate the correlations slope and its uncertainty, considering and propagating errors of the input data. Primary orogenic bends show no change of paleomagnetic declination orientations with varying structural trend, and therefore the slope is expected to be 0. In progressive oroclines, the declination variation records some fraction of the total observed orogenic strike variability, and thus the slope would range between 0 and 1, depending on the amount of primary curvature. Secondary oroclines are those in which the paleomagnetic vectors record 100% of the rotation, yielding slopes of 1, meaning that the orogenic system must have started as a roughly linear system that then underwent secondary vertical-axis rotations until its present-day curvature was acquired. The slope obtained with the orocline test can only be confidently interpreted when the chronology of fabric formation is well known.

The trend of the Variscan belt in Iberia follows a sinuous "S" shape that is especially prominent in the northwest region of the Iberian Peninsula, and then becomes more subtle due to the predominance of younger cover sequences in the central and eastern regions of the

peninsula (Fig. 1 and 2). This dramatic geometry has stimulated a century long scientific debate as to its origin (e.g. Suess, 1892; Staub, 1926; Martínez Catalán et al., 2015). To the north and convex to the west is the Cantabrian Orocline, and to the center-south and convex to the east is the Central Iberian curve. The overall trend of the Cantabrian Orocline starts in Brittany (France) and southern England and then curves through the Bay of Biscay and then south into central north Iberia (Fig. 1, 2 and 4). The Cantabrian Orocline (also known as Ibero-Armorican Orocline/ Arc, Asturian Arc or Cantabrian-Asturias Arc) is arguably the first curved orogen that was scientifically described, recognized by the change in structural trend of mapped thrusts and fold axes (Schultz, 1858, Barrois, 1882, Suess, 1892). The Cantabrian Orocline traces an arc with a curvature close to 180° within the central Cantabrian Zone (the Gondwanan foreland in Iberia, fig. 2), and opens to approximately 150° as one moves to the outer arc reaches (Fig. 1). At the crustal-scale, the Cantabrian Orocline represents a first-order vertical-axis buckle fold in planview that refolds pre-existing Variscan structures (e.g. Julivert and Marcos, 1973; Weil et al., 2001). The inner arc of the orocline, or the Cantabrian Zone is characterized by tectonic transport towards the core of the orocline, i.e., the orocline has a contractional core, where low finite strain values and locally developed cleavage occur (Pérez-Estaún et al., 1988; Gutiérrez-Alonso, 1996; Pastor-Galán et al., 2009). Within the inner core a variety of structures record non-coaxial strain, which produced complex interference folds and rotated thrust sheets (e.g. Julivert and Marcos, 1973; Julivert and Arboleya, 1984; Pérez-Estaún et al, 1988; Aller and Gallastegui, 1995: Weil, 2006, 2013; Pastor-Galán et al., 2012b; Shaw et al., 2015; 2016a; Del Greco et al., 2016). In contrast, the outer arc shows a vertical-axis fold with a ca. 150° interlimb angle vertical-axis fold that was accommodated by significant shearing, both dextral and, in lesser amounts, sinistral that was penecontemporaneous to vertical-axis rotation (Gutiérrez-Alonso et al., 2015). Weil et al. (2013, 2019) extensively review the geometry of the Cantabrian Orocline.

All kinematic data studied so far support a model in which the Cantabrian Orocline formed due to secondary vertical-axis rotation in a period of time younger than 315 Ma and older than 290 Ma. Overall, the southern limb of the orocline rotated counterclockwise (CCW) and the northern limb clockwise (CW; Fig. 4). Orocline formation happened subsequent to the main shortening phases of the orogen (C1 and C2) and late-stage orogenic collapse (E1), and therefore, it is an ideal example of a secondary orocline in the strictest sense. Development of the Cantabrian Orocline requires the existence of a roughly linear orogenic belt during early Variscan closure of the Rheic Ocean (with a roughly N-S orientation in present-day coordinates), which was subsequently bent in plan-view into an orocline during late stages of Pangea amalgamation. Such interpretation is grounded in <u>extensive</u> paleomagnetic <u>studies</u> (e.g. Hirt et al., 1992; Parés et al. 1994; Stewart, 1995; van de Voo et al., 1997; Weil, 2006; Weil et al., 2000; 2001; 2010), along with important contributions from structural (e.g. Gutiérrez-Alonso 1992; Kollmeier et al., 2000; Merino-Tomé et al., 2009; Pastor-Galán et al., 2011; 2014; Shaw et al., 2015) and geochronological studies (e.g., Tohver et al., 2008; Gutiérrez-Alonso et al., 2015). Weil et al. (2013) provides<u>d</u> a comprehensive review on the kinematic constraints, updated in 2017a by Pastor-Galán et al., and in 2019 by Weil et al.

4 The intriguing geometry of the Central Iberian curve

The more southern Central Iberian curve has a similar magnitude, but opposite curvature compared to the Cantabrian Orocline (Fig. 1 and 2B). This structure has been referred to as the Central Iberian curve, arc, bend or orocline. In this paper we use 'Central Iberian curve'. The other aforementioned terms involve still unknown parameters or are misleading: e.g., orocline impliesy kinematics (Weil and Sussman, 2004); bend refers to a mechanism of formation (e.g. Fossen, 2016); and arc could be ambiguous, since the term is commonly used for volcanic chains. The Central Iberian curve This curvature was first described by Staub (1926) and was termed the Castilian bend. Continental drift pioneers paid some attention to Staub's description (e.g. Holmes, 1929; Du Toit, 1937), but the curved structure remained largely ignored for multiple decades (e.g. Martínez Catalán et al., 2015). The hypothesis of a large-scale curvature in Central Iberia made a comeback at the beginning of the 21st century with a study of Variscan porphyroblast kinematics across Iberia by Aerden in (2004). Since then, several attempts to unveil its geometry and kinematics have been madegiven with contrasting results.

The elusive nature of the Central Iberian curve resides in the poor exposure of its putative hinge (Fig. 2). The hinge of the Cantabrian orocline crops out extensively and the changes in thrust and fold axes trend are observable at high-resolution from aerial photographs and are readily mapped using outcrop-scale observations. In contrast, the alleged hinge of the Central Iberian curve is largely covered by Mesozoic and Cenozoic basins (Fig. 2). The curvature is most recognizable at the boundary between the Galicia-Tras os Montes and Central Iberian zones (Fig. 2A; Aereden, 2004; Martínez Catalán, 2012). The thrust fault that bounds those areas traces close to a 180° of curvature and marks the emplacement of the most distal units. Before the revival of Staub's curved geometry along the entire Central Iberian Zone, there were several attempts to explain the curved shape of the Galicia Tras-os-Montes Zone. Some consider the Galicia Tras-os-Montes Zone a block that escaped during an early Variscan (C1)

non-cylindrical collision, forming a extrusion wedge towards the areas undergoing that underwent lesser amount of shortening (Martínez-Catalán, 1990, Dias da Silva, 2015; inpress2020); or alternatively a klippe of a larger allochthonous thrust sheet, or the product of an interference pattern between C2, E1 and C3 structures (e.g. Ries and Shackleton, 1971; Martínez Catalán et al., 2002; Rubio Pascual et al., 2013; Díez-Fernández et al., 2015).

In addition to the Galicia Tras-os-Montes Zone, the other areas that showing a certain degree of curvature are to the E and SE of the Central Iberian Zone. There, an approximately 20° change in strike of the Iberian ranges (NE Iberia, Fig. 2A) is observed, which represents the only known outcrop of the hinge of the Central Iberian curve's outer arc. The rest of the curvature has been deducedimplied with indirect observations leading to three competing geometric proposals for the Central Iberian curve (Fig. 2B). The main arguments used to constrain the geometry of the Central Iberian curve are: (1) the geometry of Galicia Tras-os-Montes folds and the orientation of observed garnet inclusion trails (Aerden, 2004; Fig. 2B-1); (2) the alignment ofed aeromagnetic anomalies and fold trends in the Iberian ranges and the E-SE Central Iberian Zone (Martínez-Catalán, 2012; Fig. 2A and 2B-2) and; (3) the regional distribution of paleocurrents recorded in Ordovician guartzites (Shaw et al., 2012; Fig. 2B-3 and 3). All proposed geometries share two features: (1) The curvature runs parallel to the Central Iberian Zone, and is located in the center-west of Iberia, and (2) all place the Galicia Tras-os-Montes Zone in the core of the curve with the curveds axial traces cross-cutting the Morais Complex, which is a set of mafic and ultramafic rocks that is roughly circular in shape (Fig. 2B; Dias da Silva et al., in pressDias da Silva et al., 2020).

Aerden (2004) compared the orientation of inclusions in metamorphic porphyroblasts across the Variscan allochthonous terranes of the NW Iberian Massif, and found that inclusion trails maintain a constant north–south orientation. Comparing such results with the trend of the Variscan fold axes in the <u>C</u>eentral Iberian Zone (Fig. 2A) and a <u>daring-novel</u> interpretation of the aeromagnetic anomalies of the Iberian Peninsula (Fig. 5A), Aerden suggested a geometry in which the Central Iberian curve was more prominent in the outer arc than in the inner arc (Fig. 2B-1). In Aerden's view the geometry of the Galicia Tras-os Montes Zone does not represent a large-scale curvature, but rather the original shape of the nappe, perhaps re-tightened during C3 deformation. In contrast, the Iberian Ranges and the SE Central Iberian Zone represent the more curved sector (Fig. 2B-1). In the model of Aerden (2004), the Ossa Morena and South Portuguese <u>z</u>Zones are not part of the Central Iberian curvature.

Martínez-Catalán (2012) reinterpreted Aerden's analysis of aeromagnetic map data (Fig. 5A) and the<u>ir</u> interpretied ve structural trends of C1-C2 fold axes from Central Iberian Zone

structures (Fig. 2A). In Martínez-Catalán's model, the Central Iberian curvature is a symmetric arcuate shape in which orogen trend changes equally in the inner and outer arc, and is comparable in size to the Cantabrian Orocline, but with opposite curvature and less shortening. This geometric model also excludes the Ossa Morena and the South Portuguese Zones as elements involved in the formation of the curvature (Fig. 2B-2).

Finally, Shaw et al. (2012) studied the orientation of paleocurrents in Ordovician Armorican Quartzite (e.g. Aramburu, 2002), which is one of the most prominent rocks exposed in Iberia (Fig. 3). The<u>se</u> authors found that paleocurrents fanned outward with respect to the Cantabrian Orocline curve and are approximately perpendicular to the structural trend throughout the peninsula (Fig. 3). Shaw et al. (2012) assumed that the direction and sense of paleocurrents were <u>originally</u> parallel throughout all zones, and concluded that the Central Iberia curve is <u>part of</u> a 'S' shape isoclinal structure <u>with a</u> similar in-magnitude <u>of curvature</u> to the Cantabrian Orocline (Fig. 2B-3). It is unclear from the Shaw et al. (2012) model which iswhat the involvement of the Ossa Morena and South Portuguese <u>z</u>eones in the overall curve <u>is</u> (if any), nor the prospective location of the external zones of the orogen (Cantabrian Zone) with respect to the overall curvature.

5 Move over once, move over twice: Kinematic constraints

Late Variscan kinematic data (315-290 Ma; C3, E2, C4 phases) in the Central Iberian curve were scarce prior to revival of Staub's Central Iberian curve (e.g. Vergés, 1983; Julivert et al., 1983; Parés and van der Voo, 1992). More recently, a wealth of studies have been published on the kinematics of forming the Central Iberian curve (Fig. 2B), which are reviewed below.

5.1 Structural Geology and Geochronology

Curved orogens that result from differential vertical-axis rotations develop remarkable structures within their hinges where compressive and extensive radial structures often develop in combination with tangential shear structures (e.g. Li et al., 2012; Eichelberger and McQuarrie, 2015). With the re-emergence of the Central Iberian curve debate, several studies have re-evaluated the well-documented structures from the Central Iberian Zone to constrain the origin and kinematics of curvature. The majority of studies focused on the hinge zone of the curve in the area surrounding Galicia Tras-os-Montes (e.g. Dias da Silva et al., 2014; Jacques et al., 2018a), but some <u>do</u> explored <u>areas in the more</u> outer-arc <u>areas</u> (e.g. Palero-Fernández et al., 2015; Gutiérrez-Alonso et al., 2015). The following <u>paragraphs section</u> synthesizes the findings

of new field, structural, and geochronological analyses from around the hinge of the Central Iberian curve and its surrounding regions. The reviewed studies identify several deformation events that are linked to regional Variscan deformation phases (Fig. 2A).

- An early generation of upright to overturned cylindrical folds with an associated axial planar cleavage (C1). The C1 fold axes plunge variably from horizontal to nearly vertical (e.g. Jacques et al., 2018a, 2018b). The original trend of the fold axes wereas parallel to the orogen (e.g. Pastor-Galán et al., 2019b), however interference with younger deformation events has created complicated geometries (e.g. Díez Fernández et al., 2013; Palero-Fernández et al., 2015). The emplacement of the allochthonous units of the Galicia Tras-os-Montes Zzone (commonly referred as C2) is closely associated with development of C1 folds, but is restricted to shear zones located along the boundary between the latter and the Central Iberian Zone. This phase includes orogen-parallel emplacement of the allochthonous Galicia Tras-Os Montes units and its associated thrusts (Fig. 2A). The non-coaxial nature of the emplacement of this allochthonous nappe produced folding interference and local vertical-axis rotations (Dias da Silva et al., 2020). Prograde Barrovian metamorphism (known as M1) reached its pressure peak at the end of C2 (Rubio Pascual et al., 2013).
- 2. After C1 and C2, the resulting thickened crust gravitationally collapsed (Macaya et al., 1991; Escuder Viruete et al., 1994; Díaz-Balda et al., 1995; Díez-Montes, 2010). This gravitational collapse (phase E1) formed gneiss-dome core complexes between 330 and 317 Ma (e.g. Díez Fernández and Pereira, 2016) especially at in the core of the Central Iberian curve (Fig. 2C; e.g. Martínez-Catalán, 2012). This phase formed large subhorizontal extensional detachments that exhumed to depths of the middle crust (e.g. Rubio-Pascual et al., 2013; Dias da Silva et al., in pressDias da Silva et al., 2020). General decompression produced a Buchan-type metamorphic event (M2; e.g. Rubio-Pascual et al., 2016, Solis-Alulima et al., 2019) and widespread anatectic melting (e.g. López-Moro et al., 2018; Pereira et al., 2018). The E1 phase developed a fold system with sub-horizontal axes and a penetrative subhorizontal cleavage (e.g. Dias da Silva etal., in pressDias da Silva et al., 2020). Mapped folding geometries indicate the deflection of C1 folds into overturned positions within the E1 deformation zones (e.g. Díez Fernández et al., 2013; Díez Fernández and Pereira, 2016; Pastor-Galán et al., 2019b). In addition to large-scale extensional deformation and Buchan metamorphism, E1 developed a regional dome-and-basin pattern, resulting in portions of the allochthonous terranes tectonically transported into basins (e.g. Días da Silva et al., in press2020).

- The structures developed during C1-C2 compression and E1 extension, are re-folded by a younger shortening phase (C3; syn-Cantabrian Orocline). C3 formed upright open folds and conjugate sub-vertical shear zones (e.g. Gutiérrez-Alonso et al., 2015; Díez Fernández and Pereira, 2017; Dias da Silva et al., in pressDias da Silva et al., 2020). C3 was coeval with regional retrograde metamorphism (M3) and with intrusion of mantle derived granitoids (Fing.Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a), surrounded by contact metamorphic aureoles (e.g. Yenes et al., 1999). The age of the C3 event ranges from 315 and 290 (e.g. Jacques et al., 2018a), and is concomitant with the formation of the Cantabrian Orocline (e.g. Pastor-Galán et al., 2015a). Ductile deformation, including folding with axial planar cleavage (e.g. Dias da Silva et al., 2014; Pastor-Galán et al., 2019b) and the development of conjugate shear zones, occurred at the early stages of C3 (315-305 Ma; Gutiérrez-Alonso et al., 2015; Díez-Fernández and Pereira, 2017; Jacques et al., 2018b) and was followed by brittle deformation that formed cross-joint sets and vein swarms with Sn-W mineralizations (Jacques et al., 2018a; 2018b). The conjugated shear zones, some of them with hundreds of kilometers of displacement, had activity during the period 315-305 Ma based on direct Ar-Ar dating of the shear zones (Gutiérrez-Alonso et al., 2015) and cross-cutting relationships with precisely dated igneous rocks (Díez-Fernández and Pereira, 2017). Note that these shear zones show a younger age with respect to the sinistral shear zones that bound the Ossa Morena and South Portuguese zones (340-330 Ma; e.g. Dallmeyer et al., 1993). New studies in the Central Iberian Zone have determined that several folds, previously interpreted as C1 (e.g. the Tamames-Marofa-Sátão synform) are in fact C3 structures, possibly nucleated within existing C1-C2 structures (e.g. Dias da Silva et al., 2017; Jacques et al., 2018b). The remarkable continuity along the Central Iberian Zone of these folds (Fig. 2A), previously interpreted as C1 (e.g. Díez-Balda et al., 1990; Abalos et al., 2002; Dias and Ribeiro, 1994; Dias et al., 2016), suggests the ubiguity and importance of this deformation phase.
- 4. The N-S shortening (in present day coordinates) of C3 deformation continued through the Early Permian under brittle conditions (so-called C4 event) (e.g. Dias da Silva et al., in pressDias da Silva et al., 2020) and overlapped with the formation of E2 extensional faults (Fig. 2A; Dias and Ribeiro 1991; Dias et al. 2003; Rubio Pascual et al., 2013; Arango et al., 2013; Fernández-Lozano et al. 2019; Dias da Silva et al., in pressDias da Silva et al., 2020). C4 N-S compression produced a series of NNE-SSW and NNW-SSE brittle faults (Gil Toja et al. 1985; Dias and Ribeiro 1991; Dias et al. 2003; Fernández-

Lozano et al., 2019) and associated sub-vertical and sub-horizontal widespread kinkbands (e.g. Aller et al., 2020; Dias da Silva et al., in pressDias da Silva et al., 2020) that are today exposed in Northern Iberia. E2 developed core complex-like structures with extensional shear zones that further telescoped M2 metamorphic isograds between the anatectic cores of gneiss domes and the associated hanging wall units. This event favored sub-horizontal folding, and crenulation cleavage development in the footwall together with kink-band generation in the upper low-grade structural levels.

5.2 Paleomagnetism

Paleomagnetism investigates the record of the Earth's ancient magnetic field as it is recorded in the rocks-record. Among other features, rocks can record the orientation of the magnetic field at the time of magnetization (e.g. Tauxe, 2010). The recorded magnetic vector can be geometrically defined by two components: inclination, which is a function of the paleolatitude (being 90° at the poles and 0° at the equator) at the time of magnetization acquisition; and declination, which is a measure of the horizontal angular difference between the recorded magnetic direction and true north, thereby allowing for the quantification of any vertical-axis rotations if a north reference direction paleomagnetic pole is known for the region of interest at the time of magnetization acquisition. Paleomagnetism is the best tool to quantify vertical-axis rotations in orogens due to the independence of the magnetic field from thean orogen's deformation and evolution (e.g. Butler, 1998).

Despite its uniqueness to study paleolatitudes and vertical-axis rotations, paleomagnetism is not flawless. Paleomagnetic data can yield spurious rotations when the local and regional structures are not properly <u>studied_defined_and</u> their geometries and kinematic histories not adequately corrected for (e.g. Pueyo et al., 2016). In addition, the age of magnetization acquisition is not necessarily equivalent to the age of the sampled rock. Remagnetizations are ubiquitous, especially in orogens (Weil and van der Voo, 2002; Pueyo et al., 2007; Huang et al., 2017). In remagnetized rocks, the primary magnetization is replaced or overprinted due to a set of geologic processes acting alone or in concert - usually represented by a combination of thermal or chemical reactions (<u>e.g.</u>, Jackson, 1990). Nevertheless, remagnetizations can be useful for interpreting deformation history if the relative timing of the overprint can be established and a well-constrained reference direction for that age is known (e.g. Weil et al., 2001; Izquierdo-Llavall et al. 2015; Calvín et al., 2017).

In addition to knowing the structural geology and the timing of magnetization of the studied rocks, understanding and quantifying local and regional vertical-axis rotations requires a

paleomagnetic reference pole for comparison. Permian and Mesozoic paleomagnetic studies in Iberia indicate that Iberia was a relatively stable plate from at least Guadalupian times (ca. 270 Ma) to the opening of the Bay of Biscay in the Cretaceous (e.g. Gong et al., 2008; Vissers et al., 2016). Weil et al. (2010) calculated the <u>most modernan</u> Early Permian pole for stable Iberia, which will be used herein as a reference for any vertical-axis rotation analysis (hereafter, eP pole or eP component). Weil et al.'s Virtual Geomagnetic Pole (VGP) values are Plat = 43.9; Plong = 203.3 and α_{95} = 5.4 and when transform into paleomagnetic components has a ~0° inclination (equatorial) and declinations that range from 150° to 160° (from NW to SW respectively) depending on where in Iberia you are referencing. In Fig. 6 (red arrows), a compilation of declinations that form part of this composite pole and other eP components found in recent studies are presented.

For the Central Iberian curve, the voluminous paleomagnetic database from the Cantabrian Orocline can be used to partially constrain its kinematics (e.g. Weil et al., 2013). The orocline test for the Cantabrian Orocline (Ffig. 4) quantifies the degree of differential vertical-axis rotation of variously striking Variscan segments-tectonic belts in northern Iberia. If the Central Iberian curve is a product of vertical-axis rotation, paleomagnetic declinations would bend around the Central Iberian curve opposite to that of the Cantabrian Orocline. With a well constrained orocline test, as in the Cantabrian Orocline (Fig. 4), one can use the paleomagnetic strike-test correlation slope to establish expected declinations for any along-strike portion of the orogen (Pastor-Galán et al., 2017b).

Before the resurgence of the Central Iberian curve, the only available pre-Permian paleomagnetic studies to the <u>s</u>South and west of the Cantabrian Zone in the Iberian Massif were <u>focused from on</u> the Beja Gabbroic Massif, Portugal (Perroud et al., 1985) and the Almadén syncline volcanics (Perroud et al., 1991; Pares & Van der Voo, 1992). The <u>Beja area</u> study in the Beja area showed presented varied inclinations and declinations in the gabbros, and complex overprints elsewhere. Perroud et al. (1985) did not consider any structural correction for the results, <u>as they assumeding</u> the gabbro was undeformed. Recently, Dias da Silva et al. (2018) showed that the area underwent intense deformation during the Carboniferous. Therefore, interpretation of this dataset is complicated without knowing the proper structural correction needed to restore the magnetization to its palinspastic orientation.

Several articles with new paleomagnetic studies around the Central Iberian curve have been published since 2015 (Fig. <u>56</u>). In general, these studies have reported a pervasive late Carboniferous (320 to 300 Ma) (re-)magnetization in sedimentary and igneous rocks (e.g. Pastor-Galán et al., 2015a; 2017b; Fernández-Lozano et al. 2016), which is largely

penecontemporaneous to the intrusion of E1 extensional granites (López-Moro et al., 2018) and C3 syn-orocline mantle derived granitoids (Fing.Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a). The following section describes the <u>reported magnetizations</u> from oldest to youngest.

Pastor-Galán et al. (2016) sampled for paleomagnetic analyses both E1 extensional granites (Fig. 2C; ~320 Ma; e.g. López-Moro et al., 2018) from the Tormes and Martinamor domes, and C3 mantle derived granitoids <u>in-from</u> the Central System (Fig. 2C; 305-295 Ma; e.g. Gutiérrez-Alonso et al., 2011a). Both sets of plutons are located around the Galicias Tras-os-Montes hinge of the Central Iberian curve (Fig. 6-5). The authors found an original component in E1 granities supported by a positive reversal test in both domes (Fig. 7). The magnetization has an inclination (Inc.) = 15° (paleolatitude (λ) = -7.6°) and declination (Dec.) = 81° (Fig. 7), which imply a northward movement of 700 km and a ~70° CCW rotation with respect to the C3 granites that showed an eP component (Dec. ~ 150, Inc. ~ 0). Considering the positive reversal test in E1 granites and the significant difference in inclinations with respect to C3 granitoids (eP component), a magnetization age of older than 318 Ma was proposed (pre_Kiaman superchron, 317 Ma - 267 Ma, e.g. Langereis et al., 2010), which was interpreted as a primary magnetization. The 70° CCW Pennsylvanian rotation recorded in rocks from the Central Iberian curve hinge zone is in agreement with the expected rotation of the southern limb of the Cantabrian Orocline (Fig. 4; Weil et al., 2013).

At the putative outer arc of the Central Iberian curve in_{τ} the Iberian Ranges (Fig. 2), paleomagnetic and structural studies of Devonian and Permian rocks (Pastor-Galán et al., 2018) revealed that the eP component from Permian rocks had rotated ~22° CW during the Cenozoic (Fig. 8; cf. Pastor-Galán et al., 2018). The Permian and Mesozoic rocks from the Iberian Ranges show a consistent ~22° CW rotation with respect to the Apparent Polar Wander Path for Iberia (e.g. Pastor-Galán et al. 2018). This rotation likely happened during the Alpine orogeny, in which the northern area of the Iberian Range underwent more shortening than the southern part, resulting in a regional CW vertical-axis rotation (Izquierdo-Llavall et al., 2019). After restoring the Cenozoic rotation, the Devonian rocks show a positive reversal and fold-test with inclinations that are steeper than expected from the eP component (Dec. = 85.3°, Inc. = 12.7°, λ = -6.4). This component is statistically indistinguishable from that of the E1 granites and the southern branch of the Cantabrian Zone, showing the same 70° CCW rotation from the time they were remagnetizationed (estimated in ~318 Ma) to the timing of the eP component (Fig. 8; Pastor-Galán et al., 2018). Once Cenozoic rotation is corrected for, the structural and paleomagnetic trends of the Iberian ranges become parallel to those in the southern limb of the Cantabrian Orocline, ruling out a Variscan or older origin for the outer Central Iberian curve (Fig.

The remaining paleomagnetic works published on Central and SW Iberia rocks all reveal a ubiquitous late Carboniferous to Early Permian remagnetizations during the Kiaman superchron (Fernández-Lozano et al., 2016; Pastor-Galán et al., 2015a; 2016; 2017b; Leite Mendes, in press). The authors of these papers calculated the expected declination for each site as if they were part of the Cantabrian Orocline (Fig. 9A). All localities where magnetizations pre-date the formation of the Cantabrian Orocline show the same expected rotations as the southern limb of the Cantabrian Orocline, regardless of their position within the Central Iberian curve (to the hinge: Tormes and Martinamor domes, Iberian ranges; to the southern limb: Almadén syncline and South Portuguese Zone). Other locations, especially limestones from the Central Iberian Zone, have declinations and inclinations in between the primary 318 Ma component of the E1 granites and the post-orocline eP component (Fig. 9B). Pastor-Galán et al. (2015a; 2016) interpreted these results as being caused by a remagnetization that was acquired during formation of the Cantabrian Orocline formation and therefore recorded intermediate steps between the component of the E1 granites and eP. Those authors suggest that the large amount of syn-orocline mantle derived granitoids that intruded the Central Iberian Zone (C3 granitoids) triggered the hinterland remagnetization.

Finally, two previous studies identified an earlier magnetization in the Almadén syncline region of the SE Central Iberian Zone (Perroud et al., 1991; Pares & Van der Voo, 1992). However, Leite Mendes et al. (in press) argue that these studies are likely misinterpreted. Perroud et al. (1991), applied a complicated structural correction restoring a putative plunge of the regional structural axis to all sites, including those where the syncline axis does not plunge. Leite Mendes et al. (in press) re-sampled the syncline where its axis is sub-horizontal and obtained a negative fold test, implying that the magnetization is not primary as previously interpreted. Their results, however, are similar in orientation to those components published from previous studies prior to any structural correction (Perroud et al., 1991 and Parés and van der Voo, 1992).

Two additional studies sampled Laurussian margin sequences that are today adjacent to the Cantabrian Orocline region (Fig. 10). To the north, the SW <u>area-part</u> of Ireland preserves a Late Paleozoic basin filled with Devonian red sandstone and Carboniferous limestone and siltstone, which was sampled by Pastor-Galán et al. (2015a). To the south <u>is-are</u> the aforementioned results from the South Portuguese Zone (Leite Mendes et al., in press). Both areas are interpreted to have previously been part of the Laurussian continent, <u>and therefore</u> on the opposite side of the Rheic Ocean suture at the time of Variscan collision (Fig. 10; e.g.

8).

Pastor-Galán et al., 2015b). In contrast, the rest of Iberia was part of, or proximal to, Gondwana (e.g. Franke et al., 2017). These Paleomagnetic results from the Laurussian margin suggest that the rotations involved in the formation of the Cantabrian Orocline occurred along both sides of the Rheic suture proximal to bothalong its northern and southern limbs (Fig. 10A and B). Pastor-Galán et al. (2015b) hypothesized a so-called Greater Cantabrian Orocline that would have bent the entire Appalachian/Variscan orogen around a vertical-axis, affecting at least the continental margins of both Gondwana and Laurussia.

5.3 The implications of not being a secondary orocline

The most relevant new data regarding the kinematics of the Central Iberian curve is the paleomagnetic study from the Iberian Ranges (Calvín et al., 2014; Pastor-Galán et al., 2018). These results confirm that the present-day variation in trend of the tectonostratigraphic units, generally attributed to Variscan tectonics (e.g. Weil et al., 2013; Shaw et al., 2012; 2014), is likely a product of Cenozoic Alpine orogeny. Izquierdo-Llavall et al. (2019) confirmed that the interpreted Alpine rotations correspond well with the amount of shortening reconstructed in Meso-Cenozoic basins. The best preserved and most continuous outcrop in the Central Iberian's outer arc is not a Variscan structure, casting doubt that the regionanl effects of Alpine deformation are often underestimated, especially close to the major Iberian Alpine fronts: the Pyrenees, Iberian Ranges, and the Betics.

Overall, new paleomagnetic data from the Central Iberian curve and nearby areas reveal pervasive late Carboniferous remagnetizations and regional vertical-axis rotations of the same sense and magnitude to those expected for the southern arm of the Cantabrian Orocline. The new paleomagnetic data indicate that a post ~320 Ma formation for the Central Iberia curve due to vertical-axis rotations is not supported (Pastor-Galán et al., 2016). The distribution in space and time of paleomagnetic results discards-makes it unlikely that the formation of the Central Iberian curve jas a product of Variscan gravitational collapse (E1, ~330-317 Ma) or concomitant to the Cantabrian Orocline (C3). So far, no pre-E2 paleomagnetic component has been found, and consequently, paleomagnetic data cannot reject an early orogenic origin for the inner arc of the Central Iberian curve (C1-C2, older than 330 Ma).

From a structural geology point of view, the Central Iberian curve does not display the classic geometries and structural interference patterns as found in other established oroclines (i.e., those systems that involve differential vertical-axis rotations, e.g. Li et al., 2012; van der Boon et al., 2018; Meijers et al., 2017; Rezaeian et al., <u>in press2020</u>). The geometry and

structural behaviour of oroclines should resemble, at the crustal-scale, a regional vertical-axis fold preserved in plan-view, either formed by buckling (e.g. Johnston et al., 2001) or bending (e.g. Cifelli et al., 2008) mechanisms. In oroclines, pre-existing structures tend to follow fold trends around the curvature (e.g. Rosenbaum, 2014; Li et al., 2018). In addition, orocline cores tend to preserve radial structures and shortening patterns in the inner arc and orocline parallel shear zones and extensional structures in their outer arc (e.g. Ries and Shackleton, 1976; Eichelberger and McQuarrie, 2015), similar to what is observed in multilayer folds (e.g. Fossen, 2016).

The structural geometry of the Central Iberian curve lacks such patterns. Paleomagnetism from the Iberian Ranges indicate that the Cantabrian and West Asturian Leonese zones do not follow the Central Iberian curve, instead they continue their NWNW-SESE trend into the Mediterranean in what it is now the Betic chain (Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018; van Hinsbergen et al., 2020). Structural trends in the Ossa Morena and the South Portuguese Zone do not show any change in along-strike structural trend that supports large-scale CW rotations (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019), whereas existing paleomagnetic data from those zones (Leite Mendes et al., in press) support a model of CCW rotation associated with the broader southern arm of the Cantabrian Orocline. In the Central Iberian and Galicia Tras-os-Montes zones, the trend of curvature is irregular (see C1 fold patterns in Fig. 2A) and nowhere are the expected inner and outer arc-related structures preserved (e.g. Dias da Silva et al. in press).

The curved shape of C1 fold axes in the Central Iberian Zzone is better explained by fold interference patterns than vertical-axis rotations (e.g. Pastor-Galán et al., 2019b). Moreover, the curved shape of the Galicia Tras-os-Montes allochthonous nappe, which was emplaced orogen_parallel, shows no evidence of vertical-axis rotation related structures (Fig. 2A; Dias da Silva et al., in pressDias da Silva et al., 2020). Other authors describe the changes in trend around the Central Iberian curve expressed by C1 folds (Fig. 2A) as the product of fold interference patterns (e.g. Gutiérrez-Alonso, 2009; Palero-Fernández et al., 2015; Jacques et al., 2018b; Dias da Silva et al., in pressDias da Silva et al., 2020). Pastor-Galán et al. (2019b) showed that curved C1 folds in the Central Iberian Zone around the Galicia Tras-os-Montes boundary (Fig. 2A) are coaxial with C3 folds after restoring the effects of C2 and E1 deformation phases (Fig. 11A). Both C1 and C3 structures formed under similar shortening directions. In the same area, Jacques et al. (2018b) found similar fold interference patterns, in addition they described kinematic incompatibility with the expected CW rotations that would have occurred if the Central Iberian curve waswere an orocline. In other areas of the Central Iberian Zone, the curved shape

of C1 folds has been described as an interference between C1 structures and their reorientation caused by C3 shear zones (Fig. 2A; e.g. Palero-Fernández et al., 2015; Dias et al., 2016), or alternatively the interference between C1, C3 and the E2 structures (Fig. 2A; Gutiérrez-Alonso, 2009; Arango et al., 2013; Rubio Pascual et al. 2013).

Overall, new geometric and kinematic data favor the interpretation that the Central Iberian curve is not a structure formed by differential vertical-axis rotation as was the Cantabrian Orocline, but one formed as a consequence of several competing processes. It is clear from the current data that a combination of several deformation events caused the orientation of structures that today delineate the shape of the Central Iberia curve. These include: (1) the northern part of the outer-arc <u>a</u> is the product of an Alpine rigid block rotation instead of Variscan differential vertical-axis rotation (Pastor-Galán et al., 2018); (2) the curvature of the Galicia Trasos-Montes allochthonous nappe reflects its original shape and could be defined as a primary curve (see Weil and Sussman, 2004), since it was emplaced orogen parallel and shows-preserves no sign evidence of vertical-axis rotations at any time (fig. 2A; Dias da Silva et al., in-pressDias da Silva et al., 2020); (3) Structural analysis shows that fold interference patterns explain the geometry of the curved trends of Central Iberian Zone's C1 folds (Pastor-Galán et al., 2019b), whose kinematics are incompatible with the required CW rotations expected if the curve is an orocline (Jacques et al, 2018b).

6 Get Back: Ideas flowing out and endless questions

The pioneering works in the last decade that resurrected the idea of a Central Iberian curve, speculated that both the Cantabrian and Central Iberian zones buckled together as secondary oroclines (Fig. 12Fig. 11; Martínez-Catalán 2011; Shaw et al., 2012, 2014; Shaw and Johnston, 2016; Carreras and Druguet, 2014). Later, Martínez Catalan et al. (2014) and Díez Fernández and Pereira (2017) reformulated Martínez-Catalán's 2011 hypothesis and proposed that the Central Iberian curve formed as an orocline between 315 and 305 Ma, and assigning the Cantabrian Orocline a time frame between 305 and 295 Ma (Fig. 12Fig. 11). The proposed tectonic mechanisms to support these early kinematic models are varied: (1) buckling of a ribbon 'Armorican' continent (Fig. 12Fig. 11A; Shaw et al., 2014; 2016); (2) buckling of a completely formed Variscan orogen during a putative 'Pangea B' to 'Pangea A' transition in the late Carboniferous (Fig. 12Fig. 11B; Carreras and Druguet, 2014; Martínez-Catalán et al., 2011); (3) indentation of Laurussia into Gondwana during the early stages of collision (at present day SW Iberia, South Portuguese Zone), producing first the Central Iberian curve as a mega drag-fold during Carboniferous times and then slightly later the Cantabrian Orocline as a

consequence of an indentation process (fig. 12Fig. 11C; Simancas et al., 2013).

The reviewed data in sections 4 and 5 contradict the aforementioned hypotheses. Paleomagnetism and structural patterns (section 5; Fig. 6-11) disagree with the necessary CW rotations required to support a late Carboniferous orocline origin for the Central Iberian curve (Models in Fig. 12Fig. 11A and B). In addition, the sense and magnitude of the vertical-axis rotations observed in SW Iberia (Fig. 10) imply that the South Portuguese (Avalonian segment) and Ossa Morena zones moved together with the southern limb of the Cantabrian Orocline during the Pennsylvanian and Early Permian. This means that the South Portuguese Zone was already parallel to the general trend of the Variscan orogen prior to Cantabrian Orocline formation, implying the lack of a Laurussian rigid indenter into Gondwana (e.g. Simancas et al., 2013). This discrepancy leaves orogen-parallel terrane transport as a possible explanation to the kinematics observed in the Ossa Morena and South-Portuguese Zones (e.g. Pérez-Cáceres et al., 2016). At the same time, paleomagnetism from SW Iberia backs-supports the hypothesis of a Greater Cantabrian Orocline that extended into both Gondwana and Laurussia in its northern and southern limbs (Fig. 10; Pastor-Galán et al., 2015b).

In spite of the kinematic constraints and structural patterns <u>that</u>, <u>which</u>_do not support a vertical-axis origin for the Central Iberian curve in Late Carboniferous time, <u>other there are</u> geometric constraints <u>that</u> remain challenging to account for. For example, tThe curved shape of the aeromagnetic and gravity anomalies of Iberia are real (Fig. 5). These striking patterns could be due to Variscan-Alpine structural interference, for example the previous exampleas suggested for from the Iberian Ranges, but currently there is not enough data to rigorously test this hypothesis. In addition are tThe curved traces of C1 fold-axes, whose geometry and kinematics are reasonably well constrained around Galicia Tras-os-Montes (e.g. Dias da Silva et al., 2020), but in many other areas their showing strong curvature remain largely unstudied (Fig. 2) and therefore, to date we can only especulate on their origin.

-Shaw et al. (2012) supported their hypothesis of a secondary orocline by assuming that paleocurrents were parallel through Iberia during Ordovician times. However, some of the observed deflections in the paleocurrents studied by Shaw et al. (2012; see Fig. 3) are also explained by Alpine vertical-axis rotations (the case of the Iberian ranges) and fold interference patterns (SE of the Central Iberian Zone). Others (Central and SW of the Central Iberian Zone) may be explained by a local response to basin architecture (Fig. 3), where paleo-flow directions would trend toward the deepest basin throughs. The Ordovician basin architecture of Iberia allows for opposite directed paleocurrents from both sides of such throughs (Fig. 3). However, the Early Paleozoic basin architecture in Iberia and their local deformation events require further

research (Sánchez-García et al., 2019).

Einally, aAlthough kinematic evidence is still scarce for the earliest pre-Variscan and early Variscan eventsmovemen, we argue that pre-orogenic physiographic features, such as the opening of a marginal restricted ocean between Gondwana and its distal platform at 395 Ma (Fig. 13Fig. 12A; Pin et al., 2002; Gutiérrez-Alonso et al., 2008b; Arenas et al., 2016) explains the rounded shape of the Galicia tTras-os-Montes curve as a primary arecurve. During thecollision, the latter irregularity would cause the orogen-parallel emplacement of allochthonous nappes (Fig. 13Fig. 12B; Dias da Silva et al., in pressDias da Silva et al., 2020) and the leftlateralsinistral movements of the Ossa Morena and South Portuguese zZones in SW Iberia (Fig 13A, B, C; Quesada, 2019). During the late Carboniferous, possibly due to a plate reorganization during the final amalgamation of Pangea (Fig. 13Fig. 12D; e.g. Gutiérrez-Alonso et al., 2008a; Pastor-Galán et al., 2015a), the far-field stress-field likely changed, which causedand buckled_the entire orogen to buckle around a vertical axis (Gutiérrez-Alonso et al., 2004), including both the Gondwana and Laurussia margins (Fig. 13Fig. 12E; Pastor-Galán et al., 2015b).

Acknowledgements

DPG thanks the extraordinary hard work, patience and endurance of the Utrecht University students that embraced and enjoyed studying the kinematics of the <u>Ceentral Iberian</u> curvature: Thomas Groenewegen, Bart Ursem, Daniel Brower, Mark Diederen and Bruno Leite-Mendes. DPG acknowledges FRIS and CNEAS for the continous financial support. GGA is supported by Spanish Ministry of Science, innovation and universities under the project IBERCRUST (PGC2018-096534-B-I00) and Russian Federation Government grant no. 14.Y26.31.0012. This paper is a contribution to the IGCP no. 648 "Supercontinent Cycles and Global Geodynamics". 50 years ago four fabulous guys let it be and never got back.

References

Ábalos, B., Carreras, J., Druguet, E., Escuder Viruete, J., Gómez Pugnaire, M. T., Lorenzo Álvarez, S., Quesada, C., Rodríguez Fernández, L. R. and Gil-Ibarguchi, J. I.: Variscan and pre-Variscan tectonics, Geol. Spain, 155–183, 2002.

Aerden, D.: Correlating deformation in Variscan NW-Iberia using porphyroblasts; implications for the Ibero-Armorican Arc, J. Struct. Geol., 26(1), 177–196, 2004.

Alcock, J. E., Catalán, J. R. M., Arenas, R. and Montes, A. D.: Use of thermal modeling to

assess the tectono-metamorphic history of the Lugo and Sanabria gneiss domes, Northwest Iberia, Bull. Société Géologique Fr., 180(3), 179–197, doi:10.2113/gssgfbull.180.3.179, 2009.

Aller, J., Bastida, F. and Bobillo-Ares, N. C.: On the development of kink-bands: A case study in the West Asturian-Leonese Zone (Variscan belt, NW Spain) / Sur le développement des kinkbands : un exemple dans le Zone Astur Occidentale-léonaise (chaîne varisque ibérique, nordouest de l'Espagne), Bull. Société Géologique Fr., 191(1), doi:10.1051/bsgf/2020003, 2020.

Aller, J. J. J. and Gallastegui, J. J.: Analysis of kilometric-scale superposed folding in the Central Coal Basin (Cantabrian zone, NW Spain), J. Struct. Geol., 17(7), 961–969, doi:10.1016/0191-8141(94)00115-G, 1995.

Alonso, J. L., Marcos, A. and Suárez, A.: Paleogeographic inversion resulting from large out of sequence breaching thrusts: The León Fault (Cantabrian zone, NW Iberia). A new picture of the external Variscan thrust belt in the Ibero-Armorican arc, Geol. Acta, 7(4), 451–473, doi:10.1344/105.000001449, 2009.

Álvaro, J. J., Casas, J. M., Clausen, S. and Quesada, C.: Early Palaeozoic geodynamics in NW Gondwana, J. Iber. Geol., 44(4), 551–565, doi:10.1007/s41513-018-0079-x, 2018.

Ardizone, Juan, Julio Mezcua, and Isabel Socías. Mapa aeromagnético de España peninsular. Instituto Geográfico Nacional, 1989.

Mergl,M. & Zamora,S. 2012, New and revised occurrences of... Bulletin of Geosciences, 87, 571-586., [online] Available from: http://www.geology.cz/bulletin/contents/art1327 (Accessed 6 April 2020b), n.d.

Aramburu, C., Méndez-Bedia, I. and Arbizu, M.: The Lower Palaeozoic in the Cantabrian Zone (Cantabrian Mountains, NW Spain)., edited by S. García-López and F. Bastida, pp. 35–49., 2002.

Arango, C., Díez Fernández, R. and Arenas, R.: Large-scale flat-lying isoclinal folding in extending lithosphere: Santa María de la Alameda dome (Central Iberian Massif, Spain), Lithosphere, 5(5), 483–500, doi:10.1130/L270.1, 2013.

Arenas, R., Sánchez Martínez, S., Díez Fernández, R., Gerdes, A., Abati, J., Fernández-Suárez, J., Andonaegui, P., González Cuadra, P., López Carmona, A., Albert, R., Fuenlabrada, J. M. and Rubio Pascual, F. J.: Allochthonous terranes involved in the Variscan suture of NW Iberia: A review of their origin and tectonothermal evolution, Earth-Sci. Rev., 161, 140–178, doi:10.1016/j.earscirev.2016.08.010, 2016.

Ayala, C., Bohoyo, F., Maestro, A., Reguera, M. I., Torne, M., Rubio, F., Fernàndez, M. and García-Lobón, J. L.: Updated Bouguer anomalies of the Iberian Peninsula: a new perspective to interpret the regional geology, J. Maps, 12(5), 1089–1092, doi:10.1080/17445647.2015.1126538, 2016.

Ayarza, P., Catalan, J. R. M., Gallart, J., Pulgar, J. A. and Danobeitia, J. J.: Estudio Sismico de la Corteza Iberica Norte 3.3: A seismic image of the Variscan crust in the hinterland of the NW Iberian Massif, Tectonics, 17(2), 171--+, 1998.

Azor, A., Rubatto, D., Simancas, J. F., Lodeiro, F. G., Poyatos, D. M., Parra, L. M. M. and

Matas, J.: Rheic Ocean ophiolitic remnants in Southern Iberia questioned by SHRIMP U-Pb zircon ages on the Beja-Acebuches amphibolites - art. no. TC5006, Tectonics, 27(5), C5006–C5006, 2008.

Azor, A., Dias da Silva, Í., Gómez Barreiro, J., González-Clavijo, E., Martínez Catalán, J. R., Simancas, J. F., Martínez Poyatos, D., Pérez-Cáceres, I., González Lodeiro, F., Expósito, I., Casas, J. M., Clariana, P., García-Sansegundo, J. and Margalef, A.: Deformation and Structure, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 307–348, Springer International Publishing, Cham., 2019.

Barrois, C. E.: Recherches sur le terrains anciens des Asturies et de la Galice, Six-Horemans., 1882.

Bastida, F.: Zona Cantábrica, in Geología de España, edited by J. A. Vera, pp. 25–49, SGE-IGME, Madrid., 2004.

Bayona, G., Thomas, W. A. and Van der Voo, R.: Kinematics of thrust sheets within transverse zones: A structural and paleomagnetic investigation in the Appalachian thrust belt of Georgia and Alabama, J. Struct. Geol., 25(8), 1193–1212, doi:10.1016/S0191-8141(02)00162-1, 2003.

van der Boon, A., van Hinsbergen, D. J. J. J. J., Rezaeian, M., Gürer, D., Honarmand, M., Pastor-Galán, D., Krijgsman, W. and Langereis, C. G. G.: Quantifying Arabia–Eurasia convergence accommodated in the Greater Caucasus by paleomagnetic reconstruction, Earth Planet. Sci. Lett., 482, doi:10.1016/j.epsl.2017.11.025, 2018.

Buiter, S. J. H. and Torsvik, T. H.: A review of Wilson Cycle plate margins: A role for mantle plumes in continental break-up along sutures?, Gondwana Res., 26(2), 627–653, doi:10.1016/j.gr.2014.02.007, 2014.

Butler, R.: Paleomagnetism: Magnetic domains to geologic terranes, Electron. Ed., (September), 319–319, doi:10.1006/icar.2001.6754, 1998.

Calvin, P., Casas, A. M., Villalaín, J. J., Tierz, P., Calvín, P., Casas, A. M., Villalaín, J. J., Tierz, P., Calvin, P., Casas, A. M., Villalaín, J. J. and Tierz, P.: Reverse magnetic anomaly controlled by Permian Igneous rocks in the Iberian Chain (N Spain), Geol. Acta, 12(3), 193–207, doi:10.1344/GeologicaActa2014.12.3.2, 2014.

Calvín, P., Casas-Sainz, A. M. M., Villalaín, J. J. J. and Moussaid, B.: Diachronous folding and cleavage in an intraplate setting (Central High Atlas, Morocco) determined through the study of remagnetizations, J. Struct. Geol., 97, 144–160, doi:10.1016/j.jsg.2017.02.009, 2017.

Calvín-Ballester, P. and Casas, A.: Folded Variscan thrusts in the Herrera Unit of the Iberian Range (NE Spain), Geol. Soc. Lond. Spec. Publ., 394(1), 39–52, doi:10.1144/SP394.3, 2014.

Carey, S. W.: The orocline concept in geotectonics-Part I, Pap. Proc. R. Soc. Tasman., 89, 255–288, 1955.

Carreras, J. and Druguet, E.: Framing the tectonic regime of the NE Iberian Variscan segment, Geol. Soc. Lond. Spec. Publ., 405(1), 249–264, doi:10.1144/SP405.7, 2014.

Martínez-Catalán, J. R. M., Fernández-Suárez, J., Jenner, G. A., Belousova, E. and Montes, A.:

Provenance constraints from detrital zircon U–Pb ages in the NW Iberian Massif: implications for Palaeozoic plate configuration and Variscan evolution, J. Geol. Soc., 161(3), 463–476, doi:10.1144/0016-764903-054, 2004.

Chopin, F., Schulmann, K., Skrzypek, E., Lehmann, J., Dujardin, J. R., Martelat, J. E., Lexa, O., Corsini, M., Edel, J. B., Štípská, P. and Pitra, P.: Crustal influx, indentation, ductile thinning and gravity redistribution in a continental wedge: Building a Moldanubian mantled gneiss dome with underthrust Saxothuringian material (European Variscan belt), Tectonics, 31(1), n/a-n/a, doi:10.1029/2011TC002951, 2012.

Chopin, F., Corsini, M., Schulmann, K., El Houicha, M., Ghienne, J.-F. and Edel, J.-B.: Tectonic evolution of the Rehamna metamorphic dome (Morocco) in the context of the Alleghanian-Variscan orogeny, Tectonics, 33(6), 1154–1177, doi:10.1002/2014TC003539, 2014.

Cifelli, F., Mattei, M. and Della Seta, M.: Calabrian Arc oroclinal bending: The role of subduction, Tectonics, 27(5), doi:10.1029/2008TC002272, 2008.

Dallmeyer, R. D., Fonseca, P. E., Quesada, C. and Ribeiro, A.: 40Ar/39Ar mineral age constraints for the tectonothermal evolution of a Variscan suture in southwest Iberia, Tectonophysics, 222(2), 177–194, doi:10.1016/0040-1951(93)90048-O, 1993.

Dallmeyer, R. D. D., Catalán, J. R. M., Arenas, R., Gil Ibarguchi, J. I. I., Gutiérrez^Alonso, G., Farias, P., Bastida, F. and Aller, J.: Diachronous Variscan tectonothermal activity in the NW Iberian Massif: Evidence from 40Ar/39Ar dating of regional fabrics, Tectonophysics, 277(4), 307–337, doi:10.1016/S0040-1951(97)00035-8, 1997.

Dias da Silva, Í., Gómez-Barreiro, J., Martínez Catalán, J. R., Ayarza, P., Pohl, J. and Martínez, E.: Structural and microstructural analysis of the Retortillo Syncline (Variscan belt, Central Iberia). Implications for the Central Iberian Orocline, Tectonophysics, 717, 99–115, doi:10.1016/j.tecto.2017.07.015, 2017.

Dias da Silva, Í., Pereira, M. F., Silva, J. B. and Gama, C.: Time-space distribution of silicic plutonism in a gneiss dome of the Iberian Variscan Belt: The Évora Massif (Ossa-Morena Zone, Portugal), Tectonophysics, 747–748, 298–317, doi:10.1016/j.tecto.2018.10.015, 2018.

Dias da Silva, Í., González Clavijo, E. and Díez-Montes, A.: The collapse of the Variscan belt: a Variscan lateral extrusion thin-skinned structure in NW Iberia, Int. Geol. Rev., 00(00), 1–37, doi:10.1080/00206814.2020.1719544, in press2020.

Dias da Silva, Í. F., Linnemann, U., Hofmann, M., González-Clavijo, E., Díez-Montes, A. and Catalán, J. R. M.: Detrital zircon and tectonostratigraphy of the Parautochthon under the Morais Complex (NE Portugal): implications for the Variscan accretionary history of the Iberian Massif, J. Geol. Soc., 172(1), 45–61, doi:10.1144/jgs2014-005, 2015.

Dias, R. and Ribeiro, A.: Finite strain analysis in a transpressive regime (Variscan autochthon, northeast Portugal), Tectonophysics, 191(3), 389–397, doi:10.1016/0040-1951(91)90069-5, 1991.

Dias, R. and Ribeiro, A.: CONSTRICTION IN A TRANSPRESSIVE REGIME - AN EXAMPLE IN THE IBERIAN BRANCH OF THE IBERO-ARMORICAN ARC, J. Struct. Geol., 16(11), 1543–1554, 1994.

Dias, R., Mateus, A., and Ribeiro, A., Strain partitioning in transpressive shears zones in the southern branch of the Variscan Ibero-Armorican arc: Geodinamica Acta, v. 16, no. 2–6, p. 119–129. doi:10.1016/j.geoact.2003.04.001, 2003.

Dias, R., Ribeiro, A., Romão, J., Coke, C. and Moreira, N.: A review of the arcuate structures in the Iberian Variscides; constraints and genetic models, Tectonophysics, 681, 170–194, doi:10.1016/j.tecto.2016.04.011, 2016.

Díez Balda, M. A., Vegas, R. and González Lodeiro, F.: Central-Iberian zone structure, Pre-Mesoz. Geol. Iber., 172–188, 1990.

Díez Balda, M. A., Martínez Catalán, J. R. and Ayarza Arribas, P.: Syn-collisional extensional collapse parallel to the orogenic trend in a domain of steep tectonics: the Salamanca Detachment Zone (Central Iberian Zone, Spain), J. Struct. Geol., 17(2), 163–182, doi:10.1016/0191-8141(94)E0042-W, 1995.

Díez Fernández, R. and Arenas, R.: The Late Devonian Variscan suture of the Iberian Massif: A correlation of high-pressure belts in NW and SW Iberia, Tectonophysics, 654, 96–100, doi:10.1016/j.tecto.2015.05.001, 2015.

Díez Fernández, R. and Pereira, M. F.: Extensional orogenic collapse captured by strike-slip tectonics: Constraints from structural geology and UPb geochronology of the Pinhel shear zone (Variscan orogen, Iberian Massif), Tectonophysics, 691, 290–310, doi:10.1016/j.tecto.2016.10.023, 2016.

Díez Fernández, R. and Pereira, M. F.: Strike-slip shear zones of the Iberian Massif: Are they coeval?, Lithosphere, 9(5), 726–744, doi:10.1130/L648.1, 2017.

Díez Fernández, R., Gómez Barreiro, J., Martínez Catalán, J. R. and Ayarza, P.: Crustal thickening and attenuation as revealed by regional fold interference patterns: Ciudad Rodrigo basement area (Salamanca, Spain), J. Struct. Geol., 46, 115–128, doi:10.1016/j.jsg.2012.09.017, 2013.

Díez Fernández, R., Arenas, R., Pereira, M. F., Sánchez-Martínez, S., Albert, R., Martín Parra, L.-M., Rubio Pascual, F.-J. and Matas, J.: Tectonic evolution of Variscan Iberia: Gondwana– Laurussia collision revisited, Earth-Sci. Rev., 162, 269–292, doi:10.1016/j.earscirev.2016.08.002, 2016.

Díez-Montes, A.: La Geología del Dominio "Ollo de Sapo" en las comarcas de Sanabria y Terra do Bolo., 2006.

Díez-Montes, A., Martínez-Catalán, J. R. R. and Bellido-Mulas, F.: Role of the Ollo de Sapo massive felsic volcanism of NW Iberia in the Early Ordovician dynamics of northern Gondwana, Gondwana Res., 17(2–3), 363–376, doi:10.1016/j.gr.2009.09.001, 2010.

Domeier, M. and Torsvik, T. H.: Plate tectonics in the late Paleozoic, Geosci. Front., 5(3), 303–350, doi:10.1016/j.gsf.2014.01.002, 2014.

Edel, J. B., Schulmann, K., Lexa, O. and Lardeaux, J. M.: Late Palaeozoic palaeomagnetic and tectonic constraints for amalgamation of Pangea supercontinent in the European Variscan belt, Earth-Sci. Rev., 177(September 2017), 589–612, doi:10.1016/j.earscirev.2017.12.007, 2018.

Eduard Suess: Das Antlitz der Erde, F. Tempsky; [etc., etc.]. [online] Available from: http://archive.org/details/dasantlitzderer02suesgoog (Accessed 12 March 2020), 1892.

Eichelberger, N. and McQuarrie, N.: Kinematic reconstruction of the Bolivian orocline, Geosphere, 11(2), 445–462, doi:10.1130/GES01064.1, 2015.

Escuder Viruete, J., Arenas, R. and Catalán, J. R. M.: Tectonothermal evolution associated with Variscan crustal extension in the Tormes Gneiss Dome (NW Salamanca, Iberian Massif, Spain), Tectonophysics, 238(1), 117–138, doi:10.1016/0040-1951(94)90052-3, 1994.

Fernández-Lozano, J., Pastor-Galán, D., Gutiérrez-Alonso, G. and Franco, P.: New kinematic constraints on the Cantabrian orocline: A paleomagnetic study from the Peñalba and Truchas synclines, NW Spain, Tectonophysics, 681, 195–208, doi:10.1016/j.tecto.2016.02.019, 2016.

Fernández-Lozano, J., Gutiérrez-Alonso, G., Willingshofer, E., Sokoutis, D., Vicente, G. de and Cloetingh, S.: Shaping of intraplate mountain patterns: The Cantabrian orocline legacy in Alpine Iberia, Lithosphere, 11(5), 708–721, doi:10.1130/L1079.1, 2019.

Fernández-Suárez, J., Gutiérrez-Alonso, G., Pastor-Galán, D., Hofmann, M., Murphy, J. B. and Linnemann, U.: The Ediacaran-Early Cambrian detrital zircon record of NW Iberia: Possible sources and paleogeographic constraints, Int. J. Earth Sci., 103(5), 1335–1357, doi:10.1007/s00531-013-0923-3, 2014.

Fossen, H.: Structural Geology, Cambridge University Press., 2016.

Franke, W., Cocks, L. R. M. and Torsvik, T. H.: The Palaeozoic Variscan oceans revisited, Gondwana Res., 48, 257–284, doi:10.1016/j.gr.2017.03.005, 2017.

Franke, W., Cocks, L. R. M. and Torsvik, T. H.: Detrital zircons and the interpretation of palaeogeography, with the Variscan Orogeny as an example, Geol. Mag., 157(4), 690–694, doi:10.1017/S0016756819000943, 2020.

García-Arias, M., Díez-Montes, A., Villaseca, C. and Blanco-Quintero, I. F.: The Cambro-Ordovician Ollo de Sapo magmatism in the Iberian Massif and its Variscan evolution: A review, Earth-Sci. Rev., 176, 345–372, doi:10.1016/j.earscirev.2017.11.004, 2018.

<u>Gil Toja, A., Jimenez-Ontiveros, P., and Seara Valero, J.R., La cuarta fase de deformación</u> <u>hercinica en la Zona Centroibérica del Macizo Hespérico: Cuadernos Del Laboratório Xeoloxico</u> <u>De Laxe, v. 9, p. 91–104, 1985.</u>

Gong, Z., Langereis, C. G. and Mullender, T. A. T.: The rotation of Iberia during the Aptian and the opening of the Bay of Biscay, Earth Planet. Sci. Lett., 273(1–2), 80–93, 2008.

Gozalo, R., Liñán, E., Vintaned, J. A. G., Eugenia, M., Álvarez, D., Martorell, J. B. C., Zamora, S., Esteve, J. and Mayoral, E.: The Cambrian of the Cadenas Ibéricas (ne Spain) and Its Trilobites., n.d.

Greco, K. D., Johnston, S. T., Gutiérrez-Alonso, G., Shaw, J. and Lozano, J. F.: Interference folding and orocline implications: A structural study of the Ponga Unit, Cantabrian orocline, northern Spain, Lithosphere, 8(6), 757–768, doi:10.1130/L576.1, 2016.

Gutiérrez Alonso, G.: Estratigrafía y Tectónica., in Memoria del Mapa Geológico de España 1:50000, Hoja de Mora (680), IGME, Madrid. [online] Available from: http://info.igme.es/cartografiadigital/datos/magna50/memorias/MMagna0658.pdf, 2009.

Gutiérrez-Alonso, G.: El Antiforme del Narcea y su relación con los mantos occidentales de la Zona Cantábrica., 1992.

Gutiérrez-Alonso, G.: Strain partitioning in the footwall of the Somiedo Nappe: Structural evolution of the Narcea Tectonic window, NW Spain, J. Struct. Geol., 18(10), 1217–1229, 1996.

Gutiérrez-Alonso, G., Fernández-Suárez, J. and Weil, A. B.: Orocline triggered lithospheric delamination, in Special Paper 383: Orogenic curvature: Integrating paleomagnetic and structural analyses, vol. 383, edited by A. B. Weil and A. Sussman, pp. 121–130, Geological Society of America, Boulder., 2004.

Gutiérrez-Alonso, G., Fernández-Suárez, J., Weil, A. B., Brendan Murphy, J., Damian Nance, R., Corf, F. and Johnston, S. T.: Self-subduction of the Pangaean global plate, Nat. Geosci., 1(8), 549–553, doi:10.1038/ngeo250, 2008a.

Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J. and Hamilton, M. A. A.: Rifting along the northern Gondwana margin and the evolution of the Rheic Ocean: A Devonian age for the El Castillo volcanic rocks (Salamanca, Central Iberian Zone), Tectonophysics, 461(1–4), 157–165, doi:10.1016/j.tecto.2008.01.013, 2008b.

Gutiérrez-Alonso, G., Fernández-Suárez, J., Jeffries, T. E. T. E. E., Johnston, S. T., Pastor-Galán, D., Murphy, J. B. B., Franco, M. P. P., Gonzalo, J. C. C.: Diachronous post-orogenic magmatism within a developing orocline in Iberia, European Variscides, Tectonics, 30(5), 17 PP.-17 PP., doi:10.1029/2010TC002845, 2011a.

Gutiérrez-Alonso, G., Murphy, J. B., Fernández-Suárez, J., Weil, A. B., Franco, M. P., Gonzalo, J. C.: Lithospheric delamination in the core of Pangea: Sm-Nd insights from the Iberian mantle, Geology, 39(2), 155–158, doi:10.1130/G31468.1, 2011b.

Gutiérrez-Alonso, G., Collins, A. S., Fernández-Suárez, J., Pastor-Galán, D., González-Clavijo, E., Jourdan, F., Weil, A. B. and Johnston, S. T.: Dating of lithospheric buckling:40Ar/39Ar ages of syn-orocline strike-slip shear zones in northwestern Iberia, Tectonophysics, 643, 44–54, doi:10.1016/j.tecto.2014.12.009, 2015.

Gutiérrez-Alonso, G., Fernández-Suárez, J., López-Carmona, A. and Gärtner, A.: Exhuming a cold case: The early granodiorites of the northwest Iberian Variscan belt—A Visean magmatic flare-up?, Lithosphere, 10(2), 194–216, doi:10.1130/L706.1, 2018.

Gutiérrez-Marco, J. C., Sá, A. A., García-Bellido, D. C. and Rábano, I.: The Bohemo-Iberian regional chronostratigraphical scale for the Ordovician System and palaeontological correlations within South Gondwana, Lethaia, 50(2), 258–295, doi:10.1111/let.12197, 2017.

Gutiérrez-Marco, J. C., Piçarra, J. M., Meireles, C. A., Cózar, P., García-Bellido, D. C., Pereira, Z., Vaz, N., Pereira, S., Lopes, G., Oliveira, J. T., Quesada, C., Zamora, S., Esteve, J., Colmenar, J., Bernárdez, E., Coronado, I., Lorenzo, S., Sá, A. A., Dias da Silva, Í., González-Clavijo, E., Díez-Montes, A. and Gómez-Barreiro, J.: Early Ordovician–Devonian Passive Margin Stage in the Gondwanan Units of the Iberian Massif, in The Geology of Iberia: A

Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 75–98, Springer International Publishing, Cham., 2019.

Crespo-Blanc, A. and Orozco, M.. The Southern Iberian Shear Zone: a major boundary in the Hercynian folded belt. Tectonophysics, 148(3-4), pp.221-227. <u>1988.</u>

Hindle, D., Besson, O. and Burkhard, M.: A model of displacement and strain for arc-shaped mountain belts applied to the Jura arc, J. Struct. Geol., 22(9), 1285–1296, doi:10.1016/S0191-8141(00)00038-9, 2000.

van Hinsbergen, D. J. J., Torsvik, T. H., Schmid, S. M., Maţenco, L. C., Maffione, M., Vissers, R. L. M., Gürer, D. and Spakman, W.: Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution since the Triassic, Gondwana Res., 81, 79–229, doi:10.1016/j.gr.2019.07.009, 2020.

Hirt, A. M., Lowrie, W., Julivert, M., Arboleya, M. L., Lowric, A. M. H. W. and Arboleya, M. L.: Paleomagnetic results in support of a model for the origin of the Asturian arc, Tectonophysics, 213(3–4), 321–339, 1992.

Hnat, J. S. and van der Pluijm, B. A.: Foreland signature of indenter tectonics: Insights from calcite twinning analysis in the Tennessee salient of the Southern Appalachians, USA, Lithosphere, 3(5), 317–327, doi:10.1130/L151.1, 2011.

Holmes, A.: A review of the continental drift hypothesis., Min. Mag., 40, 205–209, 1929.

Huang, W., Lippert, P. C., Zhang, Y., Jackson, M. J., Dekkers, M. J., Li, J., Hu, X., Zhang, B., Guo, Z. and van Hinsbergen, D. J. J. J.: Remagnetization of carbonate rocks in southern Tibet: Perspectives from rock magnetic and petrographic investigations, J. Geophys. Res. Solid Earth, 122(4), 2434–2456, doi:10.1002/2017JB013987, 2017.

Izquierdo-Llavall, E., Sainz, A. C., Oliva-Urcia, B., Burmester, R., Pueyo, E. L. and Housen, B.: Multi-episodic remagnetization related to deformation in the Pyrenean Internal Sierras, Geophys. J. Int., 201(2), 891–914, doi:10.1093/gji/ggv042, 2015.

Izquierdo–Llavall, E., Casas–Sainz, A. M., Oliva–Urcia, B., Villalaín, J. J., Pueyo, E. and Scholger, R.: Rotational Kinematics of Basement Antiformal Stacks: Paleomagnetic Study of the Western Nogueras Zone (Central Pyrenees), Tectonics, 37(10), 3456–3478, doi:10.1029/2018TC005153, 2018.

Izquierdo–Llavall, E., Ayala, C., Pueyo, E. L., Casas–Sainz, A. M., Oliva–Urcia, B., Rubio, F., Rodríguez–Pintó, A., Rey–Moral, C., Mediato, J. F. and García–Crespo, J.: Basement-Cover Relationships and Their Along-Strike Changes in the Linking Zone (Iberian Range, Spain): A Combined Structural and Gravimetric Study, Tectonics, 38(8), 2934–2960, doi:10.1029/2018TC005422, 2019.

Jabaloy-Sánchez, A., Talavera, C., Gómez-Pugnaire, M. T., López-Sánchez-Vizcaíno, V., Vázquez-Vílchez, M., Rodríguez-Peces, M. J. and Evans, N. J.: U-Pb ages of detrital zircons from the Internal Betics: A key to deciphering paleogeographic provenance and tectono-stratigraphic evolution, Lithos, 318–319, 244–266, doi:10.1016/j.lithos.2018.07.026, 2018.

Jackson, M.: Diagenetic sources of stable remanence in remagnetized paleozoic cratonic

carbonates: A rock magnetic study, J. Geophys. Res. Solid Earth, 95(B3), 2753–2761, doi:10.1029/JB095iB03p02753, 1990.

Jacques, D., Muchez, P. and Sintubin, M.: Superimposed folding and W-Sn vein-type mineralisation in the Central Iberian Zone associated with late-Variscan oroclinal buckling: A structural analysis from the Regoufe area (Portugal), Tectonophysics, 742–743, 66–83, doi:10.1016/j.tecto.2018.05.021, 2018a.

Jacques, D., Vieira, R., Muchez, P. and Sintubin, M.: Transpressional folding and associated cross-fold jointing controlling the geometry of post-orogenic vein-type W-Sn mineralization: examples from Minas da Panasqueira, Portugal, Miner. Deposita, 53(2), 171–194, doi:10.1007/s00126-017-0728-6, 2018b.

Johnston, S. T.: The Great Alaskan Terrane Wreck: Reconciliation of paleomagnetic and geological data in the Northern Cordillera, Earth Planet. Sci. Lett., 193(3–4), 259–272, doi:10.1016/S0012-821X(01)00516-7, 2001.

Johnston, S. T., Weil, A. B. and Gutiérrez-Alonso, G.: Oroclines: Thick and thin, GSA Bull., 125(5–6), 643–663, doi:10.1130/B30765.1, 2013.

Julivert, M.: L'évolution structurale de l'Arc Asturien, Paris., 1971.

Julivert, M. and Arboleya, M. L.: A geometrical and kinematical approach to the nappe structure in an arcuate fold belt: the Cantabrian nappes (Hercynian chain, NW Spain), J. Struct. Geol., 6(5), 499–519, doi:10.1016/0191-8141(84)90061-0, 1984.

Julivert, M. and Marcos, A.: superimposed folding under flexural conditions in the Cantabrian Zone (Hercynian Cordillera, Northwest Spain), Am. J. Sci., 273(5), 353–375, doi:10.2475/ajs.273.5.353, 1973.

Julivert, M., Fontboté, J. M., Ribeiro, A. and Nabais Conde, L. E.: Mapa Tectónico de la Península Ibérica y Baleares E: 1: 1.000. 000 y memoria explicativa, Publ IGME, 113, 1974.

Julivert, M., Vegas, R., Roiz, J. M. and Martínez Rius, A.: Laestructura de la extension SE de la Zona Centro-Iberica con metamorfismo de bajo grado, Rios-Geol. Esp. I, 339–380, 1983.

Keppie, F.: How subduction broke up Pangaea with implications for the supercontinent cycle, Geol. Soc. Lond. Spec. Publ., 424(1), 265–288, doi:10.1144/SP424.8, 2016.

Kirsch, M., Keppie, J. D., Murphy, J. B. and Solari, L. A.: Permian-Carboniferous arc magmatism and basin evolution along the western margin of Pangea: Geochemical and geochronological evidence from the eastern Acatlan Complex, southern Mexico, Geol. Soc. Am. Bull., 124(9–10), 1607–1628, doi:10.1130/B30649.1, 2012.

Kollmeier, J. M., van der Pluijm, B. A. and Van der Voo, R.: Analysis of Variscan dynamics; early bending of the Cantabria-Asturias Arc, northern Spain, Earth Planet. Sci. Lett., 181(1–2), 203–216, doi:10.1016/S0012-821X(00)00203-X, 2000.

Kroner, U. and Romer, R. L. L.: Two plates — Many subduction zones: The Variscan orogeny reconsidered, Gondwana Res., 24(1), 298–329, doi:10.1016/j.gr.2013.03.001, 2013.

Li, P. and Rosenbaum, G.: Does the Manning Orocline exist? New structural evidence from the inner hinge of the Manning Orocline (eastern Australia), Gondwana Res., 25(4), 1599–1613, doi:10.1016/j.gr.2013.06.010, 2014.

Li, P., Sun, M., Rosenbaum, G., Yuan, C., Safonova, I., Cai, K., Jiang, Y. and Zhang, Y.: Geometry, kinematics and tectonic models of the Kazakhstan Orocline, Central Asian Orogenic Belt, J. Asian Earth Sci., 153(July 2017), 42–56, doi:10.1016/j.jseaes.2017.07.029, 2018.

Li, P.-F., Rosenbaum, G. and Rubatto, D.: Triassic asymmetric subduction rollback in the southern New England Orogen (eastern Australia): the end of the Hunter-Bowen Orogeny, Aust. J. Earth Sci., 59(6), 965–981, doi:10.1080/08120099.2012.696556, 2012.

López-Carmona, A., Abati, J., Pitra, P. and Lee, J. K. W.: Retrogressed lawsonite blueschists from the NW Iberian Massif: P–T–t constraints from thermodynamic modelling and 40Ar/39Ar geochronology, Contrib. Mineral. Petrol., 167(3), 987–987, doi:10.1007/s00410-014-0987-5, 2014.

López-Moro, F. J., López-Plaza, M., Gutiérrez-Alonso, G., Fernández-Suárez, J., López-Carmona, A., Hofmann, M. and Romer, R. L.: Crustal melting and recycling: geochronology and sources of Variscan syn-kinematic anatectic granitoids of the Tormes Dome (Central Iberian Zone). A U–Pb LA-ICP-MS study, Int. J. Earth Sci., 107(3), 985–1004, doi:10.1007/s00531-017-1483-8, 2018.

Lotze, F.: Zur gliederung der Variszichen der Iberischen Meseta., Geotektonische Forschungen, 6, 78–92, 1945.

Macaya, J., González-Lodeiro, F., Martínez-Catalán, J. R. and Alvarez, F.: Continuous deformation, ductile thrusting and backfolding of cover and basement in the Sierra de Guadarrama, Hercynian orogen of central Spain, Tectonophysics, 191(3), 291–309, doi:10.1016/0040-1951(91)90063-X, 1991.

Maffione, M., Speranza, F. and Faccenna, C.: Bending of the Bolivian orocline and growth of the central Andean plateau: Paleomagnetic and structural constraints from the Eastern Cordillera (22–24°S, NW Argentina), Tectonics, 28, 23 PP.-23 PP., doi:10.1029/2008TC002402, 2009.

Marcos, A. and Pulgar, J. A. A.: An Approach to the tectonostratigraphic evolution of the Cantabrian foreland thrust and fold belt, Hercynian Cordillera of NW Spain, Neues Jahrb. Geol. Palaontologie-Abh., 163(2), 256–260, 1982.

Marcos, A., Pérez-Estaún, A., Martínez Catalán, J. R. and Gutiérrez-Marco, J. C.: Estratigrafía y Paleogeografía. Zona Asturoccidental-Leonesa, 2004.

Marshak, S.: KINEMATICS OF OROCLINE AND ARC FORMATION IN THIN-SKINNED OROGENS, Tectonics, 7(1), 73–86, 1988.

Marshak, S.: Salients, Recesses, Arcs, Oroclines, and Syntaxes — A Review of Ideas Concerning the Formation of Map-view Curves in Fold-thrust Belts, Thrust Tecton. Hydrocarb. Syst. AAPG Mem. 82, 82(1893), 131–156, 2004.

Martín-Algarra, A., García-Casco, A., Gómez-Pugnaire, M. T., Jabaloy-Sánchez, A., Laborda-López, C., López Sánchez-Vizcaíno, V., Mazzoli, S., Navas-Parejo, P., Perrone, V., RodríguezCañero, R. and Sánchez-Navas, A.: Paleozoic Basement and Pre-Alpine History of the Betic Cordillera, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 261–305, Springer International Publishing, Cham., 2019.

Martínez Catalán, J. R.: Are the oroclines of the Variscan belt related to late Variscan strike-slip tectonics?, Terra Nova, 23(4), 241–247, doi:10.1111/j.1365-3121.2011.01005.x, 2011.

Martínez Catalán, J. R.: The Central Iberian arc, an orocline centered in the Iberian Massif and some implications for the Variscan belt, Int. J. Earth Sci., 101(5), 1–16, doi:10.1007/s00531-011-0715-6, 2012.

Martinez Catalan, J. R., Rodriguez, M. P. H., Alonso, P. V., Perez-Estaun, A. and Lodeiro, F. G.: Lower Paleozoic extensional tectonics in the limit between the West Asturian-Leonese and Central Iberian Zones of the Variscan Fold-Belt in NW Spain, Geol. Rundsch., 81(2), 545–560, doi:10.1007/BF01828614, 1992.

Martínez Catalán, J. R., Martínez Poyatos, D. and Bea, F.: Zona Centroibérica, Geol. Esp., 68–133, 2004.

Martínez Catalán, J. R., Arenas, R., Abati, J., Martínez, S. S., García, F. D., Suárez, J. F., Cuadra, P. G., Castiñeiras, P., Barreiro, J. G., Montes, A. D., Clavijo, E. G., Pascual, F. J. R., Andonaegui, P., Jeffries, T. E., Alcock, J. E., Fernández, R. D. and Carmona, A. L.: A rootless suture and the loss of the roots of a mountain chain: The Variscan belt of NW Iberia, Comptes Rendus Geosci., 341(2), 114–126, doi:10.1016/j.crte.2008.11.004, 2009.

Martinez Catalan, J. R., Rubio Pascual, F. J., Diez Montes, A., Diez Fernandez, R., Gomez Barreiro, J., Dias da Silva, I., Gonzalez Clavijo, E., Ayarza, P., Alcock, J. E., Montes, A. D., Fernandez, R. D., Barreiro, J. G., Dias da Silva, I., Clavijo, E. G., Ayarza, P., Alcock, J. E., Diez Montes, A., Diez Fernandez, R., Gomez Barreiro, J., Dias da Silva, I., Gonzalez Clavijo, E., Ayarza, P., Alcock, J. E., Montes, A. D., Fernandez, R. D., Barreiro, J. G., Dias da Silva, I., Clavijo, E. G., Ayarza, P. and Alcock, J. E.: The late Variscan HT/LP metamorphic event in NW and Central Iberia: relationships to crustal thickening, extension, orocline development and crustal evolution, Geol. Soc. Lond. Spec. Publ., 405(1), 225–247, doi:10.1144/SP405.1, 2014.

Martínez Catalán, J. R., Aerden, D. G. A. M. and Carreras, J.: The "Castilian bend" of Rudolf Staub (1926): historical perspective of a forgotten orocline in Central Iberia, Swiss J. Geosci., 108(2–3), 289–303, doi:10.1007/s00015-015-0202-3, 2015.

Martínez Catalán, J. R., Gómez Barreiro, J., Dias da Silva, Í., Chichorro, M., López-Carmona, A., Castiñeiras, P., Abati, J., Andonaegui, P., Fernández-Suárez, J., González Cuadra, P. and Benítez-Pérez, J. M.: Variscan Suture Zone and Suspect Terranes in the NW Iberian Massif: Allochthonous Complexes of the Galicia-Trás os Montes Zone (NW Iberia), in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 99–130, Springer International Publishing, Cham., 2019.

Martínez-Catalán, J. R.: A non-cylindrical model for the northwest Iberian allochthonous terranes and their equivalents in the Hercynian belt of Western Europe, Tectonophysics, 179(3–4), 253–272, doi:10.1016/0040-1951(90)90293-H, 1990.

McWilliams, C. K., Kunk, M. J., Wintsch, R. P. and Bish, D. L.: Determining ages of multiple

muscovite-bearing foliations in phyllonites using the 40Ar/39Ar step heating method: Applications to the alleghanian orogeny in central new England, Am. J. Sci., 313(10), 996–1016, doi:10.2475/10.2013.02, 2013.

Meijers, M. J. M., Smith, B., Pastor-Galán, D., Degenaar, R., Sadradze, N., Adamia, S., Sahakyan, L., Avagyan, A., Sosson, M., Rolland, Y., Langereis, C. G. and Müller, C.: Progressive orocline formation in the Eastern Pontides-Lesser Caucasus, Geol. Soc. Spec. Publ., 428(1), doi:10.1144/SP428.8, 2017.

Merino-Tomé, O. A., Bahamonde, J. R., Colmenero, J. R., Heredia, N., Villa, E. and Farias, P.: Emplacement of the Cuera and Picos de Europa imbricate system at the core of the Iberian-Armorican arc (Cantabrian zone, north Spain): New precisions concerning the timing of arc closure, Bull. Geol. Soc. Am., 121(5–6), 729–751, doi:10.1130/B26366.1, 2009.

Molina Garza, R. S. and Zijderveld, J. D. a.: Paleomagnetism of Paleozoic strata, Brabant and Ardennes Massifs, Belgium: Implications of prefolding and postfolding Late Carboniferous secondary magnetizations for European apparent polar wander, J. Geophys. **Res.**, **101(B7)**, **15799–15799**, **doi:10.1029/96JB00325**, **1996**.

Moral, B. del and Sarmiento, G. N.: Conodontos del Katiense (Ordovícico Superior) del sector meridional de la Zona Centroibérica (España), Rev. Esp. Micropaleontol., 40(3), 169–245, 2008.

Munha, J.: Metamorphic Evolution of the South Portuguese/Pulo Do Lobo Zone, in Pre-Mesozoic Geology of Iberia, edited by R. D. Dallmeyer and E. M. Garcia, pp. 363–368, Springer, Berlin, Heidelberg., 1990.

Munha, J., Barriga, F. J. a. S. and Kerrich, R.: High 18 O ore-forming fluids in volcanic-hosted base metal massive sulfide deposits; geologic, 18 O/ 16 O, and D/H evidence from the Iberian pyrite belt; Crandon, Wisconsin; and Blue Hill, Maine, Econ. Geol., 81(3), 530–552, doi:10.2113/ gsecongeo.81.3.530, 1986.

Murphy, J. B., Gutierrez-Alonso, G., Fernandez-Suarez, J. and Braid, J. A.: Probing crustal and mantle lithosphere origin through Ordovician volcanic rocks along the Iberian passive margin of Gondwana, Tectonophysics, 461(1–4), 166–180, 2008.

Nance, R. D., Gutiérrez-Alonso, G., Keppie, J. D., Linnemann, U., Murphy, J. B., Quesada, C., Strachan, R. A. and Woodcock, N. H.: Evolution of the Rheic Ocean, Gondwana Res., 17(2–3), 194–222, doi:10.1016/j.gr.2009.08.001, 2010.

Oliveira, J. T., Quesada, C., Pereira, Z., Matos, J. X., Solá, A. R., Rosa, D., Albardeiro, L., Díez-Montes, A., Morais, I., Inverno, C., Rosa, C. and Relvas, J.: South Portuguese Terrane: A Continental Affinity Exotic Unit, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 173–206, Springer International Publishing, Cham., 2019a.

Oliveira, J. T., González-Clavijo, E., Alonso, J., Armendáriz, M., Bahamonde, J. R., Braid, J. A., Colmenero, J. R., Dias da Silva, Í., Fernandes, P., Fernández, L. P., Gabaldón, V., Jorge, R. S., Machado, G., Marcos, A., Merino-Tomé, Ó., Moreira, N., Murphy, J. B., Pinto de Jesus, A., Quesada, C., Rodrigues, B., Rosales, I., Sanz-López, J., Suárez, A., Villa, E., Piçarra, J. M. and Pereira, Z.: Synorogenic Basins, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 349–429, Springer International Publishing, Cham., 2019b.

P. Farias, G. G.: Aportaciones al conocimiento de la litoestratigrafía y estructura de Galicia Central, Mem. Fac. Ciênc. Universidade Porto, 411–431, 1987.

Palero-Fernández, F. J., Martin-Izard, A., Zarzalejos Prieto, M. and Mansilla-Plaza, L.: Geological context and plumbotectonic evolution of the giant Almadén Mercury Deposit, Ore Geol. Rev., 64, 71–88, doi:10.1016/j.oregeorev.2014.06.013, 2015.

Parés, J. M. and Van der Voo, R.: Paleozoic paleomagnetism of Almaden, Spain: A cautionary note, J. Geophys. Res., 97(B6), 9353–9356, doi:10.1029/91JB03073, 1992.

Parés, J. M., Van der Voo, R., Stamatakos, J. and Pérez–Estaún, A.: Remagnetizations and postfolding oroclinal rotations in the Cantabrian/Asturian arc, northern Spain, Tectonics, 13(6), 1461–1471, doi:10.1029/94TC01871, 1994.

Pastor-Galán, D., Gutiérrez-Alonso, G., Meere, P. A. and Mulchrone, K. F.: Factors affecting finite strain estimation in low-grade, low-strain clastic rocks, J. Struct. Geol., 31(12), 1586–1596, doi:10.1016/j.jsg.2009.08.005, 2009.

Pastor-Galán, D., Gutiérrez-Alonso, G. and Weil, A. B.: Orocline timing through joint analysis: Insights from the Ibero-Armorican Arc, Tectonophysics, 507(1), 31–46, doi:10.1016/j.tecto.2011.05.005, 2011.

Pastor-Galán, D., Gutiérrez-Alonso, G., Zulauf, G. and Zanella, F.: Analogue modeling of lithospheric-scale orocline buckling: Constraints on the evolution of the Iberian-Armorican arc, Bull. Geol. Soc. Am., 124(7–8), doi:10.1130/B30640.1, 2012a.

Pastor-Galán, D., Gutiérrez-Alonso, G., Mulchrone, K. F. and Huerta, P.: Conical folding in the core of an orocline. A geometric analysis from the Cantabrian Arc (Variscan Belt of NW Iberia), J. Struct. Geol., 39, 210–223, doi:10.1016/j.jsg.2012.02.010, 2012b.

Pastor-Galán, D., Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J., Hofmann, M. and Linnemann, U.: Provenance analysis of the Paleozoic sequences of the northern Gondwana margin in NW Iberia: Passive margin to Variscan collision and orocline development, Gondwana Res., 23(3), 1089–1103, doi:10.1016/j.gr.2012.06.015, 2013a.

Pastor-Galán, D., Gutiérrez-Alonso, G., Fernández-Suárez, J., Murphy, J. B. and Nieto, F.: Tectonic evolution of NW Iberia during the Paleozoic inferred from the geochemical record of detrital rocks in the Cantabrian Zone, Lithos, 182–183, 221–228, doi:10.1016/j.lithos.2013.09.007, 2013b.

Pastor-Galán, D., Martín-Merino, G. and Corrochano, D.: Timing and structural evolution in the limb of an orocline: The Pisuerga-Carrión Unit (southern limb of the Cantabrian Orocline, NW Spain), Tectonophysics, 622, 110–121, doi:10.1016/j.tecto.2014.03.004, 2014.

Pastor-Galán, D., Ursem, B., Meere, P. A. and Langereis, C.: Extending the Cantabrian Orocline to two continents (from Gondwana to Laurussia). Paleomagnetism from South Ireland, Earth Planet. Sci. Lett., 432, doi:10.1016/j.epsl.2015.10.019, 2015a.

Pastor-Galan, D., Groenewegen, T., Brouwer, D., Krijgsman, W., Dekkers, M. J., Pastor-galán, D., Groenewegen, T., Brouwer, D., Krijgsman, W. and Dekkers, M. J.: One or two oroclines in the Variscan orogen of Iberia? Implications for Pangea amalgamation, Geology, 43(6), 527–

530, doi:10.1130/G36701.1, 2015b.

Pastor-Galán, D., Dekkers, M. J., Gutiérrez-Alonso, G., Brouwer, D., Groenewegen, T., Krijgsman, W., Fernández-Lozano, J., Yenes, M. and Álvarez-Lobato, F.: Paleomagnetism of the Central Iberian curve's putative hinge: Too many oroclines in the Iberian Variscides, Gondwana Res., 39, 96–113, doi:10.1016/j.gr.2016.06.016, 2016.

Pastor-Galán, D., Mulchrone, K. F., Koymans, M. R., van Hinsbergen, D. J. J. and Langereis, C. G. : Bootstrapped total least squares orocline test: A robust method to quantify vertical-axis rotation patterns in orogens, with examples from the Cantabrian and Aegean oroclines, Lithosphere, 9(3), L547.1-L547.1, doi:10.1130/L547.1, 2017a.

Pastor-Galán, D., Gutiérrez-Alonso, G., Dekkers, M. J. M. J. and Langereis, C. G.: Paleomagnetism in Extremadura (Central Iberian zone, Spain) Paleozoic rocks: extensive remagnetizations and further constraints on the extent of the Cantabrian orocline, J. Iber. Geol., 43(4), 583–600, doi:10.1007/s41513-017-0039-x, 2017b.

Pastor-Galán, D., Pueyo, E. L., Diederen, M., García-Lasanta, C. and Langereis, C. G.: Late Paleozoic Iberian Orocline(s) and the Missing Shortening in the Core of Pangea. Paleomagnetism From the Iberian Range, Tectonics, 37(10), 3877–3892, doi:10.1029/2018TC004978, 2018.

Pastor-Galán, D., Nance, R. D., Murphy, J. B. and Spencer, C. J.: Supercontinents: myths, mysteries, and milestones, Geol. Soc. Lond. Spec. Publ., 470, 39–64, doi:10.1144/SP470.16, 2019a.

Pastor-Galán, D., Dias da Silva, Í. F., Groenewegen, T. and Krijgsman, W.: Tangled up in folds: tectonic significance of superimposed folding at the core of the Central Iberian curve (West Iberia), Int. Geol. Rev., 61(2), 240–255, doi:10.1080/00206814.2017.1422443, 2019b.

Pereira, I., Dias, R., Bento, T. and Mata, J.: Exhumation of a migmatite complex along a transpressive shear zone : inferences from the Variscan Juzbado – Penalva do Castelo Shear Zone (Central Iberian Zone), J. Geol. Soc., doi:10.1144/jgs2016-159, 2017.

Pereira, M. F., Chichorro, M., Silva, J. B., Ordóñez-Casado, B., Lee, J. K. W. W. and Williams, I. S.: Early carboniferous wrenching, exhumation of high-grade metamorphic rocks and basin instability in SW Iberia: Constraints derived from structural geology and U–Pb and 40Ar–39Ar geochronology, Tectonophysics, 558, 28–44, doi:10.1016/j.tecto.2012.06.020, 2012.

Pereira, M. F., Díez Fernández, R., Gama, C., Hofmann, M., Gärtner, A. and Linnemann, U.: Stype granite generation and emplacement during a regional switch from extensional to contractional deformation (Central Iberian Zone, Iberian autochthonous domain, Variscan Orogeny), Int. J. Earth Sci., 107(1), 251–267, doi:10.1007/s00531-017-1488-3, 2018.

Pérez–Cáceres, I., Poyatos, D. M., Simancas, J. F. and Azor, A.: The elusive nature of the Rheic Ocean suture in SW Iberia, Tectonics, 34(12), 2429–2450, doi:10.1002/2015TC003947, 2015.

Pérez-Cáceres, I., Simancas, J. F., Martínez Poyatos, D., Azor, A. and González Lodeiro, F.: Oblique collision and deformation partitioning in the SW Iberian Variscides, Solid Earth, 7(3), 857–872, doi:https://doi.org/10.5194/se-7-857-2016, 2016.

Pérez-Cáceres, I., Martínez Poyatos, D., Simancas, J. F. and Azor, A.: Testing the Avalonian affinity of the South Portuguese Zone and the Neoproterozoic evolution of SW Iberia through detrital zircon populations, Gondwana Res., 42, 177–192, doi:10.1016/j.gr.2016.10.010, 2017.

Pérez-Cáceres, I., Martínez Poyatos, D. J., Vidal, O., Beyssac, O., Nieto, F., Simancas, J. F., Azor, A. and Bourdelle, F.: Deciphering the metamorphic evolution of the Pulo do Lobo metasedimentary domain (SW Iberian Variscides), Solid Earth, 11(2), 469–488, doi:https://doi.org/10.5194/se-11-469-2020, 2020.

Pérez-Estaún, A., Bastida, F., Alonso, J. L., Marquinez, J., Aller, J., Alvarezmarron, J., Marcos, A. and Pulgar, J. A.: A THIN-SKINNED TECTONICS MODEL FOR AN ARCUATE FOLD AND THRUST BELT - THE CANTABRIAN ZONE (VARISCAN IBERO-ARMORICAN ARC), Tectonics, 7(3), 517–537, 1988.

Pérez-Estaún, A., Bastida, F., Martínez Catalán, J. R., Gutiérrez-Marco, J. C., Marcos, A. and Pulgar, J.: Stratigraphy of the West Asturian-Leonese Zone, Springer. [online] Available from: https://digital.csic.es/handle/10261/30719 (Accessed 6 April 2020), 1990.

Pérez-Estaún, A., Martinezcatalan, J. R. and Bastida, F.: CRUSTAL THICKENING AND DEFORMATION SEQUENCE IN THE FOOTWALL TO THE SUTURE OF THE VARISCAN BELT OF NORTHWEST SPAIN, Tectonophysics, 191(3–4), 243–253, 1991.

Perroud, H., Bonhommet, N. and Ribeiro, A.: Paleomagnetism of Late Paleozoic igneous rocks from southern Portugal, Geophys. Res. Lett., 12(1), 45–48, doi:10.1029/GL012i001p00045, 1985.

Perroud, H., Calza, F. and Khattach, D.: Paleomagnetism of the Silurian Volcanism at Almaden, Southern Spain, J. Geophys. Res.-Solid Earth Planets, 96(B2), 1949–1962, doi:10.1029/90JB02226, 1991.

Pin, C., Paquette, J. L., Zalduegui, J. F. S. and Ibarguchi, J. I. G.: Early Devonian suprasubduction-zone ophiolite related to incipient collisional processes in the Western Variscan Belt: The Sierra de Careón unit, Ordenes Complex, Galicia, in Variscan-Appalachian dynamics: The building of the late Paleozoic basement, Geological Society of America., 2002.

Pueyo, E. L., Mauritsch, H. J., Gawlick, H.-J., Scholger, R. and Frisch, W.: New evidence for block and thrust sheet rotations in the central northern Calcareous Alps deduced from two pervasive remagnetization events, Tectonics, 26(5), n/a-n/a, doi:10.1029/2006TC001965, 2007.

Pueyo, E. L., Sussman, A. J., Oliva-Urcia, B. and Cifelli, F.: Palaeomagnetism in fold and thrust belts: Use with caution, Geol. Soc. Spec. Publ., 425(1), 259–276, doi:10.1144/SP425.14, 2016.

Quesada, C.: The Ossa-Morena Zone of the Iberian Massif: a tectonostratigraphic approach to its evolution, Z. Dtsch. Ges. Für Geowiss., 585–595, doi:10.1127/1860-1804/2006/0157-0585, 2006.

Quesada, C. and Dallmeyer, R. D. D.: Tectonothermal evolution of the Badajoz-Cordóba shear zone (SW Iberia): characteristics and 40Ar/39Ar mineral age constraints, Tectonophysics, 231(1–3), 195–213, doi:10.1016/0040-1951(94)90130-9, 1994.

Quesada, C., Braid, J. A., Fernandes, P., Ferreira, P., Jorge, R. S., Matos, J. X., Murphy, J. B.,

Oliveira, J. T., Pedro, J. and Pereira, Z.: SW Iberia Variscan Suture Zone: Oceanic Affinity Units, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 131–171, Springer International Publishing, Cham., 2019.

Rankin, D. W.: Appalachian salients and recesses: Late Precambrian continental breakup and the opening of the lapetus Ocean, J. Geophys. Res. 1896-1977, 81(32), 5605–5619, doi:10.1029/JB081i032p05605, 1976.

Ribeiro, A., Munhá, J., Dias, R., Mateus, A., Pereira, E., Ribeiro, L., Fonseca, P., Araújo, A., Oliveira, T., Romão, J., Chaminé, H., Coke, C. and Pedro, J.: Geodynamic evolution of the SW Europe Variscides, Tectonics, 26(6), doi:10.1029/2006TC002058, 2007.

Ries, A. C. and Shackleton, R. M.: Catazonal Complexes of North-West Spain and North Portugal, Remnants of a Hercynian Thrust Plate, Nat. Phys. Sci., 234(47), 65–68, doi:10.1038/physci234065a0, 1971.

Ries, A. C. and Shackleton, R. M.: Patterns of Strain Variation in Arcuate Fold Belts, Philos. Trans. R. Soc. Lond. Ser. Math. Phys. Sci., 283(1312), 281–288, 1976.

Rodríguez-Cañero, R., Jabaloy-Sánchez, A., Navas-Parejo, P., Martín-Algarra, A., Rodríguez, R., Antonio, C., Sánchez, J., Navas, P., Agustín, P. and Algarra, M.: Linking Palaeozoic palaeogeography of the Betic Cordillera to the Variscan Iberian Massif : new insight through the first conodonts of the Nevado-Filábride Complex, Int. J. Earth Sci., 107(5), 1791–1806, doi:10.1007/s00531-017-1572-8, 2018.

Rosenbaum, G.: Geodynamics of oroclinal bending: Insights from the Mediterranean, J. Geodyn., 82, 5–15, doi:10.1016/j.jog.2014.05.002, 2014.

Rubio Pascual, F. J., Arenas, R., Martínez Catalán, J. R., Rodríguez Fernández, L. R. and Wijbrans, J. R.: Thickening and exhumation of the Variscan roots in the Iberian Central System: Tectonothermal processes and 40Ar/39Ar ages, Tectonophysics, 587, 207–221, doi:10.1016/j.tecto.2012.10.005, 2013.

Rubio Pascual, F. J., López-Carmona, A. and Arenas, R.: Thickening vs. extension in the Variscan belt: P–T modelling in the Central Iberian autochthon, Tectonophysics, 681, 144–158, doi:10.1016/j.tecto.2016.02.033, 2016.

Sánchez-García, T., Chichorro, M., Solá, A. R., Álvaro, J. J., Díez-Montes, A., Bellido, F., Ribeiro, M. L., Quesada, C., Lopes, J. C., Dias da Silva, Í., González-Clavijo, E., Gómez Barreiro, J. and López-Carmona, A.: The Cambrian-Early Ordovician Rift Stage in the Gondwanan Units of the Iberian Massif, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 27–74, Springer International Publishing, Cham., 2019.

Schulz, G.: Descripción geológica de Asturias: Publicada de Real Órden. Con un atlas, José Gonzalez., 1858.

Schwartz, S. Y. and Vandervoo, R.: PALEOMAGNETIC EVALUATION OF THE OROCLINE HYPOTHESIS IN THE CENTRAL AND SOUTHERN APPALACHIANS, Geophys. Res. Lett., 10(7), 505–508, 1983.

Shaw, J. and Johnston, S. T.: Terrane wrecks (coupled oroclines) and paleomagnetic inclination anomalies, Earth-Sci. Rev., 154, 191–209, doi:10.1016/j.earscirev.2016.01.003, 2016.

Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Weil, A. B.: Oroclines of the Variscan orogen of Iberia: Paleocurrent analysis and paleogeographic implications, Earth Planet. Sci. Lett., 329–330, 60–70, doi:10.1016/j.epsl.2012.02.014, 2012.

Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Pastor-Galán, D.: Provenance variability along the Early Ordovician north Gondwana margin: Paleogeographic and tectonic implications of U-Pb detrital zircon ages from the Armorican Quartzite of the Iberian Variscan belt, Bull. Geol. Soc. Am., 126(5–6), 702–719, doi:10.1130/B30935.1, 2014.

Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A structural study of the Cantabrian orocline, NW Iberian Massif, Lithosphere, 7(6), 653–661, doi:10.1130/L461.1, 2015.

Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A structural study of the Cantabrian orocline, NW Iberian Massif, Lithosphere, 8(1), 97–97, doi:10.1130/L461.1, 2016.

Silva, Í. D. D., Valverde-Vaquero, P., González-Clavijo, E., Díez-Montes, A. and Catalán, J. R. M.: Structural and stratigraphical significance of U–Pb ages from the Mora and Saldanha volcanic complexes (NE Portugal, Iberian Variscides), Geol. Soc. Lond. Spec. Publ., 405(1), 115–135, doi:10.1144/SP405.3, 2014.

Simancas, J. F., Carbonell, R., Lodeiro, F. G., Estaun, A., Juhlin, C., Ayarza, P., Kashubin, A., Azor, A., Poyatos, D. M., Almodovar, G. R., Pascual, E., Saez, R. and Exposito, I.: Crustal structure of the transpressional Variscan orogen of SW Iberia: SW Iberia deep seismic reflection profile (IBERSEIS), Tectonics, 22(6), 25–25, 2003.

Simancas, J. F., Ayarza, P., Azor, a., Carbonell, R., Martínez Poyatos, D., Pérez-Estaún, a. and González Lodeiro, F.: A seismic geotraverse across the Iberian Variscides: Orogenic shortening, collisional magmatism, and orocline development, Tectonics, 32(i), n/a-n/a, doi:10.1002/tect.20035, 2013.

Solís-Alulima, B. E., López-Carmona, A., Gutiérrez Alonso, G. and Álvarez Valero, A. M.: Petrologic and thermobarometric study of the Riás schists (NW Iberian Massif), Bol. Geológico Min., 130(3), 445–464, doi:10.21701/bolgeomin.130.3.004, 2019.

Stampfli, G. M. and Borel, G. D.: The TRANSMED Transects in Space and Time: Constraints from Paleotectonic Evolution of the Mediterranean Domain., edited by W. Cavazza, F. M. Roure, W. Spakman, G. M. Stampfli, and P. A. Ziegler, pp. 53–80, Springer, Berlin., 2004.

Stampfli, G. M., Hochard, C., Vérard, C., Wilhem, C. and VonRaumer, J.: The Formation of Pangea, Tectonophysics, 593, 1–19, doi:10.1016/j.tecto.2013.02.037, 2013.

Staub, R.: Gedanken zur Tektonik Spaniens., Zürich., 1926.

Stewart, S. A.: Paleomagnetic analysis of fold kinematics and implications for geological models of the Cantabrian/Asturian arc, north Spain, J. Geophys. Res. Solid Earth, 100(B10), 20079–20094, doi:10.1029/95JB01482, 1995.

Tait, J. A., Bachtadse, V. and Soffel, H. C.: Eastern Variscan fold belt : Paleomagnetic evidence for oroclinal bending, Geology, 24, 871–874, doi:10.1130/0091-7613(1996)024<0871, 1996.

Tauxe, L.: Essentials of Paleomagnetism, Univ of California Press., 2010.

Thomas, W. A.: Evolution of Appalachian-Ouachita Salients and Recesses from Reentrants and Promontories in the Continental Margin, Am. J. Sci., 277(10), 1233–1278, doi:10.2475/ajs.277.10.1233, 1977.

Thomas, W. A.: Genetic relationship of rift-stage crustal structure, terrane accretion, and foreland tectonics along the southern Appalachian-Ouachita orogen, J. Geodyn., 37(3), 549–563, doi:10.1016/j.jog.2004.02.020, 2004.

Tohver, E., Weil, A. B. B. B., Solum, J. G. G. G. and Hall, C. M. M. M.: Direct dating of carbonate remagnetization by 40Ar/39Ar analysis of the smectite–illite transformation, Earth Planet. Sci. Lett., 274(3–4), 524–530, doi:10.1016/j.epsl.2008.08.002, 2008.

Toit, A. L. Du: Our wandering continents: an hypothesis of continental drifting, Oliver and Boyd. [online] Available from: http://books.google.es/books?id=iDZEAAAAIAAJ, 1937.

Valladares, M. I., Barba, P., Ugidos, J. M., Colmenero, J. R. and Armenteros, I.: Upper Neoproterozoic-Lower Cambrian sedimentary successions in the Central Iberian Zone (Spain): sequence stratigraphy, petrology and chemostratigraphy. Implications for other European zones, Int. J. Earth Sci., 89(1), 2–20, 2000.

Van der Voo, R.: Paleomagnetism, oroclines, and growth of the continental crust, GSA Today, 14(12), 4–9, doi:10.1130/1052-5173(2004), 2004.

Vergés, J.: Estudio del Complejo volcáno-sedimentario del Devónico y de la estructura de la terminación oriental del sinclinal de Almadén (Ciudad Real), in Libro Jubilar JM Rios. Tomo 3, pp. 215–229., 1983.

Vissers, R. L. M., van Hinsbergen, D. J. J., van der Meer, D. G. and Spakman, W.: Cretaceous slab break-off in the Pyrenees: Iberian plate kinematics in paleomagnetic and mantle reference frames, Gondwana Res., 34, 49–59, doi:10.1016/J.GR.2016.03.006, 2016.

van der Voo, R., Stamatakos, J. A. and Pares, J. M.: Kinematic constraints on thrust-belt curvature from syndeformational magnetizations in the Lagos del Valle Syncline in the Cantabrian Arc, Spain, J. Geophys. Res.-Solid Earth, 102(B5), 10105–10119, 1997.

Weil, A., Pastor-Galán, D., Johnston, S. T. and Gutiérrez-Alonso, G.: Late/Post Variscan Orocline Formation and Widespread Magmatism, in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp. 527–542, Springer International Publishing, Cham., 2019.

Weil, A. B.: Kinematics of orocline tightening in the core of an arc: Paleomagnetic analysis of the Ponga Unit, Cantabrian Arc, northern Spain, Tectonics, 25(3), n/a-n/a, doi:10.1029/2005TC001861, 2006.

Weil, A. B. and Sussman, A. J.: Classifying curved orogens based on timing relationships between structural development and vertical-axis rotations, vol. 383, edited by A. J. Sussman

and A. B. Weil, pp. 1–16, Geological Society of America., 2004.

Weil, A. B., Van der Voo, R., van der Pluijm, B. A. and Parés, J. M.: The formation of an orocline by multiphase deformation: a paleomagnetic investigation of the Cantabria–Asturias Arc (northern Spain), J. Struct. Geol., 22(6), 735–756, doi:10.1016/S0191-8141(99)00188-1, 2000.

Weil, A. B., van der Voo, R. and van der Pluijm, B. A.: Oroclinal bending and evidence against the Pangea megashear: The Cantabria-Asturias arc (northern Spain), Geology, 29(11), 991–994, 2001.

Weil, A. B., Van der Voo, R. and Voo, R. V. D.: Insights into the mechanism for orogen-related carbonate remagnetization from growth of authigenic Fe-oxide: A scanning electron microscopy and rock magnetic study of Devonian carbonates from northern Spain, J. Geophys. Res.-Solid Earth, 107(B4), 14–14, 2002.

Weil, A. B., Gutiérrez-Alonso, G. and Conan, J.: New time constraints on lithospheric-scale oroclinal bending of the Ibero-Armorican Arc: a palaeomagnetic study of earliest Permian rocks from Iberia, J. Geol. Soc., 167(1), 127–143, doi:10.1144/0016-76492009-002, 2010.

Weil, A.B., Yonkee, A., and Sussman, A.: Reconstructing the kinematics of thrust sheet rotation: a paleomagnetic study of Triassic redbeds from the Wyoming Salient, U.S.A., *GSA Bulletin*, 122, ½ 2-23, 2010.

Weil, A. B. B., Gutiérrez-Alonso, G., Johnston, S. T. and Pastor-Galán, D.: Kinematic constraints on buckling a lithospheric-scale orocline along the northern margin of Gondwana: A geologic synthesis, Tectonophysics, 582, 25–49, doi:10.1016/j.tecto.2012.10.006, 2013.

Woodcock, N. H., Soper, N. J. and Strachan, R. A.: A Rheic cause for the Acadian deformation in Europe, J. Geol. Soc., 164, 1023–1036, 2007.

Yenes, M., Alvarez, F. and Gutierrez-Alonso, G.: Granite emplacement in orogenic compressional conditions: the La Alberca-Bejar granitic area (Spanish Central System, Variscan Iberian Belt), J. Struct. Geol., 21(10), 1419–1440, 1999.

Yonkee, A., Weil, A.B., and Sussman, A.: Reconstructing the kinematic evolution of curved mountain belts: internal strain patterns in the Wyoming Salient, Sevier thrust belt, U.S.A., *GSA Bulletin*, 122, ¹/₂, 24-50, 2010.

Yonkee, A. and Weil, A. B.: Quantifying vertical axis rotation in curved orogens: Correlating multiple data sets with a refined weighted least squares strike test, Tectonics, 29(3), 31 PP.-31 PP., doi:10.1029/2008TC002312, 2010b.

Zegers, T. E., Dekkers, M. J. and Bailly, S.: Late Carboniferous to Permian remagnetization of Devonian limestones in the Ardennes: Role of temperature, fluids, and deformation, J. Geophys. Res. Solid Earth, 108(B7), n/a-n/a, doi:10.1029/2002JB002213, 2003.

Captions

Fig. 1 Simplified paleogeographic map of the Variscan-Alleghanian orogeny prior to the Jurassic break-up of Pangea, with the major orogenic curves labeled. Note, this map represents

Iberian outcrops without taking into account of the post-Jurassic Alpine deformation (see text for details e.g. Gong et al., 2008; Pastor-Galán et al., 2018). Slightly darker colors in the Variscan belt indicate present-day European and African outcrops (modified after Martínez-Catalán et al., 2009; Weil et al., 2013).

Fig. 2 A) Present-day configuration of the tectonostratigraphic zones of the Iberian
Variscides and theirits major structures. White areas represent Mesozoic and Cenozoic cover.
B) Three competing geometric proposals for the Central Iberian curve. 1) A disharmonic
curvature, up to 160° at the outer arc but much less pronounced at the inner arc (Aerden, 2004);
2) A harmonic, but more open curvature as suggested by Martínez Catalán (2012); 3) aAn
isoclinal curvature model (Shaw et al., 2012). C) Distribution of the E1 (in-migmatitic domes)
and C3 to post-C3 granitoids in the NW of Iberia (modified from Pastor-Galán et al., 2016)

Fig. 3 A) Stratigraphic synthesis of the Gondwanan platform series in NW Iberia. Cantabrian Zone columns are after Aramburu et al., 2002; Bastida, 2004; Murphy et al 2008; Pastor-Galán et al., 2013a; 2013b. Iberian Range follows Gozalo et al., 2008; Mergl and Zamora, 2012 and Calvín and Casas, 2014. West Asturian-Leonese Zone stratigraphy is after Pérez-Estaún, 1990; Marcos, 2004; Martínez-Catalán et al., 2004a; Gutiérrez-Marco et al., 2019. Central Iberian Zone follows Díez-Balda,1986; Valladares et al., 2000; Díez Montes, 2007; Martínez-Catalán et al., 2004b; del Moral and Sarmiento, 2008; García-Arias et al., 2018. B) Ordovician paleocurrents orientations, modified from Shaw et al., (2012). C) Schematic basin architecture inferred from the stratigraphic compilation.

Fig. 4 A) The kinematic evolution of the Cantabrian Orocline in its core, the Cantabrian Zone, inferred from total least squares (TLS) orocline tests (Pastor-Galán et al. 2017). B) Shows there events (Strike) tests used to constrain the kinematics of the Cantabrian Orocline. The Ordovician paleocurrents, which predate any orogenic movement, recorded the complete vertical-axis rotation history and yields a slope (m) of ~1. The Moscovian paleomagnetic data (from Weil et al., 2013; Pmag.), which postdates the main orogenic phases (C1, C2 and E1) and is is coeval with C3, shows a slope of ~1. The Gzhelian joint sets (from Pastor-Galán et al., 2011) orocline test shows a slope of ~0.5, which indicates that halfpart of the orocline was already formed between at ~304 Ma and 300 Ma.

Fig. 5 A) Aeromagnetic map of Spain (Ardizone et al., 1989, for Spain and the World Digital Magnetic Anomaly Map (WDMAM project) and Portugal (modified from Martínez Catalán, 2012 and Martínez Catalán et al., 2015), showing the possible trace of the Central Iberian curve. B) Bouguer anomalies of the Iberian Peninsula, modified from Ayala et al., 2016. Gravity anomalies do not reflect the geometry of the Cantabrian Orocline nor the Central Iberian curve. but are related to the Cenozoic Alpine lithospheric structure.

Fig. 6 Paleomagnetic studies related to the Cantabrian Orocline and the Central Iberian curve: (1) Synthesis of paleomagnetism in the core of the Cantabrian Orocline (see Weil et al., 2013); (2) Permian (eP) components synthesized in Weil et al. (2010); (3) Ordovician volcanics and limestones (Laquiana) in the boundary between the West Asturian-Leonese and Central Iberian Zones (Fernández-Lozano et al., 2016); (4) Devonian sedimentary sequences and Permian subvolcanics in the Iberian ranges (Pastor-Galán et al., 2018); (5) Permian dykes and sills (Calvín et al, 2014); (5) Anatectic granites (E1) and mantle derived granitoids (C3) from Tormes Dome and Central System (Pastor-Galán et al., 2016); (6) Cambrian limestones from Tamames (N) and los Navalucillos (S) (Pastor-Galán et al., 2015a); (7) Ediacaran-Early Cambrian sedimentary rocks in the southern sector of the Central Iberian Zone (Pastor-Galán et al., 2017b); (8) Almadén volcanics from the Central Iberian Zone (Perroud et al., 1991; Parés and van der Voo, 1992; Leite Mendes et al. in press) and Volcanic rocks from southern Ossa Morena and the South Portuguese Zone (Leite Mendes et al, in press)

Fig. 7 Magnetization components with a positive reversal test in the extensional anatectic granites of Tormes (A) and Martinamor Domes (B). This component is interpreted as primary with a magnetization age of >318 Ma (Pastor-Galán et al., 2016). C) Distribution of directions and VGPs and statistical parameters from both domes combined.

Fig. 8 Cartoon depicting the different vertical_-axis rotation events that occurred in the Cantabrian Zone and the Iberian Range, modified from Pastor-Galán et al. (2018). (A) Original quasilinear Variscan Orogenic belt, B) Formation of the Cantabrian Orocline around the Carboniferous–Permian boundary after a ~70° counterclockwise rotation in the Southern branch of the Cantabrian Zone and the Iberian Range. This rotation matches the rotation for the Cantabrian Orocline, see the fit of the Iberian Range Component #2 in the orocline test for the Cantabrian Zone (below). C) Post_Permian (Cenozoic) rotation of ~22° clockwise (CW) likely produced by differential shortening during the Alpine orogeny (Izquierdo-Llavall et al., 2018). Below, the global magnetic polarity time_scale for the Pennsylvanian and Cisuralian (following Ogg et al., 2016). TLS = Total Least Squares. Note that once the 22° CW rotation in the Iberian Range is corrected, components #2, #1, and P fit as expected with the APWP for the southern limb of the <u>Cantabrian Orocline_orocline_orocline</u> (Pastor-Galán et al., 2016).

Fig. 9 Compilation of the directional distributions and average declinations with parachute of confidence (Δ Declination) in sites around the Central Iberian curve (see Fig. 6). The results show general CCW rotations in contrast to the expected CW <u>rotations</u> if the Central Iberia curve formed by vertical-axis rotations (see text). Results are compared with the expected

declinations if those sites were part of the Cantabrian Orocline following the methodology described in Pastor-Galán et al., (2017b).

Fig. 10 Orocline test of the Cantabrian Orocline (Weil et al., 2013) compared with the magnetizations found in the adjacent Laurussian segments of the orogen: Ireland (Pastor-Galán et al., 2015b) and the South Portuguese Zone (Leite Mendes et al., in press)

Fig. 11 Structural analysis of mullions in the Central Iberian Zone (after Pastor-Galán et al., 2019b) A) Photograph (with card scale, 10cm) of a bedding plane surface showing the mullions and photograph analysis. B) Interpretation of the outcrop with fold axis traces depicting the deformation phase responsible for each structure. C) Result of retro-deformation of mullions in the Mogadouro road section modified from Pastor-Galán et al. (2019b). Unfolding the effects of D3 on D2 mullions. Unfolding the effects of C3 and E1 in C1 mullions.

Fig. 12Fig. 11 Pioneering hypotheseis for the Central Iberian curve. Note that none of them fulfill the most recent geometric and kinematic criteria. A) Simplified ribbon continent model after Johnston et al. (2013) and Shaw and Johnston (2016). B) Dextral mega-shear model from Martínez-Catalán (2011). C) Kinematic model with indentation and left-lateral shearing after Simancas et al. (2013)

Fig. 13Fig. 12 Preliminary kinematic proposal for the Iberian Variscides. A) Precollisional stage after the opening of the Galicia Tras-os-Montes restricted seaway (e.g. Pin et al., 2002; Gutiérrez-Alonso et al., 2008a; Arenas et al., 2016). The irregular shape of the margin and the younging westwards deformation front (e.g. Daleyer et al., 1997) resulted in tectonic escape towards the still open Rheic Ocean (e.g. Braid et al., 2011; Murphy et al., 2016). B) After closure of the Rheic Oecean, C1 and C2 structures formed. The Galicia Tras-os-Montes was emplaced orogen parallel (e.g. Martínez-Catalán et al., 1990; Dias da Silva et al., in pressDias da Silva et al., 2020), preserving the shape of the seaway (-i.e. a primary arc). C) The gravitational collapse of the orogen produced widespread anatexis and folding interference in the hinterland and the emplacement of the foreland fold-and-thrust belt. D) At Pennsylvanian times a change in the far-field stress buckled the Variscan belt around a vertical axis (see Gutiérrez-Alonso et al., 2008; Weil et al., 2013; Pastor-Galán et al., 2015a for details), creating new interference patterns and a lithospheric-scale response (see Gutiérrez-Alonso et al., 2004, 2011a; Pastor-Galán et al., 2012a). E) When the orocline became too tight to keep rotating, new cross-cutting brittle structures (C4) formed and minor extensional collapse (E2) occurred (e.g. Fernández-Lozano et al., 2019; Dias da Silva et al., in pressDias da Silva et al., 2020).