

The enigmatic curvature of Central Iberia and its puzzling kinematics

Daniel Pastor-Galán^{1,2,3}, Gabriel Gutiérrez-Alonso^{4,5}, Arlo B. Weil⁶

¹Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

²Department of Earth Science, Tohoku University

³Center for Northeast Asian Studies, Tohoku University, 41 Kawauchi, Aoba-ku, Sendai, 980-8576, Japan. pastor.galan.daniel.a8@tohoku.ac.jp

⁴Dept. of Geology. Faculty of Sciences. University of Salamanca. Plaza de la Merced s/n. 38007, Salamanca (Spain). gabi@usal.es

⁵Geology and Geography Department, Tomsk State University, Lenin Street, 36, Tomsk 634050 Russia

⁶Department of Geology, Bryn Mawr College, PA, USA 19010

Abstract

The collision between Gondwana and Laurussia that formed the latest supercontinent, Pangea, occurred during Devonian to Early Permian times and resulted in large-scale orogeny that today transects Europe, northwest Africa and eastern North America. This orogen is characterized by an 'S' shape corrugated geometry in Iberia. The northern curve of the corrugation is the well known and studied Cantabrian (or Ibero-Armorican) Orocline and is convex to the east and towards the hinterland. Largely ignored for decades, the geometry and kinematics of the southern curvature, known as the Central Iberian curve, are still ambiguous and hotly debated. Despite the paucity of data, the enigmatic Central Iberian curvature has inspired a variety of kinematic models that attempt to explain its formation with little consensus. This paper presents the advances and milestones in our understanding of the geometry and kinematics of the Central Iberian curve from the last decade, with particular attention to structural and paleomagnetic studies.

When combined, the currently available datasets suggest that the Central Iberian curve did not undergo regional differential vertical-axis rotations during or after the latest stages of the Variscan orogeny, and did not form as the consequence of a single process. Instead, its core is likely a primary curve (i.e. inherited from previous physiographic features of the crust) whereas the curvature in areas outside the core are dominated by folding interference during the Variscan orogeny or more recent Cenozoic (Alpine) tectonics.

31 **Keywords**

32 **Central Iberia Curve, Variscan orogen, Iberia, Cantabrian Orocline, Curved orogens,**

33 **Pangea**

1 Introduction

Mountain belt systems are the most striking product of plate tectonics. In addition to their astonishing visual effect, marking the locations where ancient and modern plates collided, orogenic belts often preserve a variety of rocks that have the potential to illuminate the entirety of the systems pre- and syn-orogenic history. One of the most striking characteristics of the majority of Earth's orogens are their curvature in plan-view (e.g. van der Voo, 2004; Marshak, 2004; Rosenbaum, 2014). The degree of orogenic curvature may range from a few degrees of deflection in structural trend (e.g. Kopet Dag, Iran), to 180° of arc curvature (e.g. Kazakhstan arc and the Carpathians). The kinematics, structural and geodynamic implications of these systems are as varied as their geometries (Marshak, 2004; Weil and Sussman, 2004; Johnston et al., 2013). For example, some orogenic curvatures are hypothesized to be the consequence of physiographic features of the basement that pre-date orogen formation, such as irregular basin architectures or plate margin salients and recesses (e.g. Jura mountains, Hindle et al., 2000), which then control the growth geometry of the ensuing orogen. These systems are known as primary arcs and reflect pre-orogenic geometries and show no significant or systematic vertical-axis rotations along their structural length. On the other hand, oroclines, as classically defined by Carey in 1956, involve systematic differential vertical-axis rotations subsequent to initial orogenic shortening: different sectors of an orogen rotate by variable magnitudes or in opposite directions (e.g. Li et al., 2012). Rotations in **Oroclines** may occur at a range of scales, from thrust emplacement at upper crustal levels (e.g. Izquierdo-Llavall et al., 2018), up to a lithospheric-scale vertical-axis folding (e.g. Li et al., 2018). They can occur as single curves (e.g. Maffione et al., 2009), coupled curves (Johnston, 2001), or in trains of curves (Li and Rosenbaum, 2014). Oroclines can form during the main orogenic building event, known as progressive oroclines (Johnston et al., 2013; e.g., the Wyoming salient, Yonkee and Weil, 2010, and Weil et al., 2010) or in a subsequent tectonic pulse, so-called secondary oroclines (Weil and Sussman, 2004). Understanding the kinematics and mechanisms of curvature formation in mountain belts is a critical step to understanding orogenesis in 4D and to evaluate their geodynamic consequences and paleogeographic implications.

The Variscan-Alleghanian orogeny resulted in the suturing of Gondwana and Laurussia during Devonian-Carboniferous times, and ultimately led to the formation of the supercontinent, Pangea. This long and sinuous orogen runs for >8000 km along strike and is ca. 1000 km wide, transecting across Europe, to northwest Africa and into eastern North America. The final stages of Pangea amalgamation (e.g. Nance et al., 2010) modified the Western Europe sector of the

belt into its characteristic sinuous shape, which today traces at least one, and perhaps four arcs from Poland to Brittany, and then across the Bay of Biscay (Cantabrian Sea) into Iberia, where the system is today truncated by the Betic Alpine orogeny in southeast Iberia (Fig. 1; e.g. Weil et al., 2013). The southern truncation of the Variscan in Europe hinders a precise correlation with equivalent age outcrops in NW Africa.

Within the Iberian Peninsula, the orogen is characterized by two large-scale curves (Fig. 2): (1) to the north is the well studied and nearly 180° secondary orocline, the Cantabrian (a.k.a. Ibero-Armorican) Orocline, which buckled a segment of the Variscan belt from ~315 to ~290 Ma (e.g. Weil et al., 2019 and references therein); and (2) to the south is a curve with disputed magnitude and kinematics, and is usually referred to as the Central Iberian curve/orocline or Castillian bend (Martínez-Catalán et al., 2015). Though there remains tremendous uncertainty on the geometry and kinematics of the Central Iberia curve, multiple hypotheses exist as to its nature, and disagreements continue on its importance in the tectonic evolution of Europe during the waning stages of Paleozoic global supercontinent construction. The diversity of author's interpretations of the Central Iberian curve range from a nonexistent structure (Dias et al., 2016), to being one of the most important pieces to our understand of the late Carboniferous and Permian geodynamics of the Iberian Variscan system (e.g. Martínez-Catalán et al., 2011; 2014).

This paper reviews the most recent advances on the geometry and kinematics of the Central Iberia curve, synthesizing what we know and what we don't, and ending with a discussion of the main unsolved issues. We hope that this paper fosters novel studies that will lead to a better understanding of when and which mechanisms acted in the aftermath of the Variscan-Alleghanian orogeny.

2 The long and winding orogen

The Variscan (Europe-NW Africa)-Alleghanian (North America) orogeny is a continental-scale tectonic system (1000 km wide and 8000 km long) that sutured Gondwana and Laurussia together, forming the supercontinent Pangea (e.g. Domeier and Torsvik, 2014; Edel et al., 2018; Pastor-Galán et al., 2019a). The fragments of this system are now dispersed over three continents, Europe, Africa and North America due to the Mesozoic break-up of Pangea (Buiter and Torsvik, 2014; Keppie, 2015). This orogen formed as a consequence of a long and protracted tectonic history that involved several different events, from initial convergence (ca. 420 Ma; e.g. Franke et al., 2017), to the consumption of multiple putative oceanic tracts and/or basins that existed between Gondwana and Laurussia (ca. 280 Ma; e.g. Kirsch et al., 2012).

The Variscan-Alleghanian orogen itself represents the closing of at least one major ocean, the Rheic (e.g. Nance et al., 2010), whose axial ridge likely failed or subducted at ca. 395 Ma along its paleo-northern margin (e.g. Woodcock et al., 2007; Gutiérrez-Alonso et al., 2008a). Perhaps the orogeny involved other large oceans (Stampfli and Borel, 2002; Franke et al., 2017; 2019), but most surely involved several minor seaways and basins that existed between Gondwana, Laurussia, and several intervening micro-continents (e.g. Azor et al., 2008, Dallmeyer et al., 1997; Kroner and Romer, 2013; Díez-Fernández et al., 2016; Pérez-Cáceres et al., 2017). The final continent-continent collision began after closure of all oceans and intervening seaways. The commencement of this deformation was diachronistic and became progressively younger westwards (in present-day coordinates): with Devonian continent-continent collisions along the eastern boundary, progressing to earliest Permian ages in the westernmost sector (McWilliams et al., 2013; Chopin et al., 2014; López-Carmona et al., 2014; Franke et al., 2017).

The present-day geometry of the Variscan-Alleghanian systems has a contorted trace (Fig. 1). In Europe, from east to west, the trend starts with a prominent curve around the Bohemian massif (e.g. Tait et al., 1996), followed by a deflection in the Ardennes-Bravant (e.g. Zegers et al., 2003). In Brittany the outer curvature of the Cantabrian or Ibero-Armorican orocline begins (e.g. van der Voo et al., 1997), and wraps nearly 180° around across the Bay of Biscay as it turns in NW Iberia. The Central Iberian curve marks the final concave to the west curve (in present-day coordinates) and is the focus of this paper (e.g. Aerden, 2004; Martínez Catalán, 2011; Shaw et al., 2012). The orogen continues in North America where, from north to south, it has salients and recesses that undulate back and forth from Atlantic Maritime Canada (e.g. O'Brien, 2012) down along the Pennsylvanian and Alabama curves (e.g. Thomas, 1977).

Interpretation on the origin of these curvatures varies widely. The curvatures in North America are argued to be the result of a preexisting irregular margin of Laurentia due to the break-up of the Rodinia supercontinent, which resulted in the formation of orogenic salients and recesses during subsequent Appalachian collision (e.g. Rankin, 1976; Thomas, 1977, 2004). In this case, vertical-axis rotations affected only the upper crustal levels during orogenesis (e.g. Marshak, 1988; Bayona et al., 2003; Hnat and van der Pluijm, 2011). In Europe, the Bohemian and Ardennes-Bravant massif curvatures have poor kinematic constraints. In the Bohemian Massif, some suggest secondary rotations that formed an orocline (Tait et al., 1996), while others suggest little to no vertical-axis rotations and a primary arc (Chopin et al., 2012). The Ardennes-Bravant Massif record some vertical-axis-rotations (e.g. Molina-Garza and Zeijderveld, 1996), but it is unclear if these are a response to progressive or secondary oroclinal bending, or whether rotations only affected the upper crust. The most outstanding example of

Variscan-Alleghanian orogen curvature is exposed in the Iberian Massif, with the Cantabrian Orocline and the coupled curvature of Central Iberia.

2.1 Two of us: The Variscan orogen in Iberia

The western half of the Iberian Peninsula constitutes the Iberian massif, one of the largest exposures of the Variscan orogen and the only place that contains an almost continuous cross section of the orogen (Fig. 2; e.g. Lotze 1945, Julivert 1974, Pérez-Estaún et al., 1991; Ayarza et al., 1998; Simancas et al., 2003; Ribeiro et al. 2007, Martínez Catalán et al., 2014, 2019). The majority of the Iberian Massif contains Gondwanan affinity rocks (e.g. Murphy et al., 2008; Pastor-Galán et al., 2013a; Gutiérrez-Marco et al., 2017) and likely represents a proximal piece of the Gondwana margin until its final amalgamation with Pangea (e.g. Pastor-Galán et al., 2013b). Owing to the stratigraphic, structural and petrological styles, the Iberian Massif has been traditionally divided into six tectonostratigraphic zones (Fig. 2; Lozte, 1945; Julivert, 1971): (1) Cantabrian Zone represents a Gondwanan thin-skinned foreland fold-and-thrust belt. It has overall low-grade internal deformation and metamorphism, and represents shortening that occurred during Mississippian times (e.g. Marcos and Pulgar, 1982; Pérez Estaún et al., 1988; Gutiérrez-Alonso 1996; Alonso et al., 2009; Pastor-Galán et al., 2009; 2013b). (2) The West-Asturian Leonese Zone represents a metamorphic fold-and-thrust belt with barrovian metamorphism that collapsed coevally with thrust emplacement onto the Cantabrian Zone (e.g. Martínez-Catalán et al., 1992; Alcock et al., 2009; Martínez-Catalán et al., 2014). (3) The Central Iberian Zone represents the Gondwanan hinterland with Barrovian and Buchan metamorphism and is intruded by igneous rocks of various ages (e.g. Macaya et al., 1991; Díez Balda, 1995; Gutiérrez-Alonso et al., 2018). (4) The Ossa-Morena Zone represents the most distal zone of the Gondwana platform, and is characterized by a metamorphic fold-and-thrust belt with dominantly sinistral displacement (e.g. Robardet and Gutiérrez-Marco, 2004; Quesada, 2006). (5) The Galicia-Tras-os-Montes Zone represents a far travelled allochthonous terrane that contains high pressure units and relics of oceanic-like crust (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). (6) The South Portuguese Zone represents a foreland fold-and-thrust belt with little internal deformation and metamorphism with Avalonian affinity and a strong left-lateral component of shear (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2016; Oliveira et al., 2019). Geographically, the external zones of the Gondwana margin are nested to the north into the core of the Cantabrian Orocline, whereas the hinterland zones are to the west and center of the massif (Fig. 2; e.g. Díaz Balda, 1995; Azor et al., 2019). The southwestern-most extent of Iberia contains a putative suture of the Rheic ocean, as well as a piece of the

Laurussian margin fold-and thrust belt, today preserved in the South Portuguese Zone (e.g. Pereira et al., 2012, 2017; Oliveira et al., 2019).

The Gondwanan **autochthon** stratigraphy (Cantabrian, West Asturian-Leonese, Central Iberian and Ossa Morena Zones) consist of a Neoproterozoic arc and back-arc basin (e.g. Fernández-suárez et al., 2014), which evolved to a rift-to-drift Cambrian to Early Ordovician sequence and then to an Ordovician to Late Devonian passive margin basin sequence (e.g. Sánchez-García et al., 2019; Gutiérrez-Marco et al., 2019; Gutiérrez-Alonso et al., submitted). Overall the system transitioned from a relatively isolated Early Cambrian continental, to a restricted marine basin, to development of an open marine platform that was locally punctuated by magmatism (e.g. Gutiérrez-Alonso et al., 2008b; Palero-Fernández, 2015). The Ossa Morena zone represents the outermost platform, followed by an intermediate platform characterized by an asymmetric horst (Central Iberian Zone) and graben (West-Asturian Leonese Zone), which ends in the innermost shelf environment of the Cantabrian zone (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The differences between the West Asturian-Leonese and Central Iberian Zone are mainly deeper vs. shallower sedimentary facies (respectively) and a local Lower Ordovician unconformity in the Central Iberian Zone (Toledanian, e.g. Álvaro et al., 2018) that places Lower Ordovician strata atop pre-Cambrian to Cambrian rocks (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The Central Iberian Zone is divided into two domains: (1) The Ollo de Sapo domain, which contains abundant Lower Ordovician **calc-alkaline magmatism** (e.g. Díez Montes, 2006; Gutiérrez-Marco et al., 2019); and (2) the 'Schistose-greywacke Domain' characterized by a predominance of outcrops of Neoproterozoic to Lower Cambrian sedimentary rocks (e.g. Gutiérrez-Marco et al., 2019 and references therein).

The Galicia Tras-os-Montes Zone (Farias et al., 1987) is a complex structural stack including a basal schistose unit (Parautochthon; Dias da Silva et al., in press) structurally overlain by mafic rocks with an oceanic-like signature and other far-traveled rocks under high-pressure metamorphism (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). The oceanic rocks of this zone are classically interpreted as a Rheic Ocean suture (e.g. Martínez Catalán et al., 2009). Recent interpretations support its origin as a minor oceanic basin or seaway within the realm of Gondwana (e.g. Pin et al., 2002; Arenas et al., 2016).

The South Portuguese Zone constitutes the Laurussian foreland fold-and-thrust belt in the Iberian Variscides (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2017). It contains three units: (1) the Pulo de Lobo, a low grade metamorphic accretionary prism with clastic sedimentary rocks and basalts with MORB signature (e.g. Azor et al., 2019; Pérez-Cáceres et al., this volume); (2) The Iberian Pyrite Belt, which is a world class volcanogenic massive sulfide

deposit formed between 390 and 330 Ma (e.g. Oliveira et al., 2019a; 2019b); and (3) the Baixo Alentejo Flysch, which is located to the southwest and is a syn-orogenic composite turbiditic sequence with ages from ~330 to ~310 Ma (Oliveira et al., 2019b). The boundary between the South Portuguese and Ossa Morena zones is a sinistral shear zone (so-called Beja-Acebuches, Quesada and Dallmeyer., 1994; Pérez-Cáceres et al., 2016) that contains a strongly deformed amphibolitic belt with oceanic affinity (Munha et al., 1986; Munha, 1989; Quesada et al., 2019). This belt potentially represents dismembered relics of the Rheic ocean and/or a subsidiary seaway that opened during a Variscan transtension event in SW Iberia (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019).

Finally, Paleozoic rocks occur sporadically within the Alpine Betic chain. Their lithological monotony, paucity of fossils, and the intensity of deformation and metamorphism during Alpine orogeny, make recognizing the original features of the different successions challenging (e.g. Martín-Algarra et al., 2019). Some faunal and detrital zircon studies suggest that the Paleozoic outcrops in the Betics may be similar to the most seaward realms of the Gondwanan platform (i.e., the Cantabrian Zone; e.g. Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018). Following the latest plate reconstructions of the Mediterranean during Meso-Cenozoic times, the Paleozoic units of the Betic-Rif chain may have been located proximal to the present-day position of the Balearic Islands (van Hinsbergen et al., 2020).

The Variscan orogen in Iberia shows multiple deformation, metamorphic, and magmatic events (e.g. Martínez-Catalán et al., 2014; Azor et al., 2019; Fig. 2) that evolved diachronously from the suture towards the external zones (Dallmeyer et al., 1997): (1) An initial continent-continent collision began ca. 370-365 Ma, which produced high pressure metamorphism (e.g. Lopez-Carmona et al. 2014). (2) Between 360 and 330 Ma a protracted shortening phase occurred, frequently divided into main phases C1 and C2, that were accompanied by Barrovian type metamorphism (e.g. Dias da Silva et al., in press) and plutonism at ~340 Ma (e.g. Gutiérrez-Alonso et al., 2018). (3) An extensional collapse event, so-called E1, occurred at ~333-317 Ma, which formed core-complexes and granitic domes in the Central Iberian and West Asturian-Leonese zones (Fig. 2C; e.g. Alcock et al., 2009; Díez-Fernández and Pereira, 2016; López-Moro et al., 2018). This event is coeval and genetically linked to the formation of the foreland fold-and-thrust-belt of the Cantabrian Zone (e.g. Gutiérrez-Alonso, 1996). (4) A late Carboniferous shortening event (C3) occurred ca. 315-290 Ma and is argued to have resulted in the formation of the Cantabrian Orocline and was accompanied by the intrusion of mantle derived granitoids (Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a, 2011b; Pastor-Galán et al., 2012a). (5) A final early Permian extensional event (E2), mostly found in the Central Iberian

Zone, resulted in the formation of core complexes and regional doming (Dias da Silva et al., in press). (6) A final shortening event (C4), possibly coeval with E2, resulted in widespread brittle deformation (e.g. Azor et al., 2019; Fernández-Lozano et al., 2019).

In SW Iberia, the aforementioned Variscan deformation events are characterized by a dominant sinistral component, which contrasts with the general dextral component recognized in most other regions of the orogen (e.g. Martínez Catalán et al., 2011; Gutiérrez-Alonso et al., 2015). Early collisional structures (C1) formed NE-vergent recumbent folds in the southernmost Central Iberian Zone and SW-vergent folds and thrusts in the Ossa Morena and South Portuguese zones. This phase continued with a transtensional event that heterogeneously extended the continental lithosphere (e.g. Pérez-Cáceres et al. 2015). Coevally, an important extension-related magmatic event happened, perhaps assisted by a plume-type mantle (Simancas et al. 2006) or a slab break-off (Pin et al. 2008). After this transtensional event, significant left-lateral transpression occurred forming the extensive shear zones to the north and south of Ossa Morena Zone (Fig. 2B), which accommodated the majority of the transcurrent motion. However, left-lateral displacements are observed all along the Ossa Morena and South Portuguese zones. Pérez-Cáceres et al. (2016) estimated over 1000 km of collisional convergence in SW Iberia, most of which corresponds with left-lateral displacements parallel to terrane boundaries.

3 Synthesis on the Geometry and Kinematics of the Cantabrian Orocline

Understanding the geometry, kinematic evolution and mechanics of curved mountain systems is crucial to developing paleogeographic and tectonic reconstructions (e.g. Marshak, 2004; Van der Voo, 2004; Li et al., 2012; van Hinsbergen et al., 2020). Introduced by Carey (1955 p.257), an orocline (from Greek *opos*, mountain, and *κλινο*, bend) is “...an orogenic system, which has been flexed in plan to a horse-shoe or elbow shape.” Although sometimes used in the literature as a geometric description of any orogenic curvature, herein orocline is strictly used as a the term for map-scale bends that underwent vertical-axis rotations (Weil and Sussman, 2004; Johnston et al., 2013; Pastor-Galán et al., 2017a). The kinematic classification of curved mountain belts (Weil and Susman, 2004; Johnston et al, 2013) distinguishes two end members: (1) Primary orogenic curves, which describe those systems in which curvature is a primary feature of the orogen and formed without significant or systematic vertical-axis rotations, and (2) Secondary oroclines, where orogenic curvature was acquired due to vertical-axis rotations subsequent to primary orogenic building. Those systems whose curvature is the

product of vertical-axis rotation during the primary orogenic pulse and/or only a portion of the observed curvature is secondary are progressive oroclines.

The orocline test (or strike test), evaluates the relationship between changes in regional structural trend (relative to a reference trend for an orogen) and the orientations of a given geologic fabric element or magnetization (relative to a reference direction). In terms of evaluating developmental kinematics, the most relevant geologic marker is paleomagnetic declination, which can be used to quantitatively evaluate total and systematic rotations as a function of along-strike variability. Once acquired, data is plotted on Cartesian coordinate axes with the strike (S) of the orogen (relative to a reference) along the horizontal axis, and the fabric azimuth (F, relative to a reference) along the vertical axis. The test originally used a basic least-squares (OLS) regression (Schwartz and Van der Voo, 1983) to estimate the slope (coded m in formulas), ideally between 0 and 1, which then is interpreted with respect to vertical-axis kinematics. More recently, Yonkee and Weil (2010b) and Pastor-Galán et al. (2017a) introduced more robust statistics to estimate the slope and its uncertainty, considering and propagating errors of the input data. Primary orogenic bends show no change of paleomagnetic declination orientations with varying structural trend, and therefore the slope is expected to be 0. In progressive oroclines, the declination variation records some fraction of the total observed orogenic strike variability, and thus the slope would range between 0 and 1, depending on the amount of primary curvature. Secondary oroclines are those in which the paleomagnetic vectors record 100% of the rotation, yielding slopes of 1, meaning that the orogenic system must have started as a roughly linear system that then underwent secondary vertical-axis rotations until its present-day curvature was acquired. The slope obtained with the orocline test can only be confidently interpreted when the chronology of fabric formation is well known.

The trend of the Variscan belt in Iberia follows a sinuous “S” shape that is especially prominent in the northwest region of the Iberian Peninsula, and then becomes more subtle due to the predominance of younger cover sequences in the central and eastern regions of the peninsula (Fig. 1 and 2). This dramatic geometry has stimulated a century long scientific debate as to its origin (e.g. Suess, 1892; Staub, 1926; Martínez Catalán et al., 2015). To the north and convex to the west is the Cantabrian Orocline, and to the center-south and convex to the east is the Central Iberian curve. The overall trend of the Cantabrian Orocline starts in Brittany (France) and southern England and then curves through the Bay of Biscay and then south into central north Iberia (Fig. 1, 2 and 4). The Cantabrian Orocline (also known as Ibero-Armorican Orocline/Arc, Asturian Arc or Cantabrian-Asturias Arc) is arguably the first curved orogen that was scientifically described, recognized by the change in structural trend of mapped thrusts and fold

axes (Schultz, 1858, Barrois, 1882, Suess, 1892). The Cantabrian Orocline traces an arc with a curvature close to 180° within the central Cantabrian Zone (the Gondwanan foreland in Iberia, fig. 2), and opens to approximately 150° as one moves to the outer arc reaches (Fig. 1). At the crustal-scale, the Cantabrian Orocline represents a first order vertical-axis buckle fold in plan-view that refolds pre-existing Variscan structures (e.g. Julivert and Marcos, 1973; Weil et al., 2001). The inner arc of the orocline, or the Cantabrian Zone is characterized by tectonic transport towards the core of the orocline, i.e., the orocline has a contractional core, where low finite strain values and locally developed cleavage occur (Pérez-Estaún et al., 1988; Gutiérrez-Alonso, 1996; Pastor-Galán et al., 2009). Within the inner core a variety of structures record non-coaxial strain, which produced complex interference folds and rotated thrust sheets (e.g. Julivert and Marcos, 1973; Julivert and Arboleya, 1984; Pérez-Estaún et al., 1988; Aller and Gallastegui, 1995; Weil, 2006, 2013; Pastor-Galán et al., 2012b; Shaw et al., 2015; 2016a; Del Greco et al., 2016). In contrast, the outer arc shows a ca. 150° interlimb angle vertical-axis fold that was accommodated by significant shearing, both dextral and, in lesser amounts, sinistral penecontemporaneous to vertical-axis rotation (Gutiérrez-Alonso et al., 2015). Weil et al. (2013, 2019) extensively review the geometry of the Cantabrian Orocline.

All kinematic data studied so far support a model in which the Cantabrian Orocline formed due to secondary vertical-axis rotation in a period of time younger than 315 Ma and older than 290 Ma. Overall, the southern limb of the orocline rotated counterclockwise (CCW) and the northern limb clockwise (CW; Fig. 4). Orocline formation happened subsequent to the main shortening phases of the orogen (C1 and C2) and late-stage orogenic collapse (E1), and therefore, it is an ideal example of a secondary orocline in the strictest sense. Development of the Cantabrian Orocline requires the existence of a roughly linear orogenic belt during early Variscan closure of the Rheic Ocean (with a roughly N-S orientation in present-day coordinates), which was subsequently bent in plan-view into an orocline during late stages of Pangea amalgamation. Such interpretation is grounded in paleomagnetic studies (e.g. Hirt et al., 1992; Parés et al. 1994; Stewart, 1995; van de Voo et al., 1997; Weil, 2006; Weil et al., 2000; 2001; 2010), along with important contributions from structural (e.g. Gutiérrez-Alonso 1992; Kollmeier et al., 2000; Merino-Tomé et al., 2009; Pastor-Galán et al., 2011; 2014; Shaw et al., 2015) and geochronological studies (e.g., Tohver et al., 2008; Gutiérrez-Alonso et al., 2015). Weil et al. (2013) provides a comprehensive review on the kinematic constraints, updated in 2017a by Pastor-Galán et al., and in 2019 by Weil et al.

4 The intriguing geometry of the Central Iberian curve

The more southern Central Iberian curve has a similar magnitude, but opposite curvature compared to the Cantabrian Orocline (Fig. 1 and 2B). This structure has been referred to as the Central Iberian curve, arc, bend or orocline. In this paper we use 'Central Iberian curve'. The other aforementioned terms involve still unknown parameters or are misleading: orocline imply kinematics (Weil and Sussman, 2004); bend refers to a mechanism of formation (e.g. Fossen, 2016); and arc could be ambiguous, since the term is commonly used for volcanic chains. This curvature was first described by Staub (1926) and was termed the Castilian bend. Continental drift pioneers paid some attention to Staub's description (e.g. Holmes, 1929; Du Toit, 1937), but the curved structure remained largely ignored for multiple decades (e.g. Martínez Catalán et al., 2015). The hypothesis of a large-scale curvature in Central Iberia made a comeback at the beginning of the 21st century with a study of Variscan porphyroblast kinematics across Iberia by Aerden in 2004. Since then, several attempts to unveil its geometry and kinematics have been made with contrasting results.

The elusive nature of the Central Iberian curve resides in the poor exposure of its putative hinge (Fig. 2). The hinge of the Cantabrian orocline crops out extensively and the changes in thrust and fold axes trend are observable at high-resolution from aerial photographs and are readily mapped using outcrop-scale observations. In contrast, the alleged hinge of the Central Iberian curve is largely covered by Mesozoic and Cenozoic basins (Fig. 2). The curvature is most recognizable at the boundary between the Galicia-Tras os Montes and Central Iberian zones (Fig. 2A; Aerden, 2004; Martínez Catalán, 2012). The thrust fault that bounds those areas traces close to a 180° of curvature and marks the emplacement of the most distal units. Before the revival of Staub's curved geometry along the entire Central Iberian Zone, there were several attempts to explain the curved shape of the Galicia Tras-os-Montes Zone. Some consider the Galicia Tras-os-Montes Zone a block that escaped during an early Variscan (C1) non-cylindrical collision, forming an extrusion wedge towards the areas undergoing lesser amount of shortening (Martínez-Catalán, 1990; Dias da Silva, 2015; in press); or alternatively a klippe of a larger allochthonous thrust sheet, product of an interference pattern between C2, E1 and C3 structures (e.g. Ries and Shackleton, 1971; Martínez Catalán et al., 2002; Rubio Pascual et al., 2013; Díez-Fernández et al., 2015).

In addition to the Galicia Tras-os-Montes Zone, other areas showing a certain degree of curvature are to the E and SE of the Central Iberian Zone. There, an approximately 20° change in strike of the Iberian ranges (NE Iberia, Fig. 2A) is observed, which represents the only known outcrop of the hinge of the Central Iberian curve's outer arc. The rest of the curvature has been

deduced with indirect observations leading to three competing geometric proposals for the Central Iberia curve (Fig. 2B). The main arguments used to constrain the geometry of the Central Iberian curve are: (1) the geometry of Galicia Tras-os-Montes folds and the orientation of observed garnet inclusion trails (Aerden, 2004; Fig. 2B-1); (2) aligned aeromagnetic anomalies and fold trends in the Iberian ranges and the E-SE Central Iberian Zone (Martínez-Catalán, 2012; Fig. 2A and 2B-2) and; (3) the regional distribution of paleocurrents recorded in Ordovician quartzites (Shaw et al., 2012; Fig. 2B-3 and 3). All proposed geometries share two features: (1) The curvature runs parallel to the Central Iberian Zone, and is located in the center-west of Iberia, and (2) all place the Galicia Tras-os-Montes Zone in the core of the curve with the curves axial trace cross-cutting the Morais Complex, a set of mafic and ultramafic rocks that is roughly circular in shape (Fig. 2B; Dias da Silva et al., in press).

Aerden (2004) compared the orientation of inclusions in metamorphic porphyroblasts across the Variscan allochthonous terranes of the NW Iberian Massif, and found that inclusion trails maintain a constant north–south orientation. Comparing such results with the trend of the Variscan fold axes in the central Iberian Zone (Fig. 2A) and a daring interpretation of the aeromagnetic anomalies of the Iberian Peninsula (Fig. 5A), Aerden suggested a geometry in which the Central Iberian curve was more prominent in the outer arc than in the inner arc (Fig. 2B-1). In Aerden's view the geometry of the Galicia Tras-os Montes Zone does not represent a large-scale curvature, but rather the original shape of the nappe, perhaps re-tightened during C3 deformation. In contrast, the Iberian Ranges and the SE Central Iberian Zone represent the more curved sector (Fig. 2B-1). In the model of Aerden (2004), the Ossa Morena and South Portuguese Zones are not part of the Central Iberian curvature.

Martínez-Catalán (2012) reinterpreted Aerden's analysis of aeromagnetic map data (Fig. 5A) and the interpretive structural trends of C1-C2 fold axes from Central Iberian Zone structures (Fig. 2A). In Martínez-Catalán's model, the Central Iberian curvature is a symmetric arcuate shape in which orogen trend changes equally in the inner and outer arc, and is comparable in size to the Cantabrian Orocline, but with opposite curvature and less shortening. This geometric model also excludes the Ossa Morena and the South Portuguese Zones as elements involved in the formation of the curvature (Fig. 2B-2).

Finally, Shaw et al. (2012) studied the orientation of paleocurrents in Ordovician Armorican Quartzite (e.g. Aramburu, 2002), which is one of the most prominent rocks exposed in Iberia (Fig. 3). The authors found that paleocurrents fanned outward with respect to the Cantabrian Orocline curve and are approximately perpendicular to the structural trend throughout the peninsula (Fig. 3). Shaw et al. (2012) assumed that the direction and sense of

paleocurrents were parallel throughout all zones, and concluded that the Central Iberia curve is a 'S' shape isoclinal structure similar in magnitude to the Cantabrian Orocline (Fig. 2B-3). It is unclear from the Shaw et al. (2012) model the involvement of the Ossa Morena and South Portuguese Zones in the overall curve (if any), nor the prospective location of the external zones of the orogen (Cantabrian Zone) with respect to the overall curvature.

5 Move over once, move over twice: Kinematic constraints

Late Variscan kinematic data (315-290 Ma; C3, E2, C4 phases) in the Central Iberian curve were scarce prior to revival of Staub's Central Iberian curve (e.g. Vergés, 1983; Julivert et al., 1983; Parés and van der Voo, 1992). More recently, a wealth of studies have been published on the kinematics of forming the Central Iberian curve (Fig. 2B), which are reviewed below.

5.1 Structural Geology and Geochronology

Curved orogens that result from differential vertical-axis rotations develop remarkable structures within their hinges where compressive and extensive radial structures often develop in combination with tangential shear structures (e.g. Li et al., 2012; Eichelberger and McQuarrie, 2015). With the re-emergence of the Central Iberian curve debate, several studies have re-evaluated the well-documented structures from the Central Iberian Zone to constrain the origin and kinematics of curvature. The majority of studies focused on the hinge zone of the curve in the area surrounding Galicia Tras-os-Montes (e.g. Dias da Silva et al., 2014; Jacques et al., 2018a), but some explored more outer-arc areas (e.g. Palero-Fernández et al., 2015; Gutiérrez-Alonso et al., 2015). The following paragraphs synthesize the findings of new field, structural, and geochronological analyses from around the hinge of the Central Iberian curve and its surrounding regions. The reviewed studies identify several deformation events that are linked to regional Variscan deformation phases (Fig. 2A).

1. An early generation of upright to overturned cylindrical folds with an associated axial planar cleavage (C1). The C1 fold axes plunge variably from horizontal to nearly vertical (e.g. Jacques et al., 2018a, 2018b). The original trend of the fold axes was parallel to the orogen (e.g. Pastor-Galán et al., 2019b), however interference with younger deformation events has created complicated geometries (e.g. Díez Fernández et al., 2013; Palero-Fernández et al., 2015). The emplacement of the allochthonous units of Galicia Tras-os-Montes zone (commonly referred as C2) is closely associated with development of C1 folds, but is restricted to shear zones located along the boundary between the latter and

- the Central Iberian Zone. This phase includes orogen-parallel emplacement of the allochthonous Galicia Tras-Os Montes units and its associated thrusts (Fig. 2A). The non-coaxial nature of the emplacement of this allochthonous nappe produced folding interference and local vertical-axis rotations (Dias da Silva et al., in press). Prograde Barrovian metamorphism (known as M1) reached its pressure peak at the end of C2 (Rubio Pascual et al., 2013).
2. After C1 and C2, the resulting thickened crust gravitationally collapsed (Macaya et al., 1991; Escuder Viruete et al., 1994; Díaz-Balda et al., 1995; Díez-Montes, 2010). This gravitational collapse (phase E1) formed gneiss-dome core complexes between 330 and 317 Ma (e.g. Díez Fernández and Pereira, 2016) especially at the core of the Central Iberian curve (Fig. 2C; e.g. Martínez-Catalán, 2012). This phase formed large subhorizontal extensional detachments that exhumed to depths of the middle crust (e.g. Rubio-Pascual et al., 2013; Dias da Silva et al., in press). General decompression produced a Buchan-type metamorphic event (M2; e.g. Rubio-Pascual et al., 2016, Solís-Alulima et al., 2019) and widespread anatexis melting (e.g. López-Moro et al., 2018; Pereira et al., 2018). E1 phase developed a fold system with sub-horizontal axes and a penetrative subhorizontal cleavage (e.g. Dias da Silva et al., in press). Mapped folding geometries indicate the deflection of C1 folds into overturned positions within the E1 deformation zones (e.g. Díez Fernández et al., 2013; Díez Fernández and Pereira, 2016; Pastor-Galán et al., 2019b). In addition to large-scale extensional deformation and Buchan metamorphism, E1 developed a regional dome-and-basin pattern, resulting in portions of the allochthonous terranes tectonically transported into basins (e.g. Dias da Silva et al., in press).
 3. The structures developed during C1-C2 compression and E1 extension, are re-folded by a younger shortening phase (C3; syn-Cantabrian Orocline). C3 formed upright open folds and conjugate sub-vertical shear zones (e.g. Gutiérrez-Alonso et al., 2015; Díez Fernández and Pereira, 2017; Dias da Silva et al., in press). C3 was coeval with regional retrograde metamorphism (M3) and with intrusion of mantle derived granitoids (Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a), surrounded by contact metamorphic aureoles (e.g. Yenes et al., 1999). The age of the C3 event ranges from 315 and 290 (e.g. Jacques et al., 2018a), concomitant with the formation of the Cantabrian Orocline (e.g. Pastor-Galán et al., 2015a). Ductile deformation, including folding with axial planar cleavage (e.g. Dias da Silva et al., 2014; Pastor-Galán et al., 2019b) and shear zones, occurred at the early stages of C3 (315-305 Ma; Gutiérrez-Alonso et al., 2015; Díez-Fernández and Pereira,

2017; Jacques et al., 2018b) followed by brittle deformation that formed cross-joint sets and vein swarms with Sn-W mineralizations (Jacques et al., 2018a; 2018b). The conjugated shear zones, some of them with hundreds of kilometers of displacement, had activity during the period 315-305 based on direct Ar-Ar dating of the shear zones (Gutiérrez-Alonso et al., 2015) and cross-cutting relationships with precisely dated igneous rocks (Díez-Fernández and Pereira, 2017). Note that these shear zones show a younger age with respect to the sinistral shear zones that bound the Ossa Morena and South Portuguese zones (340-330 Ma; e.g. Dallmeyer et al., 1993). New studies in the Central Iberian Zone have determined that several folds, previously interpreted as C1 (e.g. the Tamames-Marofa-Sátão synform) are C3 structures, possibly nucleated within existing C1-C2 structures (e.g. Dias da Silva et al., 2017; Jacques et al., 2018b). The remarkable continuity along the Central Iberian Zone of these folds (Fig. 2A), previously interpreted as C1 (e.g. Díez-Peña et al., 1990; Abalos et al., 2002; Dias and Ribeiro, 1994; Dias et al., 2016), suggest the ubiquity and importance of this deformation phase.

4. Subsequent to C3 deformation, a brittle shortening event (C4) together with some late extensional faults occurred across the region (E2; Fig. 2A; Dias and Ribeiro 1991; Dias et al. 2003; Rubio Pascual et al., 2013; Arango et al., 2013; Fernández-Lozano et al. 2019; Dias da Silva et al., in press). E2 developed core complex-like structures that further telescoped the M2 metamorphic isograds between the anatectic cores of gneiss domes and the hanging wall units. This event also favoured sub-horizontal folding and kink-band generation in the upper structural levels. Post-Variscan shortening structures in Northern Iberia are characterized by a N-S compressive regime (C4) allowing the formation of brittle NNE-SSW and NNW-SSE faults and associated sub-vertical and sub-horizontal widespread kink-bands (e.g. Aller et al., 2020).

5.2 Paleomagnetism

Paleomagnetism investigates the record of the Earth's ancient magnetic field as it is recorded in the rock record. Among other features, rocks can record the orientation of the magnetic field at the time of magnetization (e.g. Tauxe, 2010). The recorded magnetic vector can be geometrically defined by two components: inclination, which is a function of the paleolatitude (being 90° at the poles and 0° at the equator) at the time of magnetization acquisition; and declination, which is a measure of the horizontal angular difference between the recorded magnetic direction and true north, thereby allowing for the quantification of any vertical-axis rotations if a north reference direction is known for the region of interest at the time

of magnetization acquisition. Paleomagnetism is the best tool to quantify vertical-axis rotations in orogens due to the independence of the magnetic field from the orogen deformation and evolution (e.g. Butler, 1998).

Despite its uniqueness to study paleolatitudes and vertical-axis rotations, paleomagnetism is not flawless. Paleomagnetic data can yield spurious rotations when the local and regional structures are not properly studied and their geometries and kinematic histories not adequately corrected for (e.g. Pueyo et al., 2016). In addition, the age of magnetization acquisition is not necessarily equivalent to the age of the sampled rock. Remagnetizations are ubiquitous, especially in orogens (Weil and van der Voo, 2002; Pueyo et al., 2007; Huang et al., 2017). In remagnetized rocks, the primary magnetization is replaced or overprinted due to a set of geologic processes acting alone or in concert - usually represented by a combination of thermal or chemical reactions (Jackson, 1990). Nevertheless, remagnetizations can be useful for interpreting deformation history if the relative timing of the overprint can be established and a well-constrained reference direction for that age is known (e.g. Weil et al., 2001; Izquierdo-Llavall et al. 2015; Calvín et al., 2017).

In addition to knowing the structural geology and the timing of magnetization of the studied rocks, understanding and quantifying local and regional vertical-axis rotations require a paleomagnetic reference pole for comparison. Permian and Mesozoic paleomagnetic studies in Iberia indicate that Iberia was a relatively stable plate from at least Guadalupian times (ca. 270 Ma) to the opening of the Bay of Biscay in the Cretaceous (e.g. Gong et al., 2008; Vissers et al., 2016). Weil et al. (2010) calculated the most modern Early Permian pole for stable Iberia, which will be used herein as a reference for any vertical-axis rotation analysis (hereafter, eP pole or eP component). Weil et al.'s Virtual Geomagnetic Pole (VGP) values are $Plat = 43.9$; $Plong = 203.3$ and $\alpha_{95} = 5.4$ and when transform into paleomagnetic components has a $\sim 0^\circ$ inclination (equatorial) and declinations that range from 150° to 160° (from NW to SW respectively) depending on where in Iberia you are referencing. In Fig. 6 (red arrows), a compilation of declinations that form part of this composite pole and other eP components found in recent studies are presented.

For the Central Iberian curve, the voluminous paleomagnetic database from the Cantabrian Orocline can be used to partially constrain its kinematics (e.g. Weil et al., 2013). The orocline test for the Cantabrian Orocline (fig. 4) quantifies the degree of differential vertical-axis rotation of variously striking Variscan segments in northern Iberia. If the Central Iberian curve is a product of vertical-axis rotation, paleomagnetic declinations would bend around the Central Iberian curve opposite to that of the Cantabrian Orocline. With a well constrained orocline test,

as in the Cantabrian Orocline (Fig. 4), one can use the paleomagnetic strike-test correlation slope to establish expected declinations for any along-strike portion of the orogen (Pastor-Galán et al., 2017b).

Before the resurgence of the Central Iberian curve, the only available pre-Permian paleomagnetic studies to the South and west of the Cantabrian Zone in the Iberian Massif were focused on the Beja Gabbroic Massif, Portugal (Perroud et al., 1985) and the Almadén syncline volcanics (Perroud et al., 1991; Pares & Van der Voo, 1992). The study in the Beja area showed varied inclinations and declinations in the gabbros, and complex overprints elsewhere. Perroud et al (1985) did not consider any structural correction for the results, assuming the gabbro was undeformed. Recently, Dias da Silva et al. (2018) showed that the area underwent intense deformation during the Carboniferous. Therefore interpretation of this dataset is complicated without knowing the proper structural correction needed to restore the magnetization to its palinspastic orientation.

Several articles with new paleomagnetic studies around the Central Iberian curve have been published since 2015 (Fig. 5). In general, these studies have reported a pervasive late Carboniferous (320 to 300 Ma) (re-)magnetization in sedimentary and igneous rocks (e.g. Pastor-Galán et al., 2015a; 2017b; Fernández-Lozano et al. 2016), which is largely penecontemporaneous to the intrusion of E1 extensional granites (López-Moro et al., 2018) and C3 syn-orocline mantle derived granitoids (Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a). The following section describes the magnetizations from oldest to youngest.

Pastor-Galán et al. (2016) sampled for paleomagnetic analyses both E1 extensional granites (Fig. 2C; ~320 Ma; e.g. López-Moro et al., 2018) from the Tormes and Martinamor domes, and C3 mantle derived granitoids in the Central System (Fig. 2C; 305-295 Ma; e.g. Gutiérrez-Alonso et al., 2011a). Both sets of plutons are located around the Galinas Montes hinge of the Central Iberian curve (Fig. 6-5). The authors found an original component in E1 granites supported by a positive reversal test in both domes (Fig. 7). The magnetization has an inclination (Inc.) = 15° (paleolatitude (λ) = -7.6°) and declination (Dec.) = 81° (Fig. 7), which imply a northward movement of 700 km and a ~70° CCW rotation with respect to the C3 granites that showed an eP component (Dec. ~ 150, Inc. ~ 0). Considering the positive reversal test in E1 granites and the significant difference in inclinations with respect to C3 granitoids (eP component), a magnetization age of older than 318 Ma was proposed (pre Kiaman superchron, 317 Ma - 267 Ma, e.g. Langereis et al., 2010), which was interpreted as a primary magnetization. The 70° CCW Pennsylvanian rotation recorded in rocks from the Central Iberian curve hinge zone is in agreement with the expected rotation of the southern limb of the

Cantabrian Orocline (Fig. 4; Weil et al., 2013).

At the putative outer arc of the Central Iberian curve, the Iberian Ranges (Fig. 2), paleomagnetic and structural studies of Devonian and Permian rocks (Pastor-Galán et al., 2018) revealed that the eP component from Permian rocks had rotated $\sim 22^\circ$ CW during the Cenozoic (Fig. 8; cf. Pastor-Galán et al., 2018). The Permian and Mesozoic rocks from the Iberian Ranges show a consistent $\sim 22^\circ$ CW rotation with respect to the Apparent Polar Wander Path for Iberia (e.g. Pastor-Galán et al. 2018). This rotation likely happened during the Alpine orogeny, in which the northern area of the Iberian Range underwent more shortening than the southern part, resulting in a regional CW vertical-axis rotation (Izquierdo-Llavall et al., 2019). After restoring the Cenozoic rotation, the Devonian rocks show a positive reversal and fold-test with inclinations that are steeper than expected from the eP component (Dec. = 85.3° , Inc. = 12.7° , $\lambda = -6.4$). This component is statistically indistinguishable from that of the E1 granites and the southern branch of the Cantabrian Zone, showing the same 70° CCW rotation from the time they were remagnetization (estimated in 318 Ma) to the timing of the eP component (Fig. 8; Pastor-Galán et al., 2018). Once Cenozoic rotation is corrected for, the structural and paleomagnetic trends of the Iberian ranges become parallel to those in the southern limb of the Cantabrian Orocline, ruling out a Variscan or older origin for the outer Central Iberian curve (Fig. 8).

The remaining paleomagnetic works published on Central and SW Iberia rocks all reveal a ubiquitous late Carboniferous to Early Permian remagnetizations during the Kiaman superchron (Fernández-Lozano et al., 2016; Pastor-Galán et al., 2015a; 2016; 2017b; Leite Mendes, in press). The authors of these papers calculated the expected declination for each site as if they were part of the Cantabrian Orocline (Fig. 9A). All localities where magnetizations pre-date the formation of the Cantabrian Orocline show the same expected rotations as the southern limb of the Cantabrian Orocline, regardless of their position within the Central Iberian curve (to the hinge: Tormes and Martinamor domes, Iberian ranges; to the southern limb: Almadén syncline and South Portuguese Zone). Other locations, especially limestones from the Central Iberian Zone, have declinations and inclinations in between the primary 318 Ma component of the E1 granites and the post-orocline eP component (Fig. 9B). Pastor-Galán et al. (2015a; 2016) interpreted these results as being caused by a remagnetization that was acquired during Cantabrian Orocline formation and therefore recorded intermediate steps between the component of the E1 granites and eP. Those authors suggest that the large amount of syn-orocline mantle derived granitoids that intruded the Central Iberian Zone (C3 granitoids) triggered the hinterland remagnetization.

Finally, two previous studies identified an earlier magnetization in the Almadén syncline region of the SE Central Iberian Zone (Perroud et al., 1991; Pares & Van der Voo, 1992). However, Leite Mendes et al. (in press) argue that these studies are likely misinterpreted. Perroud et al. (1991), applied a complicated structural correction restoring a putative plunge of the regional structural axis to all sites, including those where the syncline axis does not plunge. Leite Mendes et al. (in press) re-sampled the syncline where its axis is sub-horizontal and obtained a negative fold test, implying that the magnetization is not primary as previously interpreted. Their results, however, are similar in orientation to those components published from previous studies prior to any structural correction (Perroud et al., 1991 and Parés and van der Voo, 1992).

Two additional studies sampled Laurussian margin sequences that are today adjacent to the Cantabrian Orocline region (Fig. 10). To the north, the SW area of Ireland preserves a Late Paleozoic basin filled with Devonian red sandstone and Carboniferous limestone and siltstone, which was sampled by Pastor-Galán et al. (2015a). To the south is the aforementioned results from the South Portuguese Zone (Leite Mendes et al., in press). Both areas are interpreted to have previously been part of the Laurussian continent, on the opposite side of the Rheic Ocean suture at the time of Variscan collision (Fig. 10; e.g. Pastor-Galán et al., 2015b). In contrast, the rest of Iberia was part of, or proximal to, Gondwana (e.g. Franke et al., 2017). These Paleomagnetic results from the Laurussian margin suggest that the rotations involved in the formation of the Cantabrian Orocline occurred along both sides of the Rheic suture proximal to both its northern and southern limb (Fig. 10A and B). Pastor-Galán et al. (2015b) hypothesized a so-called Greater Cantabrian Orocline that would have bent the entire Appalachian/Variscan orogen around a vertical-axis, affecting at least the continental margins of both Gondwana and Laurussia.

5.3 The implications of not being a secondary orocline

The most relevant new data regarding the kinematics of the Central Iberian curve is the paleomagnetic study from the Iberian Ranges (Calvín et al., 2014; Pastor-Galán et al., 2018). These results confirm that the present-day variation in trend of the tectonostratigraphic units, generally attributed to Variscan tectonics (e.g. Weil et al., 2013; Shaw et al., 2012; 2014), is likely a product of Cenozoic Alpine orogeny. Izquierdo-Llavall et al. (2019) confirmed that the interpreted Alpine rotations correspond well with the amount of shortening reconstructed in Meso-Cenozoic basins. The best preserved and most continuous outcrop in the Central Iberian's outer arc is not a Variscan structure, casting doubt that Central Iberian curve's is

related to Variscan kinematics. The results are also a reminder that the regional effects of Alpine deformation are often underestimated, especially close to the major Iberian Alpine fronts: the Pyrenees, Iberian Ranges, and the Betics.

Overall, new paleomagnetic data from the Central Iberian curve and nearby areas reveal pervasive late Carboniferous remagnetizations and regional vertical-axis rotations of the same sense and magnitude to those expected for the southern arm of the Cantabrian Orocline. The new paleomagnetic data indicate that a post ~320 Ma formation for the Central Iberia curve due to vertical-axis rotations is not supported (Pastor-Galán et al., 2016). The distribution in space and time of paleomagnetic results discards the formation of the Central Iberian curve as a product of Variscan gravitational collapse (E1, ~330-317 Ma) or concomitant to the Cantabrian Orocline (C3). So far, no pre-E2 paleomagnetic component has been found, and consequently, paleomagnetic data cannot reject an early orogenic origin for the inner arc of the Central Iberian curve (C1-C2, older than 330 Ma).

From a structural geology point of view, the Central Iberian curve does not display the classic geometries and structural interference patterns as found in other established oroclines (i.e., those systems that involve differential vertical-axis rotations, e.g. Li et al., 2012; van der Boon et al., 2018; Meijers et al., 2017; Rezaeian et al., in press). The geometry and structural behaviour of oroclines should resemble, at the crustal-scale, a regional vertical-axis fold preserved in plan-view, either formed by buckling (e.g. Johnston et al., 2001) or bending (e.g. Cifelli et al., 2008) mechanisms. In oroclines, pre-existing structures tend to follow fold trends around the curvature (e.g. Rosenbaum, 2014; Li et al., 2018). In addition, orocline cores tend to preserve radial structures and shortening patterns in the inner arc and orocline parallel shear zones and extension structures in their outer arc (e.g. Ries and Shackleton, 1976; Eichelberger and McQuarrie, 2015), similar to what is observed in multilayer folds (e.g. Fossen, 2016).

The structural geometry of the Central Iberian curve lacks such patterns. Paleomagnetism from the Iberian Ranges indicate that the Cantabrian and West Asturian Leonese zones do not follow the Central Iberian curve, instead they continue their NWW-SEE trend into the Mediterranean in what it is now the Betic chain (Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018; van Hinsbergen et al., 2020). Structural trends in the Ossa Morena and the South Portuguese Zone do not show any change in along-strike structural trend that supports large-scale CW rotations (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019), whereas existing paleomagnetic data from those zones (Leite Mendes et al., in press) support a model of CCW rotation associated with the broader southern arm of the Cantabrian Orocline. In the Central Iberian and Galicia Tras-os-Montes zones, the trend of curvature is irregular (see C1

fold patterns in Fig. 2A) and nowhere are the expected inner and outer arc-related structures preserved (e.g. Dias da Silva et al. in press).

The curved shape of C1 fold axes in the Central Iberian zone is better explained by fold interference patterns than vertical-axis rotations (e.g. Pastor-Galán et al., 2019b). Moreover, the curved shape of the Galicia Tras-os-Montes allochthonous nappe, which was emplaced orogen parallel, shows no evidence of vertical-axis rotation related structures (Fig. 2A; Dias da Silva et al., in press). Other authors describe the changes in trend around the Central Iberian curve expressed by C1 folds (Fig. 2A) as the product of fold interference patterns (e.g. Gutiérrez-Alonso, 2009; Palero-Fernández et al., 2015; Jacques et al., 2018b; Dias da Silva et al., in press). Pastor-Galán et al. (2019b) showed that curved C1 folds in the Central Iberian Zone around the Galicia Tras-os-Montes boundary (Fig. 2A) are coaxial with C3 folds after restoring the effects of C2 and E1 deformation phases (Fig. 11A). Both C1 and C3 formed under similar shortening directions. In the same area, Jacques et al. (2018b) found similar fold interference patterns, in addition they described kinematic incompatibility with the expected CW rotations that would have occurred if the Central Iberian curve was an orocline. In other areas of the Central Iberian Zone, the curved shape of C1 folds has been described as an interference between C1 structures and their reorientation caused by C3 shear zones (Fig. 2A; e.g. Palero-Fernández et al., 2015; Dias et al., 2016), or alternatively the interference between C1, C3 and the E2 structures (Fig. 2A; Gutiérrez-Alonso, 2009; Arango et al., 2013; Rubio Pascual et al. 2013).

Overall, new geometric and kinematic data favor the interpretation that the Central Iberian curve is not a structure formed by differential vertical-axis rotation as was the Cantabrian Orocline, but one formed as a consequence of several competing processes. It is clear from the current data that a combination of several deformation events caused the orientation of structures that today delineate the shape of the Central Iberia curve. These include: (1) the northern part of the outer-arc is the product of an Alpine rigid block rotation instead of Variscan differential vertical-axis rotation (Pastor-Galán et al., 2018); (2) the curvature of the Galicia Tras-os-Montes allochthonous nappe reflects its original shape and could be defined as a primary curve (see Weil and Sussman, 2004), since it was emplaced orogen parallel and shows no sign of vertical-axis rotations at any time (fig. 2A; Dias da Silva et al., in press); (3) Structural analysis shows that fold interference patterns explain the geometry of the curved trends of Central Iberian Zone's C1 folds (Pastor-Galán et al., 2019b), whose kinematics are incompatible with the required CW rotations expected if the curve is an orocline (Jacques et al, 2018b).

6 Get Back: Ideas flowing out and endless questions

The pioneering works in the last decade that resurrected the idea of a Central Iberian curve, speculated that both the Cantabrian and Central Iberian zones buckled together as secondary oroclines (Fig. 12; Martínez-Catalán 2011; Shaw et al., 2012, 2014; Shaw and Johnston, 2016; Carreras and Druguet, 2014). Later, Martínez Catalan et al. (2014) and Díez Fernández and Pereira (2017) reformulated Martínez-Catalán's 2011 hypothesis and proposed that the Central Iberian curve formed as an orocline between 315 and 305 Ma, and assigning the Cantabrian Orocline a time frame between 305 and 295 Ma (Fig. 12). The proposed tectonic mechanisms to support these early kinematic models are varied: (1) buckling of a ribbon 'Armorican' continent (Fig. 12A; Shaw et al., 2014; 2016); (2) buckling of a completely formed Variscan orogen during a putative 'Pangea B' to 'Pangea A' transition in the late Carboniferous (Fig. 12B; Carreras and Druguet, 2014; Martínez-Catalán et al., 2011); (3) indentation of Laurussia into Gondwana during the early stages of collision (at present day SW Iberia, South Portuguese Zone), producing first the Central Iberian curve as a mega drag-fold during Carboniferous times and then slightly later the Cantabrian Orocline as a consequence of an indentation process (fig. 12C; Simancas et al., 2013).

The reviewed data in sections 4 and 5 contradict the aforementioned hypotheses. Paleomagnetism and structural patterns (section 5; Fig. 6-11) disagree with the necessary CW rotations required to support a late Carboniferous orocline origin for the Central Iberian curve (Models in Fig. 12A and B). In addition, the sense and magnitude of the vertical-axis rotations observed in SW Iberia (Fig. 10) imply that the South Portuguese (Avalonian segment) and Ossa Morena zones moved together with the southern limb of the Cantabrian Orocline during the Pennsylvanian and Early Permian. This means that the South Portuguese Zone was already parallel to the general trend of the Variscan orogen prior to Cantabrian Orocline formation, implying the lack of a Laurussian rigid indenter into Gondwana (e.g. Simancas et al., 2013). This discrepancy leaves orogen-parallel terrane transport as a possible explanation to the kinematics observed in Ossa Morena and South-Portuguese Zones (e.g. Pérez-Cáceres et al., 2016). At the same time, paleomagnetism from SW Iberia backs the hypothesis of a Greater Cantabrian Orocline extended into both Gondwana and Laurussia in its northern and southern limbs (Fig. 10; Pastor-Galán et al., 2015b).

In spite of the kinematic constraints and structural patterns, which do not support a vertical-axis origin for the Central Iberian curve in Late Carboniferous time, other geometric constraints remain challenging. The curved shape of the aeromagnetic and gravity anomalies of Iberia are real (Fig. 5). These striking patterns could be due to Variscan-Alpine structural

interference, for example the previous example from the Iberian Ranges, but currently there is not enough data to rigorously test this hypothesis. Shaw et al. (2012) supported their hypothesis of a secondary orocline by assuming that paleocurrents were parallel through Iberia during Ordovician times. However, some of the observed deflections in the paleocurrents studied by Shaw et al. (2012; see Fig. 3) are also explained by Alpine vertical-axis rotations (the case of the Iberian ranges) and fold interference patterns (SE of the Central Iberian Zone). Others (Central and SW of the Central Iberian Zone) may be explained by a local response to basin architecture (Fig. 3), where paleo-flow directions would trend toward the deepest basin throughs. The Ordovician basin architecture of Iberia allows for opposite directed paleocurrents from both sides of such throughs (Fig. 3). However, the Early Paleozoic basin architecture in Iberia and their local deformation events require further research (Sánchez-García et al., 2019).

Although kinematic evidence is still scarce for the earliest Variscan movements, we argue that pre-orogenic physiographic features, such as the opening of a marginal restricted ocean between Gondwana and its distal platform at 395 Ma (Fig. 13A; Pin et al., 2002; Gutiérrez-Alonso et al., 2008b; Arenas et al., 2016) explains the rounded shape of the Galicia tras-os-Montes curve as a primary arc. During the collision, the latter irregularity would cause the orogen-parallel emplacement of allochthonous nappe (Fig. 13B; Dias da Silva et al., in press) and the left-lateral movements of the Ossa Morena and South Portuguese Zones in SW Iberia (Fig 13A, B, C; Quesada, 2019). During the late Carboniferous, possibly due to a plate reorganization during the final amalgamation of Pangea (Fig. 13D; e.g. Gutiérrez-Alonso et al., 2008a; Pastor-Galán et al., 2015a), the far-field stress-field likely changed and buckled the entire orogen around a vertical axis (Gutiérrez-Alonso et al., 2004), including both the Gondwana and Laurussia margins (Fig. 13E; Pastor-Galán et al., 2015b).

Acknowledgements

DPG thanks the extraordinary hard work, patience and endurance of the Utrecht University students that embraced and enjoyed studying the kinematics of the central Iberian curvature: Thomas Groenewegen, Bart Ursem, Daniel Brower, Mark Diederer and Bruno Leite-Mendes. DPG acknowledges FRIS and CNEAS for the continuous financial support. GGA is supported by Spanish Ministry of Science, innovation and universities under the project IBERCRUST (PGC2018-096534-B-I00) and Russian Federation Government grant no. 14.Y26.31.0012. This paper is a contribution to the IGCP no. 648 “Supercontinent Cycles and Global Geodynamics”. 50 years ago four fabulous guys let it be and never got back.

769 References

- 770 Ábalos, B., Carreras, J., Druguet, E., Escuder Viruete, J., Gómez Pugnaire, M. T., Lorenzo
771 Álvarez, S., Quesada, C., Rodríguez Fernández, L. R. and Gil-Ibarguchi, J. I.: Variscan and pre-
772 Variscan tectonics, *Geol. Spain*, 155–183, 2002.
- 773 Aerden, D.: Correlating deformation in Variscan NW-Iberia using porphyroblasts; implications for
774 the Ibero-Armorican Arc, *J. Struct. Geol.*, 26(1), 177–196, 2004.
- 775 Alcock, J. E., Catalán, J. R. M., Arenas, R. and Montes, A. D.: Use of thermal modeling to
776 assess the tectono-metamorphic history of the Lugo and Sanabria gneiss domes, Northwest
777 Iberia, *Bull. Société Géologique Fr.*, 180(3), 179–197, doi:10.2113/gssgfbull.180.3.179, 2009.
- 778 Aller, J., Bastida, F. and Bobillo-Ares, N. C.: On the development of kink-bands: A case study in
779 the West Asturian-Leonese Zone (Variscan belt, NW Spain) / Sur le développement des kink-
780 bands : un exemple dans le Zone Astur Occidentale-léonaise (chaîne varisque ibérique, nord-
781 ouest de l'Espagne), *Bull. Société Géologique Fr.*, 191(1), doi:10.1051/bsgf/2020003, 2020.
- 782 Aller, J. J. J. and Gallastegui, J. J.: Analysis of kilometric-scale superposed folding in the
783 Central Coal Basin (Cantabrian zone, NW Spain), *J. Struct. Geol.*, 17(7), 961–969,
784 doi:10.1016/0191-8141(94)00115-G, 1995.
- 785 Alonso, J. L., Marcos, A. and Suárez, A.: Paleogeographic inversion resulting from large out of
786 sequence breaching thrusts: The León Fault (Cantabrian zone, NW Iberia). A new picture of the
787 external Variscan thrust belt in the Ibero-Armorican arc, *Geol. Acta*, 7(4), 451–473,
788 doi:10.1344/105.000001449, 2009.
- 789 Álvaro, J. J., Casas, J. M., Clausen, S. and Quesada, C.: Early Palaeozoic geodynamics in NW
790 Gondwana, *J. Iber. Geol.*, 44(4), 551–565, doi:10.1007/s41513-018-0079-x, 2018.
- 791 Ardizzone, Juan, Julio Mezcua, and Isabel Socías. Mapa aeromagnético de España peninsular.
792 Instituto Geográfico Nacional, 1989.
- 793 Mergl, M. & Zamora, S. 2012, New and revised occurrences of... *Bulletin of Geosciences*, 87,
794 571–586., [online] Available from: <http://www.geology.cz/bulletin/contents/art1327> (Accessed 6
795 April 2020b), n.d.
- 796 Aramburu, C., Méndez-Bedia, I. and Arbizu, M.: The Lower Palaeozoic in the Cantabrian Zone
797 (Cantabrian Mountains, NW Spain)., edited by S. García-López and F. Bastida, pp. 35–49.,
798 2002.
- 799 Arango, C., Díez Fernández, R. and Arenas, R.: Large-scale flat-lying isoclinal folding in
800 extending lithosphere: Santa María de la Alameda dome (Central Iberian Massif, Spain),
801 *Lithosphere*, 5(5), 483–500, doi:10.1130/L270.1, 2013.
- 802 Arenas, R., Sánchez Martínez, S., Díez Fernández, R., Gerdes, A., Abati, J., Fernández-
803 Suárez, J., Andonaegui, P., González Cuadra, P., López Carmona, A., Albert, R., Fuenlabrada,
804 J. M. and Rubio Pascual, F. J.: Allochthonous terranes involved in the Variscan suture of NW
805 Iberia: A review of their origin and tectonothermal evolution, *Earth-Sci. Rev.*, 161, 140–178,
806 doi:10.1016/j.earscirev.2016.08.010, 2016.

807 Ayala, C., Bohoyo, F., Maestro, A., Reguera, M. I., Torne, M., Rubio, F., Fernández, M. and
808 García-Lobón, J. L.: Updated Bouguer anomalies of the Iberian Peninsula: a new perspective to
809 interpret the regional geology, *J. Maps*, 12(5), 1089–1092,
810 doi:10.1080/17445647.2015.1126538, 2016.

811 Ayarza, P., Catalan, J. R. M., Gallart, J., Pulgar, J. A. and Danobeitia, J. J.: Estudio Sismico de
812 la Corteza Iberica Norte 3.3: A seismic image of the Variscan crust in the hinterland of the NW
813 Iberian Massif, *Tectonics*, 17(2), 171–+, 1998.

814 Azor, A., Rubatto, D., Simancas, J. F., Lodeiro, F. G., Poyatos, D. M., Parra, L. M. M. and
815 Matas, J.: Rheic Ocean ophiolitic remnants in Southern Iberia questioned by SHRIMP U-Pb
816 zircon ages on the Beja-Acebuches amphibolites - art. no. TC5006, *Tectonics*, 27(5), C5006–
817 C5006, 2008.

818 Azor, A., Dias da Silva, Í., Gómez Barreiro, J., González-Clavijo, E., Martínez Catalán, J. R.,
819 Simancas, J. F., Martínez Poyatos, D., Pérez-Cáceres, I., González Lodeiro, F., Expósito, I.,
820 Casas, J. M., Clariana, P., García-Sansegundo, J. and Margalef, A.: Deformation and Structure,
821 in *The Geology of Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T. Oliveira, pp.
822 307–348, Springer International Publishing, Cham., 2019.

823 Barrois, C. E.: Recherches sur le terrains anciens des Asturies et de la Galice, Six-Horemans.,
824 1882.

825 Bastida, F.: Zona Cantábrica, in *Geología de España*, edited by J. A. Vera, pp. 25–49, SGE-
826 IGME, Madrid., 2004.

827 Bayona, G., Thomas, W. A. and Van der Voo, R.: Kinematics of thrust sheets within transverse
828 zones: A structural and paleomagnetic investigation in the Appalachian thrust belt of Georgia
829 and Alabama, *J. Struct. Geol.*, 25(8), 1193–1212, doi:10.1016/S0191-8141(02)00162-1, 2003.

830 van der Boon, A., van Hinsbergen, D. J. J. J., Rezaeian, M., Gürer, D., Honarmand, M.,
831 Pastor-Galán, D., Krijgsman, W. and Langereis, C. G. G.: Quantifying Arabia–Eurasia
832 convergence accommodated in the Greater Caucasus by paleomagnetic reconstruction, *Earth*
833 *Planet. Sci. Lett.*, 482, doi:10.1016/j.epsl.2017.11.025, 2018.

834 Buiter, S. J. H. and Torsvik, T. H.: A review of Wilson Cycle plate margins: A role for mantle
835 plumes in continental break-up along sutures?, *Gondwana Res.*, 26(2), 627–653,
836 doi:10.1016/j.gr.2014.02.007, 2014.

837 Butler, R.: Paleomagnetism: Magnetic domains to geologic terranes, *Electron. Ed.*,
838 (September), 319–319, doi:10.1006/icar.2001.6754, 1998.

839 Calvin, P., Casas, A. M., Villalaín, J. J., Tierz, P., Calvin, P., Casas, A. M., Villalaín, J. J., Tierz,
840 P., Calvin, P., Casas, A. M., Villalaín, J. J. and Tierz, P.: Reverse magnetic anomaly controlled
841 by Permian Igneous rocks in the Iberian Chain (N Spain), *Geol. Acta*, 12(3), 193–207,
842 doi:10.1344/GeologicaActa2014.12.3.2, 2014.

843 Calvin, P., Casas-Sainz, A. M. M., Villalaín, J. J. J. and Moussaid, B.: Diachronous folding and
844 cleavage in an intraplate setting (Central High Atlas, Morocco) determined through the study of
845 remagnetizations, *J. Struct. Geol.*, 97, 144–160, doi:10.1016/j.jsg.2017.02.009, 2017.

- 846 Calvín-Ballester, P. and Casas, A.: Folded Variscan thrusts in the Herrera Unit of the Iberian
847 Range (NE Spain), *Geol. Soc. Lond. Spec. Publ.*, 394(1), 39–52, doi:10.1144/SP394.3, 2014.
- 848 Carey, S. W.: The orocline concept in geotectonics-Part I, *Pap. Proc. R. Soc. Tasman.*, 89,
849 255–288, 1955.
- 850 Carreras, J. and Druguet, E.: Framing the tectonic regime of the NE Iberian Variscan segment,
851 *Geol. Soc. Lond. Spec. Publ.*, 405(1), 249–264, doi:10.1144/SP405.7, 2014.
- 852 Martínez-Catalán, J. R. M., Fernández-Suárez, J., Jenner, G. A., Belousova, E. and Montes, A.:
853 Provenance constraints from detrital zircon U–Pb ages in the NW Iberian Massif: implications
854 for Palaeozoic plate configuration and Variscan evolution, *J. Geol. Soc.*, 161(3), 463–476,
855 doi:10.1144/0016-764903-054, 2004.
- 856 Chopin, F., Schulmann, K., Skrzypek, E., Lehmann, J., Dujardin, J. R., Martelat, J. E., Lexa, O.,
857 Corsini, M., Edel, J. B., Štípská, P. and Pitra, P.: Crustal influx, indentation, ductile thinning and
858 gravity redistribution in a continental wedge: Building a Moldanubian mantled gneiss dome with
859 underthrust Saxothuringian material (European Variscan belt), *Tectonics*, 31(1), n/a-n/a,
860 doi:10.1029/2011TC002951, 2012.
- 861 Chopin, F., Corsini, M., Schulmann, K., El Houicha, M., Ghienne, J.-F. and Edel, J.-B.: Tectonic
862 evolution of the Rehamna metamorphic dome (Morocco) in the context of the Alleghanian-
863 Variscan orogeny, *Tectonics*, 33(6), 1154–1177, doi:10.1002/2014TC003539, 2014.
- 864 Cifelli, F., Mattei, M. and Della Seta, M.: Calabrian Arc oroclinal bending: The role of subduction,
865 *Tectonics*, 27(5), doi:10.1029/2008TC002272, 2008.
- 866 Dallmeyer, R. D., Fonseca, P. E., Quesada, C. and Ribeiro, A.: $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age
867 constraints for the tectonothermal evolution of a Variscan suture in southwest Iberia,
868 *Tectonophysics*, 222(2), 177–194, doi:10.1016/0040-1951(93)90048-O, 1993.
- 869 Dallmeyer, R. D. D., Catalán, J. R. M., Arenas, R., Gil Ibarguchi, J. I. I., Gutiérrez-Alonso, G.,
870 Farias, P., Bastida, F. and Aller, J.: Diachronous Variscan tectonothermal activity in the NW
871 Iberian Massif: Evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of regional fabrics, *Tectonophysics*, 277(4),
872 307–337, doi:10.1016/S0040-1951(97)00035-8, 1997.
- 873 Dias da Silva, Í., Gómez-Barreiro, J., Martínez Catalán, J. R., Ayarza, P., Pohl, J. and Martínez,
874 E.: Structural and microstructural analysis of the Retortillo Syncline (Variscan belt, Central
875 Iberia). Implications for the Central Iberian Orocline, *Tectonophysics*, 717, 99–115,
876 doi:10.1016/j.tecto.2017.07.015, 2017.
- 877 Dias da Silva, Í., Pereira, M. F., Silva, J. B. and Gama, C.: Time-space distribution of silicic
878 plutonism in a gneiss dome of the Iberian Variscan Belt: The Évora Massif (Ossa-Morena Zone,
879 Portugal), *Tectonophysics*, 747–748, 298–317, doi:10.1016/j.tecto.2018.10.015, 2018.
- 880 Dias da Silva, Í., González Clavijo, E. and Díez-Montes, A.: The collapse of the Variscan belt: a
881 Variscan lateral extrusion thin-skinned structure in NW Iberia, *Int. Geol. Rev.*, 00(00), 1–37,
882 doi:10.1080/00206814.2020.1719544, in press.
- 883 Dias da Silva, Í. F., Linnemann, U., Hofmann, M., González-Clavijo, E., Díez-Montes, A. and
884 Catalán, J. R. M.: Detrital zircon and tectonostratigraphy of the Parautochthon under the Morais

- 885 Complex (NE Portugal): implications for the Variscan accretionary history of the Iberian Massif,
886 J. Geol. Soc., 172(1), 45–61, doi:10.1144/jgs2014-005, 2015.
- 887 Dias, R. and Ribeiro, A.: Finite strain analysis in a transpressive regime (Variscan autochthon,
888 northeast Portugal), Tectonophysics, 191(3), 389–397, doi:10.1016/0040-1951(91)90069-5,
889 1991.
- 890 Dias, R. and Ribeiro, A.: CONSTRICTION IN A TRANSPRESSIVE REGIME - AN EXAMPLE IN
891 THE IBERIAN BRANCH OF THE IBERO-ARMORICAN ARC, J. Struct. Geol., 16(11), 1543–
892 1554, 1994.
- 893 Dias, R., Ribeiro, A., Romão, J., Coke, C. and Moreira, N.: A review of the arcuate structures in
894 the Iberian Variscides; constraints and genetic models, Tectonophysics, 681, 170–194,
895 doi:10.1016/j.tecto.2016.04.011, 2016.
- 896 Díez Balda, M. A., Vegas, R. and González Lodeiro, F.: Central-Iberian zone structure, Pre-
897 Mesoz. Geol. Iber., 172–188, 1990.
- 898 Díez Balda, M. A., Martínez Catalán, J. R. and Ayarza Arribas, P.: Syn-collisional extensional
899 collapse parallel to the orogenic trend in a domain of steep tectonics: the Salamanca
900 Detachment Zone (Central Iberian Zone, Spain), J. Struct. Geol., 17(2), 163–182,
901 doi:10.1016/0191-8141(94)E0042-W, 1995.
- 902 Díez Fernández, R. and Arenas, R.: The Late Devonian Variscan suture of the Iberian Massif: A
903 correlation of high-pressure belts in NW and SW Iberia, Tectonophysics, 654, 96–100,
904 doi:10.1016/j.tecto.2015.05.001, 2015.
- 905 Díez Fernández, R. and Pereira, M. F.: Extensional orogenic collapse captured by strike-slip
906 tectonics: Constraints from structural geology and UPb geochronology of the Pinhel shear zone
907 (Variscan orogen, Iberian Massif), Tectonophysics, 691, 290–310,
908 doi:10.1016/j.tecto.2016.10.023, 2016.
- 909 Díez Fernández, R. and Pereira, M. F.: Strike-slip shear zones of the Iberian Massif: Are they
910 coeval?, Lithosphere, 9(5), 726–744, doi:10.1130/L648.1, 2017.
- 911 Díez Fernández, R., Gómez Barreiro, J., Martínez Catalán, J. R. and Ayarza, P.: Crustal
912 thickening and attenuation as revealed by regional fold interference patterns: Ciudad Rodrigo
913 basement area (Salamanca, Spain), J. Struct. Geol., 46, 115–128,
914 doi:10.1016/j.jsg.2012.09.017, 2013.
- 915 Díez Fernández, R., Arenas, R., Pereira, M. F., Sánchez-Martínez, S., Albert, R., Martín Parra,
916 L.-M., Rubio Pascual, F.-J. and Matas, J.: Tectonic evolution of Variscan Iberia: Gondwana–
917 Laurussia collision revisited, Earth-Sci. Rev., 162, 269–292,
918 doi:10.1016/j.earscirev.2016.08.002, 2016.
- 919 Díez-Montes, A.: La Geología del Dominio “Ollo de Sapo” en las comarcas de Sanabria y
920 Terra do Bolo., 2006.
- 921 Díez-Montes, A., Martínez-Catalán, J. R. R. and Bellido-Mulas, F.: Role of the Ollo de Sapo
922 massive felsic volcanism of NW Iberia in the Early Ordovician dynamics of northern Gondwana,
923 Gondwana Res., 17(2–3), 363–376, doi:10.1016/j.gr.2009.09.001, 2010.

- 924 Domeier, M. and Torsvik, T. H.: Plate tectonics in the late Paleozoic, *Geosci. Front.*, 5(3), 303–
925 350, doi:10.1016/j.gsf.2014.01.002, 2014.
- 926 Edel, J. B., Schulmann, K., Lexa, O. and Lardeaux, J. M.: Late Palaeozoic palaeomagnetic and
927 tectonic constraints for amalgamation of Pangea supercontinent in the European Variscan belt,
928 *Earth-Sci. Rev.*, 177(September 2017), 589–612, doi:10.1016/j.earscirev.2017.12.007, 2018.
- 929 Eduard Suess: *Das Antlitz der Erde*, F. Tempsky; [etc., etc.]. [online] Available from:
930 <http://archive.org/details/dasantlitzderer02suesgoog> (Accessed 12 March 2020), 1892.
- 931 Eichelberger, N. and McQuarrie, N.: Kinematic reconstruction of the Bolivian orocline,
932 *Geosphere*, 11(2), 445–462, doi:10.1130/GES01064.1, 2015.
- 933 Escuder Viruete, J., Arenas, R. and Catalán, J. R. M.: Tectonothermal evolution associated with
934 Variscan crustal extension in the Tormes Gneiss Dome (NW Salamanca, Iberian Massif, Spain),
935 *Tectonophysics*, 238(1), 117–138, doi:10.1016/0040-1951(94)90052-3, 1994.
- 936 Fernández-Lozano, J., Pastor-Galán, D., Gutiérrez-Alonso, G. and Franco, P.: New kinematic
937 constraints on the Cantabrian orocline: A paleomagnetic study from the Peñalba and Truchas
938 synclines, NW Spain, *Tectonophysics*, 681, 195–208, doi:10.1016/j.tecto.2016.02.019, 2016.
- 939 Fernández-Lozano, J., Gutiérrez-Alonso, G., Willingshofer, E., Sokoutis, D., Vicente, G. de and
940 Cloetingh, S.: Shaping of intraplate mountain patterns: The Cantabrian orocline legacy in Alpine
941 Iberia, *Lithosphere*, 11(5), 708–721, doi:10.1130/L1079.1, 2019.
- 942 Fernández-Suárez, J., Gutiérrez-Alonso, G., Pastor-Galán, D., Hofmann, M., Murphy, J. B. and
943 Linnemann, U.: The Ediacaran-Early Cambrian detrital zircon record of NW Iberia: Possible
944 sources and paleogeographic constraints, *Int. J. Earth Sci.*, 103(5), 1335–1357,
945 doi:10.1007/s00531-013-0923-3, 2014.
- 946 Fossen, H.: *Structural Geology*, Cambridge University Press., 2016.
- 947 Franke, W., Cocks, L. R. M. and Torsvik, T. H.: The Palaeozoic Variscan oceans revisited,
948 *Gondwana Res.*, 48, 257–284, doi:10.1016/j.gr.2017.03.005, 2017.
- 949 Franke, W., Cocks, L. R. M. and Torsvik, T. H.: Detrital zircons and the interpretation of
950 palaeogeography, with the Variscan Orogeny as an example, *Geol. Mag.*, 157(4), 690–694,
951 doi:10.1017/S0016756819000943, 2020.
- 952 García-Arias, M., Díez-Montes, A., Villaseca, C. and Blanco-Quintero, I. F.: The Cambro-
953 Ordovician Ollo de Sapo magmatism in the Iberian Massif and its Variscan evolution: A review,
954 *Earth-Sci. Rev.*, 176, 345–372, doi:10.1016/j.earscirev.2017.11.004, 2018.
- 955 Gong, Z., Langereis, C. G. and Mullender, T. A. T.: The rotation of Iberia during the Aptian and
956 the opening of the Bay of Biscay, *Earth Planet. Sci. Lett.*, 273(1–2), 80–93, 2008.
- 957 Gozalo, R., Liñán, E., Vintaned, J. A. G., Eugenia, M., Álvarez, D., Martorell, J. B. C., Zamora,
958 S., Esteve, J. and Mayoral, E.: The Cambrian of the Cadenas Ibéricas (ne Spain) and Its
959 Trilobites., n.d.
- 960 Greco, K. D., Johnston, S. T., Gutiérrez-Alonso, G., Shaw, J. and Lozano, J. F.: Interference

- 961 folding and orocline implications: A structural study of the Ponga Unit, Cantabrian orocline,
962 northern Spain, *Lithosphere*, 8(6), 757–768, doi:10.1130/L576.1, 2016.
- 963 Gutiérrez Alonso, G.: Estratigrafía y Tectónica., in *Memoria del Mapa Geológico de España*
964 1:50000, Hoja de Mora (680), IGME, Madrid. [online] Available from:
965 <http://info.igme.es/cartografiadigital/datos/magna50/memorias/MMagna0658.pdf>, 2009.
- 966 Gutiérrez-Alonso, G.: El Antiforme del Narcea y su relación con los mantos occidentales de la
967 Zona Cantábrica., 1992.
- 968 Gutiérrez-Alonso, G.: Strain partitioning in the footwall of the Somiedo Nappe: Structural
969 evolution of the Narcea Tectonic window, NW Spain, *J. Struct. Geol.*, 18(10), 1217–1229, 1996.
- 970 Gutiérrez-Alonso, G., Fernández-Suárez, J. and Weil, A. B.: Orocline triggered lithospheric
971 delamination, in *Special Paper 383: Orogenic curvature: Integrating paleomagnetic and*
972 *structural analyses*, vol. 383, edited by A. B. Weil and A. Sussman, pp. 121–130, Geological
973 Society of America, Boulder., 2004.
- 974 Gutiérrez-Alonso, G., Fernández-Suárez, J., Weil, A. B., Brendan Murphy, J., Damian Nance,
975 R., Corf, F. and Johnston, S. T.: Self-subduction of the Pangaeon global plate, *Nat. Geosci.*,
976 1(8), 549–553, doi:10.1038/ngeo250, 2008a.
- 977 Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J. and Hamilton, M. A. A.: Rifting
978 along the northern Gondwana margin and the evolution of the Rheic Ocean: A Devonian age for
979 the El Castillo volcanic rocks (Salamanca, Central Iberian Zone), *Tectonophysics*, 461(1–4),
980 157–165, doi:10.1016/j.tecto.2008.01.013, 2008b.
- 981 Gutiérrez-Alonso, G., Fernández-Suárez, J., Jeffries, T. E. T. E. E., Johnston, S. T., Pastor-
982 Galán, D., Murphy, J. B. B., Franco, M. P. P., Gonzalo, J. C. C.: Diachronous post-orogenic
983 magmatism within a developing orocline in Iberia, *European Variscides*, *Tectonics*, 30(5), 17
984 PP.-17 PP., doi:10.1029/2010TC002845, 2011a.
- 985 Gutiérrez-Alonso, G., Murphy, J. B., Fernández-Suárez, J., Weil, A. B., Franco, M. P., Gonzalo,
986 J. C.: Lithospheric delamination in the core of Pangea: Sm-Nd insights from the Iberian mantle,
987 *Geology*, 39(2), 155–158, doi:10.1130/G31468.1, 2011b.
- 988 Gutiérrez-Alonso, G., Collins, A. S., Fernández-Suárez, J., Pastor-Galán, D., González-Clavijo,
989 E., Jourdan, F., Weil, A. B. and Johnston, S. T.: Dating of lithospheric buckling: $^{40}\text{Ar}/^{39}\text{Ar}$ ages
990 of syn-orocline strike-slip shear zones in northwestern Iberia, *Tectonophysics*, 643, 44–54,
991 doi:10.1016/j.tecto.2014.12.009, 2015.
- 992 Gutiérrez-Alonso, G., Fernández-Suárez, J., López-Carmona, A. and Gärtner, A.: Exhuming a
993 cold case: The early granodiorites of the northwest Iberian Variscan belt—A Visean magmatic
994 flare-up?, *Lithosphere*, 10(2), 194–216, doi:10.1130/L706.1, 2018.
- 995 Gutiérrez-Marco, J. C., Sá, A. A., García-Bellido, D. C. and Rábano, I.: The Bohemo-Iberian
996 regional chronostratigraphical scale for the Ordovician System and palaeontological correlations
997 within South Gondwana, *Lethaia*, 50(2), 258–295, doi:10.1111/let.12197, 2017.
- 998 Gutiérrez-Marco, J. C., Piçarra, J. M., Meireles, C. A., Cózar, P., García-Bellido, D. C., Pereira,
999 Z., Vaz, N., Pereira, S., Lopes, G., Oliveira, J. T., Quesada, C., Zamora, S., Esteve, J.,

- 1000 Colmenar, J., Bernárdez, E., Coronado, I., Lorenzo, S., Sá, A. A., Dias da Silva, Í., González-
1001 Clavijo, E., Díez-Montes, A. and Gómez-Barreiro, J.: Early Ordovician–Devonian Passive
1002 Margin Stage in the Gondwanan Units of the Iberian Massif, in *The Geology of Iberia: A*
1003 *Geodynamic Approach*, edited by C. Quesada and J. T. Oliveira, pp. 75–98, Springer
1004 International Publishing, Cham., 2019.
- 1005 Hindle, D., Besson, O. and Burkhard, M.: A model of displacement and strain for arc-shaped
1006 mountain belts applied to the Jura arc, *J. Struct. Geol.*, 22(9), 1285–1296, doi:10.1016/S0191-
1007 8141(00)00038-9, 2000.
- 1008 van Hinsbergen, D. J. J., Torsvik, T. H., Schmid, S. M., Mañenco, L. C., Maffione, M., Vissers, R.
1009 L. M., Gürer, D. and Spakman, W.: Orogenic architecture of the Mediterranean region and
1010 kinematic reconstruction of its tectonic evolution since the Triassic, *Gondwana Res.*, 81, 79–
1011 229, doi:10.1016/j.gr.2019.07.009, 2020.
- 1012 Hirt, A. M., Lowrie, W., Julivert, M., Arboleya, M. L., Lowric, A. M. H. W. and Arboleya, M. L.:
1013 Paleomagnetic results in support of a model for the origin of the Asturian arc, *Tectonophysics*,
1014 213(3–4), 321–339, 1992.
- 1015 Hnat, J. S. and van der Pluijm, B. A.: Foreland signature of indenter tectonics: Insights from
1016 calcite twinning analysis in the Tennessee salient of the Southern Appalachians, USA,
1017 *Lithosphere*, 3(5), 317–327, doi:10.1130/L151.1, 2011.
- 1018 Holmes, A.: A review of the continental drift hypothesis., *Min. Mag.*, 40, 205–209, 1929.
- 1019 Huang, W., Lippert, P. C., Zhang, Y., Jackson, M. J., Dekkers, M. J., Li, J., Hu, X., Zhang, B.,
1020 Guo, Z. and van Hinsbergen, D. J. J. J.: Remagnetization of carbonate rocks in southern Tibet:
1021 Perspectives from rock magnetic and petrographic investigations, *J. Geophys. Res. Solid Earth*,
1022 122(4), 2434–2456, doi:10.1002/2017JB013987, 2017.
- 1023 Izquierdo-Llavall, E., Sainz, A. C., Oliva-Urcia, B., Burmester, R., Pueyo, E. L. and Housen, B.:
1024 Multi-episodic remagnetization related to deformation in the Pyrenean Internal Sierras,
1025 *Geophys. J. Int.*, 201(2), 891–914, doi:10.1093/gji/ggv042, 2015.
- 1026 Izquierdo-Llavall, E., Casas-Sainz, A. M., Oliva-Urcia, B., Villalaín, J. J., Pueyo, E. and
1027 Scholger, R.: Rotational Kinematics of Basement Antiformal Stacks: Paleomagnetic Study of the
1028 Western Noguera Zone (Central Pyrenees), *Tectonics*, 37(10), 3456–3478,
1029 doi:10.1029/2018TC005153, 2018.
- 1030 Izquierdo-Llavall, E., Ayala, C., Pueyo, E. L., Casas-Sainz, A. M., Oliva-Urcia, B., Rubio, F.,
1031 Rodríguez-Pintó, A., Rey-Moral, C., Mediato, J. F. and García-Crespo, J.: Basement-Cover
1032 Relationships and Their Along-Strike Changes in the Linking Zone (Iberian Range, Spain): A
1033 Combined Structural and Gravimetric Study, *Tectonics*, 38(8), 2934–2960,
1034 doi:10.1029/2018TC005422, 2019.
- 1035 Jabaloy-Sánchez, A., Talavera, C., Gómez-Pugnaire, M. T., López-Sánchez-Vizcaíno, V.,
1036 Vázquez-Vílchez, M., Rodríguez-Peces, M. J. and Evans, N. J.: U-Pb ages of detrital zircons
1037 from the Internal Betics: A key to deciphering paleogeographic provenance and tectono-
1038 stratigraphic evolution, *Lithos*, 318–319, 244–266, doi:10.1016/j.lithos.2018.07.026, 2018.
- 1039 Jackson, M.: Diagenetic sources of stable remanence in remagnetized paleozoic cratonic

- 1040 carbonates: A rock magnetic study, *J. Geophys. Res. Solid Earth*, 95(B3), 2753–2761,
1041 doi:10.1029/JB095iB03p02753, 1990.
- 1042 Jacques, D., Muchez, P. and Sintubin, M.: Superimposed folding and W-Sn vein-type
1043 mineralisation in the Central Iberian Zone associated with late-Variscan oroclinal buckling: A
1044 structural analysis from the Regoufe area (Portugal), *Tectonophysics*, 742–743, 66–83,
1045 doi:10.1016/j.tecto.2018.05.021, 2018a.
- 1046 Jacques, D., Vieira, R., Muchez, P. and Sintubin, M.: Transpressional folding and associated
1047 cross-fold jointing controlling the geometry of post-orogenic vein-type W-Sn mineralization:
1048 examples from Minas da Panasqueira, Portugal, *Miner. Deposita*, 53(2), 171–194,
1049 doi:10.1007/s00126-017-0728-6, 2018b.
- 1050 Johnston, S. T.: The Great Alaskan Terrane Wreck: Reconciliation of paleomagnetic and
1051 geological data in the Northern Cordillera, *Earth Planet. Sci. Lett.*, 193(3–4), 259–272,
1052 doi:10.1016/S0012-821X(01)00516-7, 2001.
- 1053 Johnston, S. T., Weil, A. B. and Gutiérrez-Alonso, G.: Oroclines: Thick and thin, *GSA Bull.*,
1054 125(5–6), 643–663, doi:10.1130/B30765.1, 2013.
- 1055 Julivert, M.: L'évolution structurale de l'Arc Asturien, Paris., 1971.
- 1056 Julivert, M. and Arboleya, M. L.: A geometrical and kinematical approach to the nappe structure
1057 in an arcuate fold belt: the Cantabrian nappes (Hercynian chain, NW Spain), *J. Struct. Geol.*,
1058 6(5), 499–519, doi:10.1016/0191-8141(84)90061-0, 1984.
- 1059 Julivert, M. and Marcos, A.: superimposed folding under flexural conditions in the Cantabrian
1060 Zone (Hercynian Cordillera, Northwest Spain), *Am. J. Sci.*, 273(5), 353–375,
1061 doi:10.2475/ajs.273.5.353, 1973.
- 1062 Julivert, M., Fontboté, J. M., Ribeiro, A. and Nabais Conde, L. E.: Mapa Tectónico de la
1063 Península Ibérica y Baleares E: 1: 1.000. 000 y memoria explicativa, Publ IGME, 113, 1974.
- 1064 Julivert, M., Vegas, R., Roiz, J. M. and Martínez Rius, A.: La estructura de la extension SE de la
1065 Zona Centro-Iberica con metamorfismo de bajo grado, *Rios-Geol. Esp. I*, 339–380, 1983.
- 1066 Keppie, F.: How subduction broke up Pangaea with implications for the supercontinent cycle,
1067 *Geol. Soc. Lond. Spec. Publ.*, 424(1), 265–288, doi:10.1144/SP424.8, 2016.
- 1068 Kirsch, M., Keppie, J. D., Murphy, J. B. and Solari, L. A.: Permian-Carboniferous arc
1069 magmatism and basin evolution along the western margin of Pangea: Geochemical and
1070 geochronological evidence from the eastern Acatlan Complex, southern Mexico, *Geol. Soc. Am.*
1071 *Bull.*, 124(9–10), 1607–1628, doi:10.1130/B30649.1, 2012.
- 1072 Kollmeier, J. M., van der Pluijm, B. A. and Van der Voo, R.: Analysis of Variscan dynamics;
1073 early bending of the Cantabria-Asturias Arc, northern Spain, *Earth Planet. Sci. Lett.*, 181(1–2),
1074 203–216, doi:10.1016/S0012-821X(00)00203-X, 2000.
- 1075 Kroner, U. and Romer, R. L. L.: Two plates — Many subduction zones: The Variscan orogeny
1076 reconsidered, *Gondwana Res.*, 24(1), 298–329, doi:10.1016/j.gr.2013.03.001, 2013.

- 1077 Li, P. and Rosenbaum, G.: Does the Manning Orocline exist? New structural evidence from the
1078 inner hinge of the Manning Orocline (eastern Australia), *Gondwana Res.*, 25(4), 1599–1613,
1079 doi:10.1016/j.gr.2013.06.010, 2014.
- 1080 Li, P., Sun, M., Rosenbaum, G., Yuan, C., Safonova, I., Cai, K., Jiang, Y. and Zhang, Y.:
1081 Geometry, kinematics and tectonic models of the Kazakhstan Orocline, Central Asian Orogenic
1082 Belt, *J. Asian Earth Sci.*, 153(July 2017), 42–56, doi:10.1016/j.jseaes.2017.07.029, 2018.
- 1083 Li, P.-F., Rosenbaum, G. and Rubatto, D.: Triassic asymmetric subduction rollback in the
1084 southern New England Orogen (eastern Australia): the end of the Hunter-Bowen Orogeny, *Aust.*
1085 *J. Earth Sci.*, 59(6), 965–981, doi:10.1080/08120099.2012.696556, 2012.
- 1086 López-Carmona, A., Abati, J., Pitra, P. and Lee, J. K. W.: Retrogressed lawsonite blueschists
1087 from the NW Iberian Massif: P–T–t constraints from thermodynamic modelling and ⁴⁰Ar/³⁹Ar
1088 geochronology, *Contrib. Mineral. Petrol.*, 167(3), 987–987, doi:10.1007/s00410-014-0987-5,
1089 2014.
- 1090 López-Moro, F. J., López-Plaza, M., Gutiérrez-Alonso, G., Fernández-Suárez, J., López-
1091 Carmona, A., Hofmann, M. and Romer, R. L.: Crustal melting and recycling: geochronology and
1092 sources of Variscan syn-kinematic anatectic granitoids of the Tormes Dome (Central Iberian
1093 Zone). A U–Pb LA-ICP-MS study, *Int. J. Earth Sci.*, 107(3), 985–1004, doi:10.1007/s00531-017-
1094 1483-8, 2018.
- 1095 Lotze, F.: Zur gliederung der Variszichen der Iberischen Meseta., *Geotektonische Forschungen*,
1096 6, 78–92, 1945.
- 1097 Macaya, J., González-Lodeiro, F., Martínez-Catalán, J. R. and Alvarez, F.: Continuous
1098 deformation, ductile thrusting and backfolding of cover and basement in the Sierra de
1099 Guadarrama, Hercynian orogen of central Spain, *Tectonophysics*, 191(3), 291–309,
1100 doi:10.1016/0040-1951(91)90063-X, 1991.
- 1101 Maffione, M., Speranza, F. and Faccenna, C.: Bending of the Bolivian orocline and growth of the
1102 central Andean plateau: Paleomagnetic and structural constraints from the Eastern Cordillera
1103 (22–24°S, NW Argentina), *Tectonics*, 28, 23 PP.-23 PP., doi:10.1029/2008TC002402, 2009.
- 1104 Marcos, A. and Pulgar, J. A. A.: An Approach to the tectonostratigraphic evolution of the
1105 Cantabrian foreland thrust and fold belt, Hercynian Cordillera of NW Spain, *Neues Jahrb. Geol.*
1106 *Palaontologie-Abh.*, 163(2), 256–260, 1982.
- 1107 Marcos, A., Pérez-Estaún, A., Martínez Catalán, J. R. and Gutiérrez-Marco, J. C.: *Estratigrafía y*
1108 *Paleogeografía. Zona Asturoccidental-Leonesa*, 2004.
- 1109 Marshak, S.: KINEMATICS OF OROCLINE AND ARC FORMATION IN THIN-SKINNED
1110 OROGENS, *Tectonics*, 7(1), 73–86, 1988.
- 1111 Marshak, S.: Salients, Recesses, Arcs, Oroclines, and Syntaxes — A Review of Ideas
1112 Concerning the Formation of Map-view Curves in Fold-thrust Belts, *Thrust Tecton. Hydrocarb.*
1113 *Syst. AAPG Mem.* 82, 82(1893), 131–156, 2004.
- 1114 Martín-Algarra, A., García-Casco, A., Gómez-Pugnaire, M. T., Jabaloy-Sánchez, A., Laborda-
1115 López, C., López Sánchez-Vizcaíno, V., Mazzoli, S., Navas-Parejo, P., Perrone, V., Rodríguez-

- 1116 Cañero, R. and Sánchez-Navas, A.: Paleozoic Basement and Pre-Alpine History of the Betic
1117 Cordillera, in *The Geology of Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T.
1118 Oliveira, pp. 261–305, Springer International Publishing, Cham., 2019.
- 1119 Martínez Catalán, J. R.: Are the oroclines of the Variscan belt related to late Variscan strike-slip
1120 tectonics?, *Terra Nova*, 23(4), 241–247, doi:10.1111/j.1365-3121.2011.01005.x, 2011.
- 1121 Martínez Catalán, J. R.: The Central Iberian arc, an orocline centered in the Iberian Massif and
1122 some implications for the Variscan belt, *Int. J. Earth Sci.*, 101(5), 1–16, doi:10.1007/s00531-
1123 011-0715-6, 2012.
- 1124 Martínez Catalan, J. R., Rodríguez, M. P. H., Alonso, P. V., Pérez-Estaún, A. and Lodeiro, F. G.:
1125 Lower Paleozoic extensional tectonics in the limit between the West Asturian-Leonese and
1126 Central Iberian Zones of the Variscan Fold-Belt in NW Spain, *Geol. Rundsch.*, 81(2), 545–560,
1127 doi:10.1007/BF01828614, 1992.
- 1128 Martínez Catalán, J. R., Martínez Poyatos, D. and Bea, F.: Zona Centroibérica, *Geol. Esp.*, 68–
1129 133, 2004.
- 1130 Martínez Catalán, J. R., Arenas, R., Abati, J., Martínez, S. S., García, F. D., Suárez, J. F.,
1131 Cuadra, P. G., Castiñeiras, P., Barreiro, J. G., Montes, A. D., Clavijo, E. G., Pascual, F. J. R.,
1132 Andonaegui, P., Jeffries, T. E., Alcock, J. E., Fernández, R. D. and Carmona, A. L.: A rootless
1133 suture and the loss of the roots of a mountain chain: The Variscan belt of NW Iberia, *Comptes*
1134 *Rendus Geosci.*, 341(2), 114–126, doi:10.1016/j.crte.2008.11.004, 2009.
- 1135 Martínez Catalan, J. R., Rubio Pascual, F. J., Díez Montes, A., Díez Fernández, R., Gómez
1136 Barreiro, J., Dias da Silva, I., González Clavijo, E., Ayarza, P., Alcock, J. E., Montes, A. D.,
1137 Fernández, R. D., Barreiro, J. G., Dias da Silva, I., Clavijo, E. G., Ayarza, P., Alcock, J. E., Díez
1138 Montes, A., Díez Fernández, R., Gómez Barreiro, J., Dias da Silva, I., González Clavijo, E.,
1139 Ayarza, P., Alcock, J. E., Montes, A. D., Fernández, R. D., Barreiro, J. G., Dias da Silva, I.,
1140 Clavijo, E. G., Ayarza, P. and Alcock, J. E.: The late Variscan HT/LP metamorphic event in NW
1141 and Central Iberia: relationships to crustal thickening, extension, orocline development and
1142 crustal evolution, *Geol. Soc. Lond. Spec. Publ.*, 405(1), 225–247, doi:10.1144/SP405.1, 2014.
- 1143 Martínez Catalán, J. R., Aerden, D. G. A. M. and Carreras, J.: The “Castilian bend” of Rudolf
1144 Staub (1926): historical perspective of a forgotten orocline in Central Iberia, *Swiss J. Geosci.*,
1145 108(2–3), 289–303, doi:10.1007/s00015-015-0202-3, 2015.
- 1146 Martínez Catalán, J. R., Gómez Barreiro, J., Dias da Silva, Í., Chichorro, M., López-Carmona,
1147 A., Castiñeiras, P., Abati, J., Andonaegui, P., Fernández-Suárez, J., González Cuadra, P. and
1148 Benítez-Pérez, J. M.: Variscan Suture Zone and Suspect Terranes in the NW Iberian Massif:
1149 Allochthonous Complexes of the Galicia-Trás os Montes Zone (NW Iberia), in *The Geology of*
1150 *Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T. Oliveira, pp. 99–130, Springer
1151 International Publishing, Cham., 2019.
- 1152 Martínez-Catalán, J. R.: A non-cylindrical model for the northwest Iberian allochthonous
1153 terranes and their equivalents in the Hercynian belt of Western Europe, *Tectonophysics*, 179(3–
1154 4), 253–272, doi:10.1016/0040-1951(90)90293-H, 1990.
- 1155 McWilliams, C. K., Kunk, M. J., Wintsch, R. P. and Bish, D. L.: Determining ages of multiple
1156 muscovite-bearing foliations in phyllonites using the $^{40}\text{Ar}/^{39}\text{Ar}$ step heating method:

- 1157 Applications to the alleghanian orogeny in central new England, *Am. J. Sci.*, 313(10), 996–1016,
1158 doi:10.2475/10.2013.02, 2013.
- 1159 Meijers, M. J. M., Smith, B., Pastor-Galán, D., Degenaar, R., Sadradze, N., Adamia, S.,
1160 Sahakyan, L., Avagyan, A., Sosson, M., Rolland, Y., Langereis, C. G. and Müller, C.:
1161 Progressive orocline formation in the Eastern Pontides-Lesser Caucasus, *Geol. Soc. Spec.*
1162 *Publ.*, 428(1), doi:10.1144/SP428.8, 2017.
- 1163 Merino-Tomé, O. A., Bahamonde, J. R., Colmenero, J. R., Heredia, N., Villa, E. and Farias, P.:
1164 Emplacement of the Cuera and Picos de Europa imbricate system at the core of the Iberian-
1165 Armorican arc (Cantabrian zone, north Spain): New precisions concerning the timing of arc
1166 closure, *Bull. Geol. Soc. Am.*, 121(5–6), 729–751, doi:10.1130/B26366.1, 2009.
- 1167 Molina Garza, R. S. and Zijdeveld, J. D. a.: Paleomagnetism of Paleozoic strata, Brabant and
1168 Ardennes Massifs, Belgium: Implications of prefolding and postfolding Late Carboniferous
1169 secondary magnetizations for European apparent polar wander, *J. Geophys. Res.*, 101(B7),
1170 15799–15799, doi:10.1029/96JB00325, 1996.
- 1171 Moral, B. del and Sarmiento, G. N.: Conodontos del Katiense (Ordovícico Superior) del sector
1172 meridional de la Zona Centroibérica (España), *Rev. Esp. Micropaleontol.*, 40(3), 169–245, 2008.
- 1173 Munha, J.: Metamorphic Evolution of the South Portuguese/Pulo Do Lobo Zone, in *Pre-*
1174 *Mesozoic Geology of Iberia*, edited by R. D. Dallmeyer and E. M. Garcia, pp. 363–368,
1175 Springer, Berlin, Heidelberg., 1990.
- 1176 Munha, J., Barriga, F. J. a. S. and Kerrich, R.: High 18 O ore-forming fluids in volcanic-hosted
1177 base metal massive sulfide deposits; geologic, 18 O/ 16 O, and D/H evidence from the Iberian
1178 pyrite belt; Crandon, Wisconsin; and Blue Hill, Maine, *Econ. Geol.*, 81(3), 530–552, doi:10.2113/
1179 gsecongeo.81.3.530, 1986.
- 1180 Murphy, J. B., Gutierrez-Alonso, G., Fernandez-Suarez, J. and Braid, J. A.: Probing crustal and
1181 mantle lithosphere origin through Ordovician volcanic rocks along the Iberian passive margin of
1182 Gondwana, *Tectonophysics*, 461(1–4), 166–180, 2008.
- 1183 Nance, R. D., Gutiérrez-Alonso, G., Keppie, J. D., Linnemann, U., Murphy, J. B., Quesada, C.,
1184 Strachan, R. A. and Woodcock, N. H.: Evolution of the Rheic Ocean, *Gondwana Res.*, 17(2–3),
1185 194–222, doi:10.1016/j.gr.2009.08.001, 2010.
- 1186 Oliveira, J. T., Quesada, C., Pereira, Z., Matos, J. X., Solá, A. R., Rosa, D., Albardeiro, L., Díez-
1187 Montes, A., Morais, I., Inverno, C., Rosa, C. and Relvas, J.: South Portuguese Terrane: A
1188 Continental Affinity Exotic Unit, in *The Geology of Iberia: A Geodynamic Approach*, edited by C.
1189 Quesada and J. T. Oliveira, pp. 173–206, Springer International Publishing, Cham., 2019a.
- 1190 Oliveira, J. T., González-Clavijo, E., Alonso, J., Armendáriz, M., Bahamonde, J. R., Braid, J. A.,
1191 Colmenero, J. R., Dias da Silva, Í., Fernandes, P., Fernández, L. P., Gabaldón, V., Jorge, R. S.,
1192 Machado, G., Marcos, A., Merino-Tomé, Ó., Moreira, N., Murphy, J. B., Pinto de Jesus, A.,
1193 Quesada, C., Rodrigues, B., Rosales, I., Sanz-López, J., Suárez, A., Villa, E., Piçarra, J. M. and
1194 Pereira, Z.: Synorogenic Basins, in *The Geology of Iberia: A Geodynamic Approach*, edited by
1195 C. Quesada and J. T. Oliveira, pp. 349–429, Springer International Publishing, Cham., 2019b.
- 1196 P. Farias, G. G.: Aportaciones al conocimiento de la litoestratigrafía y estructura de Galicia

- 1197 Central, Mem. Fac. Ciênc. Universidade Porto, 411–431, 1987.
- 1198 Palero-Fernández, F. J., Martin-Izard, A., Zarzalejos Prieto, M. and Mansilla-Plaza, L.:
 1199 Geological context and plumbotectonic evolution of the giant Almadén Mercury Deposit, Ore
 1200 Geol. Rev., 64, 71–88, doi:10.1016/j.oregeorev.2014.06.013, 2015.
- 1201 Parés, J. M. and Van der Voo, R.: Paleozoic paleomagnetism of Almadén, Spain: A cautionary
 1202 note, J. Geophys. Res., 97(B6), 9353–9356, doi:10.1029/91JB03073, 1992.
- 1203 Parés, J. M., Van der Voo, R., Stamatakis, J. and Pérez-Estaún, A.: Remagnetizations and
 1204 postfolding oroclinal rotations in the Cantabrian/Asturian arc, northern Spain, Tectonics, 13(6),
 1205 1461–1471, doi:10.1029/94TC01871, 1994.
- 1206 Pastor-Galán, D., Gutiérrez-Alonso, G., Meere, P. A. and Mulchrone, K. F.: Factors affecting
 1207 finite strain estimation in low-grade, low-strain clastic rocks, J. Struct. Geol., 31(12), 1586–1596,
 1208 doi:10.1016/j.jsg.2009.08.005, 2009.
- 1209 Pastor-Galán, D., Gutiérrez-Alonso, G. and Weil, A. B.: Orocline timing through joint analysis:
 1210 Insights from the Ibero-Armorican Arc, Tectonophysics, 507(1), 31–46,
 1211 doi:10.1016/j.tecto.2011.05.005, 2011.
- 1212 Pastor-Galán, D., Gutiérrez-Alonso, G., Zulauf, G. and Zanella, F.: Analogue modeling of
 1213 lithospheric-scale orocline buckling: Constraints on the evolution of the Iberian-Armorican arc,
 1214 Bull. Geol. Soc. Am., 124(7–8), doi:10.1130/B30640.1, 2012a.
- 1215 Pastor-Galán, D., Gutiérrez-Alonso, G., Mulchrone, K. F. and Huerta, P.: Conical folding in the
 1216 core of an orocline. A geometric analysis from the Cantabrian Arc (Variscan Belt of NW Iberia),
 1217 J. Struct. Geol., 39, 210–223, doi:10.1016/j.jsg.2012.02.010, 2012b.
- 1218 Pastor-Galán, D., Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J., Hofmann, M.
 1219 and Linnemann, U.: Provenance analysis of the Paleozoic sequences of the northern
 1220 Gondwana margin in NW Iberia: Passive margin to Variscan collision and orocline development,
 1221 Gondwana Res., 23(3), 1089–1103, doi:10.1016/j.gr.2012.06.015, 2013a.
- 1222 Pastor-Galán, D., Gutiérrez-Alonso, G., Fernández-Suárez, J., Murphy, J. B. and Nieto, F.:
 1223 Tectonic evolution of NW Iberia during the Paleozoic inferred from the geochemical record of
 1224 detrital rocks in the Cantabrian Zone, Lithos, 182–183, 221–228,
 1225 doi:10.1016/j.lithos.2013.09.007, 2013b.
- 1226 Pastor-Galán, D., Martín-Merino, G. and Corrochano, D.: Timing and structural evolution in the
 1227 limb of an orocline: The Pisuerga-Carrión Unit (southern limb of the Cantabrian Orocline, NW
 1228 Spain), Tectonophysics, 622, 110–121, doi:10.1016/j.tecto.2014.03.004, 2014.
- 1229 Pastor-Galán, D., Ursem, B., Meere, P. A. and Langereis, C.: Extending the Cantabrian
 1230 Orocline to two continents (from Gondwana to Laurussia). Paleomagnetism from South Ireland,
 1231 Earth Planet. Sci. Lett., 432, doi:10.1016/j.epsl.2015.10.019, 2015a.
- 1232 Pastor-Galán, D., Groenewegen, T., Brouwer, D., Krijgsman, W., Dekkers, M. J., Pastor-galán,
 1233 D., Groenewegen, T., Brouwer, D., Krijgsman, W. and Dekkers, M. J.: One or two oroclines in
 1234 the Variscan orogen of Iberia? Implications for Pangea amalgamation, Geology, 43(6), 527–
 1235 530, doi:10.1130/G36701.1, 2015b.

- 1236 Pastor-Galán, D., Dekkers, M. J., Gutiérrez-Alonso, G., Brouwer, D., Groenewegen, T.,
1237 Krijgsman, W., Fernández-Lozano, J., Yenes, M. and Álvarez-Lobato, F.: Paleomagnetism of
1238 the Central Iberian curve's putative hinge: Too many oroclines in the Iberian Variscides,
1239 *Gondwana Res.*, 39, 96–113, doi:10.1016/j.gr.2016.06.016, 2016.
- 1240 Pastor-Galán, D., Mulchrone, K. F., Koymans, M. R. , van Hinsbergen, D. J. J. and Langereis,
1241 C. G. : Bootstrapped total least squares orocline test: A robust method to quantify vertical-axis
1242 rotation patterns in orogens, with examples from the Cantabrian and Aegean oroclines,
1243 *Lithosphere*, 9(3), L547.1-L547.1, doi:10.1130/L547.1, 2017a.
- 1244 Pastor-Galán, D., Gutiérrez-Alonso, G., Dekkers, M. J. M. J. and Langereis, C. G.:
1245 Paleomagnetism in Extremadura (Central Iberian zone, Spain) Paleozoic rocks: extensive
1246 remagnetizations and further constraints on the extent of the Cantabrian orocline, *J. Iber. Geol.*,
1247 43(4), 583–600, doi:10.1007/s41513-017-0039-x, 2017b.
- 1248 Pastor-Galán, D., Pueyo, E. L., Diederer, M., García-Lasanta, C. and Langereis, C. G.: Late
1249 Paleozoic Iberian Orocline(s) and the Missing Shortening in the Core of Pangea.
1250 Paleomagnetism From the Iberian Range, *Tectonics*, 37(10), 3877–3892,
1251 doi:10.1029/2018TC004978, 2018.
- 1252 Pastor-Galán, D., Nance, R. D., Murphy, J. B. and Spencer, C. J.: Supercontinents: myths,
1253 mysteries, and milestones, *Geol. Soc. Lond. Spec. Publ.*, 470, 39–64, doi:10.1144/SP470.16,
1254 2019a.
- 1255 Pastor-Galán, D., Dias da Silva, Í. F., Groenewegen, T. and Krijgsman, W.: Tangled up in folds:
1256 tectonic significance of superimposed folding at the core of the Central Iberian curve (West
1257 Iberia), *Int. Geol. Rev.*, 61(2), 240–255, doi:10.1080/00206814.2017.1422443, 2019b.
- 1258 Pereira, I., Dias, R., Bento, T. and Mata, J.: Exhumation of a migmatite complex along a
1259 transpressive shear zone : inferences from the Variscan Juzbado – Penalva do Castelo Shear
1260 Zone (Central Iberian Zone), *J. Geol. Soc.*, doi:10.1144/jgs2016-159, 2017.
- 1261 Pereira, M. F., Chichorro, M., Silva, J. B., Ordóñez-Casado, B., Lee, J. K. W. W. and Williams, I.
1262 S.: Early carboniferous wrenching, exhumation of high-grade metamorphic rocks and basin
1263 instability in SW Iberia: Constraints derived from structural geology and U–Pb and 40Ar–39Ar
1264 geochronology, *Tectonophysics*, 558, 28–44, doi:10.1016/j.tecto.2012.06.020, 2012.
- 1265 Pereira, M. F., Díez Fernández, R., Gama, C., Hofmann, M., Gärtner, A. and Linnemann, U.: S-
1266 type granite generation and emplacement during a regional switch from extensional to
1267 contractional deformation (Central Iberian Zone, Iberian autochthonous domain, Variscan
1268 Orogeny), *Int. J. Earth Sci.*, 107(1), 251–267, doi:10.1007/s00531-017-1488-3, 2018.
- 1269 Pérez-Cáceres, I., Poyatos, D. M., Simancas, J. F. and Azor, A.: The elusive nature of the
1270 Rheic Ocean suture in SW Iberia, *Tectonics*, 34(12), 2429–2450, doi:10.1002/2015TC003947,
1271 2015.
- 1272 Pérez-Cáceres, I., Simancas, J. F., Martínez Poyatos, D., Azor, A. and González Lodeiro, F.:
1273 Oblique collision and deformation partitioning in the SW Iberian Variscides, *Solid Earth*, 7(3),
1274 857–872, doi:https://doi.org/10.5194/se-7-857-2016, 2016.
- 1275 Pérez-Cáceres, I., Martínez Poyatos, D., Simancas, J. F. and Azor, A.: Testing the Avalonian

- 1276 affinity of the South Portuguese Zone and the Neoproterozoic evolution of SW Iberia through
1277 detrital zircon populations, *Gondwana Res.*, 42, 177–192, doi:10.1016/j.gr.2016.10.010, 2017.
- 1278 Pérez-Cáceres, I., Martínez Poyatos, D. J., Vidal, O., Beyssac, O., Nieto, F., Simancas, J. F.,
1279 Azor, A. and Bourdelle, F.: Deciphering the metamorphic evolution of the Pulo do Lobo
1280 metasedimentary domain (SW Iberian Variscides), *Solid Earth*, 11(2), 469–488,
1281 doi:https://doi.org/10.5194/se-11-469-2020, 2020.
- 1282 Pérez-Estaún, A., Bastida, F., Alonso, J. L., Marquinez, J., Aller, J., Alvarezmarron, J., Marcos,
1283 A. and Pulgar, J. A.: A THIN-SKINNED TECTONICS MODEL FOR AN ARCUATE FOLD AND
1284 THRUST BELT - THE CANTABRIAN ZONE (VARISCAN IBERO-ARMORICAN ARC),
1285 *Tectonics*, 7(3), 517–537, 1988.
- 1286 Pérez-Estaún, A., Bastida, F., Martínez Catalán, J. R., Gutiérrez-Marco, J. C., Marcos, A. and
1287 Pulgar, J.: Stratigraphy of the West Asturian-Leonese Zone, Springer. [online] Available from:
1288 https://digital.csic.es/handle/10261/30719 (Accessed 6 April 2020), 1990.
- 1289 Pérez-Estaún, A., Martinezcatalan, J. R. and Bastida, F.: CRUSTAL THICKENING AND
1290 DEFORMATION SEQUENCE IN THE FOOTWALL TO THE SUTURE OF THE VARISCAN
1291 BELT OF NORTHWEST SPAIN, *Tectonophysics*, 191(3–4), 243–253, 1991.
- 1292 Perroud, H., Bonhommet, N. and Ribeiro, A.: Paleomagnetism of Late Paleozoic igneous rocks
1293 from southern Portugal, *Geophys. Res. Lett.*, 12(1), 45–48, doi:10.1029/GL012i001p00045,
1294 1985.
- 1295 Perroud, H., Calza, F. and Khattach, D.: Paleomagnetism of the Silurian Volcanism at Almaden,
1296 Southern Spain, *J. Geophys. Res.-Solid Earth Planets*, 96(B2), 1949–1962,
1297 doi:10.1029/90JB02226, 1991.
- 1298 Pin, C., Paquette, J. L., Zalduegui, J. F. S. and Ibarguchi, J. I. G.: Early Devonian
1299 suprasubduction-zone ophiolite related to incipient collisional processes in the Western
1300 Variscan Belt: The Sierra de Careón unit, Ordenes Complex, Galicia, in *Variscan-Appalachian*
1301 *dynamics: The building of the late Paleozoic basement*, Geological Society of America., 2002.
- 1302 Pueyo, E. L., Mauritsch, H. J., Gawlick, H.-J., Scholger, R. and Frisch, W.: New evidence for
1303 block and thrust sheet rotations in the central northern Calcareous Alps deduced from two
1304 pervasive remagnetization events, *Tectonics*, 26(5), n/a-n/a, doi:10.1029/2006TC001965, 2007.
- 1305 Pueyo, E. L., Sussman, A. J., Oliva-Urcia, B. and Cifelli, F.: Palaeomagnetism in fold and thrust
1306 belts: Use with caution, *Geol. Soc. Spec. Publ.*, 425(1), 259–276, doi:10.1144/SP425.14, 2016.
- 1307 Quesada, C.: The Ossa-Morena Zone of the Iberian Massif: a tectonostratigraphic approach to
1308 its evolution, *Z. Dtsch. Ges. Für Geowiss.*, 585–595, doi:10.1127/1860-1804/2006/0157-0585,
1309 2006.
- 1310 Quesada, C. and Dallmeyer, R. D. D.: Tectonothermal evolution of the Badajoz-Córdoba shear
1311 zone (SW Iberia): characteristics and ⁴⁰Ar/³⁹Ar mineral age constraints, *Tectonophysics*,
1312 231(1–3), 195–213, doi:10.1016/0040-1951(94)90130-9, 1994.
- 1313 Quesada, C., Braid, J. A., Fernandes, P., Ferreira, P., Jorge, R. S., Matos, J. X., Murphy, J. B.,
1314 Oliveira, J. T., Pedro, J. and Pereira, Z.: SW Iberia Variscan Suture Zone: Oceanic Affinity

- 1315 Units, in *The Geology of Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T.
1316 Oliveira, pp. 131–171, Springer International Publishing, Cham., 2019.
- 1317 Rankin, D. W.: Appalachian salients and recesses: Late Precambrian continental breakup and
1318 the opening of the Iapetus Ocean, *J. Geophys. Res.* 1896-1977, 81(32), 5605–5619,
1319 doi:10.1029/JB081i032p05605, 1976.
- 1320 Ribeiro, A., Munhá, J., Dias, R., Mateus, A., Pereira, E., Ribeiro, L., Fonseca, P., Araújo, A.,
1321 Oliveira, T., Romão, J., Chaminé, H., Coke, C. and Pedro, J.: Geodynamic evolution of the SW
1322 Europe Variscides, *Tectonics*, 26(6), doi:10.1029/2006TC002058, 2007.
- 1323 Ries, A. C. and Shackleton, R. M.: Catazonal Complexes of North-West Spain and North
1324 Portugal, Remnants of a Hercynian Thrust Plate, *Nat. Phys. Sci.*, 234(47), 65–68,
1325 doi:10.1038/physci234065a0, 1971.
- 1326 Ries, A. C. and Shackleton, R. M.: Patterns of Strain Variation in Arcuate Fold Belts, *Philos.*
1327 *Trans. R. Soc. Lond. Ser. Math. Phys. Sci.*, 283(1312), 281–288, 1976.
- 1328 Rodríguez-Cañero, R., Jabaloy-Sánchez, A., Navas-Parejo, P., Martín-Algarra, A., Rodríguez,
1329 R., Antonio, C., Sánchez, J., Navas, P., Agustín, P. and Algarra, M.: Linking Palaeozoic
1330 palaeogeography of the Betic Cordillera to the Variscan Iberian Massif : new insight through the
1331 first conodonts of the Nevado-Filábride Complex, *Int. J. Earth Sci.*, 107(5), 1791–1806,
1332 doi:10.1007/s00531-017-1572-8, 2018.
- 1333 Rosenbaum, G.: Geodynamics of oroclinal bending: Insights from the Mediterranean, *J.*
1334 *Geodyn.*, 82, 5–15, doi:10.1016/j.jog.2014.05.002, 2014.
- 1335 Rubio Pascual, F. J., Arenas, R., Martínez Catalán, J. R., Rodríguez Fernández, L. R. and
1336 Wijbrans, J. R.: Thickening and exhumation of the Variscan roots in the Iberian Central System:
1337 Tectonothermal processes and $^{40}\text{Ar}/^{39}\text{Ar}$ ages, *Tectonophysics*, 587, 207–221,
1338 doi:10.1016/j.tecto.2012.10.005, 2013.
- 1339 Rubio Pascual, F. J., López-Carmona, A. and Arenas, R.: Thickening vs. extension in the
1340 Variscan belt: P–T modelling in the Central Iberian autochthon, *Tectonophysics*, 681, 144–158,
1341 doi:10.1016/j.tecto.2016.02.033, 2016.
- 1342 Sánchez-García, T., Chichorro, M., Solá, A. R., Álvaro, J. J., Díez-Montes, A., Bellido, F.,
1343 Ribeiro, M. L., Quesada, C., Lopes, J. C., Dias da Silva, Í., González-Clavijo, E., Gómez
1344 Barreiro, J. and López-Carmona, A.: The Cambrian-Early Ordovician Rift Stage in the
1345 Gondwanan Units of the Iberian Massif, in *The Geology of Iberia: A Geodynamic Approach*,
1346 edited by C. Quesada and J. T. Oliveira, pp. 27–74, Springer International Publishing, Cham.,
1347 2019.
- 1348 Schulz, G.: Descripción geológica de Asturias: Publicada de Real Órden. Con un atlas, José
1349 Gonzalez., 1858.
- 1350 Schwartz, S. Y. and Vandervoo, R.: PALEOMAGNETIC EVALUATION OF THE OROCLINE
1351 HYPOTHESIS IN THE CENTRAL AND SOUTHERN APPALACHIANS, *Geophys. Res. Lett.*,
1352 10(7), 505–508, 1983.
- 1353 Shaw, J. and Johnston, S. T.: Terrane wrecks (coupled oroclinal) and paleomagnetic inclination

- 1354 anomalies, *Earth-Sci. Rev.*, 154, 191–209, doi:10.1016/j.earscirev.2016.01.003, 2016.
- 1355 Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Weil, A. B.: Oroclines of the Variscan
1356 orogen of Iberia: Paleocurrent analysis and paleogeographic implications, *Earth Planet. Sci.*
1357 *Lett.*, 329–330, 60–70, doi:10.1016/j.epsl.2012.02.014, 2012.
- 1358 Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Pastor-Galán, D.: Provenance variability
1359 along the Early Ordovician north Gondwana margin: Paleogeographic and tectonic implications
1360 of U-Pb detrital zircon ages from the Armorican Quartzite of the Iberian Variscan belt, *Bull. Geol.*
1361 *Soc. Am.*, 126(5–6), 702–719, doi:10.1130/B30935.1, 2014.
- 1362 Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A
1363 structural study of the Cantabrian orocline, NW Iberian Massif, *Lithosphere*, 7(6), 653–661,
1364 doi:10.1130/L461.1, 2015.
- 1365 Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A
1366 structural study of the Cantabrian orocline, NW Iberian Massif, *Lithosphere*, 8(1), 97–97,
1367 doi:10.1130/L461.1, 2016.
- 1368 Silva, Í. D. D., Valverde-Vaquero, P., González-Clavijo, E., Díez-Montes, A. and Catalán, J. R.
1369 M.: Structural and stratigraphical significance of U–Pb ages from the Mora and Saldanha
1370 volcanic complexes (NE Portugal, Iberian Variscides), *Geol. Soc. Lond. Spec. Publ.*, 405(1),
1371 115–135, doi:10.1144/SP405.3, 2014.
- 1372 Simancas, J. F., Carbonell, R., Lodeiro, F. G., Estaun, A., Juhlin, C., Ayarza, P., Kashubin, A.,
1373 Azor, A., Poyatos, D. M., Almodovar, G. R., Pascual, E., Saez, R. and Exposito, I.: Crustal
1374 structure of the transpressional Variscan orogen of SW Iberia: SW Iberia deep seismic reflection
1375 profile (IBERSEIS), *Tectonics*, 22(6), 25–25, 2003.
- 1376 Simancas, J. F., Ayarza, P., Azor, a., Carbonell, R., Martínez Poyatos, D., Pérez-Estaún, a. and
1377 González Lodeiro, F.: A seismic geotraverse across the Iberian Variscides: Orogenic
1378 shortening, collisional magmatism, and orocline development, *Tectonics*, 32(i), n/a–n/a,
1379 doi:10.1002/tect.20035, 2013.
- 1380 Solís-Alulima, B. E., López-Carmona, A., Gutiérrez Alonso, G. and Álvarez Valero, A. M.:
1381 Petrologic and thermobarometric study of the Riás schists (NW Iberian Massif), *Bol. Geológico*
1382 *Min.*, 130(3), 445–464, doi:10.21701/bolgeomin.130.3.004, 2019.
- 1383 Stampfli, G. M. and Borel, G. D.: The TRANSMED Transects in Space and Time: Constraints
1384 from Paleotectonic Evolution of the Mediterranean Domain., edited by W. Cavazza, F. M.
1385 Roure, W. Spakman, G. M. Stampfli, and P. A. Ziegler, pp. 53–80, Springer, Berlin., 2004.
- 1386 Stampfli, G. M., Hochard, C., Vérard, C., Wilhem, C. and VonRaumer, J.: The Formation of
1387 Pangea, *Tectonophysics*, 593, 1–19, doi:10.1016/j.tecto.2013.02.037, 2013.
- 1388 Staub, R.: Gedanken zur Tektonik Spaniens., Zürich., 1926.
- 1389 Stewart, S. A.: Paleomagnetic analysis of fold kinematics and implications for geological models
1390 of the Cantabrian/Asturian arc, north Spain, *J. Geophys. Res. Solid Earth*, 100(B10), 20079–
1391 20094, doi:10.1029/95JB01482, 1995.

- 1392 Tait, J. A., Bachtadse, V. and Soffel, H. C.: Eastern Variscan fold belt : Paleomagnetic evidence
1393 for oroclinal bending, *Geology*, 24, 871–874, doi:10.1130/0091-7613(1996)024<0871, 1996.
- 1394 Tauxe, L.: *Essentials of Paleomagnetism*, Univ of California Press., 2010.
- 1395 Thomas, W. A.: Evolution of Appalachian-Ouachita Salients and Recesses from Reentrants and
1396 Promontories in the Continental Margin, *Am. J. Sci.*, 277(10), 1233–1278,
1397 doi:10.2475/ajs.277.10.1233, 1977.
- 1398 Thomas, W. A.: Genetic relationship of rift-stage crustal structure, terrane accretion, and
1399 foreland tectonics along the southern Appalachian-Ouachita orogen, *J. Geodyn.*, 37(3), 549–
1400 563, doi:10.1016/j.jog.2004.02.020, 2004.
- 1401 Tohver, E., Weil, A. B. B. B., Solum, J. G. G. G. and Hall, C. M. M. M.: Direct dating of
1402 carbonate remagnetization by $^{40}\text{Ar}/^{39}\text{Ar}$ analysis of the smectite–illite transformation, *Earth*
1403 *Planet. Sci. Lett.*, 274(3–4), 524–530, doi:10.1016/j.epsl.2008.08.002, 2008.
- 1404 Toit, A. L. Du: *Our wandering continents: an hypothesis of continental drifting*, Oliver and Boyd.
1405 [online] Available from: <http://books.google.es/books?id=iDZEAAAIAAJ>, 1937.
- 1406 Valladares, M. I., Barba, P., Ugidos, J. M., Colmenero, J. R. and Armenteros, I.: Upper
1407 Neoproterozoic-Lower Cambrian sedimentary successions in the Central Iberian Zone (Spain):
1408 sequence stratigraphy, petrology and chemostratigraphy. Implications for other European
1409 zones, *Int. J. Earth Sci.*, 89(1), 2–20, 2000.
- 1410 Van der Voo, R.: Paleomagnetism, oroclines, and growth of the continental crust, *GSA Today*,
1411 14(12), 4–9, doi:10.1130/1052-5173(2004), 2004.
- 1412 Vergés, J.: Estudio del Complejo volcánico-sedimentario del Devónico y de la estructura de la
1413 terminación oriental del sinclinal de Almadén (Ciudad Real), in *Libro Jubilar JM Rios. Tomo 3*,
1414 pp. 215–229., 1983.
- 1415 Vissers, R. L. M., van Hinsbergen, D. J. J., van der Meer, D. G. and Spakman, W.: Cretaceous
1416 slab break-off in the Pyrenees: Iberian plate kinematics in paleomagnetic and mantle reference
1417 frames, *Gondwana Res.*, 34, 49–59, doi:10.1016/J.GR.2016.03.006, 2016.
- 1418 van der Voo, R., Stamatakis, J. A. and Pares, J. M.: Kinematic constraints on thrust-belt
1419 curvature from syndeformational magnetizations in the Lagos del Valle Syncline in the
1420 Cantabrian Arc, Spain, *J. Geophys. Res.-Solid Earth*, 102(B5), 10105–10119, 1997.
- 1421 Weil, A., Pastor-Galán, D., Johnston, S. T. and Gutiérrez-Alonso, G.: Late/Post Variscan
1422 Orocline Formation and Widespread Magmatism, in *The Geology of Iberia: A Geodynamic*
1423 *Approach*, edited by C. Quesada and J. T. Oliveira, pp. 527–542, Springer International
1424 Publishing, Cham., 2019.
- 1425 Weil, A. B.: Kinematics of orocline tightening in the core of an arc: Paleomagnetic analysis of
1426 the Ponga Unit, Cantabrian Arc, northern Spain, *Tectonics*, 25(3), n/a-n/a,
1427 doi:10.1029/2005TC001861, 2006.
- 1428 Weil, A. B. and Sussman, A. J.: Classifying curved orogens based on timing relationships
1429 between structural development and vertical-axis rotations, vol. 383, edited by A. J. Sussman

- 1430 and A. B. Weil, pp. 1–16, Geological Society of America., 2004.
- 1431 Weil, A. B., Van der Voo, R., van der Pluijm, B. A. and Parés, J. M.: The formation of an orocline
1432 by multiphase deformation: a paleomagnetic investigation of the Cantabria–Asturias Arc
1433 (northern Spain), *J. Struct. Geol.*, 22(6), 735–756, doi:10.1016/S0191-8141(99)00188-1, 2000.
- 1434 Weil, A. B., van der Voo, R. and van der Pluijm, B. A.: Oroclinal bending and evidence against
1435 the Pangea megashear: The Cantabria-Asturias arc (northern Spain), *Geology*, 29(11), 991–
1436 994, 2001.
- 1437 Weil, A. B., Van der Voo, R. and Voo, R. V. D.: Insights into the mechanism for orogen-related
1438 carbonate remagnetization from growth of authigenic Fe-oxide: A scanning electron microscopy
1439 and rock magnetic study of Devonian carbonates from northern Spain, *J. Geophys. Res.-Solid
1440 Earth*, 107(B4), 14–14, 2002.
- 1441 Weil, A. B., Gutiérrez-Alonso, G. and Conan, J.: New time constraints on lithospheric-scale
1442 oroclinal bending of the Ibero-Armorican Arc: a palaeomagnetic study of earliest Permian rocks
1443 from Iberia, *J. Geol. Soc.*, 167(1), 127–143, doi:10.1144/0016-76492009-002, 2010.
- 1444 Weil, A.B., Yonkee, A., and Sussman, A.: Reconstructing the kinematics of thrust sheet rotation:
1445 a paleomagnetic study of Triassic redbeds from the Wyoming Salient, U.S.A., *GSA Bulletin*,
1446 122, 1/2 2-23, 2010.
- 1447 Weil, A. B. B., Gutiérrez-Alonso, G., Johnston, S. T. and Pastor-Galán, D.: Kinematic
1448 constraints on buckling a lithospheric-scale orocline along the northern margin of Gondwana: A
1449 geologic synthesis, *Tectonophysics*, 582, 25–49, doi:10.1016/j.tecto.2012.10.006, 2013.
- 1450 Woodcock, N. H., Soper, N. J. and Strachan, R. A.: A Rheic cause for the Acadian deformation
1451 in Europe, *J. Geol. Soc.*, 164, 1023–1036, 2007.
- 1452 Yenes, M., Alvarez, F. and Gutierrez-Alonso, G.: Granite emplacement in orogenic
1453 compressional conditions: the La Alberca-Bejar granitic area (Spanish Central System, Variscan
1454 Iberian Belt), *J. Struct. Geol.*, 21(10), 1419–1440, 1999.
- 1455 Yonkee, A., Weil, A.B., and Sussman, A.: Reconstructing the kinematic evolution of curved
1456 mountain belts: internal strain patterns in the Wyoming Salient, Sevier thrust belt, U.S.A., *GSA
1457 Bulletin*, 122, 1/2, 24-50, 2010.
- 1458 Yonkee, A. and Weil, A. B.: Quantifying vertical axis rotation in curved orogens: Correlating
1459 multiple data sets with a refined weighted least squares strike test, *Tectonics*, 29(3), 31 PP.-31
1460 PP., doi:10.1029/2008TC002312, 2010b.
- 1461 Zegers, T. E., Dekkers, M. J. and Bailly, S.: Late Carboniferous to Permian remagnetization of
1462 Devonian limestones in the Ardennes: Role of temperature, fluids, and deformation, *J. Geophys.
1463 Res. Solid Earth*, 108(B7), n/a-n/a, doi:10.1029/2002JB002213, 2003.

1464 Captions

- 1465 Fig. 1 Simplified paleogeographic map of the Variscan-Alleghanian orogeny prior to the
1466 Jurassic break-up of Pangea, with the major orogenic curves labeled. Note, this map represents

1467 Iberian outcrops without taking account of the post Jurassic Alpine deformation (see text for
1468 details e.g. Gong et al., 2008; Pastor-Galán et al., 2018). Slightly darker colors in the Variscan
1469 belt indicate present-day European and African outcrops (modified after Martínez-Catalán et al.,
1470 2009; Weil et al., 2013).

1471 Fig. 2 A) Present-day configuration of the tectonostratigraphic zones of the Iberian
1472 Variscides and its major structures. White areas represent Mesozoic and Cenozoic cover. B)
1473 Three competing geometric proposals for the Central Iberian curve. 1) A disharmonic curvature,
1474 up to 160° at the outer arc but much less pronounced at the inner arc (Aerden, 2004); 2) A
1475 harmonic, but more open curvature as suggested by Martínez Catalán (2012); 3) an isoclinal
1476 curvature model (Shaw et al., 2012). C) Distribution of the E1 (in migmatitic domes) and C3 to
1477 post-C3 granitoids in the NW of Iberia (modified from Pastor-Galán et al., 2016)

1478 Fig. 3 A) Stratigraphic synthesis of the Gondwanan platform series in NW Iberia.
1479 Cantabrian Zone columns are after Aramburu et al., 2002; Bastida, 2004; Murphy et al 2008;
1480 Pastor-Galán et al., 2013a; 2013b. Iberian Range follows Gozalo et al., 2008; Mergl and
1481 Zamora, 2012 and Calvín and Casas, 2014. West Asturian Leonese Zone stratigraphy is after
1482 Pérez-Estaún, 1990; Marcos, 2004; Martínez-Catalán et al., 2004a; Gutiérrez-Marco et al.,
1483 2019. Central Iberian Zone follows Díez-Balda, 1986; Valladares et al., 2000; Díez Montes,
1484 2007; Martínez-Catalán et al., 2004b; del Moral and Sarmiento, 2008; García-Arias et al., 2018.
1485 B) Ordovician paleocurrents orientations, modified from Shaw et al., 2012. C) Schematic basin
1486 architecture inferred from the stratigraphic compilation.

1487 Fig. 4 A) The kinematic evolution of the Cantabrian Orocline in its core, the Cantabrian
1488 Zone, inferred from total least squares (TLS) orocline tests (Pastor-Galán et al. 2017). B) Shows
1489 three orocline (strike) tests used to constrain the kinematics of the Cantabrian Orocline. The
1490 Ordovician paleocurrents, which predate any orogenic movement, recorded the complete
1491 vertical-axis rotation history and yields a slope (m) of ~1. The Moscovian paleomagnetic data
1492 (from Weil et al., 2013; Pmag.), which postdates the main orogenic phases (C1, C2 and E1) and
1493 is coeval with C3, shows a slope of ~1. The Gzhelian joint sets (from Pastor-Galán et al., 2011)
1494 orocline test shows a slope of ~0.5, which indicates that half of the orocline was already formed
1495 at ~304 Ma.

1496 Fig. 5 A) Aeromagnetic map of Spain (Ardizzone et al., 1989, for Spain and the World
1497 Digital Magnetic Anomaly Map (WDMAM project) and Portugal (modified from Martínez Catalán,
1498 2012 and Martínez Catalán et al., 2015), showing the possible trace of the Central Iberian
1499 curve. B) Bouguer anomalies of the Iberian Peninsula, modified from Ayala et al., 2016. Gravity
1500 anomalies do not reflect the geometry of the Cantabrian Orocline nor the Central Iberian curve

but are related to the Cenozoic Alpine lithospheric structure.

Fig. 6 Paleomagnetic studies related to the Cantabrian Orocline and the Central Iberian curve: (1) Synthesis of paleomagnetism in the core of the Cantabrian Orocline (see Weil et al., 2013); (2) Permian (eP) components synthesized in Weil et al. (2010); (3) Ordovician volcanics and limestones (Laquiana) in the boundary between the West Asturian-Leonese and Central Iberian Zones (Fernández-Lozano et al., 2016); (4) Devonian sedimentary sequences and Permian subvolcanics in the Iberian ranges (Pastor-Galán et al., 2018); (5) Permian dykes and sills (Calvín et al., 2014); (6) Anatectic granites (E1) and mantle derived granitoids (C3) from Tormes Dome and Central System (Pastor-Galán et al., 2016); (7) Cambrian limestones from Tamames (N) and los Navalucillos (S) (Pastor-Galán et al., 2015a); (8) Ediacaran-Early Cambrian sedimentary rocks in the southern sector of the Central Iberian Zone (Pastor-Galán et al., 2017b); (9) Almadén volcanics from the Central Iberian Zone (Perroud et al., 1991; Parés and van der Voo, 1992; Leite Mendes et al. in press) and Volcanic rocks from southern Ossa Morena and the South Portuguese Zone (Leite Mendes et al. in press)

Fig. 7 Magnetization components with a positive reversal test in the extensional anatectic granites of Tormes (A) and Martinamor Domes (B). This component is interpreted as primary with a magnetization age of >318 Ma (Pastor-Galán et al., 2016). C) Distribution of directions and VGPs and statistical parameters from both domes combined.

Fig. 8 Cartoon depicting the different vertical axis rotation events that occurred in the Cantabrian Zone and the Iberian Range, modified from Pastor-Galán et al. (2018). (A) Original quasilinear Variscan Orogenic belt, B) Formation of the Cantabrian Orocline around the Carboniferous–Permian boundary after a ~70° counterclockwise rotation in the Southern branch of the Cantabrian Zone and the Iberian Range. This rotation matches the rotation for the Cantabrian Orocline, see the fit of the Iberian Range Component #2 in the orocline test for the Cantabrian Zone (below). C) Post Permian (Cenozoic) rotation of ~22° clockwise (CW) likely produced by differential shortening during the Alpine orogeny (Izquierdo-Llavall et al., 2018). Below, the global magnetic polarity time scale for the Pennsylvanian and Cisuralian (following Ogg et al., 2016). TLS = Total Least Squares. Note that once the 22° CW rotation in the Iberian Range is corrected, components #2, #1, and P fit as expected with the APWP for the southern limb of the orocline (Pastor-Galán et al., 2016).

Fig. 9 Compilation of the directional distributions and average declinations with parachute of confidence (Δ Declination) in sites around the Central Iberian curve (see Fig. 6). The results show general CCW rotations in contrast to the expected CW if the Central Iberia curve formed by vertical-axis rotations (see text). Results are compared with the expected

declinations if those sites were part of the Cantabrian Orocline following the methodology described in Pastor-Galán et al., 2017b.


Fig. 10 Orocline test of the Cantabrian Orocline (Weil et al., 2013) compared with the magnetizations found in the adjacent Laurussian segments of the orogen: Ireland (Pastor-Galán et al., 2015b) and the South Portuguese Zone (Leite Mendes et al., in press)

Fig. 11 Structural analysis of mullions in the Central Iberian Zone (after Pastor-Galán et al., 2019b) A) Photograph (with card scale, 10cm) of a bedding plane surface showing the mullions and photograph analysis. B) Interpretation of the outcrop with fold axis traces depicting the deformation phase responsible for each structure. C) Result of retro-deformation of mullions in the Mogadouro road section modified from Pastor-Galán et al. (2019b). Unfolding the effects of D3 on D2 mullions. Unfolding the effects of C3 and E1 in C1 mullions.

Fig. 12 **Pioneering hypothesis** for the Central Iberian curve. Note that none of them fulfill the most recent geometric and kinematic criteria. A) Simplified ribbon continent model after Johnston et al. (2013) and Shaw and Johnston (2016). B) Dextral mega-shear model from Martínez-Catalán (2011). C) Kinematic model with indentation and left-lateral shearing after Simancas et al. (2013)

Fig. 13 Preliminary kinematic proposal for the Iberian Variscides. A) Pre-collisional stage after the opening of the Galicia Tras-os-Montes restricted seaway (e.g. Pin et al., 2002; Gutiérrez-Alonso et al., 2008a; Arenas et al., 2016). The irregular shape of the margin and the younging westwards deformation front (e.g. Daley et al., 1997) resulted in tectonic escape towards the still open Rheic Ocean (e.g. Braid et al., 2011; Murphy et al., 2016). B) After closure of the Rheic ocean, C1 and C2 structures formed. The Galicia Tras-os-Montes was emplaced orogen parallel (e.g. Martínez-Catalán et al., 1990; Dias da Silva et al., in press), preserving the shape of the seaway, i.e. a primary arc. C) The gravitational collapse of the orogen produced widespread anatexis and folding interference in the hinterland and the emplacement of the foreland fold-and-thrust belt. D) At Pennsylvanian times a change in the far-field stress buckled the Variscan belt around a vertical axis (see Gutiérrez-Alonso et al., 2008; Weil et al., 2013; Pastor-Galán et al., 2015a for details), creating new interference patterns and a lithospheric scale response (see Gutiérrez-Alonso et al., 2004, 2011a; Pastor-Galán et al., 2012a). E) When the orocline became too tight to keep rotating, new cross-cutting brittle structures (C4) formed and minor extensional collapse (E2) occurred (e.g. Fernández-Lozano et al., 2019; Dias da Silva et al., in press).

1567 Fig. 1 Simplified paleogeographic map of the Variscan-Alleghanian orogeny prior to the
1568 Jurassic break-up of Pangea, with the major orogenic curves labeled. Note, this map represents
1569 Iberian outcrops without taking account of the post Jurassic Alpine deformation (see text for
1570 details e.g. Gong et al., 2008; Pastor-Galán et al., 2018). Slightly darker colors in the Variscan
1571 belt indicate present-day European and African outcrops (modified after Martínez-Catalán et al.,
1572 2009; Weil et al., 2013).



1573 Fig. 2 A) Present-day configuration of the tectonostratigraphic zones of the Iberian
 1574 Variscides and its major structures. White areas represent Mesozoic and Cenozoic cover. B)
 1575 Three competing geometric proposals for the Central Iberian curve. 1) A disharmonic curvature,
 1576 up to 160° at the outer arc but much less pronounced at the inner arc (Aerden, 2004); 2) A
 1577 harmonic, but more open curvature as suggested by Martínez Catalán (2012); 3) an isoclinal
 1578 curvature model (Shaw et al., 2012). C) Distribution of the E1 (in migmatitic domes) and C3 to
 1579 post-C3 granitoids in the NW of Iberia (modified from Pastor-Galán et al., 2016)







Fig. 3 A) Stratigraphic synthesis of the Gondwanan platform series in NW Iberia. Cantabrian Zone columns are after Aramburu et al., 2002; Bastida, 2004; Murphy et al 2008; Pastor-Galán et al., 2013a; 2013b. Iberian Range follows Gozalo et al., 2008; Mergl and Zamora, 2012 and Calvín and Casas, 2014. West Asturian Leonese Zone stratigraphy is after Pérez-Estaún, 1990; Marcos, 2004; Martínez-Catalán et al., 2004a; Gutiérrez-Marco et al., 2019. Central Iberian Zone follows Díez-Balda, 1986; Valladares et al., 2000; Díez Montes, 2007; Martínez-Catalán et al., 2004b; del Moral and Sarmiento, 2008; García-Arias et al., 2018. B) Ordovician paleocurrents orientations, modified from Shaw et al., 2012. C) Schematic basin architecture inferred from the stratigraphic compilation.




1589 Fig. 4 A) The kinematic evolution of the Cantabrian Orocline in its core, the Cantabrian
1590 Zone, inferred from total least squares (TLS) orocline tests (Pastor-Galán et al. 2017). B) Shows
1591 three orocline (strike) tests used to constrain the kinematics of the Cantabrian Orocline. The
1592 Ordovician paleocurrents, which predate any orogenic movement, recorded the complete
1593 vertical-axis rotation history and yields a slope (m) of ~ 1 . The Moscovian paleomagnetic data
1594 (from Weil et al., 2013; Pmag.), which postdates the main orogenic phases (C1, C2 and E1) and
1595 is coeval with C3, shows a slope of ~ 1 . The Gzhelian joint sets (from Pastor-Galán et al., 2011)
1596 orocline test shows a slope of ~ 0.5 , which indicates that half of the orocline was already formed
1597 at ~ 304 Ma.




1598 Fig. 5 A) Aeromagnetic map of Spain (Ardizzone et al., 1989, for Spain and the World
1599 Digital Magnetic Anomaly Map (WDMAM project) and Portugal (modified from Martínez Catalán,
1600 2012 and Martínez Catalán et al., 2015), showing the possible trace of the Central Iberian
1601 curve. B) Bouguer anomalies of the Iberian Peninsula, modified from Ayala et al., 2016. Gravity
1602 anomalies do not reflect the geometry of the Cantabrian Orocline nor the Central Iberian curve
1603 but are related to the Cenozoic Alpine lithospheric structure.




1604 Fig. 6 Paleomagnetic studies related to the Cantabrian Orocline and the Central Iberian
 1605 curve: (1) Synthesis of paleomagnetism in the core of the Cantabrian Orocline (see Weil et al.,
 1606 2013); (2) Permian (eP) components synthesized in Weil et al. (2010); (3) Ordovician volcanics
 1607 and limestones (Laquiana) in the boundary between the West Asturian-Leonese and Central
 1608 Iberian Zones (Fernández-Lozano et al., 2016); (4) Devonian sedimentary sequences and
 1609 Permian subvolcanics in the Iberian ranges (Pastor-Galán et al., 2018); (5) Permian dykes and
 1610 sills (Calvín et al, 2014); (5) Anatectic granites (E1) and mantle derived granitoids (C3) from
 1611 Tormes Dome and Central System (Pastor-Galán et al., 2016); (6) Cambrian limestones from
 1612 Tamames (N) and los Navalucillos (S) (Pastor-Galán et al., 2015a); (7) Ediacaran-Early
 1613 Cambrian sedimentary rocks in the southern sector of the Central Iberian Zone (Pastor-Galán et
 1614 al., 2017b); (8) Almadén volcanics from the Central Iberian Zone (Perroud et al., 1991; Parés
 1615 and van der Voo, 1992; Leite Mendes et al. in press) and Volcanic rocks from southern Ossa
 1616 Morena and the South Portuguese Zone (Leite Mendes et al, in press)




1617 Fig. 7 Magnetization components with a positive reversal test in the extensional anatectic
 1618 granites of Tormes (A) and Martinamor Domes (B). This component is interpreted as primary
 1619 with a magnetization age of >318 Ma (Pastor-Galán et al., 2016). C) Distribution of directions
 1620 and VGPs and statistical parameters from both domes combined.




1621 Fig. 8 Cartoon depicting the different vertical axis rotation events that occurred in the
 1622 Cantabrian Zone and the Iberian Range, modified from Pastor-Galán et al. (2018). (A) Original
 1623 quasilinear Variscan Orogenic belt, B) Formation of the Cantabrian Orocline around the
 1624 Carboniferous–Permian boundary after a $\sim 70^\circ$ counterclockwise rotation in the Southern branch
 1625 of the Cantabrian Zone and the Iberian Range. This rotation matches the rotation for the
 1626 Cantabrian Orocline, see the fit of the Iberian Range Component #2 in the orocline test for the
 1627 Cantabrian Zone (below). C) Post Permian (Cenozoic) rotation of $\sim 22^\circ$ clockwise (CW) likely
 1628 produced by differential shortening during the Alpine orogeny (Izquierdo-Llavall et al., 2018).
 1629 Below, the global magnetic polarity time scale for the Pennsylvanian and Cisuralian (following
 1630 Ogg et al., 2016). TLS = Total Least Squares. Note that once the 22° CW rotation in the Iberian
 1631 Range is corrected, components #2, #1, and P fit as expected with the APWP for the southern
 1632 limb of the orocline (Pastor-Galán et al., 2016)



1633 Fig. 9 Compilation of the directional distributions and average declinations with parachute of
 1634 confidence (Δ Declination) in sites around the Central Iberian curve (see Fig. 6). The results
 1635 show general CCW rotations in contrast to the expected CW if the Central Iberia curve formed
 1636 by vertical-axis rotations (see text). Results are compared with the expected declinations if
 1637 those sites were part of the Cantabrian Orocline following the methodology described in Pastor-
 1638 Galán et al., 2017b




1639 Fig. 10 Orocline test of the Cantabrian Orocline (Weil et al., 2013) compared with the
1640 magnetizations found in the adjacent Laurussian segments of the orogen: Ireland (Pastor-Galán
1641 et al., 2015b) and the South Portuguese Zone (Leite Mendes et al., in press)





1642 Fig. 11 Structural analysis of mullions in the Central Iberian Zone (after Pastor-Galán et
1643 al., 2019b) A) Photograph (with card scale, 10cm) of a bedding plane surface showing the
1644 mullions and photograph analysis. B) Interpretation of the outcrop with fold axis traces depicting
1645 the deformation phase responsible for each structure. C) Result of retro-deformation of mullions
1646 in the Mogadouro road section modified from Pastor-Galán et al. (2019b). Unfolding the effects
1647 of D3 on D2 mullions. Unfolding the effects of C3 and E1 in C1 mullions.



1648 Fig. 12 Pionering hypothesis for the Central Iberian curve. Note that none of them fulfill
 1649 the most recent geometric and kinematic criteria. A) Simplified ribbon continent model after
 1650 Johnston et al. (2013) and Shaw and Johnston (2016). B) Dextral mega-shear model from
 1651 Martínez-Catalán (2011). C) Kinematic model with indentation and left-lateral shearing after
 1652 Simancas et al. (2013)






Fig. 13 Preliminary kinematic proposal for the Iberian Variscides. A) Pre-collisional stage after the opening of the Galicia Tras-os-Montes restricted seaway (e.g. Pin et al., 2002; Gutiérrez-Alonso et al., 2008a; Arenas et al., 2016). The irregular shape of the margin and the younging westwards deformation front (e.g. Daley et al., 1997) resulted in tectonic escape towards the still open Rheic Ocean (e.g. Braid et al., 2011; Murphy et al., 2016). B) After closure of the Rheic ocean, C1 and C2 structures formed. The Galicia Tras-os-Montes was emplaced orogen parallel (e.g. Martínez-Catalán et al., 1990; Dias da Silva et al., in press), preserving the shape of the seaway, i.e. a primary arc. C) The gravitational collapse of the orogen produced widespread anatexis and folding interference in the hinterland and the emplacement of the foreland fold-and-thrust belt. D) At Pennsylvanian times a change in the far-field stress buckled the Variscan belt around a vertical axis (see Gutiérrez-Alonso et al., 2008; Weil et al., 2013; Pastor-Galán et al., 2015a for details), creating new interference patterns and a lithospheric scale response (see Gutiérrez-Alonso et al., 2004, 2011a; Pastor-Galán et al., 2012a). E) When the orocline became too tight to keep rotating, new cross-cutting brittle structures (C4) formed and minor extensional collapse (E2) occurred (e.g. Fernández-Lozano et al., 2019; Dias da Silva et al., in press).


A) Pre C1 (~380 Ma)




B) C1-C2 (~360-335 Ma)




C) E1 (~330-317 Ma)



D) C3 (~315-300 Ma)



B) C4-E2 (~300-280 Ma)



- Cantabrian Zone
- West Asturian Leonese
- Central Iberian
- Ossa-Morena
- South Portuguese-Zone
- Galicia Tras-Os-Montes and other Oceanic domains
- E1/E2 gneiss domes & granites
- C3-C4 mantle derived granitoids
- C3 Dextral shear zones
- C3 Sinistral shear zones
- C1/C2 Sinistral shear zones
- Folds trend
- E1/C3 folds Central Iberian
- C1/C2 folds Central Iberian
- ... Zone boundary
- ↗ Transport direction