

The enigmatic curvature of Central Iberia and its puzzling kinematics

Daniel Pastor-Galán^{1,2,3}, Gabriel Gutiérrez-Alonso^{4,5}, Arlo B. Weil⁶

¹Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

²Department of Earth Science, Tohoku University

³Center for Northeast Asian Studies, Tohoku University, 41 Kawauchi, Aoba-ku, Sendai, 980-8576, Japan. pastor.galan.daniel.a8@tohoku.ac.jp

⁴Dept. of Geology. Faculty of Sciences. University of Salamanca. Plaza de la Merced s/n. 38007, Salamanca (Spain). gabi@usal.es

⁵Geology and Geography Department, Tomsk State University, Lenin Street, 36, Tomsk 634050 Russia

⁶Department of Geology, Bryn Mawr College, PA, USA 19010

Abstract

The collision between Gondwana and Laurussia that formed the latest supercontinent, Pangea, occurred during Devonian to Early Permian times and resulted in a large-scale orogeny that today transects Europe, northwest Africa and eastern North America. This orogen is characterized by an 'S' shaped corrugated geometry in Iberia. The northern curve of the corrugation is the well-known and studied Cantabrian (or Ibero-Armorican) Orocline and is convex to the east and towards the hinterland. Largely ignored for decades, the geometry and kinematics of the southern curvature, known as the Central Iberian curve, are still ambiguous and hotly debated. Despite the paucity of data, the enigmatic Central Iberian curvature has inspired a variety of kinematic models that attempt to explain its formation with little consensus. This paper presents the advances and milestones in our understanding of the geometry and kinematics of the Central Iberian curve from the last decade, with particular attention to structural and paleomagnetic studies.

When combined, the currently available datasets suggest that the Central Iberian curve did not undergo regional differential vertical-axis rotations during or after the latest stages of the Variscan orogeny, and did not form as the consequence of a single process. Instead, its core is likely a primary curve (i.e. inherited from previous physiographic features of the Iberian crust) whereas the curvature in areas outside the core are dominated by folding interference during the Variscan orogeny or more recent Cenozoic (Alpine) tectonic events.

33 **Keywords**

34 **Central Iberia Curve, Variscan orogen, Iberia, Cantabrian Orocline, Curved orogens,**
35 **Pangea**

36

37

1 Introduction

Mountain belt systems are the most striking product of plate tectonics. In addition to their astonishing visual impact, marking the locations where ancient and modern plates collided, orogenic belts often preserve a variety of rocks that have the potential to illuminate the entirety of the systems pre- and syn-orogenic histories. One of the striking characteristics of most of Earth's orogens are their curvature in plan-view (e.g. van der Voo, 2004; Marshak, 2004; Rosenbaum, 2014). The degree of orogenic curvature may range from a few degrees of deflection in structural trend (e.g. Kopet Dag, Iran), to 180° of arc curvature (e.g. Kazakhstan arc and the Carpathians). The kinematics, structural and geodynamic implications of these systems are as varied as their geometries (Marshak, 2004; Weil and Sussman, 2004; Johnston et al., 2013). For example, some orogenic curvatures are hypothesized to be the consequence of physiographic features of the basement that pre-date orogen formation, such as irregular basin architectures or plate margin salients and recesses that formed from rift-to-drift processes (e.g. Jura mountains, Hindle et al., 2000), which then control the growth geometry of the ensuing orogen. These systems are known as primary arcs, and reflect pre-orogenic geometries and show no significant or systematic vertical-axis rotations along their structural length. On the other hand, oroclines, as classically defined by Carey in 1956, involve systematic differential vertical-axis rotations subsequent to initial orogenic shortening: different sectors of an orogen rotate with variable magnitudes or in opposite directions (e.g. Li et al., 2012). Rotations in oroclines may occur at a range of scales, from thrust emplacement at upper crustal levels (e.g. Izquierdo-Llavall et al., 2018), up to lithospheric-scale vertical-axis folding (e.g. Li et al., 2018). They can occur as single curves (e.g. Maffione et al., 2009), coupled curves (Johnston, 2001), or in trains of curves (Li and Rosenbaum, 2014). Oroclines can form during the main orogenic building event, known as progressive oroclines (Johnston et al., 2013; e.g., the Wyoming salient, Yonkee and Weil, 2010, and Weil et al., 2010), or during a subsequent tectonic pulse, so-called secondary oroclines (Weil and Sussman, 2004). Unraveling the kinematics and mechanisms of curvature formation in mountain belts is a critical step to understanding orogenesis in 4D and to evaluate their geodynamic consequences and paleogeographic implications.

The Variscan-Alleghanian orogeny resulted in the suturing of Gondwana and Laurussia during Devonian-Carboniferous times, and ultimately led to the formation of the supercontinent, Pangea. This long and sinuous orogen runs for >8000 km along strike and is ca. 1000 km wide, transecting across Europe, to northwest Africa and into eastern North America. The final stages

of Pangea amalgamation (e.g. Nance et al., 2010) modified the Western Europe sector of the belt into its characteristic sinuous shape. Today, this tectonic belt traces at least one, and perhaps four arcs from Poland to Brittany, and then across the Bay of Biscay (Cantabrian Sea) into Iberia, where the system is truncated by the younger Betic Alpine orogeny in southeast Iberia (Fig. 1; e.g. Weil et al., 2013). The southern truncation of the Variscan in Europe hinders a precise correlation with equivalent age outcrops in NW Africa.

Within the Iberian Peninsula, the orogen is characterized by two large-scale curves (Fig. 2): (1) to the north is the well-studied and nearly 180° secondary orocline, the Cantabrian (a.k.a. Ibero-Armorican) Orocline, which buckled a segment of the Variscan belt from ~315 to ~290 Ma (e.g. Weil et al., 2019 and references therein); and (2) to the south is a curve with disputed magnitude and kinematics that is usually referred to as the Central Iberian curve/orocline or Castillian bend (Martínez-Catalán et al., 2015). Though there remains tremendous uncertainty on the geometry and kinematics of the Central Iberian curve, multiple hypotheses exist as to its nature, and disagreements continue on its importance in the tectonic evolution of Europe during the waning stages of Paleozoic global supercontinent construction. The diversity of interpretations on the Central Iberian curve range from a nonexistent structure (Dias et al., 2016), to being one of the most important pieces in our understanding of the late Carboniferous and Permian geodynamics of the Iberian Variscan system (e.g. Martínez-Catalán et al., 2011; 2014).

This paper reviews the most recent advances on the geometry and kinematics of the Central Iberian curve, synthesizing what we know and what we don't, and ending with a discussion of the main unsolved issues. We hope that this paper fosters novel studies that will lead to a better understanding of when and which mechanisms acted in the aftermath of the Variscan-Alleghanian orogeny.

2 The long and winding orogen

The Variscan (Europe-NW Africa)-Alleghanian (North America) orogeny is a continental-scale tectonic system (1000 km wide and >8000 km long) that sutured Gondwana and Laurussia together, forming the supercontinent Pangea (e.g. Domeier and Torsvik, 2014; Edel et al., 2018; Pastor-Galán et al., 2019a). The fragments of this system are now dispersed over three continents, Europe, Africa and North America due to the Mesozoic break-up of Pangea (Buiter and Torsvik, 2014; Keppie, 2015). This orogen formed as a consequence of a long and protracted tectonic history that involved several events, from initial convergence (ca. 420 Ma; e.g. Franke et al., 2017), to the consumption of multiple putative oceanic tracts and/or basins

that existed between Gondwana and Laurussia (ca. 280 Ma; e.g. Kirsch et al., 2012). The Variscan-Alleghanian orogen itself represents the closing of at least one major ocean, the Rheic (e.g. Nance et al., 2010), whose axial ridge likely failed or subducted at ca. 395 Ma along its paleo-northern margin (e.g. Woodcock et al., 2007; Gutiérrez-Alonso et al., 2008a). Perhaps the orogeny involved other large oceans (Stampfli and Borel, 2002; Franke et al., 2017; 2019), but most surely involved several minor seaways and basins that existed between Gondwana, Laurussia, and several intervening micro-continents (e.g. Azor et al., 2008, Dallmeyer et al., 1997; Kroner and Romer, 2013; Díez-Fernández et al., 2016; Pérez-Cáceres et al., 2017). The final continent-continent collision began after closure of all oceans and intervening seaways. The commencement of this deformation was diachronistic and became progressively younger westwards (in present-day coordinates): with Devonian continent-continent collisions along the eastern boundary, progressing to earliest Permian ages in the westernmost sector (McWilliams et al., 2013; Chopin et al., 2014; López-Carmona et al., 2014; Franke et al., 2017).

The present-day geometry of the Variscan-Alleghanian systems has a contorted trace (Fig. 1). In Europe, from east to west, the trend starts with a prominent curve around the Bohemian massif (e.g. Tait et al., 1996), followed by a deflection in the Ardennes-Bravant (e.g. Zegers et al., 2003). In Brittany the outer curvature of the Cantabrian or Ibero-Armorican orocline begins (e.g. van der Voo et al., 1997), and wraps nearly 180° around and across the Bay of Biscay as it turns in NW Iberia. The Central Iberian curve marks the final concave to the west curve (in present-day coordinates) and is the focus of this paper (e.g. Aerden, 2004; Martínez Catalán, 2011; Shaw et al., 2012). The orogen continues in North America where, from north to south, it has salients and recesses that undulate back and forth from Atlantic Maritime Canada (e.g. O'Brien, 2012) down along the Pennsylvanian and into the Alabama curves (e.g. Thomas, 1977).

Interpretation on the origin of these curvatures varies widely. The curvatures in North America are argued to be the result of a preexisting irregular margin of Laurentia due to the break-up of the Rodinia supercontinent, which resulted in the formation of orogenic salients and recesses during subsequent Appalachian collision (e.g. Rankin, 1976; Thomas, 1977, 2004). In this case, vertical-axis rotations affected mainly the upper crustal levels during orogenesis (e.g. Marshak, 1988; Bayona et al., 2003; Hnat and van der Pluijm, 2011). In Europe, the Bohemian and Ardennes-Bravant massif curvatures have poor kinematic constraints. In the Bohemian Massif, some suggest secondary rotations formed an orocline (Tait et al., 1996), while others suggest little to no vertical-axis rotations and a primary arc (Chopin et al., 2012). The Ardennes-Bravant Massif recorded some vertical-axis-rotations (e.g. Molina-Garza and Zeijderveld, 1996),

but it is unclear if these are a response to progressive or secondary oroclinal bending, or whether rotations only affected the upper crust. The most outstanding example of Variscan-Alleghanian orogen curvature is exposed in the Iberian Massif, with the Cantabrian Orocline and the coupled curvature of Central Iberia.

2.1 Two of us: The Variscan orogen in Iberia

The western half of the Iberian Peninsula constitutes the Iberian massif, one of the largest exposures of the Variscan orogen and the only place that contains an almost continuous cross section of the orogen (Fig. 2; e.g. Lotze 1945, Julivert 1974, Pérez-Estaún et al., 1991; Ayarza et al., 1998; Simancas et al., 2003; Ribeiro et al. 2007, Martínez Catalán et al., 2014, 2019). The majority of the Iberian Massif contains Gondwanan affinity rocks (e.g. Murphy et al., 2008; Pastor-Galán et al., 2013a; Gutiérrez-Marco et al., 2017) and likely represents a proximal piece of the Gondwana margin until its final amalgamation with Pangea (e.g. Pastor-Galán et al., 2013b). Owing to the stratigraphic, structural and petrological styles, the Iberian Massif has been traditionally divided into six tectonostratigraphic zones (Fig. 2; Lotze, 1945; Julivert, 1971): (1) The Cantabrian Zone represents a Gondwanan thin-skinned foreland fold-and-thrust belt. It has overall low-grade internal deformation and metamorphism, and represents shortening that occurred during Mississippian times (e.g. Marcos and Pulgar, 1982; Pérez Estaún et al., 1988; Gutiérrez-Alonso 1996; Alonso et al., 2009; Pastor-Galán et al., 2009; 2013b). (2) The West-Asturian Leonese Zone represents a metamorphic fold-and-thrust belt with Barrovian metamorphism that collapsed coevally with thrust emplacement onto the Cantabrian Zone (e.g. Martínez-Catalán et al., 1992; Alcock et al., 2009; Martínez-Catalán et al., 2014). (3) The Central Iberian Zone represents the Gondwanan hinterland with Barrovian and Buchan metamorphism and is intruded by igneous rocks of various ages (e.g. Macaya et al., 1991; Díez Balda, 1995; Gutiérrez-Alonso et al., 2018). (4) The Ossa-Morena Zone represents the most distal zone of the Gondwana platform, and is characterized by a metamorphic fold-and-thrust belt with dominantly sinistral displacement (e.g. Robardet and Gutiérrez-Marco, 2004; Quesada, 2006). (5) The Galicia-Tras-os-Montes Zone represents a far-travelled allochthonous terrane that contains high-pressure units and relicts of oceanic-like crust (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). (6) The South Portuguese Zone represents a foreland fold-and-thrust belt with little internal deformation and metamorphism with Avalonian affinity and a strong sinistral component of shear (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2016; Oliveira et al., 2019). Geographically, the external zones of the Gondwana margin are nested to the north into the core of the Cantabrian Orocline, whereas the hinterland zones are to the west

and center of the massif (Fig. 2; e.g. Díaz Balda, 1995; Azor et al., 2019). The southwestern-most extent of Iberia contains a putative suture of the Rheic Ocean, as well as a piece of the Laurussian margin fold-and thrust belt, today preserved in the South Portuguese Zone (e.g. Pereira et al., 2012, 2017; Oliveira et al., 2019).

The Gondwanan autochthon stratigraphy (Cantabrian, West Asturian-Leonese, Central Iberian and Ossa Morena zones) consist of a Neoproterozoic arc and back-arc basin (e.g. Fernández-Suárez et al., 2014), which evolved to a rift-to-drift Cambrian to Early Ordovician sequence and then to an Ordovician to Late Devonian passive margin basin sequence (e.g. Sánchez-García et al., 2019; Gutiérrez-Marco et al., 2019; Gutiérrez-Alonso et al., submitted). Overall the system transitioned from a relatively isolated Early Cambrian continental basin, to a restricted marine basin, to development of an open marine platform that was locally punctuated by magmatism (e.g. Gutiérrez-Alonso et al., 2008b; Palero-Fernández, 2015). The Ossa Morena Zone represents the outermost platform, followed by an intermediate platform characterized by an asymmetric horst (Central Iberian Zone) and graben (West-Asturian Leonese Zone), which ends in the innermost shelf environment of the Cantabrian Zone (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The differences between the West Asturian-Leonese and Central Iberian Zone are mainly deeper vs. shallower sedimentary facies (respectively) and a local Lower Ordovician unconformity in the Central Iberian Zone (Toledanian, e.g. Álvaro et al., 2018), which places Lower Ordovician strata atop pre-Cambrian to Cambrian rocks (Fig. 3; e.g. Gutiérrez-Marco et al., 2019). The Central Iberian Zone is divided into two domains: (1) The Ollo de Sapo domain, which contains abundant Lower Ordovician magmatism of calc-alkaline affinity (e.g. Díez Montes, 2006; Gutiérrez-Marco et al., 2019); and (2) the 'Schistose–Greywacke domain' characterized by a predominance of outcrops of Neoproterozoic to Lower Cambrian sedimentary rocks (e.g. Gutiérrez-Marco et al., 2019 and references therein).

The Galicia Tras-os-Montes Zone (Farias et al., 1987) is a complex structural stack including a basal schistose unit (Parautochthon; Dias da Silva et al., 2020) structurally overlain by mafic rocks with an oceanic-like signature and other allochthonous rocks under high-pressure metamorphism (e.g. López-Carmona et al., 2014; Martínez-Catalán et al., 2019). The oceanic rocks of this zone are classically interpreted as a Rheic Ocean suture (e.g. Martínez Catalán et al., 2009). Recent interpretations support its origin as a minor oceanic basin or seaway within the realm of Gondwana (e.g. Pin et al., 2002; Arenas et al., 2016).

The South Portuguese Zone constitutes the Laurussian foreland fold-and-thrust belt in the Iberian Variscides (e.g. Pereira et al., 2012; Pérez-Cáceres et al., 2017). It contains three units: (1) the Pulo de Lobo, a low grade metamorphic accretionary prism with clastic

sedimentary rocks and basalts with MORB signature (e.g. Azor et al., 2019; Pérez-Cáceres et al., this volume) some times considered an independent zone; (2) The Iberian Pyrite Belt, which is a world class volcanogenic massive sulfide deposit formed between 390 and 330 Ma (e.g. Oliveira et al., 2019a; 2019b); and (3) the Baixo Alentejo Flysch, which is located to the southwest and is a syn-orogenic composite turbiditic sequence with ages from ~330 to ~310 Ma (Oliveira et al., 2019b). The boundary between the South Portuguese and Ossa Morena zones is a sinistral shear zone (the so-called Southern Iberian shear zone in the Beja-Acebuches oceanic-like unit, Crespo-Blanc and Orozco, 1988; Quesada and Dallmeyer., 1994; Pérez-Cáceres et al., 2016) that contains a strongly deformed amphibolitic belt with oceanic affinity (Munha et al., 1986; Munha, 1989; Quesada et al., 2019). This belt potentially represents dismembered relics of the Rheic Ocean and/or a subsidiary seaway that opened during a Variscan transtension event in SW Iberia (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019).

Finally, Paleozoic rocks occur sporadically within the Alpine Betic chain. Their lithological monotony, paucity of fossils, and the intensity of deformation and metamorphism during Alpine orogeny, make recognizing the original features of the different successions challenging (e.g. Martín-Algarra et al., 2019). Some faunal and detrital zircon studies suggest that the Paleozoic outcrops in the Betics may be similar to the most continentalward realms of the Gondwanan platform (i.e., the Cantabrian Zone; e.g. Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018). Following the latest plate reconstructions of the Mediterranean during Meso-Cenozoic times, the Paleozoic units of the Betic-Rif chain may have been located proximal to the present-day position of the Balearic Islands (van Hinsbergen et al., 2020).

The Variscan orogen in Iberia shows multiple deformation, metamorphic, and magmatic events (e.g. Martínez-Catalán et al., 2014; Azor et al., 2019; Fig. 2) that evolved diachronously from the suture towards the external zones (Dallmeyer et al., 1997): (1) An initial continent-continent collision began ca. 370-365 Ma, which produced high-pressure metamorphism (e.g. Lopez-Carmona et al. 2014). (2) Between 360 and 330 Ma a protracted shortening phase occurred, frequently divided into main phases C1 and C2, that were accompanied by Barrovian type metamorphism (e.g. Dias da Silva et al., 2020) and plutonism at ~340 Ma (e.g. Gutiérrez-Alonso et al., 2018). (3) An extensional collapse event, so-called E1, occurred at ~333-317 Ma, which formed core-complexes and granitic domes in the Central Iberian and West Asturian-Leonese zones (Fig. 2C; e.g. Alcock et al., 2009; Díez-Fernández and Pereira, 2016; López-Moro et al., 2018). This event is coeval and genetically linked to the formation of the foreland fold-and-thrust-belt of the Cantabrian Zone (e.g. Gutiérrez-Alonso, 1996). (4) A late

Carboniferous shortening event (C3) occurred ca. 315-290 Ma and is interpreted to have resulted in the formation of the Cantabrian Orocline and was accompanied by intrusion of mantle derived granitoids (Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a, 2011b; Pastor-Galán et al., 2012a). (5) A final early Permian extensional event (E2), mostly found in the Central Iberian Zone, resulted in the formation of core complexes and regional doming (Dias da Silva et al., 2020). (6) A final shortening event (C4), possibly coeval with E2, resulted in widespread brittle deformation (e.g. Azor et al., 2019; Fernández-Lozano et al., 2019).

In SW Iberia, the aforementioned Variscan deformation events are characterized by a dominant sinistral component, which contrasts with the general dextral component recognized in most other regions of the orogen (e.g. Martínez Catalán et al., 2011; Gutiérrez-Alonso et al., 2015). Early collisional structures (C1) formed NE-vergent recumbent folds in the southernmost Central Iberian Zone and SW-vergent folds and thrusts in the Ossa Morena and South Portuguese zones. This phase continued with a transtensional event that heterogeneously extended the continental lithosphere (e.g. Pérez-Cáceres et al. 2015). Coevally, an important extension-related magmatic event happened, perhaps assisted by a plume-type mantle (Simancas et al. 2006) or because of slab break-off (Pin et al. 2008). After this transtensional event, significant sinistral transpression occurred forming the extensive shear zones to the north and south of the Ossa Morena Zone (Fig. 2B), which accommodated the majority of transcurrent motion. However, sinistral displacements are observed all along the Ossa Morena and South Portuguese zones. Pérez-Cáceres et al. (2016) estimated over 1000 km of collisional convergence in SW Iberia, most of which corresponds with sinistral displacements parallel to terrane boundaries.

3 Synthesis on the Geometry and Kinematics of the Cantabrian Orocline

Understanding the geometry, kinematic evolution and mechanics of curved mountain systems is crucial to developing paleogeographic and tectonic reconstructions and understanding past geodynamics (e.g. Marshak, 2004; Van der Voo, 2004; Li et al., 2012; van Hinsbergen et al., 2020). Introduced by Carey (1955 p.257), an orocline (from Greek ορος, mountain, and κλivo, bend) is “...an orogenic system, which has been flexed in plan to a horse-shoe or elbow shape.” Although sometimes used in the literature as a geometric description of any orogenic curvature, herein orocline is strictly used as a the term for map-scale bends that underwent vertical-axis rotations (Weil and Sussman, 2004; Johnston et al., 2013; Pastor-Galán et al., 2017a). The kinematic classification of curved mountain belts (Weil and Susman, 2004;

Johnston et al, 2013) distinguishes two end members: (1) Primary orogenic curves, which describe those systems in which curvature is a primary feature of the orogen and formed without significant or systematic vertical-axis rotations, and (2) Secondary oroclines, where orogenic curvature was acquired due to vertical-axis rotations subsequent to primary orogenic building. Those systems whose curvature is the product of vertical-axis rotation during the primary orogenic pulse and/or only a portion of the observed curvature is secondary are termed progressive oroclines.

The orocline test (or strike test), evaluates the relationship between changes in regional structural trend (relative to a reference trend for an orogen) and the orientations of a given geologic fabric element or magnetization (relative to a reference direction). In terms of evaluating developmental kinematics, the most relevant geologic marker is paleomagnetic declination, which can be used to quantitatively evaluate total and systematic rotations as a function of along-strike variability. Once acquired, data is plotted on Cartesian coordinate axes with the strike (S) of the orogen (relative to a reference) along the horizontal axis, and the fabric azimuth (F, relative to a reference) along the vertical axis. The test originally used a basic least-squares (OLS) regression (Schwartz and Van der Voo, 1983) to estimate the slope (coded m in formulas), ideally between 0 and 1, which then is interpreted with respect to vertical-axis kinematics. More recently, Yonkee and Weil (2010b) and Pastor-Galán et al. (2017a) introduced more robust statistics to estimate the correlations slope and its uncertainty, considering and propagating errors of the input data. Primary orogenic bends show no change of paleomagnetic declination orientations with varying structural trend, and therefore the slope is expected to be 0. In progressive oroclines, the declination variation records some fraction of the total observed orogenic strike variability, and thus the slope would range between 0 and 1, depending on the amount of primary curvature. Secondary oroclines are those in which the paleomagnetic vectors record 100% of the rotation, yielding slopes of 1, meaning that the orogenic system must have started as a roughly linear system that then underwent secondary vertical-axis rotations until its present-day curvature was acquired. The slope obtained with the orocline test can only be confidently interpreted when the chronology of fabric formation is well known.

The trend of the Variscan belt in Iberia follows a sinuous “S” shape that is especially prominent in the northwest region of the Iberian Peninsula, and then becomes more subtle due to the predominance of younger cover sequences in the central and eastern regions of the peninsula (Fig. 1 and 2). This dramatic geometry has stimulated a century long scientific debate as to its origin (e.g. Suess, 1892; Staub, 1926; Martínez Catalán et al., 2015). To the north and convex to the west is the Cantabrian Orocline, and to the center-south and convex to the east is

the Central Iberian curve. The overall trend of the Cantabrian Orocline starts in Brittany (France) and southern England and then curves through the Bay of Biscay and then south into central north Iberia (Fig. 1, 2 and 4). The Cantabrian Orocline (also known as Ibero-Armorican Orocline/Arc, Asturian Arc or Cantabrian-Asturias Arc) is arguably the first curved orogen that was scientifically described, recognized by the change in structural trend of mapped thrusts and fold axes (Schultz, 1858, Barrois, 1882, Suess, 1892). The Cantabrian Orocline traces an arc with a curvature close to 180° within the central Cantabrian Zone (the Gondwanan foreland in Iberia, fig. 2), and opens to approximately 150° as one moves to the outer arc reaches (Fig. 1). At the crustal-scale, the Cantabrian Orocline represents a first-order vertical-axis buckle fold in plan-view that refolds pre-existing Variscan structures (e.g. Julivert and Marcos, 1973; Weil et al., 2001). The inner arc of the orocline, or the Cantabrian Zone is characterized by tectonic transport towards the core of the orocline, i.e., the orocline has a contractional core, where low finite strain values and locally developed cleavage occur (Pérez-Estaún et al., 1988; Gutiérrez-Alonso, 1996; Pastor-Galán et al., 2009). Within the inner core a variety of structures record non-coaxial strain, which produced complex interference folds and rotated thrust sheets (e.g. Julivert and Marcos, 1973; Julivert and Arboleya, 1984; Pérez-Estaún et al., 1988; Aller and Gallastegui, 1995; Weil, 2006, 2013; Pastor-Galán et al., 2012b; Shaw et al., 2015; 2016a; Del Greco et al., 2016). In contrast, the outer arc shows a vertical-axis fold with a ca. 150° interlimb angle that was accommodated by significant shearing, both dextral and, in lesser amounts, sinistral that was penecontemporaneous to vertical-axis rotation (Gutiérrez-Alonso et al., 2015). Weil et al. (2013, 2019) extensively review the geometry of the Cantabrian Orocline.

All kinematic data studied so far support a model in which the Cantabrian Orocline formed due to secondary vertical-axis rotation in a period of time younger than 315 Ma and older than 290 Ma. Overall, the southern limb of the orocline rotated counterclockwise (CCW) and the northern limb clockwise (CW; Fig. 4). Orocline formation happened subsequent to the main shortening phases of the orogen (C1 and C2) and late-stage orogenic collapse (E1), and therefore, it is an ideal example of a secondary orocline in the strictest sense. Development of the Cantabrian Orocline requires the existence of a roughly linear orogenic belt during early Variscan closure of the Rheic Ocean (with a roughly N-S orientation in present-day coordinates), which was subsequently bent in plan-view into an orocline during late stages of Pangea amalgamation. Such interpretation is grounded in extensive paleomagnetic (e.g. Hirt et al., 1992; Parés et al. 1994; Stewart, 1995; van de Voo et al., 1997; Weil, 2006; Weil et al., 2000; 2001; 2010), along with important contributions from structural (e.g. Gutiérrez-Alonso 1992; Kollmeier et al., 2000; Merino-Tomé et al., 2009; Pastor-Galán et al., 2011; 2014; Shaw et

al., 2015) and geochronological studies (e.g., Tohver et al., 2008; Gutiérrez-Alonso et al., 2015). Weil et al. (2013) provided a comprehensive review on the kinematic constraints, updated in 2017a by Pastor-Galán et al., and in 2019 by Weil et al.

4 The intriguing geometry of the Central Iberian curve

The more southern Central Iberian curve has a similar magnitude, but opposite curvature compared to the Cantabrian Orocline (Fig. 1 and 2B). This structure has been referred to as the Central Iberian curve, arc, bend or orocline. In this paper we use 'Central Iberian curve'. The other aforementioned terms involve still unknown parameters or are misleading: e.g., orocline implies kinematics (Weil and Sussman, 2004); bend refers to a mechanism of formation (e.g. Fossen, 2016); and arc could be ambiguous, since the term is commonly used for volcanic chains. The Central Iberian curve was first described by Staub (1926) and was termed the Castilian bend. Continental drift pioneers paid some attention to Staub's description (e.g. Holmes, 1929; Du Toit, 1937), but the curved structure remained largely ignored for multiple decades (e.g. Martínez Catalán et al., 2015). The hypothesis of a large-scale curvature in Central Iberia made a comeback at the beginning of the 21st century with a study of Variscan porphyroblast kinematics across Iberia by Aerden (2004). Since then, several attempts to unveil its geometry and kinematics have given contrasting results.

The elusive nature of the Central Iberian curve resides in the poor exposure of its putative hinge (Fig. 2). The hinge of the Cantabrian orocline crops out extensively and the changes in thrust and fold axes trend are observable at high-resolution from aerial photographs and are readily mapped using outcrop-scale observations. In contrast, the alleged hinge of the Central Iberian curve is largely covered by Mesozoic and Cenozoic basins (Fig. 2). The curvature is most recognizable at the boundary between the Galicia-Tras os Montes and Central Iberian zones (Fig. 2A; Aerden, 2004; Martínez Catalán, 2012). The thrust fault that bounds those areas traces close to a 180° of curvature and marks the emplacement of the most distal units. Before the revival of Staub's curved geometry along the entire Central Iberian Zone, there were several attempts to explain the curved shape of the Galicia Tras-os-Montes Zone. Some consider the Galicia Tras-os-Montes Zone a block that escaped during an early Variscan (C1) non-cylindrical collision, forming an extrusion wedge towards areas that underwent lesser amount of shortening (Martínez-Catalán, 1990; Dias da Silva, 2015; 2020); or alternatively a klippe of a larger allochthonous thrust sheet, or the product of an interference pattern between C2, E1 and C3 structures (e.g. Ries and Shackleton, 1971; Martínez Catalán et al., 2002; Rubio Pascual et al., 2013; Díez-Fernández et al., 2015).

In addition to the Galicia Tras-os-Montes Zone, the other areas that show a certain degree of curvature are to the E and SE of the Central Iberian Zone. There, an approximately 20° change in strike of the Iberian ranges (NE Iberia, Fig. 2A) is observed, which represents the only known outcrop of the hinge of the Central Iberian curve's outer arc. The rest of the curvature has been implied with indirect observations leading to three competing geometric proposals for the Central Iberian curve (Fig. 2B). The main arguments used to constrain the geometry of the Central Iberian curve are: (1) the geometry of Galicia Tras-os-Montes folds and the orientation of observed garnet inclusion trails (Aerden, 2004; Fig. 2B-1); (2) the alignment of aeromagnetic anomalies and fold trends in the Iberian ranges and the E-SE Central Iberian Zone (Martínez-Catalán, 2012; Fig. 2A and 2B-2) and; (3) the regional distribution of paleocurrents recorded in Ordovician quartzites (Shaw et al., 2012; Fig. 2B-3 and 3). All proposed geometries share two features: (1) The curvature runs parallel to the Central Iberian Zone, and is located in the center-west of Iberia, and (2) all place the Galicia Tras-os-Montes Zone in the core of the curve with the curved axial traces cross-cutting the Morais Complex, which is a set of mafic and ultramafic rocks that is roughly circular in shape (Fig. 2B; Dias da Silva et al., 2020).

Aerden (2004) compared the orientation of inclusions in metamorphic porphyroblasts across the Variscan allochthonous terranes of the NW Iberian Massif, and found that inclusion trails maintain a constant north–south orientation. Comparing such results with the trend of the Variscan fold axes in the Central Iberian Zone (Fig. 2A) and a novel interpretation of the aeromagnetic anomalies of the Iberian Peninsula (Fig. 5A), Aerden suggested a geometry in which the Central Iberian curve was more prominent in the outer arc than in the inner arc (Fig. 2B-1). In Aerden's view the geometry of the Galicia Tras-os Montes Zone does not represent a large-scale curvature, but rather the original shape of the nappe, perhaps re-tightened during C3 deformation. In contrast, the Iberian Ranges and the SE Central Iberian Zone represent the more curved sector (Fig. 2B-1). In the model of Aerden (2004), the Ossa Morena and South Portuguese zones are not part of the Central Iberian curvature.

Martínez-Catalán (2012) reinterpreted Aerden's analysis of aeromagnetic map data (Fig. 5A) and their interpreted structural trends of C1-C2 fold axes from Central Iberian Zone structures (Fig. 2A). In Martínez-Catalán's model, the Central Iberian curvature is a symmetric arcuate shape in which orogen trend changes equally in the inner and outer arc, and is comparable in size to the Cantabrian Orocline, but with opposite curvature and less shortening. This geometric model also excludes the Ossa Morena and the South Portuguese Zones as elements involved in the formation of the curvature (Fig. 2B-2).

Finally, Shaw et al. (2012) studied the orientation of paleocurrents in Ordovician Armorican Quartzite (e.g. Aramburu, 2002), which is one of the most prominent rocks exposed in Iberia (Fig. 3). These authors found that paleocurrents fanned outward with respect to the Cantabrian Orocline curve and are approximately perpendicular to the structural trend throughout the peninsula (Fig. 3). Shaw et al. (2012) assumed that the direction and sense of paleocurrents were originally parallel throughout all zones, and concluded that the Central Iberia curve is part of a 'S' shape isoclinal structure with a similar magnitude of curvature to the Cantabrian Orocline (Fig. 2B-3). It is unclear from the Shaw et al. (2012) model what the involvement of the Ossa Morena and South Portuguese zones in the overall curve is (if any), nor the prospective location of the external zones of the orogen (Cantabrian Zone) with respect to the overall curvature.

5 Move over once, move over twice: Kinematic constraints

Late Variscan kinematic data (315-290 Ma; C3, E2, C4 phases) in the Central Iberian curve were scarce prior to revival of Staub's Central Iberian curve (e.g. Vergés, 1983; Julivert et al., 1983; Parés and van der Voo, 1992). More recently, a wealth of studies have been published on the kinematics of forming the Central Iberian curve (Fig. 2B), which are reviewed below.

5.1 Structural Geology and Geochronology

Curved orogens that result from differential vertical-axis rotations develop remarkable structures within their hinges where compressive and extensive radial structures often develop in combination with tangential shear structures (e.g. Li et al., 2012; Eichelberger and McQuarrie, 2015). With the re-emergence of the Central Iberian curve debate, several studies have re-evaluated the well-documented structures from the Central Iberian Zone to constrain the origin and kinematics of curvature. The majority of studies focused on the hinge zone of the curve in the area surrounding Galicia Tras-os-Montes (e.g. Dias da Silva et al., 2014; Jacques et al., 2018a), but some do explore areas in the outer-arc (e.g. Palero-Fernández et al., 2015; Gutiérrez-Alonso et al., 2015). The following section synthesizes the findings of new field, structural, and geochronological analyses from around the hinge of the Central Iberian curve and its surrounding regions. The reviewed studies identify several deformation events that are linked to regional Variscan deformation phases (Fig. 2A).

1. An early generation of upright to overturned cylindrical folds with an associated axial planar cleavage (C1). The C1 fold axes plunge variably from horizontal to nearly vertical

(e.g. Jacques et al., 2018a, 2018b). The original trend of the fold axes were parallel to the orogen (e.g. Pastor-Galán et al., 2019b), however interference with younger deformation events has created complicated geometries (e.g. Díez Fernández et al., 2013; Palero-Fernández et al., 2015). The emplacement of the allochthonous units of the Galicia Tras-os-Montes Zone (commonly referred as C2) is closely associated with development of C1 folds, but is restricted to shear zones located along the boundary between the latter and the Central Iberian Zone. This phase includes orogen-parallel emplacement of the allochthonous Galicia Tras-Os Montes units and its associated thrusts (Fig. 2A). The non-coaxial nature of the emplacement of this allochthonous nappe produced folding interference and local vertical-axis rotations (Dias da Silva et al., 2020). Prograde Barrovian metamorphism (known as M1) reached its pressure peak at the end of C2 (Rubio Pascual et al., 2013).

2. After C1 and C2, the resulting thickened crust gravitationally collapsed (Macaya et al., 1991; Escuder Viruete et al., 1994; Díaz-Balda et al., 1995; Díez-Montes, 2010). This gravitational collapse (phase E1) formed gneiss-dome core complexes between 330 and 317 Ma (e.g. Díez Fernández and Pereira, 2016) especially in the core of the Central Iberian curve (Fig. 2C; e.g. Martínez-Catalán, 2012). This phase formed large subhorizontal extensional detachments that exhumed to depths of the middle crust (e.g. Rubio-Pascual et al., 2013; Dias da Silva et al., 2020). General decompression produced a Buchan-type metamorphic event (M2; e.g. Rubio-Pascual et al., 2016, Solís-Alulima et al., 2019) and widespread anatexis melting (e.g. López-Moro et al., 2018; Pereira et al., 2018). The E1 phase developed a fold system with sub-horizontal axes and a penetrative subhorizontal cleavage (e.g. Dias da Silva et al., 2020). Mapped folding geometries indicate the deflection of C1 folds into overturned positions within the E1 deformation zones (e.g. Díez Fernández et al., 2013; Díez Fernández and Pereira, 2016; Pastor-Galán et al., 2019b). In addition to large-scale extensional deformation and Buchan metamorphism, E1 developed a regional dome-and-basin pattern, resulting in portions of the allochthonous terranes tectonically transported into basins (e.g. Dias da Silva et al., 2020).
3. The structures developed during C1-C2 compression and E1 extension, are re-folded by a younger shortening phase (C3; syn-Cantabrian Orocline). C3 formed upright open folds and conjugate sub-vertical shear zones (e.g. Gutiérrez-Alonso et al., 2015; Díez Fernández and Pereira, 2017; Dias da Silva et al., 2020). C3 was coeval with regional retrograde metamorphism (M3) and with intrusion of mantle derived granitoids (Fig. 2C;

e.g. Gutiérrez-Alonso et al., 2011a), surrounded by contact metamorphic aureoles (e.g. Yenes et al., 1999). The age of the C3 event ranges from 315 and 290 (e.g. Jacques et al., 2018a), and is concomitant with the formation of the Cantabrian Orocline (e.g. Pastor-Galán et al., 2015a). Ductile deformation, including folding with axial planar cleavage (e.g. Dias da Silva et al., 2014; Pastor-Galán et al., 2019b) and the development of conjugate shear zones, occurred at the early stages of C3 (315-305 Ma; Gutiérrez-Alonso et al., 2015; Díez-Fernández and Pereira, 2017; Jacques et al., 2018b) and was followed by brittle deformation that formed cross-joint sets and vein swarms with Sn-W mineralizations (Jacques et al., 2018a; 2018b). The conjugated shear zones, some of them with hundreds of kilometers of displacement, had activity during the period 315-305 Ma based on direct Ar-Ar dating of the shear zones (Gutiérrez-Alonso et al., 2015) and cross-cutting relationships with precisely dated igneous rocks (Díez-Fernández and Pereira, 2017). Note that these shear zones show a younger age with respect to the sinistral shear zones that bound the Ossa Morena and South Portuguese zones (340-330 Ma; e.g. Dallmeyer et al., 1993). New studies in the Central Iberian Zone have determined that several folds, previously interpreted as C1 (e.g. the Tamames-Marofa-Sátão synform) are in fact C3 structures, possibly nucleated within existing C1-C2 structures (e.g. Dias da Silva et al., 2017; Jacques et al., 2018b). The remarkable continuity along the Central Iberian Zone of these folds (Fig. 2A), previously interpreted as C1 (e.g. Díez-Balda et al., 1990; Abalos et al., 2002; Dias and Ribeiro, 1994; Dias et al., 2016), suggests the ubiquity and importance of this deformation phase.

4. The N-S shortening (in present day coordinates) of C3 deformation continued through the Early Permian under brittle conditions (so-called C4 event) (e.g. Dias da Silva et al., 2020) and overlapped with the formation of E2 extensional faults (Fig. 2A; Dias and Ribeiro 1991; Dias et al. 2003; Rubio Pascual et al., 2013; Arango et al., 2013; Fernández-Lozano et al. 2019; Dias da Silva et al., 2020). C4 N-S compression produced a series of NNE-SSW and NNW-SSE brittle faults (Gil Toja et al. 1985; Dias and Ribeiro 1991; Dias et al. 2003; Fernández-Lozano et al., 2019) and associated sub-vertical and sub-horizontal widespread kink-bands (e.g. Aller et al., 2020; Dias da Silva et al., 2020) that are today exposed in Northern Iberia. E2 developed core complex-like structures with extensional shear zones that further telescoped M2 metamorphic isograds between the anatectic cores of gneiss domes and the associated hanging wall units. This event favored sub-horizontal folding, and crenulation cleavage development in the footwall together with kink-band generation in the upper low-grade structural

levels.

5.2 Paleomagnetism

Paleomagnetism investigates the Earth's ancient magnetic field as it is recorded in rocks. Among other features, rocks can record the orientation of the magnetic field at the time of magnetization (e.g. Tauxe, 2010). The magnetic vector can be geometrically defined by two components: inclination, which is a function of the paleolatitude (being 90° at the poles and 0° at the equator) at the time of magnetization acquisition; and declination, which is a measure of the horizontal angular difference between the recorded magnetic direction and true north, thereby allowing for the quantification of any vertical-axis rotations if a reference paleomagnetic pole is known for the region of interest at the time of magnetization acquisition. Paleomagnetism is the best tool to quantify vertical-axis rotations in orogens due to the independence of the magnetic field from an orogen's deformation and evolution (e.g. Butler, 1998).

Despite its uniqueness to study paleolatitudes and vertical-axis rotations, paleomagnetism is not flawless. Paleomagnetic data can yield spurious rotations when the local and regional structures are not properly defined and their geometries and kinematic histories not adequately corrected for (e.g. Pueyo et al., 2016). In addition, the age of magnetization acquisition is not necessarily equivalent to the age of the sampled rock. Remagnetizations are ubiquitous, especially in orogens (Weil and van der Voo, 2002; Pueyo et al., 2007; Huang et al., 2017). In remagnetized rocks, the primary magnetization is replaced or overprinted due to a set of geologic processes acting alone or in concert - usually represented by a combination of thermal or chemical reactions (e.g., Jackson, 1990). Nevertheless, remagnetizations can be useful for interpreting deformation history if the relative timing of the overprint can be established and a well-constrained reference direction for that age is known (e.g. Weil et al., 2001; Izquierdo-Llavall et al. 2015; Calvin et al., 2017).

In addition to knowing the structural geology and the timing of magnetization of the studied rocks, understanding and quantifying local and regional vertical-axis rotations requires a paleomagnetic reference pole for comparison. Permian and Mesozoic paleomagnetic studies in Iberia indicate that Iberia was a relatively stable plate from at least Guadalupian times (ca. 270 Ma) to the opening of the Bay of Biscay in the Cretaceous (e.g. Gong et al., 2008; Vissers et al., 2016). Weil et al. (2010) calculated the an Early Permian pole for stable Iberia, which will be used herein as a reference for any vertical-axis rotation analysis (hereafter, eP pole or eP component). Weil et al.'s Virtual Geomagnetic Pole (VGP) values are $Plat = 43.9$; $Plong = 203.3$ and $\alpha_{95} = 5.4$ and when transform into paleomagnetic components has a $\sim 0^\circ$ inclination

(equatorial) and declinations that range from 150° to 160° (from NW to SW respectively) depending on where in Iberia you are referencing. In Fig. 6 (red arrows), a compilation of declinations that form part of this composite pole and other eP components found in recent studies are presented.

For the Central Iberian curve, the voluminous paleomagnetic database from the Cantabrian Orocline can be used to partially constrain its kinematics (e.g. Weil et al., 2013). The orocline test for the Cantabrian Orocline (Fig. 4) quantifies the degree of differential vertical-axis rotation of variously striking Variscan tectonic belts in northern Iberia. If the Central Iberian curve is a product of vertical-axis rotation, paleomagnetic declinations would bend around the Central Iberian curve opposite to that of the Cantabrian Orocline. With a well constrained orocline test, as in the Cantabrian Orocline (Fig. 4), one can use the paleomagnetic strike-test correlation slope to establish expected declinations for any along-strike portion of the orogen (Pastor-Galán et al., 2017b).

Before the resurgence of the Central Iberian curve, the only available pre-Permian paleomagnetic studies to the south and west of the Cantabrian Zone in the Iberian Massif were from the Beja Gabbroic Massif, Portugal (Perroud et al., 1985) and the Almadén syncline volcanics (Perroud et al., 1991; Pares & Van der Voo, 1992). The Beja area study presented varied inclinations and declinations in the gabbros, and complex overprints elsewhere. Perroud et al. (1985) did not consider any structural correction for the results, as they assumed the gabbro was undeformed. Recently, Dias da Silva et al. (2018) showed that the area underwent intense deformation during the Carboniferous. Therefore, interpretation of this dataset is complicated without knowing the proper structural correction needed to restore the magnetization to its palinspastic orientation.

Several articles with new paleomagnetic studies around the Central Iberian curve have been published since 2015 (Fig. 6). In general, these studies have reported a pervasive late Carboniferous (320 to 300 Ma) (re-)magnetization in sedimentary and igneous rocks (e.g. Pastor-Galán et al., 2015a; 2017b; Fernández-Lozano et al. 2016), which is largely penecontemporaneous to the intrusion of E1 extensional granites (López-Moro et al., 2018) and C3 syn-orocline mantle derived granitoids (Fig. 2C; e.g. Gutiérrez-Alonso et al., 2011a). The following section describes the reported magnetizations from oldest to youngest.

Pastor-Galán et al. (2016) sampled for paleomagnetic analyses both E1 extensional granites (Fig. 2C; ~320 Ma; e.g. López-Moro et al., 2018) from the Tormes and Martinamor domes, and C3 mantle derived granitoids from the Central System (Fig. 2C; 305-295 Ma; e.g. Gutiérrez-Alonso et al., 2011a). Both sets of plutons are located around the Galicia Tras-os-

Montes hinge of the Central Iberian curve (Fig. 6-5). The authors found an original component in E1 granites supported by a positive reversal test in both domes (Fig. 7). The magnetization has an inclination (Inc.) = 15° (paleolatitude (λ) = -7.6°) and declination (Dec.) = 81° (Fig. 7), which imply a northward movement of 700 km and a $\sim 70^\circ$ CCW rotation with respect to the C3 granites that showed an eP component (Dec. ~ 150 , Inc. ~ 0). Considering the positive reversal test in E1 granites and the significant difference in inclinations with respect to C3 granitoids (eP component), a magnetization age of older than 318 Ma was proposed (pre-Kiaman superchron, 317 Ma - 267 Ma, e.g. Langereis et al., 2010), which was interpreted as a primary magnetization. The 70° CCW Pennsylvanian rotation recorded in rocks from the Central Iberian curve hinge zone is in agreement with the expected rotation of the southern limb of the Cantabrian Orocline (Fig. 4; Weil et al., 2013).

At the putative outer arc of the Central Iberian curve in the Iberian Ranges (Fig. 2), paleomagnetic and structural studies of Devonian and Permian rocks (Pastor-Galán et al., 2018) revealed that the eP component from Permian rocks had rotated $\sim 22^\circ$ CW during the Cenozoic (Fig. 8; cf. Pastor-Galán et al., 2018). The Permian and Mesozoic rocks from the Iberian Ranges show a consistent $\sim 22^\circ$ CW rotation with respect to the Apparent Polar Wander Path for Iberia (e.g. Pastor-Galán et al. 2018). This rotation likely happened during the Alpine orogeny, in which the northern area of the Iberian Range underwent more shortening than the southern part, resulting in a regional CW vertical-axis rotation (Izquierdo-Llavall et al., 2019). After restoring the Cenozoic rotation, the Devonian rocks show a positive reversal and fold-test with inclinations that are steeper than expected from the eP component (Dec. = 85.3° , Inc. = 12.7° , λ = -6.4). This component is statistically indistinguishable from that of the E1 granites and the southern branch of the Cantabrian Zone, showing the same 70° CCW rotation from the time they were remagnetized (estimated ~ 318 Ma) to the timing of the eP component (Fig. 8; Pastor-Galán et al., 2018). Once Cenozoic rotation is corrected for, the structural and paleomagnetic trends of the Iberian ranges become parallel to those in the southern limb of the Cantabrian Orocline, ruling out a Variscan or older origin for the outer Central Iberian curve (Fig. 8).

The remaining paleomagnetic works published on Central and SW Iberia rocks all reveal a ubiquitous late Carboniferous to Early Permian remagnetization during the Kiaman superchron (Fernández-Lozano et al., 2016; Pastor-Galán et al., 2015a; 2016; 2017b; Leite Mendes, in press). The authors of these papers calculated the expected declination for each site as if they were part of the Cantabrian Orocline (Fig. 9A). All localities where magnetizations pre-date the formation of the Cantabrian Orocline show the same expected rotations as the southern limb of the Cantabrian Orocline, regardless of their position within the Central Iberian

curve (to the hinge: Tormes and Martinamor domes, Iberian ranges; to the southern limb: Almadén syncline and South Portuguese Zone). Other locations, especially limestones from the Central Iberian Zone, have declinations and inclinations in between the primary 318 Ma component of the E1 granites and the post-orocline eP component (Fig. 9B). Pastor-Galán et al. (2015a; 2016) interpreted these results as being caused by a remagnetization that was acquired during formation of the Cantabrian Orocline and therefore record intermediate steps between the component of the E1 granites and eP. Those authors suggest that the large amount of syn-orocline mantle derived granitoids that intruded the Central Iberian Zone (C3 granitoids) triggered the hinterland remagnetization.

Finally, two previous studies identified an earlier magnetization in the Almadén syncline region of the SE Central Iberian Zone (Perroud et al., 1991; Pares & Van der Voo, 1992). However, Leite Mendes et al. (in press) argue that these studies are likely misinterpreted. Perroud et al. (1991), applied a complicated structural correction restoring a putative plunge of the regional structural axis to all sites, including those where the syncline axis does not plunge. Leite Mendes et al. (in press) re-sampled the syncline where its axis is sub-horizontal and obtained a negative fold test, implying that the magnetization is not primary as previously interpreted. Their results, however, are similar in orientation to those components published from previous studies prior to any structural correction (Perroud et al., 1991 and Parés and van der Voo, 1992).

Two additional studies sampled Laurussian margin sequences that are today adjacent to the Cantabrian Orocline region (Fig. 10). To the north, the SW part of Ireland preserves a Late Paleozoic basin filled with Devonian red sandstone and Carboniferous limestone and siltstone, which was sampled by Pastor-Galán et al. (2015a). To the south are the aforementioned results from the South Portuguese Zone (Leite Mendes et al., in press). Both areas are interpreted to have previously been part of the Laurussian continent, and therefore on the opposite side of the Rheic Ocean suture at the time of Variscan collision (Fig. 10; e.g. Pastor-Galán et al., 2015b). In contrast, the rest of Iberia was part of, or proximal to, Gondwana (e.g. Franke et al., 2017). These Paleomagnetic results from the Laurussian margin suggest that the rotations involved in the formation of the Cantabrian Orocline occurred along both sides of the Rheic suture along its northern and southern limbs (Fig. 10A and B). Pastor-Galán et al. (2015b) hypothesized a so-called Greater Cantabrian Orocline that would have bent the entire Appalachian/Variscan orogen around a vertical-axis, affecting at least the continental margins of both Gondwana and Laurussia.

5.3 The implications of not being a secondary orocline

The most relevant new data regarding the kinematics of the Central Iberian curve is the paleomagnetic study from the Iberian Ranges (Calvín et al., 2014; Pastor-Galán et al., 2018). These results confirm that the present-day variation in trend of the tectonostratigraphic units, generally attributed to Variscan tectonics (e.g. Weil et al., 2013; Shaw et al., 2012; 2014), is likely a product of Cenozoic Alpine orogeny. Izquierdo-Llavall et al. (2019) confirmed that the interpreted Alpine rotations correspond well with the amount of shortening reconstructed in Meso-Cenozoic basins. The best preserved and most continuous outcrop in the Central Iberian's outer arc is not a Variscan structure, casting doubt that the Central Iberian curve is related to Variscan kinematics. These results are also a reminder that the regional effects of Alpine deformation are often underestimated, especially close to the major Iberian Alpine fronts: the Pyrenees, Iberian Ranges, and the Betics.

Overall, new paleomagnetic data from the Central Iberian curve and nearby areas reveal pervasive late Carboniferous remagnetizations and regional vertical-axis rotations of the same sense and magnitude to those expected for the southern arm of the Cantabrian Orocline. The new paleomagnetic data indicate that a post ~320 Ma formation for the Central Iberia curve due to vertical-axis rotations is not supported (Pastor-Galán et al., 2016). The distribution in space and time of paleomagnetic results makes it unlikely that the formation of the Central Iberian curve is a product of Variscan gravitational collapse (E1, ~330-317 Ma) or concomitant to the Cantabrian Orocline (C3). So far, no pre-E2 paleomagnetic component has been found, and consequently, paleomagnetic data cannot reject an early orogenic origin for the inner arc of the Central Iberian curve (C1-C2, older than 330 Ma).

From a structural geology point of view, the Central Iberian curve does not display the classic geometries and structural interference patterns found in other established oroclines (i.e., those systems that involve differential vertical-axis rotations, e.g. Li et al., 2012; van der Boon et al., 2018; Meijers et al., 2017; Rezaeian et al., 2020). The geometry and structural behavior of oroclines should resemble, at the crustal-scale, a regional vertical-axis fold preserved in plan-view, either formed by buckling (e.g. Johnston et al., 2001) or bending (e.g. Cifelli et al., 2008) mechanisms. In oroclines, pre-existing structures tend to follow fold trends around the curvature (e.g. Rosenbaum, 2014; Li et al., 2018). In addition, orocline cores tend to preserve radial structures and shortening patterns in the inner arc and orocline parallel shear zones and extensional structures in their outer arc (e.g. Ries and Shackleton, 1976; Eichelberger and McQuarrie, 2015), similar to what is observed in multilayer folds (e.g. Fossen, 2016).

The structural geometry of the Central Iberian curve lacks such patterns.

Paleomagnetism from the Iberian Ranges indicate that the Cantabrian and West Asturian Leonese zones do not follow the Central Iberian curve, instead they continue their WNW-ESE trend into the Mediterranean in what it is now the Betic chain (Rodríguez-Cañero et al., 2018; Jabaloy-Sánchez et al., 2018; van Hinsbergen et al., 2020). Structural trends in the Ossa Morena and the South Portuguese Zone do not show any change in along-strike structural trend that supports large-scale CW rotations (e.g. Pérez-Cáceres et al., 2015; Quesada et al., 2019), whereas existing paleomagnetic data from those zones (Leite Mendes et al., in press) support a model of CCW rotation associated with the broader southern arm of the Cantabrian Orocline. In the Central Iberian and Galicia Tras-os-Montes zones, the trend of curvature is irregular (see C1 fold patterns in Fig. 2A) and nowhere are the expected inner and outer arc-related structures preserved (e.g. Dias da Silva et al. in press).

The curved shape of C1 fold axes in the Central Iberian Zone is better explained by fold interference patterns than vertical-axis rotations (e.g. Pastor-Galán et al., 2019b). Moreover, the curved shape of the Galicia Tras-os-Montes allochthonous nappe, which was emplaced orogen-parallel, shows no evidence of vertical-axis rotation related structures (Fig. 2A; Dias da Silva et al., 2020). Other authors describe the changes in trend around the Central Iberian curve expressed by C1 folds (Fig. 2A) as the product of fold interference patterns (e.g. Gutiérrez-Alonso, 2009; Palero-Fernández et al., 2015; Jacques et al., 2018b; Dias da Silva et al., 2020). Pastor-Galán et al. (2019b) showed that curved C1 folds in the Central Iberian Zone around the Galicia Tras-os-Montes boundary (Fig. 2A) are coaxial with C3 folds after restoring the effects of C2 and E1 deformation phases. Both C1 and C3 structures formed under similar shortening directions. In the same area, Jacques et al. (2018b) found similar fold interference patterns, in addition they described kinematic incompatibility with the expected CW rotations that would have occurred if the Central Iberian curve were an orocline. In other areas of the Central Iberian Zone, the curved shape of C1 folds has been described as an interference between C1 structures and their reorientation caused by C3 shear zones (Fig. 2A; e.g. Palero-Fernández et al., 2015; Dias et al., 2016), or alternatively the interference between C1, C3 and the E2 structures (Fig. 2A; Gutiérrez-Alonso, 2009; Arango et al., 2013; Rubio Pascual et al. 2013).

Overall, new geometric and kinematic data favor the interpretation that the Central Iberian curve is not a structure formed by differential vertical-axis rotation as was the Cantabrian Orocline, but one formed as a consequence of several competing processes. It is clear from the current data that a combination of several deformation events caused the orientation of structures that today delineate the shape of the Central Iberia curve. These include: (1) the northern part of the outer-arc as the product of an Alpine rigid block rotation instead of Variscan

differential vertical-axis rotation (Pastor-Galán et al., 2018); (2) the curvature of the Galicia Tras-os-Montes allochthonous nappe reflects its original shape and could be defined as a primary curve (see Weil and Sussman, 2004), since it was emplaced orogen parallel and preserves no evidence of vertical-axis rotations (fig. 2A; Dias da Silva et al., 2020); (3) Structural analysis shows that fold interference patterns explain the geometry of the curved trends of Central Iberian Zone's C1 folds (Pastor-Galán et al., 2019b), whose kinematics are incompatible with the required CW rotations expected if the curve is an orocline (Jacques et al., 2018b).

6 Get Back: Ideas flowing out and endless questions

The pioneering works in the last decade that resurrected the idea of a Central Iberian curve, speculated that both the Cantabrian and Central Iberian zones buckled together as secondary oroclines (Fig. 11; Martínez-Catalán 2011; Shaw et al., 2012, 2014; Shaw and Johnston, 2016; Carreras and Druguet, 2014). Later, Martínez Catalan et al. (2014) and Díez Fernández and Pereira (2017) reformulated Martínez-Catalán's 2011 hypothesis and proposed that the Central Iberian curve formed as an orocline between 315 and 305 Ma, and assigning the Cantabrian Orocline a time frame between 305 and 295 Ma (Fig. 11). The proposed tectonic mechanisms to support these early kinematic models are varied: (1) buckling of a ribbon 'Armorican' continent (Fig. 11A; Shaw et al., 2014; 2016); (2) buckling of a completely formed Variscan orogen during a putative 'Pangea B' to 'Pangea A' transition in the late Carboniferous (Fig. 11B; Carreras and Druguet, 2014; Martínez-Catalán et al., 2011); (3) indentation of Laurussia into Gondwana during the early stages of collision (at present day SW Iberia, South Portuguese Zone), producing first the Central Iberian curve as a mega drag-fold during Carboniferous times and then slightly later the Cantabrian Orocline as a consequence of an indentation process (Fig. 11C; Simancas et al., 2013).

The reviewed data in sections 4 and 5 contradict the aforementioned hypotheses. Paleomagnetism and structural patterns (section 5; Fig. 6-11) disagree with the necessary CW rotations required to support a late Carboniferous orocline origin for the Central Iberian curve (Models in Fig. 11A and B). In addition, the sense and magnitude of the vertical-axis rotations observed in SW Iberia (Fig. 10) imply that the South Portuguese (Avalonian segment) and Ossa Morena zones moved together with the southern limb of the Cantabrian Orocline during the Pennsylvanian and Early Permian. This means that the South Portuguese Zone was already parallel to the general trend of the Variscan orogen prior to Cantabrian Orocline formation, implying the lack of a Laurussian rigid indenter into Gondwana (e.g. Simancas et al., 2013). This discrepancy leaves orogen-parallel terrane transport as a possible explanation to the kinematics

observed in the Ossa Morena and South-Portuguese zones (e.g. Pérez-Cáceres et al., 2016). At the same time, paleomagnetism from SW Iberia supports the hypothesis of a Greater Cantabrian Orocline that extended into both Gondwana and Laurussia in its northern and southern limbs (Fig. 10; Pastor-Galán et al., 2015b).

In spite of the kinematic constraints and structural patterns that do not support a vertical-axis origin for the Central Iberian curve in Late Carboniferous time, there are geometric constraints that remain challenging to account for. For example, the curved shape of the aeromagnetic and gravity anomalies of Iberia are real (Fig. 5). These striking patterns could be due to Variscan-Alpine structural interference, as suggested for the Iberian Ranges, but currently there is not enough data to rigorously test this hypothesis. In addition are the curved traces of C1 fold-axes, whose geometry and kinematics are reasonably well constrained around Galicia Tras-os-Montes (e.g. Dias da Silva et al., 2020), but in many other areas their strong curvature remain largely unstudied (Fig. 2) and therefore, to date we can only speculate on their origin.

Shaw et al. (2012) supported their hypothesis of a secondary orocline by assuming that paleocurrents were parallel through Iberia during Ordovician times. However, some of the observed deflections in the paleocurrents studied by Shaw et al. (2012; see Fig. 3) are also explained by Alpine vertical-axis rotations (the case of the Iberian ranges) and fold interference patterns (SE of the Central Iberian Zone). Others (Central and SW of the Central Iberian Zone) may be explained by a local response to basin architecture (Fig. 3), where paleo-flow directions would trend toward the deepest basin troughs. The Ordovician basin architecture of Iberia allows for opposite directed paleocurrents from both sides of such troughs (Fig. 3). However, the Early Paleozoic basin architecture in Iberia and their local deformation events require further research (Sánchez-García et al., 2019).

Finally, although kinematic evidence is still scarce for the earliest pre-Variscan and early Variscan events, we argue that pre-orogenic physiographic features, such as the opening of a marginal restricted ocean between Gondwana and its distal platform at 395 Ma (Fig. 12A; Pin et al., 2002; Gutiérrez-Alonso et al., 2008b; Arenas et al., 2016) explains the rounded shape of the Galicia Tras-os-Montes curve as a primary curve. During collision, the latter irregularity would cause the orogen-parallel emplacement of allochthonous nappes (Fig. 12B; Dias da Silva et al., 2020) and the sinistral movements of the Ossa Morena and South Portuguese zones in SW Iberia (Fig 13A, B, C; Quesada, 2019). During the late Carboniferous, possibly due to a plate reorganization during the final amalgamation of Pangea (Fig. 12D; e.g. Gutiérrez-Alonso et al., 2008a; Pastor-Galán et al., 2015a), the far-field stress-field likely changed, which caused the

entire orogen to buckle around a vertical axis (Gutiérrez-Alonso et al., 2004), including both the Gondwana and Laurussia margins (Fig. 12E; Pastor-Galán et al., 2015b).

Acknowledgements

DPG thanks the extraordinary hard work, patience and endurance of the Utrecht University students that embraced and enjoyed studying the kinematics of the Central Iberian curve: Thomas Groenewegen, Bart Ursem, Daniel Brower, Mark Diederer and Bruno Leite-Mendes. DPG acknowledges FRIS and CNEAS for the continuous financial support. GGA is supported by Spanish Ministry of Science, innovation and universities under the project IBERCRUST (PGC2018-096534-B-I00) and Russian Federation Government grant no. 14.Y26.31.0012. This paper is a contribution to the IGCP no. 648 “Supercontinent Cycles and Global Geodynamics”. 50 years ago four fabulous guys let it be and never got back.

References

- Ábalos, B., Carreras, J., Druguet, E., Escuder Viruete, J., Gómez Pugnaire, M. T., Lorenzo Álvarez, S., Quesada, C., Rodríguez Fernández, L. R. and Gil-Ibarguchi, J. I.: Variscan and pre-Variscan tectonics, *Geol. Spain*, 155–183, 2002.
- Aerden, D.: Correlating deformation in Variscan NW-Iberia using porphyroblasts; implications for the Ibero-Armorican Arc, *J. Struct. Geol.*, 26(1), 177–196, 2004.
- Alcock, J. E., Catalán, J. R. M., Arenas, R. and Montes, A. D.: Use of thermal modeling to assess the tectono-metamorphic history of the Lugo and Sanabria gneiss domes, Northwest Iberia, *Bull. Société Géologique Fr.*, 180(3), 179–197, doi:10.2113/gssgfbull.180.3.179, 2009.
- Aller, J., Bastida, F. and Bobillo-Ares, N. C.: On the development of kink-bands: A case study in the West Asturian-Leonese Zone (Variscan belt, NW Spain) / Sur le développement des kink-bands : un exemple dans le Zone Astur Occidentale-léonaise (chaîne varisque ibérique, nord-ouest de l'Espagne), *Bull. Société Géologique Fr.*, 191(1), doi:10.1051/bsgf/2020003, 2020.
- Aller, J. J. J. and Gallastegui, J. J.: Analysis of kilometric-scale superposed folding in the Central Coal Basin (Cantabrian zone, NW Spain), *J. Struct. Geol.*, 17(7), 961–969, doi:10.1016/0191-8141(94)00115-G, 1995.
- Alonso, J. L., Marcos, A. and Suárez, A.: Paleogeographic inversion resulting from large out of sequence breaching thrusts: The León Fault (Cantabrian zone, NW Iberia). A new picture of the external Variscan thrust belt in the Ibero-Armorican arc, *Geol. Acta*, 7(4), 451–473, doi:10.1344/105.000001449, 2009.
- Álvaro, J. J., Casas, J. M., Clausen, S. and Quesada, C.: Early Palaeozoic geodynamics in NW Gondwana, *J. Iber. Geol.*, 44(4), 551–565, doi:10.1007/s41513-018-0079-x, 2018.

- 810 Ardizzone, Juan, Julio Mezcua, and Isabel Socías. Mapa aeromagnético de España peninsular.
811 Instituto Geográfico Nacional, 1989.
- 812 Mergl, M. & Zamora, S. 2012, New and revised occurrences of... Bulletin of Geosciences, 87,
813 571-586., [online] Available from: <http://www.geology.cz/bulletin/contents/art1327> (Accessed 6
814 April 2020b), n.d.
- 815 Aramburu, C., Méndez-Bedia, I. and Arbizu, M.: The Lower Palaeozoic in the Cantabrian Zone
816 (Cantabrian Mountains, NW Spain)., edited by S. García-López and F. Bastida, pp. 35–49.,
817 2002.
- 818 Arango, C., Díez Fernández, R. and Arenas, R.: Large-scale flat-lying isoclinal folding in
819 extending lithosphere: Santa María de la Alameda dome (Central Iberian Massif, Spain),
820 Lithosphere, 5(5), 483–500, doi:10.1130/L270.1, 2013.
- 821 Arenas, R., Sánchez Martínez, S., Díez Fernández, R., Gerdes, A., Abati, J., Fernández-
822 Suárez, J., Andonaegui, P., González Cuadra, P., López Carmona, A., Albert, R., Fuenlabrada,
823 J. M. and Rubio Pascual, F. J.: Allochthonous terranes involved in the Variscan suture of NW
824 Iberia: A review of their origin and tectonothermal evolution, Earth-Sci. Rev., 161, 140–178,
825 doi:10.1016/j.earscirev.2016.08.010, 2016.
- 826 Ayala, C., Bohoyo, F., Maestro, A., Reguera, M. I., Torne, M., Rubio, F., Fernández, M. and
827 García-Lobón, J. L.: Updated Bouguer anomalies of the Iberian Peninsula: a new perspective to
828 interpret the regional geology, J. Maps, 12(5), 1089–1092,
829 doi:10.1080/17445647.2015.1126538, 2016.
- 830 Ayarza, P., Catalan, J. R. M., Gallart, J., Pulgar, J. A. and Danobeitia, J. J.: Estudio Sísmico de
831 la Corteza Iberica Norte 3.3: A seismic image of the Variscan crust in the hinterland of the NW
832 Iberian Massif, Tectonics, 17(2), 171–+, 1998.
- 833 Azor, A., Rubatto, D., Simancas, J. F., Lodeiro, F. G., Poyatos, D. M., Parra, L. M. M. and
834 Matas, J.: Rheic Ocean ophiolitic remnants in Southern Iberia questioned by SHRIMP U-Pb
835 zircon ages on the Beja-Acebuches amphibolites - art. no. TC5006, Tectonics, 27(5), C5006–
836 C5006, 2008.
- 837 Azor, A., Dias da Silva, Í., Gómez Barreiro, J., González-Clavijo, E., Martínez Catalán, J. R.,
838 Simancas, J. F., Martínez Poyatos, D., Pérez-Cáceres, I., González Lodeiro, F., Expósito, I.,
839 Casas, J. M., Clariana, P., García-Sansegundo, J. and Margalef, A.: Deformation and Structure,
840 in The Geology of Iberia: A Geodynamic Approach, edited by C. Quesada and J. T. Oliveira, pp.
841 307–348, Springer International Publishing, Cham., 2019.
- 842 Barrois, C. E.: Recherches sur le terrains anciens des Asturies et de la Galice, Six-Horemans.,
843 1882.
- 844 Bastida, F.: Zona Cantábrica, in Geología de España, edited by J. A. Vera, pp. 25–49, SGE-
845 IGME, Madrid., 2004.
- 846 Bayona, G., Thomas, W. A. and Van der Voo, R.: Kinematics of thrust sheets within transverse
847 zones: A structural and paleomagnetic investigation in the Appalachian thrust belt of Georgia
848 and Alabama, J. Struct. Geol., 25(8), 1193–1212, doi:10.1016/S0191-8141(02)00162-1, 2003.

849 van der Boon, A., van Hinsbergen, D. J. J. J., Rezaeian, M., Gürer, D., Honarmand, M.,
850 Pastor-Galán, D., Krijgsman, W. and Langereis, C. G. G.: Quantifying Arabia–Eurasia
851 convergence accommodated in the Greater Caucasus by paleomagnetic reconstruction, *Earth*
852 *Planet. Sci. Lett.*, 482, doi:10.1016/j.epsl.2017.11.025, 2018.

853 Buiter, S. J. H. and Torsvik, T. H.: A review of Wilson Cycle plate margins: A role for mantle
854 plumes in continental break-up along sutures?, *Gondwana Res.*, 26(2), 627–653,
855 doi:10.1016/j.gr.2014.02.007, 2014.

856 Butler, R.: Paleomagnetism: Magnetic domains to geologic terranes, *Electron. Ed.*,
857 (September), 319–319, doi:10.1006/icar.2001.6754, 1998.

858 Calvin, P., Casas, A. M., Villalaín, J. J., Tierz, P., Calvin, P., Casas, A. M., Villalaín, J. J., Tierz,
859 P., Calvin, P., Casas, A. M., Villalaín, J. J. and Tierz, P.: Reverse magnetic anomaly controlled
860 by Permian Igneous rocks in the Iberian Chain (N Spain), *Geol. Acta*, 12(3), 193–207,
861 doi:10.1344/GeologicaActa2014.12.3.2, 2014.

862 Calvin, P., Casas-Sainz, A. M. M., Villalaín, J. J. J. and Moussaid, B.: Diachronous folding and
863 cleavage in an intraplate setting (Central High Atlas, Morocco) determined through the study of
864 remagnetizations, *J. Struct. Geol.*, 97, 144–160, doi:10.1016/j.jsg.2017.02.009, 2017.

865 Calvin-Ballester, P. and Casas, A.: Folded Variscan thrusts in the Herrera Unit of the Iberian
866 Range (NE Spain), *Geol. Soc. Lond. Spec. Publ.*, 394(1), 39–52, doi:10.1144/SP394.3, 2014.

867 Carey, S. W.: The orocline concept in geotectonics-Part I, *Pap. Proc. R. Soc. Tasman.*, 89,
868 255–288, 1955.

869 Carreras, J. and Druguet, E.: Framing the tectonic regime of the NE Iberian Variscan segment,
870 *Geol. Soc. Lond. Spec. Publ.*, 405(1), 249–264, doi:10.1144/SP405.7, 2014.

871 Martínez-Catalán, J. R. M., Fernández-Suárez, J., Jenner, G. A., Belousova, E. and Montes, A.:
872 Provenance constraints from detrital zircon U–Pb ages in the NW Iberian Massif: implications
873 for Palaeozoic plate configuration and Variscan evolution, *J. Geol. Soc.*, 161(3), 463–476,
874 doi:10.1144/0016-764903-054, 2004.

875 Chopin, F., Schulmann, K., Skrzypek, E., Lehmann, J., Dujardin, J. R., Martelat, J. E., Lexa, O.,
876 Corsini, M., Edel, J. B., Štípská, P. and Pitra, P.: Crustal influx, indentation, ductile thinning and
877 gravity redistribution in a continental wedge: Building a Moldanubian mantled gneiss dome with
878 underthrust Saxothuringian material (European Variscan belt), *Tectonics*, 31(1), n/a-n/a,
879 doi:10.1029/2011TC002951, 2012.

880 Chopin, F., Corsini, M., Schulmann, K., El Houicha, M., Ghienne, J.-F. and Edel, J.-B.: Tectonic
881 evolution of the Rehamna metamorphic dome (Morocco) in the context of the Alleghanian-
882 Variscan orogeny, *Tectonics*, 33(6), 1154–1177, doi:10.1002/2014TC003539, 2014.

883 Cifelli, F., Mattei, M. and Della Seta, M.: Calabrian Arc oroclinal bending: The role of subduction,
884 *Tectonics*, 27(5), doi:10.1029/2008TC002272, 2008.

885 Dallmeyer, R. D., Fonseca, P. E., Quesada, C. and Ribeiro, A.: $^{40}\text{Ar}/^{39}\text{Ar}$ mineral age
886 constraints for the tectonothermal evolution of a Variscan suture in southwest Iberia,
887 *Tectonophysics*, 222(2), 177–194, doi:10.1016/0040-1951(93)90048-O, 1993.

888 Dallmeyer, R. D. D., Catalán, J. R. M., Arenas, R., Gil Ibarguchi, J. I. I., Gutiérrez-Alonso, G.,
889 Farias, P., Bastida, F. and Aller, J.: Diachronous Variscan tectonothermal activity in the NW
890 Iberian Massif: Evidence from ⁴⁰Ar/³⁹Ar dating of regional fabrics, *Tectonophysics*, 277(4),
891 307–337, doi:10.1016/S0040-1951(97)00035-8, 1997.

892 Dias da Silva, Í., Gómez-Barreiro, J., Martínez Catalán, J. R., Ayarza, P., Pohl, J. and Martínez,
893 E.: Structural and microstructural analysis of the Retortillo Syncline (Variscan belt, Central
894 Iberia). Implications for the Central Iberian Orocline, *Tectonophysics*, 717, 99–115,
895 doi:10.1016/j.tecto.2017.07.015, 2017.

896 Dias da Silva, Í., Pereira, M. F., Silva, J. B. and Gama, C.: Time-space distribution of silicic
897 plutonism in a gneiss dome of the Iberian Variscan Belt: The Évora Massif (Ossa-Morena Zone,
898 Portugal), *Tectonophysics*, 747–748, 298–317, doi:10.1016/j.tecto.2018.10.015, 2018.

899 Dias da Silva, Í., González Clavijo, E. and Díez-Montes, A.: The collapse of the Variscan belt: a
900 Variscan lateral extrusion thin-skinned structure in NW Iberia, *Int. Geol. Rev.*, 00(00), 1–37,
901 doi:10.1080/00206814.2020.1719544, 2020.

902 Dias da Silva, Í. F., Linnemann, U., Hofmann, M., González-Clavijo, E., Díez-Montes, A. and
903 Catalán, J. R. M.: Detrital zircon and tectonostratigraphy of the Parautochthon under the Morais
904 Complex (NE Portugal): implications for the Variscan accretionary history of the Iberian Massif,
905 *J. Geol. Soc.*, 172(1), 45–61, doi:10.1144/jgs2014-005, 2015.

906 Dias, R. and Ribeiro, A.: Finite strain analysis in a transpressive regime (Variscan autochthon,
907 northeast Portugal), *Tectonophysics*, 191(3), 389–397, doi:10.1016/0040-1951(91)90069-5,
908 1991.

909 Dias, R. and Ribeiro, A.: CONSTRICTION IN A TRANSPRESSIVE REGIME - AN EXAMPLE IN
910 THE IBERIAN BRANCH OF THE IBERO-ARMORICAN ARC, *J. Struct. Geol.*, 16(11), 1543–
911 1554, 1994.

912 Dias, R., Mateus, A., and Ribeiro, A.: Strain partitioning in transpressive shears zones in the
913 southern branch of the Variscan Ibero-Armorican arc: *Geodinamica Acta*, v. 16, no. 2–6, p. 119–
914 129. doi:10.1016/j.geoact.2003.04.001, 2003.

915 Dias, R., Ribeiro, A., Romão, J., Coke, C. and Moreira, N.: A review of the arcuate structures in
916 the Iberian Variscides; constraints and genetic models, *Tectonophysics*, 681, 170–194,
917 doi:10.1016/j.tecto.2016.04.011, 2016.

918 Díez Balda, M. A., Vegas, R. and González Lodeiro, F.: Central-Iberian zone structure, *Pre-
919 Mesoz. Geol. Iber.*, 172–188, 1990.

920 Díez Balda, M. A., Martínez Catalán, J. R. and Ayarza Arribas, P.: Syn-collisional extensional
921 collapse parallel to the orogenic trend in a domain of steep tectonics: the Salamanca
922 Detachment Zone (Central Iberian Zone, Spain), *J. Struct. Geol.*, 17(2), 163–182,
923 doi:10.1016/0191-8141(94)E0042-W, 1995.

924 Díez Fernández, R. and Arenas, R.: The Late Devonian Variscan suture of the Iberian Massif: A
925 correlation of high-pressure belts in NW and SW Iberia, *Tectonophysics*, 654, 96–100,
926 doi:10.1016/j.tecto.2015.05.001, 2015.

- 927 Díez Fernández, R. and Pereira, M. F.: Extensional orogenic collapse captured by strike-slip
928 tectonics: Constraints from structural geology and UPb geochronology of the Pinhel shear zone
929 (Variscan orogen, Iberian Massif), *Tectonophysics*, 691, 290–310,
930 doi:10.1016/j.tecto.2016.10.023, 2016.
- 931 Díez Fernández, R. and Pereira, M. F.: Strike-slip shear zones of the Iberian Massif: Are they
932 coeval?, *Lithosphere*, 9(5), 726–744, doi:10.1130/L648.1, 2017.
- 933 Díez Fernández, R., Gómez Barreiro, J., Martínez Catalán, J. R. and Ayarza, P.: Crustal
934 thickening and attenuation as revealed by regional fold interference patterns: Ciudad Rodrigo
935 basement area (Salamanca, Spain), *J. Struct. Geol.*, 46, 115–128,
936 doi:10.1016/j.jsrg.2012.09.017, 2013.
- 937 Díez Fernández, R., Arenas, R., Pereira, M. F., Sánchez-Martínez, S., Albert, R., Martín Parra,
938 L.-M., Rubio Pascual, F.-J. and Matas, J.: Tectonic evolution of Variscan Iberia: Gondwana–
939 Laurussia collision revisited, *Earth-Sci. Rev.*, 162, 269–292,
940 doi:10.1016/j.earscirev.2016.08.002, 2016.
- 941 Díez-Montes, A.: La Geología del Dominio “Ollo de Sapo” en las comarcas de Sanabria y
942 Terra do Bolo., 2006.
- 943 Díez-Montes, A., Martínez-Catalán, J. R. R. and Bellido-Mulas, F.: Role of the Ollo de Sapo
944 massive felsic volcanism of NW Iberia in the Early Ordovician dynamics of northern Gondwana,
945 *Gondwana Res.*, 17(2–3), 363–376, doi:10.1016/j.gr.2009.09.001, 2010.
- 946 Domeier, M. and Torsvik, T. H.: Plate tectonics in the late Paleozoic, *Geosci. Front.*, 5(3), 303–
947 350, doi:10.1016/j.gsf.2014.01.002, 2014.
- 948 Edel, J. B., Schulmann, K., Lexa, O. and Lardeaux, J. M.: Late Palaeozoic palaeomagnetic and
949 tectonic constraints for amalgamation of Pangea supercontinent in the European Variscan belt,
950 *Earth-Sci. Rev.*, 177(September 2017), 589–612, doi:10.1016/j.earscirev.2017.12.007, 2018.
- 951 Eduard Suess: *Das Antlitz der Erde*, F. Tempsky; [etc., etc.]. [online] Available from:
952 <http://archive.org/details/dasantlitzderer02suesgoog> (Accessed 12 March 2020), 1892.
- 953 Eichelberger, N. and McQuarrie, N.: Kinematic reconstruction of the Bolivian orocline,
954 *Geosphere*, 11(2), 445–462, doi:10.1130/GES01064.1, 2015.
- 955 Escuder Viruete, J., Arenas, R. and Catalán, J. R. M.: Tectonothermal evolution associated with
956 Variscan crustal extension in the Tormes Gneiss Dome (NW Salamanca, Iberian Massif, Spain),
957 *Tectonophysics*, 238(1), 117–138, doi:10.1016/0040-1951(94)90052-3, 1994.
- 958 Fernández-Lozano, J., Pastor-Galán, D., Gutiérrez-Alonso, G. and Franco, P.: New kinematic
959 constraints on the Cantabrian orocline: A paleomagnetic study from the Peñalba and Truchas
960 synclines, NW Spain, *Tectonophysics*, 681, 195–208, doi:10.1016/j.tecto.2016.02.019, 2016.
- 961 Fernández-Lozano, J., Gutiérrez-Alonso, G., Willingshofer, E., Sokoutis, D., Vicente, G. de and
962 Cloetingh, S.: Shaping of intraplate mountain patterns: The Cantabrian orocline legacy in Alpine
963 Iberia, *Lithosphere*, 11(5), 708–721, doi:10.1130/L1079.1, 2019.
- 964 Fernández-Suárez, J., Gutiérrez-Alonso, G., Pastor-Galán, D., Hofmann, M., Murphy, J. B. and

- 965 Linnemann, U.: The Ediacaran-Early Cambrian detrital zircon record of NW Iberia: Possible
966 sources and paleogeographic constraints, *Int. J. Earth Sci.*, 103(5), 1335–1357,
967 doi:10.1007/s00531-013-0923-3, 2014.
- 968 Fossen, H.: *Structural Geology*, Cambridge University Press., 2016.
- 969 Franke, W., Cocks, L. R. M. and Torsvik, T. H.: The Palaeozoic Variscan oceans revisited,
970 *Gondwana Res.*, 48, 257–284, doi:10.1016/j.gr.2017.03.005, 2017.
- 971 Franke, W., Cocks, L. R. M. and Torsvik, T. H.: Detrital zircons and the interpretation of
972 palaeogeography, with the Variscan Orogeny as an example, *Geol. Mag.*, 157(4), 690–694,
973 doi:10.1017/S0016756819000943, 2020.
- 974 García-Arias, M., Díez-Montes, A., Villaseca, C. and Blanco-Quintero, I. F.: The Cambro-
975 Ordovician Olla de Sapo magmatism in the Iberian Massif and its Variscan evolution: A review,
976 *Earth-Sci. Rev.*, 176, 345–372, doi:10.1016/j.earscirev.2017.11.004, 2018.
- 977 Gil Toja, A., Jimenez-Ontiveros, P., and Seara Valero, J.R., La cuarta fase de deformación
978 hercínica en la Zona Centroibérica del Macizo Hespérico: Cuadernos Del Laboratório Xeoloxico
979 De Laxe, v. 9, p. 91–104, 1985.
- 980 Gong, Z., Langereis, C. G. and Mullender, T. A. T.: The rotation of Iberia during the Aptian and
981 the opening of the Bay of Biscay, *Earth Planet. Sci. Lett.*, 273(1–2), 80–93, 2008.
- 982 Gozalo, R., Liñán, E., Vintaned, J. A. G., Eugenia, M., Álvarez, D., Martorell, J. B. C., Zamora,
983 S., Esteve, J. and Mayoral, E.: The Cambrian of the Cadenas Ibéricas (ne Spain) and Its
984 Trilobites., n.d.
- 985 Greco, K. D., Johnston, S. T., Gutiérrez-Alonso, G., Shaw, J. and Lozano, J. F.: Interference
986 folding and orocline implications: A structural study of the Ponga Unit, Cantabrian orocline,
987 northern Spain, *Lithosphere*, 8(6), 757–768, doi:10.1130/L576.1, 2016.
- 988 Gutiérrez Alonso, G.: Estratigrafía y Tectónica., in *Memoria del Mapa Geológico de España*
989 1:50000, Hoja de Mora (680), IGME, Madrid. [online] Available from:
990 <http://info.igme.es/cartografiadigital/datos/magna50/memorias/MMagna0658.pdf>, 2009.
- 991 Gutiérrez-Alonso, G.: El Antiforme del Narcea y su relación con los mantos occidentales de la
992 Zona Cantábrica., 1992.
- 993 Gutiérrez-Alonso, G.: Strain partitioning in the footwall of the Somiedo Nappe: Structural
994 evolution of the Narcea Tectonic window, NW Spain, *J. Struct. Geol.*, 18(10), 1217–1229, 1996.
- 995 Gutiérrez-Alonso, G., Fernández-Suárez, J. and Weil, A. B.: Orocline triggered lithospheric
996 delamination, in *Special Paper 383: Orogenic curvature: Integrating paleomagnetic and*
997 *structural analyses*, vol. 383, edited by A. B. Weil and A. Sussman, pp. 121–130, Geological
998 Society of America, Boulder., 2004.
- 999 Gutiérrez-Alonso, G., Fernández-Suárez, J., Weil, A. B., Brendan Murphy, J., Damian Nance,
1000 R., Corf, F. and Johnston, S. T.: Self-subduction of the Pangaeon global plate, *Nat. Geosci.*,
1001 1(8), 549–553, doi:10.1038/ngeo250, 2008a.

- 1002 Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J. and Hamilton, M. A. A.: Rifting
1003 along the northern Gondwana margin and the evolution of the Rheic Ocean: A Devonian age for
1004 the El Castillo volcanic rocks (Salamanca, Central Iberian Zone), *Tectonophysics*, 461(1–4),
1005 157–165, doi:10.1016/j.tecto.2008.01.013, 2008b.
- 1006 Gutiérrez-Alonso, G., Fernández-Suárez, J., Jeffries, T. E. T. E. E., Johnston, S. T., Pastor-
1007 Galán, D., Murphy, J. B. B., Franco, M. P. P., Gonzalo, J. C. C.: Diachronous post-orogenic
1008 magmatism within a developing orocline in Iberia, *European Variscides*, *Tectonics*, 30(5), 17
1009 PP.-17 PP., doi:10.1029/2010TC002845, 2011a.
- 1010 Gutiérrez-Alonso, G., Murphy, J. B., Fernández-Suárez, J., Weil, A. B., Franco, M. P., Gonzalo,
1011 J. C.: Lithospheric delamination in the core of Pangea: Sm-Nd insights from the Iberian mantle,
1012 *Geology*, 39(2), 155–158, doi:10.1130/G31468.1, 2011b.
- 1013 Gutiérrez-Alonso, G., Collins, A. S., Fernández-Suárez, J., Pastor-Galán, D., González-Clavijo,
1014 E., Jourdan, F., Weil, A. B. and Johnston, S. T.: Dating of lithospheric buckling: $^{40}\text{Ar}/^{39}\text{Ar}$ ages
1015 of syn-orocline strike-slip shear zones in northwestern Iberia, *Tectonophysics*, 643, 44–54,
1016 doi:10.1016/j.tecto.2014.12.009, 2015.
- 1017 Gutiérrez-Alonso, G., Fernández-Suárez, J., López-Carmona, A. and Gärtner, A.: Exhuming a
1018 cold case: The early granodiorites of the northwest Iberian Variscan belt—A Visean magmatic
1019 flare-up?, *Lithosphere*, 10(2), 194–216, doi:10.1130/L706.1, 2018.
- 1020 Gutiérrez-Marco, J. C., Sá, A. A., García-Bellido, D. C. and Rábano, I.: The Bohemo-Iberian
1021 regional chronostratigraphical scale for the Ordovician System and palaeontological correlations
1022 within South Gondwana, *Lethaia*, 50(2), 258–295, doi:10.1111/let.12197, 2017.
- 1023 Gutiérrez-Marco, J. C., Piçarra, J. M., Meireles, C. A., Cózar, P., García-Bellido, D. C., Pereira,
1024 Z., Vaz, N., Pereira, S., Lopes, G., Oliveira, J. T., Quesada, C., Zamora, S., Esteve, J.,
1025 Colmenar, J., Bernárdez, E., Coronado, I., Lorenzo, S., Sá, A. A., Dias da Silva, Í., González-
1026 Clavijo, E., Díez-Montes, A. and Gómez-Barreiro, J.: Early Ordovician–Devonian Passive
1027 Margin Stage in the Gondwanan Units of the Iberian Massif, in *The Geology of Iberia: A*
1028 *Geodynamic Approach*, edited by C. Quesada and J. T. Oliveira, pp. 75–98, Springer
1029 International Publishing, Cham., 2019. Crespo-Blanc, A. and Orozco, M.: The Southern Iberian
1030 Shear Zone: a major boundary in the Hercynian folded belt. *Tectonophysics*, 148(3–4), pp.221-
1031 227., 1988.
- 1032 Hindle, D., Besson, O. and Burkhard, M.: A model of displacement and strain for arc-shaped
1033 mountain belts applied to the Jura arc, *J. Struct. Geol.*, 22(9), 1285–1296, doi:10.1016/S0191-
1034 8141(00)00038-9, 2000.
- 1035 van Hinsbergen, D. J. J., Torsvik, T. H., Schmid, S. M., Mañenco, L. C., Maffione, M., Vissers, R.
1036 L. M., Gürrer, D. and Spakman, W.: Orogenic architecture of the Mediterranean region and
1037 kinematic reconstruction of its tectonic evolution since the Triassic, *Gondwana Res.*, 81, 79–
1038 229, doi:10.1016/j.gr.2019.07.009, 2020.
- 1039 Hirt, A. M., Lowrie, W., Julivert, M., Arboleya, M. L., Lowric, A. M. H. W. and Arboleya, M. L.:
1040 Paleomagnetic results in support of a model for the origin of the Asturian arc, *Tectonophysics*,
1041 213(3–4), 321–339, 1992.
- 1042 Hnat, J. S. and van der Pluijm, B. A.: Foreland signature of indenter tectonics: Insights from

- 1043 calcite twinning analysis in the Tennessee salient of the Southern Appalachians, USA,
1044 *Lithosphere*, 3(5), 317–327, doi:10.1130/L151.1, 2011.
- 1045 Holmes, A.: A review of the continental drift hypothesis., *Min. Mag.*, 40, 205–209, 1929.
- 1046 Huang, W., Lippert, P. C., Zhang, Y., Jackson, M. J., Dekkers, M. J., Li, J., Hu, X., Zhang, B.,
1047 Guo, Z. and van Hinsbergen, D. J. J. J.: Remagnetization of carbonate rocks in southern Tibet:
1048 Perspectives from rock magnetic and petrographic investigations, *J. Geophys. Res. Solid Earth*,
1049 122(4), 2434–2456, doi:10.1002/2017JB013987, 2017.
- 1050 Izquierdo-Llavall, E., Sainz, A. C., Oliva-Urcia, B., Burmester, R., Pueyo, E. L. and Housen, B.:
1051 Multi-episodic remagnetization related to deformation in the Pyrenean Internal Sierras,
1052 *Geophys. J. Int.*, 201(2), 891–914, doi:10.1093/gji/ggv042, 2015.
- 1053 Izquierdo-Llavall, E., Casas-Sainz, A. M., Oliva-Urcia, B., Villalaín, J. J., Pueyo, E. and
1054 Scholger, R.: Rotational Kinematics of Basement Antiformal Stacks: Paleomagnetic Study of the
1055 Western Noguera Zone (Central Pyrenees), *Tectonics*, 37(10), 3456–3478,
1056 doi:10.1029/2018TC005153, 2018.
- 1057 Izquierdo-Llavall, E., Ayala, C., Pueyo, E. L., Casas-Sainz, A. M., Oliva-Urcia, B., Rubio, F.,
1058 Rodríguez-Pintó, A., Rey-Moral, C., Mediato, J. F. and García-Crespo, J.: Basement-Cover
1059 Relationships and Their Along-Strike Changes in the Linking Zone (Iberian Range, Spain): A
1060 Combined Structural and Gravimetric Study, *Tectonics*, 38(8), 2934–2960,
1061 doi:10.1029/2018TC005422, 2019.
- 1062 Jabaloy-Sánchez, A., Talavera, C., Gómez-Pugnaire, M. T., López-Sánchez-Vizcaíno, V.,
1063 Vázquez-Vílchez, M., Rodríguez-Peces, M. J. and Evans, N. J.: U-Pb ages of detrital zircons
1064 from the Internal Betics: A key to deciphering paleogeographic provenance and tectono-
1065 stratigraphic evolution, *Lithos*, 318–319, 244–266, doi:10.1016/j.lithos.2018.07.026, 2018.
- 1066 Jackson, M.: Diagenetic sources of stable remanence in remagnetized paleozoic cratonic
1067 carbonates: A rock magnetic study, *J. Geophys. Res. Solid Earth*, 95(B3), 2753–2761,
1068 doi:10.1029/JB095iB03p02753, 1990.
- 1069 Jacques, D., Muchez, P. and Sintubin, M.: Superimposed folding and W-Sn vein-type
1070 mineralisation in the Central Iberian Zone associated with late-Variscan oroclinal buckling: A
1071 structural analysis from the Regoufe area (Portugal), *Tectonophysics*, 742–743, 66–83,
1072 doi:10.1016/j.tecto.2018.05.021, 2018a.
- 1073 Jacques, D., Vieira, R., Muchez, P. and Sintubin, M.: Transpressional folding and associated
1074 cross-fold jointing controlling the geometry of post-orogenic vein-type W-Sn mineralization:
1075 examples from Minas da Panasqueira, Portugal, *Miner. Deposita*, 53(2), 171–194,
1076 doi:10.1007/s00126-017-0728-6, 2018b.
- 1077 Johnston, S. T.: The Great Alaskan Terrane Wreck: Reconciliation of paleomagnetic and
1078 geological data in the Northern Cordillera, *Earth Planet. Sci. Lett.*, 193(3–4), 259–272,
1079 doi:10.1016/S0012-821X(01)00516-7, 2001.
- 1080 Johnston, S. T., Weil, A. B. and Gutiérrez-Alonso, G.: Oroclines: Thick and thin, *GSA Bull.*,
1081 125(5–6), 643–663, doi:10.1130/B30765.1, 2013.

- 1082 Julivert, M.: L'évolution structurale de l'Arc Asturien, Paris., 1971.
- 1083 Julivert, M. and Arboleya, M. L.: A geometrical and kinematical approach to the nappe structure
1084 in an arcuate fold belt: the Cantabrian nappes (Hercynian chain, NW Spain), *J. Struct. Geol.*,
1085 6(5), 499–519, doi:10.1016/0191-8141(84)90061-0, 1984.
- 1086 Julivert, M. and Marcos, A.: superimposed folding under flexural conditions in the Cantabrian
1087 Zone (Hercynian Cordillera, Northwest Spain), *Am. J. Sci.*, 273(5), 353–375,
1088 doi:10.2475/ajs.273.5.353, 1973.
- 1089 Julivert, M., Fontboté, J. M., Ribeiro, A. and Nabais Conde, L. E.: Mapa Tectónico de la
1090 Península Ibérica y Baleares E: 1: 1.000. 000 y memoria explicativa, Publ IGME, 113, 1974.
- 1091 Julivert, M., Vegas, R., Roiz, J. M. and Martínez Rius, A.: Laestructura de la extension SE de la
1092 Zona Centro-Iberica con metamorfismo de bajo grado, *Rios-Geol. Esp. I*, 339–380, 1983.
- 1093 Keppie, F.: How subduction broke up Pangaea with implications for the supercontinent cycle,
1094 *Geol. Soc. Lond. Spec. Publ.*, 424(1), 265–288, doi:10.1144/SP424.8, 2016.
- 1095 Kirsch, M., Keppie, J. D., Murphy, J. B. and Solari, L. A.: Permian-Carboniferous arc
1096 magmatism and basin evolution along the western margin of Pangea: Geochemical and
1097 geochronological evidence from the eastern Acatlan Complex, southern Mexico, *Geol. Soc. Am.*
1098 *Bull.*, 124(9–10), 1607–1628, doi:10.1130/B30649.1, 2012.
- 1099 Kollmeier, J. M., van der Pluijm, B. A. and Van der Voo, R.: Analysis of Variscan dynamics;
1100 early bending of the Cantabria-Asturias Arc, northern Spain, *Earth Planet. Sci. Lett.*, 181(1–2),
1101 203–216, doi:10.1016/S0012-821X(00)00203-X, 2000.
- 1102 Kroner, U. and Romer, R. L. L.: Two plates — Many subduction zones: The Variscan orogeny
1103 reconsidered, *Gondwana Res.*, 24(1), 298–329, doi:10.1016/j.gr.2013.03.001, 2013.
- 1104 Li, P. and Rosenbaum, G.: Does the Manning Orocline exist? New structural evidence from the
1105 inner hinge of the Manning Orocline (eastern Australia), *Gondwana Res.*, 25(4), 1599–1613,
1106 doi:10.1016/j.gr.2013.06.010, 2014.
- 1107 Li, P., Sun, M., Rosenbaum, G., Yuan, C., Safonova, I., Cai, K., Jiang, Y. and Zhang, Y.:
1108 Geometry, kinematics and tectonic models of the Kazakhstan Orocline, Central Asian Orogenic
1109 Belt, *J. Asian Earth Sci.*, 153(July 2017), 42–56, doi:10.1016/j.jseas.2017.07.029, 2018.
- 1110 Li, P.-F., Rosenbaum, G. and Rubatto, D.: Triassic asymmetric subduction rollback in the
1111 southern New England Orogen (eastern Australia): the end of the Hunter-Bowen Orogeny, *Aust.*
1112 *J. Earth Sci.*, 59(6), 965–981, doi:10.1080/08120099.2012.696556, 2012.
- 1113 López-Carmona, A., Abati, J., Pitra, P. and Lee, J. K. W.: Retrogressed lawsonite blueschists
1114 from the NW Iberian Massif: P–T–t constraints from thermodynamic modelling and 40Ar/39Ar
1115 geochronology, *Contrib. Mineral. Petrol.*, 167(3), 987–987, doi:10.1007/s00410-014-0987-5,
1116 2014.
- 1117 López-Moro, F. J., López-Plaza, M., Gutiérrez-Alonso, G., Fernández-Suárez, J., López-
1118 Carmona, A., Hofmann, M. and Romer, R. L.: Crustal melting and recycling: geochronology and
1119 sources of Variscan syn-kinematic anatectic granitoids of the Tormes Dome (Central Iberian

- 1120 Zone). A U–Pb LA-ICP-MS study, *Int. J. Earth Sci.*, 107(3), 985–1004, doi:10.1007/s00531-017-
1121 1483-8, 2018.
- 1122 Lotze, F.: Zur gliederung der Variszischen der Iberischen Meseta., *Geotektonische Forschungen*,
1123 6, 78–92, 1945.
- 1124 Macaya, J., González-Lodeiro, F., Martínez-Catalán, J. R. and Alvarez, F.: Continuous
1125 deformation, ductile thrusting and backfolding of cover and basement in the Sierra de
1126 Guadarrama, Hercynian orogen of central Spain, *Tectonophysics*, 191(3), 291–309,
1127 doi:10.1016/0040-1951(91)90063-X, 1991.
- 1128 Maffione, M., Speranza, F. and Faccenna, C.: Bending of the Bolivian orocline and growth of the
1129 central Andean plateau: Paleomagnetic and structural constraints from the Eastern Cordillera
1130 (22–24°S, NW Argentina), *Tectonics*, 28, 23 PP.-23 PP., doi:10.1029/2008TC002402, 2009.
- 1131 Marcos, A. and Pulgar, J. A. A.: An Approach to the tectonostratigraphic evolution of the
1132 Cantabrian foreland thrust and fold belt, Hercynian Cordillera of NW Spain, *Neues Jahrb. Geol.*
1133 *Palaontologie-Abh.*, 163(2), 256–260, 1982.
- 1134 Marcos, A., Pérez-Estaún, A., Martínez Catalán, J. R. and Gutiérrez-Marco, J. C.: *Estratigrafía y*
1135 *Paleogeografía. Zona Asturoccidental-Leonesa*, 2004.
- 1136 Marshak, S.: KINEMATICS OF OROCLINE AND ARC FORMATION IN THIN-SKINNED
1137 OROGENS, *Tectonics*, 7(1), 73–86, 1988.
- 1138 Marshak, S.: Salients, Recesses, Arcs, Oroclines, and Syntaxes — A Review of Ideas
1139 Concerning the Formation of Map-view Curves in Fold-thrust Belts, *Thrust Tecton. Hydrocarb.*
1140 *Syst. AAPG Mem.* 82, 82(1893), 131–156, 2004.
- 1141 Martín-Algarra, A., García-Casco, A., Gómez-Pugnaire, M. T., Jabaloy-Sánchez, A., Laborda-
1142 López, C., López Sánchez-Vizcaíno, V., Mazzoli, S., Navas-Parejo, P., Perrone, V., Rodríguez-
1143 Cañero, R. and Sánchez-Navas, A.: Paleozoic Basement and Pre-Alpine History of the Betic
1144 Cordillera, in *The Geology of Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T.
1145 Oliveira, pp. 261–305, Springer International Publishing, Cham., 2019.
- 1146 Martínez Catalán, J. R.: Are the oroclines of the Variscan belt related to late Variscan strike-slip
1147 tectonics?, *Terra Nova*, 23(4), 241–247, doi:10.1111/j.1365-3121.2011.01005.x, 2011.
- 1148 Martínez Catalán, J. R.: The Central Iberian arc, an orocline centered in the Iberian Massif and
1149 some implications for the Variscan belt, *Int. J. Earth Sci.*, 101(5), 1–16, doi:10.1007/s00531-
1150 011-0715-6, 2012.
- 1151 Martínez Catalan, J. R., Rodriguez, M. P. H., Alonso, P. V., Perez-Estaun, A. and Lodeiro, F. G.:
1152 Lower Paleozoic extensional tectonics in the limit between the West Asturian-Leonese and
1153 Central Iberian Zones of the Variscan Fold-Belt in NW Spain, *Geol. Rundsch.*, 81(2), 545–560,
1154 doi:10.1007/BF01828614, 1992.
- 1155 Martínez Catalán, J. R., Martínez Poyatos, D. and Bea, F.: Zona Centroibérica, *Geol. Esp.*, 68–
1156 133, 2004.
- 1157 Martínez Catalán, J. R., Arenas, R., Abati, J., Martínez, S. S., García, F. D., Suárez, J. F.,

- 1158 Cuadra, P. G., Castiñeiras, P., Barreiro, J. G., Montes, A. D., Clavijo, E. G., Pascual, F. J. R.,
1159 Andonaegui, P., Jeffries, T. E., Alcock, J. E., Fernández, R. D. and Carmona, A. L.: A rootless
1160 suture and the loss of the roots of a mountain chain: The Variscan belt of NW Iberia, *Comptes*
1161 *Rendus Geosci.*, 341(2), 114–126, doi:10.1016/j.crte.2008.11.004, 2009.
- 1162 Martínez Catalan, J. R., Rubio Pascual, F. J., Diez Montes, A., Diez Fernandez, R., Gomez
1163 Barreiro, J., Dias da Silva, I., Gonzalez Clavijo, E., Ayarza, P., Alcock, J. E., Montes, A. D.,
1164 Fernandez, R. D., Barreiro, J. G., Dias da Silva, I., Clavijo, E. G., Ayarza, P., Alcock, J. E., Diez
1165 Montes, A., Diez Fernandez, R., Gomez Barreiro, J., Dias da Silva, I., Gonzalez Clavijo, E.,
1166 Ayarza, P., Alcock, J. E., Montes, A. D., Fernandez, R. D., Barreiro, J. G., Dias da Silva, I.,
1167 Clavijo, E. G., Ayarza, P. and Alcock, J. E.: The late Variscan HT/LP metamorphic event in NW
1168 and Central Iberia: relationships to crustal thickening, extension, orocline development and
1169 crustal evolution, *Geol. Soc. Lond. Spec. Publ.*, 405(1), 225–247, doi:10.1144/SP405.1, 2014.
- 1170 Martínez Catalán, J. R., Aerden, D. G. A. M. and Carreras, J.: The “Castilian bend” of Rudolf
1171 Staub (1926): historical perspective of a forgotten orocline in Central Iberia, *Swiss J. Geosci.*,
1172 108(2–3), 289–303, doi:10.1007/s00015-015-0202-3, 2015.
- 1173 Martínez Catalán, J. R., Gómez Barreiro, J., Dias da Silva, Í., Chichorro, M., López-Carmona,
1174 A., Castiñeiras, P., Abati, J., Andonaegui, P., Fernández-Suárez, J., González Cuadra, P. and
1175 Benítez-Pérez, J. M.: Variscan Suture Zone and Suspect Terranes in the NW Iberian Massif:
1176 Allochthonous Complexes of the Galicia-Trás os Montes Zone (NW Iberia), in *The Geology of*
1177 *Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T. Oliveira, pp. 99–130, Springer
1178 International Publishing, Cham., 2019.
- 1179 Martínez-Catalán, J. R.: A non-cylindrical model for the northwest Iberian allochthonous
1180 terranes and their equivalents in the Hercynian belt of Western Europe, *Tectonophysics*, 179(3–
1181 4), 253–272, doi:10.1016/0040-1951(90)90293-H, 1990.
- 1182 McWilliams, C. K., Kunk, M. J., Wintsch, R. P. and Bish, D. L.: Determining ages of multiple
1183 muscovite-bearing foliations in phyllonites using the $^{40}\text{Ar}/^{39}\text{Ar}$ step heating method:
1184 Applications to the alleghanian orogeny in central new England, *Am. J. Sci.*, 313(10), 996–1016,
1185 doi:10.2475/10.2013.02, 2013.
- 1186 Meijers, M. J. M., Smith, B., Pastor-Galán, D., Degenaar, R., Sadradze, N., Adamia, S.,
1187 Sahakyan, L., Avagyan, A., Sosson, M., Rolland, Y., Langereis, C. G. and Müller, C.:
1188 Progressive orocline formation in the Eastern Pontides-Lesser Caucasus, *Geol. Soc. Spec.*
1189 *Publ.*, 428(1), doi:10.1144/SP428.8, 2017.
- 1190 Merino-Tomé, O. A., Bahamonde, J. R., Colmenero, J. R., Heredia, N., Villa, E. and Farias, P.:
1191 Emplacement of the Cuera and Picos de Europa imbricate system at the core of the Iberian-
1192 Armorican arc (Cantabrian zone, north Spain): New precisions concerning the timing of arc
1193 closure, *Bull. Geol. Soc. Am.*, 121(5–6), 729–751, doi:10.1130/B26366.1, 2009.
- 1194 Molina Garza, R. S. and Zijdeveld, J. D. a.: Paleomagnetism of Paleozoic strata, Brabant and
1195 Ardennes Massifs, Belgium: Implications of prefolding and postfolding Late Carboniferous
1196 secondary magnetizations for European apparent polar wander, *J. Geophys. Res.*, 101(B7),
1197 15799–15799, doi:10.1029/96JB00325, 1996.
- 1198 Moral, B. del and Sarmiento, G. N.: Conodontos del Katiense (Ordovícico Superior) del sector
1199 meridional de la Zona Centroibérica (España), *Rev. Esp. Micropaleontol.*, 40(3), 169–245, 2008.

- 1200 Munha, J.: Metamorphic Evolution of the South Portuguese/Pulo Do Lobo Zone, in Pre-
1201 Mesozoic Geology of Iberia, edited by R. D. Dallmeyer and E. M. Garcia, pp. 363–368,
1202 Springer, Berlin, Heidelberg., 1990.
- 1203 Munha, J., Barriga, F. J. a. S. and Kerrich, R.: High 18 O ore-forming fluids in volcanic-hosted
1204 base metal massive sulfide deposits; geologic, 18 O/ 16 O, and D/H evidence from the Iberian
1205 pyrite belt; Crandon, Wisconsin; and Blue Hill, Maine, Econ. Geol., 81(3), 530–552, doi:10.2113/
1206 gsecongeo.81.3.530, 1986.
- 1207 Murphy, J. B., Gutierrez-Alonso, G., Fernandez-Suarez, J. and Braid, J. A.: Probing crustal and
1208 mantle lithosphere origin through Ordovician volcanic rocks along the Iberian passive margin of
1209 Gondwana, Tectonophysics, 461(1–4), 166–180, 2008.
- 1210 Nance, R. D., Gutiérrez-Alonso, G., Keppie, J. D., Linnemann, U., Murphy, J. B., Quesada, C.,
1211 Strachan, R. A. and Woodcock, N. H.: Evolution of the Rheic Ocean, Gondwana Res., 17(2–3),
1212 194–222, doi:10.1016/j.gr.2009.08.001, 2010.
- 1213 Oliveira, J. T., Quesada, C., Pereira, Z., Matos, J. X., Solá, A. R., Rosa, D., Albardeiro, L., Díez-
1214 Montes, A., Morais, I., Inverno, C., Rosa, C. and Relvas, J.: South Portuguese Terrane: A
1215 Continental Affinity Exotic Unit, in The Geology of Iberia: A Geodynamic Approach, edited by C.
1216 Quesada and J. T. Oliveira, pp. 173–206, Springer International Publishing, Cham., 2019a.
- 1217 Oliveira, J. T., González-Clavijo, E., Alonso, J., Armendáriz, M., Bahamonde, J. R., Braid, J. A.,
1218 Colmenero, J. R., Dias da Silva, Í., Fernandes, P., Fernández, L. P., Gabaldón, V., Jorge, R. S.,
1219 Machado, G., Marcos, A., Merino-Tomé, Ó., Moreira, N., Murphy, J. B., Pinto de Jesus, A.,
1220 Quesada, C., Rodrigues, B., Rosales, I., Sanz-López, J., Suárez, A., Villa, E., Piçarra, J. M. and
1221 Pereira, Z.: Synorogenic Basins, in The Geology of Iberia: A Geodynamic Approach, edited by
1222 C. Quesada and J. T. Oliveira, pp. 349–429, Springer International Publishing, Cham., 2019b.
- 1223 P. Farias, G. G.: Aportaciones al conocimiento de la litoestratigrafía y estructura de Galicia
1224 Central, Mem. Fac. Ciênc. Universidade Porto, 411–431, 1987.
- 1225 Palero-Fernández, F. J., Martin-Izard, A., Zarzalejos Prieto, M. and Mansilla-Plaza, L.:
1226 Geological context and plumbotectonic evolution of the giant Almadén Mercury Deposit, Ore
1227 Geol. Rev., 64, 71–88, doi:10.1016/j.oregeorev.2014.06.013, 2015.
- 1228 Parés, J. M. and Van der Voo, R.: Paleozoic paleomagnetism of Almadén, Spain: A cautionary
1229 note, J. Geophys. Res., 97(B6), 9353–9356, doi:10.1029/91JB03073, 1992.
- 1230 Parés, J. M., Van der Voo, R., Stamatakis, J. and Pérez-Estaún, A.: Remagnetizations and
1231 postfolding oroclinal rotations in the Cantabrian/Asturian arc, northern Spain, Tectonics, 13(6),
1232 1461–1471, doi:10.1029/94TC01871, 1994.
- 1233 Pastor-Galán, D., Gutiérrez-Alonso, G., Meere, P. A. and Mulchrone, K. F.: Factors affecting
1234 finite strain estimation in low-grade, low-strain clastic rocks, J. Struct. Geol., 31(12), 1586–1596,
1235 doi:10.1016/j.jsrg.2009.08.005, 2009.
- 1236 Pastor-Galán, D., Gutiérrez-Alonso, G. and Weil, A. B.: Orocline timing through joint analysis:
1237 Insights from the Ibero-Armorican Arc, Tectonophysics, 507(1), 31–46,
1238 doi:10.1016/j.tecto.2011.05.005, 2011.

- 1239 Pastor-Galán, D., Gutiérrez-Alonso, G., Zulauf, G. and Zanella, F.: Analogue modeling of
1240 lithospheric-scale orocline buckling: Constraints on the evolution of the Iberian-Armorican arc,
1241 *Bull. Geol. Soc. Am.*, 124(7–8), doi:10.1130/B30640.1, 2012a.
- 1242 Pastor-Galán, D., Gutiérrez-Alonso, G., Mulchrone, K. F. and Huerta, P.: Conical folding in the
1243 core of an orocline. A geometric analysis from the Cantabrian Arc (Variscan Belt of NW Iberia),
1244 *J. Struct. Geol.*, 39, 210–223, doi:10.1016/j.jsg.2012.02.010, 2012b.
- 1245 Pastor-Galán, D., Gutiérrez-Alonso, G., Murphy, J. B. B., Fernández-Suárez, J., Hofmann, M.
1246 and Linnemann, U.: Provenance analysis of the Paleozoic sequences of the northern
1247 Gondwana margin in NW Iberia: Passive margin to Variscan collision and orocline development,
1248 *Gondwana Res.*, 23(3), 1089–1103, doi:10.1016/j.gr.2012.06.015, 2013a.
- 1249 Pastor-Galán, D., Gutiérrez-Alonso, G., Fernández-Suárez, J., Murphy, J. B. and Nieto, F.:
1250 Tectonic evolution of NW Iberia during the Paleozoic inferred from the geochemical record of
1251 detrital rocks in the Cantabrian Zone, *Lithos*, 182–183, 221–228,
1252 doi:10.1016/j.lithos.2013.09.007, 2013b.
- 1253 Pastor-Galán, D., Martín-Merino, G. and Corrochano, D.: Timing and structural evolution in the
1254 limb of an orocline: The Pisuerga-Carrión Unit (southern limb of the Cantabrian Orocline, NW
1255 Spain), *Tectonophysics*, 622, 110–121, doi:10.1016/j.tecto.2014.03.004, 2014.
- 1256 Pastor-Galán, D., Ursem, B., Meere, P. A. and Langereis, C.: Extending the Cantabrian
1257 Orocline to two continents (from Gondwana to Laurussia). Paleomagnetism from South Ireland,
1258 *Earth Planet. Sci. Lett.*, 432, doi:10.1016/j.epsl.2015.10.019, 2015a.
- 1259 Pastor-Galan, D., Groenewegen, T., Brouwer, D., Krijgsman, W., Dekkers, M. J., Pastor-galán,
1260 D., Groenewegen, T., Brouwer, D., Krijgsman, W. and Dekkers, M. J.: One or two oroclines in
1261 the Variscan orogen of Iberia? Implications for Pangea amalgamation, *Geology*, 43(6), 527–
1262 530, doi:10.1130/G36701.1, 2015b.
- 1263 Pastor-Galán, D., Dekkers, M. J., Gutiérrez-Alonso, G., Brouwer, D., Groenewegen, T.,
1264 Krijgsman, W., Fernández-Lozano, J., Yenes, M. and Álvarez-Lobato, F.: Paleomagnetism of
1265 the Central Iberian curve's putative hinge: Too many oroclines in the Iberian Variscides,
1266 *Gondwana Res.*, 39, 96–113, doi:10.1016/j.gr.2016.06.016, 2016.
- 1267 Pastor-Galán, D., Mulchrone, K. F., Koymans, M. R. , van Hinsbergen, D. J. J. and Langereis,
1268 C. G. : Bootstrapped total least squares orocline test: A robust method to quantify vertical-axis
1269 rotation patterns in orogens, with examples from the Cantabrian and Aegean oroclines,
1270 *Lithosphere*, 9(3), L547.1-L547.1, doi:10.1130/L547.1, 2017a.
- 1271 Pastor-Galán, D., Gutiérrez-Alonso, G., Dekkers, M. J. M. J. and Langereis, C. G.:
1272 Paleomagnetism in Extremadura (Central Iberian zone, Spain) Paleozoic rocks: extensive
1273 remagnetizations and further constraints on the extent of the Cantabrian orocline, *J. Iber. Geol.*,
1274 43(4), 583–600, doi:10.1007/s41513-017-0039-x, 2017b.
- 1275 Pastor-Galán, D., Pueyo, E. L., Diederer, M., García-Lasanta, C. and Langereis, C. G.: Late
1276 Paleozoic Iberian Orocline(s) and the Missing Shortening in the Core of Pangea.
1277 Paleomagnetism From the Iberian Range, *Tectonics*, 37(10), 3877–3892,
1278 doi:10.1029/2018TC004978, 2018.

- 1279 Pastor-Galán, D., Nance, R. D., Murphy, J. B. and Spencer, C. J.: Supercontinents: myths,
1280 mysteries, and milestones, *Geol. Soc. Lond. Spec. Publ.*, 470, 39–64, doi:10.1144/SP470.16,
1281 2019a.
- 1282 Pastor-Galán, D., Dias da Silva, Í. F., Groenewegen, T. and Krijgsman, W.: Tangled up in folds:
1283 tectonic significance of superimposed folding at the core of the Central Iberian curve (West
1284 Iberia), *Int. Geol. Rev.*, 61(2), 240–255, doi:10.1080/00206814.2017.1422443, 2019b.
- 1285 Pereira, I., Dias, R., Bento, T. and Mata, J.: Exhumation of a migmatite complex along a
1286 transpressive shear zone : inferences from the Variscan Juzbado – Penalva do Castelo Shear
1287 Zone (Central Iberian Zone), *J. Geol. Soc.*, doi:10.1144/jgs2016-159, 2017.
- 1288 Pereira, M. F., Chichorro, M., Silva, J. B., Ordóñez-Casado, B., Lee, J. K. W. W. and Williams, I.
1289 S.: Early carboniferous wrenching, exhumation of high-grade metamorphic rocks and basin
1290 instability in SW Iberia: Constraints derived from structural geology and U–Pb and ⁴⁰Ar–³⁹Ar
1291 geochronology, *Tectonophysics*, 558, 28–44, doi:10.1016/j.tecto.2012.06.020, 2012.
- 1292 Pereira, M. F., Díez Fernández, R., Gama, C., Hofmann, M., Gärtner, A. and Linnemann, U.: S-
1293 type granite generation and emplacement during a regional switch from extensional to
1294 contractional deformation (Central Iberian Zone, Iberian autochthonous domain, Variscan
1295 Orogeny), *Int. J. Earth Sci.*, 107(1), 251–267, doi:10.1007/s00531-017-1488-3, 2018.
- 1296 Pérez-Cáceres, I., Poyatos, D. M., Simancas, J. F. and Azor, A.: The elusive nature of the
1297 Rheic Ocean suture in SW Iberia, *Tectonics*, 34(12), 2429–2450, doi:10.1002/2015TC003947,
1298 2015.
- 1299 Pérez-Cáceres, I., Simancas, J. F., Martínez Poyatos, D., Azor, A. and González Lodeiro, F.:
1300 Oblique collision and deformation partitioning in the SW Iberian Variscides, *Solid Earth*, 7(3),
1301 857–872, doi:https://doi.org/10.5194/se-7-857-2016, 2016.
- 1302 Pérez-Cáceres, I., Martínez Poyatos, D., Simancas, J. F. and Azor, A.: Testing the Avalonian
1303 affinity of the South Portuguese Zone and the Neoproterozoic evolution of SW Iberia through
1304 detrital zircon populations, *Gondwana Res.*, 42, 177–192, doi:10.1016/j.gr.2016.10.010, 2017.
- 1305 Pérez-Cáceres, I., Martínez Poyatos, D. J., Vidal, O., Beyssac, O., Nieto, F., Simancas, J. F.,
1306 Azor, A. and Bourdelle, F.: Deciphering the metamorphic evolution of the Pulo do Lobo
1307 metasedimentary domain (SW Iberian Variscides), *Solid Earth*, 11(2), 469–488,
1308 doi:https://doi.org/10.5194/se-11-469-2020, 2020.
- 1309 Pérez-Estaún, A., Bastida, F., Alonso, J. L., Marquinez, J., Aller, J., Alvarezmarron, J., Marcos,
1310 A. and Pulgar, J. A.: A THIN-SKINNED TECTONICS MODEL FOR AN ARCUATE FOLD AND
1311 THRUST BELT - THE CANTABRIAN ZONE (VARISCAN IBERO-ARMORICAN ARC),
1312 *Tectonics*, 7(3), 517–537, 1988.
- 1313 Pérez-Estaún, A., Bastida, F., Martínez Catalán, J. R., Gutiérrez-Marco, J. C., Marcos, A. and
1314 Pulgar, J.: Stratigraphy of the West Asturian-Leonese Zone, Springer. [online] Available from:
1315 https://digital.csic.es/handle/10261/30719 (Accessed 6 April 2020), 1990.
- 1316 Pérez-Estaún, A., Martinezcatalan, J. R. and Bastida, F.: CRUSTAL THICKENING AND
1317 DEFORMATION SEQUENCE IN THE FOOTWALL TO THE SUTURE OF THE VARISCAN
1318 BELT OF NORTHWEST SPAIN, *Tectonophysics*, 191(3–4), 243–253, 1991.

- 1319 Perroud, H., Bonhommet, N. and Ribeiro, A.: Paleomagnetism of Late Paleozoic igneous rocks
1320 from southern Portugal, *Geophys. Res. Lett.*, 12(1), 45–48, doi:10.1029/GL012i001p00045,
1321 1985.
- 1322 Perroud, H., Calza, F. and Khattach, D.: Paleomagnetism of the Silurian Volcanism at Almaden,
1323 Southern Spain, *J. Geophys. Res.-Solid Earth Planets*, 96(B2), 1949–1962,
1324 doi:10.1029/90JB02226, 1991.
- 1325 Pin, C., Paquette, J. L., Zalduegui, J. F. S. and Ibarguchi, J. I. G.: Early Devonian
1326 suprasubduction-zone ophiolite related to incipient collisional processes in the Western
1327 Variscan Belt: The Sierra de Careón unit, Ordenes Complex, Galicia, in *Variscan-Appalachian*
1328 *dynamics: The building of the late Paleozoic basement*, Geological Society of America., 2002.
- 1329 Pueyo, E. L., Mauritsch, H. J., Gawlick, H.-J., Scholger, R. and Frisch, W.: New evidence for
1330 block and thrust sheet rotations in the central northern Calcareous Alps deduced from two
1331 pervasive remagnetization events, *Tectonics*, 26(5), n/a-n/a, doi:10.1029/2006TC001965, 2007.
- 1332 Pueyo, E. L., Sussman, A. J., Oliva-Urcia, B. and Cifelli, F.: Palaeomagnetism in fold and thrust
1333 belts: Use with caution, *Geol. Soc. Spec. Publ.*, 425(1), 259–276, doi:10.1144/SP425.14, 2016.
- 1334 Quesada, C.: The Ossa-Morena Zone of the Iberian Massif: a tectonostratigraphic approach to
1335 its evolution, *Z. Dtsch. Ges. Für Geowiss.*, 585–595, doi:10.1127/1860-1804/2006/0157-0585,
1336 2006.
- 1337 Quesada, C. and Dallmeyer, R. D. D.: Tectonothermal evolution of the Badajoz-Cordóba shear
1338 zone (SW Iberia): characteristics and $40\text{Ar}/39\text{Ar}$ mineral age constraints, *Tectonophysics*,
1339 231(1–3), 195–213, doi:10.1016/0040-1951(94)90130-9, 1994.
- 1340 Quesada, C., Braid, J. A., Fernandes, P., Ferreira, P., Jorge, R. S., Matos, J. X., Murphy, J. B.,
1341 Oliveira, J. T., Pedro, J. and Pereira, Z.: SW Iberia Variscan Suture Zone: Oceanic Affinity
1342 Units, in *The Geology of Iberia: A Geodynamic Approach*, edited by C. Quesada and J. T.
1343 Oliveira, pp. 131–171, Springer International Publishing, Cham., 2019.
- 1344 Rankin, D. W.: Appalachian salients and recesses: Late Precambrian continental breakup and
1345 the opening of the Iapetus Ocean, *J. Geophys. Res.* 1896-1977, 81(32), 5605–5619,
1346 doi:10.1029/JB081i032p05605, 1976.
- 1347 Ribeiro, A., Munhá, J., Dias, R., Mateus, A., Pereira, E., Ribeiro, L., Fonseca, P., Araújo, A.,
1348 Oliveira, T., Romão, J., Chaminé, H., Coke, C. and Pedro, J.: Geodynamic evolution of the SW
1349 Europe Variscides, *Tectonics*, 26(6), doi:10.1029/2006TC002058, 2007.
- 1350 Ries, A. C. and Shackleton, R. M.: Catazonal Complexes of North-West Spain and North
1351 Portugal, Remnants of a Hercynian Thrust Plate, *Nat. Phys. Sci.*, 234(47), 65–68,
1352 doi:10.1038/physci234065a0, 1971.
- 1353 Ries, A. C. and Shackleton, R. M.: Patterns of Strain Variation in Arcuate Fold Belts, *Philos.*
1354 *Trans. R. Soc. Lond. Ser. Math. Phys. Sci.*, 283(1312), 281–288, 1976.
- 1355 Rodríguez-Cañero, R., Jabaloy-Sánchez, A., Navas-Parejo, P., Martín-Algarra, A., Rodríguez,
1356 R., Antonio, C., Sánchez, J., Navas, P., Agustín, P. and Algarra, M.: Linking Palaeozoic
1357 palaeogeography of the Betic Cordillera to the Variscan Iberian Massif : new insight through the

- 1358 first conodonts of the Nevado-Filábride Complex, *Int. J. Earth Sci.*, 107(5), 1791–1806,
1359 doi:10.1007/s00531-017-1572-8, 2018.
- 1360 Rosenbaum, G.: Geodynamics of oroclinal bending: Insights from the Mediterranean, *J.*
1361 *Geodyn.*, 82, 5–15, doi:10.1016/j.jog.2014.05.002, 2014.
- 1362 Rubio Pascual, F. J., Arenas, R., Martínez Catalán, J. R., Rodríguez Fernández, L. R. and
1363 Wijbrans, J. R.: Thickening and exhumation of the Variscan roots in the Iberian Central System:
1364 Tectonothermal processes and $^{40}\text{Ar}/^{39}\text{Ar}$ ages, *Tectonophysics*, 587, 207–221,
1365 doi:10.1016/j.tecto.2012.10.005, 2013.
- 1366 Rubio Pascual, F. J., López-Carmona, A. and Arenas, R.: Thickening vs. extension in the
1367 Variscan belt: P–T modelling in the Central Iberian autochthon, *Tectonophysics*, 681, 144–158,
1368 doi:10.1016/j.tecto.2016.02.033, 2016.
- 1369 Sánchez-García, T., Chichorro, M., Solá, A. R., Álvaro, J. J., Díez-Montes, A., Bellido, F.,
1370 Ribeiro, M. L., Quesada, C., Lopes, J. C., Dias da Silva, Í., González-Clavijo, E., Gómez
1371 Barreiro, J. and López-Carmona, A.: The Cambrian-Early Ordovician Rift Stage in the
1372 Gondwanan Units of the Iberian Massif, in *The Geology of Iberia: A Geodynamic Approach*,
1373 edited by C. Quesada and J. T. Oliveira, pp. 27–74, Springer International Publishing, Cham.,
1374 2019.
- 1375 Schulz, G.: Descripción geológica de Asturias: Publicada de Real Orden. Con un atlas, José
1376 Gonzalez., 1858.
- 1377 Schwartz, S. Y. and Vandervoo, R.: PALEOMAGNETIC EVALUATION OF THE OROCLINE
1378 HYPOTHESIS IN THE CENTRAL AND SOUTHERN APPALACHIANS, *Geophys. Res. Lett.*,
1379 10(7), 505–508, 1983.
- 1380 Shaw, J. and Johnston, S. T.: Terrane wrecks (coupled oroclinal) and paleomagnetic inclination
1381 anomalies, *Earth-Sci. Rev.*, 154, 191–209, doi:10.1016/j.earscirev.2016.01.003, 2016.
- 1382 Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Weil, A. B.: Oroclines of the Variscan
1383 orogen of Iberia: Paleocurrent analysis and paleogeographic implications, *Earth Planet. Sci.*
1384 *Lett.*, 329–330, 60–70, doi:10.1016/j.epsl.2012.02.014, 2012.
- 1385 Shaw, J., Johnston, S. T., Gutiérrez-Alonso, G. and Pastor-Galán, D.: Provenance variability
1386 along the Early Ordovician north Gondwana margin: Paleogeographic and tectonic implications
1387 of U–Pb detrital zircon ages from the Armorican Quartzite of the Iberian Variscan belt, *Bull. Geol.*
1388 *Soc. Am.*, 126(5–6), 702–719, doi:10.1130/B30935.1, 2014.
- 1389 Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A
1390 structural study of the Cantabrian orocline, NW Iberian Massif, *Lithosphere*, 7(6), 653–661,
1391 doi:10.1130/L461.1, 2015.
- 1392 Shaw, J., Johnston, S. T. and Gutiérrez-Alonso, G.: Orocline formation at the core of Pangea: A
1393 structural study of the Cantabrian orocline, NW Iberian Massif, *Lithosphere*, 8(1), 97–97,
1394 doi:10.1130/L461.1, 2016.
- 1395 Silva, Í. D. D., Valverde-Vaquero, P., González-Clavijo, E., Díez-Montes, A. and Catalán, J. R.
1396 M.: Structural and stratigraphical significance of U–Pb ages from the Mora and Saldanha

- 1397 volcanic complexes (NE Portugal, Iberian Variscides), *Geol. Soc. Lond. Spec. Publ.*, 405(1),
1398 115–135, doi:10.1144/SP405.3, 2014.
- 1399 Simancas, J. F., Carbonell, R., Lodeiro, F. G., Estaun, A., Juhlin, C., Ayarza, P., Kashubin, A.,
1400 Azor, A., Poyatos, D. M., Almodovar, G. R., Pascual, E., Saez, R. and Exposito, I.: Crustal
1401 structure of the transpressional Variscan orogen of SW Iberia: SW Iberia deep seismic reflection
1402 profile (IBERSEIS), *Tectonics*, 22(6), 25–25, 2003.
- 1403 Simancas, J. F., Ayarza, P., Azor, a., Carbonell, R., Martínez Poyatos, D., Pérez-Estaún, a. and
1404 González Lodeiro, F.: A seismic geotraverse across the Iberian Variscides: Orogenic
1405 shortening, collisional magmatism, and orocline development, *Tectonics*, 32(i), n/a-n/a,
1406 doi:10.1002/tect.20035, 2013.
- 1407 Solís-Alulima, B. E., López-Carmona, A., Gutiérrez Alonso, G. and Álvarez Valero, A. M.:
1408 Petrologic and thermobarometric study of the Riás schists (NW Iberian Massif), *Bol. Geológico*
1409 *Min.*, 130(3), 445–464, doi:10.21701/bolgeomin.130.3.004, 2019.
- 1410 Stampfli, G. M. and Borel, G. D.: The TRANSMED Transects in Space and Time: Constraints
1411 from Paleotectonic Evolution of the Mediterranean Domain., edited by W. Cavazza, F. M.
1412 Roure, W. Spakman, G. M. Stampfli, and P. A. Ziegler, pp. 53–80, Springer, Berlin., 2004.
- 1413 Stampfli, G. M., Hochard, C., Vérard, C., Wilhem, C. and VonRaumer, J.: The Formation of
1414 Pangea, *Tectonophysics*, 593, 1–19, doi:10.1016/j.tecto.2013.02.037, 2013.
- 1415 Staub, R.: *Gedanken zur Tektonik Spaniens.*, Zürich., 1926.
- 1416 Stewart, S. A.: Paleomagnetic analysis of fold kinematics and implications for geological models
1417 of the Cantabrian/Asturian arc, north Spain, *J. Geophys. Res. Solid Earth*, 100(B10), 20079–
1418 20094, doi:10.1029/95JB01482, 1995.
- 1419 Tait, J. A., Bachtadse, V. and Soffel, H. C.: Eastern Variscan fold belt : Paleomagnetic evidence
1420 for oroclinal bending, *Geology*, 24, 871–874, doi:10.1130/0091-7613(1996)024<0871, 1996.
- 1421 Tauxe, L.: *Essentials of Paleomagnetism*, Univ of California Press., 2010.
- 1422 Thomas, W. A.: Evolution of Appalachian-Ouachita Salients and Recesses from Reentrants and
1423 Promontories in the Continental Margin, *Am. J. Sci.*, 277(10), 1233–1278,
1424 doi:10.2475/ajs.277.10.1233, 1977.
- 1425 Thomas, W. A.: Genetic relationship of rift-stage crustal structure, terrane accretion, and
1426 foreland tectonics along the southern Appalachian-Ouachita orogen, *J. Geodyn.*, 37(3), 549–
1427 563, doi:10.1016/j.jog.2004.02.020, 2004.
- 1428 Tohver, E., Weil, A. B. B. B., Solum, J. G. G. G. and Hall, C. M. M. M.: Direct dating of
1429 carbonate remagnetization by $^{40}\text{Ar}/^{39}\text{Ar}$ analysis of the smectite–illite transformation, *Earth*
1430 *Planet. Sci. Lett.*, 274(3–4), 524–530, doi:10.1016/j.epsl.2008.08.002, 2008.
- 1431 Toit, A. L. Du: *Our wandering continents: an hypothesis of continental drifting*, Oliver and Boyd.
1432 [online] Available from: <http://books.google.es/books?id=iDZEAAAIAAJ>, 1937.
- 1433 Valladares, M. I., Barba, P., Ugidos, J. M., Colmenero, J. R. and Armenteros, I.: Upper

- 1434 Neoproterozoic-Lower Cambrian sedimentary successions in the Central Iberian Zone (Spain):
 1435 sequence stratigraphy, petrology and chemostratigraphy. Implications for other European
 1436 zones, *Int. J. Earth Sci.*, 89(1), 2–20, 2000.
- 1437 Van der Voo, R.: Paleomagnetism, oroclines, and growth of the continental crust, *GSA Today*,
 1438 14(12), 4–9, doi:10.1130/1052-5173(2004), 2004.
- 1439 Vergés, J.: Estudio del Complejo volcánico-sedimentario del Devónico y de la estructura de la
 1440 terminación oriental del sinclinal de Almadén (Ciudad Real), in *Libro Jubilar JM Rios. Tomo 3*,
 1441 pp. 215–229., 1983.
- 1442 Vissers, R. L. M., van Hinsbergen, D. J. J., van der Meer, D. G. and Spakman, W.: Cretaceous
 1443 slab break-off in the Pyrenees: Iberian plate kinematics in paleomagnetic and mantle reference
 1444 frames, *Gondwana Res.*, 34, 49–59, doi:10.1016/J.GR.2016.03.006, 2016.
- 1445 van der Voo, R., Stamatakis, J. A. and Pares, J. M.: Kinematic constraints on thrust-belt
 1446 curvature from syndeformational magnetizations in the Lagos del Valle Syncline in the
 1447 Cantabrian Arc, Spain, *J. Geophys. Res.-Solid Earth*, 102(B5), 10105–10119, 1997.
- 1448 Weil, A., Pastor-Galán, D., Johnston, S. T. and Gutiérrez-Alonso, G.: Late/Post Variscan
 1449 Orocline Formation and Widespread Magmatism, in *The Geology of Iberia: A Geodynamic*
 1450 *Approach*, edited by C. Quesada and J. T. Oliveira, pp. 527–542, Springer International
 1451 Publishing, Cham., 2019.
- 1452 Weil, A. B.: Kinematics of orocline tightening in the core of an arc: Paleomagnetic analysis of
 1453 the Ponga Unit, Cantabrian Arc, northern Spain, *Tectonics*, 25(3), n/a-n/a,
 1454 doi:10.1029/2005TC001861, 2006.
- 1455 Weil, A. B. and Sussman, A. J.: Classifying curved orogens based on timing relationships
 1456 between structural development and vertical-axis rotations, vol. 383, edited by A. J. Sussman
 1457 and A. B. Weil, pp. 1–16, Geological Society of America., 2004.
- 1458 Weil, A. B., Van der Voo, R., van der Pluijm, B. A. and Parés, J. M.: The formation of an orocline
 1459 by multiphase deformation: a paleomagnetic investigation of the Cantabria–Asturias Arc
 1460 (northern Spain), *J. Struct. Geol.*, 22(6), 735–756, doi:10.1016/S0191-8141(99)00188-1, 2000.
- 1461 Weil, A. B., van der Voo, R. and van der Pluijm, B. A.: Oroclinal bending and evidence against
 1462 the Pangea megashear: The Cantabria-Asturias arc (northern Spain), *Geology*, 29(11), 991–
 1463 994, 2001.
- 1464 Weil, A. B., Van der Voo, R. and Voo, R. V. D.: Insights into the mechanism for orogen-related
 1465 carbonate remagnetization from growth of authigenic Fe-oxide: A scanning electron microscopy
 1466 and rock magnetic study of Devonian carbonates from northern Spain, *J. Geophys. Res.-Solid*
 1467 *Earth*, 107(B4), 14–14, 2002.
- 1468 Weil, A. B., Gutiérrez-Alonso, G. and Conan, J.: New time constraints on lithospheric-scale
 1469 oroclinal bending of the Ibero-Armorican Arc: a palaeomagnetic study of earliest Permian rocks
 1470 from Iberia, *J. Geol. Soc.*, 167(1), 127–143, doi:10.1144/0016-76492009-002, 2010.
- 1471 Weil, A.B., Yonkee, A., and Sussman, A.: Reconstructing the kinematics of thrust sheet rotation:
 1472 a paleomagnetic study of Triassic redbeds from the Wyoming Salient, U.S.A., *GSA Bulletin*,

- 1473 122, ½ 2-23, 2010.
- 1474 Weil, A. B. B., Gutiérrez-Alonso, G., Johnston, S. T. and Pastor-Galán, D.: Kinematic
1475 constraints on buckling a lithospheric-scale orocline along the northern margin of Gondwana: A
1476 geologic synthesis, *Tectonophysics*, 582, 25–49, doi:10.1016/j.tecto.2012.10.006, 2013.
- 1477 Woodcock, N. H., Soper, N. J. and Strachan, R. A.: A Rheic cause for the Acadian deformation
1478 in Europe, *J. Geol. Soc.*, 164, 1023–1036, 2007.
- 1479 Yenes, M., Alvarez, F. and Gutierrez-Alonso, G.: Granite emplacement in orogenic
1480 compressional conditions: the La Alberca-Bejar granitic area (Spanish Central System, Variscan
1481 Iberian Belt), *J. Struct. Geol.*, 21(10), 1419–1440, 1999.
- 1482 Yonkee, A., Weil, A.B., and Sussman, A.: Reconstructing the kinematic evolution of curved
1483 mountain belts: internal strain patterns in the Wyoming Salient, Sevier thrust belt, U.S.A., *GSA*
1484 *Bulletin*, 122, ½, 24-50, 2010.
- 1485 Yonkee, A. and Weil, A. B.: Quantifying vertical axis rotation in curved orogens: Correlating
1486 multiple data sets with a refined weighted least squares strike test, *Tectonics*, 29(3), 31 PP.-31
1487 PP., doi:10.1029/2008TC002312, 2010b.
- 1488 Zegers, T. E., Dekkers, M. J. and Bailly, S.: Late Carboniferous to Permian remagnetization of
1489 Devonian limestones in the Ardennes: Role of temperature, fluids, and deformation, *J. Geophys.*
1490 *Res. Solid Earth*, 108(B7), n/a-n/a, doi:10.1029/2002JB002213, 2003.

1491 Captions

1492 Fig. 1 Simplified paleogeographic map of the Variscan-Alleghanian orogeny prior to the
1493 Jurassic break-up of Pangea, with the major orogenic curves labeled. Note, this map represents
1494 Iberian outcrops without taking into account post-Jurassic Alpine deformation (see text for
1495 details e.g. Gong et al., 2008; Pastor-Galán et al., 2018). Slightly darker colors in the Variscan
1496 belt indicate present-day European and African outcrops (modified after Martínez-Catalán et al.,
1497 2009; Weil et al., 2013).

1498 Fig. 2 A) Present-day configuration of the tectonostratigraphic zones of the Iberian
1499 Variscides and their major structures. White areas represent Mesozoic and Cenozoic cover. B)
1500 Three competing geometric proposals for the Central Iberian curve. 1) A disharmonic curvature,
1501 up to 160° at the outer arc but much less pronounced at the inner arc (Aerden, 2004); 2) A
1502 harmonic, but more open curvature as suggested by Martínez Catalán (2012); 3) An isoclinal
1503 curvature model (Shaw et al., 2012). C) Distribution of the E1 (migmatitic domes) and C3 to
1504 post-C3 granitoids in the NW of Iberia (modified from Pastor-Galán et al., 2016)

1505 Fig. 3 A) Stratigraphic synthesis of the Gondwanan platform series in NW Iberia.
1506 Cantabrian Zone columns are after Aramburu et al., 2002; Bastida, 2004; Murphy et al 2008;
1507 Pastor-Galán et al., 2013a; 2013b. Iberian Range follows Gozalo et al., 2008; Mergl and

Zamora, 2012 and Calvín and Casas, 2014. West Asturian-Leonese Zone stratigraphy is after Pérez-Estaún, 1990; Marcos, 2004; Martínez-Catalán et al., 2004a; Gutiérrez-Marco et al., 2019. Central Iberian Zone follows Díez-Balda, 1986; Valladares et al., 2000; Díez Montes, 2007; Martínez-Catalán et al., 2004b; del Moral and Sarmiento, 2008; García-Arias et al., 2018. B) Ordovician paleocurrents orientations, modified from Shaw et al. (2012). C) Schematic basin architecture inferred from the stratigraphic compilation.

Fig. 4 A) The kinematic evolution of the Cantabrian Orocline in its core, the Cantabrian Zone, inferred from total least squares (TLS) orocline tests (Pastor-Galán et al. 2017). B) Three orocline (strike) tests used to constrain the kinematics of the Cantabrian Orocline. The Ordovician paleocurrents, which predate any orogenic movement, recorded the complete vertical-axis rotation history and yields a slope (m) of ~ 1 . The Moscovian paleomagnetic data (from Weil et al., 2013; Pmag.), which postdates the main orogenic phases (C1, C2 and E1) and is coeval with C3, shows a slope of ~ 1 . The Gzhelian joint sets (from Pastor-Galán et al., 2011) orocline test shows a slope of ~ 0.5 , which indicates that part of the orocline was already formed between 304 Ma and 300 Ma.

Fig. 5 A) Aeromagnetic map of Spain (Ardizzone et al., 1989, for Spain and the World Digital Magnetic Anomaly Map (WDMAM project) and Portugal (modified from Martínez Catalán, 2012 and Martínez Catalán et al., 2015), showing the possible trace of the Central Iberian curve. B) Bouguer anomalies of the Iberian Peninsula, modified from Ayala et al., 2016. Gravity anomalies do not reflect the geometry of the Cantabrian Orocline nor the Central Iberian curve, but are related to the Cenozoic Alpine lithospheric structure.

Fig. 6 Paleomagnetic studies related to the Cantabrian Orocline and the Central Iberian curve: (1) Synthesis of paleomagnetism in the core of the Cantabrian Orocline (see Weil et al., 2013); (2) Permian (eP) components synthesized in Weil et al. (2010); (3) Ordovician volcanics and limestones (Laquiana) in the boundary between the West Asturian-Leonese and Central Iberian Zones (Fernández-Lozano et al., 2016); (4) Devonian sedimentary sequences and Permian subvolcanics in the Iberian ranges (Pastor-Galán et al., 2018); (5) Permian dykes and sills (Calvín et al., 2014); (6) Anatectic granites (E1) and mantle derived granitoids (C3) from Tormes Dome and Central System (Pastor-Galán et al., 2016); (7) Cambrian limestones from Tamames (N) and los Navalucillos (S) (Pastor-Galán et al., 2015a); (8) Ediacaran-Early Cambrian sedimentary rocks in the southern sector of the Central Iberian Zone (Pastor-Galán et al., 2017b); (9) Almadén volcanics from the Central Iberian Zone (Perroud et al., 1991; Parés and van der Voo, 1992; Leite Mendes et al. in press) and Volcanic rocks from southern Ossa Morena and the South Portuguese Zone (Leite Mendes et al. in press)

Fig. 7 Magnetization components with a positive reversal test in the extensional anatectic granites of Tormes (A) and Martinamor Domes (B). This component is interpreted as primary with a magnetization age of >318 Ma (Pastor-Galán et al., 2016). C) Distribution of directions and VGPs and statistical parameters from both domes combined.

Fig. 8 Cartoon depicting the different vertical-axis rotation events that occurred in the Cantabrian Zone and the Iberian Range, modified from Pastor-Galán et al. (2018). (A) Original quasilinear Variscan Orogenic belt, B) Formation of the Cantabrian Orocline around the Carboniferous–Permian boundary after a ~70° counterclockwise rotation in the Southern branch of the Cantabrian Zone and the Iberian Range. This rotation matches the rotation for the Cantabrian Orocline, see the fit of the Iberian Range Component #2 in the orocline test for the Cantabrian Zone (below). C) Post-Permian (Cenozoic) rotation of ~22° clockwise (CW) likely produced by differential shortening during the Alpine orogeny (Izquierdo-Llavall et al., 2018). Below, the global magnetic polarity time-scale for the Pennsylvanian and Cisuralian (following Ogg et al., 2016). TLS = Total Least Squares. Note that once the 22° CW rotation in the Iberian Range is corrected, components #2, #1, and P fit as expected with the APWP for the southern limb of the Cantabrian Orocline (Pastor-Galán et al., 2016).

Fig. 9 Compilation of the directional distributions and average declinations with parachute of confidence (Δ Declination) in sites around the Central Iberian curve (see Fig. 6). The results show general CCW rotations in contrast to the expected CW rotations if the Central Iberia curve formed by vertical-axis rotations (see text). Results are compared with the expected declinations if those sites were part of the Cantabrian Orocline following the methodology described in Pastor-Galán et al. (2017b).


Fig. 10 Orocline test of the Cantabrian Orocline (Weil et al., 2013) compared with the magnetizations found in the adjacent Laurussian segments of the orogen: Ireland (Pastor-Galán et al., 2015b) and the South Portuguese Zone (Leite Mendes et al., in press)

Fig. 11 Pioneering hypotheses for the Central Iberian curve. Note that none of them fulfill the most recent geometric and kinematic criteria. A) Simplified ribbon continent model after Johnston et al. (2013) and Shaw and Johnston (2016). B) Dextral mega-shear model from Martínez-Catalán (2011). C) Kinematic model with indentation and left-lateral shearing after Simancas et al. (2013)

Fig. 12 Preliminary kinematic proposal for the Iberian Variscides. A) Pre-collisional stage after the opening of the Galicia Tras-os-Montes restricted seaway (e.g. Pin et al., 2002; Gutiérrez-Alonso et al., 2008a; Arenas et al., 2016). The irregular shape of the margin and the

1576 younging westwards deformation front (e.g. Daleyer et al., 1997) resulted in tectonic escape
1577 towards the still open Rheic Ocean (e.g. Braid et al., 2011; Murphy et al., 2016). B) After closure
1578 of the Rheic Ocean, C1 and C2 structures formed. The Galicia Tras-os-Montes was emplaced
1579 orogen parallel (e.g. Martínez-Catalán et al., 1990; Dias da Silva et al., 2020), preserving the
1580 shape of the seaway (i.e. a primary arc). C) The gravitational collapse of the orogen produced
1581 widespread anatexis and fold interference in the hinterland and the emplacement of the foreland
1582 fold-and-thrust belt. D) At Pennsylvanian times a change in the far-field stress buckled the
1583 Variscan belt around a vertical axis (see Gutiérrez-Alonso et al., 2008; Weil et al., 2013; Pastor-
1584 Galán et al., 2015a for details), creating new interference patterns and a lithospheric-scale
1585 response (see Gutiérrez-Alonso et al., 2004, 2011a; Pastor-Galán et al., 2012a). E) When the
1586 orocline became too tight to keep rotating, new cross-cutting brittle structures (C4) formed and
1587 minor extensional collapse (E2) occurred (e.g. Fernández-Lozano et al., 2019; Dias da Silva et
1588 al., 2020).

1589 Fig. 1
 1590 Simplified
 1591 paleogeographic
 1592 map of the Variscan-
 1593 Alleghanian orogeny
 1594 prior to the Jurassic
 1595 break-up of Pangea,
 1596 with the major
 1597 orogenic curves
 1598 labeled. Note, this
 1599 map represents
 1600 Iberian outcrops
 1601 without taking into
 1602 account post-
 1603 Jurassic Alpine
 1604 deformation (see text
 1605 for details e.g. Gong et al., 2008; Pastor-Galán et al., 2018). Slightly darker colors in the
 1606 Variscan belt indicate present-day European and African outcrops (modified after Martínez-
 1607 Catalán et al., 2009; Weil et al., 2013).







Fig. 2 A) Present-day configuration of the tectonostratigraphic zones of the Iberian Variscides and their major structures. White areas represent Mesozoic and Cenozoic cover. B) Three competing geometric proposals for the Central Iberian curve. 1) A disharmonic curvature, up to 160° at the outer arc but much less pronounced at the inner arc (Aerden, 2004); 2) A harmonic, but more open curvature as suggested by Martínez Catalán (2012); 3) An isoclinal curvature model (Shaw et al., 2012). C) Distribution of the E1 (migmatitic domes) and C3 to post-C3 granitoids in the NW of Iberia (modified from Pastor-Galán et al., 2016)

Fig. 3 A) Stratigraphic synthesis of the Gondwanan platform series in NW Iberia. Cantabrian Zone columns are after Aramburu et al., 2002; Bastida, 2004; Murphy et al 2008; Pastor-Galán et al., 2013a; 2013b. Iberian Range follows Gozalo et al., 2008; Mergl and Zamora, 2012 and Calvín and Casas, 2014. West Asturian-Leonese Zone stratigraphy is after Pérez-Estaún, 1990; Marcos, 2004; Martínez-Catalán et al., 2004a; Gutiérrez-Marco et al., 2019. Central Iberian Zone follows Díez-Balda, 1986; Valladares et al., 2000; Díez Montes, 2007; Martínez-Catalán et al., 2004b; del Moral and Sarmiento, 2008; García-Arias et al., 2018. B) Ordovician paleocurrents orientations, modified from Shaw et al. (2012). C) Schematic basin architecture inferred from the stratigraphic compilation.



1627 Fig. 4 A) The kinematic evolution of the Cantabrian Orocline in its core, the Cantabrian
 1628 Zone, inferred from total least squares (TLS) orocline tests (Pastor-Galán et al. 2017). B) Three
 1629 orocline (strike) tests used to constrain the kinematics of the Cantabrian Orocline. The
 1630 Ordovician paleocurrents, which predate any orogenic movement, recorded the complete
 1631 vertical-axis rotation history and yields a slope (m) of ~1. The Moscovian paleomagnetic data
 1632 (from Weil et al., 2013; Pmag.), which postdates the main orogenic phases (C1, C2 and E1) and
 1633 is coeval with C3, shows a slope of ~1. The Gzhelian joint sets (from Pastor-Galán et al., 2011)
 1634 orocline test shows a slope of ~0.5, which indicates that part of the orocline was already formed
 1635 between 304 Ma and 300 Ma.






Fig. 5 A) Aeromagnetic map of Spain (Ardizzone et al., 1989, for Spain and the World Digital Magnetic Anomaly Map (WDMAM project) and Portugal (modified from Martínez Catalán, 2012 and Martínez Catalán et al., 2015), showing the possible trace of the Central Iberian curve. B) Bouguer anomalies of the Iberian Peninsula, modified from Ayala et al., 2016. Gravity anomalies do not reflect the geometry of the Cantabrian Orocline nor the Central Iberian curve, but are related to the Cenozoic Alpine lithospheric structure.



1644 Fig. 6 Paleomagnetic studies related to the Cantabrian Orocline and the Central Iberian
 1645 curve: (1) Synthesis of paleomagnetism in the core of the Cantabrian Orocline (see Weil et al.,
 1646 2013); (2) Permian (eP) components synthesized in Weil et al. (2010); (3) Ordovician volcanics
 1647 and limestones (Laquiana) in the boundary between the West Asturian-Leonese and Central
 1648 Iberian Zones (Fernández-Lozano et al., 2016); (4) Devonian sedimentary sequences and
 1649 Permian subvolcanics in the Iberian ranges (Pastor-Galán et al., 2018); (5) Permian dykes and
 1650 sills (Calvín et al, 2014); (5) Anatectic granites (E1) and mantle derived granitoids (C3) from
 1651 Tormes Dome and Central System (Pastor-Galán et al., 2016); (6) Cambrian limestones from
 1652 Tamames (N) and los Navalucillos (S) (Pastor-Galán et al., 2015a); (7) Ediacaran-Early
 1653 Cambrian sedimentary rocks in the southern sector of the Central Iberian Zone (Pastor-Galán et
 1654 al., 2017b); (8) Almadén volcanics from the Central Iberian Zone (Perroud et al., 1991; Parés
 1655 and van der Voo, 1992; Leite Mendes et al. in press) and Volcanic rocks from southern Ossa
 1656 Morena and the South Portuguese Zone (Leite Mendes et al, in press)



1657
1658
1659
1660
1661

Fig. 7 Magnetization components with a positive reversal test in the extensional anatectic granites of Tormes (A) and Martinamor Domes (B). This component is interpreted as primary with a magnetization age of >318 Ma (Pastor-Galán et al., 2016). C) Distribution of directions and VGPs and statistical parameters from both domes combined.




Fig. 8 Cartoon depicting the different vertical-axis rotation events that occurred in the Cantabrian Zone and the Iberian Range, modified from Pastor-Galán et al. (2018). (A) Original quasilinear Variscan Orogenic belt, B) Formation of the Cantabrian Orocline around the Carboniferous–Permian boundary after a $\sim 70^\circ$ counterclockwise rotation in the Southern branch of the Cantabrian Zone and the Iberian Range. This rotation matches the rotation for the Cantabrian Orocline, see the fit of the Iberian Range Component #2 in the orocline test for the Cantabrian Zone (below). C) Post-Permian (Cenozoic) rotation of $\sim 22^\circ$ clockwise (CW) likely produced by differential shortening during the Alpine orogeny (Izquierdo-Llavall et al., 2018). Below, the global magnetic polarity time-scale for the Pennsylvanian and Cisuralian (following Ogg et al., 2016). TLS = Total Least Squares. Note that once the 22° CW rotation in the Iberian Range is corrected, components #2, #1, and P fit as expected with the APWP for the southern limb of the Cantabrian Orocline (Pastor-Galán et al., 2016).




Fig. 9 Compilation of the directional distributions and average declinations with parachute of confidence (Δ Declination) in sites around the Central Iberian curve (see Fig. 6). The results show general CCW rotations in contrast to the expected CW rotations if the Central Iberia curve formed by vertical-axis rotations (see text). Results are compared with the expected declinations if those sites were part of the Cantabrian Orocline following the methodology described in Pastor-Galán et al. (2017b).





Fig. 10 Orocline test of the Cantabrian Orocline (Weil et al., 2013) compared with the magnetizations found in the adjacent Laurussian segments of the orogen: Ireland (Pastor-Galán et al., 2015b) and the South Portuguese Zone (Leite Mendes et al., in press)



1688 Fig. 11 Pioneering hypotheses for the Central Iberian curve. Note that none of them fulfill
 1689 the most recent geometric and kinematic criteria. A) Simplified ribbon continent model after
 1690 Johnston et al. (2013) and Shaw and Johnston (2016). B) Dextral mega-shear model from
 1691 Martínez-Catalán (2011). C) Kinematic model with indentation and left-lateral shearing after
 1692 Simancas et al. (2013)






Fig. 12 Preliminary kinematic proposal for the Iberian Variscides. A) Pre-collisional stage after the opening of the Galicia Tras-os-Montes restricted seaway (e.g. Pin et al., 2002; Gutiérrez-Alonso et al., 2008a; Arenas et al., 2016). The irregular shape of the margin and the younging westwards deformation front (e.g. Daleyer et al., 1997) resulted in tectonic escape towards the still open Rheic Ocean (e.g. Braid et al., 2011; Murphy et al., 2016). B) After closure of the Rheic Ocean, C1 and C2 structures formed. The Galicia Tras-os-Montes was emplaced orogen parallel (e.g. Martínez-Catalán et al., 1990; Dias da Silva et al., 2020), preserving the shape of the seaway (i.e. a primary arc). C) The gravitational collapse of the orogen produced widespread anatexis and fold interference in the hinterland and the emplacement of the foreland fold-and-thrust belt. D) At Pennsylvanian times a change in the far-field stress buckled the Variscan belt around a vertical axis (see Gutiérrez-Alonso et al., 2008; Weil et al., 2013; Pastor-Galán et al., 2015a for details), creating new interference patterns and a lithospheric-scale response (see Gutiérrez-Alonso et al., 2004, 2011a; Pastor-Galán et al., 2012a). E) When the orocline became too tight to keep rotating, new cross-cutting brittle structures (C4) formed and minor extensional collapse (E2) occurred (e.g. Fernández-Lozano et al., 2019; Dias da Silva et al., 2020).


A) Pre C1 (~380 Ma)




B) C1-C2 (~360-335 Ma)




C) E1 (~330-317 Ma)



D) C3 (~315-300 Ma)



E) C4-E2 (~300-280 Ma)



- Cantabrian Zone
- West Asturian-Leonese
- Central Iberian
- Ossa-Morena
- South Portuguese-Zone
- Galicia Tras-Os-Montes and other Oceanic domains
- E1/E2 gneiss domes & granites
- C3-C4 mantle derived granitoids
- C3 Dextral shear zones
- C3 Sinistral shear zones
- C1/C2 Sinistral shear zones
- Folds trend
- E1/C3 folds Central Iberian
- C1/C2 folds Central Iberian
- ... Zone boundary
- ↗ Transport direction