

1 **Whole-rock and zircon evidence for evolution of the Late**
2 **Jurassic high Sr/Y Zhoujiapuzi granite, Liaodong Peninsula,**
3 **North China Craton**

4
5 **Renyu Zeng**^{a, b, c *}, **Mark B. Allen**^c, **Xiancheng Mao**^d, **Jianqing Lai**^d, **Jie Yan**^{a, b},
6 **Jianjun Wan**^{a, b}

7 ^a State Key Laboratory of Nuclear Resources and Environment, East China University
8 of Technology, Nanchang, 330013, Jiangxi, China

9 ^b School of Earth Sciences, East China University of Technology, Nanchang, 330013,
10 China

11 ^c Department of Earth Sciences, Durham University, Durham DH1 3LE, UK

12 ^d School of Geosciences and Info-Physics, Central South University, Changsha 410083,
13 China

14 **Correspondence:** Renyu Zeng (zengrenyu@126.com)

15

16 **Abstract:** Middle-Late Jurassic high Sr/Y granitic intrusions are extensively exposed
17 in the Liaodong Peninsula, in the eastern part of the North China Craton (NCC).
18 However, the genesis of the high Sr/Y signature in these intrusions has not been studied
19 in detail. In this study, we report results of zircon U-Pb dating, Hf isotopic analysis and
20 zircon and whole-rock geochemical data for the Late Jurassic Zhoujiapuzi granite in
21 the middle part of the Liaodong Peninsula. The Zhoujiapuzi granite is high-K (calc-
22 alkaline) and peraluminous in nature, with high SiO₂ (68.1–73.0 wt %) and Al₂O₃

23 (14.5–16.8 wt %), low in TFe_2O_3 (1.10–2.49 wt %) and MgO (0.10–0.44 wt %), and
24 with high Sr/Y (19.9–102.0) and $\text{La}_\text{N}/\text{Yb}_\text{N}$ (14.59–80.40), characteristic of high Sr/Y I-
25 type granite. The geochemical signatures, in combination with the presence of a large
26 number of Paleoproterozoic inherited zircons, indicate that the Zhoujiapuzi granite was
27 most likely derived from partial melting of the basement in the region, and specifically
28 the Liaoji granites. The high Sr/Y signature is inherited from these source rocks. LA-
29 ICP-MS zircon U-Pb dating of the autocryst zircons from two samples (from different
30 localities) yielded consistent weighted average ages of 160.7 ± 1.1 Ma (MSWD=1.3) and
31 159.6 ± 1.1 Ma (MSWD=1.2), with $\epsilon\text{Hf}(t)$ values in the range of -26.6– -22.8.
32 Morphological and chemical studies on autocrystic zircon grains show that there are
33 two stages of zircon growth, interpreted as magmatic evolution in two distinct stages.
34 The light-CL core reflects a crystallization environment of low oxygen fugacity and
35 high $T_{\text{Zr-Ti}}$; the dark-CL rim formed with high oxygen fugacity and lower $T_{\text{Zr-Ti}}$. Based
36 on the geochemical features and regional geological data, we propose that the Liaodong
37 Peninsula in the Late Jurassic was part of a mature continental arc, with extensive
38 melting of thick crust above the Paleo-Pacific subduction zone.

39 **Keywords:** Liaodong Peninsula; Late Jurassic; Zircon U-Pb-Hf isotopes; Two stages
40 of crystal growth; High Sr/Y granite

41 **1. Introduction**

42 The Liaodong Peninsula is located in the northeast of the North China Craton
43 (NCC). The northeast NCC was influenced by three main tectonic regimes in the
44 Mesozoic, related to the subduction of the Paleo-Asian, Paleo-Pacific and Mongol-

45 Okhotsk oceans (Tang et al., 2018). The superposition of these different regimes
46 resulted in changing tectonic and magmatic patterns over time. Middle-Late Jurassic
47 granitic rocks are extensively exposed in the northern parts of the Liaodong Peninsula,
48 such as the Yutun mylonitic granite, Xiaoheishan granodiorite, Heigou monzogranite
49 (Wu et al., 2005), Wulong two-mica monzogranite (Yang et al., 2018), and Huangdi
50 biotite monzogranite (Xue et al., 2020). Most of these rocks are characterized by high
51 Sr /Y, and plot within the adakite field on Sr/Y-Y and La_N/Yb_N - Yb_N diagrams (Wu et
52 al., 2005a; Yang et al., 2015a, 2018).

53 The geodynamic settings and petrogenesis of adakite and geochemically similar
54 high Sr/Y igneous rocks have been widely discussed. The high Sr/Y rocks were
55 originally proposed to be formed by melting of young (<25 Ma) and hot subducted
56 oceanic slab in an arc setting (Defant and Drummond, 1990). However, later studies
57 have shown that the high Sr/Y rocks can form in both arc and non-arc settings by other
58 processes, such as continental interior settings (Wang et al., 2007), cold subduction
59 zones (Nakamura and Iwamori, 2013), collision or post-collision processes (Schwartz
60 et al., 2011). In addition, numerous studies have suggested that the lower continental
61 crust can also be the source of the high Sr/Y rocks (Gao et al., 2004; Ou et al., 2017).
62 However, it is debated whether crustal thickening is necessary for their formation (e.g.
63 Moyen, 2009; Kamei et al., 2009; Zhan et al., 2020). In recent years, some studies have
64 proposed that the high Sr/Y ratio in granitic rocks can be inherited from a high Sr/Y
65 crust source, regardless of pressure (Kamei et al., 2009; Ma et al., 2015; Zhan et al.,
66 2020).

67 The Middle-Late Jurassic granitic rocks in the Liaodong Peninsula are commonly
68 proposed to be the products of partial melting of thickened mafic crust with garnet in
69 the residue (Wu et al., 2005a; Yang et al., 2015a, 2018; Tang et al., 2018). However, the

70 source composition has not been fully considered in the petrogenesis of the high Sr/Y
71 rocks in the Liaodong Peninsula. Hence, the petrogenesis of the Middle-Late Jurassic
72 high Sr/Y rocks needs to be re-evaluated, based on more detailed work and a
73 consideration of possible sources. This petrogenesis is of significance for understanding
74 the Jurassic tectonics of the Liaodong Peninsula, and the NCC in general.

75 In this paper, we examined the high Sr/Y Zhoujiapuzi granite from the Xiuyan area,
76 in the middle of the Liaodong Peninsula. Zircons are analysed for U-Pb-Hf isotopes
77 and trace element geochemistry, and by Raman spectroscopy. These results are
78 integrated with whole-rock geochemistry. We focus on the zircons, because of their
79 potential to reveal the origins of the pluton (Belousova et al., 2002; Wang et al., 2007;
80 Breiter et al., 2014; Zhao et al., 2014), and so provide a case study for the evolution of
81 plutonic magma systems in general. Based on observations of the CL images and
82 chemical analysis, two zircon growth stages can be distinguished. We first determine
83 the crystallization environments of the two zircon growth stages, and then decipher the
84 petrogenesis, source characteristics and origin of the high Sr/Y signature of the pluton
85 as a whole. Integrated with previous studies, our study provides insights into the
86 tectonic evolution of the Liaodong Peninsula in the Late Jurassic.

87 **2. Geological setting**

88 The Zhoujiapuzi granite is located in the middle of the Liaodong Peninsula, at the
89 northeastern margin of the NCC (Fig. 1). The Paleoproterozoic Liaohe Group and Liaoji
90 granite are the basement in the study area. The Liaohe Group includes the Lieryu,
91 Gaojiayu, Dashiqiao and Gaixian formations. Although stratigraphic terms are used,
92 these rocks are metamorphic, and the group consists of leptynite, leptite, granulite,

93 amphibolite, marble and phyllite. The protoliths of the Liaohe Group include marine
94 volcanics, clastics, carbonates and claystones. The formation age of the
95 metasedimentary rocks in the Liaohe Group is 2.0–1.9 Ga (Wan et al., 2006; Li et al.,
96 2015). It is in unconformable contact with the overlying strata of the Mesoproterozoic
97 Cuocaogou Formation and Xiaoling Formation.

98 The study area experienced strong magmatic activity in the Paleoproterozoic,
99 which can be divided into two stages of 2.2–2.1 Ga and ~ 1.85 Ga. The 2.18–2.14 Ga
100 Liaoji granites (also called gneissic granites), which lie within an area measuring 300
101 km × 70 km, are dominated by A- and I-type granites (Li and Zhao, 2007; Yang et al.,
102 2016; Wang et al., 2020a). Metamorphosed volcanic rocks (leptynite, leptite and
103 granulite) in the Liaohe Group also formed at 2.2–2.1 Ga (Li et al., 2015). The ~1.85
104 Ga granites mainly consist of I- and S-type porphyry granites and alkaline syenites
105 (Yang et al., 2007; Yang et al., 2015b). In addition, there were small amounts of mafic
106 magmatic activity at ~2.17 Ga, ~2.1 Ga and ~1.8 Ga (Meng et al., 2014; Yuan et al.,
107 2015). There are a variety of viewpoints on the Paleoproterozoic tectono-magmatic
108 evolution in the Liaodong Peninsula, such as an intracontinental rift opening-closing
109 model (Li et al., 2005) and an arc-continent collision model (Faure et al., 2004).

110 In the Mesozoic, the region of the Liaodong Peninsula was influenced by the
111 circum-Pacific tectonic regime, the Mongol-Okhotsk tectonic regime and the Paleo-
112 Asian Ocean tectonic regime. The joint influence of multiple tectonic regimes resulted
113 in intensive magmatism during the Mesozoic (Fig. 1b). These Mesozoic magmatic
114 rocks can be divided into three stages, namely: Triassic (233–212 Ma), Jurassic (180–

115 156 Ma) and Early Cretaceous (131–117 Ma) (Wu et al., 2005b).

116 The Triassic magmatic rocks are less exposed, mainly alkaline rocks, diabase,
117 diorites and granites (Wu et al., 2005b). Among them, the granites mainly have A-type
118 affinity, and may have formed in an extensional setting (Tang et al., 2018; Wang et al.,
119 2019). Magmatism has been related to either the subduction of the Paleo-Pacific slab,
120 closure of the Paleo-Asian Ocean, or the collision between the NCC and the Yangtze
121 Craton (Tang et al., 2018; Wang et al., 2019). The majority of the Jurassic magmatic
122 rocks are monzogranite and granodiorite, which are generally calc-alkaline I-type
123 granites, and show characteristics of adakite-like rocks. Some of them, exposed near
124 later extensional structures, have undergone regional ductile deformation. These
125 Jurassic magmatic rocks are generally considered to relate to the subduction of the
126 Paleo-Pacific slab (Wu et al., 2005a; Zhai et al., 2004). In the Early Cretaceous, basic-
127 acidic-alkaline rocks were widely developed. Among them, the granites have mainly
128 A- and I-type affinities. These rocks are generally considered to have formed in an
129 intense extensional environment, which is connected with either the rollback or low-
130 angle subduction of the Paleo-Pacific slab (Wu et al., 2005c; Zheng et al., 2018).

131 **3. Samples and petrography**

132 The Zhoujiapuzi granite is located to the east of Xiuyan City, in the middle of the
133 Liaodong Peninsula (Fig. 1b). It intruded into the Lieryu Formation of the Liaohe Group.
134 Eight samples of the Zhoujiapuzi granite were collected at locations shown in Fig. 1c.

135 The Zhoujiapuzi granite is generally grey in colour and with fine-grained texture

136 (Fig. 2a). The mineral assemblage contains K-feldspar (~50 %), quartz (~25 %),
137 plagioclase (~20 %) and biotite (~5 %) as well as accessory minerals such as zircon,
138 ilmenite, magnetite and apatite. K-feldspar grains are euhedral or subhedral, and always
139 exhibit cross-hatched twinning (Fig. 2b). Quartz grains are usually xenomorphic, and
140 have indented boundaries and wavy extinction (Fig. 2b-d). Plagioclase always exhibits
141 polysynthetic twinning and have sericitization in places (Fig. 2c). Biotite mainly fills
142 in the interstices between the other minerals (Fig. 2c, d).

143 **4. Analytical methods**

144 The cathodoluminescence (CL) images of zircon were obtained by the Chengpu
145 geological Testing Co. Ltd, Langfang, China using the TIMA analysis. The LA-ICP-
146 MS zircon U-Pb analyses were performed using an Agilent Technologies 7700x ICP-
147 MS with a Teledyne Cetac Technologies Analyte Excite laser-ablation system at
148 Nanjing FocuMS Contract Testing Co. Ltd. The analyses were carried out with a 35 μm
149 spot size at 8 Hz repetition rate for 40 seconds. The ICP-MS detector has dual modes:
150 pulse for lower signal, and analog for higher signal. Pulse-analog cross calibration was
151 performed before the measurement of U-Pb isotopes, delivering a wider linear dynamic
152 range – up to 10 orders of magnitude. For a signal of ^{238}U higher than 1.2–1.4 Mio cps,
153 equivalent zircon contains U concentrations higher than 600 ppm, and are measured in
154 analog mode. 91500 was used as external standard. GJ-1 (600Ma, Jackson et al., 2004)
155 and Plešovice (337Ma, Sláma et al., 2008) were treated as quality control for
156 geochronology. During our analyses, the weighted mean age of GJ-1 and Plešovice
157 were 606.0 ± 4.8 Ma ($n=16$, MSWD = 0.50) and 340.9 ± 4.0 Ma ($n=7$, MSWD = 1.0),
158 respectively. Trace elements abundance of zircon were externally calibrated against

159 NIST SRM 610 with Si as the internal standard. The raw ICP-MS data were processed
160 using ICPMSDataCal software (Liu et al., 2010). No common-Pb correction was
161 applied to the data. Data reduction was completed using the Isoplot4.15 (Ludwig, 2003).
162 The instrument description and analytical procedure are described in detail by Zeng et
163 al. (2018).

164 The in-situ Lu-Hf isotopic analyses of zircon were performed by LA-MC-ICP-MS
165 using a Teledyne Cetac laser-ablation system and a Nu Plasma II MC-ICP-MS at
166 Nanjing FocuMS Contract Testing Co. Ltd. The 193 nm ArF excimer laser was focused
167 on zircon surface with fluence of 6.0J/cm². The ablation protocol employed a spot
168 diameter of 50 um at 8 Hz repetition rate for 40 seconds. Three standard zircons, GJ-1,
169 91500, and Penglai, were analysed for quality control at every ten unknown samples.
170 In the experiment, standard zircon GJ-1, 91500, and Penglai were analyzed, and the
171 ¹⁷⁶Hf/¹⁷⁷Hf ratios were 0.282002–0.282013, 0.282305–0.282315 and 0.282901–
172 0.282914 respectively, in accordance with their recommended values (GJ-1: 0.282012,
173 Yuan et al., 2008; 91500: 0.282307 ± 0.000031, Wu et al., 2006; Penglai: 0.282906 ±
174 0.000010, Li et al., 2010). For the calculation of εHf(t) values, we have adopted the
175 ¹⁷⁶Lu decay constant of 1.867 × 10⁻¹¹ (Söderlund et al., 2004), the present-day
176 chondritic values of ¹⁷⁶Lu/¹⁷⁷Hf = 0.0332 and ¹⁷⁶Hf/¹⁷⁷Hf = 0.282772 (Blichert-Toft and
177 Albarède 1997). To calculate one-stage model ages (T_{DM1}) relative to a depleted-mantle
178 source, we have adopted the present-day depleted-mantle values of ¹⁷⁶Lu/¹⁷⁷Hf =
179 0.0384 and ¹⁷⁶Hf/¹⁷⁷Hf = 0.28325 (Vervoort and Blichert-Toft 1999). To calculate two-
180 stage modal ages (TDM2), ‘felsic crust’ model ages are calculated using average
181 continental crust ¹⁷⁶Lu/¹⁷⁷Hf = 0.015 (Griffin et al., 2004)

182 Zircon Raman analyses were carried out using an RM2000 laser Raman
183 spectrometer at the State Key Laboratory of Nuclear Resources and Environment, East
184 China University of Technology. The selected incident wavelengths were 532 and 785
185 nm in order to clearly identify the luminescence bands due to low concentration
186 impurities. The beam power was 20 mW. The Leica 50× objective was employed.

187 Six fresh rock samples were selected for geochemical analysis. The elemental
188 analyses were conducted at Analytical Chemistry & Testing Services (ALS) Chemex
189 (Guangzhou) Ltd. Major oxides were analyzed using wave-dispersive X-ray
190 fluorescence (XRF) (ME-XRF26). Analytical precision was better than $\pm 0.01\%$.
191 Trace element abundances were measured by the lithium borate dissolution method and
192 ICP-MS (ME-MS81). The analytical uncertainties of the rare earth element (REE) and
193 high field strength element (HFSE) are $<5\%$. Analytical uncertainties are in the range
194 of 5%–10% for the other elements. Detailed analytical procedures refer to Zhang et al.
195 (2019) and Nash et al. (2020).

196 **5. Analytical results**

197 The data for major and trace elements, Raman microprobe data, zircon trace
198 elements, zircon U-Pb ages, and zircon Hf isotopes are shown in Tables S1, S2, S3, S4
199 and S5, respectively.

200 **5.1. Whole-rock major and trace element compositions**

201 SiO_2 contents range from 68.11 wt.% to 73.02 wt.% (average 71.71 wt.%).
202 Contents of Na_2O and K_2O are 3.81 – 4.65 wt.% and 4.32 – 4.71 wt.%, respectively,

203 with $\text{Na}_2\text{O}/\text{K}_2\text{O}$ ratio of 0.82 – 1.08 and total alkalis ($\text{Na}_2\text{O} + \text{K}_2\text{O}$) of 8.38 – 8.97. All
204 samples plot in the field of granite in the TAS classification except one (Fig. 3a). These
205 samples have Al_2O_3 contents of 14.49 – 16.83 wt.% (average 15.09 wt.%), CaO
206 contents of 1.04 – 1.98 wt.% (average 1.38 wt.%) and A/CNK values of 1.05 – 1.10
207 (average 1.07). In the A/NK – A/CNK diagram (Fig. 3b), all samples plot in the
208 peraluminous field (Fig. 3b). The granite samples have low TFe_2O_3 ($\text{TFe}_2\text{O}_3 = \text{all Fe}$
209 calculated as Fe_2O_3) contents and MgO contents ranging from 1.10–2.49 wt % and
210 0.10–0.44 wt %, respectively, with Mg# ($\text{Mg}\# = 100 * \text{molar Mg}/(\text{Mg} + \text{Fe})$) values of 15–
211 26.

212 The samples of the Zhoujiapuzi granite exhibit variable REEs, with total REEs
213 ranging from 59 to 302 ppm. The $\text{La}_\text{N}/\text{Yb}_\text{N}$ values of the Zhoujiapuzi granite range from
214 14.59 to 80.40 (average 38.27), showing right-declined REE patterns (Fig. 4a). The
215 samples have Eu/Eu^* of 0.62–1.94 and Ce/Ce^* of 0.94–1.16. In the primitive mantle-
216 normalized trace element diagram (Fig. 4b), the samples show negative anomalies of
217 HFSEs (e.g., Nb, Ta, Ti and P) and positive anomalies of La and LILEs (e.g., K, Rb,
218 Ba, U, La, Ce). The Zhoujiapuzi granite is characterized by high contents of Sr (309–
219 551 ppm) and low contents of Y (5.01–15.5 ppm) and Yb (0.43–1.40 ppm), with high
220 Sr/Y ratios of 19.94–102.04 (average 65.50).

221 **5.2 Zircon CL images, Raman spectra and REE elements**

222 CL images of zircons from the Zhoujiapuzi granite are shown in Fig. 5. Zircons
223 commonly have crystal sizes between 150 and 250 μm , and have length/width ratios of
224 2:1–4:1, with euhedral, stubby to elongate prisms. According to the CL images, most
225 zircons show an internal division into 2 distinct domains: light-CL core and dark-CL

226 rim. The light-CL core is characterized by bright CL intensity and widely-spaced
227 oscillatory zoning patterns. The dark-CL rim is overgrown continuously by the light-
228 CL core and is characterized by extremely low CL emission and narrowly-spaced
229 oscillatory zoning patterns. In addition, some zircons have inherited cores, which have
230 corroded and rounded shapes in contact with the light-CL core, such as 1# and 37# in
231 XY-001 and 6# and 41# in XY-008 (Fig. 5). These inherited zircons have oscillatory
232 zoning in CL images.

233 Six light-CL core spots and six dark-CL rim spots were analyzed for Raman
234 spectra. The light-CL cores have antisymmetric stretching vibration (B_{1g}) of the SiO_4
235 tetrahedra (ν_3 (SiO_4)) Raman band of 1005–1007 cm^{-1} and half-width of the ν_3 (SiO_4)
236 Raman band (b) values of 6.0–8.1 cm^{-1} , while the dark-CL rims have ν_3 (SiO_4) Raman
237 band of 1004–1007 cm^{-1} and b values of 5.4–9.0 cm^{-1} .

238 Twenty light-CL core spots, eighteen dark-CL rim spots and six inherited zircon
239 spots were analyzed for trace and rare earth elements. The light-CL core spots have
240 lower U content (28–677 ppm) than the dark-CL rim spots (U=641–3842 ppm). In the
241 chondrite-normalized REE element diagram (Fig. 6a, b), both the light-CL core and
242 dark-CL rim are characterized by HREE enrichment relative to LREE with positive Ce
243 anomalies and negative Eu anomalies. The light-CL core spots have ΣREE of 49–1115
244 ppm (average 390 ppm), $\Sigma LREE$ of 3–72 ppm (average 14 ppm) and $\Sigma HREE$ of 46–
245 1100 ppm (average 377 ppm), whereas the dark-CL rim spots have ΣREE of 327–1632
246 ppm (average 895 ppm), $\Sigma LREE$ of 2–14 ppm (average 6 ppm) and $\Sigma HREE$ of 325–
247 1627 ppm (average 889 ppm). Hence, the REE content of the light-CL core is
248 significantly lower than that of the dark-CL rim, and the difference between the two is
249 mainly in HREE content. The light-CL core spots have Eu/Eu^* of 0.07–0.60 (average
250 0.28) and Ce/Ce^* of 1.89–24.27 (average 10.03). Because the contents of La and Pr are

251 typically present very low, Ce* in this study is obtained by the formulation $(Nd_N)^2 / Sm_N$
252 (Loader et al., 2017). The dark-CL rim spots have Eu/Eu* of 0.08–0.24 (average 0.13)
253 and Ce/Ce* of 6.57–200.31 (average 79.23). These results indicate that the light-CL
254 core have a weaker negative Eu anomaly and a weaker positive Ce anomaly than those
255 of the dark-CL rim. The inherited zircon spots have ΣREE of 602–1517 ppm, and show
256 depletion of LREE, enrichment of HREE, a positive Ce anomaly (Ce/Ce* of 1.52–
257 216.08) and a negative Eu anomaly (Eu/Eu* of 0.07–0.13) (Fig. 6c).

258 **5.3 Zircon U–Pb and Hf isotope composition**

259 Seventy-seven spots were analysed for U–Pb isotope composition from samples
260 XY-001 and XY-008. In the U–Pb Concordia diagram (Fig. 7a, c), both the light-CL
261 core and dark-CL rim spots overlap within uncertainty on the Concordia curve. There
262 is a large degree of overlap between the 29 spots of dark-CL rim and 32 spots of light-
263 CL core in terms of $^{206}Pb/^{238}U$ age although the average value for $^{206}Pb/^{238}U$ age is
264 higher in the 32 spots of light-CL core (Fig. 7e). On a single zircon, the $^{206}Pb/^{238}U$ age
265 of the light-CL core is older than that of the dark-CL rim (Fig. 5), but the two values
266 are within the error range of the in-situ LA-ICP-MS analyses (individual spot of ± 3 –5%
267 relative precision, Schmitz and Kuiper, 2013). In sample XY-001, 33 spots define a
268 weighted mean $^{206}Pb/^{238}U$ age of 160.7 ± 1.1 Ma (2σ , MSWD=1.3; Fig. 7b). In sample
269 XY-008, 28 spots define a weighted mean $^{206}Pb/^{238}U$ age of 159.6 ± 1.1 Ma (2σ ,
270 MSWD=1.2; Fig. 7d). The other 10 spots with distinctly older ages ($^{207}Pb/^{206}Pb$ ages
271 ranging from 2500 to 2173 Ma) were obtained on inherited cores. Their ages are
272 discordant, suggesting that these inherited cores were variably influenced by lead loss.
273 Among these, 9 spots define a discordia line with an upper intercept age of 2163 ± 13
274 Ma (MSWD=0.45) (Fig. 7f).

275 Twenty-four zircons were analyzed for Lu-Hf isotope composition. The variation
276 in Hf isotopic data is limited, between 9 spots from light-CL core and 9 spots from dark-
277 CL rim. 18 spots exhibit a range of $^{176}\text{Hf}/^{177}\text{Hf}$ ratios from 0.281921 to 0.282030, which
278 converts to $\epsilon\text{Hf}(t)$ values between -26.6 to -22.8 (Fig. 8), and two-stage Hf model ($T_{\text{DM}2}$)
279 ages of 2650 to 2889 Ma by using the U-Pb age for each zircon. Six analytical spots,
280 which define the Concordia upper intercept age of 2163 Ma, show $^{176}\text{Hf}/^{177}\text{Hf}$ ratios
281 and $\epsilon\text{Hf}(t)$ values of 0.281443 to 0.281496 and -0.7 to 1.5, respectively, with $T_{\text{DM}2}$ age
282 of 2648 Ma to 2791 Ma by using the upper intercept age.

283 **6. Discussion**

284 **6.1 Significance of the two stages of zircon**

285 Generally, zircon with high U content can easily break down into the metamict
286 state because of the radiation damage to the lattice caused by α -particles originating
287 from the decay of uranium (Mezger and Krogstad, 1997). The physical and structural
288 changes often lead to the loss of Pb and addition of trace elements such as LREE. In
289 this study, the dark-CL rim spots have high U content, which is significantly higher than
290 the median value of zircon U content in granitic magma (350 ppm, Wang et al., 2011).
291 Hence, the metamictization degree of the zircons must be taken into consideration. Data
292 from dark-CL rim spots plot on the Concordia curve, indicating no obvious Pb loss. The
293 internal structure of dark-CL rim is relatively intact, with obvious oscillatory zoning,
294 and few cracks, implying that the physical and structural of the dark-CL rim remained
295 unchanged. Nasdala et al. (1998) suggested that the metamictization of zircon can be
296 well characterized by Raman spectroscopy. The half-width of the $\nu_3(\text{SiO}_4)$ Raman band

297 (b) of 10 cm^{-1} and 20 cm^{-1} are proposed to approximately distinguish well-crystallized,
298 intermediate and metamict zircons (Nasdala et al., 1998). The dark-CL rim have b
299 values of 5.4–9.2, characterizing them as well-crystallized. Therefore, the above
300 features indicate that the dark-CL rim are not metamict. Consequently, it can be
301 concluded that the U-Pb isotope and trace element systematics of the dark-CL rim have
302 not been changed by metamictization.

303 Both the light-CL core and dark-CL rim have oscillatory zoning patterns, and their
304 chondrite-normalized REE patterns are characterized by steeply positive slopes from
305 the LREE to HREE with strong negative Eu anomalies and pronounced positive Ce
306 anomalies. The above characteristics are consistent with those of igneous zircon
307 (Hoskin and Schaltegger, 2003). Although hydrothermal zircon can also have
308 oscillatory zoning patterns similar to magmatic zircons, there are obvious differences
309 in trace elements between the magmatic and hydrothermal zircon (Hoskin et al., 2005).
310 In the discrimination diagram (Fig. 9), both the spots of light-CL core and dark-CL rim
311 fall in or near the magmatic field, which is obviously different from hydrothermal
312 zircon. Hence, the above characteristics indicate that both the light-CL core and dark-
313 CL rim have a magmatic origin.

314 The light-CL core was overgrown continuously by the dark-CL rim. In addition,
315 the contact between the light-CL core and dark-CL rim is euhedral. Such core-mantle
316 overgrowth relationships indicate that the light-CL core domains are not inherited
317 zircons. The similar Hf isotopic data of the light-CL core and dark-CL rim is also
318 consistent with this interpretation. For the age population, the samples of XY-001 and

319 XY-008 have MSWD of 1.3 and 1.2, respectively, which are both within the expected
320 range for 95 % confidence interval (Mahon, 1996). Although the $^{206}\text{Pb}/^{238}\text{U}$ age of dark-
321 CL rim is generally younger than that of light-CL core, the ages of these two distinct
322 domains have the characteristics of continuous variation, and do not show two or more
323 distinct age populations (Fig. 7b, d). These phenomena do not support the presence of
324 antecrystic zircons (Siégel et al., 2018). Hence, both the light-CL core and dark-CL rim
325 are most likely autocrystic zircon formed in one distinct pulse of magma. The weighted
326 mean U-Pb ages of 160.7 ± 1.1 Ma and 159.6 ± 1.1 Ma can be interpreted as the
327 emplacement age of the Zhoujiapuzi granite. The obvious difference in internal
328 structure and trace element composition between the light-CL core and dark-CL rim
329 could be due to significant changes in their crystallization environments (Wang et al.,
330 2007).

331 The Zr/Hf ratio in zircon has a negative correlation with the degree of fractionation
332 in the parent melt (Claiborne et al., 2006). In this study, the Zr/Hf ratios of the dark-CL
333 rim (21–40) are obviously lower than those of the light-CL core (39–56) (Fig. 10a). In
334 addition, incompatible elements such as U and REE will become enriched in the highly
335 evolved magma (Zhao et al., 2014). In this study, the contents of U and REE of dark-
336 CL rim are significantly higher than those of light-CL core (Fig. 10a). Overall, the
337 above features reflect that the dark-CL rim crystallized from a later and more evolved
338 magma.

339 Watson and Harrison (2005) found that the Ti content of zircon has a strong
340 dependence on temperature (T), and obtained a Ti-in-zircon thermometer ($T_{\text{Zr-Ti}}$). Since

341 then, Ferry and Watson (2007) suggested that the solubility of Ti in zircon depends not
342 only on T and activity of TiO₂ (aTiO₂) but also on the activity of SiO₂ (aSiO₂), and
343 revised the T_{Zr-Ti}. We use the T_{Zr-Ti} from Ferry and Watson (2007) and the recommended
344 values (aSiO₂=1, aTiO₂ = 0.5) for the activity of SiO₂ and TiO₂ (Schiller and Finger,
345 2019), due to the presence of ilmenite and quartz in the Zhoujiapuzi granite. The T_{Zr-Ti}
346 from the light-CL core and dark-CL rim are 684–830 °C (average 761 °C) and 509–
347 712°C (average 635 °C), respectively, i.e. the light-CL core formed at higher
348 temperatures than the dark-CL rim. The T_{Zr-Ti} value shows a significant positive
349 correlation with Zr/Hf (a tracer of fractional crystallisation), and shows continual
350 fractionation and cooling (Fig. 10b). As the light-CL core and dark-CL rim formed in
351 different magmatic evolution stages, it is problematic to use the same aSiO₂ and aTiO₂
352 values to calculate both T_{Zr-Ti} values for both. For ilmenite bearing granites, Schiller
353 and Finger (2019) suggested that the variation of aTiO₂ values corresponding to
354 different zircon crystallization stages is small. In addition, Schiller and Finger (2019)
355 showed that the aSiO₂ value of the ilmenite-bearing granites at the onset of magmatic
356 zircon crystallization was more than 0.75. Even if the aSiO₂ value of the light-CL core
357 is changed from 1.0 to 0.75, the temperature will only drop by about ~27 °C, which is
358 significantly lower than the 126 °C difference between the average T_{Zr-Ti} value of the
359 light-CL core and dark-CL rim. Therefore, it is certain that the light-CL core formed at
360 higher temperatures than the dark-CL rim, although we cannot calculate the specific
361 temperature difference.

362 Cerium exists in magmas as both Ce³⁺ and Ce⁴⁺. Because the 0.84-Å radius of the

363 Zr^{4+} ion is more closely matched by the Ce^{4+} (0.97-Å radius) than the Ce^{3+} (1.143-Å
364 radius) (all ionic radii are from Shannon, 1976), Ce^{4+} is more compatible in the zircon
365 structure than the Ce^{3+} . Hence, the magnitude of Ce anomaly is a useful tool for
366 evaluating the oxygen fugacity condition of crystallization environment (e.g. Ballard et
367 al., 2002; Trail et al., 2012). Loader et al. (2017) suggested that the Ce/Ce* ratio is
368 likely to be the most robust measure of magma redox conditions, although it is only a
369 semi-quantitative measure. In this study, the Ce/Ce* ratio of the light-CL core and dark-
370 CL rim are 6.30–153.36 (average 32.51) and 21.81–5773.06 (average 787.39),
371 respectively. This result suggests that the dark-CL rim formed in a higher oxygen
372 fugacity environment than the light-CL core. As shown in the Ce/Ce*-Zr/Hf diagram
373 (Fig. 10c), Ce/Ce* has a significant negative correlation with Zr/Hf, showing that the
374 oxygen fugacity condition is increasing with the evolution of magma.

375 The absence of enclaves and disequilibrium textures in the Zhoujiapuzi granite
376 and uniform $\epsilon Hf(t)$ values of the light-CL core and dark-CL rim do not support magma
377 mixing and wall-rock assimilation. Consequently, the abrupt change between the
378 crystallization environment of the light-CL core and dark-CL rim is not due to the
379 magma mixing or contamination during magma evolution. Therefore, we propose that
380 the light-CL core was formed in a relatively deep magma chamber, which had low
381 oxygen fugacity, low Zr saturation and higher temperature. The low Th, U and REE,
382 and widely-spaced oscillatory zoning patterns indicate a low growth rate of zircon
383 (Hoskin and Schaltegger, 2003; Wang et al., 2011). In contrast, the dark-CL rim was
384 formed during the ascent and/or at the emplacement location of the magma. At this

385 stage, the oxygen fugacity significantly increased, the temperature decreased, and Zr
386 saturation increased due to the crystallization differentiation. In this environment, the
387 crystallization rate of zircon significantly increased, forming the zircons with a higher
388 content of Th, U and REE elements, low CL emission and narrowly-spaced oscillatory
389 zoning patterns.

390 Zircon U-Pb dating is the most commonly used method in geochronology,
391 especially dating the emplacement age of magmatic rocks. A weighted mean age or
392 upper intercept age is usually obtained to represent the emplacement time of a magmatic
393 rock. However, the autocrystic zircons in this study record two different magmatic
394 evolution stages. Previous studies, such as Wang et al. (2007), Zhao et al. (2014) and
395 Chen et al. (2020), also show that zircons can crystallize continually or intermittently
396 in a single phase of magmatism, showing several growth zones of clearly different
397 internal structure and distinct time difference. Therefore, autocrystic zircon can be
398 formed in two or more evolution stages during one distinct pulse or increment of
399 magma. Some scholars even regard that the age difference of different stages can be
400 more than dozens of Ma (Wang et al., 2007). Therefore, if the zircon ages in the same
401 magmatic rock have a large range of variation, this could be caused by the zircons
402 recording different stages in magmatic evolution, related to different levels of magma
403 within the crust and/or different temperature regimes. In this paper, although the
404 apparent age of the dark-CL rim is generally younger than that of the light-CL core, the
405 age difference between the two is within the error range of the in-situ LA-ICP-MS
406 analyses (individual spot of $\pm 3\text{--}5\%$ relative precision). Therefore, further work is

407 needed to verify the actual age difference between the two magmatic evolution stages.
408 Nevertheless, it is notable that the bulk petrology and geochemistry of the host pluton
409 does not record and reveal this two-stage magmatic evolution, which can only be
410 detected in the zircon analysis.

411 **6.2 Genetic type: I-type affinity**

412 The Zhoujiapuzi granite has low Zr (113 - 242 ppm), Ce (26.5 - 121.5 ppm),
413 Zr+Nb+Ce+Y (152.0 - 382.6 ppm), (Na₂O + K₂O)/CaO (4.53 - 8.31) and FeO*/MgO
414 (5.09 - 10.56), distinct from the typical A-type granites (Fig. 11a-d). Furthermore, the
415 Zhoujiapuzi granite does not contain mafic alkaline minerals, such as arfvedsonite,
416 riebeckite, etc., which is also distinctly inconsistent with typical A-type granites (Wu et
417 al., 2003). Wu et al. (2017) suggested that a high formation temperature is one of the
418 most important characteristics of A-type granite. Zircon saturation thermometry (T_{Zrn})
419 and Ti-in-zircon thermometer (T_{Zr-Ti}) are two methods for estimating magma
420 temperatures. As noted above, because the values of a_{SiO_2} and a_{TiO_2} during the early
421 zircon crystallization cannot be accurately obtained, the temperature of this period
422 cannot be accurately obtained through the Ti-in-zircon thermometer. Zircon saturation
423 thermometry was introduced by Watson and Harrison (1983) and is suitable for non-
424 peralkaline crustal source rocks. Since the zircon solubility is mainly affected by
425 temperature, major element compositions have a limited impact on calculated T_{Zrn}
426 (Miller et al., 2003). In addition, the errors introduced by crystal-rich composition tend
427 to cancel as changes in Zr concentration and M value during crystallization have

428 opposite effects on the T_{Zrn} value (Miller et al., 2003). Therefore, the composition of
429 Zhoujiapuzi granite can be used to estimate the magma temperature. The calculated T_{Zrn}
430 values for the Zhoujiapuzi granite are in the range of 803-870 °C (mean=845 ±20°C).
431 It was proposed that the T_{Zrn} suggests an upper limit on the temperature of melt
432 generation for inheritance-rich granitoid (Miller et al., 2003). Hence, the magma
433 temperature of the Zhoujiapuzi granite should be lower than or equal to the T_{Zrn} value,
434 which is significantly lower than that of typical A-type granite (>900 °C, Skjerlie and
435 Johnston, 1992; Douce, 1997). Thus, the Zhoujiapuzi granite is not an A-type granite.

436 The samples of the Zhoujiapuzi granite have $A/KNC < 1.1$, relatively high Na_2O
437 (3.96–4.65 wt.%) and lack peraluminous minerals (e.g. cordierite, andalusite,
438 muscovite and garnet), which are clearly different from S-type granites (Chappell and
439 White, 1992). With the rise of the degree of crystallization, P_2O_5 contents
440 (generally>0.1 wt.%) increase in S-type granites, accompanied by an
441 increase/immaturity in SiO_2 (Wolf and London, 1994). However, the Zhoujiapuzi
442 granite samples have low P_2O_5 contents (0.02 - 0.08 wt.%), and decrease with
443 increasing SiO_2 (Fig. 11e), which are features consistent with the I-type granite rather
444 than S-type granite (Chappell and White, 1992). Additionally, Rb has a positive
445 correlation with Y (Fig. 11f), which has been considered as an indicator of I-type granite
446 (Jiang et al., 2018). Furthermore, the composition of the Zhoujiapuzi granite fall in the
447 I-type granite field in the discrimination diagrams of granites introduced by Collins et
448 al. (1982) (Fig. 11 c-d). Therefore, we conclude that the Zhoujiapuzi granite is a I-type
449 granite.

450 **6.3 Petrogenesis of the high Sr/Y granite**

451 The samples of the Zhoujiapuzi granite have high Sr/Y and (La/Yb)_N ratios and
452 low Y and Yb contents (Fig. 12a) consistent with the geochemical signatures of modern
453 adakites (Defant and Drummond, 1990). However, other geochemical parameters of
454 the Zhoujiapuzi granite, such as the high K₂O/Na₂O ratio (0.93 –1.22), low Al₂O₃
455 content (14.49 –15.02%, except one) and Sr content (in half of the samples lower than
456 400 ppm), are obviously different from typical adakites (K₂O/Na₂O ≤0.42, Al₂O₃ ≥
457 15 %, Sr >400 ppm, Defant and Drummond., 1990; Drummond et al., 1996, Martin et
458 al., 2005). A variety of petrogenetic models have been proposed for the origin of high
459 Sr/Y magmatic rocks, such as partial melting of subducting oceanic crust (Model A,
460 Defant and Drummond, 1990), delaminated lower continental crust (LCC) (Model B,
461 Kay and Kay, 1993; Xu et al., 2002), differentiation of basaltic arc magma (Model C,
462 Castillo et al., 1999), magma mixing between mantle-derived mafic and crust-derived
463 silicic magmas (Model D, Ma et al., 2013a), partial melting of thickened basaltic LCC
464 (Model E, Gao et al., 2004; Ou et al., 2017), or melting of a high Sr/Y (and La/Yb)
465 source (Model F, Kamei et al., 2009; Ma et al., 2015).

466 **6.3.1 Model A: Partial melting of subducting oceanic crust**

467 The partial melting of the young, hot and hydrated subducted oceanic slab in the
468 garnet stability field is the classical formation model of adakite (high Sr/Y rock) (Defant
469 and Drummond, 1990). Studies have shown that the rock with this genetic model
470 generally has the characteristics of high mantle components (such as MgO, CaO and
471 Cr) because of the involvement of mantle magma (Wang et al., 2018). However, this
472 phenomenon was not seen in the Zhoujiapuzi granite. In addition, the Zhoujiapuzi
473 granite has high K₂O/Na₂O ratios (0.92–1.22, average 1.13), which is inconsistent with

474 the slab-derived adakites ($K_2O/Na_2O = \sim 0.4$, Martin et al., 2005). Moreover, the low
475 $\epsilon_{Hf}(t)$ values (-26.6 to -22.8) of the Zhoujiapuzi granite are also inconsistent with the
476 magmas derived from the partial melting of oceanic crust, which generally have
477 depleted isotopic character (Zhan et al., 2020). Furthermore, the Zhoujiapuzi granite
478 has low Ti/Eu and high Nd/Sm ratios (Fig. 13a), and markedly negative Nb-Ta
479 anomalies (Fig. 4b), which are distinct from those of oceanic basalts (Yu et al., 2012).
480 In summary, the Zhoujiapuzi granite is difficult to explain by Model A.

481 **6.3.2 Model B: Delaminated lower continental crust (LCC)**

482 High-density, garnet-bearing mafic lower crust delaminating or foundering into
483 the asthenosphere mantle and subsequent interaction with mantle peridotite could
484 produce high Sr/Y magmas (Kay and Kay 1993). Because the melt formed by partial
485 melting of the delaminated lower crust would interact with mantle peridotite during
486 magma ascent, the high Sr/Y magmas related to this petrogenetic model generally have
487 high MgO, Mg# and TiO₂ (Gao et al., 2004; Ou et al., 2017; He et al., 2021). The MgO
488 (0.10– 0.44 wt.%), Mg# (15– 26) and TiO₂ (0.09– 0.34 wt.%) values of Zhoujiapuzi
489 granite are significantly lower than the above values (Fig. 13b- d). In addition,
490 delamination of the lower crust generally occurs in within-plate extensional settings
491 (Gao et al., 2004), and will form a large number of Mg-rich (Mg#>50) rocks due to the
492 partial melting of lithospheric mantle and/or upwelling of asthenosphere (Ou et al.,
493 2017). However, these Jurassic magmatic rocks in the Liaodong Peninsula are generally
494 considered to be formed in a compressional environment related to the subduction of
495 the Paleo-Pacific slab (Li et al., 2004; Yang et al., 2015a; Zhu and Xu, 2019; Zheng et
496 al., 2018). Furthermore, the middle-late Jurassic granites are generally Mg-poor (Fig.
497 13c). Due to the high temperature of the asthenosphere (1200 °C, Parsons and

498 McKenzie, 1978; King et al., 2015), rocks formed by partial melting of the delaminated
499 lower crust should possess a high-temperature fingerprint. T_{Zm} has been used as a
500 geothermometer to estimate partial melting temperatures (e.g., Miller et al., 2003;
501 Collins et al., 2016). As mentioned before, the T_{Zm} of the Zhoujiapuzi granite is below
502 900 °C, which is markedly lower than the temperature of the asthenosphere. Therefore,
503 the petrogenetic model of delaminated lower continental crust (Model B) is also
504 inconsistent with the Zhoujiapuzi granite.

505 **6.3.3 Model C: Differentiation of basaltic arc magma**

506 Low-pressure fractional crystallization (involving olivine + clinopyroxene +
507 plagioclase + amphibole+ titanomagnetite) or high-pressure fractional crystallization
508 (involving garnet) from basaltic magmas have been proposed as two ways to generate
509 adakitic characteristics (Castillo et al., 1999; Macpherson et al., 2006).

510 However, the composition of the Zhoujiapuzi granite is relatively uniform,
511 including SiO₂, MgO and Na₂O, which does not support major fractional crystallization
512 (Xue et a., 2017). Furthermore, the Zhoujiapuzi granite has abundant inherited zircons
513 and no obvious depletion of Sr, Eu and Ba, showing that this granite has not experienced
514 extensive fractionation (Miller et al., 2003). The samples form clear partial melting
515 trends on the La/Yb versus La diagram (Fig. 13e), which also suggests that partial
516 melting was more important than fractional crystallization (Gao et al., 2007; Shahbazi
517 et al., 2021). In addition, crystal fractionation of basaltic melts can only form minor
518 volumes of granitic melts, the ratio of the two is about 9:1 (Zeng et al., 2016). However,
519 for the same age interval, no coexisting mafic-intermediate rocks have been found in
520 the research area. In the wider region of the Liaodong Peninsula, Middle-Late Jurassic
521 magmatism is dominated by felsic compositions; mafic- intermediate rocks are only

522 reported in the Huaziyu area (lamprophyre dikes, Jiang et al., 2005). Therefore, it is
523 unlikely that there are large-scale mafic- intermediate rocks contemporaneous with the
524 Zhoujiapuzi granite at depth according to the rock assemblage of Liaodong Peninsula
525 in this period. Moreover, the zircon Hf isotopic compositions of the Zhoujiapuzi granite
526 are quite different from those of the depleted mantle, but are similar to those of the
527 basement (Liaohu Group and Liaoji granite) in the study area (Fig. 8). The ancient
528 inherited zircons (2500 to 2173) with low $\epsilon_{\text{Hf}}(t)$ values also indicate older crustal
529 material in the Zhoujiapuzi granite. For these reasons, it is highly improbable that
530 Zhoujiapuzi granite was derived by differentiation of basaltic magma (Model C).

531 **6.3.4 Model D: Magma mixing between mantle-derived mafic and crust-derived** 532 **silicic magmas**

533 The Zhoujiapuzi granite has high $\text{K}_2\text{O}/\text{Na}_2\text{O}$ ratio (>1) and A/CNK value (>1),
534 together with the absence of mingling textures, mafic microgranular enclaves (MMEs),
535 felsic xenocrysts and melting texture of plagioclase, implying that the mantle-derived
536 magma is unlikely to have played an important role in the genesis of the Zhoujiapuzi
537 granite (Castro et al., 1991). In addition, the Zhoujiapuzi granite is characterized by the
538 development of biotite, but lacks amphibole and pyroxene. These features, coupled with
539 the high A/CNK value, are consistent with an origin as a crust-derived granitoid, but
540 obviously different from the granitoids formed by crust-mantle-derived magma mixing
541 (Barbarin, 1990). Moreover, granites formed by magma mixing generally have high
542 MgO, TFe_2O_3 , CaO and Cr contents and low SiO_2 content (Ma et al., 2013a; Wang et
543 al., 2018). These features are obviously inconsistent with the Zhoujiapuzi granite in this
544 study. Additionally, the $\epsilon_{\text{Hf}}(t)$ values and trace element composition of the two stages
545 of zircon also do not support magma mixing. Hence, magma mixing of mantle-derived

546 and crust-derived magmas (Model D) is also unlikely to have produced the Zhoujiapuzi
547 granite.

548 **6.3.5 Model E: Partial melting of thickened basaltic LCC**

549 Experimental studies have shown that the partial melt of basaltic LCC in the garnet
550 stabilization zone (> 40 km, i.e. ~1.2 GPa) can produce magma with a high Sr/Y ratio
551 (Rapp et al., 2003 and references therein). In these scenarios, high Sr/Y and overall
552 adakitic affinity are caused by leaving garnet as residual phases (e.g. Gao et al., 2004).
553 Based on geochemical data for the Zhoujiapuzi granites, partial melting of thickened
554 basaltic LCC is also unlikely to account for the high Sr/Y Zhoujiapuzi granite (Model
555 E). This conclusion is based on the following observations:

556 (1) This ratio of $(\text{Gd}/\text{Yb})_{\text{N}}$ is the most important feature to judge whether garnet is
557 involvement in magma genesis (Ma et al., 2012). If the $(\text{Gd}/\text{Yb})_{\text{N}}$ ratio of the source is
558 similar to the average value of the LCC (1.71, Rudnick and Gao, 2003), partial melting
559 of these crustal materials controlled by garnet at high pressure can produce melt with
560 $(\text{Gd}/\text{Yb})_{\text{N}}$ of 5.8 (Huang and He, 2010). In contrast, the $(\text{Gd}/\text{Yb})_{\text{N}}$ values (1.22–5.06,
561 average 2.69) of the Zhoujiapuzi granite are relatively low. (2) Studies of lower-crustal
562 xenoliths show that garnet may not be a common mineral in the lower crust of the NCC
563 (Ma et al., 2012). (3) As shown in the discrimination diagrams of granite sources (Fig.
564 13f, g), all samples fall in the range of metagreywacke-derived melts. Therefore, the
565 Zhoujiapuzi granite was considered to have been derived from crustal anatexis of
566 metagraywacke (or intermediate-acid igneous rock with similar mineral composition),
567 rather than basaltic lower crust.

568 **6.3.6 Model F: Melting of a high Sr/Y (and La/Yb) source**

569 Studies have shown that when a source rock has a high Sr/Y ratio, the high Sr/Y
570 signature of the derived magma can inherit from their source, regardless of pressure
571 (Kamei et al., 2009; Moyen, 2009; Ma et al., 2015). We suggest that partial melting of
572 high Sr/Y Liaoji granite was most probably the origin of the high Sr/Y Zhoujiapuzi
573 granite, as discussed below (Model F).

574 The Zhoujiapuzi granite has similar mineral assemblages (contains abundant K-
575 feldspar and lacks hornblende) and geochemical composition (Fig. 13h) to the
576 Tsutsugatake intrusion, which is explained by partial melting of arc-type tonalite or
577 adakitic granodiorite (Kamei et al., 2009). Among the inherited zircons from
578 Zhoujiapuzi granite, the $^{207}\text{Pb} / ^{206}\text{Pb}$ ages of all the spots are between 2132 and 2200
579 Ma, except one, and yield a Concordia upper intercept age of 2163 Ma. Both
580 assimilation of country-rocks and incomplete melting of source rocks can explain the
581 genesis of inherited zircon in granite. Due to the similar T_{DM2} of autocrystic zircons
582 (light-CL core and dark-CL rim) and inherited zircons, these inherited zircons most
583 likely come from the source of the Zhoujiapuzi granite. In the study area, meta-
584 sedimentary rocks and meta-volcanic rocks of the South Liaohe Group,
585 Paleoproterozoic mafic rocks, as well as the Liaoji granites, have ~2.16 Ga zircon. In
586 spite of an age peak of 2.17–2.16 Ga in detrital zircon age spectra of the metasediments
587 from the South Liaohe Group, melting of a sediment-dominated source is unlikely to
588 have occurred, as it would have also introduced other age peaks such as ~2.03 Ga and
589 ~2.50 Ga (Li et al., 2015; Wang et al., 2020b). In addition, given the I-type
590 characteristics of the Zhoujiapuzi granite, derivation from an igneous precursor is more
591 plausible rather than a metasedimentary origin (Chappell and White, 1992). Therefore,
592 these ~2.16 Ga zircons from the Zhoujiapuzi granite are unlikely to come from the

593 South Liaohe Group. As shown in the host rock discrimination diagrams (Fig. 14,
594 introduced by Belousova et al., 2002), all the ~2.16 Ga inherited zircons from
595 Zhoujiapuzi granite fall into the granitoid area (Fig. 14), precluding that these ~2.16 Ga
596 zircon come from the Paleoproterozoic mafic rocks. In addition, the ~2.16 Ga inherited
597 zircons from Zhoujiapuzi granite and the zircons from the Liaoji granites lie in a similar
598 area in the $\epsilon\text{Hf}(t)$ -age (Ma) diagram (Fig. 8). Hence, the ~2.16 Ga inherited zircon most
599 likely come from the Liaoji granites.

600 Some of the Liaoji granites, such as the Muniuhe granite (comprising granodiorite
601 and syenogranite with no distinct boundary between the two), have adakitic signatures,
602 and similar REE and trace element patterns as the Zhoujiapuzi granite (Fig. 4). Based
603 on a model of batch melting (Shaw, 1970) using the experiments of Conrad et al. (1988),
604 the high Sr/Y characteristic of the Zhoujiapuzi granite can be explained by partial
605 melting of Muniuhe granitic pluton leaving amphibole as the main residue (Fig. 12b).

606 In our modelling, we choose the XY-005 sample to approximately represent the
607 primitive melt composition. The reasons are as below: as mentioned above, the high
608 Sr/Y characteristics of the Zhoujiapuzi granite are not caused by the fractional
609 crystallization of amphibole. Furthermore, the lack of positive correlation between
610 $\text{D}_{\text{Yb}}/\text{Yb}_{\text{N}}$ ratios and $\text{La}_{\text{N}}/\text{Yb}_{\text{N}}$ ratios (Fig. 13i) also suggests that fractional
611 crystallization of amphibole was not a significant process for the Zhoujiapuzi granite.
612 On the other hand, the samples of Zhoujiapuzi granite display variable Eu and Sr
613 contents, implying that plagioclase is likely a fractional phase. The separation of
614 titanomagnetite could explain the positive trend in TFe_2O_3 with increasing TiO_2 content,
615 consistent with the occurrences of magnetite in some studied rocks. This possible
616 mineral assemblage of fractional crystallization is also reflected by the chemical
617 variations in the Sr/Y-Y diagram (Fig. 12b). Hence, the sample XY-005, which has

618 highest Sr/Y, was chosen to represent a primitive melt composition. To find the best
619 matching experimental melts, we have compared the major elements of the XY-005
620 sample with that of experimental melts and the characteristics of no garnet residue
621 discussed above. Results are shown in Fig. 12b. The Sr and Y compositions of the
622 starting material used in these experiments resemble those of the average composition
623 of the Muniuhe granitic pluton (Sr=475 ppm, Y=9.77 ppm), if the residue contains a
624 large volume of amphibole (>90 %). However, if more plagioclase is retained in the
625 residue (e.g. 18.3 %), a source region with a higher Sr content is required. Therefore, a
626 similar high Sr/Y Liaoji granite to the Muniuhe granitic pluton can produce the high
627 Sr/Y signatures of the Zhoujiapuzi granite.

628 A large number of Yanshanian adakites (or high Sr/Y rocks) are developed in the
629 NCC, which are generally considered to be derived from the thickened basaltic LCC
630 (e.g. Gao et al., 2004; Wu et al., 2005a; Ma et al., 2013b). Zhang et al. (2001, 2003)
631 suggested that these so-called “C-type adakites” indicated a large-scale crustal
632 thickening event. However, according to the studies on the Triassic and Jurassic adakitic
633 rocks near the Pingquan area, the northern part of the NCC, Ma et al. (2012, 2015)
634 suggested that the adakitic signatures of these rocks are inherited from their source
635 rocks. The research of the Zhoujiapuzi granite in this study also shows that among the
636 widely distributed Jurassic high Sr/Y granites in the Liaodong Peninsula, there is at
637 least one pluton with a high Sr/Y signature inherited from the source. Therefore, we
638 suggest that melting of a high Sr/Y (and La/Yb) source is one of the important processes
639 for the generation of Yanshanian high Sr/Y rocks in the NCC. This kind of high Sr/Y
640 granite does not need to be formed in the garnet stability field.

641 **6.4 Tectonic implications**

642 A large number of Early Jurassic arc-like igneous rocks occur in the northeast part
643 of NCC- Korean Peninsula-Hida belt, which belong to the middle-high K calc-alkaline
644 series and are characterized by enrichment in LILE and depletions in HFSE (Wu et al.,
645 2007; Tang et al., 2018 and references therein). In addition, the Early Jurassic
646 accretionary complexes in the eastern margin of the Eurasian continent and the Japan
647 islands, such as the Heilongjiang complex, the Khabarovsk complex and the Mino-
648 Tamba complex, are considered to be related to subduction (Wu et al., 2007; Tang et al.,
649 2018 and references therein). It is generally accepted that the Paleo-Pacific slab
650 subducted westwards in the Early Jurassic (Tang et al., 2018; Zhu and Xu, 2018).

651 In the middle-late Jurassic, I-type granites are dominant in the Liaodong Peninsula,
652 such as the Zhoujiapuzi granite (this study), Heigou pluton, Gaoliduntai pluton (Wu et
653 al., 2005a), Waling granite (Yang et al., 2015a) and Wulong granite (Yang et al., 2018).
654 There are not A-type granites, and mantle derived magmatism is extremely rare. These
655 granites were formed by partial melting of crustal materials without obvious
656 contribution of mantle derived magma (Wu et al., 2005a; Yang et al., 2015b, 2018; Xue
657 et al., 2020). In addition, WNW-ESE compression during 157-143 Ma was widespread
658 in the Liaodong Peninsula (Yang et al., 2004; Zhang et al., 2020). It not only
659 mylonitized the granite plutons in middle-lower crust levels, but also intensely
660 deformed the thick sedimentary cover in the upper crust (Qiu et al., 2018; Ren et al.,
661 2020). Hence, Late Jurassic magmatism in the Liaodong peninsula is most likely to be
662 related to subduction of the Paleo-Pacific plate in a mature continental arc, with crust

663 previously thickened by compressional tectonics, related to both the oceanic subduction
664 and the earlier Mesozoic collisions at the north and south margins of the NCC. This
665 setting would produce the conditions required for extensive crustal melting of pre-
666 existing basement. There is a potential resemblance to the modern arc of the Central
667 Andes (Allmendinger et al., 1997), where crustal thickening and plateau growth has
668 developed over the Cenozoic (Scott et al., 2018), and melting of older basement has
669 taken place during subduction of the Nazca plate (Miller and Harris, 1989). This model
670 is also consistent with the idea that much of eastern China was a high orogenic plateau
671 during the Mesozoic, before widespread Early Cretaceous extension and core complex
672 development (Meng, 2003; Chu et al., 2020).

673 **7. Conclusions**

674 (1) LA-ICP-MS zircon U-Pb dating indicates that the Zhoujiapuzi granite in the
675 Liaodong Peninsula formed at ~160 Ma.

676 (2) Zircon growth in Zhoujiapuzi granite can be divided into two distinct stages.
677 The light-CL core was formed in a deeper, hotter, magma chamber, which had low
678 oxygen fugacity and high temperature. The dark-CL rim formed from later, more
679 evolved, magma. Oxygen fugacity significantly increased and the temperature
680 decreased at this stage. The Zhoujiapuzi granite is a case study of multistage generation
681 and emplacement of magma, revealed by zircons, where no signals are discernible in
682 the bulk petrology or geochemistry.

683 (3) The I-type Zhoujiapuzi granite originated from partial melting of the

684 Paleoproterozoic Liaoji granites. The high Sr/Y compositions are inherited from their
685 source rocks, rather than being a direct indication of deep crustal melting, or any other
686 common mechanism for generating adakitic signatures.

687 (4) The Late Jurassic tectonic setting of the Liaodong Peninsula and the eastern
688 NCC resembled the modern orogenic plateau of the Central Andes, where silicic
689 magmatism may occur by partial melting of older continental crust in a compressional
690 environment, related to the subduction of the Paleo-Pacific plate.

691 **Acknowledgements**

692 We thank Dr. Wenzhou Xiao, Dr. Jiajie Chen and Dr. Quan Ou for constructive reviews
693 and useful suggestions. We are also grateful to Ying Liu, Chunying Guo, Jianxiong Hu,
694 Ziming Hu for their help with the field work. This research was funded by the National
695 Nature Science Foundation of China (Grants No. 42030809, 41772349, 41902075,
696 42002095; 42162013), China Scholarship Council (No. 202008360018), Geological
697 Exploration Program of China Nuclear Geology (Grant No. D1802), the research grants
698 from the East China University of Technology (Grant No. DHBK2017103), Open
699 Research Fund Program of State Key Laboratory of Nuclear Resources and
700 Environment (East China University of Technology) (Grant 2020NRE13).

701 **References**

702 Allmendinger, R. W., Jordan, T. E., Kay, S. M., and Isacks, B. L.: The evolution of the
703 Altiplano-Puna plateau of the Central Andes, *Annu. Rev. Earth Pl. Sc.*, 25, 139–
704 174, <https://doi.org/10.1146/annurev.earth.25.1.139>, 1997.

705 Altherr, R., Holl, A., Hegner, E., Langer, C. and Kreuzer, H.: High-potassium, calc-
706 alkaline I-type plutonism in the European Variscides: northern Vosges (France)
707 and northern Schwarzwald (Germany), *Lithos*, 50(1-3), 51–73,
708 [https://doi.org/10.1016/S0024-4937\(99\)00052-3](https://doi.org/10.1016/S0024-4937(99)00052-3), 2000.

709 Ballard, J. R., Palin, M. J., and Campbell, I. H.: Relative oxidation states of magmas
710 inferred from Ce(IV)/Ce(III) in zircon: application to porphyry copper deposits of
711 northern Chile. *Contrib. Mineral. Petrol.*, 144, 347–364,
712 <https://doi.org/10.1007/s00410-002-0402-5>, 2002.

713 Barbarin, B.: Granitoids: main petrogenetic classification in relation to origin and
714 tectonic setting, *Geol. J.*, 25, 227–238, 1990.

715 Belousova, E., Griffin, W., O'Reilly, S. Y., and Fisher, N.: Igneous zircon: trace
716 element composition as an indicator of source rock type, *Contrib. Mineral. Petrol.*,
717 143, 602–622, <https://doi.org/10.1007/s00410-002-0364-7>, 2002.

718 Blichert-Toft, J., and Albarède, F.: The Lu-Hf isotope geochemistry of chondrites and
719 the evolution of the mantle-crust system, *Earth. Planet. Sc. Lett.*, 148(1), 243–258,
720 [https://doi.org/10.1016/S0012-821X\(97\)00040-X](https://doi.org/10.1016/S0012-821X(97)00040-X), 1997.

721 Breiter, K., Lamarão, C. N., Borges, R. M. K., and Dall'Agnol, R.: Chemical
722 characteristics of zircon from A-type granites and comparison to zircon of S-type
723 granites, *Lithos*, 192–195, 208–225, <https://doi.org/10.1016/j.lithos.2014.02.004>,
724 2014.

725 Castillo, P. R., Janney, P. E., and Solidum, R. U.: Petrology and geochemistry of
726 Camiguin Island, southern Philippines: Insights to the source of adakites and other

727 lavas in a complex arc setting, *Contrib. Mineral. Petrol.*, 134(1), 33–51,
728 <https://doi.org/10.1007/s004100050467>, 1999.

729 Castro, A., Moreno-Ventas, I., and Rosa, J.: H-type (hybrid) granitoids: a proposed
730 revision of the granite-type classification and nomenclature, *Earth. Sci. Rev.*, 31,
731 237–253, [https://doi.org/10.1016/0012-8252\(91\)90020-G](https://doi.org/10.1016/0012-8252(91)90020-G), 1991.

732 Chappell, B. W., and White, A. J. R.: I-and S-type granites in the Lachlan Fold Belt,
733 *Geol. Soc. Am. Spec. Pap.*, 272, 1–26, <https://doi.org/10.1130/SPE272-p1>, 1992.

734 Chen, S. R., Wang, Q., Zhu, D. C., Weinberg, R. F., Zhang, L. L., and Zhao, Z. D.:
735 Reheating and magma mixing recorded by zircon and quartz from high-silica
736 rhyolite in the Coqen region, southern Tibet, *Am. Miner.*,
737 <https://doi.org/10.2138/am-2020-7426>, 2020.

738 Chu, Y., Lin, W., Faure, M., Allen, M. B., and Feng, Z.: Cretaceous exhumation of the
739 Triassic intracontinental Xuefengshan Belt: Delayed unroofing of an orogenic
740 plateau across the South China Block? *Tectonophysics*, 793, 228592,
741 <https://doi.org/1016/j.tecto.2020.228592>, 2020.

742 Claiborne, L. L., Miller, C. F., Walker, B. A., Wooden, J. L., Mazdab, F. K., and Bea,
743 F.: Tracking magmatic processes through Zr/Hf ratios in rocks and Hf and Ti
744 zoning in zircons: An example from the Spirit Mountain batholith, Nevada.
745 *Mineral. Mag.*, 70, 517–543, <https://doi.org/10.1180/0026461067050348>, 2006.

746 Collins, W. J., Beams, S. D., White, A. J. R., and Chappell, B. W.: Nature and origin
747 of A-type granites with particular reference to southeastern Australia, *Contrib.*
748 *Mineral. Petr.*, 80, 189-200, <https://doi.org/10.1007/BF00374895>, 1982.

749 Collins, W. J., Huang, H. Q., and Jiang, X. Y.: Water-fluxed crustal melting produces
750 Cordilleran batholiths, *Geology*, 44(2), 143–146,
751 <https://doi.org/10.1130/G37398.1>, 2016.

752 Conrad, W. K., Nicholls, I. A., and Wall, V. J.: Water-saturated and -undersaturated
753 melting of metaluminous and peraluminous crustal compositions at 10 kb:
754 evidence for the origin of silicic magmas in the Taupo Volcanic Zone, New
755 Zealand, and other occurrences, *J. Petrol.*, 29, 765–803,
756 <https://doi.org/10.1093/petrology/29.4.765>, 1988.

757 Defant, M. J., and Drummond, M. S.: Derivation of some modern arc magmas by
758 melting of young subducted lithosphere, *Nature*, 347, 662–665,
759 <https://doi.org/10.1038/347662a0>, 1990.

760 Douce, A. E. P.: Generation of metaluminous A-type granites by low-pressure melting
761 of calc-alkaline granitoids, *Geology*, 25, 743–746, [https://doi.org/10.1130/0091-
762 7613\(1997\)025<0743:GOMATG>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0743:GOMATG>2.3.CO;2), 1997.

763 Drummond, M. S., Defant, M. J., and Kepezhinskas, P. K.: Petrogenesis of slab-derived
764 trondhjemite–tonalite–dacite/adakite magmas, *Trans. R. Soc. Edinb.*, 87, 205–215,
765 <https://doi.org/10.1017/S0263593300006611>, 1996.

766 Faure, M., Lin, W., Moni, P., and Bruguier, O.: Paleoproterozoic arc magmatism and
767 collision in Liaodong Peninsula, NE China, *Terra. Nova.*, 16, 75–80,
768 <https://doi.org/10.1111/j.1365-3121.2004.00533.x>, 2004.

769 Ferry, J. M., and Watson, E. B.: New thermodynamic models and revised calibrations
770 for the Ti-in-zircon and Zr-in-rutile thermometers, *Contrib. Mineral. Petrol.*, 154,

771 429–437, <https://doi.org/10.1007/s00410-007-0201-0>, 2007.

772 Frost, B. R., Barnes, C. G., Collins, W. J., Arculus, R. J., Ellis, D. J., and Frost, C. D.:
773 A geochemical classification for granitic rocks, *J. Petrol.*, 42, 2033–2048,
774 <https://doi.org/10.1093/petrology/42.11.2033>, 2001.

775 Gao, S., Rudnick, R. L., Yuan, H. L., Liu, X. M., Liu, Y. S., Xu, W. L., Ling, W. L.,
776 Ayers, J. C., Wang, X. C., and Wang, Q. H.: Recycling lower continental crust in
777 the North China craton, *Nature*, 432, 892–897,
778 <https://doi.org/10.1038/nature03162>, 2004.

779 Gao, Y. F., Hou, Z. Q., Kamber, B. S., Wei, R. H., Meng, X.J., and Zhao, R. S.: Adakite-
780 like porphyries from the southern Tibetan continental collision zones: evidence for
781 slab melt metasomatism, *Contrib. Mineral. Petr.*, 153, 105–20, [10.1007/s00410-](https://doi.org/10.1007/s00410-006-0137-9)
782 [006-0137-9](https://doi.org/10.1007/s00410-006-0137-9), 2007.

783 Griffin, W. L., Belousova, E. A., Shee, S. R., Pearson, N. J., and O'Reilly, S. Y.:
784 Archean crustal evolution in the northern Yilgarn Craton: U-Pb and Hf-isotope
785 evidence from detrital zircons, *Precambrian. Res.*, 131, 231–282,
786 <https://doi.org/10.1016/J.PRECAMRES.2003.12.011>, 2004.

787 He, P. L., Huang, X. L., Yang, F., and Wang, X.: Mineralogy constraints on magmatic
788 processes controlling adakitic features of Early Permian high-magnesium diorites
789 in the Western Tianshan orogenic belt, *J. Petrol.*, 61, a114,
790 <https://doi.org/10.1093/petrology/egaa114>, 2021.

791 Hoskin, P. W. O.: Trace-element composition of hydrothermal zircon and the alteration
792 of Hadean zircon from the Jack Hills, Australia, *Geochim. Cosmochim. Ac.*, 69,

793 637–648, <https://doi.org/10.1016/j.gca.2004.07.006>, 2005.

794 Hoskin, P. W., and Schaltegger, U.: The composition of zircon and igneous and
795 metamorphic petrogenesis, *Rev. Mineral. Geochem.*, 53, 27–62,
796 <https://doi.org/10.2113/0530027>, 2003.

797 Huang, F., and He, Y.S.: Partial melting of the dry mafic continental crust: Implications
798 for petrogenesis of C-type adakites, *Chinese. Sci. Bull.*, 55, 1255–1267,
799 <https://doi.org/10.1007/s11434-010-3224-2>, 2010.

800 Jackson, S. E., Pearson, N. J., Griffin W L, and Belousova, E. A.: The application of
801 laser ablation-inductively coupled plasma-mass spectrometry to in situ U-Pb
802 zircon geochronology, *Chem. Geol.*, 211(1-2), 47–69,
803 <https://doi.org/doi:10.1016/j.chemgeo.2004.06.017>, 2004.

804 Jiang, H., Jiang, S. Y., Li, W. Q., Zhao, K. D., and Peng, N. J.: Highly fractionated
805 Jurassic I-type granites and related tungsten mineralization in the Shirenzhang
806 deposit, northern Guangdong, South China: Evidence from cassiterite and zircon
807 U-Pb ages, geochemistry and Sr-Nd-Pb-Hf isotopes, *Lithos*, S480187509,
808 <https://doi.org/10.1016/j.lithos.2018.04.030>, 2018.

809 Jiang, Y.H., Jiang, S.Y., Zhao, K.D., Ni, P., Ling, H.F., and Liu, D.Y.: SHRIMP U-Pb
810 zircon dating for lamprophyre from Liaodong Peninsula: Constraints on the initial
811 time of Mesozoic lithosphere thinning beneath eastern China, *Chinese. Sci. Bull.*,
812 50, <https://doi.org/10.1360/982005-373>, 2612–2620, 2005.

813 Kamei, A., Miyake, Y., Owada, M., and Kimura, J.: A pseudo adakite derived from
814 partial melting of tonalitic to granodioritic crust, Kyushu, southwest Japan arc,

815 Lithos, 112, 615–625, <https://doi.org/10.1016/j.lithos.2009.05.024>, 2009.

816 Kay, R. W., and Kay, S. M.: Delamination and delamination magmatism,
817 Tectonophysics, 219, 177–189, [https://doi.org/10.1016/0040-1951\(93\)90295-U](https://doi.org/10.1016/0040-1951(93)90295-U),
818 1993.

819 King, S. D., Frost, D. J., and Rubie D. C.: Why cold slabs stagnate in the transition zone,
820 Geology, 43(3), 231–234, <https://doi.org/10.1130/G36320.1>, 2015.

821 Li, C.M., Zhang, C.H., Cope, T.D., and Lin, Y.: Out-of-sequence thrusting in polycyclic
822 thrust belts: An example from the Mesozoic Yanshan belt, North China Craton,
823 Tectonics, 35, 2082–2116, <https://doi.org/10.1002/2016TC004187>, 2016.

824 Li, S. Z., and Zhao, G. C.: SHRIMP U-Pb zircon geochronology of the Liaoji granitoids:
825 Constraints on the evolution of the Paleoproterozoic Jiao-Liao-Ji belt in the
826 Eastern Block of the North China Craton, Precambrian. Res., 158, 1–16,
827 <https://doi.org/10.1016/j.precamres.2007.04.001>, 2007.

828 Li, S. Z., Zhao, G. C., Sun, M., Han, Z. Z., Hao, D. F., Luo, Y., and Xia, X. P.:
829 Deformation history of the Paleoproterozoic Liaohe Group in the Eastern Block
830 of the North China Craton, J. Asian. Earth. Sci., 24, 659–674, <https://doi.org/10.1016/j.jseaes.2003.11.008>, 2005.

832 Li, S.Z., Liu, J.Z., Zhao, G.C., Wu, F.Y., Han, Z.Z., and Yang, Z.Z.: Key
833 geochronology of Mesozoic deformation in the eastern block of the North China
834 Craton and its constraints on regional tectonics: A case of Jiaodong and Liaodong
835 Peninsula, Acta. Petrol. Sin., 2, 633–646, <https://doi.org/10.1007/BF02873097>,
836 2004.

837 Li, X.H., Long, W.G., Li, Q.L., Liu, Y., Zheng, Y.F., Yang, Y.H., Chamberlain, K.R.,
838 Wan, D.F., Guo, C.H., and Wang, X.C.: Penglai Zircon Megacrysts: A Potential
839 New Working Reference Material for Microbeam Determination of Hf-O Isotopes
840 and U-Pb Age, *Geostand. Geoanal. Res.*, 34, 117–134,
841 <https://doi.org/info:doi/10.1111/j.1751-908X.2010.00036.x>, 2010.

842 Li, Z., Chen, B., Liu, J.W., Zhang, L., and Yang, C.: Zircon U-Pb ages and their
843 implications for the South Liaohe Group in the Liaodong Peninsula, Northeast
844 China (in Chinese with English abstract), *Acta. Petrol. Sin.*, 31(6), 1589–1605,
845 2015.

846 Liu, Y.S., Hu, Z.C., Zong, K.Q., Gao, C.G., Gao, S., Xu, J., and Chen, H.H.:
847 Reappraisal and refinement of zircon U-Pb isotope and trace element analyses
848 by LA-ICP-MS, *Chinese. Sci. Bull.*, 55, 1535–1546,
849 <https://doi.org/10.1007/s11434-010-3052-4>, 2010.

850 Loader, M. A., Wilkinson, J. J., and Armstrong, R. N.: The effect of titanite
851 crystallisation on Eu and Ce anomalies in zircon and its implications for the
852 assessment of porphyry Cu deposit fertility, *Earth. Planet. Sc. Lett.*, 472, 107–119,
853 <https://doi.org/10.1016/j.epsl.2017.05.010>, 2017.

854 Ludwig, K. R.: User's manual for Isoplot 3.00: a geochronological toolkit for Microsoft
855 Excel, Berkeley. Geochronology. Center., 2003.

856 Ma, L., Jiang, S.Y., Dai, B.Z., Jiang, Y.H., Hou, M.L., Pu, W., and Xu, B.: Multiple
857 sources for the origin of Late Jurassic Linglong adakitic granite in the Shandong
858 Peninsula, eastern China: Zircon U-Pb geochronological, geochemical and Sr-Nd-

859 Hf isotopic evidence, *Lithos*, 162–163, 251–263,
860 <https://doi.org/10.1016/j.lithos.2013.01.009>, 2013b.

861 Ma, Q., Zheng, J.P., Griffin, W. L., Zhang, M., Tang, H.Y., Su, Y.P., and Ping, X.Q.:
862 Triassic “adakitic” rocks in an extensional setting (North China): Melts from the
863 cratonic lower crust, *Lithos*, 149, 159–173, [https://doi.org/](https://doi.org/10.1016/j.lithos.2012.04.017)
864 [10.1016/j.lithos.2012.04.017](https://doi.org/10.1016/j.lithos.2012.04.017), 2012.

865 Ma, Q., Zheng, J.P., Xu, Y.G., Griffin, W. L., and Zhang, R.S.: Are continental
866 “adakites” derived from thickened or foundered lower crust? *Earth. Planet. Sc.*
867 *Lett.*, 419, 125–133, <https://doi.org/10.1016/j.epsl.2015.02.036>, 2015.

868 Ma, X.H., Chen, B., and Yang, M.C.: Magma mixing origin for the Aolunhua porphyry
869 related to Mo-Cu mineralization, eastern Central Asian Orogenic Belt, *Gondwana*.
870 *Res.*, 24, 1152–1171, <https://doi.org/10.1016/j.gr.2013.02.010>, 2013a.

871 Macpherson, C. G., Dreher, S. T., and Thirlwall, M. F.: Adakites without slab melting:
872 High pressure differentiation of island arc magma, Mindanao, the Philippines,
873 *Earth. Planet. Sc. Lett.*, 243(3), 581–593,
874 <https://doi.org/10.1016/j.epsl.2005.12.034>, 2006.

875 Mahon, K. I.: The New “York” regression: Application of an improved statistical
876 method to geochemistry, *Int. Geol. Rev.*, 38(4), 293–303,
877 <https://doi.org/10.1080/00206819709465336>, 1996.

878 Maniar, P. D., and Piccoli, P. M.: Tectonic discrimination of granitoids, *Geol. Soc. Am.*
879 *Bull.*, 101, 635–643, [https://doi.org/10.1130/0016-](https://doi.org/10.1130/0016-7606(1989)101<0635:TDOG>2.3.CO;2)
880 [7606\(1989\)101<0635:TDOG>2.3.CO;2](https://doi.org/10.1130/0016-7606(1989)101<0635:TDOG>2.3.CO;2), 1989.

881 Martin, H., Smithies, R.H., Rapp, R., Moyen, J.-F., and Champion, D.: An overview of
882 adakite, tonalite-trondhjemite-granodiorite (TTG), and sanukitoid: relationships
883 and some implications for crustal evolution, *Lithos*, 79, 1–24,
884 <https://doi.org/10.1016/j.lithos.2004.04.048>, 2005.

885 Meng, E., Liu, F.L., Liu, P.H., Liu, C.H., Yang, H., Wang, F., Shi, J.R., and Cai, J.:
886 Petrogenesis and tectonic significance of Paleoproterozoic meta-mafic rocks from
887 central Liaodong Peninsula, northeast China: Evidence from zircon U-Pb dating
888 and in situ Lu-Hf isotopes, and whole-rock geochemistry, *Precambrian. Res.*, 247,
889 92–109, <https://doi.org/10.1016/j.precamres.2014.03.017>, 2014.

890 Meng, Q. R.: What drove late Mesozoic extension of the northern China-Mongolia tract?
891 *Tectonophysics*, 369, 155–174, [https://doi.org/10.1016/S0040-1951\(03\)00195-1](https://doi.org/10.1016/S0040-1951(03)00195-1),
892 2003.

893 Mezger, K., and Krogstad, E. J.: Interpretation of discordant U-Pb zircon ages: An
894 evaluation, *J. Metamorph. Geol.*, 15, 127–140, <https://doi.org/10.1111/j.1525-1314.1997.00008.x>, 1997.

896 Miller, C. F., McDowell, S. M., and Mapes, R. W.: Hot and cold granites? Implications
897 of zircon saturation temperatures and preservation of inheritance, *Geology*, 31,
898 529–532, [https://doi.org/10.1130/0091-7613\(2003\)031<0529:hacgio>2.0.co;2](https://doi.org/10.1130/0091-7613(2003)031<0529:hacgio>2.0.co;2),
899 2003.

900 Miller, J.F., and Harris, N.B.W.: Evolution of continental crust in the Central Andes;
901 constraints from Nd isotope systematics, *Geology*, 17, 615–617,
902 [https://doi.org/10.1130/0091-7613\(1989\)0172.3.CO;2](https://doi.org/10.1130/0091-7613(1989)0172.3.CO;2), 1989.

903 Moyen, J.: High Sr/Y and La/Yb ratios: The meaning of the “adakitic signature”, *Lithos*,
904 112, 556–574, <https://doi.org/10.1016/j.lithos.2009.04.001>, 2009.

905 Nakamura, H., and Iwamori, H.: Generation of adakites in a cold subduction zone due
906 to double subducting plates, *Contrib. Mineral. Petr.*, 165, 1107–1134,
907 <https://doi.org/10.1007/s00410-013-0850-0>, 2013.

908 Nasdala, L., Pidgeon, R. T., Wolf, D., and Irmer, G., Metamictization and U-Pb isotopic
909 discordance in single zircons: a combined Raman microprobe and SHRIMP ion
910 probe study, *Miner. Petrol.*, 62, 1–27, <https://doi.org/10.1007/BF01173760>, 1998.

911 Nash, D.J., Ciborowski, T., Ulliyott, J.S., Pearson, M.P. and Whitaker, K.A.: Origins of
912 the sarsen megaliths at Stonehenge, *Sci. Adv.*, 6, eabc0133,
913 <https://doi.org/10.1126/sciadv.abc0133>, 2020.

914 Ou, Q., Wang, Q., Wyman, D. A., Zhang, H.X., Yang, J.H., Zeng, J.P., Hao, L.L., Chen,
915 Y.W., Liang, and H., Qi, Y.: Eocene adakitic porphyries in the central-northern
916 Qiangtang Block, central Tibet: Partial melting of thickened lower crust and
917 implications for initial surface uplifting of the plateau, *J. Geophys. Res. Solid.*
918 *Earth.*, 122, 1025–1053, <https://doi.org/10.1002/2016JB013259>, 2017.

919 Parsons, B., and Dan, M.K.: Mantle convection and the thermal structure of the plates,
920 *J. Geophys. Res. Solid. Earth.*, 83, <https://doi.org/10.1029/JB083iB09p04485>,
921 1978.

922 Patiño Douce, A.E.P.: What do experiments tell us about the relative contributions of
923 crust and mantle to the origin of granitic magmas? *Geol. Soc. Lond. Spec. Publ.*,
924 168(1), 55–75, <https://doi.org/10.1144/GSL.SP.1999.168.01.05>, 1999.

925 Qiu, L., Kong, R.Y., Yan, D.P., Wells, M.L., Wang, A.P., Sun, W.H., Yang, W.X., Han,
926 Y.G., Li, C.M., and Zhang, Y.X.: The Zhayao tectonic window of the Jurassic
927 Yuantai thrust system in Liaodong Peninsula, NE China: Geometry, kinematics
928 and tectonic implications, *J. Asian Earth Sci.*, 164, 58–71, [https://doi.org/](https://doi.org/10.1016/j.jseaes.2018.06.012)
929 10.1016/j.jseaes.2018.06.012, 2018.

930 Rapp, R. P., Shimizu, N and Norman, M.D.: Growth of early continental crust by partial
931 melting of eclogite, *Nature*, 425(6958), 605–609,
932 <https://doi.org/10.1038/nature02031>, 2003.

933 Ren, Z.H., Lin, W., Faure, M., Meng, L.T., Qiu, H.B., and Zeng, J.P.: Triassic- Jurassic
934 evolution of the eastern North China Craton: Insights from the Lushun-Dalian area,
935 South Liaodong Peninsula, NE China, *Geol. Soc. Am. Bull.*, 133(1–2),
936 <https://doi.org/10.1130/B35533.1>, 2020.

937 Rudnick, R. L., and Gao, S.: Composition of the continental crust, *Treatise. Geochem.*,
938 3, 1–64, <https://doi.org/10.1016/b0-08-043751-6/03016-4>, 2003.

939 Schiller, D., and Finger, F.: Application of Ti-in-zircon thermometry to granite studies:
940 problems and possible solutions, *Contrib. Mineral. Petr.*, 174,
941 <https://doi.org/10.1007/s00410-019-1585-3>, 2019.

942 Schmitz, M. D., and Kuiper, K. F.: High-Precision Geochronology, *Elements.*, 9, 25–
943 30, <https://doi.org/10.2113/gselements.9.1.25>, 2013.

944 Schwartz, J. J., Johnson, K., Miranda, E. A., and Wooden, J. L.: The generation of high
945 Sr/Y plutons following Late Jurassic arc–arc collision, Blue Mountains province,
946 NE Oregon, *Lithos*, 126, 22–41, <https://doi.org/10.1016/j.lithos.2011.05.005>,

947 2011.

948 Scott, E.M., Allen, M.B., Macpherson, C.G., McCaffrey, K.J.W., Davidson, J.P.,
949 Saville, C., and Ducea, M.N.: Andean surface uplift constrained by radiogenic
950 isotopes of arc lavas, *Nat. Commun.*, 9(1), 969, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-018-03173-4)
951 018-03173-4, 2018.

952 Shahbazi, H., Maghami, Y. T., Azizi, H., Asahara, Y., Siebel, Y., Maanijou, M., and
953 Rezaei, A.: Zircon U-Pb ages and petrogenesis of late Miocene adakitic rocks from
954 the Sari Gunay gold deposit, NW Iran, *Geol. Mag.*,
955 <https://doi.org/10.1017/S0016756821000297>, 2021.

956 Shannon. R.D.: Revised effective ionic radii and systematic studies of interatomic
957 distances in halides and chalcogenides, *Acta. Crystall. A.*, 32(5), 751–767,
958 <https://doi.org/10.1107/S0567739476001551>, 1976.

959 Shaw, D.M.: Trace element fractionation during anatexis, *Geochim. Cosmochim. Ac.*,
960 34, 237–243, [https://doi.org/10.1016/0016-7037\(70\)90009-8](https://doi.org/10.1016/0016-7037(70)90009-8), 1970.

961 Siégel, C., Bryan, S. E., Allen, C. M., and Gust, D. A.: Use and abuse of zircon-based
962 thermometers: A critical review and a recommended approach to identify
963 antecrystic zircons, *Earth. Sci. Rev.*, 176, 87–116,
964 <https://doi.org/10.1016/j.earscirev.2017.08.011>, 2018.

965 Skjerlie, K. P., and Johnston, A. D.: Vapor-absent melting at 10 kbar of a biotite-and
966 amphibole-bearing tonalitic gneiss: implications for the generation of A-type
967 granites, *Geology*, 20, 263–266, [https://doi.org/10.1130/0091-](https://doi.org/10.1130/0091-7613(1992)0202.3.CO;2)
968 7613(1992)0202.3.CO;2, 1992.

969 Sláma, J., Košler, J., and Condon, D. J.: Plešovice zircon-A new natural reference
970 material for U-Pb and Hf isotopic microanalysis, *Chem. Geol.*, 249(1–2), 1–35,
971 <https://doi.org/doi:10.1016/j.chemgeo.2007.11.005>, 2008.

972 Söderlund, U., Patchett, P.J., Vervoort, J.D., and Isachsen, C.E.: The ^{176}Lu decay
973 constant determined by Lu-Hf and U-Pb isotope systematics of Precambrian mafic
974 intrusions, *Earth. Planet. Sc. Lett.*, 219(3), 311–324,
975 [https://doi.org/10.1016/S0012-821X\(04\)00012-3](https://doi.org/10.1016/S0012-821X(04)00012-3), 2004.

976 Sun, S. S., and McDonough, W. F.: Chemical and isotopic systematics of oceanic
977 basalts: implications for mantle composition and processes, *Geol. Soc. Lond. Spec.*
978 *Publ.*, 42, 313–345, <https://doi.org/10.1144/GSL.SP.1989.042.01.19>, 1989.

979 Tang, J., Xu, W. L., Wang, F., and Ge, W.C.: Subduction history of the Paleo-Pacific
980 slab beneath Eurasian continent: Mesozoic-Paleogene magmatic records in
981 Northeast Asia, *Sci. China. Earth. Sci.*, 61, 527–559,
982 <https://doi.org/CNKI:SUN:JDXG.0.2018-05-003>, 2018.

983 Trail, D., Watson, E. B., and Tailby, N. D.: Ce and Eu anomalies in zircon as proxies
984 for the oxidation state of magmas, *Geochim. Cosmochim. Ac.*, 97, 70–87,
985 <https://doi.org/10.1016/j.gca.2012.08.032>, 2012.

986 Vervoort, J.D., and Blichert-Toft, J.: Evolution of the depleted mantle: Hf isotope
987 evidence from juvenile rocks through time, *Geochim. Cosmochim. Ac.*, 63(3),
988 533–556, [https://doi.org/10.1016/S0016-7037\(98\)00274-9](https://doi.org/10.1016/S0016-7037(98)00274-9), 1999.

989 Wan, Y.S., Song, B., Liu, D.Y., Wilde, S. A., Wu, J.S., Shi, Y.R., Yin, X.Y., and Zhou,
990 H.Y.: 2006. SHRIMP U-Pb zircon geochronology of Palaeoproterozoic

991 metasedimentary rocks in the North China Craton: Evidence for a major Late
992 Palaeoproterozoic tectonothermal event, *Precambrian. Res.*, 149, 271,
993 <https://doi.org/10.1016/j.precamres.2006.06.006>, 2006.

994 Wang, F., Liu, F., Schertl, H.P., Xu, W., Liu, P., and Tian, Z.: Detrital zircon U-Pb
995 geochronology and Hf isotopes of the Liaohe Group, Jiao-Liao-Ji Belt:
996 Implications for the Paleoproterozoic tectonic evolution, *Precambrian. Res.*, 340,
997 105633, [10.1016/j.precamres.2020.105633](https://doi.org/10.1016/j.precamres.2020.105633), 2020b.

998 Wang, N., Wu, C. L, Lei, M., and Chen, H.J.: Petrogenesis and tectonic implications of
999 the Early Paleozoic granites in the western segment of the North Qilian orogenic
1000 belt, China, *Lithos*, 312–313, 89–107, <https://doi.org/10.1016/j.lithos.2018.04.023>,
1001 2018.

1002 Wang, Q., Xu, J. F., Jian, P., Bao Z. W., Zhao. Z. H., Li. C. F., Xiong. X. L., and Ma,
1003 J. L.: Petrogenesis of adakitic porphyries in an extensional tectonic setting, Dexing,
1004 South China: Implications for the genesis of porphyry copper mineralization, *J.*
1005 *Petrol.*, 47(1), 119–144, <https://doi.org/10.1093/petrology/egi070>, 2006.

1006 Wang, X. P., Oh, C. W., Peng, P., Zhai, M. G., Wang, X. H., and Lee, B. Y.:
1007 Distribution pattern of age and geochemistry of 2.18-2.14 Ga I- and A-type
1008 granites and their implication for the tectonics of the Liao-Ji belt in the North
1009 China Craton, *Lithos*, 364–365, 105518,
1010 <https://doi.org/10.1016/j.lithos.2020.105518>, 2020a.

1011 Wang, X., Griffin, W. L., Chen, J., Huang, P.Y., and Li, X.: U and Th contents and
1012 Th/U ratios of zircon in felsic and mafic magmatic rocks: Improved zircon-melt

1013 distribution coefficients, *Acta. Geol. Sin. Engl.*, 85, 164–174,
1014 <https://doi.org/10.1111/j.1755-6724.2011.00387.x>, 2011.

1015 Wang, X., Griffin, W. L., O'Reilly, S. Y., and Li, W.X.: Three stages of zircon growth
1016 in magmatic rocks from the Pingtan Complex, eastern China, *Acta. Geol. Sin.*
1017 *Engl.*, 81, 68–80, <https://doi.org/10.3321/j.issn:1000-9515.2007.01.008>, 2007.

1018 Wang, X.L., Lv, X., Liu, Y.J., Zhao, Y.Y., Li, C., Wu, W.B., Wang, Y.P., and Li, H.Y.:
1019 LA-ICP-MS zircon U-Pb ages, geochemical characteristics of Late Triassic
1020 intrusives in Xiuyan area, eastern Liaoning and their geological significances, *Ore.*
1021 *Geol. Rev.*, 65, 401–416, <https://doi.org/10.16509/j.georeview.2019.02.010>, 2019.

1022 Watson, E. B., and Harrison, T. M.: Zircon saturation revisited: temperature and
1023 composition effects in a variety of crustal magma types, *Earth. Planet. Sci. Lett.*,
1024 64, 295–304, [https://doi.org/10.1016/0012-821X\(83\)90211-X](https://doi.org/10.1016/0012-821X(83)90211-X), 1983.

1025 Watson, E. B., and Harrison, T. M.: Zircon thermometer reveals minimum melting
1026 conditions on earliest Earth, *Science*, 308, 841–844,
1027 <https://doi.org/10.1126/science.1110873>, 2005.

1028 Whalen, J. B., Currie, K. L., and Chappell, B. W.: A-type granites: geochemical
1029 characteristics, discrimination and petrogenesis, *Contrib. Mineral. Petr.*, 95, 407–
1030 419, <https://doi.org/10.1007/BF00402202>, 1987.

1031 Wolf, M. B., and London, D.: Apatite dissolution into peraluminous haplogranitic melts:
1032 An experimental study of solubilities and mechanisms, *Geochim. Cosmochim.*
1033 *Acta.*, 58, 4127–4145. [https://doi.org/10.1016/0016-7037\(94\)90269-0](https://doi.org/10.1016/0016-7037(94)90269-0), 1994.

1034 Wu, F. Y., Jahn, B. M., Wilde, S. A., Lo, C. H., Yui, T. F., Lin, Q., Ge, W. C., and Sun,

1035 D. Y: Highly fractionated I-type granites in NE China (I): geochronology and
1036 petrogenesis, *Lithos*, 66, 241–273, [https://doi.org/10.1016/s0024-4937\(02\)00222-](https://doi.org/10.1016/s0024-4937(02)00222-0)
1037 0, 2003.

1038 Wu, F. Y., Liu, X. C., Ji, W. Q., Wang, J. M., and Yang, L.: Highly fractionated granites:
1039 Recognition and research, *Sci. China. Earth. Sci.*, 60, 1201–1219,
1040 <https://doi.org/10.1007/s11430-016-5139-1>, 2017.

1041 Wu, F. Y., Yang, J. H., and Liu, X. M.: Geochronological framework of the Mesozoic
1042 granitic magmatism in the Liaodong Peninsula, Northeast China (in Chinese with
1043 English abstract), *Geol. J. China. Univ.*, 11, 305–317, 2005b.

1044 Wu, F.Y., Han, R.H., Yang, J.H., Wilde, S. A., Zhai, M.G., and Park, S. C.: Initial
1045 constraints on the timing of granitic magmatism in North Korea using U-Pb zircon
1046 geochronology, *Chem. Geol.*, 238, 232–248,
1047 <https://doi.org/10.1016/j.chemgeo.2006.11.012>, 2007.

1048 Wu, F.Y., Lin, J.Q., Wilde, S. A., Zhang, X.O., and Yang, J.H.: Nature and significance
1049 of the Early Cretaceous giant igneous event in eastern China, *Earth. Planet. Sci.*
1050 *Lett.*, 233, 103–119, <https://doi.org/10.1016/j.epsl.2005.02.019>, 2005c.

1051 Wu, F.Y., Yang, J.H., Wilde, S. A., and Zhang, X.O.: Geochronology, petrogenesis and
1052 tectonic implications of Jurassic granites in the Liaodong Peninsula, NE China,
1053 *Chem. Geol.*, 221, 127–156, <https://doi.org/10.1016/j.chemgeo.2005.04.010>,
1054 2005a.

1055 Wu, F.Y., Yang, Y.H., Xie, L.W., Yang, J.H., and Xu, P.: Hf isotopic compositions of
1056 the standard zircons and baddeleyites used in U-Pb geochronology, *Chem. Geol.*,

1057 234, 105–126, <https://doi.org/10.1016/j.chemgeo.2006.05.003>, 2006.

1058 Xu, J. F., Shinjo, R., Defant, M. J., Wang, Q., and Rapp, R. P.: Origin of Mesozoic
1059 adakitic intrusive rocks in the Ningzhen area of east China: partial melting of
1060 delaminated lower continental crust? *Geology*, 30(12), 1111–1114,
1061 [https://doi.org/10.1130/0091-7613\(2002\)0302.0.CO;2](https://doi.org/10.1130/0091-7613(2002)0302.0.CO;2), 2002.

1062 Xue, J. X., Liu, Z. H., Liu, J. X., Dong, X. J., Feng, F., and Lian, G. H.: Geochemistry,
1063 Geochronology, Hf isotope and Tectonic Significance of the Late Jurassic Huangdi
1064 Pluton in Xiuyan, Liaodong Penins (in Chinese with English abstract), *Earth
1065 Science*, 46(6), 2030–2043, <https://doi.org/10.3799/dqkx.2020.008>, 2020.

1066 Xue, S., Ling, M. X., Liu, Y. L., Zhang, H., and Sun, W.D.: The genesis of early
1067 Carboniferous adakitic rocks at the southern margin of the Alxa Block, North
1068 China, *Lithos*, 278–281, 181–194, <https://doi.org/10.1016/j.lithos.2017.01.012>,
1069 2017.

1070 Yang, F.C., Song, Y.H., Hao, L.B., and Peng, C.: Late Jurassic SHRIMP U-Pb age and
1071 Hf isotopic characteristics of granite from the Sanjiazi Area in Liaodong and their
1072 geological significance (in Chinese with English abstract), *Acta. Geol. Sin. Engl.*,
1073 89, 1773–1782, 2015a.

1074 Yang, F.C., Song, Y.H., Yang, J.L., Shen, X., and Gu, Y.C.: SHRIMP U-Pb age and
1075 geochemical characteristics of granites in Wulong-Sidaogou Gold Deposit, East
1076 Liaoning (in Chinese with English abstract), *Geotectonica et Metallogenia*, 42,
1077 940–954, <https://doi.org/10.16539/j.ddgzyckx.2018.05.010>, 2018.

1078 Yang, J. H., Wu, F. Y., Xie, L. W., and Liu, X. M.: Petrogenesis and tectonic

1079 implications of Kuangdonggou synites in the Liaodong Peninsula, east North
1080 China Craton: Constraints from in-suit zircon U-Pb ages and Hf isotopes (in
1081 Chinese with English abstract), *Acta. Petrol. Sin.*, 23, 263–276,
1082 <https://doi.org/10.1016/j.sedgeo.2006.03.028>, 2007.

1083 Yang, J.H., Wu, F.Y., Lo, C.H., Chung, S.L., Zhang, Y.B., and Wilde, S.A.:
1084 Deformation age of Jurassic granites in the Dandong area, eastern China: Ar-
1085 40/Ar-39 geochronological constraints (in Chinese with English abstract), *Acta.*
1086 *Petrol. Sin.*, 20(5), 1205–1214, 2004.

1087 Yang, M. C., Chen, B., and Yan, C.: Petrological, geochronological, geochemical and
1088 Sr-Nd-Hf isotopic constraints on the petrogenesis of the Shuangcha
1089 Paleoproterozoic megaporphyritic granite in the southern Jilin Province: Tectonic
1090 implications (in Chinese with English abstract), *Acta. Petrol. Sin.*, 31, 1573–1588,
1091 2015b.

1092 Yang, M.C., Chen, B., and Yan, C.: Paleoproterozoic Gneissic Granites in the Liaoji
1093 Mobile Belt, North China Craton: Implications for Tectonic Setting. In: Zhai M.,
1094 Zhao Y., Zhao T. (eds) *Main Tectonic Events and Metallogeny of the North China*
1095 *Craton*, Springer. Geology. Springer. Singapore., 155–180,
1096 https://doi.org/10.1007/978-981-10-1064-4_7, 2016.

1097 Yu, S., Zhang, J., and Del Real, P. G.: Geochemistry and zircon U-Pb ages of adakitic
1098 rocks from the Dulan area of the North Qaidam UHP terrane, north Tibet:
1099 constraints on the timing and nature of regional tectonothermal events associated
1100 with collisional orogeny, *Gondwana. Res.*, 21, 167–179,

1101 <https://doi.org/10.1016/j.gr.2011.07.024>, 2012.

1102 Yuan, H.L., Gao, S., Dai, M.N., Zong, C.L., Günther, D., Fontaine, G.H., Liu, X.M.,
1103 and Diwu, C.R.: Simultaneous determinations of U-Pb age, Hf isotopes and trace
1104 element compositions of zircon by excimer laser-ablation quadrupole and
1105 multiple-collector ICP-MS, *Chem. Geol.*, 247, 100–118,
1106 <https://doi.org/10.1016/j.chemgeo.2007.10.003>, 2008.

1107 Yuan, L. L., Zhang, X. H., Xue, F. H., Han, C. M., Chen, H. H., and Zhai, M. G.: Two
1108 episodes of Paleoproterozoic mafic intrusions from Liaoning province, North
1109 China Craton: Petrogenesis and tectonic implications, *Precambrian. Res.*, 264,
1110 119–139, <https://doi.org/10.1016/j.precamres.2015.04.017>, 2015.

1111 Zeng, R.Y., Lai, J.Q., Mao, X.C., Li, B., Ju, P.J., and Tao, S.L.: Geochemistry, zircon
1112 U-Pb dating and Hf isotopies composition of Paleozoic granitoids in Jinchuan,
1113 NW China: Constraints on their petrogenesis, source characteristics and tectonic
1114 implication, *J. Asian. Earth. Sci.*, 121, 20–33,
1115 <https://doi.org/10.1016/j.jseaes.2016.02.009>, 2016.

1116 Zeng, R.Y., Lai, J.Q., Mao, X.C., Li, B., Zhang, J.D., Bayless, R., and Yang, L.Z.:
1117 Paleoproterozoic Multiple Tectonothermal Events in the Longshoushan Area,
1118 Western North China Craton and Their Geological Implication: Evidence from
1119 Geochemistry, Zircon U-Pb Geochronology and Hf Isotopes, *Minerals. Basel.*, 8,
1120 <https://doi.org/10.3390/MIN8090361>, 2018.

1121 Zhai, M.G., Meng, Q.R., Liu, J.M., Hou, Q.L., Hu, S.B., Li, Z., Zhang, H.F., Liu, W.,
1122 Shao, J.A., and Zhu, R.X.: Geological features of Mesozoic tectonic regime

1123 inversion in eastern North China and implication for geodynamics, *Earth. Science.*
1124 *Frontiers.*, 11, 285–298, <https://doi.org/10.3321/j.issn:1005-2321.2004.03.027>,
1125 2004.

1126 Zhan, Q. Y., Zhu, D. C., Wang, Q., Weinberg, R. F., Xie, J. C., Li, S. M., Zhang, L.L.,
1127 and Zhao, Z. D.: Source and pressure effects in the genesis of the Late Triassic
1128 high Sr/Y granites from the Songpan-Ganzi Fold Belt, eastern Tibetan Plateau,
1129 *Lithos*, 368–369, <https://doi.org/10.1016/j.lithos.2020.105584>, 2020.

1130 Zhang, B., Guo, F., Zhang, X. B., Wu, Y.M., Wang, G.Q., and Zhao, L.: Early
1131 Cretaceous subduction of Paleo-Pacific Ocean in the coastal region of SE China:
1132 Petrological and geochemical constraints from the mafic intrusions, *Lithos*, 334–
1133 335, 8–24, <https://doi.org/10.1016/j.lithos.2019.03.010>, 2019.

1134 Zhang, Q., Qian, Q., Wang, E.Q., Wang, Y., Zhao, T.P., Hao, J., and Guo, G.J.: An
1135 East China plateau in mid-late Yanshanian period: implication from adakites,
1136 Chinese (in Chinese with English abstract). *Chinese Journal of Geology*, 36 (2),
1137 248–255, 2001.

1138 Zhang, Q., Wang, Y., Liu, H., and Wang, Y.: On the space-time distribution and
1139 geodynamic environments of adakites in China annex: controversies over differing
1140 opinions for adakites in China (in Chinese with English abstract), *Earth Science*
1141 *Frontiers*, [https://doi.org/10.1016/S0955-2219\(02\)00073-0](https://doi.org/10.1016/S0955-2219(02)00073-0), 2003.

1142 Zhang, S., Zhu, G., Xiao, S.Y., Su, N., Liu, C., Wu, X.D., Yin, H., Li, Y.J., and Lu,
1143 Y.C.: Temporal variations in the dynamic evolution of an overriding plate:
1144 Evidence from the Wulong area in the eastern North China Craton, China, *Geol.*

1145 Soc. Am. Bull., 132(9–10), 2023–2042, <https://doi.org/10.1130/B35465.1>, 2020.

1146 Zhao, K.D., Jiang, S.Y., Ling, H.F., and Palmer, M. R.: Reliability of LA-ICP-MS U-
1147 Pb dating of zircons with high U concentrations: A case study from the U-bearing
1148 Douzhashan Granite in South China, *Chem. Geol.*, 389, 110–121,
1149 <https://doi.org/10.1016/j.chemgeo.2014.09.018>, 2014.

1150 Zhao, K.D., Jiang, S.Y., Ling, H.F., and Palmer, M. R.: Reliability of LA-ICP-MS U-
1151 Pb dating of zircons with high U concentrations: A case study from the U-bearing
1152 Douzhashan Granite in South China, *Chem. Geol.*, 389, 110–121,
1153 <https://doi.org/10.1016/j.chemgeo.2014.09.018>, 2014.

1154 Zheng, J.P., and Dai, H.K.: Subduction and retreating of the western Pacific plate
1155 resulted in lithospheric mantle replacement and coupled basinmountain respond in
1156 the North China Craton, *Sci. China. Earth. Sci.*, 61, 406–424,
1157 <https://doi.org/10.1007/s11430-017-9166-8>, 2018.

1158 Zheng, Y.F., Xu, Z.F., Zhao, Z.F., and Dai, L.Q.: Mesozoic mafic magmatism in North
1159 China: Implications for thinning and destruction of cratonic lithosphere, *Sci. China.*
1160 *Earth Sci.*, 61, 353–385, <https://doi.org/10.1007/s11430-017-9160-3>, 2018.

1161 Zhu, R. X., and Xu, Y. G.: The subduction of the west Pacific plate and the destruction
1162 of the North China Craton, *Sci. China. Earth Sci.*, 62, <https://doi.org/1340-1350>,
1163 [10.1007/s11430-018-9356-y](https://doi.org/10.1007/s11430-018-9356-y), 2019.

1164 Zhu, R.X., and Xu, Y.G.: The subduction of the west Pacific plate and the destruction
1165 of the North China Craton, *Sci. China. Earth. Sci.*, 62, 1340–1350,
1166 <https://doi.org/10.1007/s11430-018-9356-y>, 2019.

1167 **Table captions**

1168 Table S1. Major element (wt. %) and trace element (ppm) compositions of the

1169 Zhoujiapuzi granite

1170 Table S2. Raman microprobe data

1171 Table S3. The zircon major element (wt. %) and trace element (ppm) from the


1172 Zhoujiapuzi granite

1173 Table S4. Zircon La-ICP-MS U-Pb isotopic data and ages of the Zhoujiapuzi granite

1174 Table S5. Zircon Hf isotopic data of the Zhoujiapuzi granite

1175

1176 **Figure captions**




1177

1178 Figure 1. (a) Simplified geological map of Northeast China (Modified from Li et al.,

1179 2016); (b) distribution of Mesozoic intrusions in the Liaodong Peninsula (Modified

1180 from Wu et al., 2005a); (c) geological map of the Zhoujiapuzi granite.

1181




1182

1183 Figure 2. Outcrop photograph (a) and corresponding micrographs (b, c, d-

1184 perpendicular polarized light). Q quartz; Kfs feldspar; Pl plagioclase; Pth perthite; Bt

1185 biotite

1186




1187

1188 Figure 3. Geochemical classification diagrams for the Zhoujiapuzi granite. (a) TAS

1189 diagram (after Frost et al., 2001); (b) A/CNK-A/NK diagram (after Maniar and

1190 Piccoli, 1989)


1191



1192

1193 Figure 4. Chondrite-normalized REE patterns and primitive mantle-normalized trace
1194 element patterns of the Zhoujiapuzi granite (chondrite and primitive mantle values are
1195 from Sun and McDonough, 1989).

1196




1197

1198 Figure 5. CL images of zircons. Circles denote U-Pb analysis spot. Numbers in the

1199 circles are the spot numbers. Numbers near the analytical spots are the U–Pb ages
1200 (Ma).


1201



1202

1203 Figure 6. Chondrite-normalized REE patterns of zircon (chondrite values are from
1204 Sun and McDonough, 1989).


1205



1206


1207 Figure 7. Concordia diagrams for zircon LA-ICP-MS U-Pb analyses.

1208




1209
 1210 Figure 8. Zircon $\epsilon\text{Hf}(t)$ -age (Ma) diagram for samples in this study and published data
 1211 for the region.

1212



1213
 1214 Fig. 9. Discrimination plots for magmatic and hydrothermal zircon (Hoskin, 2005).

1215




1216

1217 Figure 10. Covariation diagrams for zircon from the Zhoujiapuzi granite. (a) U vs.

1218 Zr/Hf; (b) T_{Zr-Ti} vs. Zr/Hf; (c) Ce/Ce* vs. Zr/Hf.

1219



○ Zhoujiapuzi granite (this study)

+ Other middle-late Jurassic granitoids in Liaodong Peninsula (Wu et al., 2005a; Yang et al., 2015b, 2018; Xue et al., 2020)

1220


1221 Figure 11. Chemical variation diagrams for the Zhoujiapuzi granite. (a and b)

1222 $\text{Zr}+\text{Nb}+\text{Ce}+\text{Y}$ vs. $(\text{Na}_2\text{O} + \text{K}_2\text{O})/\text{CaO}$ and TFeO/MgO (after Whalen et al., 1987); (c

1223 and d) SiO_2 vs. Zr and Ce (after Collins et al., 1982); (e) SiO_2 vs. P_2O_5 diagram; (f)

1224 Rb vs. Y diagram


1225



1226

1227 Figure 12. Adakite discrimination diagrams for the Zhoujiapuzi granite (after Defant
 1228 and Drummond, 1990).

1229




1230

1231 Figure 13. Source characteristics (a-d and f-h) and crystal fractionation (e and i)

1232 discrimination diagrams for the Zhoujiapuzi granite. Plots of (a) Nd/Sm vs. Ti/Eu (Yu
 1233 et al., 2012); (b-d) SiO₂ vs. MgO, Mg# and TiO₂ (after Wang et al., 2006); (e) La vs.
 1234 La/Yb (Gao et al., 2007); (f) molar Al₂O₃/(MgO+FeO_T) vs. molar CaO/(MgO+FeO_T)
 1235 (after Altherr et al., 2000); (g) (Na₂O+K₂O)/(FeO_T+MgO+TiO₂) vs.
 1236 Na₂O+K₂O+FeO_T+MgO+TiO₂ (after Patiño Douce, 1999); (h) K₂O/Na₂O vs. Al₂O₃
 1237 diagrams (after Kamei et al., 2009); (i) La_N/Yb_N vs. Dy_N/Yb_N.

1238



1239

1240 Figure 14. The fields of zircon compositions used as discriminants for different rock
 1241 types (after Belousova et al., 2002). ‘Granitoids’ include: 1 aplites and leucogranites;
 1242 2 granites; 3 granodiorites and tonalities

1243

1244