

Regional centroid MT inversion of small to moderate earthquakes in the Alps using the dense AlpArray seismic network: challenges and seismotectonic insights

Gesa Maria Petersen^{1,2}, Simone Cesca¹, Sebastian Heimann¹, Peter Niemz^{1,2}, Torsten Dahm^{1,2}, Daniela Kühn^{1,3}, Jörn Kummerow⁴, Thomas Plenefisch⁵, and AlpArray Working Group⁺

¹GeoForschungsZentrum Potsdam, Potsdam, Germany

²University of Potsdam, Institute of Geosciences, Potsdam, Germany

³NORSAR, Applied Seismology, Kjeller, Norway

⁴Freie Universität Berlin, Berlin, Germany

⁵Federal Institute for Geosciences and Natural Resources (BGR), Hanover, Germany

⁺For further information regarding the team, please visit the link which appears at the end of the paper.

Correspondence: Gesa M. Petersen (gesap@gfz-potsdam.de)

Abstract. The Alpine mountains in central Europe are characterized by a heterogeneous crust accumulating different tectonic units and blocks in close proximity to sedimentary foreland basins. Centroid moment tensor inversion provides insight into the faulting mechanisms of earthquakes and related tectonic processes, but is significantly aggravated in such an environment. Thanks to the dense AlpArray seismic network and our flexible bootstrap-based inversion tool *Grond* we are able to test different set-ups with respect to the uncertainties of the obtained moment tensors and centroid locations. We evaluate the influence of frequency bands, azimuthal gaps, input data types and distance ranges and study the occurrence and reliability of non-DC components. We infer that for most earthquakes ($M_w \geq 3.3$) a combination of time domain full waveforms and frequency domain amplitude spectra in a frequency band of 0.02-0.07 Hz is suitable. Relying on the results of our methodological tests, we perform deviatoric MT inversions for events with $M_w > 3.0$. Here, we present 75 solutions for earthquakes between 01/2016 and 12/2019 and analyse our results in the seismo-tectonic context of historical earthquakes, seismic activity of the last three decades and GNSS deformation data. We study regions of comparably high seismic activity during the last decades, namely the Western Alps, the region around Lake Garda, the eastern SouthernSE Alps, besides clusters further from the study region, in the Northern Dinarides and the Apennines. Seismicity is particularly low in the Eastern Alps and in parts of the Central Alps. We apply a clustering algorithm to focal mechanisms, considering additional focal mechanisms from existing catalogs. Related to the NS compressional regime, E-W to ENE-WSW striking thrust faulting is mainly observed in the Friuli area in the eastern SouthernSE Alps. Strike-slip faulting with a similarly oriented pressure axis is observed along the northern margin of the Central Alps and in the Northern Dinarides. NW-SE striking normal faulting is observed in the NW Alps showing a similar strike direction as normal faulting earthquakes in the Apennines. Both, our centroid depths as well as hypocentral depths in existing catalogs indicate that Alpine seismicity is predominantly very shallow; about 80% of the studied events have depths shallower than 10 km.

1 Introduction

The Alpine mountains and surrounding areas are known for their complex tectonic setting with a highly heterogeneous lithospheric structure (e.g. Handy et al., 2010, 2015; Schmid et al., 2004; Hetényi et al., 2018). The mountain range was tectonically shaped by the interaction of the Adriatic microplate and the European microplates plate in several stages of compressionconvergence between the two large converging plates, Europe and Africa (e.g. Schmid et al., 2004, 2008; Handy et al., 2010; Hetényi et al., 2018). Geological studies show that the Adriatic plate was upthrust is the upper plate in the subduction of the Alpine Tethys in the western and central Alps, while it is the lower plate of the thrust systems in the Apennines and the Dinarides (e.g. Schmid et al., 2008; Handy et al., 2015). The terranes of the Mesozoic Tethys ocean between Europe and Africa were compressed, rotated, faulted and stacked during the compression stages of the Alpine orogenesis (e.g. Handy et al., 2010). Along a distance of approximately 700 km between NW Italy and Slovenia, the Northern Alps and the Southern Alps are separated by the Periadriatic line or fault system (e.g. Handy et al., 2005, see also Fig. 1). Reversals in the subduction polarities have been proposed at the transition to both; the Apennines and the Dinarides, while the geometry and orientation of the slab is still controversial (e.g. Hetényi et al., 2018; Handy et al., 2010, 2015; Mitterbauer et al., 2011; Schmid et al., 2004). GPS measurements show that the Adriatic microplate is rotating counter-clockwise relative to Europe around an Euler pole located in the Western Alps or western Po plain (D'Agostino et al., 2008; Weber et al., 2010). Velocity anomalies in the crust and the upper mantle reflect the complexity of the crustal structure and the geodynamic setting (e.g. Diehl et al., 2009; Fry et al., 2010; Molinari et al., 2015; Kästle et al., 2018; Lu et al., 2020; Qorbani et al., 2020).

Seismic activity across the Alps is typically characterized by low to moderate magnitude earthquakes. However, large damaging earthquakes have occurred in the past, such as the 1356 Basel earthquake (Meyer et al., 1994) or the 1976 Friuli earthquake (Mw 6.45, Poli and Zanferrari, 2018). Recent seismic activity in the SE eastern Southern Alps is caused by the N-S convergence ($2-3 \text{ mm yr}^{-1}$) between the Adriatic Plate and Eurasia, which is accommodated by the ENE trending, SSE verging thrust front of the eastern Alps and by NW-SE trending right-lateral Dinaric strike-slip fault systems in western Slovenia (Moulin et al., 2016; Poli and Zanferrari, 2018). The wider Alpine region including parts of the Dinarides and the Apennines stretches across Switzerland, Austria, Liechtenstein, France, Italy, Germany, Slovenia, and Croatia. Most of these countries have national earthquake observatories, research institutes or universities that routinely monitor the regional seismicity. The Swiss Seismological Service (SED) and the Slovenian Environment Agency (ARSO) provide annual reports containing mainly first motion based focal mechanisms (e.g. Diehl et al., 2018; Ministrstvo za okolje in prostor Agencija RS za okolje, 2020) while for example INGV (Italy), GEOFON (Germany), EM-RCMT (European-Mediterranean Regional Centroid-Moment Tensors; Pondrelli, 2002), SISMOAZUR (France) as well as gGCMT (Lamont-Doherty Earth Observatory of Columbia University, USA) provide moment tensor (MT) solutions in online bulletins for magnitudes above 3.5 or larger (see also data and code availability).




Figure 1. Study area and AlpArray seismic network. Subregions that are discussed in greater detail in this study are numbered. **Arrows indicate average Neogene (20-0 Ma) Adria-Europe convergence rates from Le Breton et al. (2017).** Exposed and subsurface faults simplified from Schmid et al. (2004, 2008); Handy et al. (2010, 2015) and Patacca et al. (2008). Topographic data from SRTM-3 (Farr et al., 2007) and ETOPO1 (Amante and Eakins, 2009) datasets). **Inset: Location of the study area in Central Europe (red rectangle).** FR: France, DE: Germany, IT: Italy, SW: Switzerland, AT: Austria, SL: Slovenia, HR: Croatia.

The region can be characterised by compartments with varying tectonic movement in close proximity, as described by many studies of local seismic activity. Focal mechanism in the SW Alps indicate predominantly N-S to NNW-SSE striking normal faulting (e.g. Nicolas et al., 1998; Sue et al., 2000), while in the W Alps strike-slip earthquakes have been observed and explained as a consequence of regional NW-SE compression and NE-SW extension (Maurer et al., 1997). **A rotation of the dominant tensional axis over the bending western Alps was recently described by Mathey et al. (2020). The Albstadt Shear Zone in the NW foreland of the Alps is dominated by NNE-SSW striking sinistral strike-slip events (Mader et al., 2020).** In the central Alps, Marschall et al. (2013) observe strike-slip faulting in central Switzerland. NW-SE striking normal faulting is reported for SE Switzerland (e.g. Marschall et al., 2013; Diehl et al., 2018). Reiter et al. (2018) provide focal mechanism solutions from P and S polarities and amplitude ratios for the **Central to Eastern central to eastern northern Alps.** They report strike-slip mechanisms and oblique strike-slip mechanisms in the Brenner-Inntal transfer zone (see Brenner and Inntal fault in Fig. 1), and normal faulting is seen with a strike direction parallel to the Giudicarie fault system. Within this fault system, EW

to NE-SW striking thrust faulting with strike-slip components were described by Viganò et al. (2008). The Italian MT data set provides extensive mechanisms for N Italy. Within the Lake Garda region and in the eastern Southern (eS) Alps close to Friuli thrust faulting with ENE-WSW to ESE-WNW strike direction is dominant (Pondrelli et al., 2006; Anselmi et al., 2011; Bressan et al., 1998). East of Friuli and in the Northern Dinarides, both, (oblique) thrust and strike-slip faulting is observed (Pondrelli et al., 2006; Moulin et al., 2016). This overview is by far not complete, but provides a small glimpse into the complex seismic and tectonic activity that is not simply dominated by the main active deformation fronts at the southern and northern margin of the Alps, but also occurs in smaller fault systems across the entire region.

To study the orogenesis of the Alps and related processes like recent seismic activity, mantle dynamics, plate motion and surface processes, the AlpArray initiative was established. In this initiative, more than 35 European institutes joined resources to operate the AlpArray Seismic Network (AASN) (Hetényi et al., 2018), consisting of more than 600 temporary (AlpArray Seismic Network, 2015) and regional permanent stations with an average spacing of <60 km (Fig. 1). For comparison, in summer 2011, before the AASN planning period started and first additional permanent stations were set up, there were 234 stations in the same area (Hetényi et al., 2018). The AASN is complemented by the dense Swath-D network in the eastern Alps (Heit et al., 2017). The permanent stations of the AlpArray are part of existing European regional networks (RD, GU, CZ, ST, G, CH, OE, MN, HU, GE, RF, FR, IV, BW, SX, NI, TH, OX, see data and code availability). The dense AASN allows studying regional seismicity in new, greater detail and provides the opportunity to perform **centroid moment tensor (CMT)** inversions with a constant station coverage over the entire region. In contrast, many of the previous studies focus on specific regions or seismic sequences within the Alps, therefore not providing a broad overview. Furthermore, many of these studies relied on first motion polarities. First-motion based approaches can be used even for small earthquakes when no surface wave energy is observed. However, the obtained mechanism is only representative for the very first moment of the fracturing process. ~~Therefore, complex rupture processes cannot be represented.~~ This might introduce discrepancies when comparing first motion solutions to MT solutions (Scott and Kanamori, 1985; Guilhem et al., 2014). **The instability of take-off angles of shallow earthquakes may introduce significant errors in the polarity readings (Hardebeck and Shearer, 2002).** Additionally, first motion solutions of small earthquakes are often only based on few **data-points polarities**, which makes it difficult to assess uncertainties.

Despite the limited resolution, first motion polarity approaches are often used in the Alps, where MT inversion is particularly challenging. First, earthquake magnitudes are generally small to moderate, requiring waveform modeling at relatively high frequencies and local distances. Furthermore, structural heterogeneities, site effects and topographic effects hinder full waveform MT inversions based on 1-D velocity profiles when considering frequencies above 0.1 Hz. Signal-to-noise ratios (SNR) vary significantly across the region due to densely populated areas and environmental conditions (weather and wind, rivers, rock falls, avalanches, etc.). Each study or observatory reporting focal mechanisms uses different inversion tools, input data, as well as distance and frequency ranges. Furthermore, uncertainties are not routinely discussed, which makes it difficult to evaluate published solutions. Uncertainties can be assessed e.g. by performing a grid search over parameters like strike, dip, rake and depth (Stich et al., 2003; Cesca et al., 2010) or using independent bootstrap chains in the inversions with varying weighting of

the input data from different stations (e.g. Heimann et al., 2018; Kühn et al., 2020, and this study). The latter approach provides
100 uncertainties of all inversion parameters (e.g. MT components, centroid location) and helps to identify trade-offs between these
parameters.

Studies of focal mechanisms in the Alps have mainly considered DC solutions. Here, based on the dense seismic network,
we also attempt to consider non-DC components. The decomposition of the moment tensor allows studying the seismic source
in more detail, including not only pure tectonic dislocations represented by the DC, but also volumetric changes and tensile
105 faulting (e.g. Vavryčuk, 2015).

In this study, we predominately target small to moderate earthquakes ($M_w > 3.0$) that occurred in the Alps or surrounding
areas between 01/2016 and 12/2019, based on the operation time of most of the temporary broadband stations of the AASN.
While the temporarily densified network provides a great amount of input data to the single MT inversions, the short time span
limits the number of observed earthquakes. We test various set-ups and input types to establish work flows for a homogeneous
110 and consistent list of MT solutions for the Alps (see Supplement). We attempt to lower the magnitude threshold for inversions
compared to routinely reported solutions by optimizing the used methods and by combining different input data types (e.g. time
domain full waveforms and frequency domain amplitude spectra). Using the AASN and the bootstrap inversion framework
Grond (Heimann et al., 2018; Dahm et al., 2018; Kühn et al., 2020) allows determining most suitable set-ups for source
inversions of small to moderate earthquakes within the study area.

115 After an introduction into the inversion method, we describe the methodological tests that were performed to assess method-
ological inherent uncertainties. At the same time, we propose guidelines for MT inversion of small to moderate earthquakes
in complex tectonic settings. Subsequently, we present the MT solutions that were obtained for the Alpine region and dis-
cuss these with respect to mechanisms, spatial patterns and centroid depths. We discuss different tectonic areas in the Alps
systematically, including observations of seismicity, faulting mechanisms and GNSS deformation data.

120 2 Methodology

2.1 Moment tensor inversion using *Grond*

We use the open source software *Grond* for MT inversions (Heimann et al., 2018; Kühn et al., 2020). In a Bayesian bootstrap-
based probabilistic joint inversion scheme, solution uncertainties are retrieved along with the best fitting CMT solution (see
also Dahm et al., 2018). We invert for the six independent moment tensor components, the seismic moment, the centroid
125 location and the origin time simultaneously. The objective function is set up in a flexible way to combine different input data
types as a weighted sum. In our study, we use combinations of time domain full waveforms, time domain cross-correlations
and frequency domain amplitude spectra as an input for the inversion. Following the studies of Zahradník and Sokos (2018)
and Dahal and Ebel (2020), we implemented envelopes of time domain waveforms. The misfits of the different input data are
combined using an L1 or L2 norm. We assign the same weighting to each input data type. The misfit values of the single stations
130 within one input data type group are weighted to account for different epicentral distances (Heimann, 2011). Without applying
such a weighting, summed misfits are always dominated by the closest stations which have the highest amplitudes. In the case

of using time domain full waveforms or cross-correlation fitting procedures, we allow for small time shifts to compensate for errors in the velocity models. To avoid the mis-matching of phases, these time shifts were set to be well below a quarter of the dominant wavelength. Time shifts are regulated by a penalty function with an empirically chosen maximum of 0.05.

135 Precalculated Green's function data bases are used for rapidly computing synthetic data (Heimann et al., 2019). In our case, we used regional velocity profiles from the CRUST2.0 Earth model database (see <http://igppweb.ucsd.edu/~gabi/crust2.html>, last access June 2020; Bassin et al., 2000), which we choose according to the earthquake epicenter location. The Green's function databases were calculated with the orthonormal propagator algorithm QSEIS (Wang, 1991). *Grond* selects the appropriate time window, corrects the recorded waveforms for the instrument response and rotates to ZRT coordinate system. Filters are applied and, if specified, waveform attributes as spectra or envelopes are calculated. The inversion is performed in parallel
140 bootstrap chains (here: 100 for normal inversions, 500 for method testing), where individual bootstrap weights are applied to the single station-component-input data type combinations. Bootstrapping is applied for two reasons. Firstly, to avoid distortions due to few high misfit values resulting from a low SNR at single stations, wrong transfer functions or malfunctioning stations. ~~Further, Secondly,~~ the bootstrap chains are used to access model parameter uncertainties and trade-offs between the
145 inversion parameters. Each bootstrap chain performs an entirely independent optimization. Along with the best solution with the lowest misfit, *Grond* provides a defined number (here: 10) of best solutions of each bootstrap chain, which we call the ensemble of solutions.

Fig. 2 presents a selection of plots provided by the inversion software to assess the solution robustness, in this case for the 2019-06-14 thrust faulting earthquake in northern Italy (Mw 3.9). Fig. 2a shows the fuzzy MT, which is an illustration of
150 the MT uncertainty. It is composed of the superimposed P radiation pattern of the ensemble of solutions from the bootstrap chains. If the variability of the ensemble solutions is small and hence the uncertainties are small (as seen here), the fuzzy plot has clearly separated black and white quadrants. The red lines indicate the solution with the lowest misfit. Other plots show the station distribution (Fig. 2b) and the decomposition of the best deviatoric MT solution into the DC and the CLVD component (Fig. 2c). Fig. 2d shows the distribution of the centroid locations obtained from all ensemble solutions and Fig. 2e
155 depicts the resolution of the six independent MT components in form of probability density functions. As typical for shallow events inverted using surface waves, the m_{nd} and m_{ed} components are not as well resolved as the other components (Cesca and Heimann, 2018; Bukchin et al., 2010; Valentine and Trampert, 2012). Finally, Fig. 2f shows examples of waveform and frequency spectra fits of Z, R and T component traces at three stations. ~~Further plots, e.g. showing parameter trade-offs and the evolution of misfits and optimised parameters during the inversion, are not provided here.~~

160




Figure 2. Example of an MT inversion result, 2019-06-14, Mw 3.9, NE Italy. (a) Fuzzy beachball illustrating the MT solution uncertainty. (b) Station distribution around the epicenter. (c) Decomposition of the best MT solution into DC and CLVD component. (d) Resolution of the centroid depth, easting and northing relative to the starting position. (e) Probability density functions (PDF) showing the resolution of the six independent MT components m_{xy} normalized by the seismic moment M_0 . (f) Examples of waveform and spectral fits at three stations and three components. Red and black lines indicate synthetic and recorded waveform data, respectively. The beige shaded area represents the time window and taper function. Station name, azimuth and distance to the epicenter are indicated above each column. Numbers within the panels describe the time window and the frequency band.

The selection and joint inversion of waveform attributes can improve the stability and goodness of solutions. In the following, we want to point out advantages and drawbacks of the waveform-based input data types, which are used in the subsequent methodological tests: time domain (TD), frequency domain (FD), cross-correlation and envelopes.

165 **TD full waveform fitting:** In time domain, the misfit between a selected time window of a seismic trace and a synthetic trace in a defined frequency range is computed as the normalized sum of sample misfits. Time shifts are allowed and regulated with a penalty function. For regional MT inversion surface waves are commonly considered in full waveform approaches (e.g. Ritsema and Lay, 1995; Minson and Dreger, 2008; Sokos and Zahradnik, 2008; Dahm et al., 2018). The frequency band is magnitude dependent. While at low frequencies effects of the velocity model and topography are minor, at higher frequencies
170 the SNR is usually better. At regional distances, often magnitude-dependent frequency bands below 0.1 Hz are used to consider Rayleigh waves and Love waves, which have particularly simple waveforms at ~~this distance range regional-distanees~~ (Ritsema and Lay, 1995). However, in ~~the~~ case of very shallow sources, the resolution of the m_{xz} and m_{yz} components of the MT are limited when using surface waves (Bukchin et al., 2010; Valentine and Trampert, 2012; Cesca and Heimann, 2018, see also Fig. 2). Relying on time domain fitting only, time shifts, noisy data or distorted amplitudes can hinder finding stable initial
175 inversion solutions. It has proven to be helpful to combine time domain full waveform fitting with other input data types like frequency domain amplitude spectra which are often less affected by these issues.

FD amplitude spectra fitting: Real valued amplitude spectra of recorded Love and Rayleigh waves carry all information necessary to resolve the geometry of the MT, while neglecting the phase information and dispersion (e.g. Mendiguren, 1977).
180 This means ~~that~~ two MT solutions with common nodal planes but opposite polarities model the amplitude spectra equally well and additional information from first-motion polarity readings or time domain waveform fitting is needed to resolve this ambiguity (e.g. Cesca et al., 2010; Heimann, 2011). The misfit is computed as the misfit between amplitude spectra of recorded and synthetic waveforms in a selected frequency range and time window. Compared to fitting full waveforms in time domain, more conservative, less exact time windows can be selected. Cesca et al. (2010, 2013) propose a multistep approach to
185 stepwise combine the fitting of amplitude spectra and displacement waveforms to subsequently obtain point source parameters, the centroid location and kinematic source parameters. Compared to full waveform fitting, amplitude spectra inversion methods are less sensitive to trace misalignments and phase shifts resulting from coarse or erroneous velocity models (Cesca et al., 2010, 2013; Domingues et al., 2013). In the subsequent tests, we do not use a step-wise inversion, but use amplitude spectra and time domain full waveforms or cross-correlated waveforms simultaneously.

190

Cross-correlation waveform fitting: In the cross-correlation based fitting of full waveforms, the amplitudes of recorded and synthetic traces are normalized. The inversion searches for the maximum cross-correlation value for the selected time window in time domain, basically fitting the phase shift (Stähler and Sigloch, 2014; Kühn et al., 2020). Cross-correlations help to constrain the centroid location and centroid time in a joint inversion. We allow for small time shifts, regulated with a penalty
195 function, to compensate for imprecise velocity models. Time shifts need to be small compared to the frequency range in order to

avoid mismatching phases. Due to the amplitude normalization, this method is sensitive **for**to patterns in the waveforms, while it is not influenced by gain errors or site effects. Magnitudes cannot be resolved. Cabieces et al. (2020) used cross-correlation fitting in their MT inversion for ocean bottom stations, where absolute amplitudes could not be modelled due to the unknown coupling to the ground. When using cross-correlations to fit time domain waveforms, **both**; frequency bands as well as time
200 windows need to be selected carefully to avoid mismatching phases.

Envelope fitting: In our study, we compute the waveform envelopes by convolving the squared time series using the fast Fourier transform with a Hanning taper. This smoothens and therefore simplifies the waveforms. Hensch et al. (2019) used non-smoothed envelopes in the same inversion routine in combination with amplitude spectra, spectral ratios and time domain
205 waveforms. Envelopes, especially smoothed ones, are less influenced by small unmodelled time shifts or noisy data compared to full waveforms. Fitting envelopes of seismic waveforms can be helpful in **the** case of using high frequency bands, simplified velocity models and increased noise levels. Zahradník and Sokos (2018) stress that due to the simplification of the waveforms, the results of envelope-based inversions have a limited precision and results need an even more careful inspection of uncertainties and resolution. However, if body waves are considered, envelopes can especially help to constrain P and S phase arrivals,
210 and thus the centroid time and location. Since the envelopes are based on absolute amplitudes, they need to be combined with a method providing polarity information. Zahradník and Sokos (2018) and Dahal and Ebel (2020) have shown that envelopes can be used to derive focal mechanisms for $M < 4$ events in **the** case of unfavorable settings like sparse networks, for which full waveform fitting is not feasible. In both studies, the envelopes are combined with P-polarities of one or more nearby stations to resolve the polarity ambiguity.

215

2.2 Methodological tests

We perform methodological tests using recorded seismograms and synthetic data to investigate the resolution capacities, requirements and limits of MT inversions in the Alpine region. We use subsets of representative earthquakes that occurred between 2016 and 2019 in the Alps. The tests are particularly computationally demanding, as every single inversion of each
220 test is run in 500 bootstrap chains. The number of events in each test depends on the number of tested parameters. The proposed tests can be used as a guideline for assessing the feasibility of MT inversions in other study areas with moderate seismicity.

We benefit from the large seismic network and use more than 80 stations at distances of up to 400 km for the largest events (Fig. 3). For earthquakes with moderate magnitudes between M_w 3.5-3.9 we mostly rely on 20 to 50 stations within a radius of 200 km. The number of available stations depends on the magnitude, but also on the epicenter location within the network.
225 **Furthermore, the SNR and quality of the individual stations is variable in time and space. Before the inversions, we applied the toolbox *AutoStatsQ* to identify seismic stations with misorientations, metadata errors or gain problems (Petersen et al., 2019).**




Figure 3. Vertical component seismograms of permanent AASN stations, sorted by distance. (left) 2017-03-06, Mw 4.1, Switzerland, (right) 2017-10-27, Mw 3.6, France. Time in seconds after origin time. Waveforms bandpass filtered between 0.02 and 0.07 Hz. Rayleigh waves dominate the seismograms in this frequency range with an average phase velocity of 3 km/s (green line).

2.2.1 Double-couple, deviatoric and full moment tensor inversions

We study the stability and resolvability of non-DC components by performing full, deviatoric and pure-DC MT inversions for a subset of 32 **representative** earthquakes of the AlpArray data set. **We consider these earthquakes representative as they cover a large magnitude range (Mw 3.2-4.2), are distributed across the entire study area and comprise of different types of mechanisms.** We use a passband of 0.02-0.07 Hz and fit time domain full waveforms and frequency domain amplitude spectra simultaneously. We compared the pure-DC, the deviatoric and the full MT obtained for each earthquake with respect to the fit of the recorded data and to the uncertainties of MT components and centroid locations. Subsequently, we statistically evaluate those solutions, for which a low misfit between synthetic and observed waveforms was achieved with all three inversion types.

235 Four events with Kagan angles $> 60^\circ$ between full and deviatoric solutions were removed, since they were not well resolved.

The isotropic source component **of the seismic moment tensor** resolves volumetric changes, including processes **aslike** explosions, cavity collapses, fluid movement or ruptures on nonplanar faults (e.g. Sileny and Hofstetter, 2002; Ford et al., 2009; Miller et al., 1998; Minson and Dreger, 2008). The CLVD component is often described as the residual radiation to the best DC without geological interpretation (Dahm and Krüger, 2014), it is however required for a mathematically complete

240 decomposition (Vavryčuk, 2015). Large CLVD components are often explained by noisy data, a simplified or incorrect velocity model, neglected 3-D wave effects or insufficient station coverage (e.g. Panza and Saraò, 2000; Cesca et al., 2006), but can also be interpreted physically in combination with an isotropic component of the same sign as a product of tensile faulting

(Vavryčuk, 2015). Non-DC components are also used as an indicator of anthropogenic seismicity (Dahm et al., 2013; Cesca et al., 2013; Lizurek, 2017).

245 Despite frequent geological interpretations which propose fluid movements or tensile processes, various studies show that resolving non-DC components in MT inversions is particularly difficult. Seismic noise and inaccurate Green's functions may result in large non-DC components. Trade-offs between hypocenter location or depth and isotropic component have been observed (e.g. Dufumier and Rivera, 1997; Panza and Saraò, 2000; Křížová et al., 2013; Kühn et al., 2020). Non-DC components must therefore be evaluated carefully with respect to tectonic processes (Lizurek, 2017). **Methodological tests based on observed data as well as synthetic tests** can help to identify which non-DC components can be considered statistically significant
 250 (see also Panza and Saraò, 2000).

Fig. 4a and b show the ratio of DC and non-DC components of the full and deviatoric MT inversions **of our test dataset**. The deviatoric inversions result in 0-40% CLVD components for 70% of the events and in CLVD components of >50% for
 255 19% of the events. In **the** case of full MT inversions, we find significant isotropic components of >30% in **the** case of one third of the earthquakes. Fig. 4c indicates that the non-CLVD components of the test events scatter significantly. It is clearly visible that many events with shallow depths (dark colors) are located in the upper right and lower left quadrants of the Hudson plot, indicating isotropic and CLVD components of opposite signs. Cesca and Heimann (2018) showed that for shallow depths, isotropic and CLVD components often appear indistinguishable. Further below, we discuss this observation comparing forward
 260 calculated synthetic waveforms for one example event.




Figure 4. Histograms showing the decomposition of the full (a) and deviatoric (b) MT inversion results into isotropic (ISO), compensated linear vertical dipole (CLVD) and double couple (DC). Each bar of 10% width (x-axis) indicates for how many test earthquakes (y-axis) the proportion of decomposition is found. For example, an isotropic component of 10-20% is found for 13 test events in the full MT inversion. (c) Hudson plot showing non-DC components of individual events. Four events with Kagan angles $> 60^\circ$ between full and deviatoric solutions were removed, since they were not well resolved. Color represents depth, beachball size magnitude.

The DC component is representing the purely tectonic shear dislocation (e.g. Miller et al., 1998; Julian et al., 1998; Cesca et al., 2013), therefore, it is crucial to resolve this component unambiguously. We compare the DC component that we obtain from the decomposition of the deviatoric and the full MT with the pure-DC inversion result by computing the smallest rotation

angle (Kagan angle, Kagan, 1991) between them to assess the stability of the DC components (Fig. 5). Rotations below 30° are generally accepted as representing very similar mechanisms, while a Kagan angle $\ll 60^\circ$ is still described as corresponding (Pondrelli et al., 2006; d'Amico et al., 2011). 70% of the earthquakes have a very stable DC, with Kagan angles below 30° between the three solutions. In the case of about 10% of the events, a Kagan angle $\geq 60^\circ$ is found. These larger deviations result predominantly from large non-DC components in the full inversion result (in $>70\%$ of these events). In these cases, the CLVD combined with the isotropic component shows orientations similar to the DC component of the pure-DC and the deviatoric inversion result. Therefore, the resulting focal sphere is similar, while the DC component deviates from the pure-DC inversion result. Overall, the results of this test indicate that the DC component is in most cases very well resolved, independently of allowing for a CLVD and an isotropic component.




Figure 5. (left) Kagan angle between DC of pure-double-couple, deviatoric and full MT solutions for 30 earthquakes, Mw 3.3-4.5. The dashed and solid lines indicate Kagan angles of 30° and 60° , respectively, indicating levels of high agreement and corresponding mechanisms. (right) Histograms showing the distribution of Kagan angles between the DC of pure-DC, deviatoric and full MT inversion solutions.

In the following, we present a more detailed analysis of an exemplary earthquake with a significant non-DC components: 2019-05-28, Mw 3.9, close to Lake Geneva, France (Fig. 6a). Fig. 6b depicts the MT decompositions of the MTs obtained with a pure-DC, a deviatoric and a full MT inversion. All three inversions were performed using the same inversion set-up (full waveforms and amplitude spectra, Z, R and T components, 73 stations, 0.02-0.07 Hz). The DC component is similar for all inversion types, but the deviatoric and full inversion results indicate significant non-DC components.

To investigate whether the resolved non-DC components are unambiguous, we forward model synthetic waveforms of the three MT solutions recorded at fictional receivers in 250 km distance in azimuthal steps of 1° (Fig. 6a). We use a bandpass-filter of 0.02-0.1 Hz, which is even wider than the frequency range used in the MT inversion (0.02-0.07 Hz) to assess the similarity of the entire modelled surface wave trains. By cross-correlating the forward modelled waveforms, we find that the full and deviatoric sources produce very similar waveforms on all seismometer components in all backazimuthal directions (Fig. 6d).

Furthermore, the maximum amplitudes between deviatoric and full solution differ only slightly. This indicates that the non-DC component can be comparably well represented by a CLVD or by a combination of an isotropic plus a CLVD component.

285 A comparison of the forward modelled waveforms from a pure-DC solution with a full or deviatoric solution shows very high correlations in all azimuthal directions on the T components (Fig. 6d, lower panel). Neither the CLVD nor the isotropic component is influencing the transversal Love wave. On R and Z components, the resulting waveforms show cross-correlations below 0.9 in strike-direction only (Fig. 6d, upper and middle panel). This indicates that in the case of this event, without any station covering this ray path direction, we cannot resolve the difference between a pure-DC MT and a full or deviatoric one.

290 The true azimuthal coverage of seismic stations is much denser to the NE and E than in strike direction (Fig. 6a). Forward modelling the waveforms of the 73 used stations, results in a similar, but less well resolved pattern compared to Fig. 6d. The uneven azimuthal distribution and the lack of stations in strike directions hinders the unambiguous identification of non-DC components.

This example shows that whenever we investigate large non-DC components in a deviatoric or full MT inversion, we have to
295 ~~asses carefully teh~~ one must assess the resolution and the validity of the results. The synthetic tests indicate that including full waveforms of body waves at higher frequencies in the inversion clearly helps to improve the resolution of non-DC components. However, due to the station spacing, the relatively high noise level and the low resolution of crustal velocity models, we cannot use higher frequencies for most events in this study. Following our findings, we report deviatoric MT inversions in the result section and only perform inversions for the full MT in the case of large non-DC components for comparison.




Figure 6. Earthquake close to Lake Geneva (France-Swiss border region), Mw 3.9, 2019-05-28 08:48:06. (a) Seismic network used for the MT inversions (73 broadband sensors, red triangles) and synthetic network (circle of blue dots, in 1° steps). (b) Decomposition of the full (purple), deviatoric (orange) and pure-DC (green) MT inversion solution with the smallest misfit. (c) Hudson plots showing the ensemble of solutions for the full and the deviatoric MT inversions, colors as in (b). Larger symbols depict best solutions of b. (d) For the three MT solution (full, deviatoric and DC) synthetic data were forward computed for the fictional network shown in (a). For each component (Z,R,T), the first row shows the maximum cross-correlation values of the three synthetic traces at each station (BP filter 0.02-0.1 Hz). The second row shows the maximum absolute amplitude at each backazimuth, normalized over the three solutions. The dashed lines indicate the strike 1 and 2 directions of the two nodal planes of full, deviatoric and DC solution and their 180° equivalent.

300 2.2.2 Magnitude-distance relation

The distance range in which stations can be used for MT inversion strongly depends on the event magnitude. While we use an epicentral radius of less than 100 km for the smallest events with magnitudes Mw 3.1-3.3, epicentral distances may be as large as 300-400 km for the largest events with magnitudes greater Mw 4.0. For events between Mw 3.5-3.7 and Mw 3.8-3.9,

we use distances of up to 160 km and 200 km, respectively. This results in different inversion set-ups: the number of available stations varies between less than 10 stations for the Mw 3 events to above 80 stations for the largest events. Fig. 3 illustrates this relation. The left panel shows waveforms of an Mw 4.1 event in Switzerland at distances of up to 350 km. Even though the SNR decreases with distance, a distinct Rayleigh wave can be seen. The second event from 2017-10-27, France, has a magnitude of Mw 3.6. For distances larger than 160 km, SNR are very low for most stations. We did not remove stations with generally high noise levels from the plot to illustrate that a careful rejection of very noisy and dysfunctional stations is required. We apply the toolbox *AutoStatsQ* in advance to identify seismic stations that are misoriented, have errors in their metadata or gain problems (Petersen et al., 2019).

2.2.2 Frequency ranges and input data type

In previous studies, dependencies of the MT inversion results on the inverted frequency band have been observed and multi-step inversion workflows including several frequency bands were proposed (e.g. Barth et al., 2007). In order to find the best combination of frequency ranges and time domain or frequency domain input types for the MT inversion, we selected a subgroup of 13 earthquakes recorded by the AlpArray stations. These test events span a magnitude range of Mw 3.3 to 4.1 and are therefore considered representative. We perform MT inversions using eight different combinations of input data types (Fig. 7 and Supplement Fig. 1): time domain full waveforms (td), frequency domain amplitude spectra (fd), cross-correlations of time domain full waveforms (cc), waveform envelopes and combinations thereof. The input data is filtered using nine different bandpass filters with passbands between 0.01 and 0.7 Hz (Fig. 7 and Supplement Fig. S1). We compare the uncertainties of the resolved MTs in order to find the most appropriate parameter settings for our study and future MT studies in the Alps or similar settings.




Figure 7. Testing of input data types and frequency ranges for three earthquakes with magnitudes between Mw 3.6 to 4.1. Abbreviations: td - time domain full waveforms; fd - frequency domain amplitude spectra; cc - cross-correlation fitting of full waveforms; tdfd, tdcc, fdcc and tdfdcc are combinations of these. Color intensity represents summed uncertainty of MT components. A combination of fd and td in a frequency band between 0.02-0.07 Hz yields the best results for Mw > 3.3 (marked in green).

In Figures 7, we show the results for three exemplary events. The color intensity of each focal sphere represents the summed standard deviations of the six MT components derived from the ensemble of solutions of the bootstrap chains. Intense colors represent stable solutions with low uncertainties. The first event is a Mw 4.1 earthquake in Switzerland (Fig-7). Due to the high magnitude, the MT inversion results are stable over frequency bands ranging from 0.01-0.03 Hz up to 0.01-0.10 Hz for all input data types. The MT is not well resolved when filtering using a passband of 0.03-0.1 Hz or higher. A similar behaviour is observed for the second example, a Mw 3.9 normal faulting event close to Lake Geneva. The MT is very well resolved using bandpass filters covering the intermediate passbands between 0.02 and 0.1 Hz. In contrast to the first event and corresponding to the lower magnitude, the resolution is worse when frequencies between 0.01-0.02 Hz are included, while the frequency band between 0.03-0.1 Hz still leads to satisfying results. In general, the higher the frequency band, the lower the stability of the ensemble of solutions due to the simplified 1-D velocity model, site effects and increased noise levels.

In the case of the third example (Fig. 7), a Mw 3.6 earthquake from France, the MT solutions vary substantially. This illustrates the need for a careful selection of appropriate methods and frequency ranges and the analysis of the uncertainties of MT inversions. For both the higher frequency ranges (from 0.03-0.1 Hz and higher) as well as in the lowest frequency bands (0.01-0.03 Hz and 0.02-0.05 Hz) surface waves have insufficient SNRs. Stable results for most input types are obtained in the frequency band 0.02-0.07 Hz, in which surface waves are more distinct. A visual inspection of the recorded waveforms of various events with magnitudes of Mw 3.4-3.9 confirms that surface waves have highest SNRs for periods between 0.02 and 0.05 Hz. Extending the passband to 0.02-0.07 Hz helps to avoid mismatching monochrome phases in the inversion process.

Comparing the different input data types for all thirteen test events, we find that a combination of frequency domain amplitude spectra and time domain full waveform fitting (tdfd in Fig. 7) provides more stable results than relying on time domain waveform fitting alone. The high uncertainties of the frequency domain amplitude spectra fitting alone (fd) result from the unresolved polarity. The geometry of the nodal planes can still be determined. For most events, the other combinations (tdcc, tdfccc, fdcc) provide more stable results compared to using only time domain full waveforms (td). However, compared to the tdfd combination, they do not further improve the stability of the solution.

In addition to the presented tests of input data types, we tested waveform envelopes (Supplement Fig. S1). In order to resolve the polarity of the mechanisms, the envelopes are combined with time domain full waveforms or cross-correlation fitting of waveforms at close-by stations. This is a reasonable setting for weak events, where full waveforms may be of such low amplitudes that they can only be fitted at closer stations while envelopes of more distant stations may still be of use. We find that in the case of intermediate or large magnitude test events ($M_w \geq 3.6$), the resulting MT is well recovered, although uncertainties are larger than with a time domain-frequency domain combination. In the case of smaller events, where time domain-frequency domain combinations might fail, the envelopes may stabilize the inversion. The applicability of the combination of envelopes and close-by time domain traces depends on the data quality of the closest stations and on the careful selection of the frequency range and the smoothing of the waveform envelopes.

Following the results of our methodological tests, we routinely use a combination of frequency domain amplitude spectra and time domain full waveform fitting in a frequency band of 0.02-0.07 Hz for earthquakes with $M_w > 3.5$. In the case of smaller magnitude earthquakes, we additionally perform inversions using a frequency range of 0.03-0.10 Hz. We observe that in the case of low magnitude earthquakes the initial local magnitudes can differ significantly from our moment magnitude estimates. Furthermore, the availability of stations with a good SNR does not only depend on the event magnitude, but also on noise conditions and damping along the travel path. It is therefore necessary to adapt the approach to the individual earthquakes, but the two frequency ranges constitute reasonable guidelines.

2.2.3 Station coverage

The dense AASN provides an excellent azimuthal distribution of seismic stations for moderate to large earthquakes in the Alps. We take advantage of the large number of stations in the AASN to investigate how the stability of the deviatoric MT inversion is influenced by gaps in the azimuthal station distribution around an earthquake. This allows simulating uncertainties

of MT solutions in the marginal areas of the AASN, but the results also apply to other locations and networks (e.g. close to subduction zones). Most generally, within the AASN larger event-station distances can be taken into account for larger magnitude earthquakes and therefore both the number of stations and the azimuthal station coverage increases. In contrast, individual malfunctioning stations may already result in large azimuthal gaps for low magnitude earthquakes located within the AASN. **In theory, the DC components of a moment tensor can be resolved from a single station using 3-component data (Dufumier and Cara, 1995). However, such an analysis requires high data quality and exact knowledge about velocity structures and path effects. In practice, single-station approaches are mostly avoided as they often result in unstable solutions (Dufumier and Cara, 1995).**

Fig. 8 shows the fuzzy MTs (right panels) for decreasing azimuthal coverage of seismic stations (left panels) for three exemplary events. In **the** case of the largest event (Mw 4.1), the solution is very stable when seismic stations cover at least an azimuthal range of 90° . In the case of an even smaller coverage, the mechanism rotates slightly depending on the azimuthal direction of the remaining stations. In **the** case of the Mw 3.9 event, the uncertainties of the solutions increase with decreasing station coverage. Two examples in which the inversions were done with stations covering only an azimuthal range of 45° show significant differences between the resulting focal mechanisms. When only considering the fuzziness of the two focal mechanism plots, both ensembles of solutions seem to be well resolved and stable. This indicates that the amount and variability of input data is not sufficient to resolve the MT unambiguously. **Furthermore, we observe a clear trend of increasing non-DC components with decreasing azimuthal coverage. We find a non-DC component below 10 % for a coverage of 180° , but 40 % for the smallest tested coverage for the Mw 3.9 event.**

For the smallest earthquake (Mw 3.5), the resulting MT solutions vary even more. In **the** case of the inversions with a station coverage of 90° , the variability among the ensembles of solutions is high and depends on the location of the 90° quadrant covered with stations. **When only considering the dominant DC components of the deviatoric moment tensors, we observe the same general correlation between coverage and resolution.** It is worth noticing that even with a small number of stations covering a small azimuthal range, it is possible to resolve a MT under favorable geometrical conditions. When stations are located in strike direction and cover both tensional and compressional quadrants, they may resolve the MT correctly even when covering only 45° (Fig. 8, Mw 3.5 event, 5th row and last row).

We conclude that in **the** case of larger earthquakes with a high SNR and a sufficient number of stations at different epicentral distances even a limited **a** azimuthal coverage does not necessarily pose a problem, but lower magnitude earthquakes usually require a better azimuthal station coverage. In regard to a semi-automated MT inversion workflow, we implemented an optional minimum station distribution threshold. Based on our results and since we do not assume **any a priori** known strike direction, we limit the inversions to earthquakes with an azimuthal coverage above 90° but thoroughly evaluate all results with a coverage below 180° .




Figure 8. Resolution of the deviatoric MT depending on the azimuthal station coverage for three earthquakes with magnitude Mw , centroid depth d and event-station radius r given below **each column** (left, middle and right panels). The station coverage (blue dots in 1st, 3rd, 5th column) decreases from top to bottom as indicated in between the event-columns. The fuzzy MTs show the solution stability (Sec. 2.1 and Fig. 2). They are composed of the superimposed P radiation pattern of the ensemble of solutions from the bootstrap chains.

3 Results

400 3.1 CMT solutions for the Alpine region, 2016-2019

Based on our methodological tests, we use a combination of time domain full waveforms and frequency domain amplitude spectra as input data for the centroid MT inversion for earthquakes larger than Mw 3.0. We choose a frequency range of 0.02 to 0.07 Hz for a first inversion of each event. Depending on the event magnitude, the maximum epicentral distance varies between 80 and 300 km. In the case of poor fits, we slightly increase the frequency bands (0.03-0.1 Hz for $Mw < 3.3$) for smaller events and decrease it for the larger events (0.02-0.05 Hz for $Mw > 4.2$). Deviatoric inversions were generally favored over full moment tensors, since we demonstrated that often the isotropic and CLVD components can not be distinguished reliably. In addition, no volume changes are expected to accompany small earthquakes in the seismotectonic setting of the Alps. We obtained deviatoric

MT solutions for 75 earthquakes occurring between 01/2016 and 12/2019 in the wider Alpine region, for which we determine moment magnitudes between Mw 3.1 to 4.8 (Fig. 9, Supplement Table S1). While we were able to compute stable MTs for most
 410 **Alpine** earthquakes from regional catalogs with local magnitudes larger MI 3.3, we resolved only thirteen MTs for earthquakes with local magnitudes between MI 3.1 and 3.3, corresponding to one third of the events in this magnitude range **compared to the GEOFON catalog**. Low SNR in the tested frequency bands covering frequencies between 0.02 and 0.5 Hz and less available stations hindered successful inversions for the other small earthquakes. Furthermore, we realized that a station spacing of about 60 km is not sufficient for small earthquakes (Mw<3.3) in case a part of the data is rejected due to quality issues.




Figure 9. Moment tensor inversion results from 01/2016 to 12/2019 (focal spheres with black lines) along with MTs from 1983-2015 from bulletins of **g**GCMT, GEOFON, INGV, SED, EM-RCMT and ARSO (lighter colors). Similar colors represent clusters of comparable mechanisms, **obtained from a clustering approach based on the smallest rotation between the mechanisms** (see text). **Red and orange colors correspond to dominant normal faulting mechanisms in a cluster. Thrust faulting is indicated in blue and strike-slip faulting earthquakes are colored in green and purple.** Exposed and subsurface faults simplified from Schmid et al. (2004, 2008); Handy et al. (2010, 2015) and Patacca et al. (2008). 'PL' marks the Periadriatic line, 'SK' the Split-Karlovac Fault. (right) Pressure (P), tension (T) and null (B) axis of all focal mechanism. Topographic data from SRTM-3 (Farr et al., 2007) and ETOPO1 (Amante and Eakins, 2009) datasets.

415 For about 40% of our **own** MT solutions **for 2016 to 2019**, no MT solutions were available from regional observatories (INGV, GEOFON, EM-RCMT, SISMOAZUR, SED, ARSO). For the other earthquakes, we obtain similar MT solutions, with a median deviation Kagan angle of 21° (mean 24°). For the largest events, the deviation is 15°. In the following **discussion**, we **compare jointly analyse** our results **to with** approximately 350 moment tensors of earthquakes **that occurred** before 2016,

as reported by ~~g~~GCMT, GEOFON, INGV, EM-RCMT, SED and ARSO (Fig. 9). Whenever more than one MT solution is available from the different bulletins, we prioritize local institutes (INGV for Italian earthquakes, SED for earthquakes in Switzerland, ARSO for Slovenia), unless they indicate high uncertainties. Furthermore, EM-RCMT with great experience for the Mediterranean and surrounding areas is favored over GEOFON solutions and over GCMT.

We used a clustering algorithm (Cesca, 2020) based on the Kagan angles between all focal mechanisms obtained in this study and reported in the catalogs ~~g~~CMT, GEOFON, INGV, SED, EM-RCMT and ARSO to define classes of similar mechanisms (Fig. 9). The clustering tool uses the DBSCAN clustering algorithm (Ester et al., 1996), which is relying on two parameters, the maximum acceptable similarity distance (eps) between two events, here eps=0.14, and the minimum number of neighbouring items (nmin), here nmin=6. We choose a rather large eps value to emphasize patterns of general similarity between mechanisms. In a second step, we assign all remaining earthquakes to the cluster to which they have the smallest rotation angle.

Within the Alps, we observe four dominant groups of focal mechanisms (Fig. 9). Roughly E-W striking thrust faulting is observed in the ~~south-eastern~~ eastern Southern Alps (northern Italy) and at the central ~~northern~~ southern margin of the mountain range (blue focal spheres). A group of similar strike-slip faulting earthquakes are aligned parallel to the northern deformation front of the Alps (green focal spheres, Fig. 9). A second group of strike-slip mechanisms is situated ~~SE of the Alps in the transition of the Alps to the Dinarides and in the Northern Dinarides~~ (purple and green focal spheres). NW-SE striking normal faulting events are found in the NW Alps (red focal spheres), while mechanisms are more heterogeneous in the SW Alps.

435 3.2 Distribution of centroid depths

The resolved centroid depths within the Alps range from about 2 to 15 km, pointing at a shallow seismic activity within the mountain range. 80% of the studied events have depths shallower than 10 km (~~see also Discussion 4.2.2~~). A comparison of our inverted centroid depths to the depths published in the event catalogs is limited for two reasons. First, the event depths were fixed in some of the published moment tensor inversion results, and second, the depth estimates differ significantly among the different catalogs. We find a good correspondence with less than 3 km difference for >60 % of the events to at least one of the published solutions. For another 26% we report differences between 3-5 km.

In Fig. 10, we display the depth distribution of mechanisms depicted in Fig. 9 sorted by faulting type. In the left panels, the focal mechanisms derived from the afore mentioned bulletins are shown, while our own centroid solutions are provided in the right panels. While the depth in the catalogs may be partly fixed during the inversion, the centroid depth is determined during the MT inversion in our approach. Uncertainties of our solutions are mostly in the range of 1 to 3 km. Many events within the Alpine mountain range are shallower than 10 km (Fig. 10d-f). While we obtained depths below 5 km for all normal faulting events in the NW Alps, depths of up to 15 km are observed for thrust faulting events ~~occurring between 2016 and 2019~~ in the ~~SE eastern~~ Southern Alps.

Centroid depths in the Apennines can be significantly larger. Thrust and normal faulting events are roughly separated into two NW-SE running bands, with more thrust events in the NE and more normal faulting events in the SW part. Normal faulting and strike-slip events occur predominantly ~~in~~at shallower depths (<20 km) than the thrust faulting events with depths of often 30-50 km.




Figure 10. (a)-(c) Catalog depths of earthquakes shown in Fig. 9. (Depths **might** be fixed in some catalogs.) (d)-(e) Centroid depths of MT solutions in this study. Earthquakes are sorted according to their mechanisms: (top) thrust faulting events, (center) normal faulting events and (bottom) strike-slip events. Exposed and subsurface faults (solid and dashed lines) simplified from Schmid et al. (2004, 2008); Handy et al. (2010, 2015) and Patacca et al. (2008). 'PL' marks the Periadriatic line, 'SK' the Split-Karlovac Fault. The outlines of spatial clusters of increased seismic activity from Fig. 12a are indicated for **comparison** orientation.

4 Discussion

4.1 Dominant mechanisms and the regional stress regime

455 The distribution of ~~Our~~ mechanisms coincide well with long-term seismological and tectonic observations. The moment tensor solutions obtained from small to moderate earthquakes during 4 years with an enhanced station density in the course of the AlpArray project allows identifying multiple seismo-tectonic domains. The faulting styles of these domains are in agreement with those derived from longer term moment tensor catalogs (Fig. 9). Furthermore, new solutions for regions of sparse seismicity like the northern Central Alps provide new insights into recent activity. Thrust faulting related to the NS
460 compression between the European plate and the Adriatic plate is mainly observed in the eastern Southern SE Alps, strike-slip faulting is observed in the Northern Dinarides and along the northern Alps and normal faulting is observed in the NW Alps (Fig. 9).

The orientations of P- and T- axes of the moment tensor solutions across the Alps provide information on local deformation regimes. Their distribution of P- and T-axes across the Alps mountain belt points out both, local and regional heterogeneities
465 (Fig. 11). ~~While in general compressional regimes are dominant in the SE Alps and the Central Alps, extensional stresses regimes are observed within the Western Alps.~~ In addition to the direct interpretation of P- and T- axes ~~of the MT solutions~~, we apply a stress inversion approach based on the minimization of the seismic energy released on unfavourably oriented faults (Cesca et al., 2016) in volumes of comparably high seismic activity (Supplement Fig. S2). Stress inversion results provide the orientations of the most compressive (σ_1), the intermediate (σ_2) and the least compressive principle stresses (σ_3), and the
470 relative stress magnitude $R=(\sigma_1-\sigma_2)/(\sigma_1-\sigma_3)$. A homogeneous stress field is assumed within the selected rock volume and time period for each subregion.

In the following, we describe the patterns of P- and T- axes as well as the local stress regimes in the different seismo-tectonic domains which were inferred from the moment tensor solutions. In doing so, we first concentrate on the typical N-S compressional regime in the central to eastern Southern Alps (Region 2 and 3 in Fig. 1), before we focus on the transition
475 to the strike-slip regime in the Northern Dinarides (Region 4 in Fig. 1). Subsequently, the deformation regime in the Western Alps (Region 1 in Fig. 1) is discussed. Additionally, we discuss the findings for the neighboring northern Apenninic mountain range (Region 5 in Fig. 1).

At the southern margin of the central Southern Alps, we observe predominantly thrust mechanisms with NNW-SSE to NW-SE oriented P-axes in the central Alps, close to Lake Garda, to NNE-SSW oriented P-axes further east, ~~close to Viena~~ (Fig.
480 11a, features d and e). Our stress inversion results confirm dominating compression from ~~central to the eastern~~ the central to eastern Southern Alps with sub-horizontal σ_1 orientation (Supplement Fig. S2), which is in agreement with the stress map of the Mediterranean and Central Europe (Heidbach et al., 2016). Seismic activity at thrust faults originating from the N-S convergence of the Adriatic and Eurasian plates in the Southern Alps are well known and have been described by various studies (e.g. Pondrelli et al., 2006; Anselmi et al., 2011; Poli and Zanferrari, 2018). According to Cheloni et al. (2014), the
485 SE Alpine thrust front absorbs about 70% of the convergence between the continental plates. In the transition from the SE Southern Alps to the Northern Dinarides, ~~across a distance of 200 km~~, a rotation of the P-axes from NW-SE ~~in the western part~~

to NNE-SSW ~~in the eastern part~~ is observed (Fig. 11a, features a-c). Despite increased uncertainties due to the relatively low number of available MT solutions, we observe a ~~similar~~~~the same~~ rotation of σ_1 . ~~Although less distinct, this rotation can also be seen when looking at the stress direction obtained from thrust MTs in Heidbach et al. (2016). The changes in the orientation of the thrust mechanisms may be attributed to the bending of the southern thrust front of the Alps and to the transition to the strike-slip fault systems in the Dinarides.~~




Figure 11. Regional distribution of (a) P-, (b) T- and (c) B-axes of focal mechanisms presented in Fig. 9. Colors correspond to mechanism classes as shown in Fig. 9. Only areas with more than five events in a latitude-longitude-grid of $1^\circ \times 1^\circ$ are shown. a to h mark features which are discussed in the text.

The transition from dominant thrust faulting **in the south-eastern Alps** close to Friuli to the strike-slip events to the east and in the northern Dinarides **was also described by Pondrelli et al. (2006)** and is mapped by the change from a sub-vertical to an almost horizontal σ_3 direction (**Supplement Fig. S2**). **Moulin et al. (2016)** describe right-lateral motion ($3.8 \pm 0.6 \text{ mmyr}^{-1}$)

495 on three main Dinaric faults, and suggest that the system of NW-SE oriented right-lateral strike-slip faults might be the north-eastern boundary of the Adriatic microplate.

The W Alps show more heterogeneous faulting compared to the Southern Alps and the Northern Dinarides. We obtained four new MT solutions indicating NW-SE striking normal faulting east of Lake Geneva in the NW Alps, in agreement with the orientation of T axis at a high angle to the bending of the orogen described by Delacou et al. (2004). Our MT solutions and the catalog solutions show normal faulting, thrust faulting and some oblique strike-slip faulting events in the W to SW Alps (Region 1 on Fig. 1). Despite the small number of moment tensors, we can also infer a rotation of the T-axes in the south-western Alps. In contrast to the north-western Alps, where the tensional axis points at an extension along the mountain belt, the T-axes of the focal mechanisms are oriented roughly perpendicular to the Alpine arc (Fig. 11b, features f-g). This observation is in accordance with Mathey et al. (2020), who recently studied the stress field of the western Alps based on an new extensive dataset of P-wave first motion polarity focal mechanisms. The stress inversion results indicate an extensional regime in the Western Alps with a sub-vertical σ_1 orientation. However, across the large region the uncertainties of the stress inversion are relatively high. This is in agreement with the co-existing thrust faulting, normal faulting and strike-slip faulting also shown by Delacou et al. (2004) and Heidbach et al. (2016).

510 Normal faulting, with a similar orientation as in the NW Alps, is dominant along the central arc of the Apennines. The normal faulting events of the NW Alps are located within the strike direction of those in the Apennines. However, despite the similarity of mechanisms, the depths of the events in the NW Alps are significantly shallower (Fig. 10).

Along the Apennines, thrust faulting is dominant at the northern arc, while normal faulting earthquakes are dominant south-west of the ridge of the Apennines. The NW-SE orientations of the T axes of the normal faulting events are perpendicular to the elongation of the mountain belt as also described by Pondrelli et al. (2006). The vertical σ_1 direction and the NE-SW oriented, horizontal σ_3 direction confirm an extensional stress regime (Supplement Fig. S2). In contrast, a compressional regime is observed along the NE arc of the Apennines with P axes of the thrust faulting events oriented NW-SE to NE-SW (Fig. 11).

4.2 CMT solutions in the seismo-tectonic context

4.2 Seismicity and strain

520 Since only few focal mechanisms are available for large parts of the Alps, we additionally take into account ~~rely on~~ recent seismicity (Fig. 12a,c,d), historical large earthquakes (Fig. 12b) and GNSS data (Fig. 12e-f) ~~to further characterize the tectonic activity to analyze our results in the seismo-tectonic context and draw a more detailed picture of the seismic and tectonic activity in the study area.~~ To emphasize areas of ~~high~~ significant seismic activity (Fig. 12a) we cluster the earthquakes in the seismicity catalogs by INGV, GEOFON and ~~g~~CMT according to their epicentral locations using the DBSCAN clustering algorithm (Ester et al., 1996) implemented in the Python package *scikit-learn* (Pedregosa et al., 2011). The merged catalogs ~~comprises a total of~~ more than 50,000 earthquakes with $M_I > 2.0$ (1983-2017). The recent seismic activity ~~in the seismicity clusters~~ can be traced back to historical times. Fig. 12b shows historical earthquakes from the European Archive of Historical Earthquake Data 1000-1899 (AHEAD; Locati et al., 2014; Rovida and Locati, 2015) based on the SHARE European Earthquake Catalogue

(SHEEC; Stucchi et al., 2013) and the ISC-GEM catalog (Storchak et al., 2013, 2015; Bondár et al., 2015; Di Giacomo et al., 530 2015, 2018) with $M_w > 5.5$. ~~Additionally, we report maximum hypocenter depths and cumulative seismic moments across the study area (Fig. 12c and d).~~

In general, the seismicity along the southern margin of the Alpine mountains is higher than along its northern counterpart. Apart from the large cluster of seismicity in the Apennines, we identify five dominant clusters and several smaller ones mainly located at the margins of the Alps (Fig. 12a). ~~The largest seismicity clusters correspond to the numbered regions shown in~~ 535 ~~Fig. 1. To avoid confusion with the mechanism-based clustering we here continue writing *region* (R) when referring to these clusters of epicenters. We focus on the analysis of the largest clusters, for which we were able to obtain multiple moment tensor solutions between 2016 and 2019.~~ Following the Alpine arc from west to east, the first cluster is located in the Western Alps at the French-Italian border (R 1 in Fig. 12a and Fig. 1), two smaller clusters are found in the region around Lake Garda in the central Southern Alps (R 2) and north of it. Two more clusters of high seismicity are situated in the ~~eastern Southern SE~~ 540 ~~Alps in the border region between Italy (Friuli) and Slovenia (R 3), and in the Northern Dinarides (R 4). Fig. 12a indicates dominant faulting styles for each cluster, namely thrust faulting for regions 2 and 3 around Lake Garda and in the eastern Southern SE Alps, and strike-slip faulting in the Northern Dinarides (R 4). In the epicentral cluster of the Apennines (R 5), two representative mechanisms reflect the separation of dominant normal and thrust faulting earthquakes SW and NE of the ridge, respectively. Due to the heterogeneous faulting in the Western Alps, we assign no representative mechanism.~~

The cluster of high seismicity in the ~~eastern Southern SE~~ Alps (Fig. 12a) is located close to the epicenter of the 1976 Friuli Earthquake (M_b 6.0; Pondrelli et al., 2001). The observed E-W striking thrust events map the regional dominant stress field (Fig. 9), evolving from the underthrusting of the Friuli Plain beneath the Alps (e.g. Cipar, 1980). Focal mechanism solutions of the 1976 mainshock and aftershocks show similar ~~thrust focal mechanisms striking E-W to NE-SW; thrust mechanisms,~~ partly with a small strike-slip component, and are associated to the complex Periadriatic overthrust system (e.g. Cipar, 1980; Bressan 550 et al., 1998; Pondrelli et al., 2001, 2006; Poli and Zanferrari, 2018; Slejko, 2018). ~~In addition, o~~ Only few tenth of kilometers to the west of the Friuli area, ~~close to Forni di Sopra, Italy,~~ we observe strike-slip faulting (Fig. 9). Anselmi et al. (2011) report the occurrence of both thrust and strike-slip faulting for this area, mostly in agreement with an E-W to ENE-WSW minimum horizontal stress reported by Montone et al. (2004). Large historical events are reported ~~along the southern margin of the Alps between Lake Garda (R 2), for the eastern Southern SE Alps (R 3) and the transition to the Northern Dinarides (R 4) (Verona~~ 555 ~~1117, Slovenia 1511 and Carinthia 1348, bothall $M_w \geq 6.7$).~~ ~~and close to Lake Garda in the central S-Alps (Verona 1117, M_w 6.7).~~ Similar, high seismicity and cumulative seismic moments during the last decades (Fig. 12d) are observed here. Within the seismicity cluster close to Lake Garda, Italy, ~~we the observed thrust NE-SW striking thrust faulting. These~~ mechanisms are typical for earthquakes located in the Giudicarie region close to the Ballino-Garda fault ~~which is running through Lake Garda~~ (Viganò et al., 2008).

560 ~~Less frequent,~~ Large historical earthquakes with magnitude estimates between M_w 5 and 6 are ~~also~~ reported in the Western Alps and the Dinarides (R 2 and 4). Within the ~~last decadetime span of >30 years,~~ earthquakes with $4 < M_l < 5$ are observed across a wider part of the Western and Southern Alps, but magnitudes rarely exceed M_l 3.5 in large areas of the Central to NE Alps. ~~In contrast, t~~ The Eastern Alps north of the Periadriatic line and the area between the ~~high seismicity~~ ~~seismically active~~

regions in the Western Alps and the Central Alps have particularly low seismicity rates (Fig. 12a). ~~Therefore, only few focal~~
565 ~~mechanisms are available in this area.~~ While at least three large earthquakes occurred in the eastern part of Switzerland in
historical times, seismicity in this region appears to be relatively low in recent years (Fig. 12a and b).

The observed depth ranges ~~of our MT solutions (Fig. 10)~~ are in accordance with the maximum depths in the long-term
seismic catalogs of ~~g~~GCMT, INGV and GEOFON, including >50,000 earthquakes with $M_l > 2.0$ (Fig. 12c). ~~While the Moho~~
570 ~~depth increases gradually from less than 30 km at the northern margin of the Alps to above 50 km in the central part of the~~
~~orogen (Spada et al., 2013), we do not observe any gradual change in the event depth. The se catalogs and our own centroid~~
~~depths show that seismicity is shallow across most of the Alps with rare deeper events (<30 km) at the southern margin,~~
~~where the Moho is at about 40 km depth (Spada et al., 2013). These few deeper events are located above the Moho. ~~while~~~~
~~m~~Maximum depths of above 60 km are observed in the Apennines.




Figure 12. Characteristics of recent and historical seismicity and strain from GNSS data. (a) Seismic activity between 1978-2017, $M_I > 2.0$, from GEOFON, INGV and g GCMC catalogs, colored according to epicentral clusters of seismicity. Representative MTs from Fig. 9. Numbers refer to regions (1) W Alps, (2) Lake Garda, (3) eastern S SE Alps, (4) N Dinarides and (5) Apennines. (b) Historical earthquakes with $M_w > 5.5$ from the European Archive of Historical Earthquake Data 1000-1899 (AHEAD) (Locati et al., 2014; Rovida and Locati, 2015; Stucchi et al., 2013) and ISC-GEM catalog (1906-2016) (Storchak et al., 2013, 2015; Bondár et al., 2015; Di Giacomo et al., 2015, 2018). (c, d) Maximum event depth and the cumulative seismic moment on a grid with a spacing of $0.25 \times 0.25^\circ$ lat. and lon. (e, f) Absolute value of the spatial gradient of the relative uplift rate as a proxy for vertical strain rates and observed GNSS shear strain rate ($\text{max-shear-strain}^{2nd}$ invariant of strain tensor). Both obtained from GNSS data of the EUREF WG on European Dense Velocities (Brockmann et al., 2019). Grey dots indicate earthquakes without MTs, as in (a). Black dots indicate seismicity, $M_I > 2.0$. Events with MT solutions are color-coded as in Fig. 9. For simplicity, focal mechanisms are only plotted for representative events with $M_w \geq 4.0$. Exposed and subsurface faults (solid and dashed lines) simplified from Schmid et al. (2004, 2008); Handy et al. (2010, 2015) and Patacca et al. (2008). 'PL' marks the Periadriatic line, 'SK' the Split-Karlovac Fault. Topographic data from SRTM-3 (Farr et al., 2007) and ETOPO1 (Amante and Eakins, 2009) datasets.

The joined interpretation of seismicity, GNSS data and MT solutions shows how the overall spatial distribution of faulting styles in the study area can be interpreted in the regional tectonic regime. Fig. 12e and f present the spatial gradient of the uplift rates and the horizontal strain rates, computed from the GNSS data of the EUREF WG on European Dense Velocities (http://pnac.swisstopo.admin.ch/divers/dens_vel/index.html, Brockmann et al., 2019). Please refer to the supplement for additional methodological information. Following Keiding et al. (2015), we use the spatial derivative of the uplift rate as a proxy of vertical strain rates (Fig. 12e).

Within the Alpine mountain range, the GNSS data shows a consistent uplift relative to the surrounding areas (Supplement Fig. S3). Fig. 12e and f emphasize the relation between recent seismic activity and both; high spatial gradients of the uplift rate (e) and the shear strain rate (f) across large parts of the study area. Largest gradients of the uplift rate, and high shear strain rates, and highest seismicity rates are observed in the eastern Southern SE Alps (R 3 in Fig. 12a) and in the Apennines (R 5 in Fig. 12a) where seismicity is highest. The distribution of the cumulative seismic moment (Fig. 12d) agrees particularly well with the distribution of shear strain rates (Fig. 12f). We observe that thrust faulting events cluster at the SE margin of the Alps. We observe high seismic activity with typically E-W striking thrust faulting in the eastern Southern SE Alps, as also described in many previous studies (e.g. Pondrelli et al., 2006; Poli and Zanferrari, 2018). High horizontal velocities point at the shortening and crustal thickening in the eastern Southern SE and Eastern Alps due to the convergence of Adriatic and European plate (Supplement Fig. 3), in accordance with Serpelloni et al. (2016) and Sternai et al. (2019). South-east of this area, at the transition to the Northern Dinarides (Regions 3-4), the uplift gradients are low, while increased shear strain rates agree with our dominant strike-slip mechanisms (see also Serpelloni et al., 2016) and right-lateral motion on the Dinaric strike-slip faults (Moulin et al., 2016).

While there is no significant shear strain in the Western and Central Alps, we depict two subparallel bands of moderate spatial gradients of the uplift rate running roughly along the northern and the southern margin of the Alps. These two elongated regions bands result from the overall relative uplift of the Alps and also have higher seismicity rates compared to the central Alpine belt. In the SW Alps, the largest events cluster in the transition area between relative uplift and subsidence, indicated by the band of increased spatial gradient of the uplift in Fig. 12e. Normal faulting events are dominant. Intraplate shear strain rates are relatively low in the entire Western and Central Alps.

The Adriatic plate, which is the upper plate in the Alpine subduction zone, rotates counter-clockwise relative to Europe around an Euler pole located in the western Po plain or Western Alps (D'Agostino et al., 2008; Weber et al., 2010; Le Breton et al., 2017). The rotation results in varying convergence rates across the Alps. Le Breton et al. (2017); Le Breton et al. (2021) infer a rotation of about 5.25° during the last 20 Ma resulting in convergence rates ranging from 5.5 mmyr^{-1} in the NW Adria (Western Alps) to 7.5 mmyr^{-1} in the NE Adria (eastern Southern Alps) (Fig. 1). In comparison, kinematic reconstructions of Van Hinsbergen et al. (2020) involve less convergence but a higher rotation of Adria relative to Europe (10°), leading to 2.5 mmyr^{-1} convergence in the NW Adria to 6.25 mmyr^{-1} in the NE Adria. Recent GPS data indicates little to no horizontal movement in the Western Alps but more than 2 mmyr^{-1} NNW-ward movement of Adria in the eastern Southern Alps (see Fig. S3 in the Supplement), which is in agreement with increased seismicity rates. The Western Alps are closer to the location of the Euler pole of the rotation of the Adriatic plate, therefore convergence rates are lower. Recent GPS measurements and

the computed horizontal strain rates even indicate the absence of convergence (D'Agostino et al., 2008, , and Fig. S3 in the Supplement). Therefore, the uplift pattern of the Western and Central Alps (Fig. 12e and Fig. S3 in the Supplement) as well as the seismicity clusters in the W Alps need to be attributed to other mechanisms. Sternai et al. (2019) propose that isostatic adjustment to deglaciation and erosion, and mantle-related processes such as slab detachment or asthenospheric upwelling may jointly explain the observed uplift pattern. The assumption of a stress/strain field which is not dominantly effected by the convergence of Europe and Africa was also proposed by Delacou et al. (2004) based on focal mechanisms and stress inversion in the Western Alps. Our moment tensor solutions indicating normal and strike-slip faulting, as well as the P and T axes match these observations from GNSS data.

The seismic activity along the northern margin of the Alps is in agreement with the increased gradient of uplift in this area. However, the occurrence of strike-slip faulting earthquakes described in this study can hardly be explained by vertical strain, especially ~~for~~ favourably oriented faults. However, pre-existing faults may be unfavorably oriented, the stress field may be heterogeneous and local anomalies may not be resolved by the sparse GNSS network. Rather low seismicity is further observed in the eastern Po plain, where high spatial gradients of the uplift rate are observed. The high absolute gradient here can be attributed to the relative subsidence of the sediments in the Po plain (see supplement Fig. S3) (Carminati and Martinelli, 2002), but not to a tectonic uplift which would likely be accompanied by seismic activity.

5 Conclusions

Centroid moment tensor inversion provides insight into faulting mechanisms of earthquakes and related tectonic processes. In this study, we used the AlpArray seismic network to analyze the mechanisms of earthquakes occurring from 2016 to end of 2019. Thanks to the flexible inversion tool *Grond*, we were able to test different inversion set-ups in order to derive guidelines for MT inversions in complex tectonic settings such as the Alps. These guidelines as well as the proposed tests can, on the one hand, facilitate future studies of faulting mechanisms in the Alps and, on the other hand, help to derive workflows to obtain reliable moment tensors in other dense networks or complex study regions. ~~and to~~ We evaluated the results with respect to their uncertainties and parameter trade-offs. For subsets of events, we tested various frequency bands, distance ranges and different input data types comprising time domain full waveforms, frequency domain amplitude spectra, time domain cross-correlation fitting, waveform envelopes and combinations of these. In the case of our study area, for most earthquakes with magnitudes larger Mw 3.3, we find that a combination of time domain full waveforms and frequency domain amplitude spectra in a frequency band of 0.02-0.07 Hz is most suitable. ~~We evaluate the influence of azimuthal gaps~~ The dense deployment of the AASN is ideal to study the effect of (manually introduced) azimuthal gaps. While a higher azimuthal station coverage is in general favorable, we find that a small number of stations with little azimuthal coverage may be sufficient depending on the location of the stations with respect to the strike direction of the fault. ~~and study both the occurrence and the meaningfulness of non-DC components.~~ Performing CMT inversions constraining the solutions to a pure double-couple MT, a deviatoric MT or allowing for a full MT, indicates that for the specific Alpine context with a dense network DC components are reliably resolved independent of the applied constraint. While allowing for non-DC components reduces the overall misfit, the CLVD and the

isotropic components cannot be distinguished unambiguously. We propose to perform similar tests prior to MT inversions for other study areas, when earthquake magnitudes are small, the crustal structure is complex, the number of stations is limited or other factors might hinder straight-forward inversions.

645 ~~We observe that even in cases where non-DC components seem well resolved, these components should be carefully evaluated by forward modelling and comparing the waveforms of solutions from pure-DC, deviatoric and full inversions. In case of small events, we observe that large gaps in the azimuthal station distribution can hinder successful inversions or may lead to biased results. In contrast, under favorable geometric conditions, when strike direction and tensional as well as compressional quadrants are covered, MT inversions are possible even for small events using only few stations.~~

650 Relying on the results of the methodological tests, we performed deviatoric MT inversions for events with $M_w \geq 3.1$ ~~$M_w > 3.0$~~ . We present 75 solutions with reasonably low uncertainties ~~for earthquakes with $M_w > 3.1$~~ occurring between 2016 and 2019. With four years of acquisition of small to moderate earthquakes in the course of the AlpArray project we are able to identify seismotectonic domains which are representative in faulting styles of those derived from long term seismic observations. ~~and We compare these derived MT solutions to historical earthquakes, recent seismicity and, published focal mechanisms and~~
655 ~~GNSS deformation data.~~ Our moment tensor results indicate that while the Alps represent a rather heterogeneous study area, the region is characterised by compartments of different tectonic movement in close proximity. ~~We applied a clustering-based on the Kagan angle to identify groups of similar events.~~ Typical ENE-WSW to E-W striking thrust faulting is observed in the Friuli area in the eastern Southern SE Alps related to the N-S convergence of the Eurasian and Adriatic plate (Pondrelli et al., 2006; Poli and Zanferrari, 2018) and counter-clockwise rotation of Adria relative to Europe (e.g. D'Agostino et al., 2008;
660 Le Breton et al., 2017). Strike-slip faulting with similarly oriented P-axes are observed parallel to the northern margin of the Central Alps and in the Northern Dinarides, which is in agreement with right-lateral strike-slip faults and high shear strain rates. In contrast, NW-SE striking normal faulting events with NE-SW oriented T-axes are observed in the NW Alps. ~~with similar strike directions but at shallower depths than the dominant normal faulting events in the Apennines.~~ Faulting styles in the SW Alps are more heterogeneous with a majority of events related to an extensional stress regime. Simultaneous observations
665 of low horizontal strain rates and normal faulting earthquakes are in agreement with studies proposing that relatively high uplift rates in the Western Alps are attributed to other processes than the Europe-Adria convergence. ~~While in the Apennines mechanism types are separated along the ridge of the mountain range by depth and location, within the Alps we observe a separation of compressional stresses in the eastern Southern SE as well as in the Central Alps and a tensional regime in the Western Alps. Oblique to the bending of the Alpine mountain range, the T-axes of normal faulting earthquakes rotate from~~
670 ~~NW-SE in the NW Alps to about E-W in the SW Alps. We discussed these different tectonic areas systematically, including observations of seismicity, faulting mechanisms and GNSS deformation data.~~ Based on a clustering of epicenters, we identify five main seismically active ~~sub~~ regions, namely the Western Alps, the region around Lake Garda, the eastern Southern SE Alps, the Northern Dinarides and the Apennines. ~~These regions~~ Areas of high seismicity are mostly located in the proximity of the southern margin of the Alps, where significant vertical or horizontal strain rates are reported. Maximum observed magnitudes
675 coincide with regions of increased seismicity and significant historical earthquakes, but rarely exceed M_w 5.0. In contrast, seismicity is particularly low in the Eastern Alps and in parts of the Central Alps. ~~Both~~ The depths inferred from our moment

tensor inversions as well as the depths in seismic catalogs indicate that the seismic activity in the Alpine mountain ranges is predominantly shallow with only few events in depth greater than 15 km in the eastern Southern Alps. Significantly deeper earthquakes are observed in the Apennines.

680 ~~While we are able to obtain stable and reliable MT solutions for most earthquakes with $M_w \leq 3.3$ that occurred during the installation period of the AASN between 2016 and 2019, we struggled to invert smaller events with M_w 3.1-3.3. On one hand, the spacing of the AASN of 60 km is not sufficiently dense for smaller earthquakes and on the other hand, topographic effects, low SNRs, as well as complex subsurface structures complicated the inversions, resulting in a success rate of one third for the magnitude range M_w 3.1-3.3. We strongly recommend a data quality assessment before performing any inversion. In future,~~
685 ~~we plan to model topographic effects and include recent 1-D and 3-D velocity models developed in the course of the AlpArray research initiative.~~

Code and data availability. The moment tensor inversions were performed using the free and open-source inversion tool *Grond* (Heimann et al., 2018). Figures were plotted using *pyrocko* (Heimann et al., 2019) and *GMT* (Wessel et al., 2013).

The topographic data for the maps was taken from the SRTM-3 (Farr et al., 2007) and ETOPO1 (Amante and Eakins, 2009) (NOAA National
690 Geophysical Data Center. 2009: ETOPO1 1 Arc-Minute Global Relief Model. NOAA National Centers for Environmental Information. Accessed [December 2020].)

Seismic catalogs: Earthquake information and focal mechanisms are provided by the below mentioned institutes:

Swiss Seismological Service (SED): (Deichmann et al., 2004; Baer et al., 2005; Deichmann et al., 2006; Baer et al., 2007; Deichmann et al.,
695 2008, 2009, 2010, 2011, 2012; Diehl et al., 2013, 2014, 2015, 2018; Diehl, 2020)

Slovenian Environment Agency (ARSO): Ministrstvo za okolje in prostor Agencija RS za okolje (2018), Ministrstvo za okolje in prostor Agencija RS za okolje (2019), Ministrstvo za okolje in prostor Agencija RS za okolje (2020)

INGV (Italy): [Scognamiglio et al. \(2006\)](#), <http://terremoti.ingv.it/en> (last access 12/2020)

GEOFON (Germany): Event locations were obtained from the GEOFON program of the GFZ German Research Centre for Geosciences
700 using data from the GEVN partner networks. <https://geofon.gfz-potsdam.de/eqinfo/list.php> (last access 12/2020)

EM-RCMT (European-Mediterranean Regional Centroid-Moment Tensors): <http://rcmt2.bo.ingv.it/> (last access 12/2020), Pondrelli (2002)

SISMOAZUR (France): http://sismoazur.oca.eu/focal_mechanism_emsc (last access 12/2020), - Provide MTs obtained using FMNEAR (Delouis, 2014)

*g*GCMT (Lamont-Doherty Earth Observatory of Columbia University, USA): <https://www.globalcmt.org/> (last access 12/2020), Dziewonski
705 et al. (1981); Ekström et al. (2012)

Permanent seismic networks: The permanent stations of the AlpArray are part of existing European regional networks (RD (RESIF, 2018), GU (University Of Genova, 1967), CZ (Institute Of Geophysics, A. O. S. O. T. C. R., 1973), ST (Geological Survey-Provincia Autonoma Di Trento, 1981), G (Institut De Physique Du Globe De Paris (IPGP), & Ecole Et Observatoire Des Sciences De La Terre De Strasbourg (EOST),
710 1982), CH (Swiss Seismological Service (SED) At ETH Zurich, 1983), OE (ZAMG-Zentralanstalt Für Meteorologie Und Geodynamik, 1987), MN (MedNet Project Partner Institutions, 1990), HU (Kövesligethy Radó Seismological Observatory (Geodetic And Geophysical Institute, Research Centre For Astronomy And Earth Sciences, Hungarian Academy Of Sciences (MTA CSFK GGI KRSZO)), 1992), GE (GEOFON Data Centre, 1993), RF (University Of Trieste, 1993), FR (RESIF, 1995), IV (INGV Seismological Data Centre, 2006), BW (Department Of Earth And Environmental Sciences, Geophysical Observatory, University Of Munchen, 2001), SX (Leipzig University, 2001), NI (OGS
715 (Istituto Nazionale Di Oceanografia E Di Geofisica Sperimentale) And University Of Trieste, 2002), TH (Jena, F.S.U., 2009), OX (OGS (Istituto Nazionale Di Oceanografia E Di Geofisica Sperimentale), 2016)).

Team list

The complete member list of the AlpArray Working Group can be found at <http://www.alparray.ethz.ch>.

Author contributions. G.P., S.C. and T.D. conceptualized the study. G.P. performed the moment tensor inversions, conceptualized and developed the methodological tests, prepared the figures and wrote the original draft of the manuscript. S.C., T.D., J.K. and T.P. developed
720

the general project idea and acquired the funding. S.C., S.H., D.K. and T.D. supervised the study. S.H. et al. developed the moment tensor inversion and the software tools used in this study. S.H., S.C., T.D. and P.N. contributed to methodological developments. S.H. and P.N. helped to visualize the results. T.D and T.P. performed stress field inversions. G.P, S.C. and T.D. validated the results. The AlpArray Working Group provided access to the seismic data of the temporary network and additional background information. All Co-authors reviewed and edited the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We like to thank the three anonymous reviewers for constructive comments that helped to improve the manuscript. Eline Le Breton is thanked for providing helpful information for the geological/tectonic background. GMP is funded by DFG project ‘From Top to Bottom - Seismicity, Motion Patterns and Stress Distribution in the Alpine Crust’ (Project Number 362440331), a subproject of ‘SPP 2017: Mountain Building Processes in 4D’ (Project Number 313806092). PN is currently funded by the BMBF (German Federal Ministry of Education and Research) project SECURE (grant agreement No. 03G0872A).

References

- AlpArray Seismic Network: AlpArray Seismic Network (AASN) temporary component. AlpArray Working Group, Other/Seismic Network, https://doi.org/10.12686/alparray/z3_2015, 2015.
- 735 Amante, C. and Eakins, B.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24., National Geophysical Data Center, NOAA, <https://doi.org/10.7289/V5C8276M>, 2009.
- Anselmi, M., Govoni, A., De Gori, P., and Chiarabba, C.: Seismicity and velocity structures along the south-Alpine thrust front of the Venetian Alps (NE-Italy), *Tectonophysics*, 513, 37–48, 2011.
- Baer, M., Deichmann, N., Braunmiller, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., KRadolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2004, *Eclogae Geologicae Helvetiae*, 98, 407–418, 2005.
- 740 Baer, M., Deichmann, N., Braunmiller, J., Clinton, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2006, *Swiss Journal of Geosciences*, 100, 517–528, 2007.
- Barth, A., Wenzel, F., and Giardini, D.: Frequency sensitive moment tensor inversion for light to moderate magnitude earthquakes in eastern Africa, *Geophys. Res. Lett.*, 34, <https://doi.org/https://doi.org/10.1029/2007GL030359>, 2007.
- 745 Bassin, C., Laske, G., and Masters, G.: The current limits of resolution for surface wave tomography in North America., *EOS Trans AGU*, 81, F897, 2000.
- Bondár, I., Engdahl, E. R., Villaseñor, A., Harris, J., and Storchak, D.: ISC-GEM: Global instrumental earthquake catalogue (1900–2009), II. Location and seismicity patterns, *Physics of the Earth and Planetary Interiors*, 239, 2–13, 2015.
- Bressan, G., Snidarcig, A., and Venturini, C.: Present state of tectonic stress of the Friuli area (eastern Southern Alps), *Tectonophysics*, 292, 750 211–227, 1998.
- Brockmann, E., Lutz, S., Zurutuza, J., Caporali, A., Lidberg, M., Völksen, C., Sánchez, L., Serpelloni, E., Bitharis, S., Pikridas, C., Fotiou, A., Mathis, E., Sánchez Sobrino, J., Valdés Péres De Vargas, M., Vernant, P., Baron, A., Westerhaus, M., Legrand, J., Kreemer, C., Gianniou, M., Nykiel, G., Figurski, M., Kenyeres, A., and Kurt, A.: Towards a Dense Velocity Field in Europe as a basis for Maintaining the European Reference Frame., 27th IUGG Assembly, Montreal, 2019.
- 755 Bukchin, B., Clévédédé, E., and Mostinskiy, A.: Uncertainty of moment tensor determination from surface wave analysis for shallow earthquakes, *Journal of seismology*, 14, 601–614, 2010.
- Cabieces, R., Buforn, E., Cesca, S., and Pazos, A.: Focal parameters of earthquakes offshore Cape St. Vincent using an amphibious network, *Pure and Applied Geophysics*, pp. 1–20, 2020.
- Carminati, E. and Martinelli, G.: Subsidence rates in the Po Plain, northern Italy: the relative impact of natural and anthropogenic causation, 760 *Engineering Geology*, 66, 241–255, [https://doi.org/https://doi.org/10.1016/S0013-7952\(02\)00031-5](https://doi.org/https://doi.org/10.1016/S0013-7952(02)00031-5), 2002.
- Cesca, S.: Seiscloud, a tool for density-based seismicity clustering and visualization, *J Seismology*, 24, 443–457, <https://doi.org/10.1007/s10950-020-09921-8>, 2020.
- Cesca, S. and Heimann, S.: Challenges in regional moment tensor resolution and interpretation., in: D’Amico S. (eds) *Moment Tensor Solutions*, pp. 163–181, Springer Natural Hazards. Springer, Cham., https://doi.org/https://doi.org/10.1007/978-3-319-77359-9_7, 2018.
- 765 Cesca, S., Buforn, E., and Dahm, T.: Amplitude spectra moment tensor inversion of shallow earthquakes in Spain, *Geophysical Journal International*, 166, 839–854, 2006.
- Cesca, S., Heimann, S., Stammler, K., and Dahm, T.: Automated procedure for point and kinematic source inversion at regional distances, *Journal of Geophysical Research: Solid Earth*, 115, 2010.

- Cesca, S., Rohr, A., and Dahm, T.: Discrimination of induced seismicity by full moment tensor inversion and decomposition, *Journal of seismology*, 17, 147–163, 2013.
- Cesca, S., Grigoli, F., Heimann, S., Dahm, T., Kriegerowski, M., Sobiesiak, M., Tassara, C., and Olcay, M.: The Mw 8.1 2014 Iquique, Chile, seismic sequence: a tale of foreshocks and aftershocks, *Geophysical Journal International*, 204, 1766–1780, 2016.
- Cheloni, D., D’Agostino, N., and Selvaggi, G.: Interseismic coupling, seismic potential, and earthquake recurrence on the southern front of the Eastern Alps (NE Italy), *Journal of Geophysical Research: Solid Earth*, 119, 4448–4468, <https://doi.org/10.1002/2014JB010954>, 2014.
- Cipar, J.: Teleseismic observations of the 1976 Friuli, Italy earthquake sequence, *Bulletin of the Seismological Society of America*, 70, 963–983, 1980.
- D’Agostino, N., Avallone, A., Cheloni, D., D’anastasio, E., Mantenuto, S., and Selvaggi, G.: Active tectonics of the Adriatic region from GPS and earthquake slip vectors, *Journal of Geophysical Research: Solid Earth*, 113, <https://doi.org/https://doi.org/10.1029/2008JB005860>, 2008.
- Dahal, N. R. and Ebel, J. E.: Method for Determination of Focal Mechanisms of magnitude 2.5-4.0 Earthquakes Recorded by a Sparse Regional Seismic Network, BSSA, 2020.
- Dahm, T. and Krüger, F.: Moment tensor inversion and moment tensor interpretation, Bohrmann, P.: *New Manual of Seismological Observatory Practice 2 (NMSOP-2)*, Deutsches GeoForschungsZentrum GFZ, pp. 1–37, https://doi.org/10.2312/GFZ.NMSOP-2_IS_3.9, 2014.
- Dahm, T., Becker, D., Bischoff, M., Cesca, S., Dost, B., Fritschen, R., Hainzl, S., Klose, C. D., Kühn, D., Lasocki, S., et al.: Recommendation for the discrimination of human-related and natural seismicity, *Journal of seismology*, 17, 197–202, 2013.
- Dahm, T., Heimann, S., Funke, S., Wendt, S., Rappsilber, I., Bindi, D., Plenefisch, T., and Cotton, F.: Seismicity in the block mountains between Halle and Leipzig, Central Germany: centroid moment tensors, ground motion simulation, and felt intensities of two $M \approx 3$ earthquakes in 2015 and 2017, *Journal of Seismology*, pp. 1–19, 2018.
- d’Amico, S., Orecchio, B., Presti, D., Gervasi, A., Zhu, L., Guerra, I., Neri, G., and Herrmann, R.: Testing the stability of moment tensor solutions for small earthquakes in the Calabro-Peloritan Arc region (southern Italy), *Bollettino di Geofisica Teorica ed Applicata*, 2011.
- Deichmann, N., Baer, M., Braunmiller, J., Cornou, C., Fäh, D., Giardini, D., Gisler, M., Huber, S., Husen, S., Kästli, P., et al.: Earthquakes in Switzerland and surrounding regions during 2003, *Eclogae Geologicae Helvetiae*, 97, 447–458, 2004.
- Deichmann, N., Baer, M., Braunmiller, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2005, *Eclogae Geologicae Helvetiae*, 99, 443–452, 2006.
- Deichmann, N., Baer, M., Clinton, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2007, *Swiss Journal of Geosciences*, 101, 659–667, 2008.
- Deichmann, N., Clinton, J., Husen, S., Haslinger, F., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., Marschall, I., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2008, *Swiss Journal of Geosciences*, 102, 505, 2009.
- Deichmann, N., Clinton, J., Husen, S., Edwards, B., Haslinger, F., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., Marschall, I., et al.: Earthquakes in Switzerland and surrounding regions during 2009, *Swiss Journal of Geosciences*, 103, 535–549, 2010.
- Deichmann, N., Clinton, J., Husen, S., Edwards, B., Haslinger, F., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2010, *Swiss Journal of Geosciences*, 104, 537–547, 2011.
- Deichmann, N., Clinton, J., Husen, S., Edwards, B., Haslinger, F., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2011, *Swiss Journal of Geosciences*, 105, 463–476, 2012.

- Delacou, B., Sue, C., Champagnac, J.-D., and Burkhard, M.: Present-day geodynamics in the bend of the western and central Alps as constrained by earthquake analysis, *Geophysical Journal International*, 158, 753–774, <https://doi.org/https://doi.org/10.1111/j.1365-246X.2004.02320.x>, 2004.
- Delouis, B.: FMNEAR: Determination of focal mechanism and first estimate of rupture directivity using near-source records and a linear distribution of point sources, *Bulletin of the Seismological Society of America*, 104, 1479–1500, 2014.
- Department Of Earth And Environmental Sciences, Geophysical Observatory, University Of Munchen: BayernNetz [Data set]. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/BW>, 2001.
- Di Giacomo, D., Bondár, I., Storchak, D. A., Engdahl, E. R., Bormann, P., and Harris, J.: ISC-GEM: Global Instrumental Earthquake Catalogue (1900–2009), III. Re-computed MS and mb, proxy MW, final magnitude composition and completeness assessment, *Physics of the Earth and Planetary Interiors*, 239, 33–47, <https://doi.org/10.1016/j.pepi.2014.06.005>, 2015.
- Di Giacomo, D., Engdahl, E. R., and Storchak, D. A.: The ISC-GEM Earthquake Catalogue (1904–2014): status after the Extension Project., *Earth System Science Data*, 10, <https://doi.org/10.5194/essd-10-1877-2018>., 2018.
- Diehl, T., Husen, S., Kissling, E., and Deichmann, N.: High-resolution 3-DP-wave model of the Alpine crust, *Geophysical Journal International*, 179, 1133–1147, 2009.
- Diehl, T., Deichmann, N., Clinton, J., Husen, S., Kraft, T., Plenkens, K., Edwards, B., Cauzzi, C., Michel, C., Kästli, P., et al.: Earthquakes in Switzerland and surrounding regions during 2012, *Swiss Journal of Geosciences*, 106, 543–558, 2013.
- Diehl, T., Clinton, J., Kraft, T., Husen, S., Plenkens, K., Guilhelm, A., Behr, Y., Cauzzi, C., Kästli, P., Haslinger, F., et al.: Earthquakes in Switzerland and surrounding regions during 2013, *Swiss Journal of Geosciences*, 107, 359–375, 2014.
- Diehl, T., Deichmann, N., Clinton, J., Kästli, P., Cauzzi, C., Kraft, T., Behr, Y., Edwards, B., Guilhem, A., Korger, E., et al.: Earthquakes in Switzerland and surrounding regions during 2014, *Swiss Journal of Geosciences*, 108, 425–443, 2015.
- Diehl, T., Clinton, J., Deichmann, N., Cauzzi, C., Kästli, P., Kraft, T., Molinari, I., Böse, M., Michel, C., Hobiger, M., et al.: Earthquakes in Switzerland and surrounding regions during 2015 and 2016, *Swiss Journal of Geosciences*, 111, 221–244, 2018.
- Diehl, T. e. a.: Earthquakes in Switzerland and surrounding regions during 2017 and 2018 (in prep.), *Swiss Journal of Geosciences*, 111, 221–244, 2020.
- Domingues, A., Custodio, S., and Cesca, S.: Waveform inversion of small-to-moderate earthquakes located offshore southwest Iberia, *Geophysical Journal International*, 192, 248–259, 2013.
- Dufumier, H. and Cara, M.: On the limits of linear moment tensor inversion of surface wave spectra, pure and applied geophysics, 145, 235–257, 1995.
- Dufumier, H. and Rivera, L.: On the resolution of the isotropic component in moment tensor inversion, *Geophysical Journal International*, 131, 595–606, 1997.
- Dziewonski, A., Chou, T.-A., and Woodhouse, J. H.: Determination of earthquake source parameters from waveform data for studies of global and regional seismicity, *Journal of Geophysical Research: Solid Earth*, 86, 2825–2852, 1981.
- Ekström, G., Nettles, M., and Dziewoński, A.: The global CMT project 2004–2010: Centroid-moment tensors for 13,017 earthquakes, *Physics of the Earth and Planetary Interiors*, 200, 1–9, 2012.
- Ester, M., Kriegel, H.-P., Sander, J., Xu, X., et al.: A density-based algorithm for discovering clusters in large spatial databases with noise., in: *Kdd*, vol. 96(34), pp. 226–231, 1996.

- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., Roth, L., Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M., Burbank, D., and Alsdorf, D.: The Shuttle Radar Topography Mission, *Reviews of Geophysics*, 45, <https://doi.org/10.1029/2005RG000183>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2005RG000183>, 2007.
- 845 Ford, S. R., Dreger, D. S., and Walter, W. R.: Identifying isotropic events using a regional moment tensor inversion, *Journal of Geophysical Research: Solid Earth*, 114, 2009.
- Fry, B., Deschamps, F., Kissling, E., Stehly, L., and Giardini, D.: Layered azimuthal anisotropy of Rayleigh wave phase velocities in the European Alpine lithosphere inferred from ambient noise, *Earth and Planetary Science Letters*, 297, 95–102, 2010.
- GEOFON Data Centre: GEOFON Seismic Network. Deutsches GeoForschungsZentrum GFZ, Other/Seismic Network, 850 <https://doi.org/10.14470/TR560404>, 1993.
- Geological Survey-Provincia Autonoma Di Trento: Trentino Seismic Network. International Federation of Digital Seismograph Networks., Other/Seismic Network, <https://doi.org/10.7914/SN/ST>, 1981.
- Guilhem, A., Hutchings, L., Dreger, D., and Johnson, L.: Moment tensor inversions of $M \sim 3$ earthquakes in the Geysers geothermal fields, California, *Journal of Geophysical Research: Solid Earth*, 119, 2121–2137, 2014.
- 855 Handy, M., Babist, J., Wagner, R., Rosenberg, C., and Konrad, M.: Decoupling and its relation to strain partitioning in continental lithosphere: insight from the Periadriatic fault system (European Alps), *Geological Society, London, Special Publications*, 243, 249–276, 2005.
- Handy, M. R., Schmid, S. M., Bousquet, R., Kissling, E., and Bernoulli, D.: Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological–geophysical record of spreading and subduction in the Alps, *Earth-Science Reviews*, 102, 121–158, 2010.
- Handy, M. R., Ustaszewski, K., and Kissling, E.: Reconstructing the Alps–Carpathians–Dinarides as a key to understanding switches in 860 subduction polarity, slab gaps and surface motion, *International Journal of Earth Sciences*, 104, 1–26, 2015.
- Hardebeck, J. L. and Shearer, P. M.: A new method for determining first-motion focal mechanisms, *Bulletin of the Seismological Society of America*, 92, 2264–2276, 2002.
- Heidbach, O., Custodio, C., Kingdon, A., Mariucci, M. T., Montone, P., Müller, B., Pierdominici, S., Rajabi, M., Reinecker, J., Reiter, K., Tingay, M., Williams, J., and Ziegler, M.: Stress map of the Mediterranean and Central Europe 2016, GFZ Data Services, 865 <https://doi.org/10.5880/WSM.Europe2016>, 2016.
- Heimann, S.: A Robust Method To Estimate Kinematic Earthquake Source Parameters, Ph.D. thesis, Universität Hamburg, <http://ediss.sub.uni-hamburg.de/volltexte/2011/5357/pdf/Dissertation.pdf>, 2011.
- Heimann, S., Isken, M., Kühn, D., Sudhaus, H., Steinberg, A., Vasyura-Bathke, H., Daout, S., Cesca, S., and Dahm, T.: Grond - A probabilistic earthquake source inversion framework, <https://doi.org/10.5880/GFZ.2.1.2018.003>, <http://pyrocko.org/grond/docs/current/>, 2018.
- 870 Heimann, S., Vasyura-Bathke, H., Sudhaus, H., Paul Isken, M., Kriegerowski, M., Steinberg, A., and Dahm, T.: A Python framework for efficient use of pre-computed Green’s functions in seismological and other physical forward and inverse source problems, *Solid Earth*, 10, 1921–1935, 2019.
- Heit, B., Weber, M., Tilmann, F., Haberland, C., Jia, Y., and Pesaresi, D.: The Swath-D seismic network in Italy and Austria., GFZ Data Services. Other/Seismic Network, <https://doi.org/10.14470/mf7562601148>, 2017.
- 875 Hensch, M., Dahm, T., Ritter, J., Heimann, S., Schmidt, B., Stange, S., and Lehmann, K.: Deep low-frequency earthquakes reveal ongoing magmatic recharge beneath Laacher See Volcano (Eifel, Germany), *Geophysical Journal International*, 216, 2025–2036, 2019.
- Hetényi, G., Molinari, I., Clinton, J., Bokelmann, G., Bondár, I., Crawford, W. C., Dessa, J.-X., Doubre, C., Friederich, W., Fuchs, F., et al.: The AlpArray seismic network: a large-scale European experiment to image the Alpine orogen, *Surveys in geophysics*, 39, 1009–1033, 2018.

- 880 INGV Seismological Data Centre: Rete Sismica Nazionale (RSN). Istituto Nazionale di Geofisica e Vulcanologia (INGV), Italy, Other/Seismic Network, <https://doi.org/10.13127/SD/X0FXnH7QfY>, 2006.
- Institut De Physique Du Globe De Paris (IPGP), & Ecole Et Observatoire Des Sciences De La Terre De Strasbourg (EOST): GEOSCOPE, French Global Network of broad band seismic stations. Institut de physique du globe de Paris (IPGP), Université de Paris, Other/Seismic Network, <https://doi.org/10.18715/GEOSCOPE.G>, 1982.
- 885 Institute Of Geophysics, A. O. S. O. T. C. R.: Czech Regional Seismic Network. International Federation of Digital Seismograph Networks., Other/Seismic Network, <https://doi.org/10.7914/SN/CZ>, 1973.
- Jena, F.S.U.: Thüringer Seismologisches Netz (TSN). International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/TH>, 2009.
- Julian, B. R., Miller, A. D., and Foulger, G.: Non-double-couple earthquakes 1. Theory, *Reviews of Geophysics*, 36, 525–549, 1998.
- 890 Kagan, Y.: 3-D rotation of double-couple earthquake sources, *Geophysical Journal International*, 106, 709–716, 1991.
- Kästle, E. D., El-Sharkawy, A., Boschi, L., Meier, T., Rosenberg, C., Bellahsen, N., Cristiano, L., and Weidle, C.: Surface wave tomography of the Alps using ambient-noise and earthquake phase velocity measurements, *Journal of Geophysical Research: Solid Earth*, 123, 1770–1792, 2018.
- Keiding, M., Kreemer, C., Lindholm, C., Gradmann, S., Olesen, O., and Kierulf, H.: A comparison of strain rates and seismicity for Fennoscandia: depth dependency of deformation from glacial isostatic adjustment, *Geophysical Journal International*, 202, 1021–1028, <https://doi.org/10.1093/gji/ggv207>, 2015.
- Křížová, D., Zahradník, J., and Kiratzi, A.: Resolvability of isotropic component in regional seismic moment tensor inversion, *Bulletin of the Seismological Society of America*, 103, 2460–2473, 2013.
- Kövesligethy Radó Seismological Observatory (Geodetic And Geophysical Institute, Research Centre For Astronomy And Earth Sciences, Hungarian Academy Of Sciences (MTA CSFK GGI KRSZO)): Hungarian National Seismological Network. Deutsches Geoforschungszentrum GFZ, Other/Seismic Network, <https://doi.org/10.14470/UH028726>, 1992.
- 900 Kühn, D., Heimann, S., Isken, M. P., Ruigrok, E., and Dost, B.: Probabilistic Moment Tensor Inversion for Hydrocarbon-Induced Seismicity in the Groningen Gas Field, The Netherlands, Part 1: Testing, *Bulletin of the Seismological Society of America*, 110, 2095–2111, <https://doi.org/10.1785/0120200099>, <https://doi.org/10.1785/0120200099>, 2020.
- 905 Le Breton, E., Handy, M. R., Molli, G., and Ustaszewski, K.: Post-20 Ma motion of the Adriatic Plate: New constraints from surrounding orogens and implications for crust-mantle decoupling, *Tectonics*, 36, 3135–3154, <https://doi.org/https://doi.org/10.1002/2016TC004443>, 2017.
- Le Breton, E., Brune, S., Ustaszewski, K., Zahirovic, S., Seton, M., and Müller, R. D.: Kinematics and extent of the Piemont–Liguria Basin—implications for subduction processes in the Alps, *Solid Earth*, 12, 885–913, <https://doi.org/https://doi.org/10.5194/se-12-885-2021>, 2021.
- 910 Leipzig University: SXNET Saxon Seismic Network. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/SX>, 2001.
- Lizurek, G.: Full moment tensor inversion as a practical tool in case of discrimination of tectonic and anthropogenic seismicity in Poland, *Pure and Applied Geophysics*, 174, 197–212, 2017.
- Locati, M., Rovida, A., Albinì, P., and Stucchi, M.: The AHEAD portal: a gateway to European historical earthquake data, *Seismological Research Letters*, 85, 727–734, 2014.
- 915 Lu, Y., Stehly, L., Brossier, R., Paul, A., and Group, A. W.: Imaging Alpine crust using ambient noise wave-equation tomography, *Geophysical Journal International*, 222, 69–85, 2020.

- Marschall, I., Deichmann, N., and Marone, F.: Earthquake focal mechanisms and stress orientations in the eastern Swiss Alps, *Swiss Journal of Geosciences*, 106, 79–90, 2013.
- 920 Maurer, H., Burkhard, M., Deichmann, N., and Green, A.: Active tectonism in the central Alps: contrasting stress regimes north and south of the Rhone Valley, *Terra Nova*, 9, 91–94, 1997.
- MedNet Project Partner Institutions: Mediterranean Very Broadband Seismographic Network (MedNet). Istituto Nazionale di Geofisica e Vulcanologia (INGV), Other/Seismic Network, <https://doi.org/10.13127/SD/fBBBtDtd6q>, 1990.
- Mendiguren, J. A.: Inversion of surface wave data in source mechanism studies, *Journal of Geophysical Research*, 82, 889–894, 1977.
- 925 Meyer, B., Lacassin, R., Brulhet, J., and Mouroux, B.: The Basel 1356 earthquake: which fault produced it?, *Terra Nova*, 6, 54–63, 1994.
- Miller, A. D., Foulger, G., and Julian, B. R.: Non-double-couple earthquakes 2. Observations, *Reviews of Geophysics*, 36, 551–568, 1998.
- Ministrstvo za okolje in prostor Agencija RS za okolje: Potresi v letu 2016, Earthquakes in 2016, ARSO POTRESI, <https://www.arso.gov.si/potresi/poro%c4%8dila%20in%20publikacije/Potresi%20v%20letu%202016.pdf>, ISSN 1318 - 4792, 2018.
- Ministrstvo za okolje in prostor Agencija RS za okolje: Potresi v letu 2017, Earthquakes in 2017, ARSO POTRESI, <https://www.arso.gov.si/potresi/poro%c4%8dila%20in%20publikacije/Potresi%20v%20letu%202017.pdf>, ISSN 1318 - 4792, 2019.
- 930 Ministrstvo za okolje in prostor Agencija RS za okolje: Potresi v letu 2018, Earthquakes in 2018, ARSO POTRESI, <https://www.arso.gov.si/potresi/poro%c4%8dila%20in%20publikacije/Potresi%20v%20letu%202018.pdf>, ISSN 1318 - 4792, 2020.
- Minson, S. E. and Dreger, D. S.: Stable inversions for complete moment tensors, *Geophysical Journal International*, 174, 585–592, 2008.
- Mitterbauer, U., Behm, M., Brückl, E., Lippitsch, R., Guterch, A., Keller, G. R., Koslovskaya, E., Rumpfhuber, E.-M., and Šumanovac, F.:
935 Shape and origin of the East-Alpine slab constrained by the ALPASS teleseismic model, *Tectonophysics*, 510, 195–206, 2011.
- Molinari, I., Verbeke, J., Boschi, L., Kissling, E., and Morelli, A.: Italian and Alpine three-dimensional crustal structure imaged by ambient-noise surface-wave dispersion, *Geochemistry, Geophysics, Geosystems*, 16, 4405–4421, 2015.
- Montone, P., Mariucci, M. T., Pondrelli, S., and Amato, A.: An improved stress map for Italy and surrounding regions (central Mediterranean), *Journal of Geophysical Research: Solid Earth*, 109, 2004.
- 940 Moulin, A., Benedetti, L., Rizza, M., Jamšek Rupnik, P., Gosar, A., Bourles, D., Keddadouche, K., Aumâtre, G., Arnold, M., Guillou, V., et al.: The Dinaric fault system: Large-scale structure, rates of slip, and Plio-Pleistocene evolution of the transpressive northeastern boundary of the Adria microplate, *Tectonics*, 35, 2258–2292, <https://doi.org/https://doi.org/10.1002/2016TC004188>, 2016.
- Nicolas, M., Bethoux, N., and Madeddu, B.: Instrumental seismicity of the Western Alps: a revised catalogue, *pure and applied geophysics*, 152, 707–731, 1998.
- 945 OGS (Istituto Nazionale Di Oceanografia E Di Geofisica Sperimentale): North-East Italy Seismic Network. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/OX>, 2016.
- OGS (Istituto Nazionale Di Oceanografia E Di Geofisica Sperimentale) And University Of Trieste: North-East Italy Broadband Network. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/NI>, 2002.
- Panza, G. F. and Saraò, A.: Monitoring volcanic and geothermal areas by full seismic moment tensor inversion: Are non-double-couple
950 components always artefacts of modelling?, *Geophysical Journal International*, 143, 353–364, 2000.
- Patacca, E., Scandone, P., Di Luzio, E., Cavinato, G. P., and Parotto, M.: Structural architecture of the central Apennines: Interpretation of the CROP 11 seismic profile from the Adriatic coast to the orographic divide, *Tectonics*, 27, 2008.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine Learning in Python, *Journal of Machine Learning Research*, 12, 2825–2830, 2011.
- 955

- Petersen, G. M., Cesca, S., Kriegerowski, M., and the AlpArray Working Group: Automated Quality Control for Large Seismic Networks: Implementation and Application to the AlpArray Seismic Network, *Seismological Research Letters*, 90, 1177–1190, <https://doi.org/https://doi.org/10.1785/0220180342>, 2019.
- 960 Poli, M. and Zanferrari, A.: The seismogenic source of the 1976 Friuli earthquakes: a new seismotectonic model for the Friuli area, *Bollettino di Geofisica Teorica ed Applicata*, 59, 2018.
- Pondrelli, S.: European-Mediterranean Regional Centroid-Moment Tensors Catalog (RCMT) [Data set], Istituto Nazionale di Geofisica e Vulcanologia (INGV), <https://doi.org/10.13127/rcmt/euromed>, 2002.
- Pondrelli, S., Ekström, G., and Morelli, A.: Seismotectonic re-evaluation of the 1976 Friuli, Italy, seismic sequence, *Journal of Seismology*, 5, 73–83, 2001.
- 965 Pondrelli, S., Salimbeni, S., Ekström, G., Morelli, A., Gasperini, P., and Vannucci, G.: The Italian CMT dataset from 1977 to the present, *Physics of the Earth and Planetary Interiors*, 159, 286–303, 2006.
- Qorbani, E., Zigone, D., Handy, M. R., Bokelmann, G., AlpArray-EASI working group, et al.: Crustal structures beneath the Eastern and Southern Alps from ambient noise tomography, *Solid Earth*, 11, 1947–1968, 2020.
- Reiter, F., Freudenthaler, C., Hausmann, H., Ortner, H., Lenhardt, W., and Brandner, R.: Active seismotectonic deformation in front of the 970 Dolomites indenter, Eastern Alps, *Tectonics*, 37, 4625–4654, 2018.
- RESIF: RESIF-RLBP French Broad-band network, RESIF-RAP strong motion network and other seismic stations in metropolitan France [Data set]. RESIF - Réseau Sismologique et géodésique Français, Other/Seismic Network, <https://doi.org/10.15778/RESIF.FR>, 1995.
- RESIF: CEA/DASE broad-band permanent network in metropolitan France [Data set]. RESIF - Réseau Sismologique et géodésique Français, Other/Seismic Network, <https://doi.org/10.15778/RESIF.RD>, 2018.
- 975 Ritsema, J. and Lay, T.: Long-period regional wave moment tensor inversion for earthquakes in the western United States, *Journal of Geophysical Research: Solid Earth*, 100, 9853–9864, 1995.
- Rovida, A. and Locati, M.: Archive of historical earthquake data for the European-Mediterranean area, in: *Perspectives on European Earthquake Engineering and Seismology*, pp. 359–369, Springer, Cham, 2015.
- Schmid, S. M., Fügenschuh, B., Kissling, E., and Schuster, R.: Tectonic map and overall architecture of the Alpine orogen, *Eclogae Geologicae Helvetiae*, 97, 93–117, 2004.
- 980 Schmid, S. M., Bernoulli, D., Fügenschuh, B., Matenco, L., Schefer, S., Schuster, R., Tischler, M., and Ustaszewski, K.: The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units, *Swiss Journal of Geosciences*, 101, 139–183, 2008.
- Scognamiglio, L., Tinti, E., and Quintiliani, M.: Time Domain Moment Tensor (TDMT). Istituto Nazionale di Geofisica e Vulcanologia (INGV), [Data set], <https://doi.org/https://doi.org/10.13127/TDMT>, 2006.
- 985 Scott, D. R. and Kanamori, H.: On the consistency of moment tensor source mechanisms with first-motion data, *Physics of the Earth and planetary interiors*, 37, 97–107, 1985.
- Serpelloni, E., Vannucci, G., Anderlini, L., and Bennett, R. A.: Kinematics, seismotectonics and seismic potential of the eastern sector of the European Alps from GPS and seismic deformation data, *Tectonophysics*, 688, 157–181, 2016.
- Sileny, J. and Hofstetter, R.: Moment tensor of the 1999 Dead Sea calibration shot: Limitations in the isotropic source retrieval without a 990 detailed earth model, *Tectonophysics*, 356, 157–169, 2002.
- Slejko, D.: What science remains of the 1976 Friuli earthquake?, *Bollettino di Geofisica Teorica ed Applicata*, 59, 327–350, 2018.
- Sokos, E. N. and Zahradnik, J.: ISOLA a Fortran code and a Matlab GUI to perform multiple-point source inversion of seismic data, *Computers & Geosciences*, 34, 967–977, 2008.

- Spada, M., Bianchi, I., Kissling, E., Agostinetti, N. P., and Wiemer, S.: Combining controlled-source seismology and receiver function information to derive 3-D Moho topography for Italy, *Geophysical Journal International*, 194, 1050–1068, <https://doi.org/https://doi.org/10.1093/gji/ggt148>, 2013.
- 995
- Stähler, S. C. and Sigloch, K.: Fully probabilistic seismic source inversion—Part 1: Efficient parameterisation, *Solid Earth*, 5, 2014.
- Sternai, P., Sue, C., Husson, L., Serpelloni, E., Becker, T. W., Willett, S. D., Faccenna, C., Di Giulio, A., Spada, G., Jolivet, L., et al.: Present-day uplift of the European Alps: Evaluating mechanisms and models of their relative contributions, *Earth-Science Reviews*, 190, 589–604, <https://doi.org/https://doi.org/10.1016/j.earscirev.2019.01.005>, 2019.
- 1000
- Stich, D., Ammon, C. J., and Morales, J.: Moment tensor solutions for small and moderate earthquakes in the Ibero-Maghreb region: IBERO-MAGHREBIAN MOMENT TENSOR SOLUTIONS, *Journal of Geophysical Research: Solid Earth*, 108, <https://doi.org/10.1029/2002JB002057>, 2003.
- Storchak, D. A., Di Giacomo, D., Bondár, I., Engdahl, E. R., Harris, J., Lee, W. H., Villaseñor, A., and Bormann, P.: Public release of the ISC–GEM global instrumental earthquake catalogue (1900–2009), *Seismological Research Letters*, 84, 810–815, <https://doi.org/10.1785/0220130034>, 2013.
- 1005
- Storchak, D. A., Di Giacomo, D., Engdahl, E., Harris, J., Bondár, I., Lee, W. H., Bormann, P., and Villaseñor, A.: The ISC-GEM global instrumental earthquake catalogue (1900–2009): introduction, *Physics of the Earth and Planetary Interiors*, 239, 48–63, <https://doi.org/10.1016/j.pepi.2014.06.009>, 2015.
- 1010
- Stucchi, M., Rovida, A., Capera, A. G., Alexandre, P., Camelbeeck, T., Demircioglu, M., Gasperini, P., Kouskouna, V., Musson, R., Radulian, M., et al.: The SHARE European earthquake catalogue (SHEEC) 1000–1899, *Journal of Seismology*, 17, 523–544, 2013.
- Sue, C., Martinod, J., Tricart, P., Thouvenot, F., Gamond, J.-F., Fréchet, J., Marinier, D., Glot, J.-P., and Grasso, J.-R.: Active deformation in the inner western Alps inferred from comparison between 1972-classical and 1996-GPS geodetic surveys, *Tectonophysics*, 320, 17–29, 2000.
- 1015
- Swiss Seismological Service (SED) At ETH Zurich: National Seismic Networks of Switzerland. ETH Zürich, Other/Seismic Network, <https://doi.org/10.12686/sed/networks/ch>, 1983.
- University Of Genova: Regional Seismic Network of North Western Italy. International Federation of Digital Seismograph Networks., Other/Seismic Network, <https://doi.org/10.7914/SN/GU>, 1967.
- University Of Trieste: Friuli Venezia Giulia Accelerometric Network. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/RF>, 1993.
- 1020
- Valentine, A. P. and Trampert, J.: Assessing the uncertainties on seismic source parameters: Towards realistic error estimates for centroid-moment-tensor determinations, *Physics of the Earth and Planetary Interiors*, 210, 36–49, 2012.
- Van Hinsbergen, D. J., Torsvik, T. H., Schmid, S. M., Mañenco, L. C., Maffione, M., Vissers, R. L., Gürer, D., and Spakman, W.: Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution since the Triassic, *Gondwana Research*, 81, 79–229, <https://doi.org/https://doi.org/10.1016/j.gr.2019.07.009>, 2020.
- 1025
- Vavryčuk, V.: Moment tensor decompositions revisited, *Journal of Seismology*, 19, 231–252, 2015.
- Viganò, A., Bressan, G., Ranalli, G., and Martin, S.: Focal mechanism inversion in the Giudicarie–Lessini seismotectonic region (Southern Alps, Italy): insights on tectonic stress and strain, *Tectonophysics*, 460, 106–115, 2008.
- Wang, R.: A simple orthonormalization method for stable and efficient computation of Green’s functions., *Bulletin of the Seismological Society of America*, 89, 733–741, 1991.
- 1030

- Weber, J., Vrabec, M., Pavlovčič-Prešeren, P., Dixon, T., Jiang, Y., and Stopar, B.: GPS-derived motion of the Adriatic microplate from Istria Peninsula and Po Plain sites, and geodynamic implications, *Tectonophysics*, 483, 214–222, <https://doi.org/https://doi.org/10.1016/j.tecto.2009.09.001>, 2010.
- Wessel, P., Smith, W. H., Scharroo, R., Luis, J., and Wobbe, F.: Generic mapping tools: improved version released, *Eos, Transactions American Geophysical Union*, 94, 409–410, 2013.
- Zahradník, J. and Sokos, E.: Fitting waveform envelopes to derive focal mechanisms of moderate earthquakes, *Seismological Research Letters*, 89, 1137–1145, 2018.
- ZAMG-Zentralanstalt Für Meteorologie Und Geodynamik: Austrian Seismic Network. International Federation of Digital Seismograph Networks, Other/Seismic Network, <https://doi.org/10.7914/SN/OE>, 1987.