

Comparing global seismic tomography models using the varimax Principal Component Analysis

Olivier de Viron¹, Michel Van Camp², Alexia Grabkowiak³, and Ana M. G. Ferreira^{4,5}

¹Littoral, Environnement et Sociétés (LIENSs UMR7266 La Rochelle University – CNRS)

²Royal Observatory of Belgium

³Institut de Physique du Globe de Paris, University Paris Diderot

⁴Department of Earth Sciences, University College London

⁵CERIS, Instituto Superior Técnico, Universidade de Lisboa

Correspondence: Olivier de Viron (olivier.de_viron@univ-lr.fr)

Abstract. Global seismic tomography has greatly progressed in the past decades, with many global Earth models being produced by different research groups. Objective, statistical methods are crucial for the quantitative interpretation of the large amount of information encapsulated by the models as well as for unbiased model comparisons. We propose here to use a rotated version of the Principal Component Analysis (PCA) to compress the information, in order to ease the geological interpretation and model comparison. The method generates between 7 to 15 principal components (PC) for each of the seven tested global tomography models, capturing more than 97% of the total variance of the model. Each PC consists of a vertical profile, to which a horizontal pattern is associated by projection. The depth profiles and the horizontal patterns enable examining the key characteristics of the main components of the models. Most of the information in the models is associated with a few features: Large Low Shear Velocity Provinces (LLSVPs) in the lowermost mantle, subduction signals and low velocity anomalies likely associated with mantle plumes in the upper and lower mantle, and ridges and cratons in the uppermost mantle. Importantly, all models highlight several independent components in the lower mantle that make between 36% and 69% of the total variance, depending on the model, which suggests that the lower mantle is more complex than traditionally assumed. Overall, we find that the varimax PCA is a useful additional tool for the quantitative comparison and interpretation of tomography models.

1 Introduction

Global seismic tomography has brought a new understanding of the current state of the mantle, by inversion of massive seismic data sets to build 3-D images of the Earth's interior, both of isotropic and anisotropic structure, the latter being one of the most direct ways to constrain mantle flow (e.g., Rawlinson et al., 2014; Chang et al., 2014; McNamara, 2019). The interpretation and comparison of tomography models often include computing correlations between two models with depth and degree, analyzing power spectra (e.g., Becker and Boschi (2002)), or visual inspections and qualitative or simple descriptions of the retrieved patterns, for example of subducted slab or mantle plume candidates (e.g., Auer et al. (2014); French and Romanowicz (2014); Chang et al. (2016); Ferreira et al. (2019)). While the large-scale, upper mantle and lowermost mantle isotropic structure is fairly consistent from one model to the other, discrepancies appear when considering small-scale structures. Moreover, there

are substantial differences between existing global anisotropy models (e.g., Chang et al., 2014; Romanowicz and Wenk, 2017). Nowadays, codes or web-based tools facilitate the interpretation and visual comparison of different models (e.g., Durand et al. (2018), Hosseini et al. (2018)). This allows to identify regions with good agreement between seismic models using, e.g., vote maps (Lekic et al., 2012) or through statistical tools showing the relative frequency of seismic anomalies at specific depth ranges (Hosseini et al., 2018)). However, the large amount of information encapsulated in global tomography models, which typically involve tens of thousands of model parameters, can be difficult to mine and to interpret efficiently.

Statistical methods used in other disciplines to analyse and classify big data and models may be useful to further enhance the analysis of seismic tomography models, by providing a common ground for comparison. For example, in recent years, clustering methods have been used to partition seismic tomography models into groups of similar velocity profiles, providing an objective way of comparing the models (Lekic et al., 2012; Cottaar and Lekic, 2016). Here, we propose to ~~use~~ implement the Principal Component Analysis (PCA, Storch and Zwiers (1999)) to ~~further explore this type of approach~~ substantiate and clarify what can be learn from such comparisons, as recently proposed by Ritsema and Lekić (2020). The PCA-based method aims at approximating the tomographic models by a sum of a given number \tilde{N} of components, with \tilde{N} smaller than the actual number of slices. Each PC consists of ~~in~~ a vertical profile – the principal component (PC) – and a horizontal pattern – the load. Most of the variance of the signal being captured by a reduced number of PCs, it allows to grab all the information by analysing only the relevant components, resulting from an efficient compression.

~~However, even if~~ Although the first PC, capturing the largest variance, often corresponds to an actual physical process, ~~but~~ the others are increasingly difficult to interpret. The physical interpretation of the PCs and loads can be made easier by redistributing PCA components along other eigen-vectors. ~~In-particular,~~ We propose to apply the varimax criterion (Kaiser, 1958) that allows focusing on PCs with large values concentrated on the smallest possible subset of depths, as it is physically likely that mantle structures have a limited depth extension rather than spanning over the whole mantle depth. Previous studies in other fields of Earth sciences (see e.g. the thorough review paper by Richman (1986)) showed that, when using the varimax criterion, the redistributed components are often less sensitive to computation artifacts, for example related to data geometry, while keeping the same degree of compression as the original PCs.

The varimax analysis has previously been successfully used in various applications, such as to analyse climate models, where the different models are projected on the same set of PCs, allowing a direct comparison in terms of capture variance and retrieved features (Horel, 1981; Sengupta and Boyle, 1998; Storch and Zwiers, 1999; Tao et al., 2019; Kawamura, 1994). Motivated by these successful results, ~~in this study~~ we apply the varimax PCA to the interpretation and comparison of global tomography models. Note that we do not propose to use the varimax method as an alternative Earth model representation to the model solutions from which they are derived. Instead, we propose it as a diagnostic tool, allowing to quantify the level of independent information in the tomography models and to compare the models more easily. Projecting the models on a set of independent vertical profiles provides an optimal representation for a given number of slices (number of components), smaller than the original number of splines/boxes, or for a given portion of the information present in the model (captured variance).

In section 2, we present the seven global tomography models used, followed by a description of the statistical methods used in this study. Then, in section 4 we compare the classical and varimax PCA with a k-mean clustering approach. Sections 5–6

present and discuss the results from the application of the varimax PCA to the seven tomography models considered. We then propose a brief final discussion and conclusions in Section 7.

60 2 Seismic tomography models

We use seven 3-D global seismic tomography models: (i) S20RTS (Ritsema, 1999), 1999; (ii) S40RTS (Ritsema et al., 2011); (iii) SEISGLOB2 (Durand et al., 2017); (iv) SEMUCB-WM1 (French and Romanowicz, 2014); (v) SGLOBE-rani (Chang et al., 2015); (vi) S362WMANI+M (Moulik and Ekström, 2014); and, (vii) SAVANI (Auer et al., 2014). While the first three models are isotropic shear-wave speed models, the last four models also include lateral variations in radial anisotropy. These models were built from different data sets and using distinct modelling approaches, as summarised in Table 1. We focus on shear-wave models because the agreement of P-wave models is more limited (e.g., Cottaar and Lekic (2016)). **Nevertheless, future work may expand the analysis to recent P-wave models (Hosseini et al., 2020)** The models used show the key features in current global isotropic and radially anisotropic models, and hence are representative of the current state of global tomography. For example, all isotropic shear-wave speed models show a good correlation with tectonic features in the upper mantle, such as mid-ocean ridges and cratons (see ~ 100 km in Figure A1a). Moreover, they show the signature of subducting slabs around ~ 600 km depth, as well as the two prominent large low shear velocity provinces (LLSVPs) beneath Africa and the Pacific in the bottom of the mantle at $\sim 2,900$ km depth (Figure A1a). On the other hand, the agreement between the anisotropy models is much more limited (Figure A1b); common features between the models include a well-known positive radial anisotropy anomaly beneath the Pacific at ~ 150 km depth, negative radial anisotropy anomalies beneath the East Pacific Rise at ~ 200 km depth and negative radial anisotropy anomalies associated with the LLSVPs. The latter anomalies have been shown to be artefacts in the models due to the poor balance between SV- and SH-sensitive traveltimes in various existing body-wave data sets, which have much more data sensitive to SH- than to SV motions (e.g., Kustowski et al. (2008); Chang et al. (2014)). On the other hand, Moulik and Ekström (2014) showed that such spurious anisotropic features in even degree structure are reduced by using self-coupling normal mode splitting data in the inversions. Yet, trade-offs between isotropic and anisotropic structure in the lowermost mantle persist in odd degree structure, which is not constrained by self-coupling normal mode splitting data.

We obtained the global tomography models either directly from their authors or from the IRIS Earth model collaboration repository (REFS - <http://ds.iris.edu/ds/products/emc/>). Since some of the models have different reference 1-D models and use distinct parameterisations, for consistency we converted them into perturbations in shear-wave speed and in radial anisotropy with respect to the 1-D model PREM (Dziewonski and Anderson, 1981) on a common grid with a $1^\circ \times 1^\circ$ horizontal sampling and on 29 depth slices starting at 50 km depth and with a 100 km spacing from 100 km to 2900 km depth. **This conversion into the same 1-D reference model eases comparison. Moreover, since both the vertical structure and the horizontal patterns are normalized, this also implies that the background model does not impact our analysis.** In addition to the gridded representation allowing fair graphical presentation, we interpolate the data sets from the horizontal grids into an regular polyhedron of 9002 equiareal faces of which the vertexes were generated through the icosahedron tool of Zechmann (2019). This transformation

produces an uniform sampling of the sphere, with each vertex having a surface corresponding to that of a $250 \times 250 \text{ km}^2$ square. Considering that the statistical methods used in this study use the captured variance as a major criterion for ordering the components, the use of gridded data would overweight the contribution at the poles in the PC representation. The $180 \times 360 \times 29$ individual model data matrices are thus converted into 9002×29 matrices. As the shallowest layers contain the majority of the variability in the velocity anomalies, most of the principal components will be captured in those layers, which will be over-represented. Hence, the shear-wave speed and radial anisotropy perturbations are normalized by slice, i.e., the mean value of the slice is subtracted from each value, and each value is divided by the standard deviation of the slice. This is relevant as, in this study, we investigate relative values on a given profile; the actual magnitude can be recovered by multiplying the load patterns by the standard deviation of the layer in the original model. The normalization applied to the models does not lead to a loss of information.

3 Methods

Previous studies have compared global tomography models using k-mean clustering (e.g., Lekic et al. (2012)) or PCA (Ritsema and Lekić, 2020). Though the PCA is very different on many aspects from the clustering of the k-mean method, it is useful to start by comparing PCA and varimax PCA results with those from the k-mean, for an illustrative tomography model (S40RTS).

3.1 k-mean clustering

Considering the three-dimensional data set $D(\lambda_i, \phi_i, z_j)$, the k-mean algorithm (MacQueen, 1967) defines k clusters, corresponding to the k sets of horizontal positions (λ_i, ϕ_i) closest to their average z_i profiles. The algorithm is based on an iterative procedure. At the first iteration, it randomly chooses k horizontal positions, used as cluster centers. Each point of the data set is then associated with the cluster center to which it is the closest to. The average radial profile of the points attributed to each given cluster is computed and used as the new cluster center for the next iteration. This is repeated until convergence is achieved.

To make k-mean and PCA representations somewhat comparable, we use the clusters as horizontal patterns and the average vertical profiles of each cluster as the PCs. By construction, the variance captured by the k-mean is noticeably smaller than that for the other methods, since it is not meant to propose a compressed representation of the data set, but rather to separate the data set into subsets, which results in an important loss of information.

3.2 Principal Component Analysis (PCA)

A 2D matrix $F_{j,k}, j = 1, 2, \dots, J; k = 1, 2, \dots, K$ is transformed by the PCA into a sum of components, each component being composed of a load $\alpha_{n,j}$ and an eigen-vector $A_{n,k}$.

$$F_{j,k} = \sum_{n=1}^{N=\min(J,K)} \alpha_{n,j} A_{n,k}.$$

Model	Dataset	Parameterisation	Modelling approach
SGLOBE-rani (Chang et al., 2015)	Fundamental mode group-velocity data ($T \sim 16-150$ s), fundamental and overtone phase-velocity data (25-374 s, up to the 4th overtone), body-wave travel times.	Spherical harmonics up to degree 35 (laterally) and 21 spline functions (radially). 1-D reference model: PREM.	Ray theory. regularization: norm damping.
SAVANI (Auer et al., 2014)	Fundamental and overtone phase velocities ($T \sim 25-370$ s, up to the 4th overtone), body-wave travel times.	Variable size blocks on a $5^\circ \times 5^\circ$ base grid (laterally). 28 variable thickness depth layers. 1-D reference model: PREM.	Ray theory. regularization: vertical and horizontal smoothing.
S20RTS (Ritsema, 1999)	Fundamental mode and overtone phase velocities ($T \sim 40-275$ s, up to the 4th overtone), body-wave travel times, even-degree self-coupling normal mode splitting functions.	Spherical harmonics up to degree 20 (laterally) and 21 spline functions (radially). 1-D reference model: PREM.	Ray theory. regularization: norm damping.
S40RTS (Ritsema et al., 2011)	Expanded dataset of S20RTS (increased amount of measurements).	Spherical harmonics up to degree 40 (laterally) and 21 spline functions (radially). 1-D reference model: PREM.	Ray theory. regularization: norm damping.
SEMUCB-WM1 (French and Romanowicz, 2014)	Body waveforms ($T > 32$ s), surface waveforms ($T > 60$ s).	Spherical splines with spacing $< 2^\circ$ (laterally) and 20 cubic b-splines (radially). 1-D reference model: own model.	Spectral element method for forward modeling, non-linear asymptotic coupling theory for inverse modelling. regularization: vertical and horizontal smoothing.
S362WMANI+M (Moulik and Ekström, 2014)	Fundamental mode phase velocities ($T \sim 35-150$ s), body wave traveltimes and waveforms, normal mode splitting functions.	362 spherical splines (laterally) and 16 cubic splines (radially). 1-D reference model: STW105 (Kustowski et al. (2008)) .	Ray theory. regularization: vertical and horizontal smoothing.
SEISGLOB2 (Durand et al., 2017)	Fundamental and overtone phase velocities ($T \sim 40-360$ s, up to the 5th overtone), body-wave travel times, normal mode self- and cross-coupling coefficients.	Spherical harmonics up to degree 40 (laterally) and 21 spline functions (radially)	Ray theory. regularization: lateral smoothing controlled by an horizontal correlation length.

Table 1. Global tomography models used in this study, including a short description of the data, parameterisation and the modelling approach used in their construction. All models were built using least-squares inversions with different regularization choices.

In our case, $F_{j,k}$ corresponds to the velocity anomaly at horizontal position $p_j = (\lambda_j, \phi_j)$ and depth z_k . The $A_{n,k}$ are the eigen-vectors, or principal components (PC), of the covariance matrix. These PCs are orthogonal vertical structures representing the covariance between the slices of the model. It has large positive values if the horizontal structures from two layers are positively
120 correlated, zero values if the structures are not correlated, and large negative values if they are anti-correlated. The loading patterns $\alpha_{n,j}$ are also orthogonal to each other and each $\alpha_{i,j}$ results from the projection of the data set on the PC i , capturing the horizontal structure i associated with the vertical anomaly profile $A_{i,k}$. The loads take continuous positive and negative values. Here, those patterns correspond to horizontal maps showing where each vertical structure is more or less important in the model.

125 The components are ordered by decreasing eigen-value, as the variance captured by each PC is directly proportional to the eigen-value of the PC. Due to their orthogonality and to the mathematical properties of the transformation, the variance captured by each PC drops rapidly with the order, so that a small number of independent components ($\tilde{N} \ll N = 29$) is often sufficient to capture most of the information, allowing an efficient compression of the data set.

130 Unlike clustering methods, which are binary in that any horizontal location only belongs to one cluster, the PCA computes the amplitude of the contribution from each principal component, for every horizontal location, providing a compressed reconstruction of the data set.

The first PC corresponds to the dominant covariance, which might be physically associated with a global phenomena – in our case, a structure that would develop on the whole mantle depth – or a more local feature, i.e., associated with a limited depth range. But this covariance structure might also correlate with other features from other depths, which will also be retrieved in
135 the first PC. The second PC being orthogonal to the first, part of the physics might have been subtracted by the computation of the first PC, and it is even more so for the following principal components.

3.3 Varimax PCA

Used on space-time datasets, the PCA often produces artefacts from the domain geometry (Richman, 1986): the topographies of the PCs are primarily determined by the shape of the domain and not by the covariation among the data. In other words,
140 different correlation functions on a geometrically shaped domain have similar load patterns in a predictable sequence, which do not reflect the underlying covariation. This is the case of square domains found for example on meteorological maps, or, in our case, the unrotated PCs show a harmonic-like progression, with one maximum for the first PC, two for the second, etc. In contrast, the varimax rotation captures distinct, well-defined depth domains in the mantle, which are easier to interpret physically. Hence, the physical interpretation of the PCs and load can be made easier by redistributing PCA components
145 along other eigen-vectors, by maximizing a functional of the loads that favors some physical properties that appear physically meaningful (Storch and Zwiers, 1999; Neuhaus and Wrigley, 1954). There are several possible redistribution options for the PCs (Storch and Zwiers, 1999; Browne, 2001; Jolliffe, 2005). Among those, the varimax rotation (Kaiser, 1958) favors PCs with large values concentrated on the smallest possible subset of depths and preserves the orthogonality of the PCs.

The varimax rotation corresponds to a rotation of the basis of the same information space as that generated by the PCA; consequently, the total variance captured by \tilde{N} components is exactly the same for both types of PCs, but distributed differently

between the components, with a slower variance decrease for the varimax rotated solution. ~~Both PCs and varimax rotated PCs compress the data set to the same level.~~ Considering the variance captured by the PCA components, the number \tilde{N} of PCs to keep is selected to meet a given criterion: the total variance captured, a fixed number of components, or the minimum variance captured by a PC kept. Then we define new \tilde{N} components, so that

$$F_{j,k} \simeq \sum_{n=1}^{\tilde{N}} \alpha_{n,j} A_{n,k} = \sum_{n=1}^{\tilde{N}} \beta_{n,j} B_{n,k}.$$

$\beta_{n,j}$ are the new horizontal structures and the new PCs, $B_{n,k}$, are chosen to maximize a given objective function V , defined by the sum of the values of the objective functions V_n computed over each PC:

$$V = \sum_n^{\tilde{N}} V_n(B_{n,k}),$$

with

$$V_n = \frac{1}{K} \sum_{k=1}^K \left(\frac{B_{n,k}}{s_k} \right)^4 - \frac{1}{K^2} \sum_{k=1}^K \left(\frac{B_{n,k}}{s_k} \right)^2.$$

where the s_k are normalization factors, with $s_k = 1$, for all k in the case of the varimax rotation.

150 This transformation corresponds to a rotation of the PCs because the subspace generated by the transformation – or the reconstructed model – is the same as with the non-rotated PCA limited to \tilde{N} components. In our case, it limits the vertical extension of the PCs, i.e., each PC shows large values on only a few depths/slices.

The associated horizontal structures, $\beta_{n,j}$, are recomputed by projection of the tomography model on the rotated vertical profile. This rotation conserves the orthogonality of the eigen-vector, which is not the case for all the possible rotations, but the
 155 horizontal loads are often not orthogonal anymore (Mestas-Nuñez, 2000). The total variance captured by the rotated PCs is the same as that from unrotated PCs, but the decrease of the variance is slower than that from the original decomposition.

4 Comparing the PCA, varimax rotated PCA, and k-mean results for the model S40RTS

Figure 1 shows the PCs and loads resulting from the application of the PCA, varimax rotated PCA, and k-mean clustering methods to the tomography model S40RTS for a 6-PC decomposition. All the methods are applied on the same normalized
 160 data. ~~The x-axis, i.e. the amplitude of the vertical eigenvectors, represents the maximum absolute value of the normalized anomaly at a given depth. It must be multiplied by the horizontal loading pattern, which provides normalized loads ranging between -1 and 1.~~

Figure 1 highlights the complementary of the k-mean and varimax PCA methods. While the k-mean method allows to highlight key large-scale features, such as e.g. the distribution of lowermost mantle velocity and to compare how it appears in the different models (e.g., Lekic et al., 2012), the varimax-PCA approach provides a compressed representation of the full
 165 model. Being binary, the k-mean method does not provide amplitudes, i.e., the load for each PC is either 0 or 1: every location is part of one k-mean cluster, while it can be part of several PCs in the PCA analysis. The latter captures inherently more

complexity with fewer principal components than k-mean clustering. Hence, it is not surprising that the six k-mean components capture only 34% of the total model variance, whereas both PC-based methods recover 83.2% (Figure 1). Therefore, while the
170 k-mean method is a useful classification technique that allows a subset of data to be separated out from others, the varimax PCA is a distinct, valuable compression technique that reduces the number of parameters while minimizing loss of information. It allows identifying the most important components of tomographic models, easing their interpretation.

As ~~suggested Riehmman (1986) expected~~, the PCA profiles $A_i(z)$ show increasingly oscillating patterns with i , which may lead to nonphysical interpretations. For example, the signature of ~~tectonic patterns such as ridges, subduction zones~~ and cratons
175 spread over the whole mantle (principal components 4 and 5, purple and cyan in the top row in Figure 1) observed in the unrotated PC representation makes no physical sense. More generally, the vertical profiles retrieved by PC analysis are certainly not all associated with sound geophysical structures. Only the first PC (red) provides directly interpretable patterns: the African and Pacific Large Low Shear Velocity Provinces (LLSVPs) (Garnero and McNamara, 2008; McNamara, 2019), whose depth extent with a maximum below 1800 km can be nicely visualised in Figure 1. ~~Without normalization, every eigen-vector shows~~
180 ~~a strong contribution from the shallowest mantle (<250km), where most of the variance is located, as in Ritsema and Lekić (2020). Indeed, due to the absence of normalization, all the main four PCs obtained by Ritsema and Lekić (2020) contain a lot of energy in the shallowest zone, while our normalization allows keeping the tectonic-driven zone in essentially two components out of 6 (component 4 capturing 7.6% of the variance and component 5 capturing 6.2%).~~

On the other hand, ~~even with this reduced number of 6 components, the normalized varimax procedure recovers well-known~~
185 ~~structures. Within its first mode, we recover the tectonic patterns such as ridges, cratons, oceanic plates, subduction zone, while the LLSVPs gradually appear in the three last components~~ (for a more detailed analysis see the next section). Based on these comparisons, we find that the varimax PC method is useful to concentrate at different depths coherent information that is available in the seismic tomography models, without any preconception. The next sections will thus focus on the application of this method to the interpretation of the global tomography models considered in this study.

190 5 Compressed information from varimax PCA

5.1 Comparison of vertical profiles and horizontal patterns

We use varimax PCA to compress the seven tomography models described in Section 2 into a set of components, keeping only the most important ones, as explained below. Each component is composed of a vertical profile obtained directly from the varimax process and of an associated horizontal pattern, which is computed by projecting the model on the profile. Such data
195 compression is useful to compare the models if three major conditions hold. First, a subset of components must capture most of the variance of the signal, with the number of components being significantly smaller than the original number of depth splines/boxes, and enhancing the signal-to-noise ratio. Secondly, the relevant structures in the mantle, which will be used for comparison and for geological interpretation, should not be distorted by the compression, i.e., their shape and position must remain unchanged. Third, the power spectral densities should not be altered by the compression process.




Figure 1. Results from the PC, varimax and k-mean methods applied to the model S40RTS, using 6 principal components. The varimax PC and loads are normalized in such a way that the horizontal patterns (load) range between -1 and 1, with orange corresponding to negative, blue to positive, and white to zero, and with the intensity being proportional to the value. We note that while this normalization eases the analysis, it does not lead to a loss of information (see main text for details). Unlike the classical variance-based sorting of the principal components, we order the varimax components by the depth of the profile's maximum.

200 In order to facilitate the comparison of the horizontal structures in the models, we label the varimax components obtained from the varimax PCA using capital letters in alphabetical order from components sensitive to shallow mantle structure to components sensitive to the lowermost mantle structure. Figure 2 shows the variance captured by the varimax PCA applied to the different models. The varimax analysis is performed by considering all the components capturing more than 1% of the variance of the signal in the classical PCA. Keeping only the components with variance above 1% limits the number of maps, 205 facilitating the quest for relevant information. Tests with 5% and 10% thresholds showed that some important information is lost. When using a 10% threshold, all the models are represented by 3 components only, which misrepresent known structures such as ridges or subduction zones. Moreover, the depth distributions of the corresponding PCAs become quite broad and imprecise, each stretching over more than 1000 km depth ranges. In the 5% threshold case, the models are represented by 4 (SAVANIi) to 6 components (SEMUCB-WM1i, SEISGLOB2). Again, information is lost concerning e.g., ridges or subduction 210 zones, and the depth information is spread over a depth range greater than 500 km for all components and for all models.

The simplification brought by the varimax method is particularly efficient for tomography models with weak regularization, such as, e.g., SGLOBE-ranii, where short-scale structure is likely mixed with noise. Figure 2 shows that the number of varimax components ranges from 7 for model SAVANIi to 15 for SGLOBE-ranii, with the total variance captured by all these principal components always exceeding 97.3% (see also Table 2, which summarises the components kept in the varimax analysis). The 215 number of varimax components required by each model depends on the details of the model's construction, such as e.g. the data used (Table 1) and, importantly, on subjective choices made, such as on the level of regularization used. Increasing the strength of regularization reduces the model's effective number of free parameters and hence the number of varimax components required by the model. As the SAVANI model only needs 7 varimax PCA components, the shallowest component concentrates a lot of information (26% of the variance) that is spread into more components for the other models. Table 2 shows that the 220 number of PCs needed to explain 97.3% or more of the total information in the tomography models is always smaller than the number of splines or layers used in the models' original depth parameterisation, with 29% to 75% fewer PCs than depth splines/layers.

This fulfills the first condition for the usefulness of the data compression mentioned above. In order to check the second condition previously mentioned, Figure B1 compares 6 examples of depth slices in the original SEMUCB-WM1i model with 225 those obtained from the model's reconstruction using varimax analysis. The differences between the original and reconstructed models are small and random, highlighting that the compression process does not distort the model's features. [The residual signal is the part of the models that is not co-variant vertically.](#) Finally, We also verified that there are only very small, random differences between the power spectra of the original and of the reconstructed models (see Figures B2 and B3 in the supplementary material). Hence, the third condition of usefulness of the data compression used in this study is also satisfied.

230 Figure 3 shows the varimax PCs for all the tomography models used in this study, together with the spline functions or the variable thickness depth layers used in the models' parameterisation. The vertical profiles differ from one model to the other both in numbers of components and in the depths of their maxima. For example, as shown in Figure 2, SAVANIi requires 7 components, while SGLOBE-ranii needs 15 components. We re-emphasise that the number of principal components obtained




Figure 2. Captured variance by the PC varimax method, applied to the isotropic part of the seven tomography models used in this study. The number of components is chosen such that during the PCA, we only keep the components explaining more than 1% of the variance, which occurs after 7 (SAVANIi) to 15 (SGLOBE-ranii) components. The varimax PCs are sorted alphabetically from the shallowest one to the deepest ones.

for a given model reflects their amount of independent information, which in turn depends heavily on choices made during the model's construction, such as regularization, and amount and type of data used.

The horizontal patterns obtained from the varimax PCA also show distinct features, but this is not always in the same way as for the vertical profiles (Figures 4, C1-C6). S20RTS shows sharp vertical profiles (Figure C3), and contains even one more PC (13) in the upper mantle than the updated S40RTS model (12, Figure C4), but the horizontal patterns are smoother. This is likely due to the fact that the latter model is constrained by about 10 times more data and used a different level of regularization (Ritsema et al., 2011). SGLOBE-ranii, SEMUCB-WM1i and SEISGLOB2 (Figures C1, 4 and C6) depict sharper horizontal patterns than SAVANIi, S362WMANI+Mi and S20RTS (Figures C2, C5 and C3). SEMUCB-WM1i and SAVANIi (Figures 4 and C2) show vertical profiles concentrated closer to the surface, but their horizontal patterns are different. The PC B of SAVANIi (~200 km depth, Figure C2) corresponds to low velocity anomalies underneath all oceans, which is also the case for S20RTS and S362WMANI+Mi (Figures C3 and C5), but not for SGLOBE-ranii and SEISGLOB2 (Figures C1 and C6). Such upper mantle low velocities beneath the oceans also appear in the models S40RTS and SEMUCB-WM1i, but with a higher level of detail (Figures C4 and 4).




Figure 3. Principal components for the different individual model varimax-PCA and the combined one (see section 6), for the isotropic part of the 7 tomography models used in this study. Only the PC components above 1% are kept in this analysis (Figure 2). The dashed grey lines represent the spline functions and the grey boxes the variable thickness depth layers used by the different models. The vertical dashed lines indicate the 410 and 660 km seismic discontinuities.

We note on Figure 3 that the vertical components obtained from the varimax analysis do not fully correspond to the depth parameterisations used to build the models, especially above the 660 km discontinuity, where there are 2.3 to 3.3 lesserfewer PCs than original spline functions or boxes. This implies that the PCs do not just simply reflect the model parameterisation and inform us about the independence between the slices reconstructed from the model. The PCA objectifies the number of splines/boxes consistent with the amount of information present in the model. In the upper mantle, for all models, we end up with 3 to 4 PCs. In the lower mantle, the correspondence differs from one model to the other, whereas we observe three categories:

1. SGLOBE-rani, where the PCA reproduces quite well the original spline functions except the two deepest ones, which are recovered into one PCA. This is probably due to the relatively weak regularization used (Chang et al., 2015).
2. SEISGLOB2, S20RTS and S40RTS, for which most of the PCs reproduce the splines. For S20RTS and S40RTS, one PC encompasses the depth associated with two splines between 1500 and 2000 km, while the first spline just underneath the 660 km discontinuity is not taken over by any PC. For SEISGLOB2 there is one PC for two splines between 800 and 1000 km.
3. S362WMANI+Mi, SAVANI and SEMUCB-WM1i: 10 splines are encompassed by 5 modes for SEMUCB-WM1i and 8 splines by 5 modes for S362WMANI+Mi. For SAVANIi, the structure of the PCs seems independent from the box parameterisation.

This shows that overall the tomography models do not have a strong imprint of the depth regularization used in their construction. This is especially true above the 600 km discontinuity or in the whole mantle for the SAVANIi, S362WMANI+Mi and SEMUCB-WM1i models, where the information is recovered by fewer PCs than the number of spline functions or boxes.

One of the most striking differences between the models is the way the signal is distributed between 500 and 1,500 km depth. In this region the different tomography models require between 2 (SAVANIi [D-E]) and 7 PCs (3 for S362WMANI+Mi [C-E] and SEMUCB-WM1i [D-F]; 5 for SEISGLOB2 [D-H]; 6 for S20RTS [D-I] and S40RTS [C-H]; and 7 for SGLOBE-ranii [D-J]). The observed variability in the number of required PCs likely reflects the level of model regularization used in the construction of the various tomography models as well as the amount and variety of data used (e.g., if the model does not include constraints from overtones, structures in the transition zone and mid-mantle might not be well-retrieved). Moreover, the treatment of discontinuity topography may also matter because neglecting this topography could map directly into isotropic wave speed variations in the mid-mantle. . The PC E of SAVANIi (~1200 km depth) is dominated by low-velocity anomalies and shows a substantially different pattern to e.g. the PC F of SEMUCB-WM1i and the PC G of SEISGLOB2 (~1300 km depth), which depict alternating low and high velocity zones. The principal components G (~1000 km depth) and H (~1100 km depth) of SGLOBE-ranii present mostly low velocity anomalies, which are similar to the components F of S40RTS and G of S20RTS, both at ~1100 km depth, but in these two latter models we also observe a high-velocity anomaly under the north-west Indian ocean. In SAVANIi, this is also observed on its PC E (~1200 km depth), which is however much more broadly distributed at depth. These differences between the models reflect the high level of uncertainty for this part of the mantle, which is likely due to its limited data coverage.




Figure 4. 9 Varimax components of the SEMUCB-WM1i model. On the right, the principal components or vertical profiles, and on the left, the associated horizontal structures. The other models are shown in the supplementary material on Figures C1-C5.

5.2 Geophysical interpretation

A fully detailed geological and geophysical discussion of the models is beyond the scope of this study, and has already been performed in many previous studies (see e.g., McNamara (2019); Flament et al. (2017); Ballmer et al. (2015); Pavlis et al. (2012); van der Meer et al. (2018); Rudolph et al. (2015); Ritsema et al. (2011)). The varimax PCA recovers all the features
285 discussed in these studies. Table 3 compares how key Earth structures are captured by the varimax analysis for the isotropic part of the seven tomography models used in this study (Figures 4, C1 - C5). The table covers ridges, rifts, plateaus (low-velocity anomalies, red) and cratons (high-velocity anomalies, blue) at depths of 50 – 300 km, which corresponds to the heterosphere (Dziewonski et al., 2010), subducted slabs (high-velocity anomalies) at 300 – 1300 km depth, and the LLSVPs and the Perm low velocity zone in the lowermost mantle (red anomalies).

290 Depending on the regions, the high velocity craton signature should reach a maximum depth between 100 and 175 km (Begg et al., 2009; Heintz et al., 2005; Polet and Anderson, 1995), but tomography models often show a deeper signature, likely due to smearing effects. The model SGLOBE-ranii is the most consistent with this depth limit, as most of the cratons are concentrated in the second PC (~ 100 km depth, PC B on Figure C1), separating them from the cold oceanic crust. This is possibly due to the huge set of data sensitive to the upper mantle used in SGLOBE-ranii's construction, including massive sets of both phase
295 and group velocity measurements. Beneath Africa and the Baltic region, a high velocity zone remains visible on the third PC (~ 300 km depth, PC C on Figure C1). For the other models, the craton signatures extend from 50 to ~ 200 -300 km depth.

The low velocity zones underneath the Tibetan plateau (Legendre et al., 2015) and Hangai dome, south-west of the Baikal lake (Chen et al., 2015), are recovered by the first PC of all models with a maximum at 50 km, but with different shapes. These zones are smaller in SEMUCB-WM1i (Figure 4) than in SGLOBE-ranii and SAVANIi (Figures C1 and C2), and the Tibetan
300 plateau extends more to the south in SAVANIi and S362WMANI+Mi (Figures C2 and C5), being subdivided into 3 small zones in S40RTS and SEMUCB-WM1i (Figures C4 and 4).

From ~ 150 km to ~ 800 km depth, global tomography models often show a low velocity anomaly beneath the Pacific (e.g., Lebedev and van der Hilst, 2008). For SAVANIi, at ~ 200 km depth, it is difficult to distinguish between the Pacific and other oceans. Its PC C, with a maximum at ~ 400 km depth, is confined to the central and western Pacific, while the PC
305 C of S362WMANI+Mi, at ~ 500 km depth, is more to the south west, resembling the PC D of S20RTS (~ 600 km depth), D of S40RTS (~ 700 km depth), E of SGLOBE-ranii (~ 600 km depth), D of SEMUCB-WM1i (~ 600 km depth) and D of SEISGLOB2 (~ 700 km depth).

All models show a high velocity zone between ~ 300 and 700 km depth beneath the central Atlantic and along the Atlantic coasts of South America and Africa, similar to that from Figure 9 of Ritsema et al. (2011), and already discussed in Ritsema
310 et al. (2004). Nevertheless, this zone appears less clearly in SEMUCB-WM1i (Figure 4 [C-D]) and especially SGLOBE-ranii (Figure C1 [C-D]), where it is mixed with low velocity patches. This anomaly is a region with long transform faults, high gravity, anomalous ocean depth, and low melt production. It is thought to be the region of the Atlantic that formed during the final stages of the opening of the Atlantic because it was presumably the strongest part of the Pangean continent (Bonatti, 1996).

315 In the East African rift, the low velocity anomaly aligned with the Afar Depression and the Main Ethiopian Rift in the uppermost mantle (Benoit et al., 2006), Hansen and Nyblade (2013) appears in all the models, from the surface to the LLSVP, with narrower contours in S40RTS, SEMUCB-WM1i, and SGLOBE-ranii than for the other models. This is consistent with the presence of one or of multiple mantle plumes in the region, as proposed in previous studies (e.g., Hansen et al. (2012); Chang and Van der Lee (2011); Chang et al. (2020)).

320 All models show high-velocity subduction zones in the western Pacific, among others, notably underneath the Philippine plate over two principal components with depths ~ 400 -800 km. This complex system mixes different subduction zones (van der Meer et al., 2018). The Izu-Bonin slab subducts westerward down to ~ 870 km depth and is connected in the upper ~ 300 -400 km depth to the Marianas to the south, which plunges vertically down to ~ 1200 km depth (components [C-E] in Figure 4, [C-G] in Figure C1, [C-D] in Figure C2, [D-F] in Figure C3, [C-D] in Figure C4, [C-D] in Figure C5, and [C-F] in Figure C6)

325 More to the south, north of Papua New Guinea, the Caroline Ridge, from ~ 475 to 750 km, and, west of those zones, Manila, Sangihe, and even more west, Banda, Sumatra, and Burma, also present high velocity anomalies. This is recovered by SGLOBE-ranii (components [D-E] in Figure C1), SEMUCB-WM1i (components D in Fig. Figure 4) and SEISGLOB2 (components D in Figure C6), but is broader, especially to the north, in S20RTS (components [D-E] in Figure C3), S40RTS (components C in Figure C4), SAVANIi (components [C-D] in Figure C2), and S362WMANI+Mi (components C in Figure
330 C5).

The Tonga-Kermadec subduction zone, located below the south Fiji Basin down to a depth of ~ 1300 km in the lower mantle (van der Meer et al., 2018), is recovered by all models, but is less clear in S362WMANI+Mi (Figure C5). Conversely, SGLOBE-ranii, S40RTS, SEISGLOB2 and to a lower extent, SEMUCB-WM1i, show a narrow arc-shaped signature of this zone. On the other hand, it is difficult to assess a maximum depth of this subduction zone in SGLOBE-ranii.

335 All models evidence the LLSVPs, though they are less clear in some models, such as SAVANIi and S362WMANI+Mi (Figure C5), and they appear quite patchy in S40RTS and SGLOBE-ranii (Figures C4 and C1). All the models show low velocity anomalies spreading from the core-mantle boundary (CMB), where the LLSVPs are clearly visible, to about 1500 km depth, where low-velocity structures are less coherent (for example, components G-I in Figure C1). All together, the components encompassing the LLSVPs capture 11% (SAVANIi) to 29% (S40RTS) of the models' information. The Perm
340 anomaly is recovered in all models for the two deepest components, apart for SAVANIi, where it is recovered by the last PC only. This is because this PC is quite broad, extending from ~ 2000 km depth to the CMB. These depths are consistent with, e.g., the findings of Lekic et al. (2012) and Flament et al. (2017), which estimate that LLSVPs spread up to ~ 500 km above the CMB. Note that it is difficult to estimate the top of the LLSVPs on S362WMANI+Mi, as there are persistent low velocity zones beneath e.g. eastern Europe up to the PC E centered at ~ 1300 km depth (Figure C5)

345 Our analysis allows determining the importance of the various elements of the models. For example, for all models, principal components with maxima in their varimax PCs below 1,700 km depth and dominated by LLSVPs explain 11% (SAVANIi) to 24% (SEISGLOB2) of the model's information. On the other hand, principal components with maxima in the top 300 km and dominated by ridges, rifts and cratons explain 22% (SGLOBE-ranii) to 45% (SAVANIi) of the model's information.

Model	# splines or boxes	Single (#components)	Combined (12 components)
SGLOBE-ranii	21	98.2 (15)	92.0
SAVANIi	28	98.4 (7)	98.9
S20RTS	21	98.1 (13)	95.4
S40RTS	21	97.3 (12)	96.2
SEMUCB-WM1i	20	97.9 (9)	97.7
S362WMANI+Mi	16	98.5 (8)	97.9
SEISGLOB2	21	97.7 (13)	94.7

Table 2. Variance [%] obtained from the individual varimax analysis of each model. In parenthesis, the number of components capturing more than 1% of the variance is shown. The last column provides the variance captured by 12 components, for the combined analysis of the 7 models, as discussed in Section 6. The second column provides the number of splines or boxes originally used in the models.

6 Combined PCA

350 The horizontal patterns associated with each PC result from the projection of the tomography model on the varimax PCs, which differ from one model to the other. As suggested by Sengupta and Boyle (1998) in another context, it is interesting to compare the different models using a common PCA, which removes the inconsistencies between the representations. Thus, we apply a varimax PC analysis to the seven models stacked on the horizontal axis, i.e., to a $7 \times 9002 = 63014$ by 29 matrix, and refer to the results as a combined analysis in the remainder of this paper. Using the same 1% threshold limit as used before, 355 this analysis generates 12 varimax PCs, i.e., 12 vertical profiles (A-L) common to the seven models. Then, we compute the horizontal structures associated with each PC by projecting each of the 7 models on those vertical profiles (Figures 5 and D1 to D11).

Components A-D (at $\sim 50, 200, 300, 600$ km depths) are mostly confined in the upper mantle, while lower mantle structure is represented by components E-L ($\sim 800, 1000, 1200, 1400, 1700, 2000, 2300, 2800$ km depths), as shown in the vertical 360 profiles from the combined varimax analysis presented on the last column of Figure 3. As expected, the vertical profiles from the combined analysis are smoother than those from the individual model analysis for the more detailed models (SGLOBE-ranii, SEMUCB-WM1i, S40RTS and SEISGLOB2), and sharper for the smoothest tomography models (SAVANIi, S20RTS, SEMUCB-WM1i and S362WMANI+Mi). Note that the 1% criterion is applied globally, and not on the individual models, as was done in the previous section. The last column of Table 2 shows the total variance captured by the 12 components for each 365 model. It shows that the combined analysis is very efficient for the smooth SAVANIi model, capturing 98.9% of its variance, whereas the variance captured for the other models lies between 92.0% and 97.9%. SGLOBE-ranii model is only resolved at the 92.0% level, which is not surprising as it is more detailed than the other models (likely due to the use of less regularization), with the individual analysis requiring 15 PCs and allowing a finer localisation of the models' patterns (Figure C1).

Most of the patterns described in Table 3 and discussed in the previous section are also recovered by the combined analysis. 370 This common projection makes it easier to compare the components E (~ 800 km depth) to G (~ 1200 km depth) in the

Model	SGLOBE -ranii	SAVANIi	S20RTS	S40RTS	SEMUCB -WM1i	S362W MANI+Mi	SEIS GLOB2
# of components	15	7	13	12	9	8	13
Ridges, rifts and plateaus							
African Rift LV ¹	✓	✓	✓	✓	✓	✓	✓
Fast spreading Pacific zone	50-100	50-200	50	50	50	50	50
Tibetan plateau, Hangai dome LV ²	50	50	50	50	50	50	50
Craton HV zones							
African	100-200	50-200	50-300	50-300	50-200	50-300	50-200
Antarctic	100	50-200	50-300	50-300	50-200	50-300	50-200
Arabian	100	50-200	50-300	50-300	50-200	50-300	200
Australian	100	50-200	50-300	50-300	50-200	50-300	50-400
Baltic	100-200	50-200	50-300	50-300	50-200	50-300	50-200
Siberia	100	50-200	50-300	50-300	50-200	50-300	50-200
Indian	100	50-200	50-300	50-300	50-200	50-300	50
North American	100	50-200	50-200	50-300	50-200	50-300	50-400
South American	100	50-200	50-200	50-300	50-200	50-300	50-200
Back arc LV							
Japan	50-100	50-200	50	50	50-200	50	50
Philippines	50-100	50-200	50	50	50-200	50	50
Tonga-Kermadec	50-100	50-200	50	50	50-200	50	50
Subducted slabs							
Izu Bonin-Mariana HV	300	200	200	300 Poor	400	?	400
East Pacific HV	600-700	400-800	600-800	700	600	500	700
Tonga-Kermadec HV ³	100-1300	400-1200	600-1300	800-1500	400-1300	300-1800?	200-1000
North Pacific, Sunda HV ³	800-1100	400-1200	1000-1300	1000-1100	?	500-800?	400-1000
Others							
Pacific LV	300-600	200-400	200-300	300	200-600	300-500	200-700
Central Atlantic HV ⁴	-	400	300-600	300-700	400-600	500	400-700
Lower mantle structures							
LLSVPs	1400-2700	1900-2800	1500-2800	1500-2800	1300-2800	1800-2800	1900-2800
Perm LV	2300-2700	2800	2300-2800	2000-2800	2300-2800	2200-2800	2100

Table 3. Examples of key geophysical patterns recovered in the mantle by the varimax analysis (see Figures 4 and C1 to C6). HV = high velocity zone, LV = low velocity.

¹No clear interruption from the surface down to the CMB; ²Legendre et al. (2015); Chen et al. (2015); ³van der Meer et al. (2018); ⁴Ritsema et al. (2011).

Component F, var. capt=6.8%




Figure 5. PC F (maximum at 1000 km depth) of the combined analysis of the isotropic models. On the right, the principal components or vertical profiles, and on the left, the associated horizontal structures. The other components are shown in the supplementary material on Figures D1 to D11.

lower mantle. These components (Figures 5 and D5–D6) capture $\sim 20.1\%$ of the information in the models and display a similar pattern in SGLOBE-ranii, S20RTS, S40RTS and SAVANIi. On the other hand, SEMUCB-WM1i, S362WMANI+Mi and SEISGLOB2 show different features, such as e.g. fewer high-velocity zones in this depth range. Regarding principal component H (~ 1400 km depth, Figure D7), it describes 7.2% of the information. All the models show a similar large-scale pattern except for S362WMANI+Mi, which shows isolated low-velocity zones in the Pacific, especially in the south.

As shown in the previous section, all models display several independent components in the lower mantle, making between 36% (SAVANIi) and 69% (SEISGLOB2) of the total components, depending on the model. This highlights complexity in the lower mantle and supports recent studies suggesting that the region of the lower mantle above the lowermost D'' layer is more complex than previously thought. For example, slab stagnation and lateral deflection of mantle plumes have been proposed in the uppermost lower mantle (Fukao and Obayashi, 2013; French and Romanowicz, 2015). Moreover, intriguing observations of seismic discontinuities (Kawakatsu and Niu, 1994; Jenkins et al., 2017) and of scatterers (Kaneshima, 2016) have also been reported at these depths. Compositional layering (Ballmer et al., 2015), a viscosity increase (Marquardt and Miyagi, 2015;

Rudolph et al., 2015), and spin transitions that seem to occur in Fe-bearing mantle minerals (Lin et al., 2013) have been proposed in the lower mantle, which can potentially influence the region's elasticity and transport properties.

385 6.1 Anisotropic structure

In addition to isotropic shear-wave speed anomalies, four of the models considered in our study also include radial anisotropy perturbations, that is, speed differences between vertically and horizontally polarised shear waves: SGLOBE-rania, SEMUCB-WM1a, SAVANIA, and S362WMANI+Ma. The seismic imaging of anisotropy is more challenging than that of isotropic structure because the sensitivity of seismic data to anisotropy is weaker (e.g., Chang et al., 2014; Beghein and Trampert, 2004; 390 Romanowicz and Wenk, 2017). Moreover, it has been shown that if crustal effects are not properly modelled, this can lead to substantial errors in the estimated mantle anisotropy (Panning et al., 2010; Ferreira et al., 2010; Chang et al., 2016; Bozdağ and Trampert, 2008; Lekić et al., 2010). These difficulties are at least partly responsible for the strong differences between existing radial anisotropy models (e.g., Figure A1b). Varimax PC analysis is thus a natural candidate to analyse and compare the models since it enhances their robust information. Figures E3 and E1–E10 show the vertical profiles (varimax PCs) and 395 horizontal patterns from the combined varimax analysis on the anisotropic part of the four radially anisotropic models, for which 10 components capture more than 1% of variance. As expected, there is poorer agreement between the radial anisotropy structure in the models than between the isotropic structure discussed in the previous sections, though some common features can be identified.

The two LLSVPs appear on the deepest PC J of SGLOBE-rania and SEMUCB-WM1a (Figure E10), which captures 10% 400 of the models' information. The Pacific LLSVP appears barely in the last PC for SAVANIA and S362WMANI+Ma. However, as explained in section 3, e.g. Kustowski et al. (2008) and Chang et al. (2014, 2015) showed that the signature of LLSVPs in radial anisotropy models is an artefact due to leakage of isotropic structure into artificial anisotropic structure in the lowermost mantle.

For PC B (with a maximum at ~ 100 km depth, Figure E2), a positive zone appears beneath the Pacific and the Nazca 405 plates in SGLOBE-rania and SAVANIA, and to a lesser extent in SEMUCB-WM1a, while no clear pattern is evidenced in S362WMANI+Ma. The same holds true for mid-ocean ridges. A positive anomaly is observed on PC C under the Pacific plate for all models with a maximum around a depth of 200 km (Figure E3). A broad positive radial anisotropy anomaly beneath the Pacific at these upper mantle depths has been well documented in previous studies, and may be due to horizontal mantle flow in the region and/or thin layers of partial melt in the asthenosphere (e.g., Ekström and Dziewonski (1998); Gung et al. (2003)).

410 Components B (maximum depth 100 km, Figure E2), D (maximum depth 400 km, Figure E4) and especially C (Figure E3) evidence subduction patterns on SGLOBE-rania (Alaska, Izu-Bonin, Fiji-Tonga-Kermadec). This is also observed, but less clearly, for SAVANIA and S362WMANI+Ma on PC C. We also distinguish subduction signatures deeper in the mantle on PC F along Cascadia, Central and South America, Tonga, the western Pacific and the north of the Mediterranean Sea (Figure E6). An overall red negative anisotropy anomaly for the first PC A is common to SEMUCB-WM1a and SAVANIA (Figure E1). 415 SGLOBE-rani shows such red anomaly only under the oceans, which is probably due to the different way the crust is treated in

this model, with crustal thickness perturbations being jointly inverted for along with isotropic and anisotropic structure (Chang et al., 2015).

7 Conclusions

Global seismic tomography models typically involve thousands to tens of thousands of parameters, which can be cumbersome to handle and difficult to interpret. This is also true for model comparison, where we lack a common basis for comparing models built with different parameterisations. In this study we used a rotated version of the Principal Component Analysis to compress the information, to ease the geological interpretation and model comparison. The varimax PC analysis results in a separation of the information into different components associated with depth distributions, which are linked to a horizontal pattern obtained by orthogonal projection. We tested the analysis on seven global tomography models: S20RTS, S40RTS, SEISGLOB2, SEMUCB-WM1, SAVANI, SGLOBE-rani, and S362WMANI+M, where the latter four include laterally varying radial anisotropy. We analysed the models both individually as well as jointly.

We found that using the varimax method we reduce by 29% to 75% the number of independent depth components needed to describe more than 97% of the total information in the tomography models. We note that the scale of heterogeneity is not relevant for the varimax PCA method, which is only based on the vertical covariance. Considering the low amount of variance lost in the reconstruction (e.g., Figure 4) and the spectrum shown in the Supplementary Online Material, we capture most of the information, and we do not change the spectrum of the signal. Thus, the method is valid for any scale, as long as the signal is robust. In the varimax comparison, what is called noise is not the small-scale features, but rather the part of the models that is not covariant vertically. Hence, the varimax analysis simplifies the number of patterns that needs to be analyzed without any significant loss of information and by ensuring the orthogonality of the depth components it eased the detection and comparison of the relevant information. Overall, the large majority of depth components and horizontal maps obtained from the varimax analysis are different from the original parameterisations used for building the models. This is especially true above the 600 km discontinuity or in the whole mantle for the SAVANIi, S362WMANI+Mi and SEMUCB-WM1i models, where the spline functions over-sample the available information the information is recovered by fewer PCs than the spline functions. This implies that the PCs do not only reflect the model parameterisation and inform us about the independence between the slices reconstructed from the model. This also shows that the tomography models do not have a strong imprint of their model parameterisation. The varimax technique, being a data compression method, allows an easier view of the whole information present in the tomography models, and is complementary to clustering methods, such as the k-mean technique, which allow to evidence zones of comparable properties. Both can help users get a better understanding of the complex Earth's interior structure.

Being data-based, the varimax method is neutral with respect to the assumptions made in the model's construction. The combined multi-model varimax analysis allows the comparison of the various tomography models on a neutral set of modes, determined by the level of compression fixed by the user. Based on the vertical consistency between the various tomography models, it provides a set of data-based vertical distribution functions. Those functions represent the information present in the


PC-based reconstruction, and how the models relate to each other. It is fast and simple to implement, and, as we maximize the captured variance, the level of compression is lower for a given number of components/depths than would be required by other methods such as k-mean. As the truncation level is a free parameter, the user controls the amount of signal suppressed from the compression to an arbitrary level.

When applying the varimax analysis to isotropic tomography models, we found that the most important elements of the models contributing to most of the information are: (i) Large Low Shear Velocity Provinces (LLSVPs) in the lowermost mantle; (ii) subducted slabs and low velocity anomalies probably associated with mantle plumes in the upper and lower mantle; and, (iii) ridges and cratons in the uppermost upper mantle. The analysis highlights several independent components in the lower mantle that make between 36% and 69% of the total components depending on the model, which supports recent studies suggesting that the lower mantle is more complex than previously thought. The reasons of this complexity remain a very active field of research. On the other hand, we find limited agreement between the radial anisotropy structure of the models, with common features mainly in the asthenosphere and to some extent in the lower mantle beneath the Pacific and beneath subduction zones.

Choices such as data types and amounts, and the strength of regularization used in the construction of tomography models are probably key controls of the number of varimax components required by each model. Hence, the PCA-based model compression preserves the impact of the choices made in the construction of the tomography models, and facilitates their interpretation in terms of geophysical objects. Future work will expand this analysis for the interpretation and comparison of local and regional models, which tend to use more diverse underlying datasets than in global models, and have highly variable spatial resolutions. Moreover, we will also focus on comparisons with other sources of information, such as e.g. geodynamical models, gravity anomalies, magnetic anomalies or heat fluxes.

Code and data availability. A python function for computing PCA and varimax PCA is provided at <https://gitlab.univ-lr.fr/odevtron/varimax/>. With our configuration, i.e., 29 slices with 9000 points per profile, it analyses an individual model in about 0.15 CPU sec, whereas the combined analysis with 7 models takes about 1.3 CPU sec, using a 2.3 GHz 8 cores Intel Core i9.

Appendix A: Models






Figure A1. Depth slices of the isotropic and anisotropic models used in this study at the depths of 100, 200, 300, 400, 600, 800, 1000, 1400, 2000, and 2900 km. The isotropic (a) and anisotropic (b) models show perturbations in shear-wave speed and in $\xi = V_{SH}^2/V_{SV}^2$ with respect to PREM (Dziewonski and Anderson, 1981), respectively. The colour scale is normalised to vary from -max to +max, with the range of model amplitude variations shown at the left of each row. For simplicity, in the remainder of this paper we add "i" letters to the tomography models' names when referring to their isotropic part and we add "a" letters when referring to their radially anisotropic part; in the different figures "SGLOBE" will stand for SGLOBE-rani, and "SEMUCB" for SEMUCB-WM1

Appendix B: Comparison of the original and the reconstructed models

480 B1 Spatial domain




Figure B1. Examples of 6 depth slices in the original SEMUCB-WM1i model (top), in the model recovered without the principal components with [less](#)fewer than 1% of the variance, i.e., with 9 components (middle), and differences between the two (bottom).

B2 Spectral domain




Figure B2. PSDs of the original models.




Figure B3. PSDs of the reconstructed models, from the varimax modes. The number of modes is given in Table 2.

Appendix C: Varimax PCA of isotropic models




Figure C1. 15 Varimax components of the isotropic part of the SGLOBE-rani model. On the right, the principal components or vertical profiles, and on the left, the associated horizontal structures.




Figure C2.7 Varimax components of the isotropic part of the SAVANI model.




Figure C3. 13 varimax components of the S20RTS model.




Figure C4.12 Varimax components of the S40RTS model.




Figure C5. 8 Varimax components of the isotropic part of the S362WMANI+M model.




Figure C6.13 Varimax components of the isotropic part of the SEISGlob2 model.

Appendix D: Combined PCA of isotropic models




Figure D1. PC A (maximum at 50 km depth) of the combined analysis of the isotropic parts of the models. On the right, the principal components or vertical profiles, and on the left, the associated horizontal structures.




Figure D2. PC B (maximum at 200 km depth) of the combined analysis of the isotropic parts of the models.




Figure D3. PC C (maximum at 300 km depth) of the combined analysis of the isotropic parts of the models.




Figure D4. PC D (maximum at 600 km depth) of the combined analysis of the isotropic parts of the models.




Figure D5. PC E (maximum at 800 km depth) of the combined analysis of the isotropic parts of the models.




Figure D6. PC G (maximum at 1200 km depth) of the combined analysis of the isotropic parts of the models.




Figure D7. PC H (maximum at 1400 km depth) of the combined analysis of the isotropic parts of the models.




Figure D8. PCI (maximum at 1700 km depth) of the combined analysis of the isotropic parts of the models.




Figure D9. PC J (maximum at 2000 km depth) of the combined analysis of the isotropic parts of the models.




Figure D10. PC K (maximum at 2300 km depth) of the combined analysis of the isotropic parts of the models.




Figure D11. PC L (maximum at 2800 km depth) of the combined analysis of the isotropic parts of the models.

Appendix E: Combined PCA of anisotropic models




Figure E1. PC A (maximum at 50 km depth) of the combined analysis of the anisotropic parts of the models. On the right, the principal components or vertical profiles, and on the left, the associated horizontal structures.




Figure E2. PC B (maximum at 100 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E3. PC C (maximum at 200 km depth) of the combined analysis of the anisotropic models.




Figure E4. PC D (maximum at 400 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E5. PC E (maximum at 600 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E6. PC F (maximum at 800 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E7. PC G (maximum at 1300 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E8. PC H (maximum at 1900 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E9. PC I (maximum at 2300 km depth) of the combined analysis of the anisotropic parts of the models.




Figure E10. PC J (maximum at 2800 km depth) of the combined analysis of the anisotropic parts of the models.

485 *Author contributions.* OdV and AG performed the formal analysis, OdV and MVC developed and adapted the methodologies used, OdV made the visualization, all the authors contribute to investigation and validation of the results

Competing interests. The authors have no competing interests

Acknowledgements. A.M.G.F. thanks support from NERC Grant NE/N011791/1. We greatly thank Lapo Boschi, Barbara Romanowicz and Raj Moulik for providing us files with their models' parameterisations. OdV and AG work was financially supported by CNES as
490 methodological development/test for analyzing and interpreting the geodesy mission data, in particular for GRACE satellite gravity. The two reviewers of a previous version of this manuscript are gratefully acknowledged for their insightful comments.

References

- Auer, L., Boschi, L., Becker, T. W., Nissen-Meyer, T., and Giardini, D.: *Savani* : A variable resolution whole-mantle model of anisotropic shear velocity variations based on multiple data sets, *Journal of Geophysical Research: Solid Earth*, 119, 3006–3034, <https://doi.org/10.1002/2013JB010773>, 2014.
- Ballmer, M. D., Schmerr, N. C., Nakagawa, T., and Ritsema, J.: Compositional mantle layering revealed by slab stagnation at ~1000-km depth, *Science Advances*, 1, e1500815, <https://doi.org/10.1126/sciadv.1500815>, 2015.
- Becker, T. W. and Boschi, L.: A comparison of tomographic and geodynamic mantle models, *Geochemistry, Geophysics, Geosystems*, 3, n/a–n/a, <https://doi.org/10.1029/2001GC000168>, 2002.
- Begg, G., Griffin, W., Natapov, L., O'Reilly, S. Y., Grand, S., O'Neill, C., Hronsky, J., Djomani, Y. P., Swain, C., Deen, T., and Bowden, P.: The lithospheric architecture of Africa: Seismic tomography, mantle petrology, and tectonic evolution, *Geosphere*, 5, 23–50, <https://doi.org/10.1130/GES00179.1>, 2009.
- Beghein, C. and Trampert, J.: Probability density functions for radial anisotropy: implications for the upper 1200 km of the mantle, *Earth and Planetary Science Letters*, 217, 151–162, [https://doi.org/10.1016/S0012-821X\(03\)00575-2](https://doi.org/10.1016/S0012-821X(03)00575-2), 2004.
- Benoit, M. H., Nyblade, A. A., Owens, T. J., and Stuart, G.: Mantle transition zone structure and upper mantle S velocity variations beneath Ethiopia: Evidence for a broad, deep-seated thermal anomaly, *Geochemistry, Geophysics, Geosystems*, 7, n/a–n/a, <https://doi.org/10.1029/2006GC001398>, 2006.
- Bonatti, E.: Anomalous opening of the Equatorial Atlantic due to an equatorial mantle thermal minimum, *Earth and Planetary Science Letters*, 143, 147–160, [https://doi.org/10.1016/0012-821X\(96\)00125-2](https://doi.org/10.1016/0012-821X(96)00125-2), 1996.
- Bozdağ, E. and Trampert, J.: On crustal corrections in surface wave tomography, *Geophysical Journal International*, 172, 1066–1082, <https://doi.org/10.1111/j.1365-246X.2007.03690.x>, 2008.
- Browne, M. W.: An Overview of Analytic Rotation in Exploratory Factor Analysis, *Multivariate Behavioral Research*, 36, 111–150, https://doi.org/10.1207/S15327906MBR3601_05, 2001.
- Chang, S., Kendall, E., Davaille, A., and Ferreira, A. M. G.: The Evolution of Mantle Plumes in East Africa, *Journal of Geophysical Research: Solid Earth*, 125, <https://doi.org/10.1029/2020JB019929>, 2020.
- Chang, S.-J. and Van der Lee, S.: Mantle plumes and associated flow beneath Arabia and East Africa, *Earth and Planetary Science Letters*, 302, 448–454, <https://doi.org/10.1016/j.epsl.2010.12.050>, 2011.
- Chang, S.-J., Ferreira, A. M., Ritsema, J., van Heijst, H. J., and Woodhouse, J. H.: Global radially anisotropic mantle structure from multiple datasets: A review, current challenges, and outlook, *Tectonophysics*, 617, 1–19, <https://doi.org/10.1016/j.tecto.2014.01.033>, 2014.
- Chang, S.-J., Ferreira, A. M. G., Ritsema, J., van Heijst, H. J., and Woodhouse, J. H.: Joint inversion for global isotropic and radially anisotropic mantle structure including crustal thickness perturbations, *Journal of Geophysical Research: Solid Earth*, 120, 4278–4300, <https://doi.org/10.1002/2014JB011824>, 2015.
- Chang, S.-J., Ferreira, A. M. G., and Faccenda, M.: Upper- and mid-mantle interaction between the Samoan plume and the Tonga–Kermadec slabs, *Nature Communications*, 7, <https://doi.org/10.1038/ncomms10799>, 2016.
- Chen, M., Niu, F., Liu, Q., and Tromp, J.: Mantle-driven uplift of Hangai Dome: New seismic constraints from adjoint tomography, *Geophysical Research Letters*, 42, 6967–6974, <https://doi.org/10.1002/2015GL065018>, 2015.
- Cottaar, S. and Lekic, V.: Morphology of seismically slow lower-mantle structures, *Geophysical Journal International*, 207, 1122–1136, <https://doi.org/10.1093/gji/ggw324>, 2016.

- Durand, S., Debayle, E., Ricard, Y., Zanolli, C., and Lambotte, S.: Confirmation of a change in the global shear velocity pattern at around
530 1000 km depth, *Geophysical Journal International*, 211, 1628–1639, <https://doi.org/10.1093/gji/ggx405>, 2017.
- Durand, S., Abreu, R., and Thomas, C.: SeisTomoPy: Fast Visualization, Comparison, and Calculations in Global Tomographic Models, *Seismological Research Letters*, 89, 658–667, <https://doi.org/10.1785/0220170142>, 2018.
- Dziewonski, A. and Anderson, D. L.: Preliminary reference Earth model, *Physics of the Earth and Planetary Interiors*, 25, 297–356, [https://doi.org/10.1016/0031-9201\(81\)90046-7](https://doi.org/10.1016/0031-9201(81)90046-7), 1981.
- 535 Dziewonski, A. M., Lekic, V., and Romanowicz, B. A.: Mantle Anchor Structure: An argument for bottom up tectonics, *Earth and Planetary Science Letters*, 299, 69–79, <https://doi.org/10.1016/j.epsl.2010.08.013>, 2010.
- Ekström, G. and Dziewonski, A. M.: The unique anisotropy of the Pacific upper mantle, *Nature*, 394, 168–172, <https://doi.org/10.1038/28148>, 1998.
- Ferreira, A. M. G., Woodhouse, J. H., Visser, K., and Trampert, J.: On the robustness of global radially anisotropic surface wave tomography,
540 *Journal of Geophysical Research*, 115, <https://doi.org/10.1029/2009JB006716>, 2010.
- Ferreira, A. M. G., Faccenda, M., Sturgeon, W., Chang, S.-J., and Scharong, L.: Ubiquitous lower-mantle anisotropy beneath subduction zones, *Nature Geoscience*, 12, 301–306, <https://doi.org/10.1038/s41561-019-0325-7>, 2019.
- Flament, N., Williams, S., Müller, R. D., Gurnis, M., and Bower, D. J.: Origin and evolution of the deep thermochemical structure beneath Eurasia, *Nature Communications*, 8, <https://doi.org/10.1038/ncomms14164>, 2017.
- 545 French, S. W. and Romanowicz, B.: Broad plumes rooted at the base of the Earth’s mantle beneath major hotspots, *Nature*, 525, 95–99, <https://doi.org/10.1038/nature14876>, 2015.
- French, S. W. and Romanowicz, B. A.: Whole-mantle radially anisotropic shear velocity structure from spectral-element waveform tomography, *Geophysical Journal International*, 199, 1303–1327, <https://doi.org/10.1093/gji/ggu334>, 2014.
- Fukao, Y. and Obayashi, M.: Subducted slabs stagnant above, penetrating through, and trapped below the 660 km discontinuity, *Journal of*
550 *Geophysical Research: Solid Earth*, 118, 5920–5938, <https://doi.org/10.1002/2013JB010466>, 2013.
- Garnero, E. J. and McNamara, A. K.: Structure and Dynamics of Earth’s Lower Mantle, *Science*, 320, 626–628, <https://doi.org/10.1126/science.1148028>, 2008.
- Gung, Y., Panning, M., and Romanowicz, B.: Global anisotropy and the thickness of continents, *Nature*, 422, 707–711, <https://doi.org/10.1038/nature01559>, 2003.
- 555 Hansen, S. E. and Nyblade, A. A.: The deep seismic structure of the Ethiopia/Afar hotspot and the African superplume, *Geophysical Journal International*, 194, 118–124, <https://doi.org/10.1093/gji/ggt116>, 2013.
- Hansen, S. E., Nyblade, A. A., and Benoit, M. H.: Mantle structure beneath Africa and Arabia from adaptively parameterized P-wave tomography: Implications for the origin of Cenozoic Afro-Arabian tectonism, *Earth and Planetary Science Letters*, 319-320, 23–34, <https://doi.org/10.1016/j.epsl.2011.12.023>, 2012.
- 560 Heintz, M., Debayle, E., and Vauchez, A.: Upper mantle structure of the South American continent and neighboring oceans from surface wave tomography, *Tectonophysics*, 406, 115–139, <https://doi.org/10.1016/j.tecto.2005.05.006>, 2005.
- Horel, J. D.: A Rotated Principal Component Analysis of the Interannual Variability of the Northern Hemisphere 500 mb Height Field, *Monthly Weather Review*, 109, 2080–2092, [https://doi.org/10.1175/1520-0493\(1981\)109<2080:ARPCAO>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<2080:ARPCAO>2.0.CO;2), 1981.
- Hosseini, K., Matthews, K. J., Sigloch, K., Shephard, G. E., Domeier, M., and Tsekhmistrenko, M.: SubMachine: Web-Based Tools for
565 Exploring Seismic Tomography and Other Models of Earth’s Deep Interior, *Geochemistry, Geophysics, Geosystems*, 19, 1464–1483, <https://doi.org/10.1029/2018GC007431>, 2018.

- Hosseini, K., Sigloch, K., Tsekhmistrenko, M., Zaheri, A., Nissen-Meyer, T., and Igel, H.: Global mantle structure from multifrequency tomography using P, PP and P-diffracted waves, *Geophysical Journal International*, 220, 96–141, <https://doi.org/10.1093/gji/ggz394>, 2020.
- 570 Jenkins, J., Deuss, A., and Cottaar, S.: Converted phases from sharp 1000 km depth mid-mantle heterogeneity beneath Western Europe, *Earth and Planetary Science Letters*, 459, 196–207, <https://doi.org/10.1016/j.epsl.2016.11.031>, 2017.
- Jolliffe, I.: Principal Component Analysis, in: *Encyclopedia of Statistics in Behavioral Science*, edited by Everitt, B. S. and Howell, D. C., John Wiley & Sons, Ltd, Chichester, UK, <https://doi.org/10.1002/0470013192.bsa501>, 2005.
- Kaiser, H. F.: The varimax criterion for analytic rotation in factor analysis, *Psychometrika*, 23, 187–200, <https://doi.org/10.1007/BF02289233>,
575 1958.
- Kaneshima, S.: Seismic scatterers in the mid-lower mantle, *Physics of the Earth and Planetary Interiors*, 257, 105–114, <https://doi.org/10.1016/j.pepi.2016.05.004>, 2016.
- Kawakatsu, H. and Niu, F.: Seismic evidence for a 920-km discontinuity in the mantle, *Nature*, 371, 301–305, <https://doi.org/10.1038/371301a0>, 1994.
- 580 Kawamura, R.: A Rotated EOF Analysis of Global Sea Surface Temperature Variability with Interannual and Interdecadal Scales, *Journal of Physical Oceanography*, 24, 707–715, [https://doi.org/10.1175/1520-0485\(1994\)024<0707:AREAOG>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<0707:AREAOG>2.0.CO;2), 1994.
- Kustowski, B., Ekström, G., and Dziewoński, A. M.: Anisotropic shear-wave velocity structure of the Earth’s mantle: A global model, *Journal of Geophysical Research*, 113, <https://doi.org/10.1029/2007JB005169>, 2008.
- Lebedev, S. and van der Hilst, R. D.: Global upper-mantle tomography with the automated multimode inversion of surface and *S*-wave
585 forms, *Geophysical Journal International*, 173, 505–518, <https://doi.org/10.1111/j.1365-246X.2008.03721.x>, 2008.
- Legendre, C., Zhao, L., and Chen, Q.-F.: Upper-mantle shear-wave structure under East and Southeast Asia from Automated Multimode Inversion of waveforms, *Geophysical Journal International*, 203, 707–719, <https://doi.org/10.1093/gji/ggv322>, 2015.
- Lekic, V., Cottaar, S., Dziewoński, A., and Romanowicz, B.: Cluster analysis of global lower mantle tomography: A new class of structure and implications for chemical heterogeneity, *Earth and Planetary Science Letters*, 357–358, 68–77, <https://doi.org/10.1016/j.epsl.2012.09.014>,
590 2012.
- Lekić, V., Panning, M., and Romanowicz, B.: A simple method for improving crustal corrections in waveform tomography: Crustal corrections in waveform tomography, *Geophysical Journal International*, pp. no–no, <https://doi.org/10.1111/j.1365-246X.2010.04602.x>, 2010.
- Lin, J.-F., Speziale, S., Mao, Z., and Marquardt, H.: Effects of the electronic spin transitions of iron in lower-mantle minerals: Implications to deep-mantle geophysics and geochemistry, *Reviews of Geophysics*, 51, 244–275, <https://doi.org/10.1002/rog.20010>, 2013.
- 595 MacQueen, J.: Some methods for classification and analysis of multivariate observations, in: *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability*, vol. 1, pp. 281–297, U. California, Berkeley, 1967.
- Marquardt, H. and Miyagi, L.: Slab stagnation in the shallow lower mantle linked to an increase in mantle viscosity, *Nature Geoscience*, 8, 311–314, <https://doi.org/10.1038/ngeo2393>, 2015.
- McNamara, A. K.: A review of large low shear velocity provinces and ultra low velocity zones, *Tectonophysics*, 760, 199–220, <https://doi.org/10.1016/j.tecto.2018.04.015>, 2019.
600
- Mestas-Nuñez, A. M.: Orthogonality properties of rotated empirical modes, *International Journal of Climatology*, 20, 1509–1516, [https://doi.org/10.1002/1097-0088\(200010\)20:12<1509::AID-JOC553>3.0.CO;2-Q](https://doi.org/10.1002/1097-0088(200010)20:12<1509::AID-JOC553>3.0.CO;2-Q), 2000.
- Moulik, P. and Ekström, G.: An anisotropic shear velocity model of the Earth’s mantle using normal modes, body waves, surface waves and long-period waveforms, *Geophysical Journal International*, 199, 1713–1738, <https://doi.org/10.1093/gji/ggu356>, 2014.

- 605 Neuhaus, J. O. and Wrigley, C.: The quartimax method: an analytic approach to orthogonal simple structure, *British Journal of Statistical Psychology*, 7, 81–91, <https://doi.org/10.1111/j.2044-8317.1954.tb00147.x>, 1954.
- Panning, M. P., Lekić, V., and Romanowicz, B. A.: Importance of crustal corrections in the development of a new global model of radial anisotropy, *Journal of Geophysical Research*, 115, <https://doi.org/10.1029/2010JB007520>, 2010.
- Pavlis, G. L., Sigloch, K., Burdick, S., Fouch, M. J., and Vernon, F. L.: Unraveling the geometry of the Farallon plate: Synthesis of three-
610 dimensional imaging results from USArray, *Tectonophysics*, 532–535, 82–102, <https://doi.org/10.1016/j.tecto.2012.02.008>, 2012.
- Polet, J. and Anderson, D. L.: Depth extent of cratons as inferred from tomographic studies, *Geology*, 23, 205, [https://doi.org/10.1130/0091-7613\(1995\)023<0205:DEOCAI>2.3.CO;2](https://doi.org/10.1130/0091-7613(1995)023<0205:DEOCAI>2.3.CO;2), 1995.
- Rawlinson, N., Fichtner, A., Sambridge, M., and Young, M. K.: Seismic Tomography and the Assessment of Uncertainty, in: *Advances in Geophysics*, vol. 55, pp. 1–76, Elsevier, <https://doi.org/10.1016/bs.agph.2014.08.001>, 2014.
- 615 Richman, M. B.: Rotation of principal components, *Journal of Climatology*, 6, 293–335, <https://doi.org/10.1002/joc.3370060305>, 1986.
- Ritsema, J.: Complex Shear Wave Velocity Structure Imaged Beneath Africa and Iceland, *Science*, 286, 1925–1928, <https://doi.org/10.1126/science.286.5446.1925>, 1999.
- Ritsema, J. and Lekić, V.: Heterogeneity of Seismic Wave Velocity in Earth’s Mantle, *Annual Review of Earth and Planetary Sciences*, 48, 377–401, <https://doi.org/10.1146/annurev-earth-082119-065909>, 2020.
- 620 Ritsema, J., van Heijst, H. J., and Woodhouse, J. H.: Global transition zone tomography, *Journal of Geophysical Research: Solid Earth*, 109, <https://doi.org/10.1029/2003JB002610>, 2004.
- Ritsema, J., Deuss, A., van Heijst, H. J., and Woodhouse, J. H.: S40RTS: a degree-40 shear-velocity model for the mantle from new Rayleigh wave dispersion, teleseismic traveltime and normal-mode splitting function measurements: S40RTS, *Geophysical Journal International*, 184, 1223–1236, <https://doi.org/10.1111/j.1365-246X.2010.04884.x>, 2011.
- 625 Romanowicz, B. and Wenk, H.-R.: Anisotropy in the deep Earth, *Physics of the Earth and Planetary Interiors*, 269, 58–90, <https://doi.org/10.1016/j.pepi.2017.05.005>, 2017.
- Rudolph, M. L., Lekić, V., and Lithgow-Bertelloni, C.: Viscosity jump in Earth’s mid-mantle, *Science*, 350, 1349–1352, <https://doi.org/10.1126/science.aad1929>, 2015.
- Sengupta, S. and Boyle, J. S.: Using Common Principal Components for Comparing GCM Simulations, *Journal of Climate*, 11, 816–830, [https://doi.org/10.1175/1520-0442\(1998\)011<0816:UCPCFC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<0816:UCPCFC>2.0.CO;2), 1998.
- 630 Storch, H. v. and Zwiers, F. W.: *Statistical analysis in climate research*, Cambridge University Press, Cambridge ; New York, 1999.
- Tao, W., Qiu, X., Wu, R., and Zhou, K.: Role of Differences in Surface Diurnal–Nocturnal Thermodynamics over Complex Terrain in a Squall Line Process, *Journal of Meteorological Research*, 33, 1–17, <https://doi.org/10.1007/s13351-019-8052-y>, 2019.
- van der Meer, D. G., van Hinsbergen, D. J., and Spakman, W.: Atlas of the underworld: Slab remnants in the mantle, their sinking history, and a new outlook on lower mantle viscosity, *Tectonophysics*, 723, 309–448, <https://doi.org/10.1016/j.tecto.2017.10.004>, 2018.
- 635 Zechmann, E.: Make Icosahedron - File Exchange - MATLAB Central File Exchange, <https://www.mathworks.com/matlabcentral/fileexchange/19169-make-icosahedron>, 2019.