

# 1 Covid-19 lockdown effects on the seismic recordings of Central 2 America

3 Mario Arroyo-Solórzano<sup>1</sup>, Diego Castro-Rojas<sup>2</sup>, Frédéric Massin<sup>3</sup>, Lepolt Linkimer<sup>1</sup>, Ivonne  
4 Arroyo<sup>1</sup> & Robin Yani<sup>2</sup>

5  
6 1: Escuela Centroamericana de Geología y Red Sismológica Nacional, Universidad de Costa Rica, San José, Costa  
7 Rica.

8 2: Instituto Nacional de Sismología, Vulcanología, Meteorología e Hidrología, INSIVUMEH, Ciudad de  
9 Guatemala, Guatemala.

10 3: ETHZ-SED, Swiss Seismological Service, Zurich, Switzerland.

11 *Correspondence to:* Mario Arroyo-Solórzano ([mario.arroyosolorzano@ucr.ac.cr](mailto:mario.arroyosolorzano@ucr.ac.cr))

12  
13 **Abstract.** A noticeable decrease in seismic noise was registered worldwide during the lockdown measures  
14 implemented in 2020 to prevent the spread of Covid-19. In Central America, strong lockdown measures started  
15 during March of 2020. We have used seismic stations from Costa Rica, Guatemala, El Salvador, and Nicaragua  
16 to study the effects of these measures on seismic records by characterizing temporal variations in the high-  
17 frequency band (4-14 Hz) via spectral and amplitude analyses. In addition, we study the link between the reduction  
18 of seismic noise and the number of earthquake detections and felt reports in Costa Rica and Guatemala. We found  
19 that seismic stations near the capitals of Costa Rica, Guatemala, and El Salvador, presented a decrease in their  
20 typical seismic noise levels, from 200 to 140 nm, 100 to 80 nm, and 120 to 80 nm, respectively. Our results showed  
21 that the largest reduction of ~ 50% in seismic noise was observed at seismic stations near main airports, busy  
22 roads, and densely populated cities. In Nicaragua, the seismic noise levels remained constant (~ 40 nm) as no  
23 lockdown measures were applied. We suggest that the decrease in seismic noise levels may have increased  
24 earthquake detections and the number of felt reports of low magnitude earthquakes. However, the variations  
25 observed in several seismic parameters before and after the lockdown are not significant enough to easily link our  
26 observations or separate them to other contributing factors. Our results imply that the study of seismic noise levels  
27 can be useful to verify the compliance of lockdown measures and to explore their effects in earthquake detection  
28 and felt reports.

## 29 1. Introduction

30 The seismic noise recorded by seismometers includes microseisms and atmospheric, anthropogenic or cultural  
31 noise (Nimiya, 2020). The anthropogenic seismic noise in urban areas tends to be rowdier and more complex than  
32 elsewhere. This includes seismic signals generated by human activities such as transportation and industrial  
33 activities (Gross and Ritter, 2009; Diaz et al., 2017; Hong et al., 2020). It is difficult to identify precisely at what  
34 frequencies and how different human activities are represented in seismic records (McNamara and Buland, 2004;  
35 Green et al., 2017; Lecoq et al., 2020a). Indeed, the seismic noise includes various anthropogenic noises as a  
36 function of frequency, time, and distance in a range usually between 1-40 Hz (Kuzma et al., 2009; Riahi and  
37 Gerstoft, 2015; Diaz et al., 2017).

38 A reduction in the seismic noise worldwide has been observed coinciding with the lockdown measures to prevent  
39 the spread of Covid-19, whose outbreak was declared pandemic in March 2020 by the World Health Organization  
40 (Sohrabi et al., 2020). This effect has been first described for Shillong (India) by Somalia (2020), for Northern  
41 Italy by Poli et al. (2020), and by Lecocq et al. (2020a) at a global scale. Governments have tried to prevent or  
42 delay the spread of Covid-19 by forcing the social distancing through measures like limiting non-essential  
43 activities, closing schools and universities, restriction of the mobility of the citizens, and shutdown of workplaces  
44 (Piccinini, et al., 2020).

45 Central America has been severely affected by the of Covid-19 pandemic with an estimated death toll of 18,145  
46 by the end of 2020 (SICA, 2020). This small land bridge (1400 km long, 80–400 km wide) between the Americas  
47 is home to about 50 million inhabitants. It is located mostly in the Caribbean Plate and the Panama Microplate,  
48 surrounded by four major tectonic plates: the Cocos plate to the southwest, the Nazca plate to the south, and the  
49 North American and South American plates to the north and southeast, respectively. The boundary between the  
50 Cocos and Caribbean plates occurs at the Middle America Trench (MAT), where the Cocos Plate subducts  
51 underneath the Caribbean Plate and the Panama Microplate. The North Panama Deformed Belt (NPDB)  
52 constitutes the Caribbean Plate-Panama Microplate boundary, and the Polochic-Motagua Fault System (PMFS)  
53 marks the Caribbean-North American Plate boundary (Figure 1) (e.g. Adamek et al., 1988; Kellogg and Vega,  
54 1995; Trenkamp et al., 2002; Vargas and Mann, 2013). This complex and active tectonic setting in Central  
55 America generates high seismicity rates and volcanic activity. For instance, some of the deadliest earthquakes  
56 (Figure 1) were the 1910 M 6.4 Cartago earthquake in Costa Rica, the 1972 M 6.3 Managua earthquake in  
57 Nicaragua, the 1976 M 7.5 Guatemala earthquake, and the 1986 M 5.7 San Salvador earthquake in El Salvador,  
58 with ~600, ~20,000, ~23,000, and ~6,000 fatalities, respectively (Espinosa, 1976; Mann et al., 1990; Harlow et  
59 al., 1993; Alonso-Henar et al., 2013).

60 Seismometers in urban settings optimize the spatial coverage of seismic networks at these areas, and warn of local  
61 geological hazards, for example the amplification of seismic waves (Ashenden et al., 2011). Some of the main  
62 institutions in charge of the permanent monitoring of seismicity in Central America are: the National Institute of  
63 Seismology, Volcanology, Meteorology and Hydrology (INSIVUMEH) in Guatemala, the National Service for  
64 Territorial Studies (SNET in El Salvador), the Nicaraguan Institute for Territorial Studies (INETER) in Nicaragua,  
65 and in Costa Rica the National Seismological Network of the University of Costa Rica (RSN-UCR) and the  
66 Volcanological and Seismological Observatory of the National University of Costa Rica (OVSICORI-UNA).  
67 Belize and Honduras lack an official seismic service and Panama has several local networks, such as the Chiriquí  
68 and the University of Panama (UPA).

69 Because high levels of the root mean square (RMS) of the high-frequency seismic anthropogenic noise  
70 displacement (HFSAND-RMS) hampers the ability to detect signals from earthquakes and volcanic eruptions, its  
71 analysis and delimitation is of the utmost importance (Lecoq et al., 2020a). The objective of this work is to present  
72 the first study of HFSAND-RMS during Covid-19 in Central America. We have used seismic stations in Central  
73 America (Figure 1) to evaluate the effects of lockdown measures in the seismic record near urban centers of four  
74 countries in the region: Costa Rica, Guatemala, El Salvador, and Nicaragua. In addition, specific sites near  
75 populated centers and airports in Costa Rica and Guatemala are also analyzed. Finally, we explore the possible

76 impact of the low noise levels on the capability of detecting earthquakes and on the number of felt reports during  
77 the lockdown.

78 **2. Data and methods**

79 **2.1. Seismic Stations and Data**

80 We consider data from vertical-component seismometers from 10 seismic stations located in Costa Rica,  
81 Guatemala, El Salvador, and Nicaragua. Four of them are operated by the RSN (Costa Rica), four more by the  
82 INSIVUMEH (Guatemala), one belongs to the MARN (El Salvador), and one to the INETER (Nicaragua) (Figures  
83 1 and A1). Most of these stations have been selected within or near the capitals of each country, to obtain an  
84 overview of the changes in the seismic records induced by the lockdown measures. In Costa Rica and Guatemala,  
85 where we had more access to the records, we also selected three stations close-by other populated centers or  
86 airports.

87

88 The selected stations include both broadband seismometers (BB) and short-period geophones (SP). For Costa  
89 Rica, we use the TC.SJS1 station (BB, Guralp CMG-6TD) located at the main campus of University of Costa Rica  
90 in San Jose. This station is 3 meters below the ground level, and the sensor is installed in a concrete pillar. We  
91 also inspected the stations TC.BELE, TC.ERIA, and TC.ZEDO (SP, Sixaola instruments manufactured by  
92 Raspberry Shake), located in urban areas at fire station buildings at ground level, close to Juan Santamaria  
93 International airport, Liberia city in Northwestern Costa Rica, and in San Isidro city (Perez Zeledon) in  
94 Southeastern Costa Rica, respectively. For Guatemala, we used the GI.GCG4 station (SP, OSOP Sixaola) located  
95 in an urban area, close to the Aurora International airport. This station is 3 meters below the ground level, and the  
96 sensor is also installed in a concrete pillar. Furthermore, we analyzed the stations GI.HUEH, GI.RETA, and  
97 GI.CHIE (BB-Guralp CMG-3ESP), located in urban areas, close to local airports and Huehuetenango, Retalhuleu,  
98 and Esquipulas downtowns, respectively. These stations are at ground level inside a dedicated vault with a  
99 concrete pillar.

100

101 For El Salvador and Nicaragua, we use the stations SV.CEDA (BB, Nanometrics Trillium-120p), located close to  
102 the Panamerican highway at the city of La Libertad and NU.MGAN (BB, Streckeisen STS-2), located at INETER  
103 in Managua downtown. The continuous seismic data for Costa Rica and Guatemala were obtained directly from  
104 the seismological networks of each country (INSIVUMEH, 2013; RSN, 2017) and from IRIS for El Salvador and  
105 Nicaragua, via FDSN (International Federation of Digital Seismograph Network) web services (INETER, 1975;  
106 SNET, 2004). We analyzed data over a time span covering one year from November 1, 2019 to October 31, 2020.

107 **2.2. Seismic noise analyses**

108 A spectral and amplitude analysis was carried out to characterize temporal variations in high-frequency (4-14 Hz)  
109 seismic signals dominated by the anthropogenic noise via RMS. We computed the high-frequency seismic noise  
110 amplitude prior to and during the pandemic lockdown measures (before March 16, 2020). The results have been  
111 compared to the lockdown measures implemented by the governments for each country, as documented in Table  
112 A1 (SICA, 2020), and their compliance by the population.

113  
114 The high-frequency seismic noise amplitudes have been computed following the method described by Lecoq et  
115 al. (2020a) and using the code provided by Lecoq et al. (2020b). In this technique, the method of Welch (1967) is  
116 applied, in which a power spectral density (PSD) is calculated for each 30-minute time-windows with a 50 percent  
117 overlap, converted into equivalent displacement, and combined into a single RMS value per time-window  
118 (Blackman and Tukey, 1958). This reduces the numerical noise in the power spectra at the expense of reducing  
119 the frequency resolution due to frequency binning, but this effect is minimized with a conservative smoothing  
120 parameterization (Lecoq et al., 2020a). Moreover, to highlight the general temporal pattern of the amplitude of  
121 the seismic noise, the displacement RMS time series with four samples per hour were averaged each day between  
122 6h and 16h, according to the local time (UTC-6). Additionally, for some stations in Costa Rica and Guatemala, the  
123 median displacement RMS of each hour was computed, for each day, comparing the result before and during the  
124 lockdown. Finally, an analysis of the percentage of change in the HFSAND-RMS was performed to obtain a  
125 median amplitude value for the whole period during the major restrictive measures, from March 16 to May 15,  
126 2020, and to compare it with the median of the period before lockdown measures from January 15 to March 15,  
127 2020.

128 **2.3. Earthquake detection capability and felt reports**

129 For Costa Rica and Guatemala, the effect of the reduction of HFSAND-RMS on earthquake detection was  
130 explored during the lockdown period, using the seismic catalogs of the RSN (Costa Rica) and the INSIVUMEH  
131 (Guatemala). In both observatories, earthquakes are located automatically by SeisComP3 (Gempa, 2019) and  
132 manually using the software SeisAn (Havskov et al. 2020). With this objective, a period before lockdown from  
133 November 1, 2019 to March 15, 2020 (4.5 months) was taken as a reference to compare with a proportional period  
134 during the confinement measures, from March 16 as of July 31, 2020 (4.5 months).

135  
136 Using curves of earthquake magnitude-frequency distribution, we inspected earthquakes with moment magnitude  
137 ( $M_w$ )  $\leq 4.0$  before and during lockdown, because small events reflect better the variations in the detection  
138 capability. Furthermore, we quantified the average number of seismic phases per earthquake as function of the  
139 magnitude, obtaining linear regressions for the two data sets. In addition, to explore the increase in the number of  
140 earthquakes detected as a function of magnitude, we calculated the Gutenberg-Richter relationship (Gutenberg  
141 and Richter, 1944) before and during the lockdown. To calculate these seismic parameters and their uncertainties,  
142 we used the classical maximum likelihood technique of Aki (1965) modified by Weichert (1980). To run this  
143 methodology, we used the OpenQuake software (GEM, 2020). The magnitude of completeness ( $M_c$ ) was  
144 estimated by the MAXC method, which corresponds to the maximum point in the non-cumulative graph of the  
145 Gutenberg-Richter relationship (e.g. Wiemer and Wyss, 2000; Woessner and Wiemer, 2005).

146  
147 We have also investigated the effect of lockdown measures on the number of earthquakes felt by the population.  
148 For both, Costa Rica and Guatemala, we counted the number of felt earthquakes in different magnitude intervals.  
149 In Costa Rica and Guatemala, the population reports via smartphones and social networks. Further, to analyze if  
150 there is a correlation between the decrease in HFSAND-RMS and the increase in low-magnitude felt events, and

151 between the low-magnitude felt events and the hour of the day, we explored the spatial context of these events  
152 with a map of felt seismicity and its occurrence hour, before and during lockdown.

153  
154 Additionally, the RSN maintains an interactive application for smartphones called “RSN”, which includes the  
155 module “*¿Lo Sentiste?*” (Linkimer and Arroyo, 2020), also available at the RSN website. This app was developed  
156 by the RSN based on the questionnaire “*Did you feel it?*” of the United States Geological Service (USGS)  
157 (Atkinson and Wald, 2007; Wald et al., 2011), which was translated to Spanish and simplified and adapted to  
158 Costa Rica. The users access the app and answer 12 simple questions (Table A2) and obtain a quick estimation of  
159 the intensity determined by the community decimal intensity (CDI), which is an aggregate of the average sums of  
160 the indexes associated with the questions (Dengler and Dewey, 1998). All the reports are shown in an emoticon  
161 map that updates continuously and can be accessed in real time in the app or in the RSN website. Finally, after  
162 enough ( $> \sim 300$ ) reports and outliers have been manually removed, an average intensity map is generated  
163 (Linkimer and Arroyo, 2020). We use this tool to complement the felt earthquakes analysis in Costa Rica,  
164 collecting the number of felt earthquakes reported through this app, before and during lockdown, including only  
165 the events with at least three reports and with  $Mw < 5.0$ . These events were also averaged by magnitude intervals.

166 **3. Results and discussion**

167 **3.1. Seismic noise and lockdown measures**

168 The stronger lockdown measures in Central America started on March 16. In Costa Rica, some of the main  
169 restrictions implemented by the governments were the closure of borders, schools, non-essential stores, and  
170 beaches, as well as massive public events prohibition (concerts, soccer games, etc.) and home-office  
171 implementation. Although in Costa Rica no curfew was imposed on citizens, a strict vehicle mobility restriction  
172 has been maintained during the whole pandemic. For Guatemala and El Salvador, the lockdown measures were  
173 very similar to those implemented in Costa Rica, but in some cases included restrictions on citizen mobility and  
174 curfews. These measures have been softened or hardened as the pandemic evolved in each of these countries  
175 (Table A1). Very few lockdown measures were taken in Nicaragua to prevent the spread of Covid-19 and there  
176 were no specific measures applied to restrict social mobility.

177  
178 An important decrease in the HFSAND-RMS is shown in seismic stations located near the capitals of Costa Rica  
179 (University of Costa Rica campus, San José), Guatemala (close to the Aurora International airport, Guatemala  
180 City), and El Salvador (close to the Panamerican highway) (Figure A1). This has been observed during the  
181 lockdown measures (Table A1 and Figures 2 and 3), except for Nicaragua (urban area, at INETER, Managua).

182  
183 Figure 2 shows the time of day on the vertical axis, the period analyzed on the horizontal axis, and the high-  
184 frequency displacement RMS in colors, blue for the lowest level and yellow for the maximum. The graph clearly  
185 shows the stillness of the night (blue colors between 22h and 5h), the relative quiet of weekends (vertical blue  
186 bars that alternate periodically), and the hustle and bustle from day to day (in yellow colors). Figure 3 shows the  
187 displacement on the vertical axis and time on the horizontal axis. The orange line represents the median recorded  
188 displacement, which usually has its maximum during the hours of the day, when there is more seismic noise, and

189 its minimum during the nights. In addition, Figure 3 also show that the noise level is lower during the weekends.  
190 In these figures, we marked the beginning of the social distancing measures on March 16 (red line), the period of  
191 the end of the year holidays 2019 (1), Easter 2020 (2), and a brief period in July 2020 when there was a strict  
192 return of lockdown measures in Costa Rica (3) (Table A1).

193  
194 In both types of graphs (Figures 2 and 3), for the stations in Costa Rica, Guatemala, and El Salvador, the effect of  
195 the social distancing measures can be clearly seen beginning on March 16, as a notable drop in seismic noise  
196 (displacement). At the station in Managua, Nicaragua, where no important measures were adopted to limit urban  
197 mobility and economic activities, there is no change in the seismic record (Figures 2d and 3d). In Costa Rica,  
198 Guatemala, and El Salvador, the seismological stations show that the measures of social distancing produced a  
199 decrease in the seismic noise levels similar to those observed in the 2019 New Year holidays.

200  
201 For the TC.SJS1 station in San Jose, Costa Rica (Figure 2a and 3a), the displacement during a typical working  
202 day before the lockdown used to be up to 200 nm, while during the social distancing measures, these values  
203 decreased to 140 nm on average. In the case of the GI.GCG4 station in Guatemala City (Figure 2b and 3b), the  
204 usual displacement before the lockdown was 100 nm on average, and during social distancing measures, these  
205 values decreased to 80 nm on average. At the SV.CEDA station, near San Salvador (Figure 2c and 3c), the usual  
206 displacement before the lockdown was on average about 120 nm, while these values decreased to 80 nm on  
207 average during the lockdown.

208  
209 The lower displacement values, tended to return to their usual averages values as the restrictive measures were  
210 progressively eliminated. Even so, the values have not yet returned to the usual pre-pandemic levels by November  
211 2020, when the average values were ~ 160 nm for San Jose, ~ 85 nm for Guatemala City, and ~ 110 nm near San  
212 Salvador. This shows that some of the social distancing measures were still in place by November 2020 (Table  
213 A1) or other factors are affecting the station environment (permanent or long-term activity loss, e.g., company  
214 shutdown) (Figures 2 and 3). The place location where the level is closer to the usual averages is at La Libertad  
215 near San Salvador. The most drastic effect of the limitation in the mobility of the inhabitants was observed during  
216 Easter in April of 2020. The values observed were: ~60 nm in San José and ~50 nm in Guatemala City and near  
217 San Salvador.

218  
219 In the case of the NU.MGAN station in Managua, Nicaragua (Figures 2d and 3d), the registered displacement  
220 values have remained constant before and during the pandemic, without any variation with respect to the usual  
221 displacement records of this station (40 nm). It is a low-noise site compared to the other capitals because it has  
222 particular conditions that favor its isolation from the environmental noise.

223  
224  
225 Results for Costa Rican stations TC.BELE, TC.ERIA, and TC.ZEDO are presented in Figure 4a and Guatemalan  
226 stations GI.HUEH, GI.RETA, and GI.CHIE in Figure 4b. All these stations also show a decrease in displacement  
227 RMS since the application of lockdown measures. However, the reduction, as well as the pattern of the  
228 displacement RMS time series is very specific to each station. For Costa Rica (Figure 4a), the largest percentage

229 difference in the station record is found in the TC.BELE station, located near (~3 km) the country's main airport  
230 and in the Great Metropolitan Area (GAM) (Figure 1). This station varied from ~40 nm before lockdown to ~20  
231 nm during the lockdown measures. For Guatemala (Figure 4b), a similar behavior was observed in the GI.RETA  
232 station, towards the western part of this country, which is one of the most touristic regions. This station varied  
233 from ~100 nm before lockdown to ~50 nm during the lockdown measures.

234

235 In addition, the daily variation of the average seismic noise per weekday, before and during the pandemic, was  
236 inspected for the station TC.SJS1 in Costa Rica (Figure 5a) and for station GI.RETA in Guatemala (Figure 5b).  
237 In both stations, the typical pattern of HFSAND-RMS is also shown, with minimums during weekends and nights,  
238 and maximums during the week and day (Figure 5). This is also highlighted during the lockdown, due to the  
239 measures adopted by each country (Table A1). Costa Rica, despite not imposing a curfew, established measures  
240 of vehicular restriction from 19h to 5h at its most strict stage. This is highlighted in the station TC.SJS1, showing  
241 a great decrease in HFSAND-RMS during these hours (Figure 5a). For Guatemala, the station GI.RETA clearly  
242 shows the effect of the curfew in its most restrictive stage, imposed from 17h to 5h (Figure 5b).

243

244 The percentage of HFSAND-RMS decrease was determined for the 10 stations analyzed here, in the same  
245 frequency band (4-14 Hz). Figure 6a shows the change obtained for the stations in Guatemala and El Salvador  
246 and Figure 6b shows the percentage obtained for the stations in Nicaragua and Costa Rica. Four categories of  
247 seismic noise reduction were identified: very high (36-49%), high (26-35%), intermediate (16-25%), and low (6-  
248 15%). The most outstanding seismic noise reduction (very high) due to the lockdown measures were obtained at  
249 the GI.RETA and TC.BELE stations. These values are explained by the site characteristics described above, strong  
250 sources of noise at stations close to major cities, highways, and high-traffic airports. The stations that show a high  
251 decrease were GI.GCG4, SV.CEDA, and TC.SJS1. These changes are closely related to the location of these sites  
252 at the populated capital cities of San José and Guatemala City, and to the Panamerican highway near San Salvador  
253 (Figure 1). Intermediate decrease values were identified in the stations GI.HUEH, GI.CHIE, and TC.ERIA. In this  
254 case, these values are associated with cities with less population density (Figure 1) but significant activity and  
255 proximity to touristic airports, such as Liberia city, in northwestern Costa Rica. Finally, low changes were  
256 identified in the NU.MGAN and TC.ZEDO stations. In Managua, this is due to the lack of social distancing  
257 measures. In the case of Perez Zeledon (Costa Rica), it could be related to lower population density, a station site  
258 building more isolated from population and environment dynamics, or lack of lockdown measures compliance.

### 259 **3.2. Possible effects of the lockdown in earthquake detection and felt reports**

260 There were no significant earthquakes in Costa Rica and Guatemala during the time this study was conducted.  
261 Before lockdown in Costa Rica, the biggest earthquakes had a Mw of 5.6 on 21-01-2020 and in Guatemala a Mw  
262 of 6.2 on 19-11-2019. During lockdown, the largest events were a Mw 5.5 earthquake on 15-04-2020 in Costa  
263 Rica and a Mw 5.7 on 26-03-2020 in Guatemala. All these events were offshore earthquakes related to the  
264 interplate seismogenic zone. We conclude that the seismic rates during the time periods considered in this work  
265 were not affected by any specific large event. In Table 1 we summarize the main observations when comparing  
266 the earthquake detections and felt reports from the time before and during the lockdown for Costa Rica and  
267 Guatemala.

268

269 **Costa Rica**

270 Since 2018, the RSN network consists of around 160 sites, qualifying as a robust network with a high capacity for  
271 detecting low-magnitude earthquakes (Linkimer et al., 2018). Figure 7 shows the comparison of the number of  
272 seismic events recorded by the RSN and the reports of earthquakes felt before and during the lockdown measures.

273

274 Figure 7a shows the number of earthquakes on the vertical axis, and the range of magnitude on the horizontal axis,  
275 grouped in intervals of 0.5, from Mw 2.0 to 5.5. The graph shows that there is no increase in the capacity to detect  
276 earthquakes during the lockdown measures, and it even seems that more seismicity was detected in the period  
277 before lockdown, possibly due to higher seismic productivity in that period. The Mc shows that the impact of the  
278 lockdown measures in the detected earthquakes is not significant as Mc varies from 2.9 before to 3.0 during  
279 lockdown (Figure 8a and 8b). Also, the a- and b-values obtained were very consistent and similar to recent studies  
280 for Costa Rica (i.e., Arroyo and Linkimer, 2021). The a-value decreased slightly during the lockdown from 3.62  
281 to 3.56 (Figure 8a and 8b), showing a general decrease in the seismic rate. On the other hand, the increase in the  
282 b-value from 0.76 to 0.77 (Figure 8a and 8b) could be explained as an increment in the number of low-magnitude  
283 earthquakes compared to the number of higher magnitudes. This b-value could support the idea that more low-  
284 magnitude earthquakes were detected during the lockdown period (Table 1). However, as it can be seen the  
285 variations are very small to allow strong conclusion and this needs extra seismic noise analyses or specific site  
286 noise characterization. The observed variations could also be explained due to other contributing factors or even  
287 as random coincidence.

288

289 In Figure 7b we show the average number of P-wave phases per earthquake for earthquakes with  $M \leq 4.0$ .  
290 Although the difference in the number of P-wave arrivals before and during the pandemic is not too much (between  
291 1 and 5 picks for magnitudes between 1.8 and 4.0), the values are consistently higher (on average ~20%) during  
292 the pandemic, especially for the lower magnitudes ( $M \leq 3.5$ ). This additional observation also favors the idea that  
293 the decrease in HFSAND-RMS during lockdown may have had an effect on the earthquake detection capability  
294 of the RSN (Table 1).

295

296 Figure 7c shows the number of felt earthquakes reported through social networks or telephone calls with respect  
297 to a range of Mw from 2.0 to 5.5. This graph shows that there were a greater number of earthquakes with  $Mw >$   
298 3.5 reported as felt before the lockdown measures, but during the confinement, a greater number of reports for  
299 low magnitude earthquakes ( $M < 3.5$ ) were collected (Table 1). In addition, based on reports through the RSN  
300 application "*¿Lo Sentiste?*", we present in Figure 7d the average number of reports for  $M < 5.0$  by magnitude  
301 intervals. The trend lines in this graph show how the application "*¿Lo Sentiste?*" collected, on average, more felt  
302 reports for magnitudes  $M < 5.0$  during the lockdown measures. These figures suggest a greater sensitivity of the  
303 population to low-magnitude earthquakes, possibly because the longer stays in their homes and the  
304 implementation of home office and restrictions of mobility, allowed them to perceive more events and also to  
305 report them to seismic agencies (Table 1).

306

307 Figure 9 shows the geographical distribution of felt events reported by the RSN (Figure 7c) before (1 Nov 2019-  
308 15 Mar 2020) and during (16 Mar-31 Jul 2020) lockdown measures in Costa Rica. Even though the number of  
309 earthquake reports was higher before (99) than during (74) the lockdown measures, the percentage of low  
310 magnitude ( $M < 3.5$ ) felt earthquakes was clearly higher (46%) during the lockdown than before it (only 27%).  
311 These numbers suggest that the quiescence of the environment could be an important contributing factor to more  
312 small earthquakes being reported (Table 1). Another aspect is that there were a greater number of higher-  
313 magnitude earthquakes ( $M > 4.5$ ) before the lockdown (20) than during it (11), and these events were located  
314 onshore and closer to population centers (Figure 9).

315  
316 We also checked the correlation between felt events, before and during lockdown, and the decrease in the  
317 HFSAND-RMS of the four seismic stations analyzed in Costa Rica. There seems to exist a correlation in three of  
318 them, two located in the metropolitan area of Central Costa Rica (TC.SJS1 and TC.BELE) and the other in an  
319 urban area in Southeastern Costa Rica (TC.ZEDO). There were 25 felt earthquakes for Central Costa Rica before  
320 the lockdown and 17 of them had a low magnitude ( $M < 3.5$ ). These numbers are lower than those during  
321 lockdown, when there were 36 felt earthquakes, including 28 of low magnitude (Figure 9). For Southeastern Costa  
322 Rica, near the Perez Zeledon urban area (TC.ZEDO), just five felt earthquakes were reported before the lockdown,  
323 all of them with  $M > 3.5$ , but during lockdown seven events were reported as felt, four of them of low magnitude  
324 (Figure 9 and Table 1). Then, again, these observations suggest an increase in felt reports from small earthquakes  
325 in these regions as well. These felt events, as expected, were more perceived during the night hours, from 18h to  
326 6h (Figure 9).

327  
328 **Guatemala**  
329 The INSIVUMEH seismic network is still under development. This network consists of 24 seismic stations, most  
330 of them Guralp broadband sensors, some installed inside the main military detachments or national airports, while  
331 others are installed in the main tourist cities of the country. For this reason, these stations can reach high levels of  
332 seismic noise, which is why detecting low-magnitude earthquakes under pre-pandemic conditions were  
333 challenging.

334  
335 In Figure 10a, we present a comparison between the statistics before and during the lockdown measures. Similar  
336 than Costa Rica, for Guatemala this graph shows that a higher number of events was recorded during the lockdown  
337 measures (Table 1). However, the analysis of other seismic parameters are not conclusive:  $M_c$  varies only slightly  
338 from 3.7 to 3.8 from before to during the lockdown, which does not imply a significant increase in earthquake  
339 detection (Figure 8c and 8d). But the a- and b-values seem to suggest an increment in low magnitude earthquake,  
340 changing from 4.43 to 4.63 for the a-value and from 0.87 to 0.91 for the b-value (Figure 8c and 8d), for the periods  
341 of before and during the lockdown.

342  
343 In Guatemala, the average number of seismic phases per event of the same magnitude increases clearly for  
344 magnitudes lower than 3.0 during the lockdown (on average ~40%) (Figure 10b). Then, the possible effect of the  
345 lower levels in seismic noise for detecting more low-magnitude earthquakes could be stronger in Guatemala than  
346 in Costa Rica (Table 1). This may be related to the much lower seismic station density of Guatemala. Furthermore,

347 it should be noted that the lockdown measures implemented by the Guatemalan government were more drastic  
348 than in Costa Rica. Moreover, the number of felt events reported by the population shows again the interesting  
349 trend of a general increase during the lockdown period, but in the case of Guatemala this is visible for  $M \geq 3.5$   
350 (Figure 10c and Table 1).

351  
352 Figure 11 shows the distribution of the felt earthquakes reported to INSIVUMEH (Figure 10c) before and during  
353 the lockdown. Before the lockdown, a total of 34 seismic felt events were reported, while during lockdown that  
354 number increased to 47. The earthquake magnitudes were higher before lockdown (Figure 11a) with 21  
355 earthquakes above  $M 4.5$  (62% of the total felt events in this period) than during lockdown (Figure 11b), with 19  
356 earthquakes above  $M 4.5$  (40% of the total felt events in this period). Hence, during the lockdown there were more  
357 felt earthquakes of lower magnitude ( $M < 4.5$ ), including three earthquakes below  $M 3.5$ , in contrast to just one  
358 before the lockdown. Once more, this trend suggests a correlation between the quiescence of the environment and  
359 an increment in low magnitude felt events during the lockdown (Table 1). These felt events, as expected, were  
360 more perceived during the night hours, from 18h to 6h (Figure 11).

361  
362 Finally, from the spatial distribution of felt earthquakes during lockdown, we observed that, unlike the time before  
363 the pandemic, there are more events originated onshore and mainly close to the populated places, such as  
364 Guatemala City and Huehuetenango. These are epicentral locations near two of the seismic stations for which the  
365 decrease in the HFSAND-RMS was observed: GI.GCG4 and GI.HUEH, respectively. In a seismic network under  
366 development like the INSIVUMEH, with fewer stations, a high percentage of the low-magnitude seismicity is  
367 likely not detected due to ambient noise, but the observations provided above seems to support the idea that the  
368 lockdown measures cause an improvement in the detection capacity of this network.

#### 369 **4. Conclusions**

370 An important decrease in the high-frequency seismic noise was detected at stations of three Central American  
371 countries during the lockdown measures adopted to prevent the spread of Covid-19. In Costa Rica, Guatemala,  
372 and El Salvador, the measures of social distancing produced seismic noise levels comparable to those observed  
373 during the New Year holidays from previous years. The displacement observed decreased from 200 to 140 nm in  
374 San Jose, from 100 to 80 nm in Guatemala City, and from 120 to 80 nm in the city of La Libertad near San  
375 Salvador. In Nicaragua, with very few measures in place, there were no effects on the seismic noise levels of the  
376 station analyzed, which also happens to be a quiet site (40 nm).

377  
378 The decrease in the high-frequency seismic anthropogenic noise displacement is strongly dependent on the  
379 location of the station and on the lockdown measures. Four categories of seismic noise reduction were identified  
380 (very high, high, intermediate, and low), with significant values of ~50% decrease were observed in stations near  
381 airports, busy roads, and densely populated cities. The greatest impact in the noise levels started on March 16 and  
382 was specially related to the closure of educational centers and non-essential stores, the curfew from 17h to 5h in  
383 Guatemala, and the restriction on vehicular mobility from 19h to 5h in Costa Rica. The most drastic effect of the  
384 limitation in the mobility of the inhabitants was observed during Easter in April of 2020, when the values observed  
385 were as low as ~60 nm in San José and ~50 nm in Guatemala City and near San Salvador.

386

387 We suggest that in Costa Rica, the lower levels in seismic noise allowed the detection of a higher number of  
388 seismic phases (~20% more per magnitude) and therefore, the location of more small earthquakes than usual, as  
389 well as more low magnitude earthquakes ( $M < 3.5$ ) reported as felt. Similarly, in Guatemala, where the seismic  
390 network is still under development, we suggest a similar effect with an increase of ~40% in the amount of the  
391 phase picking for low magnitude events and, the detection of a higher number of low magnitude earthquakes (<  
392 2.5) than before the lockdown. In addition, there were more felt reports during lockdown in both countries, mainly  
393 in the urban areas of Central and Southeastern Costa Rica, and in Guatemala City and Huehuetenango. Although  
394 we present some evidences to link the reduction in seismic noise and the increase in earthquake detections and  
395 felt reports, the variations in the magnitude of completeness and the a- and b- values are not significant enough  
396 before and during the lockdown to confirm that connection or to discriminate other possible contributing factors.  
397

398 Finally, we suggest that there could be a connection between the lockdown measures and the number of felt reports  
399 for smaller earthquakes ( $M < 3.5$ ) in Costa Rica and Guatemala. This possible effect may have been induced by a  
400 more quiescence environment, more people with home office measures and longer time at homes, which might  
401 have stimulated a higher sensitivity in the population to feel low magnitude earthquakes and to report them to the  
402 seismic agencies. This work demonstrates that seismic networks can monitor population mobility and  
403 consequently can be used to verify the compliance of lockdown measures and to explore the consequences of  
404 reducing seismic noise in earthquake detection and felt reports.

405

406

#### 407 **Code availability**

408 All the codes used to analyze the seismic data are available in Lecocq et al. (2020b).

409

#### 410 **Data availability**

411 The data is available by FDSN web services for INSIVUMEH and RSN-UCR seismic networks.

412

#### 413 **Author contributions**

414 MA designed the study and wrote the paper with contributions from all co-authors. MA and DC processed the  
415 seismic data and made figures. All the authors interpreted and analyzed the results and revised the article.  
416

#### 417 **Competing interests**

418 There are no competing interests.

419

#### 420 **Acknowledgments**

421 This work was partially supported by the University of Costa Rica (UCR) projects 113-B5-704 “Vigilancia  
422 sísmica de Costa Rica” and 113-B9-911 “Programa de Investigación Red Sismológica Nacional”. Financial  
423 support at the RSN for seismic instrumentation has been mainly provided by the UCR and the Law Number 8488  
424 of the Republic of Costa Rica. Financial support at the INSIVUMEH for seismic instrumentation has been mainly  
425 provided by the government of Guatemala. We thank Dr. Koen Van Noten (Topical Editor), Dr. Valerio de Rubeis  
426 and Dr. Alan Kafka (referees), for their comments and suggestions to improve the manuscript. Also, special thanks

427 to all the personnel of all Central American seismic networks for maintaining the seismic instruments, locating  
428 systems, and computer programs in operation, most of the time under very difficult conditions.

429

430 **References**

431 Adamek, S., Frohlich, C., and Pennington, W.: Seismicity of the Caribbean-Nazca Boundary: Constraints on  
432 Microplate Tectonics of the Panama Region. *J. Geophys. Res.*, 93, 2053-2075,  
433 <https://doi:10.1029/JB093iB03p02053>, 1998.

434

435 Aki, K.: Maximum likelihood estimated of  $b$  in the formula  $\log N = A - b^*M$  and its confidence limits. *Rev.*  
436 *Earthquakes Res. Inst. Tokyo Univ.*, 43, 237-239, 1965.

437

438 Alonso-Henar, J., Montero, W., Martínez-Díaz, J., Álvarez-Gómez, J., Insua-Arévalo, J., and Rojas, W.: The  
439 Aguacaliente Fault, source of the Cartago 1910 destructive earthquake (Costa Rica). *Terra Nova*. 25(5), 368-373.  
440 doi: 10.1111/ter.12045, 2013.

441

442 Alvarado, G., Benito, B., Staller, A., Climent, Á., Camacho, E., Rojas, W., and Lindholm, C.: The new Central  
443 American seismic hazard zonation: Mutual consensus based on up to day seismotectonic framework.  
444 *Tectonophysics*, 721 462–476, <https://dx.doi.org/10.1016/j.tecto.2017.10.013>, 2017.

445

446 Arroyo-Solórzano, M., and Linkimer, L.: Spatial variability of the  $b$ -value and seismic potential in Costa Rica.  
447 *Tectonophysics*, 814 (2021) 228951, <https://doi.org/10.1016/j.tecto.2021.228951>, 2021.

448

449 Ashenden, C., Lindsay, J., Sherburn, S., Smith, I., Miller, C., and Malin, P.: Some challenges of monitoring a  
450 potentially active volcanic field in a large urban area: Auckland volcanic field, New Zealand. *Nat. Hazards*, 59,  
451 507–528. <https://doi:10.1007/s11069-011-9773-0>, 2011.

452

453 Atkinson, G. and Wald, D.: “Did You Feel It?” intensity data: A surprisingly good measure of earthquake ground  
454 motion. *Seismological Research Letters*, 78(3), 362-368, 2007.

455

456 Blackman, R., Tukey, J., and Tukey, W.: The measurement of power spectra from the point of view of  
457 communications engineering—Part I. *Bell Syst. Tech. J.*, 37, 185–282, <https://doi:10.1002/j.1538-7305.1958.tb03874.x>, 1958.

459

460 Dengler, L. A. and Dewey, J. W.: An Intensity Survey of Households Affected by the Northridge, California,  
461 Earthquake of 17 January 1994. *Bulletin of the Seismological Society of America*, 88, 441-462, 1998.

462

463 Díaz, J., Ruiz, M., Sánchez-Pastor, P. S., and Romero, P.: Urban Seismology: On the origin of earth vibrations  
464 within a city. *Sci. Rep.*, 7, 1–11, 2017.

465

466 Espinosa, A.F.: The Guatemalan Earthquake of February 4, 1976. *US Geol. Surv. Prof. Pap.* 1002, 90, 1976.

467

468 GEM.: The OpenQuake-engine User Manual, Global Earthquake Model (GEM) OpenQuake Manual for Engine  
469 version 3.9.0, doi: 10.13117/GEM.OPENQUAKE.MAN.ENGINE.3.9.0, 183 pages, 2020.

470

471 Gempa: SeisComp 3 Real time data acquisition and processing [software computacional].  
472 <https://www.seiscomp.de/doc/index.html>, last access : 20 Dicembre 2020, 2019.

473

474 Green, D. N., and Bowers, D.: Seismic raves: Tremor observations from an electronic dance music festival.  
475 *Seismological Research Letters*, <https://doi: 10.1785 / gssrl.79.4.546> Page 15/22, 2008.

476  
477 Green, D., Bastow, I., Dashwood, B., and Nippes, S.: Characterizing Broadband Seismic Noise in Central  
478 London. *Seismological Research Letters*, 88, 113–124, <https://doi:10.1785/0220160128>, 2017.

479  
480 Groos, J. C., and Ritter, J. R. R.: Time domain classification and quantification of seismic noise in an urban  
481 environment. *Geophys. J.*, <https://doi: 10.1111 / j.1365-246X.2009.04343.x>, 2009.

482  
483 Gutenberg, B., and Richter, C: Frequency of earthquakes in California. *Bulletin of the Seismological Society of*  
484 *America*, 34, 185-188, 1944.

485  
486 Harlow, D.H., White, R.A., Rymer, M.J., and Alvarez, S.: The San Salvador earthquake of 10 October 1986 and  
487 its historical context. *Bull. Seismol. Soc. Am.* 83 (4), 1143–1154, 1993.

488  
489 Havskov, J., Voss, P. H., and Ottemöller, L.: Seismological observatory software: 30 Yr of SEISAN.  
490 *Seismological Research Letters*, 91(3), 1846-1852, <https://doi.org/10.1785/0220190313>, 2018.

491  
492 Hong, T.-K., Lee, J., Lee, G., Lee, J., and Park, S.: Correlation between Ambient Seismic Noises and Economic  
493 Growth. *Seismological Research Letters*, 91, 2343–2354, <https://doi: 10.1785 / 0220190369>, 2020.

494  
495 Información de la Red Sismológica Nacional de Costa Rica (RSN-UCR): Universidad de Costa Rica,  
496 <https://doi.org/10.15517/TC>, 2017.

497 Instituto Nacional De Sismología, Vulcanología, Meteorología e Hidrología de Guatemala (INSIVUMEH): Red  
498 Sismológica Nacional [Data set]. International Federation of Digital Seismograph Networks.  
499 <https://doi.org/10.7914/SN/GI>, 2013.

500 Instituto Nicaragüense De Estudios Territoriales (INETER): Nicaraguan Seismic Network. Instituto Nicaragüense  
501 de Estudios Territoriales (INETER), <https://doi.org/10.7914/SN/NU>, 1975.

502  
503 Kellogg, J.N. and Vega, V.: Tectonic development of Panama, Costa Rica and the Colombian Andes: constraints  
504 from global positioning geodetic systems and gravity. In: Mann, P. (Ed.), *Geologic and Tectonic Development of*  
505 *the Caribbean Plate Boundary in Southern Central America*, pp. 75–90 (GSA Special Paper, 295), 1995

506  
507 Kuzma, H. A.: Vehicle traffic as a source for near-surface passive seismic imaging. *Symposium on the Application*  
508 *of Geophysics to Engineering and Environmental Problems*, 2009 609–615, <https://doi.org/10.4133/1.3176748>,  
509 2009.

510  
511 Lecocq, T. et al.: Global quieting of high-frequency seismic noise due to COVID-19 pandemic lockdown  
512 measures. *Science*, <http://science.sciencemag.org/content/early/2020/07/22/science.abd2438>, 2020a.

513  
514 Lecocq, T., Massin, F., Satriano, C., Vanstone, M., and Megies, T.: Seismo RMS - A simple Python/Jupyter  
515 Notebook package for studying seismic noise changes. <https://zenodo.org/record/3820046#.YBaXbS3pM1K>,  
516 2020b.

517 Linkimer, L., Arroyo, I. G., Alvarado, G. E., Arroyo, M., and Bakkar, H.: The National Seismological Network  
518 of Costa Rica (RSN): An Overview and Recent Developments. *Seismological Research Letters*, 89 (2A), 392–  
519 398, <https://doi.org/10.1785/0220170166>, 2018.

520 Linkimer, L and Arroyo, I.: Ciencia ciudadana y herramientas de comunicación en la Red Sismológica Nacional  
521 de la Universidad de Costa Rica. *Revista Comunicación*, 29 (2), 5-21, ISSN: 0379-3974 / e-ISSN1659-3820,  
522 2020.

523 Mann, P., Schubert, C. and Burke, K.: Review of Caribbean neotectonics. In: Dengo, G., Case, J.E. (Eds.), The  
524 Caribbean Region. The Geology of North America. Geol. Soc. Amer, Boulder, Colorado, pp. 307–338, 1990.

525 McNamara, D., and Buland, R.: Ambient Noise Levels in the Continental United States. Bull. Seismol. Soc. Am.,  
526 94, 1517–1527, <https://doi:10.1785/012003001>, 2004.

527

528 Nimiya, H., Ikeda, T., and Tsuji, T.: Temporal changes in anthropogenic seismic noise levels associated with  
529 economic and leisure activities during the COVID-19 pandemic. Research Square,  
530 <https://doi.org/10.21203/rs.3.rs-77786/v1>, 2020.

531

532 Piccinini, D., Giunchi, C., and Olivieri, M.: COVID-19 lockdown and its latency in Northern Italy: seismic  
533 evidence and socio-economic interpretation. Sci Rep 10, 16487, <https://doi.org/10.1038/s41598-020-73102-3>,  
534 2020.

535

536 Poli, P., Boaga, J., Molinari, I., Cascone, V., and Boschi, L.: The 2020 coronavirus lockdown and seismic  
537 monitoring of anthropic activities in Northern Italy. Sci Rep 10, 9404, <https://doi.org/10.1038/s41598-020-66368-0>, 2020.

539

540 Riahi, N., and Gerstoft, P.: The seismic traffic footprint: Tracking trains, aircraft, and cars seismically. Geophys.  
541 Res. Lett., <https://doi: 10.1002 / 2015GL063558>, 2015.

542

543 Servicio Nacional de Estudios Territoriales (SNET), El Salvador (SNET-BB).: International Federation of Digital  
544 Seismograph Networks, <https://www.fdsn.org/networks/detail/SV/>, 2004.

545

546 Sistema de Integración Centroamericana (SICA) (2020). Observatorio Regional SICA-COVID 19.  
547 <https://www.sica.int/coronavirus/observatorioSICACOVID19/medidas/Costarica>, last access: December 20,  
548 2020.

549

550 Sohrabi, C., Alsafi, Z., O'Neill, N., Khan, M., Kerwan, A., Al-Jabir, A., Iosifidis, C., Agha, R.: World Health  
551 Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-19). Int. J. Surg.,  
552 76:71-76, <https://doi: 10.1016/j.ijsu.2020.02.034>, 2020.

553

554 Somala, S. N.: Seismic noise changes during COVID-19 pandemic: a case study of Shillong, India. Nat. Hazards,  
555 103, 1623–1628, <https://doi.org/10.1007/s11069-020-04045-1>, 2020.

556

557 Trenkamp, R., Kellogg, J.N., Freymueller, J.T. and Mora, H.P.: Wide plate margin deformation, southern Central  
558 America and northwestern South America, CASA GPS observations. J. S. Am. Earth Sci. 15, 157–171, 2002.

559

560 Vargas, C.A. and Mann, P.: Tearing and breaking off of subducted slabs as the result of collision of the Panama  
561 arc-indenter with northwestern South America. Bull. Seismol. Soc. Am. 103 (3), 2025–2046, 2013.

562

563 Wald, D.J., Quitoriano, V., Worden, C.B., Hopper, M. and Dewey, J. W.: USGS “Did You Feel It?” Internet-  
564 based Macroseismic Intensity Maps. Annals of Geophysics. 54 (6), 688-707, doi: 10.4401/ag-5354, 2011.

565

566 Weichert, D.: Estimation of the Earthquake Recurrence Parameters for Unequal Observation Periods for Different  
567 Magnitudes. Bulletin of the Seismological Society of America, 70.4, 1337 – 1346, 1980.

568

569 Welch, P.: The use of fast Fourier transform for the estimation of power spectra: A method based on time  
570 averaging over short, modified periodograms. IEEE Trans. Audio Electroacoust, 15, 70–73,  
571 <https://doi:10.1109/TAU.1967.1161901>, 1967.

572

573 Wiemer, S., and Wyss, M.: Minimum Magnitude of Completeness in Earthquake Catalogs: Examples from  
574 Alaska, the Western United States, and Japan. *Bulletin of the Seismological Society of America*, 90 (4), 859–869,  
575 2000.

576  
577 Woessner, J., and Wiemer, S.: Assessing the Quality of Earthquake Catalogues: Estimating the Magnitude of  
578 Completeness and Its Uncertainty. *Bulletin of the Seismological Society of America*, 95 (2), 684–698, 2005.

579  
580  
581  
582  
583

## 584 Tables

585  
586 **Table 1.** Summary of observations when comparing the earthquake detections and felt reports from the time  
587 before the lockdown (BL) and during lockdown (DL) for Costa Rica and Guatemala (see also Figures 7, 8, 9, 10  
588 and A2). The observations that favor an increase in lower magnitude earthquakes (LM) are marked.


589

Observations	Costa Rica		Guatemala	
	$M \leq 3.5$	$M > 3.5$	$M \leq 3.5$	$M > 3.5$
<i>Number of detected earthquakes</i>	Slightly decrease	No change	Slightly increase (LM)	Increase
<i>Number of picked phases</i>	~20% increase (LM)	Slightly increase	~40% increase (LM)	No change
<i>Number of Felt earthquakes</i>	Increase (LM)	Decrease	No change	Increase
<i>Number of felt earthquakes reports</i>	No change	Increase	-----	
<i>Mc</i>	Slightly increase		Slightly increase	
<i>a-value</i>	Slightly decrease		Increase (LM)	
<i>b-value</i>	Slightly increase (LM)		Slightly increase (LM)	

590


591 **Figures**

592




593

594 **Figure 1.** Map of Central American and location of the selected seismic stations from Costa Rica, Guatemala, El  
 595 596 597 598 599 600  
 599 San Salvador, El Salvador. © ESRI and its data partners (ArcGis Services).




604 **Figure 2.** High-frequency seismic anthropogenic noise displacement (HFSAND-RMS) evolution near capitals of  
 605 the countries studied in an hourly grid representation. a) TC.SJS1 station in San Jose, Costa Rica. b) GI.GCG4  
 606 station in Guatemala City, Guatemala. c) SV.CEDA station, near San Salvador in the city of La Libertad. d)  
 607 NU.MGAN station in Managua, Nicaragua. Gaps correspond to periods for which seismic data were unavailable  
 608 and the vertical red lines indicate the time when the first lockdown measures started in Central America. The  
 609 numbers 1, 2, and 3 show the New Year holidays of 2019, Easter of 2020, and a brief period in July 2020 when a  
 610 strict return of lockdown measures was implemented in Costa Rica, respectively.  
 611




612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623

**Figure 3.** High-frequency seismic anthropogenic noise displacement (HFSAND-RMS) evolution at the seismic stations near capitals of the countries studied. a) TC.SJS1 station in San Jose, Costa Rica. b) GI.GCG4 station in Guatemala City, Guatemala. c) SV.CEDA station, near San Salvador in the city of La Libertad. d) NU.MGAN station in Managua, Nicaragua. The blue line corresponds to the RMS amplitude time series of the vertical component, filtered between 4 and 14 Hz, and the orange line corresponds to median day-time, between the 6 and 16 hours of local time. Gaps correspond to periods for which seismic data were unavailable and the vertical red lines indicate the time when the first lockdown measures started in Central America. The numbers 1, 2, and 3, show the New Year holidays of 2019, Easter of 2020, and a brief period in July 2020 when a strict return of lockdown measures was implemented in Costa Rica, respectively.




624  
625  
626  
627  
628  
629  
630  
631  
632  
633


**Figure 4.** High-frequency seismic anthropogenic noise displacement (HFSAND-RMS) evolution for specific sites in Costa Rica and Guatemala in an hourly grid representation. a) TC.BELE (central Costa Rica, near Juan Santamaria international airport), TC.ERIA (northern Costa Rica, urban area), and TC.ZEDO (southern Costa Rica, urban area). b) GI.HUEH (northwest of Guatemala, urban area), GI.RETA (southwest of Guatemala, urban area), and GI.CHIE (east of Guatemala, urban area). The gaps correspond to the time periods for which seismic data were unavailable and the vertical red lines indicate the time when the first lockdown measures started in Central America. The numbers 1 and 2 show the New Year holidays of 2019 and Easter of 2020, respectively.




**Figure 5.** Daily variation of the median seismic noise per weekday, before and during the lockdown (LD). a) TC.SJS1 station in San Jose, Costa Rica. b) GI.RETA station in Retalhuleu, southwest of Guatemala.



**Figure 6.** Percent of change of the high-frequency seismic anthropogenic noise displacement (HFSAND-RMS) in the band 4-14Hz during the period from January 15 to March 15, 2020 (before lockdown measures) with respect to the time interval from March 16 to May 15, 2020 (after the lockdown measures were applied). a) Percent of change in seismic stations of Guatemala and El Salvador. b) Percent of change in seismic stations of Nicaragua and Costa Rica. © OpenStreetMap contributors 2020. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.




**Figure 7.** Event detection and felt earthquakes reported in Costa Rica before (1 Nov 2019-15 Mar 2020) and during (16 Mar-31 Jul 2020) lockdown (LD) measures. a) Number of detected earthquakes. b) Earthquake magnitude versus the average number of P-wave picks per Mw for the events with Mw < 4.0, and the corresponding linear fits. The inset graph shows the average difference in the number of picks for magnitude bins for the time periods before (BL) and during (DL) the lockdown. c) Number of felt earthquakes in Costa Rica. d) Magnitude versus the number of reports from the population through the RSN application for smartphones "¿Lo Sentiste?" for events with Mw < 5.0, and the corresponding linear fits.




658  
659  
660  
661  
662  
663  
664  
665

**Figure 8.** Magnitude-frequency distribution for earthquakes. a) Costa Rica before lockdown. b) Costa Rica during lockdown. c) Guatemala before lockdown. d) Guatemala during lockdown. Green bars represent the incremental (non-cumulative) number of earthquakes and yellow circles the cumulative distribution of events. The grey solid line fits the data points for the cumulative distribution above the magnitude of completeness (Mc). Vertical lines indicate the Mc estimated from the maximum curvature (MAXC) method.




666  
667 **Figure 9.** Map of felt earthquakes reported in Costa Rica. a) Before lockdown measures (1 Nov 2019-15 Mar  
668 2020). b) During lockdown measures (16 Mar-31 Jul 2020).  
669



670  
671  
672  
673  
674  
675

**Figure 10.** Event detection and felt earthquakes reported in Guatemala before (1 Nov 2019-15 Mar 2020) and during (16 Mar-31 Jul 2020) lockdown (LD) measures. a) Number of detected earthquakes. b) Earthquakes magnitude versus the average number of P-wave picks per Mw for the events with  $Mw < 4.0$ , and the corresponding linear fits. The inset graph shows the average difference in the number of picks for magnitude bins for the periods before (BL) and during (DL) lockdown. c) Number of felt earthquakes in Guatemala.





**Figure A1.** Images from Open Street Maps of the site areas where the seismic stations (orange triangles) used are located. © OpenStreetMap contributors 2020. Distributed under the Open Data Commons Open Database License (ODbL) v1.0.

685 **Table A1.** Main mobility lockdown measures between March and October 2020, in Costa Rica, Guatemala, and  
 686 El Salvador during the Covid-19 pandemic according with the “Sistema de Integracion Centroamericana” (SICA),  
 687 (2020). Nicaragua did not establish lockdown measures for social mobility. On March 18 the first positive case of  
 688 Covid-19 was found in Nicaragua.  
 689

Month	Costa Rica	Guatemala	El Salvador
<i>March 2020</i>	<p><b>Day 6</b>, the first positive case for Covid-19 is detected.</p> <p><b>Day 9</b>, the National Emergency Commission (CNE) and the Ministry of Health declare a yellow health alert.</p> <p><b>Day 12</b>, closure of schools at risk; 50% reduction in capacity in meeting spaces; suspension of trips abroad for public employees.</p> <p><b>Day 16</b>, a state of national emergency is declared.</p> <p><b>Day 17</b>, closure of public and private educational centers, closure of non-essential stores, prohibition of mass events and total closure of beaches throughout the country.</p> <p><b>Day 24</b>, vehicle mobility restriction from 22h to 5h.</p>	<p><b>Day 6</b>, red alert is extended to the entire country after the first positive case for Covid-19 is detected</p> <p><b>Day 16</b>, classes are suspended and the borders with El Salvador are closed.</p> <p><b>Day 17</b> interruption of work activities in the public and private sectors, suspension of public transport, prohibition of meetings of any kind, religious and sport activities, closing of shopping centers. The total closure of air and land borders is implemented.</p>	<p><b>Day 11</b>, interruption of educational activities</p> <p><b>Day 16</b>, discontinuation of public and sport shows, closing of bars and gyms and non-essential shops.</p> <p><b>Day 17</b>, El Salvador International Airport closes operations.</p> <p><b>Day 18</b>, first positive case for Covid-19 detected.</p> <p><b>Day 22</b>, mandatory 30-day quarantine and ban on crowds.</p>
<i>April 2020</i>	<p><b>Day 1</b>, during Easter the daytime vehicle restriction was in place from 5h-17h. Vehicular traffic was allowed only one day per week according to the license plate number to make essential purchases.</p> <p><b>Day 11</b>, a nighttime vehicle restriction is established 19h-5h until May 15.</p>	<p><b>Day 1</b>, Greater air monitoring at Borders.</p> <p><b>Day 9</b>, break in school lessons for the whole month.</p>	<p><b>Day 3</b>, no circulation on beaches, rivers, lakes, spas, or tourist centers in the country.</p> <p><b>Day 13</b>, entire population to be kept in compulsory home security with some exceptions, extended until April 28.</p>
<i>May 2020</i>	<p><b>Day 1</b>, all public spaces and shops work at 50%; gradual reactivation of work centers from 5h to 19h.</p> <p><b>Day 16</b>, vehicle restriction is maintained from 5h to 19h. Opening of hotels with a capacity of 50%. Opening of some national parks (50%). Beaches are open Monday-Friday from 5h to 8h.</p>	<p><b>Day 4</b>, opening of shopping malls with few stores. Activities and public transport, as well as classes are suspended. Curfew from 18h to 4h. Prohibition of transit between territorial departments.</p> <p><b>Day 14</b>, markets can open Monday, Wednesday, and Thursday from 6h to 13h, as well as supermarkets and convenience stores, from 9h to 16h. Closure of the country under restriction of total mobilization.</p> <p><b>Day 25</b>, curfew 17h-5h with vehicle restriction between territorial departments. Total closure of the country.</p>	<p><b>Day 7</b>, restriction of mobility between municipalities, people can circulate according to their occupation.</p> <p><b>Day 10</b>, public transport may only circulate to mobilize duly identified health personnel.</p>

<i>June 2020</i>	<p><b>Day 1</b>, tourist transport is enabled with restrictions and special measures. Operation of gyms, restaurants, and museums with a capacity of 50%</p> <p><b>Day 20</b>, access to beaches from 5h-9:30h</p> <p><b>Day 26</b>, mandatory use of masks is established.</p>	<p><b>Day 15</b>, mobilization according to the last digit of the car plate in the territorial departments.</p>	<p><b>Day 16</b>, restriction of mobility according to the last number of the identity document.</p>
<i>July 2020</i>	<p><b>Day 3</b>, public parks, bars and massive events closed.</p> <p><b>Day 11</b>, vehicle restriction for the whole country from 5h to 17h according to license plate number.</p> <p><b>Day 14</b>, increase in the capacity in public transport from 20% to 50% in routes shorter than 75 km.</p> <p><b>Day 20</b>, vehicle restriction from 17h-5h according to license plates, but only in areas of orange alert.</p>	<p><b>Day 13</b>, curfew is established from Monday to Friday from 18h to 5h and Saturdays from 14h to 5h.</p> <p><b>Day 26</b>, vehicle restriction by license plate is eliminated and the curfew is modified from 21h to 4h.</p>	<p><b>Day 6</b>, high restrictions are extended for 15 days, due to the increase in infections and deaths from Covid-19.</p> <p><b>Day 29</b>, beginning of a new phase of economic reopening.</p>
<i>August 2020</i>	<p><b>Day 1</b>, opening of commercial flights with limited routes from Europe.</p> <p><b>Day 31</b>, Costa Ricans who return to the country may not quarantine. New vehicle restriction for weekdays 5h-22h and weekends 5h-20h.</p>	<p><b>Day 24</b>, religious activities are reestablished with a maximum of one hour and a limit of attendees.</p> <p><b>Day 26</b>, reopening of the La Aurora Zoo.</p>	<p><b>Day 9</b>, personal clusters are established with groups not exceeding ten people.</p> <p><b>Day 24</b>, beginning of the "Transitory Phase" in the process of the gradual reactivation of the economy.</p>
<i>September 2020</i>	<p><b>Day 9</b>, temporary suspension of activities that involve massive movements of people. Controlled opening of economic activities.</p>	<p><b>Day 6</b>, public servants work hours from 7h to 15h.</p> <p><b>Day 18</b>, international airport reopens. Entry to the country conditioned to a negative PCR test.</p> <p><b>Day 28</b>, vehicle restriction from 9h to 16h. Prohibition of alcohol sale between 19h and 5h.</p>	<p><b>Day 1</b>, start of a new phase of economic reopening which includes public transportation and most of economic activities.</p>
<i>October 2020</i>	<p><b>Day 1</b>, opening of borders for travelers from California and Ohio and Mexico and Jamaica.</p> <p><b>Day 15</b>, opening of flights from Central America.</p> <p><b>Day 26</b>, foreign travelers are not required to present a negative Covid-test.</p> <p><b>Day 27</b>, permission to use outdoor spaces for recreation.</p>	<p><b>Day 1</b>, reopening of higher education centers and technical training centers.</p> <p><b>Day 7</b>, hotel and tourism workers are allowed to hold events with restricted capacity.</p>	<p><b>Day 3</b>, cultural spaces reopen. Opening of stadiums and public shows with social distancing (two meters) and beach trips at restricted hours. Opening of museums, cinemas, and hotels with capacity reduced to 50%. The public sector in general returns to work.</p>

**Table A2.** Questions on the RSN module "*¿Lo Sentiste?*" (Linkimer and Arroyo, 2020).

Number	Question
1	Did you feel it?
2	What were you doing?
3	Where were you?
4	Did others nearby feel it?
5	How would you describe the shaking?
6	How did you react?
7	Was it difficult to stand and/or walk?
8	Did light objects move or fall from the shelves?
9	Did pictures on walls move or get knocked askew?
10	Did the furniture fall, overturn or fall?
11	Was there any damage to the buildings?
12	Additional comments on effects in nature, such as landslides, cracks in the ground, among others?