

Elastic anisotropies of rocks in a subduction and exhumation setting

Michael J. Schmidtke¹, Ruth Keppler¹, Jacek Kossak-Glowczewski¹, Nikolaus Froitzheim¹, Michael Stipp²

¹ Institut für Geowissenschaften, Rheinische Friedrich-Wilhelms-Universität, Bonn, 53115, Germany

² Institute of Geosciences and Geography, Martin-Luther University, Halle (Saale), 06099, Germany

Correspondence to: Michael J. Schmidtke (Michael.schmidtke@uni-bonn.de)

Abstract. Subduction and exhumation are key processes in the formation of orogenic systems across the world, for example, in the European Alps. For geophysical investigations of these orogens, it is essential to understand the petrophysical properties of the rocks involved. These are the result of a complex interaction of mineral composition and rock fabric including mineral textures (i.e. crystallographic preferred orientations). In this study we present texture-derived elastic anisotropy data for a representative set of different lithologies involved in the Alpine orogeny. Rock samples were collected in the Lago di Cignana area in Valtournenche, in the Italian Northwestern Alps. At this locality a wide range of units of continental and oceanic origin with varying paleogeographic affiliations and tectono-metamorphic histories are accessible.

Their mineral textures were determined by time-of-flight neutron diffraction ~~at the Frank Laboratory of Neutron Physics at the JINR in Dubna, Russia.~~ From these data the elastic properties of the samples were calculated. The data set includes representative lithologies from a subduction-exhumation-setting. In subducted lithologies originating from the oceanic crust, the ~~elastic-P-wave~~ anisotropies (~~AVP [%]~~) range from 1.4 to ~~5.03.7~~ % with average P-wave velocities of ~~7.0420~~-8.24 km/s and VP/VS-ratios of ~~1.7470~~-~~1.7675~~. In the metasediments of the former accretionary prism the ~~elastic-AVP anisotropies~~ range from ~~4.73.7~~ to ~~8.27.1~~ %-.%. ~~This tectonic setting displays~~ average P-wave velocities ~~of are~~ ~~6.4766~~-7.23 km/s and VP/VS-ratios ~~of are~~ ~~1.610~~-1.76. Continental crust which is incorporated in the collisional orogen shows ~~elastic anisotropies~~AVP ranging from ~~1.84~~ to ~~2.81~~ % with average P-wave velocities of ~~6.4252~~-~~6.5162~~ km/s and VP/VS-ratios of 1.56-1.60. Our results suggest that mafic and felsic rocks in subduction zones at depth may be discriminated by a combination of seismic signatures: lower anisotropy and higher VP/VS ratio for mafic rocks, higher anisotropy and lower VP/VS ratio for felsic rocks and metasediments.

1. Introduction

During subduction, exhumation and continental collision, rocks of different paleogeographic affiliations are ~~brought juxtaposed together~~, strongly deformed and submitted to variable metamorphic conditions. Oceanic crust and its sedimentary cover, as well as continental fragments are frequently subducted before or during continental collision. This leads to a large variety of lithologies within the subduction zone from ultramafic rocks originating from mantle material, ~~over-through~~ mafic rocks originating from oceanic crust, to felsic rocks which were either part of continental fragments, ~~continental margins~~, or

sedimentary cover of the oceanic rocks. Deformation during subduction and exhumation ~~causes the formation of~~ produces crystallographic preferred orientation (CPO), which in turn leads to elastic anisotropy of the rocks involved. The metamorphic overprint also influences their seismic velocities and Poisson's ratios. With increasing depth, the rocks undergo blueschist- to eclogite-facies metamorphic conditions. During exhumation the rocks are frequently exposed to further metamorphic overprint, depending on various factors such as exhumation speed and available water content (e.g., Peacock, 1993). These rocks ~~which that~~ can now be found in outcrops at the Earth's surface yield insights into ~~the variable various~~ stages of subduction, collision, and exhumation, which can be compared to the geophysical investigations of such rocks at depth.

Such high-resolution geophysical imaging of 3D structures is currently taking place within the AlpArray initiative using a high-end seismological array within the Alpine orogen. For this project, knowledge of the ~~elastic seismic~~ anisotropy of the imaged rocks is of great interest, because in conjunction with the VP/VS-ratio it is a key parameter for the interpretation of geological structures in the subsurface. Receiver-function analysis allows separating crustal from mantle anisotropy in the Alps (Link and Rumpker, 2019). Because of the larger lithological heterogeneity of the crust, as compared to the mantle, information on the seismic properties of various crustal rocks is required for interpreting the crustal anisotropy pattern.

Elastic anisotropies of rocks can be calculated from mineral compositions, single-crystal mineral properties, and the CPOs of all involved mineral phases (Christoffel, 1874; Mainprice and Humbert, 1994). They are also influenced by other factors, such as shape preferred orientation (SPO) of the minerals, their grain sizes, and microcracks in the rocks (Siegesmund and Vollbrecht, 1991; Siegesmund et al., 1993; Ullemeyer et al., 2011). With increasing depth, ~~the importance of~~ porosity and microcracks ~~are closed is diminished~~, leading to CPO as the most important factor influencing elastic anisotropies in the deeper parts of subduction zones and collisional orogens.

In this study, we collected a set of rock samples from the Lago di Cignana area in Valtournenche, western Alps (figure 1a and 1b). The samples are representative of various stages within a subduction- and exhumation-cycle, including rocks of oceanic and continental origin and variable degrees of metamorphism and deformation (figure 2). The CPOs of the mineral phases of the samples were acquired by time-of-flight (TOF) neutron diffraction. Measurements were conducted at the SKAT diffractometer of the Frank Laboratory of Neutron Physics at the JINR in Dubna, Russia (Ullemeyer et al., 1998, 2006, 2010; Ivankina et al., 2000; Ivankina et al., 2001; Nikitin et al. 2004; Kern et al., 2008; Keppler et al., 2014). This method has the advantage of being able to measure bulk CPOs of all mineral phases in large, polyphase, coarse-grained samples simultaneously (Keppler et al., 2014). From the CPO and single-crystal elastic anisotropy data the seismic properties of all collected samples were calculated. These data provide a catalogue for seismic imaging and geodynamic models of the Alpine orogen as well as other subduction zones and continental collision zones around the world. Among the texture-derived properties presented are the P- and S-wave velocities, the VP/VS-ratio and the ~~elastic P-wave~~ anisotropies of each sample.

2. Geological overview of the study area

The samples in this study were collected in the western Alps in northwestern Italy (figure 1a). In this area the Alpine orogen
65 consists of tectonic nappes of continental and oceanic origin with various paleogeographic provenances and tectono-
metamorphic histories. ~~This-The~~ nappe stack formed from continent-continent- collision of Europe and Adria occurring in
the Palaeogene. The units in ~~this-the~~ nappe stack were accreted to the Adriatic margin during SE-directed subduction. The
units from which the samples were collected are derived from the Piemont-Ligurian ocean and the Margna-Sesia continental
fragment (Schmid et al, 2004), also termed Cervinia (Pleuger, et al., 2007).

70 At Lago Di Cignana (figure 1b), a cross-section through continental and oceanic units is ~~very well accessible~~ exposed.
Generally, the units exhibit a shallowly NW dipping foliation and SE-NW to E-W trending lineation (figure 1c). The oceanic
Zermatt-Saas zone is found in the footwall, separated from the continental hanging wall unit of the Dent Blanche Nappe by
the oceanic and metasedimentary Combin zone. These three units are separated by shear zones and faults; ~~and clear jumps~~
differences in metamorphic grade can be found. For this study, metabasites (eclogite, blueschist, amphibolite and
75 greenschist) were collected from the Zermatt-Saas zone. Furthermore, several metasedimentary rocks of ~~varying various~~
compositions were collected from the Combin zone as well as two gneiss samples from the Dent Blanche Nappe.

The rocks in this study followed different P-T paths. The Dent Blanche unit has experienced peak P-T conditions of 1.4-1.6
GPa and 450-520 °C (Ballèvre et al., 1986, Oberhänsli and Bucher, 1987; Angiboust et al., 2014; Manzotti et al., 2014). The
age of blueschist-facies metamorphism was dated ~~to-at~~ 48-43 Ma by Rb-Sr geochronology (Angiboust et al., 2014) in the
80 Arolla mylonites at the base of the Dent Blanche unit.

The Combin zone displays P-T conditions of 1.2 GPa and 450°C (Bousquet, 2008). The age of blueschist- to greenschist-
facies metamorphism in the Combin zone has been dated between 48 and 36 Ma (Reddy et al., (1999, 2003)) by Rb-Sr and
Ar/Ar methods.

In the Zermatt-Saas zone the P-T path involved prograde blueschist-facies conditions, High Pressure (HP) to Ultra-High
85 Pressure (UHP) peak conditions and retrograde amphibolite- and greenschist-facies overprints. Peak conditions reached 2.5-
3 GPa and 550-600 °C (Bucher et al., 2005), with locally UHP peak conditions of 3.2 GPa and over 600 °C (Groppo et al.,
2009; Frezzotti et al., 2011). The UHP-peak UHP metamorphism was dated ~~to-at~~ 40.6±2.6 Ma by Sm-Nd garnet (Amato et
al., 1999), to 44-43 Ma by the Ar/Ar step-heating method on phengite (Gouzu et al., 2006) and ~~to-at~~ 48.8 ± 2.1 Ma by Lu-Hf
on garnet-omphacite-whole-rock (Lapen et al., 2003). The greenschist-facies retrograde overprint was dated ~~to-at~~ 38 ± 2 Ma
90 by Rb-Sr whole-rock-phengite chronology (Amato et al., 1999).

3. Methods

CPOs were measured by the TOF neutron diffractometer SKAT at the Frank Laboratory of Neutron Physics at JINR in
Dubna, Russia (Keppler et al., 2014; Ullemeyer et al., 1998). Neutron diffraction has several advantages over other texture
analysis methods such as optical measurement and electron and ~~*X~~-ray diffraction analysis (Wenk et al. 1998; Ullemeyer et

95 al. 2000; Nikitin and Ivankina, 2004). Among these advantages is the low absorption of thermal neutrons in matter. This allows the measurement of large sample volumes (up to 65 cm³) of complex material such as coarse-grained rocks containing multiple low-symmetry mineral phases (Ullemeyer et al., 2006; Ullemeyer et al., 2010; Ivankina et al., 2005; Kern et al., 2008; Keppler et al., 2014).

100 The diffraction data are processed by Rietveld Texture Analysis (RTA) (Von Dreele, 1997; Matthies et al., 1997, Lutterotti et al., 1997). This method allows for the determination of the textures of all mineral phases in the sample by calculation of the orientation distribution function (ODF) using the E-WIMV algorithm (Lutterotti et al. 2004, Chateigner et al. 2019). The RTA is performed in the software Mineral Analysis Using Diffraction (MAUD: Lutterotti, 2010, Wenk et al., 2010). For further discussion on the functions and limitations of MAUD, see the publication of Wenk et al. (2012). For comparability of texture strength within the sample set we calculated F2 indexes (Bunge et al., 1982; Matthies, et al. 1997) of all mineral phases in the samples of this study (see appendix A). The F2 index is an OFD based texture index in which a completely random texture as found in a powder sample would result in an index of 1, while a perfect single crystal would result in an index of ∞ (Matthies et al., 1997).

105 In RTA, amphiboles are generally very difficult to distinguish, since they are extremely similar crystallographically very similar. For this reason, the crystallographically best fitting amphibole structure was picked for refinement and not, which is not always the chemically the compositional best fitting one in the sample. This procedure has an insignificant/negligible influence on the calculated CPO due to the crystallographic similarity of the amphibole group (C2/m space group; Reynard et al. 1989). For RTA, knowledge/the analysis of the mineral assemblage, is of great importance. For this reason, thin sections parallel to the stretching lineation and perpendicular to the foliation of all samples were made and analyzed by optical polarization microscopy.

115 To calculate the P-wave velocities and other elastic properties of the samples, the program BEARTEX (Wenk et al. 1998) was used. To calculate the bulk-rock elastic moduli, the orientation distribution function (ODF) of the mineral phases, knowledge of their single-crystal elastic constants and their volume fractions is necessary. The elastic moduli were then calculated using the Christoffel equation (1) (Christoffel, 1874/1877; Mainprice and Humbert, 1994):

$$Det[\Gamma_{ij} - V^2\delta_{ij}] = 0 \quad (1)$$

120 In it the phase velocity of P-, S1- or S2-waves is represented by V , ~~the~~ The Kronecker delta is represented by δ_{ij} and the Christoffel Tensor is represented by Γ_{ij} . The Equation of the Christoffel Tensor (2) is as follows:

$$\Gamma_{ij} = a_{ijkl}x_kx_l \quad (2)$$

In it a_{ijkl} represents the density-corrected elastic (stiffness) moduli and x_kx_l represent the direction cosines of the wave propagation direction.

125 ~~For the averaging,~~ It has been shown by Hill (1952) that the widely used averaging schemes of Voigt (1928) and Reuss (1929) give upper and lower bounds of the elastic ~~moduli~~ stiffness, respectively. Furthermore, other averaging schemes have been proposed by Matthies and Humbert (1995) and Matthies (2012). However, since the values of the parameters a_{ijkl} are

130 still uncertain, the Voigt averaging scheme was used consistently throughout this study. It ~~is known~~ should be kept in mind that the recalculated velocities are therefore maximum velocities only. In order to gain these uppermost possible VP and avoid overestimation as well as uncertainties concerning the values of the a_{ijk} -parameters the Voigt averaging scheme was selected.

135 The ODF of all phases are exported from MAUD. The volume percentages of the phases are estimated from thin-sections and calculated in MAUD by RTA. The single-crystal elastic constants were taken from literature (quartz: Heyliger et al., 2003; albite: Brown et al., 2006; muscovite: Vaughan and Guggenheim, 1986; calcite: Chen et al., 2001 ~~Dandekar, 1968~~; dolomite: Humbert & Plique, 1972; hornblende: Aleksandrov and Ryzhova, 1961; epidote/zoisite/clinozoisite: Aleksandrov et al., 1974; garnet: Zhang et al., 2008; glaucophane: Bezacier et al., 2010; omphacite: Bhagat et al., 1992; chlorite: Mookherjee and Mainprice, 2014; actinolite: Brown and Abramson, 2016). In some cases, single-crystal elastic constants are not available for all phases (~~chlorite, actinolite, barroisite and clinozoisite~~), so similar minerals were selected as approximations. ~~For samples containing chlorite we have selected the single crystal elastic constant of muscovite (Vaughan and Guggenheim, 1986) to approximate chlorite.~~ For clinozoisite the single-crystal elastic constants of epidote (Aleksandrov et al., 1974) were selected, for the blueschist-facies amphibole barroisite those of glaucophane (Bezacier et al., 2010) ~~and for actinolite those of hornblende (Aleksandrov and Ryzhova, 1961)~~. In most cases, the contents of the phases for which approximations were necessary, ~~were are~~ below 10% of the total sample volume. At this low content, the CPOs of these minerals are of low relevance for the physical properties of the rock (Mainprice and Ildefonse, 2009). ~~In RTA, amphiboles are generally very difficult to distinguish, since they are extremely similar crystallographically. For this reason, the crystallographically best fitting amphibole structure was picked for refinement, which is not always the chemically fitting one in the sample. This has an insignificant influence on the calculated CPO due to the crystallographic similarity of the amphibole group (C2/m space group; Reynard et al. 1989).~~

140
145
150
155 It is known that at low confining pressures, close to the Earth's surface, the elastic properties of rocks are greatly influenced by microcracks and crack fabrics, as well as a shape preferred orientation (SPO). This has been shown in multiple studies (Kern et al., 2002, 2008, Pros et al., 2003, Ivankina et al., 2005, Ullemeyer et al., 2006). Further studies have even shown that the influence of microcracks is still visible up to a confining pressure of 1 GPa (Christensen, 1974, Ullemeyer et al., 2011). In this study the elastic properties are calculated solely on the basis of mineral texture and single-crystal data without taking microcracks into account. Since the microcracks are closed or almost closed at great depth, the calculated elastic properties can be seen as representative of these conditions.

4. Composition and microfabrics of the samples studied

In this study we have related the subduction and exhumation settings to several structural positions, each containing characteristic lithologies (figure 2). These positions are (1) the oceanic crust of the subducting plate, (2) the sedimentary cover of the subducting plate as well as the accretionary prism and ~~finally~~ (3) the continental crust of a microcontinent. The

160 eclogite, blueschist, amphibolite and greenschist of the Zermatt-Saas zone belong to the first of these. The metasediments of
the Combin zone represent the second position and ~~finally~~ the gneisses of the Dent-Blanche represent the position of the
deformed continental crust. However, the samples do not always represent the peak conditions of metamorphism but contain
minerals grown during various steps of the P-T path. In most samples presented here a clear chlorite-forming greenschist
overprint is visible, even if the rock originated at higher metamorphic conditions. This is especially true for the metabasic
165 rocks in this study, which often contain hornblende, actinolite and chlorite, indicating the amphibolite- and greenschist-facies
overprint, respectively.

4.1 Eclogite

The eclogite (MJS17) is composed of omphacite (30 vol.-%), garnet (30 vol.-%), clinozoisite (15 vol.-%), glaucophane (10
vol.-%), albite (10 vol.-%), and rare white mica (< 5 vol.-%), and retrograde chlorite (< 5 vol.-%) (table 1; figure 3a).
170 Mesoscopically the fabric is strongly dominated by a clear foliation defined by omphacite and glaucophane, surrounding
larger grains of garnet. Elongated omphacite and glaucophane grains form the stretching lineation and are visible in the
foliation plane. On the microscopic level the eclogite is strongly foliated with idiomorphic garnet grains surrounded by
layers of omphacite, glaucophane, clinozoisite and feldspar. The grain size distribution is bimodal with a mostly fine-grained
matrix of omphacite, glaucophane and some feldspar surrounding larger garnet grains. The presence of albite shows a
175 greenschist-facies overprint.

4.2 Blueschist

The blueschist (MJS16) comprises barroisite (40 vol.-%), albite (30 vol.-%), clinozoisite (15 vol.-%), garnet (10 vol.-%) and
epidote (5 vol.-%). The foliation in the blueschist is clearly pronounced on the mesoscopic scale (table 1; figure 3f). The
stretching lineation is formed by elongated barroisite and feldspar grains in the foliation plane. On the microscopic scale a
180 foliation formed by barroisite can clearly be seen. The grain-size distribution is bimodal, with a finer grained matrix of
amphibole and feldspar, and layers of larger, strongly retrogressed, inclusion-rich and fractured garnets therein.

4.3 Amphibolite

The composition of the amphibolite (MJS26; table 1; figure 3e) is dominated by hornblende (40 vol.-%). Feldspar (25 vol.-%),
clinozoisite (15 vol.-%), and chlorite (10 vol.-%) are also present in the matrix. Some large, fractured garnet (5 vol.-%)
185 grains are found as accessory minerals, which display strong greenschist-facies retrogression, with abundant chlorite
surrounding them. Hornblende defines a weak foliation in MJS26. Both hornblende and feldspar grains form a stretching
lineation on the foliation plane.

4.4 Greenschist

In the greenschist MJS36, the main constituent minerals are chlorite (30 vol.-%), clinozoisite (30 vol.-%), and actinolite (20 vol.-%). Feldspar is visible throughout the matrix (table 1, figure 3b). Actinolite defines the foliation of the sample, while chlorite grains show a weaker preferred orientation. In the foliation, amphibole grains can be seen forming a stretching lineation. The greenschist is fine-grained and shows a very homogeneous mineral distribution.

4.5 Metasediments

The composition of the metasediments (MJS18, MJS20, MJS22) is variable (table 1), but they all display a clear foliation and stretching lineation (figure 3c). In sample MJS18 the matrix consists of dolomite (60 vol.-%), quartz (20 vol.-%), calcite (10 vol.-%) and white mica (10 vol.-%). The foliation is pronounced and defined by dolomite and calcite grains. Layers of finer and coarser grained calcite, dolomite and quartz alternate. Mica fishes are found interlayered in the matrix. In the layers of coarser grained calcite, dolomite and quartz, a weak oblique foliation, in addition to the main foliation can be seen. The stretching lineation is visible on the foliation plane, in elongated calcite and dolomite grains. In the calc-micaschist (MJS20) a clear foliation is defined by white mica. Alternating layers of coarser and finer grained calcite (15 vol.-%), and quartz (40 vol.-%) are separated by bands of white mica (30 vol.-%) and chlorite (10 vol.-%). Larger grains of strongly fractured garnet (5 vol.-%) are present. These garnet grains give the fabric a knotty appearance, with the other minerals flowing around them. The calcschist MJS22 is made up of quartz (55 vol.-%), calcite (30 vol.-%) and chlorite (15 vol.-%). Compositional layering is clearly visible, with thick bands of very coarse-grained quartz, alternating with bands of fine-grained quartz and calcite with chlorite fishes. The coarser grains of both quartz and calcite form a weak oblique foliation.

4.6 Gneiss

The composition of the two gneisses (MJS34, MJS35) is quite similar with-as the main mineral components in both being-are quartz (40-57 vol.-%), feldspar (27-35 vol.-%), white mica (7-15 vol.-%), and chlorite (9-10 vol.-%) (table 1, figure 3d). The fabric of both samples is also very similar. They are strongly foliated with a lineation formed by quartz and feldspar. Sample MJS35 contains a 5 mm thick quartz band, which is a feature quite common in this unit in general. The other sample has a more homogeneous distribution of minerals. Microscopically the foliation is formed by mica, quartz, and feldspar. Mica shows a pronounced SPO of mineral-fishes-grains aligned in the foliation plane. Where chlorite is present it is interwoven with white mica between the quartz- and feldspar-rich layers.

5. Results

215 5.1 Crystallographic preferred orientations

5.1.1 Eclogite

In the eclogite (MJS17) omphacite shows a distinct CPO and has the highest texture strength of all phases in the sample with an F2 index (Bunge, 1982; see complete list in appendix A) of 3.49. The c-axes display the highest intensity clusters with a maximum parallel to the stretching lineation. The a-axes display maxima in a girdle perpendicular to the stretching lineation.

220 The omphacite b-axes form a weak girdle perpendicular to the stretching lineation with a clear maximum normal to the foliation plane (figure 4a). The glaucophane in eclogite has a weak CPO with an F2 index of 1.19. The c-axes display a maximum parallel to the stretching lineation. The a-axes develop a girdle perpendicular to the stretching lineation with a maximum paralleling the kinematic y-axis. The glaucophane b-axes display two maxima, the stronger of which ~~being is~~ normal to the foliation plane and a secondary maximum parallel to the stretching lineation (figure 4a). Clinzoisite has weak textures with an F2 index of 1.01. The c- axes display a maximum parallel~~ing to~~ the kinematic y-axis. The main b-axes maximum is parallel to the stretching lineation (figure 4a). The texture of white mica in the eclogite is quite weak (F2 index of 1.02). However, the basal planes show an alignment to the foliation plane (figure 4a). Garnet and albite show random textures.

5.1.2 Blueschist

230 In the blueschist (MJS16), barroisite displays the strongest textures with an F2~~-~~index of 2.36. The strongest textures are seen in the c-axes, which exhibit a maximum parallel to the stretching lineation. The a-axes in barroisite display ~~maxima in a~~ girdle distribution perpendicular to the lineation direction with a maximum parallel to the kinematic y-axis. The b-axes display maxima in a similar girdle perpendicular to the stretching lineation with highest densities normal to the foliation plane (figure 4b). Clinzoisite and epidote CPOs are very similar and relatively weak. Both show an F2 index of 1.02 and b-axes with a maximum almost parallel to the stretching lineation. A-axes and c-axes show girdles perpendicular to the lineation (figure 4b). Garnet and albite exhibit random textures.

5.1.3 Amphibolite

240 In the amphibolite MJS26, the strongest textures are found in the hornblende, with an F2 index of 1.62. The hornblende c-axes show the strongest CPO with a distinct maximum parallel to the stretching lineation. The b-axes are distributed in a girdle normal to the stretching lineation, with a maximum normal to the foliation plane. The a-axes also display a girdle normal to the stretching lineation (figure 5a). For chlorite, the F2 index is 1.10. Chlorite c-axes show a girdle perpendicular to the lineation with a maximum parallel to the foliation normal (figure 5a). In albite the texture strength is low, with an F2 index of 1.07. However, a-axes display a girdle perpendicular to the stretching lineation and a weak maximum normal to the

foliation plane while b-axes display a broad, weak maximum parallel to the stretching lineation (figure 5a). The textures in
245 garnet and clinozoisite are nearly random.

5.1.4 Greenschist

In the greenschist MJS36, the strongest textures are found in chlorite with an F2 index of 1.34. Chlorite basal planes are
oriented parallel to the foliation plane (figure 5b). The textures of actinolite are of intermediate strength with an F2 index of
1.21. The c-axes display a strong maximum parallel to the stretching lineation and a weak girdle distribution in the foliation
250 plane. The a-axes show highest intensities with a maximum normal to the foliation plane. The b-axes display a slightly
weaker texture with a maximum parallel to the foliation normal, connected by a weak, irregular girdle normal to the
stretching lineation. Furthermore, secondary maxima are slightly oblique to the stretching lineation (figure 5b). Albite has an
F2 index of 1.18 and a-axes display the strongest CPO with a maximum parallel to the foliation normal. The b-axes are
concentrated in a girdle in the foliation plane with weak maxima therein. The c-axes display a weak CPO with diffuse
255 maxima (figure 5b). Clinozoisite shows a random CPO.

5.1.5 Metasediments

In the metasediment samples (MJS18, MJS20, MJS22) the CPOs are quite distinct and often stronger than in the samples
described above.

In the dolomitic micaschist MJS18, the textures of muscovite are strongest with an F2 index of 2.18. Its basal planes display
260 a strong alignment with the foliation plane (figure 6a). The textures of dolomite are of intermediate strength with an F2 index
of 1.14. The c-axes display a maximum normal to the foliation plane. The a-axes show a girdle distribution in the foliation
plane with a maximum parallel to the stretching lineation (figure 6a). In calcite the textures are slightly stronger than in
dolomite and have an F2 index of 1.28. The c-axes display a strong maximum perpendicular to the foliation plane and a-axes
display a girdle distribution in the foliation plane (figure 6a). The CPO of quartz is weak with an F2 index of 1.04. The c-
265 axes display small girdles around the foliation normal. A-axes show a weak diffuse maximum oblique to the stretching
lineation. Prism and rhomb planes display almost random distributions (figure 6a).

In the calcareous micaschist (MJS20), muscovite and chlorite show strong textures with F2 indices of 2.83 and 2.17,
respectively. Both minerals yield a strong alignment of their basal planes in the foliation (figure 6b). Calcite textures are
weak, with an F2 index of 1.08. The c-axes display a maximum normal to the foliation and the a-axes are concentrated in a
270 girdle in the foliation plane (figure 6b). The texture of quartz is of intermediate strength with an F2 index of 1.14. The c-
axes display small circle distributions around the foliation normal, while the a-axes display two maxima at the periphery,
each at low angle to the stretching lineation. [01-11] shows one maximum normal to the foliation and a second set of
maxima in the foliation plane. [10-10] displays a diffuse maximum nearly parallel to the stretching lineation, while [10-11]
displays maxima clustered around the foliation normal and in the foliation plane parallel to the y-direction. Garnet shows a
275 random CPO.

In the calc schist MJS22 all mineral phases display medium texture strength (appendix A). In calcite the F2 index is 1.34. The c-axes display a strong maximum parallel to the foliation normal. A-axes are concentrated in the foliation plane with a maximum parallel to the stretching lineation (figure 6c). In quartz the textures are stronger than in calcite, with an F2 index of 1.54. The quartz c-axes display a weak sinistrally rotated girdle with a strong maximum offset from the kinematic y-axis by ca. 30°. A weaker density cluster is found perpendicular to this girdle. The a-axis maximum is located close to the periphery of the pole figure slightly sinistrally rotated away from the x-direction. The poles of the prism and rhomb planes display weaker maxima in girdles paralleling the c-axes girdle (figure 6c). Chlorite has an F2 index of 1.52 and displays a strong alignment of its basal plane in the foliation (figure 6c).

5.1.6 Gneiss

In the gneisses (MJS34, MJS35) CPOs are similar to those of the metasediments (figures 6 and 7).

In MJS34 the muscovite and chlorite display basal planes parallel to the foliation plane. The textures are strongest for chlorite with an F2 index of 2.15. Muscovite has an F2 index of 1.27. Quartz shows an F2 index of 1.18 and its c-axes are distributed in small circles around the foliation normal. The a-axes display a weak and diffuse girdle distribution in the foliation plane. In albite a very weak (F2 index of 1.03) but distinct CPO can be seen. The a-axes display a maximum parallel to the stretching lineation (figure 7a).

In MJS35 both muscovite and chlorite display clear alignment of the basal planes in the foliation plane. The F2 indices of these two minerals are 1.13 and 1.53, respectively. The CPOs in quartz are weaker (F2 index of 1.06) than those in MJS34. C-axes display an asymmetric crossed girdle distribution. The a-axes display diffuse intensity clusters at the periphery around the stretching lineation. In albite the CPO is nearly random (F2 index = 1.01), with a faint maximum of the a-axes in the lineation direction (figure 7b).

5.2 Elastic anisotropy properties

P-wave anisotropies for the various lithologies were calculated from the ODFs of constituent mineral phases, single-crystal data of the respective phase and their volume fraction within the sample using the software BEARTEX (Wenk et al., 1998). Graphical representations of S-wave velocity distributions are not shown in this study, since the differences between maximum and minimum velocities in a sample are on the order of 0.01 km/s, relatively small and deemed insignificant.

5.2.1 Eclogite

In the eclogite (MJS17) the average P-wave velocity (VP) is 8.2 km/s and the average S-wave velocity (VS) is 4.8 km/s resulting in a VP/VS-ratio of 1.72. The elastic anisotropy AVP is 1.4%, which is defined by omphacite (30 vol-%; table 1) and glaucophane (10 vol-%; table 1). The fast velocities correlate with the c-axis maximum of omphacite as well as of glaucophane parallel to the stretching lineation. The lowest P-wave velocities are found parallel to the foliation normal and intermediate velocities can be found parallel to the kinematic y-axis (figures 4a and 8a).

5.2.2 Blueschist

In the blueschist the average VP is 7.6 km/s, with a maximum of 7.8 km/s and minimum of 7.5 km/s. The average VS is 4.5 km/s, resulting in a VP/VS-ratio of 1.71. The P-wave anisotropy is of medium strength with 3.7 % and defined by barroisite (40 vol.-%, table 1) and clinozoisite (15 vol.-%, table 1). The distribution of the fast velocities matches the orientation of the barroisite c-axes and the clinozoisite b-axes. Lowest P-wave velocities can be found in a girdle distribution perpendicular to the stretching lineation (figures 4b and 8b).

5.2.3 Amphibolite

The amphibolite has an elastic anisotropy ΔVP of 3-4.2.8 %. The VP range from 7.0-14 km/s to 7.34 km/s, with an average of 7.4-20 km/s. The average VS is 4.0-11 km/s, resulting in a VP/VS-ratio of 1.7675. Maximum VP are distributed is aligned parallel to the stretching lineation, with a girdle of low velocities in the kinematic y-z-plane, perpendicular to the stretching lineation (figure 8c). The maximum velocity correlates well with the c-axes of hornblende (40 vol.-%, table 1) in the amphibolite (figures 5a and 8c).

5.2.4 Greenschist

In the greenschist MJS36, the average VP is 7.30 km/s, with a maximum of 7.2-43 km/s and a minimum of 6.87.16 km/s. The average VS is 4.1-28 km/s, resulting in a VP/VS-ratio of 1.7370. The P-wave anisotropy is high with a value of 5-3.7 %. The highest VP are found in the foliation plane, with a maximum parallel to the stretching lineation. Lowest VP are found parallel to the foliation normal (figure 8d). The highest velocities are parallel to the actinolite (20 vol.-%, table 1) c-axes and distributed similarly to the chlorite (30 vol.-%, table 1) basal planes (figures 5b and 8d).

5.2.5 Metasediments

The metasedimentary schist samples (MJS18, MJS20 and MJS22) show an average VP of 6.566-7.23 km/s and an average VS of 4.012-4.1-16 km/s. The P-wave anisotropies range from 4.73.7-8.27.1 % and VP/VS-ratios average at 1.66 range from 1.61-1.76 (figure 8e-g).

In the dolomitic micaschist (MJS18) the average VP of 7.2 km/s is high for a metasedimentary sample. The VP anisotropy ΔVP is 6.7%, the highest determined in the metasediments. The average VS is 4.1 km/s, resulting in a VP/VS-ratio of 1.76. Maximum VP is parallel to the stretching lineation lowest VP is found parallel to the foliation normal (figure 8e).

In the calcareous micaschist (MJS20) the average velocities are lower than in the previously described sample. The average VP is 6.77 km/s and the average VS is 4.16 km/s, resulting in a VP/VS-ratio of 1.63. The P-wave anisotropy in the calcareous micaschist is very strong the highest determined in the metasediments at 8.27.1%. Highest VP is concentrated in the foliation plane, lowest VP is found parallel to the foliation normal (figure 8f).

In the calcschist (MJS22) the average VP is ~~6.5-66~~ km/s with an ~~anisotropy ΔVP~~ of ~~4.73.7~~%. The average VS is ~~4.0-15~~ km/s, resulting in a VP/VS-ratio of ~~1.6061~~. ~~Highest-The highest~~ P-wave velocities are concentrated in a broad girdle parallel to the foliation plane with lowest P-wave velocities perpendicular to it (figures ~~6c~~ and ~~8g~~).

340 5.2.6 Gneiss

In the gneiss samples (MJS34, MJS35) the P-wave anisotropies ~~are relatively low~~ and ~~the~~ velocity-distributions ~~are quite similar~~ ~~show highest VP in the foliation plane~~ (figure ~~8h-i~~).

In MJS34 the average VP is ~~6.5-62~~ km/s and the average VS is ~~4.14~~ km/s resulting in a VP/VS-ratio of 1.6. The P-wave anisotropy of ~~2.81~~% is low, and the velocity-distribution displays fast velocities in the foliation plane and a minimum velocity perpendicular to it (figure ~~8h~~).

In MJS35 the average VP is slightly lower than in MJS34 at ~~6.4-52~~ km/s. The average VS is ~~4.18~~ km/s, resulting in a VP/VS-ratio of 1.56. The ~~P-wave anisotropy ΔVP~~ is ~~1.84~~%. Similar to MJS34, the highest velocities are found in the foliation plane. However, ~~instead of the continuous girdle of higher VP in the foliation plane, several distinct maxima occur in the foliation plane, connected by a weak girdle a weak maximum is visible almost parallel to the stretching lineation~~ (figure ~~8i~~).

350 6. Discussion

The elastic anisotropy of rocks is influenced by several factors, ~~among which are its including~~ mineral composition, ~~its~~ grain fabric, possible microcracks, and the CPO of its constituent minerals. The CPO ~~constrains-contributes~~ a large part of the anisotropy and mostly results from ~~ductile~~ rock deformation. According to Mainprice and Ildefonse (2009) only mineral phases which make up more than 10% of the sample, have an important influence on the elastic properties of the rock in question. In our calculations we have, however, included all mineral phases down to a volume fraction of 5 %.

6.1 Metabasic rocks

6.1.1 CPO development

In the metabasic rocks of eclogite, blueschist, amphibolite and greenschist facies metamorphic conditions the compositions are dominated by pyroxenes (omphacite), amphiboles (glaucophane, barroisite, hornblende and actinolite) and chlorite (table 1).

The CPOs of omphacite, glaucophane, barroisite, ~~actinolite, chlorite~~ and hornblende have a strong effect on the ~~elastic seismic~~ anisotropy of the ~~eclogite, the blueschist and the amphibolite~~ ~~metabasic rocks~~. In previous studies, the CPO geometry of omphacite has been interpreted as being the result of corresponding strain geometry (Helmstaedt et al., 1972; Kurz et al., 2004; Neufeld et al., 2008; Müller et al., 2011; Keppler et al., 2016; ~~Zhang, et al. 2007~~). L-type (lineation dominated) fabrics are interpreted as the result of constrictional strain, S-type (foliation dominated) fabrics as the result of flattening strain and the transitional fabrics as indicators of plain strain deformation (e.g. Helmstaedt et al., 1972; ~~Zhang, et al. 2007~~). SL- and

LS-type fabrics have been described by Godard and Van Roermund (1995). Omphacite in MJS17 displays an L- to LS-type fabric, with c-axes ~~paralleling-parallel to~~ the stretching lineation and a- and b-axes ~~paralleling-parallel to~~ the kinematic y- and z-axes, respectively (figure 4). Keppler (2018) suggests that this correlation of strain geometry to CPO geometry found in omphacite is applicable for amphiboles as well. The amphibole CPO in three of the four metabasites (eclogite, blueschist, and amphibolite; figure 4 and 5) can therefore also be interpreted as L-type or LS-type fabrics, since the c-axes are aligned in lineation direction, while a-axes and b-axes are closer to a girdle distribution perpendicular to the lineation. In the greenschist the actinolite c-axes display a maximum ~~paralleling-parallel to~~ the stretching lineation with a weak girdle extending into the foliation plane, while the a-axes display a maximum perpendicular to the foliation plane (figure 5), which points to a transitional SL-type fabric.

Since the omphacite, glaucophane and barroisite c-axes are parallel to both the stretching lineation and one another, the assumption can be made that they formed in the same prograde deformation event. A further possibility would be retrograde mimetic overgrowth of omphacite by glaucophane (McNamara et al., 2012). There ~~is-are~~ considerable debates on timing of CPO-formation in eclogites, with proponents of formation during subduction, during exhumation, or both, including concurrent switches in strain geometry (Zulauf 1997; Kurz et al., 2004; Kurz, 2005; Neufeld et al., 2008; Müller et al., 2011; Keppler et al., 2016; Keppler, 2018). The CPOs of clinozoisite and epidote frequently have similarities to those of omphacite and the amphiboles. The clinozoisite b-axes display a maximum paralleling the stretching lineation, matching the omphacite and amphibole c-axes. Keppler et al. (2017) suggest concordant CPOs due to opposing crystallographic axis definitions and further conclude that clinozoisite CPOs can likewise be classified in S-type, L-type and intermediate fabrics (Keppler et al. 2018; Puelles et al. 2017). In this case the CPOs of clinozoisite would suggest L-type fabrics hinting at constrictional strain and matching those already described for omphacite and the amphiboles. Albite preferentially displays weak CPOs in the metabasites. However, in the amphibolite and greenschist the weak albite texture of the b- and c-axes corresponds to the stretching lineation, while the a-axis maximum is aligned nearly normal to the foliation plane (figure 5). The slip-systems in plagioclase are highly complex and a correlation with the strain geometry is difficult to infer (Hacker and Christie, 1990; Prior and Wheeler, 1999). When chlorite or white mica are present, they mostly display a strong alignment of the basal planes in the foliation plane. The similarities in the CPOs of chlorite, amphibole and albite in the greenschist suggest a formation during the same deformation, which likely took place during ~~the retrograde~~ exhumation.

6.1.2 Elastic ~~anisotropies~~ properties

The ~~elastic-seismic~~ anisotropies and the VP-distribution in the metabasites are primarily controlled by the influences of omphacite, amphibole and chlorite. These minerals result primarily in low to intermediate AVP (1.4-3.7 %) and VP-distributions with highest VP parallel to the stretching lineation. In the greenschist, the influence of chlorite, is significant, resulting in a girdle-type distribution of higher VP in the foliation plane and a weaker maximum parallel to the stretching lineation.

400 ~~in two variations of elastic anisotropies and VP distributions. On the one hand there is a group of results with lower anisotropies (eclogite: 1.4 %; blueschist 3.7%; amphibolite: 3.1 %) and highest VP paralleling the lineation direction and on the other hand there is the greenschist, with a higher anisotropy (5%) and VP distributed in a girdle in the foliation plane.~~

The compositions, ~~anisotropies-AVP and VP-distributions of the first group~~ are primarily controlled by omphacite and amphibole and secondarily by chlorite, clinozoisite, epidote and garnet. In the amphiboles, highest P-wave velocities in the single-crystals correlate almost perfectly with the c-axes, only deviating by a few degrees. Since the c-axes of the

405 amphiboles in these metabasites parallel the stretching lineation, this results in a velocity maximum in lineation direction. The influence of omphacite is slightly more complicated, since in the omphacite single-crystals the highest velocity is not parallel to the c-axis and instead can be found between the c- and a-axes. This discrepancy has little influence in this case since it also results in a velocity maximum parallel to the stretching lineation, but it should be kept in mind when considering anisotropy strength and distribution. Epidote and clinozoisite CPOs in the blueschist likely contribute to the same pattern of

410 high velocities in lineation direction. Even though garnet has a near random CPO and does not contribute directly to the anisotropy, the high garnet content in the eclogite sample (30 vol.-%, table 1) and the very high ~~isotropic~~ P-wave velocities in garnet single-crystals increase the average seismic velocities in MJS17. In the blueschist and the amphibolite in which the garnet is only present to 5-10% this effect is negligible. Albite can also influence the bulk anisotropy of the metabasites due to its strong single-crystal velocity anisotropy (Brown et al., 2006; Brown et al., 2016). The highest velocities in albite are

415 parallel to the b- and c-axes (Brown et al., 2006), both of which display weak maxima parallel to the stretching lineation in the metabasites, further contributing to the high velocities parallel to the stretching lineation (figure 4 and 5). In the amphibolite, the chlorite basal planes display a girdle distribution perpendicular to the stretching lineation with a maximum perpendicular to the foliation (figure 5). This CPO reflects a rotation of the chlorite basal planes around an axis ~~paralleling parallel to~~ the stretching lineation. Due to the lowest velocities in sheet silicates normal to the basal plane, the CPO of chlorite in this sample contributes to the low velocity girdle perpendicular to the stretching lineation. The eclogite contains white mica ~~-, however~~ However, due to its very low volume fraction of 5 % it is considered insignificant for the bulk anisotropy (Mainprice and Ildefonse, 2009).

In the greenschist, ~~the influence of~~ chlorite dominates the composition ~~and~~ However the ~~anisotropies-anisotropy~~ as well as the VP-distribution are primarily controlled by actinolite in the sample. Actinolite CPOs in the greenschist display a

425 transitional SL-type fabric, resulting in highest VP parallel to the stretching lineation and contributing to the weak girdle of slightly higher VP in the foliation plane. In sheet silicates the highest velocities in the single crystals are distributed parallel to the basal planes. The strong alignment of these basal planes in the foliation plane results in ~~high elastic anisotropy and highest velocity-higher VP~~ in the foliation plane ~~very similar but far weaker than to that the pattern~~ observed in the metasediments (figures 5, 6 and 8 d-g). Albite in the greenschist also contributes to this pattern due to the maxima of b- axes and c-axes within the foliation plane. ~~Actinolite CPOs in the greenschist display a transitional SL-type fabric and similarly contribute to the high VP distribution in the foliation plane.~~

430

When comparing the ~~elastic anisotropy~~AVP results to anisotropy data from literature (table 3) we find a very wide range of anisotropy values for these lithologies. For the eclogite our data (1.4 %) fits well inside the bounds of the texture derived AVP data of 0.6% to 5.0% by Llana-Fúnez and Brown (2012), 0.4-3.7 % by Keppler et al. (2015), 2.9% by Bezacier et al. (2010), 1.2-2.9 % by Bascou et al. (2001), 1.1-2.8 % by Worthington et al. (2013), 1.4-10.2 % by Cao et al. (2013) and 0.99-2.46 % by Cao et al. (2020), ~~and~~In the case of experimentally derived results, our data fits well into the range of 0.6-1.7 % reported by Wang et al. (2009), while it is significantly lower than the results of 7-11 % reported in Zertani et al. (2019) or the value of 8.3 % from Keppler et al. (2015). In a study by Cao et al. (2013) P wave anisotropies of 1.4-10.2 % were calculated from CPOs into which our value of 1.4% would also fit. Similarly, our data from the amphibolite (A: ~~3-12.8~~ %) is ~~good~~in good agreement with texture derived data from previous studies on amphibolites. Ivankina et al. (2005) report AVP of 0.59-5.51 %, Ullemeyer et al. (2006) of 4.4 %, and Keppler et al. (2015) of 2.6 % and Keppler et al. (submitted to same issue) AVP of 2.1-4.5 %. As with the eclogite our texture derived data is lower than the experimental AVP reported by Siegesmund et al. (1989) ~~report an elastic anisotropy of with~~ 6.4 %, Barruol and Kern, (1996) with 5.69-12.74 % and Kästner et al. (2020) with 7.75-10.90 %. However the studies by Ivankina et al. (2005) and Keppler et al. (2015) also report lower experimental AVP of 0.86-8.27 % and 4.4 %, respectively. These are quite similar to the data of our study. Ivankina et al. (2005) of 0.59-5.51 %, Ullemeyer et al. (2006) of 4.4 % and Keppler et al. (2015) of 2.6 %. In the case of the blueschist, our data (AVP: 3.7%), fit very well into the range of texture derived data by Fujimoto et al. (2010) with AVP of 1.7-6.2 %, however~~However, the elastic anisotropy (A: 3.7%) our value~~ is significantly lower than that presented in other studies. Bezacier et al. (2010) report texture derived a P wave anisotropyAVP of 16.1 % for a blueschist, while Cao et al. (2013) ~~report even higher elastic anisotropies of 13.4-24.6 % and Kim et al. (2013) of 7.1-29.5 %.~~ There are several possible explanations for the discrepancy to the three later studies. The first of these is that in many studies, blueschist samples are collected from highly deformed bands resulting in stronger anisotropies. In our study, however, a sample with a fabric that is representative for the lithology within the Alps was selected, resulting in weaker overall mineral CPOs and a lower anisotropy. Furthermore, in the two studies mentioned the amphibole in the blueschist was glaucophane, while in the case of the sample in this study it is mostly barroisite. This was addressed in part when calculating the elastic-seismic properties of the blueschist since no single-crystal data for barroisite were available so the single-crystal data of glaucophane (Bezacier et al. 2010) were selected as a suitable substitute. However, glaucophane may still react differently to deformation than barroisite, possibly resulting in stronger CPOs leading to higher elastic-seismic anisotropies. Finally, few studies are available on the elastic-seismic anisotropies anisotropy of greenschists. Due to the high chlorite-content (30 vol.-%; table 1) which contains a lot of water, measurements by neutron diffraction are very difficult. Measurements by synchrotron diffraction are possible, but limited to very fine grain sizes. However with an AVP of 3.7 % and an actinolite content of 20 % in the greenschist of this study (figure 8d), it can be compared well to the amphibolite MJS26. ~~When comparing the elastic anisotropy of 5 % in the greenschist of this study, it can be compared to other lithologies containing such high sheet silicate fractions like the metasediments and gneisses mentioned below (figure 8).~~

465 The VP/VS-ratios in the metabasites range from 1.70 in the greenschist, 1.71 in the blueschist, over 1.72 in the eclogite, ~~4.73~~
~~in the greenschist~~ and up to 1.76-75 in the amphibolite. These VP/VS-ratios are the direct result of the VP/VS-ratios of their
constituent minerals and their respective volume fractions (table 1). The lower ratios of 1.69 in the glaucophane are balanced
out by the slightly higher ratios of pyroxene, garnet, clinozoisite, epidote, chlorite, actinolite and albite whose ratios range
between 1.73-1.745. The ratio in the amphibolite is particularly high due to the high hornblende content of 40 vol.-% (table
470 1) with its very high VP/VS-ratio of 1.84.

6.2 Metasediments

6.2.1 CPO development

The small circles of quartz c-axes around the foliation normals of samples MJS18 and MJS20 might indicate flattening strain
(Lister and Hobbs, 1980; Schmid and Casey, 1986) (figure 6). This is also supported by the distribution of quartz a-axes, but
475 the textures are quite weak. The c-axis pole figures of MJS18 and MJS20 are indicative of basal <a> slip and rhomb <a> slip
in quartz (cf. Schmid and Casey, 1986) (figure 6). In MJS22 the quartz c-axes display an inclined single girdle pointing to
simple shear. This distribution is also reflected by the quartz a-axes, prism and rhomb planes and indicative of multi-slip
activity, possibly a combination of rhomb <a> slip, prism <a> slip and perhaps some basal <a> slip (e.g., Schmid and Casey,
1986; Stipp et al., 2002) (figure 6). In all metasediments the basal planes of white mica and chlorite display a strong
480 alignment in the foliation plane. In all three metasedimentary samples the CPOs of calcite c- and a-axes exhibit axial
symmetry with respect to the foliation normal as a symmetry axis (figure 6). This symmetry in calcite has first been
interpreted as indicative of pure shear deformation (Wenk et al., 1987; Kern and Wenk, 1983). However, more recently,
there was considerable debate on this topic (De Bresser, 1989; Ratschbacher et al., 1991; Erskine et al., 1993; Burlini et al.,
1998; Leiss et al., 1999; Bestmann et al., 2000). The maximum of the calcite a-axes can either be parallel to the stretching
485 lineation or to the kinematic y-axis or even display multiple maxima between the two (Ratschbacher et al., 1991; Punturo et
al., 2005; Bestmann et al., 2000; Bestmann et al., 2006; Trullenque et al., 2006). ~~Very f~~Few studies have focused on
dolomite CPOs. Delle Piane et al. (2009) show that dolomite CPOs very closely reflect those of calcite, but deformation
itself strongly influences the texture development when both phases deform together. Dolomite in MJS18 displays very
similar CPOs to calcite, with the same axial symmetry in the c- and a-axes (figure 6).

490 6.2.2 Elastic anisotropiesproperties

The ~~elastic anisotropies~~AVP of the metasediments (MJS18, MJS20 and MJS22) range from 43.7-8.27.1 % and are among
the highest presented in this study (figure 8e-g). These anisotropies directly correlate to the pronounced foliation and high
mica and chlorite contents found in the metasediments, as well as the strong influence of calcite and dolomite CPOs on the
samples. This corresponds well to texture derived elastic anisotropyAVP data from similar metasediments reported in other
495 studies (see overview in table 3) such as Weiss et al. (1999) with 4.56-6.53 %, Erdman et al. (2013) with 2.3-11.4 %, ~~and~~

Keppler et al. (2015) with 5.1-7.4 % and Keppler et al. (submitted to same issue) 7.9 %. Our data fall into the range of experimental AVP reported by Kästner et al. (2020) with its AVP of 2.95-26.41%.

In our study, the orientation of white mica and chlorite (10-40 vol.-%) basal planes and the very high single-crystal anisotropy of white mica, which is used for both sheet silicates, mainly contribute to the ~~elastic-seismic~~ anisotropy in the metasediments. The VP distribution in muscovite single crystals is dominated by ~~its-the~~ highest velocity parallel to and the lowest velocity normal to the basal plane (figures 6 and 8e-g). As micas preferentially align with increasing strain in the foliation, the result is a strong tendency for high P-wave velocities in the foliation plane in deformed mica-rich rocks.

The composition of the metasediments varies substantially. Calcite and dolomite combined range from 15 vol.-% up to 70 vol.-% (table 1). The CPOs of both ~~of these~~ minerals strongly influence the P-wave anisotropy of the metasediments. They result in high velocities in the foliation plane with a maximum parallel to the stretching lineation. Further they contribute to the velocity minima normal to the foliation plane. This is the result of the calcite single crystal data in which the c- and a-axes are the slow and fast directions, respectively (Punero et al., 2005). The single-crystal anisotropy of dolomite is quite similar to that of calcite. Hence, both similarly contribute to the bulk anisotropy of sample MJS18. The contribution of quartz to the P-wave anisotropy is weak due to the nearly random CPO observed for the poles of the rhombs, which represent the fastest direction in the quartz crystal. Furthermore, the three-fold symmetry of the quartz single crystal makes any influence on the velocity distribution complicated to assess. The garnet present in MJS20 displays a random texture and therewith has no significant influence on the ~~elastic-seismic~~ anisotropy.

The VP/VS-ratios in the metasediments range from 1.61 to 1.76~~1.60 to 1.76~~. The highest ratio is found in sample MJS18 at 1.76, while the ratios of the other two samples range from 1.60~~1.61~~-1.63. The high ratio of MJS18 is the result of the very high dolomite content of 60 vol.-% and the 10 vol.-% of calcite in this sample. ~~Both-d~~olomite and calcite have very high single crystal VP/VS-ratios of 1.851 and 1.918~~922~~, respectively. ~~These-They~~ are significantly higher than those of quartz (20 vol.-%) and white mica (vol.-10%) at 1.476 and 1.726, ~~respectively~~. For the other two samples the velocity ratio is dominated by quartz (40-55 vol.-%), resulting in their low ratios of 1.61 to 1.63.

6.3 Gneiss

6.3.1 CPO development

The small circles around the foliation normal, visible in the quartz c-axes and the distribution of a-axes of MJS34 indicate flattening strain with combined activity of rhomb <a> slip and basal <a> slip (Lister and Hobbs, 1980; Schmid and Casey, 1986) (figure 7). In MJS35, the type I crossed girdle in the quartz c-axes is indicative of plane strain with combined basal <a>, rhomb <a> slip and prism <a> slip (e.g., Schmid and Casey, 1986). White mica and chlorite CPO in both MJS34 and MJS35 bear a strong relation to the foliation plane (figure 7). Albite in MJS34 and MJS35 shows very weak CPOs, but albite a-axes in both display a weak but visible maximum paralleling-parallel to the stretching lineation (figure 7). Due to the

random distribution of the b-axes and c-axes a clear assignment cannot be made, however this CPO pattern is closest to an axial A type, indicating that the activated plagioclase slip system is [100](010) and/or [100](001) (Satsukawa et al., 2013).

6.3.2 Elastic ~~anisotropies~~ properties

530 The ~~elastic anisotropy AVP~~ in the gneisses MJS34 ~~and MJS35~~ is primarily constrained by the CPOs of mica and chlorite (25 vol.-% ~~and 15 vol.-% respectively~~, table 1) ~~while in the gneiss and MJS35 secondarily by it is primarily constrained by quartz and albite. However, the influence of quartz is far weaker than that of the sheet silicates as the pole figure maxima of quartz do not match well to the bulk elastic anisotropy diagram. In MJS34~~ The strong single-crystal anisotropy of muscovite and chlorite in combination with their strong CPO, results in the high velocities found in the foliation plane and the
535 corresponding perpendicular velocity minima (figures 7 and 8h). In MJS35, the effect of the sheet silicates is far weaker and only results in a weak girdle of higher VP in the foliation plane. However, the three-fold symmetry of quartz likely results in the distinct VP maxima visible in the foliation plane. ~~The VP/VS-ratio of the gneiss samples ranges from 1.56 to 1.60. As in the metasediments this low ratio is the result of the very low ratio of 1.48 in quartz (40 vol.-% and 60 vol.-% respectively) and the intermediate ratios in albite (1.73) and the sheet silicates (1.73 and 1.72; 25 vol.-%) in the samples (appendix 2;~~
540 ~~table 1). The velocity distribution is very similar to that of the metasediments, with higher velocities in the foliation plane and low velocities perpendicular to this plane (figure 8h-i), but the elastic anisotropy AVP in both samples is significantly lower. When comparing the results of our study to those previously published (table 3), we see a very wide range of P-wave anisotropy data by different methods reported in gneisses. Values-Texture derived AVP range from low to intermediate anisotropies-values such as 1.76-8.73 % by Weiss et al. (1999), 1.72-2.99 % by Ivankina et al. (2005), 0.9-3.2 % by Ullemeyer et al. (2006), and 7.9 % by Kern et al. (2008), up to high to very high anisotropies-values such as 4.1-20.1 % by Erdman et al. (2013), ~~and~~ 8.0-11.3 % by Llana-Fúnez and Brown (2012) and 1.3-20.5 % by Keppler et al. (submitted to same issue). Experimental studies equally display a very wide range of AVP for gneisses. AVP of 2.71-2.74 % by Ivankina et al. (2005), 15.1 by Kern et al. (2008), 4.7-13.6 % by Khazanehdari et al. (2000) 9.6 % by Vasin et al. (2017) and 3.63-14.56 % by Kästner et al. (2020) are reported. ~~As gneisses predominantly display weak CPOs similar to our results, our low elastic anisotropy AVP values are common for gneisses.~~~~

550 elastic anisotropy AVP values are common for gneisses.

6.4 Elastic anisotropy data in the context of a subduction and exhumation setting

In this study we examine the elastic seismic properties of a wide array of lithologies in a subduction and exhumation setting. As previously mentioned and is visible in figure 2, we have divided this setting into several structural positions. These are the oceanic crust of the subducting plate undergoing different metamorphic facies, the sedimentary cover of the subducting
555 plate, which is at least partly scraped off and transferred into the accretionary prism and finally the deformed continental crust. This continental crust can be part of a microcontinent (as in the case of the Dent Blanche unit) or can be the continental crust on the opposite side of the ocean, which enters the subduction zone subsequently to the oceanic crust leading to continental collision. This scenario is presented in the schematic cross-section in figure 2. The structural positions

560 vary in a range of criteria such as degree of deformation and mineral composition. The different degrees of deformation and metamorphism lead to different texture strengths and variable mineral content. When comparing the different structural positions, samples can be related in order of SiO₂-content, with quartz- and feldspar-rich felsic rocks on one end of the spectrum, and the olivine-rich ultramafic rocks on the opposite end of the spectrum. This has been discussed extensively ~~in a very insightful study~~ by Almqvist and Mainprice (2017) for the earth's crust. Generally, a trend of lower velocities in the felsic and higher velocities in the mafic to ultramafic rocks can be observed. In this study, we further discuss another important criterion ~~of great importance on for~~ the ~~elastic-seismic~~ properties, which is the influence of CPO of the constituent minerals on the ~~elastic-seismic~~ anisotropy. This might differ from the compositional effect in the spectrum from felsic, over mafic to ultramafic. Concerning ~~elastic-seismic anisotropies~~ anisotropy, the sheet-silicate content and its alignment caused by deformation show a pronounced influence.

570 During the prograde metamorphism in subduction zones the basaltic oceanic crust undergoes blueschist-facies metamorphism leading to the formation of glaucophane (in some cases barroisite), epidote, omphacite and some accessory minerals. Glaucophane and feldspar usually dominate the composition, of which glaucophane greatly influences the anisotropy and the velocity distribution. Blueschists display strongly varying ~~elastic-anisotropies~~AVP of ~~31.7-24.6~~ % (Cao et al., 2013; Bezacier et al., 2010; Kim et al., 2013; Fujimoto et al., 2010). The deformation during subduction leads to CPO formation in amphibole. In most cases plane strain is dominant. The orientation of the c-axes of amphibole parallel to the stretching lineation results in the highest velocity parallel to the lineation. Dominant constriction in the blueschist of the present study produces a similar pattern (see figures 4 and 8b). At greater depth the oceanic crust reaches eclogite facies metamorphic conditions. There, omphacite and garnet are the major rock-forming minerals (table 1) and omphacite has the strongest ~~influence on contribution to~~ the ~~elastic-seismic anisotropies~~ anisotropy. The ~~elastic-anisotropies~~AVP in eclogites are lower than those in the blueschists and display values of 0.4-~~10.211~~ %, due to the higher ~~elastic-anisotropy~~AVP of amphiboles compared to pyroxenes (Llana-Fúnez and Brown, 2012; Keppler et al., 2015; Cao et al., 2013; Bezacier et al., 2010; Wang et al., 2009; Bascou et al., 2001; Zertani et al., 2019; Worthington et al., 2013; Cao et al. 2020). The highest P-wave velocities remain parallel to the stretching lineation, ~~due to because~~ the omphacite CPO ~~yielding-yields~~ an alignment of [001] in lineation direction during plane strain ~~as well as and~~ constriction (see figures 4 and 8b). Especially strongly deformed eclogites could be detected by seismic imaging due to their composition and their CPO-related specific seismic properties (e.g. Boundy et al. 1992; Fountain et al. 1994).

585 During exhumation the high pressure rocks frequently experience retrogression as they pass through the amphibolite facies, where they are overprinted or completely retrogressed. In the amphibolite-facies rocks, hornblende together with feldspar and epidote dominates the composition. These minerals result in anisotropies of 0.59-~~6.412.74~~ % (Siegesmund et al., 1989; Ivankina et al., 2005; Ullemeyer et al., 2006; Keppler et al., 2015; Barruol and Kern, 1996; Kästner et al., 2020; Keppler et al., submitted to same issue), of which the amphiboles have the strongest influence. The velocity distribution in the amphibolites is very similar to those observed in the blueschists and eclogites, with highest velocities found parallel to the stretching lineation (figure 8c). Further along the exhumation path, the rocks enter greenschist facies conditions, where

further retrogression is possible. In the greenschist facies, chlorite is formed and together with actinolite, epidote and feldspar, makes up the greenschists. ~~The high chlorite~~ Chlorite and actinolite content results in intermediate to high anisotropies- ΔVP of 5-3.7 %, presented in this study, and a The velocity distribution with displays highest VP parallel to the stretching lineation slightly extending into the foliation plane caused by chlorite. Very few studies have been carried out on the seismic properties of greenschists and further research is necessary to better constrain this lithology. high velocities in the foliation plane (figure 8). This is directly caused by the sheet silicate chlorite and provides greenschist VP patterns very similar to those of the metasediments and the deformed continental crust. Very few studies have been made on the elastic properties of greenschists and further research is necessary to better constrain this lithology.

~~The present study, however, shows that greenschist facies metabasites constitute a special case concerning their elastic properties. While VP is still higher than that of metasediments and comparable to the other metabasites, the velocity distribution resembles that of the metasediments. Due to their frequent occurrence in subduction settings and collisional orogens this needs to be considered in seismic investigations.~~

In the overlying metasediments VP anisotropies display values of 2.3-~~26.41~~4.4 % (Erdman et al., 2013; Keppler et al., 2015; Weiss et al., 1999; Kästner et al., 2020; Keppler et al., submitted to same issue). The composition of these metasediments in the present study is highly variable. However, quartz, feldspar, mica, chlorite, calcite and dolomite are the most abundant mineral phases and also those controlling control the elastic seismic properties. The velocity distribution shows the highest velocities in the foliation plane with a velocity-minimum normal to it, as seen in the greenschists, due to the high contents in sheet silicates as well as of calcite and dolomite.

In the deformed continental crust the composition is dominated by feldspar, quartz, mica and chlorite, represented by the gneisses in this study. In the gneisses the elastic anisotropies ΔVP are similar to those observed in the metasediments, with values of 0.9-20.54-~~72~~20.4 % (Ivankina et al., 2005; Ullemeyer et al., 2006; Erdman et al., 2013; Llana-Fúnez and Brown, 2012; Kern et al., 2008; Weiss et al., 1999; Vasin et al., 2017; Khazanehdari et al., 2000; Kästner et al., 2020; Keppler et al., submitted to same issue) even though our samples are at the lower end of this range. The velocity distribution of the gneisses displays highest velocities in a girdle in the foliation plane, similar to the greenschists and metasediments. This results from the high sheet silicate-contents of this lithology (figure 8h-i). In both the metasediments and the gneisses the metamorphic facies does not change the elastic properties beyond the greenschist facies up to HP conditions, since the single-crystal anisotropies in most of the constituent minerals are constant at higher metamorphic conditions (e.g. quartz, feldspar, calcite and dolomite). Mica shows changes in chemistry with rising pressures and quartz transitions to is replaced by coesite at UHP conditions, but due to limited single-crystal anisotropy data of these minerals their influence cannot be determined.

With these structural positions and their characteristic elastic properties in mind, a better interpretation of seismic data is feasible. Lower VP/VS-ratios with high elastic anisotropies hint at felsic compositions and high sheet silicate content, while the higher VP/VS-ratios and overall lower to intermediate elastic anisotropies ΔVP indicate more mafic lithologies, with exception of the zones of both high velocity ratios and high anisotropies characteristic of deformed greenschists (Almqvist and Mainprice, 2017). In the majority of the mafic rocks a dominance of high velocities parallel to the stretching lineation

can be observed. In comparison, the metasediments and gneisses display a distinctly different velocity distribution with the highest velocities in the foliation plane. ~~But again, greenschists are an exception. This mafic lithology displays high velocities in the foliation plane with an intermediate to high P-wave anisotropy of 5% reminiscent of the metasediments.~~ The knowledge of the characteristic properties of the different structural positions can ~~help to better~~ help to identify thickness and boundaries of the predominantly mafic to ultramafic subducting plate ~~and the felsic continental margins~~. It ~~can further assist~~ might also allow to better constrain the anatomy of subduction channels.

6.5 Common issues for the calculation and application of seismic properties~~Common issues for calculated elastic properties~~

Precise single-crystal data are not available for all mineral phases, but our fitting represents a good approximation. Firstly, the selected substitute minerals are very similar to those in the samples, especially with respect to their crystallography. Secondly, the goal of our models is not to simulate specific rock samples, but the rock types that are representative of distinct sections of the subduction and exhumation cycle. Hence, for the missing mineral phases substitute single crystal data were selected, which we considered to be acceptable matches due to the metamorphic and petrological context they can be found in. As mentioned above, this was done for clinozoisite, for which we selected the single-crystal data of epidote (Aleksandrov et al., 1974), ~~as well as for chlorite, for which we selected the data of muscovite (Vaughan and Guggenheim, 1986).~~ Furthermore, the almandine garnet single-crystal data of Zhang et al. (2008) was used for all garnet-bearing samples, even if different garnet compositions were present in the samples. A similar simplification was made for the amphiboles. In samples with glaucophane or barroisite, we used glaucophane single-crystal data by Bezacier et al. (2010), ~~while for all other amphiboles we utilized the single-crystal elastic constants for hornblende from Aleksandrov and Rhyzova (1961).~~

Throughout this study the Voigt averaging scheme was used ~~consistently~~. This model is based on the “equal strain” assumption and represents the upper bound for the stiffness of a polycrystalline material. As a result, the velocities presented are ~~therefore~~ maximum velocities ~~only~~. This does not correspond to the “equal stress” assumption of the Reuss model that represents the lower stiffness boundary. Hence, the use of the Voigt method should be kept in mind for the interpretation of the calculated P-wave velocity distributions. Neither method takes grain shape, layering, microcracks, fractures or pores into account ~~(Vasin, et al. 2017). A study by Cyprych et al. (2017) showed that particularly in rocks with weak textures, the microstructural arrangement of minerals can influence the anisotropy. There are~~ Other methods such as the self-consistent method GeoMIXself (GMS) by Matthies (2010; 2012) or the ESP toolbox of Vel et al. (2016), which can take CPOs, morphologies and SPOs of the mineral grains into account. However, only based on CPO data the Voigt averaging scheme is ~~deemed assumed to be~~ appropriate to estimate representative samples. ~~Furthermore, the Voigt averaging scheme has widely been used on the subject and ensures therefore comparability with most other published texture-derived elastic anisotropy data.~~

In addition to CPO based anisotropy of individual lithological units, local tectonic structures can affect the regional anisotropy. Folds and local shear zones can have a strong impact on the velocity patterns and the large-scale structural

Formatiert: Englisch (USA)

660 geometry has to be accounted for (e.g. Bleibinhaus and Gebrande, 2006; Okaya et al., 2018). The possibility to detect these structures is strongly dependent on the resolution of seismic imaging. In addition, the propagation of seismic waves is usually not parallel to the symmetry axes of geological structures within the crust. Such non-axial propagation has to be taken into account in any seismic model (e.g. Okaya and Christensen, 2002).

665 In this study all volume proportions of the different mineral phases were estimated from thin-section microscopy and calculated by RTA in MAUD. Related errors need to be considered, when calculating the elastic properties. However, as we do not wish to model specific rocks rather general lithologies, the resulting compositional variations fall within the spectrum of the lithologies of interest.

670 As the single crystal properties were measured at ambient conditions and our calculations were based on these, a degree of uncertainty results from changes of these properties with metamorphic conditions. In our calculations we cannot take these changes into account, yet as noted in the example of the barrosite-bearing sample MJS16 this is a possibility for errors.

A further issue, which requires discussion is the influence of microcracks on the elastic properties of ~~the crustal~~ rocks (Siegesmund and Vollbrecht, 1991; Siegesmund et al., 1993; Ullemeyer et al. 2011, Vasin et al., 2014; 2017). The ~~elastic seismic~~ anisotropies presented in this study are texture-derived and do not take the presence of microcracks into account. Instead, they depend on the mineral composition, single-crystal mineral properties and the CPOs of the respective phases.

675 This means that the calculated data reflect the ~~elastic-seismic~~ anisotropies of rocks at great depth, in which microcracks are closed due to high confining pressure. The influence of microcracks can well be observed in studies in which CPO-derived calculated results have been compared to ~~elastic-seismic~~ anisotropy measurements in laboratory experiments at high confining pressure (some examples listed in table 3). Some of these studies have demonstrated that the influence of microcracks is still detectable up to confining pressures of 1 GPa (Christensen, 1974; Ullemeyer et al., 2011). However, with increasing pressure and closing of the microcracks, the results far better resemble the CPO-derived results and the influence of microcracks decreases significantly (table 3; Ivankina et al., 2005; Lokajicek et al., 2017; Vasin et al., 2017).

7. Conclusion

Using time-of-flight neutron diffraction, texture data from a wide range of compositionally variable and complex rocks were collected. These are representative of different positions for subduction and exhumation in a collisional orogen. From these data texture-derived ~~elastic-seismic~~ anisotropies and other petrophysical properties were modelled. The following conclusions can be made:

1. Composition and CPO have a strong effect on the rock texture and the related elastic properties in subduction and exhumation settings. Felsic rocks with high sheet silicate and calcite or dolomite contents preferentially display the fastest P-wave velocities in the foliation plane independently of the orientation of the stretching lineation. The mafic rocks with high amphibole and pyroxene contents in this study mostly display highest P-wave velocities parallel to the stretching lineation, with exception of greenschist, in which the high chlorite content results in the fast velocities in the foliation plane.

2. Among mafic rocks the elastic anisotropies vary ~~strongly~~ by composition. The elastic anisotropies increase from a composition dominated by pyroxenes (1.4 %), to those dominated by ~~ever~~ amphiboles and chlorite (3-12.8-3.7 %), and are highest in the chlorite dominated greenschist (5 %).

3. The described structural positions have varying elastic properties (~~elastic anisotropy~~ $\Delta V/P$, VP/VS-ratio, P-wave-velocity distribution), often overlapping each other. However, in many cases the combination of properties can be utilized to distinguish these positions from one another. In this study the mafic rocks of oceanic crustal origin such as eclogite, blueschist, greenschist and amphibolite display lower anisotropies (1.4-3.7 %) with higher VP/VS-ratios (1.~~7470~~-1.~~7675~~) and highest velocities parallel to the stretching lineation. An exception to this is the greenschist (5 %: 1.73), which displays properties more characteristic to the felsic rocks of the continental crust and the metasedimentary cover. These gneisses, micaschists and calcareous schists display higher elastic anisotropies (1.~~84~~-~~8.27~~1 %), lower VP/VS-ratios (1.56-1.76) and fast P-wave-velocities distributed in the foliation plane.

Acknowledgments

This study was funded by the German research foundation (DFG-grant No. KE 2268/2-1) as part of the DFG priority programme “Mountain Building Processes in 4 Dimensions”. Fruitful discussions within the priority programme are gratefully acknowledged. We wish to thank the Frank Laboratory of Neutron Physics (FLNP) at the Joint Institute of Neutron Research (JINR) in Dubna, Russia for the texture measurements made possible at SKAT. We acknowledge very constructive and helpful comments by Bjarne Almqvist and an anonymous reviewer, which have greatly improved the quality of the manuscript.

References:

- Aleksandrov, K.S., and Ryzhova, T.V.: The elastic properties of rock forming minerals, Izvestija Academy of Science USSR, Geophys. Ser. 12, 1799–1804, 1961.
- Aleksandrov, K.S., Alchikov, U.V., Belikov, B.P., Zaslavski, B.I., and Krupny, A.I.: Elastic wave velocities in minerals at atmospheric pressure and increasing precision of elastic constants by means of EVM, Izvestija Academy of Science USSR, Geol. Ser. 10, 15–24, 1974.
- Almqvist, B.S.G., and Mainprice, D: Seismic properties and anisotropy of the continental crust: Predictions based on mineral texture and rock microstructure, Rev. Geophys., 55, 367–433, 2017.
- Amato, J.M., Johnson, C.M., Baumgartner, L.P., and Beard, B.L.: Rapid exhumation of the Zermatt–Saas ophiolite deduced from high precision Sm–Nd and Rb–Sr geochronology, Earth Planet Sci Lett 171:425–438, 1999.
- Angiboust, S., Glodny, J., Oncken, O., and Chopin, C.: In search of transient subduction interfaces in the Dent Blanche–Sesia Tectonic System (W. Alps), Lithos 205:298–321, 2014.

- Ballèvre, M., Kienast, J.R., Vuichard, and J.P.: La “nappe de la Dent-Blanche” (Alpes occidentales): deux unités austroalpines indépendantes, *Eclogae Geol Helv* 79:57–74, 1986.
- 725 [Barruol, G. and Kern, H.: Seismic anisotropy and shear-wave splitting in lower-crustal and upper-mantle rocks from the Ivrea Zone experimental and calculated data. *Physics of the Earth and Planetary Interiors* 95, 175-194, 1996.](#)
- [Bascou, J., Barruol, G., Vauchez, A., Mainprice, D., Egydio-Silva, M.: EBSD-measured lattice-preferred orientations and seismic properties of eclogites, *Tectonophysics* 342, 61– 80, 2001.](#)
- Bestmann, M., Kunze, K., and Matthews, A.: Evolution of a calcite marble shear zone complex on Thassos Island, Greece; microstructural and textural fabrics and their kinematic significance, *Journal of Structural Geology* 22 (11–12), 1789–1807,
- 730 2000.
- Bestmann, M., Prior, D.J., and Grasemann, B.: Characterisation of deformation and flow mechanics around porphyroclasts in a calcite marble ultramylonite by means of EBSD analysis, *Tectonophysics* 413, 185–200, 2006.
- Bezacier, L., Reynard, B., Bass, J.D., Wang, J., and Mainprice, D.: Elasticity of glaucophane, seismic velocities and anisotropy of the subducted oceanic crust, *Tectonophysics* 494, 201–210, 2010.
- 735 Bhagat, S.S., Bass, J.D., Smyth, and J.R.: Single-crystal elastic properties of omphacite-C2/C by Brillouin spectroscopy, *J. Geophys. Res. Solid Earth* 97, 6843–6848, 1992.
- [Bleibinhaus, F., and Gebrande, H.: Crustal structure of the Eastern Alps along the TRANSALP profile from wide-angle seismic tomography: *Tectonophysics*, v. 414, p. 51–69, 2006.](#)
- [Boundy, T.M., Fountain, D.M. and Austrheim, H.: Structural development and petrofabrics of eclogite facies shear zones, Bergen Arcs, western Norway: implications for deep crustal deformational processes. *J. metamorphic Geol.*, 10, 127-146, 1992.](#)
- 740 Bousquet, R.: Metamorphic heterogeneities within a single HP unit: overprint effect or metamorphic mix? *Lithos* 103:46–69, 2008.
- Brown, J.M., Abramson, E.H., and Angel, R.J.: Triclinic elastic constants for low albite, *Phys. Chem. Miner.* 33, 256–265,
- 745 2006.
- [Brown, J. M., & Abramson, E. H.: Elasticity of calcium and calciumsodium amphiboles. *Physics of the Earth and Planetary Interiors*, 261, 161-171, 2016. doi:10.1016/j.pepi.2016.10.010.](#)
- Brown, J. M., Angel, R. J., and Ross, N. L.: Elasticity of plagioclase feldspars, *J. Geophys. Res. Solid Earth*, 121, 663–675, 2016.
- 750 Bucher, K., Fazis, Y., de Capitani, C., and Grapes, R.: Blueschists, eclogites, and decompression assemblages of the Zermatt–Saas ophiolite: high-pressure metamorphism of subducted Tethys lithosphere, *Am Mineral* 90:821–835, 2005.
- Bunge, H.: *Texture Analysis in Materials Science: Mathematical Models*. Butterworths, London, 1982.
- Burlini, L., Marquer, D., Challandes, N., Mazzola, S., and Zangarini, N.: Seismic properties of highly strained marbles from the Splügenpass, central Alps, *Journal of Structural Geology* 20, 277–292, 1998.

- 755 Cao, Y., Jung, H., and Song, S.: Petro-fabrics and seismic properties of blueschist and eclogite in the North Qilian suture zone, NW China: Implications for the low-velocity upper layer in subducting slab, trench-parallel seismic anisotropy, and eclogite detectability in the subduction zone, *J. Geophys. Res. Solid Earth*, 118, 3037–3058. doi:10.1002/jgrb.50212, 2013.
[Cao, Y., Jung, H., & Ma, J.: Seismic properties of a unique olivine-rich eclogite in the Western Gneiss Region, Norway, *Minerals*, 10\(9\), 774, 2020.](#)
- 760 Chateigner, D., Lutterotti, L. and Morales, M.: “Chapter 5.3: Quantitative texture analysis and combined analysis”, *International Tables of Crystallography*, volume H: Powder Diffraction. International Union of Crystallography, doi:10.1107/97809553602060000968, 2019.
[Chen, C.-C., Lin, C.-C., Liu, L.G., Sinogeikin, S.V., Bass, J.D.: Elasticity of single-crystal calcite and rhodochrosite by Brillouin spectroscopy, *American Mineralogist*, Volume 86, pages 1525–1529, 2001.](#)
- 765 Christensen, N.I.: Compressional wave velocities in possible mantle rocks to pressures of 30 kbars, *J. Geophys. Res.* 79, 407–412, 1974.
 Christoffel, E.B.: Über die Fortpflanzung von Stößen durch elastische, feste Körper, *Annali di Matematica* 8, 193–243, ~~1874~~[1877](#).
[Cyprych, D., Piazzolo, S., Almqvist, B.: Seismic anisotropy from compositional banding in granulites from the deep magmatic arc of Fiordland, New Zealand, *Earth and Planetary Science Letters*, 477, 156-167, 2017.](#)
~~Dandekar, D.P.: Variation in the elastic constants of calcite with pressure, *Am. Geophys. Union Trans.* 49, 323 pp., 1968.~~
~~De Bresser, J.H.P.: Calcite c axis textures along the Gavarnie thrust zone, central Pyrenees, *Geologie en Mijabouw* 68, 367–375, 1989.~~
- 770 Delle Piane, C., Burlini, L., and Kunze, K., 2009. The influence of dolomite on the plastic flow of calcite Rheological, microstructural and chemical evolution during large strain torsion experiments. *Tectonophysics* 467, 145–166.
- Erdman, M. E., Hacker, B.R., Zandt, G., and Seward, G.: Seismic anisotropy of the crust: Electron-backscatter diffraction measurements from the Basin and Range, *Geophys. J. Int.*, doi:10.1093/gji/ggt287, 2013.
- Erskine, B.G., Heidelbach, F., and Wenk, H.R.: Lattice preferred orientations and microstructures of deformed Cordilleran marbles; correlation of shear indicators and determination of strain path, *J. Struct. Geol.*, 15, 9–10, 1189–1205, 1993.
- 780 Frezzotti, M.L., Selverstone, J., Sharp, Z.D., and Compagnoni, R.: Carbonate dissolution during subduction revealed by diamond-bearing rocks from the Alps, *Nat Geosci* 4:703–706, 2011.
[Fujimoto, Y., Kono, Y., Hirajima, T., Kanagawa, K., Ishikawa, M., Arima, M.: Pwave velocity and anisotropy of lawsonite and epidote blueschists: constraints on water transportation along subducting oceanic crust. *Phys. Earth Planet. Inter.* 183, 219–228, 2010.](#)
- 785 Godard, G., and Van Roermund, H.L.M.: Deformation-induced clinopyroxene fabrics from eclogites, *J. Struct. Geol.* 17, 1425–1443, 1995.
 Gouzu, C., Itaya, T., Hyodo, H., and Matsuda, T.: Excess ⁴⁰Ar-free phengite in ultrahigh-pressure metamorphic rocks from the Lago di Cignana area, Western Alps, *Lithos* 92:418–430, 2006.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

- Groppo, C., Beltrando, M., and Compagnoni, R.: The P–T path of the ultra-high pressure Lago di Cignana and adjoining meta-ophiolitic units: insights into the evolution of the subducting Tethyan slab, *J. Met. Geol.* 27:207–231, 2009.
- 790 Hacker, B.R., and Christie, J.M.: Brittle/ductile and plastic/cataclastic transitions in experimentally deformed and metamorphosed amphibolite, *AGU Monograph* 56, 127-147, 1990.
- Helmstaedt, H., Anderson, D.L., and Gavasci, A.T.: Petrofabric studies of eclogite, spinel websterite, and spinel-lherzolite xenoliths from kimberlite-bearing breccia pipes in southeastern Utah and northeastern Arizona, *J. Geophys. Res.* 77, 4350–
- 795 4365, 1972.
- Heyliger, P., Ledbetter, H., Kim, S.: Elastic constants of natural quartz, *J. Acoust. Soc. Am.* 114, 644–650, 2003.
- Hill, R.: The elastic behavior of a crystalline aggregate, *Proc. Phys. Soc. A.* 65, 349–354, 1952.
- Humbert, P., and Plique, F.: Propriétés élastiques de carbonates rhomboédriques monocristallins calcite, magnésite, dolomite, *C.R. Acad. Sci. Paris*, 275, 391–394, 1972.
- 800 Ivankina, T.I., Kern, H., and Nikitin, A.N.: Directional dependence of P- and S-wave propagation and polarization in foliated rocks from the Kola superdeep well: evidence from laboratory measurements and calculations based on TOF neutron diffraction, *Tectonophysics* 407, 25–42, 2005.
- Ivankina, T.I., Nikitin, A.N., and Telepnev, A.: Textures and physical properties of marbles deformed at 20–250 °C, *High Pres. Res.* 17, 335, 2000.
- 805 Ivankina, T.I., Nikitin, A.N., and Telepnev, A.S.: The effect of temperature and long-term mechanical stresses on the deformational, thermal, and textural characteristics of marble, *Izvestiya, Phys. Sol. Earth* 37, 1, 46, 2001.
- [Kästner, F., Pierdominici, S., Elger, J., Zappone, A., Kück, J., Berndt, C.: Correlation of core and downhole seismic velocities in high-pressure metamorphic rocks: a case study for the COSC-1 borehole, Sweden, *Solid Earth*, 11, 607–626, 2020.](#)
- 810 Kern, H., Ivankina, T.I., Nikitin, A.N., Lokajicek, T., and Pros, Z.: The effect of oriented microcracks and crystallographic and shape preferred orientation on bulk elastic anisotropy of a foliated biotite gneiss from Outokumpu, *Tectonophysics* 457, 143–149, 2008.
- Kern, H., and Wenk, H.R.: Calcite texture development in experimentally induced ductile shear zones, *Contributions to Mineralogy and Petrology* 83 (3–4), 231–236, 1983.
- 815 Kern, H., Jin, Z., Gao, S., Popp, T., and Xu, Z.: Physical properties of ultrahigh-pressure metamorphic rocks from the Sulu terrain, eastern central China: implications for the seismic structure at the Donghai (CCSD) drilling site, *Tectonophysics* 354, 315–330, 2002.
- Keppeler, R.: Crystallographic preferred orientations in eclogites – A review, *J. Struct. Geol.*, 115, 284-296, 2018.
- Keppeler, R., Stipp, M., Behrmann, J.H., Ullemeyer, K., and Heidelbach, F.: Deformation inside a paleosubduction channel—
- 820 insights from microstructures and crystallographic preferred orientations of eclogites and metasediments from the Tauern Window, Austria, *J. Struct. Geol.* 82, 60–79, 2016.

- Keppler, R., Ullemeyer, K., Behrmann, J.H., Stipp, M.: Potential of full pattern fit methods for the texture analysis of geological materials: implications from texture measurements at the recently upgraded neutron time-of-flight diffractometer SKAT, *J. Appl. Crystallogr.* 47. DOI:10.1107/S1600576714015830, 2014.
- 825 Keppler, R., Behrmann, J.H., Stipp, M.: Textures of eclogites and blueschists from Syros island, Greece: inferences for elastic anisotropy of subducted oceanic crust, *Geophys. Res. Solid Earth* DOI:10.1002/2017JB014181, 2017.
- Keppler, R., Ullemeyer, K., Behrmann, J.H., Stipp, M., Kurzawski, R.M., and Lokajíček, T.: Crystallographic preferred orientations of exhumed subduction channel rocks from the Eclogite Zone of the Tauern Window (Eastern Alps, Austria), and implications on rock elastic anisotropies at great depth, *Tectonophysics*, 647-648, 89-104, 2015.
- 830 [Keppler R., Vasin R.N., Stipp, M., Lokajíček, T., Petruzálek, M., Froitzheim, N., Elastic anisotropies of deformed upper crustal rocks in the Alps, submitted to same issue.](#)
- [Kim, D., Katayama, I., Michibayashi, K., Tsujimori, T.: Deformation fabrics of natural blueschists and implications for seismic anisotropy in subducting oceanic crust, *Physics of the Earth and Planetary Interiors*, Volume 222, Pages 8-21, 2013.](#)
- Kirst, F., and Leiss, B.: Kinematics of syn- and post-exhumational shear zones at Lago di Cignana (Western Alps, Italy): constraints on the exhumation of Zermatt–Saas (ultra)high-pressure rocks and deformation along the Combin Fault and Dent Blanche Basal Thrust, *Int. J. Earth Sci.*, 106, 1, 215-236, 2016.
- 835 Kurz, W.: Constriction during exhumation: evidence from eclogite microstructures, *Geology* 33, 1, 37–40, 2005.
- Kurz, W., Jansen, E., Hundenborn, R., Pleuger, J., Schäfer, W., and Unzog, W.: Microstructures and crystallographic preferred orientations of omphacite in Alpine eclogites: implications for the exhumation of (ultra-) high-pressure units, *J. Geodyn.* 37, 1–55, 2004.
- 840 Lapen, T.J., Johnson, C.M., Baumgartner, L.P., Mahlen, N.J., Beard, B.L., Amato, J.M.: Burial rates during prograde metamorphism of an ultrahigh-pressure terrane: an example from Lago di Cignana, Western Alps, Italy, *Earth Planet. Sci. Lett.*, 215, 57–72, 2003.
- Leiss, B., Siegesmund, S., and Weber, K.: Texture asymmetries as shear sense indicators in naturally deformed mono- and poly-phase carbonate rocks, *Textures and Microstructures* 33, 61–74, 1999.
- 845 Link, F., and Rümpler, G.: Crustal anisotropy in the European Alps inferred from crustal phases in receiver functions and first implications for the mantle dynamics, *Geophysical Research Abstracts* 21, EGU2019-7235, 2019.
- Lister, G.S., and Hobbs, B.E. : The simulation of fabric development during plastic deformation and its application to quartzite: the influence of deformation history, *J. Struct. Geol.* 2, 355–371, 1980.
- 850 Llana-Fúnez, S., and Brown, D.: Contribution of crystallographic preferred orientation to seismic anisotropy across a surface analog of the continental Moho at Cabo Ortegal, Spain, *Geol. Soc. Am. Bull.*, 124, 1495–1513, 2012.
- Lokajíček, T., Svitek, T., Kern, H., and Wenk, H.-R.: Rock Elastic Anisotropy Study by X-ray Synchrotron, Neutron Diffraction and Ultrasonic Methods. *Procedia Engineering*, 191, 467-475, 2017.
- Lutterotti, L., Chateigner, D., Ferrari, S., and Ricote, J.: Texture, residual stress and structural analysis of thin films using a combined X-ray analysis, *Thin Solid Films*, 450, 34-41, 2004.
- 855

- Lutterotti, L., Matthies, S., Wenk, H.-R., Schultz, A.J., and Richardson, J.W.: Combined texture and structure analysis of deformed limestone from time-of-flight neutron diffraction spectra, *J. Appl. Phys.* 81, 594–600, 1997.
- Lutterotti, L.: Total pattern fitting for the combined size-strain-stress-texture determination in thin film diffraction, *Nuclear Inst. and Methods in Physics Research, B*, 268, 334–340, 2010.
- 860 Mainprice, D., and Ildefonse, B.: Seismic anisotropy of subduction zone minerals - contribution of hydrous phases, In: Lallemand, S., Funiciello, F. (Eds.), *Subduction Zone Geodynamics*. Springer-Verlag, Berlin Heidelberg, 63–84, 2009.
- Mainprice, D., and Humbert, M.: Methods of calculating petrophysical properties from lattice preferred orientation data, *Surv. Geophys.* 15, 575–592, 1994.
- Manzotti, P., Zucali, M., Ballèvre, M., Robyr, M., and Engi, M.: Geometry and kinematics of the Roisan-Cignana Shear
865 Zone, and the orogenic evolution of the Dent Blanche Tectonic System (Western Alps), *Swiss J. Geosci.*, 107, 23–47, 2014.
- Matthies, S.: On the combination of self-consistent and geometric mean elements for the calculation of the elastic properties of textured multi-phase samples, *Solid State Phenom.*, 160, 87–93, 2010.
- Matthies, S.: GEO-MIX-SELF calculations of the elastic properties of a textured graphite sample at different hydrostatic pressures, *J. appl. Crystallogr.*, 45, 1–16, 2012.
- 870 Matthies, S., and Humbert, M.: On the principle of a geometric mean of even-rank symmetric tensors for textured polycrystals, *J. Appl. Crystallogr.* 28, 254–266, 1995.
- Matthies, S., Lutterotti, and L., Wenk, H.R.: Advances in Texture Analysis from Diffraction Spectra, *J. Appl. Cryst.* 30, 31–42, 1997.
- McNamara, D.D., Wheeler, J., Pearce, M., Prior, and D.J.: Fabrics produced mimetically during static metamorphism in
875 retrogressed eclogites from the Zermatt-Saas zone, Western Italian Alps, *J. Struct. Geol.* 44, 167–178, 2012.
- [Mookherjee, M., & Mainprice, D.: Unusually large shear wave anisotropy for chlorite in subduction zone settings, *Geophysical Research Letters*, 41\(5\), 1506-1513, 2014. doi:10.1002/2014gl059334.](#)
- Müller, W.F., Xu, Z., and Walte, N.P.: Transmission electron microscopy of omphacite and other minerals in eclogites from the CCSD borehole, China: indications for their deformation and temperature history, *Eur. J. Mineral* 23, 645–659, 2011.
- 880 Neufeld, K., Ring, U., Heidelbach, F., Dietrich, S., and Neuser, R.D.: Omphacite textures in eclogites of the Tauern Window: implications for the exhumation of the eclogite zone, eastern Alps, *J. Struct. Geol.* 30, 976–992, 2008.
- Nikitin, A.N., and Ivankina, T.I.: Neutron scattering in geosciences, *Part. Nucl. Sci. Rev. J.* 35, part 2, 348–407, 2004.
- Nikitin, A.N., Ivankina, T.I., and Sobolev, G.A.: Neutron diffraction study of intracrystalline strains and stresses in a marble sample at high temperatures and external load, *Izvestiya. Phys. Sol. Earth*, 40, 1, 83, 2004.
- 885 Oberhänsli, R., and Bucher K.: Tectonometamorphic evolution of the Dent Blanche nappe, *Terra Cognita*, 7, 2-3, 95, 1987.
- Peacock, S.M.: The importance of blueschist -> eclogite dehydration reactions in subducting oceanic crust, *Geological Society of America Bulletin*, 105, 684 – 694, 1993.
- [Okaya, D.A. Vel, S.S., Johnson, S.E., and Song, W.J.: Modification of crustal seismic anisotropy by geological structures \("structural geometric anisotropy"\), *Geosphere*, 15, 2018.](#)

- 890 [Okaya, D. A. and Christensen, N.I.: Anisotropic effects of non-axial seismic wave propagation in foliated
crustal rocks. *Geophysical Research Letters*, 29, 11, 1507-1511, 2002.](#)
- Pleuger, J., Roller, S., Walter, J.M., Jansen, E., and Froitzheim, N.: Structural evolution of the contact between two Penninic nappes (Zermatt-Saas zone and Combin zone, Western Alps) and implications for the exhumation mechanism and palaeogeography, *Int. J. Earth Sci.*, 96, 229-252, 2007.
- 895 Prior, D.J., and Wheeler, J.: Feldspar fabrics in a greenschist facies albite-rich mylonite from electron backscatter diffraction, *Tectonophysics*, 303, 29–49, 1999.
- Puelles, P., Beranoaguirre, A., Ábalos, B., Gil Ibarguchi, J.I., García de Madinabeitia, S., Rodríguez, J., and Fernández-Armas, S.: Eclogite inclusions from subducted metagneous continental crust (Malpica-Tui Allochthonous Complex, NW Spain): petrofabric, geochronology and calculated seismic properties, *Tectonics* 36, 1376–1406, 2017.
- 900 Puntero, G., Kunze, K., Heilbronner R., Stünitz, H., and Schmid, S.M.: Microfabrics of calcite ultramylonites as records of coaxial and non-coaxial deformation kinematics: Examples from the Rocher de l'Yret shear zone (Western Alps), *Tectonophysics*, 424, 69–97, 2005.
- Ratschbacher, L., Wenk, H.R., and Sintubin, M.: Calcite textures; examples from nappes with strain-path partitioning, *J. Struc. Geol.*, 13, 4, 369–384, 1991.
- 905 Reddy, S.M., Wheeler, J., and Cliff, R.A.: The geometry and timing of orogenic extension: an example from the Western Italian Alps, *J. Met. Geol.* 17, 573–589, 1999.
- Reddy, S.M., Wheeler, J., Butler, R.W.H., Cliff, R.A., Freeman, S., Inger, S., Pickles, C., and Kelley, S.P.: Kinematic reworking and exhumation within the convergent Alpine Orogen, *Tectonophysics* 365, 77–102, 2003.
- Reuss, A.: Berechnung der Fließgrenze von Mischkristallen auf Grund der Plastizitätsbedingung für Einkristalle, *Z. Angew. Math. Mech.* 9, 49–58, 1929.
- 910 Satsukawa, T., Ildefonse, B., Mainprice, D., Morales, L.F.G., Michibayashi, K., and Barou, F.: A database of plagioclase crystal preferred orientations (CPO) and microstructures – implications for CPO origin, strength, symmetry and seismic anisotropy in gabbroic rocks, *Solid Earth*, 4, 511-542, 2013.
- Siegesmund, S., Takeshita, T., and Kern H.: Anisotropy of V_p and V_s in an amphibolite of the deeper crust and its relationship to the mineralogical, microstructural and textural characteristics of the rock, *Tectonophysics*, 157, 25–38, 1989.
- 915 Siegesmund, S., and Vollbrecht, A.: Complete seismic properties obtained from microcrack fabrics and textures in an amphibolite from the Ivrea zone, Western Alps, Italy, *Tectonophysics* 1999, 13–24, 1991.
- Siegesmund, S., Vollbrecht, A., Chlupac, T., Nover, G., Dürrast, H., Müller, J., and Weber, K.: Fabric-controlled anisotropy of petrophysical properties observed in KTB core samples, *Sci. Drill.* 4, 31–54, 1993.
- 920 Schmid, S.M., Fügenschuh, B., Kissling, E., and Schuster, R.: Tectonic map and overall architecture of the Alpine orogeny, *Eclogae geol. Helv.*, 97, 93-117, 2004.
- Schmid, S.M., and Casey, M.: Complete fabric analysis of some commonly observed quartz c-axis patterns, *Geophys. Monogr.* 36, 263–286, 1986.

- Trullenque, G., Kern, H., Cirrincione, R., Mazzoleni, P., and Pezzino, A.: P- and S-wave velocities and densities in silicate and calcite rocks from the Peloritani Mountains, Sicily (Italy): The effect of pressure, temperature and the direction of wave propagation, *Tectonophysics*, 409, 55–72, 2006.
- Ullemeyer, K., Braun, G., Dahms, M., Kruhl, J.H., Olesen, N.Ø., and Siegesmund, S.: Texture analysis of a muscovite-bearing quartzite: a comparison of some currently used techniques, *J. Struct. Geol.* 22, 1541–1557, 2000.
- Ullemeyer, K., Leiss, B., and Stipp, M.: Textures and Microstructures in Peridotites from the Finero Complex (Ivrea Zone, Alps) and its Influence on the Elastic Rock Properties, *Solid State Phenomena* 160, 183-188, 2010.
- Ullemeyer, K., Spalhoff, P., Heinitz, J., Isakov, N.N., Nikitin, A.N., and Weber, K.: The SKAT texture diffractometer at the pulsed reactor IBR-2 at Dubna: experimental layout and first measurements, *Nucl. Inst. Methods Phys. Res.* 412, 80–88, 1998.
- Ullemeyer, K., Siegesmund, S., Rasolofosaon, P.N.J., and Behrmann, J.H.: Experimental and texture-derived P-wave anisotropy of principal rocks from the TRANSALP traverse: an aid for the interpretation of seismic field data, *Tectonophysics* 414, 97–116, 2006.
- Ullemeyer, K., Nikolayev, D.I., Christensen, N.I., and Behrmann, J.H.: Evaluation of intrinsic velocity–pressure trends from low-pressure P-wave velocity measurements in rocks containing microcracks, *Geophys. J. Int.* 185, 1312–1320, 2011.
- Vasin, R.N., Lebensohn, R.A., Matthies, S., Tome, C.N., and Wenk, H.-R.: The influence of grain shape and volume fraction of sheet silicates on elastic properties of aggregates: biotite platelets in an isotropic matrix, *Geophysics*, 79, 433–441, 2014.
- Vasin, R.N., Kern, H., Lokajčák, T., Svitek, T., Lehmann, E., Mannes, D.C., Chaouche, M., and Wenk, H.-R.: Elastic anisotropy of Tambo gneiss from Promontogno, Switzerland: a comparison of crystal orientation and microstructure-based modelling and experimental measurements, *Geophys. J. Int.*, 209, 1–20., 2017.
- Vaughan, M.T., and Guggenheim, S.: Elasticity of muscovite and its relationship to crystal structure, *J. Geophys. Res.* 91, 4657–4664, 1986.
- [Vel, S., Cook, A.C., Johnson, S.E., Gerbi, C.: Computational homogenization and micromechanical analysis of textured polycrystalline materials, *Comput. Methods Appl. Mech. Eng.*, 310, 749–779, 2016.](#)
- Voigt, W.: *Lehrbuch der Kristallphysik*. Teubner, Leipzig, 1928.
- Von Dreele, R.B.: Quantitative texture analysis by rietveld refinement, *J. Appl. Cryst.* 30, 517–525, 1997.
- Weiss, T., Siegesmund, S., Rabbel, W., Bohlen, T., and Pohl, M.: Seismic Velocities and Anisotropy of the Lower Continental Crust: A Review, *Pure appl. geophys.*, 156, 97–122, 1999.
- Wenk, H.-R., Lutterotti, L., and Vogel, S.C.: Rietveld texture analysis from TOF neutron diffraction data, *Powder Diffraction* 25, 283–296, 2010.
- Wenk, H.-R., Matthies, S., Donovan, J., and Chateignier, D.: BEARTEX, a Windows-based program system for quantitative texture analysis, *J. Appl. Cryst.* 31, 262–269, 1998.

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)

Formatiert: Englisch (USA)


Wenk, H.-R., Vasin, R.N., Kern, H., Matthies, S., Vogel, S.C., and Ivankina, T.I.: Revisiting elastic anisotropy of biotite gneiss from the Outokumpu scientific drill hole based on new texture measurements and texture-based velocity calculations, *Tectonophysics* 570–571, 123–134, 2012.

960 [Worthington, J.R., Hacker, B.R., Zandt, G.: Distinguishing eclogite from peridotite: EBSD-based calculations of seismic velocities. *Geophys. J. Int.*, 193,489–505, 2013.](#)

[Zertani, S., John, T., Tilmann, F., Motra, H. B., Keppler, R., Andersen, T. B., & Labrousse, L.: Modification of the seismic properties of subducting continental crust by eclogitization and deformation processes. *Journal of Geophysical Research: Solid Earth*, 124, 9731–9754, 2019. <https://doi.org/10.1029/2019JB017741>.](#)

965 [Zhang, J. F., & Green, H. W.: On the deformation of UHP eclogite: From laboratory to nature, *International Geology Review*, 49\(6\), 487-503 2007.](#)

Zhang, J.F., Wang, Y.F., and Jin, Z.M.: CPO-induced seismic anisotropy in UHP eclogites, *Sci China Ser D-Earth Sci*, Vol. 51, No. 1, 11-21, 2008.








Figure 1: (a) Miniature map displaying approximate location of the study area in the European alpine orogen. (b) Geological map of the study area at Lago di Ciganan, in the Western Italian Alps modified after the map of Kirst and Leiss, 2016. Sample locations indicated by stars. (c) Stereoplots of foliation normal and stretching lineation as equal area projections in the lower hemisphere; data compiled from own measurements (diamonds) and from Kirst and Leiss (2016) (circles); Colors match those used in the map.



975

Figure 2: Schematic Cross section of subduction and exhumation in a collisional orogeny. The stars show locations which the samples represent within this setting. Colors correlate to map in figure 1.







Figure 3: Thin-section photographs of typical microstructures in four representative samples. (a) eclogite (MJS17) with foliation of glaucophane (Gln) and omphacite (Omp) flowing around larger grains of garnet (Grt) with some retrograde chlorite (Chl) visible at its rim. (b) greenschist (MJS36) with actinolite (Act) and chlorite (Chl) visible in the foliation and larger chlorite grains found

985 unoriented grown over the foliation. (c) calcareous micaschist (MJS20) with quartz (Qz), calcite (Calc) and white mica (Wmca)
oriented in the foliation. (d) gneiss (MJS35) displaying strongly foliated fine grained quartz (Qz) and white mica (Wmca) lar ger
coarse grained quartz. (e) amphibolite (MJS26) displaying abundant hornblende (Hbl) with some chlorite (Chl) and smaller
fragmented garnet (Grt) grains. (f) garnet (Grt) rich band in the blueschist (MJS36) surrounded by barroisite (Brs) and
clinozoisite (Czo). Sections are oriented perpendicular to the foliation and parallel to the stretching lineation. (a), ~~and (b)~~, (e) and
(f) are shown under plane polarized lighte, while (c) and (d) are shown under x-polarized light. For sample descriptions see table 1.


990

Eclogite - MJS17


Glaucophane




Omphacite




Clinozoisite




White Mica




Barroisite




Clinozoisite



Epidote




Blueschist - MJS16



995

Figure 4: Calculated pole figures of important mineral phases in the eclogite (**a**: MJS17) and the blueschist (**b**: MJS16). The stretching lineation (kinematic x-direction) is oriented EW in the pole figures, the foliation normal (kinematic z-direction) is oriented NS and the kinematic y-direction (perpendicular to x- and z-directions) is normal to the pole figure plane. The pole figures are lower hemisphere equal area projections. The contour levels display multiples of random distribution and minimum and maximum values are listed below the pole figure.








Figure 5: Calculated pole figures of important mineral phases in the amphibolite (a; MJS26) and the greenschist (b; MJS36). The stretching lineation (kinematic x-direction) is oriented EW in the pole figures, the foliation normal (kinematic z-direction) is oriented NS and the kinematic y-direction (perpendicular to x- and z-directions) is normal to the pole figure plane. The pole figures are lower hemisphere equal area projections. The contour levels display multiples of random distribution and minimum and maximum values are listed below the pole figure.


1005




Dolomitic Micaschist - MJS18



Calcareous Micaschist - MJS20




Calcschist - MJS22




1015 | Figure 6: Calculated pole figures of important mineral phases in the dolomitic micaschist ([a](#); MJS18), the calcareous micaschist ([b](#); MJS20) and the calcschist ([c](#); MJS22). The stretching lineation (kinematic x-direction) is oriented EW in the pole figures, the foliation normal (kinematic z-direction) is oriented NS and the kinematic y-direction (perpendicular to x- and z-directions) is normal to the pole figure plane. The pole figures are lower hemisphere equal area projections. The contour levels display multiples of random distribution and minimum and maximum values are listed below the pole figure.

Gneiss - MJS34



Gneiss - MJS35







Figure 7: Calculated pole figures of important mineral phases in the gneisses (**a**: MJS34) and (**b**: MJS35). The stretching lineation (kinematic x-direction) is oriented EW in the pole figures, the foliation normal (kinematic z-direction) is oriented NS and the kinematic y-direction (perpendicular to x- and z-directions) is normal to the pole figure plane. The pole figures are lower hemisphere equal area projections. The contour levels display multiples of random distribution and minimum and maximum values are listed below the pole figure.


1020

1025


Eclogite - MJS17




Blueschist - MJS16




Amphibolite - MJS26




Greenschist - MJS36




Dolomitic Micaschist - MJS18




Calcareous Micaschist - MJS16




Calcschist - MJS22




Gneiss - MJS34




Gneiss - MJS35





1030 **Figure 8:** Texture derived calculated P-wave velocity distributions of all samples in the study. All velocities in [km/s] with maximum and minimum values as well as P-wave anisotropy (in %) given below the pole figure.



1035

Figure 9: AVP [%] and VP/VS-ratios of all samples from this study, divided into the three categories of metabasic rocks, metasediment and gneisses.

Sample	Lithology and discription	Composition
MJS16	Strongly foliated blueschist	40 % barroisite, 30% albite, 10% almandine, 15% clinozoisite, 5% epidote 1040
MJS17	(U)HP eclogite with a clear foliation and lineation and large garnets	30% omphacite, 30% garnet, 15% clinozoisite, 10% glaucophane, 10% albite, 5% white mica
MJS18	Dolomitic micaschist with calcite and a strong mica-related foliation	60% dolomite, 20% quartz, 10% calcite, 10% muscovite
MJS20	Garnet-bearing calcareous micaschist	40% quartz, 30% muscovite, 15% calcite, 10% chlorite, 5% garnet 1045
MJS22	Well foliated calcschist	55% quartz, 30% calcite, 15% chlorite
MJS26	Weakly retrogressed amphibolite	40% hornblende, 25% albite, 20% clinozoisite, 10% chlorite, 5% garnet
MJS34	Weakly retrogressed gneiss	40% quartz, 35% albite, 15% muscovite, 10% chlorite
MJS35	Quartz-rich weakly retrogressed gneiss	57% quartz, 27% albite, 9% chlorite, 7% muscovite 1050
MJS36	Actinolite-bearing greenschist	30% chlorite, 30% clinozoisite, 20% albite, 20% actinolite

Table 1: Description and modal composition of investigated samples ~~in the study~~.

1055

Sample	Lithology	average VP [km/s]	min VP [km/s]	max VP [km/s]	VP-anisotropy [%]	average VS [km/s]	min VS1 [km/s]	max VS1 [km/s]	VS1-anisotropy [%]	min VS2 [km/s]	max VS2 [km/s]	VS2-anisotropy [%]	VP/VS
MJS16	Blueschist	7.63	7.53	7.81	3.7	4.47	4.49	4.52	0.7	4.42	4.49	1.6	1.70
MJS17	Eclogite	8.24	8.21	8.32	1.4	4.78	4.78	4.81	0.8	4.77	4.78	0.2	1.72
MJS18	Dolomitic micaschist	7.23	6.96	7.45	6.7	4.12	4.06	4.24	4.4	4.02	4.10	1.9	1.76
MJS20	Calcareous micaschist	6.77	6.51	6.99	7.1	4.16	4.05	4.31	6.1	4.02	4.18	3.9	1.63
MJS22	Calcschist	6.66	6.51	6.76	3.7	4.15	4.12	4.23	2.7	4.08	4.16	1.8	1.61
MJS26	Amphibolite	7.20	7.14	7.34	2.8	4.11	4.11	4.15	0.9	4.08	4.12	1.1	1.75
MJS34	Gneiss	6.62	6.55	6.69	2.1	4.14	4.10	4.18	2.0	4.07	4.16	2.0	1.60
MJS35	Gneiss	6.52	6.48	6.57	1.4	4.18	4.17	4.21	0.9	4.14	4.20	1.5	1.56
MJS36	Greenschist	7.30	7.16	7.43	3.7	4.28	4.25	4.35	2.4	4.22	4.29	1.6	1.70


 Formatierte Tabelle

Table 2: P-wave (VP) and S-wave (VS) velocities, **elastic seismic** anisotropies (**VP-, VS1- and VS2-anisotropies** %) and VP/VS-ratios of all samples, modelled from CPOs.

1060

<u>Study</u>	<u>AVP [%]</u>	<u>Average VP [km/s]</u>	<u>Average VS [km/s]</u>	<u>VP/VS</u>	<u>Method</u>
Eclogite					
<u>Our Study</u>	<u>1.4</u>	<u>8.24</u>	<u>4.78</u>	<u>1.72</u>	<u>Neutron diff.</u>
<u>Llana-Fúnez and Brown [2012]</u>	<u>0.6-5.0</u>	<u>n.a.</u>	<u>n.a.</u>	<u>n.a.</u>	<u>EBS</u>
<u>Keppeler et al. [2015]</u>	<u>8.3</u> <u>0.4-3.7</u>	<u>7.56</u> <u>7.40-8.27</u>	<u>n.a.</u> <u>4.23-4.81</u>	<u>n.a.</u> <u>1.70-1.77</u>	<u>-multi-anvil apparatus</u> <u>-Neutron diff.</u>
<u>Bezacier et al. [2010]</u>	<u>2.9</u>	<u>7.92</u>	<u>4.70</u>	<u>1.685</u>	<u>EBS</u>
<u>Wang et al. [2009]</u>	<u>0.6-1.7</u>	<u>8.50-8.53</u>	<u>n.a.</u>	<u>n.a.</u>	<u>pulse transmission technique (500 MPa /20 °C)</u>
<u>Cao et al. [2013]</u>	<u>1.4-10.2</u>	<u>7.73-8.28</u>	<u>4.45-4.74</u>	<u>1.706-1.751</u>	<u>EBS</u>
<u>Bascou et al. [2001]</u>	<u>1.2-2.9</u>	<u>n.a.</u>	<u>n.a.</u>	<u>n.a.</u>	<u>EBS</u>
<u>Zertani et al. [2019]</u>	<u>7-11</u>	<u>n.a.</u>	<u>n.a.</u>	<u>n.a.</u>	<u>-multi-anvil apparatus (600 MPa/600 °C)</u>
<u>Worthington et al. [2013]</u>	<u>1.1-2.8</u>	<u>8.0-8.5</u>	<u>4.5-4.8</u>	<u>n.a.</u>	<u>EBS</u>
<u>Cao et al. [2020]</u>	<u>0.99-2.46</u>	<u>8.00-8.33</u>	<u>4.55-4.72</u>	<u>1.758-1.770</u>	<u>EBS</u>
Blueschist					
<u>Our Study</u>	<u>3.7</u>	<u>7.63</u>	<u>4.47</u>	<u>1.70</u>	<u>Neutron diff.</u>
<u>Cao et al. [2013]</u>	<u>13.4-24.6</u>	<u>6.91-7.89</u>	<u>4.20-4.60</u>	<u>1.645-1.722</u>	<u>EBS</u>
<u>Bezacier et al. [2010]</u>	<u>16.1</u>	<u>7.71</u>	<u>4.49</u>	<u>1.717</u>	<u>EBS</u>
<u>Kim et al. [2013]</u>	<u>7.1-29.5</u>	<u>7.67</u>	<u>4.44</u>	<u>n.a.</u>	<u>Averages modelled from EBS</u>
<u>Fujimoto et al. [2010]</u>	<u>1.7-6.2</u> <u>n.a.</u>	<u>7.19</u> <u>6.0-7.4</u>	<u>n.a.</u> <u>n.a.</u>	<u>n.a.</u> <u>n.a.</u>	<u>-EBS</u> <u>-piston-cylinder apparatus (0GPa and 20°C)</u>
Amphibolite					
<u>Our Study</u>	<u>2.8</u>	<u>7.20</u>	<u>4.11</u>	<u>1.75</u>	<u>Neutron diff.</u>
<u>Siegesmund et al. [1989]</u>	<u>9.5</u>	<u>n.a.</u>	<u>n.a.</u>	<u>n.a.</u>	<u>multi-anvil apparatus</u>
<u>Barruol and Kern [1996]</u>	<u>5.69-12.74</u>	<u>6.92-7.02</u>	<u>3.93-3.94</u>	<u>1.76-1.78</u>	<u>multi-anvil apparatus</u>
<u>Ivankina et al. [2005]</u>	<u>0.86-8.27</u> <u>0.59-5.51</u>	<u>6.68-6.78</u> <u>6.73-6.77</u>	<u>n.a.</u> <u>n.a.</u>	<u>n.a.</u> <u>n.a.</u>	<u>-multi-anvil apparatus</u> <u>-Neutron diff.</u>

Ullemeyer et al. [2006]	4.4	6.87	n.a.	n.a.	-Neutron diff.
Keppler et al. [2015]	4.4 2.6	7.25 7.14	- 4.01	- 1.78	-multi-anvil apparatus -Neutron diff.
Kästner et al. [2020]	7.75-10.90	6.86-6.94	n.a.	n.a.	pulse-transmission method
Keppler et al. [2021]	2.1-4.5	6.7-6.94	3.81-3.88	1.76-1.79	Neutron diff.
Metasediments					
Our Study	3.7-7.1	6.66-7.23	4.12-4.16	1.61-1.76	Neutron diff.
Weiss et al. [1999]	4.56-6.53	n.a.	n.a.	n.a.	Calculated averages
Erdman et al. [2013]	2.3-11.4	6.02-7.26	3.50-4.05	1.72-1.88	EBSD
Keppler et al. [2015]	5.1-7.4	6.24-6.86	3.72-4.22	1.53-1.84	Neutron diff.
Kästner et al. [2020]	2.95-26.41	6.17-6.35	n.a.	n.a.	pulse-transmission method
Keppler et al. [2021]	7.9	7.22	3.96	1.82	Neutron diff.
Gneiss					
Our Study	1.4-2.1	6.62-6.52	4.14-4.18	1.65-1.60	Neutron diff.
Weiss et al. [1999]	1.76-8.73	n.a.	n.a.	n.a.	Calculated averages
Ivankina et al. [2005]	2.71-2.74 1.72-2.99	6.08-6.12 6.35-6.40	n.a. n.a.	n.a. n.a.	-multi-anvil apparatus -Neutron diff.
Ullemeyer et al. [2006]	0.9-3.2	6.28-6.55	n.a.	n.a.	-Neutron diff.
Kern et al. [2008]	15.1 7.9	n.a. n.a.	n.a. n.a.	n.a. n.a.	-multi-anvil apparatus -Neutron diff.
Erdman et al. [2013]	4.1-20.1	5.80-6.12	3.54-4.17	1.46-1.66	EBSD
Llana-Fúnez and Brown [2012]	8.0-11.3	n.a.	n.a.	n.a.	EBSD
Khazanchdari et al. [2000]	4.7-13.6	5.84-6.06	3.48-3.51	1.66-1.70	multi-anvil apparatus (550 MPa/20 °C)
Vasin et al. [2017]	9.6 8.4	n.a. n.a.	n.a. n.a.	n.a. n.a.	-multi-anvil apparatus -hydrostatic pressure-device (sphere)

Kästner et al. [2020]	3.63-14.56	5.99-6.81	n.a.	n.a.	pulse-transmission method
Keppler et al. [2021]	1.3-20.5	6.18-6.59	3.73-4.27	1.51-1.67	Neutron diff.

Table 3: AVP [%], average VP, average VS and VP/VS-ratios from literature in comparison to the data from this study.

1065

Sample	Phase	F2 index
MJS16	almandine	1.01
	albite	1.01
	barroisite	2.36
	clinozoisite	1.02
	epidote	1.02
MJS17	omphacite	3.49
	almandine	1.00
	glaucophane	1.19
	albite	1.02
	clinozoisite	1.01
	white mica	1.02
MJS18	quartz	1.04
	dolomite	1.14
	calcite	1.28
	muscovite	2.18
MJS20	quartz	1.14
	muscovite	2.83
	calcite	1.08
	chlorite	2.17
	almandine	1.02
MJS22	calcite	1.34
	quartz	1.54
	chlorite	1.52
MJS26	almandine	1.01
	hornblende	1.63
	albite	1.07
	clinozoisite	1.01
	chlorite	1.09
MJS34	quartz	1.18
	albite	1.03
	chlorite	2.15

	muscovite	1.27
MJS35	quartz	1.06
	muscovite	1.13
	albite	1.01
	chlorite	1.53
MJS36	albite	1.18
	actinolite	1.21
	clinozoisite	1.03
	chlorite	1.34

Appendix A: F2-indicies after Bunge (1982) for all mineral phases ~~in~~of the study.