

# Supplementary Material

## **Structure of Moho and mantle lithosphere in the greater Alpine area from S-to-P converted waves**

Rainer Kind<sup>1,2</sup>, Stefan M. Schmid<sup>3</sup>, Xiaohui Yuan<sup>2</sup>, Ben Heit<sup>2</sup>, Thomas Meier<sup>4</sup> and the AlpArray and AlpArray-SWATH-D Working Groups

<sup>1</sup> Freie Universität Berlin

<sup>2</sup> GeoForschungsZentrum Potsdam



<sup>3</sup> Eidgenössische Technische Hochschule Zürich

<sup>4</sup> Christian-Albrechts-Universität zu Kiel



Most seismic profiles in the main paper have used seismic traces within a bin of one degree latitude or longitude (depending on profile direction) perpendicular to the profile direction. The bin width in profile direction was in all cases  $0.2^\circ$  in latitude or longitude. The bins did not overlap. This leads to a high resolution in profile direction but a lower resolution perpendicular to the profile direction. The quality of the summation trace of all traces within a bin depends on the number of traces within one bin. Therefore the choice of the bin size depends on the number of available traces and one has to find a compromise between both. Of course, if the station density is higher and recording time longer, smaller bins resulting in higher resolution are possible. In the main paper we discarded bins with less than 200 traces. The purpose of the following figures is to show that also profile widths of half a degree still produce good results in within our network. The widths of the profiles are marked at the bottom of the figures. Here we accepted only bins with more than 80 traces. The signal-to-noise ratio is therefore reduced, but we think the results are still very useful.

The profiles on the left (A) use bins located at 50 km depth with theoretical piercing point depths there. The profiles on the right (B) are for piercing points at 100 km depth. That means the left side figures are approximately focused at the Moho, whereas the profiles on the right are more focused on the uppermost mantle. In some of the figures we have marked examples of Moho observations by a dashed black line. Examples of the lower boundary of the NVGs are marked by less clear defined grey lines. This is to point out that the onsets of the NVG (Negative Velocity Gradient) converted signal are usually much less correlated laterally than for example the Moho signals.



# **North-South Profiles**





**Fig. S1**





**Fig. S2**





**Fig. S3**





**Fig. S4**





**Fig. S5**



**Fig. S6**





**Fig. S7**





**Fig. S8**



# **East-West Profiles**





**Fig. S9**





**Fig. S10**





**Fig. S11**





**Fig. S12**





**Fig. S13**





**Fig. S14**



**Fig. S15**



**Fig. S16**



**Fig. S17**