

Deep vs shallow – two contrasting theories? A tectonically activated Late Cretaceous deltaic system in the axial part of the Mid-Polish Trough: a case study from southeast Poland

Zbyszek Remin¹, Michał Cyglicki¹⁺², Mariusz Niechwedowicz¹

5 ¹Faculty of Geology, University of Warsaw, Al. Żwirki i Wigury 93, PL-02-089 Warsaw, Poland; ²Polish Geological Institute – National Research Institute, Rakowiecka 4, PL-00-975 Warsaw, Poland; Corresponding Author: Zbyszek Remin; zremin@uw.edu.pl

Abstract:

10 The Polish Basin – a part of the trans-European sedimentary basin, i.e. Central European Basin System – stretching from Denmark through Germany to southeastern Poland, and even further to the south into Ukraine – experienced uplift during the Late Cretaceous, which consequently resulted in the inversion of its axial part (i.e. Mid-Polish Trough) and development into the Mid-Polish Anticlinorium. In many existing paleotectonic interpretations, southeast Poland – i.e., the subsurface San 15 Anticlinorium and the present-day Roztocze Hills – was included into the Mid-Polish Trough during the Late Cretaceous, representing its axial, most subsiding part. This paleotectonic model was the basis for facies and bathymetric interpretations that assumed Upper Cretaceous sediments, deposited close to the axial part of the Mid-Polish Trough (e.g., in the Roztocze Hills), constituted the deepest facies. In recent years, several studies have contradicted this notion. A growing body of data suggests that the 20 this area (southeast Poland) was already a landmass by the Coniacian and Santonian – and certainly the Campanian and Maastrichtian – rather than forming the deepest part of the Polish Basin.

The shallow marginal marine, cyclic middle Campanian, marly to sandy deposits, recently discovered in the Roztocze Hills, were interpreted to be of deltaic origin. It has led to the adoption of new facies and bathymetric model, which – notably – are in stark opposition to most previous interpretations. 25 Notably, the new interpretation implies the presence of a landmass in the place where, in prior frameworks, the deepest and most subsiding part of the Polish Basin (i.e. Mid-Polish Trough) was located during the Cretaceous.

Here, we document this Late Cretaceous depositional system – the Szozdy delta developed in the axial 30 part of the Mid-Polish Trough. These middle Campanian deposits crop out extensively in the Roztocze Hills region, close to the village of Szozdy, and exhibit coarsening-upward tripartite cycloths. Three facies associations have been distinguished: (1) dark grey calcareous mudstones, (2) a yellow 35 calcareous sandstone, and (3) calcareous gaize lithofacies. The sequence, as a whole, was accumulated via the repeated progradation and abandonment of deltaic complexes on the delta front platform setting, with the primary transport direction originating from the axis of the inverting Mid-Polish Trough (thus from the subsurface San Anticlinorium) toward the northeast. This interpretation is supported by a suite of sedimentological, palynofacies, and heavy mineral data.

The development of the Szozdy delta system is framed in the context of the dynamic tectonic processes 40 operating contemporaneously in southeast Poland: that is, tectonic inversion (uplift) on one hand, and the generation of new accommodation space via enhanced subsidence on the other. This discovery

sheds new light on our understanding of Late Cretaceous facies distribution, bathymetry, paleogeography, and potentially might suggest a different burial history than thought so far.



Key words: Late Cretaceous, inversion tectonics, Szozdy delta, sedimentology, palynofacies, heavy minerals, provenance

1. Introduction

During the Late Cretaceous, the central and western Europe has been affected by compressional tectonics, which resulted in ~~the~~ ¹ ~~the~~ ² ~~the~~ ³ ~~the~~ ⁴ ~~the~~ ⁵ ~~the~~ ⁶ ~~the~~ ⁷ ~~the~~ ⁸ ~~the~~ ⁹ ~~the~~ ¹⁰ ~~the~~ ¹¹ ~~the~~ ¹² ~~the~~ ¹³ ~~the~~ ¹⁴ ~~the~~ ¹⁵ ~~the~~ ¹⁶ ~~the~~ ¹⁷ ~~the~~ ¹⁸ ~~the~~ ¹⁹ ~~the~~ ²⁰ ~~the~~ ²¹ ~~the~~ ²² ~~the~~ ²³ ~~the~~ ²⁴ ~~the~~ ²⁵ ~~the~~ ²⁶ ~~the~~ ²⁷ ~~the~~ ²⁸ ~~the~~ ²⁹ ~~the~~ ³⁰ ~~the~~ ³¹ ~~the~~ ³² ~~the~~ ³³ ~~the~~ ³⁴ ~~the~~ ³⁵ ~~the~~ ³⁶ ~~the~~ ³⁷ ~~the~~ ³⁸ ~~the~~ ³⁹ ~~the~~ ⁴⁰ ~~the~~ ⁴¹ ~~the~~ ⁴² ~~the~~ ⁴³ ~~the~~ ⁴⁴ ~~the~~ ⁴⁵ ~~the~~ ⁴⁶ ~~the~~ ⁴⁷ ~~the~~ ⁴⁸ ~~the~~ ⁴⁹ ~~the~~ ⁵⁰ ~~the~~ ⁵¹ ~~the~~ ⁵² ~~the~~ ⁵³ ~~the~~ ⁵⁴ ~~the~~ ⁵⁵ ~~the~~ ⁵⁶ ~~the~~ ⁵⁷ ~~the~~ ⁵⁸ ~~the~~ ⁵⁹ ~~the~~ ⁶⁰ ~~the~~ ⁶¹ ~~the~~ ⁶² ~~the~~ ⁶³ ~~the~~ ⁶⁴ ~~the~~ ⁶⁵ ~~the~~ ⁶⁶ ~~the~~ ⁶⁷ ~~the~~ ⁶⁸ ~~the~~ ⁶⁹ ~~the~~ ⁷⁰ ~~the~~ ⁷¹ ~~the~~ ⁷² ~~the~~ ⁷³ ~~the~~ ⁷⁴ ~~the~~ ⁷⁵ ~~the~~ ⁷⁶ ~~the~~ ⁷⁷ ~~the~~ ⁷⁸ ~~the~~ ⁷⁹ ~~the~~ ⁸⁰ ~~the~~ ⁸¹ ~~the~~ ⁸² ~~the~~ ⁸³ ~~the~~ ⁸⁴ ~~the~~ ⁸⁵ ~~the~~ ⁸⁶ ~~the~~ ⁸⁷ ~~the~~ ⁸⁸ ~~the~~ ⁸⁹ ~~the~~ ⁹⁰ ~~the~~ ⁹¹ ~~the~~ ⁹² ~~the~~ ⁹³ ~~the~~ ⁹⁴ ~~the~~ ⁹⁵ ~~the~~ ⁹⁶ ~~the~~ ⁹⁷ ~~the~~ ⁹⁸ ~~the~~ ⁹⁹ ~~the~~ ¹⁰⁰ ~~the~~ ¹⁰¹ ~~the~~ ¹⁰² ~~the~~ ¹⁰³ ~~the~~ ¹⁰⁴ ~~the~~ ¹⁰⁵ ~~the~~ ¹⁰⁶ ~~the~~ ¹⁰⁷ ~~the~~ ¹⁰⁸ ~~the~~ ¹⁰⁹ ~~the~~ ¹¹⁰ ~~the~~ ¹¹¹ ~~the~~ ¹¹² ~~the~~ ¹¹³ ~~the~~ ¹¹⁴ ~~the~~ ¹¹⁵ ~~the~~ ¹¹⁶ ~~the~~ ¹¹⁷ ~~the~~ ¹¹⁸ ~~the~~ ¹¹⁹ ~~the~~ ¹²⁰ ~~the~~ ¹²¹ ~~the~~ ¹²² ~~the~~ ¹²³ ~~the~~ ¹²⁴ ~~the~~ ¹²⁵ ~~the~~ ¹²⁶ ~~the~~ ¹²⁷ ~~the~~ ¹²⁸ ~~the~~ ¹²⁹ ~~the~~ ¹³⁰ ~~the~~ ¹³¹ ~~the~~ ¹³² ~~the~~ ¹³³ ~~the~~ ¹³⁴ ~~the~~ ¹³⁵ ~~the~~ ¹³⁶ ~~the~~ ¹³⁷ ~~the~~ ¹³⁸ ~~the~~ ¹³⁹ ~~the~~ ¹⁴⁰ ~~the~~ ¹⁴¹ ~~the~~ ¹⁴² ~~the~~ ¹⁴³ ~~the~~ ¹⁴⁴ ~~the~~ ¹⁴⁵ ~~the~~ ¹⁴⁶ ~~the~~ ¹⁴⁷ ~~the~~ ¹⁴⁸ ~~the~~ ¹⁴⁹ ~~the~~ ¹⁵⁰ ~~the~~ ¹⁵¹ ~~the~~ ¹⁵² ~~the~~ ¹⁵³ ~~the~~ ¹⁵⁴ ~~the~~ ¹⁵⁵ ~~the~~ ¹⁵⁶ ~~the~~ ¹⁵⁷ ~~the~~ ¹⁵⁸ ~~the~~ ¹⁵⁹ ~~the~~ ¹⁶⁰ ~~the~~ ¹⁶¹ ~~the~~ ¹⁶² ~~the~~ ¹⁶³ ~~the~~ ¹⁶⁴ ~~the~~ ¹⁶⁵ ~~the~~ ¹⁶⁶ ~~the~~ ¹⁶⁷ ~~the~~ ¹⁶⁸ ~~the~~ ¹⁶⁹ ~~the~~ ¹⁷⁰ ~~the~~ ¹⁷¹ ~~the~~ ¹⁷² ~~the~~ ¹⁷³ ~~the~~ ¹⁷⁴ ~~the~~ ¹⁷⁵ ~~the~~ ¹⁷⁶ ~~the~~ ¹⁷⁷ ~~the~~ ¹⁷⁸ ~~the~~ ¹⁷⁹ ~~the~~ ¹⁸⁰ ~~the~~ ¹⁸¹ ~~the~~ ¹⁸² ~~the~~ ¹⁸³ ~~the~~ ¹⁸⁴ ~~the~~ ¹⁸⁵ ~~the~~ ¹⁸⁶ ~~the~~ ¹⁸⁷ ~~the~~ ¹⁸⁸ ~~the~~ ¹⁸⁹ ~~the~~ ¹⁹⁰ ~~the~~ ¹⁹¹ ~~the~~ ¹⁹² ~~the~~ ¹⁹³ ~~the~~ ¹⁹⁴ ~~the~~ ¹⁹⁵ ~~the~~ ¹⁹⁶ ~~the~~ ¹⁹⁷ ~~the~~ ¹⁹⁸ ~~the~~ ¹⁹⁹ ~~the~~ ²⁰⁰ ~~the~~ ²⁰¹ ~~the~~ ²⁰² ~~the~~ ²⁰³ ~~the~~ ²⁰⁴ ~~the~~ ²⁰⁵ ~~the~~ ²⁰⁶ ~~the~~ ²⁰⁷ ~~the~~ ²⁰⁸ ~~the~~ ²⁰⁹ ~~the~~ ²¹⁰ ~~the~~ ²¹¹ ~~the~~ ²¹² ~~the~~ ²¹³ ~~the~~ ²¹⁴ ~~the~~ ²¹⁵ ~~the~~ ²¹⁶ ~~the~~ ²¹⁷ ~~the~~ ²¹⁸ ~~the~~ ²¹⁹ ~~the~~ ²²⁰ ~~the~~ ²²¹ ~~the~~ ²²² ~~the~~ ²²³ ~~the~~ ²²⁴ ~~the~~ ²²⁵ ~~the~~ ²²⁶ ~~the~~ ²²⁷ ~~the~~ ²²⁸ ~~the~~ ²²⁹ ~~the~~ ²³⁰ ~~the~~ ²³¹ ~~the~~ ²³² ~~the~~ ²³³ ~~the~~ ²³⁴ ~~the~~ ²³⁵ ~~the~~ ²³⁶ ~~the~~ ²³⁷ ~~the~~ ²³⁸ ~~the~~ ²³⁹ ~~the~~ ²⁴⁰ ~~the~~ ²⁴¹ ~~the~~ ²⁴² ~~the~~ ²⁴³ ~~the~~ ²⁴⁴ ~~the~~ ²⁴⁵ ~~the~~ ²⁴⁶ ~~the~~ ²⁴⁷ ~~the~~ ²⁴⁸ ~~the~~ ²⁴⁹ ~~the~~ ²⁵⁰ ~~the~~ ²⁵¹ ~~the~~ ²⁵² ~~the~~ ²⁵³ ~~the~~ ²⁵⁴ ~~the~~ ²⁵⁵ ~~the~~ ²⁵⁶ ~~the~~ ²⁵⁷ ~~the~~ ²⁵⁸ ~~the~~ ²⁵⁹ ~~the~~ ²⁶⁰ ~~the~~ ²⁶¹ ~~the~~ ²⁶² ~~the~~ ²⁶³ ~~the~~ ²⁶⁴ ~~the~~ ²⁶⁵ ~~the~~ ²⁶⁶ ~~the~~ ²⁶⁷ ~~the~~ ²⁶⁸ ~~the~~ ²⁶⁹ ~~the~~ ²⁷⁰ ~~the~~ ²⁷¹ ~~the~~ ²⁷² ~~the~~ ²⁷³ ~~the~~ ²⁷⁴ ~~the~~ ²⁷⁵ ~~the~~ ²⁷⁶ ~~the~~ ²⁷⁷ ~~the~~ ²⁷⁸ ~~the~~ ²⁷⁹ ~~the~~ ²⁸⁰ ~~the~~ ²⁸¹ ~~the~~ ²⁸² ~~the~~ ²⁸³ ~~the~~ ²⁸⁴ ~~the~~ ²⁸⁵ ~~the~~ ²⁸⁶ ~~the~~ ²⁸⁷ ~~the~~ ²⁸⁸ ~~the~~ ²⁸⁹ ~~the~~ ²⁹⁰ ~~the~~ ²⁹¹ ~~the~~ ²⁹² ~~the~~ ²⁹³ ~~the~~ ²⁹⁴ ~~the~~ ²⁹⁵ ~~the~~ ²⁹⁶ ~~the~~ ²⁹⁷ ~~the~~ ²⁹⁸ ~~the~~ ²⁹⁹ ~~the~~ ³⁰⁰ ~~the~~ ³⁰¹ ~~the~~ ³⁰² ~~the~~ ³⁰³ ~~the~~ ³⁰⁴ ~~the~~ ³⁰⁵ ~~the~~ ³⁰⁶ ~~the~~ ³⁰⁷ ~~the~~ ³⁰⁸ ~~the~~ ³⁰⁹ ~~the~~ ³¹⁰ ~~the~~ ³¹¹ ~~the~~ ³¹² ~~the~~ ³¹³ ~~the~~ ³¹⁴ ~~the~~ ³¹⁵ ~~the~~ ³¹⁶ ~~the~~ ³¹⁷ ~~the~~ ³¹⁸ ~~the~~ ³¹⁹ ~~the~~ ³²⁰ ~~the~~ ³²¹ ~~the~~ ³²² ~~the~~ ³²³ ~~the~~ ³²⁴ ~~the~~ ³²⁵ ~~the~~ ³²⁶ ~~the~~ ³²⁷ ~~the~~ ³²⁸ ~~the~~ ³²⁹ ~~the~~ ³³⁰ ~~the~~ ³³¹ ~~the~~ ³³² ~~the~~ ³³³ ~~the~~ ³³⁴ ~~the~~ ³³⁵ ~~the~~ ³³⁶ ~~the~~ ³³⁷ ~~the~~ ³³⁸ ~~the~~ ³³⁹ ~~the~~ ³⁴⁰ ~~the~~ ³⁴¹ ~~the~~ ³⁴² ~~the~~ ³⁴³ ~~the~~ ³⁴⁴ ~~the~~ ³⁴⁵ ~~the~~ ³⁴⁶ ~~the~~ ³⁴⁷ ~~the~~ ³⁴⁸ ~~the~~ ³⁴⁹ ~~the~~ ³⁵⁰ ~~the~~ ³⁵¹ ~~the~~ ³⁵² ~~the~~ ³⁵³ ~~the~~ ³⁵⁴ ~~the~~ ³⁵⁵ ~~the~~ ³⁵⁶ ~~the~~ ³⁵⁷ ~~the~~ ³⁵⁸ ~~the~~ ³⁵⁹ ~~the~~ ³⁶⁰ ~~the~~ ³⁶¹ ~~the~~ ³⁶² ~~the~~ ³⁶³ ~~the~~ ³⁶⁴ ~~the~~ ³⁶⁵ ~~the~~ ³⁶⁶ ~~the~~ ³⁶⁷ ~~the~~ ³⁶⁸ ~~the~~ ³⁶⁹ ~~the~~ ³⁷⁰ ~~the~~ ³⁷¹ ~~the~~ ³⁷² ~~the~~ ³⁷³ ~~the~~ ³⁷⁴ ~~the~~ ³⁷⁵ ~~the~~ ³⁷⁶ ~~the~~ ³⁷⁷ ~~the~~ ³⁷⁸ ~~the~~ ³⁷⁹ ~~the~~ ³⁸⁰ ~~the~~ ³⁸¹ ~~the~~ ³⁸² ~~the~~ ³⁸³ ~~the~~ ³⁸⁴ ~~the~~ ³⁸⁵ ~~the~~ ³⁸⁶ ~~the~~ ³⁸⁷ ~~the~~ ³⁸⁸ ~~the~~ ³⁸⁹ ~~the~~ ³⁹⁰ ~~the~~ ³⁹¹ ~~the~~ ³⁹² ~~the~~ ³⁹³ ~~the~~ ³⁹⁴ ~~the~~ ³⁹⁵ ~~the~~ ³⁹⁶ ~~the~~ ³⁹⁷ ~~the~~ ³⁹⁸ ~~the~~ ³⁹⁹ ~~the~~ ⁴⁰⁰ ~~the~~ ⁴⁰¹ ~~the~~ ⁴⁰² ~~the~~ ⁴⁰³ ~~the~~ ⁴⁰⁴ ~~the~~ ⁴⁰⁵ ~~the~~ ⁴⁰⁶ ~~the~~ ⁴⁰⁷ ~~the~~ ⁴⁰⁸ ~~the~~ ⁴⁰⁹ ~~the~~ ⁴¹⁰ ~~the~~ ⁴¹¹ ~~the~~ ⁴¹² ~~the~~ ⁴¹³ ~~the~~ ⁴¹⁴ ~~the~~ ⁴¹⁵ ~~the~~ ⁴¹⁶ ~~the~~ ⁴¹⁷ ~~the~~ ⁴¹⁸ ~~the~~ ⁴¹⁹ ~~the~~ ⁴²⁰ ~~the~~ ⁴²¹ ~~the~~ ⁴²² ~~the~~ ⁴²³ ~~the~~ ⁴²⁴ ~~the~~ ⁴²⁵ ~~the~~ ⁴²⁶ ~~the~~ ⁴²⁷ ~~the~~ ⁴²⁸ ~~the~~ ⁴²⁹ ~~the~~ ⁴³⁰ ~~the~~ ⁴³¹ ~~the~~ ⁴³² ~~the~~ ⁴³³ ~~the~~ ⁴³⁴ ~~the~~ ⁴³⁵ ~~the~~ ⁴³⁶ ~~the~~ ⁴³⁷ ~~the~~ ⁴³⁸ ~~the~~ ⁴³⁹ ~~the~~ ⁴⁴⁰ ~~the~~ ⁴⁴¹ ~~the~~ ⁴⁴² ~~the~~ ⁴⁴³ ~~the~~ ⁴⁴⁴ ~~the~~ ⁴⁴⁵ ~~the~~ ⁴⁴⁶ ~~the~~ ⁴⁴⁷ ~~the~~ ⁴⁴⁸ ~~the~~ ⁴⁴⁹ ~~the~~ ⁴⁵⁰ ~~the~~ ⁴⁵¹ ~~the~~ ⁴⁵² ~~the~~ ⁴⁵³ ~~the~~ ⁴⁵⁴ ~~the~~ ⁴⁵⁵ ~~the~~ ⁴⁵⁶ ~~the~~ ⁴⁵⁷ ~~the~~ ⁴⁵⁸ ~~the~~ ⁴⁵⁹ ~~the~~ ⁴⁶⁰ ~~the~~ ⁴⁶¹ ~~the~~ ⁴⁶² ~~the~~ ⁴⁶³ ~~the~~ ⁴⁶⁴ ~~the~~ ⁴⁶⁵ ~~the~~ ⁴⁶⁶ ~~the~~ ⁴⁶⁷ ~~the~~ ⁴⁶⁸ ~~the~~ ⁴⁶⁹ ~~the~~ ⁴⁷⁰ ~~the~~ ⁴⁷¹ ~~the~~ ⁴⁷² ~~the~~ ⁴⁷³ ~~the~~ ⁴⁷⁴ ~~the~~ ⁴⁷⁵ ~~the~~ ⁴⁷⁶ ~~the~~ ⁴⁷⁷ ~~the~~ ⁴⁷⁸ ~~the~~ ⁴⁷⁹ ~~the~~ ⁴⁸⁰ ~~the~~ ⁴⁸¹ ~~the~~ ⁴⁸² ~~the~~ ⁴⁸³ ~~the~~ ⁴⁸⁴ ~~the~~ ⁴⁸⁵ ~~the~~ ⁴⁸⁶ ~~the~~ ⁴⁸⁷ ~~the~~ ⁴⁸⁸ ~~the~~ ⁴⁸⁹ ~~the~~ ⁴⁹⁰ ~~the~~ ⁴⁹¹ ~~the~~ ⁴⁹² ~~the~~ ⁴⁹³ ~~the~~ ⁴⁹⁴ ~~the~~ ⁴⁹⁵ ~~the~~ ⁴⁹⁶ ~~the~~ ⁴⁹⁷ ~~the~~ ⁴⁹⁸ ~~the~~ ⁴⁹⁹ ~~the~~ ⁵⁰⁰ ~~the~~ ⁵⁰¹ ~~the~~ ⁵⁰² ~~the~~ ⁵⁰³ ~~the~~ ⁵⁰⁴ ~~the~~ ⁵⁰⁵ ~~the~~ ⁵⁰⁶ ~~the~~ ⁵⁰⁷ ~~the~~ ⁵⁰⁸ ~~the~~ ⁵⁰⁹ ~~the~~ ⁵¹⁰ ~~the~~ ⁵¹¹ ~~the~~ ⁵¹² ~~the~~ ⁵¹³ ~~the~~ ⁵¹⁴ ~~the~~ ⁵¹⁵ ~~the~~ ⁵¹⁶ ~~the~~ ⁵¹⁷ ~~the~~ ⁵¹⁸ ~~the~~ ⁵¹⁹ ~~the~~ ⁵²⁰ ~~the~~ ⁵²¹ ~~the~~ ⁵²² ~~the~~ ⁵²³ ~~the~~ ⁵²⁴ ~~the~~ ⁵²⁵ ~~the~~ ⁵²⁶ ~~the~~ ⁵²⁷ ~~the~~ ⁵²⁸ ~~the~~ ⁵²⁹ ~~the~~ ⁵³⁰ ~~the~~ ⁵³¹ ~~the~~ ⁵³² ~~the~~ ⁵³³ ~~the~~ ⁵³⁴ ~~the~~ ⁵³⁵ ~~the~~ ⁵³⁶ ~~the~~ ⁵³⁷ ~~the~~ ⁵³⁸ ~~the~~ ⁵³⁹ ~~the~~ ⁵⁴⁰ ~~the~~ ⁵⁴¹ ~~the~~ ⁵⁴² ~~the~~ ⁵⁴³ ~~the~~ ⁵⁴⁴ ~~the~~ ⁵⁴⁵ ~~the~~ ⁵⁴⁶ ~~the~~ ⁵⁴⁷ ~~the~~ ⁵⁴⁸ ~~the~~ ⁵⁴⁹ ~~the~~ ⁵⁵⁰ ~~the~~ ⁵⁵¹ ~~the~~ ⁵⁵² ~~the~~ ⁵⁵³ ~~the~~ ⁵⁵⁴ ~~the~~ ⁵⁵⁵ ~~the~~ ⁵⁵⁶ ~~the~~ ⁵⁵⁷ ~~the~~ ⁵⁵⁸ ~~the~~ ⁵⁵⁹ ~~the~~ ⁵⁶⁰ ~~the~~ ⁵⁶¹ ~~the~~ ⁵⁶² ~~the~~ ⁵⁶³ ~~the~~ ⁵⁶⁴ ~~the~~ ⁵⁶⁵ ~~the~~ ⁵⁶⁶ ~~the~~ ⁵⁶⁷ ~~the~~ ⁵⁶⁸ ~~the~~ ⁵⁶⁹ ~~the~~ ⁵⁷⁰ ~~the~~ ⁵⁷¹ ~~the~~ ⁵⁷² ~~the~~ ⁵⁷³ ~~the~~ ⁵⁷⁴ ~~the~~ ⁵⁷⁵ ~~the~~ ⁵⁷⁶ ~~the~~ ⁵⁷⁷ ~~the~~ ⁵⁷⁸ ~~the~~ ⁵⁷⁹ ~~the~~ ⁵⁸⁰ ~~the~~ ⁵⁸¹ ~~the~~ ⁵⁸² ~~the~~ ⁵⁸³ ~~the~~ ⁵⁸⁴ ~~the~~ ⁵⁸⁵ ~~the~~ ⁵⁸⁶ ~~the~~ ⁵⁸⁷ ~~the~~ ⁵⁸⁸ ~~the~~ ⁵⁸⁹ ~~the~~ ⁵⁹⁰ ~~the~~ ⁵⁹¹ ~~the~~ ⁵⁹² ~~the~~ ⁵⁹³ ~~the~~ ⁵⁹⁴ ~~the~~ ⁵⁹⁵ ~~the~~ ⁵⁹⁶ ~~the~~ ⁵⁹⁷ ~~the~~ ⁵⁹⁸ ~~the~~ ⁵⁹⁹ ~~the~~ ⁶⁰⁰ ~~the~~ ⁶⁰¹ ~~the~~ ⁶⁰² ~~the~~ ⁶⁰³ ~~the~~ ⁶⁰⁴ ~~the~~ ⁶⁰⁵ ~~the~~ ⁶⁰⁶ ~~the~~ ⁶⁰⁷ ~~the~~ ⁶⁰⁸ ~~the~~ ⁶⁰⁹ ~~the~~ ⁶¹⁰ ~~the~~ ⁶¹¹ ~~the~~ ⁶¹² ~~the~~ ⁶¹³ ~~the~~ ⁶¹⁴ ~~the~~ ⁶¹⁵ ~~the~~ ⁶¹⁶ ~~the~~ ⁶¹⁷ ~~the~~ ⁶¹⁸ ~~the~~ ⁶¹⁹ ~~the~~ ⁶²⁰ ~~the~~ ⁶²¹ ~~the~~ ⁶²² ~~the~~ ⁶²³ ~~the~~ ⁶²⁴ ~~the~~ ⁶²⁵ ~~the~~ ⁶²⁶ ~~the~~ ⁶²⁷ ~~the~~ ⁶²⁸ ~~the~~ ⁶²⁹ ~~the~~ ⁶³⁰ ~~the~~ ⁶³¹ ~~the~~ ⁶³² ~~the~~ ⁶³³ ~~the~~ ⁶³⁴ ~~the~~ ⁶³⁵ ~~the~~ ⁶³⁶ ~~the~~ ⁶³⁷ ~~the~~ ⁶³⁸ ~~the~~ ⁶³⁹ ~~the~~ ⁶⁴⁰ ~~the~~ ⁶⁴¹ ~~the~~ ⁶⁴² ~~the~~ ⁶⁴³ ~~the~~ ⁶⁴⁴ ~~the~~ ⁶⁴⁵ ~~the~~ ⁶⁴⁶ ~~the~~ ⁶⁴⁷ ~~the~~ ⁶⁴⁸ ~~the~~ ⁶⁴⁹ ~~the~~ ⁶⁵⁰ ~~the~~ ⁶⁵¹ ~~the~~ ⁶⁵² ~~the~~ ⁶⁵³ ~~the~~ ⁶⁵⁴ ~~the~~ ⁶⁵⁵ ~~the~~ ⁶⁵⁶ ~~the~~ ⁶⁵⁷ ~~the~~ ⁶⁵⁸ ~~the~~ ⁶⁵⁹ ~~the~~ ⁶⁶⁰ ~~the~~ ⁶⁶¹ ~~the~~ ⁶⁶² ~~the~~ ⁶⁶³ ~~the~~ ⁶⁶⁴ ~~the~~ ⁶⁶⁵ ~~the~~ ⁶⁶⁶ ~~the~~ ⁶⁶⁷ ~~the~~ ⁶⁶⁸ ~~the~~ ⁶⁶⁹ ~~the~~ ⁶⁷⁰ ~~the~~ ⁶⁷¹ ~~the~~ ⁶⁷² ~~the~~ ⁶⁷³ ~~the~~ ⁶⁷⁴ ~~the~~ ⁶⁷⁵ ~~the~~ ⁶⁷⁶ ~~the~~ ⁶⁷⁷ ~~the~~ ⁶⁷⁸ ~~the~~ ⁶⁷⁹ ~~the~~ ⁶⁸⁰ ~~the~~ ⁶⁸¹ ~~the~~ ⁶⁸² ~~the~~ ⁶⁸³ ~~the~~ ⁶⁸⁴ ~~the~~ ⁶⁸⁵ ~~the~~ ⁶⁸⁶ ~~the~~ ⁶⁸⁷ ~~the~~ ⁶⁸⁸ ~~the~~ ⁶⁸⁹ ~~the</del~~

the Mid-Polish Trough), like those in the Roztocze Hills (southeast Poland) would constitute the deepest facies (Fig. 2c).

85 A series of studies performed over the past few years have supplied data in stark contradiction to the above interpretation, showing that already by the Coniacian and Santonian (and possibly even by the Late Turonian) the axial part of the Polish Basin (i.e. the Mid-Polish Trough) should be considered an emergent landmass or paleomorphological barrier, rather than as the deepest part of the basin, especially in the southeastern Poland (e.g. Krzywiec et al., 2009; Remin et al., 2015a; Walaszczyk and Remin, 2015; Remin et al., 2016; Krzywiec et al., 2018; Remin, 2018). **This view is supported by the presence of clearly shallow-water facies along the northeastern edge of the present-day Mid-Polish Anticlinorium, especially in the southeastern Poland, thus in area where deepest facies should be expected according to the model of Kutek and Głażek (1972)** **These are the shallow water Turonian Janików Limestone and deltaically influenced sedimentation of the Roztocze Hills formed during the Campanian and Maastrichtian (Remin et al., 2015a; Walaszczyk and Remin, 2015).** Additionally, seismic data show progradational bodies trending from the SW toward the NE (Krzywiec et al., 2009, 2018), thus extending from the supposedly axial part of the basin toward the northeast. However, all these data did not provide definitive proof for the existence of an emerged landmass in southeast Poland during the Late Cretaceous, and when treated individually their value and insights might be undermined.

100 In 2015a, Remin et al. provided a conceptual interpretation that the middle Campanian and Maastrichtian siliciclastic deposits of the Roztocze Hills were of deltaic origin with the main source area located to the SW: this in a place where, according to the widely accepted paleotectonic model of Kutek and Głażek (1972) and several later authors, the axial and deepest part of the Mid-Polish Trough was located (Fig. 2c). The implementation of the new concept suggested a need for a dramatic revision to the existing paleotectonic model of southeastern part of the Polish Basin, and drove the adoption of a new facies and bathymetric model for several Late Cretaceous facies, one that was essentially the opposite of most previous interpretations (compare Figs. 2c and 2d; and discussion in: Krzywiec et al., 2009; Remin et al., 2015a; Walaszczyk and Remin, 2015; Krzywiec et al., 2018). Since this theoretical interpretation lacked precise, stratigraphically and sedimentologically-grounded corroboration, the present paper fills this gap by providing evidence to a wider audience that the development of the Szozdy deltaic system was induced by active inversion tectonics.

115 The study area in the Roztocze Hills gives direct insight into this special Campanian and Maastrichtian sedimentary succession, thus allowing for precise bio- and chronostratigraphic dating, essential for detailed sedimentological, facies, and paleotectonic interpretations.

120 These new datasets provide the foundation for the first direct sedimentological, petrographic, and palynofacies evidence for the presence of an emerged landmass in an area now devoid of Mesozoic cover – that is, the subsurface San Anticlinorium (the SE part of the Mid-Polish Anticlinorium; Fig. 1b) – which had previously been considered to be deeply submerged during the Late Cretaceous. These data, at least in part confirm the earlier concepts of e.g. Krzywiec et al. (2009, 2018) based on subsurface seismic data.

The objective of this paper is to show how a dynamic tectonic regime, including spatially variable inversion processes (uplift) and rapid subsidence, coupled with possible wave and/or bottom current

125 transport processes and significant terrigenous sediment input, drove the development of the Szozdy
deltaic system. Its subaqueous remnants, originally developed in the axial part of the Mid-Polish Trough,
are preserved in the rock record of the Roztocze Hills.

130




Figure 1. (a) Regional schematic map of western and central Europe (after Krzywiec et al., 2018 with further references) with the schematic extent of the Late Cretaceous - Paleocene inversion (patterned

area); red rectangle – area shown in (b) and in figure 2; (b) General position of the Roztocze Hills and San Anticlinorium in the geological map of Poland excluding Cenozoic deposits, adopted from Dadlez et al. (2000) – except for the recent-day extension of the Carpathian Foredeep; MVVS = Middle Vistula Valley Section.




Figure 2. (a) A ~~higher-resolution~~ view of the Roztocze Hills area, relative to the position of the Holy Cross Mountain  and Lower San Anticlinorium; (b) The general distribution of Campanian lithofacies in southeast Poland (adopted from Świdrowska, 2007): sandy limestone = deep; chalk = shallow; red dashed line – an interpreted cross-section of two competing tectonic concepts (c) and (d); (c) A facies and bathymetric sedimentation model based on traditional interpretations, assuming the deep-water character of deposits close to the axial part of the Mid-Polish Trough; vertical dashed-lines indicate areas devoid of Cretaceous deposits at present; (d) An alternative depositional, structural, and environmental interpretation of facies and bathymetry in a cross-section perpendicular to the axis of the Mid-Polish Anticlinorium (adopted from Remin et al., 2015a; Walaszczuk and Remin, 2015).

150 2. The current state of the art: a short story of two contrasting theories

Broadly speaking, the current literature hosts two theories regarding the onset of Mid-Polish Trough inversion tectonics and, subsequently, the formation of the Mid Polish-Anticlinorium (for a thorough review, see: Krzywiec et al., 2009; Walaszczyk and Remin, 2015; Krzywiec et al., 2018).

155 The first – and more widely accepted, at least historically – concept holds that the inversion of the Mid-Polish Trough began during the Late Maastrichtian and Paleogene (e.g., Kutek and Głazek, 1972; Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska, 2007; Świdrowska et al., 2008). Kutek and Głazek's (1972) paper was indisputably a benchmark for interpretations of the Late Cretaceous paleogeographic and paleotectonic evolution of southeastern segment of the Polish Basin, and especially its axial part, i.e. Mid-Polish Trough.

160 Kutek and Głazek (1972) provided palinspastic restoration of a paleothickness of particular stratigraphic units in the cross-sections to the Mid-Polish Anticlinorium in the Holy Cross Mountains (to the NW from the present study area – compare Fig. 1b and 2a). In their framework, the increase in thickness of Mesozoic sedimentary successions toward the present-day Mid-Polish Anticlinorium (the area devoid of Mesozoic cover), (Fig. 1b and 2a), constituted their key argument for highest burial and 165 the presence of a depocentre in the axial part of the Mid-Polish Trough, at least until the early Maastrichtian. This also implies that the area currently devoid of Mesozoic remnants was deeply buried by those deposits.

170 The general framework of Kutek and Głazek (1972) was widely accepted, and for decades dominated thinking about the Mesozoic paleotectonic evolution of the Polish Basin. In their model, the southeastern part of the Polish Basin, i.e. Mid-Polish Trough would represent the axial, most subsiding part of the Mid-Polish Trough during the Late Cretaceous (compare Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska, 2007; Świdrowska et al., 2008). The acceptance of this paleotectonic model has some foundational implications: for instance, it was widely assumed that the facies developed within the Mid-Polish Trough would represent the deepest sedimentary environment (e.g., Kutek and Głazek, 1972; Hakenberg and Świdrowska, 1998; Świdrowska 2007). As 175 such, the facies and bathymetric interpretation was evidently fit to the presumptive paleotectonic model, rather than *vice versa*. It is noteworthy, however, that Upper Cretaceous mixed carbonate-siliceous facies elude easy environmental interpretation, and the adopted paleobathymetric model and the spatial distribution of particular facies (Fig. 2c) were not supported by any sedimentological features.

180 Alternatively, another hypothesis has argued for a much earlier onset of inversion tectonics: in this framework, the inversion may have started in the Coniacian and/or Santonian, or perhaps even earlier, in the Late Turonian (e.g., Pożaryski, 1960, 1962; Jaskowiak-Schoeneichowa and Krassowska, 1988; Walaszczyk, 1992; Krzywiec et al., 2009; Leszczyński, 1997, 2010, 2012; Leszczyński and Dadlez, 1999; 185 Walaszczyk and Remin, 2015; Remin et al., 2016; Krzywiec et al., 2018; Remin, 2018; Łuszczak et al., 2020).

Based on new sedimentological and chronostratigraphic data, Remin et al. (2015a) and Walaszczyk and Remin (2015) proposed a diametrically opposed sedimentological model (Fig. 2d). Their interpretation suggests the presence of a land area or paleomorphological barrier in the heart of Mid-Polish Trough, 190 where the deepest part of the basin would have been located according to the model of Kutek and Głazek (1972). Furthermore, their proposed model (Fig. 2d) envisions that the deposits adjacent to the area of epigenetic erosion represent the shallowest facies, and pass into deeper facies towards the northeast (Fig. 2d) (Remin et al., 2015a; Walaszczyk and Remin, 2015).

A confirmation of this model (Fig. 2d) is provided by the presence of some specific facies developed 195  ~~near~~ along the northern edge of the Mid-Polish Anticlinorium: the Turonian Janików Limestone (sandy limestone) characterized by progradational sedimentary pattern and transport direction towards NE to the flank of the basin (Walaszczyk, 1992) and Campanian/Maastrichtian deltaically/sandy influenced sedimentation in the area of the modern-day Roztocze Hills; both are clearly indicative of a shallow-water environment. It is worth noting that the sandy facies of the 200 Roztocze Hills have been mentioned by several authors (e.g., Kamieński, 1925; Jaskowiak-Schoeneichowa and Krassowska, 1988; Świdrowska et al., 2007; Leszczyński, 1997, 2010, 2012; and literature therein), but this prominent terrigenous input has never been explored sedimentologically to reveal their possible paleobathymetry and environment and source area.

The above interpretation is amply supported by new sedimentological and biotic data (Remin et al. 205 2015a; Cyglicki and Remin, 2016, 2018; Niechwedowicz et al., 2016; Remin et al., 2016; Remin, 2018). Additionally, extraordinarily preserved plant debris, including complete compounds leaves (Halamski, 2013) and characteristic shallow-marine palynofacies (this paper), are strong indica  of proximity to land. In further support of the existence of a paleobiogeographic barrier is its potential impact on 210 paleocirculation, paleotemperature (as derived from oxygen isotopes), and the distribution of ammonite fauna (Remin et al., 2016), and the distribution pattern of belemnites during the early Maastrichtian (Remin, 2018).

All those surface observations are in line with seismic data interpretation that show an overall thinning 215 Upper Cretaceous sedimentary cover towards SW – i.e. toward  uplifted (at least partly) Mid-Polish Anticlinorium, especially in SE Poland (Krzywiec, 2009; Krzywiec et al., 2009, 2018). Additionally, those data also demonstrate local unconformities (Turonian – Maastrichtian) and progradational bodies directed  from the axial part of the basin towards its NE flank (Krzywiec, 2009; Krzywiec et al., 2009, 2018).

It is noteworthy that these two concepts are based more or less on similar input data set – and in 220 particular, the geometry and thickness patterns of sedimentary successions in cross-sections perpendicular to the Mid-Polish Anticlinorium (Fig. 2), however, the processing of those data and final interpretations were different.

Intriguing, ~~this~~ is not a novel interpretation. At the beginning of the 20th century, and indeed up until 225 the 1960s  Rogala, 1909; Nowak, 1907, 1908; Kamieński, 1925; Samsonowicz, 1925; Pożaryski, 1960, 1962), the area of southeast Poland was interpreted as an emerged landmass during the Late Cretaceous. A proliferation of different names were adopted for this landmass, including the "Łysogóry-Dobrogea Land" of Samsonowicz (1925) and Jurkowska et al. (2019a, b); the "Maastrichtian island" (Jurkowska and Barski, 2017); the "Krukienic Island" of Pasternak (1959), Pasternak et al. (1968,

1987), Walaszczyk (1992), Dubicka et al. (2014) and the “Świętokrzyski Land” or “Małopolska Land” of Pożaryski (1960, 1962) and Jaskowiak-Schoeneichowa and Krassowska (1988), among others.

230

3. Regional setting

3.1 The fill of the Polish Basin

235 The Polish Basin with its most subsiding axial part – i.e. Mid-Polish Trough represented the eastern part of the Central European Basin System (CEBS) formed during the Permian and Mesozoic in the epicontinental basins of western and central Europe (e.g. Pharaoh et al., 2010). In the Permian times the Polish Basin constituted the eastern part of the Southern Permian Basin (Kierszak et al., 1995; Krzywiec et al., 2017).

240 From Permian until the Late Cretaceous the Polish Basin experienced long term subsidence (Dadlez et al., 1995) what resulted in its filling of c. 8000 m of the Permian and Mesozoic deposits, especially along the NW-SE trending Mid-Polish Trough (e.g. Dadlez 1998) in the central part of the basin.

245 During the Late Cretaceous times and after the episode of long term Mesozoic subsidence, the Polish Basin and especially its axial part – the Mid-Polish Trough underwent uplift and inversion. The inversion tectonics started in the Late Turonian and lasted until the end of Maastrichtian (post-Maastrichtian) times (e.g. Krzywiec, 2000, 2002, 2006, 2009; Resak et al., 2008; Krzywiec et al., 2009, 2018).

250 The axial part of the Polish Basin (i.e. Mid-Polish Trough) has been transformed into a prominent structural unit – Mid-Polish Anticlinorium (Fig. 1b). In the same time two marginal troughs were formed: Szczecin-Miechów Synclinorium to the SW and Kościerzyna – Puławy Synclinorium to the NE (Fig. 1b), feed by the elevated areas in the axial zone of the Polish Basin. The marginal troughs surrounding the Mid-Polish Anticlinorium were formed in a manner initially proposed by Nielsen and Hansen (2000) for other inverted basins across Europe. As a result of the inversion procedure the Upper Cretaceous deposits are only preserved along both flanks of the Mid-Polish Anticlinorium (Fig. 1b).

255 The sedimentary cover of the southeastern part of the Polish Basin (SE Poland) is different in comparison to its central part where thick Permian, Triassic, Jurassic and Cretaceous strata accumulated. Namely, only Middle/Upper Jurassic and Cretaceous deposits are well-documented (e.g. Świdrowska et al., 2008 for an overview) whereas the missing part of succession is ascribed to denudation.

260 In the SE Poland (study area), the Middle/Upper Albian (Lower Cretaceous) and the whole Upper Cretaceous succession reach c. 1000 m of deposits formed in rather shallow epicontinental basin (e.g. Jaskowiak-Schoeneich and Krassowska, 1988). Outcrops and well data enabled precise biostratigraphic subdivision based on inoceramids, ammonites, belemnites, echinoids, foraminifera, and to a lesser extent on other groups (Walaszczyk et al., 2016). The Upper Cretaceous succession is best exposed in the NE peripheries of the elevated Mid-Polish Anticlinorium where is available in series of natural and artificial exposures, e.g. in the Middle Vistula Valley Section and in the Lubotocze area (Fig. 1b and 2a). Starting from the Cenomanian until the Maastrichtian the marine deposits cover most of

southeastern Poland and were ascribed as a siliciclastic-carbonate shelf system (e.g. Leszczyński, 1997, 2010, 2012).



270

3.2 The Roztocze Hills

The Roztocze Hills form a prominent range of hills (approximately 185 x 25 km) that extends from Kraśnik (southeast Poland) to Lviv in western Ukraine along the southwesterly margin of the Kościerzyna–Puławy Synclinorium in southeast Poland (Figs 1b and 2a). The current day Roztocze Hills are delineated along the southwest by a prominent escarpment, which constitutes a border zone with the subsurface northeastern edge of the San Anticlinorium underneath the Miocene cover of the present-day Carpathian Foredeep (Figs 1b and 2a) (e.g. Dziadzio et al., 2006 for further references). In the field, this boundary is manifested as an up to 50–80 m high escarpment, clearly visible from the almost flat surface of the Carpathian Foredeep towards the northeast. The pre-Miocene cover of the San Anticlinorium (southeastern part of the Mid-Polish Anticlinorium) is almost entirely devoid of Mesozoic remnants (with a few exceptions; Fig. 2a) and is represented mainly by ~~strongly~~ tectonized and weakly metamorphosed flysch-type sedimentary rocks of unknown thickness, in addition to anchimetamorphic rocks of Cambrian and Neoproterozoic age (e.g., Dziadzio and Jachowicz, 1996; Buła et al., 2008; Żelaźniewicz et al., 2009; Buła and Habryń, 2011).

275

This prominent tectonic line has been variously interpreted by different workers (e.g., Kutek and Głażek, 1972; Narkiewicz et al., 2015; Narkiewicz and Petecki, 2017). It is now considered that the southwestern edge of the Roztocze Hills ~~were founded on~~ crustally-rooted fault zones marked as nearly vertical crustal-scale conductive zones (review in Narkiewicz et al., 2015; Narkiewicz and Petecki, 2017), and were repeatedly reactivated during the Late Cretaceous inversion of the Mid-Polish Trough as a ~~reverse~~ ^{normal} faults (Krzywiec, 1999; Krzywiec et al., 2009) and the Carpathian tectonics during the Miocene (Tatarska et al., 2000; Buła et al., 2008; Buła and Habryń, 2011).

280

In the Polish part of the Roztocze Hills, the range is made up of Campanian (up to 550 meters thick) and Maastrichtian (c. 250 m or more, but incomplete) deposits; older deposits of the Lower/Upper Cretaceous succession are not exposed. From place to place the Upper Cretaceous is unconformably overlain by Miocene sediments (e.g., Pożaryski, 1956). Depending on the section, the Upper Cretaceous deposits of the Roztocze Hills are represented by various opoka facies (= siliceous chalk/limestone with a variable admixture of biogenic silica), gaizes (= siliceous limestone with a considerably admixture of detrital quartz, glaucony and clay), marls, calcareous sandstones, and calcareous mudstones, in addition to argillaceous mudstone or clays; the latter sometimes devoid of any CaCO_3 admixture. Regionally, the succession dips gently to the northeast.

285

3.3 The Szozdy section

The Szozdy section is situated within a railroad cutting of the Broad Gauge Metallurgical Railway Line, about 4 km southwest of Zwierzyniec close to the small village of Szozdy in southeast Poland (Fig. 3). The studied interval has yielded a rich fossil assemblage, comprising ammonites, belemnites, inoceramid and non-inoceramid bivalves, echinoids, diverse gastropods, and a suite of microfossils

(Remin et al., 2015a, b) in addition to plant debris (partially carbonized) like leaves or boughs (Figs. 3e-f).

310




Figure 3. (a) Location of the studied site in Europe; (b) Simplified geological sketch-map of extra-Carpathian Poland, excluding the Cenozoic cover (adopted from Pożaryski, 1974); tectonic units after Żelaźniewicz et al. (2011); (c) Detailed location of the studied section; (d) General view of the section – the exposures (white dashed lines) are at both sides of the railroad cutting at distance of approximately 800m; (e) Carbonized plant fragment (arrow) – the specimen is c. 10cm across; (f) The imprint of tree bough (arrow) – the specimen is c. 25cm across.

315

This fossil assemblage defines a middle Campanian age for the Szozdy section (Remin et al., 2015a):
 320 specifically, the lower/middle *Didymoceras donezianum* ammonite Zone (*sensu* Błaszkiewicz, 1980) and the middle "*Inoceramus*" *tenuilineatus* inoceramid Zone (Walaszczyk, 2004). The interval available at Szozdy, together with other sections in the Roztocze Hills, is easily correlatable with the equivalent interval in the Middle Vistula River Valley section (for location compare Figs 1b and 2a), a reference

section for the Upper Cretaceous of the southeastern Polish Basin (e.g. Walaszczyk, 2004; Walaszczyk et al., 2016).
 325




Figure 4. The stratigraphic position of the Szozdy section (right column; scale not included relative to the Middle Vistula River composite section (left column; scale bar = 2 m). A full description of the other

330 sections of the Roztocze Hills and their stratigraphic position will be provided elsewhere. The biostratigraphic subdivision follows Błaszkiewicz (1980), Walaszczyk (2004), Remin (2012, 2015), Remin et al. (2015b), and Remin (2018).

4. Material and Methods

335 The Szozdy section was sampled for 1) sedimentological evaluation, including both macro- and micro-scale observations; 2) heavy minerals, and 3) palynofacies analysis; the latter two mainly to examine hydrodynamic properties and paleoenvironmental context, respectively.

340 Of 22 total samples, seventeen (1-17) were subjected to detailed analyses: all of these were collected from the lower portion of the section (Fig. 5). This part of the section (cycles II-IV; Fig. 5), as being most densely sampled, was chosen to examine the differences and/or regularities governing the distribution of heavy minerals (Fig. 6) and palynofacies indicators, complementing sedimentological observations.

In the present study, the Szozdy section is placed within the wider context of CaCO_3 , quartz sand, clay content, plant debris, and thickness distribution patterns for the Campanian strata of the Middle and Eastern Roztocze Hills (Fig. 7).

345 The thickness pattern of the Campanian deposits in the Roztocze Hills area used in this study is based on 16 boreholes with documented Campanian deposits: Chrzanów IG-1, Dyle IG-1, Izbica IG-1, Jarczów IG-2, Jarczów IG-4, Komarów IG-1 – IG4, Narol PIG-1, Narol PIG-2, Rachanie IG-1, Stróża IG-1, Tarnawatka IG-1, Tomaszów Lubelski IG-1, and Ulhówek IG-1 (Fig. 8). The well-log data are stored in the Central Geological Data Base (CBDG; <http://baza.pgi.gov.pl/>) of the Polish Geological Institute - Polish Research Institute (PIG-PIB). The stratigraphic data, especially the lower and upper limit of the Campanian stage (where available) are based on data stored in the CBDG database for particular boreholes. Where necessary, well-log data were interpolated using the kriging and TopoToRaster geostatistic tools of ArcGIS.

355 **Sedimentology:** The scope of sedimentological analysis included both macro- and microscale (i.e., thin sections) observations. During fieldwork, the main lithofacies and thicknesses of the successive units were observed. Persistent sedimentary structures – i.e., a coarsening upward succession within the recognized cycles, trace fossils (bioturbation), macrofossil content, and micro-bioturbation (the latter within thin sections) – were also noted.

360 CaCO_3 contents were measured in the Polish Geological Institute (PIG-PIB) at over 50 localities (and c. three times more measurements). Thin section analysis enabled a volumetric estimation of the main granular components, including quartz, glauconite, and biogenic calcium carbonate. Silt and clay content were roughly estimated based on the volume of the granular and CaCO_3 components.

365 **Heavy minerals analysis:** In the present study, heavy mineral analysis includes both qualitative and quantitative analyses of heavy minerals assemblages (here, operationally defined as having a density $> 2.9 \text{ g/cm}^3$). A combination of optical microscopy, scanning electron microscope - energy dispersive X-ray spectroscopy (SEM-EDS), and electron microprobe (EPMA) analyses were applied (Mange and Maurer, 1992; Velbel, 1999, 2007; Velbel et al., 1996, 2007; Turner and Morton, 2007; Andò et al., 2012; Garzanti and Andò, 2019). EPMA analyses included an examination of tourmaline and garnet geochemistry (e.g., Wright, 1938; Henry and Guidotti, 1985; Preston et al., 2002; Mange and Morton, 2007; Meres, 2008; Andò et al., 2013).

370 **Palynofacies analysis:** Following Tyson (1993, 1995), kerogen particles were subdivided into phytoclasts (opaque phytoclasts, translucent phytoclasts, cuticle), palynomorphs (terrestrial, marine), and amorphous organic matter (AOM). A minimum of 300 kerogen particles per sample were counted for palynofacies analysis. A preliminary analysis of palynomorph assemblages is focused on both 375 sporomorphs (spores, pollen, saccate pollen) and aquatic palynomorphs (dinoflagellate cysts, foraminiferal test linings, acritarchs, algae, and freshwater algae), based on at least 250 counts per sample. Additionally, the terrestrial to marine (T/M) palynomorph index and peridinioid to gonyaulacoid (P/G) dinoflagellate cyst ratio (see, e.g., Olde et al., 2015) were also ascertained.

380 5. Results

385 5.1 Sedimentology – macro-and microscopic observations in the Szozdy section

The most prominent feature of the Szozdy section is the presence of tripartite cycloths. Where 385 complete, an individual cyclothem consists of three units: from bottom to top, calcareous mudstone, calcareous sandstone, and calcareous gaize (Fig. 5). Not every cyclothem is complete within the study section: for instance, cyclothem II lacks the calcareous sandstone unit (Fig. 5).

Presently, we cannot estimate the total thickness of the Szozdy deposits represented by tripartite cycloths, since neither the lower nor upper limit of the cyclic succession is known. The exposure 390 available at Szozdy (Fig. 3d) gives access to approximately 30-35 meters of the succession at a distance 800 m (the direct access to the section need some field works). Other natural or artificial exposures are not present; furthermore, well-log data are not available in close proximity to the Szozdy section. In the more distal boreholes, the Late Cretaceous rocks were cored only as cuttings: therefore, the existing well-log data preclude any firm conclusions. For the present study we choose 12-meters-thick interval that consists of six cycles including both complete and incomplete ones (Fig. 5).

395 The **calcareous mudstone lithofacies** (Fig. 5) is dark grey and poorly indurated. The color is most likely derived from disseminated carbonized organic matter. Of the distinguished lithofacies, the clay and silt content is highest in the calcareous mudstone, and ranges from 31% to 44% (mean = 38%). The sand fraction is composed mainly of quartz, and ranges from 16% to 26% (mean = 22%): it is the lowest within the whole section. The quartz is fine-grained, with a mean value of c. 0.1 mm; the grains are sub-angular to angular, with, less commonly, sub-rounded grains. In successive cycloths, the CaCO_3 400 content of this unit varies over a relatively narrow interval (c. 34 – 46%; mean = 38%), and is ascribed mainly to broken biocomponents. In outcrops, small bioturbation structures are visible: other sedimentary structures are absent. In this unit, macrofauna are rare: only some bivalves and badly preserved echinoids were found.

405 The **calcareous sandstone lithofacies** (Fig. 5) is yellow to yellow-brownish and poorly indurated. The clay and silt content is c. 10% lower than in the underlying calcareous mudstone, ranging from 24% to 36% (mean = 29%). The sand fraction, largely consisting of quartz and, to a subordinate extent, glauconite, varies in a narrow interval (36 – 43%; mean = 40%): these values are the highest of the lithofacies recognized in the Szozdy section. The quartz sand is still fine-grained: however, the size of 410 the grains is up to approximately 0.2 mm, twice as large relative to the underlying calcareous mudstones. The quartz grains are sub-angular to angular – sub-rounded grains are somewhat less

common. The CaCO_3 content is c. 10% lower relative to calcareous mudstone, ranging from 26% to 38% (mean = 31%). In outcrop, this unit seems to be bioturbated at least to some degree; correspondingly, sedimentary structures are absent. Macrofauna are extremely rare in this lithofacies: similarly to the underlying calcareous mudstone, only some badly preserved bivalves were found.

415 The **calcareous gaize lithofacies** (Fig. 5) is white-gray, fully indurated, and can be extremely hard; in the field, it is expressed as protruding horizons between more erosion-prone deposits (Fig. 5). This unit is composed of a limited clay and silt fraction, ranging from 8% to 18% of the bulk rock (mean = 13%): these are the lowest clay and silt fractions of the lithofacies recognized in the Szozdy section. The sand fraction, consisting of quartz sand with, to a subordinate extent, glauconite, ranges from 21% to 35% (mean = 24%): the sand fraction is similar to that recognized in the underlying unit, but is fully cemented by a calcareous cement, with only a subordinate siliceous cement. The CaCO_3 content is the highest out of the recognized lithofacies: 60 – 75% (mean = 64%). Of the three lithofacies of the full cyclothem, only this lithofacies yielded well preserved marine fauna, consisting of relatively frequent ammonites (including baculites), inoceramids and, extremely rarely, belemnites and solitary corals.

420 Other fauna include non-inoceramid bivalves and different snail species. It is worth mentioning that ammonites and other aragonitic shell material frequently retain remnants of the original shell structure, including iridescence. This unit also yielded abundant macroscopic plant debris, including tree boughs, tree branches, and relatively well-preserved compound leaves and/or their fragments.

425




Figure 5. Lithological column of the Szozdy section. Triangles on the left side of the column indicate coarsening-upward cyclothsems. Top right: typical thin sections from successive lithofacies of a single cyclothem; scale bar = 500 μ m; mos of the white particles represent quartz grains. Bottom right: field photo of a part of the section with prominent layers of calcareous gaize, indicated on the section by red dashed lines. The horizontal scale represents relative resistance to weathering of particular units.

5.2 Heavy minerals

The heavy-mineral assemblages from the mixed calcareous siliciclastic deposits of the Szozdy section are characterized by a stable composition, with little differentiation. The analyzed associations included ultrastable and stable phases: in order of decreasing resistance to weathering under surface conditions, these include rutile, zircon, tourmaline, sillimanite, kyanite, staurolite, garnet, authigenic

Usunięte: Tourmaline and garnet geochemistry

Tourmaline and garnet electron microprobe (EPMA) analyses are summarized on the ternary diagrams below (Figs. 6, 7A,B). In these diagrams, the vast majority of tourmalines occupy fields 4 and 5, following the classification of Henry and Guidotti (1985) (Fig. 6). ¶




Figure 6. Al - Fe(tot)50 - Mg50 ternary diagrams for tourmaline from various igneous and metamorphic lithologies (after Henry and Guidotti, 1985). 1 = Li-rich granitoid pegmatites and aplites; 2 = Li-poor granitoids and associated pegmatites and aplites; 3 = ferric iron-rich quartz-tourmaline rocks (hydrothermally altered granites); 4 = metapelites and metapsammites coexisting with an Al-saturating phase; 5 = metapelites and metapsammites not coexisting with an Al-saturating phase; 6 = ferric iron-rich quartz-tourmaline rocks, calc-silicate rocks, and metapelites; 7 = low-Ca metatulframas and Cr- and V-rich metasediments; 8 = metacarbonates and meta-pyroxenites. The blue dots indicate the geochemistry of tourmalines from the Szozdy section. ¶

The results of the EPMA analyses allowed us to divide garnet grains into four genetic groups, based on their compositions. The largest group consists of grains containing a high almandine content (60-70%), a somewhat lower pyrope content (20-30%), and low spessartine (<2%) and grossular (<5%) contents. The second group consists of grains with higher almandine composition (>70%), lower pyrope content (<20%), and comparatively higher spessartine (2-10%) and grossular abundance (>7%) (Figs 7A,B). The third group is characterized by an increased grossular content (about 30%) and a reduced almandine content (50-60%). The last group, in relation to the third group, is distinguished by an increased spessartine content (>8%) (Figs 7A,B). ¶

tu widać tylko kontynuację chemii w zakresie szerokim

¶

515 glauconite, pyrite, and other subordinate species such as chlorite, Cr-spinel, epidote, and single apatite grains.

The zircon-tourmaline-rutile (ZTR) maturity index (Hubert, 1962) ranges between 55-85% (mean = 67%). The highest ZTR values are derived from the calcareous mudstone lithofacies, and driven mainly by the tourmaline abundance (dravite to shorl ratio, 4:1). In general, of the three ZTR index mineral species, tourmaline dominated in all of the studied samples. Amongst the ZTR minerals, subrounded and rounded crystals dominate (> 65%); to a much lesser extent, other ZTR crystals are angular or even euhedral (c. 4%). On the other hand, garnet, kyanite, and staurolite crystals are almost exclusively angular (> 70-80%), with almost no oval and/or rounded ones.

In the lower part of the Szozdy section (which, as described above, was chosen for detailed studies), a distinct rutile and tourmaline abundance pattern can be observed - namely, that these two mineral phases are inversely correlated (Fig. 6). It is worth noting that these mineral phases are of similar size c. 100 µm in each unit. This is best observed in cycloths III and IV (Fig. 6), which are the most densely sampled. In the calcareous mudstone unit of cyclothem III, the abundance of rutile is markedly reduced, whereas the tourmaline content is significantly higher than average. The opposite pattern is observed in the calcareous sandstone overlying the dark mudstone: here, rutile abundance is substantially elevated, while the tourmaline content is lower than average. A similar pattern as in the sandstone lithofacies is observed within the calcareous gaize lithofacies (Fig. 6). The only exception to this general trend is the calcareous gaize unit at the top of the (incomplete) cyclothem II, in which the sandstone lithofacies is absent. In this case, the rutile – tourmaline abundance trends follow the pattern recognized in the underlying mudstone.

To emphasize the relative changes in abundance of these two mineral species, a standardized Z-score statistic was calculated (Fig. 6). This statistic quantifies how many standard deviations each sample is from the mean value for the whole group for the entire Szozdy section (for comparison: Ryan et al., 2007).

540




Figure 6. Vertical chart of Z-score standardized rutile and tourmaline abundance in the Szozdy section. The vertical line represents the mean abundance of the analyzed minerals over the whole section. The bars are associated with the equivalent samples shown on the stratigraphic column; SD – standard deviation.

5.3 Palynofacies

Terrestrially sourced organic matter, represented by the phytoclast group (opaque phytoclasts, translucent phytoclasts, and cuticle), is the dominant palynofacies component throughout the studied succession. The AOM content is generally low. Among the terrestrially-derived palynomorphs, 550 sporomorphs (spores, pollen, saccate pollen) are well-represented, while freshwater algae are rare. Marine palynomorphs are represented by dinoflagellate cysts, foraminiferal test linings, acritarchs, and algae. Dinoflagellate cysts and foraminiferal linings are the most common marine palynomorphs, although their abundances fluctuate throughout the succession; acritarchs and algae are less common, but have relatively consistent abundances. The dinoflagellate cyst assemblages are rich and diverse; 555 gonyaulacoid dinoflagellate cysts are generally more common than peridiniods.

Calcareous mudstone. In this lithofacies, palynofacies are dominated by phytoclasts (c. 85–93% of total kerogen), of which translucent phytoclasts (c. 33–44% of total kerogen) and cuticle (c. 22–42% of total kerogen) are the main constituents. The concentration of cuticle in calcareous mudstones is visibly 560 higher than in other lithofacies; however, the cuticle debris is largely degraded. The content of AOM fluctuates, but at low concentrations (0.4–5% of total kerogen). The terrestrial/marine ratio is relatively high, ranging between 26% and 41%, and is ascribed mainly to the abundance of non-saccate pollen (up to 82% of sporomorphs). Foraminiferal linings (c. 44–64%) are clearly more abundant than the other marine palynomorphs. The P/G ratio is variable (11–39%), and indicates that peridiniod dinoflagellate cysts are comparatively less common in calcareous mudstones relative to calcareous 565 sandstones.

Calcareous sandstone. This lithofacies is characterized by the dominance of phytoclasts (up to 79% of total kerogen), which are mainly represented by translucent phytoclasts (c. 43–63% of total kerogen). Of note are the very low concentrations of cuticle, which constitute only up to 3% of the total kerogen. The AOM content is generally low (c. 0.5–2% of total kerogen), but may occasionally reach higher values 570 (c. 15%). Calcareous sandstones are characterized by the highest T/M ratios (up to 49%) in the entire section; elevated saccate pollen concentrations (up to 51% of sporomorphs) are notable. Marine palynomorphs are dominated by dinoflagellate cysts; peridiniods are visibly more common (P/G = 28–45%) in calcareous sandstones compared to other lithofacies. Conspicuously, foraminiferal linings are almost entirely absent.


575 **Calcareous gaize.** Phytoclast concentrations, although still relatively high (c. 44–67% of total kerogen) and dominated by translucent phytoclasts (c. 18–44% of total kerogen), are visibly lower compared to the other lithofacies. Cuticle abundance is low to moderate (6–22% of total kerogen) and the cuticle debris is mostly degraded. The AOM content is generally high (c. 35–47% of total kerogen); however occasionally it may be somewhat lower (c. 7%). The T/M ratio (27–33%) is visibly lower compared to other distinguished lithofacies, with non-saccate pollen dominant over sporomorphs. Dinoflagellate cysts predominate over the other marine palynomorphs, and the P/G ratio exhibits moderate values (19–32%).

5.4 Distribution of CaCO_3 and thickness patterns in Campanian deposits

585 To place the middle Campanian succession of the Szozdy section in a wider sedimentological context, two maps were constructed. The first (Fig. 7) shows the distribution pattern of CaCO_3 content coupled

to terrigenous material (quartz sand and clay) content, and is based on more than 50 discrete outcrop observations (green triangles on Fig. 7) and c. three times as much of measurements. In most of the outcrops, two types of lithologies were available – less and more clayly. The map is constructed based 590 on the lowest available measurements from particular outcrops. CaCO_3 content is more or less inversely correlated to quartz sand and clay content. Additionally, the distribution pattern of macroscopic plant debris is provided (Fig. 7), including imprints of tree boughs, tree branches, and leaves. The frequency of plant debris is described by two subjective classes – rare and common – based on field observations in particular outcrops (Fig. 7).

595 The lowest CaCO_3 values are concentrated in a relatively narrow belt, bordered to the southwest by the present Carpathian Foredeep (the subsurface northeastern edge the San Anticlinorium) (Figs. 2A and 7): correspondingly, this is the area with the highest abundance of terrigenous material (quartz sand and clay) and plant debris. Along a SW-NE transect, the CaCO_3 content progressively increases, whereas the quartz sand and clay content tend to decline towards the northeast. It is worth observing 600 that in the southeast part of the studied area – i.e., the Horynec Zdrój area (Fig. 7) – pure clay intervals may also occur in couplets with clayey opokas.



605 **Figure 7. (a)** Distribution of CaCO_3 in the Roztocze Hills area between Zwierzyniec and the Polish/Ukrainian border to the east. Note that CaCO_3 is consistently lowest at the edge of the Roztocze Hills, at the geologic border with the Carpathian Foredeep. Green triangles indicate findings of macroscopic plant fragments; **(b)** General map of Poland with study area. Squares with numbers represent the associated geological map sheets (www.geologia.pgi.gov.pl).

The second map shows the thickness pattern of Campanian deposits in the Roztocze Hills, based on 16 boreholes with appropriate interpolations (see methods for further details). It is worth mentioning

610 that the original thicknesses, especially in the most southwesterly part which borders the Carpathian
 611 Foredeep (subsurface northeastern edge of the San Anticlinorium – compare Fig. 2a and 8) may be
 even higher, since the entire Campanian is not preserved locally. The highest values are located in the
 vicinity of the Narol boreholes (Fig. 8), where the Campanian reaches c. 550 m and along the
 612 southeastern edge of the Roztocze Hills. To the northeast, the thickness of the Campanian deposits
 progressively decreases, declining to c. 150-200 m in the Komarów and Ulhówek boreholes (Fig. 8).




Figure 8. (a) Thickness pattern of Campanian deposits in the Roztocze Hills based on well data; (b) General map of Poland with study area. Squares with numbers represent the associated geological map sheets (www.geologia.pgi.gov.pl).

6. Interpretation and discussion

The study section is special: similar lithologies have not been described previously neither from the
 Roztocze area nor from other part of the Polish Basin. Even the boreholes from the vicinity of
 625 the study section did not provide similar sedimentological pattern. The Szozdy section is characterized by
 coarsening-upward tripartite cyclothsems represented by mixed carbonate siliciclastic sediment rich in
 terrestrial components. The presence of frequent plant debris, i.e. imprint, carbonized fragments and
 disseminated organic matter of phytogenic origin, and scarcity of macroscopic, fully marine fauna in
 selected units of the cyclothsems (the calcareous sandstone) point to its origination in close proximity
 630 to land areas and at least some interplay of marine and fresh waters (compare below). For those
 reasons, Remin et al. (2015a) suggested a deltaically influenced sedimentation in the Szozdy area. Such
 an interpretation and especially its shallow, marginal marine sedimentary environment (Fig. 2d) were
 recently independently confirmed by Jurkowska et al. (2019a, b) and Machalski (2019). Additionally
 the prominent quartz sand input and its progressive disappearance towards the NE direction (Fig. 7),

635 thus outside the uplifted Mid-Polish Anticlinorium also suggest proximity of land which is considered
to be present in a place of recent-day San Anticlinorium. Although the sandy-rich facies in the Roztocze
Hills has been mentioned by several authors (e.g., Kamieński, 1927; Szaryski, 1956, 1960, 1962;
Jaskowiak-Schoeneichowa and Krassowska, 1988; Hakenberg and Świdrowska, 2001; Świdrowska,
2007; Świdrowska et al., 2008; Leszczyński, 1997, 2010, 2012), these quartz/clay-rich deposits have
640 never been explored sedimentologically, to place them within a paleobathymetric or
paleoenvironmental and sedimentological context.

Although similar cyclic sedimentation can be recognized in several marginal marine environments
(lower shoreface, storm deposits, tide-dominated deltas and more offshore shelf deposits) we stay in
645 opinion that the Szozdy section is most easily evaluated if a deltaic origin of those tripartite cycloths
is accepted. The upper and lower shoreface together with storm deposits, if any, will produce cyclicity
with normal grading, thus opposite to tide-dominated case; the tide-dominated deltas are mainly characterized
by mm- to cm-scale rhythmites (Klaes, 2012); the more offshore settings might be preclude since there
is no chance for the influence of fresh water suggested by the palynofacies analysis (compare below).

650 Besides macro- and microscopic sedimentological characteristics, the deltaic origin of the Szozdy
section is here independently supported by the palynofacies analysis and hydrodynamic properties
derived from heavy mineral analysis (as elaborated below) in addition to the distribution patterns of
siliciclastic material and CaCO₃ in wider context (Fig. 7).

6.1 Palynofacies analysis

655 The characteristic palynofacies features of this succession include abundant phytoclasts (particularly
translucent phytoclasts and cuticle), sporomorphs, peridinoid dinoflagellate cysts, and foraminiferal
linings, all of which are indicative of a relatively proximal depositional setting. In particular, the
abundance of translucent phytoclasts – that is, fresh, unoxidized particles – are suggestive of a very
660 short and rapid transportation (Tyson, 1993). A similar inference may be made from the abundance of
cuticle particles, which can easily be degraded and, thus, may serve as an indicator of rapid
transportation (e.g., Tyson, 1993). The relatively high concentrations of sporomorphs, and particularly
the presence of spores and non-saccate pollen, support this interpretation, as these sporomorphs are
665 preferentially deposited in close proximity to land (Tyson, 1993). The elevated percentages of
peridinoid dinoflagellate cysts and foraminiferal linings are likely related to an increased availability
of nutrients, which may either originate from upwelling (e.g., Wall et al., 1977; Lewis et al., 1990;
Powell et al., 1990, 1992; Eshet et al., 1994), or riverine discharge (e.g., Downie et al., 1971; Wall et
al., 1977; Powell et al., 1990, 1992; Hardy and Wrenn, 2009). In the present case, the latter source of
670 nutrients seems more probable, judging from the composition of the phytoplankton assemblage (e.g.,
the absence of typically oceanic dinoflagellate cyst genera, such as *Impagidinium* and
Nematosphaeropsis, and the presence of estuarine gymnodinoid dinocysts).

The cyclic fluctuations in the palynofacies and palynomorph assemblages characteristic of particular
lithofacies suggest highly dynamic conditions. The most evident differences exist between the
calcareous mudstones and sandstones (Fig. 9), as revealed by the changing relative percentages of
675 phytoclasts (mainly cuticle), pollen, peridinoid dinoflagellate cysts, and foraminiferal linings. Such a
sharp transition between the palynofacies patterns could be explained by either varying salinity level,

the different hydrodynamic properties of palynofacies components, or a mixture of both. Both dinoflagellate cysts and foraminiferal linings are indicative of relatively normal salinity conditions, although some dinoflagellates might have been tolerant of abnormal salinities in nearshore settings (Tyson, 1993). Hence, the rarity of foraminiferal linings in the calcareous sandstone units could possibly be explained by the influence of brackish conditions. The almost complete absence of cuticles in calcareous sandstones (Fig. 9) may also indicate higher flow conditions, resulting in the continued suspension of the most buoyant particles, such as cuticle (see Tyson, 1993), and subsequent bypassing via floatation and deposition in more distal, lower-energy settings (e.g., calcareous mudstones and gaizes) (Fig. 10). On the other hand, the elevated percentages of saccate pollen in calcareous sandstones may suggest the opposite – a decrease in water energy, since saccate pollen are preferentially concentrated in low-energy environments (see Tyson, 1995). Nevertheless, the slightly higher T/M and P/G ratios recorded in calcareous sandstones (Fig. 9) suggest that this lithofacies was apparently deposited more proximally than the calcareous mudstones and gaizes (Figs. 9 and 10).

The palynofacies patterns and palynomorph assemblages documented from calcareous mudstones and gaizes (Fig. 9) are less spectacular. In both lithofacies, the phytoclast content (including cuticle) is comparable and relatively high. Dinoflagellate cysts and foraminiferal linings are also well represented in both units, although the proportion of the latter is visibly lower in calcareous gaizes. Gaizes are also characterized by elevated AOM concentrations (the highest values recorded throughout the succession), suggestive of a more distal setting for this lithofacies (see Tyson, 1993, 1995). Considering the above, it is reasonable to suggest that calcareous sandstones represent the most proximal setting, while gaizes were apparently deposited in a more offshore environment (or eventually experienced less material supply); mudstones occupied an intermediate position (compare Fig. 10).

6.2 Hydrodynamic properties of the environment

The strong inverse relationship between tourmaline and rutile can be readily recognized throughout the whole section (Figs. 6 and 9), indicating systematic changes in hydrodynamic power in the sedimentary environment.

The Z-score calculations, normalized to the whole study section, underline the changes in the proportion of respective minerals in succeeding samples, enabling a better understanding of the processes governing their distribution pattern within the section (Fig. 6) (e.g., Ryan et al., 2007). Both minerals are of similar shape, durability and size; however, they possess markedly different densities – namely, 3.03-3.18 g/cm³ and 4.23 g/cm³ for tourmaline (dravite) and rutile, respectively. Since the weight of the two analyzed minerals is high, it might be expected that these two mineral phases will be strongly dependent, both vertically and spatially, on the local sedimentary environment – and by extension, the hydrodynamic power that actually existed during the deposition of the cyclothem. The recurring increase in tourmaline abundance, coupled to a simultaneous decrease in rutile abundance in muddy units, most likely resulted from a decrease in hydrodynamic power in the depositional environment. Sedimentologically, this may be interpreted as a transition to a comparatively more distal depositional setting from the main source area, e.g. a river discharge – that is, the prodelta or analogous environment (compare Figs. 9 and 10) (Komar, 2007; Omran, 2007).

Analogously, the increase in the proportion of rutile, and the simultaneous decrease of tourmaline share, in the sandy units can be linked to an increased flow rate, which might be interpreted as a transition to an environment more proximal to the river discharge, thus representing the main delta lobe and/or slope setting (compare Figs. 9 and 10). Simply put, the (lighter) tourmaline will be transported further to the prodelta environment, rendering the prodelta facies overrepresented in this mineral phase, whereas the (markedly heavier) rutile will be deposited closer to the river discharge.




Figure 9. The proposed environment for the successive lithofacies within an individual cyclothem in the Szozdy delta, coupled with the vertical distribution of selected heavy minerals and palynofacies indicators: cyclothem III adopted from Fig. 6. The values in brackets show the range for the lithofacies across all studied cyclothems. SD = standard deviation.

6.3 Sedimentary environment – looking for recent counterpart

We found that the modern Mahakam deltaic system (Indonesia) may serve as a good modern counterpart for the Szozdy Delta sedimentary environment. The similarities are, of course, general – the Mahakam Delta is huge in comparison to the Szozdy deltaic system (although the spatial and vertical distribution of the latter is unknown); however, beyond those differences, several analogies can be found in their respective sedimentary environments.

The presence and relative importance of physical processes (e.g., water flow, sediment input, ocean currents, wind, waves, and tides) regulate the morphology and internal geometry of deltas. Accordingly, three main classes are commonly distinguished: i) tide-dominated, ii) wave-dominated, and iii) river-dominated deltas (e.g., Coleman and Wright, 1975; Galloway, 1975).

The modern Mahakam delta is considered a textbook example of a fluvial-tide-dominated delta system built across a narrow shelf (Galloway, 1975; Allen et al., 1977). The shelf and deltaic deposits border the N-S oriented Makassar Trough, which acts as a throughflow pathway for Pacific waters to the Indian Ocean (Wyrtki, 1987); as such, it must also regulate, at least in part, the redistribution of suspended sediments. The Mahakam River discharge is characterized by a distinct absence of flood surges: therefore, the avulsion of distributary channels does not typically occur (e.g., Allen et al., 1977; Storms et al., 2005). Additionally, the extremely wide, submerged delta front platform, which extends as a 5-meter isobath up to 15-20 km offshore, works as an ideal platform to dissipate wave energy (Roberts and Sydow, 2003). As a consequence, the fluvial distributaries efficiently push suspended

750 sediment basinward, resulting in extremely fast progradation (Storms et al., 2005, and references therein).

755 For the Szozdy Delta, it is difficult to ascertain which process(es) were dominant in the development of this deltaic system (fluvial-, tide-, and/or wave-dominated): at a minimum, such assessments are speculative to at least some degree. Although relatively large, the exposure at Szozdy – a 800 m long railroad cutting that provides access to approximately 30-35 vertical meters of the succession, still represents a single locality, with such a specific sedimentation style that can be best ascribed as deltaic in origin.

760 On the other hand, it is worth accounting for the fact that successive cycloths are very regular, and particular units can be traced ^{over} dozens to few hundreds of meters. Strong wave and/or tidal action would most likely result in a more chaotic development of the successive lithofacies, and ultimately entire cycloths. Accordingly, at the current stage, we can only speculate that fluvially-dominated processes are predominant, with possibly some additional wave and/or tidal action.

765 The limited thickness of the recurring cycloths – 1 – 5 m (Fig. 5) – and upwards grain-size coarsening patterns in each cycloth indicate that the progradation of succeeding facies – i.e., the transition between the more muddy and more sandy lithofacies of the cycloths (Figs. 5 and 9) – proceeded relatively quickly. This also suggest that accommodation space was highly limited (Fig. 10).

770 Since there is no evidence for the presence of delta plain deposits, it is suggested that the entirety of sedimentation occurred in the subaqueous part of the Szozdy deltaic system, as exemplified in figure 10.

775 The implication is that, at least locally, the deposition of the submerged deltaic facies of the Szozdy delta took place along a relatively flat, wide, shallow to extremely shallow delta front platform (Figs. 10 and 11), similar to the delta front platform of the modern Mahakam delta, where progradation is fastest. The platform was wide enough to dissipate most of the wave-energy action, rendering the influence of waves low to moderate, at a maximum.




Figure 10. Block diagram illustrating the relationships between the subaerial (not available in the study area) and subaqueous deltaic environments of the Szozdy delta in relation to active and abundant

distributary channels. Arrows 1 and 2 indicate two different situations i.e. 1) – a full tripartite cyclothem and 2) an incomplete cyclothem; compare to Fig. 5. Not to scale.

780

The differing thickness of successive lithofacies in the cycloths (Fig. 5) suggests that specific environmental conditions had different durations before a threshold was reached that drove a change in paleoenvironmental regime from prodelta to delta lobe/slope facies. This suggests that rather dynamic tectonic conditions existed, resulting in changes of the delta architecture, the migration and 785 avulsion of distributary channels, and simultaneous changes in the sedimentary environment at particular localities, resulting in the repetition of cycloths.

785

According to this proposed environmental interpretation, the lowest unit of each cyclothem – that is, the calcareous mudstone – would represent the prodelta environment (Figs. 10 and 11), deposited on the submerged delta front platform in a comparatively distal region from the river discharge (Figs. 10 790 and 11). This unit possesses the highest clay abundances and the finest sand fraction. Upsection, the calcareous mudstone passes quickly into yellowish calcareous sandstone, with markedly less disseminated organic matter of phytogenic origin. In additional support of such an environmental interpretation is the scarcity of macroscopic, fully marine fauna: while some bivalves and echinoids 795 were recognized in the mudstone lithofacies, in the sandstone lithofacies such fauna are almost absent.

795

The origin of the calcareous gaize is more problematic. This lithofacies is considered to originate in an area cut off from the sediment supply. Once the siliciclastic sediment supply is inhibited – for instance, via tectonic activity and/or changes in delta architecture (e.g., the avulsion of distributary channels) – 800 a relatively quick calcification process is thought to quickly lithify and “freeze” the upper portion of the sediment.

800

Three observations support such an interpretation. First, in all studied cases, the dominant grain size is always the same as the grain size observed in the underlying unit. In complete cycloths [i.e., cycloths I and III-VI], the gaize possess a grain size similar to the sandy unit (Fig. 5). In cyclothem II, where gaize directly overlies the muddy unit, the grain size is smaller and follows that observed in the mudstone. Second, in this unit, fully marine fauna are quite common, which indicates that the 805 freshwater supply was comparatively limited or redirected. Additionally, terrestrial components are also abundant. In particular, macroscopic plant debris, including tree boughs, tree branches, and compound leaves, are commonly found: it is suggested that buoyant fragments were transported, sunk, and subsequently “frozen” by calcification. Third, ammonites and other aragonitic shelled 810 animals still retain remnants of the original shell structure, including their iridescence. This indicates that they were cut off from the surrounding environment, which would have assurdly been corrosive to aragonite – in other words, the original aragonite could not have been preserved without syndepositional and/or early diagenetic calcification and chemical stabilization.

815

6.4 Towards an understanding of Late Cretaceous facies distribution

The very existence of a middle Campanian deltaic system developed in the axial part of the inverting Mid-Polish Trough (into the Mid-Polish Anticlinorium) has some fundamental consequences for our overall understanding of certain Late Cretaceous facies. In particular, this concerns mainly opoka, chalk, and their respective variants, in addition to their mutual spatial and bathymetric relationships in transects perpendicular to the Łysogóry-Dobrogea Land (Fig. 2d).

The evidence presented here for a shallow water, deltaic origin of these mixed carbonate siliciclastic deposits confirms the conceptual proposition of deltaically influenced sedimentation in the Roztocze Hills during the Campanian initially proposed by Remin et al. (2015a) and Walaszczyk and Remin (2015), and fully supports their general sedimentological and bathymetrical model (compare Fig. 2d). This bathymetric model and a shallow water position of the opoka facies (coastal environment) has been recently confirmed independently by Jurkowska et al. (2019a, b) and Machalski (2019).

In this respect a continuously growing San Anticlinorium during the Late Cretaceous acted as a source area for the siliciclastics of the Roztocze Hills. In the present study, for the first time, we can directly demonstrate that these terrigenous-rich deltaic sediments interfinger with gaize and opoka variants located slightly northeast of the recognized Szozdy deltaic system. It clearly shows that this specific facies – opoka – should be considered as having been deposited in a shallow-water environment, at least in this specific case, in opposition to previous notions rooted in different tectonic and sedimentological models, indicating its rather deep-origination. In the SW-NE transect perpendicular to the Szozdy deltaic system and the Łysogóry-Dobrogea Land (in a place of recent day San Anticlinorium), as defined here, the more carbonate-rich deposits, including chalk, were deposited in more offshore and deeper zones located to the northeast of the Szozdy deltaic system. The presence of such a deeper zone is confirmed by the discovery of contourites (Krzewiec et al., 2009, 2018): the contourites current must have operated along a regional slope located northeast of both the Łysogóry-Dobrogea Land and the Szozdy deltaic system (Fig. 12).

Quite similar facies and structural situation we might find in the Subhercynian Cretaceous Basin which undergone inversion c. in the same time as the Mid-Polish Trough. The vast majority of the Coniacian to Campanian deposits, dominated by sandy and marly deposits, were derived from adjacent source areas, i.e. an elevated Harz Mountains (Voigt et al., 2004, 2006, 2008; von Eynatten et al., 2008). The Coniacian storm-dominated deposits and Santonian deposits of a low-energy delta plain to tidal-flat environment consisting of sands were deposited close to the areas elevated during inversion tectonics and pass into deeper facies in more offshore zones.




Figure 11. (a) CaCO_3 distribution pattern (blue area, compare Fig. 7) against the isolines of Campanian deposit thickness. Red dots = boreholes (compare Fig. 8); yellow dots = interpolation points; (b) General map of Poland with study area; (c) Subaerial view of the Szozdy deltaic environment in relation to the distribution pattern of selected facies bordering the deltaic environment – not to scale. Squares with numbers represent the associated geological map sheets (www.geologia.pgi.gov.pl).

860 For several decades, our understanding of facies distribution in the Upper Cretaceous of extra-
 855 Carpathian Poland was heavily influenced by the paleotectonic model developed and proposed by
 Kutek and Głażek (1972). Those authors *a priori* synonymized the axial part of the Mid-Polish Trough
 (in the present day represented as the Mid Polish Anticlinorium; Fig. 2b) with the most subsiding,
 deepest portion of the basin. It was consequently assumed that the facies deposited closest to the
 865 axial part of the Mid-Polish Trough (e.g., in the Roztocze Hills area) would represent the deepest
 sedimentary environment. As such, facies distributions and bathymetric interpretations were fit to this
 paleotectonic model (see also discussion in Walaszczyk and Remin, 2015). It is noteworthy, however,

that Late Cretaceous mixed carbonate siliceous facies elude easy environmental interpretation, and the adopted paleobathymetric model and the spatial distribution of particular facies (compare Figs 2c),
870 were not supported by any relevant sedimentological and/or biological indicators.

Nevertheless, the notion that opokas were largely of deep water origin (Fig. 2c) has been widely accepted, and for decades dominated thinking concerning the spatial distribution of facies and their relation to paleobathymetry (Leszczyński, 1997, 2010, 2012; Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska 2007; Świdrowska et al., 2008; Świerczewska-Gładysz, 2006). Accordingly, the Late Cretaceous opoka facies, the dominant facies of the Roztocze Hills, were considered as deep-water sediments until the present study, despite a considerable amount of terrigenous material (mainly sand-grained quartz, as well as clays). In light of the new data presented herein, this conception cannot be feasibly upheld, at least for the opoka facies of the Roztocze Hills.
875

880 **6.5 Redistribution of terrigenous material**

In a wider context, ocean currents (contourite currents) acting parallel to the shoreline of the identified Łysogóry-Dobrogea Land may have some influence on the redistribution of (mainly) the finest terrigenous material, pushing it away from the northern part of the Szozdy delta towards southeastern direction (Figs. 11 and 12).

885 The existence of contourite currents operating along this putative landmass during the Campanian was revealed based on seismic profiles perpendicular to the landmass postulated here (Krzywiec et al., 2009; 2018) (Fig. 12). In general, their appearance within the basin might be related to oceanographic, bathymetric condition or enhanced tectonic activity (inversion), regulating the morphology of the seafloor (e.g. Rebesco et al., 2014). Similar structures, based on seismic data, has been also proved from the Danish Basin (e.g. Surlyk and Lykke-Andersen, 2007). Their recognition provides independent evidence for the paleo-circulation within the Polish Basin as based on surface data.
890

Intriguingly, presence of currents flowing from the north and along the northern shore of the supposed paleobiogeographic barrier, interpreted here as a Łysogóry-Dobrogea Land (i.e., the area presently occupied by the San Anticlinorium), during the Coniacian and Santonian was suggested by Remin et al. (2016), who outlined its potential impact on paleocirculation, paleotemperature (as derived from oxygen isotopes), and the distribution of ammonite fauna. In further support of the existence of a paleobiogeographic barrier is the distribution pattern of belemnites during the early Maastrichtian (Remin, 2018).

900 The existence of contourite currents indicates that the Szozdy deltaic system was limited to the northeast by a regional slope in the sea-bottom; currents flowing parallel to the strike redistributed terrigenous material towards the southeast. A potential manifestation of this current-driven redistribution of the finest terrigenous material may be pure clay deposits of approximately the same age located to the southeast of the Szozdy delta – that is, in the Horyniec Zdrój area (Figs. 11 and 12). This area is additionally characterized by the lowest amount of CaCO_3 in the Campanian deposits of the Roztocze Hills, and high terrestrial input.
905

Such a paleogeographic situation is, again, somewhat similar to the modern Mahakam deltaic system that borders the Makassar Trough: that is a throughflow channel facilitating a contourite curren

(Shanmugam, 2017; Brackenridge et al., 2020), which redistributes material from the Mahakam prodelta to the south.

910




Figure 12. Schematic paleogeographic map and location of the Roztocze Hills and the Łysogóry-Dobrogea Land. The direction of ocean current flow is based on Remin et al. (2016) and Remin (2018); the presence of contourite currents (= CC) is according to Krzywiec et al. (2009, 2018).

915

6.6 Drivers of accommodation space generation

920

Inversion-related tectonic instability along the deeply rooted fault zone (the southwestern boundary of the Roztocze Hills; Fig. 2a) led to the development of the Szozdy deltaic system. To the southwest of this fault zone, thus in the area of the subsurface San Anticlinorium (Figs. 2a and 2d), there existed a landmass during the Late Cretaceous – i.e., the Łysogóry-Dobrogea Land – that was a source area for the siliciclastics of the Szozdy delta. During the process of tectonic inversion, this area underwent uplift, which surely resulted in changes to detritial weathering, weatherability, erosion, and provenance. Correspondingly, on the other side of this fault zone (that is, to the northeast) the Upper Cretaceous strata of the present Roztocze Hills were deposited, including the cycloths of the 925 subaqueous delta front platform of the Szozdy deltaic system. This area, being a part of forming marginal trough, experienced enhanced subsidence, which resulted in the deposition of thick Campanian strata, especially along this active fault zone (compare Figs. 8 and 11).

935

Subsidence of the Roztocze Hills area, on the one hand, and uplift and erosion of the Łysogóry-Dobrogea Land on the other (in the area of the modern San Anticlinorium) during the Late Cretaceous inversion, could have conceivably driven the deltaic architecture, supplying additional accommodation space for the development of succeeding Szozdy delta cycloths (Figs. 10 and 11). Furthermore, tectonically-induced migration and avulsion of distributary channels led to changes in the local sedimentary environment, which resulted in a predictable repetition of Szozdy cycloths.

6.7 Speculative comments on burial history and the rate of subsequent erosion

Since the paper of Kutek and Glazek (1972), the Holy Cross Mountains, and the San Anticlinorium to the southeast (Figs. 1b and 2a), have been traditionally included by most workers into the axial part of the former Mid-Polish Trough, which subsequently underwent inversion, uplift, erosion, and transformation into the Mid-Polish Anticlinorium. Apart from the concept of inversion tectonics onset itself, there was a general agreement that the axial part of the Mid-Polish Anticlinorium (e.g. area of the recent-day Holy Cross Mountains and the San Anticlinorium) were deeply buried by thick Permian and Mesozoic sediments (e.g., Kutek and Glazek, 1972). Kutek and Glazek (1972) suggested that c. 3000m of sedimentary rocks have been removed by erosion from the axial part of the Mid-Polish Anticlinorium, especially from the Holy Cross Mountains and the San Anticlinorium (Figs. 1b and 2a). More recent estimations varies between 2000-2500m (Resak et al., 2008; Narkiewicz et al., 2010). Similar values of overburden (1700-2300) were obtained by Luszczak et al. (2020).

Since still little is known about possible Mesozoic overburden over the San Anticlinorium (this area is almost devoid of Mesozoic remnants; Figs. 1b and 2a), the heavy mineral assemblages from the Szozdy delta were analyzed to look for possible indicators of provenance. It provides intriguing data regarding the potential source rocks for the siliciclastics of the Szozdy deltaic system.

The heavy mineral assemblages from the Szozdy section exhibit a truly polycyclic nature. On the one hand, high values of the ZTR index (ascribed to ultrastable mineral phases), the high degree of roundness of (most) mineral grains, and the presence of some abrasive structures suggest multiple rounds of redeposition (Hubert, 1962; Mange and Wright, 2007; Garzanti, 2017; Garzanti and Andò, 2007, 2019). On the other hand, the presence of mineral groups weakly resistant to erosion – garnets, kyanites, and staurolites – that consists almost exclusively of angular crystals (> 70-80%) with almost no oval and/or rounded representatives is inconsistent with repeated redeposition; instead, they support a multi-source provenance for the Szozdy deltaic deposits, as further elaborated below.

Accordingly, the above data suggest the presence of at least two distinct sources. The first yielded multi-recycled mineral phases dominated by minerals from the ZTR group, for which the source might be metapelites/metapsamites. The second, “fresh” source delivered angular-shaped crystals of garnets, kyanites, and staurolites, for which amphibolite facies were the main source.

This data suggest that the source rocks for the heavy mineral assemblages of Szozdy delta are represented by both (i) multi-recycled sedimentary rocks and (ii) freshly weathered metamorphic and igneous rocks. These characteristics would be rather difficult to obtain from the Mesozoic sedimentary cover of the San Anticlinorium (if it was even present at all), necessitating a search for other sources. A tempting hypothesis for the potential source rocks, although highly speculative, would be a comparison of mineral assemblages from the Szozdy delta to the Cambrian and Neoproterozoic rocks subcropping beneath the Miocene deposits of the Carpathian Foredeep, i.e. recent-day San Anticlinorium, represented by metamorphosed (anachimamorphic) flysch-type siliciclastics.

975

Usunięte: This part is markedly shorten and moved to “Regional setting”[¶]

Presently, we cannot estimate the total thickness of the Szozdy deltaic deposits, since neither the lower nor upper limit of the cyclic succession is known. The exposure available at Szozdy gives access to approximately 30-35 meters of the succession. Other natural or artificial exposures are not present; furthermore, well-log data are not available in close proximity to the Szozdy section. In the more distal boreholes, the Late Cretaceous rocks were cored only as cuttings: therefore, the existing well-log data może to powinno trafić do opisu geologicznego[¶]

675-680. This is very important and should also mentioned in the introduction.[¶]

The current day Roztocze Hills are delineated along the southwest by a prominent escarpment, which constitutes a border zone with the subsurface northeastern edge of the Łysogóry and Małopolska Blocks underneath the Miocene cover of the present-day Carpathian Foredeep (Fig. 2). This prominent tectonic line has been variously interpreted by different workers: the original interpretations considered it to be a prolongation of the Holy Cross Mountain Fault (e.g., Kutek and Glazek, 1972). Others (e.g., Narkiewicz et al., 2015; Narkiewicz and Petek, 2017) synonymized the southwestern edge of the Roztocze Hills with the Janów Fault (Fig. 2) and placed the prolongation of the Holy Cross Mountain Fault (i.e., the Cieszanów

Usunięte: Kutek and Glazek (1972) suggested that c. 3000 m of sedimentary rocks have been removed by erosion from the axial part of the Mid-Polish Anticlinorium, especially from the Holy Cross Segment (Fig. 2). Later estimations, like those of Belka (1990), suggested that in the Holy Cross area (to the northwest of the San Anticlinorium; Fig. 2), the thickness of post-Devonian overburden varied between 1500 and 3800-5500 m depending on the specific locality, and was generally higher close to the Holy Cross Fault. Similar values of overburden (1700 – 2300 m) were recently obtained by Luszczak et al. (2020) for the western slopes of the Holy Cross Mountains.[¶]

Those values, however, cannot be simply transposed to the San Anticlinorium, because the regional geological background is different. First of all, there are no Permian, Triassic, and Lower Jurassic deposits northeast of the San Anticlinorium (e.g., Świdrowska et al., 2008). The oldest known sedimentary strata originated in the Middle Jurassic: as such, the stratigraphic record is markedly reduced in comparison to the Holy Cross segment and the Kujawy region to the northwest, where thick Permian, Triassic, and Jurassic deposits are present (Kutek and Glazek, 1972; Świdrowska et al., 2008; Krzywic et al., 2018 for overview).[¶]

Usunięte: As suggested by the reviewer 3 (Thomas Voigt) we should avoid most of the provenance data since we don't have anything for comparison and there is additional lack of the zircon age spectra from the basement of the San Anticlinorium. I concur and most of the geochemistry of the heavy minerals are avoided – only some actual highlights are left (as a base for future studies).[¶]
The tourmalines belong to the alkaline subgroup, in which the ratio of dravite to shorl is 4:1, which in turn suggests that the tourmaline grains potentially came from metamorphic rocks. Some of the tourmalines correspond to metamorphosed sedimentary rocks rich in aluminum phases, such as kyanite and staurolite, which together define the studied mineral complexes (Fig. 6). A further portion of the analyzed grains correspond to tourmalines derived from Li-poor granitoids and their vein counterparts, based on their chemical composition. A potential source of two tourmaline grains (Fig. 6) might be ultramafic rocks and metasediments enriched in Cr and V, which is additionally underpinned by the presence of Cr-spinels (Fig. 6).[¶]

In the ternary diagrams of Meres (2008), most garnet grains occupy fields C1 and C2 (Figs 7A,B). These fields correspond to garnets th...

7. Summary

The interpretation presented herein implies the presence of a landmass area – that is, the Łysogóry-Dobrogea Land – in a place where, according to many previous interpretations, the deepest part of the Mid-Polish Trough was located. For the first time, clear and conclusive evidence for this iconoclastic paleogeographic and paleobathymetric model are provided, based on sedimentological data coupled with heavy mineral and palynofacies analyses. We commend that the name Łysogóry-Dobrogea Land should be used, as it has priority (vide Samsonowicz, 1925).



The recognition of the Szozdy deltaic system in the supposedly axial part of the Mid-Polish Trough grants a unique perspective into the sedimentary environment, hydrodynamics, facies architecture, and bathymetric position of several Late Cretaceous facies. For instance, various styles of opoka facies are located adjacent to deltaic deposits, consistent with a shallow-water origin. A deeper sedimentary environment is suggested for chalk facies.



Inversion-related tectonic instability along the deeply rooted fault zone (the southwestern boundary of the Roztocze Hills) led to the development of the Szozdy deltaic system. Uplift of the present-day San Anticlinorium area during the Late Cretaceous inversion, on the one hand, and enhanced subsidence of the present-day area of the Roztocze Hills on the other, rearranged the delta architecture, supplying additional accommodation space for the development of Szozdy delta cycloths.


The Łysogóry-Dobrogea Land, located to the south and southwest to the Roztocze Hills – thus, where the San Anticlinorium is presently situated – is the only reasonable source for the siliciclastics of the Szozdy deltaic system. This area is currently almost completely devoid of Cretaceous and other Mesozoic remnants and is now hidden under the Miocene cover of the Carpathian Foredeep.

The geochemistry of tourmalines and garnets (compare also Cyglicki and Remin, 2016, 2018) may suggest the strata subcropping beneath the Miocene series of the Carpathian Foredeep (to the southwest of the Roztocze Hills) were already emerged and eroding during the Late Cretaceous, at least to some degree.

8. Current challenges and future roadmap

The results obtained in this study and from previous research raise several crucial issues that should be addressed in the near future, and furthermore allow us to formulate a working hypothesis that the modern San Anticlinorium was an elevated structural element during the Late Cretaceous. Such a hypothesis, although highly speculative, is supported by the initial geochemical data of heavy minerals we obtained, which might suggest that at least part of the material is derived from freshly weathered metamorphosed rocks. Further evidence and verification of these postulations would allow the following issues to be examined:

1. What was the source region for the Late Cretaceous Szozdy deltaic system and siliciclastics of the Roztocze Hills? Was it the Mesozoic cover of the San Anticlinorium? Or, alternatively, were the



Usunięte: Its sedimentary cover, that subcropping beneath the Carpathian Foredeep, was already actively eroding, supplying terrigenous material for the siliciclastics of the Roztocze Hills.

1150 Cambrian and Neoproterozoic rocks beneath the Miocene cover of the Carpathian Foredeep already eroding during the Cretaceous?

2. Did the San Anticlinorium constitute a homogeneous part (structurally and paleogeographically) of the Mid-Polish Trough during the Cretaceous?

3. In this context, what was the real nature of the southeastern extension of this trans-European sedimentary basin, the Mid-Polish Trough, during the Cretaceous?

4. Accordingly, what was the burial history of the San Anticlinorium, if burial had occurred at all?

1155 Deciphering these crucial objectives, based on a revised paleofacies and paleobathymetric model, is of fundamental significance for understanding the Late Cretaceous paleotectonic evolution of the supposed southeastern edge of the former Mid-Polish Trough.

1160 In turn, a precise recognition of the stratigraphic succession and facies distribution (work which is currently ongoing) in the Roztocze Hills will allow for the reliable identification of the main phases of terrigenous material input from the nearby landmass, along with the estimation of subsidence gradients and the identification of these events on high-resolution seismic profiles (also ongoing). In this framework, a reduction of the subsidence rate should lead to the progradation of the siliciclastic facies toward the northeast, thus trending from more proximal to more offshore zones.

1165 Solving the issues outlined above, especially with regards to resolving whether or not the San Anticlinorium constitutes a structural part of the Mid-Polish Trough, may fundamentally change our understanding of regional geology and burial history, and more broadly speaking the paleogeography and paleobiogeography of Central Europe during the Late Cretaceous, with far-reaching interpretative consequences.

1170 *Author contributions.* ZR: idea, conceptualization, data compilation, discussion and validation, original draft, revision, funding acquisition (NCN; grant number UMO-2018/29/B/ST10/02947); MC: conceptualization, heavy minerals analyses, discussion and validation, revision; MN: conceptualization, palynofacies analyses, discussion and validation, revision.

Competing interests. The authors declare that they have no conflict of interest.

1180 **Acknowledgements:** Our thanks are extended to Bartosz Gościński and Joanna Roszkowska-Remin for help during the field work. Special thanks go to Robert Banaś, the chief forester of the Lubaczów district, and local forest officers who granted access to several outcrops and served with help in the field. Financial support was provided by the Polish National Science Centre [Narodowe Centrum Nauki, Polska] grant number UMO-2018/29/B/ST10/02947.

1185

9. References

1190 Allen, G. P., Laurier, D., and Thouvenin, J.: Sediment distribution patterns in the modern Mahakam delta, Proceedings Indonesian Petroleum Association, 5, 159–178, 1977.

Andò, S., Garzanti, E., Padoan, M., and Limonta, M.: Corrosion of heavy minerals during weathering and diagenesis: A catalog for optical analysis, *Sedimentary Geology*, 280, 165–178, 2012.

1195 Andò, S., Morton, A. C., and Garzanti, E.: Metamorphic grade of source rocks revealed by chemical fingerprints of detrital amphibole and garnet, *Geological Society, London, Special Publications*, 386, 351–371, 2013.

Błaszkiewicz, A.: Campanian and Maastrichtian ammonites of the Middle Vistula River valley, Poland: a stratigraphic and paleontological study, *Prace Instytutu Geologicznego*, 92, 1–63, 1980.

1200 Brackenridge, R. E., Nicholson, U., Sapiie, B., Stow, D., and Tappin, D. R.: Indonesian Throughflow as a preconditioning mechanism for submarine landslides in the Makassar Strait. *Geological Society, London, Special Publications*, 500, 195–217, 2020.

1205 Buła, Z., Byś, I., Florek, R., Habryn, R., Jachowicz, M., Kwarciński, J., Laskowicz, R., Liszka, B., Madej, K., Maksym, A., Markowiak, M., Pietrusiak, M., Probuski, J., Ryłko, W., Salwa, S., Sikora, R., Staryszak, G., Tabol-Wójcik, P., Tomaś, A., Zacharski, J.: Geological-structural atlas of the Palaeozoic basement of the Outer Carpathians and Carpathian Foredeep, Polish Ministry of Environment, Warszawa, 104, 775–796, 2008.

1210 Buła, Z. and Habryn, R.: Precambrian and Palaeozoic basement of the Carpathian Foredeep and the adjacent Outer Carpathians (SE Poland and western Ukraine), *Ann Soc Geol Pol*, 81, 221–239, 2011.

Coleman, J. M. and Wright, L. D. (Eds). *Modern river deltas: variability of processes and sand bodies*. 1975.

1215 Cyglicki, M. and Remin, Z.: Provenance of heavy minerals in the middle Campanian (Cretaceous) siliciclastic deposits of the Roztocze Hills, SE Poland, in: X Baltic Stratigraphic Conference, Checiny, 28–29, 2016.

1220 Cyglicki, M. and Remin, Z.: HRHMA (high-resolution heavy mineral analysis) zastosowana dla górnokampeńskich skał silikoklastycznych krawędziowej części Roztocza Środkowego, SE Polska, *Pokos*, VII, 67, 2018.

Dadlez, R., Marek, S., and Pokorski, J. (Eds.): *Geological map of Poland without Cenozoic deposits at 1:1000 000 scale*, Warszawa, Polish Geological Institute, 2000.

1225 Dadlez, R.: Mesozoic thickness pattern in the Mid-Polish Trough, *Geological Quarterly*, 47, 223–240, 2003.

Dadlez, R., Narkiewicz, M., Stephenson, R. A. Visser, M. T. M., and van Wess, J. D.: Tectonic evolution of the Mid-Polish Trough: modelling implications and significance for central European geology. *Tectonophysics*, 252, 179–195, 1995.

1230 Dadlez, R., Marek, S., and Pokorski, J. (Eds.): *Palaeogeographical Atlas of the Epicontinental Permian and Mesozoic in Poland (1:2 500 000)*. Polish Geological Institute, Warszawa, Poland [in Polish with English summary], 1998.

Downie, C., Hussain, M. A., and Williams, G. L.: Dinoflagellate cyst and acritarch associations in the Paleogene of southeast England, *Geoscience and Man* 3, 29–35, 1971.

1235 Dubicka, Z., Peryt, D., and Szuszkiwicz, M.: Foraminiferal evidence for paleogeographic and paleoenvironmental changes across the Coniacian–Santonian boundary in western Ukraine, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 401, 43–56, 2014.

1240 Dziadzio, P. and Jachowicz, M.: Budowa podłożu utworów mioceńskich na SW od wyniesienia Lubaczowa, *Przegląd Geologiczny*, 44, 1124–1130, 1996.

1245 Dziadzio, P., Maksym, A., and Olszewska, B.: Miocene deposition in the eastern part of the Carpathian Foredeep in Poland, *Przegląd Geologiczny*, 54, 413–420, 2006.

Eshet, Y., Almogi-Labin, A., and Bein, A.: Dinoflagellate cysts, paleoproductivity and upwelling systems: A Late Cretaceous example from Israel, *Marine Micropaleontology* 23, 231–240, 1994.

1250 Eynatten, von H., Voigt, T., Meier, A., Franzke, H.-J., and Gaupp, R.: Provenance of the clastic Cretaceous Subhercynian Basin fill: constraints to exhumation of the Harz Mountains and the timing of inversion tectonics in the Central European Basin. *Int. J. Earth Sci.*, 97, 1315–1330, 2008.

1255 Galloway, W. E.: Process framework for describing the morphologic and stratigraphic evolution of deltaic depositional systems, in: *Deltas, Models for Exploration*, edited by: Broussard, M.L., Houston Geological Society, Houston, TX, pp. 87–98, 1975.

Garzanti, E.: The maturity myth in sedimentology and provenance analysis, *Journal of Sedimentary Research*, 87, 353–365, 2017.

1260 Garzanti, E. and Andò, S.: Heavy mineral concentration in modern sands: implications for provenance interpretation, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, pp. 517–545, 2007.

Garzanti, E. and Andò, S.: Heavy minerals for Junior Woodchucks, *Minerals*, 9(3), 148, 1–25, 2019

1265 Hakenberg, M. and Świdrowska, J.: Evolution of the Holy Cross segment of the Mid-Polish Trough during the Cretaceous, *Geological Quarterly*, 42, 239–262, 1998.

Hakenberg, M. and Świdrowska, J.: Cretaceous basin evolution in the Lublin area along the Teisseyre-Tornquist Zone (SE Poland), *Annales Societatis Geologorum Poloniae*, 71 (1), 1–20, 2001.

1270 Halamski, A. T.: Latest Cretaceous leaf floras from Southern Poland and Western Ukraine, *Acta Paleontologica Polonica*, 58 (2), 407–433, 2013.

Hardy, M. J. and Wrenn, J. H.: Palynomorph distribution in modern tropical deltaic and shelf sediments – Mahakam Delta, Borneo, Indonesia, *Palynology* 33, 19–42, 2009.

1275 Henry, D. J. and Guidotti, C. V.: Tourmaline as a petrogenetic indicator mineral: an example from the staurolite-grade metapelites of NW Maine, *American Mineralogist*, 70, 1–15, 1985.

1280 Hubert, J. F.: A zircon-tourmaline-rutile maturity index and interdependence of the composition of heavy mineral assemblages with the gross composition and textures of sandstones, *Journal of Sedimentary Research*, 32, 440–450, 1962.

Jaskowiak-Schoeneichowa, M. and Krassowska, A.: Paleomiąższości, litofacje i paleotektonika epikontynentalnej kredy górnej w Polsce, *Geological Quarterly*, 32, 177–198, 1988.

1285 Jurkowska, A. and Barski, M.: Maastrichtian island in the Central European Basin-new data inferred from palynofacies analysis and inoceramid stratigraphy, *Facies* 63, 1–20, 2017.

Jurkowska, A., Barski, M., and Worobiec, E.: The relation of a coastal environment to early diagenetic clinoptilolite (zeolite) formation-New data from the Late Cretaceous European Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 524, 166–182, 2019a.

1290 Jurkowska, A., Świerczewska-Gładysz, E., Bąk, M., and Kowalik, S.: The role of biogenic silica in the formation of Upper Cretaceous pelagic carbonates and its palaeoecological implications, *Cretaceous Research*, 93, 170–187, 2019b.

Kamieński, M.: Przyczynki do znajomości kredy żurawieńskiej. Contribution à la connaissance du facies Sablonneux des caches cre'tace's de Żurawno (Pologne), *Kosmos*, 50, 1408–1425, 1925.

Kiersnowski, H., Paul, J., Peryt, T. M., and Smith, D. B.: 1995. Facies, paleogeography, and sedimentary history of the Southern Permian Basin in Europe. In: Scholle, P., Peryt, T. M., and Ulmer-Scholle, D. (Eds.), *The Permian of Northern Pangea. Volume 2: Sedimentary Basins and Economic Resources*. Springer, Berlin, 119–136.

1295 Kley, J. and Voigt, T.: Late Cretaceous intraplate thrusting in Central Europe: Effect of Africa-Iberia-Europe convergence, not Alpine collision, *Geology*, 36, 839–842, 2008.

1300 Komar, P., D.: The entrainment, transport, and sorting of heavy minerals by waves and currents, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, 3-48, 2007.

Kowalska, S., Kranc, A., Maksym, A., and Śmist, P.: Geology of the north-eastern part of the Carpathian Foredeep basement, the Lubaczów-Biszcza Region, *Nafta-Gaz* (in Polish), 54, 158–178, 2000.

1305 Krzywiec, P.: Miocene tectonic evolution of the Eastern Carpathian Foredeep Basin (Przemyśl–Lubaczów) in light of seismic data interpretation, *Prace Państwowego Instytutu Geologicznego* (in Polish), 168, 249–276, 1999.

1310 Krzywiec, P.: O mechanizmach inwersji brudzy śródkowopolskiej – wyniki interpretacji danych sejsmicznych (On mechanism of the Mid-Polish Trough inversion), *Bulletin of the Geological Institute of Poland*, 393, 135–166, 2000.

Krzywiec, P.: Mid-Polish Trough inversion – Seismic examples, main mechanisms and its relationship to the Alpine Carpathian collision. Continental collision and the tectonosedimentary evolution of forelands, *European Geophysical Society Special Publication*, 1, 151–165, 2002.

1315 Krzywiec, P.: Structural inversion of the Pomeranian and Kujawian segments of the Mid-Polish Trough–lateral variations in timing and structural style, *Geol. Q.*, 50, 151–168, 2006.

1320 Krzywiec, P.: Devonian–Cretaceous repeated subsidence and uplift along the Teisseyre–Tornquist zone in SE Poland—Insight from seismic data interpretation, *Tectonophysics*, 475, 142–159, 2009.

Krzywiec, P. and Stachowska, A.: Late Cretaceous inversion of the NW segment of the Mid-Polish Trough – how marginal troughs were formed, and does it matter at all?, *Z. Dtsch. Ges. Geowiss.*, 167, 107–119,

1325 Krzywiec, P., Gutkowski, J., Walaszczyk, I., Wróbel, G., and Wybraniec, S.: Tectonostratigraphic model of the Late Cretaceous–inversion along the Nowe Miasto–Zawichost fault zone, SE Mid-Polish Trough, *Kwartalnik Geologiczny*, 53, 27–48, 2009.

1330 Krzywiec, P., Peryt, T. M., Kiersnowski, H., Pomiczowski, P., Czapowski, G., and Kwolek, K.: Permo-Triassic evaporites of the Polish Basin and their bearing on the tectonic evolution and hydrocarbon system, an overview. In: Soto, J. I., Flinch, J. F., and Tari, G. (Eds.), *Permo-Triassic Salt Provinces of Europe, North Africa and the Central Atlantic: Tectonics and Hydrocarbon Potential*. Elsevier, Amsterdam, 243–261, 2017.

Krzywiec P., Stachowska A., and Stypa A.: The only way is up – on Mesozoic uplifts and basin inversion events in SE Poland, in: *Mesozoic Resource Potential in the Southern Permian Basin*, edited by: Kilhams, B., Kukla, P. A., Mazur, S., McKie, T., Mijnlieff, H. F., and van Ojik, K., Geological Society, London, Special Publications, 469, 2018.

1335 Kutek, J. and Głazek, J.: The Holy Cross area, Central Poland, in the Alpine cycle, *Acta Geologica Polonica*, 22, 603–652, 1972.

Kvale, E.P.: Tidal Constituents of Modern and Ancient Tidal Rhythmites: Criteria for Recognition and Analyses. In: Davis, Jr., R. A. and Dalrymple, R. W. (Eds.), *Principles of Tidal Sedimentology*, Springer, 1–17, 2012.

1340 Leszczyński, K.: The Upper Cretaceous carbonate-dominated sequences of the Polish Lowlands, *Geological Quarterly*, 41,

521–532, 1997.

1345 Leszczyński, K.: Lithofacies evolution of the Late Cretaceous basin in the Polish Lowlands, *Buletyn Państwowego Instytutu Geologicznego*, 443, 33–54, 2010 [in Polish with English summary].

Leszczyński, K.: The internal geometry and lithofacies pattern of the Upper Cretaceous–Danian sequence in the Polish Lowlands, *Geological Quarterly*, 56, 363–386, 2012.

1350 Leszczyński, K. and Dadlez, R. 1999. Subsidence and the problem of incipient inversion in the Mid-Polish Trough based on thickness maps and Cretaceous lithofacies analysis – discussion. *Przegląd Geologiczny*, 47, 625–628 [in Polish with English summary].

Lewis, J., Dodge, J. D., and Powell, A. J.: Quaternary dinoflagellate cysts from the upwelling system offshore Peru, Hole 686B, ODP Leg 112, *Proceedings of the Ocean Drilling Program, Scientific Results* 112, 323–328, 1990.

1355 Łuszczak, K., Wyglądała, M., Śmigielski, M., Waliczek, M., Matyja, B. A., Konon, A., and Ludwiniak, M.: How to deal with missing overburden - Investigating exhumation of the fragment of the Mid-Polish Anticlinorium by a multi-proxy approach, *Marine and Petroleum Geology*, 114, 104–229, 2020.

Machalski, M. and Malchuk, O.: Relative bathymetric position of the opoka and chalk in the Late Cretaceous European Basin, *Cretaceous Research*, 102, 30–36, 2019.

Mange, M. A. and Maurer, H. F. W.: *Heavy mineral in colour*, Springer Science & Business Media, 1992.

1360 Mange, M. A. and Morton, A. C.: Geochemistry of heavy minerals, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, pp. 345 - 391, 2007.

Mange, M. A. and Wright, D. T.: High-resolution heavy mineral analysis (HRHMA): a brief summary, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, pp. 433-436, 2007.

1365 Meres, Š.: Garnets - important information resource about source area and parental rocks of the siliciclastic sedimentary rocks, in: Conference “Cambelove dni 2008”. Abstract Book, edited by : Jurkovič, L., Comenius University, Bratislava, pp. 37-43, 2008 (in Slovak with English abstract).

1370 Narkiewicz, M., Resak, M., Littke, R., and Marynowski, L.: New constraints on the Middle Palaeozoic to Cenozoic burial and thermal history of the Holy Cross Mts. (Central Poland): results from numerical modelling, *Geologica Acta*, 8, 189–205, 2010.

1375 Narkiewicz, M., Maksym, A., Malinowski, M., Grad, M., Guterch, A., Petecki, Z., Probulski, J., Janik, T., Majdański, M., Środa, P., Czuba, W., Gaczyński, E., and Jankowski, L.: Transcurrent nature of the Teisseyre–Tornquist Zone in Central Europe: results of the POLCRUST-01 deep reflection seismic profile, *International Journal of Earth Sciences*, 104(3), 775–796, 2015.

Narkiewicz, M. and Petecki, Z.: Basement structure of the Paleozoic Platform in Poland, *Geological Quarterly*, 61(2), 502–520, 2017.

1380 Niechwedowicz, M., Cyglicki, M., and Remin, Z.: Zmiany zespołów cyst dinoflagellata z osadów deltowych środkowego kampanu Roztocza, SE Polska – implikacje środowiskowe, in: III Polski Kongres Geologiczny, Wrocław, 267–268, 2016.

Nielsen, S. B. and Hansen, D. L.: Physical explanation of the formation and evolution of inversion zones and marginal troughs, *Geology*, 28, 875–878, 2000.

1385 Nowak, J.: Przyczynek do znajomości kredy Lwowsko-Rawskiego Roztocza, *Kosmos*, 32, 160–169, 1907.

Nowak, J.: Spostrzeżenia w sprawie wieku kredy zachodniego Podola, *Kosmos*, 33, 279–285, 1908.

1390 Olde, K., Jarvis, I., Uličný, D., Pearce, M. A., Trabuco-Alexandre, J., Čech, S., Gröcke, D. R., Laurin, J., Švábenická, L., and

Tocher, B. A.: Geochemical and palynological sea-level proxies in hemipelagic sediments: A critical assessment from the Upper Cretaceous of the Czech Republic, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 435, 222–243, 2015.

1395 Omran, E. F.: The Nile delta: Processes of heavy mineral sorting and depositional patterns, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 49–74, 2007.

Pharaoh, T. C., Dusar, M., et al.: Tectonic evolution. In: Doornenbal, J. C. and Stevenson, A. A. (Eds.). *Petroleum Geological Atlas of the Southern Permian Basin Area*, European Association of Geoscientists and Engineers (EAGE), Houten, The Netherlands, 25–57, 2010.

1400 Pasternak, S. I.: *Biostratygrafiya kreydovykh vidkladiv Volyno-Podilskoi plyty*, Akademia Nauk Ukrainskoi RSR, Kiev, 3–98, 1959.

1405 Pasternak, S. I., Gavrylyshyn, V. I., Ginda, V. A., Kotsyubinsky, S. P., and Senkovskyi, Y., M.: *Stratygrafia i fauna kredowych vidkladiv zachodu Ukrayny*, Naukova Dumka, Kiev, 1–272, 1968.

Pasternak, S. I., Senkovskyi, Y. M., and Gavrylyshyn, V. I.: *Volyno-Podillya u kreydovomu periodi*, Naukova Dumka, Kiev, 3–258, 1987.

1410 Powell, A. J., Dodge, J. D., and Lewis, J.: Late Neogene to Pleistocene palynological facies of the Peruvian continental margin upwelling, Leg 112, *Proceedings of the Ocean Drilling Program, Scientific Results* 112, 297–321, 1990.

Powell, A. J., Lewis, J., and Dodge, J. D.: The palynological expressions of post-Palaeogene upwelling: a review, in: *Upwelling Systems: Evolution Since the Early Miocene*, edited by: Summerhayes, C. P., Prell, W. L., and Emeis, K. C., Geological Society Special Publication 64, 215–226, 1992.

1415 Pożaryski, W.: *Kreda*, in: *Regionalna geologia Polski*, 2, edited by: Książkiewicz, M. and Dżułyński, S., Polskie Towarzystwo Geologiczne, Państwowe Wydawnictwo Naukowe, Kraków, 1956.

Pożaryski, W.: *Zarys stratygrafii i paleogeografii na Niżu Polskim*, Prace Instytutu Geologicznego, 30, 377–418, 1960.

1420 Pożaryski, W.: *Kreda*, in: *Atlas geologiczny Polski. Zagadnienia stratygraficzno-facialne*, Instytut Geologiczny, 10, 1962.

Pożaryski, W.: *Tectonics. Part 1. Polish Lowlands*, in: Pożaryski, W. (Ed.), *Geology of Poland IV*, Wydawnictwa Geologiczne, Warszawa, 2–34, 1974.

1425 Preston, J., Hartley, A., Mange-Rajetzky, M., Hole, M., May, G., Buck, S. and Vaughan, L.: The provenance of Triassic continental sandstones from the Beryl Field, northern North Sea: mineralogical, geochemical, and sedimentological constraints, *Journal of Sedimentary Research*, 72, 18–29, 2002.

1430 Rebisco, M., Hernández-Molina, F. J., Van Rooij, D., and Wåhlén, A.: Contourites and associated sediments controlled by deep-water circulation processes: State-of-the-art and future considerations. *Marine Geology*, 352, 111–154, 2014.

Remin, Z.: The Belemnella stratigraphy of the Campanian–Maastrichtian boundary; a new methodological and taxonomic approach, *Acta Geologica Polonica*, 62, 495–533, 2012.

1435 Remin, Z.: The Belemnitella stratigraphy of the Upper Campanian–basal Maastrichtian of the Middle Vistula section, central Poland, *Geological Quarterly*, 59, 783–813, 2015.

Remin, Z., Cyglicki, M., Cybula, M., and Roszkowska-Remin, J.: Deep versus shallow? Deltaically influenced sedimentation and new transport directions – case study from the Upper Campanian of the Roztocze Hills, SE Poland, in: *Proceedings of the 31st IAS Meeting of Sedimentology, Kraków*, Poland, p. 438, 2015a.

1440 Remin, Z., Machalski, M., and Jagt, J. W. M.: The stratigraphically earliest record of *Diplomoceras cylindraceum* (heteromorph

ammonite) – implications for Campanian/Maastrichtian boundary definition, *Geological Quarterly*, 59, 843–848, 2015b.

Remin, Z., Gruszczyński M., and Marshall J. D.: Changes in paleo-circulation and the distribution of ammonite faunas at the Coniacian–Santonian transition in central Poland and Western Ukraine, *Acta Geologica Polonica*, 66 (1), 107–124, 2016.

1445 Remin, Z.: Understanding coleoid migration patterns between eastern and western Europe–belemnite faunas from the upper lower Maastrichtian of Hrebenne, southeast Poland, *Cretaceous Research*, 87, 368–384, 2018.

1450 Resak, M., Narkiewicz, M., and Littke, R.: New basin modelling results from the Polish part of the Central European Basin system: implications for the Late Cretaceous–Early Paleogene structural inversion, *International Journal of Earth Sciences*, 97, 955–972, 2008.

1455 Roberts, H. H. and Sydow, J.: Late Quaternary stratigraphy and sedimentology of the offshore Mahakam delta, East Kalimantan (Indonesia), in: *Tropical Deltas of Southeast Asia – Sedimentology, Stratigraphy, and Petroleum Geology*, edited by: Sidi, F. H., Nummedal, D., Imbert, P., Darman, H., and Posamentier, H. W., SEPM Special Publication, 76, 125–145, 2003.

Rogala, W.: O stratygrafii utwórzów kredowych Podola, *Kosmos*, 34, 1160–1164, 1909.

1460 Ryan, P. D., Mange, M. A., and Dewey, J. F.: Statistical analysis of high-resolution heavy minerals stratigraphic data from Ordovician of western Ireland and its tectonic consequences, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 465–489, 2007.

Samsonowicz, J.: Szkic geologiczny okolic Rachowa nad Wisłą, *Sprawozdania Państwowego Instytutu Geologicznego*, 3, 45–118, 1925.

1465 Shanmugam, G. Contourites: Physical oceanography, process sedimentology, and petroleum geology, *Petroleum exploration and development*, 44, 183–216, 2017.

1470 Storms, J. E., Hoogendoorn, R. M., Dam, R. A., Hoitink, A. J. F., and Kroonenberg, S. B.: Late-holocene evolution of the Mahakam delta, East Kalimantan, Indonesia, *Sedimentary Geology*, 180, 149–166, 2005.

Surlyk, F. and Lykke-Anderson, H.: Contourite drifts, moats and channels in the Upper Cretaceous chalk of the Danish Basin, *Sedimentology*, 54.2, 405–422, 2007.

1475 Świdrowska, J.: Kreda w regionie lubelskim–sedimentacja i jej tektoniczne uwarunkowania, *Buletyn Instytutu Geologicznego*, 422, 63–78, 2007.

Świdrowska, J. and Hakenberg, M.: Subsydencja i początki inwersji brzudzy śródłódzkiej na podstawie analizy map miąższości i litofacji osadów górnokredowych, *Przegląd Geologiczny*, 47, 61–68, 1999.

1480 Świdrowska, J., Hakenberg, M., Poluhić, B., Seghedi, A., and Višnákov, I.: Evolution of the Mesozoic basins on the southwestern edge of the East European Craton (Poland, Ukraine, Moldova, Romania), *Studia Geologica Polonica*, 130, 3–130, 2008.

1485 Świerczewska-Gładysz, E.: Late Cretaceous siliceous sponges from the Middle Vistula River Valley (Central Poland) and their palaeoecological significance, *Annales Societatis Geologorum Poloniae*, 76, 227–296, 2006.

Turner, G. and Morton, A. C.: The effects of burial diagenesis on detrital heavy mineral grain surface textures, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 393–412, 2007.

1490 Tyson, R. V.: Palynofacies analysis, in: *Applied micropaleontology*, edited by: Jenkins, D.G., Kluwer Academic Publishers, Dordrecht, The Netherlands, 153–191, 1993.

Tyson, R. V.: *Sedimentary organic matter. Organic facies and palynofacies*, Chapman and Hall, London, 615 pp., 1995.

1495 Velbel, M. A.: Bond strength and the relative weathering rates of Simple orthosilicates, *Amer. J. Sci.* 299, 679–696, 1999.

Velbel, M. A.: Surface textures and dissolution processes of heavy minerals in the sedimentary cycle: examples from pyroxenes and amphiboles, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 113-150, 2007.

1500 Velbel, M. A., Basso Ch. L., and Zieg, M. J.: The natural weathering of staurolite: crystal-surface textures, relative stability, and the rate-determining step, *American J. Sc.* 296, 453–472, 1996.

1505 Velbel, M. A., McGuire, J. T., and Madden, A. S.: Scanning electron microscopy of garnet from southern Michigan soils: etching rates and inheritance of pre-glacial and pre-pedogenic grain-surface textures, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 413-432, 2007.

Voigt, T., von Eynatten, H. and Franzke H-J. Late Cretaceous unconformities in the Subhercynian Cretaceous Basin (Germany). *Acta Geologica Polonica*, 54: 675–696, 2004.

1510 Voigt, T., Wiese, F., von Eynatten, H., Franzke, H-J., and Gaupp, R.: Facies evolution of syntectonic Upper Cretaceous Deposits in the Subhercynian Cretaceous Basin and adjoining areas (Germany). *Z. Dt. Geol. Ges.*, 157/2, 203-244, 2006.

1515 Voigt, T., Reicherter, K., von Eynatten, H., Littke, R. Voigt, S., and Kley, J.: Sedimentation during basin inversion. In: Littke, R., Bayer, U., Gajewski, D., and Nelskamp, S. (ed.): *Dynamics of complex sedimentary basins. The example of the Central European Basin System*, 211–232, 2008.

Voigt, T., Kley, J., and Voigt, S.: Dawn and Dusk of Late Cretaceous Basin Inversion in Central Europe, *Solid Earth*, 12, 1443–1471, 2021.

1520 Walaszczyk, I.: Turonian through Santonian deposits of the Central Polish Upland; their facies development, inoceramid paleontology and stratigraphy, *Acta Geologica Polonica*, 42, 1–122, 1992.

1525 Walaszczyk, I.: Inoceramids and inoceramid biostratigraphy of the Upper Campanian to basal Maastrichtian of the Middle Vistula River section, central Poland, *Acta Geologica Polonica*, 54, 95–168, 2004.

Walaszczyk, I. and Remin, Z.: Kreda obrzeżenia Góra Świętokrzyskich, *Przewodnik LXXXIV Zjazdu Polskiego Towarzystwa Geologicznego, Chęciny*, September 9–11, 41–50, 2015.

1530 Walaszczyk, I., Dubicka, Z., Olszewska-Nejbert, D., and Remin, Z.: Integrated biostratigraphy of the Santonian through Maastrichtian (Upper Cretaceous) of extra-Carpathian Poland, *Acta Geologica Polonica*, 66, 313–350, 2016.

1535 Wall, D., Dale, B., Lohmann, G. P., and Smith, W. K.: The environmental and climatic distribution of dinoflagellate cysts in modern marine sediments from regions in the North and South Atlantic Oceans and adjacent seas, *Marine Micropaleontology* 2, 121–200, 1977.

Wright, W. I.: The composition and occurrence of garnets. *American Mineralogist*, 23, 436-449, 1938.

1540 Wyrtki, K.: Indonesian throughflow and the associated pressure gradient, *Journal of Geophysical Research: Oceans*, 92(C12), 12941–12946, 1987.

Zigler, P. A.: *Geological Atlas of Western and Central Europe*, 2nd edn. Shell Internationale Petroleum Maatschappij, The Hague. Geological Society, London, 1990.

1545 Żelaźniewicz, A., Aleksandrowski, P., Buła, Z., Karnkowski, P., Konon, A., Oszczypko, N., Ślączka, A., Żaba, J., and Żytko, K.:

Regionalizacja tektoniczna Polski, 1–60, Komitet Nauk Geologicznych, Wrocław, 2011.

Żelaźniewicz, A., Buła, Z., Fanning, M., Seghedi, A., and Żaba, J.: More evidence on Neoproterozoic terranes in southern Poland and southeastern Romania, Geological Quarterly, 53, 93–124, 2009.

1550