

## Deep vs shallow – two contrasting theories? A tectonically activated Late Cretaceous deltaic system in the axial part of the Mid-Polish Trough: a case study from southeast Poland

Zbyszek Remin<sup>1</sup>, Michał Cyglicki<sup>1+2</sup>, Mariusz Niechwedowicz<sup>1</sup>

5 <sup>1</sup>Faculty of Geology, University of Warsaw, Al. Żwirki i Wigury 93, PL-02-089 Warsaw, Poland; <sup>2</sup>Polish Geological Institute – National Research Institute, Rakowiecka 4, PL-00-975 Warsaw, Poland; Corresponding Author: Zbyszek Remin; [zremin@uw.edu.pl](mailto:zremin@uw.edu.pl)

### Abstract:

10 The Polish Basin – a part of the trans-European sedimentary basin, i.e. Central European Basin System – stretching from Denmark through Germany to southeastern Poland, and even further to the south into Ukraine – experienced uplift during the Late Cretaceous, which consequently resulted in the inversion of its axial part (i.e. Mid-Polish Trough) and development into the Mid-Polish Anticlinorium. In many existing paleotectonic interpretations, southeast Poland – i.e., the subsurface San Anticlinorium and the present-day Roztocze Hills – was included into the Mid-Polish Trough during the  
15 Late Cretaceous, representing its axial, most subsiding part. This paleotectonic model was the basis for facies and bathymetric interpretations that assumed Upper Cretaceous sediments, deposited close to the axial part of the Mid-Polish Trough (e.g., in the Roztocze Hills), constituted the deepest facies. In recent years, several studies have contradicted this notion. A growing body of data suggests that the  
20 this area (southeast Poland) was already a landmass by the Coniacian and Santonian – and certainly in the Campanian and Maastrichtian – rather than forming the deepest part of the Polish Basin.

The shallow marginal marine, cyclic middle Campanian, marly to sandy deposits, recently discovered in the Roztocze Hills, were interpreted to be of deltaic origin. It has led to the adoption of new facies and bathymetric model, which – notably – are in stark opposition to most previous interpretations.  
25 Notably, the new interpretation implies the presence of a landmass in the place where, in prior frameworks, the deepest and most subsiding part of the Polish Basin (i.e. Mid-Polish Trough) was located during the Cretaceous.

Here, we document this Late Cretaceous depositional system – the Szozdy delta developed in the axial part of the Mid-Polish Trough. These middle Campanian deposits crop out extensively in the Roztocze  
30 Hills region, close to the village of Szozdy, and exhibit coarsening-upward tripartite cyclothems. Three facies associations have been distinguished: (1) dark grey calcareous mudstones, (2) a yellow calcareous sandstone, and (3) calcareous gale lithofacies. The sequence, as a whole, was accumulated via the repeated progradation and abandonment of deltaic complexes on the delta front platform setting, with the primary transport direction originating from the axis of the inverting Mid-Polish  
35 Trough (thus from the subsurface San Anticlinorium) toward the northeast. This interpretation is supported by a suit of sedimentological, palynofacies, and heavy mineral data.

The development of the Szozdy delta system is framed in the context of the dynamic tectonic processes operating contemporaneously in southeast Poland: that is, tectonic inversion (uplift) on one hand, and  
40 the generation of new accommodation space via enhanced subsidence on the other. This discovery

sheds new light on our understanding of Late Cretaceous facies distribution, bathymetry, paleogeography, and potentially might suggest a different burial history than thought so far.

45 Key words: Late Cretaceous, inversion tectonics, Szozdy delta, sedimentology, palynofacies, heavy minerals, provenance

## 1. Introduction

During the Late Cretaceous, the central and western Europe has been affected by compressional tectonics, which resulted in deformations observed across the whole Central European Basin System. This tectonic event induced the uplift and inversion of the basement units. The uplifted structures were eroded and material were redeposited in newly formed flexural basin (marginal troughs) (e.g. Zigler, 1990; Nielsen and Hansen, 2000; Krzywiec, 2006; Klay and Voigt, 2008; Voigt et al., 2021). Due to Late Cretaceous inversion tectonics, the paleogeographic and facies architecture of the Central European Basin System were markedly rebuild especially in a belt along the margin of the East European Platform (Fig. 1a).

The Polish Basin – a large, Mesozoic trans-European sedimentary basin alternatively known as the Mid-Polish Basin – underwent similar tectonic stress (inversion tectonics) at the whole Central European Basin System (for review compare: Krzywiec, 2002; Krzywiec et al., 2009, 2018). The Late Cretaceous paleotectonic evolution of the Polish Basin has been of considerable interest to geologists for more than a hundred years. The tectonic evolution and timing of the final tectonic inversion of the Mid-Polish Trough, especially its southeastern segment (Figs. 1b and 2a), is still a matter of active study, with no clear consensus reached to date (see reviews in Krzywiec et al., 2009; Walaszczyk and Remin, 2015; Krzywiec et al., 2018). The most important conclusion reached from these debates is that existing models developed over the past 50 years do not and cannot explain the entire geological history of this fragment of Poland (a part of the supposed Mid-Polish Trough), during the Mesozoic and, especially, the Late Cretaceous.

Over the past few decades, two independent paleotectonic models have been developed, focused mainly on the onset (and timing) of the inversion (uplift) of the Mid-Polish Trough (including its southeastern segment), and the consequent implications for facies and bathymetric (environmental) interpretations. Ultimately, the inversion-related uplift and subsequent erosion of the former axial part of the basin led to the formation of a prominent structural unit - the Mid-Polish Anticlinorium (Figs. 1b and 2a) (for reviews see Krzywiec et al., 2009; Walaszczyk and Remin, 2015; Krzywiec et al., 2018) with two synclinoria (troughs) bordering the Mid-Polish Anticlinorium toward NE (Kościerzyna-Puławy Synclinorium) and SW (Szczecin-Łódź-Miechów Synclinorium). The surrounding troughs (synclinoria) were formed in a way inially proposed by Nielsen and Hansen (2000).

According to the more traditional and so far widely-accepted interpretations, the southeastern edge of the Mid-Polish Trough (an axial part of the Polish Basin) (Fig. 2a), which is currently almost devoid of Mesozoic overburden (Fig. 2a) – represented its axial part and contained the deepest, most subsiding sedimentary environments during the Late Cretaceous (e.g., Kutek and Głazek, 1972; Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska 2007; Świdrowska et al., 2008). In turn, this paleotectonic model formed the basis for facies and bathymetric interpretations that largely suggest sediments deposited close to the axial part of the Polish Basin (i.e.

the Mid-Polish Trough), like those in the Roztocze Hills (southeast Poland) would constitute the deepest facies (Fig. 2c).

85 A series of studies performed over the past few years have supplied data in stark contradiction to the  
above interpretation, showing that already by the Coniacian and Santonian (and possibly even by the  
Late Turonian) the axial part of the Polish Basin (i.e. the Mid-Polish Trough) should be considered an  
emergent landmass or paleomorphological barrier, rather than as the deepest part of the basin,  
90 especially in the southeastern Poland (e.g. Krzywiec et al., 2009; Remin et al., 2015a; Walaszczyk and  
Remin, 2015; Remin et al., 2016; Krzywiec et al., 2018; Remin, 2018). This view is supported by the  
presence of clearly shallow-water facies along the northeastern edge of the present-day Mid-Polish  
Anticlinorium, especially in the southeastern Poland, thus in area where deepest facies should be  
95 expected according to the model of Kutek and Głazek (1972): these are the shallow water Turonian  
Janików Limestone and deltaically influenced sedimentation of the Roztocze Hills formed during the  
Campanian and Maastrichtian (Remin et al., 2015a; Walaszczyk and Remin, 2015). Additionally, seismic  
data show progradational bodies trending from the SW toward the NE (Krzywiec et al., 2009, 2018),  
thus extending from the supposedly axial part of the basin toward the northeast. However, all these  
data did not provide definitive proof for the existence of an emerged landmass in southeast Poland  
during the Late Cretaceous, and when treated individually their value and insights might be  
100 undermined.



In 2015a, Remin et al. provided a conceptual interpretation that the middle Campanian and  
?Maastrichtian siliciclastic deposits of the Roztocze Hills were of deltaic origin with the main source  
area located to the SW: this in a place where, according to the widely accepted paleotectonic model  
of Kutek and Głazek (1972) and several later authors, the axial and deepest part of the Mid-Polish  
105 Trough was located (Fig. 2c). The implementation of the new concept suggested a need for a dramatic  
revision to the existing paleotectonic model of southeastern part of the Polish Basin, and drove the  
adoption of a new facies and bathymetric model for several Late Cretaceous facies, one that was  
essentially the opposite of most previous interpretations (compare Figs. 2c and 2d; and discussion in:  
Krzywiec et al., 2009; Remin et al., 2015a; Walaszczyk and Remin, 2015; Krzywiec et al., 2018). Since  
110 this theoretical interpretation lacked precise, stratigraphically and sedimentologically-grounded  
corroboration, the present paper fills this gap by providing evidence to a wider audience that the  
development of the Szozdy deltaic system was induced by active inversion tectonics.

The study area in the Roztocze Hills gives direct insight into this special Campanian and Maastrichtian  
sedimentary succession, thus allowing for precise bio- and chronostratigraphic dating, essential for  
115 detailed sedimentological, facies, and paleotectonic interpretations.


These new datasets provide the foundation for the first direct sedimentological, petrographic, and  
palynofacies evidence for the presence of an emerged landmass in an area now devoid of Mesozoic  
cover – that is, the subsurface San Anticlinorium (the SE part of the Mid-Polish Anticlinorium; Fig. 1b)  
– which had previously been considered to be deeply submerged during the Late Cretaceous. These  
120 data, at least in part confirm the earlier concepts of e.g. Krzywiec et al. (2009, 2018) based on  
subsurface seismic data.

The objective of this paper is to show how a dynamic tectonic regime, including spatially variable  
inversion processes (uplift) and rapid subsidence, coupled with possible wave and/or bottom current

125 transport processes and significant terrigenous sediment input, drove the development of the Szody deltaic system. Its subaquatic remnants, originally developed in the axial part of the Mid-Polish Trough, are preserved in the rock record of the Roztocze Hills.



**Figure 1. (a)** Regional schematic map of western and central Europe (after Krzywiec et al., 2018 with further references) with the schematic extent of the Late Cretaceous - Paleocene inversion (patterned area); red rectangle – area shown in (b) and in figure 2; **(b)** General position of the Roztocze Hills and San Anticlinorium in the geological map of Poland excluding Cenozoic deposits, adopted from Dadlez et al. (2000) – except for the recent-day extension of the Carpathian Foredeep; MVVS = Middle Vistula Valley Section.



**Figure 2.** (a) A higher-resolution view of the Roztocze Hills area, relative to the position of the Holy Cross Mountains and Lower San Anticlinorium; (b) The general distribution of Campanian lithofacies in southeast Poland (adopted from Świdrowska, 2007): sandy limestone = deep; chalk = shallow; red dashed line – an interpreted cross-section of two competing tectonic concepts (c) and (d); (c) A facies and bathymetric sedimentation model based on traditional interpretations, assuming the deep-water character of deposits close to the axial part of the Mid-Polish Trough; vertical dashed-lines indicate areas devoid of Cretaceous deposits at present; (d) An alternative depositional, structural, and environmental interpretation of facies and bathymetry in a cross-section perpendicular to the axis of the Mid-Polish Anticlinorium (adopted from Remin et al., 2015a; Walaszczyk and Remin, 2015).

140

145

150 **2. The current state of the art: a short story of two contrasting theories**

Broadly speaking, the current literature hosts two theories regarding the onset of Mid-Polish Trough inversion tectonics and, subsequently, the formation of the Mid Polish-Anticlinorium (for a thorough review, see: Krzywiec et al., 2009; Walaszczyk and Remin, 2015; Krzywiec et al., 2018).

155 The first – and more widely accepted, at least historically – concept holds that the inversion of the Mid-Polish Trough began during the Late Maastrichtian and Paleogene (e.g., Kutek and Głazek, 1972; Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska, 2007; Świdrowska et al., 2008). Kutek and Głazek’s (1972) paper was indisputably a benchmark for interpretations of the Late Cretaceous paleogeographic and paleotectonic evolution of southeastern segment of the Polish Basin, and especially its axial part, i.e. Mid-Polish Trough.

160 Kutek and Głazek (1972) provided palinspastic restoration of a paleothickness of particular stratigraphic units in the cross-sections to the Mid-Polish Anticlinorium in the Holy Cross Mountains (to the NW from the present study area – compare Fig. 1b and 2a). In their framework, the increase in thickness of Mesozoic sedimentary successions toward the present-day Mid-Polish Anticlinorium (the area devoid of Mesozoic cover), (Fig. 1b and 2a), constituted their key argument for highest burial and  
165 the presence of a depocentre in the axial part of the Mid-Polish Trough, at least until the early Maastrichtian. This also implies that the area currently devoid of Mesozoic remnants was deeply buried by those deposits.

The general framework of Kutek and Głazek (1972) was widely accepted, and for decades dominated thinking about the Mesozoic paleotectonic evolution of the Polish Basin. In their model, the  
170 southeastern part of the Polish Basin, i.e. Mid-Polish Trough would represent the axial, most subsiding part of the Mid-Polish Trough during the late Cretaceous (compare Hakenberg and Świdrowska, 1998, 2001; Świdrowska and Hakenberg, 1999; Świdrowska, 2007; Świdrowska et a., 2008). The acceptance of this paleotectonic model has some foundational implications: for instance, it was widely assumed that the facies developed within the Mid-Polish Trough would represent the deepest sedimentary  
175 environment (e.g., Kutek and Głazek, 1972; Hakenberg and Świdrowska, 1998; Świdrowska 2007). As such, the facies and bathymetric interpretation was evidently fit to the presumptive paleotectonic model, rather than *vice versa*. It is noteworthy, however, that Upper Cretaceous mixed carbonate siliceous facies elude easy environmental interpretation, and the adopted paleobathymetric model and the spatial distribution of particular facies (Fig. 2c) were not supported by any sedimentological  
180 features.

Alternatively, another hypothesis has argued for a much earlier onset of inversion tectonics: in this framework, the inversion may have started in the Coniacian and/or Santonian, or perhaps even earlier, in the Late Turonian (e.g., Pożaryski, 1960, 1962; Jaskowiak-Schoeneichowa and Krassowska, 1988; Walaszczyk, 1992; Krzywiec et al., 2009; Leszczyński, 1997, 2010, 2012; Leszczyński and Dadlez, 1999;  
185 Walaszczyk and Remin, 2015; Remin et al., 2016; Krzywiec et al., 2018; Remin, 2018; Łuszczak et al., 2020).

Based on new sedimentological and chronostratigraphic data, Remin et al. (2015a) and Walaszczyk and Remin (2015) proposed a diametrically opposed sedimentological model (Fig. 2d). Their interpretation suggests the presence of a land area or paleomorphological barrier in the heart of Mid-Polish Trough, where the deepest part of the basin would have been located according to the model of Kutek and Głazek (1972). Furthermore, their proposed model (Fig. 2d) envisions that the deposits adjacent to the area of epigenetic erosion represent the shallowest facies, and pass into deeper facies towards the northeast (Fig. 2d) (Remin et al., 2015a; Walaszczyk and Remin, 2015).

A confirmation of this model (Fig. 2d) is provided by the presence of some specific facies developed locally along the northern edge of the Mid-Polish Anticlinorium: the Turonian Janików Limestone (sandy limestone) characterized by progradational sedimentary pattern and transport direction towards NE to the flank of the basin (Walaszczyk, 1992) and Campanian/Maastrichtian deltaically/sandy influenced sedimentation in the area of the modern-day Roztocze Hills; both are clearly indicative of a shallow-water environment. It is worth noting that the sandy facies of the Roztocze Hills have been mentioned by several authors (e.g., Kamieński, 1925; Jaskowiak-Schoeneichowa and Krassowska, 1988; Świdrowska et al., 2007; Leszczyński, 1997, 2010, 2012; and literature therein), but this prominent terrigenous input has never been explored sedimentologically to reveal their possible paleobathymetry and environment and source area.

The above interpretation is amply supported by new sedimentological and biotic data (Remin et al. 2015a; Cyglicki and Remin, 2016, 2018; Niechwedowicz et al., 2016; Remin et al., 2016; Remin, 2018). Additionally, extraordinarily preserved plant debris, including complete compounds leaves (Halamski, 2013) and characteristic shallow-marine palynofacies (this paper), are strong indicators of proximity to land. In further support of the existence of a paleobiogeographic barrier is its potential impact on paleocirculation, paleotemperature (as derived from oxygen isotopes), and the distribution of ammonite fauna (Remin et al., 2016), and the distribution pattern of belemnites during the early Maastrichtian (Remin, 2018).

All those surface observations are in line with seismic data interpretation that show an overall thinning Upper Cretaceous sedimentary cover towards SW – i.e. toward uplifted (at least partly) Mid-Polish Anticlinorium, especially in SE Poland (Krzywiec, 2009; Krzywiec et al., 2009, 2018). Additionally, those data also demonstrate local unconformities (Turonian – Maastrichtian) and progradational bodies directed from the axial part of the basin towards its NE flank (Krzywiec, 2009; Krzywiec et al., 2009, 2018).

It is noteworthy that these two concepts are based more or less on similar input data set – and in particular, the geometry and thickness patterns of sedimentary successions in cross-sections perpendicular to the Mid-Polish Anticlinorium (Fig. 2), however, the processing of those data and final interpretations were different.

Intriguing, this is not a novel interpretation. At the beginning of the 20th century, and indeed up until the 1960s (e.g., Rogala, 1909; Nowak, 1907, 1908; Kamieński, 1925; Samsonowicz, 1925; Pożaryski, 1960, 1962), the area of southeast Poland was interpreted as an emerged landmass during the Late Cretaceous. A proliferation of different names were adopted for this landmass, including the "Łysogóry-Dobrogea Land" of Samsonowicz (1925) and Jurkowska et al. (2019a, b); the "Maastrichtian island" (Jurkowska and Barski, 2017); the "Krukienic Island" of Pasternak (1959), Pasternak et al. (1968,

1987), Walaszczyk (1992), Dubicka et al. (2014) and the “Świętokrzyski Land” or “Małopolska Land” of Pożaryski (1960, 1962) and Jaskowiak-Schoeneichowa and Krassowska (1988), among others.

230

### 3. Regional setting

#### 3.1 The fill of the Polish Basin

235 The Polish Basin with its most subsiding axial part – i.e. Mid-Polish Trough represented the eastern part of the Central European Basin System (CEBS) formed during the Permian and Mesozoic in the epicontinental basins of western and central Europe (e.g. Pharaoh et al., 2010). In the Permian times the Polish Basin constituted the eastern part of the Southern Permian Basin (Kiersnowski et al., 1995; Krzywiec et al., 2017).

240 From Permian until the Late Cretaceous the Polish Basin experienced long term subsidence (Dadlez et al., 1995) what resulted in its filling of c. 8000 m of the Permian and Mesozoic deposits, especially along the NW-SE trending Mid-Polish Trough (e.g. Dadlez 1998) in the central part of the basin.

During the Late Cretaceous times and after the episode of long term Mesozoic subsidence, the Polish Basin and especially its axial part - the Mid-Polish Trough undergone uplift and inversion. The inversion tectonics started in the Late Turonian and lasted until the end of Maastrichtian (post-Maastrichtian) times (e.g. Krzywiec, 2000, 2002, 2006, 2009; Resak et al., 2008; Krzywiec et al., 2009, 2018).

245 The axial part of the Polish Basin (i.e. Mid-Polish Trough) has been transformed into prominent structural unit – Mid-Polish Anticlinorium (Fig. 1b). In the same time two marginal troughs were formed: Szczecin – Miechów Synclinorium to the SW and Kościerzyna – Puławy Synclinorium to the NE (Fig. 1b), feed by the elevated areas in the axial zone of the Polish Basin. The marginal troughs surrounding the Mid-Polish Anticlinorium were form in a manner initially proposed by Nielsen and Hansen (2000) for other inverted basins across Europe. As a result of the inversion procedure the Upper Cretaceous desposits are only preserved along both flanks of the Mid-Polish Anticlinorium (Fig. 1b).

255 The sedimentary cover of the southeastern part of the Polish Basin (SE Poland) is different in comparison to its central part where thick Permian, Triassic, Jurassic, and Cretaceous strata accumulated. Namely, only Middle/Upper Jurassic and Cretaceous deposits are well-documented (e.g. Świdrowska et al., 2008 for an overview) whereas the missing part of succession is ascribed to denudation.

260 In the SE Poland (study area), the Middle/Upper Albian (Lower Cretaceous) and the whole Upper Cretaceous succession reach c. 1000 m of deposits formed in rather shallow epicontinental basin (e.g. Jaskowiak-Schoeneich and Krassowska, 1988). Outcrops and well data enabled precise biostratigraphic subdivision based on inoceramids, ammonites, belemnites, echinoids, foraminifera, and to a lesser extent on other groups (Walaszczyk et al., 2016). The Upper Cretaceous succession is best exposed in 265 the NE peripheries of the elevated Mid-Polish Anticlinorium where is available in series of natural and artificial exposures, e.g. in the Middle Vistula Valley Section and in the Roztocze Hills area (Fig. 1b and 2a). Starting from the Cenomanian until the Maastrichtian the marine deposits cover most of

southeastern Poland and were ascribed as a siliciclastic-carbonate shelf system (e.g. Leszczyński, 1997, 2010, 2012).

270

### 3.2 The Roztocze Hills

The Roztocze Hills form a prominent range of hills (approximately 185 x 25 km) that extends from Kraśnik (southeast Poland) to Lviv in western Ukraine along the southwesterly margin of the Kościerzyna–Puławy Synclinorium in southeast Poland (Figs 1b and 2a). The current day Roztocze Hills are delineated along the southwest by a prominent escarpment, which constitutes a border zone with the subsurface northeastern edge of the San Anticlinorium underneath the Miocene cover of the present-day Carpathian Foredeep (Figs 1b and 2a) (e.g. Dziadzio et al., 2006 for further references). In the field, this boundary is manifested as an up to 50–80 m high escarpment, clearly visible from the almost flat surface of the Carpathian Foredeep towards the northeast. The pre-Miocene cover of the San Anticlinorium (southeastern part of the Mid-Polish Anticlinorium) is almost entirely devoid of Mesozoic remnants (with a few exceptions; Fig. 2a) and is represented mainly by strongly tectonized and weakly metamorphosed flysch-type sedimentary rocks of unknown thickness, in addition to anchimetamorphic rocks of Cambrian and Neoproterozoic age (e.g., Dziadzio and Jachowicz, 1996; Buła et al., 2008; Żelaźniewicz et al., 2009; Buła and Habryń, 2011).

275

280

285

290

This prominent tectonic line has been variously interpreted by different workers (e.g., Kutek and Głazek, 1972; Narkiewicz et al., 2015; Narkiewicz and Petecki, 2017). It is now considered that the southwestern edge of the Roztocze Hills were founded on crustally-rooted fault zones marked as nearly vertical crustal-scale conductive zones (review in Narkiewicz et al., 2015; Narkiewicz and Petecki, 2017), and were repeatedly reactivated during the Late Cretaceous inversion of the Mid-Polish Trough as a reversed faults (Krzywiec, 1999; Krzywiec et al., 2009) and the Carpathian tectonics during the Miocene (Kowalska et al., 2000; Buła et al., 2008; Buła and Habryń, 2011).

295

300

In the Polish part of the Roztocze Hills, the range is made up of Campanian (up to 550 meters thick) and Maastrichtian (c. 250 m or more, but incomplete) deposits; older deposits of the Lower/Upper Cretaceous succession are not exposed. From place to place the Upper Cretaceous is unconformably overlain by Miocene sediments (e.g., Pożaryski, 1956). Depending on the section, the Upper Cretaceous deposits of the Roztocze Hills are represented by various opoka facies (= siliceous chalk/limestone with a variable admixture of biogenic silica), gaizes (= siliceous limestone with a considerably admixture of detritical quartz, glaucony and clay), marls, calcareous sandstones, and calcareous mudstones, in addition to argillaceous mudstone or clays; the latter sometimes devoid of any CaCO<sub>3</sub> admixture. Regionally, the succession dips gently to the northeast.


### 3.3 The Szozdy section

The Szozdy section is situated within a railroad cutting of the Broad Gauge Metallurgical Railway Line, about 4 km southwest of Zwierzyniec close to the small village of Szozdy in southeast Poland (Fig. 3). The studied interval has yielded a rich fossil assemblage, comprising ammonites, belemnites, inoceramid and non-inoceramid bivalves, echinoids, diverse gastropods, and a suite of microfossils

305

(Remin et al., 2015a, b) in addition to plant debris (partially carbonized) like leaves or boughs (Figs. 3e-f).

310



315

**Figure 3.** (a) Location of the studied site in Europe; (b) Simplified geological sketch-map of extra-Carpathian Poland, excluding the Cenozoic cover (adopted from Pożaryski, 1974); tectonic units after Żelaźniewicz et al. (2011); (c) Detailed location of the studied section; (d) General view of the section – the exposures (white dashed lines) are at both sides of the railroad cutting at distance of approximately 800m; (e) Carbonized plant fragment (arrow) – the specimen is c. 10 cm across; (f) The imprint of tree bough (arrow) – the specimen is c. 25cm across.

320

This fossil assemblage defines a middle Campanian age for the Szozdy section (Remin et al., 2015a): specifically, the lower/middle *Didymoceras donezianum* ammonite Zone (*sensu* Błaszczewicz, 1980) and the middle "*Inoceramus tenuilineatus* inoceramid Zone (Walaszczyk, 2004). The interval available at Szozdy, together with other sections in the Roztocze Hills, is easily correlatable with the equivalent interval in the Middle Vistula River Valley section (for location compare Figs 1b and 2a), a reference

325 section for the Upper Cretaceous of the southeastern Polish Basin (e.g. Walaszczyk, 2004; Walaszczyk et al., 2016).



**Figure 4.** The stratigraphic position of the Szozdy section (right column; scale not included relative to the Middle Vistula River composite section (left column; scale bar = 2 m). A full description of the other sections of the Roztocze Hills and their stratigraphic position will be provided elsewhere. The biostratigraphic subdivision follows Błaszkiwicz (1980), Walaszczyk (2004), Remin (2012, 2015), Remin et al. (2015b), and Remin (2018).

#### 4. Material and Methods

The Szozdy section was sampled for 1) sedimentological evaluation, including both macro- and micro-scale observations; 2) heavy minerals, and 3) palynofacies analysis; the latter two mainly to examine hydrodynamic properties and paleoenvironmental context, respectively.

Of 22 total samples, seventeen (1-17) were subjected to detailed analyses: all of these were collected from the lower portion of the section (Fig. 5). This part of the section (cycles II-IV; Fig. 5), as being most densely sampled, was chosen to examine the differences and/or regularities governing the distribution of heavy minerals (Fig. 6) and palynofacies indicators, complementing sedimentological observations.

In the present study, the Szozdy section is placed within the wider context of CaCO<sub>3</sub>, quartz sand, clay content, plant debris, and thickness distribution patterns for the Campanian strata of the Middle and Eastern Roztocze Hills (Fig. 7).

The thickness pattern of the Campanian deposits in the Roztocze Hills area used in this study is based on 16 boreholes with documented Campanian deposits: Chrzanów IG-1, Dyle IG-1, Izbica IG-1, Jarczów IG-2, Jarczów IG-4, Komarów IG-1 – IG4, Narol PIG-1, Narol PIG-2, Rachanie IG-1, Stróża IG-1, Tarnawatka IG-1, Tomaszów Lubelski IG-1, and Ułhówek IG-1 (Fig. 8). The well-log data are stored in the Central Geological Data Base (CBDG; <http://baza.pgi.gov.pl/>) of the Polish Geological Institute - Polish Research Institute (PIG-PIB). The stratigraphic data, especially the lower and upper limit of the Campanian stage (where available) are based on data stored in the CBDG database for particular boreholes. Where necessary, well-log data were interpolated using the kriging and TopoToRaster geostatistic tools of ArcGIS.

**Sedimentology:** The scope of sedimentological analysis included both macro- and microscale (i.e., thin sections) observations. During fieldwork, the main lithofacies and thicknesses of the successive units were observed. Persistent sedimentary structures – i.e., a coarsening upward succession within the recognized cycles, trace fossils (bioturbation), macrofossil content, and micro-bioturbation (the latter within thin sections) – were also noted.

CaCO<sub>3</sub> contents were measured in the Polish Geological Institute (PIG-PIB) at over 50 localities (and c. three times more measurements). Thin section analysis enabled a volumetric estimation of the main granular components, including quartz, glauconite, and biogenic calcium carbonate. Silt and clay content were roughly estimated based on the volume of the granular and CaCO<sub>3</sub> components.

**Heavy minerals analysis:** In the present study, heavy mineral analysis includes both qualitative and quantitative analyses of heavy minerals assemblages (here, operationally defined as having a density > 2.9 g/cm<sup>3</sup>). A combination of optical microscopy, scanning electron microscope - energy dispersive X-ray spectroscopy (SEM-EDS), and electron microprobe (EPMA) analyses were applied (Mange and Maurer, 1992; Velbel, 1999, 2007; Velbel et al., 1996, 2007; Turner and Morton, 2007; Andò et al.,

2012; Garzanti and Andò, 2019). EPMA analyses included an examination of tourmaline and garnet geochemistry (e.g., Wright, 1938; Henry and Guidotti, 1985; Preston et al., 2002; Mange and Morton, 2007; Meres, 2008; Andò et al., 2013).

**Palynofacies analysis:** Following Tyson (1993, 1995), kerogen particles were subdivided into phytoclasts (opaque phytoclasts, translucent phytoclasts, cuticle), palynomorphs (terrestrial, marine), and amorphous organic matter (AOM). A minimum of 300 kerogen particles per sample were counted for palynofacies analysis. A preliminary analysis of palynomorph assemblages is focused on both sporomorphs (spores, pollen, saccate pollen) and aquatic palynomorphs (dinoflagellate cysts, foraminiferal test linings, acritarchs, algae, and freshwater algae), based on at least 250 counts per sample. Additionally, the terrestrial to marine (T/M) palynomorph index and peridinioid to gonyaulacoid (P/G) dinoflagellate cyst ratio (see, e.g., Olde et al., 2015) were also ascertained.

## 380 5. Results

### 5.1 Sedimentology – macro-and microscopic observations in the Szozdy section

The most prominent feature of the Szozdy section is the presence of tripartite cyclothem. Where complete, an individual cyclothem consists of three units: from bottom to top, calcareous mudstone, calcareous sandstone, and calcareous gyaize (Fig. 5). Not every cyclothem is complete within the study section: for instance, cyclothem II lacks the calcareous sandstone unit (Fig. 5).

Presently, we cannot estimate the total thickness of the Szozdy deposits represented by tripartite cyclothem, since neither the lower nor upper limit of the cyclic succession is known. The exposure available at Szozdy (Fig. 3d) gives access to approximately 30-35 meters of the succession at a distance of c. 800 m (the direct access to the section need some field works). Other natural or artificial exposures are not present; furthermore, well-log data are not available in close proximity to the Szozdy section. In the more distal boreholes, the Late Cretaceous rocks were cored only as cuttings: therefore, the existing well-log data preclude any firm conclusions. For the present study we choose 12-meters-thick interval that consists of six cycles including both complete and incomplete ones (Fig. 5).


The **calcareous mudstone lithofacies** (Fig. 5) is dark grey and poorly indurated. The color is most likely derived from disseminated carbonized organic matter. Of the distinguished lithofacies, the clay and silt content is highest in the calcareous mudstone, and ranges from 31% to 44% (mean = 38%). The sand fraction is composed mainly of quartz, and ranges from 16% to 26% (mean = 22%): it is the lowest within the whole section. The quartz is fine-grained, with a mean value of c. 0.1 mm; the grains are sub-angular to angular, with, less commonly, sub-rounded grains. In successive cyclothem, the CaCO<sub>3</sub> content of this unit varies over a relatively narrow interval (c. 34 – 46%; mean = 38%), and is ascribed mainly to broken biocomponents. In outcrops, small bioturbation structures are visible: other sedimentary structures are absent. In this unit, macrofauna are rare: only some bivalves and badly preserved echinoids were found.

The **calcareous sandstone lithofacies** (Fig. 5) is yellow to yellow-brownish and poorly indurated. The clay and silt content is c. 10% lower than in the underlying calcareous mudstone, ranging from 24% to 36% (mean = 29%). The sand fraction, largely consisting of quartz and, to a subordinate extent,

glaucanite, varies in a narrow interval (36 – 43%; mean = 40%): these values are the highest of the lithofacies recognized in the Szozdy section. The quartz sand is still fine-grained: however, the size of the grains is up to approximately 0.2 mm, twice as large relative to the underlying calcareous mudstones. The quartz grains are sub-angular to angular – sub-rounded grains are somewhat less common. The CaCO<sub>3</sub> content is c. 10% lower relative to calcareous mudstone, ranging from 26% to 38% (mean = 31%). In outcrop, this unit seems to be bioturbated at least to some degree; correspondingly, sedimentary structures are absent. Macrofauna are extremely rare in this lithofacies: similarly to the underlying calcareous mudstone, only some badly preserved bivalves were found.

The **calcareous gaize lithofacies** (Fig. 5) is white-gray, fully indurated, and can be extremely hard; in the field, it is expressed as protruding horizons between more erosion-prone deposits (Fig. 5). This unit is composed of a limited clay and silt fraction, ranging from 8% to 18% of the bulk rock (mean = 13%): these are the lowest clay and silt fractions of the lithofacies recognized in the Szozdy section. The sand fraction, consisting of quartz sand with, to a subordinate extent, glauconite, ranges from 21% to 35% (mean = 24%): the sand fraction is similar to that recognized in the underlying unit, but is fully cemented by a calcareous cement, with only a subordinate siliceous cement. The CaCO<sub>3</sub> content is the highest out of the recognized lithofacies: 60 – 75% (mean = 64%). Of the three lithofacies of the full cyclothem, only this lithofacies yielded well preserved marine fauna, consisting of relatively frequent ammonites (including baculites), inoceramids and, extremely rarely, belemnites and solitary corals. Other fauna include non-inoceramid bivalves and different snail species. It is worth mentioning that ammonites and other aragonitic shell material frequently retain remnants of the original shell structure, including iridescence. This unit also yielded abundant macroscopic plant debris, including tree boughs, tree branches, and relatively well-preserved compound leaves and/or their fragments.

430



435

**Figure 5.** Lithological column of the Szozdy section. Triangles on the left side of the column indicate coarsening-upward cyclothems. Top right: typical thin sections from successive lithofacies of a single cyclothem; scale bar = 500 μm; most of the white particles represent quartz grains. Bottom right: field photo of a part of the section with prominent layers of calcareous gaize, indicated on the section by red dashed lines. The horizontal scale represents relative resistance to weathering of particular units.

## 5.2 Heavy minerals


440 The heavy-mineral assemblages from the mixed calcareous siliciclastic deposits of the Szozdy section  
are characterized by a stable composition, with little differentiation. The analyzed associations  
included ultrastable and stable phases: in order of decreasing resistance to weathering under surface  
conditions, these include rutile, zircon, tourmaline, sillimanite, kyanite, staurolite, garnet, authigenic  
glauconite, pyrite, and other subordinate species such as chlorite, Cr-spinel, epidote, and single apatite  
445 grains.

The zircon-tourmaline-rutile (ZTR) maturity index (Hubert, 1962) ranges between 55-85% (mean =  
67%). The highest ZTR values are derived from the calcareous mudstone lithofacies, and driven mainly  
by the tourmaline abundance (dravite to shorl ratio, 4:1). In general, of the three ZTR index mineral  
species, tourmaline dominated in all of the studied samples. Amongst the ZTR minerals, subrounded  
450 and rounded crystals dominate (> 65%); to a much lesser extent, other ZTR crystals are angular or even  
euhedral (c. 4%). On the other hand, garnet, kyanite, and staurolite crystals are almost exclusively  
angular (> 70-80%), with almost no oval and/or rounded ones.

In the lower part of the Szozdy section (which, as described above, was chosen for detailed studies), a  
distinct rutile and tourmaline abundance pattern can be observed - namely, that these two mineral  
455 phases are inversely correlated (Fig. 6). It is worth noting that these mineral phases are of similar size  
c. 100  $\mu\text{m}$  in each unit. This is best observed in cyclothem III and IV (Fig. 6), which are the most densely  
sampled. In the calcareous mudstone unit of cyclothem III, the abundance of rutile is markedly  
reduced, whereas the tourmaline content is significantly higher than average. The opposite pattern is  
observed in the calcareous sandstone overlying the dark mudstone: here, rutile abundance is  
460 substantially elevated, while the tourmaline content is lower than average. A similar pattern as in the  
sandstone lithofacies is observed within the calcareous gaize lithofacies (Fig. 6). The only exception to  
this general trend is the calcareous gaize unit at the top of the (incomplete) cyclothem II, in which the  
sandstone lithofacies is absent. In this case, the rutile – tourmaline abundance trends follow the  
pattern recognized in the underlying mudstone.

465 To emphasize the relative changes in abundance of these two mineral species, a standardized Z-score  
statistic was calculated (Fig. 6). This statistic quantifies how many standard deviations each sample is  
from the mean value for the whole group for the entire Szozdy section (for comparison: Ryan et al.,  
2007).

470



**Figure 6.** Vertical chart of Z-score standardized rutile and tourmaline abundance in the Szozdy section. The vertical line represents the mean abundance of the analyzed minerals over the whole section. The bars are associated with the equivalent samples shown on the stratigraphic column; SD – standard deviation.

### 475 5.3 Palynofacies

Terrestrially sourced organic matter, represented by the phytoclast group (opaque phytoclasts, translucent phytoclasts, and cuticle), is the dominant palynofacies component throughout the studied succession. The AOM content is generally low. Among the terrestrially-derived palynomorphs, sporomorphs (spores, pollen, saccate pollen) are well-represented, while freshwater algae are rare.

480 Marine palynomorphs are represented by dinoflagellate cysts, foraminiferal test linings, acritarchs, and algae. Dinoflagellate cysts and foraminiferal linings are the most common marine palynomorphs, although their abundances fluctuate throughout the succession; acritarchs and algae are less common, but have relatively consistent abundances. The dinoflagellate cyst assemblages are rich and diverse; gonyaulacoid dinoflagellate cysts are generally more common than peridinioids.

485 **Calcareous mudstone.** In this lithofacies, palynofacies are dominated by phytoclasts (c. 85–93% of total kerogen), of which translucent phytoclasts (c. 33–44% of total kerogen) and cuticle (c. 22–42% of total kerogen) are the main constituents. The concentration of cuticle in calcareous mudstones is visibly higher than in other lithofacies; however, the cuticle debris is largely degraded. The content of AOM fluctuates, but at low concentrations (0.4–5% of total kerogen). The terrestrial/marine ratio is relatively high, ranging between 26% and 41%, and is ascribed mainly to the abundance of non-saccate pollen (up to 82% of sporomorphs). Foraminiferal linings (c. 44–64%) are clearly more abundant than the other marine palynomorphs. The P/G ratio is variable (11–39%), and indicates that peridinioid dinoflagellate cysts are comparatively less common in calcareous mudstones relative to calcareous sandstones.

495 **Calcareous sandstone.** This lithofacies is characterized by the dominance of phytoclasts (up to 79% of total kerogen), which are mainly represented by translucent phytoclasts (c. 43–63% of total kerogen). Of note are the very low concentrations of cuticle, which constitute only up to 3% of the total kerogen. The AOM content is generally low (c. 0.5–2% of total kerogen), but may occasionally reach higher values (c. 15%). Calcareous sandstones are characterized by the highest T/M ratios (up to 49%) in the entire section; elevated saccate pollen concentrations (up to 51% of sporomorphs) are notable. Marine palynomorphs are dominated by dinoflagellate cysts; peridinioids are visibly more common (P/G = 28–

500


45%) in calcareous sandstones compared to other lithofacies. Conspicuously, foraminiferal linings are almost entirely absent.

505 **Calcareous gaize.** Phytoclast concentrations, although still relatively high (c. 44–67% of total kerogen) and dominated by translucent phytoclasts (c. 18–44% of total kerogen), are visibly lower compared to the other lithofacies. Cuticle abundance is low to moderate (6–22% of total kerogen) and the cuticle debris is mostly degraded. The AOM content is generally high (c. 35–47% of total kerogen): however occasionally it may be somewhat lower (c. 7%). The T/M ratio (27–33%) is visibly lower compared to other distinguished lithofacies, with non-saccate pollen dominant over sporomorphs. Dinoflagellate  
510 cysts predominate over the other marine palynomorphs, and the P/G ratio exhibits moderate values (19–32%).

#### 5.4 Distribution of CaCO<sub>3</sub> and thickness patterns in Campanian deposits

To place the middle Campanian succession of the Szozdy section in a wider sedimentological context,  
515 two maps were constructed. The first (Fig. 7) shows the distribution pattern of CaCO<sub>3</sub> content coupled to terrigenous material (quartz sand and clay) content, and is based on more than 50 discrete outcrop observations (green triangles on Fig. 7) and c. three times as much of measurements. In most of the outcrops, two types of lithologies were available – less and more clayly. The map is constructed based on the lowest available measurements from particular outcrops. CaCO<sub>3</sub> content is more or less  
520 inversely correlated to quartz sand and clay content. Additionally, the distribution pattern of macroscopic plant debris is provided (Fig. 7), including imprints of tree boughs, tree branches, and leaves. The frequency of plant debris is described by two subjective classes – rare and common – based on field observations in particular outcrops (Fig. 7).

The lowest CaCO<sub>3</sub> values are concentrated in a relatively narrow belt, bordered to the southwest by  
525 the present Carpathian Foredeep (the subsurface northeastern edge the San Anticlinorium) (Figs. 2A and 7): correspondingly, this is the area with the highest abundance of terrigenous material (quartz sand and clay) and plant debris. Along a SW-NE transect, the CaCO<sub>3</sub> content progressively increases, whereas the quartz sand and clay content tend to decline towards the northeast. It is worth observing that in the southeast part of the studied area – i.e., the Horyniec Zdrój area (Fig. 7) – pure clay intervals  
530 may also occur in couplets with clayey opokas.



**Figure 7. (a)** Distribution of  $\text{CaCO}_3$  in the Roztocze Hills area between Zwierzyniec and the Polish/Ukrainian border to the east. Note that  $\text{CaCO}_3$  is consistently lowest at the edge of the Roztocze Hills, at the geologic border with the Carpathian Foredeep. Green triangles indicate findings of macroscopic plant fragments; **(b)** General map of Poland with study area. Squares with numbers represent the associated geological map sheets ([www.geologia.pgi.gov.pl](http://www.geologia.pgi.gov.pl)).


535

540

The second map shows the thickness pattern of Campanian deposits in the Roztocze Hills, based on 16 boreholes with appropriate interpolations (see methods for further details). It is worth mentioning that the original thicknesses, especially in the most southwesterly part which borders the Carpathian Foredeep (subsurface northeastern edge of the San Anticlinorium – compare Fig. 2a and 8) may be even higher, since the entire Campanian is not preserved locally. The highest values are located in the vicinity of the Narol boreholes (Fig. 8), where the Campanian reaches c. 550 m and along the

southeastern edge of the Roztocze Hills. To the northeast, the thickness of the Campanian deposits progressively decreases, declining to c. 150-200 m in the Komarów and Ulhówek boreholes (Fig. 8).

545



**Figure 8. (a)** Thickness pattern of Campanian deposits in the Roztocze Hills based on well data; **(b)** General map of Poland with study area. Squares with numbers represent the associated geological map sheets ([www.geologia.pgi.gov.pl](http://www.geologia.pgi.gov.pl)).

550

## 6. Interpretation and discussion

The study section is special: similar lithologies have not been described previously neither from the Roztocze Hills area nor from other part of the Polish Basin. Even the boreholes from the vicinity of study section did not provide similar sedimentological pattern. The Szozdy section is characterized by coarsening-upward tripartite cyclothems represented by mixed carbonate siliciclastic sediment rich in terrestrial components. The presence of frequent plant debris, i.e. imprints, carbonized fragments and disseminated organic matter of phytogenic origin, and scarcity of macroscopic, fully marine fauna in selected units of the cyclothems (the calcareous sandstone) point to its origination in close proximity to land areas and at least some interplay of marine and fresh waters (compare below). For those

555

560 reasons, Remin et al. (2015a) suggested a deltaically influenced sedimentation in the Szozdy area. Such  
an interpretation and especially its shallow, marginal marine sedimentary environment (Fig. 2d) were  
recently independently confirmed by by Jurkowska et al. (2019a, b) and Machalski (2019). Additionally  
the prominent quartz sand input and its progressive disappearance towards the NE direction (Fig. 7),  
thus outside the uplifted Mid-Polish Anticlinorium also suggest proximity of land which is considered  
565 to be present in a place of recent-day San Anticlinorium. Although the sandy-rich facies in the Roztocze  
Hills has been mentioned by several authors (e.g., Kamieński, 1925; Pożaryski, 1956, 1960, 1962;  
Jaskowiak-Schoeneichowa and Krassowska, 1988; Hakenberg and Świdrowska, 2001; Świdrowska,  
2007; Świdrowska et al., 2008; Leszczyński, 1997, 2010, 2012), these quartz/clay-rich deposits have  
never been explored sedimentologically, to place them within a paleobathymetric or  
570 paleoenvironmental and sedimentological context.

Although similar cyclic sedimentation can be recognized in several marginal marine environments  
(lower shorface, storm deposits, tide-dominated deltas and more offshore shelf deposits) we stay in  
opinion that the Szozdy section is most easily evaluated if a deltaic origin of those tripartite cyclothems  
is accepted. The upper and lower shorface together with storm deposits, if any, will produce cyclicity  
575 with normal grading, thus opposite to Szozdy case; the tide-dominated deltas are mainly characterized  
by mm- to cm-scale rhythmites (Klave, 2012); the more offshore settings might be preclude since there  
is no chance for the influence of fresh water suggested by the palynofacies analyses (compare below).

Besides macro- and microscopic sedimentological characteristics, the deltaic origin of the Szozdy  
section is here independently supported by the palynofacies analysis and hydrodynamic properties  
580 derived from heavy mineral analysis (as elaborated below) in addition to the distribution patterns of  
siliciclastic material and CaCO<sub>3</sub> in wider context (Fig. 7).

### 6.1 Palynofacies analysis

The characteristic palynofacies features of this succession include abundant phytoclasts (particularly  
585 translucent phytoclasts and cuticle), sporomorphs, peridinioid dinoflagellate cysts, and foraminiferal  
linings, all of which are indicative of a relatively proximal depositional setting. In particular, the  
abundance of translucent phytoclasts – that is, fresh, unoxidized particles – are suggestive of a very  
short and rapid transportation (Tyson, 1993). A similar inference may be made from the abundance of  
cuticle particles, which can easily be degraded and, thus, may serve as an indicator of rapid  
590 transportation (e.g., Tyson, 1993). The relatively high concentrations of sporomorphs, and particularly  
the presence of spores and non-saccate pollen, support this interpretation, as these sporomorphs are  
preferentially deposited in close proximity to land (Tyson, 1993). The elevated percentages of  
peridinioid dinoflagellate cysts and foraminiferal linings are likely related to an increased availability  
of nutrients, which may either originate from upwelling (e.g., Wall et al., 1977; Lewis et al., 1990;  
595 Powell et al., 1990, 1992; Eshet et al., 1994), or riverine discharge (e.g., Downie et al., 1971; Wall et  
al., 1977; Powell et al., 1990, 1992; Hardy and Wrenn, 2009). In the present case, the latter source of  
nutrients seems more probable, judging from the composition of the phytoplankton assemblage (e.g.,  
the absence of typically oceanic dinoflagellate cyst genera, such as *Impagidinium* and  
*Nematosphaeropsis*, and the presence of estuarine gymnodinioid dinocysts).

600 The cyclic fluctuations in the palynofacies and palynomorph assemblages characteristic of particular lithofacies suggest highly dynamic conditions. The most evident differences exist between the calcareous mudstones and sandstones (Fig. 9), as revealed by the changing relative percentages of phytoclasts (mainly cuticle), pollen, peridinioid dinoflagellate cysts, and foraminiferal linings. Such a sharp transition between the palynofacies patterns could be explained by either varying salinity level, 605 the different hydrodynamic properties of palynofacies components, or a mixture of both. Both dinoflagellate cysts and foraminiferal linings are indicative of relatively normal salinity conditions, although some dinoflagellates might have been tolerant of abnormal salinities in nearshore settings (Tyson, 1993). Hence, the rarity of foraminiferal linings in the calcareous sandstone units could possibly be explained by the influence of brackish conditions. The almost complete absence of cuticles in 610 calcareous sandstones (Fig. 9) may also indicate higher flow conditions, resulting in the continued suspension of the most buoyant particles, such as cuticle (see Tyson, 1993), and subsequent bypassing via floatation and deposition in more distal, lower-energy settings (e.g., calcareous mudstones and gaizes) (Fig. 10). On the other hand, the elevated percentages of saccate pollen in calcareous sandstones may suggest the opposite – a decrease in water energy, since saccate pollen are 615 preferentially concentrated in low-energy environments (see Tyson, 1995). Nevertheless, the slightly higher T/M and P/G ratios recorded in calcareous sandstones (Fig. 9) suggest that this lithofacies was apparently deposited more proximally than the calcareous mudstones and gaizes (Figs. 9 and 10).

The palynofacies patterns and palynomorph assemblages documented from calcareous mudstones and gaizes (Fig. 9) are less spectacular. In both lithofacies, the phytoclast content (including cuticle) is 620 comparable and relatively high. Dinoflagellate cysts and foraminiferal linings are also well represented in both units, although the proportion of the latter is visibly lower in calcareous gaizes. Gaizes are also characterized by elevated AOM concentrations (the highest values recorded throughout the succession), suggestive of a more distal setting for this lithofacies (see Tyson, 1993, 1995). Considering the above, it is reasonable to suggest that calcareous sandstones represent the most proximal setting, 625 while gaizes were apparently deposited in a more offshore environment (or eventually experienced less material supply); mudstones occupied an intermediate position (compare Fig. 10).


## 6.2 Hydrodynamic properties of the environment

The strong inverse relationship between tourmaline and rutile can be readily recognized throughout 630 the whole section (Figs. 6 and 9), indicating systematic changes in hydrodynamic power in the sedimentary environment.

The Z-score calculations, normalized to the whole study section, underline the changes in the proportion of respective minerals in succeeding samples, enabling a better understanding of the processes governing their distribution pattern within the section (Fig. 6) (e.g., Ryan et al., 2007). Both 635 minerals are of similar shape, durability and size: however, they possess markedly different densities – namely, 3.03-3.18 g/cm<sup>3</sup> and 4.23 g/cm<sup>3</sup> for tourmaline (dravite) and rutile, respectively. Since the weight of the two analyzed minerals is high, it might be expected that these two mineral phases will be strongly dependent, both vertically and spatially, on the local sedimentary environment – and by extension, the hydrodynamic power that actually existed during the deposition of the cyclothem. The 640 recurring increase in tourmaline abundance, coupled to a simultaneous decrease in rutile abundance

in muddy units, most likely resulted from a decrease in hydrodynamic power in the depositional environment. Sedimentologically, this may be interpreted as a transition to a comparatively more distal depositional setting from the main source area, e.g. a river discharge – that is, the prodelta or analogous environment (compare Figs. 9 and 10) (Komar, 2007; Omran, 2007).

645 Analogously, the increase in the proportion of rutile, and the simultaneous decrease of tourmaline share, in the sandy units can be linked to an increased flow rate, which might be interpreted as a transition to an environment more proximal to the river discharge, thus representing the main delta lobe and/or slope setting (compare Figs. 9 and 10). Simply put, the (lighter) tourmaline will be transported further to the prodelta environment, rendering the prodelta facies overrepresented in this  
650 mineral phase, whereas the (markedly heavier) rutile will be deposited closer to the river discharge.



**Figure 9.** The proposed environment for the successive lithofacies within an individual cyclothem in the Szozdy delta, coupled with the vertical distribution of selected heavy minerals and palynofacies indicators: cyclothem III adopted from Fig. 6. The values in brackets show the range for the lithofacies across all studied cyclothem. SD = standard deviation.

### 6.3 Sedimentary environment – looking for recent counterpart

660 We found that the modern Mahkanam deltaic system (Indonesia) may serve as a good modern counterpart for the Szozdy Delta sedimentary environment. The similarities are, of course, general – the Mahakam Delta is huge in comparison to the Szozdy deltaic system (although the spatial and vertical distribution of the latter is unknown); however, beyond those differences, several analogies can be found in their respective sedimentary environments.

665 The presence and relative importance of physical processes (e.g., water flow, sediment input, ocean currents, wind, waves, and tides) regulate the morphology and internal geometry of deltas. Accordingly, three main classes are commonly distinguished: i) tide-dominated, ii) wave-dominated, and iii) river-dominated deltas (e.g., Coleman and Wright, 1975; Galloway, 1975).

670 The modern Mahakam delta is considered a textbook example of a fluvial-tide-dominated delta system built across a narrow shelf (Galloway, 1975; Allen et al., 1977). The shelf and deltaic deposits border the N-S oriented Makassar Trough, which acts as a throughflow pathway for Pacific waters to the Indian Ocean (Wyrтки, 1987); as such, it must also regulate, at least in part, the redistribution of suspended sediments. The Mahakam River discharge is characterized by a distinct absence of flood

675 surges: therefore, the avulsion of distributary channels does not typically occur (e.g., Allen et al., 1977; Storms et al., 2005). Additionally, the extremely wide, submerged delta front platform, which extends as a 5-meter isobath up to 15-20 km offshore, works as an ideal platform to dissipate wave energy (Roberts and Sydow, 2003). As a consequence, the fluvial distributaries efficiently push suspended sediment basinward, resulting in extremely fast progradation (Storms et al., 2005, and references therein).


680 For the Szozdy Delta, it is difficult to ascertain which process(es) were dominant in the development of this deltaic system (fluvial-, tide-, and/or wave-dominated): at a minimum, such assessments are speculative to at least some degree. Although relatively large, the exposure at Szozdy – a 800 m long railroad cutting that provides access to approximately 30-35 vertical meters of the succession, still  
685 represents a single locality, with such a specific sedimentation style that can be best ascribed as deltaic in origin.

On the other hand, it is worth accounting for the fact that successive cyclothems are very regular, and particular units can be traced over dozens to few hundreds of meters. Strong wave and/or tidal action would most likely result in a more chaotic development of the successive lithofacies, and ultimately  
690 entire cyclothems. Accordingly, at the current stage, we can only speculate that fluvially-dominated processes are predominant, with possibly some additional wave and/or tidal action.

The limited thickness of the recurring cyclothems – 1 – 5 m (Fig. 5) – and upwards grain-size coarsening patterns in each cyclothem indicate that the progradation of succeeding facies – i.e., the transition between the more muddy and more sandy lithofacies of the cyclothems (Figs. 5 and 9) – proceeded  
695 relatively quickly. This also suggest that accommodation space was highly limited (Fig. 10).

Since there is no evidence for the presence of delta plain deposits, it is suggested that the entirety of sedimentation occurred in the subaquatic part of the Szozdy deltaic system, as exemplified in figure 10.

700 The implication is that, at least locally, the deposition of the submerged deltaic facies of the Szozdy delta took place along a relatively flat, wide, shallow to extremely shallow delta front platform (Figs. 10 and 11), similar to the delta front platform of the modern Mahakam delta, where progradation is fastest. The platform was wide enough to dissipate most of the wave-energy action, rendering the influence of waves low to moderate, at a maximum.



705 **Figure 10.** Block diagram illustrating the relationships between the subaerial (not available in the study area) and subaqueous deltaic environments of the Szozdy delta in relation to active and abundant distributary channels. Arrows 1 and 2 indicate two different situations i.e. 1) – a full tripartite cyclothem and 2) an incomplete cyclothem; compare to Fig. 5. Not to scale.

710 The differing thickness of successive lithofacies in the cyclothem (Fig. 5) suggests that specific environmental conditions had different durations before a threshold was reached that drove a change in paleoenvironmental regime from prodelta to delta lobe/slope facies. This suggests that rather dynamic tectonic conditions existed, resulting in changes of the delta architecture, the migration and avulsion of distributary channels, and simultaneous changes in the sedimentary environment at particular localities, resulting in the repetition of cyclothem.

720 According to this proposed environmental interpretation, the lowest unit of each cyclothem – that is, the calcareous mudstone – would represent the prodelta environment (Figs. 10 and 11), deposited on the submerged delta front platform in a comparatively distal region from the river discharge (Figs. 10 and 11). This unit possesses the highest clay abundances and the finest sand fraction. Upsection, the calcareous mudstone passes quickly into yellowish calcareous sandstone, with markedly less disseminated organic matter of phytogenic origin. In additional support of such an environmental interpretation is the scarcity of macroscopic, fully marine fauna: while some bivalves and echinoids were recognized in the mudstone lithofacies, in the sandstone lithofacies such fauna are almost absent.

725 The origin of the calcareous gaize is more problematic. This lithofacies is considered to originate in an area cut off from the sediment supply. Once the siliciclastic sediment supply is inhibited – for instance, via tectonic activity and/or changes in delta architecture (e.g., the avulsion of distributary channels) –

a relatively quick calcification process is thought to quickly lithify and “freeze” the upper portion of the sediment.

730 Three observations support such an interpretation. First, in all studied cases, the dominant grain size is always the same as the grain size observed in the underlying unit. In complete cyclothems [i.e., cyclothems I and III-VI], the gaize possess a grain size similar to the sandy unit (Fig. 5). In cyclothem II, where gaize directly overlies the muddy unit, the grain size is smaller and follows that observed in the mudstone. Second, in this unit, fully marine fauna are quite common, which indicates that the  
735 freshwater supply was comparatively limited or redirected. Additionally, terrestrial components are also abundant. In particular, macroscopic plant debris, including tree boughs, tree branches, and compound leaves, are commonly found: it is suggested that buoyant fragments were transported, sunk, and subsequently “frozen” by calcification. Third, ammonites and other aragonitic shelled animals still retain remnants of the original shell structure, including their iridescence. This indicates  
740 that they were cut off from the surrounding environment, which would have absurdly been corrosive to aragonite – in other words, the original aragonite could not have been preserved without syndepositional and/or early diagenetic calcification and chemical stabilization.

745

#### **6.4 Towards an understanding of Late Cretaceous facies distribution**


The very existence of a middle Campanian deltaic system developed in the axial part of the inverting Mid-Polish Trough (into the Mid-Polish Anticlinorium) has some fundamental consequences for our overall understanding of certain Late Cretaceous facies. In particular, this concerns mainly opoka,  
750 chalk, and their respective variants, in addition to their mutual spatial and bathymetric relationships in transects perpendicular to the Łysogóry-Dobrogea Land (Fig. 2d).

The evidence presented here for a shallow water, deltaic origin of these mixed carbonate siliciclastic deposits confirms the conceptual proposition of deltaically influenced sedimentation in the Rostocze  
755 Hills during the Campanian initially proposed by Remin et al. (2015a) and Walaszczyk and Remin (2015), and fully supports their general sedimentological and bathymetrical model (compare Fig. 2d). This bathymetric model and a shallow water position of the opoka facies (coastal environment) has been recently confirmed independently by Jurkowska et al. (2019a, b) and Machalski (2019).

760 In this respect a continuously growing San Anticlinorium during the Late Cretaceous acted as a source area for the siliciclastics of the Rostocze Hills. In the present study, for the first time, we can directly demonstrate that these terrigenous-rich deltaic sediments interfinger with gaize and opoka variants located slightly northeast of the recognized Szozdy deltaic system. It clearly shows that this specific facies – opoka – should be considered as having been deposited in a shallow-water environment, at  
765 least in this specific case, in opposition to previous notions rooted in different tectonic and sedimentological models, indicating its rather deep-origination. In the SW-NE transect perpendicular to the Szozdy deltaic system and the Łysogóry-Dobrogea Land (in a place of recent day San Anticlinorium), as defined here, the more carbonate-rich deposits, including chalk, were deposited in

770 more offshore and deeper zones located to the northeast of the Szozdy deltaic system. The presence  
of such a deeper zone is confirmed by the discovery of contourites (Krzywiec et al., 2009, 2018): the  
contourites current must have operated along a regional slope located northeast of both the Łysogóry-  
Dobrogea Land and the Szozdy deltaic system (Fig. 12).

775 Quite similar facies and structural situation we might find in the Subhercynian Cretaceous Basin which  
undergone inversion c. in the same time as the Mid-Polish Trough. The vast majority of the Coniacian  
to Campanian deposits, dominated by sandy and marly deposits, were derived from adjacent source  
areas, i.e. an elevated Harz Mountains (Voigt et al., 2004, 2006, 2008; von Eynatten et al., 2008). The  
Coniacian storm-dominated deposits and Santonian deposits of a low-energy delta plain to tidal-flat  
environment consisting of sands were deposited close to the areas elevated during inversion tectonics  
780 and pass into deeper facies in more offshore zones.



785 **Figure 11. (a)** CaCO<sub>3</sub> distribution pattern (blue area, compare Fig. 7) against the isolines of Campanian  
deposit thickness. Red dots = boreholes (compare Fig. 8); yellow dots = interpolation points; **(b)**  
General map of Poland with study area; **(c)** Subaerial view of the Szozdy deltaic environment in relation  
to the distribution pattern of selected facies bordering the deltaic environment – not to scale. Squares  
with numbers represent the associated geological map sheets ([www.geologia.pgi.gov.pl](http://www.geologia.pgi.gov.pl)).

790 For several decades, our understanding of facies distribution in the Upper Cretaceous of extra-  
Carpathian Poland was heavily influenced by the paleotectonic model developed and proposed by  
Kutek and Głazek (1972). Those authors *a priori* synonymized the axial part of the Mid-Polish Trough  
(in the present day represented as the Mid Polish Anticlinorium; Fig. 2b) with the most subsiding,  
deepest portion of the basin. It was consequently assumed that the facies deposited closest to the  
795 axial part of the Mid-Polish Trough (e.g., in the Roztocze Hills area) would represent the deepest  
sedimentary environment. As such, facies distributions and bathymetric interpretations were fit to this  
paleotectonic model (see also discussion in Walaszczyk and Remin, 2015). It is noteworthy, however,  
that Late Cretaceous mixed carbonate siliceous facies elude easy environmental interpretation, and  
the adopted paleobathymetric model and the spatial distribution of particular facies (compare Figs 2c),  
were not supported by any relevant sedimentological and/or biological indicators.

800 Nevertheless, the notion that opokas were largely of deep water origin (Fig. 2c) has been widely  
accepted, and for decades dominated thinking concerning the spatial distribution of facies and their  
relation to paleobathymetry (Leszczyński, 1997, 2010, 2012; Hakenberg and Świdrowska, 1998, 2001;  
Świdrowska and Hakenberg, 1999; Świdrowska 2007; Świdrowska et al., 2008; Świerczewska-Gładysz,  
2006). Accordingly, the Late Cretaceous opoka facies, the dominant facies of the Roztocze Hills, were  
805 considered as deep-water sediments until the present study, despite a considerable amount of  
terrigenous material (mainly sand-grained quartz, as well as clays). In light of the new data presented  
herein, this conception cannot be feasibly upheld, at least for the opoka facies of the Roztocze Hills.

### 6.5 Redistribution of terrigenous material


810 In a wider context, ocean currents (contourite currents) acting parallel to the shoreline of the identified  
Łysogóry-Dobrogea Land may have some influence on the redistribution of (mainly) the finest  
terrigenous material, pushing it away from the northern part of the Szozdy delta towards southeastern  
direction (Figs. 11 and 12).

The existence of contourite currents operating along this putative landmass during the Campanian was  
815 revealed based on seismic profiles perpendicular to the landmass postulated here (Krzywiec et al.,  
2009; 2018) (Fig. 12). In general, their appearance within the basin might be related to oceanographic,  
bathymetric condition or enhanced tectonic activity (inversion), regulating the morphology of the  
seafloor (e.g. Rebesco et al., 2014). Similar structures, based on seismic data, has been also proved  
from the Danish Basin (e.g. Surlyk and Lykke-Andersen, 2007). Their recognition provides independent  
820 evidence for the paleo-circulation within the Polish Basin as based on surface data.

Intriguingly, presence of currents flowing from the north and along the northern shore of the supposed  
paleobiogeographic barrier, interpreted here as a Łysogóry-Dobrogea Land (i.e., the area presently  
occupied by the San Anticlinorium), during the Coniacian and Santonian was suggested by Remin et al.  
(2016), who outlined its potential impact on paleocirculation, paleotemperature (as derived from  
825 oxygen isotopes), and the distribution of ammonite fauna. In further support of the existence of a  
paleobiogeographic barrier is the distribution pattern of belemnites during the early Maastrichtian  
(Remin, 2018).

830 The existence of contourite currents indicates that the Szozdy deltaic system was limited to the northeast by a regional slope in the sea-bottom; currents flowing parallel to the strike redistributed terrigenous material towards the southeast. A potential manifestation of this current-driven redistribution of the finest terrigenous material may be pure clay deposits of approximately the same age located to the southeast of the Szozdy delta – that is, in the Horyniec Zdrój area (Figs. 11 and 12). This area is additionally characterized by the lowest amount of CaCO<sub>3</sub> in the Campanian deposits of the Roztocze Hills, and high terrestrial input.

835 Such a paleogeographic situation is, again, somewhat similar to the modern Mahakam deltaic system that borders the Makassar Trough: that is a throughflow channel facilitating a contourite current (Shanmugam, 2017; Brackenridge et al., 2020), which redistributes material from the Mahakam prodelta to the south.



840 **Figure 12.** Schematic paleogeographic map and location of the Roztocze Hills and the Łysogóry-Dobrogea Land. The direction of ocean current flow is based on Remin et al. (2016) and Remin (2018); the presence of contourite currents (= CC) is according to Krzywiec et al. (2009, 2018).

## 6.6 Drivers of accommodation space generation

845 Inversion-related tectonic instability along the deeply rooted fault zone (the southwestern boundary of the Roztocze Hills; Fig. 2a) led to the development of the Szozdy deltaic system. To the southwest of this fault zone, thus in the area of the subsurface San Anticlinorium (Figs. 2a and 2d), there existed a landmass during the Late Cretaceous – i.e., the Łysogóry-Dobrogea Land – that was a source area for the siliciclastics of the Szozdy delta. During the process of tectonic inversion, this area underwent uplift, which surely resulted in changes to detrital weathering, weatherability, erosion, and provenance. Correspondingly, on the other side of this fault zone (that is, to the northeast) the Upper Cretaceous strata of the present Roztocze Hills were deposited, including the cyclothems of the

855 subaquatic delta front platform of the Szozdy deltaic system. This area, being a part of forming marginal trough, experienced enhanced subsidence, which resulted in the deposition of thick Campanian strata, especially along this active fault zone (compare Figs. 8 and 11).

860 Subsidence of the Roztocze Hills area, on the one hand, and uplift and erosion of the Łysogóry-Dobrogea Land on the other (in the area of the modern San Anticlinorium) during the Late Cretaceous inversion, could have conceivably driven the deltaic architecture, supplying additional accommodation space for the development of succeeding Szozdy delta cyclothems (Figs. 10 and 11). Furthermore, tectonically-induced migration and avulsion of distributary channels led to changes in the local sedimentary environment, which resulted in a predictable repetition of Szozdy cyclothems.

865

### **6.7 Speculative comments on burial history and the rate of subsequent erosion**

870 Since the paper of Kutek and Głazek (1972), the Holy Cross Mountains, and the San Anticlinorium to the southeast (Figs. 1b and 2a), have been traditionally included by most workers into the axial part of the former Mid-Polish Trough, which subsequently underwent inversion, uplift, erosion, and transformation into the Mid-Polish Anticlinorium. Apart from the concept of inversion tectonics onset itself, there was a general agreement that the axial part of the Mid-Polish Anticlinorium (e.g. area of the recent-day Holy Cross Mountains and the San Anticlinorium) were deeply buried by thick Permian and Mesozoic sediments (e.g., Kutek and Głazek, 1972). Kutek and Głazek (1972) suggested that c. 875 3000m of sedimentary rocks have been removed by erosion from the axial part of the Mid-Polish Anticlinorium, especially from the Holy Cross Mountains and the San Anticlinorium (Figs. 1b and 2a). More recent estimations varies between 2000-2500m (Resak et al., 2008; Narkiewicz et al., 2010). Similar values of overburden (1700-2300) were obtained by Łuszczak et al. (2020).

880 Since still little is known about possible Mesozoic overburden over the San Anticlinorium (this area is almost devoid of Mesozoic remnants; Figs. 1b and 2a), the heavy mineral assemblages from the Szozdy delta were analyzed to look for possible indicators of provenance. It provides intriguing data regarding the potential source rocks for the siliciclastics of the Szozdy deltaic system.

885 The heavy mineral assemblages from the Szozdy section exhibit a truly polycyclic nature. On the one hand, high values of the ZTR index (ascribed to ultrastable mineral phases), the high degree of roundness of (most) mineral grains, and the presence of some abrasive structures suggest multiple rounds of redeposition (Hubert, 1962; Mange and Wright, 2007; Garzanti, 2017; Garzanti and Andò, 2007, 2019). On the other hand, the presence of mineral groups weakly resistant to erosion – garnets, kyanites, and staurolites – that consists almost exclusively of angular crystals (> 70-80%) with almost no oval and/or rounded representatives is inconsistent with repeated redeposition; instead, they 890 support a multi-source provenance for the Szozdy deltaic deposits, as further elaborated below.

Accordingly, the above data suggest the presence of at least two distinct sources. The first yielded multi-recycled mineral phases dominated by minerals from the ZTR group, for which the source might be metapelites/metapsamites. The second, “fresh” source delivered angular-shaped crystals of garnets, kyanites, and staurolites, for which amphibolite facies were the main source.

895 This data suggest that the source rocks for the heavy mineral assemblages of Szozdy delta are represented by both (i) multi-recycled sedimentary rocks and (ii) freshly weathered metamorphic and igneous rocks. These characteristics would be rather difficult to obtain from the Mesozoic sedimentary cover of the San Anticlinorium (if it was even present at all), necessitating a search for other sources. A tempting hypothesis for the potential source rocks, although highly speculative, would be a comparison of mineral assemblages from the Szozdy delta to the Cambrian and Neoproterozoic rocks subcropping beneath the Miocene deposits of the Carpathian Foredeep, i.e. recent day San Anticlinorium, represented by metamorphosed (anchimetamorphic) flysch-type siliciclastics.

905

## 7. Summary

The interpretation presented herein implies the presence of a landmass area – that is, the Łysogóry-Dobrogea Land – in a place where, according to many previous interpretations, the deepest part of the Mid-Polish Trough was located. For the first time, clear and conclusive evidence for this iconoclastic paleogeographic and pelebathymetric model are provided, based on sedimentological data coupled with heavy mineral and palynofacies analyses. We recommend that the name Łysogóry-Dobrogea Land should be used, as it has priority (*vide* Samsonowicz, 1925).

915 The recognition of the Szozdy deltaic system in the supposedly axial part of the Mid-Polish Trough grants a unique perspective into the sedimentary environment, hydrodynamics, facies architecture, and bathymetric position of several Late Cretaceous facies. For instance, various styles of opoka facies are located adjacent to deltaic deposits, consistent with a shallow-water origin. A deeper sedimentary environment is suggested for chalk facies.

920

Inversion-related tectonic instability along the deeply rooted fault zone (the southwestern boundary of the Roztocze Hills) led to the development of the Szozdy deltaic system. Uplift of the present-day San Anticlinorium area during the Late Cretaceous inversion, on the one hand, and enhanced subsidence of the present-day area of the Roztocze Hills on the other, rearranged the delta architecture, supplying additional accommodation space for the development of Szozdy delta cyclothems.

925 The Łysogóry-Dobrogea Land, located to the south and southwest to the Roztocze Hills – thus, where the San Anticlinorium is presently situated – is the only reasonable source for the siliciclastics of the Szozdy deltaic system. This area is currently almost completely devoid of Cretaceous and other Mesozoic remnants and is now hidden under the Miocene cover of the Carpathian Foredeep.

930 The geochemistry of tourmalines and garnets (compare also Cyglicki and Remin, 2016, 2018) may suggest the strata subcropping beneath the Miocene series of the Carpathian Foredeep (to the

935 southwest of the Roztocze Hills) were already emerged and eroding during the Late Cretaceous, at  
least to some degree.

## 8. Current challenges and future roadmap

940 The results obtained in this study and from previous research raise several crucial issues that should  
be addressed in the near future, and furthermore allow us to formulate a working hypothesis that the  
modern San Anticlinorium was an elevated structural element during the Late Cretaceous. Such a  
hypothesis, although highly speculative, is supported by the initial geochemical data of heavy minerals  
we obtained, which might suggest that at least part of the material is derived from freshly weathered  
metamorphosed rocks. Further evidence and verification of these postulations would allow the  
945 following issues to be examined:

1. What was the source region for the Late Cretaceous Szozdy deltaic system and siliciclastics of  
the Roztocze Hills? Was it the Mesozoic cover of the San Anticlinorium? Or, alternatively, were the  
Cambrian and Neoproterozoic rocks beneath the Miocene cover of the Carpathian Foredeep already  
eroding during the Cretaceous?
- 950 2. Did the San Anticlinorium constitute a homogeneous part (structurally and  
paleogeographically) of the Mid-Polish Trough during the Cretaceous?
3. In this context, what was the real nature of the southeastern extension of this trans-European  
sedimentary basin, the Mid-Polish Trough, during the Cretaceous?
4. Accordingly, what was the burial history of the San Anticlinorium, if burial had occurred at all?

955 Deciphering these crucial objectives, based on a revised paleofacies and paleobathymetric model, is of  
fundamental significance for understanding the Late Cretaceous paleotectonic evolution of the  
supposed southeastern edge of the former Mid-Polish Trough.

In turn, a precise recognition of the stratigraphic succession and facies distribution (work which is  
currently ongoing) in the Roztocze Hills will allow for the reliable identification of the main phases of  
960 terrigenous material input from the nearby landmass, along with the estimation of subsidence  
gradients and the identification of these events on high-resolution seismic profiles (also ongoing). In  
this framework, a reduction of the subsidence rate should lead to the progradation of the siliciclastic  
facies toward the northeast, thus trending from more proximal to more offshore zones.

965 Solving the issues outlined above, especially with regards to resolving whether or not the San  
Anticlinorium constitutes a structural part of the Mid-Polish Trough, may fundamentally change our  
understanding of regional geology and burial history, and more broadly speaking the paleogeography  
and paleobiogeography of Central Europe during the Late Cretaceous, with far-reaching interpretative  
consequences.

970 *Author contributions.* ZR: idea, conceptualization, data compilation, discussion and validation, original  
draft, revision, funding acquisition (NCN; grant number UMO-2018/29/B/ST10/02947); MC:  
conceptualization, heavy minerals analyses, discussion and validation, revision; MN:  
conceptualization, palynofacies analyses, discussion and validation, revision.

975

*Competing interests.* The authors declare that they have no conflict of interest.

**Acknowledgements:** Our thanks are extended to Bartosz Gościmski and Joanna Roszkowska-Remin for help during the field work. Special thanks go to Robert Banaś, the chief forester of the Lubaczów district, and local forest officers who granted access to several outcrops and served with help in the field. Financial support was provided by the Polish National Science Centre [Narodowe Centrum Nauki, Polska] grant number UMO-2018/29/B/ST10/02947.

985

## 9. References

Allen, G. P., Laurier, D., and Thouvenin, J.: Sediment distribution patterns in the modern Mahakam delta, Proceedings Indonesian Petroleum Association, 5, 159–178, 1977.

990

Andò, S., Garzanti, E., Padoan, M., and Limonta, M.: Corrosion of heavy minerals during weathering and diagenesis: A catalog for optical analysis, *Sedimentary Geology*, 280, 165–178, 2012.

995

Andò, S., Morton, A. C., and Garzanti, E.: Metamorphic grade of source rocks revealed by chemical fingerprints of detrital amphibole and garnet, Geological Society, London, Special Publications, 386, 351–371, 2013.

Błaszkiwicz, A.: Campanian and Maastrichtian ammonites of the Middle Vistula River valley, Poland: a stratigraphic and paleontological study, *Prace Instytutu Geologicznego*, 92, 1–63, 1980.

1000

Brackenridge, R. E., Nicholson, U., Sapiie, B., Stow, D., and Tappin, D. R.: Indonesian Throughflow as a preconditioning mechanism for submarine landslides in the Makassar Strait. Geological Society, London, Special Publications, 500, 195–217, 2020.

1005

Buła, Z., Byś, I., Florek, R., Habryn, R., Jachowicz, M., Kwarcirski, J., Laskowicz, R., Liszka, B., Madej, K., Maksym, A., Markowiak, M., Pietrusiak, M., Probulski, J., Ryłko, W., Salwa, S., Sikora, R., Staryszak, G., Tabor-Wójcik, P., Tomasz, A., Zacharski, J.: Geological-structural atlas of the Palaeozoic basement of the Outer Carpathians and Carpathian Foredeep, Polish Ministry of Environment, Warszawa, 104, 775–796, 2008.

1010

Buła, Z. and Habryn, R.: Precambrian and Palaeozoic basement of the Carpathian Foredeep and the adjacent Outer Carpathians (SE Poland and western Ukraine), *Ann Soc Geol Pol*, 81, 221–239, 2011.

Coleman, J. M. and Wright, L. D. (Eds). *Modern river deltas: variability of processes and sand bodies*. 1975.

1015

Cyglicki, M. and Remin, Z.: Provenance of heavy minerals in the middle Campanian (Cretaceous) siliciclastic deposits of the Rostocze Hills, SE Poland, in: X Baltic Stratigraphic Conference, Chęciny, 28–29, 2016.

Cyglicki, M. and Remin, Z.: HRHMA (high-resolution heavy mineral analysis) zastosowana dla górno-kampańskich skał

silikoklastycznych krawędziowej części Roztocza Środkowego, SE Polska, Pokos, VII, 67, 2018.

- 1020 Dadlez, R., Marek, S., and Pokorski, J. (Eds.): Geological map of Poland without Cenozoic deposits at 1:1000 000 scale, Warszawa, Polish Geological Institute, 2000.
- Dadlez, R.: Mesozoic thickness pattern in the Mid-Polish Trough, *Geological Quarterly*, 47, 223–240, 2003.
- 1025 Dadlez, R., Narkiewicz, M., Stephenson, R. A. Visser, M. T. M., and van |Wess, J. D.: Tectonic evolution of the Mid-Polish Trough: modelling implications and significance for central European geology. *Tectonophysics*, 252, 179–195, 1995.
- Dadlez, R., Marek, S., and Pokorski, J. (Eds.): Palaeographical Atlas of the Epicontinental Permian and Mesozoic in Poland (1:2 500 000). Polish Geological Institute, Warszawa, Poland [in Polish with English summary], 1998.
- 1030 Downie, C., Hussain, M. A., and Williams, G. L.: Dinoflagellate cyst and acritarch associations in the Paleogene of southeast England, *Geoscience and Man* 3, 29–35, 1971.
- 1035 Dubicka, Z. Peryt, D., and Szuszkiewicz, M.: Foraminiferal evidence for paleogeographic and paleoenvironmental changes across the Coniacian–Santonian boundary in western Ukraine, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 401, 43–56, 2014.
- Dziadzio, P. and Jachowicz, M.: Budowa podłoża utworów miocenijskich na SW od wyniesienia Lubaczowa, *Przegląd Geologiczny*, 44, 1124–1130, 1996.
- 1040 Dziadzio, P., Maksym, A., and Olszewska, B.: Miocene deposition in the eastern part of the Carpathian Foredeep in Poland, *Przegląd Geologiczny*, 54, 413–420, 2006.
- 1045 Eshet, Y., Almogi-Labin, A., and Bein, A.: Dinoflagellate cysts, paleoproductivity and upwelling systems: A Late Cretaceous example from Israel, *Marine Micropaleontology* 23, 231–240, 1994.
- Eynatten, von H., Voigt, T., Meier, A., Franzke, H-J., and Gaupp, R.: Provenance of the clastic Cretaceous Subhercynian Basin fill: constraints to exhumation of the Harz Mountains and the timing of inversion tectonics in the Central European Basin. *Int. J. Earth Sci.*, 97, 1315-1330, 2008.
- 1050 Galloway, W. E.: Process framework for describing the morphologic and stratigraphic evolution of deltaic depositional systems, in: *Deltas, Models for Exploration*, edited by: Broussard, M.L., Houston Geological Society, Houston, TX, pp. 87–98, 1975.
- 1055 Garzanti, E.: The maturity myth in sedimentology and provenance analysis, *Journal of Sedimentary Research*, 87, 353-365, 2017.
- 1060 Garzanti, E. and Andò, S.: Heavy mineral concentration in modern sands: implications for provenance interpretation, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, pp. 517- 545, 2007.
- Garzanti, E. and Andò, S.: Heavy minerals for Junior Woodchucks, *Minerals*, 9(3), 148, 1-25, 2019
- 1065 Hakenberg, M. and Świdrowska, J.: Evolution of the Holy Cross segment of the Mid-Polish Trough during the Cretaceous, *Geological Quarterly*, 42, 239–262, 1998.
- Hakenberg, M. and Świdrowska, J.: Cretaceous basin evolution in the Lublin area along the Teisseyre-Tornquist Zone (SE Poland), *Annales Societatis Geologorum Poloniae*, 71 (1), 1–20, 2001.
- Halamski, A. T.: Latest Cretaceous leaf floras from Southern Poland and Western Ukraine, *Acta Paleontologica Polonica*, 58 (2), 407–433, 2013.

- 1070 Hardy, M. J. and Wrenn, J. H.: Palynomorph distribution in modern tropical deltaic and shelf sediments – Mahakam Delta, Borneo, Indonesia, *Palynology* 33, 19–42, 2009.
- 1075 Henry, D. J. and Guidotti, C. V.: Tourmaline as a petrogenetic indicator mineral: an example from the staurolite-grade metapelites of NW Maine, *American Mineralogist*, 70, 1–15, 1985.
- Hubert, J. F.: A zircon-tourmaline-rutile maturity index and interdependence of the composition of heavy mineral assemblages with the gross composition and textures of sandstones, *Journal of Sedimentary Research*, 32, 440-450, 1962.
- 1080 Jaskowiak-Schoeneichowa, M. and Krassowska, A.: Paleomiąższości, litofacje i paleotektonika epikontynentalnej kredy górnej w Polsce, *Geological Quarterly*, 32, 177–198, 1988.
- Jurkowska, A. and Barski, M.: Maastrichtian island in the Central European Basin-new data inferred from palynofacies analysis and inoceramid stratigraphy, *Facies* 63, 1–20, 2017.
- 1085 Jurkowska, A., Barski, M., and Worobiec, E.; The relation of a coastal environment to early diagenetic clinoptilolite (zeolite) formation-New data from the Late Cretaceous European Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 524, 166-182, 2019a.
- Jurkowska, A., Świerczewska-Gładysz, E., Bąk, M., and Kowalik, S.: The role of biogenic silica in the formation of Upper Cretaceous pelagic carbonates and its palaeoecological implications, *Cretaceous Research*, 93, 170–187, 2019b.
- 1090 Kamieński, M.: Przyczynek do znajomości kredy żurawieńskiej. Contribution a'la connaissance du facies Sablonneux des cauches cre'tace'es de Żurawno (Pologne), *Kosmos*, 50, 1408–1425, 1925.
- Kiersnowski, H., Paul, J., Peryt, T. M., and Smith, D. B.:1995. Facies, paleogeography, and sedimentary history of the Southern Permian Basin in Europe. In: Scholle, P., Peryt, T. M., and Ulmer-Scholle, D. (Eds.), *The Permian of Northern Pangea. Volume 2: Sedimentary Basins and Economic Resources*. Springer, Berlin, 119–136.
- 1095 Kley, J. and Voigt, T.: Late Cretaceous intraplate thrusting in Central Europe: Effect of Africa-Iberia-Europe convergence, not Alpine collision, *Geology*, 36, 839–842, 2008.
- Komar, P., D.: The entrainment, transport, and sorting of heavy minerals by waves and currents, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, 3-48, 2007.
- 1100 Kowalska, S., Kranc, A., Maksym, A., and Śmist, P.: Geology of the north-eastern part of the Carpathian Foredeep basement, the Lubaczów-Biszczka Region, *Nafta-Gaz (in Polish)*, 54, 158–178, 2000.
- 1105 Krzywiec, P.: Miocene tectonic evolution of the Eastern Carpathian Foredeep Basin (Przemyśl–Lubaczów) in light of seismic data interpretation, *Prace Państwowego Instytutu Geologicznego (in Polish)*, 168, 249–276, 1999.
- Krzywiec, P.: O mechanizmach inwersji bruzdy śródkowopolskiej – wyniki interpretacji danych sejsmicznych (On mechanism of the Mid-Polish Trough inversion), *Biuletyn Państwowego Instytutu Geologicznego*, 393, 135–166, 2000.
- 1110 Krzywiec, P.: Mid-Polish Trough inversion – Seismic examples, main mechanisms and its relationship to the Alpine Carpathian collision. Continental collision and the tectonosedimentary evolution of forelands, *European Geophysical Society Special Publication*, 1, 151–165, 2002.
- 1115 Krzywiec, P.: Structural inversion of the Pomeranian and Kuiavian segments of the Mid-Polish Trough–lateral variations in timing and structural style, *Geol. Q.*, 50, 151–168, 2006.
- Krzywiec, P.: Devonian–Cretaceous repeated subsidence and uplift along the Teisseyre–Tornquist zone in SE Poland—Insight from seismic data interpretation, *Tectonophysics*, 475, 142–159, 2009.

- 1120 Krzywiec, P. and Stachowska, A.: Late Cretaceous inversion of the NW segment of the Mid-Polish Trough – how marginal troughs were formed, and does it matter at all?, *Z. Dtsch. Ges. Geowiss.*, 167, 107–119,
- Krzywiec, P., Gutkowski, J., Walaszczyk, I., Wróbel, G., and Wybraniec, S.: Tectonostratigraphic model of the Late Cretaceous-inversion along the Nowe Miasto–Zawichost fault zone, SE Mid-Polish Trough, *Kwartalnik Geologiczny*, 53, 27–48, 2009.
- 1125 Krzywiec, P., Peryt, T. M., Kiersnowski, H., Pomianowski, P., Czapowski, G., and Kwolek, K.: Permo-Triassic evaporites of the Polish Basin and their bearing on the tectonic evolution and hydrocarbon system, an overview. In: Soto, J. I., Flinch, J. F., and Tari, G. (Eds.), *Permo-Triassic Salt Provinces of Europe, North Africa and the Central Atlantic: Tectonics and Hydrocarbon Potential*. Elsevier, Amsterdam, 243–261, 2017.
- 1130 Krzywiec P., Stachowska A., and Stypa A.: The only way is up – on Mesozoic uplifts and basin inversion events in SE Poland, in: *Mesozoic Resource Potential in the Southern Permian Basin*, edited by: Kilhams, B., Kukla, P. A., Mazur, S., McKie, T., Mijnlief, H. F., and van Ojik, K., Geological Society, London, Special Publications, 469, 2018.
- 1135 Kutek, J. and Głazek, J.: The Holy Cross area, Central Poland, in the Alpine cycle, *Acta Geologica Polonica*, 22, 603–652, 1972.
- Kvale, E.P.: Tidal Constituents of Modern and Ancient Tidal Rhythmites: Criteria for Recognition and Analyses. In: Davis, Jr., R. A. and Dalrymple, R. W. (Eds.), *Principles of Tidal Sedimentology*, Springer, 1–17, 2012.
- 1140 Leszczyński, K.: The Upper Cretaceous carbonate-dominated sequences of the Polish Lowlands, *Geological Quarterly*, 41, 521–532, 1997.
- Leszczyński, K.: Lithofacies evolution of the Late Cretaceous basin in the Polish Lowlands, *Biuletyn Państwowego Instytutu Geologicznego*, 443, 33–54, 2010 [in Polish with English summary].
- Leszczyński, K.: The internal geometry and lithofacies pattern of the Upper Cretaceous–Danian sequence in the Polish Lowlands, *Geological Quarterly*, 56, 363–386, 2012.
- 1145 Leszczyński, K. and Dadlez, R. 1999. Subsidence and the problem of incipient inversion in the Mid-Polish Trough based on thickness maps and Cretaceous lithofacies analysis – discussion. *Przegląd Geologiczny*, 47, 625–628 [in Polish with English summary].
- Lewis, J., Dodge, J. D., and Powell, A. J.: Quaternary dinoflagellate cysts from the upwelling system offshore Peru, Hole 686B, ODP Leg 112, *Proceedings of the Ocean Drilling Program, Scientific Results* 112, 323–328, 1990.
- 1150 Łuszczak, K., Wyglądała, M., Śmigieński, M., Waliczek, M., Matyja, B. A., Konon, A., and Ludwiniak, M.: How to deal with missing overburden - Investigating exhumation of the fragment of the Mid-Polish Anticlinorium by a multi-proxy approach, *Marine and Petroleum Geology*, 114, 104–229, 2020.
- Machalski, M. and Malchuk, O.: Relative bathymetric position of the opoka and chalk in the Late Cretaceous European Basin, *Cretaceous Research*, 102, 30–36, 2019.
- 1155 Mange, M. A. and Maurer, H. F. W.: *Heavy mineral in colour*, Springer Science & Business Media, 1992.
- Mange, M. A. and Morton, A. C.: Geochemistry of heavy minerals, in: *Heavy minerals in use*, edited by: Mange, M. and Wright, D. T., Elsevier, pp. 345 - 391, 2007.
- 1160 Mange, M. A. and Wright, D. T.: High-resolution heavy mineral analysis (HRHMA): a brief summary, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, pp. 433-436, 2007.
- Meres, Š.: Garnets - important information resource about source area and parental rocks of the siliciclastic sedimentary rocks, in: *Conference “Cambelove dni 2008”*. Abstract Book, edited by : Jurkovič, L., Comenius University, Bratislava, pp. 37-43, 2008 (in Slovak with English abstract).
- 1165 Narkiewicz, M., Resak, M., Littke, R., and Marynowski, L.: New constraints on the Middle Palaeozoic to Cenozoic burial and thermal history of the Holy Cross Mts. (Central Poland): results from numerical modelling, *Geologica Acta*, 8, 189–205, 2010.

- 1170 Narkiewicz, M., Maksym, A., Malinowski, M., Grad, M., Guterch, A., Petecki, Z., Probulski, J., Janik, T., Majdański, M., Środa, P., Czuba, W., Gaczyński, E., and Jankowski, L.: Transcurrent nature of the Teisseyre–Tornquist Zone in Central Europe: results of the POLCRUST-01 deep reflection seismic profile, *International Journal of Earth Sciences*, 104(3), 775–796, 2015.
- 1175 Narkiewicz, M. and Petecki, Z.: Basement structure of the Paleozoic Platform in Poland, *Geological Quarterly*, 61(2), 502–520, 2017.
- Niechwedowicz, M., Cyglicki, M., and Remin, Z.: Zmiany zespołów cyst dinoflagellata z osadów deltowych środkowego kampanu Rostocza, SE Polska – implikacje środowiskowe, in: III Polski Kongres Geologiczny, Wrocław, 267–268, 2016.
- 1180 Nielsen, S. B. and Hansen, D. L.: Physical explanation of the formation and evolution of inversion zones and marginal troughs, *Geology*, 28, 875–878, 2000.
- Nowak, J.: Przyczynek do znajomości kredy Lwowsko-Rawskiego Rostocza, *Kosmos*, 32, 160–169, 1907.
- 1185 Nowak, J.: Spostrzeżenia w sprawie wieku kredy zachodniego Podola, *Kosmos*, 33, 279–285, 1908.
- Olde, K., Jarvis, I., Uličný, D., Pearce, M. A., Trabucho-Alexandre, J., Čech, S., Gröcke, D. R., Laurin, J., Švábenická, L., and Tocher, B. A.: Geochemical and palynological sea-level proxies in hemipelagic sediments: A critical assessment from the Upper Cretaceous of the Czech Republic, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 435, 222–243, 2015.
- 1190 Omran, E. F.: The Nile delta: Processes of heavy mineral sorting and depositional patterns, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 49-74, 2007.
- 1195 Pharaoh, T. C., Duser, M., et al.: Tectonic evolution. In: Doornenbal, J. C. and Stevenson, A. A. (Eds.). *Petroleum Geological Atlas of the Southern Permian Basin Area*, European Association of Geoscientists and Engineers (EAGE), Houten, The Netherlands, 25–57, 2010.
- 1200 Pasternak, S. I.: Biostratygrafia kreydovykh vidkladiv Volyno-Podilskoi plyty, *Akademia Nauk Ukrainskoi RSR*, Kiev, 3–98, 1959.
- Pasternak, S. I., Gavrylyshyn, V. I., Ginda, V. A., Kotsyubinsky, S. P., and Senkovskiy, Y.,M.: *Stratygrafia i fauna kredowych vidkladiv zachodu Ukrainy*, Naukova Dumka, Kiev, 1–272, 1968.
- 1205 Pasternak, S. I., Senkovskiy, Y. M., and Gavrylyshyn, V. I.: Volyno-Podillya u kreydovomu periodi, *Naukova Dumka*, Kiev, 3–258, 1987.
- Powell, A. J., Dodge, J.D., and Lewis, J.: Late Neogene to Pleistocene palynological facies of the Peruvian continental margin upwelling, Leg 112, *Proceedings of the Ocean Drilling Program, Scientific Results 112*, 297–321, 1990.
- 1210 Powell, A. J., Lewis, J., and Dodge, J. D.: The palynological expressions of post-Palaeogene upwelling: a review, in: *Upwelling Systems: Evolution Since the Early Miocene*, edited by: Summerhayes, C. P., Prell, W. L., and Emeis, K. C., Geological Society Special Publication 64, 215–226, 1992.
- 1215 Pożaryski, W.: Kreda, in: *Regionalna geologia Polski*, 2, edited by: Książkiewicz, M. and Dżułyński, S., Polskie Towarzystwo Geologiczne, Państwowe Wydawnictwo Naukowe, Kraków, 1956.
- Pożaryski, W.: *Zarys stratygrafii i paleogeografii na Niżu Polskim*, Prace Instytutu Geologicznego, 30, 377–418, 1960.
- Pożaryski, W.: Kreda, in: *Atlas geologiczny Polski. Zagadnienia stratygraficzno-facjalne*, Instytut Geologiczny, 10, 1962.
- 1220 Pożaryski, W.: Tectonics. Part 1. Polish Lowlands, in: Pożaryski, W. (Ed.), *Geology of Poland IV*, Wydawnictwa Geologiczne,

Warszawa, 2–34, 1974.

- 1225 Preston, J., Hartley, A., Mange-Rajetzky, M., Hole, M., May, G., Buck, S. and Vaughan, L.: The provenance of Triassic continental sandstones from the Beryl Field, northern North Sea: mineralogical, geochemical, and sedimentological constraints, *Journal of Sedimentary Research*, 72, 18-29, 2002.
- 1230 Rebesco, M., Hernández-Molina, F. J., Van Rooij, D., and Wåhlin, A.: Contourites and associated sediments controlled by deep-water circulation processes: State-of-the-art and future considerations. *Marine Geology*, 352, 111-154, 2014.
- Remin, Z.: The Belemnella stratigraphy of the Campanian–Maastrichtian boundary; a new methodological and taxonomic approach, *Acta Geologica Polonica*, 62, 495–533, 2012.
- Remin, Z.: The Belemnella stratigraphy of the Upper Campanian–basal Maastrichtian of the Middle Vistula section, central Poland, *Geological Quarterly*, 59, 783–813, 2015.
- 1235 Remin, Z., Cyglicki, M., Cybula, M., and Roszkowska-Remin, J.: Deep versus shallow? Deltaically influenced sedimentation and new transport directions – case study from the Upper Campanian of the Roztocze Hills, SE Poland, in: *Proceedings of the 31st IAS Meeting of Sedimentology*, Kraków, Poland, p. 438, 2015a.
- 1240 Remin, Z., Machalski, M., and Jagt, J. W. M.: The stratigraphically earliest record of *Diplomoceras cylindraceum* (heteromorph ammonite) – implications for Campanian/Maastrichtian boundary definition, *Geological Quarterly*, 59, 843–848, 2015b.
- Remin, Z., Gruszczński M., and Marshall J. D.: Changes in paleo-circulation and the distribution of ammonite faunas at the Coniacian–Santonian transition in central Poland and Western Ukraine, *Acta Geologica Polonica*, 66 (1), 107–124, 2016.
- 1245 Remin, Z.: Understanding coleoid migration patterns between eastern and western Europe–belemnite faunas from the upper lower Maastrichtian of Hrebenne, southeast Poland, *Cretaceous Research*, 87, 368–384, 2018.
- Resak, M., Narkiewicz, M., and Littke, R.: New basin modelling results from the Polish part of the Central European Basin system: implications for the Late Cretaceous–Early Paleogene structural inversion, *International Journal of Earth Sciences*, 97, 955–972, 2008.
- 1250 Roberts, H. H. and Sydow, J.: Late Quaternary stratigraphy and sedimentology of the offshore Mahakam delta, East Kalimantan (Indonesia), in: *Tropical Deltas of Southeast Asia – Sedimentology, Stratigraphy, and Petroleum Geology*, edited by: Sidi, F. H., Nummedal, D., Imbert, P., Darman, H., and Posamentier, H. W., *SEPM Special Publication*, 76, 125–145, 2003.
- 1255 Rogala, W.: O stratygrafii utworów kredowych Podola, *Kosmos*, 34, 1160–1164, 1909.
- Ryan, P. D., Mange, M. A., and Dewey, J. F.: Statistical analysis of high-resolution heavy minerals stratigraphic data from ordovician of western Ireland and its tectonic consequences, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 465-489, 2007.
- 1260 Samsonowicz, J.: Szkic geologiczny okolic Rachowa nad Wisłą, *Sprawozdania Państwowego Instytutu Geologicznego*, 3, 45–118, 1925.
- 1265 Shanmugam, G. Contourites: Physical oceanography, process sedimentology, and petroleum geology, *Petroleum exploration and development*, 44, 183–216, 2017.
- Storms, J. E., Hoogendoorn, R. M., Dam, R. A., Hoitink, A. J. F., and Kroonenberg, S. B.: Late-holocene evolution of the Mahakam delta, East Kalimantan, Indonesia, *Sedimentary Geology*, 180, 149–166, 2005.
- 1270 Surlyk, F. and Lykke-Anderson, H.: Contourite drifts, moats and channels in the Upper Cretaceous chalk of the Danish Basin, *Sedimentology*, 54.2, 405–422, 2007.

- Świdrowska, J.: Kreda w regionie lubelskim-sedymentacja i jej tektoniczne uwarunkowania, *Biuletyn Instytutu Geologicznego*, 422, 63–78, 2007.
- 1275 Świdrowska, J. and Hakenberg, M.: Subsycjencja i początki inwersji bruzdy śródpolskiej na podstawie analizy map miąższości i litofacji osadów górnokredowych, *Przegląd Geologiczny*, 47, 61–68, 1999.
- 1280 Świdrowska, J., Hakenberg, M., Poluhtovič, B., Seghedi, A., and Višňakov, I.: Evolution of the Mesozoic basins on the southwestern edge of the East European Craton (Poland, Ukraine, Moldova, Romania), *Studia Geologica Polonica*, 130, 3–130, 2008.
- Świerczewska-Gładysz, E.: Late Cretaceous siliceous sponges from the Middle Vistula River Valley (Central Poland) and their palaeoecological significance, *Annales Societatis Geologorum Poloniae*, 76, 227–296, 2006.
- 1285 Turner, G. and Morton, A. C.: The effects of burial diagenesis on detrital heavy mineral grain surface textures, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 393–412, 2007.
- 1290 Tyson, R. V.: Palynofacies analysis, in: *Applied micropaleontology*, edited by: Jenkins, D.G., Kluwer Academic Publishers, Dordrecht, The Netherlands, 153–191, 1993.
- Tyson, R. V.: *Sedimentary organic matter. Organic facies and palynofacies*, Chapman and Hall, London, 615 pp., 1995.
- 1295 Velbel, M. A.: Bond strength and the relative weathering rates of Simple orthosilicates, *Amer. J. Sci.* 299, 679–696, 1999.
- Velbel, M. A.: Surface textures and dissolution processes of heavy minerals in the sedimentary cycle: examples from pyroxenes and amphiboles, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 113–150, 2007.
- 1300 Velbel, M. A., Basso Ch. L., and Zieg, M. J.: The natural weathering of staurolite: crystal-surface textures, relative stability, and the rate-determining step, *American J. Sc.* 296, 453–472, 1996.
- Velbel, M. A., McGuire, J. T., and Madden, A. S.: Scanning electron microscopy of garnet from southern Michigan soils: etching rates and inheritance of pre-glacial and pre-pedogenic grain-surface textures, in: *Heavy minerals in use*, edited by: Mange M. and Wright D. T., Elsevier, 413–432, 2007.
- 1305 Voigt, T., von Eynatten, H. and Franzke H-J. Late Cretaceous unconformities in the Subhercynian Cretaceous Basin (Germany). *Acta Geologica Polonica*, 54: 675–696, 2004.
- 1310 Voigt, T., Wiese, F., von Eynatten, H., Franzke, H-J., and Gaupp, R.: Facies evolution of syntectonic Upper Cretaceous Deposits in the Subhercynian Cretaceous Basin and adjoining areas (Germany). *Z. Dt. Geol. Ges.*, 157/2, 203–244, 2006.
- Voigt, T., Reicherter, K., von Eynatten, H., Littke, R. Voigt, S., and Kley, J.: Sedimentation during basin inversion. In: Littke, R., Bayer, U., Gajewski, D., and Nelskamp, S. (ed.): *Dynamics of complex sedimentary basins. The example of the Central European Basin System*, 211–232, 2008.
- 1315 Voigt, T., Kley, J., and Voigt, S.: Dawn and Dusk of Late Cretaceous Basin Inversion in Central Europe, *Solid Earth*, 12, 1443–1471, 2021.
- 1320 Walaszczyk, I.: Turonian through Santonian deposits of the Central Polish Upland; their facies development, inoceramid paleontology and stratigraphy, *Acta Geologica Polonica*, 42, 1–122, 1992.
- Walaszczyk, I.: Inoceramids and inoceramid biostratigraphy of the Upper Campanian to basal Maastrichtian of the Middle Vistula River section, central Poland, *Acta Geologica Polonica*, 54, 95–168, 2004.

- 1325 Walaszczyk, I. and Remin, Z.: Kreda obrzeżenia Gór Świętokrzyskich, Przewodnik LXXXIV Zjazdu Polskiego Towarzystwa Geologicznego, Chęciny, September 9–11, 41–50, 2015.
- Walaszczyk, I., Dubicka, Z., Olszewska-Nejbert, D., and Remin, Z.: Integrated biostratigraphy of the Santonian through Maastrichtian (Upper Cretaceous) of extra-Carpathian Poland, *Acta Geologica Polonica*, 66, 313–350, 2016.
- 1330 Wall, D., Dale, B., Lohmann, G. P., and Smith, W. K.: The environmental and climatic distribution of dinoflagellate cysts in modern marine sediments from regions in the North and South Atlantic Oceans and adjacent seas, *Marine Micropaleontology* 2, 121–200, 1977.
- 1335 Wright, W. I.: The composition and occurrence of garnets. *American Mineralogist*, 23, 436-449, 1938.
- Wyrtki, K.: Indonesian throughflow and the associated pressure gradient, *Journal of Geophysical Research: Oceans*, 92(C12), 12941–12946, 1987.
- 1340 Zigler, P. A.: *Geological Atlas of Western and Central Europe*, 2nd edn. Shell Internationale Petroleum Maatschappij, The Hague. Geological Society, London, 1990.
- Żelaźniewicz, A., Aleksandrowski, P., Buła, Z., Karnkowski, P., Konon, A., Oszczytko, N., Ślącza, A., Żaba, J., and Żytko, K.: Regionalizacja tektoniczna Polski, 1–60, Komitet Nauk Geologicznych, Wrocław, 2011.
- 1345 Żelaźniewicz, A., Buła, Z., Fanning, M., Seghedi, A., and Żaba, J.: More evidence on Neoproterozoic terranes in southern Poland and southeastern Romania, *Geological Quarterly*, 53, 93–124, 2009.