

Late Cretaceous – early Palaeogene inversion-related tectonic structures at the NE margin of the Bohemian Massif (SW Poland and northern Czechia)

Andrzej Głuszyński¹, Paweł Aleksandrowski^{2,1}

5 ¹Polish Geological Institute, Rakowiecka 4, PL 00-975 Warszawa, Poland

²University of Wrocław, Institute of Geological Sciences, Cybulskiego 32, PL 50-205 Wrocław, Poland

Correspondence to: Andrzej Głuszyński (andrzej.gluszynski@pgi.gov.pl)

Abstract. A brief, regional-scale review of the Late Cretaceous – early Palaeogene inversion-related tectonic structures affecting the Sudety Mts and their foreland at the NE margin of the Bohemian Massif is presented and complemented with
10 results of new seismic studies. The Sudetes expose Variscan-deformed basement, partly overlain by post-orogenic Permo-Mesozoic cover, containing a wide spectrum of tectonic structures, both brittle and ductile, in the past in this area referred to as young Saxonian or Laramide. We have used newly reprocessed legacy seismics to study these structures in the two main post-Variscan structural units of the area, the North-Sudetic and Intra-Sudetic synclinoria. The results were discussed together with regionally-distributed examples of tectonic structures coming from quarries and underground mines as well as
15 those known from the literature. The Late Cretaceous –early Palaeogene tectonic structures in consecutively reviewed Sudetic tectonic units, from the north to south, typically include gentle to moderate buckle folds, locally of detachment type, or fault-related, high-angle reverse and normal faults, as well as low-angle thrusts – often rooted in the crystalline basement. The structures termed grabens in the literature, are at the same time frequently interpreted as bounded by reverse faults (hence we use here the term ‘reverse grabens’) and typically reveal a strongly synclinal pattern of their sedimentary fill. The
20 top of crystalline basement, as imaged by seismic data in the North Sudetic Synclinorium below the faulted and folded cover, is synformally down-warped with a wavelength of up to 30 km, whereas on the elevated areas, where the basement top is exposed at the surface, it is up-warped (i.e. tectonically buckled). The compressional structures typically show an orientation fitting the regionally-known Late Cretaceous – early Palaeogene tectonic shortening direction of NE-SW to NNE-SSW. The same applies to the regional jointing pattern, typically comprising an orthogonal system of steep joints of c.
25 NW-SE and NE-SW strikes. All the reviewed structures are considered as due to the Late Cretaceous – early Palaeogene tectonic shortening episode, although some of the discussed faults with a strike-slip component of motion may have been modified, or even produced, by late Cenozoic, tectonism.

1 Introduction

30 During Late Cretaceous to early Palaeogene times, compressional tectonic structures developed over vast areas of western and north-central Europe (e.g. Kley and Voigt, 2008; Navabpour et al., 2017; Kley, 2018; Nádaskay et al., 2019; Malz et al., 2020; Voigt et al., 2021). They are currently interpreted as due to collisional interaction of the Iberian mass with continental Europe and the resultant propagation of far-field compressional stress (Kley & Voigt 2008), which brought about widespread inversion of Permo-Mesozoic basins. In Poland and easternmost Germany (e.g. Krzywiec, 2002, 2006; Mazur et al., 2005; 35 Krzywiec and Stachowska, 2016; Kley 2018) the resultant map-scale tectonic structures are mostly NW-SE-trending, large-scale, gentle folds up to 1000 km long, c. 150 km wide and up to at least 3 km high (e.g. Pożaryski, 1979). The Mid-Polish Swell (or Anticlinorium), spanning the entire breadth of Poland's NW-SE diagonal from the Baltic Sea to westernmost Ukraine, is the most spectacular example of such structures. To the SW of the Mid-Polish Swell there is the Szczecin-Łódź Trough (Synclinorium), and a successive elevated structural element, occurring at the NE margin of the Bohemian Massif, in 40 the borderland between Poland and Czechia. This complex element combines the Fore-Sudetic Homocline and the Sudetic area, composed of the Fore-Sudetic and Sudetic blocks (Fig. 1). The latter tectonic units are the principal object to be presented in this paper, whose goal is to briefly overview the wide spectrum of structural effects produced by the Late Cretaceous to early Cenozoic trans-European compressional event at the NE margin of the Bohemian Massif.

An important part of the review is based on recently reprocessed legacy seismic data from the central areas of the Sudetic 45 region. The other presented examples of tectonic structures supply the wide regional context for the seismic results and come either from our own field work or from descriptions made by other authors, though sometimes interpreted by us in a different way. Although the tectonic structures that formed at the turn of Cretaceous and Palaeogene times in the Permo-Mesozoic strata obviously must have developed also in the Variscan basement, they are – in general - difficult to distinguish from the older ones. Therefore, the scope of this paper is mostly limited to the tectonic phenomena that occur in the post-orogenic 50 Permo-Mesozoic strata, though, in places also in the late-orogenic upper Carboniferous. An overview of the Late Cretaceous – early Palaeogene deformation structures in the Sudetic area is presented in this paper, mostly in their regional context and in reference to particular structural units of the Sudetes, moving from the north to south across the area.

2 Geological outline

The Sudetes (Polish and Czech name: *Sudety*), low-topography mountains located on the northeastern margin of the 55 Bohemian Massif, together with the southern part of the Silesian-Lusatian Plain share the same pre-Cenozoic geological composition and structure and – in the geological sense – are for decades usually considered collectively as the Sudetic area or region, or, still, simply the Sudetes (e.g. Aleksandrowski and Mazur, 2002; Mazur et al., 2006, Kroner et al., 2008). The differences in topography – mountainous versus lowland - are due to splitting of the Sudetic area into two tectonic blocks: the currently downthrown and mostly flat or hilly Fore-Sudetic Block in the northeast and the elevated, mountainous Sudetic 60 Block in the southwest (Fig. 1). The Sudetes constitute the northeasternmost segment of the Central European Variscan

internides, exposing crystalline basement of strongly deformed and metamorphosed Late Neoproterozoic to Carboniferous rocks abundantly intruded by Carboniferous granites (e.g. Mazur et al., 2006, 2007, 2020). The Variscan Orogenic Belt formed in Europe at approximately Middle Devonian to late Carboniferous times and currently extends from Portugal and Spain, across Britain, France and Germany, to Czechia and Poland. It continues further to the southeast to Romania and
65 Anatolia and is also represented by numerous dismembered internal massifs within the Alpine-Carpathian orogenic system. Its formation was contemporaneous with the Acadian and Alleghenian orogenies in the Appalachians, whose particular major tectonostratigraphic zones define links to those of the European Variscan Belt across the Atlantic (e.g. Matte, 2001; Martinez Catalan et al., 2002; Kroner et al., 2008). In the Sudetes, the main tectonostratigraphic domains of the European Variscides: the Moldanubian, Tepla-Barrandian and – particularly – Saxothuringian find their continuation (e.g. Martinez-
70 Catalan et al., 2021).

The Variscan basement rocks are partly overlain by syn- to late-tectonic Carboniferous intramontane basins and remnants of the now much eroded post-orogenic Permo-Mesozoic sedimentary-volcanic cover. To the north, in the North Sudetic Synclinorium, there occur lower Permian red beds with volcanics and upper Permian Zechstein facies carbonates, sulphates and clastics, averaging c. 1000 m in thickness. The Lower Triassic Buntsandstein of variegated sandstones, totalling c. 600
75 m in thickness, are capped by Röt dolomites and later Keuper evaporites and Muschelkalk limestones (e.g. Baranowski et al. 1990; Chrzastek and Wojewoda 2011). To the south, in the Intra-Sudetic Synclinorium, lower Permian red beds and shales with volcanics up to 1250 m thick are followed by upper Permian, mostly fluviatile sedimentary rocks. The Lower Triassic is represented by the Buntsandstein facies, which does not exceed a few tens of meters in thickness (e.g. Dziedzic and Teisseyre 1990). The Sudetic area is described in the literature to have been emergent in its most part during the Triassic
80 (Feist-Burkhardt et al. 2008). In the Jurassic and Lower Cretaceous times the Sudetes must have also remained emergent, as no Jurassic and Lower Cretaceous strata are known from the area. In the Late Cretaceous, however a system of sedimentary basins formed along reactivated Variscan, mostly NW-SE-trending shear and fault zones over the northern Bohemian Massif, accumulating at least 1000 m-thick succession of shallow marine sediments (e.g. Scheck et al., 2002; Uličný et al., 2009; Wojewoda, 1997). The Intra- and North Sudetic synclinoria were filled with Cenomanian, Turonian, Coniacian and
85 Santonian sandstones, marls and siltstones (Wojewoda, 1997).

The Sudety Mountains acquired their present-day mountainous relief due to Neogene uplift of the Sudetic Block in front of the then actively growing Alps and Carpathians (e.g. Żelaźniewicz and Aleksandrowski, 2008; Jarosiński et al., 2009; Żelaźniewicz et al., 2011). This uplift affected an area mostly planated during the Palaeogene and early Miocene times, following the earlier, Late Cretaceous – early Palaeogene more prominent uplift, which at that time had occurred over a
90 broader area, including also the Fore-Sudetic Block (Fig. 1), which at present is downthrown along the Sudetic Boundary Fault (e.g. Cloos, 1922; Teisseyre, 1957; Oberc, 1972).

The Late Cretaceous – early Palaeogene uplift, that occurred concurrently with or slightly postdated the trans-European compressional event, related to the Africa-Iberia-Western/Central Europe convergence at ca 86-70 Ma (Kley and Voigt, 2008), exhumed the Variscan basement from below the post-Variscan, uppermost Carboniferous through Permo-Mesozoic

95 cover and left the entire Sudetic area tectonically elevated with respect to the adjoining depressed areas: the Fore-Sudetic Homocline to the northeast and the North Bohemian Cretaceous Basin to the southwest. Within both these tectonic units much of the deformed Permo-Mesozoic post-Variscan sedimentary fill has been preserved. Among the Late Cretaceous – early Palaeogene contractional structures in the Sudetes that have escaped erosion, are the North and Intra-Sudetic synclinoria, representing the largest objects of that kind (Fig. 2). For over a century, the Sudetes have been considered a part of the classical area in Central Europe, in which the term “Saxonische Tektonik” (cf. Kley, 2013) was applied to deformation structures, mostly fault-related, observed in the Permo-Mesozoic sedimentary basins. In the older literature these contractional structures were most often ascribed specifically to the “young Saxonian” (Closs, 1922; Beyer, 1939; Oberc, 1972, 1977) or “Laramide” tectonism (Oberc 1972, 1977).

3 Data and methods

105 The concise regional, though not fully systematic, review of the tectonic structures of likely Late Cretaceous – early Palaeogene age presented in this paper is based (1) on authors’ own analysis and structural interpretation of the newly reprocessed seismic reflection data located in the two key areas in the Sudetes containing the most complete post-Variscan sedimentary record and (2) on structural field data of the present authors or (3) on data coming from critically assessed literature descriptions. We have also used data from exploration boreholes drilled in the last five decades, prospecting for copper and coal deposits in the North Sudetic and Intra-Sudetic synclinoria, respectively.

The analysed seismic data were acquired by mining companies in the end 1970s (1976-1980) in the North Sudetic Synclinorium and in the early 1990s (1991-1993) in the Intra-Sudetic Synclinorium (Fig. 2). We received them as raw seismic data (field data, partly pre-processed, reloaded to discs from original magnetic tapes) and the accompanying paper prints. These seismic data are unique at the scale of the entire Sudetes as the source of valuable information on the structural geology of the Permo-Mesozoic succession in the both main Sudetic synclinorial units.

115 The reprocessing of, in total, c. 650 km of the seismic profiles was carried out in 2019, using an up-to-date oil-industry software (Głuszyński and Smajdor, 2020). The new processing included post-stack time migration (PostSTM), while to some of the profiles also the pre stack time migration stage (PreSTM) procedure was applied. No time-depth conversion was, however, attempted because of too scarce coverage of the area with appropriate drillhole data. As complementary material we used also an analogue/paper print version of a seismic profile from the vicinity of Broumov in Czechia and also some seismic profiles from the Fore-Sudetic Homocline. The most part of the structurally interpreted seismic profiles are presented in this paper. Our structural interpretation of these profiles provides an entirely new material showing the tectonic style and geometry of the compressional tectonic structures affecting the post-Variscan sedimentary cover along the NE margin of the Bohemian massif and allows for some inferences as to the mechanics of their formation. An equally important part of the paper, aimed at giving its readers an overall information on the distribution and genetic and geometrical diversity of the tectonic structures that formed or evolved under the Late Cretaceous – early Palaeogene compressional regime, are

depiction and short description of a selection of such structures exposed in natural outcrops and in active mines and quarries throughout the entire Sudetic area or its direct vicinities. This material is either our own or is based on critically evaluated literature accounts.

130

4. Products of Late Cretaceous – early Palaeogene tectonic shortening: examples from particular structural domains

4.1. The north-eastern margin of the Sudetes: transition to the Fore-Sudetic Homocline

135 Directly northeast of the Sudetic area, separated by the Middle Odra Fault Zone, the Fore-Sudetic Homocline extends (e.g. Kłapciński et al., 1984; Kroner et al., 2008; Żelaźniewicz et al. 2011). It is defined by gently (1-5°) NE-dipping Permian to Mesozoic strata (Figs. 2 and 3) on top of Variscan-folded Carboniferous (Mazur et al., 2006, 2010), representing the SW limb of the Szczecin-Miechów Synclinorium of the Polish-German Basin. The Permian-Mesozoic succession constitutes the fill of the extensive Polish-German Basin and unconformably overlies the Carboniferous fold-and-thrust flysch belt of the Variscan externides (e.g. Mazur et al., 2010, 2020).

140 The Permo-Mesozoic strata in the southern part of the Fore-Sudetic Homocline are very gently folded and in few places affected by steep to low-angle reverse faults (Fig. 4). In the close vicinity of the NW-SE-trending Middle Odra Fault system, however, in galleries of c. 0.5-1.2 km-deep mines of the Legnica-Głogów Copper District, a rich inventory of contractional and, also, extensional tectonic structures occur in the base Zechstein/top Rotliegend ore-bearing formation, containing clayey shale, carbonates, anhydrite and rock-salt, apart from sandstones. These structures typically include thrust faults, accompanied by bedding-parallel décollements, as well as folds, all indicative of NE-SW directed tectonic shortening. The meso-scale structures can be studied directly on the galleries' walls (Fig. 5; see e.g. Salski, 1965, 1968; Oberc and Salski, 1968; Dumicz and Don, 1977, 1990; Markiewicz and Szarowski, 1990; Żelaźniewicz and Markiewicz, 1991). Larger structures appear on mining maps as a complex network of faults, whose pattern reflects reactivation of a few major and numerous minor NW-SE trending fractures propagating upwards from the Variscan basement, accompanied by a number of relatively large WSW-ENE en-echelon fractures (cf. Markiewicz, 2007), the latter most probably formed due to a Late Cretaceous – early Palaeogene (or late Cenozoic?) sinistral strike-slip activity of the major NW-SE faults.

150 The southwestern margin of the Fore-Sudetic Homocline is the only area in the scope of interest of this paper, in which, due to the presence of rock salt in the Zechstein strata, the formation of Late Cretaceous – early Palaeogene structures may have been locally affected by salt tectonics.

155 4.2. The Fore-Sudetic Block

At the base Cenozoic level, the Fore-Sudetic Block (Figs. 2 and 3) exposes Variscan basement rocks of various metamorphic grades, varying in age from the Neoproterozoic to probable Devonian (Pożaryski, 1979; Kotański, 1997; Cwojdzński and Żelaźniewicz, 1995; Cymerman, 2010), and variable igneous plutonic rocks, mostly representing elements of a basic to

ultrabasic Silurian/Early Devonian ophiolitic suite and Carboniferous to early Permian granitoids (e.g. Mazur et al., 2006, 2007; Kroner et al., 2008). As the Fore-Sudetic Block is (except at its easternmost and westernmost peripheries) devoid of Permo-Mesozoic deposits and its crystalline basement is eroded deeper than that on the mountainous Sudetic Block, an idea was conceived long ago (Cloos, 1922; Teisseyre, 1957) that the former block had been uplifted with respect to the present-day Sudety Mountains across the Sudetic Boundary Fault, following the Late Cretaceous – early Palaeogene inversion. This situation was subsequently reversed in the Late Miocene, when the Sudety Mountains were uplifted whereas the Sudetic Block and the Fore-Sudetic Block were downthrown simultaneously with the formation of the Carpathian-Alpine forebulge (e.g. Żelaźniewicz et al., 2011). The lack of Permo-Mesozoic sedimentary cover hampers the direct recognition of the effects of the Late Cretaceous – early Palaeogene tectonism on the Fore-Sudetic Block, which, nevertheless, can be partly assessed thanks to the presence of the so-called Żary Pericline (Fig. 2) in the geological map intersection pattern. This pericline consists in map-view enveloping of the NW-projecting basement “peninsula” by the Permo-Triassic outcrop zone, which continues onto the Fore-Sudetic Block from the area of the Fore-Sudetic Homocline. This basement/cover intersection pattern illustrates the likely up-warping (updoming) effect of the Late Cretaceous – early Palaeogene compression, which seems to have undulated the roof surface of the crystalline basement, producing a very gentle NW-SE trending antiform. This antiform is analogous to the large, gentle synform that occurs to the south of it and is observable on seismic profiles – in the floor of the North-Sudetic Synclinorium (see below). The uplift related to the formation of the antiform in question, must have been one of the important factors leading to the erosion of the original Permo-Mesozoic cover over the most part of the Fore-Sudetic Block.

4.3. The North Sudetic Synclinorium

The North Sudetic Synclinorium (Figs. 2, 3 and 6) is a fault-bounded, NE-SE elongated tectonic structure, c. 60 km long and up to 24 km wide (Beyer, 1939; Oberc, 1972, 1977; Solecki, 1994; Kiersnowski *in* McCann et al., 2008a; Żelaźniewicz and Aleksandrowski, 2008; Żelaźniewicz et al., 2011). It is filled with post-Variscan, end Carboniferous-Permian-Triassic and Upper Cretaceous (Cenomanian to Coniacian) continental and shallow marine sedimentary rocks, including also a Lower Permian volcanic member (e.g. Śliwiński et al., 2003; Chrzastek and Wojewoda, 2011). The succession overlies a subsided top surface of the Variscan epimetamorphic basement, termed the Kaczawa Slate Belt in this region (e.g. Aleksandrowski and Mazur, 2002; Mazur et al., 2006; Żelaźniewicz and Aleksandrowski, 2008). The synclinorium acquired its geometry due to the Santonian-early Campanian to late Maastrichtian-Palaeocene (Walaszczyk *in* Voigt et al., 2008) tectonism. The North Sudetic Synclinorium in its western and central parts defines, in general, a single synclinal structure, while at its eastern and southern extremities it splits into several NW-SE elongated second-order synclinal elements, grabens and half-grabens, some of them separated by basement horsts (Figs. 2 and 7; Beyer, 1939; Śliwiński et al., 2003).

The end Mesozoic – early Cenozoic compressional deformation affected both the crystalline basement and post-Variscan cover. The cover must have once been relatively thick and must have extended over much wider areas of the Sudetic area

than it does today (cf. e.g. Migoń and Danišik, 2012; Sobczyk et al., 2015, 2020). It was subsequently eroded from the most uplifted areas, but preserved in places where the basement/cover interface had undergone compression-driven down-warping. According to Beyer (1939), this compression in the North Sudetic Synclinorium brought about inversion of some of the earlier normal faults, which were then transformed into reverse ones (Fig. 7a). The Variscan basement at the upthrown sides of the former inverted faults was believed by this author to have acted like rigid jaws of a vice, which horizontally squeezed the Permo-Mesozoic succession in the process of a “frame-controlled folding”(German *Rahmenfaltung*, Fig. 7b).

The structural geometry of the Late Cretaceous – early Palaeogene folds and related thrusts in the North Sudetic Synclinorium has been known so far mostly from geological map intersection patterns combined with drilling results (Beyer, 1939; Oberc, 1972, 1977; Leśniak, 1979; Solecki, 1986, 1994; Cymerman, 1998) and from supposed smaller-scale analogues occurring in active quarries (Fig. 8). The seismic data acquired in 1976-1980 made it possible to better constrain the structure of the post-Variscan cover, which was attempted by Bałazinska and Bossowski (1979; their Fig. 3) who assumed a dominance of nearly vertical faults.

A much better insight into the geometry and structural style of the deformed Permo-Mesozoic succession was recently enabled by the reprocessing of the same legacy seismic data (Głuszyński and Smajdor, 2019). The structural interpretation results of these data are presented in Figure 9 and depict the along-strike changing geometry of the Permo-Mesozoic succession of the North-Sudetic Synclinorium. This geometry, in general, seems to have resulted from gentle buckle downfolding of the basement/cover interface, producing a single syncline, up to c. 20 km wide, in the Permo-Mesozoic strata. The syncline is locally complicated by reverse faults (or their clusters), cutting across the Cretaceous, Triassic and Zechstein strata, and being hard to follow in the Rotliegend clastics and volcanics, though in some cases probably penetrating the epi-metamorphic basement, while shallowing their dips. The reverse faults are often related to folds, some of them likely having been initiated as fault-propagation folds. In places, decoupling/decollement phenomena occur along weak, clayey horizons in the Zechstein and Muschelkalk strata affected by local thrusting, accompanied by spectacular, but rather infrequent zones, where meso-scale detachment folds have developed (Figs. 8), apparently due to locally increased horizontal shortening in the footwall of steep reverse faults (Fig. 9g).

On neither of the seismic profiles studied by us any structures modified by salt tectonics were identified, in spite of a single rock salt intercalation in boreholes relatively close the profile shown in Fig. 9a.

The down-warping of the metamorphic basement below the Permo-Mesozoic cover may have facilitated the development of local low-angle thrusts in the basement (not visible in Fig. 9), where some of the thrusts affecting the Permo-Mesozoic are likely rooted. The *Rahmenfaltung* at the synclinorium edges (Fig. 7), invoked by Beyer (1939) and understood as upthrowing the relatively rigid metamorphic basement on reverse (inverted?) faults, may have also had its effect on the deformation process through exerting a horizontal-component tectonic force on the adjacent horizontal sedimentary strata. The local folds in the synclinorium interpreted from both the reprocessed seismic sections and from map intersection patterns, show, in general, near-parallel geometry, and shallow WNW-plunging axes (Solecki, 1986, 1994, 2011). Our analysis of the reprocessed seismic profiles made possible a subsurface mapping of a number of high-angle fault zones. They are, as a rule,

225 trending NW-SE and often continue over distances of tens of kilometres (up to at least 40 km). Other identified major faults represent low-angle thrusts, continuing into the Variscan basement. The polarity of the local thrust in the Permo-Mesozoic succession of the North-Sudetic Synclinorium is bimodal and directed both to the NE and SW. This seems typical of folding affecting a detached succession (the Permo-Mesozoic sedimentary rocks detached from the underlying low-grade metamorphic rocks). The interpreted fault zones cut across the Permo-Mesozoic strata from the Zechstein at the bottom to
230 the Upper Cretaceous at the top, which confirms their activity continued until at least the Late Cretaceous.

The structural analysis of the reprocessed seismics makes it possible to reinterpret the hitherto widely held concept of the internal structure for the North-Sudetic Synclinorium that assumed the dominance of high-angle fault block tectonics, although Solecki (1986, 1994) already earlier postulated that the main factor controlling the formation of the North Sudetic Synclinorium might have been a compressive inversion of hypothetical normal listric faults in the basement. In our opinion,
235 of primary significance in this structure are effects of compressional down-warping of the top basement surface due to buckling and of lesser significance - those related to both high-angle and low-angle thrust faulting, and to local detachment folding affecting the Permo-Mesozoic cover.

A recent mapping and structural study by Kowalski (2021) of one of the already mentioned second-order synclinal/graben elements branching off from the SE rim of the North Sudetic Synclinorium, the Wleń Graben (Figs. 2 and 7) offers an
240 opportunity to have a closer look on another example of a Late-Cretaceous – early Palaeogene macro-scale tectonic structure in the NW part of the Sudetes. The Wleń Graben is a narrow structure, c. 17.5 km long and up to 3.5 km wide, downthrown into the low-grade metamorphics of the Kaczawa Slate Belt, filled with Permian, Triassic and Upper Cretaceous (up to lower Coniacian) mostly sedimentary, shallow marine or continental deposits and bounded by steep, NW–SE-oriented, normal and reverse faults (Kowalski, 2021; Fig. 10). The graben was earlier studied by Gorczyca-Skała (1977) and discussed by Solecki
245 (1994, 2011), who stressed the particular significance of NNE-SSW to NE-SW directed compression in its formation. Kowalski (2021) advocates a multistage evolution of the graben, from Late Cretaceous times (post-Santonian?) onwards, including the first, extensional stage of its formation with active NW-SE striking normal faults, followed by compressional event of “probably latest Cretaceous to early Palaeogene(?)” age, re-activating and inverting the latter discontinuities and producing new reverse faults and overthrusts, particularly in the central and southern parts of the graben. This was postdated
250 by minor sinistral displacements of the graben fragments along cross-cutting faults of NE-SE trend, and the evolution is believed by this author to have ended with an extensional episode of limited significance, presumably already during the mid- to late Cenozoic.

The cross-sections of the Wleń Graben elaborated by Kowalski (2021) on the basis on his mapping fieldwork and scarce borehole data (Fig. 10) show a rather regular syncline, only slightly modified by the graben’s boundary faults and locally
255 complicated by faults striking obliquely to the syncline’s axis (Fig. 10). The metamorphic basement top is depicted as concordantly adherent to the overlying, synclinally bent base Rotliegend surface. The concordant folding of the basement and cover, together with the above mentioned apparent lack of a dip-slip displacement on the boundary faults of the Wleń Graben, suggest that the “graben’s” origin may have been rather due to downfolding (buckling) than downthrowing on

reverse faults. It is no wonder the cited author suggests a seismic survey to be made in order to better understand the structural geometry of the graben.

4.4. The Intra-Sudetic Synclinorium

The Intra-Sudetic Synclinorium is another, apart from the North Sudetic one, extensive Late Cretaceous – early Palaeogene tectonic structure of NW-SE trend in the Sudetes (Figs. 2, 3 and 11). It has a comparable, though somewhat larger size of c. 80 km in length and up to 30 km in width. It, similarly, affects the post-Variscan continental to shallow marine Permian-Mesozoic succession, which represents the upper structural level of the Intra-Sudetic Synclinorium (Żelaźniewicz and Aleksandrowski, 2008; Żelaźniewicz et al., 2011), unconformably deposited on top of a thick succession of intramontane Carboniferous syn- to late-orogenic clastics deformed by Variscan tectonism, corresponding to the lower structural level of the Intra-Sudetic Synclinorium (see also: Nemec et al., 1982; Dziedzic and Teisseyre, 1990; Bossowski et al., 1995; McCann, 2008b). The Carboniferous deposits of the Intra-Sudetic Synclinorium rest on the Variscan-deformed, mostly crystalline basement of the Sudetic Block. To the SE, the axial zone of the Intra-Sudetic Synclinorium, composed of Upper Cretaceous deposits, merges with the Upper Nysa-Králiky Graben (Figs. 2 and 11). Another similarity of the Intra-Sudetic to the North Sudetic Synclinorium is its generally simple, open structure of a single syncline in its northwestern and central parts, which is replaced by a number of synclinal grabens separated by basement horsts on its eastern flank (Figs. 2 and 11). Some of the synclinal grabens are bounded by reverse faults.

The Late Cretaceous to early Palaeogene tectonic structures of the Intra-Sudetic Synclinorium imaged by reprocessed seismic data, can be only seen in the NE limb of the synclinorium, located between Mioszów in the NW and Ścinawka in the SE. In that area it is the Permian rocks that are exposed at the surface, while the Cretaceous succession of the Intra-Sudetic Synclinorium axial zone occurs to the SW and has not been imaged by seismic data. The seismic profiles (location in Fig. 11) are mostly trending SW-NE, i.e. roughly perpendicular to the synclinorium's axis. They are consecutively presented, from NW to SE, in Figures 12 and 13, in order to illustrate in a systematic way the tectonic structures affecting the Permian and Carboniferous strata.

In the NW part of the area covered by seismic data (Figs. 11 and 12), three consecutive profiles show a group of planar faults in Carboniferous clastics, which do not continue upwards into the Permian, although in some profiles (Figs. 12 b and c) these faults manifest themselves also in the Permian strata as gentle folds above the faults. These folds are geometrically entirely consistent with fault-related folds in the Carboniferous. The most likely interpretation of this case seems to be a post-early Permian reverse-slip activity of faults in the Carboniferous succession, associated with gentle folding that locally affected also the overlying Permian strata, though in places the displacement on them entirely ceases still at a level below the Permian (Fig. 12a). The fault zone in the Carboniferous succession can be followed on seismic profiles over a distance of 7 km, but probably continues further on both to the NW and SE beyond the seismically explored area.

290 Further to the SE, near Broumov in Czechia (Figs. 11 and 13a), a low-angle, NE-vergent thrust zone (which may represent a more developed continuation of the fault zone from Fig. 12) rooted in the crystalline basement, produces a reverse-slip displacement of the Carboniferous strata by at least 2 km, associated with the formation of fault-bend folds. The thrust seems to continue to the NE as a décollement within the Permian strata, overlain by an anticline-syncline pair, which can be recognized in the surficial geological map outcrop pattern.

295 Moving further to the SE along the NE flank of the Intra-Sudetic Synclinorium axial zone, in the area of Radków, on the successive seismic profiles (Fig. 13b-e) one can follow along-strike geometrical changes of a conspicuous low-angle, NE-vergent thrust fault, responsible for up to 2.5 km of displacement of the crystalline basement and the overlying Carboniferous and Permian over their footwall. The overthrust is related – similar to the case at Broumov – with an anticline-syncline pair of fault-bend or fault-propagation(?) folds.

300 The conspicuous thrust fault identified at depth in the area of Radków – Ścinawka (Figs. 13b-e) and – most probably – continuing into the vicinities of Broumov (Fig. 13a), seems to correspond to the major NW-SE trending Ścinawka-Krosnowice Fault (Fig. 11) exposed at the surface. The latter fault has been, however, so far consistently interpreted as a NE-throwing normal (Grocholski and Augustyniak, 1971) or vertical fault (Bossowski and Ihnatowicz, 2006). Therefore, our interpretation of the reprocessed seismic data implies an important change in understanding the geometry and kinematics of

305 this well-known mapped fault, so far regarded as a steep fracture at least in its NW, Ścinawka segment, as a low angle Late Cretaceous – early Palaeogene thrust. In the light of their geological context, there can be little doubt about the timing of the tectonic structures identified on the seismic profiles. Our interpretation is corroborated by their occurrence near the axial zone of the Intra-Sudetic Synclinorium characterised by thick preserved Upper Cretaceous deposits. It is also confirmed by the SE-ward surficial extension of the Ścinawka-Krosnowice Fault, which at its Krosnowice segment defines a tectonic

310 boundary between the Permo-Mesozoic succession of the Intra-Sudetic Basin and Nysa-Králíky Graben on one side and the Śnieżnik – Krowiarki crystalline Massif on the other side.

An interesting Late Cretaceous – early Palaeogene structure at the NE periphery of the Intra-Sudetic Synclinorium is the NW-SE trending Czerwieńczyce Reverse Graben (Figs. 11 and 14). According to Oberc (1972, 1977), it is bounded by high-angle reverse faults (and, hence, we apply to it the term “reverse graben”) and filled with Lower Permian Rotliegend and

315 Upper Carboniferous sedimentary rocks (Oberc, 1972, 1977). On approaching the graben’s boundary faults, the strata are strongly dragged upwards, which renders a synclinal geometry to the graben’s fill. This ‘synclinal’ structural characteristics of the Czerwieńczyce Reverse Graben – if correct, being based on surficial geological mapping and involving much extrapolation at depth – is, nevertheless, quite symptomatic of a number of Sudetic grabens and to some degree it resembles cross-sections of e.g. the Wleń Graben by Kowalski (2020; Fig. 10) and – still more – by Gorczyca-Skała (1977, not reproduced here), or of the Upper Nysa – Králíky Graben by e.g. Jerzykiewicz (1970) and Don and Gotowała (2008) – partly

320 with reverse boundary faults.

4.5. The Upper Nysa - Králiky Graben

The Upper Nysa Nysa – Králiky Graben (Figs. 2 and 15) is a distinctive tectonic and topographic fault-bounded feature of approximate N-S trend, c. 45 km long and from 3 to 12 km wide, merging to the north with the Intra-Sudetic Synclinorium. It is composed of shallow marine sedimentary rocks, including a time span from the early Cenomanian, through Turonian to Coniacian and Santonian (e.g. Don and Don, 1960; Jerzykiewicz, 1970, 1971; Radwański, 1975; Don and Gotowała, 2008; Badura and Rauch, 2014), downthrown with respect to the medium-grade metamorphic rocks of the Orlica-Śnieżnik Massif of the graben's shoulders. Most of the earlier authors (except Jerzykiewicz, 1970, 1971; Radwański, 1975 and Oberc 1972) explained the origin of the Upper Nysa – Králiky Graben as a result of a Late Cretaceous rifting modified later by Cenozoic subsidence (e.g. Don and Don, 1960; Don, 1996; Wojewoda, 1997; Don and Gotowała, 2008). The early rifting was inferred to have occurred on stratigraphic and sedimentological premises – not convincing in our opinion, as they ignored the striking resemblance of the Cretaceous stratigraphic columns from within the graben and from its shoulders (Don and Gotowała, 2008; except for the columns' upper parts – see below) and the probable absence of fault-controlled coarse-grained deposits along the graben's edges. The assumed rifting was explained by most authors as due to compression- or, on the contrary, as due to extension-driven updoming of the Orlica-Śnieżnik Massif during the Cretaceous or, still, by pull-apart graben formation due to strike-slip displacements on NW-SE trending structural discontinuities in the crystalline basement. In contrast, we interpret the mostly N-S trending Upper Nysa – Králiky Graben to represent a Late Cenozoic feature, whose boundary faults cut out and downthrown a strip of the Cretaceous (Cenomanian, Turonian, Coniacian through to Santonian) shallow marine succession previously much more widespread over the area. In this way, the Cretaceous is now preserved within the graben. The uniqueness of the Cretaceous deposits in the Upper Nysa – Králiky Graben consists only in the preservation of the thick Coniacian to Santonian succession, absent from the uplifted areas nearby (Don and Gotowała, 2008) from where it must have been eroded. The Upper Nysa – Králiky Graben is imposed on and cuts across earlier, Late Cretaceous –early Palaeogene, tectonic structures of consistent NW-SE strike, which are in most places oblique to and cross-cut by the graben's boundaries (Fig. 15). These structures are represented by differentially dip-slip displaced fault blocks, often bounded by reverse faults, and by gentle folds (Badura & Rauch, 2014; Don and Gotowała, 2008). Locally, shallow-dipping thrust faults, most probably of Late Cretaceous-early Palaeogene age were mapped, drilled or directly observed in outcrops within or in direct vicinity of the Upper Nysa – Králiky Graben. A relatively extensive, at least 25 km-long, NW-SE trending trace of the Zieleniec Thrust Fault was mapped by Cymerman (1990) and extrapolated on maps by other authors within the SW flank of the Upper Nysa –Králiky Graben. On this c. 40-50° SW-dipping fault a 500 Ma orthogneiss mass from the Góry Bystrzyckie Mts was overthrust, displaced by at least 5300 m toward the NE (Cymerman, 1990) and emplaced on top of the Turonian clastics, equivalent to those from the lower part of the Upper Nysa - Králiky Graben fill (Fig. 16) and resting themselves on analogous orthogneiss basement. Cymerman (1990) interpreted this thrust as produced by Late Alpine (Miocene) compression. The Zieleniec Thrust was later confirmed by drilling, as reported by Kozdrój (2014), who interpreted the thrusting as the result of a Cenozoic gigantic, gravity-driven landslide, at the same

355 time referring to inversion of an original normal fault into a reverse one. One of the most important arguments for this interpretation was finding a subhorizontal position of the Cretaceous/gneiss contact in the drillcore. In our opinion, taking into account its significant extent and the involvement in the major NW-SE trending fault system (Fig. 11), termed the South-Sudetic Shear Zone by Wojewoda (e.g. in Wojewoda and Kowalski, 2016) which also comprises the major Pořici-Hronov Fault (see below), there cannot be much doubt that the Zieleniec thrust represents a Late Cretaceous – early
360 Palaeogene thrust, similar to that of the Łużyce Thrust (see below), though of opposite vergence.

To the NW the Zieleniec Thrust merges with the southeastern extension of the Pořici-Hronov major fault system (Fig. 11), partly representing the SW boundary to the Intra-Sudetic Synclinorium. It was active, as well, in the Late Cretaceous - early Palaeogene as a high-angle reverse fault of complex structure, but of opposite, SW, polarity (Prouza et al., 2014), which – according to some authors - is combined with the strike-slip component of motion (Wojewoda, 2007, 2009; Nováková,
365 2014) (Fig. 17a). To the SE, the Pořici-Hronov fault continues into the Duszniki-Gorzanów/Krosnowice Fault, at the northern end of the Upper Nysa – Králiky Graben.

A presumable splay fault, branching off from the Zieleniec Thrust (Figs. 11 and 17b) was described directly in a c. 1 km-long chain of outcrops along a road cut between Duszniki Zdrój and Lewin Kłodzki near the pass of Polskie Wrota on the same SW flank of the Upper Nysa - Králiky Graben (Wojewoda et al., 2010; Aleksandrowski and Wojewoda, 2010). A c.
370 20° NW-dipping detachment can be seen to occur there roughly along the bedding in middle Turonian mudstones and calcareous claystone. On a closer inspection, the detachment was found to represent a shear zone, attaining 7 to 25 cm in thickness and composed of interconnected, kinematically linked surfaces, confined between two distinct boundaries of mostly intact rock (Fig. 17b). The shear zone shows internal flaser bedding and is composed of cataclastic flow products of host-rock composition, which coexist with crush breccia and calcite veins. Based on evidence from the orientation of the
375 accompanying joints, the detachment was then interpreted as SE-vergent and probably genetically related to a supposed strike slip motion on the nearby major NW-SE-trending Duszniki-Gorzanów Fault, being a SE continuation of the Pořici-Hronov fault system (cf. Petrascheck, 1933; Prouza et al., 2014). It seems likely, however, that the joints can represent the youngest, Miocene, deformation and episode of motion on the detachment in question, whereas, its very origin and principal displacement were related to the Zieleniec Thrust.

380 **4.6. The south-western margin of the Sudetes: transition to the North Bohemian Cretaceous Basin**

The NW segment of the Sudetic Block, represented by the Karkonosze-Izera Massif, to the SW adjoins the North Bohemian Cretaceous Basin (Fig. 2). At its Sudetic margin, the basin is composed of deltaic to shallow-marine and hemipelagic Cenomanian to Santonian clastic deposits, not exceeding 600 to 700 m in thickness that rest subhorizontally on scarce
385 subcrops of Permian and Carboniferous, but mostly on the crystalline Variscan basement of the Bohemian Massif (e.g. Klein and Soukup, 1966; Malkovský, 1987; Uličný et al., 2009; Wilmsen et al., 2014). Its interior is only little affected by the Late Cretaceous –early Palaeogene tectonism, however at its boundary with the Sudetic Block the deformation is concentrated at

the spectacular and at places well exposed Lusatian Thrust (Fig. 2 and 18 a-c), which continues NW up to the vicinities of Dresden (e.g. Wagenbreth, 1967), whereas to the east it continues to the western Karkonosze Piedmont area (Prouza et al., 2013). The Lusatian Thrust has been excellently described and analysed in detail by Coubal et al. (2014).

390 The Lusatian Thrust (e.g. Malkovský, 1976), similarly to other main NW-SE trending fault zones of the Sudetic area at the Bohemian NE margin (the Intra-Sudetic Fault, the Sudetic Boundary Fault, the Odra Fault Zone) derives its origin from a major Variscan fracture, rooted deeply in the basement and likely having a primary strike-slip characteristics (cf. Aleksandrowski et al., 1997). It must have been reactivated during the Late Cretaceous – early Palaeogene compressional episode and propagated into the Permo-Mesozoic cover. The Lusatian Thrust emplaces the Variscan crystalline complexes of
395 Lusatia and of the West Sudetes on top of the Mesozoic strata of the North Bohemian Cretaceous Basin. This fault zone reveals significant along-strike changes in the dip angle of its displacement surface, between low-angle or subhorizontal in the northwest (Figs. 2 and 18a) through medium-angle (Figs. 2 and 18b) and high-angle attitude (Figs. 2 and 18c). The angle made by the displacement zone/fault core, apparently representing a reactivated fracture, with respect to the – supposedly generally subhorizontal - tectonic shortening direction had serious influence on the structural style and complexity of the
400 brittle structures developed around the fault core and on the width of the damage zone (Coubal et al., 2014). The smaller the angle between the two, the more smoothly the displacement occurred (Fig. 18a-c). An increase of the fault plane dip angle made the displacement more difficult to be achieved, which resulted in widening a zone of damage and of fault drag and favoured splitting the displacement into several slip surfaces (Fig. 18b and c).

405 **5 Regional jointing pattern and deformation bands – likely products of Late Cretaceous – early Palaeogene deformation**

The regional pattern of tectonic joints over the Sudetic area or that of SW Poland can be inferred from a number of local studies on jointing, most of which were conducted decades ago. Not attempting to make a systematic review of such studies, for the needs of a brief review made in this paper, several stereograms and rose diagrams of the dominant joint sets coming from various sources (Jerzykiewicz et al., 1974; Aleksandrowski, 1976; Solecki, 2011; Selerowicz et al., 2014) and various
410 locations in the Polish Sudetes have been shown on the geological map (Fig. 19). The data come mostly from the Permo-Mesozoic strata exposed in outcrops in the both Sudetic synclinoria, in underground galleries of the deep copper mines at the Fore-Sudetic Homocline and in the Opole Cretaceous Basin.

Two stereograms (Fig. 19 f, g) show, additionally the jointing pattern in the Carboniferous granitic rocks of the Karkonosze Massif. From recent low-temperature geochronology studies (Migoń and Danišík 2012, Sobczyk et al. 2015) it follows that
415 the present-day exposure level of at least some parts of the Karkonosze pluton was still at a depth of a few kilometres at the end of Cretaceous, and, hence the joints we observe recently in granites on the earth's surface may, indeed, be Late Cretaceous- Cenozoic, in respect of their initiation and opening (cf. e.g. Price (1966), Jaroszewski (1984), Suppe (1985), Engelder (1985, 1987, 1993) and Price and Cosgrove (1994), similarly as are the joints in the Mesozoic sedimentary rocks.

We suppose that the initiation of the dominant regional tectonic joints pattern in the Sudetic rock complexes may have occurred at a depth of a few kilometres, under a significant overburden, due to the Late Cretaceous–early Palaeogene compression. The initiation involved processes of subcritical crack growth under directional stress (e.g. Atkinson, 1982; Atkinson and Meredith, 1987), leading to the formation of anisotropy defined by systematically oriented microfractures. The latter mechanical anisotropy acquired by the rocks on compression, may have subsequently controlled the massive joint opening during the Cenozoic regional uplift and concomitant unloading and extension.

All the diagrams presented in Fig. 19 show most steep joints to be concentrated in two mutually perpendicular maxima, in which the joint planes are approximately parallel and perpendicular, respectively, to the inferred Late Cretaceous-early Palaeogene shortening direction (cf. e.g. Solecki, 2011, Nováková, 2014). As, moreover, they occur in the Permo-Mesozoic strata, a conclusion that they are genetically related to that shortening event seems plausible. We suppose that the formation of the dominant jointing pattern in the post-Variscan (but also partly in the Variscan) Sudetes and, in general, in the SW part of Poland resulted from the Late Cretaceous-early Palaeogene compressional event.

A more direct record of the Late Cretaceous-early Palaeogene compressional event is contained in the well-developed systems of silicified complementary deformation bands (Aydin and Johnson, 1983; Fossen et al., 2007), described by Solecki (1988, 1994, 2011) as cataclastic bands from the Buntsandstein and Coniacian sandstones of the North-Sudetic Synclinorium (see also Kowalski, 2021). The bisectors of the acute angle between the best developed two complementary sets of the deformation bands, dipping at moderate angles to the NE and SW, respectively, correspond to the maximum principal compressive stress axis during their formation, and, as a rule, are show subhorizontal, NE-SW oriented position.

6 Conclusion

In our brief review of the Late Cretaceous – early Palaeogene tectonic structures affecting the NE margin of the Bohemian Massif we have shown their common and widespread occurrence all over the region and, to some degree, shortly discussed their style and mechanisms of formation. These structures are easily recognizable in the Permo-Mesozoic post-Variscan cover which is devoid of other, earlier formed contractional deformations, but, though undoubtedly present, they are much less obvious in the Variscan basement that was heavily tectonized before. The structures developed in the post-Variscan cover, particularly those due to fault-displacements – and, among them, the strike-slip related ones - can still be difficult to be separated from the structures that might have resulted from the younger Cenozoic events.

Our structural analysis of newly reprocessed legacy seismic profiles, complemented with outcrop and drillhole data, has revealed the structural geometry and some interesting details of the tectonic structures produced by contractional deformation in the Permo-Mesozoic (partly also in the Carboniferous) strata of the two main Sudetic synclinoria. The deformation included gentle buckling of the basement/cover interface in the North Sudetic Synclinorium, with local complications of thus produced wide synclinal folds by relatively minor thrust, often rooted in the Variscan metamorphic basement and accompanied by thrust-related folds. Locally, rather infrequent decoupling/decollement of some packages

within the Permo-Mesozoic succession along weak, clayey horizons in e.g. Triassic Muschelkalk strata was also recorded. As the low- or medium-angle thrust faults identified in the Permo-Mesozoic strata are often rooted in the top parts of the Variscan basement, particularly in the Intra_Sudetic Synclinorium, therefore some elements of the thick-skinned style of shortening can be expected in the deep crystalline basement. It is, therefore, likely that also outside the areas covered with

455 Permo-Mesozoic sedimentary sequences, over the crystalline basement exposure areas some (even major) faults interpreted from surficial geological maps and outcrop relationships as vertical or high-angle structures, may have at depth a geometry of low-angle thrusts. This may have important practical implications for e.g. geothermal prospecting in the Sudetes, where the deep underground waters in the crystalline basement circulate, almost exclusively, along structural discontinuities, such as fault zones and fracture corridors (e.g. Dowgiałło, 2002).

460 The Late Cretaceous – early Palaeogene compressional event seems to be also responsible for the formation of the regionally dominant tectonic joints pattern in both Permo-Mesozoic and older rocks of the Sudetic area, characterized by an orthogonal joint system with two sub-vertical sets of c. NW-SE and NE-SW strikes, as is shown by a comparison of results from a few local studies.

465 **Data availability**

The seismic data from Poland are available at the National Geological Archive, Polish Geological Institute in Warsaw, and the seismic profile from Broumov area is available at the Czech Geological Survey Archive in Prague.

Author contributions

470 Both AG and PA wrote the text, prepared the figures, and compiled the paper.

Competing interests

The authors declare they have no conflict of interest.

475 **Acknowledgements.**

The consent of the Polish Oil and Gas Company (PGNiG) to publish their seismic data is gratefully acknowledged. Similarly, Janusz Badura, Aleksander Kowalski, Marta Rauch and Andrzej Solecki, together with the geological institutes of the Polish Academy of Sciences and of the University of Wrocław, as well as the Polish Geological Society and the Polish Geological Institute – National Research Institute are thanked for the permission to reproduce their illustrative materials. We

480 also express our gratitude to Jonas Kley, Andrzej Solecki and Christoph von Hagke for their constructive and helpful reviews, and, last but not least, to Piotr Krzywiec for the inspiration and encouragement to write this paper as well as for valuable editorial hints.

Financial support:

485 This work was supported by Polish Geological Institute grant no 62.9012.2046.00.0. The reprocessing of the key reflection seismic data from the North- and Intra-Sudetic synclinoria was commissioned by the Polish Geological Institute as part of its statutory duties, financed by the National Fund of Environmental Protection and Water Management in Warsaw under contract no 44/2017/Wn-07/FG-SM-DN/D and performed by GK Processing Sp. z o.o. in Cracow.

References

- 490 Aleksandrowski P.: Geological mapping survey of the Karkonosze main range between Mt. Śnieżka and Mt. Skalny Stół. Unpubl. MSc thesis, University of Wrocław (in Polish), 1976.
- Aleksandrowski P., Kryza, R., Mazur, S. and Žaba J.: Kinematic data on major Variscan fault and shear zones in the Polish Sudetes, NE Bohemian Massif, Geological Magazine, 134, 727-39, 1997.
- Aleksandrowski P. and Mazur S.: Collage tectonics in the northeasternmost part of the Variscan Belt: the Sudetes, Bohemian
- 495 Massif. In: Winchester J.A., Pharaoh T.C. and Verniers J. (eds): Palaeozoic Amalgamation of Central Europe, vol 201. Geological Society, Special Publications, London, 237–277. <https://doi.org/10.1144/GSL.SP.2002.201.01.12>, 2002.
- Aleksandrowski, P. and Wojewoda, J.: Low-angle detachment related to strike-slip faulting in Late Cretaceous mudstones of the Table Mountains (SW Poland). In: 11th Czech-Polish Workshop On Recent Geodynamics of the Sudetes and Adjacent Areas, Třešť, November 4-6th, 2010, Academy of Sciences of the Czech Republic (Institute of Rock Mechanics) and
- 500 Wrocław University of Environmental Sciences (Institute of Geodesy and Geoinformatics), Abstracts, 30-31, 2010.
- Atkinson, B.K.: Subcritical Crack Growth in Geological Materials. Journal of Geophysical Research, 89 (B6), 4077–4114, 1982.
- Atkinson, B.K., Meredith, P.G: The theory of subcritical crack growth with applications to minerals and rocks. In: Fracture Mechanics of Rocks (ed. B. K., Atkinson): 111-166. Academic Press. London, 1987.
- 505 Badura, J. and Rauch, M.: Tectonics of the Upper Nysa Kłodzka Graben. Geologia Sudetica, 42, 137-148, 2014.
- Aydin, A., Johnson, A.M.: Analysis of faulting in porous sandstones. Journal of Structural Geology, 5, 19-31, 1983.
- Fossen H., Schultz R.A., Shipton Z.K. and Mair K.: Deformation bands in sandstone: a review. Journal of the Geological Society, London, Vol. 164, 755–769, 2007.
- Bałażyńska, J. and Bossowski, A.: Wgłębna budowa geologiczna środkowej i zachodniej części synklinorium
- 510 północnosudeckiego w świetle nowych danych (Deep geological structure of central and western parts of the North-Sudetic Synclinorium; some new data), Kwartalnik Geologiczny, 23, 2, 309-321, 1979.
- Baranowski, Z., Haydukiewicz, A., Kryza, R., Lorenc, S., Muszynski, A.: Outline of the geology of the Góry Kaczawskie (Sudetes, Poland). Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, 179: 223-257, 1990.
- Beyer, K.: Die nordsudetische Rahmenfaltung, Abh. Naturforsch. Ges. Górlitz, 32, 121-172, 1939.
- 515 Bossowski, A. and Bałażyńska, J.: Ewolucja tektoniczno-strukturalna synklinorium północnosudeckiego (Tectonic-structural evolution of the North-Sudetic Synclinorium), Biuletyn Instytutu Geologicznego, 341, 163-167, 1982.

- Bossowski, A. and Ihnatowicz, A.: Atlas geologiczny Dolnośląskiego Zagłębia Węglowego (Geological atlas of the Lower Silesian coal Basin), 1 : 100 000, Państwowy Instytut Geologiczny, Warszawa, 2006.
- Bossowski, A., Ihnatowicz, A., Mastalerz, K., Kurowski, L. and Nowak, G.: Lithostratigraphy and sedimentologic-paleogeographic development, Intra-Sudetic Depression. In: Zdanowski A. and Żakowa H. (eds), The Carboniferous system in Poland, *Prace Państwowego Instytutu Geologicznego* 148, 142-147, 1995.
- Brada, Z., Fejfar, M., Ibrmajer, I., Jihlavecová, R., Petřík, A.: Study of coal-bearing formations in the Bohemian Massif - Partial report on seismic reflection survey of the Broumov promontory. Geofyzika Brno, (Unpublished report in Czech language; Archive, Czech Geological Survey), 1982.
- Chrzastek, A., Wojewoda, J.: Mezozoik południowo-zachodniej Polski - synklinorium północnosudeckie (Mesozoic of SW Poland, the North Sudetic Synclinorium). In: Żelaźniewicz, A., Wojewoda, J., Ciężkowski, W. (eds), *Mezozoik i kenozoik Dolnego Śląska*. Polskie Towarzystwo Geologiczne, Wrocław, 1-10 (in Polish, English abstract), 2011.
- Cloos, H.: Der Gebirgsbau Schlesiens und die Stellung seiner Bodenschätze. Gebrüder Borntraeger, Berlin, 1-107, 1922.
- Coubal, M., Adamovič, J., Málek, J., and Prouza, V.: Architecture of thrust faults with alongstrike variations in fault-plane dip: anatomy of the Lusatian Fault, Bohemian Massif, *Journal of Geosciences*, 59, 183–208, DOI: 10.3190/jgeosci.174, 2014.
- Coubal, M., Malek, J., Adamovič, J. and Štěpančíková, P.: Late Cretaceous and Cenozoic dynamics of the Bohemian Massif inferred from the paleostress history of the Lusatian Fault Belt, *Journal of Geodynamics*, 87. 10.1016/j.jog.2015.02.006, 2015.
- Cwojdzinski, S. and Żelaźniewicz, A.: Podłoże krystaliczne bloku przedsudeckiego (Crystalline basement of the Fore-Sudetic block). In: *Annales Societatis Geologorum Poloniae*, Special Volume: Przewodnik 66 Zjazdu Polskiego Towarzystwa Geologicznego (Guide to excursions, 66th Annual Meeting of the Polish Geological Society), Wrocław 21-24 September 1995, 11-28 (in Polish with English summary), 1995.
- Cymerman, Z.: Młodo-alpejskie nasunięcie Zieleńca w Górach Orlickich (Late-Alpine Zieleniec overthrust in the Orlickie Mts), *Przegląd Geologiczny*, 10, 422–427, 1990.
- Cymerman, Z.: Młodoalpejskie fałdy w depresji północnosudeckiej: przykłady z wapienia muszlowego z Raciborowic, *Przegląd Geologiczny*, 4, 348-354, 1998.
- Cymerman, Z.: Alpejska transpresja w Sudetach (Alpine transpression in the Sudety Mts), *Przegląd Geologiczny*, 47, 942–945, 1999.
- Cymerman, Z.: Tectonic map of the Sudetes and the Fore-Sudetic Block. 1 : 200 000, 2nd Ed. Państwowy Instytut Geologiczny, Warszawa, 2010.
- Don, J.: The Late Cretaceous Nysa Graben: implications for Mesozoic – Cenozoic fault – block tectonics of the Sudetes, *Zeitschrift für geologische Wissenschaften*, 24, 317–424, 1996.
- Don, J. and Gotowała, R.: Tectonic evolution of the late Cretaceous Nysa Kłodzka Graben, Sudetes, SW Poland, *Geologia Sudetica*, 40, 51–63, 2008.

- Don, B. and Don, J.: Geneza rowu Nysy na tle badań wykonanych w okolicach Idzikowa. [Notes on the origin of the Nysa graben], *Acta Geologica Polonica*, 10, 71–106 (in Polish), 1960.
- Dowgiałło J.: The Sudetic geothermal region of Poland, *Geothermics*, 31: 343-359, 2002.
- Dumicz, M. and Don, J.: Analiza strukturalna monokliny przedsudeckiej w rejonie Polkowic (Structural analysis of the Fore-Sudetic Homocline in vicinities of Polkowice), *Acta Universitatis Wratislaviensis* 378, *Prace Geologiczno-Mineralogiczne*, 6, 279-297, 1977.
- Dumicz, M. and Don, J., Próba odtworzenia następstwa zjawisk tektonicznych w osadach cechsztynu obszaru Polkowic na podstawie obserwacji drobnych struktur tektonicznych (An attempt at reconstruction of the sequence of tectonic phenomena in the Zechstein sediments at the Polskowice area on the basis of study on minor tectonic structures). In: Materiały konferencji komisji tektoniki Komitetu Nauk Geologicznych PAN: Problemy tektoniki Legnicko-Głogowskiego Okręgu Miedziowego, część II – wycieczki geologiczne, Lubin, Wydawnictwo CUPRUM, Wrocław, 39-45, (in Polish), 1990.
- Dziedzic, K. and Teisseyre, A. K.: The Hercynian molasse and younger deposits in the Intra-Sudetic Depression, SW Poland, *Neues Jahrb. Geol. Paläont. ABH*, 179, 285–305, 1990.
- Engelder, T.: Loading paths to joint propagation during a tectonic cycle: an example of the Appalachian Plateau, USA. *J. Struct. Geol.*, 7, 459-476, 1985.
- Engelder, T.: Joints and shear fractures in rocks. In: *Fracture Mechanics of Rocks* (ed. B. K. Atkinson), 27-65, Academic Press. London, 1987. Engelder, T.: *Stress regimes in the lithosphere*, Princeton University Press, Princeton, New Jersey, 457 p. 1993.
- Feist-Burkhardt, S., Götz, A., Szulc, J., Borkhataria, R., Geluk, M., Haas, J., Hornung, J., Jordan, P., Kempf, O., Michalík, J., Nawrocki, J., Reinhardt, L., Ricken, W., Röhling, H-G., Rüffer, T., Török, Á., Zühlke, R.: Triassic. In: McCann (ed.), *The Geology of Central Europe, Volume 2, Mesozoic and Cenozoic*. Geological Society, London, 749-821. 2008.
- Głuszyński, A., and Smajdor, Ł.: Analiza archiwalnych danych głębokiej sejsmiki poszukiwawczej na obszarze synklinoriów sudeckich (Analysis of legacy prospecting seismic data from the Sudetic synclinoria). In: Aleksandrowski P. (ed): Unpubl. report of Polish Geological Survey project *Młode strefy tektoniczne a warunki geotermalne w Sudetach w świetle badań geochronologicznych, strukturalnych i termometrycznych – etap II*, 259-269 (in Polish), National Geological Archive, Inw.#9548/2021, Warszawa, 2020.
- Gorczyca-Skała, J.: Budowa geologiczna rowu Wlenia (Geological structure of the Wleń Graben), *Geologia Sudetica*, 12, 1, 71–100, 1977.
- Grocholski, A. and Augustyniak, K.: *Atlas geologiczny Dolnośląskiego Zagłębia Węglowego* (Geological atlas of the Lower Silesian coal Basin), 1 : 50 000. Pt. I, Instytut Geologiczny, Warszawa, 1971.
- Jarosiński, M., Poprawa, P., Ziegler, P. A.: Cenozoic dynamic evolution of the Polish Platform. *Geological Quarterly*, 53 (1), 3–26, 2009. Jerzykiewicz, T.: The Upper Cretaceous turbidite sequence in the Sudetes (South-western Poland), *Bulletin de l'Academie Polonaise des Sciences, Série des scis géol. et géogr.*, 18, 3, 149-159, 1970.
- Jaroszewski, W.: *Fault and fold tectonics*. Polskie Wydawnictwo Naukowe, Warszawa, 565p., 1984

- 585 Jerzykiewicz, T.: A flysch/lithoral succession in the Sudetic Upper Cretaceous. *Acta Geologia Polonica*, 21, 165–199, 1971.
- Jerzykiewicz, T., Mierzejewski, M. and Żelaźniewicz, A.: Joint and Fracture Patterns in Basement and Sedimentary Rocks in the Sudetes Mountains, Proceedings of the First International Conference on the New Basement Tectonics (Utah Geol. Assoc. Publication no 5), 295–306, https://archives.datapages.com/data/uga/data/091/091001/295_ugs0910295.htm, 1974.
- Klein, V. and Soukup, J.: The Bohemian Cretaceous Basin. *In*: Svoboda J. et al. (Ed), Regional Geology of Czechoslovakia, Pt 1, The Bohemian Massif, Geological Survey of Czechoslovakia, Prague, 487–512, 1966.
- 590 Kley J.: Saxonische Tektonik im 21. Jahrhundert. *Z. Dt. Ges. Geowiss. (German J. Geosci.)*, 164 (2), 295–311, 2013.
- Kley, J.: Timing and spatial patterns of Cretaceous and Cenozoic inversion in the Southern Permian Basin, in: Mesozoic Resource Potential of the Southern Permian Basin, edited by: Kilhams, B., Kukla, P. A., Mazur, S., McKie, T., Munlieff, H. F., and van Ojik, K., Geological Society, London, Special Publications, 469, 19–31, <https://doi.org/10.1144/SP469.12>, 2018.
- 595 Kley, J. and Voigt, T.: Late Cretaceous intraplate thrusting in central Europe: effect of Africa-Iberia-Europe convergence, not Alpine collision, *Geology*, 36, 839–842, 2008.
- Kotański, Z. (ed.): Geological Atlas of Poland, Geological maps of horizontal section, 1 : 750 000, Państwowy Instytut Geologiczny, Warszawa, 1997.
- Kowalski, A.: Fault geometry and evidence of depocentre migration within a transtensional intra-basinal high : a case study from the Łączna Anticline (Intrasudetic Synclitorium, SW Poland) *Geological Quarterly*, 2017, 61 (4): 779–794, 2017.
- 600 Kowalski, A.: Multistage structural evolution of the end-Cretaceous–Cenozoic Wleń Graben (the Sudetes, NE Bohemian Massif) – a contribution to the post-Variscan tectonic history of SW Poland, *Annales Societatis Geologorum Poloniae*, 91, 37–66, 2021.
- Kozdrój, W.: Results of shallow scientific drillings in the Upper Nysa Kłodzka Graben and the Zieleniec area, Sudetes, *Geologia Sudetica*, Wrocław, 42, 149–159, 2014.
- 605 Kroner, U., Mansy, J.-L., Mazur, S., Aleksandrowski, P., Hann, H.P., Huckriede, F., Lacquement, F., Lamarche, J., Ledru, P., Pharaoh, T.C., Zedler, H., Zeh, A., Zulauf, G.: Variscan Tectonics. *In*: McCann, T. (ed.): *The Geology of Central Europe*. The Geological Society, London: 599–664, 2008.
- Krzywiec, P.: Mid-Polish Trough inversion – seismic examples, main mechanisms and its relationship to the Alpine-Carpathian collision, *Stephan Mueller Special Publication Series*, 1, 151–165, <https://doi.org/10.5194/smsps-1-151>, 2002.
- 610 Krzywiec, P.: Structural inversion of the Pomeranian and Kuiavian segments of the Mid-Polish Trough – lateral variations in timing and structural style, *Geological Quarterly*, Warszawa, 51, 151–168, 2006.
- Krzywiec, P. and Stachowska, A.: Late Cretaceous inversion of the NW segment of the Mid-Polish Trough – how marginal troughs were formed, and does it matter at all?, *Zeitschrift der deutschen geologischen Gesellschaft*, 167, 107–119, 2016.
- 615 Krzywiec, P., Peryt, T., Kiersnowski, H., Pomianowski, P., Czapowski, G. and Kwolek, K.: Permo-Triassic evaporites of the Polish Basin and their bearing on the tectonic evolution and hydrocarbon system, an overview, Chapter 11 in Soto, J.I., Flinch, J., Tari, G. (Eds) - Permo-Triassic salt provinces of Europe, North Africa and the Atlantic margins: tectonics and hydrocarbon potential, 243–261, Elsevier, <http://dx.doi.org/10.1016/B978-0-12-809417-4.00012-4>, 2017.

- Leśniak, T.: Tektonika obszaru między Raciborowicami a Łaziskami w północno-wschodniej części depresji
620 północnosudeckiej, Zeszyty Naukowe AGH, Geologia, 5, 4, Kraków, 87-107, (in Polish), 1979.
- Malkovský, M.: Saxonische Tektonik der Böhmischen Masse, Geologische Rundschau, 65, 10.1007/BF01808459,
1976. Malkovský, M.: Important faults of the platform cover of the northern part of the Bohemian Massif. Výzk Práce Ústř
Úst geol, 14, 1–32, 1977.
- Malkovský, M.: The Mesozoic and Tertiary basins of the Bohemian Massif and their evolution, Tectonophysics, 137, 31–42,
625 10.1016/0040-1951(87)90311-8, 1987.
- Malz, A., Nachtweide, Ch., Emmerlich, S., and Schimpf, L.: Mesozoic intraplate deformation in the southern part of the
Central European Basin - Results from large-scale 3D modelling, Tectonophysics, 776. 10.1016/j.tecto.2019.228315, 2020.
- Markiewicz, A.: Tektonika obszaru złoża (The tectonics of the ore-deposit area), Chapter 2.11 in: Piestrzyński A., (ed.),
Monografia KGHM Polska Miedź S.A., KGHM CUPRUM, Wrocław, 115-132, 2007.
- 630 Markiewicz, A., Szarowski, W.: Zjawiska tektoniczne w południowej części kopalni Lubin, In: Materiały konferencji
komisji tektoniki Komitetu Nauk Geologicznych PAN: Problemy tektoniki Legnicko-Głogowskiego Okręgu Miedziowego,
część II – wycieczki geologiczne, Lubin, Wydawnictwo CUPRUM, Wrocław, 4-8, 1990.
- Martínez Catalán, J.R., Hatcher Jr, R.D., Arenas, R., Díaz García, F (eds): *Variscan-Appalachian dynamics: the building
of the Late Paleozoic Basement*. Geol. Soc. Am. Spec. Pap., Denver, vol. 364, 305 p., 2002.
- 635 Martínez Catalán J.R., Collett S., Schulmann K., Aleksandrowski P. and Mazur S.: Correlation of allochthonous terranes and
major tectonostratigraphic domains between NW Iberia and the Bohemian Massif, European Variscan belt, International
Journal of Earth Sciences (Geol Rundsch), 109, 4, 1105-1131, <https://doi.org/10.1007/s00531-019-01800-z>, 2020.
- Mazur, S., Aleksandrowski, P., Kryza, R. and Oberc-Dziedzic, T.: The Variscan orogen in Poland, *Geological Quarterly*, 50,
89-118, 2006.
- 640 Matte, P.: The Variscan collage and orogeny (480–290 Ma) and the tectonic definition of the Armorica microplate: a review.
Terra Nova 13:122–128, 2001.
- Mazur, S., Aleksandrowski, P., Turniak, K. and Awdankiewicz, M.: Geology, tectonic evolution and Late Palaeozoic
magmatism of Sudetes – an overview, In: A. Kozłowski and J. Wiszniewska (eds), Granitoids in Poland, AM (Archivum
Mineralogiae) Monograph No. 1, Komitet Nauk Mineralogicznych PAN & Wydział Geologii Uniwersytetu Warszawskiego,
645 59-87, 2007.
- Mazur, S., Aleksandrowski, P., Turniak, K., Krzemiński, L., Mastalerz, K., Górecka-Nowak, A., Kurowski, L., Krzywiec, P.,
Żelaźniewicz, A. and Fanning, M.C.: Uplift and late orogenic deformation of the Central European Variscan belt as revealed
by sediment provenance and structural record in the Carboniferous foreland basin of western Poland, *International Journal
of Earth Sciences*, 99, 47-64, 2010.
- 650 Mazur, S., Aleksandrowski, P., Gągała, Ł., Krzywiec P., Żaba J., Gaidzik K. and Sikora R.: Late Palaeozoic strike-slip
tectonics versus oroclinal bending at the SW outskirts of Baltica: case of the Variscan belt's eastern end in Poland,
International Journal of Earth Sciences, 109, 1133-1160, 2020.

- McCann, T., Kiersnowski, H., Krainer, K., Vozárová, A., Peryt, T.M., Oplustil, S., Stollhofen, H., Schneider, J., Wetzel, A., Boulvain, F., Dugar, M., Török, Á., Haas, J., Tait, J. and Körner, F.: Permian. *In* McCann T. (ed.): The Geology of Central Europe, The Geological Society, London, 531-597, 2008a.
- 655 Mazur S., Scheck-Wenderoth M. and Krzywiec P.: Different modes of the Late Cretaceous-Early Tertiary inversion in the North German and Polish basins. *International Journal of Earth Sciences (Geologische Rundschau)*, 94, 782-798, 2005.
- McCann, T., Skompski, S., Poty, E., Dugar, M., Vozárová, A., Schneider, J., Wetzel, A., Krainer, K., Kornpihl, K., Schäfer, A., Krings, M., Oplustil, S. and Tait, J.: Carboniferous. *In* McCann T. (ed.): The Geology of Central Europe, The Geological
- 660 Society, London, 411-529, 2008b.
- McClay, K.R.: Glossary of thrust tectonics terms in: McClay, K.R. (Ed), Thrust tectonics, Chapman & Hall, London, 419-433, 1992.
- Migoń, P. and Danišík, M.: Erosional history of the Karkonosze Granite Massif – constraints from adjacent sedimentary basins and thermochronology. *Geological Quarterly*, 56, 3, 440–454, doi: 10.7306/gq.1032, 2012.
- 665 Milewicz, J.: The geological structure of the North-Sudetic Depression, *Biuletyn Instytutu Geologicznego*, 227, 5–27, 1968.
- Milewicz, J.: Górna kreda depresji północnosudeckiej, *Acta Universitatis Wratislaviensis, Prace Geologiczno-Mineralogiczne*, 61, 1-58, 1997.
- Navabpour, P., Malz, A., Kley, J., Sieburg, M., Kasch, N. and Ustaszewski, K.: Intraplate brittle deformation and states of paleostress constrained by fault kinematics in the central German platform, *Tectonophysics*, 694, 146–163,
- 670 10.1016/j.tecto.2016.11.033, 2017.
- Nádaskay, R., Žák, J., Sláma, J., Sidorinová, T., and Valečka, J.: Deciphering the Late Paleozoic to Mesozoic tectonosedimentary evolution of the northern Bohemian Massif from detrital zircon geochronology and heavy mineral provenance, *International Journal of Earth Sciences*, 108, 2653–2681, 10.1007/s00531-019-01781-z, 2019.
- Nemec, W., Porębski, S. J. and Teisseyre, A. K.: Explanatory notes to the lithotectonic molasse profile of the Intra-Sudetic
- 675 Basin, Polish part (Sudety Mts, Carboniferous-Permian), *Veröffentlichungen Zentralinstituts für Physik der Erde, Akad. der Wissenschaften der DDR*, 66, 267–278, 1982.
- Nováková, L.: Evolution of paleostress fields and brittle deformation in Hronov-Poříčí Fault Zone, Bohemian Massif, *Stud. Geophys. Geodet.*, 58, 269–288, 2014.
- Oberc, J.: Sudety i obszary przyległe, [In:] *Budowa geologiczna Polski. T. 4, Tektonika, cz. 2.* Wydawnictwa Geologiczne, Warszawa, 1972.
- 680 Oberc, J.: The Early Alpine epoch in South-West Poland: The North Sudetic Synclinorium, In: Pożaryski W. (Ed), *Geology of Poland, Vol.4. Tectonics*, Wydawnictwa Geologiczne, Warszawa, 419-424, 1977.
- Oberc, J. and Salski, W.: Fałdy i spękania w skałach dolnozecheńskich na obszarze szybu wschodniego kopalni Lubin (Fold and joints in lower Zechstein rocks around the eastern shaft of the Lubin mine), *Kwartalnik Geologiczny*, 3, 519-536,
- 685 1968.

- Petrascheck, W.: Der böhmische Anteil der Mittelsudeten und sein Vorland. Mitteilungen der Geol. Gesellschaft in Wien, 24, 1933.
- Požaryski, W.: Mapa geologiczna Polski i krajów ościennych bez utworów kenozoicznych 1 : 1 000 000 (Geological map of Poland and adjacent countries without Cenozoic deposits), Instytut Geologiczny, Warszawa (in Polish), 1979.
- 690 Prouza, V., Coubal, M. and Adamovič, J.: Specifika architektury hronovsko-poříčského zlomu (Specific architecture fo the Hronov –Poříčí Fault), Zprávy o geologických výzkumech v roce 2014/A – Regionální geologie a stratigrafie, Česká geologická služba, Praha, 13-18, 2014.
- Price, N.J.: Fault and joint development in brittle and semibrittle rock. Pergamon Press, 1—176, 1966.
- Price, N.J., Cosgrove, J.W.: Analysis of Geological Structures. Cambridge University Press, Cambridge, 502 p.,
- 695 1990.Radwański, S.: Kreda Sudetów Środkowych w świetle wyników nowych otworów wiertniczych [Upper Cretaceous of the central part of the Sudetes in the light of new borehole materials], Biuletyn Instytutu Geologicznego, 187, 5–59, 1975.
- Salski, W.: Problemy małej tektoniki w rejonie Lubina (Problems of minor tectonics in vicinities of Lubin). Rudy i Metale Nieżelazne, 4, 485-489 (in Polish), 1965.
- Salski, W.: Charakterystyka litologiczna i drobne struktury łupków miedzionośnych monokliny przedsudeckiej (Lithological
- 700 characteristics and minor structures of copper-bearing shales in Fore-Sudetic Homocline), Kwartalnik Geologiczny, 12/4, 855-873 (in Polish), 1968.
- Scheck, M., Bayer, U., Otto, V., Lamarche, J., Banka, D., Pharaoh, T.: The Elbe Fault System in North Central Europe – a basement controlled zone of crustal weakness, Tectonophysics 360, 281–299, 2002.
- Schroder, B.: Inversion tectonics along the Western margin of the Bohemian Massif, Tectonophysics, 137, 93-100,
- 705 10.1016/0040-1951(87)90316-7, 1987.
- Selerowicz, T., Głuszyński, A., Niedbał, M.: Zechstein limestone (Ca1) joint orientation analysis in mining shafts of “Polkowice-Sieroszowice” copper and silver mine (SE Poland): field study results, Geologia Sudetica, 42, 84-84, 2014.
- Sobczyk, A., Danišík, M., Aleksandrowski, P. and Anczkiewicz, A.: Post-Variscan cooling history of the central Western Sudetes (NE Bohemian Massif, Poland) constrained by apatite fission-track and zircon (U-Th)/He thermochronology,
- 710 Tectonophysics 649, 47–57, <http://dx.doi.org/10.1016/j.tecto.2015.02.021>, 2015.
- Sobczyk, A., Sobel, E. R. and Georgieva, V.: Meso–Cenozoic cooling and exhumation history of the Orlica–Śnieżnik Dome (Sudetes, NE Bohemian Massif, Central Europe): Insights from apatite fission–track thermochronometry, Terra Nova, 32, 122–133, 2020.
- Śliwiński, W., Raczyński, P., Wojewoda, J.: Sedymentacja utworów epiwaryscyjskiej pokrywy osadowej w basenie północnosudeckim (Sedimentation of the epi-Variscan cover in the North Sudetic Basin). In: Ciężkowski, W., Wojewoda, J., Żelaźniewicz, A. (eds), *Sudety Zachodnie: od wendy do czwartorzędu*. Polskie Towarzystwo Geologiczne, Wrocław, 119–126 (in Polish, English abstract), 2003.
- Solecki, A.T.: Tektonika dysjunktywna i jej wpływ na warunki występowania kopalin w synklinorium północnosudeckim, PhD thesis, University of Wrocław, 1-152 (in Polish), 1986.

- 720 Solecki, A.: Conjugate cataclastic zones in the sandstones of the North-Sudetic Synclinorium, *Przegląd Geologiczny*, 36, 577–581 (in Polish, English summary), 1988.
- Solecki, A.: Tectonics of the North Sudetic Synclinorium, *Acta Universitatis Wratislaviensis No 1618, Prace Geologiczno-Mineralogiczne*, 45, 1-60, University of Wrocław, 1994.
- Solecki, A.: Joints and shears of the N-Sudetic Synclinorium. In: Rossmanith, H-P. (ed), *Mechanics of jointed and faulted*
725 *rocks*. Balkema, 341-346, 1995.
- Solecki, A.: Rozwój strukturalny epiwaryscyjskiej pokrywy platformowej w obszarze synklinorium północnosudeckiego (Structural development of the epi-variscan cover in the North Sudetic Synclinorium area). In: Żelaźniewicz, A. et al. (eds), *Mezozoik i kenozoik Dolnego Śląska*, Polskie Towarzystwo Geologiczne, Wrocław, 19-36, 2011.
- Suppe, J.: *Principles of Structural Geology*. Prentice-Hall, 537p.m 1985.
- 730 Teisseyre, H.: Ważniejsze dyslokacje ramowe Sudetów (Major boundary faults of the Sudetes), In: Teisseyre H. (Ed.), *Regionalna geologia Polski, T. III: Sudety, Z.1: Utwory przedtrzeciorzędowe*, Polskie Towarzystwo Geologiczne, Kraków, 25-29 (in Polish), 1957.
- Uličný, D., Špičáková, L., Grygar, R., Svobodová, M., Čech, S. and Laurin, J.: Palaeodrainage systems at the basal unconformity of the Bohemian Cretaceous Basin: roles of inherited fault systems and basement lithology during the onset of
735 basin filling, *Bulletin of Geosciences*, 84, 4, 577–610, 2009.
- Voigt, S., Wagreeich, M., Surlyk, F., Walaszczyk, I., Uličny, D., Čech, S., Voigt, T., Wiese, F., Wilmsen, M., Niebuhr, B., Reich, M., Funk, H., Michalík, J., Jagt, J.W.M., Felder, P.J. and Schulp, A.S.: Cretaceous, *In* McCann T. (ed.): *The Geology of Central Europe*, The Geological Society, London, 923-997, 2008.
- Voigt, T., Kley, J. and Voigt, S.: Dawn and dusk of Late Cretaceous basin inversion in central Europe, *Solid Earth*, 12,
740 1443–1471, <https://doi.org/10.5194/se-12-1443-2021>, 2021.
- Wilmsen, M., Uličný, D. and Košťák, M.: Cretaceous basins of Central Europe: deciphering effects of global and regional processes – a short introduction, *Zeitschrift der Deutschen, Gesellschaft für. Geowissenschaften*, 165, 4, 495–499, 2014.
- Wojewoda, J.: Sukcesja litoralno-szelfowa górnej kredy na obszarze niecki śródsudeckiej i rowu górnej Nysy w Sudetach, In: Wojewoda, J. (Ed.) *Obszary źródłowe: Zapis w osadach*, I, 81–96, WIND, Wrocław (in Polish), 1997.
- 745 Wojewoda, J.: Neotectonic aspect of the Intrasudetic Shear Zone. *Acta Geodynamica et Geomaterialia*, Vol. 4, No. 4 (148), 31-41, 2007.
- Wojewoda, J.: Žďárky-Pstrážna Dome: a strike-slip fault-related structure at the eastern termination of the Poříčí-Hronov Fault Zone (Sudetes), *Acta Geodynamica et Geomaterialia*, 6, 273–290, 2009.
- Wojewoda, J., Koszela, S. and Aleksandrowski, P.: A kilometre-scale low-angle detachment related to strike-slip faulting in
750 Late Cretaceous mudstones of the Table Mountains (Central Sudetes, SW Poland). In: *Central European Tectonic Studies Group (CETEG) 8th Meeting*, 22-25 April 2010, Mąchocice Kapitulne, *Conference Proceedings* (ed. by M. Ludwiniak, A. Konon & A. Żylińska), University of Warsaw and Polish Geological Institute - National Research Institute, 127-128, 2010.

- Wojewoda, J. and Kowalski, A.: Rola południowosudeckiej strefy ścinania w ewolucji Sudetów (The role of the South-Sudetic Shear Zone in the evolution of the Sudetes)., In: Wojewoda, J. and Kowalski, A. (eds): Wyzwania Polskiej Geologii, 3. Polski Kongres Geologiczny, Przewodnik do wycieczek kongresowych, Wycieczka 2.2., Polskie Towarzystwo Geologiczne, Wrocław, 28-43, 2016.
- Ziegler, P.A.: Late Cretaceous and Cenozoic intra-plate compressional deformations in the Alpine foreland-a geodynamic model, *Tectonophysics*, 137, 389-420, 10.1016/0040-1951(87)90330-1, 1987.
- Ziegler, P. A., Cloetingh, S. and van Wees, J. D.: Dynamics of intra-plate compressional deformation: the Alpine foreland and other examples, *Tectonophysics*, 252, 7–59, 1995.
- Żelaźniewicz, A. and Aleksandrowski, P.: Regionalizacja tektoniczna Polski: Polska południowo-zachodnia (Tectonic subdivision of Poland: southwestern Poland), *Przegląd Geologiczny*, 56, 904-911 (in Polish, English abstract), 2008.
- Żelaźniewicz, A., Aleksandrowski, P., Buła, Z., Kamkowski, P.H., Konon, A., Oszcypko, N., Ślaczka, A., Żaba, J. and Żytko, K.: Regionalizacja geologiczna Polski, Komitet Nauk Geologicznych PAN, Wrocław, 60, 2011.
- Żelaźniewicz, A. and Markiewicz, A.: Struktury ekstensyjne w cechsztyńskich ewaporatach monokliny przedsudeckiej a strefa tektoniczna Odry, *Przegląd Geologiczny*, 10, 463-471, 1991.




Figure 1: The main Late Cretaceous – early Palaeogene tectonic structures in Central Europe. Modified after Kley and Voigt 2008.. Abbreviations: F – Flechtingen High, FL – Franconian Line, FSB - Fore-Sudetic Block, FSH – Fore-Sudetic Homocline, Ha-Harz Block, HCM - Holy Cross Mountains, La – Lusatian Thrust, LB - Lusatian Block, Lf - Lusatian Main Fault, NBCB - North Bohemian Cretaceous Basin, Od - Odra Fault, TW - Thuringian Forest, SB - Sudetic Block, SMF - _Sudetic Boundary Fault.




Figure 2: Geological map without Cenozoic of NE margin of the Bohemian Massif and its surroundings in SW Poland and N Czechia (simplified after Dadlez et al., 2000). NKG – Nysa-Králiky Graben. Red boxes show location of Figs. 6, 10a, 11 and 14.




Figure 3: Schematic regional cross sections across the Bohemian Massif NE margin (highly vertically exaggerated). Location in Fig. 2.




Figure 4: Two seismic profiles from the Fore-Sudetic Homocline (location in Fig. 2) showing very gentle folds in Triassic and Permian (Rotliegend not marked). Note steep to low-angle mainly reverse faults. The anticlines are structural traps for hydrocarbons, with gas fields known in Rotliegend strata (Grochowice gas field in Fig. 4a). Please note the different scales of the two profiles.




Figure 5: Examples of tectonic structures observed on galleries' walls in copper and silver mines on the Fore-Sudetic Homocline at the base Zechstein (Upper Permian) ore-bearing formation at its contact with the Rotliegend (Lower Permian) strata (location in Fig. 2). (a) Inverted steep fault, originally normal. The inversion caused fault-drag of Ca1 Zechstein Limestone. (b) Thrust fault in carbonate rock (orientation recorded as dip direction/dip angle). Abbreviations: Bs- Weissliegend Sandstone (Lower Permian), T1-Kupferschiefer Shale (Upper Permian), Ca1-Zechstein Limestone (Upper Permian).




Figure 6: Geological map (without Cenozoic) of the North Sudetic Synclinorium (based on Bałazińska and Bossowski, 1979; Solecki, 1994; Krentz et al., 2001 and Cymerman, 2010), showing locations of the interpreted seismic profiles. For full-length location of seismic profile of Fig. 9a - see Fig. 2.




Figure 7: Kurt Beyer's (1939) interpretation of the "young Saxonian" tectonics in the North Sudetic Synclinorium (Basin). (a) Main tectonic deformation stages in the North Sudetic Basin. (b) Block-diagram showing simple open structure of the western part of the North Sudetic Synclinorium (Bolesławiec Trough) and complex pattern of blocks and synclinal grabens (Jerzmanice Syncline and Wleń Trough) in its eastern part (cf. Figs. 2 and 6).



Figure 8: Folds in Muschelkalk in a quarry at Raciborowice (location in Fig. 6). (a) photo courtesy of Andrzej Solecki.




Figure 9: Interpreted seismic profiles from the North-Sudetic Synclinorium (location in Fig. 6).









Figure 10: Simplified geological map and three selected cross-sections of the Wleń Graben according to Kowalski (2020)



830 **Figure 11: Geological map of the Intra-Sudetic Synclinorium (based on Grocholski and Augustyniak, 1971, Bossowski and Ihnatowicz, 2006, and Cymerman, 2010). Location of the interpreted seismic profiles and described localities is shown. CZT – Czerwieńczyce Trough (reverse graben).**



835 **Figure 12:** Seismic profiles from vicinity of Mioszów arranged successively along-strike of the structures, showing probable effects of Late Cretaceous – early Palaeogene reactivation of originally Carboniferous faults. The Permian volcanic member (in red) continues to the NNE to outcrop massively in the Suche Mountains. Location in Fig. 11. The faults in (a) do not show effects of post-Carboniferous reactivation, but further to SE the motion on faults resulted in very gentle folding of the overlying Permian strata.




Figure 13: Seismic profiles (paper print version – Brada et al., 1982) from vicinity of Broumov (a) and from vicinities of Radków (b-e). Location in Fig. 11. The section (a) shows that the Broumov well was drilled in the hanging-wall of a thrust fault and, hence, the Carboniferous in the neighbourhood may occur deeper than found in the well. The thrust continues to the NE parallel to the bedding – most probably along the Lower Permian Anthracosia Shale member.




Figure14: Geological cross- section of the Czerwieńczyce Reverse Graben after Oberc (1972). Location in Figs 2 and 11.




Figure 15: Perspective view towards the north onto the base Cretaceous relief (known from boreholes) in the Upper Nysa-Králiky Graben (from Badura and Rauch, 2014). The oblique-striking troughs, steps and horsts reflect NW-SE trending Late-Cretaceous – early Palaeogene structural grain defined by inverted blocks underlying the Cretaceous graben floor and folds in its fill (removed from the picture), once continuing laterally, before the graben formed in Miocene times. Location given in Fig 2.




Figure16: Schematic section across the Zieloniec Thrust (based on data from Cymerman, 1990, and Kozdrój, 2014). Location in Figs 2 and 11.




Figure 17: Examples of outcrop-scale brittle structures related to major faults at the SW boundary of the Intra-Sudetic Synclinorium.

(a) The Pořiči-Hronov Fault, interpreted as high-angle reverse structure, near the village of Malé Svatoňovice: tectonic contact between nearly vertically dipping Upper Cretaceous sandstones in the SW and the Permian sandstone in the NE.

(b) Shallow NW-dipping detachment along the bedding of Middle Turonian mudstones and calcareous claystones in a road cut near the village of Lewin Klodzki, probably related to the NW continuation of the Zieloniec Thrust. Location in Figs 2 and 11.




Figure 18: Schematic sections across the Lusatian Thrust at three localities, showing along-strike stucture changes (based on Coubal et al., 2014). Location in Fig 2.




Figure 19: Comparison of stereograms and rose diagrams of the dominant steep joint sets coming from several local studies (by Jerzykiewicz et al., 1974, Aleksandrowski, 1976, Solecki, 2011, Selerowicz et al., 2014) made at various locations in the Polish Sudetes. Background: the geological map of Fig. 2.