

1 **Plume-ridge interactions: Ridge-ward versus plate-drag plume flow**

2 **Fengping Pang¹, Jie Liao^{1,2,3}, Maxim D. Ballmer⁴, Lun Li^{1,2,3}**

3 ¹School of Earth Sciences and Engineering, Sun Yat-Sen University, Guangzhou 510275, China

4 ²Guangdong Provincial Key Lab of Geodynamics and Geohazards, Guangzhou 510275, China

5 ³Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519000, China

6 ⁴Department of Earth Sciences, University College London, London, United Kingdom

7
8 Correspondence: Jie Liao (liaojie5@mail.sysu.edu.cn)

9
10 **Abstract**

11 The analysis of mid-ocean ridges and hotspots that are sourced by deep-rooted mantle plumes allows
12 us to get a glimpse of mantle structure and dynamics. Dynamical interaction between ridge and
13 plume processes have been widely proposed and studied, particularly in terms of ridge-ward plume
14 flow. However, the effects of plate drag on plume-lithosphere and plume-ridge interaction remain
15 poorly understood. In particular, the mechanisms that control plume flow towards vs. away from the
16 ridge have not yet been systematically studied. Here, we use 2D thermomechanical numerical models
17 of plume-ridge interaction to systematically explore the effects of (i) ridge spreading rate, (ii) initial
18 plume head radius, and (iii) plume-ridge distance. Our numerical experiments suggest two different
19 geodynamic regimes: (1) plume flow towards the ridge is favored by strong buoyant mantle plumes,
20 slow spreading rates and small plume-ridge distances; (2) plume drag away from the ridge is in turn

删除[pangfengping]:
删除[pangfengping]:

21 promoted by fast ridge spreading, for small-to-intermediate plumes and large plume-ridge distance.

22 We find that the pressure gradient between the buoyant plume and spreading ridge at first drives

23 ridge-ward flow, but eventually the competition between plate drag and the gravitational force of

24 plume flow along the base of the sloping lithosphere controls the fate of plume (spreading towards vs.

25 away from the ridge). Our results highlight that fast-spreading ridges exert strong plate dragging

26 force, which sheds new light on natural observations of largely absent plume-lithosphere interaction

27 along fast-spreading ridges, such as the East Pacific Rise.

删除[pangfengping]: at least

删除[pangfengping]:

删除[pangfengping]:

1 Introduction

Mid-ocean ridges (MORs) and hotspots are two main regions for deep material recycling to the surface of the Earth. However, these two units are not always isolated, but rather show strong interactions in some cases, termed as plume-ridge interaction (Morgan, 1978). Of up to 50 mantle plumes revealed by seismic tomography (French and Romanowicz, 2015; Montelli et al., 2004), more than 20 plumes are found to be associated with nearby ridges (Fig.1a; Ito et al., 2003). Plume-ridge interaction is manifested by geophysical and geochemical anomalies along the ridge axis, e.g., high mantle potential temperature (Dalton et al., 2014), enriched radiogenic isotope anomalies (Cushman et al., 2004; Douglass and Schilling, 1999; Yang et al. 2017), and adjacent lineations of seamounts (Ballmer et al., 2013b; Geissler et al., 2020; Lénat and Merle, 2009). Furthermore, plumes may promote migration of MOR spreading centers (Müller et al., 1998; Mittelstaedt et al., 2008, 2011; Whittaker et al., 2015), as evidenced by successive ridge jumps towards mantle plumes, e.g., at Iceland, Amsterdam-Saint Paul and Galapagos hotspots (Hardarson et al., 1997; Maia et al., 2011; Mittelstaedt et al., 2012). The interaction dynamics of a ridge with a ridge-centered and off-ridge plume has been widely studied and modeled in analogue and numerical experiments, revealing that the major controlling factors involve the ridge spreading rate, plume buoyancy flux and their spatial distance (François et al., 2018; Ito et al., 1997; Kincaid et al., 1996; Ribe et al., 1995; Ribe, 1996; Sleep, 1997). Indeed, most plume-ridge interaction systems are associated with slow-spreading ridges and small mantle plumes and short plume-ridge distances (Fig.1b). However, numerical studies systematically investigating the effects of these parameters on plume-ridge interaction and quantify the controlling forces remain scarce.

删除[pangfengping]:

删除[pangfengping]:

51 As has been noted previously, buoyant plumes tend to spread ridge-ward along the sloping base
52 of the lithosphere (Morgan, 1978; Schilling, 1991; Small, 1995). Regions of divergent mantle flow
53 beneath MORs represent the lowest dynamic-pressure regions in the oceanic asthenosphere, and thus
54 tend to suck ambient asthenospheric and plume materials towards the spreading center (Niu, 2014).
55 On the other hand, the viscous drag at the base of the plate tends to convey the spreading plume
56 material away from the MOR (Ribe and Christensen, 1994, 1999). Indeed, plume spreading at the
57 base of the lithosphere is governed by the competition of trench-ward viscous plate drag vs.
58 ridge-ward gravitational and pressure-driven forces (Kincaid et al., 1996). These gravitational and
59 tectonic forces compete with other to control the regime of plume-ridge interaction, but their balance
60 remains to be quantified.

61 The different distribution of hotspots with classified as plume-ridge interaction (ridge-ward
62 spreading) vs. no interaction (plate-drag spreading) also still remains enigmatic. Plume-ridge
63 interaction is much more common near the Mid-Atlantic ridge (MAR) than near the East Pacific Rise
64 (EPR) (Fig. 1a). Near the EPR, only the Pukapuka and Sojourn ridges display clear evidence of
65 ridge-ward flow of the magmatic source, but these volcanic ridges have been attributed to a
66 horizontally viscous differences or small-scale convection in uppermost mantle, and not a mantle
67 plume (Ballmer et al., 2013b; Clouard and Bonneville, 2005; Harmon et al., 2011). A previous study
68 (Jellinek et al., 2003) proposed that fast-spreading ridges guide upwelling mantle flow towards the
69 spreading center to convey the surrounding plumes from deep depth entirely into the MOR melting
70 zone (Fig. 1c), resulting in the absence of hotspots adjacent to the EPR (see also Rowley et al., 2016;
71 Rowley and Forte, 2022). However, fast plate spreading also tends to drag mantle plumes away from
72 the MOR (Kincaid et al., 1995, 1996), leading to the typically parabolic shapes of hotspot swells

73 such as near Hawaii (Ribe and Christensen, 1994). Whether the increased spreading rates in the
74 Pacific vs. Atlantic promote ridge-ward vs. plate-drag plume flow remains an intriguing question.

75 The principal goal of this study is to investigate the process of plume-ridge interaction, with an
76 emphasis on the effects of model parameters on the ridge-ward vs. plate-drag plume spreading. We
77 explore the effects of various model parameters, such as the size of the plume, ridge spreading rate,
78 and plume-ridge distance. Finally, we use our model results to interpret the difference of natural
79 plume-ridge interaction systems in different oceans, particularly the striking difference between the
80 East Pacific and Atlantic in this regard.

81




Figure 1. Global plume-ridge interaction systems. **(a)** Global distribution of mid-ocean ridges and mantle plumes. Residual bathymetry of the ocean basins come from Straume et al. (2019). Mid-ocean ridges are painted in color solid lines corresponding to half-spreading rate. Plumes not interacting with a ridge are shown by green circles, and hotspots linked to ridges are in red dots (Ito et al., 2003); size refers to the plume buoyancy flux from Hoggard (2020). Black lines denote the regions of two LLSVPs under the South Africa and Pacific Ocean (Torsvik et al., 2006). **(b)** Histograms of influential factors of plume-ridge interaction systems. Half spreading rate and plume-ridge distance is taken from GPlates (Müller et al., 2016; Whittaker et al., 2015). Plume-ridge interaction systems link to slow-spreading ridge and small mantle plumes and short plume-ridge distance. **(c)** Sketches of ridge-ward (top panel) and plate-drag plume flow (bottom panel) mode proposed, respectively.




Figure 2. Model setup. **(a)** Initial composition and boundary conditions. The oceanic plate consists of half-space cooling part and the thermal equilibrium part. A 50-Myrs-old mid-ocean ridge sets in the middle of the model based on half-space cooling temperature structure. A thermal and chemical anormal mantle plume locates at 660 km. Different colors indicate the initial rock types and corresponding newly formed molten rock types. Yellow arrows are the half-spreading rates imposed internal in the lithosphere (i.e., from 20 km to 120 km in depth) to simulate ridge spreading. **(b)** Initial tested ridge and plume configurations. **(c)** Initial tested plume-ridge distances.

2 Numerical modelling

2.1 Modelling methods

To explore plume-lithosphere and plume-ridge interaction, we conduct numerical simulations

删除[pangfengping]:

删除[pangfengping]:

utilizing the 2D thermo-mechanical code I2VIS, which is based on staggered finite difference method combined with marker-in-cell techniques (Gerya and Yuen, 2003, 2007). The equations of conservation of mass, momentum and energy (Eq. (1)-(3), respectively) are solved in a fully staggered grid assuming an incompressible media:

删除[pangfengping]: This modeling framework uses both Eulerian grids and randomly-distributed Lagrangian markers to jointly solve

删除[pangfengping]:):

$$\nabla \cdot \vec{v} = 0 \quad (1)$$

$$\frac{\partial \sigma'_{ij}}{\partial x_j} - \frac{\partial P}{\partial x_i} + \rho g_i = 0 \quad (2)$$

$$\rho C_p \left(\frac{DT}{Dt} \right) = -\nabla \cdot \vec{q} + H_r + H_a + H_s + H_l \quad (3)$$

where v refers to the velocity, σ'_{ij} the deviatoric stress tensor, P the pressure, ρ the density, g the gravity acceleration, $\frac{D}{Dt}$ the Lagrangian time derivative, C_p the heat capacity, and q the heat flux. Additionally, H_r , H_a , H_s , and H_l are the radioactive, adiabatic, shear, and latent heat productions, respectively.

We employ a non-Newtonian visco-plastic rheology (Gerya and Yuen, 2007) in the models. The viscous rheology depends on stress, temperature and pressure. The appropriate viscosity is expressed as that of a composite diffusion and dislocation-creep material (Eq. (4)).

$$\frac{1}{\eta_{vis}} = \frac{1}{\eta_{diff}} + \frac{1}{\eta_{disl}} \quad (4)$$

in which η_{diff} and η_{disl} are the diffusion and dislocation creep viscosity, respectively, and can be further computed as Eq. (5) and Eq. (6):

$$\eta_{diff} = \frac{1}{2} A \sigma_{crit}^{1-n} \exp \left(\frac{PV_a + E_a}{RT} \right) \quad (5)$$

$$\eta_{disl} = \frac{1}{2} A \frac{1}{n} \dot{\epsilon}_{II}^{\frac{1-n}{n}} \exp \left(\frac{PV_a + E_a}{nRT} \right) \quad (6)$$

where P is the pressure, T is the temperature, $\dot{\epsilon}_{II}$ is the second invariant of the strain rate tensor, σ_{crit} is the diffusion-dislocation creep transition stress, and A , E_a , V_a , and n are the strain rate pre-exponential factor, activation energy, activation volume, and stress exponent, respectively. The

删除[pangfengping]:

删除[pangfengping]:

128 plastic behavior η_{pla} is described by the Drucker-Prager yield criterion (Byerlee, 1978; Ranalli,
129 1995) according to Eq. (7) and Eq. (8):

130
$$\sigma_y = C + P\varphi \quad (7)$$

131
$$\eta_{pla} = \frac{\sigma_y}{2\dot{\epsilon}_{II}} \quad (8)$$

132 in which σ_y is the yield stress, C is the rock cohesion and φ is the effective friction coefficient.
133 The effective viscosity η_{eff} of rocks is thus constrained by both viscous and plastic deformation,
134 where the rheological behavior depends on the minimum viscosity attained between ductile and
135 brittle fields:

136
$$\eta_{eff} = \min(\eta_{vis}, \eta_{pla}) \quad (9)$$

137 Partial melting, melt extraction and percolation are also considered in the model in a simplified
138 way (Gerya, 2013). The melt fraction (M_0) of the crust are assumed to increase with temperature and
139 are calculated according to Eq. (10):

140
$$M_0 = 0 \text{ when } T \leq T_{solidus}$$

141
$$M_0 = \frac{(T - T_{solidus})}{(T_{liquidus} - T_{solidus})} \quad \text{when } T_{solidus} < T < T_{liquidus} \quad (10)$$

142
$$M_0 = 1 \text{ when } T \geq T_{liquidus}$$

143 where $T_{solidus}$ and $T_{liquidus}$ are the solidus and liquidus temperature of different rock types,
144 respectively, taken from Katz et al. (2003).

145 In our model, melt extraction is modeled indirectly and considered as an instantaneous process
146 (Gerya et al., 2015). The extracted melt is assumed to move vertically from the molten source and
147 then added to the bottom of the crust. Partial melt is extracted from the mantle and instantaneously
148 displaced to the bottom of the crust and converted into hot mafic magma, obeying the conservation
149 of material. The amount of extracted melt during the evolution of each experiment is traced by the

150 Lagrangian markers (Gerya, 2013). The total amount of melt, M , for every marker excludes the
151 amount of previously extracted melt according to Eq. (11):

152
$$M = M_0 - \Sigma_n M_{ext} \quad (11)$$

153 where $\Sigma_n M_{ext}$ refers to the total melt fraction extracted during the previous n melt extraction
154 timesteps.

155 The effective density of mafic magma and molten crust depends on its melt fraction and is
156 calculated as (Gerya et al., 2015; Gülcher et al., 2020):

157
$$\rho_{eff} = \rho_{solid}(1 - M + M \frac{\rho_{0,molten}}{\rho_{0,solid}}) \quad (12)$$

158 where $\rho_{0,molten}$ and $\rho_{0,solid}$ are the reference densities of the molten and solid crust. ρ_{solid} is the
159 crust density at given pressure and temperature, which can be computed as:

160
$$\rho_{solid} = \rho_{0,solid}[1 - \alpha(T - 298)][1 + \beta(P - 0.1)] \quad (13)$$

161 with thermal expansion $\alpha = 3 \times 10^{-5} K^{-1}$ and compressibility $\beta = 10^{-11} Pa^{-1}$.

162 Surface processes, such as erosion and sedimentation, are considered by solving the transport
163 equation on the Eulerian nodes at each time step (Gerya and Yuen, 2003). Our erosion/sedimentation
164 model uses gross-scale erosion/sedimentation rates which are independent of local elevation and
165 topography (Burov and Cloetingh, 1997). We use constant and moderate rates of erosion (0.315
166 mm/yr) and sedimentation (0.0315 mm/yr), respectively, which falls within naturally observed
167 ranges.

168

169 2.2 Model setup

170 The size of the model box is 6600×1200 km, with a nonuniform grid of 501×301
171 computational nodes in length and depth, respectively (Fig. 2). The densest grid is located in the

center of the model domain (i.e., grid size decreases linearly from 20 km at the edges to 2 km at the ridge axis, and increases linearly from 1 km at the top to 7 km at the bottom), where plume-ridge interaction would happen. The model consists of a 20 km thick sticky air layer to accommodate crustal surface deformation. To reproduce the oceanic lithosphere, we choose a typical layered model, where the crust is composed of a water level (2 km), a sediment layer (1.5 km), and a basalt layer (7.5 km). The oceanic lithosphere and asthenosphere in the model are both modeled as dry olivine (the different colors for the mantle lithosphere and asthenosphere in the figures of this paper are only for better visualization). Besides, a 50-Myrs-old mid-ocean ridge is set on central part of the lithosphere, splitting the model domain into two parts. At the depth of 660 km, a 200-km-wide semicircular plume is located on the left of model domain, corresponding to the onset of plume-ridge interaction from the mantle transition zone. Detailed rock parameters are listed in Table 1.

The thermal conditions at the top and bottom boundaries are fixed at 273 and 2513 K, respectively. The left and right boundaries are both insulating, with no external heat flow across them. The initial temperature structure of the mantle is adiabatic (0.5 K km^{-1}), which results in a temperature at 660 km depth of 1843 K. The initial temperature structure of the oceanic plate consists of half-space cooling part and thermal equilibrium part (Fig. 2a). The half-space cooling model is used to describe the oceanic plate younger than 50 Myr, and the thermal equilibrium structure is used to describe older oceanic parts. In other words, the thermal age of the lithosphere far away from the ridge is fixed at 50 Myr with a constant plate thickness (i.e., $\sim 100\text{ km}$). The hot plume is set a circular thermal and compositional (see Table 1) anomaly with an excess temperature of 250 K to trigger a rising thermochemical plume. All the velocity boundaries are free slip boundaries. Additional internal boundary velocities are imposed at 500 km from each side boundary in the

删除[pangfengping]:
删除[pangfengping]:

194 lithosphere to maintain the imposed half spreading rate (Fig. 2a).

195

196 **Table 1.** Rock physical properties used in the numerical models.

Parameters	Sediments	Ocean Crust	Mantle	Plume	Reference ^a
Flow law	Wet quartz	Basalt	Dry olivine	Wet olivine	
Preexponential factor $A(\text{Pa}^n\text{s})$	1.97×10^{17}	4.80×10^{22}	3.98×10^{16}	5.01×10^{20}	1
Activation energy $E_a(\text{KJ mol}^{-1})$	154	238	532	470	1
Activation volume $V_a(\text{J bar}^{-1}\text{mol}^{-1})$	0	0	1	0.8	1
Exponent n	2.3	3.2	3.5	4	1
Cohesion $C(\text{Pa})$	2×10^7	2×10^7	2×10^7	2×10^7	1
Effective friction coefficient φ	0.6/0.3	0.6/0.3	0.6/0.3	0.6/0.3	1
Density $\rho(\text{Kg m}^{-3})$	2600	3000	3300	3270	2
Radioactive heating $H_r(\text{W m}^{-3})$	2×10^{-6}	2.2×10^{-7}	2.2×10^{-8}	2.5×10^{-8}	2

197 a: 1-(Ranalli, 1995), 2-(Turcotte and Schubert, 2014)

198 Other physical parameters used for all rocks include: gas constant $R=8.314\text{ J K}^{-1}\text{mol}^{-1}$, thermal
199 expansion $\alpha=3\times10^{-5}\text{ K}^{-1}$, compressibility $\beta=1\times10^{-11}\text{ Pa}^{-1}$, heat capacity $Cp=1000\text{ J kg}^{-1}\text{K}^{-1}$.

200

201 **3 Model Results**

202 We conduct a series of numerical experiments to investigate ridge suction versus plate drag acts
203 on plumes. The effects of three major model parameters (i.e., the spreading rate of mid-ocean ridge,
204 the initial plume head radius, and the plume-ridge distance) are systematically studied. The typical
205 dynamic evolution of models with ridge-ward vs. plate-drag plume flow are demonstrated.

206 **3.1 Model evolution with ridge-ward plume flow**

207 For models with dominant ridge-ward flow, the typical model evolution is shown in Fig. 3 (the
208 major model parameters used in this case are: the half spreading rate of 8 mm yr^{-1} , the initial plume

删除[pangfengping]:

删除[pangfengping]:

head radius of 200 km, and the off-axis distance of 800 km). In the early plume head stage, the buoyant mantle plume rises up rapidly in a mushroom-like shape and imposes dynamic stresses at the base of the overriding oceanic plate, leading to significant surface uplift (Figs. 3a,b). The ascending plume experiences extensive decompression melting at the base of the overriding plate, and due to the dynamic overpressure, spreads laterally, forming two branches that flow in opposite directions (Fig. 3c). A large amount of plume material is eventually entrained towards the spreading center, ponding underneath the ridge axis, and significantly affecting the ridge dynamics. The entrainment of hot plume material promotes decompression melting (Figs. 3d, e) and increases the temperature beneath the ridge (Fig. S2). Within the overlying lithosphere, the buoyant mantle plume leads to stress localization and strongly weakens the oceanic plate (Figs. S1, S3). As the plume eventually flows upward along the base of the gradually sloping plate near MOR, melting and crust production occurs (Fig. S1), forming an oceanic plateau of thickened crust. In addition to this gravitational force that guides plume material of the right branch ridge-ward, plate spreading drags both branches in the opposite direction. Moreover, convective and tectonic stresses (“plume push” and “ridge suction”) affect both branches of the plume in a different way. As a consequence, the two branches evolve asymmetrically: the right branch that flows towards the ridge axis is more vigorous than the left branch, and the plume tail is also tilted towards the spreading center (Figs. 3c-e).

删除[pangfengping]: (Fig. 3b)

删除[pangfengping]: Fig

删除[pangfengping]: increasingly sloping

删除[pangfengping]: the

删除[pangfengping]: For a more detailed discussion of the underling controlling forces, see below.

删除[pangfengping]:

删除[pangfengping]:




Figure 3. The evolution of the reference model M12 (see Table S1 in supplementary material) with dominant ridge-ward plume flow. The main model parameters employed in this case are: half spreading rate of 8 mm yr^{-1} , an initial plume head radius of 200 km, and an off-axis distance of 800 km. **(a)** surface topography over time along the flow path. **(b-e)** Snapshots of composition for the reference model (M12). **(f)** Profiles of the horizontal velocity component over time at the sections as indicated (color-coded) in panel (c-e).

The mantle flow horizontal velocity profiles (Fig. 3f) further demonstrate the dominance of ridge-ward plume flow, showing that plume flow is faster towards the spreading ridge than away from it. The velocity profiles elucidate dominant Poiseuille flow, with the maximum flow velocities in the middle of the asthenospheric channel. Such velocity profiles are well consistent with observations of seismic anisotropy at the Reunion plume (Barruol et al. 2019). The branches of the spreading plume head move significantly faster than the overriding plate. Because of the asymmetrical spreading of the plume head, the buoyancy flux carried by the right branch of the

删除[pangfengping]:

删除[pangfengping]: Therefore, plate drag actually slows down the spreading of the plume branches in this model case.

删除[pangfengping]:

删除[pangfengping]:

241 plume (density anomaly multiplied by horizontal velocity from Figure 3f) is also much larger than
242 that carried by the left branch.

243

244 **3.2 Model evolution with plate-drag plume flow**

245 For models with dominant plume flow away from the ridge (“plate-drag flow”), the typical
246 model evolution is shown in Fig. 4. The controlling parameters of the representative model shown in
247 Figure 4 are the same as for the model shown in Figure 3, except for a smaller radius (100 km) and
248 faster spreading ridge (half spreading rate: 45 mm yr⁻¹). At first, the ascending plume head spreads
249 out similarly as in the case described above and interacts with the overriding oceanic lithosphere.
250 The largest surface uplift is sustained just above the plume head (Fig. 4a), slightly different from the
251 previous model in which the highest surface elevation is observed on both sides of the plume conduit
252 (Fig. 3a). Related to this spreading and uplift, the mantle plume changes the stress state of the
253 overlying oceanic plate (Fig. S4), resulting in a slightly weakening (Fig. S6). The plume head
254 undergoes significant decompression melting near the deflection point (Fig. 4c). However, it is
255 difficult for small plume to extract molten plume due to the low melt flux. (Fig. S4). As the plume
256 cools, partially molten plume gets solidified speedily (Figs. 4d-e and S5). In contrast to the reference
257 model from section 3.1, this model displays most plume material flowing away from the ridge, likely
258 due to dominant plate drag (Figs. 4c-e). Indeed, the left branch of the plume consistently displays
259 larger buoyancy fluxes and maximum velocities than the right side over time (Fig. 4f).

260

删除[pangfengping]: divergent stresses are sustained in

删除[pangfengping]: lithosphere

删除[pangfengping]: but no

删除[pangfengping]: or yielding occurs

删除[pangfengping]: thick and cold lithosphere prevents
magma from extracting

删除[pangfengping]:

删除[pangfengping]:




Figure 4. Same as Figure 3 for case M77 (i.e., the reference model for the plate-drag plume flow regime). The main model parameters employed in this case are: half spreading rate of 45 mm yr^{-1} , an initial plume head radius of 100 km, and an off-axis distance of 800 km.

The underlying mechanism for dominant plate-drag plume flow is the frictional shear force of the moving plate, which is further demonstrated by the plume flow velocity profiles (Fig. 4f). In the early plume head stage ($\sim 1.08 \text{ Myr}$), the plume spreads out faster than plate velocity, which is primarily driven by the overpressure of the ponding plume head at this stage. After a short amount of time ($\sim 2.32 \text{ Myr}$), however, plume spreading becomes significantly slower than plate velocity, and hence plate drag drives and controls the plume flow. Indeed, the flow mode in the asthenosphere rapidly shifts from Poiseuille flow (i.e., active plume flow) to Couette flow (i.e., passive plume flow) (Fig. 4f), indicating the increasing role of plate drag on plume flow, soon after an initial of plume-head spreading.

删除[pangfengping]:

删除[pangfengping]:




Figure 5. Comparison between models with ridge-ward vs. plate-drag plume flow. **(a)** Ridge-ward flow with downwelling beneath the MOR (results from case M12 as in Figure 3). White dashed lines are streamlines; black arrows visualize the flow field. Schematic of flow in the sub-panel on the right-hand side. **(b)** Plate-drag flow with upwelling mantle corner flow beneath the MOR (results from case M77 as in Figure 4). **(c)** The dynamic pressure and gravitational gradient of plume marker (i.e. green circle in (a)) over time. **(d)** The dynamic pressure and gravitational gradient of plume marker (i.e. green circle in (b)) over time. The yellow box marks the location for the computation of average dynamic pressure at the ridge, needed for the calculation of the dynamic pressure gradient (see text). The (c) and (d) are representing the results of ridge-ward and plate drag flow, respectively.

3.3 Two modes of plume-lithosphere interaction

删除[pangfengping]:

删除[pangfengping]: The yellow box in (b)

删除[pangfengping]: (d) The dynamic pressure and gravitational gradient of plume marker (i.e. green circle in (b)) over time.

删除[pangfengping]:

删除[pangfengping]:

288 The dominant ridge-ward and dominant plate-drag plume flow regimes are two distinct modes of
289 plume-plate interaction. The differences between these two regimes are further demonstrated in
290 terms of mantle flow (Figs. 5a,b), driving forces (Figs. 5c,d).

291 In the ridge-ward dominated models, clockwise mantle develops from the plume to the spreading
292 ridge (Fig. 5a). Molten plume material flows to the spreading ridge and occupies the space
293 underneath the ridge axis, sustaining significant asymmetry of mid-ocean ridge melting (Conder et
294 al., 2002). As a consequence to the continuous supply of the plume material, downward mantle flow
295 forms beneath the ridge axis. This flow pattern dramatically differs from that shown in the plate-drag
296 dominated models, which show upward mantle flow underneath the ridge axis (Fig. 5b), as typical
297 for the flow beneath a MOR without the influence of a plume.

298 The distinct modes of plume-ridge interaction (ridge-ward vs. plate-drag flow) are controlled by
299 the competition of the tectonic (plate drag, ridge suction) and gravitational (plume buoyancy) driving
300 forces. On one hand, the moving plate drags sub-lithospheric plume material away from the ridge.
301 On the other hand, the mechanism of ridge-ward flow is twofold. First, the buoyant plume material
302 flows along the sloping base of the lithosphere towards the shallow ridge along the gravitational
303 gradient. Second, the plume is driven along the dynamic-pressure gradient from the pressure
304 maximum (e.g., where the plume sustains dynamic topograph) towards the pressure minimum
305 beneath the diverging ridge. These gravitational (G_{gv}) and pressure-driven (G_{dp}) gradients are
306 calculated by tracing plume markers (Figs. 5c,d) as follows:

307
$$G_{dp} = (P_{mk} - P_r)/L \quad (12)$$

308
$$G_{gv} = (\rho_0 - \rho_{mk}) * g * k \quad (13)$$

309 where P_{mk} is the dynamic pressure of plume marker and P_r is the averaged pressure in a 50 km box
310 at ridge center (Fig. 5b); L is the horizontal distance from plume marker to ridge axis; ρ_{mk} and ρ_0
311 are the plume marker density and initial density, respectively; g is the gravitational acceleration; k is
312 the local slope of the base of the lithosphere.

313 In the early stage of model evolution, the plume head's dynamic overpressure is dominant,
314 driving plume spreading in both directions (Figs.5c,d), in particular in the direction of the
315 low-pressure ridge. However, this pressure gradient systematically diminishes over time as the plume
316 (head) spreads. Once the spreading plume approaches the ridge, the lithospheric slope increases. At
317 some point, the gravitational gradient exceeds the dynamic pressure gradient, taking over as the
318 major driving force of guiding plume material towards the ridge. However, for low flux plume, the
319 dynamic pressure and gravitation are not enough to support the flow to the ridge. Consequently, one
320 of the essential conditions for plume-ridge interaction is that the plume must be able to reach the
321 critical zone near the ridge, where the slope is sufficiently steep to take over for the ever diminishing
322 pressure gradient. This implies that the plume buoyancy must (1) overcome the shearing force of
323 plate drag, and (2) the pressure-gradient must be sustained long enough to reach the critical zone, in
324 which the gravitational gradient can take over. The (1) shearing force scales with the rate of ridge
325 spreading, and the (2) plume with high buoyancy flux is more likely to reach the critical region at a
326 given plume-ridge distance.

329 **3.4 Influence of model parameters**

330 We have systematically investigated the effect of the three main model parameters (i.e., the

删除[pangfengping]: Fig.

设置格式[pangfengping]: 字体颜色: 背景 2

删除[pangfengping]: critical zone is more readily reached for

设置格式[pangfengping]: 字体颜色: 背景 2

删除[pangfengping]: fluxes

设置格式[pangfengping]: 字体颜色: 背景 2

删除[pangfengping]:

删除[pangfengping]:

spreading rate of the mid-ocean ridge, initial plume head radius and plume-ridge distance) on plume-ridge interaction. We explored half spreading rates of the ridge of 8, 15, 30, and 45 mm yr⁻¹, corresponding to ultra-slow, slow, medium, and fast-spreading mid-ocean ridges, respectively (Gerya, 2012). We varied initial plume head radii in the range of 100 km to 300 km. Further, we tested plume-ridge distance in the range of 600 to 1400 km.

3.4.1 Plume head radius

The size of the buoyant plume exerts an important control on plume-ridge interaction. Small plumes tend to be dragged away from the ridge, with typically larger lateral fluxes of the left branch than the right branch of the spreading plume (Figs. 6a,b). The buoyancy flux, defined as the integral over a horizontal plane of the product of the vertical velocity and the density deficit within the plume, in each branch is calculated by multiplying the velocity of the markers in plume pipe (Figs. 6d-f) by the density. The dynamic pressure decreases with decreasing plume size (Fig. S8a), and the pressures gradient is thus not strong enough for small plumes to reach the ridge. Plate shearing dominates plume flow soon after plume head spreading, and the moving plate then drags plume head material, leaving a tilted plume tail (Fig. 6d). In contrast, with larger initial plume head radius or buoyancy flux, the ponding plume spreads more vigorously (Fig. 6c) and sustains much higher overpressures at the base of the plate (Fig. S8a). This vigorous spreading can overcome plate drag to drive Poiseuille flow in both directions. Once the right plume branch approaches the spreading center, it is attracted and further accelerated by ridge suction. The plume tail is also markedly tilted towards the ridge axis due to asymmetric spreading (Fig. 6f). The larger the plume is, the more plume material gets entrained by the spreading center.





Figure 6 Models varying initial plume head radii (model M53, M58, and M63, Table S1 in supplementary material) shown by buoyancy flux and viscosity. **(a-c)** Buoyancy flux in spreading plume branches over time. Green and red triangles are markers used for buoyancy flux calculations of left and right plume branches, respectively. U and R stand for half of spreading rate and plume radius. **(d-f)** Viscosity snapshots of models with different plume head radii. Models with green circle represent plate-drag plume flow and ridge-ward plume flow in red.

3.4.2 Plume-ridge distance

Plume-ridge distance also controls the regime of plume-ridge interaction. A plume at large distances spreads similarly as a plume at a small distance, but is less likely to get affected by ridge

suction (Fig. 7). The pressure gradient between the plume and ridge drives the ridge-ward plume flow. However, the larger the plume-ridge distance, the smaller the pressure gradient would be (Fig. S8b), resulting in a lower buoyancy flux across the plume pipe (Figs. 7a-c). In the cases of distant plumes, the spreading of the plume head is strongly affected by plate drag (Figs. 7b, c).

删除[pangfengping]: Figs. 7e,f

删除[pangfengping]: On the other hand, the difficulty in sustaining ridge-ward plume flow may also link to the heat transfer between the cold plate and the hot plume rocks. With gradually cooling from upper plate by heat conduction and diffusion, the viscosity of plume increases as it cools. Such increasing viscosity slows the plume down, stopping the ridge-ward plume flow eventually (Figs. 7e, f). Hence, for cases with large plume-ridge distances and hence travel times, the ponding plume head cools and is ultimately carried away by the moving plate.





Figure 7. Models varying plume-ridge distances (model M57-M59, Table S1 in supplementary material) shown by buoyancy flux and viscosity. (a-c) Buoyancy flux in spreading plume branches over time. Green and red triangles are markers used for buoyancy flux calculations of left and right plume branches, respectively. U and D stand for half of spreading rate and plume-ridge distance. (d-f) Viscosity snapshots of models with different plume-ridge distances. Models with green circle represent plate-drag plume flow and ridge-ward plume flow in red.



删除[pangfengping]: calculation.

删除[pangfengping]: calculation.

删除[pangfengping]:



删除[pangfengping]:

3.4.3 Half spreading rate of ridge

Another parameter that is worth investigating is the spreading rate of the ridge. The modeling results indicate that fast-spreading ridges promote plume flow away from the ridge due to the friction (Figs.8 and 9a). With increasing spreading rate, the effect of plate shearing on plume-lithosphere interaction increases, as quantified by the spreading fraction. The spreading fraction γ (Eq.(14)) is defined here as the ratio of ridge-ward vs. plate-drag plume volume fluxes. We integrated the ridge-ward plume volume flux (right branch), V_{rw} , and plate-drag plume volume flux (left branch), V_{tw} . V_p is the total plume volume flux in the model. Ridge-ward plume spreading is dominant for positive γ ; plate-drag plume spreading is dominant for negative γ .

$$\gamma = (V_{rw} - V_{tw})/V_p \quad (14)$$

In the early stage (~ 1 Myr), pressure-driven flow dominates in all models and spreading fractions are positive, mainly driven by the expansion of the overpressured plume heads along the pressure gradient. After a certain time, the spreading fractions decrease dramatically with the decay of the mantle plume activity, representing the transition from the ridge-ward to the plate-drag regime in some cases. The characteristic spreading fractions after 8 Myr model time as a function of our model parameters are shown in Fig. 8. This compilation of our results reveals that the dominance of ridge-ward flow decreases with increasing spreading rate and off-axis distance, but significantly increases with plume size. For models with fast-spreading ridges, the parameter range of plate-drag flow dominated models is expanded, indicating the critical role of plate drag in restricting ridge-ward flow and plume-ridge interaction.



删除[pangfengping]:

Figure 8. Parameter regime diagram of the contrasting modes of plume-ridge interaction. Spreading fractions γ (Eq. (14)) at ~ 8 Myr model time. Each of the circles represents one of the numerical experiments, and sizes refer to γ . Circles in red and green represent models with dominant ridge-ward plume flow and plate drag, respectively.

The transition from ridge-ward (positive γ) to plate-drag (negative γ) flow in some of our cases is mainly determined by the competition between the effects of pressure-driven plume head spreading and plate shearing. The overpressure in the plume head drives plume materials towards the lower pressure spreading center, while the moving plate shears plume away. Hence, we quantify the shear force of the overriding oceanic plate on the plume head using an integral approach:

$$F_s = \int \sigma_{xz} dA \quad (15)$$

Equation (15) is employed to calculate the shear force, where F_s is the total shear force the spreading oceanic plate exerts on the uppermost part of the plume. σ_{xz} is the shear stress on each mantle plume grid cell, A refers to the area of each grid cell. The pressure gradients, both

删除[pangfengping]:

删除[pangfengping]:

gravitational and dynamic pressure, are calculated by tracing the plume markers according to equations (12-13). As the plume material rises to the base of the lithosphere, the shear force exerted by the plate increases over time. We find that the integrated shear force between the spreading plate and the plume increases significantly as half spreading rate increases (Fig. 9c).

Conversely, ridge spreading rates control gravitational and pressure-driven plume driving forces (Fig. 9d). Increasing the spreading rate of the ridge implies a smaller dynamic pressure gradient, because the pressure gradient is related to the plate thickness difference at the ridge and plume, which is dependent on the spreading rate. A fast-spreading ridge also implies a smaller gravitational gradient, because it leaves a more shallowly-dipping lithospheric base. Thus, relatively strong plate shearing combined with relatively small pressure and gravitational gradients tend to advance plate-drag plume flow for high spreading rates.





Figure 9. Model results influenced by different half spreading rates. (a) Effect of half spreading rate and plume-ridge distance on ridge-ward vs. plate-drag. Viscosity snapshots are shown (model M7-M9, M82-M84, Table S1 in supplementary material). Fast-spreading ridge promotes plume material dragged when plume near the ridge. Models with green circle represent plate-drag plume flow and



删除[pangfengping]:

删除[pangfengping]: flow

删除[pangfengping]: .

删除[pangfengping]: flow

删除[pangfengping]: .

删除[pangfengping]:

删除[pangfengping]:

ridge-ward plume flow in red. **(b)** Dynamic evolutions of ridge-ward and plate-drag plume flow, revealed by defined ridge spreading fraction (eq.14). **(c)** Shear force (F_s) between moving plate and plume material under different spreading rates. **(d)** Pressure gradient between plume head and ridge center in different half spreading rate models. The solid and dash lines are the plume gravitation and dynamic pressure gradient, respectively. R and D stand for half of plume radius and plume-ridge distance.

4 Discussion


Natural observations show that there are only very few hotspots indicative of ridge-ward plume flow close to the East Pacific Rise (EPR) (Fig. 10a), in contrast to many such hotspots in the Atlantic and Indian oceans. A previous study (Jellinek et al., 2003) proposed that fast-spreading ridges such as the EPR efficiently convey any surrounding plumes into the spreading center from the deep mantle (Fig. 1c), which leads to fewer hotspots nearby fast-spreading ridges. However, based on our modeling results, fast-spreading ridges tend to promote plate-drag flow of the spreading plume material, providing an alternative explanation to the relatively absence of hotspots along the EPR. We discuss the possibility of this potential explanation combined with geological and geophysical observations (Fig. 10).

Firstly, the plate drag effect of fast-spreading ridges on plumes is evidenced by geophysical observations. We locate the positions of the mantle plumes at the core-mantle boundary (CMB) and the associated hot spots on the surface based on global seismic tomography (Jackson et al., 2021; Koppers et al., 2021). A lateral offset between the deep and surface positions of plumes is a common

删除[pangfengping]: viability
删除[pangfengping]: by comparing

删除[pangfengping]:
删除[pangfengping]:

449 feature, indicating the deflection of plumes due to mantle flow. Specifically, a large portion of (7 of
450 14) plumes located in the Atlantic are tilted towards the mid-ocean ridge. However, only very few
451 plumes (6 of 16) in the Pacific are tilted towards the mid-ocean ridge. Indeed, the majority of plumes
452 are tilted away from the ridges, indicating the significant effect of plate drag on plumes beneath fast
453 plates. Such observations are consistent with the predictions of our models with dominant plate-drag
454 plume spreading.



455 **Figure 10.** A compilation of hotspots along with spreading ridges in the Atlantic and the Pacific. (a)

457 Distribution of surface hotspots (circles) together with depth-projected source locations at CMB
458 (blue dots) of the plumes based on (Jackson et al., 2021). Plumes in magenta circles are mantle
459 plumes interacted with ridges (Ito et al., 2003), and plumes not interacted with ridges are shown as
460 green circles, whose size refers to the plume buoyancy flux (Hoggard et al., 2020). Blue arrows
461 indicate the changes in the position of the plume at the top and bottom of the mantle. Yellow dots are
462 MORB samples mapped in (b). **(b)** Plot of radioactive isotopes ratios along ridge MORB samples.
463 The data are downloaded from the PetDB Database (<http://portal.earthchem.org/>). The colored
464 symbols refer to samples in different mid-ocean ridge. Main hotspots influencing MORBs are
465 labeled with shaded bands. The black dash lines are the mean MORB isotopes ratio from Gale (2013).
466 Red and green lines are the mean ratios of the samples in Mid-Atlantic ridge and EPR, respectively.

467
468 Geochemical studies suggest that mantle plumes are enriched in light rare earth elements
469 (LREEs) and radiogenic isotopes of Sr and Pb but depleted in Nd isotopes. These geochemical
470 anomalies are evident in MORB at the sites of active plume-ridge interaction (Cushman et al., 2004;
471 Douglass and Schilling, 1999; Yang et al. 2017). We find that MORB sampled along both the
472 Mid-Atlantic ridge and the EPR indeed display geochemical anomalies (Fig. 10b), indicating
473 ridge-ward flow of plume material at specific locations. However, the Mid-Atlantic MORB dataset is
474 slightly more heterogeneous than the East Pacific Rise in terms of geochemical isotopes. The EPR is
475 basically characterized as normal oceanic basalt, along which only very few regions show
476 composition associated with nearby plumes. This contradicts the view (Jellinek et al., 2003) that
477 mantle plumes are fully entrained into the central MOR melting zone at fast-spreading ridges.

478 Based on our modeling results, initial plume head radius and plume-ridge distance also control

删除[pangfengping]:
删除[pangfengping]:

the mode of plume-ridge interaction. However, for different buoyancy fluxes B , there is only a small difference in the proportion of interacting plumes to non-interacting plumes. A small majority of major plumes (5 of 8 with $B > 1.6$ Mg/s), and a small minority of small-to-intermediate plumes (11 of 25 for $B < 1.6$ Mg/s) display interaction with the ridge (Fig. 11a). The underlying cause for this observation remains unclear, but may be related to the distribution of large plumes globally with many of them being located very far from MORs. Also note that our 2D models are limited in that plume material cannot spread in the out-of-plane direction, hence somewhat exaggerating the effects of buoyancy flux. In any case, the distribution of observed plume buoyancy fluxes (Hoggard et al., 2020) varies little across different oceans (Fig. 11a). Therefore, the effects of plume size are not a good candidate to explain the notable difference between the Atlantic and Pacific in terms of plume-ridge interaction mode.

On the other hand, compared with the Atlantic and Indian Oceans, Pacific plumes are located significantly further from the mid-ocean ridge (Fig. 11b). Plume-ridge distances in the Pacific are mostly >2000 km, which exceeds the maximum plume-ridge interaction distance of 1400 km (Schilling, 1991). Most plumes in the Pacific exhibit the typical signatures of plume flow away from the ridge, such as parabolic swell shapes (e.g., Society, Marquesas and Hawaii plumes; Ballmer et al., 2013a; Ballmer et al., 2015; Cheng et al., 2015; Wolfe et al., 2009), and linear volcanic chains (Buff et al., 2021; Clouard and Bonneville, 2005; Jackson et al., 2010). Age-progressive hotspots trails indicate an absence of dominant ridge-ward flow. By contrast, most plumes in the Atlantic have been close to the ridge since the opening of the ocean. These mantle plumes (e.g., Discovery, Iceland, Tristan-Gough; O'Connor et al., 2012) did not move much since the breakup of the Atlantic. One factor may be that the underlying plume generation zone (i.e., the edge of the African LLSVP) round

设置格式[pangfengping]: 字体颜色: 背景 2
设置格式[pangfengping]: 字体颜色: 背景 2
删除[pangfengping]: terms of
设置格式[pangfengping]: 字体颜色: 背景 2
删除[pangfengping]: fraction
设置格式[pangfengping]: 字体颜色: 背景 2
删除[pangfengping]: vs.
设置格式[pangfengping]: 字体颜色: 背景 2
删除[pangfengping]: for different buoyancy fluxes B : a
设置格式[pangfengping]: 字体颜色: 背景 2
删除[pangfengping]:) vs.
设置格式[pangfengping]: 字体颜色: 背景 2
设置格式[pangfengping]: 字体颜色: 红色

删除[pangfengping]:
删除[pangfengping]:

501 largely parallel to the Mid-Atlantic Ridge (Fig. 1) (Torsvik et al., 2006). In this case, plume-ridge
502 distance may play a critical role in the plume-ridge interaction, and could explain the striking
503 difference between the Pacific and Atlantic in terms of the proportion of plume-ridge interacting vs.
504 non-interacting systems. In addition, the rapid movement of the Pacific plate tends to inhibit
505 ridge-ward plume flow at a given plume-ridge distance. The distribution of interacting (stars) vs
506 non-interacting systems in Figure 11b is almost exactly as predicted by our models for the coupled
507 effects of plume-ridge distance and plate velocity. For example, we note that fast-spreading ridges
508 can still interact with adjacent plumes under the appropriate conditions. In the case of very short
509 plume-ridge distances, there is good evidence of plume-ridge interaction in the southern Pacific
510 ocean (e.g., Louisville plume; Conder et al., 2002; Toomey et al., 2002; Vlastélic and Dosso, 2005).
511 Based on a series of numerical modeling as well as geological and geophysical observations, we
512 conclude that mantle plumes in the Pacific are more likely to spread away from the ridge and into the
513 direction of plate motion than in the Atlantic and Indian Oceans. The tendency of fast plate velocities
514 to promote plume spreading away from the MOR through viscous drag may depend, however, on the
515 details of lithosphere-asthenosphere rheological coupling such as the presence of a weak decoupling
516 (e.g., melt) layer (Rychert et al., 2020). Further studies of plume spreading and plume-ridge
517 interaction are needed to shed light on the coupling of the plate-mantle system.

删除[pangfengping]: number

删除[pangfengping]:

删除[pangfengping]:




Figure 11. Buoyancy flux, plate speed and plume-ridge distance of mantle plumes in different oceans. Mantle plumes in the Pacific, Atlantic and Indian Ocean are shown in green, red and pink circles, respectively. Blue stars mark the ridge-interacted plumes according to Ito et al. (2003). **(a)** Plot of plume-ridge distance and plume buoyancy flux. Data are from Hoggard et al. (2020). **(b)** Plot of plume-ridge distance and plate speed at the location of plumes. Plume-ridge distance comes from GPlates (Müller et al., 2016; Whittaker et al., 2015), and plate speed data comes from Becker et al. (2015)

5 Conclusion

In this study, we explore the evolution of plume-ridge interaction with 2D thermomechanical numerical models. Based on model results, we find that:

- (1) Plume-ridge interaction is mainly governed by the competition between the effects of plume spreading (overpressure in the plume-head stage), upward gravitationally-driven flow of the plume along the base of the sloping lithosphere and plate shearing. These driving forces are controlled by plume size, plume-ridge distance and the spreading rate of the mid-ocean ridge.

534 (2) MOR spreading does not only draw upwelling plumes into the spreading center, but also tends to
535 drag mantle plumes away from the ridge. Plume flow away from the ridge is favored by small
536 and/or distant plumes as well as fast spreading rates, whereas plume flow towards the ridge is
537 promoted by large and/or nearby plumes, as well as slow spreading rates.

538 (3) Considering the high plate velocity and typically large plume-ridge distances, mantle plumes in
539 the Pacific are more likely to be dragged away from the EPR than being drawn towards the ridge
540 center.

541

542

Code availability

The source numerical modeling code in this study is available from the corresponding author upon reasonable request.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author contribution

Fengping Pang performed all numerical models, interpreted results and wrote the manuscript. Jie Liao proposed the study, modify the code and contributed to rewriting and scientific discussion. Maxim D. Ballmer contributed with significant help in rewriting and scientific discussion. Lun Li participated in discussion and interpretations. All authors have read and edited draft versions of the paper and have approved the final version.

Competing interest

The authors declare that they have no conflict of interest.

Acknowledgement

This research is financially supported by NSFC projects (U1901214, 41974104, 91855208) and Guangdong project 2017ZT07Z066. We are grateful to Prof. Taras Gerya for his long-lasting guidance on our geodynamical modeling. We gratefully acknowledge Hongjian Fang for insightful

删除[pangfengping]:
删除[pangfengping]:

565 discussions. Numerical simulations were performed on the clusters of National Supercomputer

566 Center in Guangzhou (Tianhe-II).

567

568 **Reference**

569 Ballmer, M. D., Ito, G., Wolfe, C. J. and Solomon, S. C.: Double layering of a thermochemical

570 plume in the upper mantle beneath Hawaii, *Earth Planet. Sci. Lett.*, 376, 155–164,

571 doi:10.1016/j.epsl.2013.06.022, 2013a.

572 Ballmer, M. D., Conrad, C. P., Smith, E. I. and Harmon, N.: Non-hotspot volcano chains produced

573 by migration of shear-driven upwelling toward the East Pacific Rise, *Geology*, 41(4), 479–482,

574 doi:10.1130/G33804.1, 2013b.

575 Ballmer, M. D., Ito, G. and Cheng, C.: Asymmetric dynamical behavior of thermochemical plumes

576 and implications for hawaiian lava composition, *Geophys. Monogr. Ser.*, 208, 35–57,

577 doi:10.1002/9781118872079.ch3, 2015.

578 Barruol, G., Sigloch, K., Scholz, J. R., Mazzullo, A., Stutzmann, E., Montagner, J. P., Kiselev, S.,

579 Fontaine, F. R., Michon, L., Deplus, C. and Dymant, J.: Large-scale flow of Indian Ocean

580 asthenosphere driven by Réunion plume, *Nat. Geosci.*, 12(12), 1043–1049,

581 doi:10.1038/s41561-019-0479-3, 2019.

582 Becker, T. W., Schaeffer, A. J., Lebedev, S. and Conrad, C. P.: Toward a generalized plate motion

583 reference frame, *Geophys. Res. Lett.*, 42(9), 3188–3196, doi:10.1002/2015GL063695, 2015.

584 Buff, L., Jackson, M. G., Konrad, K., Konter, J. G., Bizimis, M., Price, A., Rose-Koga, E. F.,

585 Blusztajn, J., Koppers, A. A. P. and Herrera, S.: “Missing Links” for the Long-lived Macdonald and

586 Arago Hotspots, South Pacific Ocean, *Geology*, 49(5), 541–544, doi:10.1130/G48276.1, 2021.

587 Burov, E. and Cloetingh, S.: Erosion and rift dynamics: New thermomechanical aspects of post-rift

588 evolution of extensional basins, *Earth Planet. Sci. Lett.*, 150(1–2), 7–26,

589 doi:10.1016/s0012-821x(97)00069-1, 1997.

删除[pangfengping]:

删除[pangfengping]:

590 Byerlee, J.: Friction of rocks, Pure Appl. Geophys. PAGEOPH, 116(4–5), 615–626,
591 doi:10.1007/BF00876528, 1978.

592 Cheng, C., Allen, R. M., Porritt, R. W. and Ballmer, M. D.: Seismic constraints on a double-layered
593 asymmetric whole-mantle plume beneath Hawai’i, Hawaiian Volcanoes From Source to Surf., 19–34,
594 doi:10.1002/9781118872079.ch2, 2015.

595 Clouard, V. and Bonneville, A.: Ages of seamounts, islands, and plateaus on the Pacific plate, Spec.
596 Pap. Geol. Soc. Am., 388, 71–90, doi:10.1130/0-8137-2388-4.71, 2005.

597 Conder, J. A., Forsyth, D. W. and Parmentier, E. M.: Asthenospheric flow and asymmetry of the East
598 Pacific Rise, MELT area, J. Geophys. Res. Solid Earth, 107(B12), ETG 8-1-ETG 8-13,
599 doi:10.1029/2001jb000807, 2002.

600 Cushman, B., Sinton, J., Ito, G. and Dixon, J. E.: Glass compositions, plume-ridge interaction, and
601 hydrous melting along the Galapagos spreading center, 90.5 °wto 98 ° W, Geochemistry, Geophys.
602 Geosystems, 5(8), doi:10.1029/2004GC000709, 2004.

603 Dalton, C. A., Langmuir, C. H. and Gale, A.: Geophysical and geochemical evidence for deep
604 temperature variations beneath mid-ocean ridges, Science (80-.), 344(6179), 80–83,
605 doi:10.1126/science.1249466, 2014.

606 Douglass, J., Schilling, J. G. and Fontignie, D.: Plume-ridge interactions of the Discovery and Shona
607 mantle plumes with the southern Mid-Atlantic Ridge (40°-55° S), J. Geophys. Res. Solid Earth,
608 104(B2), 2941–2962, doi:10.1029/98jb02642, 1999.

609 François, T., Koptev, A., Cloetingh, S., Burov, E. and Gerya, T.: Plume-lithosphere interactions in
610 rifted margin tectonic settings: Inferences from thermo-mechanical modelling, Tectonophysics,
611 746(October 2015), 138–154, doi:10.1016/j.tecto.2017.11.027, 2018.

删除[pangfengping]:
删除[pangfengping]:

612 French, S. W. and Romanowicz, B.: Broad plumes rooted at the base of the Earth’s mantle beneath
613 major hotspots, *Nature*, 525(7567), 95–99, doi:10.1038/nature14876, 2015.

614 Gale, A., Dalton, C. A., Langmuir, C. H., Su, Y. and Schilling, J. G.: The mean composition of ocean
615 ridge basalts, *Geochemistry, Geophys. Geosystems*, 14(3), 489–518, doi:10.1029/2012GC004334,
616 2013.

617 Geissler, W. H., Wintersteller, P., Maia, M., Strack, A., Kammann, J., Eagles, G., Jegen, M.,
618 Schloemer, A. and Jokat, W.: Seafloor evidence for pre-shield volcanism above the Tristan da Cunha
619 mantle plume, *Nat. Commun.*, (2020), doi:10.1038/s41467-020-18361-4, 2020.

620 Gerya, T.: Origin and models of oceanic transform faults, *Tectonophysics*, 522–523, 34–54,
621 doi:10.1016/j.tecto.2011.07.006, 2012.

622 Gerya, T. V.: Three-dimensional thermomechanical modeling of oceanic spreading initiation and
623 evolution, *Phys. Earth Planet. Inter.*, 214, 35–52, doi:10.1016/j.pepi.2012.10.007, 2013.

624 Gerya, T. V. and Yuen, D. A.: Characteristics-based marker-in-cell method with conservative
625 finite-differences schemes for modeling geological flows with strongly variable transport properties,
626 *Phys. Earth Planet. Inter.*, 140(4), 293–318, doi:10.1016/j.pepi.2003.09.006, 2003.

627 Gerya, T. V. and Yuen, D. A.: Robust characteristics method for modelling multiphase
628 visco-elasto-plastic thermo-mechanical problems, *Phys. Earth Planet. Inter.*, 163(1–4), 83–105,
629 doi:10.1016/j.pepi.2007.04.015, 2007.

630 Gerya, T. V., Stern, R. J., Baes, M., Sobolev, S. V. and Whattam, S. A.: Plate tectonics on the Earth
631 triggered by plume-induced subduction initiation, *Nature*, 527(7577), 221–225,
632 doi:10.1038/nature15752, 2015.

633 Gülcher, A. J. P., Gerya, T. V., Montési, L. G. J. and Munch, J.: Corona structures driven by

删除[pangfengping]:
删除[pangfengping]:

634 plume–lithosphere interactions and evidence for ongoing plume activity on Venus, *Nat. Geosci.*,
635 13(8), 547–554, doi:10.1038/s41561-020-0606-1, 2020.

636 Hardarson, B. S., Fitton, J. G., Ellam, R. M. and Pringle, M. S.: Rift relocation - A geochemical and
637 geochronological investigation of a palaeo-rift in northwest Iceland, *Earth Planet. Sci. Lett.*,
638 153(3–4), 181–196, doi:10.1016/s0012-821x(97)00145-3, 1997.

639 Harmon, N., Forsyth, D. W., Weeraratne, D. S., Yang, Y. and Webb, S. C.: Mantle heterogeneity and
640 off axis volcanism on young Pacific lithosphere, *Earth Planet. Sci. Lett.*, 311(3–4), 306–315,
641 doi:10.1016/j.epsl.2011.09.038, 2011.

642 Hoggard, M. J., Parnell-turner, R. and White, N.: Hotspots and mantle plumes revisited: Towards
643 reconciling the mantle heat transfer discrepancy, *Earth Planet. Sci. Lett.*, 542, 116317,
644 doi:10.1016/j.epsl.2020.116317, 2020.

645 Ito, G., Lin, J. and Gable, C. W.: Interaction of mantle plumes and migrating mid-ocean ridges:
646 Implications for the Galapagos plume-ridge system, , v, 1–3, 1997.

647 Ito, G., Lin, J. and Graham, D.: Observational and theoretical studies of the dynamics of mantle
648 plume-mid-ocean ridge interaction, *Rev. Geophys.*, 41(4), doi:10.1029/2002RG000117, 2003.

649 Jackson, M. G., Hart, S. R., Konter, J. G., Koppers, A. A. P., Staudigel, H., Kurz, M. D., Blusztajn, J.
650 and Sinton, J. M.: Samoan hot spot track on a “hot spot highway”: Implications for mantle plumes
651 and a deep Samoan mantle source, *Geochemistry, Geophys. Geosystems*, 11(12),
652 doi:10.1029/2010GC003232, 2010.

653 Jackson, M. G., Becker, T. W. and Steinberger, B.: Spatial Characteristics of Recycled and
654 Primordial Reservoirs in the Deep Mantle, *Geochemistry, Geophys. Geosystems*, 22(3),
655 doi:10.1029/2020GC009525, 2021.

656 Jellinek, A. M., Gonnermann, H. M. and Richards, M. A.: Plume capture by divergent plate motions:
657 Implications for the distribution of hotspots, geochemistry of mid-ocean ridge basalts, and estimates
658 of the heat flux at the core-mantle boundary, *Earth Planet. Sci. Lett.*, 205(3–4), 361–378,
659 doi:10.1016/S0012-821X(02)01070-1, 2003.

660 Jiang, Q., Jourdan, F., Olierook, H. K. H., Merle, R. E. and Whittaker, J. M.: Longest continuously
661 erupting large igneous province driven by plume-ridge interaction, *Geology*, 1–3,
662 doi:10.1130/G47850.1, 2020.

663 Kincaid, C., Ito, G. and Gable, C.: Laboratory investigation of the interaction of off-axis mantle
664 plumes and spreading centres, *Nature*, 376(6543), 758–761, doi:10.1038/376758a0, 1995.

665 Kincaid, C., Schilling, J.-G. and Gable, C.: The dynamics of off-axis plume-ridge interaction in the
666 uppermost mantle, *Earth Planet. Sci. Lett.*, 137(1–4), 29–43, doi:10.1016/0012-821X(95)00201-M,
667 1996.

668 Koppers, A. A. P., Becker, T. W., Jackson, M. G., Konrad, K., Müller, R. D., Romanowicz, B.,
669 Steinberger, B. and Whittaker, J. M.: Mantle plumes and their role in Earth processes, *Nat. Rev.*
670 *Earth Environ.*, 2(6), 382–401, doi:10.1038/s43017-021-00168-6, 2021.

671 Lénat, J. F., Merle, O. and Lespagnol, L.: La réunion: An example of channeled hot spot plume, *J.*
672 *Volcanol. Geotherm. Res.*, 184(1–2), 1–13, doi:10.1016/j.jvolgeores.2008.12.001, 2009.

673 Maia, M., Pessanha, I., Courrges, E., Patriat, M., Gente, P., Hémond, C., Janin, M., Johnson, K.,
674 Roest, W., Royer, J. Y. and Vatteville, J.: Building of the Amsterdam-Saint Paul plateau: A 10 Myr
675 history of a ridge-hot spot interaction and variations in the strength of the hot spot source, *J. Geophys.*
676 *Res. Solid Earth*, 116(9), 1–19, doi:10.1029/2010JB007768, 2011.

677 Mittelstaedt, E., Ito, G. and Behn, M. D.: Mid-ocean ridge jumps associated with hotspot magmatism,

删除[pangfengping]:
删除[pangfengping]:

678 Earth Planet. Sci. Lett., 266(3–4), 256–270, doi:10.1016/j.epsl.2007.10.055, 2008.

679 Mittelstaedt, E., Ito, G. and Van Hunen, J.: Repeat ridge jumps associated with plume-ridge
680 interaction, melt transport, and ridge migration, J. Geophys. Res. Solid Earth, 116(1), 1–20,
681 doi:10.1029/2010JB007504, 2011.

682 Mittelstaedt, E., Soule, S., Harpp, K., Fornari, D., McKee, C., Tivey, M., Geist, D., Kurz, M. D.,
683 Sinton, C. and Mello, C.: Multiple expressions of plume-ridge interaction in the Galápagos: Volcanic
684 lineaments and ridge jumps, Geochemistry, Geophys. Geosystems, 13(5),
685 doi:10.1029/2012GC004093, 2012.

686 Montelli, R., Nolet, G., Dahlen, F. A., Masters, G., Engdahl, E. R. and Hung, S. H.: Supporting
687 OnlineMaterial Timing, Science, 303(5656), 338–343, doi:10.1126/science.1092485, 2004.

688 Morgan, W. J.: Rodriguez, Darwin, Amsterdam, ..., A second type of Hotspot Island, J. Geophys.
689 Res. Solid Earth, 83(8), 5355–5360, 1978.

690 Müller, R. D., Roest, W. R. and Royer, J.-Y.: Asymmetric sea-floor spreading caused by
691 ridge–plume interactions, Nature, 396(6710), 455–459, doi:10.1038/24850, 1998.

692 Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., Shephard, G.
693 E., Maloney, K. T., Barnett-Moore, N., Hosseinpour, M., Bower, D. J. and Cannon, J.: Ocean Basin
694 Evolution and Global-Scale Plate Reorganization Events since Pangea Breakup, Annu. Rev. Earth
695 Planet. Sci., 44, 107–138, doi:10.1146/annurev-earth-060115-012211, 2016.

696 Niu, Y.: Ridge suction drives plume-ridge interactions, , (October), doi:10.13140/2.1.4728.0961,
697 2014.

698 O’Connor, J. M., Jokat, W., Le Roex, A. P., Class, C., Wijbrans, J. R., Keßling, S., Kuiper, K. F. and
699 Nebel, O.: Hotspot trails in the South Atlantic controlled by plume and plate tectonic processes, Nat.

700 Geosci., 5(10), 735–738, doi:10.1038/ngeo1583, 2012.

701 Ranalli: Rheology of the Earth, 1995.

702 Ribe, N. M.: The dynamics of plume-ridge interaction: 2. Off-ridge plumes, , 101, 1996.

703 Ribe, N. M. and Christensen, U. R.: Three-dimensional modeling of plume-lithosphere interaction, ,

704 99, 669–682, 1994.

705 Ribe, N. M. and Christensen, U. R.: The dynamical origin of Hawaiian volcanism, Earth Planet. Sci.

706 Lett., 171(4), 517–531, doi:10.1016/S0012-821X(99)00179-X, 1999.

707 Ribe, N. M., Christensen, U. R. and Theiβing, J.: The dynamics of plume-ridge interaction, 1:

708 Ridge-centered plumes, Earth Planet. Sci. Lett., 134(1–2), 155–168,

709 doi:10.1016/0012-821X(95)00116-T, 1995.

710 Rowley, D. B. and Forte, A. M.: Kinematics of the East Pacific Rise Retrodicted From Pacific and

711 Farallon/Nazca Subduction-Related Torques: Support for Significant Deep Mantle Buoyancy

712 Controlling EPR Spreading, J. Geophys. Res. Solid Earth, 127(2), 1–24, doi:10.1029/2020JB021638,

713 2022.

714 Rowley, D. B., Forte, A. M., Rowan, C. J., Glišović, P., Moucha, R., Grand, S. P. and Simmons, N.

715 A.: Kinematics and dynamics of the east pacific rise linked to a stable, deep-mantle upwelling, Sci.

716 Adv., 2(12), 1–19, doi:10.1126/sciadv.1601107, 2016.

717 Rychert, C. A., Harmon, N., Constable, S. and Wang, S.: The Nature of the

718 Lithosphere-Asthenosphere Boundary, J. Geophys. Res. Solid Earth, 125(10), 1–39,

719 doi:10.1029/2018JB016463, 2020.

720 Schilling, J. G.: Fluxes and excess temperatures of mantle plumes inferred from their interaction with

721 migrating mid-ocean ridges, Nature, 352(6334), 397–403, doi:10.1038/352397a0, 1991.

删除[pangfengping]:

删除[pangfengping]:

722 Sleep, N. H.: Lateral flow and ponding of starting plume material, J. Geophys. Res. Solid Earth,
723 102(B5), 10001–10012, doi:10.1029/97jb00551, 1997.

724 Small, C.: Observations of ridge-hotspot interactions in the Southern Ocean, J. Geophys. Res.,
725 100(B9), doi:10.1029/95jb01377, 1995.

726 Straume, E. O., Gaina, C., Medvedev, S., Hochmuth, K., Gohl, K., Whittaker, J. M., Abdul Fattah, R.,
727 Doornenbal, J. C. and Hopper, J. R.: GlobSed: Updated Total Sediment Thickness in the World’s
728 Oceans, Geochemistry, Geophys. Geosystems, 20(4), 1756–1772, doi:10.1029/2018GC008115,
729 2019.

730 Toomey, D. R., Wilcock, W. S. D., Conder, J. A., Forsyth, D. W., Blundy, J. D., Parmentier, E. M.
731 and Hammond, W. C.: Asymmetric mantle dynamics in the MELT region of the East Pacific Rise,
732 Earth Planet. Sci. Lett., 200(3–4), 287–295, doi:10.1016/S0012-821X(02)00655-6, 2002.

733 Torsvik, T. H., Smethurst, M. A., Burke, K. and Steinberger, B.: Large igneous provinces generated
734 from the margins of the large low-velocity provinces in the deep mantle, Geophys. J. Int., 167(3),
735 1447–1460, doi:10.1111/j.1365-246X.2006.03158.x, 2006.

736 Turcotte, D. and Schubert, G.: Geodynamics, Cambridge University Press., 2014.

737 Vlastélic, I. and Dosso, L.: Initiation of a plume-ridge interaction in the South Pacific recorded by
738 high-precision Pb isotopes along Hollister Ridge, Geochemistry, Geophys. Geosystems, 6(5), 1–13,
739 doi:10.1029/2004GC000902, 2005.

740 Whittaker, J. M., Afonso, J. C., Masterton, S., Müller, R. D., Wessel, P., Williams, S. E. and Seton,
741 M.: Long-term interaction between mid-ocean ridges and mantle plumes, Nat. Geosci., 8(6),
742 doi:10.1038/NGEO2437, 2015.

743 Wolfe, C. J., Solomon, S. C., Laske, G., Collins, J. A., Detrick, R. S., Orcutt, J. A., Bercovici, D. and

删除[pangfengping]:
删除[pangfengping]:

744 Hauri, E. H.: Mantle shear-wave velocity structure beneath the Hawaiian hot spot, Science (80-.),
745 326(5958), 1388–1390, doi:10.1126/science.1180165, 2009.

746 Yang, A. Y., Zhao, T. P., Zhou, M. F. and Deng, X. G.: Isotopically enriched N-MORB: A new
747 geochemical signature of off-axis plume-ridge interaction—A case study at 50°28'E, Southwest
748 Indian Ridge, J. Geophys. Res. Solid Earth, 122(1), 191–213, doi:10.1002/2016JB013284, 2017.

749

Supplementary material for

**Plume-ridge interactions: Ridge-ward versus plate-drag plume
flow**

Fengping Pang¹, Jie Liao^{1,2,3}, Maxim D. Ballmer⁴, Lun Li^{1,2,3}

¹School of Earth Sciences and Engineering, Sun Yat-Sen University, Guangzhou 510275, China

²Guangdong Provincial Key Lab of Geodynamics and Geohazards, Guangzhou 510275, China

³Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai 519000, China

⁴Department of Earth Sciences, University College London, London, United Kingdom

Contents of this file

Figures S1 to S8
Table S1




Figure S1. Reference model (M12, see Table S1, same as Fig. 3) evolution of ridge-ward plume flow shown by (a) crust and sediment thickness, (b) normal stress. The mantle plume weakens the overlying oceanic plate and changes the stress state of the overlying oceanic plate. Molten plume material beneath the lithosphere is extracted to the crust.

删除[pangfengping]:

删除[pangfengping]:




Figure S2. Temperature evolution of reference ridge-ward plume flow model (M12, see Table S1, same as Fig. 3).





Figure S3. Viscosity evolution of reference ridge-ward plume flow model (M12, see Table S1, same as Fig. 3).

删除[pangfengping]:

删除[pangfengping]:



删除[pangfengping]:

删除[pangfengping]:




Figure S4. Reference model (M77, see Table S1, same as Fig. 4) evolution of plate-drag plume flow shown by (a) crust and sediment thickness, (b) normal stress. The mantle plume weakens the overlying oceanic plate and changes the stress state of the overlying oceanic plate. Molten plume material beneath the lithosphere is extracted to the crust.




Figure S5. Temperature evolution of reference plate-drag plume flow model (M77, see Table S1, same as Fig. 4).




Figure S6. Viscosity evolution of reference plate-drag plume flow model (M77, see Table S1, same as Fig. 4).

删除[pangfengping]:

删除[pangfengping]:




Figure S7. Shear force (F_s) between plate and plume in different plume head size models. The shear force imposed on the plume increases with plume size. The negative shear force indicates stronger friction imposed on the ridge-ward flowing (right) plume branch than that on the plate-drag flowing (left) plume branch.




Figure S8. Pressure gradient between plume head and ridge center. The plume gravitation (solid lines) and dynamic pressure gradient (dash lines) of **(a)** different size plumes, **(b)** different plume-ridge distances are shown by solid and dash lines, respectively.

删除[pangfengping]: gravitation
删除[pangfengping]: pressure

删除[pangfengping]:
删除[pangfengping]:

Table S1. Description of experiments.

Model	Ridge half spreading rate (mm/yr)	Plume radius (km)	Plume ridge distance (km)	Plume excess temperature (K)	Regimes
M1	8	100	600	250	Ridge-ward flow
M2	8	100	800	250	Plate-drag flow
M3	8	100	1000	250	Plate-drag flow
M4	8	100	1200	250	Plate-drag flow
M5	8	100	1400	250	Plate-drag flow
M6	8	150	600	250	Ridge-ward flow
M7	8	150	800	250	Ridge-ward flow
M8	8	150	1000	250	Ridge-ward flow
M9	8	150	1200	250	Ridge-ward flow
M10	8	150	1400	250	Ridge-ward flow
M11	8	200	600	250	Ridge-ward flow
M12	8	200	800	250	Ridge-ward flow
M13	8	200	1000	250	Ridge-ward flow
M14	8	200	1200	250	Ridge-ward flow
M15	8	200	1400	250	Ridge-ward flow
M16	8	250	600	250	Ridge-ward flow
M17	8	250	800	250	Ridge-ward flow
M18	8	250	1000	250	Ridge-ward flow
M19	8	250	1200	250	Ridge-ward flow
M20	8	250	1400	250	Ridge-ward flow
M21	8	300	600	250	Ridge-ward flow
M22	8	300	800	250	Ridge-ward flow
M23	8	300	1000	250	Ridge-ward flow
M24	8	300	1200	250	Ridge-ward flow
M25	8	300	1400	250	Ridge-ward flow
M26	15	100	600	250	Ridge-ward flow
M27	15	100	800	250	Plate-drag flow
M28	15	100	1000	250	Plate-drag flow
M29	15	100	1200	250	Plate-drag flow
M30	15	100	1400	250	Plate-drag flow
M31	15	150	600	250	Ridge-ward flow
M32	15	150	800	250	Ridge-ward flow
M33	15	150	1000	250	Ridge-ward flow
M34	15	150	1200	250	Plate-drag flow
M35	15	150	1400	250	Plate-drag flow
M36	15	200	600	250	Ridge-ward flow
M37	15	200	800	250	Ridge-ward flow

M38	15	200	1000	250	Ridge-ward flow
M39	15	200	1200	250	Ridge-ward flow
M40	15	200	1400	250	Ridge-ward flow
M41	15	250	600	250	Ridge-ward flow
M42	15	250	800	250	Ridge-ward flow
M43	15	250	1000	250	Ridge-ward flow
M44	15	250	1200	250	Ridge-ward flow
M45	15	250	1400	250	Ridge-ward flow
M46	15	300	600	250	Ridge-ward flow
M47	15	300	800	250	Ridge-ward flow
M48	15	300	1000	250	Ridge-ward flow
M49	15	300	1200	250	Ridge-ward flow
M50	15	300	1400	250	Ridge-ward flow
M51	30	100	600	250	Ridge-ward flow
M52	30	100	800	250	Plate-drag flow
M53	30	100	1000	250	Plate-drag flow
M54	30	100	1200	250	Plate-drag flow
M55	30	100	1400	250	Plate-drag flow
M56	30	150	600	250	Ridge-ward flow
M57	30	150	800	250	Ridge-ward flow
M58	30	150	1000	250	Plate-drag flow
M59	30	150	1200	250	Plate-drag flow
M60	30	150	1400	250	Plate-drag flow
M61	30	200	600	250	Ridge-ward flow
M62	30	200	800	250	Ridge-ward flow
M63	30	200	1000	250	Ridge-ward flow
M64	30	200	1200	250	Ridge-ward flow
M65	30	200	1400	250	Plate-drag flow
M66	30	250	600	250	Ridge-ward flow
M67	30	250	800	250	Ridge-ward flow
M68	30	250	1000	250	Ridge-ward flow
M69	30	250	1200	250	Ridge-ward flow
M70	30	250	1400	250	Ridge-ward flow
M71	30	300	600	250	Ridge-ward flow
M72	30	300	800	250	Ridge-ward flow
M73	30	300	1000	250	Ridge-ward flow
M74	30	300	1200	250	Ridge-ward flow
M75	30	300	1400	250	Ridge-ward flow
M76	45	100	600	250	Ridge-ward flow
M77	45	100	800	250	Plate-drag flow
M78	45	100	1000	250	Plate-drag flow
M79	45	100	1200	250	Plate-drag flow
M80	45	100	1400	250	Plate-drag flow

删除[pangfengping]:

删除[pangfengping]:

M81	45	150	600	250	Ridge-ward flow
M82	45	150	800	250	Ridge-ward flow
M83	45	150	1000	250	Plate-drag flow
M84	45	150	1200	250	Plate-drag flow
M85	45	150	1400	250	Plate-drag flow
M86	45	200	600	250	Ridge-ward flow
M87	45	200	800	250	Ridge-ward flow
M88	45	200	1000	250	Ridge-ward flow
M89	45	200	1200	250	Plate-drag flow
M90	45	200	1400	250	Plate-drag flow
M91	45	250	600	250	Ridge-ward flow
M92	45	250	800	250	Ridge-ward flow
M93	45	250	1000	250	Ridge-ward flow
M94	45	250	1200	250	Ridge-ward flow
M95	45	250	1400	250	Ridge-ward flow
M96	45	300	600	250	Ridge-ward flow
M97	45	300	800	250	Ridge-ward flow
M98	45	300	1000	250	Ridge-ward flow
M99	45	300	1200	250	Ridge-ward flow
M100	45	300	1400	250	Ridge-ward flow