

Reply to Anonymous referee

Dear Sir, Madam,

thank you very much for your input on the manuscript, it is highly appreciated. Here is our reply to your comments. We hope the changes we implemented improve the shortcomings of the manuscript highlighted by your comments and suggestions. Please do not hesitate to contact us shall this not be the case for some comments.

1. Comments from Anonymous referee

Comment 1: I suggest that this paper be accepted/reconsidered after major revisions. In general, I enjoyed this paper: it is a good, well-written paper with a structurally interesting dataset from a major transform plate boundary fault zone. The dataset is collected from a transpressional uplift within the San Andreas fault zone, then compared to other similar features along strike. As such, the paper stands to be a good contribution for those trying to understand the internal structure, along-strike complexity, and tectonic evolution of transform plate boundary fault zones, and more specifically the along-strike complexity of the southeastern terminus of the San Andreas transform plate boundary fault.

Comment 2: The overwhelming majority of my comments are minor, albeit numerous. However, there are a few major points concerning the figures that need to be addressed should the manuscript be accepted for publication. These few major points concerning the figures may take some time to complete, and are my only reason for listing the revision as major, not minor. These include: Figure 1 needs to be redone to include a regional map with all the features discussed in the text plotted on that map and, in general, showing the study area in the regional context (southern California, southwestern USA). An updated figure could take the form of a two-panel figure, where Fig. 1a is the regional map showing major features discussed in text, and Fig. 1b is the close-up map that is currently presented as the sole Fig. 1. At present, the reader has no regional context for the features discussed in-text, and some features and faults are not shown on any map, making their comparison and importance to the study area difficult and unclear.

Comment 3: All maps in the figures (Figs. 2, 3, 5, and 6) should have coordinates of some sort, whether as points or a grid. Additionally, I suggest that un-interpreted images of all of the map

areas should be added to the supplemental material (an un-interpreted Fig. 6 is already in the SM).

Comment 4: Folds and faults mapped on Fig. 2 appear continuous across some parts of the Northwestern, Central, and Southeastern domains. However, in Figs. 3 and 6, the folds and faults appear short and discontinuous. These figures should be updated to reflect the full extent of the structure(s) within the figure's frame to be consistent with their geology on the ground and as shown on Fig. 2. Should these changes be addressed, I think the paper will make a good contribution. Good luck, and I hope to see this in print in the near future.

Comment 5: Specific Comments - Title: Should a broader geographic description be applied to the title, given this is European journal but the study area is in the USA? Perhaps "Tectonic evolution of the Indio Hills segment of the San Andreas fault in southern California, southwestern USA"

Comment 6: Line 46-47 - What about this continuation in to the ECSZ? The sentence needs more description about the significance of the Indio Hills fault with the ECSZ.

Comment 7: Line 68 - I am curious about the use of the term "culmination" - I am only familiar with this term in fold-thrust systems. As defined at <https://link.springer.com/content/pdf/bbm%3A978-94-011-3066-0%2F1.pdf> : "Culmination: An anticline or dome with four way closure generated by movement of the thrust sheet over underlying ramps." I understand you have transpressional folding/thrusting going on in your study area, so the term could be used, but does the Indio Hills exhibit folding over underlying thrust ramps? Or are you simply referring to a variety of distinct tectonic elements all observed together in one place? If the latter, I think a different term is warranted. If you choose to keep the term culmination, I think you need to explicitly define it, either here or in your Tectonic Culminations section below. Perhaps it is best to simply call it the Indio Hills uplift here on Line 68, as you do in the Fig. 1 caption, and leave the use of culmination (if you keep it) for the section below.

Comment 8: Line 69 - You state the Indio Hills are a transpressional uplift, but consider it analogous to a rift feature (which would suggest transtension)? See next comment.

Comment 9: Line 68-70 - I think what you mean is that the Indio Hills and Mecca Hills are analogous in that they are both inverted basins? If that is correct, be more explicit here. For example, you could say: "The Indio Hills uplift is an inverted Miocene–Pliocene sedimentary

basin lying upon Mesozoic granitic basement rocks. Further to the southeast, the Mecca Hills are also shown to be an inverted Miocene–Pliocene sedimentary basin (Keller et al., 1982; Damte, 1997; McNabb et al., 2017; Bergh et al., 2019).”

Comment 10: Lines 84-86 – You state “We consider” but then list references. Are you interpreting that these units are The Mecca Formation, or did the cited authors interpret these units to be the Mecca Formation in the Indio Hills. The former is slightly problematic, as it is an interpretation before the data section (but understandably a necessary one to make for your study).

Comment 11: Line 103-104 - How would sediment accumulation rates define the age of a formation? More than likely, the dates of those stratigraphic members were used to calculate the sediment accumulation rates. Did the lower and upper members of the Palm Springs Formation show increased rates of sediment accumulation during these intervals? If so, specify that.

Comment 12: Lines 114-117 - See above comment on use of the term culmination. You only use the term four times in the paper, here three times and once in the former section. I suspect the term should be changed, given the formal definition I pasted in the Line 68 comment above, but if you choose to keep the term then define what you mean by “culmination” either here or at Line 68.

Comment 13: Line 119-124 - Your broad-scope description of tectonic elements here shows the necessity of adding a regional map to your Figure 1. At present, the reader has no context for the Eastern California shear zone (which is a much broader region than you show it in Fig. 1), the San Bernardino and San Jacinto faults, and San Geronio Pass. These features need added to a map with the location of the study area clearly shown so the reader can see their relationship and importance to the work presented here.

Comment 14: Line 155-156 – Delete the word “off-fault” – damage zones typically encompass principal slip surfaces but are technically part of the fault zone, too, so it seems kind of like a misnomer to say off-fault

Comment 15: Lines 160-163 – Is there a reference for this statement?

Comment 16: Line 219 - Are you saying that the open upright fold geometry is the result of (via) the kink/chevron styles? If so, no change is really necessary, but perhaps it could be described more clearly? If not, and instead you are describing a sequence of changing fold patterns, then I'd replace "via" with "to"

Comment 17: Line 241 – Here again with “via”. Do you mean the something is the result of the kink/chevron geometry, or are you saying it is spatially changing from symmetric style, to then changing to a kink/chevron style, to then changing to isoclinal? If so, I'd suggest replacing "via" with "to"

Comment 18: Line 273 – What do you mean by monocline-like? It seems the fold would either be a monocline or an anticline, not a mix of the two. According to your Fig. 3C the fold closest to the Indio Hills fault very much looks to be an asymmetric anticline, with 20NE dip on the northeast limb and 45SW dip on the southwest limb, in which case I would delete “monocline-like” from the sentence.

Comment 19: Line 299-305 – What kind of folds are these? Anticline? Syncline? Both? Note that hinge lines are not mapped on Figure 5 like they are on Figure 2.

Comment 20: Lines 307-319; Lines 310-311 – If you are discussing faults and fractures in the basement, then that is not a fold-related fault (unless the basement is folded). Perhaps the section should be renamed “Major and minor faults, fractures, and fold-related faults”

Comment 21: Line 388 – Cite Figure 2 stereonet at the end of the sentence. Also, these fracture sets look to be $\sim 90^\circ$ to one another; I'd expect conjugates to be $\sim 60^\circ$ ($40-70^\circ$) to one another. It might be best to delete the “possibly representing conjugate sets” from the sentence, as I don't think these are conjugates. This shouldn't pose a problem, as you don't discuss these features any further in the manuscript.

Comment 22: Line 416 – “...indicates a younger phase of deformation.” Saying younger slip event makes it sound like only one slip event caused the present-day observed deformation pattern.

Comment 23: Line 419 – You could delete “strain” after shortening; since shortening is a strain term it is a little redundant.

Comment 24: Lines 433-440 – I think it would help the reader here to remind them stratigraphically which unit overlies/underlies which unit, or which unit is older and which unit is younger. E.g., “the Mecca Formation and overlying Palm Springs Formation.” or something to that effect.

Comment 25: Line 435 – would the fault be below the contact between the PS and MH formations, or would the fault be at/near the contact of the PS and MH formations?

Comment 26: Line 456-457 – dip-slip fault-parallel fold: wouldn't this just be a fault-propagation fold? I suppose it could also be a fault-bend fold by that description, but I get the impression it is fault-propagated.

Comment 27: Line 509 – stress, or strain?

Comment 28: Line 516 – I'd be more satisfied if these features were rigorously measured and restored back to a discrete bedding orientation through stereonet analysis. As presented in Supplement S6, you are “restoring” apparent dips at the outcrop face to an approximate horizontal based on the apparent dip of bedding in the picture/outcrop face. I absolutely agree with what you are saying and interpreting, but wonder if you should not refer to this as a restoration, per se, but rather that these features “appear to define a low-angle fold and thrust system (Supplement S6).”

Comment 29: Lines 558-559 – Concerning use of axial surfaces (i.e., axial planes), wouldn't a surface/plane be E-W-striking (not trending)? Perhaps it is better to just say E-W-trending folds.

Comment 30: Line 667 – Transpressional plate regime. Are you suggesting that the plate is entirely under transpression in this area (in which case, which plate – or both?), or are you saying San Andreas fault zone transpression? Depending which you mean could be important, as just to the east ~100 km some of us are arguing for late Miocene–Pliocene (and possibly ongoing) transtension in the lower Colorado River corridor. This is all the more complicated in that the ECSZ does seem to be overwhelmingly transpressive. Looking through your cited reference (Bergh et al., 2019), I think you mean San Andreas fault zone transpression – if so, please modify the “transpressional plate regime” part of the sentence to instead reflect SAFZ transpression. If you indeed mean transpression across the plate(s), I think you need to be more specific of the extent of this transpressional plate regime, and possibly even reconcile your claims by looking into recent literature for Pacific-North America plate boundary transtension inboard of the SAFZ just next door to the east (e.g., Singleton et al., 2019; Thacker et al., 2020; Dorsey et al., 2021), albeit ca. 3 to 1 Ma earlier than you propose the Indio Hills to have formed.

Comment 31: Lines 713-722; Point 4 in Conclusions (Lines 738-740) – I am a bit confused about how the Indio Hills and Durmid Hills are shown as initially different in Fig. 8a, but in this paragraph you suggest that the two areas might be similar in that the Indio Hills might be an early phase of a ladder structure like the Durmid Hills. In Fig. 8a you clearly show an inherent difference between the two areas: Indio Hills has E-W folds, Durmid Hills has NE-SW left-lateral

faults - am I to assume the E-W folds had already formed, or did E-W folds not form, which would again suggest an inherent difference between the two areas? I am also confused how these two areas are potentially similar when the proposed timing of fault activation for the oblique dextral-reverse fault in both locations is opposite: The Indio Hills fault (what became the oblique dextral-reverse fault) formed before the Banning fault, while the Eastern Shoreline fault (what became the oblique dextral-reverse fault) formed after the main San Andreas fault, according to your figure.

Comment 32: Technical Corrections – Line 33 – A geographic description is required. For example: “...San Andreas fault zone (SAFZ: Fig. 1; California, southwestern USA), ...”

Comment 33: Line 34 – add “the” (“...deformation compared to the Mecca Hills...”)

Comment 34: Lines 41-42 – Note that “Eastern California shear zone” is commonly written with “shear zone” not capitalized. Change here, and throughout the manuscript to be “Eastern California shear zone”

Comment 35: Line 60 – delete “transform” and remove “s” from movements so it reads “...North American plates and movement along the SAFZ...” Also, should it be North America plate or North American plate?

Comment 36: Line 65 (end of paragraph) – I think a final sentence is needed here that brings it all back into perspective. Perhaps something akin to: "This recent work provides the opportunity to explore the understudied Indio Hills segment in order to compare its structural development with other along-strike uplifted features on a major transform plate boundary fault zone."

Comment 37: Line 128 – Gorgonio is misspelled

Comment 38: Line 127-130 – Suggest breaking this one sentence into two different sentences.

Comment 39: Line 134 – Eastern California shear zone (decapitalize shear zone) – change here and throughout the manuscript and figure captions.

Comment 40: Line 137 – Delete “attitude and” so the sentence reads “Farther southeast, however, the geometry of the...”

Comment 41: Line 138 – Add an “s” to remains

Comment 42: Line 140-142 - Suggest separating these into two sentences: "The transpressional character of the Indio Hills uplift was suggested by Parrish (1983) and Sylvester and Smith (1987). Recent work, however, has not been conducted, and detailed structural analyses have not been published from this segment of the SAFZ."

Comment 43: Line 143 – perhaps change focusing to “that focused”

Comment 44: Line 149 – Be explicit here with who you are referring to. I think you mean Keller et al. (1982). If so, I suggest replacing "Their" with the reference.

Comment 45: Line 173 - I think you mean main San Andreas fault strand, based on the abbreviation, but that is not totally clear as written. Suggest saying "main San Andreas fault (mSAF) strand..."

Comment 46: Line 178 and throughout the manuscript and figures – Make sure to decapitalize "fault" after all formal names. E.g., East Shoreline fault, Banning fault, etc., even for San Andreas fault.

Comment 47: Lines 204-205 – I think this sentence needs reworked: "The study area comprises three major fold systems that are oblique to the SAFZ. These fold systems are E–W trending, moderately west-plunging, and contain multiple smaller-scale parasitic folds (Fig. 2)."

Comment 48: Lines 243–244 – This is more of an editorial preference by EGU, but I don't think forelimb and backlimb need dashed? If not, change throughout the manuscript. If so, ignore.

Comment 49: Line 267 – I think you mean southeastern here, not southwestern

Comment 50: Line 314 – Change offset to displacement.

Comment 51: Lines 327-329 – Add “for a damage zone of a”: “The granite there is highly fractured and cut by vein and joint networks (see description below), as is expected for a damage zone of a major brittle fault.”

Comment 52: Line 377 – minor-scale (needs a dash I think)

Comment 53: Line 386 – in other places I think you refer to it as a leucogranite. Be consistent, whether you choose simply granite or leucogranite.

Comment 54: Lines 396-397 – Suggested rewording: “The folds are arranged in a right-stepping pattern, and are increasingly asymmetric and sigmoidal (Z-shaped) to the northeast as they approach the Indio Hills fault.” Change as you see fit, but at present the sentence is difficult to understand.

Comment 55: Line 429 – as inferred for other parts of the SAFZ

Comment 56: Line 430 – remove en dash (–) in front of to

Comment 57: Line 506 – perhaps just say slip here, not “the last slip event”

Comment 58: Lines 560-562 – Your sentence is in present tense (“this is observed”) but you refer to the Banning fault as you interpret it to have been at a former time. Perhaps say “what was then a precursory Banning fault.”

Comment 59: Lines 601-602 – Should be Eastern California shear zone (says East, not Eastern, and shear zone needs decapitalized)

Comment 60: Lines 607-608 – These two faults do not appear to be on Figure 1

Comment 61: Line 610 – delete comma after “enhanced”

Comment 62: Lines 633-634 – Earlier in the paper (and in Fig. 8) you define main San Andreas fault as mSAF, whereas here you say main SAFZ. Is there a reason for the difference (e.g., one refers to a discrete/singular fault plane, whereas the other refers to the main fault zone)? Should mSAF just be changed to main SAFZ, or vice versa? Also do this at Lines 40, 117, 608, 637, 654, 690, 695, 698, and in various figures.

Comment 63: Line 639 – the Indio Hills fault (missing “the”)

Comment 64: Lines 634 and 649 – On line 634 you reference Fig. 8c before referencing 8a and 8b, and on line 649 you reference Fig. 8c before referencing Fig. 8b. You do reference Fig. 8 in its entirety at line 627 – this is more of an editorial decision by EGU if subfigures can be referenced out of sequence.

Comment 65: Line 652 – missing a reference

Comment 66: Line 680 – Eastern Shoreline fault (combine Shore and line)

Comment 67: Line 689 – Here I think you mean Eastern Shoreline fault

Comment 68: Line 692 – see comment above about main SAFZ and mSAF. Here you say main SAF, which you defined earlier in the paper as mSAF - should this one be mSAF or main SAFZ?

Comment 69: Line 695 and 697 – Eastern Shoreline fault

Comment 70: Line 736 – delete “in”

Comment 71: **Detailed comments on figures** – Figure 1 – Figure 1 needs a regional scope. At the very least, a regional map showing California and the study area should be squeezed onto to Figure 1. However, I’d suggest a more detailed regional map showing structural relationships in the area and the numerous features mentioned in the text that are not on any of the maps (e.g., San Geronio Pass). For example, from Figure 1, the reader at present would have no context to the extent of the Eastern California shear zone. This can be done as a two-panel figure, where

Fig. 1a is a regional map showing major features discussed in the text and the field area, and Fig. 1b can be the present Fig. 1 map.

Comment 72: Line 982 – Brawley Seismic Zone needs defined as BSZ in the caption.

Comment 73: Lines 985-986 – As in the manuscript, decapitalize shear zone in “Eastern California shear zone” in this caption and in all figure captions. It is okay, of course, for the abbreviation to be ECSZ.

Comment 74: All fault names here, in all figure captions, and throughout the manuscript should not have “fault” capitalized as part of the name. E.g., Banning Fault should be Banning fault, etc.

Comment 75: Figure 2 – I hate to be a stickler here because these are GoogleEarth images, but all maps (Figs. 2, 3, 5, 6) should technically have at least a few coordinates, whether as a grid of lat/long or UTM, or a few lat/long coordinate points.

Comment 76: Note that your typed words have the spell check wiggle line underneath them. Make sure your final image does not have these.

Comment 77: On your stereonets labels, I suggest adding that the first two are bedding: “SAFZ-oblique bedding planes” and “SAFZ-parallel bedding planes”

Comment 78: In the text I think you say faults and fractures, but here you only say fractures. If both were measured, both should be specified here: “Sediments faults and fractures”; “Basement faults and fractures”

Comment 79: What program did you use to make the stereonets? Allmendinger’s? You should probably cite the program, unless it is a script you wrote.

Comment 80: Figures 2, 3, 5, and 6 – In the supplemental file you have an un-interpreted figure 6; it would be good to also put un-interpreted images of figures 2, 3, and 5 in the supplemental file as well.

Comment 81: Figures 3, 5, and 6 – Your mapped features (fold hinges and faults) commonly end before the end of the figure’s frame, whereas on Figure 2 many of these same features are shown to be continuous across the frame of the figure. I would suggest mapping the features along their full extent and ending them at the end of the figure frame, instead of cutting them short within the figure frame. As currently drawn, it gives the impression to the reader that these folds and faults are short and discontinuous only within the frame of the figure, but Figure 2 shows clearly that many of these features are continuous from one domain into the other. For example, the southeastern corner of Fig. 3b is also the northwestern corner of Fig. 3c – from southwest to

northeast there is an anticline then syncline then overturned anticline then overturned syncline/syncline – I think that these are the same folds in both figures, but as currently drawn Figs. 3b and 3c give the impression these are different folds.

Comment 82: Figure 3 – What are the yellow dots? I think these are photograph locations; if so state this in the Figure 3 caption.

Comment 83: In each panel (a, b, and c), you could place the domain name right above the scale bar. For example: in Fig. 3a, label “Northwestern” above the scale bar, “Central” in 3b, and “Southeastern” in 3c. This would make it easier for the reader.

Comment 84: Figure 4 – Line 1019 – The stereonet represents the cm-scale folds, correct? If so, add “cm-scale” to the caption so it is clear to the reader that these are the small cm-scale folds that cannot be seen in the photos.

Comment 85: Figure 5 – Label “Banning fault” on the main figure.

Comment 86: The fold hinge should be mapped on this figure like it is in Fig. 2.

Comment 87: Figure 7 – Line 1040 – You say Tentative model here; tentative on what? Perhaps just say “Model illustrating...”

Comment 88: Figure 8 – I make this point at Lines 633-634, but in your figure, how does SAFZ differ from mSAF? Is one a discrete fault that is considered the main strand (mSAF) and the other is a zone of deformation (SAFZ)? Is using mSAF necessary?

2. Author’s reply

Comment 1: agreed.

Comment 2: agreed.

Comment 3: agreed.

Comment 4: agreed. See response to comment 81.

Comment 5: agreed.

Comment 6: agreed.

Comment 7: agreed. Yes, the Indio Hills exhibit folding over underlying thrust ramps, as proposed for the SAFZ-parallel anticline near the Indio Hills fault (see also Supplement S3a for an example in the field). We agree though that it is certainly more appropriate to use the term “uplift” instead of “culmination” to avoid confusion.

Comment 8: agreed. See response to comment 9.

Comment 9: agreed.

Comment 10: agreed.

Comment 11: agreed.

Comment 12: agreed. See response to comment 7.

Comment 13: agreed. See response to comment 2.

Comment 14: agreed.

Comment 15: agreed.

Comment 16: agreed.

Comment 17: agreed.

Comment 18: agreed.

Comment 19: agreed.

Comment 20: agreed.

Comment 21: agreed.

Comment 22: agreed.

Comment 23: agreed.

Comment 24: agreed.

Comment 25: the fault would be below the contact, not at/near the contact.

Comment 26: agreed.

Comment 27: agreed.

Comment 28: agreed.

Comment 29: agreed.

Comment 30: agreed.

Comment 31: the Durmid Hills and Indio Hills uplifts are located on either sides of the main San Andreas fault and the E–W-trending macro-folds in the Durmid Hills seem to have formed slightly (but perhaps not significantly) after those in the Indio Hills (see new Table 1 for the timing of the main geological events in the Coachella Valley as suggested by the other reviewer). These uncertainties around the timing of geological events in the various uplifted areas along the main San Andreas fault are, no doubt, related to the sparsity of geochronological ages of structures along the fault. As depicted by the similar timing of uplift in all three uplifted areas, it is probable that all areas evolved (almost) synchronously, but even if it were the case, it is not yet possible to argue

for such a scenario. More geochronological data and absolute ages are needed in this part of California.

Comment 32: agreed.

Comment 33: deformation is meant in a general sense.

Comment 34: agreed.

Comment 35: agreed. However, “North American plate” is the correct term.

Comment 36: agreed.

Comment 37: agreed.

Comment 38: agreed.

Comment 39: agreed. See response to comment 34.

Comment 40: agreed.

Comment 41: agreed.

Comment 42: agreed.

Comment 43: the sentence was deleted and the paragraph reworked according to the other reviewer’s comments.

Comment 44: agreed. However, inserted reference earlier than suggested by the anonymous reviewer’s comment.

Comment 45: agreed.

Comment 46: agreed.

Comment 47: agreed.

Comment 48: agreed. However, the authors of the present manuscript will wait for proof-reading comments by the editorial team to make the suggested correction in case it is not required by the journal.

Comment 49: agreed.

Comment 50: agreed.

Comment 51: agreed.

Comment 52: agreed.

Comment 53: agreed.

Comment 54: agreed.

Comment 55: agreed.

Comment 56: agreed.

Comment 57: agreed.

Comment 58: agreed.

Comment 59: agreed. Also see response to comment 34.

Comment 60: agreed.

Comment 61: agreed.

Comment 62: disagreed. “main SAFZ” should be changed to “main San Andreas fault”.

Comment 63: agreed.

Comment 64: agreed.

Comment 65: agreed.

Comment 66: agreed.

Comment 67: agreed. Also did this throughout the manuscript.

Comment 68: agreed.

Comment 69: agreed. See response to comment 67.

Comment 70: agreed.

Comment 71: agreed. See response to comment 2.

Comment 72: agreed.

Comment 73: agreed. See response to comment 34.

Comment 74: agreed.

Comment 75: agreed. See response to comment 3.

Comment 76: agreed.

Comment 77: agreed.

Comment 78: disagreed. The term “fracture” is general and applies both to “faults” and “fractures”.

Comment 79: agreed.

Comment 80: agreed. See response to comment 3.

Comment 81: agreed.

Comment 82: agreed.

Comment 83: since the three macro-folds are shown in order from northwestern to southeastern, it is unnecessary to specify the name of the “domain” on each part of Figure 3. In addition, this could give the impression to the reader that, e.g., only the southeastern macro-fold may be observed on Figure 3c, which is not the case since the central macro-fold is also shown there.

Comment 84: agreed.

Comment 85: agreed.

Comment 86: agreed.

Comment 87: agreed.

Comment 88: agreed.

3. Changes implemented

Comment 1: none commanded by the reviewer's comment.

Comment 2: designed new figure 1a and b.

Comment 3: added coordinates to Figure 2, on which all Google Earth images are located. Also added uninterpreted version of all Google Earth images to the supplements and reorganized the supplement numbers in the manuscript.

Comment 4: see response to comment 81.

Comment 5: added “, southwestern USA” in the title.

Comment 6: added “and its role as possible transfer fault” lines 51–52.

Comment 7: replaced “culmination” by “uplift” lines 74, 76, 198, 229, 680, 720 and 980, and by “tectonic uplifts” line 840.

Comment 8: see response to comment 9.

Comment 9: changed “culmination” into “uplift” line 122. Changed “analogous” into “an analog” and “rift” into “inverted” line 124.

Comment 10: replaced “We consider” by “Previous mapping in the area (Dibblee, 1954; Lancaster et al., 2012) considered” lines 139–140, and deleted reference to Dibblee (1954) line 142.

Comment 11: deleted “) are consistent with sediment-accumulation rate estimates (” lines 165–166.

Comment 12: see response to comment 7.

Comment 13: see response to comment 2.

Comment 14: deleted “off-fault” line 249.

Comment 15: added reference to Sylvester and Smith (1976, 1979, 1987) and Bergh et al. (2019) lines 256–257.

Comment 16: replaced “via” by “to” line 319.

Comment 17: replaced “via” by “to” line 345.

Comment 18: deleted “to monocline-like” lines 379–380.

Comment 19: replaced “folds” by “synclines” line 407, and added “(*synclines*)” line 1321 and the hinge line of the folds in Fig. 5.

Comment 20: deleted “fold-related” line 416.

Comment 21: added “(see stereoplot in Fig. 2)” line 506. Deleted “, possibly representing, conjugate sets” lines 505–506.

Comment 22: replaced “a younger slip event” by “younger deformation along this fault” lines 548–549.

Comment 23: deleted “strain “ line 552.

Comment 24: replaced “Palm Spring and Mecca foramtions” by “Mecca Formation and overlying Palm Spring Formation” lines 570–571.

Comment 25: none.

Comment 26: deleted “dip-slip” and replaced “parallel” by “propagation” lines 591–592.

Comment 27: replaced “stress” by “strain” line 673.

Comment 28: replaced “restored” by “rotated” line 497, and “restoring” by “rotating” lines 500 and 686.

Comment 29: replaced “trending” by “oriented” line 744.

Comment 30: deleted “in a changing transpressional plate regime” line 884.

Comment 31: added “The *en echelon* folds formed at a comparable time, i.e., < 0.76 Ma in the Indio Hills and at ca. 0.5 Ma in the Durmid Hills (Table 1).” lines 919–920.

Comment 32: added “in California, southwestern USA” lines 34–35.

Comment 33: none.

Comment 34: changed “Eastern California Shear Zone” into “Eastern California shear zone” throughout the manuscript.

Comment 35: replaced “transform movements” by “movement” line 67.

Comment 36: added “These recent works call for further characterization of the understudied Indio Hills segment in order to compare its structural development with other uplifted features along a major transform plate boundary fault zone.” lines 72–75.

Comment 37: corrected into “San Gorgonio Pass” line 216.

Comment 38: split the sentence into two line 217.

Comment 39: see response to comment 34.

Comment 40: deleted “attitude and” line 226.

Comment 41: changed “remain” into “remains” line 227.

Comment 42: split the sentence into two and replaced “, but” by “However” line 231.

Comment 43: the sentence was deleted and the paragraph reworked according to the other reviewer’s comments.

Comment 44: replaced “their study” by “Keller et al. (1982)” lines 240–241.

Comment 45: added “San Andreas ” and deleted “strand ” line 273.

Comment 46: decapitalized “fault” throughout the manuscript.

Comment 47: added a comma before and after “SAFZ-oblique” line 307 and changed “with” into “having” line 308.

Comment 48: none for the moment. Awaiting comments by the editorial team.

Comment 49: changed “southwestern” into “southeastern” line 379.

Comment 50: replaced “offset” by “displacement” line 429.

Comment 51: replaced “near” by “in the damage zone of” line 446.

Comment 52: added an hyphen between “minor” and “scale” line 501.

Comment 53: deleted “leuco-” line 444.

Comment 54: changed sentence into “In map view (Fig. 2), the folds are right-stepping, and each fold set is increasingly asymmetric (Z-shaped) and sigmoidal towards the Indio Hills fault in the northeast.” lines 523–525.

Comment 55: added “other ” line 570.

Comment 56: deleted en dash line 571.

Comment 57: replaced “slip event” by “episode of movement” line 675.

Comment 58: deleted “precursory” line 753.

Comment 59: replaced “East” by “Eastern” line 810. Also see response to comment 34.

Comment 60: added the Camp Rock and Calico faults to Figure 1.

Comment 61: deleted comma after “enhanced” line 825.

Comment 62: replaced “SAFZ” by “San Andreas fault” lines 853–854.

Comment 63: replaced “in” by “along the” line 860.

Comment 64: added “a–c” line 844.

Comment 65: replaced missing figure reference by “Figs 2 & 3c and Supplement S3a” line 875.

Comment 66: combined “Shore” and “line” line 904.

Comment 67: replaced “Shore” by “Shoreline” lines 920, 928, and 929.

Comment 68: replaced “main SAF” by “main San Andreas fault” line 930.

Comment 69: see response to comment 67.

Comment 70: deleted “in” line 976.

Comment 71: see response to comment 2.

Comment 72: added “; **BSZ: Brawley seismic zone**” line 1285.

Comment 73: see response to comment 34.

Comment 74: uncapitalized “Fault” throughout the manuscript.

Comment 75: see response to comment 3.

Comment 76: adjusted text in figures 2, 7, and 8.

Comment 77: added “(bedding)” twice in figure 2.

Comment 78: none.

Comment 79: added “**via the Orient software (Vollmer, 2015)**” line 1304 and Vollmer (2015) to the reference list.

Comment 80: see response to comment 3.

Comment 81: adjusted the hinge line of structures in Figures 3a–c, 5, and 6.

Comment 82: added “**The yellow dots show the location of field photographs.**” line 1319.

Comment 83: none.

Comment 84: added “, **centimeter-scale**” line 1329.

Comment 85: added “main San Andreas fault” to Figure 5.

Comment 86: re-drew the fold hinge in Figure 6 as it appears in Figure 2.

Comment 87: deleted “Tentative” line 1351.

Comment 88: changed “SAFZ” into “mSAF” in Figure 8.

Additional revisions by the author of the present manuscript

- Added “(we refrain from using the name “Indio strand” given to this fault by Gold et al., 2015 to avoid confusion with the Indio Hills fault)” lines 39–41.
- Moved “Atwater and Stock, 1998;” before “Spotila et al., 2007;” line 64.
- Changed “uppermost members” into “upper member” line 118.
- Replaced “marks the” by “is a” line 201.
- Deleted “-“ line 480.
- Deleted en-dash line 506.

- Added “(probably southwest-dipping)” line 753.
- Replaced “show” by “suggest” line 786.
- Added a comma line 831.
- Corrected “Janecke et al., 2019” into “Janecke et al., 2018” line 842.
- Deleted “and main SAFZ” line 863.
- Moved “basement-seated” from line 899 to line 891.
- Deleted “The Indio Hills fault acted as a SW-dipping, normal fault in Miocene time, i.e., prior to inversion as an oblique-slip, right-lateral-reverse fault during mid (–late?) Pleistocene times” lines 899–901.
- Moved “, whereas the main San Andreas fault initiated probably as a dominantly right-slip fault during the later stages of uplift in the late Pleistocene.” from lines 903–904 to lines 894–896.
- Deleted “portion of the SAFZ” line 912.
- Deleted “in Durmid Hills” line 1235.

Reply to Jonathan Matti

Dear Dr. Matti,

thank you very much for your input on the manuscript, it is highly appreciated. Here is our reply to your comments. We hope the changes we implemented improve the shortcomings of the manuscript highlighted by your comments and suggestions. Please do not hesitate to contact us shall this not be the case for some comments.

1. Comments from Dr. Matti

Manuscript se-2022-9 consists of three parts:

- a detailed structural analysis of macro- and micro-folds and associated faults that deform a sequence of Pliocene-lower Pleistocene sedimentary rocks exposed in the tectonically uplifted Indio Hills;
- a comparison of the Indio Hills structural geology with that of two similar uplifted and inverted late Cenozoic basin fills occurring farther SE within the San Andreas Fault zone (SAFZ);
- integration of the structural data into a synthesis that interprets coeval uplift of the various inverted basins in the context of Quaternary dextral-oblique transpressive tectonics within the southern San Andreas Fault system writ large.

The manuscript explores these three themes with mixed success:

- The discussion of fold and fault structures in the Indio Hills is robust and comprehensive, including appropriate analytical data and exceptional aerial and outcrop photographs that nicely illustrate structural features and relationships. One concern I have is that the structural terminology and technical language used in the manuscript are pitched toward the structural specialist—not toward general geologists like myself. I address this point below.
- The manuscript's comparison of the Indio Hills structural setting with that farther to the southeast within the SAFZ is moderately successful. The report depends heavily on results of other published investigations, and provides only cursory discussion of structural correlations and comparisons among the three inverted basins. The report would benefit from expanded discussion of these correlations, including one or more new map-type figures that better summarize geologic structures SE of the Indio Hills (otherwise, the reader has to chase the

other publications down in order to evaluate manuscript se-2022-9's proposed structural comparisons and correlations).

- By comparison with the preceding two themes, the manuscript's regional synthesis in my opinion is the weakest link in the three themes. In my review I raise some technical questions and issues that I believe need to be addressed more completely—and in some cases explained or corrected. These are not deal-breakers, but should be addressed by the authors.

I do not know whether Copernicus Publications provides an extensive review by a science editor, but I think that manuscript se-2022-9 needs a heavy editorial hand—either by Copernicus staff or by the authors themselves based on peer-review feedback. In part, problems with the narrative structure may stem from the fact that English may not be the first language of two of the three authors. But in addition, I sense that the narrative is too cursory and includes logic jumps that need to be explained more fully. My marginal comments on the manuscript identify many specific instances where I think the narrative can be improved both content-wise and in terms of organization.

All of this said, I enjoyed reading the manuscript. First, it adds to the body of detailed structural analysis so critical to documenting and understanding the geologic history of the southern San Andreas Fault zone and associated depositional basins; and second, it provides a testable regional synthesis for dextral and contractional events within the SAFZ writ large—including possible interactions with the Eastern California Shear Zone and the sequential development of discrete SAFZ strands in the Salton Trough.

My recommendation: The manuscript needs work, but it should be published by Copernicus Solid Earth.

My review consists of two parts:

- General comments contained in this memo
- Detailed comments, questions, and suggested edits integrated into the .pdf version of Manuscript se-2022-9.

NOTE: For my review I separated the manuscript into four discrete documents: (1) the text without references, (2) references alone, (3) figures alone, and (4) supplemental material.

I enjoyed reviewing this manuscript, although I have questions and comments that may (or may not) improve the paper. I trust that the authors will receive my comments and critique in the spirit with which they are offered: to refine and clarify an important contribution our understanding of the tectonic evolution of the southern San Andreas Fault system.

Good luck with forward progress of the manuscript.

Behr, W.M., Rood, D.H., Fletcher, K.E., Guzman, N., Finkel, R., Hanks, T.C., Hudnut, K.W., Kendrick, K.J., Platt, J.P., Sharp, W.D., Weldon, R.J., Yule, J.D., 2010, Uncertainties in slip-rate estimates for the Mission Creek strand of the southern San Andreas Fault at Biskra Palms Oasis, Southern California: *GSA Bulletin* v. 122, no. 9-10, p. 1360-1377.

Cohen, K.M., Finney, S.C., Gibbard, P.L., and Fan, J.-X., 2013 (updated 5/2021), The ICS International Chronostratigraphic Chart, version 2021-5: Episodes, v. 36, p. 199-204, accessed 7/16/21 at <http://www.stratigraphy.org/ICSchart/ChronostratChart2021-05.pdf>.

Dokka, R.K., and Travis, C.J., 1990a, Late Cenozoic strike-slip faulting in the Mojave Desert, California: *Tectonics*, v. 9, p. 311-340, accessed 3/8/2022 at <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/TC009i002p00311>.

Dokka, R.K., and Travis, C.J., 1990b, Role of the eastern California shear zone in accommodating Pacific-North American plate motion: *Geophysical Research Letters*, v. 17, p. 1323-1326, accessed 3/8/2022 at <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/GL017i009p01323>.

Fattaruso, L.A., Cooke, M.L., Dorsey, R.J., Housen, B.A., 2016, Response of deformation patterns to reorganization of the southern San Andreas fault system since ca. 1.5 Ma: *Tectonophysics*, v. 693, Part B, p. 474–488. doi: 10.1016/j.tecto.2016.05.035.

Fuis, G.S., Scheirer, D.S., Langenheim, V.E., and Monica D. Kohler, M.D., 2012, A new perspective on the geometry of the San Andreas Fault in southern California and its relationship to lithospheric structure: *Bulletin of the Seismological Society of America*, v. 102, no. 1, p. 236–251, doi: 10.1785/0120110041.

Fuis, G.S., Bauer, K., Goldman, MR., Ryberg, T., Langenheim, V.E., Scheirer, D.S., Rymer, M.J., Stock, J.M., Hole, J.A., Catchings, R.D., Graves, R.W., and Aagaard, B., 2017, Subsurface geometry of the San Andreas Fault in southern California: Results from the Salton Seismic

Imaging Project (SSIP) and strong ground motion expectations: *Bulletin of the Seismological Society of America*, Vol. 107, No. 4, p. 1642–1662, doi: 10.1785/0120160309

Gibbard, P.L., Head, M.J., Walker, M.J.C., and the Subcommission on Quaternary Stratigraphy, 2010, Formal ratification of the Quaternary System/Period and the Pleistocene Series/Epoch with a base at 2.58 Ma: *Journal of Quaternary Science*, v. 25, no. 2, p. 96–102.
<https://doi.org/10.1002/jqs.1338>.

Gold, P.O., Behr, W.M., Rood, D., Sharp, W.D., Rockwell, T.K., Kendrick, K., and Salin, A., 2015, Holocene geologic slip rate for the Banning strand of the southern San Andreas Fault, southern California: *Journal of Geophysical Research, Solid Earth*, v. 120, 25 p., accessed at <https://doi.org/10.1002/2015JB012004>.

Guns, K.A., Bennett, R.A., Spinler, J.C., and McGill, S.F., 2020, New geodetic constraints on southern San Andreas fault-slip rates, San Geronio Pass, California: *Geosphere*, v. 17, p. 39–68, accessed 3/8/2022 at <https://doi.org/10.1130/GES02239.1>.

Kendrick, K.J., Matti, J.C., and Mahan, S.A., 2015, Late Quaternary slip history of the Mill Creek strand of the San Andreas fault in San Geronio Pass, southern California: The role of a subsidiary left-lateral fault in strand switching: *Geological Society of America Bulletin*, v. 127, p. 825–849, accessed at <https://doi.org/10.1130/B31101.1>

Lancaster, J.T., Hayhurst, C.A., and Bedrossian, T.L., 2012, Preliminary geologic map of Quaternary surficial deposits in southern California: Palm Springs 30' x 60' quadrangle, in Bedrossian, T.L., Roffers, P., Hayhurst, C.A., Lancaster, J.T., and Short, W.R., *Geologic compilation of Quaternary surficial deposits in southern California December 2012*
<https://www.conservation.ca.gov/cgs/fwgp/Pages/sr217.aspx#palm Springs>.

Matti, J.C., and Morton, D.M., 1993, Paleogeographic evolution of the San Andreas fault in southern California: a reconstruction based on a new cross-fault correlation, in Powell, R.E., Weldon, R.J., and Matti, J.C., eds., *The San Andreas fault system: displacement, palinspastic reconstruction, and geologic evolution: Geological Society of America Memoir 178*, p. 107–159.

Matti, J.C., Morton, D.M. and Cox, B.F., 1992, *The San Andreas fault system in the vicinity of the central Transverse Ranges province, southern California: U.S. Geological Survey Open-File Report 92-354*, 40 p., scale 1:250,000. <https://pubs.er.usgs.gov/publication/ofr92354>

Matti, J.C., Kendrick, K.J., Yule, J.D., and Heermance, R.K., 2019, The Mission Creek Fault in the San Geronio Pass region—A long-abandoned strand of the San Andreas Fault, or a major

player in the latest Quaternary San Andreas strain budget? Geological Society of America Cordilleran Section Meeting, Paper 329432

<https://gsa.confex.com/gsa/2019CD/meetingapp.cgi/Paper/329432>

McCaffrey, R., 2005, Block kinematics of the Pacific–North America plate boundary in the southwestern United States from inversion of GPS, seismological, and geologic data: *Journal of Geophysical Research Solid Earth*, v. 110, p. B07401, doi: 10.1029/2004JB003307, accessed 3/8/2022 at <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2004JB003307>.

Nur, A., Hagai, R., and Beroza, G., 1993a, Landers-Mojave earthquake line: a new fault system?: *GSA Today*, v. 3, p. 253, 256-258, accessed 3/9/2022 at <https://www.geosociety.org/gsatoday/archive/3/10/pdf/i1052-5173-3-10-sci.pdf>.

Nur, A., Hagai, R. and Beroza, G. C., 1993b, The Nature of the Landers-Mojave Earthquake Line: *Science*, v. 261, p. 201–203, <https://www.science.org/doi/10.1126/science.261.5118.201>.

Passchier, C.W., and Platt, J.P., 2017, Shear zone junctions: Of zippers and freeways: *Journal of Structural Geology*, v. 95, p. 188-202, accessed 3/9/2022 at <https://doi.org/10.1016/j.jsg.2016.10.010>.

Pillans, B., and Gibbard, P., 2012, The Quaternary Period, in Gradstein, F.M., Ogg, J.G., Schmitz, M.D., and Ogg, G.M., eds., *The Geologic Time Scale*: Elsevier, p. 979–1010, accessed 7/2/2021 at <https://doi.org/10.1016/B978-0-444-59425-9.00030-5>.

Platt, J.P., and Passchier, C.W., 2016, Zipper junctions: A new approach to the intersections of conjugate strike-slip faults: *Geology*, v. 44, p. 795-798, accessed 3/9/2022 at <https://doi.org/10.1130/G38058.1>.

Powell, R.E., Matti, J.C., and Cossette, P.M., 2015, *Geology of the Joshua Tree National Park geodatabase*: U.S. Geological Survey Open-File Report 2015–1175, GIS database, accessed 3/9/2022 at <https://pubs.er.usgs.gov/publication/ofr20151175>.

Rymer, M.J., 2000, Triggered surface slips in the Coachella Valley area associated with the 1992 Joshua Tree and Landers, California, earthquakes: *Bulletin of the Seismological Society of America*, v. 90, p. 832-848, DOI: 10.1785/0119980130, accessed 3/8/2022 at <https://pubs.geoscienceworld.org/ssa/bssa/article/90/4/832/120523/Triggered-Surface-Slips-in-the-Coachella-Valley>.

Spotila, J.A., and Garvue, M.M., 2021, Kinematics and evolution of the southern Eastern California shear zone, based on analysis of fault strike, distribution, activity, roughness, and

secondary deformation: *Tectonics*, v. 40, 32 p., e2021TC006859, accessed 3/9/2022 at <https://doi.org/10.1029/2021TC006859>.

Treiman, J.A., 1992a, Eureka Peak and related faults, San Bernardino and Riverside Counties, California: California Geological Survey [formerly California Division of Mines and Geology] Fault Evaluation Report FER-230, scale 1:24,000, accessed 3/11/2022 at <https://maps.conservation.ca.gov/geologichazards/DataViewer/index.html>.

Treiman, J.A., 1992b, Eureka Peak and Burnt Mountain faults, two "new" faults in Yucca Valley, San Bernardino County, California, in Ebersold, D.B., ed., Landers earthquake of June 28, 1992, San Bernardino County, California: Los Angeles, Association of Engineering Geologists, southern California section, Annual Field Trip Guidebook, p. 19-22.

Walker, M., Head, M.J., Lowe, J., Berkelhammer, M., Björck, S., Cheng, H., Cwynar, L.C., Fisher, D., Gkinis, V., Long, A., Rewi, N., Rasmussen, S.O., and Weiss, H., 2019, Subdividing the Holocene Series/Epoch: formalization of stages/ages and subseries/subepochs, and designation of GSSPs and auxiliary stratotypes: *Journal of Quaternary Science*, v. 34, no. 3, p. 173-186, accessed 7/13/2021 at <https://doi.org/10.1002/jqs.3097>.

Yule, D., Matti, J., Kendrick, K., and Heermance, R., 2019, Evidence for inactivity since ~100 ka on the northern route of the San Andreas fault, southern California: Geological Society of America Cordilleran Section Meeting, Paper 16-8
<https://gsa.confex.com/gsa/2019CD/meetingapp.cgi/Paper/329441>

Matti, J.C., and Yule, J.D., 2020, Does a “structural knot” in the Quaternary San Andreas Fault exist in San Geronio Pass? Traditional and recent views provide conflicting answers: *Geological Society of America Abstracts with Programs*, v. 52, no. 4. doi: 10.1130/abs/2020CD-347317

Comment 1: General comment #1: Who is your audience?—In my opinion it is not clear who manuscript se-2022-9 is trying to reach: the specialist in structural geology? Or the regional geologist who primarily is interested in reconstructing the tectonic history of the SAFZ and related faults over the last 6 ma?

I assert this because the structural analysis of fold and faults in the Indio Hills and their kinematic interpretation (theme 1, above) is laden with specialized structural terms with which the average geologist will not be familiar. This easily can be solved by the author's sensitivity to those

geologists that are interested in the paper but become irritated when the technical language stands in the way of understanding local and regional structures.

This easily can be addressed—not by dumbing down and diluting the structural contributions—but rather by using techniques like the following example:

Instead of “Farther southeast along strike, the Indio Hills and Banning faults merged along a dextral freeway junction (Platt and Passchier, 2016) that may have enhanced....” (manuscript lines 610-611), consider the following:

“Farther southeast along strike, the Indio Hills and Banning faults presumably merge along a dextral freeway junction—a type of fault intersection where the faults have similar shear sense in all three branches (see Platt and Passchier, 2016; Passchier and Platt, 2017). This configuration may have enhanced.....”. (BTW, you may want to add the Passchier and Platt [2017] citation to your list of references).

I recommend you use this type of narrative format to speak both to the structural geologist (probably familiar with the term already) and to the regional geologist like me (inquiring minds want to know).

Specialized structural terms are scattered throughout the manuscript. Here are a few that could be explained:

- fore-limb (of folds)

- back-limb (of folds)

- ladder structure

- shear-folding (as opposed to other fold drivers)

Comment 2: General comment #2: Discussion of faults in the greater Coachella Valley

region—Manuscript se-2022-9 discusses faults of the greater San Andreas system (writ large) in three separate segments of the report: lines 39-47, lines 116-124, and lines 320-336. Not only are these lines scattered throughout various parts of the report (thus making it hard for the reader to keep track of which faults are doing what and when), but the scattered text contain assertions and interpretations that the reader has to remember and appreciate from isolated sections, and then relate within a total picture of tectonic history stretching over 6-7 million years.

This is difficult to do without a well-organized and complete section at the front of the manuscript that summarizes regionally-important faults throughout the greater Coachella Valley region. Absent this introductory summary, the reader reaches no sense of structural complexity

within the SAFZ in southern California—both in terms of discrete faults strands throughout the region and how they evolved through time and space.

Why is a coherent introductory regional statement needed?

The manuscript ostensibly focuses on structural relations from the latitude of the southern Indio Hills south. However, the report integrates certain regional faults, concepts, and nomenclature not only into its concluding tectonic synthesis but also into its use of fault names locally. This especially is apparent with how the authors use the name “Banning Fault” (General Comment 3 below) and with how they integrate late Quaternary strain history in the Indio Hills with modern strain patterns in the Eastern California Shear Zone (General Comment 5 below).

Absent a coherent introductory summary of regional fault relations, the reader can’t help but believe that the manuscript’s findings in the southern Indio Hills (and similar domains to the southeast) resolve all issues related to strain distribution in the southern SAFZ throughout the last 6-7 Ma.

Recommendation:

To address this, I recommend that all discussion of regional faults be moved into a single section under “Geologic Setting”, following an outline like this (or something like it):

Geologic Setting

Regional faults (including what is known about fault ages; see figure 2 in Kendrick and others, 2015)

Regional stratigraphy (already discussed in the manuscript)

Regional Tectonic Culminations (already discussed in the manuscript)

This new section hopefully will incorporate (and resolve) issues and questions identified in General Comments 3, 4, and 5 (below).

Comment 3: General comment #3: Use of the term “Banning Fault”—The manuscript applies this fault name from the San Gorgonio Pass region southeast beyond the southern Indio Hills (see figures 1 and [especially] 2, and lines 39-42, 633-634). This runs counter to the way most workers interpret faults and fault names.

The problem: Because the manuscript lacks a coherent discussion of fault nomenclature, distribution, and movement history in the greater Coachella Valley region, the reader reaches no sense of structural complexity within the southern California SAFZ—both in terms of discrete faults strands throughout the Coachella Valley region, how they evolved through time and space,

and how they interacted together. Although the manuscript ostensibly focuses on structural relations from the latitude of the southern Indio Hills south, it nevertheless brings certain concepts (and attendant nomenclature) southward from the northern Coachella Valley where structural relations are more complex than implied in the manuscript. The reader can't help but believe that the findings in manuscript se-2022-9 resolve all remaining issues related to strain distribution in the southern SAFZ throughout the last 6-7 Ma.

Manuscript se-2022-9's use of the term "Banning Fault" inadvertently (but unfortunately) contributes to this problem

Recommendation:

If manuscript se-2022-9 retains its current nomenclatural approach regarding the Banning Fault, at a minimum the report needs to address how its usage differs from that of other workers (discussed below). It would be better if the authors evaluated regional tectonic implications of extending the name "Banning Fault" as far south as they do—especially because northwest of their study area the fault has been shown to have a very limited time during which it functioned as a discrete strand of the SAFZ, and this time frame is incompatible with that the authors propose for the "Banning Fault" in the southern Indio Hills.

In short: Manuscript se-2022-9 needs to acknowledge that the timing they propose for movement on the "Banning Fault"—and the important structural role it plays in their analysis of how the tectonic culmination evolved—is not compatible with what is known about slip on the Banning strand of the SAF in the northern Coachella Valley.

Nomenclatural and fault-reconstruction precedents:

Matti and others (1992; Matti and Morton, 1993) have addressed nomenclature problems for strands of the SAFZ in the Coachella Valley region. They applied the name "Banning Fault" southward from eastern San Geronimo Pass to the fault's junction with the Mission Creek Fault midway along the Indio Hills. In addition, they alluded to "Coachella Valley segments" of the various faults, anticipating future nomenclatural refinements that have emerged over the last decade or so.

Recent investigators follow this precedent regarding the spatial extent of the "Banning Fault". Behr and others (2010), Fuis and others (2012, 2017), Fattaruso and others (2014, 2016), Gold and others (2015), Kendrick and others (2015), Beyer and others (2018), and most other workers do not apply the term "Banning Fault" southeast of its junction with the Mission Creek strand.

Beyond that juncture, some workers apply the name “Indio strand” to the SAF (Behr and others, 2010; Gold and others, 2015; see fig. 1 of both reports). Other workers apply the name “Coachella Valley segment of the SAF” to the fault southeast of the juncture (see Fattaruso and others, 2014, 2016).

You are not obliged to use the more common usage of “Banning Fault”. However, it is incumbent on you to address the nomenclature issue.

If you choose to revise your nomenclatural approach, then you need to come up with a name that you can apply to the “San Andreas Fault” strand southeast of the Banning/Mission Creek junction. Given that—in the Indio Hills region—the Mission Creek Fault was the major SAF strand during the period 4 Ma to Holocene (the critical period during which you require a bounding dextral fault on the SW side of the Indio Hills culmination), I personally would apply the name “Mission Creek Fault” in place of your universal application of the name “Banning Fault” to this structure.

Finally, almost all workers agree that the Banning strand of the SAF between San Geronio Pass and its juncture with the Mission Creek strand evolved during the late Quaternary time (last 200 ka???) as the result of a left step from the Mission Creek to a newly evolving strand to the west (i.e., the Banning Fault) (Matti and others, 1985; 1992; Matti and Morton, 1993) (especially see the important paper by Gold and others, 2015, that explores exhumation and uplift rates in the northwestern Indio Hills—a story that sounds a bit like your own, only younger?).

Comment 4: General comment #4: Indio Hills Fault—Manuscript se-2022-9 proposes that the Indio Hills Fault plays a significant structural role in both (1) evolution of the local tectonic culmination and (2) the evolution of the SAFZ. The authors identify three phases of movement history for the Indio Fault:

- An initial role as a southwest-dipping normal fault (late Miocene);
- An intermediate role as a dextral strike-slip fault;
- A penultimate and current role as an active transpressive dextral-oblique thrust fault.

I found it difficult to understand its polyphase role (normal fault followed by dextral-slip fault followed by oblique reverse-thrust fault)—especially the timing of activity during each tectonic phase. Comments and observations and interpretations about this structure are scattered throughout the manuscript, so it was hard for me to keep these three phases in mind and to appreciate when each was active.

New to my awareness is the manuscript's proposal that the Indio Hills Fault initially was a late Miocene normal-slip fault (1, above). This assertion needs to be supported with evidence. The authors at some places in the report point to previous workers who propose a southwest-dipping "basin-and-range" type of structure that had to exist early in the evolution of the San Andreas Fault system in the Salton Trough, but I am not aware that the Indio Hills structure was part of that "basin-and-range"-type system. Please explain and elaborate, including where this regional "basin-and-range"-type system can be recognized NW of the Indio Hills

Comment 5: General comment #5: Landers-Mojave Line connection with Indio Hills

Fault—I have problems with how the manuscript projects the Indio Hills Fault into a seismic trend that Nur and others (1993a, b) identified as the "Landers-Mojave Line". That "concept" was defined to represent a seismicity belt that was observed following the 1992 Landers earthquake in the Mojave Desert (note that a recent paper by Spotila and Garvue (2021) challenge some of the assertions by Nur and others, 1993a, b). Manuscript se-2022-9 asserts that the Indio Hills Fault can be connected structurally with the Landers-Mojave Line via faults in the Little San Bernardino Mountains (LSBM).

My concerns include:

- The manuscript (fig. 1) connects the Indio Hills Fault northwestward to a presumed fault at the south edge of the LSBM. Although this fault is depicted in some publications, its distribution, structural role, and age have not been documented. Therefore any reference to this fault in manuscript se-2022-9 needs to acknowledge this reality.

I recommend that you cite the recent digital geologic-map database of Joshua Tree National Park by Powell and others (2015) for a more recent and detailed rendering of geologic units and faults. The report can be viewed only in a GIS (ArcMap, for example), but once loaded into a GIS platform the files reveal much more about JTNP geology than was known previously.

- The unnamed fault is depicted by Rymer (2000, fig. 1) who plots it east of his West Deception Canyon Fault. Although Rymer (his figs. 1 and 2) shows the epicenter of the 1992 Joshua Tree earthquake located a few km north of the unnamed fault, he did not report any ground rupture on it. Instead, Rymer documented ground rupture on the West Wide Canyon Fault (see his figure 2). This structure is well to the NW of where manuscript se-2022-9 speculates that the "active" Indio Hills Fault would intersect the LSBM and connect with the "Landers-Mojave Line".

- In lines 600-603 the authors reference Dokka and Travis (1993a, b) in support of the hypothesis that the Indio Hills Fault [strand of the San Andreas Fault] “connects” with the Eastern California Shear zone (ECSZ) and the “Landers-Mojave Line of Nur and others (1993). Reference to Dokka and Travis as supportive evidence for this hypothesis is not appropriate because those authors (1990a, b) do not show the ECSZ extending south of the left-lateral Pinto Mountain Fault (see figs. 2, 14, 15, and 18 of Dokka and Travis, 1990a, and figs. 2 and 3 of Dokka and Travis, 1990b). In fact, Dokka and Travis, 1990b, p. 1325 clearly explore the notion that connection between the ECSZ and the San Andreas Fault (in this case, the Indio Hills strand) is based on slip budgets for the North American plate margin—the physical and kinematic basis for this connection is not obvious.
- Occurrence of (a) ground rupture triggered by the 1992 Joshua Tree earthquake and (b) ground rupture on the Eureka Peak fault south of the Pinto Mountain Fault during the 1992 Landers earthquake (Treiman, 1992a, b) is tempting evidence that strain probably is transferred kinematically between the southern San Andreas Fault and the ECSZ. Note, however, that ground rupture associated with the Joshua Tree event was not coextensive with the Eureka Peak Fault. Thus, it is unlikely that transfer of strain between the ECSZ and the SAFZ occurs along a single fault trace (the authors don’t claim that it does, but their figure 1 implies as much. Better to clarify).
- The California Geological Survey classifies the Indio Hills Fault as a late Pleistocene and older feature, with no evidence for Holocene displacement. For current interpretation of fault activity, see California’s interactive geologic-hazards map at <https://maps.conservation.ca.gov/geologichazards/DataViewer/index.html> and also the Quaternary Fault and Fold Database at <https://www.usgs.gov/programs/earthquake-hazards/faults>). These data call into question the notion that the Indio Hills Fault is a Holocene extension of the Holocene and Recent ECSZ and Nur’s Landers-Mojave Line.

The authors probably will protest that the scope and purpose of manuscript se-2022-9 is much broader and regional in scope to address details of the kind that I provide here in General Comment 5. I agree. I provide my analysis mainly to remind the authors that any model that incorporates latest Quaternary activity on the NE-bounding structure of their Indio Hills tectonic culmination (in this case, the Indio Hills Fault) needs to be compatible with what is known about the distribution and geologic history of fault elements that might (or might not) connect their

tectonic model for the SAFZ with the ECSZ—or with SAFZ structures northwest of the Indio Hills culmination.

Comment 6: General comment #6: Ages for fault activity—In general, I found it quite difficult to determine the sequencing and ages for faults the manuscript discusses and integrates into their concluding time-space model. This is very frustrating because the timing of fault movements (1) relative to overall history of SAF history in the greater Coachella Valley region and (2) relative to when and how the Indio Hills tectonic culmination evolved is a critical part of the author's tectonic model.

Recommendations:

- Develop a new section called “Summary of fault ages”, and consolidate all the disparate observations about age of faulting currently scattered throughout the report.
- I recommend developing a new diagram like figure 2 of Kendrick and others (2015).
- Make certain that the manuscript's use of “Pliocene”, “Pleistocene”, and “Holocene” conform to current international standards (see Pillans and Gibbard, 2012; Cohen and others, 2013; Gibbard and others, 2013; Walker and others, 2019; and other references on this subject). The boundary between Pliocene and Pleistocene now is ~ 2.6 Ma.

Comment 7: General comment #7: “Possibly”, “or”, and “may have”—The manuscript frequently has sentences like the following:

“Structural feature X formed by process Y, and (or) it [possibly, may have] formed by process Z”.

As a reader, I asked myself “are the authors not committing themselves to structural feature X formed by process Y—their first choice”? Adding caveats like “or may have” makes sentences containing this kind of grammatical structure [presentation] sound like the authors are not sure about their assertions, and are covering themselves.

Recommendation:

- Examine the narrative, find those kind of grammatical instances, and design a more appropriate way of expressing the level of confidence the authors have in conclusions X and Y—in other words, include [discuss] the error bars that prevent complete confidence.

In short: the authors need to choose more definitively among the suite of interpretive possibilities, and not just cover their hypotheses with “or alternatively it could be a different way”! As Gozer challenged the Ghostbusters: “Choose the form of the Destructor”.

Comment 8: General comment #8: Identity, position, and age of “SW master bounding

fault”—It may just be me, but I had trouble understanding what the SW-bounding master fault was throughout the evolution of the Indio Hills uplift, where was it positioned throughout this evolution, when did the uplift start, how long did it last, and is it still active? Comments here not only are relevant to the narrative but also to figures—especially figure 7.

Regarding the what: In general comment #3 I questioned your application of the name “Banning Fault” to the SAF strand on the SW flank of the Indio Hills uplift. My comment there pointed out that (according to current understanding) the Banning strand of the SAF in the northwestern Coachella Valley became active only in the last few hundred thousand years (late Pleistocene and slightly older). That “fact” calls into question whether the “Banning Fault” could have been the SW-bounding “master fault” during (say) 500 ka? 1.0 Ma? 1.5 Ma? 2.0 Ma? So, together with my concerns in Comment #3 I question whether you should use the term “Banning Fault” for whatever SAF strand may have formed the “master fault” bounding the SW side of the long-evolving Indio Hills uplift.

But if current thinking regarding the age of the Banning strand of the SAF in the northwestern Coachella Valley is correct, then during the last few hundred thousand years that strand was feeding slip SE along the Pacific margin of the Indio Hills uplift, so at that time application of the name “Banning” to that SW-bounding master fault may have been appropriate (as implied in fig. 7c) (but only during that slip episode).

Bottom line: Multiple SAF strands northwest of the Indio Hills have been active throughout the last 6-8 Ma, those strands have evolved sequentially (Kendrick and others, 2015, fig. 2), each of those strands has a different name, and each of those strands sequentially has fed dextral slip southeast toward the Indio Hills uplift. So the SAF bounding the SW margin of the Indio Hills has had a “changing name” throughout the total 6-8 Ma of southern SAF evolution in the Coachella Valley.

This is why application by Behr and others (2010) and Gold and others (2015) of the term “Indio strand” or “segment” to the SAF southeast of the junction between the Mission Creek and Banning Faults is so appealing: throughout time, all the messy SAF strands NW of the Indio Hills have sequentially evolved northwestward of the Indio strand—presumably NW of the current junction between the Mission Creek and Banning strands of the SAF.

Regarding the where: Your figures 7a and 7b position a queried “Banning Fault” west of the trace of the “Banning” shown in fig. 7c. Why do you do this? What is the basis for the location difference?

Regarding the when and for how long: In my General Comment #6 I recommended a new section that consolidates all fault-age information currently scattered throughout the report—or not addressed clearly. I also recommended a new figure like figure 2 of Kendrick and others (2015). Such a figure easily could add a “range-bar” for the Indio Hills tectonic culmination, thereby resolving my questions about the when and how long.

Regarding the still active?: In my General Comment #5 I questioned your correlation of the Indio Hills Fault with the “Landers-Mojave Line” of Nur and others (1993a, b). Depending on how you address my comments there, the Indio Hills fault may (or may not) still be active—and the tectonic culmination may (or may not) still be actively growing.

Relevant to the question of “is it still growing”—I can’t remember whether your manuscript discusses the evidence for reverse-dextral slip on the SW-bounding SAF strand (whatever its name). Do you have fault-plane evidence or other evidence that the SW-bounding fault has generated up-on-the-NE displacement (other than the fact that the landscape is higher to the NE than the SW)? Is it possible that the Indio Hills uplift tilted SW away from a high landscape adjacent to the Indio Hills Fault toward a low landscape to the SW? In other words: has uplift on both NE- and SW-bounding master faults been equal? I think this is an important question to address.

Recommendation:

- The manuscript needs to expand and clarify questions about the what, where, and when aspects of tectonic culmination of the Indio Hills. I recommend a new section, or at least addressing these questions in a single part of the narrative.

Comment 9: General comment #9: Character of folds as they approach the Banning Fault—

In lines 336 and 508 (and elsewhere) you discuss the geometry of folds as they “approach” the Banning Fault. But these folds presumably are older than a few hundred thousand years, and their axes never reach the position of the “Banning Fault” as you depict it in figures 7a and 7b (and folds closest to the “modern” Banning Fault” in figure 7c area fault-parallel and are not relevant to folds that are oblique to the master faults. Therefore, for the latter, how can you comment on the structural style, morphology, and configuration of the fold sets depicted in figs 7a and 7b?

They do not reach the queried and discontinuous “Banning Fault” trace that you show more valleyward in figures 7a and 7b. Please clarify.

Comment 10: **General comment #10: Update references**—Lines 140-150 need to cite Gold and others (2015), and do a more thorough job of describing what Keller and others (1982). At the end of this memo I include many references that you should at least consider for inclusion and evaluation for your report.

Comment 11: Lines 37–38: You absolutely need to have a section that discusses more thoroughly the distribution and nomenclature of major faults in the greater Coachella Valley/Mojave Desert/Peninsular Ranges region. Your readers for the most part are going to need such a regional summary----and the few lines here don't do the job.

Comment 12: Line 39: Careful with this statement. In the literature, the name "Banning Fault" typically is applied from San Geronimo Pass SE to its junction with the Mission Creek Fault. The literature is mixed about what name to apply to the singular dextral fault SE of this juncture various names are applied to the singular fault, but no author (to my knowledge) uses the name "Banning Fault" here

Comment 13: Lines 39–40: No. I know of no investigator who applies the name "Banning Fault" to a structure in the Mecca Hills and Durmid Hills. Cite source (other than Janecke----see my next comment.

Comment 14: Line 40: No, Janecke (2018) uses the name "San Andreas Fault main trace". She does not apply the name "Banning Fault" to any structure in the Durmid Hills. In a couple of places she refers to "the Banning fault, the most active strand of the mSAF nearby"----by which she means that "nearby to the Durmid Hills the Banning Fault is the main SAF trace." In no way was she extending the name "Banning Fault" to the Durmid Hills.

Comment 15: Line 41: No. Allen (1957, fig. 1) does not apply any name to the SAF southeast of its juncture with the Mission Creek fault.

Comment 16: Line 42: This statement is a hypothesis, not an axiom. I comment on this later in the manuscript. Also, careful with the names, as my comments on lines 38-42 indicate.

Comment 17: Lines 46–47: You really need to refer to this as a POSSIBILITY----not as a proved and documented fact.

Comment 18: Lines 56–58: Actually, this is not accurate. The "Coachella Valley segment of the SAFZ" is "expressed by multiple dextral fault strands of the zone----the uplifts themselves (and

their inverted sedimentary fills) are a product of the dextral-oblique transpressional structural evolution you propose in this report. Modify.

Comment 19: Line 71: Palm Spring Formation (singular "Spring")

Comment 20: Lines 79–80: Useful information. However, given that the Indio Hills Fault intervenes between the leuco-granitic rocks and the majority of the Indio Hills stratigraphy, a comment here is needed to the effect that you interpret the leucocratic rocks as "basement" for the inverted Indio Hills basin. Or not, eh?

Comment 21: Line 84: Are the granitic clasts similar to the "disconnected" leuco-granites?

Comment 22: Lines 93–100: The narrative in lines 93-100 is a little hard to follow, and the implications are not completely clear. I recommend *first* discussing the stratigraphy and lithology and member relations of the PS Formation in the Indio Hills, then compare and contrast with the PS Formation in the Mecca Hills. Intermingling observations in the two areas makes it difficult for the reader to follow.

Comment 23: Line 98: In contrast, in the Indio Hills the nature.....

Comment 24: Lines 94–112: Upon reading lines 94-112, I see that they incorporate two subjects: (1) the physical stratigraphy of the Palm Spring Formation in the Indio Hills in comparison with that in the Mecca Hills, and (2) evidence for the age of the PS Formation in the Mecca Hills----with implications for the formation's age in the Indio Hills. Given the importance of linkage between stratigraphy, unconformities, and unit ages, I would rewrite lines 89-112 to more clearly separate (1) stratigraphy and unconformities (or not) in the two areas, and (2) age relations for the PS Formation based on Mecca Hills data with extrapolation to Indio Hills. The narrative should emphasize Indio Hills data, with contrasts and similarities with Mecca Hills.

Comment 25: Lines 104–106: This applies to the Mecca Hills, not your study area----another example of intermixing content about the two areas. Moreover, these are interpretive statements that follow from the geochronologic data----not indicators of age (see my previous two comments). I would place this at the end of the paragraph as a conclusion derived from lines 93-112.

Comment 26: Line 107: In the Mecca Hills only? Or also in the Indio Hills? Add this for specificity, OK?

Comment 27: Line 114: After I read through this section, in my opinion the heading should be "Tectonic culminations" similarities and differences regionally"

Comment 28: Lines 119–120: To be fair, Matti and others (1985, 1992; Matti and Morton, 1993) proposed the left step at San Gorgonio Pass, Dair and Cooke (2009) simply used the USGS geology. See references in comment for line 128

Comment 29: Line 120: You need to define what the term "proto-SAFZ" means. You have not used it previously in the manuscript, and its introduction out of the blue catches the reader by surprise. Also, if you are referring to the old-school use of "proto-SAF" as an "early SAF", that term largely has fallen out of use. Please provide meaning and literature source of usage.

Comment 30: Line 121: Meaning of "low-topographic relief SAFZ segment" is unclear. What segment?

Comment 31: Line 121: What? are these

Comment 32: Line 122: uplifted San Bernardino "Mountains" (yes?)

Comment 33: Line 120–124: Lines 120-124 apparently are intended to provide regional structural relations surrounding the Indio Hills. As such, the narrative both (1) has some unclear structural statements and (2) some errors. First, "proto-SAF" needs to be explained. Second, the "low-relief SAFZ segment" is completely unclear, and is not identified on figures 1 and 2. Third, the left-slip faults are not identified: are these the sinistral faults of the eastern Transverse Ranges? If so, cite Powell (1993) as a useful reference for their description and interpretation. Fourth, I don't understand how the "left-slip splay faults" (unspecified) merge into the San Jacinto Fault strands. Fifth, inclusion of the West Salton Detachment in the same single sentence that begins with "north of the Coachella Valley" implies that the WSD is north of the study area and not southwest.

Comment 34: Line 128: Matti and others (1992) is more accessible to readers. Also see Matti and Morton, 1993).

Matti, J.C., Morton, D.M. and Cox, B.F., 1992, The San Andreas fault system in the vicinity of the central Transverse Ranges province, southern California: U.S. Geological Survey Open-File Report 92-354, 40 p., scale 1:250,000. <https://pubs.er.usgs.gov/publication/ofr92354>

Matti, J.C., and Morton, D.M., 1993, Paleogeographic evolution of the San Andreas fault in southern California: a reconstruction based on a new cross-fault correlation, in Powell, R.E., Weldon, R.J., and Matti, J.C., eds., The San Andreas fault system: displacement, palinspastic reconstruction, and geologic evolution: Geological Society of America Memoir 178, p. 107-159.

Comment 35: Line 129: Rarely is the gouge actually exposed where the fault "line" traverses the desert. Better to use "trace" in place of "main fault gouge"

Comment 36: Line 131: There is something wrong with this reference. In your references cited you identify Parrish (1983) as:

"Parrish, J.G., Geological compilation of Quaternary surficial deposits in southern California, Palm Springs 30' x 60' quadrangle, CGS Special Report 217, Plate 24, 1983"

I don't know what the 1983 report is that you actually want to cite (other than the Palm Springs sheet) or where you got "1983" from, but the "J.G. Parrish" that you cite actually was the California State Geologist when the Palm Springs 30' x 60' sheet (surficial geology) was published by CGS in 2012. The CGS Special report 217 that you refer to is the umbrella that covers all of the individual 30' x 60' surficial maps, including the Palm Springs sheet.

I provide a correct bibliographic citation for the PS sheet in your references cited section, and in my general comments memo.

Comment 37: Line 131: Assuming you want to refer to the 2012 Palm Springs sheet (yes?), throughout the manuscript you should search and replace "Parrish, 1983" with "Lancaster et al, 2012".

Comment 38: Line 137: "Southeast" of what? The LSBM? Your study area? clarify

Comment 39: Line 143: Need to cite Gold and others (2015) and summarize what they contributed to uplift of the NW Indio Hills.

Comment 40: Line 145: As I recall, Blisniuk et al 2021 explore local uplift as a driver for channel avulsion, but I don't see a discussion of regional uplift as a driver for the Indio Hills as a "tectonic culmination". Eliminate reference, or clarify.

Comment 41: Line 146: Antecedent unclear. Refers to Keller and others, 1982? See my proposed edit.

Comment 42: Lines 159–160: How does your "steep SAF" compare to the seismic-profiling implications of a moderately east-dipping SAF (Fuis and others that you cite in your regional studies section)

Comment 43: Line 171: Is the "elongate ridge" aligned with the Brawley Seismic Zone, or is the SAF itself? Moreover, is "aligned" the correct term, or should it be "projects toward"?

Comment 44: Line 176: It is worth citing Janecke et al 2018 again because their study is so detailed and exhaustive, and is the most recent?

Comment 45: Lines 194–195: Is this based on field investigation or imagery analysis>

Comment 46: Line 234: "When" is a time term.

Comment 47: Lines 239: It is a little unclear why you use the phrase "SAFZ-parallel" anticline here instead of "Indio Hills-parallel anticline". Explain

Comment 48: Line 242: Antecedent unclear. What does the plural pronoun "they" refer to previously in the sentence?

Comment 49: Lines 253–254: Does this imply detachment between strongly and weakly folded sedimentary materials?

Comment 50: Line 285: Not needed, unless you specify applicable ambiguities

Comment 51: Line 294: grammar unclear (do you mean subparallel?)

Comment 52: Line 303: Antecedent unclear: what does pronoun "its" refer back to? If the "couple of major folds", then pronoun needs to be "their".

Comment 53: Line 310: Are these "brittle faults" the same as the "meter-wide" fold-related faults? Or a different category? The adjective "brittle" at the beginning of this sentence is puzzling

Comment 54: Line 322: If this is your criterion for fault dip (reverse, thrust), then in figure 2 why don't you extend the thrust barbs NW along the entire Indio Hills?

Comment 55: Line 325: You don't need this, as nearly all workers view the two faults as merging to the SE.

Comment 56: Line 328: Do you know whether these features formed due to movement on the Indio Hills Fault versus some older unrelated deformation that affected crystalline rocks on this part of the greater Little San Bernardino Mountain front?

Comment 57: Line 331: Have you actually observed the fault plane of the Banning? If not, I would begin the paragraph by stating "Like the Indio Hills Fault, fault-plane dip and strike of the Banning Fault must be inferred indirectly".

Comment 58: Lines 331–332: Couldn't this truncation be produced by up on the east dip-slip displacement on the Banning?

Comment 59: Line 337: As with your section describing folds, I would use a different heading format for this fault section, specifically:

Faults in folded sedimentary strata:

Minor strike-slip faults

Reverse faults

Comment 60: Line 338: If you are going to use NW-SE previously in this sentence I would use NE here.

Comment 61: Line 340: By definition, a mudstone consists of silt and clay-sized grains. So if you observed both "mudstones" and "siltstones", list both of them as separate discrete lithologies

Comment 62: Line 378: But I thought this section dealt with reverse faults?

Comment 63: Line 386: See my comment above questioning correlation of these structures with late Cenozoic SAF displacements

Comment 64: Lines 396–397: This sentence is difficult to wade through. See my edit for whether it changes your meaning.

Comment 65: Line 400: Show "right-lateral" only once in this sentence, either several words earlier or here, but not twice?

Comment 66: Line 400: Can you provide evidence as to why this is "simple shear" (i.e., it has to be because.....).

Comment 67: Line 408: This has to be the Banning and Indio Hills faults, yes?

Comment 68: Lines 412–414: There is a logic jump here. If I understand your model, the folds tighten and become more complex toward the Indio Hills Fault. Thus, would it not be the fold structural geometry (rather than the deposition of the Palm Spring Formation) that would support "folds propagating outward from the Indio Hills Fault"? The formation deposition (all of the PSF) determines the age of folding, but does it determine fold- propagation style?

Comment 69: Line 418–419: How about structural tilting to the SW in the "hangingwall" of the Indio Hills fault?

Comment 70: Line 424: What is this? Reference to such a thing in the literature? Or is it simply a generic term for convergence in a fault zone?

Comment 71: Line 427: I recommend a paragraph break here.

Comment 72: Line 430: I recommend that you introduce figure 7 at this early point so that readers can better follow your structural logic.

Comment 73: Line 450: label this fold on figure 3C

Comment 74: Line 461: Yes, but doesn't he focus on extensional folds, rather than contractional folds as here in the Indio Hills?

Comment 75: Lines 472–473: Not clear to me how the fault-parallel anticline under discussion could "act" like a SW-dipping thrust fault. Either the structure was a fault or it was not. I would allow the fold's vergence to express its tectonic-transport role.

Comment 76: Line 488: I think you need an introductory paragraph here that sets up the important interpretive synthesis of the section that follows. Line 488 is not a very helpful topic sentence for either its paragraph----you just plunge into assertions about the history without providing any leadup or context. I recommend a paragraph that states something like:

"In this section we use the geometry and kinematics of folds and faults in the southern Indio Hills to reconstruct the story these structures tell us about the geologic and tectonic history not only of the inverted late Cenozoic basin but also about strike-slip and dip-slip faults that bound the basin. Essential elements of this story are (1), (2), (3), and (4)----etc."

Comment 77: Lines 488–489: See comment at the end of line 493

Comment 78: Line 490: As best I can tell, figure 3Sb does not illustrate "normal faults".

Comment 79: Line 491: See my comment on S6

Comment 80: Line 493: It probably is just me, but I don't think you have made a convincing case for the Indio Hills fault originally being a normal fault. If other reviewers agree, then I think you need to do a better job documenting the "normal-fault" interpretation.

Comment 81: Line 497: I may have touched on this in an earlier comment, but I think that you need to build a more thorough case for a "precursory Indio Hills fault". Simply asserting its existence, as you do here, without reminding us (or informing us) that an early history of the Indio Hills fault is required by your structural analysis leaves me asking the question "what is this precursory episode, and why do these guys propose it? What did they tell me earlier in the manuscript?"

Comment 82: Line 497: It only "gradually changed" if you make a better case for a normal-fault origin for the Indio Hills Fault prior to "changing" structural style.

Comment 83: Lines 503–504: Margin of what? The tectonic culmination of the Indio Hills block? The original depositional basin? Clarify

Comment 84: Line 505: Do you mean: "while during this period the Banning Fault was generating significant dextral slip"???? Clarify

Comment 85: Line 506: "Slip event" generally is viewed as an earthquake event in a moment of time. Better here to use ""recent slip style"

Comment 86: Line 509: This must be "off-fault deformation", yes?

Comment 87: Line 510: Why "early"? Wouldn't off-fault deformation (your "distributed stress") have continued throughout the entire history of dextral slip on the Banning Fault? (both early stage and late stage?)

Comment 88: Line 511: "post-dating slip on the Indio Hills Fault", yes? Otherwise, late-stage has no contextual meaning. "Late" relative to what? Geologic time? (i.e., recency in terms of the geologic time scale).

Comment 89: Line 513: In line 511 you already indicated the footprint of fold-fault interaction.

Comment 90: Lines 514–515: Why is the occurrence of "minor fault-related folds" on steeply north-dipping beds important? How about moderately-dipping or shallowly-dipping beds? Do minor folds not occur on these strata? Or if they do, are those occurrences irrelevant to your story? The caveat of "steeply dipping" catches my eye and causes me to stumble.

Comment 91: Line 516: Supplement S6 doesn't appear to address restoration of "steep dips" to "horizontal" orientation. I must be missing something.

Comment 92: Lines 516–521: The giant run-on sentence in lines 516-521 needs to be reshaped into more than one sentence.

Comment 93: Line 522–523: This is confusing: line 517 indicates that you cannot discriminate whether "minor folds and faults" "pre-date (or were coeval with) the macro-folding event. But here in lines 522-523 you indicate that "minor right-slip faults evolved synchronously" with the en echelon fold limbs. Clarify. In line 517 maybe you mean "all minor faults other than right-slip faults"????

Comment 94: Line 531: Line 529 begins with "out-of-the syncline contractional faults, yet here you indicate a specific "anticline". Text is hard to follow.

Also, as indicated in a comment elsewhere, label "the related anticline" in figure 3c.

Comment 95: Line 537: What are these? You have not discussed these elsewhere in the manuscript, and figure 4 caption does not refer to them.

Comment 96: Line 539: You absolutely have to choose between these two different scenarios---- state what you conclude, not just what two contrasting choices exist without telling us what you favor.

Comment 97: Line 546: This section needs an introductory paragraph indicating what you will do with the section. I recommend something like:

"In this section we will use detailed structural analysis of folds and faults in the southern Indio Hills to (1) outline the structural history of the tectonic culmination itself, (2) compare and contrast this structural evolution with that of nearby culmination (Mecca Hills, Durmid Hills), and (3) integrate these local structural histories into a structural synthesis for the southern San Andreas Fault zone from X my to Y my. Finally, we evaluate this proposed structural synthesis in terms of what is known about strain budgets within the southern San Andreas Fault system writ large."

Something like this is needed in order to allow the reader to follow how you stitch together all the disparate structural details you have documented so far in the report.

It really would help if you began this section with a bullet outline---or even better, a table---outlining the sequence of depositional/tectonic events together with their age---especially for the complicated and sequential structural evolution of the Indio Hills Fault.

Comment 98: Line 549: I don't understand this phrase: If "inversion" is defined to be destruction and tectonic uplift of the depositional basin---and if the basinal episode involved accumulation of the Miocene Mecca Formation and the Pliocene-Pleistocene Palm Spring Formation---then how could "inversion" occur in Miocene time as line 549 asserts?

Comment 99: Lines 549–550: See my earlier comments about the Indio Hills fault having an early history of dip-slip displacement.

Comment 100: Line 552: This statement is puzzling: **First**, If the granitic basement is exposed only on the east side of the Indio Hills Fault (your figure 2), **second**, if that fault intervenes between the granitic rocks and the Indio Hills and the inverted sedimentary basin of the southern Indio Hills, and **third** if the Indio Hills Fault has a significant normal-slip, dextral-slip, and thrust-slip history, then how and why do you include erosion of the isolated granitic rocks and overlap by the Miocene Mecca Formation in your discussion of the structural history of the tectonic culmination? The depositional overlap you refer to occurred **beyond** (outside) the depositional basin that subsequently was "inverted", yes? Clarify

Comment 101: Line 552: Did you present evidence for this statement earlier in the manuscript? I don't remember reading this. You need to present evidence for this age range.

Comment 102: Line 558: Which strand of the SAF? The Indio Hills Fault? The nascent Banning Fault? Clarify

Comment 103: Line 559: Why do you say "probably"? Either the fold axes trended at a high angle to the bounding faults or they didn't. Fix.

Comment 104: Line 559: You refer to "bounding master faults". But you are discussing ONLY the Indio Hills Fault. What is the *other* "bounding fault"???? In figure 7 you do not identify a coeval bounding fault (the Banning Fault is questionably identified, but in line 511 you identify the Banning as a "late stage" structure. You need to clarify this issue of "bounding faults" (plural).

Comment 105: Line 560: See my comment on line 400

Comment 106: Line 562: Precursory to what? Precursory to slip on the Indio Hills Fault? Clarify.

Comment 107: Lines 564–565: You can't just identify two very different structural styles, without favoring one or the other. Choose one style and discount the other, then explain and justify your choices.

Comment 108: Line 566: Again: what is your choice? There is a significant difference between minor-fold deformation "prior to or together with" macro-fold development, eh?

Comment 109: Line 567: But your statement in line 566 indicates that you don't know when the minor folds formed, hence how can you use these structures to assign "minor partitioning"? And I presume you mean "strain partitioning"?

Comment 110: Line 568: "Partly partitioned" shortening already is going on based on line 566, so in line 568 how can there be a *gradual change to* "partly partitioned" shortening?

Comment 111: Line 570: What is the nature of the "oblique-slip"? Is there a dextral-slip component? Do you ever actually document dextral slip on the Indio Hills Fault? Please clarify.

Comment 112: Line 571: By this do you mean that----prior to this time----the Banning was NOT a major player, although it would become so later in the structural evolution of the culmination?

Comment 113: Line 572: By this do you mean that formerly "open" folds became "more tightly closed" as they were squeezed by "increased shear folding"?

Comment 114: Line 573: What is this fold style? You have not discussed it previously in your structural analysis.

Comment 115: Line 578: Because youthful dextral slip on the later truncated culmination folds and displaced their truncated counterparts NW away from the culmination----yes?

It is important to inform the reader that you actually have no idea about the structural configuration of the macro folds SW of where you actually observe them.

Comment 116: Lines 576–578: I do not understand how the interpretation in lines 579-581 follows from lines 576-578

Comment 117: Line 582: Age of "late stage" needs to be specified. In other words, if basin inversion and tectonic uplift occurred in stages, you need to specify the age ranges of terms like "early-stage" and "late-stage".

Comment 118: Line 582: Does this mean "spatially involved"? "kinematically evolved"? Clarify

Comment 119: Line 587: label this structure on figures 3c and 7c.

Comment 120: Line 588: margin-parallel to what? The tectonic culmination? Clarify

Comment 121: Lines 592–593: Not clear to me what the interpretive boundary is between when "although overlapping and synchronous formation....may have occurred" (that is, if some unspecified percentage of structures may have overlapped timing wise, when does that percentage become a deal-breaker for your structural paradigm?)

Comment 122: Line 594: Antecedent unclear: what is the "latter"?

Comment 123: Line 594: Does this mean "we don't have enough field data to confirm X, Y, and Z"? Or does it mean "we have not observed a sufficient number of specific fold and fault types to confirm "progressive evolution from distributed to partly distributed deformation" (see line 591)? Please clarify

Comment 124: Line 602: See my comments earlier in this report. In the Little San Bernardino Mountains this fault zone does not exist at the surface. Revise and explain

Comment 125: Line 606: No. You can only relate recent earthquakes in the vicinity of the "Landers-Mojave Line" in the ECSZ if you can show that the Indio Hills Fault is active as an earthquake generator. You have not discussed this possibility, and no researcher to my knowledge has shown that the Indio Hills Fault is an active player.

Comment 126: Line 609: To compare alleged active faulting on the Indio Hills Fault with far-flung faults like the Calico and Camp Rock is inappropriate. Better to compare with nearby faults investigated by Rymer (2000) in the Little San Bernardino Mountains, or with the 1992 Landers and Joshua Tree events themselves. Revise.

Comment 127: Line 609: See my comment earlier about regional extent of the term "Banning Fault".

Comment 128: Line 610: You absolutely need to explain what this is. You can't just point us to Platt and Passchier and find out for ourselves.

By the way, reference to P&P (2016) implies that those two authors investigated the SAFZ southeast of the Indio Hills. Did they? I am not going to go look. You need to assert that you are adopting the dextral freeway junction (sensu P&P, 2016) who invented the term based on work elsewhere.

Comment 129: Line 610: Something wrong with this grammar (phrase)

Comment 130: Line 611: What does (late) mean? In comparison with "early"? You really need a table or figure that shows the age and movement history of the Banning and Indio Hills Faults.

See my previous comment.

Comment 131: Lines 615–616: Have you stated this age and discussed it earlier in your manuscript? I cannot find such a discussion. You need to document ages for the Banning and Indio Hills faults, with basis for age calls. Given the complexity of fault relations, the diverse tectonic interpretations in the literature, and uncertainty about fault names throughout the region, you absolutely need to clarify and document age relations for specific faults.

Comment 132: Line 625: Well, almost certainly the genesis of the Indio Hills tectonic culmination (resulting in basin inversion) was gradual and progressive. Is it better to state that you have documented what is intuitive to most investigators in this region? Compare with Gold and others (2015) and Blisniuk and others (2021). Do they elaborate on progressive uplift and exhumation?

Comment 133: Lines 633–634: I don't know what you mean by "the SAFZ main strand".

According to previous statements in this manuscript, the Banning Fault is the "main strand" SE of the Indio Hills (but see my comment to that effect earlier in this manuscript). So what does the Banning Fault merge into as it projects southeastward? See how Behr and others (2010) deal with this using an "Indio strand".

Comment 134: Line 639: Antecedent unclear. What does "them" refer to?

Comment 135: Lines 639–640: Do you mean "Indio Hills" footprint? Or "Indio Hills Fault"?.

Clarify. If you refer to the fault, then how do you know it has thin versus thick gouge if you can't see the fault in any outcrops? (Everywhere concealed by unconsolidated surficial deposits).

Comment 136: Lines 641–642: Also, the relatively massively structured Mecca Formation intervenes between the Indio Hills Fault and the highly deformed Palm Spring Formation. Would that make a difference?

Comment 137: Liens 645–646: This is new information for me. McNabb and others (2017) do not refer to this (contrary to your statement in line 648), nor do Fattaruso and others (2014, 2016). For benefit to readers like myself, it would be helpful if you were to review the proposals for----and basis for----large normal-slip faults bounding the east margins of Miocene-Pliocene sedimentary basins that evolved in the Indio-Mecca Hills footprint of the ancient Salton Trough. I have read the summary on p. 4 of Bergh and others (2019), and I see that some workers posit that basin fills like those in the Mecca and Indio Hills accumulated in a Miocene-Pliocene depocenter formed (in the hangingwall?) adjacent to a large Basin-and-Range style "normal fault". Is this structure documented, or is model-dependent? Does this large "normal fault" crop out anywhere, or has it been so transformed by younger activity that its normal-slip origin can only be inferred? A better summary by you would be helpful to regional geologists like myself.

Comment 138: Line 648: In McNabb et al I do not see any reference to a normal fault on the NE side of the Mecca Hills, that later was reactivated as an oblique reverse fault

Comment 139: Line 649: I see on p. 5 of Bergh and others (2019) that Sylvester and company view the Painted Canyon Fault as an "oblique reverse strike-slip fault"; but might this fault originally have had a normal-slip origin. Please help us here.

Comment 140: Lines 699–700: Do you actually observe the width of the Banning Fault damage zone in the southern Indio Hills? I don't remember you discussing this in your manuscript. Where you might observe the "damage zone" isn't it typically obscured by young surficial deposits and wind-blown sand?

Comment 141: Line 725: I don't remember you defining this or referring to it elsewhere in the manuscript. What is the distribution of this structure in figure 1? Is this the same as the Banning Fault + Indio Hills Fault? Most workers refer to the combined traces of the SAFZ in the Coachella Valley as the "Coachella Valley segment of the SA" following Matti and others, 1992 (see

Comment 142: Line 731–732: If the Indio Hills and Banning faults are not coeval, then what was the southwest bounding fault that worked together with the Indio Hills Fault to produce the tectonic culmination of the Indio Hills?

Comment 143: Line 738: Your conclusions should emphasize that the Indio Hills Fault has a polyphase history: (1) It started its life as a Miocene normal fault according to your interpretation; (2) it evolved into a reverse/thrust fault that initiated inversion of the Pliocene sedimentary basin;

(3) it now (?) is a dextral strand of the SAFZ according to conclusion (4). This polyphase history should go near the top of the conclusions section in my opinion.

Comment 144: Lines 738–739: As I repeat throughout the manuscript, this is a hypothesis only. You must state it as such.

Comment 145: Line 881: Insert "Lancaster and coauthors, 2012", here in place of Parrish, 1983 as discussed below and in my memo

Comment 146: Line 928: This reference is incorrect (I should know, as I should have been lead author). The actual citation should be:

Lancaster, J.T., Hayhurst, C.A., and Bedrossian, T.L., 2012, Preliminary geologic map of Quaternary surficial deposits in southern California: Palm Springs 30' x 60' quadrangle, in Bedrossian, T.L., Roffers, P., Hayhurst, C.A., Lancaster, J.T., and Short, W.R., Geologic compilation of Quaternary surficial deposits in southern California December 2012

<https://www.conservation.ca.gov/cgs/fwgp/Pages/sr217.aspx#palmsprings>.

Comment 147: Line 930: You may want to cite Passchier and Platt (2016) in the Journal of Structural Geology in addition to Platt and Passchier

Comment 148: Line 978: Most workers would apply the name "Mission Creek Fault" to what you term the "Banning Fault". The point is: which fault is the dominant throughgoing structure at the latitude of your study area?

Comment 149: Line 1002: On this figure, it would be useful to show the geologic contact between the Mecca Formation and the Palm Spring Formation.

Comment 150: Line 1002: Label this fold so that readers of lines 586-587 can identify it and associate it with your kinematic model.

Comment 151: Line 1002: You apparently refer to this "fault-parallel anticline in lines 450 and 454. If so, I would label it here.

Comment 152: Line 1041: What is this time interval, and how long was it? Episode in Pliocene? Pleistocene? Clarify

Comment 153: Line 1055: Literature source for extending the Garnet Hill strand this far SE

Comment 154: Line 1062: cite reference (Janecke et al, 2018)????

Comment 155: Line 1063: Singular "Hill"

Comment 156: Supplement S6: Not clear from the photographs that these are "normal" faults. Please explain how you make this interpretation.

2. Author's reply

Comment 1: agreed, we trying to reach both specialists in structural geology and regional geologists. See also response to comments 33, 47, 128 and 147. However, the term “ladder” was defined upon first occurrence in the text lines 272–274: “as defined by Davis (1999) and Schulz and Balasko (2003), where overlapping, E–W- to NW–SE-striking step-over faults rotated along multiple connecting cross faults”.

Comment 2: agreed.

Comment 3: agreed.

Comment 4: agreed.

Comment 5: we agree that reference to Dokka et al. (1990a, 1990b) to support connection of the Eastern California Shear Zone and Landers Mojave Line with the Indio Hills fault is not enough. Therefore we add reference to Nur et al. (1993a, 1993b) who connect the 1992 Joshua Tree earthquake with the series of earthquakes that occurred along the Landers Mojave Line between 1947–1992 (see their figure 1). We also agree that the manuscript should clarify that it is unlikely that transfer of strain between the ECSZ and the SAFZ occurs along a single fault trace. However, we do not understand the comment of the reviewer claiming that the epicenter of the 1992 Joshua Tree earthquake is located a few km north of the West Deception Canyon fault. Looking at Rymer (2000, his figures 1 and 2), it is clear that the earthquake's epicenter is located exactly along the trace of the West Deception Canyon fault. Rymer (2000) even mentions that although he observed rupture along the East Wide Canon fault, that it is clear from the location of the epicenter (east of this fault) and from this fault's western dip and very small amount of rupture-related offset (ca. 6 mm) that it cannot have triggered the 1992 Joshua Tree earthquake. Rymer (1992) further specifies that structural fieldwork and interpretation of aerial images suggest that the West Deception Canyon fault is connected to the Eureka Peak fault, which moved during the June 1992 earthquake.

Comment 6: agreed.

Comment 7: the “grammatical structures” mentioned in the comment are the way chosen to express uncertainty. In some cases, the data allow us to “favor” one or the other possibility (e.g., lines 473, 559, and 691), whereas in other cases they do not. In the instances the data do not allow us to favor one alternative or the other(s), it would be inappropriate and unethical to arbitrarily choose any of the alternatives. We prefer to remain open and admit that we do not “know” everything for certain,

leaving ample room to future workers in the area to further validate, reject, or rework the proposed model.

Comment 8: agreed. See response to comments 5, 12, and 125. Uplift along the Indio Hills fault seems to have been superior to uplift along the main San Andreas fault as suggested by the intensely folded geometries of sedimentary strata and higher topographic relief in the vicinity of the Indio Hills fault. This needs to be mentioned in the manuscript. However, the present manuscript does not address present-day growth or not of the Indio Hills.

Comment 9: agreed.

Comment 10: agreed. Also see response to comments 39 and 40.

Comment 11: agreed. See response to comment 2.

Comment 12: agreed. Instead of calling the main splay of the San Andreas fault the “Banning Fault”, we change it to the “main San Andreas fault” (mSAF), in accordance to the recent study by Janecke et al. (2018) in the area.

Comment 13: agreed. See response to comment 12.

Comment 14: agreed. See response to comment 12.

Comment 15: agreed. See response to comment 12.

Comment 16: agreed. Also see response to comment 12.

Comment 17: agreed.

Comment 18: agreed.

Comment 19: agreed.

Comment 20: agreed.

Comment 21: agreed. See response to comment 20.

Comment 22: agreed.

Comment 23: see response to comment 22.

Comment 24: agreed.

Comment 25: agreed. Also see response to comment 6.

Comment 26: agreed.

Comment 27: partly agreed.

Comment 28: agreed.

Comment 29: agreed.

Comment 30: agreed.

Comment 31: agreed. See response to comment 30.

Comment 32: agreed. See response to comment 30.

Comment 33: agreed. See response to comment 30.

Comment 34: agreed. See response to comment 28.

Comment 35: agreed.

Comment 36: agreed.

Comment 37: see response to comment 36.

Comment 38: agreed.

Comment 39: agreed.

Comment 40: agreed.

Comment 41: agreed.

Comment 42: the study of Fuis et al. (2017) suggests that the SAFZ dips steeply at shallow depth both in the Indio Hills and Mecca Hills, but more moderate dips are probable at higher depth.

Comment 43: agreed.

Comment 44: agreed.

Comment 45: agreed. The study is based on both field study and imagery analysis.

Comment 46: agreed.

Comment 47: the term “Indio Hills-parallel” does not bear any structural meaning, whereas “SAFZ-parallel” suggests that the anticline is the product of dip-slip to oblique-slip thrusting rather than strike-slip fault movements as inferred for the SAFZ-oblique macro-folds.

Comment 48: agreed.

Comment 49: this does not necessarily imply detachment between strongly and weakly folded strata, but local detachment/décollement may occur as suggested by the next sentence (see also possible detachment folds in figure 4 in the present manuscript). We concede that these local detachment folds and décollements should be discussed, but not in the Result section since it is too interpretative. We instead mention this interpretation in the Discussion section lines 583–584 and 613.

Comment 50: agreed.

Comment 51: agreed.

Comment 52: agreed.

Comment 53: agreed.

Comment 54: agreed.

Comment 55: agreed.

Comment 56: the fractures in the granitic basement have not been dated and may therefore be older than the deformation of sedimentary strata in the Indio Hills. However, the strike of brittle fault sets in the granite basement matches that of fractures in sedimentary strata (see stereoplots in Figure 2), which suggests that the fractures formed due to similarly oriented stress. The authors of the present manuscript suggest to mention this explanation in the manuscript if also judged convenient by the referee.

Comment 57: agreed.

Comment 58: yes indeed, the truncation of the vertical folds and *en echelon* SAF-oblique folds may result from dip-slip reverse movement along the main San Andreas fault. However, no direct evidence was encountered to support reverse dip-slip movement along this fault, and the geometry of vertical (shear) folds in the direct vicinity of the fault (Figure 5) and the anticlockwise rotation of the axis of the three *en echelon* macro-folds towards the fault (Figure 2) suggest a significant component of lateral movement along the main San Andreas fault at some point.

Comment 59: agreed. However, this would require an additional level of sub-title/sub-section, which is not allowed by the journal's standards.

Comment 60: disagreed because of the journal's standard.

Comment 61: agreed.

Comment 62: agreed.

Comment 63: see response to comment 56.

Comment 64: agreed.

Comment 65: agreed.

Comment 66: agreed.

Comment 67: agreed.

Comment 68: agreed.

Comment 69: agreed.

Comment 70: agreed.

Comment 71: agreed.

Comment 72: agreed.

Comment 73: agreed.

Comment 74: agreed.

Comment 75: agreed.

Comment 76: agreed.

Comment 77: see response to comment 80.

Comment 78: agreed.

Comment 79: see response to comments 80 and 156.

Comment 80: agreed.

Comment 81: see response to comment 80.

Comment 82: agreed.

Comment 83: agreed.

Comment 84: agreed.

Comment 85: partly agreed.

Comment 86: yes.

Comment 87: the truncating relationship of the main San Andreas fault and the en echelon macro-folds in the study area suggest that deformation was distributed at first, but more localized during the activation of the main San Andreas fault in the late stage of deformation since no major reworking of the macro-folds is observed along this fault (apart from some shear folding near the fault; see figure 5 in the present manuscript).

Comment 88: agreed. The term “late” was meant as “late for the study area”. However, we concede that its meaning is not appropriate since there is probably ongoing slip along the Mission Creek and main San Andreas faults at present (Gold et al., 2015), and possibly along the Indio Hills fault if it is part of the Landers Mojave line (e.g., Nur et al., 1993a, 1993b). Clarification of the meaning is therefore needed in the sentence.

Comment 89: agreed. However, this paragraph analyzes the fold-fault relationships for a different purpose, i.e., to show that strain partitioning already occurred in the early phase of deformation of the area despite the occurrence of distributed deformation (e.g., SAFZ-oblique macro-folds).

Comment 90: we want to draw the attention of the reader on the fact that, in their current position, it is not possible to explain the formation of the observed small folds and faults with simple transpression and/or contraction. The unorthodox geometries of the observed folds and faults is however simply due to the steep dip of the strata in which they were observed. If restoring the strata

to horizontal (i.e., prior to deformation), the small folds and faults turn into structures with geometries typical of contractional folds and thrusts.

Comment 91: agreed. The sentence is not clear. Supplement S6 addresses restoration of the steep of the sedimentary strata to horizontal (i.e., prior to deformation).

Comment 92: agreed.

Comment 93: precisely, agreed.

Comment 94: agreed.

Comment 95: agreed.

Comment 96: if the reviewer's comment is meant for distributed versus partitioned deformation, we specify that we favor synchronous distributed and partitioned deformation. If the comment is targeting the progressive versus synchronous bit, the end-member we favor is specified line 677.

Comment 97: agreed.

Comment 98: completely agreed.

Comment 99: agreed. See response to comment 80.

Comment 100: agreed. This is a mix up from the multiple re-writing phases of the manuscript. The meaning of the sentence needs to be adjusted. The Mecca Formation is not found in the footwall of the fault (e.g., map by Lancaster et al., 2012).

Comment 101: this statement is based on the previous study by McNabb et al. (2017) in the Mecca Hills, not on the present study.

Comment 102: agreed.

Comment 103: agreed.

Comment 104: agreed. See response to comment 102.

Comment 105: agreed.

Comment 106: agreed.

Comment 107: disagreed. The present dataset and observations do not allow to favor one or the other. It is therefore important to mention both possibilities to account for uncertainties, which are a very important part of the research itself from a science ethics perspective.

Comment 108: disagreed. There is not much difference between "prior to" and "together with" in the present scenario/model, and, again, the dataset does not allow to favor one or the other alternative.

Comment 109: yes, we meant “strain partitioning”. It is true that the minor folds may have formed either prior to or together with the *en echelon* macro-folds. However, this is irrelevant with regards to the present sentence because we argued that distributed deformation occurred in the early stages of deformation during formation of the macro-folds. Thus, if the minor folds indicate minor strain partitioning, it suggests that both distributed and partitioned deformation occurred at the same time.

Comment 110: agreed.

Comment 111: agreed. We connect SAFZ-oblique *en echelon* macro-folds in the study area to right-lateral slip along the Indio Hills fault.

Comment 112: absolutely. Agreed that the sentence need rewording rto better reflect its intended meaning.

Comment 113: yes, agreed.

Comment 114: agreed, the term is confusing.

Comment 115: the macro-folds were not observed southwest of the main San Andreas fault. But it is irrelevant to mention this in the present section. We agree, however, to mention it in the result section where it is more appropriate.

Comment 116: agreed. It does not. The sentence needs rewording.

Comment 117: agreed.

Comment 118: agreed. We mean kinematically evolved.

Comment 119: it is probably not wise to overcrowd the figures with excessive labelling.

However, we agree that we should be more specific to help the reader follow.

Comment 120: agreed.

Comment 121: see response to comment 105.

Comment 122: agreed.

Comment 123: see response to comment 105.

Comment 124: see response to comment 5.

Comment 125: the work of Nur et al. (1993a, 1993b) demonstrates that the Landers Mojave Line is a through-going fault that crosscuts the Pinto Mountain fault. They add that “segments of this fault were identified in the field before 1992 (M. Rymer, personal communication)”, but that “it was not recognized as a through-going, coherent and seismogenic fault” then. Rymer had been working with the Indio Hills fault at that time and the location of this fault matches that of the southern segment of the Landers Mojave Line near the Joshua Tree earthquake. We therefore

argue that the Indio Hills fault is a segment of this through-going active fault. Nevertheless, we concede that it is necessary to add up to this paragraph for clarification.

Comment 126: partly agreed. The contribution by Rymer (2000) is appropriate to mention here. Rymer shows that the 1992 Joshua Tree earthquake occurred along the NNW–SSE-striking, west-dipping West Deception Canyon fault in the Little San Bernardino Mountains, and that this fault merges to the south with the Indio Hills fault (see his figure 1). It is therefore relevant and paramount to discuss the role of the Indio Hills fault with regards to the various fault systems it seems to connect, including NNW–SSE-striking faults of the Landers Mojave Line (e.g., West Deception Canyon fault), NW–SE-striking faults of the Eastern California Shear Zone (e.g., Calico and Camprock faults), and the main San Andreas fault.

Comment 127: agreed. See response to comment 12.

Comment 128: agreed.

Comment 129: agreed.

Comment 130: agreed.

Comment 131: agreed. See response to comment 117.

Comment 132: partly agreed. Gold et al. (2015) and Blisniuk et al. (2021) investigate recent (Holocene) slip rates for the Mission Creek and Banning faults, whereas we study older deformation history in the southeastern Indio Hills (Pleistocene). However, we concede that a few sentences are needed in order to compare our findings to those of Keller et al. (1982) in the northwestern Indio Hills.

Comment 133: agreed.

Comment 134: agreed.

Comment 135: partly agreed. However, gouge and anastomosing geometries should probably be observed along minor faults that strike parallel to, show comparable kinematics, and are located in the vicinity of the Indio Hills fault (e.g., S3a). This is not the case.

Comment 136: yes, it could, but the conglomerates of the Mecca Formation are only weakly folded.

Comment 137: disagreed. McNabb et al. (2017) inferred “SW-side down slip” along the Painted Canyon fault based on the presence of the Mecca Formation conglomerate in the hanging wall and its absence in the footwall of the fault, and on sedimentary facies (conglomerate fining up upwards; see first paragraph in their discussion pp. 81 and pp., their figs. 15 and 16, and the second paragraph of the conclusion pp. 84). Also see response to comment 80.

Comment 138: see response to comment 137.

Comment 139: see response to comment 137.

Comment 140: agreed.

Comment 141: agreed. See response to comment 12.

Comment 142: we did not imply that the main San Andreas and Indio Hills faults were not coeval. They were coeval at least in the later stages of our model (i.e., in the late Pleistocene). See also response to comment 117.

Comment 143: agreed.

Comment 144: see response to comments 125 and 126.

Comment 145: agreed.

Comment 146: agreed. See response to comment 145.

Comment 147: agreed. See response to comment 128.

Comment 148: agreed.

Comment 149: the transition between the two units is gradual and is not very well constrained. In addition, the transition between the two units is not critical to the present contribution. The reader may have a look at the map by Lancaster et al. (2012) to get an approximative idea of the location of the transition.

Comment 150: agreed. See response to comment 73.

Comment 151: partly agreed. See response to comment 119.

Comment 152: agreed.

Comment 153: agreed.

Comment 154: the wide arrows indicating main shortening direction are not from Janecke et al. (2018), but from the present contribution for the Indio Hills, and from Bergh et al. (2019) for the Mecca Hills.

Comment 155: agreed.

Comment 156: a simple rotation of sedimentary and truncating micro faults by 52 degrees counterclockwise to restore the sedimentary bed to the horizontal (i.e., prior to macro-folding deformation) shows that some micro faults display reverse offsets of bedding surfaces and most likely formed as micro thrusts (see Supplement S6a–b), whereas other micro faults display normal offsets of bedding surfaces and form micro graben structures with associated syn-kinematic growth strata (see Supplement S6c).

3. Changes implemented

Comment 1: see also response to comments 33, 47, 128 and 147. Also added “(stretched long limb in an overturned fold)” line 319, “(the shortened, inverted limb indicating the direction of tectonic transport in an overturned fold)” lines 339–340, and “folds involving shearing along a plane that is parallel to subparallel to the fold’s axial plane; Groshong, 1975; Meere et al., 2013;” lines 302–304 and Groshong (1975) and Meere et al. (2013) to the reference list.

Comment 2: rewrote lines 40–47 and moved some information to the new section about “Regional faults” as suggested by the referee. Moved first paragraph of the section about Tectonic culminations to the new section about “Regional faults”. Wrote section about Regional faults as follows: “***Regional faults***

The southeastern Indio Hills are a WNW–ESE–trending tectonic culmination situated in a small restraining bend northeast of the main San Andreas fault (Figs. 1 and 2). The studied culmination is located along strike about 25–50 kilometers northwest of the Mecca Hills and Durmid Hills, and to the southeast of the major left bend in the SAFZ trace near San Gorgonio Pass (Matti et al., 1985, 1992; Matti and Morton, 1993; Dair and Cooke, 2009).

The main faults in the southeastern Indio Hills include the Indio Hills fault in the northeast (Allen, 1957; Tyley, 1974), and the main San Andreas fault in the southwest. Regionally, the Indio Hills fault possibly merges with the Landers Mojave Line and the eastern California Shear Zone in the north (Dokka and Travis, 1990a, 1990b; Nur et al., 1993a, 1993b; Thatcher et al., 2016). The Landers Mojave Line is believed to be the locus of several recent earthquakes aligned along a NNW–SSE-trending axis, including the 1992 Joshua Tree earthquake (Fig. 1; Nur et al., 1993a, 1993b). These earthquakes were tentatively ascribed to movement along a through-going NNW–SSE-striking fault, possibly the west-dipping, Quaternary West Deception Canyon fault (Sieh et al., 1993; Rymer, 2000). This fault is thought to crosscut the E–W- to ENE–WSW-striking, left-lateral, Holocene Pinto Mountain fault, which merges with the main strand of the San Andreas fault in the west at the intersection of the right-lateral Mission Creek and Mill Creek strands (Allen, 1957; Bryant, 2000; Kendrick et al., 2015; Blisniuk et al., 2021). The former is thought to correspond to the continuation of the main San Andreas fault to the northwest (Gold et al., 2015) and may have accommodated ca. 89 km of right slip in the past 4 million years,

whereas the latter accommodated about 8 km right slip at 0.5–0.1 Ma and is offset ca. 1 km by the Pinto Mountain fault (Kendrick et al., 2015).

The main San Andreas fault continues to the southeast where it bounds the Mecca Hills to the southwest, whereas the Painted Canyon fault, a previous (late Miocene?–) Pliocene southwest-dipping normal fault reactivated as a right-lateral-reverse oblique-slip fault in the Pleistocene–present-day bounds the Mecca basin to the northeast (Sylvester and Smith, 1987; McNabb et al., 2017; Bergh et al., 2019). Farther southeast, the main San Andreas fault proceeds along the northeast flank of the Durmid Hills opposite the Pleistocene (ca. 1 Ma), right-lateral East Shoreline fault (Babcock, 1969, 1974; Bürgmann, 1991; Janecke et al., 2018). There, the main San Andreas fault merges with the Brawley seismic zone (Lin et al., 2007; Hauksson et al., 2012; Lin, 2013) and, together with the right-lateral San Jacinto fault zone, they merge into the right-lateral Imperial fault (Rockwell et al., 2011). In the north, the main San Andreas fault splays into the Banning strand and the Mission Creek fault in the northwestern part of the Indio Hills (Keller et al., 1982; Gold et al., 2015). The Banning strand is much younger than the Mission Creek fault and may have accommodated approximately 3 km of right slip in the past 0.1 million years (Kendrick et al., 2015).

Northwest and west of the Coachella Valley, Miocene–Pleistocene sedimentary strata are structurally bounded by the San Bernardino and San Jacinto fault strands of the SAFZ (Bilham and Williams, 1985; Matti et al., 1985; Morton and Matti, 1993; Spotila et al., 2007). To the southwest, Miocene–Pleistocene strata are bounded by the West Salton Detachment fault (Dorsey et al., 2011). The San Jacinto fault is typically believed to have slipped ca. 25 km right-laterally in the past 1.5 million years (Matti and Morton, 1993; Kendrick et al., 2015), whereas the West Salton Detachment fault is a low-angle normal fault that accumulated ca. 8–10 km of normal-oblique movement starting in the mid Miocene and is related to the opening of the Gulf of California (Prante et al., 2014 and references therein).” lines 73–118.

Also added “Below we summarize local and regional fault nomenclature, distribution, and fault movement history (Table 1) throughout the greater Coachella Valley region (Fig. 1), the stratigraphy of the Indio Hills area, and previous structural work in the main Indio Hills, Mecca Hills, and Durmid Hills uplift areas.” lines 75–79.

Comment 3: see response to comment 12.

Comment 4: see response to comments 80, 117, 130, 131, and 143.

Comment 5: added reference to Nur et al. (1993a, 1993b) line 698. Added “be one of several faults to” lines 808–809. Changed “transfers” into “contributes to transfer” line 966.

Comment 6: developed a new summary table with the ages of relevant geological events in the Coachella Valley and nearby and added reference to the table lines 79, 114, 128, 139, 206, . In addition, adjusted ages throughout the manuscript after latest findings as suggested, including “Miocene–Pliocene” into “Pliocene–Pleistocene” lines 13, 61, and 71, “2.2–” into “or later than” line 18, “2.2” into “2.6” line 18, “Miocene–Pliocene” into “mid to upper Pliocene” line 88, “is an angular unconformity that signals” into “is marked by two angular unconformities that signal” line 103, “3.7–2.6” into “3.0–2.3”, and “mid–late Pliocene” into “latest Pliocene–early Pleistocene” line 107, “2.8–1.0” into “2.6–0.76” and “late Pliocene–mid Pleistocene” into “earliest Pleistocene to earliest late Pleistocene” line 108, “Pliocene” lines 152 and 156 into “Pleistocene”, “4.0–3.7” into “3.7–3.0” line 675, “3.7–2.8” into “3.0–2.3” line 676, “2.8–1.0” into “2.6–0.76” line 677, “mid” into “late” line 695, “1” into “0.76” line 766, and “3.0–2.2” into “2.6–0.76” and “at 1.0–0.76” into “after 0.76” line 840.

Also added reference to McNabb et al. (2017) line 112. Added “In contrast to other uplift areas in Coachella Valley, the Ocotillo Formation has not been mapped in the Indio Hills in the present study. However, based on the occurrence of the Bishop Ash at the northwestern edge of the study area and on the occurrence of the volcanic deposit within the uppermost Palm Spring Formation or at the base of the overlying Ocotillo Formation in the Mecca Hills, it is likely that the Ocotillo Formation occurs just northwest of the area mapped (Fig. 2). In addition, it is deposited on the flank northeast of the Indio Hills fault, and southwest of the main San Andreas fault (Figs. 1 and 2), indicating that this unit was either not deposited or eroded in the area that recorded the most uplift in Indio Hills.” lines 116–123. Deleted “Mecca Hills and Durmid Hills” lines 124–125 and replaced by “the Coachella Valley”. Added “. The volcanic deposit is found within” line 127, “(which is unconformably overlain by the Ocotillo Formation) in the hanging wall of the Painted Canyon fault away from the fault, and within the base of the Ocotillo Formation in the hanging wall of the Painted Canyon fault near the fault (Ocotillo and uppermost Palm Spring formations interfingering near the fault) and in the footwall of the fault” lines 128–132, and “. The unconformable contact between the Palm Spring and Ocotillo formations away from the Painted Canyon fault towards the southwest and their interfingering relationship near the fault suggest that uplift had already initiated prior to deposition of the Ocotillo Formation (i.e., before 0.76 Ma,

in the mid Pleistocene), possibly during the formation of the lower unconformity between the lower and upper members of the Palm Spring Formation (McNabb et al., 2017). In addition, the involvement of the Bishop Ash in deformation suggest that deformation continued past 0.76 Ma (in the late Pleistocene).” lines 132–139. Deleted “Janecke et al., 2018” line 132. Deleted “In contrast to other uplift areas in Coachella Valley, the Ocotillo Formation has not been mapped in the Indio Hills, but rather is deposited on the flank northeast of the Indio Hills fault, and southwest of the main San AndreasBanning fault (Figs. 1 and 2), indicating that the Ocotillo Formation was either not deposited, or eroded in the area of uplift.” lines 140–143. Added “because of the involvement in folding of the Bishop Ash and of adjacent strata possibly of the Ocotillo Formation (i.e., maximum age of 0.76 Ma – earliest late Pleistocene; Fig. 2 and Table 1)” lines 483–485 and “Should the whole Ocotillo Formation be folded in the Indio Hills, the maximum age constraints could be narrowed to < 0.6–0.5 Ma based on magnetostratigraphic ages for the upper part of the Ocotillo Formation (Kirby et al., 2007).” lines 486–489. Added “ in the late Pleistocene” line 538. Added “in the (earliest?) late Pleistocene (Table 1)” lines 546–547. Added “-stage (i.e., late Pleistocene)” lines 563–564. Added “in the mid-Miocene–Pliocene (ca. 15–3.0 Ma)” lines 570–571, “in the (earliest?) late Pleistocene to present-day (< 0.76 Ma)” line 572, and “in the late Pleistocene to present-day (< 0.76 Ma; Table 1)” lines 573–574. Added “in the late Pleistocene” lines 606–607, 615, and 617. Added “in the – earliest? – late Pleistocene” line 618. Added “(see phases 1 and 2 in Table 1)” line 654. Added “(< 0.76 Ma)” line 659. Added “late” line 671. Added “(Miocene?–) Pliocene” line 672. Replaced “they were overlain by” by “strata of” line 674. Added “were deposited in the Pliocene” line 675, “members of the” line 676 and reference to Chang et al. (1987) and Boley et al. (1994) lines 677–678. Deleted “by” and “ succeeding,” line 675. Added “(earliest?) late” line 679. Added “after the latter was deposited (< 0.76 Ma), i.e., probably in earliest late Pleistocene time (Table 1)” lines 682–683. Deleted “Late-stage ” line 711. Added “and phase 3 in Table 1” line 717. Added “(overlapping of phases 1 and 2 in Table 1)” line 727 and “; phase 3 in Table 1” line 728. Added “late” line 736 and “to present-day” lines 736–737. Added “in the late Pleistocene” and “and Table 1” line 761. Added “–Pliocene” line 803. Added “(late)” line 805. Added “(i.e., earliest to mid Pleistocene) with partial and local erosion of the Palm Spring Formation (see lower and upper unconformities in McNabb et al., 2017)” lines 840–842. Added “(see unconformity between the uppermost Palm Spring Formation and base of the Ocotillo Formation southwest of the Painted Canyon fault in

McNabb et al., 2017)” lines 842–844. Added “whole” line 844. Deleted “A comparable time frame and ongoing activity are expected for the Indio Hills.” lines 846–847 and replaced by “Fault activity and tectonic uplift of the Mecca Hills therefore most likely initiated earlier (earliest Pleistocene) than in the Indio Hills (earliest late Pleistocene; Table 1), where the transition from the lower to the upper member of the Palm Spring Formation is gradual and does not show any major unconformity.” lines 847–850. Added “(ca. 1 Ma – early/mid Pleistocene)” line 870. Added “(i.e., probably in the earliest or middle part of the late Pleistocene)” lines 874–875. Added “mid” line 892. Added “in the late Pleistocene” line 896. Added “The initiation of right-lateral to right-lateral-reverse slip along major SAFZ-parallel faults and the main San Andreas fault in the Coachella Valley is younger towards the northwest (Pliocene in the Durmid Hills, early Pleistocene in the Mecca Hills and late Pleistocene in the Indio Hills). The onset of transpressional uplift, however, appears to be coeval in all tectonic culminations (late to latest Pleistocene.” Lines 918–922.

Comment 7: none.

Comment 8: see response to comments 5, 12, and 125. Also adjusted the position of the main San Andreas fault in figure 7a and b to match that in figure 7c. Added “Possibly as a consequence of a longer period of activity, and as suggested by relatively higher topographic relief and more intensely folded geometries of sedimentary strata in the vicinity and along the Indio Hills fault than along the main San Andreas fault, it is probable that the former accommodated significantly larger amounts of uplift than the latter. This implies a southwest-tilted geometry for the Indio Hills culmination.” lines 587–591.

Comment 9: deleted “where they approach the main San Andreas fault” lines 456–457, and added this to the Discussion chapter. Deleted “main San Andreas and”, added “rather” and “a single”, and replaced “faults” by “fault” twice lines 540–541. Deleted “active” and added “more active” line 541. Deleted “(i.e., near the main San Andreas fault)” line 588. Replaced “main San Andreas fault” by “the southwest”, replaced “strain” by “off-fault deformation”, and added “main San Andreas” lines 681–683. Replaced “with the main San Andreas fault” by “farther southwest” lines 776–777.

Comment 10: added Gold et al. (2015) to the reference list. Added “The study also showed that drainage systems were offset recently (at ca. 0.03–0.02 Ma) and indicate relatively high slip rates along the Mission Creek fault in the order of 23–35 cm.y⁻¹, i.e., comparable to the more recent c.

23 cm.y⁻¹ estimate by Blisniuk et al. (2021).” lines 238–240. Also see response to comments 39 and 40.

Comment 11: see response to comment 2.

Comment 12: replaced “Banning” by “main San Andreas” lines 17, 23, 25, 39, 42, 96, 116, 131, 132, 145, 210, 220, 223, 233, 239, 244, 248, 284, 317, 323, 331, 337, 340, 346, 351, 354, 356, 358, 392, 422, 436, 445, 475, 517, 539, 540, 543, 546, 602, 611, 618, 626, 630, 635, 641, 653, 655, 658, 660, 665, 678, 723, 743, 754, 760, 773, 777, 783, 786, 1043, 1074, 1093, and 1103. Deleted “thought to correspond to the main SAFZ in” line 40. Added “, which merge into the main San Andreas fault” lines 154–155.

Comment 13: see response to comment 12.

Comment 14: see response to comment 12.

Comment 15: see response to comment 12.

Comment 16: replaced “marges” by “may merge” line 41–42. Also see response to comment 12.

Comment 17: added “potential” line 46.

Comment 18: rewrote the sentence lines 56–58 as “The Coachella Valley segment of the SAFZ in southern California is expressed as multiple, right-lateral fault strands, which uplifted blocks in the Indio Hills, Mecca Hills, and Durmid Hills”.

Comment 19: corrected “Palm Springs Formation” into “Palm Spring Formation” lines 73 and 471.

Comment 20: deleted “basement” lines 81 and 86, and added “and that at least part of the clasts are from the leuco-granitic rocks, which must correspond to basement rocks of the inverted Indio Hills basin” lines 88–90.

Comment 21: see response to comment 20.

Comment 22: added “gradual” line 99 and “By contrast, “ line 101. Deleted “In the Indio Hills, however, the nature of the transition between the lower and upper member of the Palm Spring Formation and the presence of an angular unconformity is unknown.” lines 103–105. Moved “respectively” earlier in the sentence for clarity line 107. Split the last sentence into a new paragraph lines 117–120 because dealing with a different stratigraphic unit (Ocotillo Formation).

Comment 23: see response to comment 22.

Comment 24: separated the narrative about age relationships into a discrete paragraph (lines 106–116).

Comment 25: deleted “Inversion of the Mecca basin started and lasted beyond the early/mid Pleistocene (< 0.76 Ma).” lines 111–112. Also see response to comment 6.

Comment 26: replaced “Mecca Hills and Durmid Hills” by “the Coachella Valley” lines 112–113.

Comment 27: rewrote the section title into “Major tectonic culminations in the Coachella Valley” line 124.

Comment 28: added reference to Matti et al. (1985, 1992) and Matti and Morton (1993) lines 129–130 and Matti et al. (1992) and Matti and Morton (1993) to the reference list.

Comment 29: deleted “proto-SAFZ” lines 130–131 and replaced by “sedimentary”.

Comment 30: rewrote sentence lines 130–136 into “Northwest and west of the Coachella Valley, the Miocene–Pliocene sedimentary strata are structurally bounded by the San Bernardino and San Jacinto fault strands of the SAFZ (Bilham and Williams, 1985; Matti et al., 1985; Morton and Matti, 1993; Spotila et al., 2007). To the southwest, Miocene–Pliocene strata are bounded by the West Salton detachment fault (Dorsey et al., 2011).”. Also added Morton and Matti (1993) to the reference list.

Comment 31: see response to comment 30.

Comment 32: see response to comment 30.

Comment 33: see response to comment 30.

Comment 34: see response to comment 28.

Comment 35: replaced “fault gouge” by “trace” line 141.

Comment 36: deleted reference to Parrish (1983) throughout the manuscript (lines 35, 94, 144 and 154) and from the reference list. Also added reference to Dibblee and Mich (2008) line 94.

Comment 37: see response to comment 36.

Comment 38: replaced “Farther southeast” by “Southeast of the Indio Hills” line 150.

Comment 39: changed phrase lines 150–151 to “Gold et al. (2015) explore tectonogeomorphic evidence for dextral-oblique uplift and Keller et al. (1982) focus on landscape evolution”. Also added Gold et al. (2015) to the reference list.

Comment 40: moved reference to Blisniuk et al. (2021) together with Keller et al. (1982) they both studied landscape evolution in the Indio Hills.

Comment 41: replaced “Besides studying” by “In addition to investigating” line 153 and “their study” by “Keller et al. (1982)” lines 154–155.

Comment 42: added “(shallow)” and “(Fuis et al., 2012, 2017)” line 178.

Comment 43: rewrote lines 189–192 into “The Durmid Hills are an elongate ridge that parallels the main strand of the SAFZ at the south edge of the Salton Sea in Imperial Valley (Fig. 1). To the south, this deformation zone and the SAFZ project towards the Brawley seismic zone, an oblique, transtensional rift area with particularly high seismicity”.

Comment 44: deleted reference to Janecke et al. (2018) line 194 and added it lines 196 and 198.

Comment 45: added “both in the field and via imagery analysis” lines 215–216.

Comment 46: changed “when approaching” into “as they approach” line 234.

Comment 47: none.

Comment 48: replaced “they” by “the folds of the central macro-fold” line 254.

Comment 49: none.

Comment 50: deleted “with some confidence” line 300.

Comment 51: deleted parenthesis.

Comment 52: changed “The fold geometry is” by “Fold geometries are” line 329. Replaced “Its” by “The” line 330 and added “of these folds” line 331.

Comment 53: moved the following sentence “Brittle faults exist both in granitic basement and in sedimentary rocks of the Mecca and Palm Spring formations.” to the beginning of the paragraph lines 336–337.

Comment 54: extended the barbs to the northwest along the Indio Hills fault in Figure 2.

Comment 55: deleted “Rymer, unpublished data” line 340.

Comment 56: added “The fault sets in granitic basement rocks trend parallel to fault sets in sedimentary strata southeast of the Indio Hills fault (see stereoplots in Figure 2) and are therefore suggested to have formed due to similarly oriented stress.” lines 427–430.

Comment 57: added “Like the Indio Hills Fault, fault-plane dip and strike of the main San Andreas fault must be inferred indirectly.” lines 361–362.

Comment 58: none.

Comment 59: none because not allowed by the journal’s standards.

Comment 60: none.

Comment 61: changed “mud–silt–stone” into “[mudstone–siltstone](#)” line 356.

Comment 62: added “and thrust” to the title of the sub-section line 400 to better reflect the content.

Comment 63: see response to comment 56.

Comment 64: changed sentence to “In map view (Fig. 2), the folds are right-stepping, and each fold set is increasingly asymmetric (Z-shaped) and sigmoidal towards the Indio Hills fault in the northeast” lines 414–416.

Comment 65: deleted “, right-lateral” line 421.

Comment 66: deleted “due to distributed simple shear” lines 443–444.

Comment 67: added “(main San Andreas and Indio Hills faults)” line 430.

Comment 68: added “folds propagating outward from the Indio Hills fault is supported by the increased structural complexity of the fold geometries towards the Indio Hills fault.” lines 457–459, and deleted “, thus favoring folds propagating outward from the Indio Hills fault” lines 460–461.

Comment 69: added “, and/or to structural tilting in the hanging wall of the Indio Hills fault” lines 469–470.

Comment 70: replaced “convergent tectonic” by “transpressional uplift” line 564, and added “(i.e., a contractional uplift formed synchronously with successively with simple shear transpression to balance internal forces in a crustal-scale critical taper; Dahlen, 1990)” lines 564–566 and Dahlen (1990) to the reference list.

Comment 71: added a paragraph break line 449.

Comment 72: added reference to Figure 7 line 478.

Comment 73: labelled the parasitic folds in Figure 3C.

Comment 74: replaced reference to Schlische (1995) by Suppe and Medwedeff (1990) line and in the reference list.

Comment 75: split the sentence into two for clarity. Replaced “, which” by “. The fault” line 522.

Comment 76: added “In this section we use the geometry and kinematics of folds and faults in the southern Indio Hills to reconstruct the tectonic history of the area, not only of the inverted late Cenozoic basin but also about strike-slip and dip-slip faults that bound the basin. Essential tectonic events include (1) extensional normal faulting along the Indio Hills fault, (2) reactivation of the Indio Hills fault as a right-lateral to right-lateral-reverse fault, and (3) right-lateral movement along the main San Andreas fault.” Lines 539–544

Comment 77: see response to comment 80.

Comment 78: modified reference to Supplement S3b to Supplement S3d line 541.

Comment 79: see response to comments 80 and 156.

Comment 80: added “, by the deposition and preservation of sedimentary strata of the Palm Spring and Mecca formations southwest of the Indio Hills, whereas they were eroded or never deposited northeast of the fault, and by fining up upwards of the stratigraphic units from conglomerates in the Mecca Formation to coarse-grained sandstone in the lower parts of the Palm Spring Formation” lines 548–550. Added “In addition, the flat geometry of micro thrust faults (e.g., Supplements S3b–c) suggests that they were intensely rotated during macro-folding. Restoration of all micro faults in their initial position prior to macro-folding shows that some of these faults exhibit normal kinematics with associated syn-tectonic growth strata (Supplements S3d and S6).” lines 550–554. Added “basin geometry and formation similar to that of the Mecca Hills, where down-SW slip along the Painted Canyon fault was inferred in the (Miocene?–) Pliocene (McNabb et al., 2017), and of the transtensional Ridge Basin though with opposite vergence (Crowell, 1982; Ehman et al., 2000) with a” lines 556–558 and Crowell (1982) and Ehman et al. (2000) to the reference list. Added “during basin inversion” line 559. Also added “Formation of the Indio Hills fault probably occurred in mid-Miocene times during extension related to the opening of the Gulf of California (Stock and Hodges, 1989; Stock and Lee, 1994) as proposed for the Salton Trough (Dorsey et al., 2011 and references therein).” And Stock and Hodges (1989) and Stock and Lee (1994) to the reference list.

Comment 81: see response to comment 80.

Comment 82: deleted “, which gradually changed to a dominantly right-lateral-reverse fault” lines 562–563.

Comment 83: replaced “margin” by “convergent plate boundary” line 569.

Comment 84: added “during this period” line 572.

Comment 85: replaced “slip event” by “episode of movement along” line 573 and replaced “is clearly younger than the episode of” by “clearly postdates” line 574.

Comment 86: none commanded by the referee’s comment.

Comment 87: none.

Comment 88: added “(i.e., after the initial transpressional slip events along the Indio Hills fault).” lines 580–581.

Comment 89: none.

Comment 90: none.

Comment 91: split the sentence lines 584–587 into two and changed the second sentence to “. However, when restoring the sedimentary strata to horizontal (Supplement S6), the fault-related folds define a low-angle fold-and-thrust system”.

Comment 92: split the sentence lines 587–593 into two and added “. This implies” line 590.

Comment 93: added “(other than right-slip faults)” line 588.

Comment 94: added commas lines 602 and 603, added “upright” line 603, and changed “suggests” into “suggest” line 604.

Comment 95: replaced “layer-parallel” by “low-angle”. Added “These disharmonic folds are interpreted as intra-detachment folds.” lines 307–308. Added “intra-detachment” in figure 4 caption line 1126.

Comment 96: none.

Comment 97: added “In this section we use detailed structural analysis of folds and faults in the southeastern Indio Hills to outline the structural history of the tectonic culmination itself, evaluate it in terms of what is known about strain budgets within the southern San Andreas fault system, link it to nearby structures (Eastern California Shear Zone and Landers Mojave Line), and integrate the local structural history into a structural synthesis for the southern San Andreas Fault zone in the past 4 Myr.” lines 621–626. Added “Here we compare and contrast the structural evolution of the southeastern Indio Hills with that of nearby culminations (Mecca Hills and Durmid Hills).” lines 718–719.

Comment 98: replaced “Miocene” by “Pleistocene” line 629. Deleted “early” line 117. Replaced “mid” by “late” line 107. Deleted “Pliocene–” line 737. Deleted “Pliocene and” line 821.

Comment 99: see response to comment 80.

Comment 100: replaced “bounded” by “downthrew Miocene sedimentary strata against” lines 630–631. Added “in the footwall of the fault. In the hanging wall of the fault they were” lines 632–633.

Comment 101: none.

Comment 102: changed “master fault(s)” into “” lines 643–644.

Comment 103: changed the phrase lines 641–643 to “The fold set evolved oblique to the main San Andreas strand of the SAFZ and formed a right-stepping pattern of E–W-oriented axial surfaces that trend”.

Comment 104: see response to comment 102.

Comment 105: added “in between two active strike slip faults” line 533 and “consistently” line 534. Replaced “near the main San Andreas fault” by “in the southwest” line 537. Added “, where the macro-folds still display their initial non-plunging geometries” lines 757–758.

Comment 106: deleted “precursory” line 646.

Comment 107: none.

Comment 108: none.

Comment 109: added “strain” line 651.

Comment 110: added “from mostly distributed with minor partitioned deformation” lines 653–654.

Comment 111: added “right-lateral-“ line 656.

Comment 112: replaced “seems to have still played a minor role” by “did not yet play a major role” line 657.

Comment 113: rewrote “attenuation of the macro-folds toward the Indio Hills fault, increased shear folding, and clockwise rotation of fold axes to a steeper westerly plunge” into “tightening of the macro-folds toward the Indio Hills fault and clockwise rotation of fold axes to a steeper westerly plunge due to increased shear folding” lines 658–660.

Comment 114: replaced “buckle-” by “*en echelon* upright” line 660.

Comment 115: added “These folds were not observed northeast of the Indio Hills fault, nor southwest of the main San Andreas fault.” lines 250–251.

Comment 116: added “Furthermore, “ line 666.

Comment 117: added “in the Pliocene” line 634, “in the Pleistocene” line 638, “, i.e., probably in mid Pleistocene time” line 642, “in the mid Pleistocene” line 654, “in the late Pleistocene (earliest late Pleistocene 0.765 Ma Bishop Ash involved in folding; Sarna-Wojcicki et al., 2000; Zeeden et al., 2014)” lines 670–672, and “mid (–late?)” line 849.

Comment 118: added “kinematically” line 780.

Comment 119: added “see anticline closest to Indio Hills fault in” lines 678–679.

Comment 120: deleted “margin-parallel” and added “parallel to the convergent plate boundary” lines 679–680.

Comment 121: see response to comment 105.

Comment 122: replaced “latter” by “overlapping and synchronous formation of structures” line 623.

Comment 123: see response to comment 105.

Comment 124: see response to comment 5.

Comment 125: added “that a through-going NNW–SSE-striking fault crosscuts the Pinto Mountain fault (e.g., 1992 Joshua Tree earthquake near the study area) and” line 708–709, “its segment in the study area,” lines 709, “, NW–SE-striking” line 712, and “in the south” lines 7013–714.

Comment 126: added “. Notably, the 1992 Joshua Tree earthquake occurred along the NNW–SSE-striking, west-dipping West Deception Canyon fault (Rymer, 2000 and references therein), which merges with the Indio Hills fault in the south (see figure 1 in Rymer, 2000). Therefore, we propose” lines 708–711, and Rymer (2000) to the reference list.

Comment 127: see response to comment 12.

Comment 128: added “, i.e., a junction of three dextral fault branches” and “sensu” line 716, replaced “that” by “, which” line 717, reference to Passchier and Platt (2017) lines 717 and 721, and Passchier and Platt (2017) to the reference list.

Comment 129: deleted comma line 717.

Comment 130: added “late Pleistocene” lines 718–719.

Comment 131: added “, i.e., late Pleistocene” lines 724–725. Also see response to comment 117.

Comment 132: added “Our observations of mostly lateral movement along the main San Andreas fault (i.e., southeastern continuation of the Mission Creek fault) and the proposed Pleistocene age for deformation in the southeastern Indio Hills are consistent with work by Keller et al. (1982). A major difference between the northwestern and southeastern Indio Hills is the relatively higher amount and more intense character of deformation in between the two bounding faults in the latter with tighter macro-folding over a narrower area (Figs. 2 and 3; Keller et al., 1982; Lancaster et al., 2012).” lines 694–700.

Comment 133: rewrote the sentence into “in the southeasternmost Indio Hills and proceed as the main San Andreas fault” lines 746–747.

Comment 134: replaced “them” by “these faults” line 753.

Comment 135: none.

Comment 136: added “Another possible explanation may be the presence of coarse-grained deposits of the Mecca Formation, which may have partitioned/decoupled deformation along the Indio Hills fault from that in overlying Palm Spring sedimentary strata.” lines 758–761.

Comment 137: added reference to McNabb et al. (2017) line 184. Also see response to comment 80.

Comment 138: see response to comment 137.

Comment 139: see response to comment 137.

Comment 140: deleted “The increasing width of damage zones adjacent to SAFZ-related faults southward in Coachella Valley, and increased number of strike-slip and oblique to orthogonal cross faults in the Durmid Hills compared with Indio Hills and Mecca Hills may be due to closeness and transition to a transtensional rift setting around the Brawley seismic zone (Janecke et al., 2018).” lines 934–938. Replaced “one–two” by “several” line 545.

Comment 141: see response to comment 12.

Comment 142: see response to comment 117.

Comment 143: added “The Indio Hills fault probably initiated as a SW-dipping normal fault during the opening of the Gulf of California in the Miocene, and was later inverted as an right-lateral reverse, oblique-slip fault in the mid (–late?) Pleistocene due to transpression along the convergent plate boundary.” lines 843–846.

Comment 144: see response to comments 125 and 126.

Comment 145: added Lancaster et al. (2012) to the reference list.

Comment 146: see response to comment 145.

Comment 147: see response to comment 128.

Comment 148: see response to comment 12.

Comment 149: none.

Comment 150: see response to comment 73.

Comment 151: see response to comment 119.

Comment 152: added ages of the three tectonic phases to figures 7 and 8.

Comment 153: decreased length of Garnet Hill fault in figure 8.

Comment 154: none.

Comment 155: correct “Hills” into “Hill” line 1230.

Comment 156: see response to comment 80.

Revisions by the authors of the present manuscript based on editing comments in supplementary pdf by the referee

-Replaced “late” by “mid/upper” line 74 and added reference to Kirby et al. (2007) lines 76–77.

-Deleted “the” line 81.

- Replace “disconnected” by “segmented” line 81.
- Replaced “turns into” by “is succeeded by” line 87.
- Added “southwestwards” line 94.
- Deleted “was” and “and” and moved reference to Bergh et al. (2019) line 95.
- Deleted “Absolute dating revealed an” line 102.
- Deleted “and” and replaced by “are consistent with” line 105.
- Changed “1.0” to “0.76” lines 106 and 642 for the minimum age of the Palm Spring Formation for consistency with dating of the Bishop Ash by Sarna-Wojcicki et al. (2000).
- Added “-“ line 106.
- Added “We infer a similar age range for the Palm Spring Formation in the southern Indio Hills.” Lines 106–107.
- Replaced “emerges from the involvement” by “include tephrochronology”, and added “, which is involved in deformation” line 111.
- Moved “either” early in sentence and added “structures like” lines 142–143.
- Deleted “modern data remain scarce, and” line 146.
- Replaced “have not been published from this segment of the SAFZ” by “documenting this hypothesis for the culmination as a whole have not been conducted” lines 147–148.
- Added “, i.e., northwest of the study area” lines 152–153.
- Moved reference to Bergh et al. (2014, 2019) from line 168 to line 163.
- Replaced “exist” by “occur”.
- Deleted “whereas” and moved “northeast of the SAF” earlier in the sentence, and split sentence into two lines 189–190.
- Replaced “the present study” by “our investigation of the Indio Hills” line 202.
- Deleted “in the Indio Hills” line 204.
- Replaced “when” by “for” line 207.
- Replaced “with” by “and nearby” lines 208–209.
- Replaced “with” by “having” line 216.
- Changed “an approximately two kilometers wide zone” into “a zone approximately two kilometers wide” lines 217–218.
- Changed “with” into “showing” line 223.
- Deleted “in map view” line 229.

- Replaced “, whereas” by “. In contrast,” line 235.
- Deleted “in the southwest” line 236.
- Replaced “when approaching” by “as they approach” lines 245–246.
- Replaced “turn” by “transform” line 248.
- Changed the phrase lines 251–253 into “From southwest to northeast, the central macro-fold hinge zone displays a corresponding change in geometry”.
- Replaced “with” by “having” line 261.
- Changed “units” into “parts” line 265.
- Changed “the Mecca Formation conglomerates” into “conglomerates of the underlying Mecca Formation” lines 265–266.
- Added “(especially)” and moved “in relatively weak clayish–silty dark mudstone layers” earlier in sentence lines 267–268.
- Replaced “On the contrary” by “By contrast” line 270.
- Replaced “with” by “showing” line 273.
- Changed the phrase lines 276–278 to “Combined with a relatively narrow hinge zone, these attitudes define”.
- Changed phrase line 285 into “conglomerates of the underlying Mecca Formation”.
- Added “surfaces” line 285 and changed “is” into “are” line 286.
- Replaced “with” by “having” line 290.
- Replaced “units” by “strata” line 291.
- Changed phrase lines 297–298 into “the conglomerates of the Mecca Formation”.
- Replaced “until” by “to where” line 300.
- Replaced “almost” by “nearly” line 304.
- Added “strata of the” line 304.
- Replaced “with” by “showing” and “is” by “are” line 313.
- Changed phrase line 327 into “, where preserved, they display”.
- Changed sentence lines 335–338 into “Along the Indio Hills fault, poor exposures make it difficult to measure fault strike and dip directly, but DEM images suggest a rectilinear geometry in map view relative to the uplifted sedimentary strata to the southwest”.
- Replaced “possibly” by “probably” line 339.
- Moved “by the Banning fault” from line 350 to 349.

- Changed “when approaching” into “where they approach” lines 351–352.
- Replaced “impact” by “developed” line 355.
- Added “-“ line 361.
- Replaced “with” by “displaying” line 368.
- Replaced “operate together defining” by “appear to form” line 376.
- Replaced “formed” by “developed” line 377.
- Added “-“ lines 395 and 402.
- Changed “50 meters” into “50-m” line 403.
- Added “Structural” lines 412 and 471.
- Changed phrase lines 413–414 into “We mapped and analyzed three macro-scale fold systems that occur”.
- Changed the phrase lines 416–417 into “Based on these properties, we interpret the fold sets”.
- Changed the sentence lines 418–422 into “Various investigators (Babcock, 1974; Miller, 1998; Titus et al., 2007; Janecke et al., 2018; Bergh et al., 2019) describe similar fold geometries in sedimentary strata from many other segments of the SAFZ and are interpreted as structures formed by right-lateral displacement between two major fault strands due to distributed simple shear.”.
- Changed the phrase lines 422–423 into “However, the present fold-orientation data in the Indio Hills (Fig. 2)”.
- Replaced “e.g.,” by “compare with” line 432.
- Added “-“ line 447.
- Added “other” line 452.
- Replaced “hidden” by “blind” line 457.
- Deleted “(Fig. 3c)” line 478.
- Added “(NE-vergent)” line 482.
- Added “(not fully understood)” line 492.
- Replaced “with” by “having” line 521.
- Added “(and decreasing right-lateral)” line 522.
- Replaced “acted” by “ultimately functioned” line 525.
- Added “simultaneously” line 528.
- Replaced “However” by “In addition” line 531.
- Replaced “when approaching” by “towards” line 532.

- Deleted “also” line 533.
- Changed the phrase lines 536–537 into “spatial, temporal and kinematic” and deleted “in the Indio Hills” line 537
- Changed phrase lines 550–551 into “escaped from the mudstone beds and propagated”.
- Replaced “events with” by “, incorporating” line 573.
- Added “of the Indio Hills” line 610.
- Changed phrase lines 611–613 into “was accommodated by right-lateral-oblique, top-NE thrusting along the Indio Hills fault and major strike-slip movement”.
- Added “3c and” line 616.
- Changed the phrase lines 629–631 to “The right-lateral-reverse character of the Indio Hills Fault and its role in our kinematic model for basin inversion in the southern Indio Hills”.
- Replaced “East” by “Eastern” line 632.
- Replaced “simultaneously” by “coevally” line 699.
- Added “right-lateral” line 700.
- Rewrote “main SAFZ” into “main San Andreas fault” line 751.

Additional revisions by the authors of the present manuscript

- Added “(we refrain from using the name “Indio strand” given to this fault by Gold et al., 2015 to avoid confusion with the Indio Hills fault)” lines 39–41.
- Moved “Atwater and Stock, 1998;” before “Spotila et al., 2007;” line 64.
- Changed “uppermost members” into “upper member” line 118.
- Replaced “marks the” by “is a” line 201.
- Deleted “-“ line 480.
- Deleted en-dash line 506.
- Added “(probably southwest-dipping)” line 753.
- Replaced “show” by “suggest” line 786.
- Added a comma line 831.
- Corrected “Janecke et al., 2019” into “Janecke et al., 2018” line 842.
- Deleted “and main SAFZ” line 863.
- Moved “basement-seated” from line 899 to line 891.

-Deleted “The Indio Hills fault acted as a SW-dipping, normal fault in Miocene time, i.e., prior to inversion as an oblique-slip, right-lateral-reverse fault during mid (–late?) Pleistocene times” lines 899–901.

-Moved “, whereas the main San Andreas fault initiated probably as a dominantly right-slip fault during the later stages of uplift in the late Pleistocene.” from lines 903–904 to lines 894–896.

-Deleted “portion of the SAFZ” line 912.

-Deleted “in Durmid Hills” line 1235.

Tectonic evolution of the Indio Hills segment of the San Andreas fault in southern California, southwestern USA

Jean-Baptiste P. Koehl^{1,2,3,4}, Steffen G. Bergh^{2,3}, Arthur G. Sylvester⁵

1) Centre for Earth Evolution and Dynamics, (CEED), University of Oslo, N-0315 Oslo, Norway.

2) Department of Geosciences, UiT The Arctic University of Norway in Tromsø, N-9037 Tromsø, Norway.

3) Research Center for Arctic Petroleum Exploration (ARCEX), UiT The Arctic University of Norway.

4) CAGE – Centre for Arctic Gas Hydrate, Environment and Climate, UiT The Arctic University of Norway.

5) Department of Earth Science, University of California, Santa Barbara, USA.

Correspondence: jeanbaptiste.koehl@gmail.com

Abstract

Transpressional uplift domains of inverted ~~Miocene~~–Pliocene–Pleistocene basin fill along the San Andreas fault zone in Coachella Valley, southern California, are characterized by fault linkage and segmentation and deformation partitioning. The Indio Hills wedge-shaped uplift block is located in between two boundary fault strands, the Indio Hills fault to the northeast and the ~~Banning~~–main San Andreas fault to the southwest, which merge to the southeast. Uplift commenced about or later than 2.2–0.76 million years ago and involved progressive fold and faulting stages caused by a change from distributed strain to partly partitioned right-slip and reverse/thrust displacement on the bounding faults when approaching the fault junction. Major fold structures in the study area include oblique, right-stepping, partly overturned *en echelon* macro-folds that tighten and bend into parallelism with the Indio Hills fault to the east and become more open towards the main San Andreas~~Banning~~ fault to the west, indicating an early and close relationship of the macro-folds with the Indio Hills fault and a late initiation of the main San Andreas~~Banning~~ fault. Sets of strike-slip to reverse step-over and right- and left-lateral cross faults and conjugate kink bands affect the entire uplifted area, and locally offset the *en echelon* macro-folds. Comparison with the Mecca Hills and Durmid Hills uplifts farther southeast in Coachella Valley reveals notable similarities, but also differences in fault architectures, spatial and temporal evolution, and deformation mechanisms.

Introduction

This paper describes and evaluates structural patterns of the Indio Hills uplift in the northwestern part of Coachella Valley along the San Andreas Fault Zone (SAFZ) in

California, southwestern USA; (Fig. 1), where the fold–fault architecture, evolution, and partitioning of deformation compared to Mecca Hills and Durmid Hills are not well understood (e.g., Keller et al., 1982, ~~Parrish, 1983~~; Dibblee and Minch, 2008). The main goal of this study is to analyze internal macro- and meso-scale folds and related faults and to outline the kinematic evolution in relation to major SAFZ-related fault strands in the area (Fig. 1: Keller et al., 1982; Guest et al., 2007). These include the Indio Hills fault in the northeast (Allen, 1957; Tyley, 1974), and the main San Andreas~~Banning~~ fault along the southwest flank of the Indio Hills (we refrain from using the name “Indio strand” ascribed to this fault by Gold et al., 2015 to avoid confusion with the Indio Hills fault); ~~and of the thought to correspond to the main SAFZ in~~ Mecca Hills, and along the northeast flank of the Durmid Hills (Janecke et al., 2018); ~~and the Indio Hills fault in the northeast (Allen, 1957; Tyley, 1974; Fig. 1), which may merges with the Eastern California Shear Zone to the north and with the main San Andreas Banning fault in the southeast.~~ The progressive tectonic evolution model for the Indio Hills uplift is then compared and correlated with other major uplifts and SAFZ-related fault strands along strike in the Mecca Hills and Durmid Hills (Sylvester and Smith, 1987; McNabb et al., 2017; Janecke et al., 2018; Bergh et al., 2019). We also discuss briefly the potential northwestward continuation of the Indio Hills fault into the East California Shear Zone and its role as possible transfer fault (Dokka and Travis, 1990a, 1990b; Thatcher et al., 2016). The variable fault and fold architectures and associated ongoing seismic activity in these uplift areas underline the need for persistent along-strike studies of the SAFZ to characterize the fundamental geometry, resolve the kinematic development, and correlate regionally major fault strands (cf. Janecke et al., 2018). Such studies are essential to explain the observed lateral variations in fold and fault architectures and to resolve mechanisms of transpression, fault linkage, and areal segmentation in continental transform settings.

Geological setting

The Coachella Valley segment of the SAFZ in southern California is expressed as ~~three uplifted~~multiple, right-lateral, ~~transpressional domains~~ fault strands, which uplifted blocks~~located~~ in the Indio Hills, Mecca Hills, and Durmid Hills (Fig. 1; Sylvester, 1988). These domains comprise thick successions of ~~Pliocene–Pleistocene~~Pliocene–Pleistocene ~~Miocene–Pliocene~~ sedimentary strata uplifted and deformed in Pleistocene–Holocene time due to oblique convergence of the Pacific and North American plates and ~~transform~~ movements along the SAFZ and related faults (e.g., Atwater and Stock, 1998; Spotila et al., 2007; ~~Atwater and~~

~~Stock, 1998~~; Dorsey et al., 2011). Recent structural studies in the Mecca Hills (McNabb et al., 2017; Bergh et al., 2019), and Durmid Hills at the southern termination of the SAFZ (Janecke et al., 2018), show that individual fault strands are linked, and that the deformation splits into abruptly changing fold and fault geometries (Fuis et al., 2012, 2017). These recent works call for further characterization of the understudied Indio Hills segment in order to compare its structural development with other uplifted features along a major transform plate boundary fault zone. Below we summarize local and regional fault nomenclature, distribution, and fault movement history (Table 1) throughout the greater Coachella Valley region (Fig. 1), the stratigraphy of the Indio Hills area, and previous structural work in the main Indio Hills, Mecca Hills, and Durmid Hills uplift areas.

Regional faults

The southeastern Indio Hills are a WNW–ESE–trending tectonic ~~uplift~~ ~~culmination~~ situated in a small restraining bend northeast of the main San Andreas fault (Figs. 1 and 2 and Supplement S1). The studied ~~culmination~~ ~~uplift~~ is located along strike about 25–50 kilometers northwest of the Mecca Hills and Durmid Hills, and to the southeast of the major left bend in the SAFZ trace near San Gorgonio Pass (Matti et al., 1985, 1992; Matti and Morton, 1993; Dair and Cooke, 2009).

The main faults in the southeastern Indio Hills include the Indio Hills fault in the northeast (Allen, 1957; Tyley, 1974), and the main San Andreas fault in the southwest. Regionally, the Indio Hills fault possibly merges with the Landers Mojave Line and the eastern California shear zone in the north (Dokka and Travis, 1990a, 1990b; Nur et al., 1993a, 1993b; Thatcher et al., 2016). The Landers Mojave Line is believed to be the locus of several recent earthquakes aligned along a NNW–SSE-trending axis, including the 1992 Joshua Tree earthquake (Fig. 1b; Nur et al., 1993a, 1993b). These earthquakes were tentatively ascribed to movement along a through-going NNW–SSE-striking fault, possibly the west-dipping, Quaternary West Deception Canyon fault (Sieh et al., 1993; Rymer, 2000). This fault is thought to crosscut the E–W- to ENE–WSW-striking, left-lateral, Holocene Pinto Mountain fault, which merges with the main strand of the San Andreas fault in the west at the intersection of the right-lateral Mission Creek and Mill Creek strands (Allen, 1957; Bryant, 2000; Kendrick et al., 2015; Blisniuk et al., 2021). The former is thought to correspond to the continuation of the main San Andreas fault to the northwest (Gold et al., 2015) and may have accommodated ca. 89 km of right slip in the past 4 million years, whereas the latter

accommodated about 8 km right slip at 0.5–0.1 Ma and is offset ca. 1 km by the Pinto Mountain fault (Kendrick et al., 2015).

The main San Andreas fault continues to the southeast where it bounds the Mecca Hills to the southwest, whereas the Painted Canyon fault, a previous (late Miocene?)–Pliocene southwest-dipping normal fault reactivated as a right-lateral-reverse oblique-slip fault in the Pleistocene–present-day bounds the Mecca basin to the northeast (Sylvester and Smith, 1987; McNabb et al., 2017; Bergh et al., 2019). Farther southeast, the main San Andreas fault proceeds along the northeast flank of the Durmid Hills opposite the Pleistocene (ca. 1 Ma), right-lateral East Shoreline fault (Babcock, 1969, 1974; Bürgmann, 1991; Janecke et al., 2018). There, the main San Andreas fault merges with the Brawley seismic zone (Lin et al., 2007; Hauksson et al., 2012; Lin, 2013) and, together with the right-lateral San Jacinto fault zone, they merge into the right-lateral Imperial fault (Rockwell et al., 2011). In the north, the main San Andreas fault splays into the Banning strand and the Mission Creek fault in the northwestern part of the Indio Hills (Keller et al., 1982; Gold et al., 2015). The Banning strand is much younger than the Mission Creek fault and may have accommodated approximately 3 km of right slip in the past 0.1 million years (Kendrick et al., 2015).

Northwest and west of the Coachella Valley, Miocene–Pleistocene sedimentary strata are structurally bounded by the San Bernardino and San Jacinto fault strands of the SAFZ (Bilham and Williams, 1985; Matti et al., 1985; Morton and Matti, 1993; Spotila et al., 2007). To the southwest, Miocene–Pleistocene strata are bounded by the West Salton Detachment fault (Dorsey et al., 2011). The San Jacinto fault is typically believed to have slipped ca. 25 km right-laterally in the past 1.5 million years (Matti and Morton, 1993; Kendrick et al., 2015), whereas the West Salton Detachment fault is a low-angle normal fault that accumulated ca. 8–10 km of normal-oblique movement starting in the mid Miocene and is related to the opening of the Gulf of California (Prante et al., 2014 and references therein).

Stratigraphy of the Indio Hills and adjacent areas

The Indio Hills ~~culmination-uplift~~ is an inverted ~~Pliocene–Pleistocene~~~~Miocene–Pliocene~~ sedimentary basin lying upon Mesozoic granitic basement rocks, which we regard as ~~an~~ analogous to the ~~inverted~~ Mecca ~~rift~~~~inverted~~ basin farther southeast (Keller et al., 1982; Damte, 1997; McNabb et al., 2017; Bergh et al., 2019). In the Mecca basin, alluvial, fluvial and lacustrine deposits of the Mecca and Palm Springs formations are truncated unconformably by the ~~mid to upper~~~~late~~ Pleistocene–~~Quaternary~~ Ocotillo Formation (Dibblee, 1954; Sylvester and Smith, 1976, 1987; Boley et al., 1994; Rymer, 1994; Sheridan et al.,

1994; Sheridan and Weldon, 1994; Winker and Kidwell, 1996; Kirby et al., 2007; McNabb et al., 2017; Table 1). Similar uplifted strata at Durmid Hills (Fig. 1) belong to the Pliocene–Pleistocene Borrego Formation, and are overlain by mid to upper Pleistocene deposits of the Brawley and Ocotillo formations (Dibblee, 1997; Herzig et al., 1988; Lutz et al., 2006; Kirby et al., 2007; Dibblee and Minch, 2008).

Leuco-granitic basement rocks crop out near gently SW-dipping conglomerates along the northeastern flank of the Indio Hills, near the trace of the Indio Hills fault (Fig. 2). Despite the proximity of the conglomerates with disconnected-segmented granite outcrops, the contact itself is not exposed. The conglomerates are the lowermost stratigraphic unit exposed in the Indio Hills and are characterized by a succession of meter-thick beds of very coarse, poorly sorted blocks of gneissic and granitic basement rocks more than a meter in size. Previous mapping in the area (Dibblee, 1954; Lancaster et al., 2012) We considered the conglomerates as stratigraphic equivalents to the Miocene–mid to upper Pliocene Mecca Formation in the Mecca Hills (Dibblee, 1954; Sylvester and Smith, 1987; McNabb et al., 2017; Bergh et al., 2019) and that at least part of the clasts are from the leuco-granitic rocks, which must correspond to basement rocks of the inverted Indio Hills basin. Up-section toward the southwest the conglomerate gradually turns into is succeeded by coarse-grained sandstone, which defines the transition from the Mecca Formation to the lower Palm Spring Formation.

The Palm Spring Formation in the Indio Hills consists of moderately- to well-consolidated alluvial fan deposits (Dibblee and Minch, 2008; Parrish, 1983), with some interbedded gypsum layers and red-colored calcareous mudstone, as in the Mecca Hills (Sylvester and Smith, 1987). The main rock types include beds of light-colored, medium- to coarse-grained sandstone, gray–brown silty sandstone, and dark biotite-rich mudstone. The southwestwards increase in silt–clay toward the main San Andreas Banning fault was (also recorded in the Mecca Hills; Bergh et al., 2019) and may indicate a gradual transition from the lower to the upper member of the Palm Spring Formation (Bergh et al., 2019).

By contrast, the transition between the lower and upper members of the Palm Spring Formation in the Mecca Hills is marked by two angular unconformities that signals further steps in uplift and inversion of the Mecca basin (Table 1; McNabb et al., 2017; Bergh et al., 2019). In the Indio Hills, however, the nature of the transition between the lower and upper member of the Palm Spring Formation and the presence of an angular unconformity is unknown.

Absolute dating revealed an ages of 3.07–2.36 Ma (mid-late Pliocene–latest Pliocene–early Pleistocene) and 2.68–1.07 Ma (late Pliocene–mid Pleistocene–earliest Pleistocene to

earliest late Pleistocene), were obtained respectively for the lower and upper member of the Palm Spring Formation, respectively, in the Mecca Hills, based on reversed magnetic polarity data (Chang et al., 1987; Boley et al., 1994;), and re-consistent with sediment accumulation rate estimates (McNabb, 2013; McNabb et al., 2017; Table 1). We infer a similar age range for the Palm Spring Formation in the southern Indio Hills. Inversion of the Mecca basin started and lasted beyond the early/mid Pleistocene (<0.76 Ma).

In contrast to other uplift areas in Coachella Valley, the Ocotillo Formation has not been mapped in the Indio Hills in the present study. However, based on the occurrence of the Bishop Ash at the northwestern edge of the study area and on the occurrence of the volcanic deposit within the uppermost Palm Spring Formation or at the base of the overlying Ocotillo Formation in the Mecca Hills, it is likely that the Ocotillo Formation is present just northwest of the area mapped (Fig. 2). In addition, it is deposited on the flank northeast of the Indio Hills fault, and southwest of the main San Andreas fault (Figs. 1 and 2), indicating that this unit was either not deposited or eroded in the area that recorded the most uplift in Indio Hills.

Additional dating limits constraints on the transpressional uplift in Mecca Hills and Durmid Hills the Coachella Valley emerges include from tephrochronology the involvement of the 0.765 million year old Bishop Ash layer (Sarna-Wojcicki et al., 2000; Zeeden et al., 2014; Table 1). This volcanic deposit, which is found involved in deformation within the uppermost members of the Palm Spring Formation (which is unconformably overlain by the Ocotillo Formation) in the hanging wall of the Painted Canyon fault away from the fault, and within the base of the Ocotillo Formation in the hanging wall of the Painted Canyon fault near the fault (Ocotillo and uppermost Palm Spring formations interfingering near the fault) and in the footwall of the fault (McNabb et al., 2017; Bergh et al. 2019; Janecke et al., 2018). The unconformable contact between the Palm Spring and Ocotillo formations away from the Painted Canyon fault towards the southwest and their interfingering relationship near the fault suggest that uplift had already initiated prior to deposition of the Ocotillo Formation (i.e., before 0.76 Ma, in the mid Pleistocene), possibly during the formation of the lower unconformity between the lower and upper members of the Palm Spring Formation (McNabb et al., 2017; Table 1). Complementarily, the involvement of the Bishop Ash in deformation suggest that deformation continued past 0.76 Ma (in the late Pleistocene).

In contrast to other uplift areas in Coachella Valley, the Ocotillo Formation has not been mapped in the Indio Hills, but rather is deposited on the flank northeast of the Indio Hills fault, and southwest of the main San Andreas Banning fault (Figs. 1 and 2), indicating that the Ocotillo Formation was either not deposited, or eroded in the area of uplift.

Major tectonic uplifts/culminations in the Coachella Valley

Indio Hills

The Indio Hills are a WNW–ESE trending tectonic culmination situated in a small restraining bend northeast of the main SAFZ trace (Figs. 1 and 2). The culmination is located along strike about 25–50 kilometers northwest of the Mecca Hills and Durmid Hills, and to the southeast of the major left bend in the SAFZ trace near San Geronio Pass (Dair and Cooke, 2009). The Miocene–Pliocene proto-SAFZ strata are structurally bounded north of the Coachella Valley by a low topographic relief SAFZ segment and several left-slip splay faults that merge into the uplifted San Bernardino and San Jacinto fault strands (Bilham and Williams, 1985; Spotila et al., 2007), and the West Salton detachment fault in the southwest (Dorsey et al., 2011).

The southeastern end of the Indio Hills is an uplifted domain of deformed strata of the Mecca and Palm Spring formations situated in between the main San Andreas Banning and Indio Hills fault (Fig. 2). The main San Andreas Banning fault corresponds to a major oblique strike-slip fault segment at the eastern end of San Geronio Pass (Matti et al., 1985; Morton et al., 1987), and it is easily traced to Indio Hills (Figs. 1 and 2) since its main fault gouge trace provides preferential pathways for ground water flow and growth of wild palm trees along strike.

The Indio Hills fault was mapped north of the study area (Parrish, 1983; Dibblee and Minch, 2008) extending into the Landers–Mojave Line (Nur et al., 1993a, 1993b), a NNW–SSE-striking right-lateral fault system extending hundreds of kilometers northward from the southeastern Indio Hills into the East California Shear Zone and related fault segments such as the Calico and Camp Rock faults (Fig. 1; Dokka et al., 1990a; Nur et al. 1993b). The Indio Hills fault may correspond to a major fault splay of the SAFZ (Dokka and Travis, 1990a, 1990b; Thatcher et al., 2016). Farther-southeast of the Indio Hills, however, the attitude and geometry of the Indio Hills fault remains elusive, and the fault either dies out or merges either with structures like the main San Andreas Banning fault, the Skeleton Canyon fault, and/or the Painted Canyon fault in the Mecca Hills (Fig. 1).

The transpressional character of the Indio Hills uplift was suggested by Parrish (1983) and Sylvester and Smith (1987), but However, modern data remain scarce, and detailed structural analyses have not been published from this segment of the SAFZ documenting this hypothesis for the culmination uplift as a whole have not been conducted. Gold et al. (2015) explore tectonogeomorphic evidence for dextral-oblique uplift and Keller et al. (1982) and

[Blisniuk et al. \(2021\)](#) focus on landscape evolution. An exception is the study of Keller et al. (1982) focusing on an area northwest of our study area and aimed at investigating the tectonic geomorphology near the intersection of the Banning [strand](#) and Mission Creek faults (northwest of the study area), which merge into the main San Andreas fault (Fig. 1; [Blisniuk et al., 2021](#)). Besides studying [In addition to investigating](#) soil profiles, offset drainage systems, and recent (a few thousand years old) displacement along the SAFZ, ~~their~~ Keller et al. (1982) study called attention to a strong dominance of gently plunging and upright macrofolds in bedrock strata along the Mission Creek fault and at the southeastern end of the Banning ~~fault~~ [strand](#) where these faults merge. Their study showed that bends and steps along the main fault traces were consistently located near brittle fault segments and zones of uplift. [The study also showed that drainage systems were offset recently \(at ca. 0.03–0.02 Ma\) and indicate relatively high slip rates along the Mission Creek fault in the order of 23–35 cm.y⁻¹, i.e., comparable to the more recent c. 23 cm.y⁻¹ estimate by Blisniuk et al. \(2021\).](#)

Mecca Hills

Farther south, the Mecca Hills uplift was previously defined as a classic flower-structure (Sylvester and Smith, 1976, 1987; Sylvester, 1988), in which all folds and faults formed synchronously and merged at depth. Recent analyses ([Bergh et al., 2014, 2019](#)) indicate that a modified flower-like structure, consisting of a steep SAFZ fault core zone to the southwest, a surrounding ~~off fault~~ approximately one–two kilometers wide damage zone expressed by *en echelon* folds and faults oblique to the SAFZ (including left-slip cross faults), steeply plunging folds, and SAFZ-parallel fold and thrust belt features (including right- and left-slip and oblique-reverse faults) formed in kinematic succession (~~Bergh et al., 2014, 2019~~). In addition to the steep [\(shallow\)](#) SAFZ ([Fuis et al., 2012, 2017](#)), two other, major NW–SE-striking faults ~~exist-occur~~ in the Mecca Hills (Fig. 1). One is the Skeleton Canyon fault, which initiated as a steep SAFZ-parallel strike-slip fault and was reactivated as a reverse and thrust fault dipping gently northeastwards in the late kinematic stages ([Sylvester and Smith, 1976, 1979, 1987; Bergh et al., 2019](#)). The other is the Painted Canyon fault, which ~~marks the~~ [is a](#) former Miocene–Pliocene basin-bounding normal fault ([McNabb et al., 2017](#)) and is now reactivated as a NE-directed thrust fault with dip to the southwest (Bergh et al., 2019; [Table 1](#)). The polyphase evolution and reactivation of internal oblique, step-over faults, and SAFZ-parallel faults, were explained by a series of successive–overlapping events involving a change from distributed, locally partitioned, into fully partitioned strain in a changing, oblique-plate convergence regime (Bergh et al., 2019).

Durmid Hills

The Durmid Hills are an elongate ridge that parallels the main strand of the SAFZ at the south edge of the Salton Sea in Imperial Valley (Fig. 1). ~~Farther south, this deformation zone and the SAFZ is aligned~~ project towards to the south with the Brawley seismic zone, an oblique, transtensional rift area with particularly high seismicity (Lin et al., 2007; Hauksson et al., 2012; Lin, 2013). The main San Andreas fault-strand (mSAF) is located on the northeast side of the Durmid Hills (~~Janecke et al., 2018~~) and has been thoroughly studied (Dibblee, 1954, 1997; Babcock, 1969, 1974; Bilham and Williams, 1985; Bürgmann, 1991; Sylvester et al., 1993; Lindsey and Fialko, 2013; Janecke et al., 2018). The rocks southwest of the mSAF consist of highly folded Pliocene–Pleistocene deposits (Babcock, 1974; Bürgmann, 1991; Markowski, 2016; Janecke et al., 2018) bounded to the southwest by the subsidiary East Shoreline Fault strand of the SAFZ. Northeast of the mSAF, ~~whereas~~ the formations are much less deformed ~~northeast of the mSAF~~ (Janecke et al., 2018). The overall structure (Fig. 1) resembles a right-lateral strike-slip duplex (Sylvester, 1988), but the geometry is not fully consistent with a duplex model due to abundant left-lateral cross faults and internal block rotations. Instead, the Durmid Hills structure was interpreted as a ladder structure (Janecke et al., 2018), as defined by Davis (1999) and Schulz and Balasko (2003), where overlapping, E–W- to NW–SE-striking step-over faults rotated along multiple connecting cross faults. The one–three kilometers wide Durmid ladder structure consists of multiple internal, clockwise-rotating blocks bounded by major *en echelon* folds and right- and left-lateral cross faults in between the right-slip mSAF and Eastern Shoreline ~~f~~Fault strand, indicating a complex termination of the SAFZ around the Brawley Seismic Zone to the southeast (Fig.1).

Methods and data

In ~~the present study~~ your investigation of the Indio Hills, we used high-resolution Google Earth DEM images and aerial photographs (© Google Earth 2011) as a basis for detailed field and structural analyses ~~in the Indio Hills~~ (Fig. 2). We mapped and analyzed individual macro- and meso-scale folds and associated faults in Miocene–Pliocene strata both in the field and via imagery analysis. Key horizons of light-colored quartz sandstone and carbonate rocks in the Palm Spring Formation provide structural markers, notably ~~when for~~ restoring bed offsets and fault–fold geometries and kinematics. We address crosscutting relations of the Banning–main San Andreas and Indio Hills faults ~~with and nearby~~ fold structures. Structural orientation data are obtained from meso-scale folds and faults and are

integrated between the areal segments to link a prevalent pattern of deformation into a wider structural architecture (Fig. 2).

Results

Structural overview of the Indio Hills

The study area comprises three major, SAFZ-oblique, asymmetric, E–W-trending, moderately west-plunging fold systems ~~with having~~ multiple smaller-scale parasitic folds (Fig. 2). The main folds affect most of the Palm Spring Formation in a ~~an~~ zone approximately two kilometers wide ~~zone~~ between the ~~main San Andreas~~Banning and Indio Hills faults (Fig. 2). The northeastern flank of the Indio Hills is structurally different by consisting of a sub-horizontal, NW–SE-trending, open, upright anticline, which trends parallel to the Indio Hills fault (Fig. 2). Similarly, close to the ~~main San Andreas~~Banning fault, tilted strata of the Palm Spring Formation are folded into a tight, steeply plunging shear fold (folds involving shearing along a plane that is parallel to subparallel to the fold’s axial plane; Groshong, 1975; Meere et al., 2013; Fig. 2). At smaller scale, several subsidiary reverse faults and mostly right-slip, step-over faults ~~with having~~ orientations both parallel with (E–W to NW–SE) and perpendicular (NNE–SSW) to the bounding faults exist within the macro-folded domain. Most of these faults truncate individual SAFZ-oblique folds.

SAFZ-oblique macro-folds

SAFZ-oblique macro-folds are consistently asymmetric and mostly south-verging, and their axial surfaces are arcuate and right-stepping ~~in map view~~ (Fig. 2). Fold geometries change from open and nearly upright near the ~~main San Andreas~~Banning fault, ~~via to~~ kink/chevron styles in the middle part, to very tight (isoclinal) and overturned fold styles adjacent to the Indio Hills fault (Fig. 3a–c and Supplement S2a–c). These changes in geometry correspond to a change in obliquity of the fold axial surface trace from approximately 60–70° to less than 20° with the Indio Hills fault (Fig. 2). All three macro-folds have axial trends that bend and partly merge into parallelism with the Indio Hills fault. ~~In contrast, whereas~~ moderate to steeply WSW-dipping strata of the Palm Spring Formation are obliquely truncated by the ~~main San Andreas~~Banning fault ~~in the southwest~~. Tighter fold hinges are mapped in the central macro-fold and on the back-limb (stretched long limb in an overturned fold) of the Z-shaped, southeastern macro-fold (Fig. 2). These folds were not observed northeast of the Indio Hills fault, nor southwest of the main San Andreas fault.

Northwestern and central macro-folds

The northwestern and central macro-folds define two major, compound and arcuate fold systems that affect the entire Palm Spring Formation between the main San Andreas and Indio Hills faults (Fig. 3a–b). They consist of eight subsidiary Z- and S-shaped, south-verging anticline-syncline pairs, and show fold axes plunging variably but mostly about 30° to the west (Fig. 2). At large scale, both folds tighten northeastward and display clockwise bend of axial traces from ENE–WSW near the main San Andreas fault, to E–W and NW–SE ~~when-as they~~ approaching the Indio Hills fault (Fig. 2 and 3c). Fold hinges in the west are typically symmetric, concentric, and open (Supplement S34a–b), and become gradually tighter and dominantly Z-shaped kink folds eastward (Supplement S34c). The folds ~~turn~~ transform into tight, isoclinal, and inverted geometries (Supplement S34d–e) when approaching the central macro-fold back-limb (Fig. 3b), and they potentially merge with the SAFZ-parallel anticline less than 200 meters from the Indio Hills fault (Fig. 2). From southwest to northeast, A the central macro-fold hinge zone displays a corresponding change in ~~the geometry of the central macro-fold hinge zone is observed~~ northeastward, i.e., from symmetric, ~~via-to~~ kink/chevron, and to isoclinal overturned styles (Supplement S24a–b), until they folds of the central macro-fold flank the back-limb of the southeastern macro-fold (Supplement S24c–d). Bedding surfaces on the fore-limb (the shortened, inverted limb indicating the direction of tectonic transport in an overturned fold) of the central macro-fold dip steeply or are inverted, whereas strata on the back-limb mostly dip gently to the north or northwest, i.e., at a high angle to the bounding faults, and gradually change to northward dip when approaching the Indio Hills fault (Fig. 3c).

Another feature of the central macro-fold is that it is offset by a system of both layer-parallel and bed-truncating faults (Fig. 3b). Strata east of the fault system are affected by a large shear fold ~~with~~ having thickened hinges and thinned limbs. The next fold to the north-northeast changes from open to tight, overturned, and locally isoclinal (Supplement S24a–c), and merges with the inverted, NE-dipping back-limb of the southeastern macro-fold (Fig. 3c). Notably, the consistent eastward tightening of fold hinges occurs within the lower stratigraphic ~~units-parts~~ of the Palm Spring Formation, whereas conglomerates of the underlying Mecca Formation ~~conglomerates~~ are only weakly folded (see section about the southeastern macro-fold). Furthermore, beds in tighter folds (especially in relatively weak clayish-silty dark mudstone layers) are commonly accompanied by disharmonic folds and internal structural disconformities ~~in relatively weak clayish-silty dark mudstone layers. On~~ the ~~By~~ contrast, more rigid, and thicker sandstone beds are more commonly fractured.

Southeastern macro-fold

The southeastern macro-fold is expressed as a kilometer-wide, Z-shaped, open to tight, south-verging syncline-anticline pair ~~with showing~~ moderately west-plunging axes and steeply north-dipping axial surfaces (Fig. 3c). Most of the Palm Spring Formation strata on the back-limb trend parallel to the Indio Hills fault and dip about 50–70° to the north, whereas strata in the hinge and fore-limb dip about 40–70° to the west/southwest (Fig. 3c). ~~Combined with a relatively narrow hinge zone, T~~hese attitudes ~~combined with a relatively narrow hinge zone classify define~~ the southeastern macro-fold as a chevron type. The axial trend of the syncline-anticline pair is at a low angle ($< 20^\circ$) to the Indio Hills fault but bends into a NE–SW trend westward with a much higher (oblique) angle to the ~~main San Andreas~~~~Banning~~ fault, which cuts off the fore-limb strata (Fig. 2). The south~~we~~astern macro-fold is very tight in the north and east and has several smaller-scale, tight to isoclinal, strongly attenuated folds on the main back-limb that merge from the central macro-fold, thus indicating increasing strain intensity northeastward (see discussion). In contrast to the tightly folded beds of the Palm Spring Formation, bedding ~~surfaces~~ in ~~conglomerates of~~ the underlying Mecca Formation ~~conglomerate is are~~ only weakly folded northeastward and becomes part of the open ~~to monocline-like~~ SAFZ-parallel anticline close to the Indio Hills fault.

A macro-folded siltstone layer of the lower Palm Spring Formation more than 200 meters southwest of the Indio Hills fault (Fig. 4a) contains centimeter-scale, upright (sub-horizontal) and disharmonic folds ~~with having~~ E–W trend and western plunge (Fig. 4b). These intra-layer folded ~~units-strata~~ are cut by low-angle reverse faults yielding a NE-directed sense-of-shear. The upright geometry and the sub-horizontal fold axes (about 5° plunge) of these intra-bed minor folds differ from the SAFZ-oblique folds but resemble those of the macro-scale, SAFZ-parallel NW–SE-trending anticline near the Indio Hills fault. ~~These disharmonic folds are interpreted as intra-detachment folds (see disucssion).~~

SAFZ-parallel macro-folds

About 100–200 meters southwest of the trace of the Indio Hills fault, the ~~conglomerates of the~~ Mecca Formation ~~conglomerate is are~~ folded into a major open anticline, whose axis is parallel to slightly oblique ($< 20^\circ$) to the Indio Hills fault. This macro-fold is traceable ~~with some confidence~~ northwestward ~~until to where~~ the Indio Hills fault bends northward (Fig. 1). The southwestern limb of the fold marks the transition from the Mecca Formation conglomerate with the overlying Palm Spring Formation on the back-limb of the southeastern and central macro-folds (Fig. 2 and Supplement S24c). The conglomerate beds

are thicker, ~~almost-nearly~~ unconsolidated, and much less internally deformed than the ~~strata of~~
~~the~~ Palm Spring Formation ~~strata~~. The major anticline displays an open, symmetric, partly
box-shaped, NW–SE-trending, upright geometry with 2–3° plunge of the fold axis to the
northwest. Outcrops on the SW-dipping limb of the anticline (Fig. 3c) are cut by a SW-
dipping reverse fault system that is ~~(sub-)~~parallel to the Indio Hills fault (Supplement S35a).
These reverse faults may be linked with the reverse fault in folded strata of the Palm Spring
Formation on the southeastern macro-fold back-limb described above (Fig. 4). The upright
geometry and sub-horizontal NW–SE-trending axes of related small-scale folds in a mudstone
layer (Fig. 4) resembles that of the SAFZ-parallel anticline.

A couple of major ~~fold-synclines~~ ~~with-showing~~ axial traces parallel to the ~~main San~~
~~Andreas~~Banning fault ~~is-are~~ also well displayed on DEM images (Fig. 5 ~~and Supplement S6~~).
These folds affect WSW-dipping strata of the Palm Spring Formation on the broadened
western part of the northwest and central macro-folds. ~~F~~The fold geometries ~~ies isare~~ tight and
asymmetric, with wavelengths less than 200 meters, and presumably steep NW-plunging
axes. ~~Its-~~The local appearance and sheared geometry ~~of these folds~~ contrast both with the
broad SAFZ-oblique folds near the ~~main San Andreas~~Banning fault, and with that of the
upright, SAFZ-parallel anticline near the Indio Hills fault.

Major and minor ~~fold-related~~ faults

~~Brittle faults exist both in granitic basement and in sedimentary rocks of the Mecca~~
~~and Palm Spring formations.~~ Fold-related brittle faults ~~exist both in granitic basement and in~~
~~sedimentary rocks of the Mecca and Palm Spring formations~~ in the study area. ~~Such faults are~~
~~mostly display~~ narrow damage zones less than one meter wide and are geometrically either
related to SAFZ-oblique or SAFZ-parallel macro- and meso-scale folds, or are orthogonal to
the SAFZ and related faults. ~~Brittle faults exist both in granitic basement and in sedimentary~~
~~rocks of the Mecca and Palm Spring formations.~~ With exception of the main ~~San~~
~~Andreas~~Banning and Indio Hills faults, brittle faults are generally difficult to trace laterally
but, ~~arewhere~~ preserved, ~~in places they display with~~ centimeter- to meter-scale strike-slip
and/or reverse dip-slip ~~offset~~displacement. Large-scale fault orientations and kinematics in
sedimentary rocks are more variable than in basement rocks, but strike commonly WNW–
ESE to N–S and show moderate–steep dips to the northeast (Fig. 2). Subsidiary meso-scale
faults include high-angle SW- and SE-dipping strike-slip faults, and low-angle SW-dipping
thrust faults. We describe the Indio Hills and ~~main San Andreas~~Banning faults, strike-slip

faults, and thrust faults in sedimentary strata, and fractures in basement rocks northeast of the Indio Hills fault.

Indio Hills and main San Andreas Banning faults

Along the Indio Hills fault, poor exposures make it difficult to measure fault strike and dip directly ~~Direct field observations of the strike and dip of the Indio Hills fault were not possible~~, but DEM images suggest a rectilinear geometry in map view relative to the uplifted sedimentary strata to the southwest (Fig. 2). The fault strikes mainly NW–SE and is subparallel to the northeastern flank of the Indio Hills. Farther southeast, it ~~possibly-probably~~ merges with the main San Andreas Banning fault (Fig. 1; Tyler, 1974; ~~Rymer, unpublished data~~). In the southeastern part of the study area (Fig. 2), the Indio Hills fault is most likely located between an outcrop of basement ~~leuco~~-granite and the first outcrops of overlying strata of the Palm Spring Formation. The granite there is highly fractured and cut by vein and joint networks (see description below), as may be expected near-in the damage zone of a major brittle fault.

Like the Indio Hills Fault, fault-plane dip and strike of the main San Andreas fault must be inferred indirectly. The main San Andreas Banning fault in the study area strikes WNW–ESE and is sub-vertical based on its consistent rectilinear surficial trace, and because it truncates both back- and fore-limb strata on most of the SAFZ-oblique macro-folds (Fig. 2). Thus, the main San Andreas Banning fault does not seem to have had major impact on the initial geometry and development of the macro-folds in the Indio Hills. However, notable exceptions include displacement by the main San Andreas fault of the two shear folds on the southern flank of the macro-folds ~~by the Banning fault~~ (Fig. 5), and a consistent anticlockwise bend of most axial traces of the macro-folds when they approaching the main San Andreas Banning fault (Fig. 2).

Strike-slip faults in folded sedimentary strata

One major brittle fault set striking NW–SE and dipping steeply to the northeast has impact-developed on the central macro-fold (Figs. 3b and 6). The faults splay out from a bedding-parallel core zone subparallel to steeply SW-dipping mudstone–silt-stone layers on the southern limb of the central macro-fold, and then proceed to truncate NW-dipping sedimentary strata and offset the hinge of a macro-fold by c. 70 meters right-laterally before dying out (Supplement S74a–b). The fault damage zone is traceable for more than one kilometer along strike as a right-slip fault which displaces the hinge of a major, tight, asymmetric, shear-like (similar style) fold (Fig. 6 and Supplement S58). The shear-folded sedimentary strata bend clockwise toward the main fault, thus supporting dominant right-

lateral slip (Fig. 6). Minor faults branch out from the fault core zone and either die out in the macro-fold hinge, and/or persist as bedding-parallel faults for some distance on the southern limb of the macro-fold (Fig. 6).

At smaller scale, the folded and tilted strata of the Palm Spring Formation are commonly truncated by sets of steep NW–SE-striking right-lateral and NNE–SSW-striking left-lateral faults, ~~with displaying~~ meter- to centimeter-scale offsets (Supplement S47b–d). These minor faults generally dip steeply to the northeast to east-northeast, i.e., opposite to most bedding surfaces, which dip southwest (Fig. 3b), and, in places, develop reddish fault gouge along strike. Furthermore, these minor faults typically cut sandstone beds and flatten, and/or die out within, mudstone beds, which restricts their lateral extent to a few decimeters–meters. Kinematic indicators, such as offset of bedding surfaces and fold axial surfaces, yield mostly right-slip displacements, in places with minor reverse components. In some localities, on fold limbs within thick and competent sandstone beds, such minor right- and left-slip faults ~~operate together defining appear to form~~ conjugate sets (Supplement S47b and d) that may have ~~formed-developed~~ simultaneously. In addition, NNE–SSW-striking, ESE-dipping faults and/or semi-brittle kink bands sub-orthogonal to the SAFZ are well displayed in the southeastern macro-fold (Fig. 3c and Supplement S47e) and cut bedding surfaces at high angles with left-slip displacement, therefore potentially representing cross faults between segments/splays of the SAFZ system.

Reverse ~~and thrust~~ faults in folded sedimentary strata

Reverse and thrust faults are common and traceable on the back-limb of the central and southeastern macro-folds near the SAFZ-parallel anticline and the Indio Hills fault, but not recorded in areas close to the ~~main San Andreas~~**Banning** fault. Reverse faults strike mainly NW–SE and dip gently to the southwest, although subsidiary gently NE-dipping faults exist. An example is the low-angle reverse fault that propagates out-of-the syncline on the southeastern macro-fold (Fig. 4) and yields a NE-directed sense-of-shear. This thrust fault may continue westward into the central macro-fold (Fig. 3b), where reverse offset of SW-dipping strata of the Palm Spring Formation constrains vertical displacement from about 10–15 meters (Supplement S35a), though offset is only of a few centimeters in the southeast (Fig. 4). This fault system has a listric geometry, and internal splay faults die out in thick silt- to mud-stone layers. The low-angle faults seem to develop almost consistently near major fold hinge zones and propagate northeastward as out-of-the syncline thrusts (Fig. 4 and Supplement S53a).

In sandstone beds on the north-dipping limb of the major syncline, minor-scale thrust faults, offset asymmetric fold hinges (Supplement S74c) and yield down-to-the-north (normal) sense of shear if the strata are ~~restored-rotated~~ to a horizontal position (Supplement S96). An opposite effect is apparent for a conjugate set of minor normal faults in a small-scale graben structure on the steep, north-dipping layer, which defines a set of reverse faults when ~~restoring-rotating~~ the sedimentary strata to horizontal (Supplements S47d and S96).

Fractures and faults in basement rocks north of the Indio Hills fault

Basement-rock exposures in the Indio Hills are limited to a single, approximately 50-~~m meters~~-long chain of outcrops located in the southeasternmost part of the study area (Fig. 2). These outcrops of massive granite are heavily fractured with mostly steep to sub-vertical sets that strike dominantly NE-SW to ENE-WSW and subsidiary NW-SE to NNW-SSE, ~~possibly representing, conjugate sets (see stereoplot in Fig. 2).~~ Kinematic indicators are generally lacking, but in highly fractured areas, centimeter-thick lenses of unconsolidated reddish gouge are present, comparable to fault rocks observed in Palm Spring Formation sedimentary rocks and corresponding to similar small-scale strike-slip and reverse faults in the basement granite. The fault sets in granitic basement rocks trend parallel to fault sets in sedimentary strata southeast of the Indio Hills fault (see stereoplots in Figure 2) and are therefore suggested to have formed due to similarly oriented stress.

Discussion

Structural eEvolution of SAFZ-oblique folds

~~We mapped and analyzed t~~Three macro-scale fold systems ~~are mapped and analyzed in that occur~~ between the Indio Hills and ~~main San Andreas~~Banning faults. In map view (Fig. 2), ~~T~~the folds are ~~arranged in a~~right-stepping, and ~~each fold set is~~ increasingly asymmetric (Z-shaped); and sigmoidal ~~towards the Indio Hills fault in the~~ northeastward (Fig. 2). ~~Thus~~Based on these properties, we ~~classify-interpret them~~ fold sets as modified SAFZ-oblique *en echelon* macro-folds. Various investigators (Babcock, 1974; Miller, 1998; Titus et al., 2007; Janecke et al., 2018; Bergh et al., 2019) describe sSimilar fold geometries in sedimentary strata ~~are described~~ from many other segments of the SAFZ and are interpreted as structures formed by right-lateral displacement between two major fault strands ~~due to distributed, right-lateral simple shear (Babcock, 1974; Miller, 1998; Titus et al., 2007; Bergh et al., 2019).~~ However, ~~T~~the present fold-orientation data in the Indio Hills; however (Fig. 2), do not correspond with a uniform simple shear model in between two active strike slip faults because the long axis of the strain ellipse is not consistently about 45° to the shear zone as

expected (Sanderson and Marchini, 1984; Sylvester, 1988). Instead, fold geometries vary both
 across and along strike, e.g., axial surface traces of dying-out macro-fold hinges are at high
 obliquity angles ($> 50\text{--}65^\circ$) ~~near the main San Andreas~~ Banning fault in the southwest,
 whereas they are at much lower angles ($< 20\text{--}30^\circ$) and merge with sigmoidal-shaped patterns
 against the Indio Hills fault (Fig. 2). Thus, we propose that the SAFZ-oblique macro-folds in
 Indio Hills rather evolved from a single boundary faults ~~(main San Andreas and Indio Hills~~
~~faults)~~ being ~~active~~ progressively more active through time. For example, a model in which
 the folds initially splayed out from an early active Indio Hills fault through right-lateral
 distributed displacement (~~e.g., compare with~~ Titus et al., 2007) is consistent with fold hinges
 extending outward south of the Indio Hills fault and dying out (broadening) away from the
 fault in a ~~one-two~~ several kilometer-wide damage zone (Fig. 2). Fold propagation outward
from the Indio Hills fault is supported by the increased structural complexity of the fold
geometries towards the Indio Hills fault. Furthermore, the initial upright, *en echelon* folding
 clearly occurred after deposition of the entire Palm Spring Formation because of the
involvement in folding of the Bishop Ash and of adjacent strata possibly of the Ocotillo
Formation (i.e., maximum age of 0.76 Ma – earliest late Pleistocene; Fig. 2 and Table 1), thus
~~favoring folds propagating outward from the Indio Hills fault~~. Should the whole Ocotillo
Formation be folded in the Indio Hills, the maximum age constraints could be narrowed to <
0.6–0.5 Ma based on magnetostratigraphic ages for the upper part of the Ocotillo Formation
(Kirby et al., 2007). By contrast, the ~~main San Andreas~~ Banning fault truncates both limbs of
 the open-style, *en echelon* folds (Fig. 2), which therefore indicates ~~a~~ younger ~~slip~~
~~event~~ deformation along this fault.

The moderate–steep westward plunge of all three macro-folds ($\geq 30^\circ$), however,
 shows that the presumed initial horizontal fold hinges rotated into a steeper plunge. Such
 steepening may be due to, e.g., progressive shortening ~~strain~~ above a deep-seated fault, a
 hidden splay of the Indio Hills fault, or to an evolving stage of distributed shortening (folding)
 adjacent to the master strike-slip faults (e.g., Bergh et al., 2019), with gradually changing
 stress–strain orientation through time, and/or due to structural tilting in the hanging wall of
the Indio Hills fault. This kind of fold reworking favors a situation where the northwestern
 and central macro-folds were pushed up and sideways (right-laterally), following the
 topography and geometry of an evolving ~~convergent tectonic~~ transpressional uplift wedge
(i.e., a contractional uplift formed synchronously with successively with simple shear
transpression to balance internal forces in a crustal-scale critical taper; Dahlen, 1990). The
 corresponding eastward–tightening, enhanced shear folding, and recurrent SW-directed

574 overturned geometries of the central macro-fold on the back-limb of the southeastern macro-
575 fold near the Indio Hills fault (Fig. 3b) support this idea.

576 -We propose a progressive model that changes from distributed (*en echelon* folding) to
577 partly partitioned, i.e., pure shear (shortening) plus simple shear (strike-slip) deformation
578 (Fig. 7), as inferred for other parts of the SAFZ, e.g., in the Mecca Hills (Bergh et al., 2019).
579 In this model, the tight to isoclinal fold geometries to the northeast (Fig. 3b) may account for
580 progressively more intense shortening near the Indio Hills fault, whereas coeval strike-slip
581 faulting affected the already folded and steeply dipping strata of the lower Palm Spring
582 Formation (Fig. 6). This model would favor shortening strain to have evolved synchronously
583 with renewed strike-slip shearing adjacent to the Indio Hills fault, and/or on a ~~hidden-blind~~
584 fault below the contact between the ~~Palm Spring and Mecca Formations~~ and overlying Palm
585 Spring Formation, because the Mecca Formation is much less deformed (Fig. 3c).

586 Alternatively, the more mildly deformed character of the Mecca Formation conglomerate may
587 arise from its homogeneity, which contrasts with alternating successions of mudstone–
588 siltstone and sandstone of the Palm Springs Formation prone to accommodating large
589 amounts of deformation and to strain partitioning. Regardless, such reshaping of *en echelon*
590 folds is supported by analog modelling (McClay et al., 2004; Leever et al., 2011a, 2011b)
591 suggesting that partly partitioned strain may lead to a narrowing of fold systems near a major
592 strike-slip fault (i.e., Indio Hills fault), whereas widening away from the fault indicates still
593 ongoing distributed deformation (~~i.e., near the main San Andreas Banning fault~~). Partly
594 partitioned deformation is supported by the tight to isoclinal and consistent Z-like geometry of
595 smaller-scale folds present on the back-limb of the central and southeastern macro-folds (Fig.
596 3b–c), indicating that they are all parasitic folds and related to the same partly partitioned
597 shear-folding event. Where S- and Z-like fold geometries are present, these minor folds may
598 have formed by buckling in an early stage of *en echelon* folding. An alternative interpretation
599 is that the tight, reshaped parasitic folds are temporally linked to the SAFZ-parallel macro-
600 fold south of the Indio Hills fault (Fig. 3c; see next section).

602 **Structural eEvolution of SAFZ-parallel folds**

603 The SAFZ-parallel anticline differs significantly in geometry from the *en echelon*
604 macro-folds and associated parasitic folds by having an upright and symmetric geometry <
605 20° oblique to the Indio Hills fault (~~Fig. 3e~~). Thus, it resembles that of a ~~dip-slip~~ fault-
606 ~~parallel propagation~~ fold in a more advanced partitioned transpressional segment of the SAFZ
607 (e.g., Titus et al., 2007; Bergh et al., 2019). We suggest that this fold formed by dominant

NE–SW-oriented horizontal shortening, i.e., at high obliquity to the main Indio Hills fault (near-orthogonal pure shear), and/or as a fault-related fold above a buried, major reverse (SW-dipping NE-vergent) oblique-slip splay of the Indio Hills fault at depth (e.g., Schlische, 1995; Suppe and Medwedeff, 1990). The timing might be after the tight reworking of *en echelon* folds in the late Pleistocene, i.e., comparable to other settings (e.g., western Svalbard; Bergh et al., 1997; Braathen et al., 1999). The idea of a late-stage, highly oblique pure-shear overprint onto the macro-folds is supported by small-scale upright folds located within the tight *en echelon* syncline on the back-limb of the modified central macro-fold system (Fig. 4). The NW–SE trend, upright style, and negligible plunge of the fold axes indicate that these folds may be superimposed on the steeper plunging and reshaped *en echelon* folds, and/or that they formed in progression to an increased component of NE–SW shortening on the Indio Hills fault. Nonetheless, ~~it is possible that~~ these folds may have formed simultaneously with the *en echelon* macro-folds in the (earliest?) late Pleistocene (Table 1) due to uncertain (not fully understood) crosscutting relationships.

Progressive NE–SW-oriented contraction may have triggered formation of the upright SAFZ-parallel anticline adjacent to the Indio Hills fault (Fig. 2 and 3c), ~~which~~ The fault then acted as a SW-dipping thrust fault with top-NE displacement. The oblique shortening then led to a certain amount of uplift near the Indio Hills fault, and possibly also accomplished the overturning of folds on the northeastern back-limb of the central and southeastern macro-fold. A similar mode of advanced partitioned shortening was proposed for SAFZ-parallel fold structures in central and southern California (Mount and Suppe, 1987; Titus et al., 2007; Bergh et al., 2019). Our results are supported by stress orientation data acquired by Hardebeck and Hauksson (1999) along a NE–SW-trending profile across the Indio Hills. They recorded an abrupt change in the maximum horizontal stress direction from about 40° oblique to the SAFZ around the main San Andreas Banning fault, to about 70° oblique (i.e., sub-orthogonal) farther northeast, near the Indio Hills fault, which supports the change in attitude and shape of macro-fold geometries that we have outlined. Shortening and strike-slip partitioning, however, would require synchronous right slip on another major fault strand, e.g., the main San Andreas Banning fault, a hypothesis that is supported by the recorded late-stage (i.e., late Pleistocene) shear folding there (Fig. 5).

Fold and fault interaction, evolution, and relative timing

In this section we use the geometry and kinematics of folds and faults in the southern Indio Hills to reconstruct the tectonic history of the area, not only of the inverted late

Cenozoic basin but also about strike-slip and dip-slip faults that bound the basin. Essential tectonic events include (1) extensional normal faulting along the Indio Hills fault in the mid-Miocene–Pliocene (ca. 15–3.0 Ma), (2) reactivation of the Indio Hills fault as a right-lateral to oblique-reverse fault in the (earliest?) late Pleistocene to present-day (< 0.76 Ma), and (3) right-lateral movement along the main San Andreas fault in the late Pleistocene to present-day (< 0.76 Ma; Table 1).

Prior to inversion and uplift of the Indio Hills, the Indio Hills fault most likely acted as a SW-dipping, extensional, basin-bounding normal fault. Evidence-Indications of an early-stage episode of extension ~~is are preserved shown by~~ micro-fault grabens in steeply dipping layers (Supplements S53db and S6), ~~and by the deposition and preservation of sedimentary strata of the Palm Spring and Mecca formations southwest of the Indio Hills, whereas they were eroded or never deposited northeast of the fault, and by fining upwards of the stratigraphic units from conglomerates in the Mecca Formation to coarse-grained sandstone in the lower parts of the Palm Spring Formation.~~ In addition, the flat geometry of micro thrust faults (e.g., Supplements S53b–c) suggests that they were ~~intensely~~ rotated during macro-folding. Restoration of all micro faults in their initial position prior to macro-folding shows that some of these faults exhibit normal kinematics with associated syn-tectonic growth strata (Supplements S53d and S96). Alternatively, the Indio Hills fault dips northeast and uplifted the granitic basement rocks in the hanging wall to the northeast, followed by erosion of the overlying Mecca, Palm Spring and Ocotillo formations there (Fig. 1). We favor a basin geometry and formation similar to that of the Mecca Hills, where down-SW slip along the Painted Canyon fault was inferred in the (Miocene?–) Pliocene (McNabb et al., 2017), and of the transtensional Ridge Basin though ~~with~~ having opposite vergence (Crowell, 1982; Ehman et al., 2000) with a steep, SW-dipping normal fault that was progressively reactivated as an oblique-slip reverse/thrust fault during basin inversion. Formation of the Indio Hills fault as a normal fault probably occurred in mid-Miocene times during extension related to the opening of the Gulf of California (Stock and Hodges, 1989; Stock and Lee, 1994) as proposed for the Salton Trough (Dorsey et al., 2011 and references therein).

Right-lateral to right-lateral-reverse movement along the Indio Hills fault that led to the formation of the SAFZ-oblique *en echelon* macro-folds also ~~indicates supports~~ a steeply dipping character for the precursory Indio Hills fault, ~~which gradually changed to a dominantly right-lateral-reverse fault.~~ The change to a right-lateral-reverse fault is further supported by the presence of both meso-scale strike-slip and thrust faults ~~with~~ having similar NW–SE strikes (Fig. 4, and Supplements S24c and S53a). The increased reverse ~~(and~~

~~decreasing right-lateral~~ component of faulting may have triggered rotation of the *en echelon* macro-fold axes to a steeper plunge, reshaped the open asymmetric folds into tight overturned folds, and caused gentle buckling of strata in the nearby SAFZ-parallel anticline. Hence, the Indio Hills fault ~~acted~~ ultimately functioned as an oblique-slip thrust oblique to the convergent plate boundary margin in the late Pleistocene, which is supported by oblique maximum horizontal stress near the Indio Hills fault (c. 70°; Hardebeck and Hauksson, 1999), while the main San AndreasBanning fault simultaneously accommodated right slip during this period.

By contrast, the last ~~slip event~~ one episode of movement along the main San AndreasBanning fault ~~is clearly younger than the episode of~~ clearly postdates *en echelon* folding, from its truncating attitude (Fig. 2). ~~However~~ In addition, the anticlockwise bending of the axial traces into an ENE–WSW trend ~~when approaching towards~~ the southwest the main San AndreasBanning fault suggests that a distributed component of off-fault deformation stress ~~also~~ affected the area around the main San Andreas fault in its early kinematic stages in the late Pleistocene. The refolding of the southwest limb of the central macro-fold near the main San AndreasBanning fault (Fig. 5) also favors a late-stage activation of this fault in the late Pleistocene (i.e., after the initial transpressional slip events along the Indio Hills fault in the – earliest? – late Pleistocene). Possibly as a consequence of a longer period of activity, and as suggested by relatively higher topographic relief and more intensely folded sedimentary strata in the vicinity of and along the Indio Hills fault than along the main San Andreas fault, the former probably accommodated significantly larger amounts of uplift than the latter. This implies a southwest-tilted geometry for the Indio Hills culmination uplift.

Minor faults in the Indio Hills provide additional input to resolve the spatial, ~~and~~ temporal and kinematic relations between macro-fold and fault interaction ~~in the Indio Hills~~. We analyzed minor fault-related folds (Supplement S53c), which, in their current position on steep north-dipping beds, define down-to-the north displacement. However, when restoring rotating the sedimentary strata to horizontal (Supplement S69), but the fault-related folds define a low-angle fold ~~and~~ – thrust system ~~when restored to horizontal (Supplement S6)~~. These geometric relationships suggest that the minor folds and faults (other than right-slip faults) pre-date (or were coeval with) the SAFZ-oblique macro-folding event, and that they formed initially as internal fractures due to N–S-oriented shortening when the sedimentary strata were still horizontal. i.e., This implies that some partitioning (e.g., SAFZ-parallel small-scale thrust faults) occurred simultaneously with distributed deformation (e.g., SAFZ-oblique *en echelon* macro-folds).

Further, our field data suggest that minor right-slip faults evolved synchronously and parallel with the E–W-trending *en echelon* fold limbs, propagating through rheologically weaker mudstone beds that flowed plastically and acted as slip surfaces during distributed deformation. Later or simultaneously, these faults ~~propagated beyond~~ escaped from the mudstone beds and propagated as NW–SE-striking right-slip faults adjacent to tightened shear folds during partly partitioned deformation, and finally ended up with truncation of the SAFZ-oblique folds (Fig. 6 and Supplement S74a–c).

The presence of out-of-the syncline reverse/thrust faults relative to the reshaped and tightened SAFZ-oblique macro-folds (Fig. 4 and Supplement S53a and d), where SW-dipping thrust faults formed (sub-) parallel to the Indio Hills fault, and the related upright anticline (Fig. 3c) suggests successive distributed and partly partitioned strain in the study area. The proximity and superimposed nature of reverse/thrust faults relative to the reshaped *en echelon* folds suggest that they utilized modified fold hinges and steeply tilted limbs as preexisting zones of weakness. Despite the uncertainty around the crosscutting relationship between the SAFZ-parallel anticline and the SAFZ-oblique *en echelon* macro-folds, the ~~layer-parallel~~ low-angle thrust and intra-detachment folds in the southeastern macro-fold (Fig. 4) indicate that such thrust detachments may have already formed during (early?) distributed deformation, i.e., that distributed and partitioned deformation occurred simultaneously and/or progressively (see phases 1 and 2 in Table 1).

The conjugate WNW–ESE- to NNW–SSE-striking right-slip and NNE–SSW-striking left-slip faults and kink band features truncate strata on both macro-fold limbs (Fig. 3b–c) with an acute angle perpendicular to the macro-folded and tilted Palm Spring Formation strata (e.g., Supplement S74e). Thus, they formed together with or after the *en echelon* macro-folding (< 0.76 Ma).

Tectonic model

In this section we use detailed structural analysis of folds and faults in the southeastern Indio Hills to outline the structural history of the tectonic ~~culmination~~ uplift itself, evaluate it in terms of what is known about strain budgets within the southern San Andreas fault system, link it to nearby structures (Eastern California shear zone and Landers Mojave Line), and integrate the local structural history into a structural synthesis for the southern San Andreas Fault zone in the past 4 Myr.

Our field and structural data support inversion and uplift of the Indio Hills involving progressive or stepwise stages of folding and faulting, ~~events with~~ incorporating a switch

from distributed to partly partitioned transpression (Fig. 7). Prior to inversion in late
Pleistocene time, the Indio Hills fault may have been a steep, SW-dipping normal fault that
~~bounded-downthrew (Miocene?) Pliocene~~ sedimentary strata against granitic basement rocks
in its footwall to the northeast. These basement rocks were partly eroded in the footwall of the
fault. In the hanging wall of the fault, they were ~~and overlain by strata of~~ the Mecca Formation
were deposited in the Pliocene, most likely at 4.03.7–3.73.0 Ma, and ~~by the succeeding~~, lower
and upper members of the Palm Spring Formation ~~strata~~ respectively at 3.07–2.38 Ma and
2.68–10.760 Ma, as suggested from paleomagnetic studies in the Mecca Hills (Chang et al.,
1987; Boley et al., 1994; McNabb et al., 2017).

Early inversion involved distributed transpressional strain triggered by right-lateral
slip along the Indio Hills fault (Fig. 7a). Three macro-scale, upright *en echelon* folds and
associated parasitic folds formed in loosely consolidated sedimentary rocks of the Mecca and
Palm Spring formations after the latter was deposited (< 0.76 Ma), i.e., probably in earliest
late Pleistocene time (Table 1). These ~~SAFZ-oblique fold set~~ evolved oblique to the main
~~strand of the SAFZ displayed and formed~~ a right-stepping pattern ~~with of~~ E–W-trending
~~oriented~~ axial surfaces, ~~probably that trend~~ at a high angle (45°) to the bounding Indio Hills
~~faultmaster fault(s)~~ due to uniform simple shear (e.g., Sanderson and Marchini, 1984;
Sylvester, 1988). This is notably observed in the less deformed southwestern part of the study
area (Fig. 2) near the ~~precursory-main San AndreasBanning~~ fault, where the macro-folds still
display their initial non-plunging geometries. Bed-internal minor fold and fault systems in
weak mudstone beds (Fig. 4 and Supplement S35a) may have formed parallel to the E–W-
trending *en echelon* fold traces, either as thrust detachments due to oblique N–S shortening
when strata were horizontal, and/or as strike-slip faults on the fold limbs. In addition, minor
(bed-internal) SAFZ-parallel thrusts and folds formed prior to or together with the *en echelon*
macro-folds (Supplements S42b–c and S69a–b), thus suggesting minor strain partitioning.

Further deformation in the late Pleistocene led to gradual change from mostly
distributed with minor partitioned deformation to partly partitioned shortening and right-
lateral faulting and folding (Fig. 7b), probably since the Indio Hills fault started to
accommodate an increasing amount of reverse slip, thus acting as an oblique-slip right-lateral-
reverse fault, and where the ~~main San AndreasBanning~~ fault ~~seems to have still played a~~
~~minor role~~ did not yet play a major role. The main result was attenuation-tightening of the
macro-folds toward the Indio Hills fault, ~~increased shear folding~~, and clockwise rotation of
fold axes to a steeper westerly plunge due to increased shear folding, whereas *en echelon*
upright buckle folding continued in the southwest (Fig. 7b). Increased shortening and

778 shearing reshaped the macro-folds and their back-limb folds to tight, isoclinal, and partly
779 overturned folds with consistent Z-style and sigmoidal axial-surface traces near the Indio
780 Hills fault (Fig. 7b). The sigmoidal pattern of the WNW–ESE-trending *en echelon* macro-
781 folds formed at a much lower angle with the Indio Hills fault ($< 20\text{--}30^\circ$) than [farther](#)
782 [southwest with the main San AndreasBanning fault](#) ($60\text{--}70^\circ$). [Furthermore, t](#)The incremental
783 component of lateral strain is recorded as progressively crosscutting NW–SE-striking, strike-
784 slip shear faults terminating with local truncation of the central macro-fold (see Fig. 7c and
785 section below).

786 [Late-stage uUplift of the Indio Hills in the late Pleistocene \(because the earliest late](#)
787 [Pleistocene 0.765 Ma Bishop Ash is involved in folding; Sarna-Wojcicki et al., 2000; Zeeden](#)
788 [et al., 2014\)](#) was marked by a gradual switch to more [kinematically](#) evolved transpressional
789 strain partitioning, where the dominant shortening component ~~affected was accommodated by~~
790 ~~the Indio Hills fault as a~~ right-lateral-oblique, [top-NE thrusting along the Indio Hills](#) fault and
791 ~~the main major~~ strike-slip ~~component movement was centered~~ along the [main San](#)
792 [AndreasBanning](#) fault (Fig. 7c [and phase 3 in Table 1](#)). NE-directed oblique thrusting on the
793 Indio Hills fault and related minor, reverse, out-of-the syncline faults led to uplift, which
794 resulted in formation of a major anticline parallel to the Indio Hills fault in sediments of the
795 Mecca Formation ([see anticline closest to Indio Hills fault in Fig. 3c and 7c](#)). With increasing
796 partitioning, ~~margin-parallel-slip~~ [parallel to the convergent plate boundary](#) was accommodated
797 by right-slip along the linear [main San AndreasBanning](#) fault, where subvertical folds formed
798 locally, and presumed antithetic conjugate kink band sets of right- and left-slip cross faults
799 affected the entire uplifted area.

800 We favor a progressive evolution from distributed to partly partitioned deformation as
801 presented in Fig. 7a–c, although overlapping and synchronous formation of various structures
802 may have occurred ([overlapping of phases 1 and 2 in Table 1](#)), at least locally (except for the
803 late-stage [main San AndreasBanning](#) fault and related shear folds; [phase 3 in Table 1](#)). The
804 ~~overlapping and synchronous formation of structureslatter~~ is based on uncertainties in our
805 field data, e.g., variable cross-cutting relations of early, bedding-parallel strike-slip and thrust
806 faults and *en echelon* macro-folds (Figs. 4 and 6, and Supplements [S53c–d](#) and [S47](#)), and
807 from the spatial variations in the direction of maximum horizontal stress across the Indio Hills
808 at present, from 40° oblique to the boundary faults near the [main San AndreasBanning](#) fault to
809 70° oblique near the Indio Hills fault (Hardebeck and Hauksson, 1999).

810 [Our observations of mostly lateral movement along the main San Andreas fault \(i.e.,](#)
811 [southeastern continuation of the Mission Creek fault\) and the proposed late Pleistocene to](#)

present-day age for deformation in the southeastern Indio Hills are consistent with work by Keller et al. (1982). A major difference between the northwestern and southeastern Indio Hills is the relatively tighter macro-folding over a narrower area and more intense character of deformation in between the two bounding faults in the southeastern Indio Hills (Figs. 2 and 3; Keller et al., 1982; Lancaster et al., 2012).

~~The present model and right-lateral-reverse character of the~~ The right-lateral-reverse character of the Indio Hills Fault and its role in our kinematic model for basin inversion in the southern Indio Hills ~~fault~~ are further supported by the relationship of the Indio Hills fault with the Eastern California ~~s~~Shear ~~z~~Zone, which merge together north of the study area where the Indio Hills fault bends into a NNW–SSE strike along the Landers–Mojave Line (Dokka and Travis, 1990a, 1990b; Nur et al., 1993a, 1993b; Thatcher et al., 2016). Recent activity along the Landers–Mojave Line recorded as six–seven earthquakes with $M > 5$ between 1947 and 1999 (Fig. 1; Nur et al., 1993a, 1993b; Du and Aydin, 1996; Spinler et al., 2010) indicates that a through-going NNW–SSE-striking fault crosscuts the Pinto Mountain fault (Nur et al., 1993a, 1993b; Rymer, 2000). Notably, the 1992 Joshua Tree earthquake occurred along the NNW–SSE-striking, west-dipping West Deception Canyon fault (Rymer, 2000 and references therein), which merges with the (probably southwest-dipping) Indio Hills fault in the south (see figure 1 in Rymer, 2000). Therefore, we propose that the Indio Hills fault, may be one of several faults to transfer displacement from unsuitably oriented, NW–SE-striking right-slip faults in the north, such as the Calico and Camp Rock faults, to the main SAFZ strand in the south (Fig. 1).

Farther southeast along strike, the Indio Hills and ~~main San AndreasBanning~~ faults merged along a dextral freeway junction, i.e., a junction of three dextral fault branches (sensu Platt and Passchier, 2016 and Passchier and Platt, 2017), ~~that-which~~ may have enhanced, wedge-shaped transpressional uplift of the Indio Hills after the (late) formation of the ~~main San AndreasBanning~~ fault in the late Pleistocene (Fig. 8a–c and Table 1). However, anticlockwise rotation of the Indio Hills block and related structures in map view as predicted in a dextral freeway junction (Platt and Passchier, 2016; Passchier and Platt, 2017) was not recorded by our field data (except along the ~~main San AndreasBanning~~ fault due to localized right-slip along the fault; cf. sub-vertical shear fold in Fig. 5). This may be due in part to the late formation of the ~~main San AndreasBanning~~ fault (< 0.764 Ma, i.e., late Pleistocene), i.e., clockwise rotation (in map view) of the fold and fault structures due to right-lateral slip along the Indio Hills fault, and to the oblique-slip character of the Indio Hills fault. Thus, the dextral freeway junction in the Indio Hills may be more of a transitional nature. Instead of major

anticlockwise rotation of the Indio Hills block in map view, the accretion of material toward the fault junction due to right slip along the main San Andreas~~Banning~~ fault is probably partly accommodated by the dominant vertical slip component along the Indio Hills fault, leading to further uplift near the junction (i.e., clockwise rotation in cross section).

Regional comparison and implications

The proposed progressive tectonic model for the Indio Hills uplift has wide implications when compared and correlated with other fault strands of the SAFZ bounding uplifted domains along strike in the Coachella and Imperial ~~V~~ valleys (Fig. 8a–c), and in explaining lateral variations in fault architectures, kinematic evolution and timing, deformation mechanisms and areal segmentation (Sylvester and Smith 1987; McNabb et al., 2017; Janecke et al., 2018; Bergh et al., 2019). Here we compare and contrast the structural evolution of the southeastern Indio Hills with that of nearby tectonic culminations uplifts (Mecca Hills and Durmid Hills).

Comparison with the Mecca Hills

Previous studies of SAFZ-related uplifts between the Indio Hills and Durmid Hills in Coachella Valley ~~suggests~~show that the Indio Hills and main San Andreas~~Banning~~ faults link up in the southeasternmost Indio Hills and proceed as ~~directly with~~ the main San Andreas ~~FZ~~fault strand in the Mecca Hills (Fig. 8c) which then, together with the subsidiary Skeleton Canyon and Painted Canyon faults, bounds a much wider flower-like uplift area than in the Indio Hills (Fig. 8c; Sylvester and Smith, 1976, 1987; Sylvester, 1988; McNabb et al., 2017; Bergh et al., 2019). In contrast to the Indio Hills fault, however, the main San Andreas ~~fault~~FZ in Mecca Hills has an anastomosing geometry with thick (10–500 m), red-stained fault gouge. Regardless, we consider these faults ~~to~~ to be correlative and infer the lack of fault gouge ~~in~~ along the Indio Hills fault to be due to more localized strain on the Indio Hills fault than on the SAFZ in Mecca Hills. This is supported by a more rectilinear geometry and lack of fold–fault linkage in Indio Hills, which may have allowed initial lubrication of the fault surface in basement rocks with high contrasting rheology (e.g., Di Toro et al., 2011; Fagereng and Beall, 2021), and which hampered fluid circulation and extensive cataclasis. Another possible explanation may be the presence of coarse-grained deposits of the Mecca Formation, which may have partitioned/decoupled deformation along the Indio Hills fault from that in overlying Palm Spring sedimentary strata.

Both the Indio Hills and Mecca Hills uplift areas are bounded to the northeast by a presumed Miocene–Pliocene, SW-dipping normal fault (Fig. 8a), which later acted as major

SAFZ-parallel oblique-reverse faults, and which significantly contributed to the uplift of these areas in ~~Pliocene~~-(late) Pleistocene time (Sylvester and Smith, 1976, 1987; McNabb et al., 2017; Bergh et al., 2019). In the Mecca Hills (Fig. 8c), the Painted Canyon fault is flanked in the hanging-wall to the southwest by a basement-cored, macro-fold (Mecca anticline), which is similar to the upright anticline that parallels the Indio Hills fault and adjacent minor thrust faults (~~Error! Reference source not found.~~Figure 2 & Figure 3c and Supplement S5a). Similar folds appear adjacent to the Hidden Springs–Grotto Hills fault (Sheridan et al., 1994; Nicholson et al., 2010), a NW–SE-striking, now reverse splay fault of the main SAFZ between the Mecca Hills and Durmid Hills (Fig. 8c). It is, however, unlikely that these marginal faults link up directly along strike. Rather, they merge or splay with the SAFZ and SAFZ-oblique faults.

The inversion and main uplift history of the Mecca Hills segment of the SAFZ (Bergh et al., 2019) initiated with right-lateral slip on a steep SAFZ, from where SAFZ-oblique *en echelon* folds and dominantly right-slip faults splayed out in a one–two kilometers wide damage zone on either side of the SAFZ (Fig. 8a). The subsidiary Skeleton Canyon fault initiated as a steep right-lateral and SAFZ-parallel strike-slip fault along a small restraining bend (Fig. 8b). Successive lateral shearing reshaped the *en echelon* folds into steeply plunging folds with axial traces parallel to the SAFZ. The final kinematic stage generated SW-verging fold and thrust structures parallel to the SAFZ (Fig. 8c), which truncated the *en echelon* folds and the NE-dipping Skeleton Canyon fault. The resulting wedge-like flower structure thus records a polyphase kinematic evolution from distributed, through locally partitioned, to fully partitioned strain-in-a-changing-transpressional-plate-regime (Bergh et al., 2019).

Based on the geometric similarities, we consider that the *en echelon* macro-folds in both Indio Hills and Mecca Hills formed ~~simultaneously~~coevally, but not on the same regional ~~right-lateral~~ fault strand (Fig. 8a). In both areas, the *en echelon* folds and faults are strongly reworked and tightened into sigmoidal shapes where they merge with the Indio Hills and Skeleton Canyon faults respectively (Fig. 8b; Bergh et al., 2019), and SAFZ-parallel thrust faults formed early (i.e., prior to macro-folding) both in the Indio Hills (Supplement S35c–d) and in the Mecca Hills (Rymer, 1994), thus supporting continuous, partly partitioned strain field in both areas. Strain partitioning caused major uplift of the Mecca Hills block along the Skeleton Canyon, Painted Canyon, and Hidden Springs–Grotto Hills faults (Fig. 8c), all acting as SAFZ-parallel oblique-slip thrust faults (Sheridan et al., 1994; Bergh et al., 2019). The partitioned right-slip component was partly transferred to the main San AndreasBanning fault in Indio Hills, and/or to an unknown hidden fault southwest of the

SAFZ (e.g., in Mecca Hills; Hernandez Flores, 2015; Fuis et al., 2017), possibly the Eastern Shore-line fault (Janecke et al., 2018).

Based on paleomagnetic and structural field studies, uplift of the SAFZ-related Mecca basin started at ca. 2.63–0–2.20.76 Ma (i.e., earliest to mid Pleistocene) with partial and local erosion of the Palm Spring Formation (see lower and upper unconformities in McNabb et al., 2017) and culminated after 1.0–0.76 Ma (see unconformity between the uppermost Palm Spring Formation and base of the Ocotillo Formation southwest of the Painted Canyon fault in McNabb et al., 2017), i.e., after deposition of the whole Palm Spring Formation (McNabb et al., 2017; Janecke et al., 2018). Uplift is still ongoing at present (Fattaruso et al., 2014; Janecke et al., 2018). A comparable time frame and ongoing activity are expected for the Indio Hills. Fault activity and tectonic uplift of the Mecca Hills therefore most likely initiated earlier (earliest Pleistocene) than in the Indio Hills (earliest late Pleistocene; Table 1), where the transition from the lower to the upper member of the Palm Spring Formation is gradual and does not show any major unconformity.

Comparison with Durmid Hills

The Durmid ladder structure along the southern 30 kilometers of the SAFZ in Imperial Valley defines a similar but oppositely merging, one–three kilometers wide wedge-shaped uplift as in Indio Hills, bounded by the right-lateral and reverse Eastern Shoreline fault to the southwest and the main SAFZ to the northeast (Fig. 8c; Janecke et al., 2018). Internally, the ladder structure comprises *en echelon* folds (Babcock, 1974; Bürgmann, 1991) that merge in a sigmoidal pattern with the main SAF, and subsidiary sets of conjugate SAFZ-parallel right-lateral and SAFZ-oblique E–W-striking, left-slip cross faults, which accommodated clockwise rotation of internal blocks (Janecke et al., 2018). The *en echelon* folds formed at a comparable time, i.e., < 0.76 Ma in the Indio Hills and at ca. 0.5 Ma in the Durmid Hills (Table 1). By assuming a northwest continuation of the main SAFZ with the SAFZ in Mecca Hills, the Eastern Shoreline fault has no exposed correlative fault in the Mecca Hills and Indio Hills (Fig. 8c; Damte, 1997; Bergh et al., 2019). Nevertheless, the Eastern Shoreline fault may continue at depth southwest of the main San Andreas Banning fault and main SAFZ (Janecke et al., 2018).

~~The increasing width of damage zones adjacent to SAFZ-related faults southward in Coachella Valley, and increased number of strike-slip and oblique to orthogonal cross faults in the Durmid Hills compared with Indio Hills and Mecca Hills may be due to closeness and transition to a transtensional rift setting around the Brawley seismic zone (Janecke et al., 2018).~~ A significant difference between the Indio Hills–Mecca Hills and the Durmid Hills,

however, is the large number of cross faults in the Durmid ladder structure. Such faults are interpreted as early-stage (ca. 1 Ma – early/mid Pleistocene), NE–SW-striking, left-lateral, faults (Fig. 8a), which were rotated clockwise by progressive right-lateral motion into sigmoidal parallelism with the SAFZ and Eastern Shoreline fault (Fig. 8b–c; Janecke et al. 2018). In contrast, cross faults in Indio Hills are much less common and, where present, ~~possibly-probably~~ formed late, but prior to the main San AndreasBanning fault (i.e., in the earliest or middle part of the late Pleistocene). Thus, in the Indio Hills, there is no evidence of clockwise rotation of early-stage cross faults as in the Durmid Hills, but rather clockwise rotation of fold axial traces is common, which may be a first step in the formation of ladder-like fault blocks (e.g., Davis, 1999; Schultz and Balasko, 2003).

A major outcome of the comparison with Durmid Hills is that the wedge-shaped uplift block between the Indio Hills and main San AndreasBanning faults may represent a failed uplift and/or the early stage of formation of a ladder structure. This idea is supported by presence of similar master faults and structures with comparable kinematics in both the Indio Hills and Durmid Hills, including oblique *en echelon* macro-folds, strike-slip faults acting as step-over faults, and reverse faults. Younger, non-rotated, conjugate cross faults exist in the Indio Hills but not in the Durmid Hills where such faults are more evolved features due to larger strain and more advanced stage of ladder structure formation. From these observations, one should expect to find ladder structures operating at different evolution stages among the many, yet unexplored uplifts in Coachella Valley.

Conclusions

1) The Indio Hills fault likely initiated as a SW-dipping, basement-seated normal fault during the opening of the Gulf of California in the mid Miocene, and was later inverted as a right-lateral reverse, oblique-slip fault in the (earliest?–) late Pleistocene due to transpression along the convergent plate boundary, whereas the main San Andreas fault initiated probably as a dominantly right-slip fault during the later stages of uplift in the late Pleistocene.

1)2) The Indio Hills segment of the SAFZ in Coachella Valley, southern California evolved as a wedge-shaped uplift block between two major SAFZ-related fault strands, the Indio Hills and main San AndreasBanning faults, which merge in a dextral freeway junction of a transitional nature to the southeast.

2) The Indio Hills fault acted as a SW-dipping, basement-seated normal fault in Miocene time, i.e., prior to inversion as an oblique-slip, right lateral reverse fault during Pliocene and

Pleistocene times, whereas the main San Andreas Banning fault initiated probably during the later stages of uplift as a dominantly right-slip fault.

3) Transpressive deformation triggered uplift and inversion of the Indio Hills through a progressive change from distributed *en echelon* folding to partly partitioned right-slip thrusting. We favor a progressive rather than stepwise model in which the main uplift was related to late shortening ~~in~~ at the freeway junction where the Indio Hills and main San Andreas Banning faults merge.

4) The Indio Hills fault is a splay fault of the SAFZ that merges to the north with the Landers–Mojave Line and contributes to transfers slip from unsuitably oriented faults of the Eastern California sShear zZone to the main San Andreas Banning fault ~~portion of the SAFZ~~ in the southeast.

5) A significant difference of the Indio Hills with the Durmid Hills is that left-lateral step-over and cross faults in the Durmid Hills rotated subparallel with the mSAF, whereas in Indio Hills, all cross faults are oblique with the SAFZ and, thus, may reflect an earlier stage of a still evolving ladder structure.

5)6) The initiation of right-lateral to right-lateral-reverse slip along major SAFZ-parallel faults and the main San Andreas fault in the Coachella Valley is younger towards the northwest (Pliocene in the Durmid Hills, early Pleistocene in the Mecca Hills and late Pleistocene in the Indio Hills). The onset of uplift, however, appears to be coeval in all tectonic ~~culminations~~ uplifts (late to latest) Pleistocene.

Data availability

The structural dataset and field photographs used in the present study are available on DataverseNO (Open Access repository) at <https://doi.org/10.18710/TM18UZ>. DEM images are from Google Earth (© Google Earth 2011).

Authors contribution

All authors contributed to collect structural measurements in the Indio Hills. JBPK wrote the first draft of the manuscript and designed half the figures and supplements (workload: 35%). Prof. SGB made major revision to the initial draft and designed half the figures and supplements (workload: 35%). Prof. AGS also revised the manuscript and provided major input about the local geology (workload: 30%).

Competing interests

The authors declare that they have no known competing interests.

Acknowledgments

The staff at the University of California–Santa Barbara and San Diego State University provided great hospitality during Steffen Bergh’s sabbatical leaves in 2011–2012 and 2016–2017 while working with the San Andreas fault. We thank all the persons from these institutions that were involved in this project. The authors thank Prof. Emeritus Arild Andresen (University of Oslo) and Prof. Holger Stunitz (UiT) for helpful comments and Jack Brown (San Diego State University) for fieldwork collaboration. Prof. Susanne Janecke (Utah State University) and Dr. Miles Kenney (Kenney Geoscience) provided fruitful discussion.

Financial support

The present study is part of the CEED (Centre for Earth Evolution and Dynamics) and ARCEX projects (Research Centre for Arctic Petroleum Exploration), which are funded by grants from UiT The Arctic University of Norway in Tromsø and the Research Council of Norway (grant numbers 223272 and 228107) together with eight academic and six industry partners.

References

- Allen, C. R.: San Andreas fault zone in San Geronimo Pass, southern California, *GSA Bull.*, 68, 315–360, 1957.
- Atwater, T. and Stock, J.: Pacific-North America Plate Tectonics of the Neogene Southwestern United States: An Update, *International Geology Review*, 40:5, 375–402, 1998.
- Babcock, E. A.: Structural Geology and Geophysics of the Durmid Area, Imperial Valley, California, Ph.D. Thesis, University of California, Riverside, 149 pp., 1969.
- Babcock, E. A.: Geology of the northeast margin of the Salton Trough, Salton Sea, California, *GSA Bull.*, 85, 321–332, 1974.
- Bergh, S. G., Braathen, A. and Andresen, A.: Interaction of basement-involved and thin-skinned tectonism in the tertiary fold-and thrust belt of Central Spitsbergen, Svalbard, *AAPG Bull.*, 81, 637–661, 1997.
- Bergh S. G., Sylvester, A. G., Damte, A. and Indrevær, K.: Evolving transpressional strain fields along the San Andreas fault in southern California: implications for fault

branching, fault dip segmentation and strain partitioning, *Geophys. Res. Abs.*, 16, EGU General Assembly, 24th April–2nd May, Vienna, Austria, 2014.

Bergh, S. G., Sylvester, A. G., Damte, A. and Indrevær, K.: Polyphase kinematic history of transpression along the Mecca Hills segment of the San Andreas fault, southern California, *Geosphere*, 15, 34 pp., 2019.

Bilham, R. and Williams, P.: Sawtooth segmentation and deformation processes on the southern San Andreas fault, California, *Geophys. Res. Lett.*, 12, 9, 557–560, 1985.

Blisniuk, K., Scharer, K., Sharp, W. D., Burgmann, R., Amos, C. and Rymer, M.: A revised position for the primary strand of the Pleistocene-Holocene San Andreas fault in southern California, *Sci. Adv.*, 7, eaaz5691, 2021.

Boley, J.-L., Stimac, J. P., Weldon II, R. J. and Rymer, M. J.: Stratigraphy and paleomagnetism of the Mecca and Indio Hills, southern California, in: *Geological Investigations of an Active Margin*, edited by: McGill, S. F. and Ross, T. M., GSA, Cordilleran Section Guidebook, 27th Annual Meeting, San Bernardino County Museum Associations, USA, 336–344, 1994.

Braathen, A., Bergh, S. G. and Maher Jr., H. D.: Application of a critical wedge taper model to the Tertiary transpressional fold-and thrust belt on Spitsbergen, Svalbard, *GSA Bull.*, 111, 1468–1485, 1999.

[Bryant, W. A.: Fault number 118, Pinto Mountain fault zone \(includes Morongo Valley fault\). Quaternary fault and fold database of the United States, USGS website, 2000. https://earthquakes.usgs.gov/hazards/qfaults, accessed 05/03/2022 09:32 a.m.](https://earthquakes.usgs.gov/hazards/qfaults)

Bürgmann, R.: Transpression along the southern San Andreas fault, Durmid Hills, California, *Tectonics*, 10, 1152–1163, 1991.

Chang, S.-B. R., Allen, C. R. and Kirschvink, J. L.: Magnetic stratigraphy and a test for block rotation of sedimentary rocks within the San Andreas fault zone, Mecca Hills, southeastern California, *Quat. Res.*, 27, 30–40, 1987.

[Crowell, J. C.: The tectonics of the Ridge Basin, southern California, in: *Geologic History of Ridge Basin, Southern California, SEPM, Pacific Section*, 22, 25–42, 1982.](#)

[Dahlen, F. A.: Critical Taper Model of Fold-and-Thrust Belts and Accretionary Wedges, *Annu. Rev. Earth Planet. Sci.*, 18, 55–99, 1990.](#)

Dair, L. and Cooke, M. L.: San Andreas fault geometry through the San Gorgonio Pass, California, *Geology*, 37, 2, 119–122, 2009.

- Damte, A.: Styles of deformation in zones of oblique convergence: Examples from the Mecca Hills, southern San Andreas fault, Unpublished Ph.D. thesis, University of California, Santa Barbara, 164 pp., 1997.
- Davis, G. H.: Structural geology of the Colorado Plateau region of southern Utah, with emphasis on deformation bands, GSA Spec. Pap., 342, 157 pp., 1999.
- Di Toro, G., Han, R., Hirose, T., De Paola, N., Nielsen, S., Mizoguchi, K., Ferri, F., Cocco, M. and Shimamoto, T.: Fault lubrication during earthquakes, *Nature*, 471, 494–498, 2011.
- Dibblee, T. W. Jr.: Geology of the Imperial Valley, California, Californian Division of Mines Bulletin, 170, 21–28, 1954.
- Dibblee, T. W. Jr.: Geology of the southeastern San Andreas fault zone in the Coachella Valley area, Southern California, in: Southern San Andreas fault, Whitewater to Bombay Beach, Salton Trough, California, edited by: Baldwin, J., Lewis, L., Payne, M. and Rocquemore, G., South Coast Geological Society, Annual Field Trip Guide Book, 25, 35–56, 1997.
- Dibblee, T. W. Jr. and Minch, J. A.: Geological map of the Thousand Palms & Lost Horse Mountain 15 minutes quadrangles, Riverside County, California, Dibblee Foundation, Map DF-372, 2008.
- Dokka, R. K. and Travis, C. J.: Late Cenozoic strike-slip faulting in the Mojave Desert, California, *Tectonics*, 9, 311–340, 1990a.
- Dokka, R. K. and Travis, C. J.: Role of the Eastern California Shear Zone in accommodating Pacific-North American plate motion, *Geophys. Res. Lett.*, 17, 1323–1326, 1990b.
- Dorsey, R. J., Housen, B. A., Janecke, S. U., Fanning, C. M. and Spears, A. L. F.: Stratigraphic record of basin development within the San Andreas fault system: Late Cenozoic Fish Creek–Vallecito basin, southern California, *GSA Bull.*, 123, 5/6, 771–793, 2011.
- Du, Y. and Aydin, A.: Is the San Andreas big bend responsible for the Landers earthquake and the eastern California shear zone, *Geology*, 24, 3, 219–222, 1996.
- Ehman, K. D., Sullivan, M. D. and May, S. R.: Ridge Basin Field Trip Guidebook, Tectonic Controls on Facies Distribution and Stacking Patterns, Ridge Basin, southern California, SEPM, Pacific Section, Los Angeles, California, USA, 87, 50 pp., 2000.
- Fagereng, Å. and Beall, A.: Is complex fault zone behavior a reflection of rheological heterogeneity, *Phil. Trans. R. Soc. A329*: 20190421, 2021.

1114 Fattaruso L.A., M.L. Cooke & R.J. Dorsey. 2014. Sensitivity of uplift patterns to dip of the
1115 San Andreas fault in the Coachella Valley, California. *Geosphere*, vol. 10, pp. 1.12.

1116 Fuis, G. S., Scheirer, D. S., Langenheim, V. E. and Kohler, M. D.: A New Perspective on the
1117 Geometry of the San Andreas Fault in Southern California and Its Relationship to
1118 Lithospheric Structure, *BSSA*, 102, 236–251, 2012.

1119 Fuis G. S., Bauer, K., Goldman, M. R., Ryberg, T., Langenheim, V. E., Scheirer, D. S.,
1120 Rymer, M. J., Stock, J. M., Hole, J. A., Catchings, R. D., Graves, R. W. and Aagaard,
1121 B.: Subsurface Geometry of the San Andreas Fault in Southern California: Results
1122 from the Salton Seismic Imaging Project (SSIP) and Strong Ground Motion
1123 Expectations, *BSSA*, 107, 4, 1642–1662, 2017.

1124 [Gold, P. O., Behr, W. M., Rood, D., Sharp, W. D., Rockwell, T. K., Kendrick, K. and Salin,](#)
1125 [A.: Holocene geologic slip rate for the Banning strand of the southern San Andreas](#)
1126 [Fault, southern California, *J. Geophys. Res. Solid Earth*, 120.](#)
1127 [doi:10.1002/2015JB012004](#)

1128 [Groshong, R. H.: Strain, fractures, and pressure solution in natural single-layer folds, *GSA*](#)
1129 [Bull., 86, 1363–1376, 1975.](#)

1130 Guest, B., Niemi, N. and Wernicke, B.: Stateline fault system: A new component of the
1131 Miocene-Quaternary Eastern California shear zone, *GSA Bull.*, 119, 11/12, 1337–
1132 1346, 2007.

1133 Hardebeck J.L. & E. Hauksson. 1999. Role of fluids in faulting inferred from stress field
1134 signatures. *Science*, 285, pp. 236-239.

1135 Hauksson, E., Yang, W. and Schearer, P. M.: Waveform Relocated Earthquake Catalog for
1136 Southern California (1981 to June 2011), *BSSA*, 102, 5, 2239–2244, 2012.

1137 Hernandez Flores, A. P.: Paleosismologia del sistema de fallas imbricado en la Sierra
1138 Cucapah, Baja California, Mexico, Master's Thesis, Centre for Scientific Research and
1139 Higher Education, Ensenada, Mexico, 254 pp., 2015.

1140 Herzig, C. T., Mehegan, J. M. and Stelling, C. E.: Lithostratigraphy of the State 2-14
1141 Borehole: Salton Sea Scientific Drilling Project, *J. Geophys. Res.*, 93, B11, 12969–
1142 12980, 1988.

1143 Janecke, S. U., Markowski, D. K., Evans, J. P., Persaud, P. and Kenney, M.: Durmid ladder
1144 structure and its implications for the nucleation sites of the next $M > 7.5$ earthquake on
1145 the San Andreas fault or Brawley seismic zone in southern California, *Lithosphere*, 10,
1146 5, 602–631, 2018.

- Keller, E. A., Bonkowski, M. S., Korsch, R. J. and Schlemmon, R. J.: Tectonic geomorphology of the San Andreas fault zone in the southern Indio Hills, Coachella Valley, California, GSA Bull., 93, 46–56, 1982.
- Kirby, S. M., Janecke, S. U., Dorsey, R. J., Housen, B. A., Langenheim, V. E., McDougall, K. A. and Stealy, A. N.: Pleistocene Brawley and Ocotillo Formations: Evidence for Initial Strike-Slip Deformation along the San Felipe and San Jacinto Fault Zones, Southern California, J. of Geology, 115, 43–62, 2007.
- [Lancaster, J. T., Hayhurst, C. A. and Bedrossian, T. L.: Preliminary geologic map of Quaternary surficial deposits in southern California: Palm Springs 30' x 60' quadrangle, in: Geologic compilation of Quaternary surficial deposits in southern California, edited by: Bedrossian, T. L., Roffers, P., Hayhurst, C. A., Lancaster, J. T. and Short, W. R., California Geological Survey, Sacramento, December 2012. \[https://www.conservation.ca.gov/cgs/fwgp/Pages/sr217.aspx#palmsprings.\]\(https://www.conservation.ca.gov/cgs/fwgp/Pages/sr217.aspx#palmsprings\)](https://www.conservation.ca.gov/cgs/fwgp/Pages/sr217.aspx#palmsprings)
- Leever, K., Gabrielsen, R. H., Sokoutis, D. and Willingshofer, E.: The effect of convergence angle on the kinematic evolution of strain partitioning in transpressional brittle wedges: Insight from analog modeling and high-resolution digital image analysis, Tectonics, 30, 2011a.
- Leever, K., Gabrielsen, R. H., Faleide, J. I. and Braathen, A.: A transpressional origin for the West Spitsbergen fold- and thrust belt: Insight from analog modeling, Tectonics, 30, 1–24, 2011b.
- Lin, G.: Three-Dimensional Seismic Velocity Structure and Precise Earthquake Relocations in the Salton Trough, Southern California, BSSA, 103, 5, 2694–2708, 2013.
- Lin, G., Schearer, P. M. and Hauksson, E.: Applying a three-dimensional velocity model, waveform cross correlation, and cluster analysis to locate southern California seismicity from 1981 to 2005, J. Geophys. Res., 112, B12309, 2007.
- Lindsey, E. O. and Fialko, Y.: geodetic slip rates in the southern San Andreas Fault system: Effects of elastic heterogeneity and fault geometry, J. Geophys. Res., 118, 689–697, 2013.
- Lutz, A. T., Dorsey, R. J., Housen, B. A. and Janecke, S. U.: Stratigraphic record of Pleistocene faulting and basin evolution in the Borrego Badlands, San Jacinto fault zone, Southern California, GSA Bull., 118, 11/12, 1377–1397, 2006.
- Markowski, D. K.: Confirmation of a New Geometric and Kinematic Model of the San Andreas Fault at Its Southern Tip, Durmid Hill, Southern California, Master's Thesis, Utah State University, Logan, USA, 151 pp., 2016.

- Matti, J. C., Morton, D. M. and Cox, B. F.: Distribution and geologic relations of fault systems in the vicinity of the Central Transverse Ranges, southern California, USGS Report, 85-365, 31 pp., 1985.
- [Matti, J. C., Morton, D. M. and Cox, B. F.: The San Andreas fault system in the vicinity of the central Transverse Ranges province, southern California, U.S. Geological Survey Open-File, Report 92-354, 40 pp., 1992.](#)
- [Matti, J. C., and Morton, D. M.: Paleogeographic evolution of the San Andreas fault in southern California: a reconstruction based on a new cross-fault correlation, in: The San Andreas fault system: displacement, palinspastic reconstruction, and geologic evolution, edited by: Powell, R. E., Weldon, R. J. and Matti, J. C., GSA Mem., 178, 107–159, 1993.](#)
- McClay, K. R., Whitehouse, P. S., Dooley, T. and Richards, M.: 3D evolution of fold and thrust belts formed by oblique convergence, Mar. Petrol. Geol. Bull., 21, 857–877, 2004.
- McNabb, J. C.: Stratigraphic record of Pliocene-Pleistocene basin evolution and deformation along the San Andreas fault, Mecca Hills, California, Unpublished Master's Thesis, University of Oregon, 70 pp., 2013.
- McNabb, J. C., Dorsey, R. J., Housen, B. A., Dimitroff, C. and Messé, G. T.: Stratigraphic record of Pliocene–Pleistocene basin evolution and deformation within the Southern San Andreas Fault Zone, Mecca Hills, California, Tectonophys., 719–720, 66–85, 2017.
- [Meere, P. A., Mulchrone, K. F. and Timmerman, M.: Shear folding in low-grade metasedimentary rocks: Reverse shear along cleavage at a high angle to the maximum compressive stress, geology, 41, 8, 879–882, 2013.](#)
- Miller, D.D.: Distributed shear, rotation, and partitioned strain along the San Andreas fault, central California, Geology, 26, 867–870, 1998.
- Morton, D., Matti, J. and Tinsley, J.: Banning fault, Cottonwood canyon, San Geronio pass, southern California, in: Cordilleran Section of the Geological Society of America, edited by: Hill, M., 191–192, Geol. Soc. of Am., Boulder, Colorado, USA, 1987.
- [Morton, D. and Matti, J. C.: Extension and contraction within an evolving divergent strike-slip fault complex: The San Andreas and San Jacinto fault zones at their convergence in southern California, in: The San Andreas fault system: displacement, palinspastic reconstruction, and geologic evolution, edited by: Powell, R. E., Weldon, R. J. and Matti, J. C., GSA Mem., 178, 217–230, 1993.](#)

- Mount, V. S. and Suppe, J.: State of stress near the San Andreas fault: implications for wrench tectonics, *Geology*, 15, 1143–1146, 1987.
- Nicholson, C., Hauksson, E. and Plesch, A.: Revised 3D Fault Models for the Southern San Andreas Fault System Extending from San Geronimo Pass to the Salton Sea, 106th Annual Meeting AAPG, 27th–29th May, 2010.
- Nur, A., Hagai, R. and Beroza, G. C.: The Nature of the Landers-Mojave Earthquake Line, *Science*, 261, 201–203, 1993a.
- Nur, A., Hagai, R. and Beroza, G. C.: Landers-Mojave earthquake Line: A New Fault System?, *GSA Today*, 3, 10, 255–258, 1993b.
- [Passchier, C. W. and Platt, J. P.: Shear zone junctions: Of zipper and freeways, *J. Structural Geol.*, 95, 188–202, 2017.](#)
- ~~[Parrish, J.G.: Geological compilation of Quaternary surficial deposits in southern California, Palm Springs 30' x 60' quadrangle, CGS Special Report 217, Plate 24, 1983.](#)~~
- Platt, J. P. and Passchier, C. W.: Zipper junctions: A new approach to the intersections of conjugate strike-slip faults, *Geology*, 44, 10, 795–798, 2016.
- Rymer, M. J.: Quaternary Fault-Normal Thrusting in the Northwestern Mecca Hills, Southern California, *GSA Cordilleran Section Guidebook, Trip 15*, 268–272, 1994.
- [Rymer, M. J.: Triggered Surface Slips in the Coachella Valley Area Associated with the 1992 Joshua Tree and Landers, California, Earthquakes, *BSSA*, 90, 4, 832–848, 2000.](#)
- Sanderson, D. J. and Marchini, W. R. D.: Transpression, *J. Structural Geol.*, 6, 5, 449–458, 1984.
- Sarna-Wojcicki, A. M., Pringle, M. S. and Wijbrans, J.: New ⁴⁰Ar/³⁹Ar age of the Bishop Tuff from multiple sites and sediment rate calibration for the Matuyama-Brunhes boundary, *J. Geophys. Res.*, 105, B9, 21431–21443, 2000.
- ~~[Schlische, R. W.: Geometry and origin of fault-related folds in extensional settings, *AAPG Bull.*, 79, 1661–1678, 1995.](#)~~
- Schultz, R. A. and Balasko, C. M.: Growth of deformation bands into echelon and ladder geometries, *Geophys. Res. Lett.*, 30, 2033, 2003.
- Sheridan, J. M. and Weldon II, R. J.: Accommodation of compression in the Mecca Hills, California, *GSA Cordilleran Section Guidebook, Trip 15*, 273–279, 1994.
- Sheridan, J. M., Weldon II, R. J., Thornton, C. A. and Rymer, M. J.: Stratigraphy and Deformational History of the Mecca Hills, Southern California, *GSA Cordilleran Section Guidebook, Trip 15*, 262–268, 1994.


- [Sieh, K., Jones, L., Hauksson, E., Hudnut, K., Eberhart-Phillips, D., Heaton, T., Hough, S., Hutton, K., Kanamori, H., Lilje, A., Lindvall, S., McGill, S. F., Mori, J., Rubin, C., Spotila, J. A., Stock, J., Kie Thio, H., Treiman, J., Wernicke, B. and Zachariasen, J.: Near-Field Investigations of the Landers Earthquake Sequence, April to July 1992, *Science*, 260, 171–176, 1993.](#)
- Spinler, J. C., Bennett, R. A., Anderson, M. L., McGill, S. F., Hreinsdottir, S. and McCallister, A.: Present-day strain accumulation and slip rates associated with southern San Andreas and eastern California shear zone faults, *J. Geophys. Res.*, 115, B11407, 2010.
- Spotila, J. A., Niemi, N., Brady, R., House, M., Buscher, J and Oskin, M.: Long-term continental deformation associated with transpressive plate motion: The San Andreas fault, *Geology*, 35, 11, 967–970, 2007.
- [Stock, J. M. and Hodges, K. V.: Pre-Pliocene extension around the Guld of California and the transfer of Baja California to the Pacific plate, *Tectonics*, 8, 1, 99–115, 1989.](#)
- [Stock, J. M. and Lee, J.: Do microplates in subduction zones leave a geological record?, *Tectonics*, 13, 6, 1472–1487, 1994.](#)
- [Suppe, J. and Medwedeff, D. A.: Geometry and kinematics of fault-propagation folding, *Ecolgae geol. Helv.*, 83/3, 409–454, 1990.](#)
- Sylvester, A. G.: Strike-slip faults, *GSA Bull.*, 100, 1666–1703, 1988.
- Sylvester, A. G. and Smith, R. R.: Tectonic Transpression and Basement-Controlled Deformation in San Andreas Fault Zone, Salton Trough, California, *Treatise of Petroleum Geology*, 60, 12, 2081–2102, 1976.
- Sylvester, A. G. and Smith, R. R.: Structure section in Painted Canyon, Mecca Hills, southern California, *GSA Centennial Field Guide, Cordilleran Section*, 103–108, 1987.
- Sylvester, A. G., Bilham, R., Jackson, M. and Barrientos, S.: Aseismic Growth of Durmid Hill, Southeasternmost San Andreas Fault, California, *J. Geophys. Res.*, 98, B8, 14233–14243, 1993.
- Titus, S. J., Housen, B. and Tikoff, B.: A kinematic model for the Rinconada fault system in central California based on structural analysis of en echelon folds and paleomagnetism, *J. Struct. Geol.*, 29, 961–982, 2007.
- Thatcher, W., Savage, J. C. and Simpson, R. W.: The Eastern California Shear Zone as the northward extension of the southern San Andreas Fault, *J. Geophys. Res., Solid Earth*, 121, 2904–2914, 2016.


1281 Tyley, S. J.: Analog model study of the ground-water basin of the upper Coachella Valley,
 1282 California, U.S. Geological Survey Water-Supply, 2027, 1974.

1283 [Vollmer, F. W.: Orient 3: a new integrated software program for orientation data analysis,](#)
 1284 [kinematic analysis, spherical projections, and Schmidt plots, GSA Abstracts with](#)
 1285 [Programs, 47, 7, 49, 2015.](#)

1286 Winker, C. D. and Kidwell, S. M.: Stratigraphy of a marine rift basin: Neogene of the western
 1287 Salton Trough, California, in: Field Conference Guide, edited by: Abbott, P. L. and
 1288 Cooper, J. D., AAPG National Convention, San Diego, California, USA, 295–336,
 1289 1996.

1290 Zeeden, C., Rivera, T. A. and Storey, M.: An astronomical age for the Bishop Tuff and
 1291 concordance with radioisotopic dates, Geophys. Res. Lett., 41, 3478–3484, 2014.








Figure 11: (a) Map of the main geological features of southern California, Baja California and the Gulf of California. The location of (b) is shown as a fuchsia polygon. Modified after Janecke et al. (2018). (b) Simplified geological map of the Coachella Valley and Salton Trough, southern California, showing the three main transpressional uplift areas along the SAFZ: the Indio Hills (IH), Mecca Hills (MH), and Durmid Hills (DH). Note the link of the SAFZ with the Brawley seismic zone to the south. The study area is shown in a ~~black-green~~ rectangle. Recent earthquakes (≤ 75 years) along the Landers–Mojave Line (LML) are shown as yellow ~~dot~~stars with associated year of occurrence ~~in parenthesis~~. Faults are drawn after Rymer (2000), Guest et al. (2007), Janecke et al. (2018), and Bergh et al. (2019). Earthquakes after Nur et al. (1993a, 1993b). Abbreviations: 1947M: 1947 Manix earthquake; 1965C: 1965 Calico earthquake; 1975GL: 1975 Galway Lake earthquake; 1979HV: 1979 Homestead Valley earthquake; 1992JT: 1992 Joshua Tree earthquake; 1992L: 1992 Landers earthquake; BFBS: Banning Fault~~strand~~; BSZ: Brawley seismic zone; BP: Biskra Palms; DL: Durmid-ladder; CF: Calico fault; CRF: Camp Rock fault; DH: Durmid Hills uplift; ECSZ: East California Shear Zone; ESF: Eastern Shoreline Ffault; GF: Garlock fault; GHF: Garnet Hill fault; GOC: Gulf of California; HSGHF: Hidden Springs–Grotto Hills fault; IF: Imperial fault; IH: Indio Hills uplift; IHF: Indio Hills fault; LML: Landers–Mojave Line; MCF: Mission Creek Ffault; MF: Mill Creek fault; MH: Mecca Hills uplift; mSAF: main San Andreas fault; PCF: Painted Canyon Ffault; PMF: Pinto Mountain fault; SBF: San Bernardino fault; SCF: Skeleton Canyon Ffault; SJFZ: San Jacinto fault zone; WDCF: West Deception Canyon fault; WSDF: West Salton detachment fault. ~~Modified after Bergh et al. (2019).~~





Table 1: Summary of the timing of the main events in the Coachella Valley and Gulf of California. Note the presumed timing phases (1-3) of fold-faulting and uplift events in the Indio Hills (this work). The stratigraphy is common to the Mecca Hills and Indio Hills, although some features are only observed in one area (e.g., unconformities 1 and 2 in the Mecca Hills but not in the Indio Hills).











Figure 2: Interpreted DEM image in the southeasternmost part of the Indio Hills uplift area. Three main SAFZ-oblique macro-folds (northwestern, central, southeastern) are mapped in between the bounding Indio Hills and main San Andreas faults, whereas one SAFZ-parallel anticline is present close to the Indio Hills fault. More detailed figures are numbered and framed. Structural datasets are plotted in lower hemisphere Schmidt stereonets [via the Orient software \(Vollmer, 2015\)](#). Bedding surfaces are shown as pole to plane with frequency contour lines, with average π S great circle (red great circles), fold axial surface (blue great circles) and fold axis (red dots). Brittle fractures in sedimentary strata and basement rocks are plotted as great circles. Source: Google Earth historical imagery 09-2011. [Uninterpreted version of the image available as Supplement S1](#). © Google Earth 2011.







1342




Figure 3: Detailed structural maps showing the architecture and outline of anticline-syncline pairs, traces of bedding and strike and dip orientation, axial surface traces, and fold-related faults in (a) the northwestern, (b) central, and (c) southeastern macro-folds. Note tighter and consistently asymmetric (Z-shaped) geometries of the macro-folds to the east, whereas folds to the west are more open and symmetric. Traces and orientation of bedding show a back-limb composed of attenuated shear folds merging from the central macro-fold in the north, whereas the fore-limb is much shorter and more regularly folded. [The yellow dots show the location of field photographs.](#) See fig. 2 for legend and location. [Uninterpreted version of the images available as Supplement S2a–c.](#) © Google Earth 2011.




Figure 4: Meso-scale folds and related faults on the back-limb of the southeastern macro-fold. See location in fig. 3c. (a) Syncline in upper Palm Spring Formation units adjacent to the SAFZ-parallel macro-fold near the Indio Hills fault. (b) Close-up view of the synclinal fold hinge in (a), where a meter thick sandstone bed is slightly offset by a minor, low-angle thrust fault (red line) with NE-directed sense-of-shear. The minor thrust faults die out in the overlying sandstone bed. The mudstone bed below acts as a décollement layer with internal, plastically folded lamination, including disharmonic, intra-detachment folds. Structural orientation data of minor, centimeter-scale fold limbs in the décollement zone are plotted in a lower hemisphere Schmidt stereonet, indicating E–W-trending fold axes and a sub-horizontal axial surface (average great circle in red and fold axis as a red dot).





Figure 5: Interpreted SAFZ-parallel macro-folds ([synclines](#)) adjacent to the [main San Andreas](#) fault, which affect the southern limb of earlier (*en echelon*) macro-folded and tilted strata of the Palm Spring Formation. Note shear fold geometry in inset map with a thickened hinge zone and thinned limb to the south, ~~and a~~ steeply plunging axis, ~~and axial trace parallel to the main San Andreas fault~~. See fig. 2 for location. [Uninterpreted version of the image available as Supplement S4](#). © Google Earth 2011.






Figure 6: Interpreted satellite image of the central macro-fold showing right-lateral offset of the entire fold hinge/axial surface (upper left dashed blue line) by a NNW–SSE-trending, NE-dipping strike-slip fault (red lines). Note that the fault merges out from a layer in the southern limb of the macro-fold (black lines) and continues as a right-lateral fault. Offset geological markers include thick sandstone beds (yellow, white, light brown lines) and the fold axial surfaces of a second syncline fold farther south (lower right, dashed blue lines). Note that the syncline axial trace dies out to the southwest, and that kink bands acting as cross faults crop out in the eastern part of image (dashed pink

1384 lines). Uninterpreted version of the image available as Supplement S58. © Google Earth
1385 2011.







Figure 7: Tentative model illustrating the progressive uplift/inversion history of the Indio Hills presuming a narrow time interval between formation of all structures in the area, except for the main San Andreas fault and associated folds. (a) Early distributed transpressional strain and formation of three major, *en-echelon* oriented macro-folds, several subsidiary parasitic anticline-syncline fold pairs, and bed-parallel strike-slip and reverse (décollement) faults initiating at a high angle (c. 45°) to the Indio Hills fault. (b) Incremental partly partitioned transpression when the Indio Hills fault started to accommodate oblique-reverse movement forcing previous horizontal *en-echelon* macro-folds and parasitic folds to tighten, overturn, and rotate into steeper westward plunges. Note also sigmoidal rotation of axial traces on the back-limbs of the macro-folds to low angle (< 20–30°) with the Indio Hills fault. (c) Late-stage advanced

1399 strain partitioning with dominant shortening component on the oblique-reverse Indio
1400 Hills fault, and right-lateral slip on the main San Andreas~~Banning~~ Fault. Notice the
1401 formation of the anticline parallel to the Indio Hills fault, subsidiary fold-internal strike-
1402 slip faults, and conjugate cross faults and kink bands that overprinted the macro-folds.
1403 Legend as in fig. 2.







Figure 8: Kinematic evolution, timing, and along-strike correlation of the Indio Hills, Mecca Hills, and Durmid Hills uplift domains and bounding master faults in the Coachella valley, southern California. We present a progressive kinematic evolution from (a) distributed, through (b) partly partitioned, to (c) advanced partitioned strain events. See text for further explanation. Black lines are faults (full or stippled). Blue lines are fold axial traces. Wide arrows indicate main shortening direction, half-arrows lateral (strike-slip) shearing. Abbreviations: BSF: Banning fault strand; ESF: Eastern Shoreline fault; GHF: Garnet Hills fault; HSGHF: Hidden Springs–Grotto Hills fault; IHF: Indio Hills fault; mSAF: main San Andreas fault in Durmid Hills; MCF: Mission Creek fault; PCF: Painted Canyon fault; ~~SAFZ: San Andreas transform fault~~; SCF: Skeleton Canyon fault.

Supplements




S1: Uninterpreted Fig. 2. See Fig. 1 for location. © Google Earth 2011.






S2: Uninterpreted Fig. 3a–c. See Fig. 2 for location. © Google Earth 2011.




S31: Examples of subsidiary fold styles in the northwestern macro-fold. For location, see fig. 3a. (a) DEM image showing an upright and west-plunging anticline-syncline pair. © Google Earth 2011. (b) Symmetric and concentric anticline, same as in (a) viewed to the

east. (c) Kink-style syncline changed along strike southeastward from symmetric in (a). (d) Tight to isoclinal, steeply west-plunging anticline in the northern part of the macro-fold. Note folded quartz-rich sandstone layer used as stratigraphic marker in the upper Palm Spring Formation. (e) Same quartz-sandstone layer as in (d) repeated by tight/isoclinal folding.




S42: Examples of macro- and meso-scale fold styles in the central macro-fold (location in fig. 3b). (a) Open to slightly asymmetric syncline fold hinge plunging moderately west. (b) Outcrop of the fold hinge of the central macro-fold. The hinge zone is relatively tight, and the fold partly overturned to the SW. Note how the mudstone bed (white lines) thickens into


the fold hinge. Dashed yellow line represents the fold axial surface. (c) Panorama view of the central macro-fold, showing change in geometry and tightness of subsidiary anticline-syncline pairs toward northeast. Note presence of the major SAFZ-parallel, open anticline in the northeast, and the location of Indio Hills fault. (d) DEM image of the same outline as in (c). © Google Earth 2011.



S53: (a) Cliff view of a reverse/thrust fault system (black lines) in upper Palm Spring Formation strata that truncates and offset bedding surfaces (white stippled lines). Note the presence of fault-related drag folds that reveal top-NE (right) sense of shear. See fig. 3c for location. (b) Minor reverse fault in SW-dipping sandstone bed. Note fault movement top-SW. Location is shown in fig. 3c. (c) Outcrop photograph viewed in section on NNE-dipping sandstone beds, comprising E-W-trending, north-verging minor, asymmetric folds and faults. Note that the low-angle minor faults (dashed red lines) formed within the minor fold hinges. Location is shown in fig. 3c. (d) Field photograph on NNE-dipping sandstone layers showing a conjugate set of minor reverse faults (dashed red lines) that offset normally, thin sandstone beds (green, blue and yellow). Location is shown in fig. 3c.



S6: Uninterpreted Fig. 5. See Fig. 2 for location. © Google Earth 2011.




S74: (a) Outcrop photograph showing right-lateral offset (c. 70 m) of the central macro-fold axial surface trace (yellow stippled line) by a steep, NNW-SSE trending, NE-dipping brittle

fault (red line). Note partly overturned bedding (white line) in the hinge zone to the west. See fig. 3b for location. (b) Sketch of the steep right-lateral strike-slip fault that decapitates the entire hinge of the central macro-fold. Note also subsidiary NW–SE and NNE–SSW trending, NE-dipping, right- and left-lateral faults, respectively that offsets the macro-fold limbs. See fig. 3b for location. (c) Outcrop photograph of a meter-thick sandstone bed crosscut and offset by minor steep, NW–SE to NNW–SSE trending right-lateral strike-slip fault. See fig. 3b for location. (d) Outcrop in vertical view showing a minor conjugate fault set defined by N-S and NNE–SSE trending right- and left-lateral strike-slip faults. Locality shown in fig. 3c. (e) Satellite image illustrating large scale kink bands arranged as cross faults at high angle to bedding on the southeastern macro-fold. See fig. 2 for location. © Google Earth 2011.



S85: Uninterpreted Fig. 6. See Fig. 2 for location. ~~From Google Earth.~~ © Google Earth 2011.



S96: Field photographs of centimeter-scale faults in the Indio Hills. The photographs were rotated anticlockwise by 52° to better analyze the microstructures they display. Low-angle, bending normal faults in (a) and (b) become similar to micro thrust-faults and planar reverse faults in (c) resemble extensional, graben-bounding fault. Notice the potential thickened wedges of syn-tectonic sediments in the extensional micro-graben in (c) (cf. green, blue and yellow lines). Location is shown in fig. 3c.